January 2023 Volume 30 Number 0

Editors:

Kaye Wilson, & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz
@Pharmaceutical Management Agency
ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written from and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	•	
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	36
	Cardiovascular System	46
	Dermatologicals	60
	Genito Urinary System	70
	Hormone Preparations – Systemic	77
	Infections – Agents For Systemic Use	89
	Musculoskeletal System	109
	Nervous System	120
	Oncology Agents & Immunosuppressants	147
	Respiratory System & Allergies	234
	Sensory Organs	243
	Various	248
Section C	Extemporaneous Compounds (ECPs)	250
Section D	Special Foods	253
Section I	National Immunisation Schedule	274

Introducing Pharmac

Index

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION BY ALIMENTARY TRACT AND METAROLISM

	SECTION B: ALIMENTARY TRACT AND ME	Subsidy (Manufacturer's Price)		Fully Subsidised	
		(Manufacturer's Frice)	Per	Jubsidised	
Α	ntacids and Antiflatulents				
Α	ntacids and Reflux Barrier Agents				
AL	GINIC ACID				
	Sodium alginate 225 mg and magnesium alginate 87.5 mg posachet		30	✓	Gaviscon Infant
SO	DIUM ALGINATE				
*	Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
*	Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m	nl	Acidex
P	hosphate Binding Agents				
	JMINIUM HYDROXIDE				
	Tab 600 mg	12.56	100	•	Alu-Tab
CA	LCIUM CARBONATE				
	Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) — Subsidy by endorsement Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed according	cium carbonate table	500 m		Roxane ium carbonate tablets are
٨	ntidiarrhoeals				
	midial mocals				
A	gents Which Reduce Motility				
*	PERAMIDE HYDROCHLORIDE – Up to 30 cap available on a Tab 2 mg	10.75	400 400	_	Nodia Diamide Relief

LOPERAMIDE HYDROCHLORIDE - Up to 30 c	ap available on a PSO		
* Tab 2 mg	10.75	400	✓ Nodia
* Cap 2 mg	7.25	400	✓ Diamide Relief

Rectal and Colonic Anti-inflammatories

BUDESONIDE

Cap 3 mg - Special Authority see SA1886 below - Retail 90 ✓ Entocort CIR pharmacy......166.50

⇒SA1886 Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP 21.1 g OP	✓ Cortifoam S29✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam §29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
OLSALAZINE Tab 500 mg	56.02	60	•	Atnahs Olsalazine \$29
Cap 250 mg	93.37 53.00	100 100	_	Dipentum Dipentum
PREDNISOLONE SODIUM Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential Prednisolone 629
SODIUM CROMOGLICATE Cap 100 mg	92.91	100		Nalcrom Ralicrom
(Nalcrom Cap 100 mg to be delisted 1 April 2023) SULFASALAZINE				
* Tab EC 500 mg	14.00 15.53	100 100		Salazopyrin Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g11.06	30 g OP	✓ Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg7.30	12	✓ Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on PSO		10	✓ Max Health
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO	6.35	5	✓ Buscopan
MEREVERINE HYDROCHI ORIDE			
* Tab 135 mg	9.20	90	✓ Colofac
HYOSCINE BUTYLBROMIDE * Tab 10 mg* * Inj 20 mg, 1 ml – Up to 5 inj available on a PSO MEBEVERINE HYDROCHLORIDE	6.35 6.35	100 5	✓ Busc ✓ Busc

	ALIMENTARY	/ TRAC	T AND	METABOLISM
	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Antiulcerants				
Antisecretory and Cytoprotective				
MISOPROSTOL * Tab 200 mcg - Up to 120 tab available on a PSO	41.50	120	✓ 0	cytotec
Helicobacter Pylori Eradication				
CLARITHROMYCIN Tab 500 mg — Subsidy by endorsement	ori eradication and presc		endorsed	
H2 Antagonists				
FAMOTIDINE - Only on a prescription * Tab 20 mg	4.91	100	√ F	amotidine Hovid §29
* Tab 40 mg	8.48	100	√ F	amotidine Hovid \$29
* Inj 10 mg per ml, 4 ml - Subsidy by endorsement Subsidy by endorsement - Subsidised for patients re		10 t of pallia		lylan S29
Proton Pump Inhibitors				
LANSOPRAZOLE * Cap 15 mg * Cap 30 mg OMEPRAZOLE For omeprazole suspension refer Standard Formulae, pa	5.26	100 100		anzol Relief anzol Relief
* Cap 10 mg		90	√ <u>C</u>	Omeprazole actavis 10
* Cap 20 mg	1.86	90	√ <u>C</u>	Omeprazole actavis 20
* Cap 40 mg	3.11	90	√ <u>C</u>	meprazole actavis 40
Powder – Only in combination Only in extemporaneously compounded omeprazole		5 g	✓ N	lidwest
* Inj 40 mg ampoule with diluent		5	✓ [<u>Pr Reddy's</u> Omeprazole
PANTOPRAZOLE * Tab EC 20 mg* * Tab EC 40 mg		100 100		Panzop Relief
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mg	14.51	50	√ 0	Gastrodenol S29

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
SUCRALFATE Tab 1 g	35.50 (48.28)	120		Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail Tab 550 mg SA1461 Special Authority for Subsidy Initial application only from a gastroenterologist, hepatological application only from a gastroenterologist.	625.00		mendation	
hepatologist. Approvals valid for 6 months where the patiel tolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Prahepatologist. Approvals valid without further renewal unles benefiting from treatment.	actitioner on the recomme	endatio	n of a gastr	oenterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE - Special Authority see SA1320 below - Reta Cap 25 mg Cap 100 mg Oral liq 50 mg per ml	110.00	100 100 30 ml 0	✓ OP ✓	Proglicem \$29 Proglicem \$29 Proglycem \$29 e5 Pharma \$29
➤ SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approval hypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid with appropriate and the patient is benefiting from treatment.			d for the tre	eatment of confirmed
GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	✓	Glucagen Hypokit
Insulin - Short-acting Preparations				
INSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml (Actrapid Humulin B
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓.	Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
INSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	•	NovoMix 30 FlexPen
INSULIN ISOPHANE ▲ Inj human 100 u per ml	17.68	10 ml (Humulin NPH Protaphane

▲ Inj human 100 u per ml, 3 ml......29.86

5

✓ Humulin NPH

✓ Protaphane Penfill

	Cubaidu		Fully Drand or
	Subsidy (Manufacturer's F	Price) Subs	Fully Brand or idised Generic
	\$	Per	✓ Manufacturer
ISULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
		_	✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
			✓ PenMix 30✓ PenMix 50
			▼ Penwix 50
ISULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml, 3 ml		5	✓ Humalag Miy 25
		5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml, 3 ml		5	✓ Humalog Mix 50
O III			- Hamaiog inix oo
Insulin - Long-acting Preparations			
SULIN GLARGINE			•
Inj 100 u per ml, 10 ml		1	✓ Lantus
Inj 100 u per ml, 3 ml		5 5	✓ Lantus✓ Lantus SoloStar
IIIJ 100 u per IIII, 3 IIII disposable peri	94.50	5	V Lantus SoloStar
nsulin - Rapid Acting Preparations			
SULIN ASPART			
Inj 100 u per ml, 10 ml		1	✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill ✓ NovoRapid FlorBox
Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
ISULIN GLULISINE			4.4.1
Inj 100 u per ml, 10 ml		1	✓ Apidra
Inj 100 u per ml, 3 ml		5 5	✓ Apidra✓ Apidra SoloStar
ISULIN LISPRO		3	- Apidia odiootai
isolin liseno . Inj 100 u per ml, 10 ml	34 92	10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml		5	✓ Humalog
Alpha Glucosidase Inhibitors			J
CARBOSE			
: Tab 50 mg	8.95	90	✓ Accarb
Tab 100 mg		90	✓ Accarb
Oral Hypoglycaemic Agents			
LIBENCLAMIDE			
: Tab 5 mg	7.50	100	✓ Daonil
LICLAZIDE			
: Tab 80 mg	15.18	500	✓ Glizide
LIPIZIDE			_
: Tab 5 mg	4.58	100	✓ Minidiab
ETFORMIN HYDROCHLORIDE			
Tab immediate-release 500 mg	14.74	1,000	✓ Metformin Mylan
		.,500	✓ Metformin Viatris
Tab immediate-release 850 mg	11.28	500	Metformin Mylan
•			-

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
	\$	Per	•	Manufacturer
PIOGLITAZONE				
* Tab 15 mg	6.80	90	✓ V	/exazone
* Tab 30 mg	7.30	90	✓ V	/exazone
* Tab 45 mg		90	✓ V	/exazone
VILDAGLIPTIN				
Tab 50 mg	35.00	60	√ (alvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	√ (Salvumet
Tab 50 mg with 850 mg metformin hydrochloride		60	√ (Salvumet

GLP-1 Agonists

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Maori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*: and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

DULAGLUTIDE - Special Authority see SA2065 above - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*: or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 10 mg58.56	30	Jardiance
	Tab 25 mg58.56	00	Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

✓ CareSens Dual 1 OP

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes; or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

1 OP ✓ CareSens N ✓ CareSens N POP 20.00 ✓ CareSens N Premier

Note: Only 1 meter available per PSO

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips	26.20	50 test OP	SensoCard
---------------------------	-------	------------	-----------------------------

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

*	29 g × 12.7 mm10.9	5 100	✓ B-D Micro-Fine
	31 g × 5 mm12.2		✓ B-D Micro-Fine
	31 g × 6 mm9.5		✓ Berpu
	31 g × 8 mm10.9		✓ B-D Micro-Fine
	32 g × 4 mm10.9		✓ B-D Micro-Fine

15

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	
_		\$	Per		Manufacturer
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	- Maximum of 200	dev p	oer prescrip	otion
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.36	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.36	10		
		(1.99)			B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — **(severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

1 OP

1 OP

✓ TruSteel

✓ TruSteel

	ALIMENTALLI	1117	OI AILD	METABOLISM
	Subsidy (Manufacturer's Price)	S Per	Fully ubsidised	Brand or Generic Manufacturer
continued than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mr 3 The patient has not had an increase in severe unexplaine				ne; and
 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within the 	ir vocational scope.			
INSULIN PUMP CARTRIDGE - Special Authority see SA1985	on page 19 – Retail ph	armac	y	
 a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded pe 	r year.			
Cartridge 300 U, t:lock × 10		1 OP	✓ Ta	andem Cartridge
 INSULIN PUMP INFUSION SET (STEEL CANNULA) – Special a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 	Authority see SA1985	on pa	ge 19 – Re	tail pharmacy
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-866A
8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-874A
8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-876A
6 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock	130.00	1 OP	√ S	ure-T MMT-863
8 mm steel needle; 29 G; manual insertion; 60 cm tubing \times 10 with 10 needles; luer lock		1 OP	_	ure-T MMT-873
INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGH' Retail pharmacy a) Maximum of 3 sets per prescription	Г INSERTION) – Spe	cial Au	thority see	SA1985 on page 19 –
b) Only on a prescriptionc) Maximum of 13 infusion sets will be funded per year.				
6 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles	130.00	1 OP	✓ T	ruSteel
8 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	130.00	1 OP	✓ Ti	ruSteel
6 mm steel cannula; straight insertion; 60 cm line × 10 with				

10 needles......130.00

8 mm steel cannula; straight insertion; 60 cm line × 10 with

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing x 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing \times 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing \times 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-386A

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION WITH IN	SERT	ION DEVIC	E) - Special Authority see
SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c				
line × 10 with 10 needles		1 OP		AutoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cm			_	
line × 10 with 10 needles		1 OP		AutoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION) - Speci	al Aut	thority see S	SA1985 on page 19 -
Retail pharmacy				
a) Maximum of 3 sets per prescription Only on a prescription				
b) Only on a prescriptionc) Maximum of 13 infusion sets will be funded per year.				
17 mm teflon cannula; angle insertion; 60 cm line × 10 with				
10 needles; luer lock	130.00	1 OP	1	Silhouette MMT-373
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH		INS	ERTION DE	FVICE) - Special Authority
see SA1985 on page 19 – Retail pharmacy	T INCENTION WITH	1 11 401		evice) openial realionty
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; insertion device;			_	
110 cm line × 10 with 10 needles		1 OP	•	AutoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 cr		1 00		AutoSoft 90
line x 10 with 10 needles9 mm teflon cannula; straight insertion; insertion device;	140.00	1 OP	• /	AU105011 90
110 cm line × 10 with 10 needles	140.00	1 OP	•	AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 cr		. 0.		-utocont 50
line × 10 with 10 needles		1 OP	1	AutoSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH	T INSERTION) - S	necial	Authority s	see SA1985 on page 19 -
Retail pharmacy	1 11021111011, 0	poolai	riamonty o	oo on nood on page 10
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with	1	4.00		
10 needles; luer lock		1 OP	✔ (Quick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles; luer lock		1 OP		Quick-Set MMT-392
•				Juick-Set MiM1-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 or	1 page 19 – Retail pr	narma	ıcy	
a) Maximum of 3 sets per prescriptionb) Only on a prescription				
c) Maximum of 13 packs of reservoir sets will be funded per	vear			
10 × luer lock conversion cartridges 1.8 ml for Paradigm pum		1 OP	1	ADR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 10		1 OP		MiniMed
				1.8 Reservoir
				MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP	✓ I	MiniMed
				3.0 Reservoir
				MMT-332A

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Digestives Including Enzymes

_			. ~	_	_		_	_	_		_		
Ρ	Δ	Λ	Ю	ж	н	Δ	ш	(:	ы	N	'	ΥN	ЛF

1 / WOTE/WIO ENZIME			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))	94.40	100	✓ Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)	34.93	20 g OP	✓ Creon Micro
(Panzytrat Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amy	lase, 1,250 U p	rotease)) to	be delisted 1 June 2023)
URSODEOXYCHOLIC ACID – Special Authority see SA1739 below		nacy 100	✓ <u>Ursosan</u>

⇒SA1739 Special Authority for Subsidy

Initial application — (Alaquille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Fither:
 - 1 Patient has been diagnosed with Alagille syndrome; or
 - 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults: and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- Both:
 - 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
 - 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN): and
- 2 Liver function has not improved with modifying the TPN composition.

	Subsidy	Fully		Brand or
(Man	ufacturer's Price)	Subsidi	sed	Generic
	\$	Per	•	Manufacturer

continued...

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Laxatives

Bulk-forming Agents

* Powder for oral soin	5.00	250 g OP	Psyllium Husk
	12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS			
* Dry	6.02	500 g OP	
	(17.32)	_	Normacol Plus
	2.41	200 g OP	
	(8.72)	Ü	Normacol Plus

Faecal Softeners

* Tab 50 mg 2.3 * Tab 120 mg 3.1			✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u>
DOCUSATE SODIUM WITH SENNOSIDES Tab 50 mg with sennosides 8 mg	50 :	200	✓ <u>Laxsol</u>
POLOXAMER — Only on a prescription Not funded for use in the ear.			
* Oral drops 10%	98 30	ml OP	✓ Coloxyl

Opioid Receptor Antagonists - Peripheral

DOCUSATE SODIUM - Only on a prescription

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below - Retail p	harmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy		Fully Brand or
	(Manufacturer's Price)	Per Subsi	idised Generic Manufacturer
Osmotic Laxatives			
GLYCEROL			
* Suppos 3.6 g – Only on a prescription		20	✓ PSM
* Suppos 4 g - Only on a prescription	10.39	20	✓ Lax-suppositories Glycerol
(PSM Suppos 3.6 g to be delisted 1 February 2023)			diyeeror
LACTULOSE – Only on a prescription			
* Oral liq 10 g per 15 ml	3.61	500 ml	✓ Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BI	CARBONATE AND	SODIUM CI	HLORIDE
Powder for oral soln 13.125 g with potassium chloride 46.6 r	ng,		
sodium bicarbonate 178.5 mg and sodium chloride 350.	7 mg6.70	30	✓ <u>Molaxole</u>
SODIUM ACID PHOSPHATE - Only on a prescription			
Enema 16% with sodium phosphate 8%	2.50	1	✓ Fleet Phosphate
CODULINA OLT DATE MUTILI CODULINA LALIDVIL OLU DI IOACETATE	- 0.1	to at a co	Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml	, ,	iption	
5 ml	,	50	✓ Micolette
			✓ Micolette-S29 S29
Stimulant Laxatives			
BISACODYL - Only on a prescription			
* Tab 5 mg		200	✓ Bisacodyl Viatris
* Suppos 10 mg	3.69	10	✓ <u>Lax-Suppositories</u>
SENNA – Only on a prescription	0.47	400	
* Tab, standardised	(8.21)	100	Senokot
	0.43	20	Seriokot
	(2.06)		Senokot
SODIUM PICOSULFATE - Special Authority see SA2053 below	– Retail pharmacy		
Oral soln 7.5 mg per ml		0 ml OP	Dulcolax SP Drop
⇒SA2053 Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals valid	d for 6 months for ap	plications n	neeting the following criteria:
Both:	o on adoquato trial a	f other oral	nharmanatharanian ingledian
 The patient is a child with problematic constipation despit macrogol where practicable; and 	e an auequate (fiai 0	i other oral	pharmacomerapies including
The matient would although a married a bight wall was because			at a disease of a se

Metabolic Disorder Agents

is benefiting from treatment.

2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ADCININE	 Special Authority see 	CA2042 bolow	Potail pharmacy
ABUILINE	- Special Aumoniv See	SAZU4Z DEIOW -	- Belali Dharmacv

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg	CBS	50	✓ Solgar
Powder		400 g	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE - Special Authority see SA19	87 on the next page – Retail pharmacy	
_ , , , ,		

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5.10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - S	pecial Authority see SA2039 below – Retail pharmacy		
Cap 120 mg	CBS	30	✓ Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
 \$	Per	✓	Manufacturer	

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assav in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE – Special Authority see SA2040 below	 Retail pharmacy 		
Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral liq 1 g per 10 ml	CBS	118 ml	✓ Carnitor S29
Oral liq 500 mg per 10 ml		300 ml	✓ Balance

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN - Special Authority see SA2041 on the next p	age - Retail pharmacy		
Tab 100 mg	CBS	100	✓ Country Life
Cap 100 mg	CBS	100	✓ Solgar

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE - Special Authority see SA1989 below - Retail pharmacy

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy
Soln 100 mg per mlCBS

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 on the next page - Retail pharmacy

Grans 483 mg per g......2,016.00 174 g OP ✓ Pheburane

100 ml

✓ Amzoate S29

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	 Life Extension

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA2137 below - Retail pharmacy		
Inj 200 unit vial1,072.00	1	✓ Elelyso

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and

	Subsidy (Manufacturer's Price) \$	Sub: Per	Fully sidised	Brand or Generic Manufacturer
continued				
 3 Radiological (MRI) signs of bone activity performed at tw demonstrate no deterioration shown by the MRI, compar or adjusted dose; and 4 Patient has not developed another medical condition tha ERT; and 5 Patient is adherent with regular treatment and taliglucera every other week rounded to the nearest whole vial (200 	red with MRI taken imn at might reasonably be ase alfa is to be admini	nediately expected	orior to o	commencement of therapy promise a response to
Mouth and Throat				
Agents Used in Mouth Ulceration				
BENZYDAMINE HYDROCHLORIDE Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with		500 ml		

Endorsement	9.00	500 ml	
	(20.31)		Difflam
Additional subsidy by endorsement for a patient wh prescription is endorsed accordingly.	o has oral mucositis a	s a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	· ·	Orabase
	`1.52 [′]	5 g OP	
	(3.60)	Ü	Orabase
Powder	, ,	28 g OP	
	(10.95)	3 -	Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE	` ,		
* Adhesive gel 8.7% with cetalkonium chloride 0.01%		15 g OP	
* Adilesive gel 6.7 /6 with Cetalkorlium Chloride 0.01 /6	(6.00)	13 g OF	Bonjela
	(0.00)		Donjela
TRIAMCINOLONE ACETONIDE			_
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
		20	· rungiiii
MICONAZOLE			
Oral gel 20 mg per g	4.74	40 g OP	✓ <u>Decozol</u>
NYSTATIN			

Vit	am	Πî	e

Vitamin B

HY	DROXOCOBALAMIN		
*	Inj 1 mg per ml, 1 ml ampoule - Up to 6 inj available on a PSO2.46	3	✓ <u>Hydroxocobalamin</u> Panpharma

Oral liq 100,000 u per ml1.76

24 ml OP

✓ Nilstat

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	
PYRIDOXINE HYDROCHLORIDE	*			
a) No more than 100 mg per dose				
b) Only on a prescription				
* Tab 25 mg - No patient co-payment payable		90		Vitamin B6 25
Tab 50 mg	23.45	500	/	Pyridoxine
				multichem
THIAMINE HYDROCHLORIDE – Only on a prescription	4.65	100		Thiamine multichem
* Tab 50 mg	7.09	100		Max Health
(Max Health Tab 50 mg to be delisted 1 April 2023)	7.00		•	max ricuiti
VITAMIN B COMPLEX				
* Tab, strong, BPC	7.15	500	1	Bplex
				•
Vitamin C				
ASCORBIC ACID				
a) No more than 100 mg per dose				
b) Only on a prescription	40.50	500	,	0.4.
* Tab 100 mg	12.50	500	•	Cvite
Vitamin D				
ALFACALCIDOL				
* Cap 0.25 mcg	26.32	100		One-Alpha
* Cap 1 mcg	87.98	100		One-Alpha
Nr. Ovel draws O seem and	00.00	00 1 0		One-Alpha S29 S29
* Oral drops 2 mcg per ml	00.08	20 ml O	Ρ •	One-Alpha
CALCITRIOL	7.00	100	./	Coloitrial AET
* Cap 0.25 mcg * Cap 0.5 mcg		100 100		Calcitriol-AFT Calcitriol-AFT
COLECALCIFEROL		100		<u>odiolitioi Ai i</u>
* Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescript	ion2.95	12	1	Vit.D3
* Oral liq 188 mcg per ml (7,500 iu per ml)		4.8 ml C		Puria
Multivitamin Preparations				
MULTIVITAMIN RENAL - Special Authority see SA1546 below	Potail pharmaou			
* Cap	, ,	30	1	Clinicians Renal Vit
⇒SA1546 Special Authority for Subsidy		50	-	
Initial application from any relevant practitioner. Approvals vali	d without further rer	newal ur	nless notif	ied for applications meeting
the following criteria:				11
Either:				
1 The patient has chronic kidney disease and is receiving e				
2 The patient has chronic kidney disease grade 5, defined a	as patient with an es	stimated	glomerul	ar tiltration rate of <
15 ml/min/1.73 m ² body surface area (BSA).				

200 g OP

✓ Paediatric Seravit

\$ Per Manufacturer		Subsidy (Manufacturer's Price) \$	Fu Subsidise Per	,	
---------------------	--	---	------------------------	---	--

⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINS

- ✓ Mvite 1.000
- * Cap (fat soluble vitamins A, D, E, K) Special Authority see 60 ✓ Vitabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

Minerals

Calcium

CAI	_CIUM CARBONATE		
	Tab 1.25 g (500 mg elemental)	250	✓ Calci-Tab 500
	Tab eff 1.25 g (500 mg elemental) – Subsidy by endorsement260.00	100	✓ Calcium 500 mg
			Hexal S29
	Subsidy by endorsement – Only when prescribed for paediatric patients (<	5 years) wh	nere calcium carbonate oral liquid is

$\triangle VI$	IN A	α	LICONATE	

	considered unsultable.				
CA	LCIUM GLUCONATE				
*	Inj 10%, 10 ml ampoule	32.00	10	✓ Max Health -	

64.00 20 ✓ Max Health S29

Hameln \$29

Fluoride

SODIUM FLUORIDE		
* Tab 1.1 mg (0.5 mg elemental)	100	PSM
(PSM Tab 1.1 mg (0.5 mg elemental) to be delisted 1 March 2023)		

lodine

POTASSILIM IODATE

10	I AGGIUW IODA I L			
*	Tab 253 mcg (150 mcg elemental iodine)4.58	90	✓ NeuroTabs

Iron

FERROUS FUMARATE			
* Tab 200 mg (65 mg elemental)	3.04	100	✓ Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID			

✓ Ferro-F-Tabs 100

	Subsidy (Manufacturer's Price) \$	Sub:	Fully sidised	Brand or Generic Manufacturer
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental) * Oral liq 30 mg (6 mg elemental) per 1 ml		30 500 ml		<u>Ferrograd</u> <u>Ferodan</u>
IRON (AS FERRIC CARBOXYMALTOSE) — Special Authority se Inj 50 mg per ml, 10 ml vial		Retail phar 1	•	Ferinject

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE * Inj 50 mg per ml, 2 ml ampoule34.50	5	✓ Ferrosig
Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%33.60	355 ml	✓ Phillips Milk of Magnesia \$29
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule25.53	10	✓ <u>Martindale</u>
Zinc		
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)11.00	100	✓ Zincaps

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

BLOOD AND BLOOD FORMING ORGANS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA1775 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe	100.00	6	Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	✓ Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	✓ Binocrit
Inj 6,000 iu in 0.6 ml, syringe		6	✓ Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	✓ Binocrit
Inj 10,000 iu in 1 ml, syringe		6	✓ Binocrit
Inj 40,000 iu in 1 ml, syringe		1	✓ Binocrit

	Subsidy (Manufacturer's Price) \$		ully sed	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID Tab 0.8 mg	26.60	1,000		olic Acid multichem
* Tab 5 mg Oral liq 50 mcg per ml		100 5 ml OP		olic Acid Mylan iomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

reaters Group in conjunction with the National F	iaemopnilia ivianagement grou	ıр.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial		1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 be Wastage claimable	elow – Retail pharmacy		
Tab 25 mg	1,550.00	28	Revolade
Tab 50 mg	3,100.00	28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy. Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	1	Manufacturer

continued...

and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

Inj 30 mg in 1 ml vial	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	7,138.00	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial		1	✓ Hemlibra
Inj 150 mg in 1 ml vial	17,846.00	1	✓ Hemlibra

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Either:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	✓	

continued...

- 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
 - 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	·	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria

Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	✓ Xyntha
Inj 1,000 iu prefilled syringe		1	✓ Xyntha
Inj 2,000 iu prefilled syringe		1	✓ Xyntha
Inj 3,000 iu prefilled syringe		1	Xyntha

NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group

man are manerial macricepinna management on	, ap.		
Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial	870.00	1	✓ RIXUBIS
Inj 2,000 iu vial		1	✓ RIXUBIS
Ini 3.000 iu vial	,	1	✓ RIXUBIS

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [Xpharm]

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

managed by the naemophilia Treaters Group in conjunction	n with the national	паетнорпіна	Management G
Inj 250 iu vial	210.00	1	✓ Advate
Inj 500 iu vial	420.00	1	Advate
Inj 1,000 iu vial	840.00	1	Advate
Inj 1,500 iu vial	1,260.00	1	✓ Advate
Inj 2,000 iu vial	1,680.00	1	Advate
Inj 3,000 iu vial	2,520.00	1	Advate

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Sı Per	ubsidised •	Generic Manufacturer
OTOGOG ALEA IDECOMPINIANT FACTOR VIIII (KOCENIATE	<u> </u>	1 01		Widifulactarci
CTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE For patients with haemophilia. Rare Clinical Circumstances		a racom	hinant fa	ictor VIII Access to funda
treatment is managed by the Haemophilia Treaters Group in subject to criteria.				
Inj 250 iu vial	237.50	1	✓	Kogenate FS
Inj 500 iu vial	475.00	1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial	,	1		Kogenate FS
Inj 3,000 iu vial	2,850.00	1	•	Kogenate FS
URIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]				
For patients with haemophilia A receiving prophylaxis treatment		d treatn	nent is m	anaged by the Haemoph
Treaters Group in conjunction with the National Haemophilia				
Inj 250 iu vial		1		Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial		1		Adynovate
Inj 2,000 iu vial	2,400.00	1	•	Adynovate
ODIUM TETRADECYL SULPHATE				
5 Inj 3% 2 ml		5		
	(73.00)			Fibro-vein
RANEXAMIC ACID				
Tab 500 mg	10.45	60	✓	Mercury Pharma
Vitamin K				
HYTOMENADIONE				
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO	8.00	5	✓	Konakion MM
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO	9.21	5	✓	Konakion MM
Antithrombotic Agents				
Antiplatelet Agents				
SPIRIN • Tab 100 mg	14.95	990	✓	Ethics Aspirin EC
LOPIDOGREL				·
Tab 75 mg	4.60	84	1	Clopidogrel
- az - 5 - 19		٠.		Multichem
	5.07		1	Arrow - Clopid
Clopidogrel Multichem Tab 75 mg to be delisted 1 May 2023)				
PYRIDAMOLE				
Tab long-acting 150 mg	10.90	60	✓ 1	Pytazen SR
ICAGRELOR - Special Authority see SA1955 below - Retail p		50	- '	,
Tab 90 mg	,	56	.	Ticagrelor Sandoz
i au au iliy	90.00	50		Brilinta
Brilinta Tab 90 mg to be delisted 1 March 2023)	30.00		• 1	Dimina
 Special Authority for Subsidy itial application — (acute coronary syndrome) from any rel 				

continued...

meeting the following criteria:

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic
 \$	Per 🗸	Manufacturer

continued...

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA2152 be	elow - Retail pharmacy		
Inj 20 mg in 0.2 ml syringe	31.28	10	Clexane
Inj 40 mg in 0.4 ml syringe	42.49	10	Clexane
Inj 60 mg in 0.6 ml syringe	60.67	10	Clexane
Inj 80 mg in 0.8 ml syringe	80.89	10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe	125.87	10	 Clexane Forte
Inj 150 mg in 1 ml syringe	143.86	10	Clexane Forte

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HEPARIN SODIUM				
Inj 1,000 iu per ml, 5 ml ampoule	72.84	50	√ P	fizer
Inj 5,000 iu per ml, 1 ml	32.66	5	✓ D	BL Heparin
				Sodium S29
	70.33		✓ H	ospira
Inj 5,000 ju per ml, 5 ml ampoule	289.05	50	✓ P	•
Inj 25,000 iu per ml, 0.2 ml		5	✓ H	ospira
, ., , ,	42.40			eparin DBL S29
	482.20	50		eparin DBL S29
HEDADINIOED CALINE	402.20	50	• 11	cpariii DDL 22
HEPARINISED SALINE	05.40			
Inj 10 iu per ml, 5 ml	65.48	50	✓ P	fizer
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	76.36	60	✓ P	radaxa
Cap 110 mg	76.36	60	✓ P	radaxa
Cap 150 mg	76.36	60	✓ P	radaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83 10	30	√ X	arelto
Tab 15 mg - Up to 14 tab available on a PSO		28		arelto
Tab 20 mg		28		arelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	2.46	50	1 C	oumadin
* Tab Tilly	6.46	100	•	arevan
* Tab 2 mg	****	50		oumadin
* Tab 2 mg		100	-	arevan
* Tab 5 mg		50		oumadin
* Tab Jing	11.48	100	-	arevan
	11.40	100	→ IV	arcvari

FILGRASTIM - Special Authority see SA1259 below - Retail pl	harmacy		
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓ <u>Nivestim</u>
Inj 480 mcg per 0.5 ml prefilled syringe	148.58	10	✓ Nivestim

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×109/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic r ✓ Manufacturer
PEGFILGRASTIM - Special Authority see SA1912 below - Reta	il pharmacy		
Inj 6 mg per 0.6 ml syringe	65.00	1	✓ Ziextenzo
	1,080.00		✓ Neulastim

(Neulastim Inj 6 mg per 0.6 ml syringe to be delisted 1 June 2023)

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

GLUCOSE [DEXTROSE]			
* Inj 50%, 10 ml ampoule - Up to 5 inj available on a PSO30	0.65	5	Biomed
* Inj 50%, 90 ml bottle - Up to 5 inj available on a PSO	5.00	1 🗸	Biomed
POTASSIUM CHLORIDE			
* Inj 75 mg per ml, 10 ml65	5.00 5	50	Juno
SODIUM BICARBONATE			
Inj 8.4%, 50 ml21	1.40	1 🗸	Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			
Inj 8.4%, 100 ml21	1.95	1 🗸	Biomed
a) Un to Fini available on a DCO			

a) Up to 5 inj available on a PSOb) Not in combination

SODIUM CHI ORIDE

Not funded for use as a nasal drop. Not funded for nebuliser use except when used in conjunction with an antibiotic intended for nebuliser use.

	1.36	1,000 ml	✓ Baxter	
Only if prescribed on a prescription for renal dialysis, maternity	or post-nata	I care in the h	home of the patient, or on a PSO	
for emergency use. (500 ml and 1,000 ml packs)				
Inj 23.4% (4 mmol/ml), 20 ml ampoule	.35.50	5	✓ Biomed	

For Sodium chloride oral liquid formulation refer Standard For	mulae, page	e 250	
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO	4.00	20	✓ Fresenius Kabi
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PSO	5.25	50	✓ Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	✓ Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)			

1) (On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical
Ç	Schedule requiring a solvent or diluent: or

- 2) On a bulk supply order; or
- 3) When used in the extemporaneous compounding of eye drops; or

4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 10 ml ampoule - Up to 5 inj available on a PSO7.19	50	✓ Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO5.00	20	✓ Fresenius Kabi

500 ml

1 OP

✓ Baxter

✓ TPN

WATER

	Subsidy (Manufacturer's F \$	Price) Subsi Per		
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	✓ Calciur	m Resonium
COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO		50	✓ Electra	<u>I</u>
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)		1,000 ml OP	✓ Pedialy Bubb	rte - olegum
PHOSPHORUS			4.5.	
Tab eff 500 mg (16 mmol)	82.50	100	Phospi	nate Phebra
POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26	60	Chlorve	escent
* Tab long-acting 600 mg (8 mmol)	, ,	200	✓ Span-k	(
SODIUM BICARBONATE				
Cap 840 mg	8.52	100	✓ Sodibio ✓ Sodibio	
SODIUM POLYSTYRENE SULPHONATE				
Powder	84.65	454 a OP	✓ Resoni	um-A

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	Generic
		\$	Per	1	Manufacturer
Alpha-Adren	oceptor Blockers				
Alpha Adrend	oceptor Blockers				
DOXAZOSIN					
		17.35	500	1	Doxazosin Clinect
			500		Doxazosin Clinect
J		20.04	500	•	DOXUZUSIII OIIIICCI
PHENOXYBENZA	MINE HYDROCHLORIDE				
* Cap 10 mg		65.00	30	✓	BNM S29
		216.67	100	✓	Dibenzyline S29
PRAZOSIN					•
		E E0	100	./	Arrotex-Prazosin
★ Tab Ting			100	•	
				_	S29 S29
* Tab 2 mg		7.00	100	✓	Arrotex-Prazosin
					S29 S29
* Tab 5 mg		11.70	100	/	Arrotex-Prazosin
					S29 S29
					329 023
Agonto Affoo	ting the Renin-Angiotensin Systen	•			
Agents Anec	ung me nemin-Angiotensin System				
A OF 1-1-1-1-1-1					
ACE Inhibitor	rs .				
CAPTOPRIL					
		04.00	0	n /	0
	per ml	94.99 95	5 ml O	Ρ •	Capoten
Oral liquid	restricted to children under 12 years of age.				
CILAZAPRIL - Su	bsidy by endorsement				
	dorsement - Subsidised for patients who were	taking cilazapril prior	r to 1 I	May 2021	and the prescription is
	ordingly. Pharmacists may annotate the presc				
dispensing of d					
, ,		2 69	90	1	Zapril
0			90		Zapril
•			90		Zapril
ŭ		10.00	50	•	Zapin
ENALAPRIL MALE				_	
			100		Acetec
* Tab 10 mg		2.02	100	✓	Acetec
* Tab 20 mg		2.42	100	✓	Acetec
LISINOPRIL					
		11 07	90	1	Ethics Lisinopril
* Tub o mg			00		Teva Lisinopril
* Tab 10 mg		11.67	90		Ethics Lisinopril
• 1 ab 10 1119		11.01	90		
₩ Tab 00 m=		14.60	00		Teva Lisinopril
* Tab 20 mg		14.69	90		Ethics Lisinopril
				•	Teva Lisinopril
PERINDOPRIL					
				_	

30

30

30

✓ Coversyl

✓ Coversyl

✓ Coversyl

Tab 10 mg5.18 90 ✓ Arrow	
Tab 10 mg	
Tab 20 mg	Quinapril 5
RAMIPRIL * Cap 1.25 mg	Quinapril 10
 ★ Cap 1.25 mg	Quinapril 20
 Cap 2.5 mg	
★ Cap 5 mg	
★ Cap 10 mg	
ACE Inhibitors with Diuretics QUINAPRIL WITH HYDROCHLOROTHIAZIDE − Subsidy by endorsement Subsidy by endorsement − Subsidised for patients who were taking quinapril with hydrochlorothiazide pri 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsexists a record of prior dispensing of quinapril with hydrochlorothiazide. Tab 10 mg with hydrochlorothiazide 12.5 mg	
QUINAPRIL WITH HYDROCHLOROTHIAZIDE – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who were taking quinapril with hydrochlorothiazide pri 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsexists a record of prior dispensing of quinapril with hydrochlorothiazide. Tab 10 mg with hydrochlorothiazide 12.5 mg	
Subsidy by endorsement – Subsidised for patients who were taking quinapril with hydrochlorothiazide pri 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsexists a record of prior dispensing of quinapril with hydrochlorothiazide. Tab 10 mg with hydrochlorothiazide 12.5 mg	
Angiotensin II Antagonists	
CANDESARTAN CILEXETIL	
* Tab 4 mg2.00 90 ✓ Cande	star
★ Tab 8 mg	
K Tab 16 mg	
≰ Tab 32 mg5.26 90 ✓ Cande	star
OSARTAN POTASSIUM	•
	an Actavis
	an Actavis
	an Actavis
k Tab 100 mg3.50 84 ✓ Losart	an Actavis
Angiotensin II Antagonists with Diuretics	
OSARTAN POTASSIUM WITH HYDROCHI OROTHIAZIDE	

LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE

* Tab 50 mg with hydrochlorothiazide 12.5 mg......4.00

✓ <u>Arrow-Losartan &</u> Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

✓ Entresto 24/26

✓ Entresto 49/51

30

56

✓ Entresto 97/103

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 2.1 Patient is in NYHA/WHO functional class II; or
- 2.2 Patient is in NYHA/WHO functional class III; or
- 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local,	page 120	
AMIODARONE HYDROCHLORIDE	00	. Austra
▲ Tab 100 mg	30 30	✓ <u>Aratac</u> ✓ Aratac
▲ Tab 200 mg4.49 Inj 50 mg per ml, 3 ml ampoule — Up to 10 inj available on a PSO9.12	30 6	✓ Aratac ✓ Cordarone-X
15.22	10	✓ Max Health
	10	▼ IVIAX FICAILII
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a	40	4.11
PSO15.09	10	✓ <u>Martindale</u>
DIGOXIN		_
* Tab 62.5 mcg – Up to 30 tab available on a PSO	240	✓ <u>Lanoxin PG</u>
* Tab 250 mcg – Up to 30 tab available on a PSO	240	Lanoxin
* Oral liq 50 mcg per ml	60 ml	Lanoxin
		✓ Lanoxin Paediatric Elixir \$29
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan
FLECAINIDE ACETATE	100	- Tryamioudii
. ==	60	✓ Flecainide BNM
▲ Tab 50 mg	90	✓ Flecainide BNM ✓ Flecainide
Cap to 19-acting 100 trig	90	Controlled Release Teva
▲ Cap long-acting 200 mg167.92	90	✓ Flecainide
2 Cap (orig doding 200 mg)	00	Controlled Release Teva
Inj 10 mg per ml, 15 ml ampoule100.00	5	✓ Tambocor
MEXILETINE HYDROCHLORIDE	Ü	- 1411100001
▲ Cap 150 mg	100	✓ Teva S29
▲ Cap 250 mg202.00	100	✓ Teva S29
PROPAFENONE HYDROCHLORIDE		
▲ Tab 150 mg40.90	50	✓ Rytmonorm

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

Antihypotensives

MIDODRINE - Special Authority see SA1474 below - Retail p	harmacy		
Tab 2.5 mg	53.00	100	Gutron
Tab 5 mg	79.00	100	Gutron

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

AT	ENOLOL Tob 50 mg	0.22	500	✓ Mylan Atenolol
不	Tab 50 mg	9.33	300	✓ Mylan Atendiol
*	Tab 100 mg	14.20	500	✓ Mylan Atenolol
*	Oral liq 25 mg per 5 ml		300 ml OP	✓ Atenolol AFT
	3,1			S29 S29
		38.20		✓ Essential
				Generics S29
		49.85		✓ Atenolol AFT
	Restricted to children under 12 years of age.			
BIS	SOPROLOL FUMARATE			
*	Tab 2.5 mg	1.84	90	✓ Bisoprolol Mylan
*	Tab 5 mg	2.55	90	✓ Bisoprolol Mylan
*	Tab 10 mg	3.62	90	✓ Bisoprolol Mylan
CA	RVEDILOL			
*	Tab 6.25 mg	2.24	60	✓ Carvedilol Sandoz
*	Tab 12.5 mg	2.30	60	✓ Carvedilol Sandoz
*	Tab 25 mg	2.95	60	Carvedilol Sandoz
LA	BETALOL			
*	Tab 100 mg	14.50	100	✓ Trandate
*	Tab 200 mg	27.00	100	✓ Trandate
*	Inj 5 mg per ml, 20 ml ampoule	59.06	5	
		(88.60)		Trandate
*	inj 5 mg per ml, 20 ml vial		1	
		(48.20)		Alvogen S29
ME	TOPROLOL SUCCINATE			
*	Tab long-acting 23.75 mg	1.45	30	✓ Betaloc CR
*	Tab long-acting 47.5 mg		30	✓ Betaloc CR
*	Tab long-acting 95 mg		30	✓ Betaloc CR
*	Tab long-acting 190 mg	4.27	30	✓ Betaloc CR
ME	TOPROLOL TARTRATE			
*	Tab 50 mg	5.66	100	✓ IPCA-Metoprolol
*	Tab 100 mg		60	✓ IPCA-Metoprolol
*	Tab long-acting 200 mg		28	✓ Slow-Lopresor
*	Inj 1 mg per ml, 5 ml vial	26.50	5	Metoprolol IV Mylan

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
NADOLOL				
* Tab 40 mg	19.19	100	✓	Nadolol BNM
* Tab 80 mg	30.39	100	1	Nadolol BNM
PROPRANOLOL				
Tab 10 mg	7.04	100	✓	Drofate
* Tab 40 mg	8.75	100	✓	IPCA-Propranolol
* Cap long-acting 160 mg		100	✓	Cardinol LA
* Oral liq 4 mg per ml - Special Authority see SA1327 below -	_			
Retail pharmacy	CBS	500 n	nl 🗸	Roxane- Propranolol \$29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

AMI ODIPINE

*	Tab 80 mg37.50	500	✓ Mylan
*	Tab 160 mg14.00	100	✓ Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

ΑIV	LODIFINE		
*	Tab 2.5 mg1.08	90	✓ Vasorex
*	Tab 5 mg	90	✓ Vasorex
*	Tab 10 mg1.19	90	✓ Vasorex
FE	LODIPINE		
*	Tab long-acting 2.5 mg	30	✓ Plendil ER
*	Tab long-acting 5 mg4.07	90	✓ Felo 5 ER
*	Tab long-acting 10 mg4.32	90	✓ Felo 10 ER
NIF	EDIPINE		
*	Tab long-acting 10 mg18.80	56	✓ Tensipine MR10 S29
*	Tab long-acting 20 mg9.12	50	✓ Mylan (12 hr
			release) \$29
	17.72	100	Nyefax Retard
*	Tab long-acting 30 mg4.78	14	Mylan Italy (24 hr
			release) S29
	34.10	100	✓ Mylan (24 hr
			release) S29
*	Tab long-acting 60 mg52.81	100	✓ Mylan (24 hr
			release) \$29

V	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
Cap extended-release 120 mg	44.40	100	1	Accord S29
* Cap long-acting 120 mg		500	✓	Apo-Diltiazem CD
	65.35		✓	Diltiazem CD Clinect
* Cap long-acting 180 mg	7.00	30	✓	Cardizem CD
* Cap long-acting 240 mg	9.30	30	✓	Cardizem CD
Accord 229 Cap extended-release 120 mg to be delisted 1 June (Apo-Diltiazem CD Cap long-acting 120 mg to be delisted 1 June 2				
PERHEXILINE MALEATE				
米 Tab 100 mg	62.90	100	•	Pexsig
/ERAPAMIL HYDROCHLORIDE				
* Tab 40 mg	7.01	100	1	Isoptin
* Tab 80 mg		100		Isoptin
* Tab long-acting 120 mg	36.02	100	/	Isoptin Retard \$29
				Isoptin SR
* Tab long-acting 240 mg	15.12	30	✓	Isoptin SR
* Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a				
PSO	25.00	5	✓	Isoptin
Centrally-Acting Agents				
CLONIDINE ★ Patch 2.5 mg, 100 mcg per day – Only on a prescription	10.34	4	/	<u>Mylan</u>
 Patch 2.5 mg, 100 mcg per day – Only on a prescription Patch 5 mg, 200 mcg per day – Only on a prescription 	13.18	4		Mylan Mylan
* Patch 2.5 mg, 100 mcg per day - Only on a prescription	13.18		1	
 Patch 2.5 mg, 100 mcg per day – Only on a prescription Patch 5 mg, 200 mcg per day – Only on a prescription 	13.18	4	1	Mylan
 Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription 	13.18 16.93	4	1	Mylan
 Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription CLONIDINE HYDROCHLORIDE 	13.18 16.93 29.32	4	<i>y y</i>	Mylan Mylan
Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription CLONIDINE HYDROCHLORIDE Tab 25 mcg	13.18 16.93 29.32 37.07	4 4 112	<i>y y y</i>	Mylan Mylan Clonidine Teva
Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription CLONIDINE HYDROCHLORIDE Tab 25 mcg	13.18 16.93 29.32 37.07	4 4 112 100	<i>y y y</i>	Mylan Mylan Clonidine Teva Catapres
Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription CLONIDINE HYDROCHLORIDE Tab 25 mcg	13.18 16.93 29.32 37.07 29.68	4 4 112 100	\ \ \ \	Mylan Mylan Clonidine Teva Catapres
Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription CLONIDINE HYDROCHLORIDE Tab 25 mcg Tab 150 mcg Inj 150 mcg per ml, 1 ml ampoule	13.18 16.93 29.32 37.07 29.68	4 4 112 100 10	\ \ \ \	Mylan Mylan Clonidine Teva Catapres Medsurge
Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription CLONIDINE HYDROCHLORIDE Tab 25 mcg Tab 150 mcg Inj 150 mcg per ml, 1 ml ampoule	13.18 16.93 29.32 37.07 29.68	4 4 112 100 10 100	\ \ \ \	Mylan Mylan Clonidine Teva Catapres Medsurge Methyldopa Mylan
Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription CLONIDINE HYDROCHLORIDE Tab 25 mcg Inj 150 mcg per ml, 1 ml ampoule	13.18 16.93 29.32 37.07 29.68	4 4 112 100 10 100	\ \ \ \	Mylan Mylan Clonidine Teva Catapres Medsurge Methyldopa Mylan Methyldopa Mylan
Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription CLONIDINE HYDROCHLORIDE Tab 25 mcg Tab 150 mcg Inj 150 mcg per ml, 1 ml ampoule	13.18 16.93 29.32 37.07 29.68	4 4 112 100 10 100	\ \ \ \	Mylan Mylan Clonidine Teva Catapres Medsurge Methyldopa Mylan Methyldopa Mylan
Patch 2.5 mg, 100 mcg per day — Only on a prescription Patch 5 mg, 200 mcg per day — Only on a prescription Patch 7.5 mg, 300 mcg per day — Only on a prescription CLONIDINE HYDROCHLORIDE Tab 25 mcg Inj 150 mcg per ml, 1 ml ampoule	13.18 16.93 29.32 37.07 29.68	4 4 112 100 10 100	\ \ \ \	Mylan Mylan Clonidine Teva Catapres Medsurge Methyldopa Mylan Methyldopa Mylan
* Patch 2.5 mg, 100 mcg per day — Only on a prescription * Patch 5 mg, 200 mcg per day — Only on a prescription * Patch 7.5 mg, 300 mcg per day — Only on a prescription * CLONIDINE HYDROCHLORIDE * Tab 25 mcg * Tab 150 mcg * Inj 150 mcg per ml, 1 ml ampoule	13.18 16.93 29.32 37.07 29.68	4 4 112 100 10 100	\ \ \ \	Mylan Mylan Clonidine Teva Catapres Medsurge Methyldopa Mylan Methyldopa Mylan
* Patch 2.5 mg, 100 mcg per day — Only on a prescription * Patch 5 mg, 200 mcg per day — Only on a prescription * Patch 7.5 mg, 300 mcg per day — Only on a prescription * Patch 7.5 mg, 300 mcg per day — Only on a prescription * CLONIDINE HYDROCHLORIDE * Tab 25 mcg * Tab 150 mcg * Inj 150 mcg per ml, 1 ml ampoule	13.18 29.32 37.07 29.68 15.10 52.85	4 4 112 100 10 100 500	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Mylan Mylan Clonidine Teva Catapres Medsurge Methyldopa Mylan Methyldopa Mylan S29 \$29
* Patch 2.5 mg, 100 mcg per day — Only on a prescription * Patch 5 mg, 200 mcg per day — Only on a prescription * Patch 7.5 mg, 300 mcg per day — Only on a prescription * Patch 7.5 mg, 300 mcg per day — Only on a prescription * CLONIDINE HYDROCHLORIDE * Tab 25 mcg * Tab 150 mcg	13.18 29.32 37.07 29.68 15.10 52.85	4 4 112 100 10 100		Mylan Mylan Clonidine Teva Catapres Medsurge Methyldopa Mylan Methyldopa Mylan

	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer
UROSEMIDE [FRUSEMIDE]				
Tab 40 mg - Up to 30 tab available on a PSO	8.00	1,000	1	IPCA-Frusemide
★ Tab 500 mg		50	1	Urex Forte
	89.48		1	Furosemid-
				Ratiopharm S29
	169.96	100	1	Furosemid-
				Ratiopharm §29
Oral liq 10 mg per ml	11.20	30 ml OP	1	Lasix
foliation in the last section in the last sect	60.65	6	1	Lasix
Inj 10 mg per ml, 2 ml ampoule - Up to 5 inj available o	on a PSO2.40	5	1	Furosemide-Baxter
Potassium Sparing Diuretics				
MILORIDE HYDROCHLORIDE				
Oral liq 1 mg per ml	32.10	25 ml OP	1	Biomed
PLERENONE - Special Authority see SA1728 below - Re				
Tab 25 mg		30	1	Inspra
Tab 50 mg		30		Inspra
■ SA1728 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals the following criteria: toth: 1 Patient has heart failure with ejection fraction less the Either: 2.1 Patient is intolerant to optimal dosing of spiror 2.2 Patient has experienced a clinically significant	an 40%; and			·
IETOLAZONE				
Tab 5 mg	CBS	1	1	Metolazone S29
		50	1	Zaroxolyn S29
PIRONOLACTONE				-
€ Tab 25 mg	3.68	100	1	Spiractin
€ Tab 100 mg		100		Spiractin
Oral liq 5 mg per ml	30.60	25 ml OP		Biomed
Potassium Sparing Combination Diuretics				
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE				

 28

50

✓ Frumil

✓ Moduretic

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
Thiazide and Related Diuretics			
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	✓ <u>Arrow-</u> Bendrofluazide
May be supplied on a PSO for reasons other than emerg * Tab 5 mg		500	✓ <u>Arrow-</u> Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml CHLORTALIDONE [CHLORTHALIDONE]	27.82	25 ml OP	✓ Biomed
Tab 25 mg	3.90 6.95	30 50	✓ Igroton S29 ✓ Hygroton
(Igroton ©29 Tab 25 mg to be delisted 1 April 2023) INDAPAMIDE * Tab 2.5 mg		90	✓ <u>Dapa-Tabs</u>
	11.61	100	Mylan Indapamide \$29
Vasopressin receptor antagonists			
TOLVAPTAN - Special Authority see SA2166 below - Retail pha	armacy		•

IOLVAPTAN – Special Authority see SA2166 below – F	łetail pharmacy		
Tab 15 mg	873.50	28 OP	Jinarc
Tab 30 mg	873.50	28 OP	Jinarc
Tab 45 mg + 15 mg		56 OP	Jinarc
Tab 60 mg + 30 mg		56 OP	Jinarc
Tab 90 mg + 30 mg		56 OP	Jinarc

⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Either:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

	Subsidy (Manufacturer's Price) \$	F Subsid Per	ully ised	Brand or Generic Manufacturer
Lipid-Modifying Agents				
Fibrates				
BEZAFIBRATE * Tab 200 mg * Tab long-acting 400 mg		90 30	_	Bezalip Bezalip Retard
Other Lipid-Modifying Agents				
ACIPIMOX	21.56	30		Dibetam Dibetam S29 S29
Resins				
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	32.89	30	✓ C	Colestid
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN * Tab 10 mg	9.24 26.54 2.11 3.61 il pharmacy 1.70 2.42 3.92 5.28	500 500 500 500 28 28 30 30 30 30		orstat orstat orstat orstat orstat ravastatin Mylan ravastatin Mylan ravastatin Mylan rosuvastatin Viatris rosuvastatin Viatris rosuvastatin Viatris
Initial application — (cardiovascular disease risk) from any runless notified for applications meeting the following criteria: Either: 1 Both: 1.1 Patient is considered to be at risk of cardiovascula 1.2 Patient is Māori or any Pacific ethnicity; or 2 Both:	·	Approvals va	alid w	ithout further renewal

- 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
- 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	•	

continued...

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
 - 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIMVASTATIN

*	Tab 10 mg	90	Simvastatin Mylan
	Tab 20 mg	90	✓ Simvastatin Mylan
	Tab 40 mg	90	✓ Simvastatin Mylan
	Tab 80 mg7.12	90	✓ Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Special Authority see SA1045 below – Retail	pharmacy		
* Tab 10 mg	1.95	30	✓ Ezetimibe Sandoz

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 x normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 on the next page - Retail pharmacy

Tab 10 mg with simvastatin 10 mg	5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg	7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	✓ Zimybe

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
<u> </u>	Per	✓	Manufacturer

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

.....6.09

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

*	Oral pump spray, 400 mcg per dose – Up to 250 dose
	available on a PSO

*	Patch 25 mg, 5 mg per day15.73	30
*	Patch 50 mg, 10 mg per day18.62	30

✓ Nitrolingual Pump Spray✓ Nitroderm TTS

•	·	
ISC	SORBIDE MONONITRATE	
*	Tab 20 mg	100

✓ Nitroderm TTS
✓ Ismo 20

*	Tab 20 mg	19.55
*	Tab long-acting 40 mg	8.20
*	Tab long-acting 60 mg	9.25

✓ Ismo 40 Retard
✓ Duride

Sympathomimetics

Α	n	R	F	N	Α	Ш	IN	F

Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO4.98	
10.76	
Ini 1 in 10,000, 10 ml ampoule – Up to 5 ini available on a PSO27.00	

✓ Aspen Adrenaline✓ DBL Adrenaline

...27.00 5

250 dose OP

30

90

5

✓ Hospira

49.00 10 ✓ Aspen Adrenaline

Vasodilators

HYDRALAZINE HYDROCHLORIDE

*	Tab 25 mg - Special Authority see SA1321 below - Retail
	pharmacy

1 ✓ Hydralazine
56 ✓ Onelink \$29

56 **✓ One** 84 **✓ AM**

100 5 ✓ AMDIPHARM S29

Onelink S29

Apresoline

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

.....CBS

Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIL

▲ Tab 10 mg70.00 100 **✓ Loniten**

Reddy's

Reddy's

✓ Bosentan Dr

60

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	
NICORANDIL				
▲ Tab 10 mg	25.57	60	✓	Ikorel
▲ Tab 20 mg	32.28	60	•	Ikorel
PAPAVERINE HYDROCHLORIDE				
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	✓	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]				
Tab 400 mg	42.26	50	1	Trental 400
Endothelin Receptor Antagonists				
AMBRISENTAN – Special Authority see SA1702 below – Retail Tab 5 mg Tab 10 mg	1,550.00	30 30	√	Ambrisentan Mylan Ambrisentan Mylan Ambrisentan Viatris Mylan
(Ambrisentan Mylan Tab 10 mg to be delisted 1 March 2023)				•
Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertensic Notes: Application details may be obtained from Pharmac's websithe Coordinator, PAH Panel Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac BOSENTAN − Special Authority see SA1991 below − Retail pharmac	site <u>schedule.pharma</u> .govt.nz rmacy			
Tab 62.5 mg	119.00	60	V	Bosentan Dr

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and

- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and

continued...

- 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1992 below – Retail ph	armacy		
Tab 25 mg	0.85	4	✓ Vedafil
Tab 50 mg	1.70	4	✓ Vedafil
Tab 100 mg		12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy	e)	Fully	Brand or
(Manufacturer's Pric		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV: and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- Both:
 - 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
 - 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL – Special Authority see SA1696 below – Retail p	harmacy		
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri
TO CA1606 Chaolal Authority for Cubaidy			

| ⇒SA1696 | Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 below - Retail pharmacy

(Ventavis Nebuliser soln 10 mcg per ml, 2 ml to be delisted 1 March 2023)

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

ADAPAI FNF

- a) Maximum of 30 g per prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA2023 below - Ret	ail pharmacy		
Cap 5 mg	11.26	60	Oratane
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
- 2 Patient is not of child bearing potential.

TRFTINOIN

✓ ReTrieve Crm 0.5 mg per g - Maximum of 50 g per prescription15.57 50 g OP

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

HYDROGEN PEROXIDE

* Crm 1%	8.56	0	✓ Crystaderm✓ Crystaderm
MUPIROCIN	2.22	45.00	
Oint 2%	6.60 (11.50)	15 g OP	Bactroban

- a) Only on a prescription
- b) Not in combination

	Subsidy		Fully	Brand or
	(Manufacturer's Pri		Subsidised	
	\$	Per		Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID]			_	
Crm 2%	1.59	5 g OP	•	<u>Foban</u>
a) Maximum of 5 g per prescription				
b) Only on a prescriptionc) Not in combination				
Oint 2%	1.59	5 g OP	1	Foban
a) Maximum of 5 g per prescription		og o.		<u> </u>
b) Only on a prescription				
c) Not in combination				
SULFADIAZINE SILVER				
Crm 1%	10.80	50 g OF	•	Flamazine
a) Up to 250 g available on a PSO				
b) Not in combination				
Audiformatic Tourism				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals, pa	ge 96			
AMOROLFINE	90 00			
a) Only on a prescription				
b) Not in combination				
Nail soln 5%	14.93	5 ml OF	•	MycoNail
CLOTRIMAZOLE				
* Crm 1%	1.10	20 g OF	•	Clomazol
a) Only on a prescription		-		
b) Not in combination				
* Soln 1%		20 ml O	•	•
	(7.55)			Canesten
a) Only on a prescriptionb) Not in combination				
•				
ECONAZOLE NITRATE Crm 1%	1.00	20 a OE)	
OIII 176	(7.48)	20 g OF		Pevaryl
a) Only on a prescription	(7.40)			1 ovaryi
b) Not in combination				
Foaming soln 1%, 10 ml sachets	9.89	3		
	(17.23)			Pevaryl
a) Only on a prescription				
b) Not in combination				
MICONAZOLE NITRATE				
* Crm 2%	0.81	15 g OF	•	Multichem
a) Only on a prescription				
b) Not in combination * Lotn 2%	4 36	30 ml Ol	5	
A LOUI 2 /0	(10.03)	30 1111 01		Daktarin
a) Only on a prescription	(.0.00)			
b) Not in combination				
* Tinct 2%	4.36	30 ml O	-	
	(12.10)			Daktarin
a) Only on a prescription				
b) Not in combination				

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antipruritic Preparations

CALAMINE

a) Only on a prescription

b) Not in combination

.08 100 g

✓ Calamine-AFT

CROTAMITON

a) Only on a prescription

b) Not in combination

3.29 20 g OP

✓ Itch-Soothe

MENTHOL - Only in combination

1) Only in combination with a dermatological base or proprietary Topical Corticosteriod - Plain

2) With or without other dermatological galenicals.

 25 g 100 g ✓ MidWest
✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 78

Corticosteroids - Plain

BE	TAMETHASONE DIPROPIONATE			
	Crm 0.05%	2.96	15 g OP	✓ <u>Diprosone</u>
		36.00	50 g OP	✓ Diprosone
	Oint 0.05%	2.96	15 g OP	✓ Diprosone
		36.00	50 g OP	✓ Diprosone
	Oint 0.05% in propylene glycol base	4.33	30 g OP	✓ Diprosone OV
BE	TAMETHASONE VALERATE			
*	Crm 0.1%	4.53	50 g OP	✓ Beta Cream
*	Oint 0.1%	5.84	50 g OP	✓ Beta Ointment
*	Lotn 0.1%		50 ml OP	✓ Betnovate
CL	OBETASOL PROPIONATE			
*	Crm 0.05%	2.40	30 g OP	✓ Dermol
*	Oint 0.05%	2.33	30 g OP	✓ Dermol
CL	OBETASONE BUTYRATE		-	
	Crm 0.05%	5.38	30 g OP	
		(10.00)	-	Eumovate
HY	DROCORTISONE			
*	Crm 1% - Only on a prescription	1.78	30 g OP	✓ Ethics
	, , ,	17.15	500 g	✓ Hydrocortisone (PSM)
*	Powder – Only in combination	49.95	25 g	✓ ABM
	Up to 5% in a dermatological base (not proprietary Topica galenicals			or without other dermatological
HY	DROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
	Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only on	ı		
	4			_

a prescription.......10.57

✓ DP Lotn HC

250 ml

Subsidy (Manufacturer's Price) Subsidy (Manufacturer's Price) Subsidy (Manufacturer's Price) Subsidy (Manufacturer's Price) Subside Per	
HYDROCORTISONE BUTYRATE Lipocream 0.1%	Fully Brand or
HYDROCORTISONE BUTYRATE Lipocream 0.1%	ubsidised Generic
Lipocream 0.1%	✓ Manufacturer
Oint 0.1% 10.28 100 g OP Milky emul 0.1% 12.33 100 ml OP METHYLPREDNISOLONE ACEPONATE 2.46 15 g OP Oint 0.1% 4.46 15 g OP MOMETASONE FUROATE 1.95 15 g OP Oint 0.1% 1.95 15 g OP Oint 0.1% 1.95 15 g OP Oint 0.1% 1.95 15 g OP 2.90 50 g OP	
Milky emul 0.1% 12.33 100 ml OP METHYLPREDNISOLONE ACEPONATE 4.46 15 g OP Oint 0.1% 4.46 15 g OP MOMETASONE FUROATE 1.95 15 g OP Crm 0.1% 1.95 15 g OP Oint 0.1% 1.95 15 g OP 2.90 50 g OP	Locoid Lipocream
METHYLPREDNISOLONE ACEPONATE Crm 0.1%	
Crm 0.1% 4.46 15 g OP Oint 0.1% 4.46 15 g OP MOMETASONE FUROATE 1.95 15 g OP Crm 0.1% 1.95 15 g OP Oint 0.1% 1.95 15 g OP 2.90 50 g OP	✓ <u>Locoid Crelo</u>
Oint 0.1%	
MOMETASONE FUROATE Crm 0.1%	✓ <u>Advantan</u>
Crm 0.1% 1.95 15 g OP 3.10 50 g OP Oint 0.1% 1.95 15 g OP 2.90 50 g OP	✓ Advantan
3.10 50 g OP Oint 0.1%	
Oint 0.1%	✓ Elocon Alcohol Free
2.90 50 g OP	✓ Elocon Alcohol Free
=:	✓ Elocon
Lotn 0.1% 4.50 30 ml OP	✓ Elocon
0	✓ <u>Elocon</u>
TRIAMCINOLONE ACETONIDE	
Crm 0.02%	✓ Aristocort
Oint 0.02%	✓ Aristocort
Corticosteroids - Combination	
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID]	
Crm 0.1% with sodium fusidate (fusidic acid) 2%3.49 15 g OP	
(10.45)	Fucicort
a) Maximum of 15 g per prescription	
b) Only on a prescription	
HYDROCORTISONE WITH MICONAZOLE – Only on a prescription	
* Crm 1% with miconazole nitrate 2%	✓ Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN – Only on a prescription	
Crm 1% with natamycin 1% and neomycin sulphate 0.5%	✓ Pimafucort
Oint 1% with natamycin 1% and neomycin sulphate 0.5%	✓ Pimafucort
(Pimafucort Crm 1% with natamycin 1% and neomycin sulphate 0.5% to be delisted 1 April 20.	
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN. NEOMYCIN AND NYSTATIN	-5)
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g - Only on a prescription 3.49 15 g OP	
(9.28)	Viaderm KC
(0.20)	Viddominio
Barrier Creams and Emollients	
Barrier Creams	
DIMETHICONE	
* Crm 5% pump bottle	
* Crm 10% pump bottle	Dimethicone 5%
	Dimethicone 5% ✓ healthE
ZINC AND CASTOR OIL	Dimethicone 5%
* Oint	Dimethicone 5% ✓ healthE

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

(AM	Subsidy anufacturer's	Price) Subsi	Fully Brand or dised Generic
(1/1	\$	Per Per	✓ Manufacturer
Emollients			
AQUEOUS CREAM			
₭ Crm	1.73	500 g	✓ Evara
			✓ GEM Aqueous
France Com to be delicted 1 April 2002)			<u>Cream</u>
Evara Crm to be delisted 1 April 2023)			
CETOMACROGOL ★ Crm BP	1.00	500 g	✓ Cetomacrogol-AFT
	1.99	500 g	• Celoinacrogor-AFT
CETOMACROGOL WITH GLYCEROL Crm 90% with glycerol 10%	2.25	500 ml OP	✓ Boucher
Citil 90% with gryceror 10%	2.33	500 IIII OF	✓ Evara
			✓ Pharmacy Health
			Sorbolene with
			Glycerin
	3.10	1,000 ml OP	✓ Boucher
			✓ Evara
Boucher Crm 90% with glycerol 10% to be delisted 1 March 2023)			
Boucher Crm 90% with glycerol 10% to be delisted 1 March 2023)			
EMULSIFYING OINTMENT * Oint BP	2.40	E00 a	√ Emulaifuina
K UIII Dr	3.40	500 g	Emulsifying Ointment ADE
DIL IN WATER EMULSION			Omtinent ABL
★ Crm	2 04	500 g	✓ Fatty Cream AFT
PARAFFIN	2.07	000 g	rutty orcani Ar I
Oint liquid paraffin 50% with white soft paraffin 50%	4 94	500 g OP	✓ White Soft Liquid
Sint liquid parallili 50 /5 With Winto 551t parallili 50 /5		000 g 0.	Paraffin AFT
Oint liquid paraffin 50% with white soft paraffin 50%,	5.35	500 ml OP	✓ healthE
healthE Oint liquid paraffin 50% with white soft paraffin 50%, to be o	delisted 1 Ma	ay 2023)	
JREA			
₭ Crm 10%	1.37	100 g OP	✓ healthE Urea Cream
VOOL FAT WITH MINERAL OIL - Only on a prescription			
★ Lotn hydrous 3% with mineral oil	5.60	1,000 ml	
	(11.95)	050 100	DP Lotion
	1.40 (4.53)	250 ml OP	DP Lotion
	5.60	1,000 ml	DF LOUOII
	(20.53)	1,000 1111	Alpha-Keri Lotion
	(23.91)		BK Lotion
	1.40	250 ml OP	
	(7.73)		BK Lotion
Other Dermatological Bases			
•			
PARAFFIN White soft - Only in combination	⊿ 00	450 g	✓ healthE
Willie 30tt - Offig in combination	19.99	2,500 g	✓ healthE
Only in combination with a dermatological galenical or as a			

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer
Minor Skin Infections				
POVIDONE IODINE Oint 10% a) Maximum of 130 g per prescription b) Only on a prescription	7.40 6	65 g OP	✓ <u>B</u>	<u> Betadine</u>
Antiseptic Solution 10% Antiseptic soln 10%	3.83	100 ml 15 ml 500 ml	✓ F	R <u>iodine</u> Riodine Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	(3.48)	100 ml	В	Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63 (7.78)	100 ml	P	Pfizer
Parasiticidal Preparations				

אווע	/1⊏	ΙН	ICONE	

IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy

Tab 3 mg − Up to 100 tab available on a PSO......17.20 4 ✓ Stromectol

- PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables); or

DERMATOLOGICALS

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
- 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

- Any of the following:
 - 1 Filaricides; or
 - 2 Cutaneous larva migrans (creeping eruption); or
 - 3 Strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5%5.75	30 g OP	 Lyderm
Lotn 5%	30 ml OP	✓ A-Scabies

Psoriasis and Eczema Preparations

ACITRETIN – Special Authority see SA2024 on the next page –	Retail pharmacy		
Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

DERMATOLOGICALS

\$ Per ✔ Manufacturer	Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic
	 \$	Per 🗸	Manufacturer

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around actiretin and is competent to prescribe actiretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential

2 Patient is not of child bearing potential.			
BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Foam spray 500 mcg with calcipotriol 50 mcg per g	50 05	60 a OP	✓ Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g		60 g OP	✓ Daivobet
0 1		J	
Oint 500 mcg with calcipotriol 50 mcg per g	15.90	30 g OP	✓ <u>Daivobet</u>
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR		Ü	
Soln BP - Only in combination	36.25	200 ml	✓ Midwest
 Up to 10% only in combination with a dermatological ba With or without other dermatological galenicals. 	se or proprie	tary Topical Co	rticosteriod – Plain
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUF	?		
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and			
allantoin crm 2.5%	6 59	75 g OP	
and to the Lio / v	(8.00)	70 g 01	Egopsoryl TA
	3.43	20 a OB	Lgopsoryi 1A
		30 g OP	Г.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
	(4.35)		Egopsoryl TA

	3.43 (4.35)	30 g OP	Egopsoryl T
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ Coco-Scalp
,	7.95	40 g OP	✓ Coco-Scalp

PIMECROLIMUS - Special Authority see SA1970 on the next page - Retail pharmacy

- a) Maximum of 15 g per prescription
- b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks.

2) Trotor a maximum or ro g por procemption and no more than	5o p. 000p	pooc	,
Cream 1%	28.50	15 g OP	Elidel



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1970 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN – Only on a prescription * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium..........4.44 500 ml

SALICYLIC ACID

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain or collodion flexible
- 2) With or without other dermatological galenicals.

SULPHUR

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain
- 2) With or without other dermatological galenicals.

TACROLIMUS

Oint 0.1% − Special Authority see SA2074 below − Retail pharmacy.......33.00 30 g OP ✓ Zematop

- a) Maximum of 30 g per prescription
- b) Note: a maximum of 30 g per prescription and no more than one prescription per 12 weeks.

⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

BETAMETHASONE VALERATE			
* Scalp app 0.1%	9.84	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%	6.26	30 ml OP	✓ <u>Dermol</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	6.57	100 ml OP	✓ <u>Locoid</u>
KETOCONAZOLE Shampoo 2%	3.23	100 ml OP	✓ <u>Sebizole</u> ✓ <u>Sebizole</u>

- a) Maximum of 100 ml per prescription
- b) Only on a prescription

Pinetarsol

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 66

IMIQUIMOD

PODOPHYLLOTOXIN

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

GENITO-URINARY SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Subsidy (Manufacturer's Price) \$ Per

Fully Subsidised Brand or Generic Manufacturer

Contraceptives - Non-hormonal

Condoms

റ	NDOMS			
-	49 mm - Up to 144 dev available on a PSO	11.42	144	✓ Moments
	53 mm		10	✓ Moments
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
	53 mm, 0.05 mm thickness	0.95	10	✓ Moments
		11.42	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	53 mm, chocolate, brown	0.95	10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	53 mm, strawberry, red	0.95	10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm	0.97	10	✓ Moments
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
	56 mm, 0.05 mm thickness	1.30	12	Gold Knight
		15.57	144	Gold Knight
	 a) Up to 60 dev available on a PSO 			
	b) Maximum of 60 dev per prescription			
	56 mm, 0.05mm thickness (bulk pack)	14.61	144	Gold Knight
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
	56 mm, 0.08 mm thickness	0.97	10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, 0.08 mm thickness, red	0.97	10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, chocolate	1.30	12	Gold Knight
		15.57	144	Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, strawberry	1.30	12	Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	60 mm	1.42	12	Gold Knight XL
		14.87	144	✓ Shield XL
		17.02		Gold Knight XL

GENITO-URINARY SYSTEM

		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
*	60 mm (bulk pack)	14.87	144	✓	Gold Knight XL
	a) Maximum of 60 dev per prescription				
	b) Up to 60 dev available on a PSO				

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

* IUD 29.1 mm length × 23.2 mm width	29.80	1	✓ 7 MED NSHA Silver/ Copper Short
			✓ Choice 380 7med Nsha Silver/ copper Short
			✓ Choice TT380 Short
* IUD 33.6 mm length × 29.9 mm width	29.80	1	✓ Choice TT380 Standard
* IUD 35.5 mm length × 19.6 mm width	33.00	1	✓ Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit: and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
 - 1 Patient is on a Social Welfare benefit; or
 - 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO 10.00 84 ✓ Mercilon 28

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	I Generic
	\$	Per	•	Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets -	-			
Up to 112 tab available on a PSO	2.18	84	✓	Microgynon 20 ED
·	6.45	112		Femme-Tab ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
	(16.50)			Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Auth	nority see SA0500 on	the	orevious p	age
b) Up to 63 tab available on a PSO	•			
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	1.77	84	1	Levlen ED
	6.45	112	✓	Femme-Tab ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to	1			
84 tab available on a PSO		84	/	Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U		•		
to 84 tab available on a PSO		84	/	Norimin
to on tab available on a 1 oo	29.32	112		Norimin
	23.32	112	•	Norman

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

* Tab 30 mcg -	Up to 84 tab available on a PSO	16.50 22.00	84 112	✓ Microlut✓ Microlut
	olant (2 × 75 mg rods) - Up to 3 pack availabl		1	✓ <u>Jadelle</u>
	ESTERONE ACETATE ml, 1 ml syringe – Up to 5 inj available on a F	°SO7.98	1	✓ Depo-Provera

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84	✓ <u>N</u>	loriday 28
Emergency Contraceptives				
LEVONORGESTREL * Tab 1.5 mg	1.75	1	√ L	evonorgestrel BNM
a) Maximum of 2 tab per prescription	4.95		√ P	ostinor-1

- b) Up to 5 tab available on a PSO
 - c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

(Postinor-1 Tab 1.5 mg to be delisted 1 June 2023)

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEI	C ACID		
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulph	nate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with ap	plicator8.43	100 g OP	
,	(24.15)	Ů	Aci-Jel
CLOTRIMAZOLE			
* Vaginal crm 1% with applicators	3.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators	3.85	20 g OP	Clomazol
MICONAZOLE NITRATE			
* Vaginal crm 2% with applicator	6.89	40 g OP	✓ <u>Micreme</u>
NYSTATIN			
Vaginal crm 100 000 u per 5 g with applicator(s)	4 00	75 a OP	✓ Nilstat

Myometrial and Vaginal Hormone Preparations

	Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		
	PSO160.00	5	✓ DBL Ergometrine
OE	STRIOL		
*	Crm 1 mg per g with applicator6.62	15 g OP	✓ Ovestin
*	Pessaries 500 mcg6.86	15	✓ Ovestin

ERGOMETRINE MALEATE

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
OXYTOCIN – Up to 5 inj available on a PSO	4.00	_		
Inj 5 iu per ml, 1 ml ampoule		5		Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	5.98	5	•	Oxytocin BNM
			•	Oxytocin GH S29
OXYTOCIN WITH ERGOMETRINE MALEATE — Up to 5 inj avail Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampou		5	•	Syntometrine

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

40 test OP

16.00

 Smith BioMed Rapid **Pregnancy Test**

✓ David One Step Cassette **Pregnancy Test**

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 106

5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy 100 ✓ Ricit

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Fither:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy ✓ Tamsulosin-Rex

100

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN 100 ✓ Alchemy Oxybutynin S29

GENITO-URINARY SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

POTASSIUM CITRATE

Oral liq 3 mmol per ml - Special Authority see SA1083 below -

Retail pharmacy......31.80 200 ml OP **✓ Biomed**

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE

*	Grans eff 4 g sachets	2.22	28	✓ <u>Ural</u>
SOI	LIFENACIN SUCCINATE			
	Tab 5 mg	2.05	30	✓ Solifenacin Mylan
	Tab 10 mg	3.72	30	✓ Solifenacin Mylan

Detection of Substances in Urine

ORTHO-TOLIDINE

*	Compound diagnostic sticks7.50		50 test OP		
	(8.25))		H	Hemastix

TETRABROMOPHENOL

*	Blue diagnostic strips	13.92	100 test OP	✓ Albustix

Obstetric Preparations

Antiprogesterones

M	FEF	PRI	I P	\cap	NF
IVII		וח־	ΟI	U	NE.

✓ Mifegyne	1	O	Tab 200 mg - Up to 15 tab available on a PSO.
✓ Mifegyne	3	180.00	

Subsit (Manufacture	. ,	,
\$	Per	✓ Manufacturer

Calcium Homeostasis

CALCITONIN			_
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	Miacalcic
CINACALCET - Special Authority see SA2170 below - Re	etail pharmacy		
Tab 30 mg - Wastage claimable	42.06	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable	84.12	28	✓ Cinacalet Devatis

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (**primary hyperparathyroidism**) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
 \$	Per	1	Manufacturer

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial − Special Authority see SA2109 below −

Retail pharmacy.......18.00 1

✓ Zoledronic acid

Mylan

✓ Zoledronic acid

Viatris

⇒SA2109 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement: and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Roth:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer*) from any relevant practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 3 years.

Note: Indications marked with * are unapproved indications.

Initial application — (symptomatic hypercalcaemia*) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has symptomatic hypercalcaemia.

Note: Indications marked with * are unapproved indications.

Corticosteroids and Related Agents for Systemic Use

RETAMETHASONE SODILIM PHOSPHATE WITH RETAMETHASONE ACETATE

* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	19.20 (36.96)	5	Celestone Chronodose
Tab 0.5 mg – Up to 60 tab available on a PSO Tab 4 mg – Up to 30 tab available on a PSO Oral lig 1 mg per ml	2.65	30 30 25 ml OP	✓ <u>Dexmethsone</u> ✓ <u>Dexmethsone</u> ✓ Biomed

	Subsidy		Fully Brand or
	(Manufacturer's Price)	Subsidised Generic
	\$	Per	✓ Manufacturer
DEXAMETHASONE PHOSPHATE			
Dexamethasone phosphate injection will not be funded for o	oral use		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a F		10	✓ Hameln
The first tring per fills, i fill ampould to p to 5 mg available off a f	9.25	10	✓ Dexamethasone
	3.23		Phosphate
			•
No. 1 of A consequent O colores and a little to E fed a collection on E	10.40	40	Panpharma
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a F		10	✓ Hameln
	16.37		✓ Dexamethasone
			Phosphate
			Panpharma
(Dexamethasone Phosphate Panpharma Inj 4 mg per ml, 1 ml a			
(Dexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml a	impoule to be delisted	d 1 Fe	bruary 2023)
FLUDROCORTISONE ACETATE			
* Tab 100 mcg	11 46	100	✓ Florinef
•		100	<u> </u>
HYDROCORTISONE	0.40	400	45
* Tab 5 mg		100	3
* Tab 20 mg		100	3
* Inj 100 mg vial	4.38	1	✓ <u>Solu-Cortef</u>
 a) Up to 5 inj available on a PSO 			
b) Only on a PSO			
METHYLPREDNISOLONE			
* Tab 4 mg	112.00	100	✓ Medrol
* Tab 100 mg		20	✓ Medrol
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
,	00.00		Calu Madral Ast
Inj 40 mg vial	22.30	1	✓ Solu-Medrol-Act-
			O-Vial
Inj 125 mg vial	24.10	1	✓ Solu-Medrol-Act-
IIIJ 125 IIIg viai	34.10		O-Vial
			O-viai
Inj 500 mg vial	26.88	1	✓ Solu-Medrol-Act-
iiij 300 iiig vidi	20.00	'	O-Vial
			O-Viai
Inj 1 g vial	32 84	1	✓ Solu-Medrol
, ,			5 Ocia incaror
METHYLPREDNISOLONE ACETATE	47.00	_	45
Inj 40 mg per ml, 1 ml vial	47.06	5	✓ Depo-Medrol
PREDNISOLONE			
* Oral liq 5 mg per ml - Up to 30 ml available on a PSO	6.00 3	30 ml C	OP ✓ Redipred
Restricted to children under 12 years of age.			
PREDNISONE			
* Tab 1 mg	18 50	500	✓ Prednisone Clinect
* Tab 1 flig* * Tab 2.5 mg		500	
		500	
* Tab 5 mg - Up to 30 tab available on a PSO			
* Tab 20 mg - Up to 30 tab available on a PSO	1 6.06	500	Prednisone Clinect

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully Brand or sidised Generic Manufacturer
TETRACOSACTRIN			
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	✓ AU Synacthen✓ Synacthen✓ UK Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓ Synacthen Depot ✓ Synacthene Retard \$29
(AU Synacthen Inj 250 mcg per ml, 1 ml ampoule to be delisted 1	March 2023)		notal a
TRIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule		5	✓ Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule	51.10	5	✓ Kenacort-A 40
Sex Hormones Non Contraceptive			
Androgen Agonists and Antagonists			
CYPROTERONE ACETATE			
Tab 50 mg	14.37	50	✓ <u>Siterone</u>
Tab 100 mg	28.03	50	✓ <u>Siterone</u>
TESTOSTERONE			
Patch 5 mg per day	225.00	30	✓ Androderm
TESTOSTERONE CIPIONATE			
Inj 100 mg per ml, 10 ml vial		1	✓ Depo-Testosterone
	393.00		✓ Taro-
			Testosterone S29
TESTOSTERONE ESTERS			
Inj 250 mg per ml, 1 ml	12.98	1	✓ Sustanon Ampoules
TESTOSTERONE UNDECANOATE			-
Cap 40 mg - Subsidy by endorsement	21.00	60	✓ Andriol Testocaps
	35.00	100	✓ Steril-Gene S29
Subsidy by endorsement – subsidised for patients who w 1 November 2021 and the prescription is endorsed accor where there exists a record of prior dispensing of testost Inj 250 mg per ml, 4 ml vial	rdingly. Pharmacists erone undecanoate c	may ann	otate the prescription as endorsed

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

Hormone Replacement Therapy - Systemic

Ш	ormone replacement rherapy bystemio			
C	estrogens			
OE	STRADIOL			
*	Tab 1 mg	4.12	28 OP	
		(11.10)		Estrofem
*	Tab 2 mg		28 OP	
	Datab 50 man now 04 hours	(11.10)	4	Estrofem
	Patch 50 mcg per 24 hours	7.04	4	✓ Climara
	a) No more than 1 patch per week			
	b) Only on a prescription Patch 25 mcg per day	6 12	8	✓ Estradot
	1 alcii 25 mog per day	13.50	U	✓ Estrador MX S29
	a) No more than 2 patch per week	10.50		- Estraderiii MX
	b) Only on a prescription			
	Patch 50 mcg per day	7.04	8	✓ Estradot 50 mcg
		9.22	-	✓ Estradiol TDP
				Mylan S29
		14.50		✓ Estraderm MX S29
	a) No more than 2 patch per week			
	b) Only on a prescription			
	Patch 75 mcg per day	7.91	8	✓ Estradot
		10.60		Estradiol TDP
				Mylan S29
	a) No more than 2 patch per week			
	b) Only on a prescription			•
	Patch 100 mcg per day		8	✓ Estradot
		15.50		✓ Estraderm MX S29
	No more than 2 patch per week			
	b) Only on a prescription			
	STRADIOL VALERATE	40.00	0.4	4.5
	Tab 1 mg		84	✓ Progynova
	Tab 2 mg	12.36	84	✓ Progynova
-	STROGENS	0.04	00	
*	Conjugated, equine tab 300 mcg		28	Premarin
*	Conjugated, equine tab 625 mcg	(17.50)	28	Premann
*	Conjugated, equine tab 625 mcg	(17.50)	20	Premarin
		(17.00)		r romain
Р	rogestogens			
	DROXYPROGESTERONE ACETATE	4.00	00	/ Duamana
木	Tab 2.5 mg	4.69 8.75	30 56	✓ Provera✓ Provera
*	Tab 5 mg		56	✓ Provera ✓ Provera
-10	142 5 mg	17.50	100	✓ Provera
*	Tab 10 mg		30	✓ Provera
-	•			

	Subsidy		Fully Brand or
	(Manufacturer's Pric		osidised Generic
	\$	Per	✓ Manufacturer
Progestogen and Oestrogen Combined Prepar	ations		
DESTRADIOL WITH NORETHISTERONE			
* Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP	
	(18.10)		Kliovance
* Tab 2 mg with 1 mg norethisterone acetate		28 OP	
	(18.10)		Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	F 40	00 OD	
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	Trianantana
	(18.10)		Trisequens
Other Oestrogen Preparations			
ETHINYLOESTRADIOL - Subsidy by endorsement			
Subsidy by endorsement – Subsidised for patients who wer	e taking ethinyloesti	adiol prior t	o 1 March 2022 and the
prescription is endorsed accordingly. Pharmacists may ann	notate the prescription	on as endor	sed where there exists a record o
prior dispensing of ethinyloestradiol.			_
Tab 10 mcg	17.60	100	✓ NZ Medical and
			Scientific
NZ Medical and Scientific Tab 10 mcg to be delisted 1 Februar	y 2023)		
DESTRIOL			
米 Tab 2 mg	7.00	30	✓ Ovestin
Other Bregestagen Brenerations			
Other Progestogen Preparations			
EVONORGESTREL			
* Intra-uterine device 52 mg		1	✓ Mirena
* Intra-uterine device 13.5 mg	215.60	1	✓ Jaydess
MEDROXYPROGESTERONE ACETATE			
Tab 100 mg	116.15	100	✓ Provera HD
IORETHISTERONE			
* Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Primolut N
PROGESTERONE			
* Cap 100 mg	14.85	30	✓ Utrogestan
Thyroid and Antithyroid Agents			
CARBIMAZOLE			
* Tab 5 mg	7.56	100	✓ Neo-Mercazole
.EVOTHYROXINE			
* Tab 25 mcg	5.55	90	✓ Synthroid
* Tab 50 mcg	1.71	28	✓ Mercury Pharma
	5.79	90	✓ Synthroid
	64.28	1,000	✓ Eltroxin
★ Tab 100 mcg		28	✓ Mercury Pharma
	6.01	90	✓ Synthroid
	66.78	1,000	✓ Eltroxin
PROPYLTHIOURACIL - Special Authority see SA1199 on the	next page – Retail p	harmacy	

Su	ubsidy Ful	y Brand or
(Manufac	cturer's Price) Subsidise	d Generic
	\$ Per •	Manufacturer

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA2032 below	v – Retail pharm	асу	
*	Inj 5 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 15 mg cartridge	.139.50	1	✓ Omnitrope
	, ,			✓ Omnitrope S29 S29

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — **(short stature without growth hormone deficiency)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Fither:

|--|

continued...

- 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
- 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria;
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life

Subsidy (Manufacturer's Price)	Cubo	Fully	Brand or	
(Manufacturer's Price)	Per	idised •	Generic Manufacturer	

continued...

questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GOSERELIN			
Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe	122.37	1	✓ Teva

LEUPRORELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of			
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy			
of \$591.68 per 1 ini with Endorsement	177.50	1	

(591.68)

Vasopressin Agonists

DESMOPRESSIN Wafer 120 mcg	47.00	30	✓ Minirin Melt
DESMOPRESSIN ACETATE			
Tab 100 mcg	25.00	30	✓ Minirin
Tab 200 mcg	54.45	30	✓ Minirin
▲ Nasal spray 10 mcg per dose	27.95	6 ml OP	✓ <u>Desmopressin-</u> <u>PH&T</u>
Ini 4 mca per ml. 1 ml	67.18	10	✓ Minirin

Other Endocrine Agents

CABERGOI INF

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
Dostinex	2	waived by Special Authority see SA2070 on the next page 3.75
Dostinex	8	15.20

Lucrin Depot 3-month

Subsidy		Fully	Brand or
(Manufacturer's Price	,	Subsidised	Generic
\$	Per		Manufacturer

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLOMIFENE CITRATE Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE Cap 250 mg	558.00	50	✓ Metopirone

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Anthelmintics

ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

Tab 100 mg	7.97	6	✓ Vermox
Oral liq 100 mg per 5 ml		15 ml	
	(7.53)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	Biltricide

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 60
- b) For anti-infective eve preparations, refer to SENSORY ORGANS, page 243

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE			
Cap 250 mg	24.70	100	✓ Ranbaxy-Cefaclor S29 S29
	25.85		✓ Ranbaxy-Cefactor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.53	100 ml	✓ Ranbaxy-Cefactor S29 S29
	3.75		✓ Ranbaxy-Cefactor
(Ranbaxy-Cefaclor S29 S29 Cap 250 mg to be delisted 1 April 2023)			
(Ranbaxy-Cefaclor S29 S29 Grans for oral liq 125 mg per 5 ml to be of	delisted 1 Ap	ril 2023)	
CEFALEXIN			
Cap 250 mg	3.85	20	 Cephalexin ABM
Cap 500 mg	5.85	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	7.88	100 ml	✓ Flynn
Grans for oral liq 50 mg per ml – Wastage claimable	10.38	100 ml	✓ Flynn

CEFAZOLIN - Subsidy by endorsement

Only if prescribed for dialysis or cellulitis in accordance with a Health NZ Hospital approved protocol and the prescription is endorsed accordingly.

Inj 500 mg vial	3.39	5	✓ <u>AFT</u>
Ini 1 g vial	3.49	5	✓ AFT

CEFTRIAXONE - Subsidy by endorsement

- a) Up to 10 ini available on a PSO
- b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Inj 500 mg vial	0.79	1	✓ Ceftriaxone-AFT
lnj 1 g vial	3.59	5	✓ Ceftriaxone-AFT

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	(Manufacturer's Price)	Sub	Fully osidised	Brand or Generic	
	\$	Per	•	Manufacturer	
CEFUROXIME AXETIL – Subsidy by endorsement					
Only if prescribed for prophylaxis of endocarditis and the pre	escription is endorsed	accordin	gly.		
Tab 250 mg	45.93	50	✓ Z	innat	
(Zinnat Tab 250 mg to be delisted 1 March 2024)					

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO2.57	2	✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage		
claimable16.97	15 ml	Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN – Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 on the next page
Tab 250 mg8.53 14 Klacid
Grans for oral lig 250 mg per 5 ml – Wastage claimable.......192.00 50 ml

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Atypical mycobacterial infection; or

FRYTHROMYCIN (AS LACTORIONATE)

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Inj 1 g vial10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE		
Tab 400 mg16.95	100	E-Mycin
a) Up to 20 tab available on a PSO		
b) Up to 2 x the maximum PSO quantity for RFPP		
Grans for oral liq 200 mg per 5 ml	100 ml	E-Mycin
a) Up to 300 ml available on a PSO		
b) Up to 2 x the maximum PSO quantity for RFPP		
c) Wastage claimable	100	/ E Musin
Grans for oral liq 400 mg per 5 ml	100 ml	✓ E-Mycin
a) Up to 200 ml available on a PSO		
b) Wastage claimable		
ROXITHROMYCIN		<i>.</i>
Tab disp 50 mg	10	✓ Rulide D
Restricted to children under 12 years of age.	50	
Tab 150 mg29.50	50	✓ Arrow- Roxithromycin
		Hoxidiloniyoni
Tab 300 mg35.50	50	✓ Arrow-
		Roxithromycin

(Rulide D Tab disp 50 mg to be delisted 1 March 2023)

	Subsidy (Manufacturer's Price	·) S	Fully	
	\$	Per		Manufacturer
Penicillins				
AMOXICILLIN				
Cap 250 mg	22.50	500	1	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	36.98	500	•	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	•	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable	4.70	400	,	Alb050
Grans for oral liq 250 mg per 5 ml	1./3	100 ml	•	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPPc) Wastage claimable				
Inj 250 mg vial	15 97	10	1	Ibiamox
Inj 500 mg vial		10		Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO		10		Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab				
available on a PSO	0.89	10	1	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 r		10	•	Outum Duo 000/120
per ml	-	100 ml	1	Augmentin
a) Up to 200 ml available on a PSO				-
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r	ma			
per ml – Up to 200 ml available on a PSO		00 ml O	P 🗸	Curam
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj				
available on a PSO	375.97	10	1	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS	SO 11.09	10	1	Sandoz
FLUCLOXACILLIN	30 11.00	10	•	<u>Juliu02</u>
Cap 250 mg - Up to 30 cap available on a PSO	15 70	250	1	Flucloxacillin-AFT
Cap 500 mg – Up to 30 cap available on a PSO		500		Flucloxacillin-AFT
Grans for oral lig 25 mg per ml		100 ml		AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 50 mg per ml	3.68	100 ml	1	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Inj 250 mg vial		10		Flucloxin
Inj 500 mg vial		10	_	Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.70	5	/	<u>Flucil</u>

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	S	ubsidised	Generic
	\$	Per	1	Manufacturer
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg - Up to 30 cap available on a PSO	3.84	50	1	Cilicaine VK
Cap 500 mg		50	1	Cilicaine VK
a) Up to 20 cap available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	3.40	100 ml	✓ <u>I</u>	<u>AFT</u>
a) Up to 200 ml available on a PSOb) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	4.24	100 ml	√ <u>!</u>	<u>AFT</u>
 a) Up to 300 ml available on a PSO 				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
PROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe - Up to 5 inj available on a PSO	123.50	5	✓ (Cilicaine
(Cilicaine Inj 1.5 g in 3.4 ml syringe to be delisted 1 February 202				

Tetracyclines

DO)	///	\sim	N 18	ıΕ
עטע	11/	ハし	ı∟II	٧C

*	Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	✓ Doxine
MIN	IOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	19.32	100	
		(52.04)		Minomycin

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 60

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
CLINDAMYCIN				
Cap hydrochloride 150 mg	4.61	24	✓	Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	•	Dalacin C
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S	Subsidy by endorseme	ent		
Only if prescribed for dialysis or cystic fibrosis patient and th	e prescription is endo	rsed	accordingl	y.
Inj 150 mg	65.00	1	1	Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement.	95.00	5	1	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection	and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement.	91.00	5	1	Wockhardt S29
	182.00	10	1	Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection	and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement.	17.50	10	1	Pfizer
	87.50	50	1	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	r trac	t infection	and the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Retai No patient co-payment payable	l pharmacy			
Tab 400 mg	42.00	5	✓	Avelox

⇒SA1740 Special Authority for Subsidy

Initial application — (**Tuberculosis**) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*: and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications: or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Fither:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and

	ii Lonono A	<u> </u>		OTOTE MIO GOL
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
continued				
3 Treatment is only for 7 days.				
Initial application — (Penetrating eye injury) only from an opht requires prophylaxis following a penetrating eye injury and treatments. Note: Indications marked with * are unapproved indications.	ent is for 5 days only		alid for 1 mo	onth where the patient
PAROMOMYCIN - Special Authority see SA1689 below - Retail				
Cap 250 mg	126.00	16	✓ F	lumatin S29
▶SA1689 Special Authority for Subsidy Initial application only from an infectious disease specialist, clinic month for applications meeting the following criteria: Either:	cal microbiologist or	gastro	enterologis	st. Approvals valid for 1
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 				
Renewal only from an infectious disease specialist, clinical microbapplications meeting the following criteria: Either:	piologist or gastroent	terolog	jist. Approv	vals valid for 1 month for
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 				
PYRIMETHAMINE - Special Authority see SA1328 below - Reta	il pharmacy			
Tab 25 mg		30	✓ [Daraprim (\$29)
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of	a period of 3 month		nless notifie	d for applications meeting
SODIUM FUSIDATE [FUSIDIC ACID] Tab 250 mg	67.85	36	√ F	ucidin
SULFADIAZINE SODIUM - Special Authority see SA1331 below Tab 500 mg		56	√ V	Vockhardt \$29
■ SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of	without further rene a period of 3 month	ewal ur		
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement		5	√ √	<u>obramycin Mylan</u> /iatris
Only if prescribed for dialysis or cystic fibrosis patient and	I the prescription is	endors	ed accordin	ngly.
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by endorsement	395.00 5	56 dos	e 🗸 <u>T</u>	obramycin BNM
a) Wastage claimableb) Only if prescribed for a cystic fibrosis patient and the p	orescription is endor	sed ac	cordingly.	
TRIMETHOPRIM	10.55	-	, -	·MD
* Tab 300 mg - Up to 30 tab available on a PSO	18.55	50	✓ Ī	<u>WP</u>

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	AZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – I to 30 tab available on a PSO		500	✓ <u>T</u> I	risul
Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 available on a PSO		100 ml	✓ D	eprim
VANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or for difficile following metronidazole failure and the prescription is Inj 500 mg vial	endorsed according			ment of Clostridium

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 61
- b) For topical antifungals refer to GENITO URINARY, page 74

FLUCONAZOLE

Cap 50 mg	2.75	28	✓ Dizole✓ Mylan
Cap 150 mg	0.65	1	✓ Mylan
Cap 200 mg		28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority			-
see SA1359 below – Retail pharmacy Wastage claimable	109.34	35 ml	✓ Diflucan

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg4.27	15	✓ Itrazole
Oral liq 10 mg per ml - Special Authority see SA1322 on the		
next page – Retail pharmacy141.80	150 ml OP	✓ Sporanox

((Subsidy Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
<u></u>	\$	Per	✓	Manufacturer

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCT	CBS	30	✓ Link Healthcare S29
			✓ Nizoral S29
		100	✓ Strides Shasun S29
			✓ Taro S29
(Link Healthcare S29 Tab 200 mg to be delisted 1 July 2023)			
(Nizoral S29 Tab 200 mg to be delisted 1 July 2023)			
(Strides Shasun S29 Tab 200 mg to be delisted 1 July 2023)			
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Re	tail pharmacy		
Tab modified-release 100 mg	206.00	24	✓ Posaconazole Juno
•	869.86		✓ Noxafil
Oral liq 40 mg per ml	342.51	105 ml OP	✓ Devatis
· · · · · · · · · · · · · · · · · · ·	761.13		✓ Noxafil

(Noxafil Tab modified-release 100 mg to be delisted 1 April 2023) (Noxafil Oral lig 40 mg per ml to be delisted 1 May 2023)

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

*	Tab 250 mg8.	.15 8	34 🗸	Deolate 2

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	d Generic
VORICONAZOLE - Special Authority see SA1273 below - Retail	l pharmacy			
Tab 50 mg	91.00	56	✓	Vttack
Tab 200 mg	350.00	56	✓	Vttack
Powder for oral suspension 40 mg per ml - Wastage				
claimable	1,523.22	70 m	· •	Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised: and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient is immunocompromised; and
 - 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
 - 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE - Special Authority see SA1684 below - Retail	pharmacy		
Tab 15 mg	400.00	100	✓ Sanofi
			Primaguine \$29

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

- Both:
 - 1 The patient has vivax or ovale malaria; and
 - 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ <u>Metrogyl</u>
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl

	INI ECTIONS - A	<u> </u>		1 3131EWIIC USE
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
ORNIDAZOLE Tab 500 mg	00.10	10	,	Away Owidanala
Tab 500 mg	36.16	10	•	Arrow-Ornidazole
Antituberculotics and Antileprotics				
Note: There is no co-payment charge for all pharmaceuticals	listed in the Antitubercul	otics	and Antile	protics group regardless of
immigration status.				h
CLOFAZIMINE - Retail pharmacy-Specialist				
a) No patient co-payment payable				
 b) Prescriptions must be written by, or on the recommen dermatologist. 		iseas	e physicia	n, clinical microbiologist or
* Cap 50 mg	442.00	100	•	Lamprene S29
CYCLOSERINE - Retail pharmacy-Specialist				
a) No patient co-payment payable				
 b) Prescriptions must be written by, or on the recommen respiratory physician. 	dation of, an infectious d	iseas	e physicia	n, clinical microbiologist or
Cap 250 mg	344.00	60	✓	Cyclorin S29
DAPSONE - Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommen	dation of, an infectious d	iseas	e physicia	n, clinical microbiologist or
dermatologist				
Tab 25 mg		100		Dapsone
Tab 100 mg		100	•	Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Spec	ialist			
a) No patient co-payment payable				
 Prescriptions must be written by, or on the recommen respiratory physician 		iseas	e physicia	n, clinical microbiologist or
Tab 100 mg	85.73	100	/	EMB Fatol S29
Tab 400 mg	49.34	56	•	Myambutol S29
ISONIAZID - Retail pharmacy-Specialist				
 a) No patient co-payment payable 				
b) Prescriptions must be written by, or on the recommen		dicine	e physiciar	ı, paediatrician, clinical
microbiologist, dermatologist or public health physician		400	,	2011
* Tab 100 mg		100	•	<u>PSM</u>
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist				
No patient co-payment payable				
b) Prescriptions must be written by, or on the recommen		dicine	e pnysiciar	i, paediatrician, clinical
microbiologist, dermatologist or public health physicial * Tab 100 mg with rifampicin 150 mg		100	_	Rifinah
* Tab 100 mg with rifampicin 300 mg		100	_	Rifinah
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialis		100	•	<u> </u>
	OL .			
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommen	dation of, an infectious d	iseas	e specialis	st, clinical microbiologist or
respiratory physician	addon on, an imodicad a	10040	o opoolane	n, omnour moroprorogict or
Grans for oral liq 4 g sachet	280.00	30	1	Paser S29
PROTIONAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommen	dation of, an infectious d	iseas	e specialis	st, clinical microbiologist or
respiratory physician			•	ŭ
Tab 250 mg	305.00	100	✓	Peteha S29
▲ Three months supply may be dispensed at one time if endorsed "co				

99

		Subsidy (Manufactured a Prior	۰\	Fully	Brand or
		(Manufacturer's Price \$	e) S Per	Subsidised	Generic Manufacturer
PY	RAZINAMIDE - Retail pharmacy-Specialist	-			
	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommendati	on of, an infectious	disease	physician	, clinical microbiologist or
*	respiratory physician	64.05	100	.1	A ET_Durazinamida
	Tab 500 mg	04.93	100	•	AFT-Pyrazinamide
HIF	ABUTIN – Retail pharmacy-Specialist a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommendati	on of, an infectious	disease	physician	, respiratory physician or
	gastroenterologist			. ,	, , ,,,
*	Cap 150 mg	299.75	30	✓ I	Mycobutin
RIF	AMPICIN – Subsidy by endorsement				
	a) No patient co-payment payable	in combination with	athar of	factive on	ti atanbulaaaaal
	 For confirmed recurrent Staphylococcus aureus infection antimicrobial based on susceptibilities and the prescription 				
	Retail pharmacy - Specialist. Specialist must be an interr				
	paediatrician, or public health physician.			_	
*	Cap 150 mg		100	-	Rifadin Difadin
*	Cap 300 mg Oral liq 100 mg per 5 ml		100 60 ml		<u>Rifadin</u> Rifadin
*	Cital liq 100 mg per 5 mi	12.00	00 1111	• !	<u>Illiaulli</u>
A	ntivirals				
Гол	aug proporations refer to Fue Drangrations. Anti-Infective Dra	narationa naga 04	2		
	eye preparations refer to Eye Preparations, Anti-Infective Pre	parations, page 24	J		
Н	epatitis B Treatment				
	TECAVIR				
*	Tab 0.5 mg	52.00	30		Entecavir Mylan
	ADVIDING O CLASS O CASOCILL D. C.			• 1	Entecavir Sandoz
LAI	MIVUDINE - Special Authority see SA1685 below - Retail ph. Tab 100 mg		28	1	Zetlam
	Oral lig 5 mg per ml		20 240 ml O		Zeffix
>	SA1685 Special Authority for Subsidy				
	ial application only from a relevant specialist or medical prac	titioner on the reco	mmenda	tion of a re	elevant specialist.
	provals valid for 1 year where used for the treatment or preven				
	newal from any relevant practitioner. Approvals valid for 2 year	ars where used for	the treat	ment or pr	evention of hepatitis B.
ΙĿΙ	NOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for the tre	aatment of HIV is in	odudad ir	the coun	t of up to 1 subsidised
	antiretrovirals for the purposes of Special Authority SA2139.,		iciuu c u ii	i iiie couri	t of up to 4 subsidised
*	Tab 245 mg (300 mg as a maleate)		30	1	Tenofovir Disoproxil
					<u>Mylan</u>
Н	erpesvirus Treatments				
	·				
	CLOVIR	1.70	0.5		avi.
* *	Tab dispersible 200 mg Tab dispersible 400 mg		25 56		Lovir Lovir
*	Tab dispersible 800 mg		35		Lovir
	ACICLOVIR				
	Tab 500 mg	6.50	30	✓ 1	/aclovir
	Tab 1,000 mg	13.76	30	✓ <u>\</u>	<u>Vaclovir</u>

	Subsidy (Manufacturer's Price)	Sub Per	Fully bsidised	Brand or Generic Manufacturer
VALGANCICLOVIR – Special Authority see SA1993 below – F Tab 450 mg	' '	60	_	alganciclovir Mvlan

⇒SA1993 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Roth:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Roth:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Fither:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
- 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below No patient co-payment payable

28 Tab 90 mg with sofosbuvir 400 mg.......24,363.46 ✓ Harvoni

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/mayiret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac. PO Box 10-254. WELLINGTON Tel: (04) 460 4990.

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV. prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 103 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a

maleate).......15.45 30 ✓ Tenofovir Disoproxil **Emtricitabine**

Mylan

⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion: and

Subsidy		Fully	Brand or
(Manufacturer's F	Price) Su	ubsidised	Generic
\$	Per	1	Manufacturer

continued...

2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical quidelines:

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose

Subsidy		Fully	Brand or
(Manufacturer's Price)		osidised	Generic
	Per	✓	Manufacturer

continued...

ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (**Percutaneous exposure**) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA2139 on the previous p	age – Retail pharm	nacy		
Tab 200 mg	190.15	90	✓ Stocrin	
Tab 600 mg	63.38	30	✓ Stocrin	
ETRAVIRINE - Special Authority see SA2139 on the previous	page – Retail pharr	macy		
Tab 200 mg	770.00	60	✓ Intelence	

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully Brand or idised Generic Manufacturer
EVIRAPINE – Special Authority see SA2139 on page 103 – I Tab 200 mg		60	✓ <u>Nevirapine</u> Alphapharm
Oral suspension 10 mg per ml	203.55	240 ml OP	Nevirapine ViatrisViramuneSuspension
Nucleosides Reverse Transcriptase Inhibitors			
BACAVIR SULPHATE - Special Authority see SA2139 on pa		harmacy	
Tab 300 mg		60	✓ Ziagen
Oral liq 20 mg per ml		240 ml OP	✓ Ziagen
BACAVIR SULPHATE WITH LAMIVUDINE – Special Author Note: abacavir with lamivudine (combination tablets) coun anti-retroviral Special Authority.	ts as two anti-retro		ns for the purposes of the
Tab 600 mg with lamivudine 300 mg	29.50	30	✓ Abacavir/ Lamivudine Viatris
	63.00		✓ Kivexa
Kivexa Tab 600 mg with lamivudine 300 mg to be delisted 1 M	lay 2023)		
narmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disopr 245 mg (300 mg as a maleate)	roxil	inti-retroviral me 30	dications for the purposes of t Mylan
MTRICITABINE – Special Authority see SA2139 on page 100		•	/ Fundadora
Cap 200 mg		30	✓ Emtriva
AMIVUDINE - Special Authority see SA2139 on page 103 -	Rotail nharmacu		
Tab 150 mg	, ,	60	✓ <u>Lamivudine</u> <u>Alphapharm</u>
, , , , , , , , , , , , , , , , , , , ,	84.50	60 240 ml OP	
Tab 150 mg		240 ml OP	Alphapharm
Tab 150 mg Oral liq 10 mg per ml IDOVUDINE [AZT] — Special Authority see SA2139 on page		240 ml OP macy	Alphapharm ✓ 3TC
Tab 150 mg Oral liq 10 mg per ml		240 ml OP macy 100 200 ml OP ge 103 – Retail	Alphapharm ✓ 3TC ✓ Retrovir ✓ Retrovir pharmacy
Tab 150 mg Oral liq 10 mg per ml IDOVUDINE [AZT] — Special Authority see SA2139 on page Cap 100 mg Oral liq 10 mg per ml IDOVUDINE [AZT] WITH LAMIVUDINE — Special Authority s		240 ml OP macy 100 200 ml OP ge 103 – Retail	Alphapharm ✓ 3TC ✓ Retrovir ✓ Retrovir pharmacy
Tab 150 mg		240 ml OP macy 100 200 ml OP ge 103 – Retail anti-retroviral m	Alphapharm 3TC Retrovir Retrovir pharmacy ledications for the purposes of
Tab 150 mg		240 ml OP macy 100 200 ml OP ge 103 – Retail anti-retroviral m	Alphapharm 3TC Retrovir Retrovir pharmacy ledications for the purposes of
Tab 150 mg		240 ml OP macy 100 200 ml OP ge 103 – Retail anti-retroviral m	Alphapharm ✓ 3TC ✓ Retrovir ✓ Retrovir pharmacy nedications for the purposes of ✓ Alphapharm ✓ Atazanavir Mylan
Oral liq 10 mg per ml		240 ml OP macy 100 200 ml OP ge 103 – Retail anti-retroviral m 60 il pharmacy 60	Alphapharm 3TC Retrovir Retrovir pharmacy nedications for the purposes of Alphapharm Atazanavir Mylan Teva
Tab 150 mg		240 ml OP macy 100 200 ml OP ge 103 – Retail anti-retroviral m 60	Alphapharm ✓ 3TC ✓ Retrovir ✓ Retrovir pharmacy nedications for the purposes of ✓ Alphapharm ✓ Atazanavir Mylan

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	Fully Brand or ised Generic Manufacturer
DARUNAVIR – Special Authority see SA2139 on page 103 – Re Tab 400 mg Tab 600 mg	132.00	60 60	✓ <u>Darunavir Mylan</u> ✓ <u>Darunavir Mylan</u> ✓ <u>Darunavir Viatris</u>
LOPINAVIR WITH RITONAVIR – Special Authority see SA2139 Tab 100 mg with ritonavir 25 mg		pharmacy 60	✓ <u>Lopinavir/Ritonavir</u> Mylan
Tab 200 mg with ritonavir 50 mg	295.00	120	✓ Lopinavir/Ritonavir Mylan
Oral liq 80 mg with ritonavir 20 mg per ml		0 ml OP	✓ Kaletra
Tab 100 mg		30	✓ Norvir

Strand Transfer Inhibitors

DOLUTEGRAVIR - Special Authority see SA2139 on page 103	- Retail pharmacy		
Tab 50 mg	1,090.00	30	✓ Tivicay
RALTEGRAVIR POTASSIUM - Special Authority see SA2139	on page 103 – Retail p	harmacy	
Tab 400 mg	1,090.00	60	✓ Isentress
Tab 600 mg	1,090.00	60	✓ Isentress HD

Immune Modulators

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the

Subsidy (Manufacturer's Pric	e)	Fully Subsidised	Brand or Generic	
\$	Per	· •	Manufacturer	

continued...

following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	bsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either
 - 3.1 Patient has a cutaneous T cell lymphoma*: or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Initial application — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE * Tab 1 g	19.95	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg - Up to 15 cap available on a			
PSO	86.40	100	✓ Macrobid
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement		100	

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per	✓	Manufacturer
And Make Proceedings				
Anticholinesterases				
EOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	√ i	Max Health
YRIDOSTIGMINE BROMIDE				
Tab 60 mg	45.79	100	√ I	Mestinon
Non-Steroidal Anti-Inflammatory Drugs				
ICLOFENAC SODIUM				
* Tab EC 25 mg	1 99	50	√ [Diclofenac Sandoz
Tab 50 mg dispersible		20	_	Voltaren D
Tab EC 50 mg		50		Diclofenac Sandoz
Tab long-acting 75 mg		100	_	Voltaren SR
Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a F		5		Voltaren Voltaren
Suppos 12.5 mgSummanipodie — Op to 3 inj available on a r		10		Voltaren
Suppos 25 mg		10		Voltaren
Suppos 50 mg - Up to 10 supp available on a PSO		10		Voltaren
Suppos 100 mg		10		Voltaren
	7.00	10	• '	VOILAIEII
UPROFEN				
Tab 200 mg		1,000	_	Relieve
Tab long-acting 800 mg		30	_	Brufen SR
Oral liq 20 mg per ml		200 m		<u>Ethics</u>
	11.29		✓ [Fenpaed 100 mg per 5 ml
ETOPROFEN				·
Cap long-acting 200 mg	12.07	28	✓ (Oruvail SR
EFENAMIC ACID				
	1.05	E0		
Cap 250 mg		50		Dt
	(9.16)	20	ı	Ponstan
	0.50	20		Damatan
ADDOVEN	(7.50)		ŀ	Ponstan
APROXEN Tab 250 mg	32 69	500	1	Noflam 250
Tab 500 mg		250	_	Noflam 500
Tab long-acting 750 mg		28	_	Naprosyn SR 750
Tab long-acting 1 g		28	_	Naprosyn SR 1000
0 0 0		20	٠ ١	Tapiooyii oii 1000
ENOXICAM	40.50	400	, -	T11 411
Tab 20 mg		100	_	<u> Tilcotil</u>
Inj 20 mg vial	9.95	1	•	AFT
NSAIDs Other				
ELECOXIB				
Cap 100 mg	3.45	60		Celebrex
			_	Celecoxib Pfizer
Cap 200 mg	3.20	30		Celebrex
			✓ (Celecoxib Pfizer

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	/	Manufacturer	

Topical Products for Joint and Muscular Pain

CAPSAICIN

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary)*, and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroguine. Note: Indication marked with a * is an unapproved indication.

* Tab 200 mg			✓ Plaquenil
LEFLUNOMIDE			
Tab 10 mg	6.00	30	✓ Arava
Tab 20 mg	6.00	30	✓ Arava
PENICILLAMINE			
Tab 125 mg	67.23	100	D-Penamine
Tab 250 mg	110.12	100	D-Penamine

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODIUM			
* Tab 70 mg	.2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL			
* Tab 70 mg with colecalciferol 5,600 iu	.1.51	4	✓ Fosamax Plus

Other Treatments

DENOSUMAB – Special Authority see SA1777 below – R	letail pharmacy		
Inj 60 mg prefilled syringe	326.00	1	Prolia

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or

continued...

- 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	27.53	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial		1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial		1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA1	779 below – Retail	pharmacy	
* Tab 60 mg	53.76	28	✓ Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Tab 35 mg		4	•	Risedronate Sandoz
ΓERIPARATIDE – Special Authority see SA1139 on the next page -	- Retail pharma	acy		
Inj 250 mcg per ml, 2.4 ml	490.00	1	1	Forteo

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, bag – Special Authority see SA2110 on page 115 – Retail pharmacy	. 22.53	100 ml OP	✓ Zoledronic Acid Viatris
Inj 0.05 mg per ml, 100 ml, vial – Special Authority see SA2110 below – Retail pharmacy	. 60.00	100 ml OP	✓ Aclasta

⇒SA2110 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Any of the following:

- 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
- 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	1	Manufacturer

continued...

densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or

- 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It

Subsidy (Manufacturer's Price)	•		Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- is unlikely that this provision would apply to many patients under 75 years of age; or
- 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (spinal cord injury*) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has experienced an acute traumatic spinal cord injury in the last six months; and
- 2 Patient is being managed by a specialist spinal acute care and rehabilitation unit; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Note: Indications marked with * are unapproved indications.

Renewal — (spinal cord injury*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period; and
- 2 The patient has not received more than two doses of zoledronic acid for this indication.

The patient must not have had more than 1 prior approval.

Notes: No further renewals will be subsidised. A maximum of 2 vials of zoledronic acid treatment for spinal cord injury will be subsidised. Indications marked with * are unapproved indications.

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

⇒SA2110 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity: or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or

Subsidy (Manufacturer's Price)	Fı Subsidis	ully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 2.5 Preparation for orthopaedic surgery; and
 - 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

,		Fully	Brand or Generic		
\$	Per	1	Manufacturer		

continued...

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (**spinal cord injury***) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has experienced an acute traumatic spinal cord injury in the last six months; and
- 2 Patient is being managed by a specialist spinal acute care and rehabilitation unit; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Note: Indications marked with * are unapproved indications.

Renewal — (spinal cord injury*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period; and
- 2 The patient has not received more than two doses of zoledronic acid for this indication.

The patient must not have had more than 1 prior approval.

Notes: No further renewals will be subsidised. A maximum of 2 vials of zoledronic acid treatment for spinal cord injury will be subsidised. Indications marked with * are unapproved indications.

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

	(Manufacturer's Price)	Per	⊅sidised ✓	Manufacturer
Hyperuricaemia and Antigout				
ALLOPURINOL * Tab 100 mg	11.47	500	√ D	P-Allopurinol
* Tab 300 mg		500	_	P-Allopurinol
BENZBROMARONE - Special Authority see SA1963 below -	Retail pharmacy			
Tab 50 mg	22.50	100	✓ N	arcaricin mite S29
Tab 100 mg	13.50	30	✓ D	esuric S29

Subsidy

45.00

100

Fully

Brand or

✓ Urinorm S29

100 S29

✓ Benzbromaron AL

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg	6.00	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA2054 below - Retail pharma	асу		
Tab 80 mg	20.00	28	✓ Febuxostat
			multichem
Tab 120 mg	20.00	28	✓ <u>Febuxostat</u>
			multichem

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

PROBENECID

100

✓ Norflex

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Muscle Relaxants				
BACLOFEN				
* Tab 10 mg	4.20	100	✓ P	acifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsemer	ıt11.55	1	√ L	ioresal Intrathecal
Subsidised only for use in a programmable pump in pati		oastic ag	ents have	e been ineffective or have
caused intolerable side effects and the prescription is er	0,			
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement		5	_	<u>ledsurge</u>
Subsidised only for use in a programmable pump in pat caused intolerable side effects and the prescription is er		oastic ag	ents hav	e been ineffective or have
DANTROLENE				
Cap 25 mg	97.50	100	✓ D)antrium
			✓ D	Oantrium S29 S29
Cap 50 mg	77.00	100	✓ D)antrium
ORPHENADRINE CITRATE				

Tab 100 mg20.76

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic

\$ Per ✔ Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE		
▲ Cap 100 mg	60	 Symmetrel
63.73	100	Symmetrel
APOMORPHINE HYDROCHLORIDE		
▲ Inj 10 mg per ml, 2 ml ampoule	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule121.84	5	✓ Movapo
ENTACAPONE		
▲ Tab 200 mg18.04	100	✓ Comtan
LEVODOPA WITH BENSERAZIDE		
* Tab dispersible 50 mg with benserazide 12.5 mg13.25	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg13.75	100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg15.80	100	Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg22.85	100	Madopar HBS
* Cap 200 mg with benserazide 50 mg26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA		
* Tab 100 mg with carbidopa 25 mg21.11	100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg43.65	100	✓ Sinemet CR
* Tab 250 mg with carbidopa 25 mg38.39	100	✓ Sinemet
PRAMIPEXOLE HYDROCHLORIDE		
▲ Tab 0.25 mg	100	✓ Ramipex
▲ Tab 1 mg18.66	100	✓ Ramipex
RASAGILINE		
* Tab 1 mg53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE		
▲ Tab 0.25 mg	84	✓ Ropin
▲ Tab 1 mg	84	Ropin
▲ Tab 2 mg	84	Ropin
▲ Tab 5 mg14.50	84	✓ Ropin

SELEGILINE HYDROCHLORIDE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of selegiline hydrochloride.

	prescription is chaorsed accordingly. I flamfacists may afficiate the	prescription	is chacksca	WHERE THERE CAISIS & record to
	prior dispensing of selegiline hydrochloride.			
*	Tab 5 mg48	8.00	100	✓ Eldepryl S29
TO	LCAPONE			

Anticholinergics

BENZATROPINE MESYLATE			
Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
a) Up to 10 ini available on a PSO			

b) Only on a PSO

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg7.40 100 ✓ Kemadrin

100

✓ Tasmar

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE - Special Authority see SA1403 below - Retail pharmacy

Wastage claimable

⇒SA1403 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Anaesthetics

Local

LIDOCAINE [LIGNOCAINE]

- a) Up to 150 ml available on a PSO
- b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

- a) Up to 5 each available on a PSO
- Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.

	Subsidy (Manufacturer's Price			Generic
	\$	Per		Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 ml	1	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	9.50	25	1	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25	25	1	Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20	5	1	Lidocaine-Baxter
			1	Lidocaine-Claris
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO(Lidocaine-Claris Inj 1%, 20 ml vial to be delisted 1 June 2023)	6.45	5	1	Lidocaine-Baxter
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –	100.00	40	,	D#
Subsidy by endorsement	103.32	10	•	Pfizer
 a) Up to 5 each available on a PSO 				

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority se	ee SA0906 above – Retail pharr	nacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE -	- Special Authority see SA0906	above - Reta	ail pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

Analgesics

A ODIDIN

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 109

Non-opioid	Analgesics
------------	-------------------

ASPIRIN * Tab dispersible 300 mg - Up to 30 tab available on a PSO	4.50	100	✓ Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or diabe accordingly.	tic periphera	al neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.14	57 g OP	✓ Rugby Capsaicin Topical Cream ^{©29}
NEFOPAM HYDROCHLORIDE Tab 30 mg	23.40	90	✓ Acupan

NERVOUS SYSTEM

_					
		Subsidy	` .	Fully	Brand or
		(Manufacturer's Pri		sidised	Generic
_		\$	Per		Manufacturer
PA	RACETAMOL				
	Tab 500 mg - blister pack	19.75	1,000	√ P	acimol
	a) Maximum of 300 tab per prescription; can be waiv		1,000	-	
	b) Up to 30 tab available on a PSO	ca by chaolochich			
	c)				
	Subsidy by endorsement for higher quantitie	e ie available for nati	onte with lone	torm o	anditions who require
	regular daily dosing for one month or greate				
	annotate the prescription as endorsed where				
	2) Maximum of 100 tab per dispensing for non-				
	(for non-endorsed patients), then dispense in	repeat dispensings	not exceedin	g 100 ta	ab per dispensing.
	Tab 500 mg - bottle pack - Maximum of 300 tab per	47.00	4 000		
	prescription; can be waived by endorsement	17.92	1,000	→ V	loumed
					<u>Paracetamol</u>
	 Subsidy by endorsement for higher quantities is 	available for patients	s with long ter	m cond	litions who require regular
	daily dosing for one month or greater, and the p	rescription is annotat	ed according	ly. Pha	rmacists may annotate the
	prescription as endorsed where dispensing histo	ory supports a long-te	erm condition		
	Maximum of 100 tab per dispensing for non-end	lorsed patients. If qu	antities preso	ribed fo	or more than 100 tabs (for
	non-endorsed patients), then dispense in repeat				
	, , , , , , , , , , , , , , , , , , ,	3 · · · ·	3		-
	Oral lig 120 mg per 5 ml	3 98	200 ml	√ P	aracetamol
			200		(Ethics)
		5.45	1 000 ml	./ 0	aracare
			1,000 ml		raracare Avallon
		10.50	200 ml OP	• 4	wallon
	 a) Maximum of 600 ml per prescription; can be waive 	ed by endorsement			
	b) Up to 200 ml available on a PSO				
	c) Not in combination				
	d)				
	 Maximum of 200 ml per dispensing for non-e 				
	non-endorsed patients), then dispense in rep				
	Subsidy by endorsement for higher quantitie	s is available for pation	ents with long	term c	onditions who require
	regular daily dosing for one month or greater	r and the prescription	is endorsed	or anno	tated accordingly.
	Pharmacists may annotate the prescription a	as endorsed where di	spensing hist	ory sup	ports a long-term
	condition.				
	Oral liq 250 mg per 5 ml	3.35	200 ml	✓ P	amol
		6.25	1,000 ml	✓ P	aracare Double
			,		Strength
	a) Maximum of 600 ml per prescription; can be waive	ad hy andorsament			g
	b) Up to 200 ml available on a PSO	od by ondoroomone			
	c) Not in combination				
	d)				
	,	andorood notionto If	auantitica pr	ooribo	d avaged 200 ml (for
	1) Maximum of 200 ml per dispensing for non-e				,
	non-endorsed patients), then dispense in rep				
	2) Subsidy by endorsement for higher quantitie			•	•
	regular daily dosing for one month or greater				
	Pharmacists may annotate the prescription a	as endorsed where di	spensing his	ory sup	ports a long-term
	condition.		46		
	Suppos 125 mg		10		iacet
	Suppos 250 mg		10		acet
*	Suppos 500 mg	12.40	50	√ (lacet
(Pa	aracare Oral liq 120 mg per 5 ml to be delisted 1 June 2023)			

(Paracare Double Strength Oral liq 250 mg per 5 ml to be delisted 1 April 2023)

	Subsidy		Fully Brand or
	(Manufacturer's Price)	Per	sidised Generic Manufacturer
Opioid Analgesics			
CODEINE PHOSPHATE – Safety medicine; prescriber may	determine dispensing fro	aniency	
Tab 15 mg	, ,	100	✓ Noumed
140 10 mg	6.25	100	✓ PSM
Tab 30 mg		100	✓ Aspen
			✓ Noumed
	7.45		✓ PSM
Tab 60 mg	13.89	100	✓ Noumed
	14.25		✓ PSM
(PSM Tab 15 mg to be delisted 1 May 2023)			
(PSM Tab 30 mg to be delisted 1 April 2023)			
(PSM Tab 60 mg to be delisted 1 April 2023)			
DIHYDROCODEINE TARTRATE			
Tab long-acting 60 mg	8.60	60	✓ DHC Continus
FENTANYL			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensin	g frequency		
Inj 50 mcg per ml, 2 ml ampoule		10	 Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule	9.41	10	✓ Boucher and Muir
Patch 12.5 mcg per hour	6.99	5	✓ Fentanyl Sandoz
Patch 25 mcg per hour	7.99	5	✓ Fentanyl Sandoz
Patch 50 mcg per hour		5	✓ Fentanyl Sandoz
Patch 75 mcg per hour		5	✓ Fentanyl Sandoz
Patch 100 mcg per hour	18.59	5	✓ Fentanyl Sandoz
METHADONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
 c) Safety medicine; prescriber may determine dispensing 			
d) Extemporaneously compounded methadone will only	be reimbursed at the rat	te of the ch	neapest form available
(methadone powder, not methadone tablets).			
e) For methadone hydrochloride oral liquid refer Standa		10	/ Mathataha
Tab 5 mg	1.40 1.45	10	✓ Methatabs✓ Methadone BNM
Oral lig 2 mg per ml		200 ml	✓ Methadone BNM ✓ Biodone
Oral liq 5 mg per ml		200 ml	✓ Biodone Forte
Oral liq 10 mg per ml		200 ml	✓ Biodone Extra Forte
Inj 10 mg per ml, 1 ml		10	✓ AFT
(Methatabs Tab 5 mg to be delisted 1 February 2023)			
MORPHINE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing	a frequency		
Oral lig 1 mg per ml		200 ml	✓ RA-Morph
Oral liq 2 mg per ml		200 ml	✓ RA-Morph
Oral liq 5 mg per ml		200 ml	✓ Ordine S29
70			✓ RA-Morph
Oral liq 10 mg per ml	27.74	200 ml	✓ Ordine S29
1 - 91-			✓ RA-Morph
			r

Subsidy

Fully

Brand or

				· ·
	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	· ·	Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fi	requency			
Tab immediate-release 10 mg		10	✓	Sevredol
Tab immediate-release 20 mg		10		Sevredol
Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 30 mg		10		m-Eslon
Cap long-acting 60 mg		10		m-Eslon
Cap long-acting 100 mg		10	1	m-Eslon
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a F		5	1	Medsurge
., p	6.99			DBL Morphine
				Sulphate
Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO 4.68	5	1	Medsurge
ing to mg por mi, i mi ampoule — op to o mg available on a	5.61	Ü		DBL Morphine
	0.01		•	Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO 5.53	5	1	Medsurge
ing 13 mg per mi, 1 mi ampoule — op to 3 mg available on a	7.08	5		DBL Morphine
	7.00		•	Sulphate
Ini 20 mg normi 1 ml amnaula . Un ta E ini available an a	DCO 6.00	-	./	Medsurge
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a	7.28	5		
	1.20		•	DBL Morphine Sulphate
(DBL Morphine Sulphate Inj 5 mg per ml, 1 ml ampoule to be de (DBL Morphine Sulphate Inj 10 mg per ml, 1 ml ampoule to be de (DBL Morphine Sulphate Inj 15 mg per ml, 1 ml ampoule to be de (DBL Morphine Sulphate Inj 30 mg per ml, 1 ml ampoule to be de	lelisted 1 March 2023) Ielisted 1 March 2023)			
, , , , , , , , , , , , , , , , , , , ,	onotod i maion 2020)			
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing for		20	./	Ovvendene Canden
Tab controlled release 5 mg		20 20		Oxycodone Sandoz
Tab controlled-release 10 mg Tab controlled-release 20 mg		20		Oxycodone Sandoz Oxycodone Sandoz
3		20		Oxycodone Sandoz
Tab controlled release 40 mg				
Tab controlled-release 80 mg		20 20		Oxycodone Sandoz
Cap immediate-release 5 mg		20		OxyNorm OxyNorm
Cap immediate release 10 mg		20		OxyNorm OxyNorm
Cap immediate-release 20 mg		250 n		OxyNorm
Oral liq 5 mg per 5 ml		250 II 5		Hameln
Inj 10 mg per ml, 1 ml ampoule		5 5		Hameln
Inj 50 mg per ml, 2 ml ampoule		5 5		Hameln
, , , , , , , , , , , , , , , , , , , ,				
PARACETAMOL WITH CODEINE – Safety medicine; prescribe				
* Tab paracetamol 500 mg with codeine phosphate 8 mg	27.50	1,000	/	Paracetamol +

Codeine (Relieve)

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensin	g frequency			
Tab 50 mg		10		PSM
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available or	n a PSO29.88	5	•	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available or	n a PSO30.72	5	/	DBL Pethidine Hydrochloride
FRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg	1.52	20	1	Tramal SR 100
Tab sustained-release 150 mg	2.10	20	✓	Tramal SR 150
Tab sustained-release 200 mg	2.75	20	✓	Tramal SR 200
Cap 50 mg	2.80	100	✓	Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine	ne dispensing frequency			
Tab 10 mg		100	1	Arrow-Amitriptyline
Tab 25 mg	1.51	100	1	Arrow-Amitriptyline
Tab 50 mg	2.51	100	1	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; pre	escriber may determine d	ispen	sina freau	encv
Tab 10 mg		30		Clomipramine Teva
		30		Clomipramine Teva
1 au 2J 1114		30	•	Cionniprannile reva
Tab 25 mg DOSUIL EPIN IDOTHIEPINI HYDROCHI ORIDE — Subsidy b		30	•	Cionipramine reva
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by	y endorsement	30	•	Ciompramme reva
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by a) Safety medicine; prescriber may determine dispensin	y endorsement g frequency			
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement – Subsidised for patients wh	y endorsement g frequency no were taking dosulepin	[doth	epin] hydr	ochloride prior to 1 June
a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement – Subsidised for patients where 2019 and the prescription is endorsed accordingly.	y endorsement g frequency no were taking dosulepin Pharmacists may annotate	[doth	epin] hydr	ochloride prior to 1 June
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride.	[doth the	epin] hydr orescriptio	ochloride prior to 1 June n as endorsed where ther
a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement – Subsidised for patients where 2019 and the prescription is endorsed accordingly.	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride.	[doth	epin] hydr orescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan
Safety medicine; prescriber may determine dispensin Safety medicine; prescriber may determine dispensin Subsidy by endorsement – Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothing the content of the	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride. 3.85	[doth the	epin] hydr orescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride. 3.85	[doth the	epin] hydr orescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin
Safety medicine; prescriber may determine dispensin Safety medicine; prescriber may determine dispensin Subsidy by endorsement – Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothing the content of the	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride. 3.85	[doth the	epin] hydr orescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride	[doth the 30 50	epin] hydr orescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride3.857.83 iber may determine dispe	[doth the 30 50	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride3.857.83 liber may determine dispe5.48	[doth the 30 50 nsing	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride3.857.83 liber may determine dispe5.48 10.96	[doth the 30 50 nsing 50 100	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 // Tofranil Tofranil
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride	30 50 nsing 50 100 50	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 Tofranil Tofranil Tofranil
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothic Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride3.857.83 iber may determine dispe5.48 10.968.80 rescriber may determine of	(doth) the 30 50 nsing 50 100 50	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 Tofranil Tofranil Tofranil Iency
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride3.857.83 iber may determine dispe5.48 10.968.80 rescriber may determine of	30 50 nsing 50 100 50	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 // Tofranil Tofranil Tofranil tency Norpress
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothic Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride	(doth) the 30 50 nsing 50 100 50	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 Tofranil Tofranil Tofranil Iency
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg	y endorsement g frequency no were taking dosulepin rharmacists may annotate epin] hydrochloride3.857.83 iber may determine dispe5.4880 rescriber may determine of	30 50 50 nsing 50 100 50 dispee	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 // Tofranil Tofranil Tofranil tency Norpress
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothic Tab 75 mg Cap 25 mg Cap 25 mg MIPRAMINE HYDROCHLORIDE — Safety medicine; prescri Tab 10 mg Tab 25 mg NORTRIPTYLINE HYDROCHLORIDE — Safety medicine; pr Tab 10 mg Tab 25 mg Tab 25 mg	y endorsement g frequency no were taking dosulepin rharmacists may annotate epin] hydrochloride3.857.83 iber may determine dispe5.4880 rescriber may determine of	30 50 50 nsing 50 100 50 dispee	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 // Tofranil Tofranil Tofranil tency Norpress
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride	30 50 50 nsing 50 100 50 dispee	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 // Tofranil Tofranil Tofranil tency Norpress
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023) MIPRAMINE HYDROCHLORIDE — Safety medicine; prescri Tab 10 mg Tab 25 mg NORTRIPTYLINE HYDROCHLORIDE — Safety medicine; pr Tab 10 mg Tab 25 mg Monoamine-Oxidase Inhibitors (MAOIs) - Noi	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride	30 50 50 100 50 4isper 180	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 / Tofranil Tofranil Tofranil tency Norpress Norpress
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE — Subsidy by a) Safety medicine; prescriber may determine dispensin b) Subsidy by endorsement — Subsidised for patients wh 2019 and the prescription is endorsed accordingly. P exists a record of prior dispensing of dosulepin [dothie Tab 75 mg	y endorsement g frequency no were taking dosulepin Pharmacists may annotate epin] hydrochloride	30 50 50 50 50 50 50 50 50 100 180	epin] hydrorescriptio	ochloride prior to 1 June n as endorsed where ther Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan \$29 / Tofranil Tofranil Tofranil tency Norpress Norpress

			INL	1V003 3131EW
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg	11.80	60	1	<u>Aurorix</u>
₭ Tab 300 mg	19.25	60	✓	<u>Aurorix</u>
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE				
★ Tab 20 mg	1.91	84	✓	PSM Citalopram
	2.86		✓	Celapram
PSM Citalopram Tab 20 mg to be delisted 1 March 2023)				
SCITALOPRAM				
★ Tab 10 mg	1.07	28	1	<u>Escitalopram</u>
				(Ethics)
★ Tab 20 mg	1.92	28	✓	Escitalopram
				(Ethics)
LUOXETINE HYDROCHLORIDE				
★ Tab dispersible 20 mg, scored – Subsidy by endorsement.	2.50	28	✓	Fluox
When prescribed in a daily dose that is not a multi- endorsed. Note: Tablets should be combined with				
Cap 20 mg	2 91	84	/	Fluox
ουρ 20 mg	3.13	90		Arrow-Fluoxetine
Fluox Cap 20 mg to be delisted 1 June 2023)	00	•	•	and a reconstant
PAROXETINE				
# Tab 20 mg	4 11	90	1	Loxamine
-		50	•	LOXAIIIIIC
ERTRALINE ₭ Tab 50 mg	0.00	30	1	Setrona
r Tab 50 Hig	0.99	30		Setrona AU
₭ Tab 100 mg	1 74	30		Setrona
- 100 mg		00		Setrona AU
Setrona AU Tab 50 mg to be delisted 1 April 2023) Setrona AU Tab 100 mg to be delisted 1 April 2023)				
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2.60	28	1	Noumed
Tab 45 mg		28		Noumed
/ENLAFAXINE				
K Cap 37.5 mg	6.38	84	1	Enlafax XR
V 0 75	0.44	04	,	Lilialax XII

84

84

✓ Enlafax XR
✓ Enlafax XR

Cap 75 mg......8.11

* Cap 150 mg......11.16

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Antiepilepsy Drugs

Agents for Control of Status Epilepticus

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement23.66	5	✓ Hospira
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
 c) PSO must be endorsed "not for anaesthetic procedures". 		
Rectal tubes 5 mg - Up to 5 tube available on a PSO54.58	5	Stesolid
PHENYTOIN SODIUM		
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a		
PSO104.58	5	Hospira
* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a		
PSO154.01	5	Hospira

Control of Epilepsy

CARBAMAZEPINE

* Tab 200 mg	14.53	100	✓ Tegretol
* Tab long-acting 200 mg		100	✓ Tegretol CR
	33.96	200	✓ Tegretol CR
* Tab 400 mg	34.58	100	✓ Tegretol
* Tab long-acting 400 mg		100	✓ Tegretol CR
* Oral liq 20 mg per ml		250 ml	✓ Tegretol
CLOBAZAM - Safety medicine; prescriber may determine disp			-
Tab 10 mg	. ,	50	✓ Frisium
CLONAZEPAM - Safety medicine; prescriber may determine of	lispensing frequen	icy	
Oral drops 2.5 mg per ml	7.38	10 ml OP	✓ Rivotril
ETHOSUXIMIDE			
Cap 250 mg	78.89	56	✓ Essential
			Ethosuximide S29
	140.88	100	✓ Zarontin
Oral liq 250 mg per 5 ml	56.35	200 ml	✓ Zarontin
GABAPENTIN			
Note: Not subsidised in combination with subsidised prega	ıbalin		
* Cap 100 mg	6.45	100	✓ Nupentin
* Cap 300 mg	8.45	100	✓ Nupentin
* Cap 400 mg	10.00	100	/ Nunontin
	10.26	100	✓ Nupentin
LACOSAMIDE – Special Authority see SA1125 on the next pa			Nupenun
	ge – Retail pharma		✓ Nuperiun ✓ Vimpat
LACOSAMIDE - Special Authority see SA1125 on the next pa	ge – Retail pharma 25.04	асу	
LACOSAMIDE – Special Authority see SA1125 on the next pa A Tab 50 mg	ge – Retail pharma 25.04	acy 14	✓ Vimpat

▲ Tab 200 mg400.55

56 56

300.40

✓ Vimpat

✓ Vimpat

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

⇒SA1125 Special Authority for Subsidy

I AMOTRIGINE

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see note).

Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

	VIOTRIGINE			_	
	Tab dispersible 2 mg		30		Lamictal
lack	Tab dispersible 5 mg	.50.00	30	•	Lamictal
*	Tab dispersible 25 mg	2.76	56	•	Logem
*	Tab dispersible 50 mg	3.31	56	1	Logem
*	Tab dispersible 100 mg	4.40	56	1	Logem
LE	/ETIRACETAM				
	Tab 250 mg	4.99	60	1	Everet
	Tab 500 mg		60	1	Everet
	Tab 750 mg		60		Everet
	Tab 1,000 mg		60		Everet
	Oral lig 100 mg per ml		300 ml OP	1	Levetiracetam-AFT
ДЦ	ENOBARBITONE	•			
ГΠ	For phenobarbitone oral liquid refer Standard Formulae, page 250				
*	Tab 15 mg	40.00	500	_	PSM
	· ·		500		PSM
	Tab 30 mg	.40.00	500	•	POW
	ENYTOIN SODIUM			_	
*	Tab 50 mg		200		Dilantin Infatab
	Cap 30 mg	.74.00	200		Dilantin
	Cap 100 mg		200		Dilantin
*	Oral liq 30 mg per 5 ml	.22.03	500 ml	/	Dilantin
PR	EGABALIN				
	Note: Not subsidised in combination with subsidised gabapentin				
	Cap 25 mg	2.25	56	1	Pregabalin Pfizer
	, ,	7.80			Milpharm S29
*	Cap 75 mg		56		Pregabalin Pfizer
-		8.10			Milpharm S29
	Cap 150 mg		56		Lyrica
	Oap 100 mg	7.01	30		Pregabalin Pfizer
	Cap 300 mg	7 38	56		Pregabalin Pfizer
-		7.00	30	٠	i regulatii i iizei
	IMIDONE	07.05	400	,	B : : : : : : : :
*	Tab 250 mg	.37.35	100	•	Primidone Clinect
SO	DIUM VALPROATE				
	Tab 100 mg	.13.65	100	1	Epilim Crushable
	Tab 200 mg EC	.27.44	100	1	Epilim
	Tab 500 mg EC	.52.24	100	1	Epilim
*	Oral liq 200 mg per 5 ml	.20.48	300 ml	1	Epilim S/F Liquid
	•			1	Epilim Syrup
*	Inj 100 mg per ml, 4 ml	.41.50	1	•	Epilim IV

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
STIRIPENTOL - Special Authority see SA1330 below - Retail ph	armacy			
Cap 250 mg	509.29	60	✓	Diacomit \$29
Powder for oral liq 250 mg sachet	509.29	60	✓	Diacomit S29
⇒SA1330 Special Authority for Subsidy				

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE

▲ Tab 500 mg	,	100	✓ Sabril
VIGABATRIN - Special Authority see SA2088 below - Retail pharm.			- p
▲ Sprinkle cap 25 mg	26.04	60	✓ Topamax
▲ Sprinkle cap 15 mg	20.84	60	✓ Topamax
	129.85		✓ Topamax
·			✓ Topiramate Actavis
▲ Tab 200 mg	55.19	60	✓ Arrow-Topiramate
	75.25		✓ Topamax
			Topiramate Actavis
▲ Tab 100 mg	31.99	60	Arrow-Topiramate
	44.26		✓ Topamax
			Topiramate Actavis
▲ Tab 50 mg	18.81	60	Arrow-Topiramate
	26.04		✓ Topamax
			Topiramate Actavis
▲ Tab 25 mg	11.07	60	Arrow-Topiramate

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
 - 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 109

Acute Migraine Treatment

RIZATRIPTAN Tab orodispersible 10 mg3.65	30	✓ Rizamelt
SUMATRIPTAN		
Tab 50 mg14.41	90	✓ Sumagran
Tab 100 mg22.68	90	✓ Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per		
prescription34.00	2 OP	✓ Imigran

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 49

PIZOTIFEN

* Tab 16 mg

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT - Special Authority see SA0987 below - Retail pharmacy

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

4 62

100

/ Sarc

Viatris

$\neg \neg \neg$	LUOTINE	DILIVEDOCIN	ODIDE
BFIA	AHISTINE	DIHYDROCHI	ORIDE

	100	<u> </u>
0.49	10	✓ Nausicalm
16.36	10	✓ HameIn
2.85	100	✓ Pharmacy Health
4.00		✓ Domperidone
	16.36	16.36 10

(Pharmacy Health Tab 10 mg to be delisted 1 June 2023)

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
HYOSCINE HYDROBROMIDE					
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	√	Martindale S29	
Patch 1.5 mg - Special Authority see SA1998 below - Retai	l				
pharmacy	17.70	2	1	Scopoderm TTS	

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE

*	Tab 10 mg - Up to 30 tab available on a PSO1.30	100	✓ Metoclopramide Actavis 10
*	Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO7.00	10	✓ Baxter
ON	DANSETRON		
*	Tab 4 mg2.68	50	✓ Onrex
	Tab disp 4 mg - Up to 10 tab available on a PSO0.76	10	✓ <u>Ondansetron</u> <u>ODT-DRLA</u>
*	Tab 8 mg4.57	50	✓ Onrex
	Tab disp 8 mg - Up to 10 tab available on a PSO1.13	10	✓ <u>Ondansetron</u> <u>ODT-DRLA</u>
PR	OCHLORPERAZINE		
*	Tab 3 mg buccal	50	
	(30.00)		Buccastem
*	Tab 5 mg - Up to 30 tab available on a PSO8.00	250	✓ Nausafix
*	Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine d	ispensing frequency	1	
Tab 100 mg	5.15	30	✓ Sulprix
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg	29.78	60	✓ Sulprix
ARIPIPRAZOLE - Safety medicine; prescriber may determine	dispensing frequenc	су	
Tab 5 mg	10.50	30	Aripiprazole Sandoz
Tab 10 mg	10.50	30	Aripiprazole Sandoz
Tab 15 mg	10.50	30	Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz
Tab 30 mg	10.50	30	✓ Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; p	rescriber may deter	mine disper	nsing frequency
Tab 10 mg - Up to 30 tab available on a PSO	14.83	100	✓ Largactil
Tab 25 mg - Up to 30 tab available on a PSO	15.62	100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO	36.73	100	✓ Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO	30.79	10	✓ Largactil

	Subsidy		Fully	
	(Manufacturer's Pric	ce) Per	Subsidised	Generic Manufacturer
	5	Per		Manufacturer
LOZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing freq			_	
Tab 25 mg	6.69	50		Clopine
				Clozaril
	13.37	100		Clopine
				Clozaril
Tab 50 mg		50		Clopine
	17.33	100		Clopine
Tab 100 mg	17.33	50		Clopine
				Clozaril
	34.65	100		Clopine
				Clozaril
Tab 200 mg		50		Clopine
0	69.30	100		Clopine
Suspension 50 mg per ml	67.62	100 m	/	Versacloz
ALOPERIDOL - Safety medicine; prescriber may determine of	dispensing frequency	y		
Tab 500 mcg - Up to 30 tab available on a PSO	6.23	100	✓	Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO	9.43	100	✓	Serenace
Tab 5 mg - Up to 30 tab available on a PSO	14.86	50	✓	Serenace
•	29.72	100	✓	Serenace
Oral liq 2 mg per ml - Up to 200 ml available on a PSO	23.84	100 m	· •	Serenace
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	PSO21.55	10	✓	Serenace
VOMEPROMAZINE - Safety medicine; prescriber may dete	ermine disnensina fr	eallency	,	
Tab 25 mg (33.8 mg as a maleate)		100		Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan
Tab 100 mg (135 mg as a maleate)		100		Nozinan (Swiss)
Tab 100 mg as a maleate		100		Nozinan (Swiss)
G				
VOMEPROMAZINE HYDROCHLORIDE - Safety medicine;				
Inj 25 mg per ml, 1 ml ampoule	16.75	5	/	Neuraxpharm §29
			✓	Nozinan S29 S29
	24.48	10	✓	Wockhardt
	33.50		✓	Nozinan
ozinan Inj 25 mg per ml, 1 ml ampoule to be delisted 1 April 2	2023)			
THIUM CARBONATE - Safety medicine; prescriber may dete	ermine dispensina fr	eauency	,	
Tab long-acting 400 mg	, ,	100		Priadel
Cap 250 mg		100		Douglas
ANZAPINE – Safety medicine; prescriber may determine dis				3 * *
Tab 2.5 mg		28	J	Zypine
•		28 28		
Tab oradisparsible 5 mg		28 28		Zypine Zypine ODT
Tab 10 mg		28 28		Zypine ODT Zypine
Tab 10 mg		28 28		
Tab orodispersible 10 mg			•	Zypine ODT
ERICYAZINE - Safety medicine; prescriber may determine d			_	
Tab 2.5 mg		84		Neulactil
	12.49	100		Neulactil
T-1, 40	27 24	84	/	Neulactil
Tab 10 mg	44.45	100		Neulactil

	Subsidy		Fully	
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
	Ψ	rei		Manuacturer
QUETIAPINE – Safety medicine; prescriber may determine disp			_	
Tab 25 mg	2.15	90	/	<u>Quetapel</u>
Tab 100 mg	5.06	90	•	<u>Quetapel</u>
Tab 200 mg	8.90	90	✓	Quetapel
Tab 300 mg	12.86	90	✓	Quetapel
RISPERIDONE - Safety medicine; prescriber may determine di	spensing frequency			
Tab 0.5 mg	1.86	60	✓	Risperidone (Teva)
Tab 1 mg	2.06	60	✓	Risperidone (Teva)
Tab 2 mg		60	✓	Risperidone (Teva)
Tab 3 mg	2.50	60	✓	Risperidone (Teva)
Tab 4 mg	3.42	60	✓	Risperidone (Teva)
Oral liq 1 mg per ml	8.90	30 m	· •	Risperon
ZIPRASIDONE - Safety medicine; prescriber may determine dis	spensing frequency			
Cap 20 mg	17.90	60	✓	Zusdone
Cap 40 mg	27.41	60	✓	Zusdone
Cap 60 mg	38.39	60	✓	Zusdone
Cap 80 mg		60	1	Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre	escriber may determin	e dis	pensing fre	equency
Tab 10 mg	31.45	100	*	Clopixol

Depot Injections

FLUPENTHIXOL DECANOATE – Safety medicine; prescriber Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO	13.14	ensing freq 5 5 5	uency Fluanxol Fluanxol Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber	may determine dispe	nsing frequ	ency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO		5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	55.90	5	 Haldol Concentrate
			✓ Haldol
			Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - Retail	pharmacy		
Safety medicine; prescriber may determine dispensing fre	quency		
Inj 210 mg vial	252.00	1	Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial	504.00	1	Zyprexa Relprevy

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

	Subsidy (Manufacturer's Price)		Fully Subsidised		
	\$	Per	•	Manufacturer	
PALIPERIDONE - Special Authority see SA1429 below - Retai	l pharmacy				
Safety medicine; prescriber may determine dispensing frequ	iency				
Inj 25 mg syringe	194.25	1	✓	Invega Sustenna	
Inj 50 mg syringe	271.95	1	✓	Invega Sustenna	
Inj 75 mg syringe	357.42	1	✓	Invega Sustenna	
Inj 100 mg syringe	435.12	1	✓	Invega Sustenna	
Inj 150 mg syringe	435.12	1	✓	Invega Sustenna	

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

Inj 175 mg syringe815.85	1	Invega Trinza
Inj 263 mg syringe	1	✓ Invega Trinza
Inj 350 mg syringe	1	✓ Invega Trinza
Inj 525 mg syringe	1	✓ Invega Trinza

⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg vial		1	✓ Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.



	Subsidy (Manufacturer's Price)	S Per	Fully ubsidised	Brand or Generic Manufacturer
ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescribe	r may determine disp	ensing	frequency	1
Inj 200 mg per ml, 1 ml - Up to 5 inj available on a PSO	19.80	5	✓ 0	Clopixol
Anxiolytics				
BUSPIRONE HYDROCHLORIDE				
* Tab 5 mg	18.50	100	✓ <u>B</u>	Buspirone Viatris
* Tab 10 mg	12.50	100	✓ <u>B</u>	Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 500 mcg	5.64	100	✓ P	axam
Tab 2 mg	10.78	100	✓ P	axam
DIAZEPAM - Safety medicine; prescriber may determine dispen	sing frequency			
Tab 2 mg		500	✓ A	rrow-Diazepam
Tab 5 mg	73.60	500	✓ A	rrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 1 mg	9.72	250	✓ A	<u>itivan</u>

Multiple Sclerosis Treatments

⇒SA2176 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 2 Patients has an EDSS score between 0 6.0; and

- 3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 4 All of the following:
 - 4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 4.5 Either:
 - 4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 6 Any of the following:
 - 6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium

continued...

100

Ativan

			14=11	V003 3131LW
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
continued				
enhancing lesion; or				
6.2 A sign of that new inflammatory activity is a lesion s	0		,	
6.3 A sign of that new inflammatory is a T2 lesion with6.4 A sign of that new inflammatory activity is a promin				a for the clinical features o
a recent attack that occurred within the last 2 years		uny ic	теоропові	
6.5 A sign of that new inflammatory activity is new T2 le		h a pr	evious MRI	scan.
Note: Treatment on two or more funded multiple sclerosis treatm				
Renewal — (Multiple sclerosis) only from a neurologist or gene				
had an EDSS score of 0 to 6.0 (inclusive) with or without the use (i.e. the patient has walked 100 metres or more with or without a				e in the last six months
Note: Treatment on two or more funded multiple sclerosis treatm				
DIMETHYL FUMARATE - Special Authority see SA2176 on the	•		•	
a) Wastage claimable			,	
b) Note: Treatment on two or more funded multiple sclerosis				
Cap 120 mg		14		ecfidera
Cap 240 mg	•	56	V 1	ecfidera
FINGOLIMOD – Special Authority see SA2176 on the previous p	age – Retail pharma	су		
a) Wastage claimableb) Note: Treatment on two or more funded multiple sclerosis	s treatments simultan	eous	ly is not ner	mitted
Cap 0.5 mg		28	, ,	ilenya
GLATIRAMER ACETATE - Special Authority see SA2176 on the	previous page – Re	tail ph	narmacy	
Note: Treatment on two or more funded multiple sclerosis tre	eatments simultaneou	usly is	not permitt	
Inj 40 mg prefilled syringe	•	12	_	opaxone
INTERFERON BETA-1-ALPHA – Special Authority see SA2176				
Note: Treatment on two or more funded multiple sclerosis tre Inj 6 million iu prefilled syringe		usiy is 4		ea. . vonex
Injection 6 million iu per 0.5 ml pen injector		4	-	vonex Pen
INTERFERON BETA-1-BETA - Special Authority see SA2176 or		- Reta	ail pharmacy	1
Note: Treatment on two or more funded multiple sclerosis tre			not permitt	ed.
Inj 8 million iu per 1 ml	·	15	✓ B	etaferon
NATALIZUMAB - Special Authority see SA2176 on the previous				
Note: Treatment on two or more funded multiple sclerosis tre		usly is 1		ed. 'ysabri
Inj 20 mg per ml, 15 ml vial	•		• 1	ysabii
OCRELIZUMAB – Special Authority see SA2176 on the previous Note: Treatment on two or more funded multiple sclerosis tre			not nermitt	ed.
Inj 30 mg per ml, 10 ml vial		1		Ocrevus
TERIFLUNOMIDE – Special Authority see SA2176 on the previo	us page – Retail pha	rmacy		
a) Wastage claimable			,	
b) Note: Treatment on two or more funded multiple sclerosis				
Tab 14 mg	659.90	28	✓ <u>A</u>	ubagio
Sedatives and Hypnotics				

Sedatives and Hypnotics

MELATONIN − Special Authority see SA1666 on the next page − Retail pharmacy
Tab modified-release 2 mg − No more than 5 tab per day......11.50 30 ✓ Vigisom



Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	_	Manufacturer

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

MIDAZOLAM Cofety modicine: properitor may determine dispensing frequency

Note: Indications marked with * are unapproved indications.

MIDAZOLAM – Safety medicine; prescriber may determine dispens	sing frequency		
Inj 1 mg per ml, 5 ml ampoule	6.10	10	✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available			
on a PSO	17.28	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be en	dorsed for statu	s epilepticu	us use only.
Inj 5 mg per ml, 3 ml ampoule	5.00	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available on	1		
a PSO	13.09	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be en	dorsed for statu	ıs epilepticı	us use only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 bel	ow – Retail pha	rmacy	
Inj 200 mg per ml, 1 ml ampoule	103.30	10	✓ Max Health S29
SA1386 Special Authority for Subsidy			

SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM – Safety medicine; prescriber may determine Tab 10 mg	1 0 1 ,	25	✓ Normison
TRIAZOLAM – Safety medicine; prescriber may determine d			
Tab 125 mcg	5.10	100	
·	(9.85)		Hypam
Tab 250 mcg	4.10 [°]	100	
Ç	(11.20)		Hypam
ZOPICLONE - Safety medicine; prescriber may determine of	lispensing frequency		
Tab 7.5 mg		500	✓ Zopiclone Actavis

NERVOUS SYSTEM

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Spinal Muscular Atrophy

⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.



	(Manufacturer's Price)		Subsidised	Generic
	\$	Per		Manufacturer
Stimulants/ADHD Treatments				
ATOMOXETINE			_	
Cap 10 mg	18.41	28		APO-Atomoxetine
			•	APO-Atomoxetine
				S29 S29
	407.00			Generic Partners
0 10	107.03	00		Strattera
Cap 18 mg	27.06	28		APO-Atomoxetine
	107.03			Generic Partners Strattera
Cap 25 mg		28		APO-Atomoxetine
Oap 25 Hg	29.22	20		Generic Partners
Cap 40 mg	29 22	28		APO-Atomoxetine
σαρ 40 mg		20		Generic Partners
	107.03			Strattera
Cap 60 mg		28	/	APO-Atomoxetine
			1	APO-Atomoxetine
				S29 S29
			1	Generic Partners
Cap 80 mg	56.45	28	1	APO-Atomoxetine
			1	APO-Atomoxetine
				S29 S29
			1	Generic Partners
Cap 100 mg	58.48	28	1	APO-Atomoxetine
			1	APO-Atomoxetine
				S29 S29
			1	Generic Partners
(Strattera Cap 10 mg to be delisted 1 November 2023)				
(Strattera Cap 18 mg to be delisted 1 November 2023)				
(Strattera Cap 40 mg to be delisted 1 November 2023)				
DEXAMFETAMINE SULFATE - Special Authority see SA1149 be	elow – Retail pharma	СУ		
a) Only on a controlled drug form	•	•		
b) Safety medicine; prescriber may determine dispensing free	quency			
Tab 5 mg		100	1	<u>PSM</u>
	28.50		1	Aspen

Subsidy

Fully

Brand or

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg		30	✓ Ritalin
•			✓ Rubifen
Tab extended-release 18 mg	7.75	30	✓ Methylphenidate ER - Teva
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg	10.95	30	✓ Rubifen SR
Tab extended-release 27 mg	11.45	30	Methylphenidate ERTeva
Tab extended-release 36 mg	15.50	30	Methylphenidate ERTeva
Tab extended-release 54 mg	22.25	30	✓ Methylphenidate ER - Teva

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.



bsidy urer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Roth:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	Concerta
Tab extended-release 36 mg	71.93	30	✓ Concerta
Tab extended-release 54 mg		30	✓ Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Fither:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or

NERVOUS SYSTEM

Subsid	dy Fi	ılly Brand or	
(Manufacture	r's Price) Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Retail pharmac	y		
Tab 100 mg	29.13	60	✓ Modavigil

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below	/ - Retail pharmacy		
Patch 4.6 mg per 24 hour	38.00	30	✓ Rivastigmine Patch
			<u>BNM 5</u>
Patch 9.5 mg per 24 hour	38.00	30	 Rivastigmine Patch
			BNM 10

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
` ¢ ′	Dor ./	Manufacturar

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg11.76

✓ Buprenorphine 28 Naloxone BNM

Tab sublingual 8 mg with naloxone 2 mg34.00 28 ✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone);
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

30 Zyban

./ Habitual

	Subsidy (Manufacturer's Price \$		Fully bsidised	Brand or Generic Manufacturer
DISULFIRAM				
Tab 200 mg	236.40	100	✓ <u>P</u>	Antabuse S29
NALTREXONE HYDROCHLORIDE - Special Authority	see SA1408 below - Retail	pharmacy	,	
Tab 50 mg	133.33	30	✓ V	Naltraccord
Tab 50 mg	133.33	30	• [vaitraccor

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Compliance with the medication (prescriber determined); and

Lin to 00 noteh available on a DCO

- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Patch / mg - Up to 28 patch available on a PSO18.14	28	Habitrol
Patch 7 mg for direct distribution only - [Xpharm]3.94	7	✓ Habitrol
Patch 14 mg - Up to 28 patch available on a PSO19.95	28	Habitrol
Patch 14 mg for direct distribution only - [Xpharm]4.52	7	Habitrol
Patch 21 mg - Up to 28 patch available on a PSO22.86	28	Habitrol
Patch 21 mg for direct distribution only - [Xpharm]5.18	7	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO19.18	216	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	36	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO21.02	216	Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	36	Habitrol
Gum 2 mg (Fruit) – Up to 384 piece available on a PSO38.21	384	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]8.64	96	Habitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO38.21	384	Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]8.64	96	Habitrol
Gum 4 mg (Fruit) – Up to 384 piece available on a PSO24.17	204	Habitrol
44.17	384	Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]10.01	96	Habitrol
Gum 4 mg (Mint) - Up to 384 piece available on a PSO44.17	384	Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]10.01	96	Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	53 OF	✓ <u>Varenicline Pfizer</u>
Tab 1 mg17.6	52 56	✓ Varenicline Pfizer



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA2153 below

Inj 25 mg vial77.00	•	✓ Ribomustin
Inj 100 mg vial308.00) 1	✓ Ribomustin
Inj 1 mg for ECP	3 1 mc	✓ Baxter

⇒SA2153 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+): and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

booder Air — i o i — netaii phannacy-opecialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
, ,	45.20		 Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	710.00	1	✓ BiCNU
Inj 100 mg for ECP		100 mg OP	✓ Baxter
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist		-	
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	✓ Cisplatin Ebewe
.,	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP	0.31	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE		· ·	
Tab 50 mg - PCT - Retail pharmacy-Specialist	145.00	50	✓ Cyclonex
Inj 1 g vial – PCT – Retail pharmacy-Specialist		1	✓ Endoxan
, 3 , ,	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist	71.25	1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist		1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
Inj 1 g	96.00	1	✓ Holoxan
Inj 2 g		1	✓ Holoxan
Inj 1 mg for ECP		1 mg	✓ Baxter
<i>,</i>		Ü	

1 ma

✓ Baxter

	Subsidy (Manufacturer's Price)		Fully	
	\$	Per	✓ Cubsidised	
LOMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	✓	CeeNU
Cap 40 mg	399.15	20	✓	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	/	Alkeran
Inj 50 mg - PCT only - Specialist		1	1	Melpha
	67.80		✓	Alkeran
			1	Alkeran S29 S29
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
,				100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1		Oxaliplatin Accord
Inj 1 mg for ECP	0.48	1 mg	✓	Baxter
THIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	/	Bedford \$29
11j 10 11g via		•		Max Health S29
				THIO-TEPA \$29
				Tepadina S29
Let 400 mm aled	ODO	_		•
Inj 100 mg vial	CBS	1		Max Health S29
			•	Tepadina S29
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see SA				
Inj 100 mg vial	75.06	1	•	Azacitidine Dr
				Reddy's

⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome: or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder);
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- - 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and patient is benefitting from treatment.

				_
	Subsidy		Fully Brand or	
	(Manufacturer's Price		bsidised Generic	
	\$	Per	✓ Manufacturer	_
CALCIUM FOLINATE				
Tab 15 mg - PCT - Retail pharmacy-Specialist	114.69	10	DBL Leucovorin	
			Calcium	
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist		5	✓ Hospira	
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Special	ist7.28	1	✓ Calcium Folinate	
			Sandoz	
			Calcium Folinate	
			Sandoz S29 S29	
Inj 50 mg - PCT - Retail pharmacy-Specialist	72.80	10	✓ Leucovorin	
			Pharmacia S29	
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	Calcium Folinate	
			Sandoz	
Inj 100 mg - PCT only - Specialist	7.33	1	Calcium Folinate	
			Ebewe	
	94.90	10	✓ Leucovorin	
			Pharmacia S29	
Inj 300 mg - PCT only - Specialist	22.51	1	Calcium Folinate	
			Ebewe	
	25.14		✓ Leucovorin DBL S29	
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	✓ Calcium Folinate	
ing forming per fini, 33 thii viai = 1 of only = Specialist	25.14	'	Sandoz	
			✓ Calcium Folinate	
			Sandoz S29 S29	
Inj 1 g - PCT only - Specialist	67 51	1	✓ Calcium Folinate	
ing right of only openials	07.01	'	Ebewe	
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	✓ Calcium Folinate	
ing to mg por init, roo ini nai in on only opeoiainet initiani		•	Sandoz	
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓ Baxter	
CAPECITABINE - Retail pharmacy-Specialist		3		
Tab 150 mg	10.00	60	✓ Capercit	
Tab 500 mg		120	✓ Capercit	
CLADRIBINE – PCT only – Specialist				
Inj 2 mg per ml, 5 ml	7/0 06	1	✓ Litak \$29	
Inj 1 mg per ml, 10 ml		1	✓ Leustatin	
Inj 10 mg for ECP		10 mg OP	✓ Baxter	
CYTARABINE			24	
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special	ist 400.00	5	✓ Pfizer	
Inj 100 mg per ml, 20 ml vial – PCT – Retail	101	O	1 HZGI	
pharmacy-Specialist	41.36	1	✓ Pfizer	
Inj 1 mg for ECP - PCT only - Specialist		10 mg	✓ Baxter	
Inj 100 mg intrathecal syringe for ECP - PCT only - Special		100 mg OP		
FLUDARABINE PHOSPHATE		ŭ		
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	✓ Fludara Oral	
Inj 50 mg vial – PCT only – Specialist		5	✓ Fludarabine Ebewe	
Inj 50 mg for ECP - PCT only - Specialist		50 mg OP	✓ Baxter	
		-		

	Subsidy		Fully	Brand or
	(Manufacturer's Price		sidised	
	\$	Per		Manufacturer
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	✓	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	29.44	1	✓	Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist	0.62	100 mg	✓	Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	1	DBL Gemcitabine
Inj 1 g		1	1	Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg	✓	Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist		•		
Inj 20 mg per ml, 5 ml vial	52.57	1	1	Accord
, g p,	71.44	•	1	Irinotecan Actavis
				100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	1	Baxter
MERCAPTOPURINE		ŭ		
Tab 50 mg - PCT - Retail pharmacy-Specialist	25 90	25	1	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialist			•	1 411 11041101
Special Authority see SA1725 below		100 ml OP	1	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist9.98	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist33.71	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	Methotrexate Sandoz
*	Inj 10 mg prefilled syringe14.66	1	Methotrexate Sandoz
*	Inj 15 mg prefilled syringe14.77	1	Methotrexate Sandoz
*	Inj 20 mg prefilled syringe14.88	1	Methotrexate Sandoz
*	Inj 25 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 30 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ Methotrexate DBL Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist79.99	1	✓ Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PEMETREXED - PCT only - Specialist - Special Authority see	SA1679 below			
Inj 100 mg vial	60.89	1	✓	Juno Pemetrexed
Inj 500 mg vial	217.77	1	✓	Juno Pemetrexed
Inj 1 mg for ECP	0.55	1 mg	✓	Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 40 mg126.31

· ·			
Other Cytotoxic Agents			
AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-S	Specialist		
Cap 0.5 mg	1,175.87	100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter

I anvis

25

	Subsidy (Manufacturer's P \$	rice) Subs	Fully sidised	Brand or Generic Manufacturer
BLEOMYCIN SULPHATE - PCT only - Specialist				
Inj 15,000 iu, vial	185.16	1	✓ D	BL Bleomycin Sulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	✓ B	Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see \$	SA1889 below			
Inj 3.5 mg vial	74.93 105.00	1	_	BL Bortezomib Bortezomib Dr-Reddy's
Inj 1 mg for ECP(Bortezomib Dr-Reddy's Inj 3.5 mg vial to be delisted 1 May 2023		1 mg	✓ B	Baxter

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	62.70	1	DBL Dacarbazine
, 3	580.60	10	✓ Dacarbazine
			APP S29
Inj 200 mg for ECP	62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist			
Inj 2 mg per ml, 10 ml	149.50	1	✓ Pfizer
Inj 20 mg vial		10	✓ Daunorubicin
, ,	•		Zentiva S29
Inj 20 mg for ECP	149.50	20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist			
Inj 20 mg	48.75	1	✓ Docetaxel Sandoz
lnj 10 mg per ml, 8 ml vial	46.89	1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	✓ Docetaxel
. •			Accord S29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP	0.65	1 mg	✓ Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	10.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1	Doxorubicin Ebewe
, •	17.00		✓ Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	65.00	1	✓ Arrow-Doxorubicin
	69.99		Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	✓ Baxter

\$ Per		Subsidy		Fully	
Inj 2 mg per ml, 5 ml vial		` • '		Subsidised	
Inj 2 mg per ml, 5 ml vial	EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 100 ml vial		25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial	Inj 2 mg per ml, 25 ml vial	30.00	1	✓	Epirubicin Ebewe
Inj 1 mg for ECP			1	✓	Epirubicin Ebewe
Cap 50 mg - PCT - Retail pharmacy-Specialist			1 mg	/	Baxter
Cap 100 mg - PCT - Retail pharmacy-Specialist	ETOPOSIDE				
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist7.90 Inj 1 mg for ECP – PCT only – Specialist	Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	1	Vepesid
Inj 1 mg for ECP − PCT only − Specialist	Cap 100 mg - PCT - Retail pharmacy-Specialist	340.73	10	1	Vepesid
ETOPOSIDE PHOSPHATE – PCT only – Specialist Inj 100 mg (of etoposide base)	Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Special	ist7.90	1	1	Rex Medical
Inj 100 mg (of etoposide base)	Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	/	Baxter
Inj 1 mg (of etoposide base) for ECP	ETOPOSIDE PHOSPHATE - PCT only - Specialist				
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pharmacy-Specialist	Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
	Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	/	Baxter
	HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail phar	macy-Specialist	_		
Cap 500 mg	Cap 500 mg		100	1	Devatis
IBRUTINIB – Special Authority see SA2168 below – Retail pharmacy					
Tab 140 mg3,217.00 30 ✓ Imbruvica		•	30	1	Imbruvica
Tab 420 mg9,652.00 30 ✓ Imbruvica			30	1	Imbruvica

⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
 - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
 - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
 - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

Inj 5 mg vial - PCT only - Specialist	109.74	1	✓ Zavedos
Inj 10 mg vial - PCT only - Specialist	233.64	1	Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacture	r
LENALIDOMIDE - Retail pharmacy-Specialist - Special Autho	rity see SA2047 below			
Wastage claimable	•			
Cap 5 mg	5,122.76	28	✓ Revlimid	
Cap 10 mg	4,655.25	21	✓ Revlimid	
	6,207.00	28	✓ Revlimid	
Cap 15 mg	5,429.39	21	✓ Revlimid	
, •	7,239.18	28	✓ Revlimid	
Cap 25 mg	7,627.00	21	✓ Revlimid	

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Fither:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MFSNA

Tab 400 mg - PCT - Retail pharmacy-Specialist314.00	50	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist448.50	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.45	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist407.40	15	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist2.96	100 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
MITOMYCIN C - PCT only - Specialist				
Inj 5 mg vial	641.70	1	1	Accord S29
Inj 20 mg vial		1	✓.	Teva
Inj 1 mg for ECP	269.85	1 mg	✓	Baxter
MITOZANTRONE - PCT only - Specialist				
Inj 2 mg per ml, 10 ml vial	97.50	1	✓	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	✓	Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority se	ee SA2163 below			
Tab 100 mg	3,701.00	56	✓	Lynparza
Tab 150 mg		56	✓	Lynparza

⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Fither
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment

Subsidy	F	ully	Brand or
(Manufacturer's	Price) Subsid	sed	Generic
\$	Per	✓	Manufacturer

continued...

period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or

5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg	24.00	1	✓ Paclitaxel Ebewe
, -	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
, -	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
, -	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see SA	1979 below		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMII F)

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 The patient has relapsed acute lymphoblastic leukaemia; and

DENITOCIATINI IDEOVVCOEODMVCINI DOT only Consistint

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE – PCT – Retail pharmacy-Sp		'	• Ripent 425
Cap 50 mg		50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 on the next page	– Retail pha	rmacy	
Cap 5 mg	9.13	5	✓ Temaccord
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
Cap 100 mg	35.98	5	✓ Temaccord
	40.20		✓ Apo-Temozolomide
Cap 140 mg	50.12	5	✓ Temaccord
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ Temaccord

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special	Authority see SA1124 on the	next page	
Cap 50 mg	378.00	28	✓ Thalomid
Can 100 mg	756.00	28	✓ Thalomid

Subsidy (Manufacturer's F	rice) Su	Fully	Brand or Generic	
\$	Per	✓	Manufacturer	

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist479.50	100	✓ Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA1868 be	elow	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	42 OP	✓ Venclexta
Tab 10 mg95.78	14 OP	✓ Venclexta
Tab 50 mg239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable8,209.41	120	✓ Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax: and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

Subsidy		Fully	
(Manufacturer's Price		Subsidised	
<u></u>	Per		Manufacturer
VINBLASTINE SULPHATE			
Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37	5	✓	Hospira
Inj 1 mg for ECP - PCT only - Specialist6.00	1 mg	· ·	Baxter
VINCRISTINE SULPHATE			
Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Specialist74.52	5	•	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist102.73	5	•	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	·	Baxter
VINORELBINE - PCT only - Specialist			
Inj 10 mg per ml, 1 ml vial12.00	1	✓	Navelbine
42.00		✓	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00	1	✓	Navelbine
210.00		✓	Vinorelbine Ebewe
328.65		✓	Sagent S29
Inj 1 mg for ECP1.25	1 mg	·	Baxter
	50 mg	OP 🗸	Baxter (Sagent)

Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable			
Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel
•	,		

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and

(1)	Subsidy	Fully		Brand or
	Manufacturer's Price)	Subsidised		Generic
	\$	Per	1	Manufacturer

continued...

- 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day: or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB – Retail pharmacy-Specialist – Special /	Authority see SA2115 below		
Tab 100 mg	329.70	30	✓ Alchemy
•	764.00		✓ Tarceva
Tab 150 mg	569.70	30	✓ Alchemy
•	1.146.00		✓ Tarceva

(Tarceva Tab 100 mg to be delisted 1 February 2023) (Tarceva Tab 150 mg to be delisted 1 February 2023)

⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
GEFITINIB - Retail pharmacy-Specialist - Special Authority see	SA2116 below			
Tab 250 mg	918.00	30	✓ Ir	essa

SA2116 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	Tab 100 mg - [Xpharm] - Special Authority see SA1460		
	below2,400.00	60	✓ Glivec
*	Cap 100 mg58.23	60	✓ Imatinib-Rex
*	Cap 400 mg84.79	30	✓ Imatinib-Rex

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 on the next page - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage	claimable
---------	-----------

Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6,532.00	120	✓ Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

Wastage claimable	ļ
Tab 75 mg	

Tab 75 mg4,000.00	21	✓ Ibrance
Tab 100 mg4,000.00	21	✓ Ibrance
Tab 125 mg4,000.00	21	✓ Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

state: and

4.2.2 Either:

- 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
- 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB – Special Authority see SA1190 below – Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal: or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised		
RUXOLITINIB – Special Authority see SA1890 below – Re Wastage claimable	tail pharmacy				
Tab 5 mg	2,500.00	56	1	Jakavi	
Tab 10mg	5,000.00	56	1	Jakavi	
Tab 15 mg	5,000.00	56	1	Jakavi	
Tab 20 mg	5 000 00	56	/	Jakavi	

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

		SA2117 below – Retail pharmacy	SUNITINIB – Special Authority see SA2117
 Sunitinib Pfizer 	28	208.38	Cap 12.5 mg
✓ Sunitinib Pfizer	28	416.77	Cap 25 mg
✓ Sunitinib Pfizer	28	694.62	Cap 50 mg

⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:

				_
Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
- 5.2 Haemoglobin level < lower limit of normal; or
- 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
- 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
- 5.5 Karnofsky performance score of less than or equal to 70; or
- 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 83

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

Tab 250 mg4,276.19 120 ✓ Zytiga

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE Tab 50 mg	4.21	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
· ·	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Specia	Authority see SA1895 on t	he next pag	е
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL ACETATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking megestrol acetate prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of megestrol acetate.

Tab 160 mg	48.80	30	Megace S29
(Megace S29) Tab 160 mg to be delisted 1 February 2023)			
OCTREOTIDE			
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	Max Health
		•	Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	Max Health
Inj 500 mcg per ml, 1 ml ampoule1	13.10	5	Max Health
OCTREOTIDE LONG-ACTING - Special Authority see SA2119 below -	- Retail pharma	CV	
Inj depot 10 mg prefilled syringe4	39.97	i 🗸	Octreotide Depot
			Teva
Inj depot 20 mg prefilled syringe6	47.03	1	Octreotide Depot
			Teva
Inj depot 30 mg prefilled syringe7	18.55	1	Octreotide Depot
			Teva

⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

continued...

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Fither:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed: or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has acromegaly; and

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
continued				
2 Patient has a large pituitary tumour, greater than 10 mm a3 Patient is scheduled to undergo pituitary surgery in the ne	,			
TAMOXIFEN CITRATE				
* Tab 10 mg * Tab 20 mg		60 60	_	amoxifen Sandoz amoxifen Sandoz
Aromatase Inhibitors				
ANASTROZOLE				
* Tab 1 mg	4.55	30	✓ <u>A</u>	natrole
# Tab 25 mg	14.50	30	√ P	fizer Exemestane
LETROZOLE				

Immunosuppressants

* Tab 2.5 mg

Cytotoxic Immunosuppressants

AZATHIOPRINE			
* Tab 25 mg	7.36	60	Azamun
* Tab 50 mg	8.10	100	Azamun
MYCOPHENOLATE MOFETIL			
Tab 500 mg	35.90	50	✓ Cellcept
Cap 250 mg	35.90	100	✓ Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement	187.25	165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

30

✓ Letrole

Fusion Proteins

ETANERCEPT – Special Authority see SA2103 below -	- Retail pharmacy		
Inj 25 mg	690.00	4	✓ Enbrel
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Inj 50 mg prefilled syringe		4	✓ Enbrel

⇒SA2103 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
- 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD: or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Fither:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per Manufacturer

continued...

55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:

Subsidy (Manufacturer's I	Price) Sub	Fully	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

- 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
- 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*: and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g.
 - prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:

Subsidy (Manufacturer's Price)	,	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses unless contraindicated); and
- 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

Subsidy	,	Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Fither:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Fither:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	1	Manufacturer

continued...

- 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Sp	Decialist			
Inj 50 mg per ml, 5 ml	2,774.48	5	✓ ATGAM	
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT	only – Specialist			
Subsidised only for bladder cancer.				
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE	
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG \$29	

Monoclonal Antibodies

	у	2177 below - Retail pharmac	ADALIMUMAB (AMGEVITA) - Special Authority see SA2
✓ Amgevita	1	190.00	Inj 20 mg per 0.4 ml prefilled syringe
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled pen
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled syringe

⇒SA2177 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions: and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline: and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

Subsidy (Manufacturer's Price)		Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.2 Either:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has pyoderma gangrenosum*; and
 - 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Roth:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
- 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
- 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or

Subsidy	Fi	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	/	Manufacturer	

continued...

- 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
 - 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

Subsidy		Fully	Brand or	П
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or

2 All of the following:

- 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
- 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis: or

2 All of the following:

- 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
- 2.5 Fither:

(Mar	Subsidy nufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

- 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:
Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
 - 12 Fither
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on adalimumab; or
- 2 The PUCAI score has reduced by 30 points or more from the PUCAI score since initiation on adalimumab.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs: and
- 4 Patient has bilateral sacroiliitis demonstrated by radiological imaging; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not responded to at least three months of methotrexate, or azathioprine at a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of sulfasalazine at a maximum tolerated dose; and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see \$A2157 below - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe	1,599.96	2	✓ Humira
Inj 40 mg per 0.4 ml prefilled syringe	1,599.96	2	Humira
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	1,599.96	2	Humira

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease - severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

from baseline: and

- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	idised	Generic
\$	Per	1	Manufacturer

continued...

4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
 - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
 - 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Fither
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacv

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Fither:

Subsidy	Fully	/ Brand or
(Manufacturer's Price)	Subsidised	d Generic
 \$	Per 🗸	Manufacturer

continued...

- 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
- 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or

Subsidy		Fully	Brand or	
Gabblay		. ,	Diana oi	
(Manufacturer's Price)		Subsidised	Generic	
(Wandacturer 3 Frice)		Oubsidisca	acricio	
\$	Per	_	Manufacturer	
Ψ	1 01	•	Manadatarer	

continued...

- 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Fither:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

⇒SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated: and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 below

_ 1 O / (1141) (D	1 0 1 0 1 11 9	poolanot opoolai	ridinolity dod orthoor bolon		
Inj 5 mg p	er ml, 20 ml via	d	364.00	1	Erbitux
Inj 5 mg p	er ml, 100 ml v	ial	1,820.00	1	✓ Erbitux
Ini 1 ma fo	or ECP		3.82	1 ma	✓ Baxter

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

	Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic	
	\$	Per	✓	Manufacturer	
GEMTUZUMAB OZOGAMICIN - PCT only - Specialist - Specialis	ial Authority see SA21	58 below			
Ini 5 mg vial	12.973.00	1	✓ N	Ivlotarg	

⇒SA2158 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2082 below

 Inj 100 mg
 806.00
 1
 ✓ Remicade

 Inj 1 mg for ECP
 8.29
 1 mg
 ✓ Baxter

⇒SA2082 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Crohn's disease (children))** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy	,	Fully	Brand or	
(Manufacturer's Price	·) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the qut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...
Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis: and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation; or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis: or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plague psoriasis; or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 1 The patient 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids,

Subsidy	Fı	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	_
(Manufacturer's Price))	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Fither
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65: and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Fither:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment: and
- 3 A maximum of 8 doses.

MEPOLIZUMAB - Special Authority see SA2154 below - Retail pharmacy

✓ Nucala	1	1,638.00	 Inj 100 mg prefilled pen.
✓ Nucala	1	1,638.00	 Inj 100 mg vial

⇒SA2154 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	Manufacturer

continued...

- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Fither:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Fither:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA2155 below

Inj 25 mg per ml, 40 ml vial	5,910.00	1	✓ Gazyva
Inj 1 mg for ECP	6.21	1 mg	Baxter

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pha	armacy		
Inj 150 mg prefilled syringe	450.00	1	✓ Xolair
Ini 150 mg vial		1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Specialist - Special Authority see SA2143 below

(Synagis Inj 100 mg per ml, 1 ml vial to be delisted 1 January 2024)

⇒SA2143 Special Authority for Subsidy

Initial application — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Infant was born in the last 2 years and has severe lung, airway, neurological or neuromuscular disease that requires ongoing, life-sustaining community ventilation; or
- 2 Both:
 - 2.1 Infant was born in the last 12 months; and
 - 2.2 Any of the following:
 - 2.2.1 Patient was born at less than 28 weeks gestation; or

Subsidy		Fully	Brand or
(Manufacturer's	Price)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

2.2.2 Both:

2.2.2.1 Patient was born at less than 32 weeks gestation; and

2.2.2.2 Either:

2.2.2.2.1 Patient has chronic lung disease: or

2.2.2.2.2 Patient is Māori or any Pacific ethnicity; or

2.2.3 Both:

2.2.3.1 Patient has haemodynamically significant heart disease; and

2.2.3.2 Any of the following:

2.2.3.2.1 Patient has unoperated simple congenital heart disease with significant left to right shunt (see note a); or

2.2.3.2.2 Patient has unoperated or surgically palliated complex congenital heart disease; or

2.2.3.2.3 Patient has severe pulmonary hypertension (see note b); or

2.2.3.2.4 Patient has moderate or severe LV failure (see note c).

Notes:

- a) Patient requires/will require heart failure medication, and/or patient has significant pulmonary hypertension, and/or patient will require surgical palliation/definitive repair within the next 3 months.
- b) Mean pulmonary artery pressure more than 25 mmHg.
- c) LV Ejection Fraction less than 40%.

Renewal — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months where patient still meets initial criteria.

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

 Inj 500 mg per 50 ml vial
 2,688.30
 1
 ✓ Mabthera

 Inj 1 mg for ECP
 5.64
 1 mg
 ✓ Baxter (Mabthera)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated: and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin: or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Fither:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and
- 2 Fither:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

5	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special A	uthority see SA2114 b	elow	
Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2114 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — **(ANCA associated vasculitis)** from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
- 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
- 3.3 Cyclophosphamide and methotrexate are contraindicated; or
- 3.4 Patient is a female of child-bearing potential; or
- 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia: and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax: or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL;
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment: and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsic	lised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
 - 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy;
 - 2.2 To be used for a maximum of 6 treatment cycles.

Subsidy (Manufacturer's Price)			Brand or Generic
 \$	Per	1	Manufacturer

continued...

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*: and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Fither:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's F	Price) Su	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*: and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Fither
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Fither
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and

Subsidy	,	Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2084 below - Retail pharmacy

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
- 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Sylvant	1	ial770.57	Inj 100 mg vial
✓ Sylvant	1	ial3,082.33	Inj 400 mg vial

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

4. Francisco de La della collega describa 400 como conse

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Inj 100 mg per mi, 1.5 ml viai with cilgavimab 100 mg per ml,1.5 ml viai	0.00	1	✓ Evusheld
TOCILIZUMAB - PCT only - Special Authority see SA2159 on the next	page		
Inj 20 mg per ml, 4 ml vial22	20.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
88	30.00	4	✓ RoActemra S29 S29
Inj 20 mg per ml, 10 ml vial55	50.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
Inj 20 mg per ml, 20 ml vial1,10	00.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
4,40	00.00	4	✓ RoActemra S29 S29
Inj 1 mg for ECP	2.85	1 mg	✓ Baxter

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
` c ′	Por 🗸	Manufacturer	

⇒SA2159 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis: or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and

Subsid	dy Fully	Brand or
(Manufacture	er's Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Tocilizumab is to be used as monotherapy; and
- 3 Fither:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Fither:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Fither:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	0 1	Herceptin
Inj 440 mg vial	0 1	✓ Herceptin
Inj 1 mg for ECP	6 1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:

 1. The patient has metastatic breast cancer ever
 - 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
 - 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
 - 4 Trastuzumab not to be given in combination with lapatinib; and
 - 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — **(early breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib: or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Fither:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2144 below

Inj 100 mg vial	2,320.00	1	✓ Kadcyla
Inj 160 mg vial	3.712.00	1	✓ Kadcvla
Inj 1 mg for ECP		1 mg	✓ Baxter

⇒SA2144 Special Authority for Subsidy

Initial application — **(early breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

DURVALUMAB - PCT only - Specialist - Special Authority s	ee SA2164 below		
Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
Inj 50 mg per ml, 2.4 ml vial		1	Imfinzi
Inj 1 mg for ECP	9.59	1 mg	✓ Baxter

⇒SA2164 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Fither:

Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
 \$	Per	•	Manufacturer

continued...

- 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
- 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

	 PCT only – Specialist – 	Chanial Authority and	C 1 2 1 2 0 holow
INIVULUIVIAD	- FUT ONLY - SUPCIAIISI -	ODECIAI AUTHORITY SEE	OAZ IZU DEIUW

Opdivo	1	per ml, 4 ml vial1,051.98	Inj 10 mg per ml, 4 ml v
Opdivo	1	per ml, 10 ml vial2,629.96	Inj 10 mg per ml, 10 ml
✓ Baxter	1 mg	r ECP27.62	Inj 1 mg for ECP

⇒SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised Per 🗸	Generic Manufacturer
Ψ		a.iaiaaiaia

continued...

- 2.2 Patient has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special A	uthority see SA2121 below		
Inj 25 mg per ml, 4 ml vial	4,680.00	1	✓ Keytruda
Inj 1 mg for ECP	49.14	1 mg	✓ Baxter

⇒SA2121 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	1	Manufacturer

continued...

- 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
- 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg		50	✓ Neoral
Cap 100 mg		50	✓ Neoral
Oral lig 100 mg per ml		50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail phar	macy		
Wastage claimable			
Tab 10 mg	.6,512.29	30	Afinitor
Tab 5 mg	.4,555.76	30	Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

	Subsidy (Manufacturer's Price) \$ F		Brand or Generic Manufacturer
SIROLIMUS – Special Authority see SA2005 below – Retail pha	•		
Tab 1 mg		100	Rapamune
Tab 2 mg Oral liq 1 mg per ml	,	100 60 ml C	Rapamune Rapamune

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy: or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsic	lised	Generic
\$	Per	1	Manufacturer

continued...

- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg		100	✓ Tacrolimus Sandoz
Cap 1 mg		100	✓ Tacrolimus Sandoz
Cap 5 mg		50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – Special Authority see SA2079 or	the next page – Retail pharmacy		
Tab 15 mg	1,271.00	28	✓ RINVOQ

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00

8.00 1 **✓** Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

benefiting from troutinent.		
BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above -	Retail pharma	асу
Initiation kit - 5 vials freeze dried venom with diluent305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with		
diluent285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent		
9 ml, 3 diluent 1.8 ml305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent 305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see SA1367 above	- Retail pharr	macy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		
dried venom, with diluent305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze		
dried venom, with diluent305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze		
dried venom, with diluent305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze	4.00	4 All
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze	4 OD	/ Vanamil 200
dried venom, with diluent305.00	1 OP	✓ Venomil S29

	Subsidy		Fully	Brand or
	(Manufacturer's Pric		sidised	
	\$	Per		Manufacturer
Authistoning				
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg	7.50	100	1	Zista
* Oral liq 1 mg per ml	2.84	200 ml	1	Histaclear
CHLORPHENIRAMINE MALEATE				
* Oral lig 2 mg per 5 ml	9.37	500 ml	1	Histafen
		000 1111	•	inotaton
DEXTROCHLORPHENIRAMINE MALEATE	0.00	40		
* Tab 2 mg		40		Polaramine
	(8.40)	20		Polaramine
	1.01	20		Polaramine
* Oral liq 2 mg per 5 ml	(5.99) 1.77	100 ml		rolatattille
* Oral liq 2 mg per 5 mi	(10.29)	100 1111		Polaramine
	(10.29)			1 Olaraninie
FEXOFENADINE HYDROCHLORIDE	4.04	00		
* Tab 60 mg		20		T.16
W. Tal- 100 mm	(8.23)	10		Telfast
* Tab 120 mg		10		Talfact
	(8.23)	20		Telfast
	14.22 (26.44)	30		Telfast
	(20.44)			Tellast
LORATADINE	4.70	400	,	
* Tab 10 mg		100		Lorafix
* Oral liq 1 mg per ml	1.43	100 ml	•	Haylor syrup
PROMETHAZINE HYDROCHLORIDE				
* Tab 10 mg		50		Allersoothe
* Tab 25 mg		50		Allersoothe
* Oral liq 1 mg per 1 ml		100 ml		Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a	PSO 17.87	5		Hospira
Interted Applications in				
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose	14.01 20	00 dose OP	1	Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		00 dose OP	1	Beclazone 50
Aerosol inhaler, 100 mcg per dose		00 dose OP		Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		00 dose OP	1	Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free		00 dose OP	1	Beclazone 250
BUDESONIDE				
Powder for inhalation, 100 mcg per dose	17.00 20	00 dose OP	1	Pulmicort
Total for initial and it, 100 may per addo		00 0000 01	-	Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00 20	00 dose OP	1	Pulmicort
1 Owder for illinatation, 200 they per dose	13.00 20	UU UUSE OF	•	Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 20	00 dose OP	1	Pulmicort
i owder for illialation, 400 mby per dose	02.00 2	UU UUSE OF	•	Turbuhaler

	Subsidy	Drico) C	Fully Brand or
	(Manufacturer's \$	Price) Subsi Per	dised Generic Manufacturer
LUTICASONE			
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ <u>Flixotide</u>
Aerosol inhaler, 250 mcg per dose		120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose	13.60	60 dose OP	✓ Flixotide Accuhaler
nhaled Long-acting Beta-adrenoceptor Agoni	sts		
FORMOTEROL FUMARATE			
Powder for inhalation, 12 mcg per dose, and monodose de	evice20.64 (35.80)	60 dose	Foradil
Foradil Powder for inhalation, 12 mcg per dose, and monodose	e device to be de	listed 1 July 2023	3)
FORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dos	se)10.32	60 dose OP	
(- 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	(16.90)		Oxis Turbuhaler
IDACATEROL	(/		
Powder for inhalation 150 mcg	61.00	30 dose OP	✓ Onbrez Breezhaler
Powder for inhalation 300 mcg		30 dose OP	✓ Onbrez Breezhaler
ALMETEROL		00 0000 01	• Olibioz Biooznaloi
ALMETEROL Aerosol inhaler CFC-free, 25 mcg per dose	26.25	120 dose OP	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated		60 dose OP	✓ Serevent Accuhaler
, 31			• Serevent Accumater
Inhaled Corticosteroids with Long-Acting Beta	a-Adrenocept	or Agonists	
UDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide	with		
6 mcg eformoterol fumarate metered dose)	41.50	120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fuma			
per dose (equivalent to 400 mcg budesonide with 12 m	ncg		
eformoterol fumarate metered dose) - No more than 2			
dose per day		120 dose OP	✓ DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6	mcg33.74	120 dose OP	✓ Symbicort Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
	mcg33.74	120 dose OP	✓ Symbicort
Powder for inhalation 200 mcg with eformoterol fumarate 6			Turbuhaler 200/6
Powder for inhalation 200 mcg with eformoterol fumarate 6			
Powder for inhalation 200 mcg with eformoterol fumarate 6 Powder for inhalation 400 mcg with eformoterol fumarate			
· ·	33.74	60 dose OP	✓ Symbicort
Powder for inhalation 400 mcg with eformoterol fumarate	33.74	60 dose OP	✓ Symbicort Turbuhaler 400/12
Powder for inhalation 400 mcg with eformoterol fumarate	33.74	60 dose OP	

20

✓ Duolin

	Subsidy (Manufacturer's I	Price) Subs	Fully Brand or idised Generic Manufacturer
FLUTICASONE WITH SALMETEROL			
Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.79	120 dose OP	✓ Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - No			
more than 2 dose per day	33.74	60 dose OP	 Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg - No)		
more than 2 dose per day	44.08	60 dose OP	Seretide Accuhaler
Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Oral lig 400 mcg per ml	40.00	150 ml	✓ Ventolin
Infusion 1 mg per ml, 5 ml		10	✓ Ventolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO	53.00	5	✓ Ventolin
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000			
dose available on a PSO	3 80	200 dose OP	✓ Respigen
doco avalladio on a 1 do		200 0000 01	✓ SalAir
	(6.20)		Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb)		
available on a PSO	8.96	20	✓ <u>Asthalin</u>
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule - Up to 30 neb			
available on a PSO	9.43	20	✓ <u>Asthalin</u>
TERBUTALINE SULPHATE			
Powder for inhalation, 200 mcg per dose (equivalent to			
250 mcg metered dose), breath activated	22.20	120 dose OP	Bricanyl Turbuhaler
Anticholinergic Agents			
PRATROPIUM BROMIDE			
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dos	SA		
available on a PSO		200 dose OP	✓ Atrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 nc		_00 0000 01	/ • · · · · · · · · · · · · · · · · ·
available on a PSO		20	✓ Univent
	28.20		✓ Accord S29
Inhaled Beta-Adrenoceptor Agonists with Antic	cholinergic A	gents	
SALBUTAMOL WITH IPRATROPIUM BROMIDE		-	
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg	nor		
dose CFC-free		200 dose OP	✓ Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per	12.10	_00 0000 01	PWVIII 111 A
wiel 0.5 ml amounts. The to 00 male and "to the control DOC	11.01	00	✓ Dualin

vial, 2.5 ml ampoule - Up to 20 neb available on a PSO 11.04

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg81.00 30 dose OP

Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg.......81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Antifibrotics

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

Subsidy		Fully	Brand or	
(Manufacturer's Price)	:	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib

Troto: I monidono lo not cabolaleca in combination with cabol	alood milloddino.		
Tab 801 mg	3,645.00	90	Esbriet
Tab 267 mg	1,215.00	90	Esbriet

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su Per	bsidised	Generic Manufacturer	
Ψ	1 01		Manadada et	

Leukotriene Receptor Antagonists

MC	NTELUKAST		
*	Tab 4 mg3.10	28	✓ Montelukast Mylan
*	Tab 5 mg3.10	28	✓ Montelukast Mylan
			✓ Montelukast Viatris
*	Tab 10 mg	28	✓ Montelukast Mylan
	·		✓ Montelukast Viatris

Methylxanthines

AMINOPHYLLINE

*	Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available on a		
	PSO180.00	5	✓ DBL Aminophylline
TH	EOPHYLLINE		
*	Tab long-acting 250 mg23.02	100	✓ Nuelin-SR
*	Oral lig 80 mg per 15 ml	500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA − Special Authority see SA1978 below − Retail pharmacy
Nebuliser soln, 2.5 mg per 2.5 ml ampoule250.00 6 ✓ Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

${\sf IVACAFTOR\ -PCT\ only-Specialist-Special\ Authority\ see\ SA2017\ below}$

Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at

	RESPIRAT	ORY SYSTE	EM AI	ND ALLERGIES
	Subsidy (Manufacturer's I	Price) Subs Per	Fully idised	Brand or Generic Manufacturer
continued				
least 1 allele; or 2.2 Patient must have other gating (class III) mutation and S549R) in the CFTR gene on at least 1 allele; 3 Patients must have a sweat chloride value of at least 60 n sweat collection system; and 4 Treatment with ivacaftor must be given concomitantly with 5 Patient must not have an acute upper or lower respiratory (including antibiotics) for pulmonary disease in the last 4 v 6 The dose of ivacaftor will not exceed one tablet or one sar 7 Applicant has experience and expertise in the manageme SODIUM CHLORIDE	and nmol/L by quant a standard thera infection, pulmoveeks prior to co chet twice daily;	py for this condi onary exacerbat ommencing treat and	e ionto	phoresis or by Macrodu ad changes in therapy
Not funded for use as a nasal drop. Soln 7%	24.50	90 ml OP	✓ B	liomed
Nasal Preparations				
Allergy Prophylactics				
BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose Metered aqueous nasal spray, 100 mcg per dose		200 dose OP 200 dose OP	_	teroClear teroClear
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ <u>F</u>	lixonase Hayfever & Allergy
PRATROPIUM BROMIDE Aqueous nasal spray, 0.03%	5.23	15 ml OP	√ <u>U</u>	nivent
Respiratory Devices				
MASK FOR SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO c) Only for children aged six years and under Small	2.20	1	√ e	-chamber Mask
PEAK FLOW METER a) Up to 25 dev available on a PSO b) Only on a PSO			·	
Low range	9.54	1	✓ N	lini-Wright AFS Low Range
Normal range	9.54	1	✓ N	lini-Wright Standard
SPACER DEVICE				- Candala
a) Up to 50 dev available on a PSO b) Only on a PSO				
220 ml (single patient)	2.95	1	√ e	-chamber Turbo
E10 ml (cinale notiont)	T 10	4		alaamalaan I a

e-chamber La Grande ✓ Volumatic

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per Manufacturer

Respiratory Stimulants

CAFFEINE CITRATE

Oral liq 20 mg per ml (10 mg base per ml)......15.10 25 ml OP **✔ Biomed**

	Subsidy (Manufacturer's F \$	rice) Subsi Per	Fully Brand or dised Generic Manufacturer
Ear Preparations			
FLUMETASONE PIVALATE Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform ED's ✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g		TIN 7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml	4.50 (9.27)	8 ml OP	Sofradex
FRAMYCETIN SULPHATE Ear/Eye drops 0.5%	4.13 (8.65)	8 ml OP	Soframycin
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expli	citly stated other	wise.	
Anti-Infective Preparations			
ACICLOVIR * Eye oint 3% CHLORAMPHENICOL	14.88	4.5 g OP	✓ <u>ViruPOS</u>
Eye oint 1% Eye drops 0.5%Funded for use in the ear*. Indications marked with * au	7.50	5 g OP 10 ml OP dications.	✓ <u>Devatis</u> ✓ Chlorafast
CIPROFLOXACIN Eye drops 0.3% — Subsidy by endorsement When prescribed for the treatment of bacterial keratitis of for the second line treatment of chronic suppurative otiting the second line treatment of chronic suppurative of the second line treatment of th	or severe bacteria s media (CSOM)		
GENTAMICIN SULPHATE Eye drops 0.3%(Genoptic Eye drops 0.3% to be delisted 1 August 2023) PROPAMIDINE ISETHIONATE	11.40	5 ml OP	✓ Genoptic
* Eye drops 0.1%	2.97 (14.55)	10 ml OP	Brolene

SODIUM FUSIDATE [FUSIDIC ACID]

TOBRAMYCIN

✓ Fucithalmic

✓ Tobrex

✓ Tobrex

5 g OP

3.5 g OP 5 ml OP



Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	•	Manufacturer	
				ě

Corticosteroids and Other Anti-Inflammatory Preparations

DE	XAMETHASONE			
*	Eye oint 0.1%	5.86	3.5 g OP	Maxidex
*	Eye drops 0.1%	4.50	5 ml OP	Maxidex
	Ocular implant 700 mcg - Special Authority see SA1680 below			
	- Retail pharmacy1,4	44.50	1	Ozurdex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin		·	
b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM			
Eye drops 0.1%	8.80	5 ml OP	✓ Voltaren Ophtha
FLUOROMETHOLONE			
* Eve drops 0.1%	3.09	5 ml OP	✓ FML
•	5.20		✓ Flucon

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price) Subs	sidised	Generic
	\$	Per	✓	Manufacturer
EVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
, , ,	(10.34)		L	ivostin
ODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	√ L	.omide
PREDNISOLONE ACETATE				
Eye drops 1%	6.92	10 ml OP	✓ P	rednisolone-AFT
, ,	7.00	5 ml OP	✓ P	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Author	ritv see SA1715 belov	v – Retail phar	macv	
Eye drops 0.5%, single dose (preservative free)		20 dose		Minims
, , , ,				Prednisolone

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

5 ml OP

✓ Betoptic S

SODIUM CROMOGLICATE	
---------------------	--

Eye drops 2%	1.79	5 ml OP	Rexacrom
	2.62	10 ml OP	Allerfix
(Daysaram Five drang 00/ to be delicted 1 March 0000)			

(Rexacrom Eye drops 2% to be delisted 1 March 2023)

DET	AXOL	\cap I
DEI	AXUL	_UL

ACETAZOL AMIDE

不	Eye drops 0.5%7.50	5 IIII OP	• веторис
TIN	10LOL		
*	Eye drops 0.25%	5 ml OP	✓ Arrow-Timolol
*	Eye drops 0.5%	5 ml OP	✓ Arrow-Timolol
*	Eye drops 0.5%, gel forming	2.5 ml OP	✓ Timoptol XE

Glaucoma Preparations - Carbonic Anhydrase Inhibitors

Eye drops 0.25%11.80

	Tab 250 mg	17.03	100	✓ Diamox
BR	INZOLAMIDE			

DORZOLAMIDE WITH TIMOLOL (17.44) Trusopt

Glaucoma Preparations - Prostaglandin Analogues

BIMATOPROST

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's P	rice) Subsi	Fully idised	Brand or Generic
	\$	Per	1	Manufacturer
LATANOPROST				
* Eye drops 0.005%	1.82	2.5 ml OP	1	<u>Teva</u>
TRAVOPROST				
* Eye drops 0.004%	9.75	2.5 ml OP	1	Travatan
			-	
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE				
* Eye drops 0.2%	4.29	5 ml OP	1	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE				
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	1	Combigan
LATANOPROST WITH TIMOLOL				Ū
Eye drops 0.005% with timolol 0.5%	2 49	2.5 ml OP	1	Arrow - Lattim
PILOCARPINE HYDROCHLORIDE			-	
* Eye drops 1%	4 26	15 ml OP	√ I	sopto Carpine
* Eye drops 2%		15 ml OP		sopto Carpine
* Eye drops 4%		15 ml OP		sopto Carpine
Subsidised for oral use pursuant to the Standard Formul		* ···· * ·	•	P
* Eye drops 2% single dose - Special Authority see SA0895				
below – Retail pharmacy	31.95	20 dose	√ [Minims Pilocarpine
. ,				•

⇒SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics		
* Eye drops 1%	15 ml OP	✓ Atropt
* Eye drops 1%	15 ml OP	✓ Cyclogyl
* Eye drops 0.5%	15 ml OP 15 ml OP	✓ Mydriacyl✓ Mydriacyl
Preparations for Tear Deficiency		
For acetylcysteine eye drops refer Standard Formulae, page 250 HYPROMELLOSE		
* Eye drops 0.5%	15 ml OP	✓ Methopt
HYPROMELLOSE WITH DEXTRAN ★ Eye drops 0.3% with dextran 0.1%	15 ml OP	✓ Poly-Tears

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Preservative Free Ocular Lubricants

⇒SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA2134 above – Retail pha	armacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL -	Special Authority	see <mark>SA213</mark> 4	above – Retail pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	 Systane Unit Dose
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml	10.78	30	 Systane Unit Dose
(Systane Unit Dose Eye drops 0.4% and propylene glycol 0.3%,	0.4 ml to be deliste	ed 1 June 20	023)
SODIUM HYALURONATE [HYALURONIC ACID] - Special Auth	nority see SA2134	above - Re	tail pharmacy
Eye drops 1 mg per ml	13.85	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Ph	armacy Procedure	s Manual re	estriction allowing one bottle per
month is not relevant and therefore only the prescribed of	dosage to the near	est OP may	be claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%4.15 (Naphcon Forte Eye drops 0.1% to be delisted 1 September 2023)	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%2.17	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%3.63	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g3.80	5 g OP	✓ VitA-POS



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

10

✓ Hameln

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE Inj 200 mg per ml, 10 ml ampoule52.88	10	✓ Martindale Pharma
NALOXONE HYDROCHLORIDE		
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
* Inj 400 mcg per ml, 1 ml ampoule22.60	5	✓ DBL Naloxone
		Hydrochloride

35.26

(DBL Naloxone Hydrochloride Inj 400 mcg per ml, 1 ml ampoule to be delisted 1 February 2023)

Removal and Elimination

CHARCOAL

* Oral lig 50 g per 250 ml	250 ml OF	Carbosorb-X
----------------------------	-----------	-------------

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	Exjade
Tab 500 mg dispersible	1,105.00	28	Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE – Special Authority see SA1480 on the next page – Retail pharmacy					
Tab 500 mg	533.17	100	✓ Ferriprox		
Oral liq 100 mg per 1 ml		250 ml OP	✓ Ferriprox		



Subsidy	ı	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE ★ Inj 500 mg vial	151.31	10	✓ DBL Desferrioxamine Mesylate for Inj BP ✓ Deferoxamine Pfizer S29 S29
SODIUM CALCIUM EDETATE * Inj 200 mg per ml, 5 ml	EQ 01	6	
* III] 200 IIIg pei IIII, 5 IIII		0	0.1.
	(156.71)		Calcium Disodium
			Versenate



Standard Formulae

Statitual di Fortificiae			
ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol	60 mg 40 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	
Preservative Water	qs to 100 ml	Phenobarbitone Sodium Glycerol BP Water	400 mg 4 ml to 40 ml
CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative Water	300 mg 40 ml qs to 100 ml	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative Water	qs qs to 500 ml
FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Water	1 tab qs to 500 ml	(Preservative should be used if quantity supplied is than 5 days.) SALIVA SUBSTITUTE FORMULA Methylcellulose Preservative	5 g
(Preservative should be used if quantity supplied is f than 5 days. Maximum 500 ml per prescription.)	oi more	Water (Preservative should be used if quantity supplied is	qs to 500 ml for more
METHADONE MIXTURE Methadone powder Glycerol Water	qs qs to 100 ml	than 5 days. Maximum 500 ml per prescription.) SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml	qs
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqu	10 g to 100 ml	Water (Only funded if prescribed for treatment of hyponatra VANCOMYCIN ORAL SOLUTION (50 mg per ml) Vancomycin 500 mg injection	qs aemia) 10 vials
OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml	Glycerol BP Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	40 ml to 100 ml

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or Generic (Manufacturer's Price) Subsidised Per Manufacturer \$

Extemporaneously Compounded Preparat	tions and Galenica	Is		
CODEINE PHOSPHATE - Safety medicine; prescriber m	nay determine dispensing	frequency		
Powder - Only in combination	63.09	25 g		
·	(90.09)	-	Douglas	
Only in extemporaneously compounded codeine	linctus		_	

COLLODION FLEXIBLE Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined. ✓ PSM 100 ml COMPOUND HYDROXYBENZOATE - Only in combination Only in extemporaneously compounded oral mixtures. 100 ml ✓ Midwest GLYCERIN WITH SODIUM SACCHARIN - Only in combination Only in combination with Ora-Plus. ✓ Ora-Sweet SF 473 ml GLYCERIN WITH SUCROSE - Only in combination Only in combination with Ora-Plus. 473 ml ✓ Ora-Sweet GI YCFROI

500 ml

✓ healthE Glycerol BP

✓ AET

METHADONE HYDROCHLORIDE

Powder

- a) Only on a controlled drug form
- b) No patient co-payment payable
- c) Safety medicine; prescriber may determine dispensing frequency

Only in extemporaneously compounded oral liquid preparations.

d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).

7 9/

Powder	ı g	♥ AFI
METHYL HYDROXYBENZOATE		
Powder8.98	25 g	✓ Midwest
METHYLCELLULOSE		
Powder36.95	100 g	✓ MidWest
Suspension - Only in combination30.95	473 ml	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN - Only in C	combination	
Suspension30.95	473 ml	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only in combination		
Suspension30.95	473 ml	Ora-Blend
PHENOBARBITONE SODIUM		
Powder - Only in combination52.50	10 g	✓ MidWest
325.00	100 g	✓ MidWest
Only in children up to 12 years		
PROPYLENE GLYCOL		
Only in extemporaneously compounded methyl hydroxybenzoate 10% solutio	n.	
Liq11.25	500 ml	✓ Midwest
SODIUM BICARBONATE		
Powder BP - Only in combination10.05	500 g	✓ Midwest
Only in extemporaneously compounded omeprazole and lansoprazole su	spension.	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatior	ns.			
Liq WATER	14.95	500 ml	✓ M	lidwest
Tap – Only in combination	0.00	1 ml	✓ Ta	ap water

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...



Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	ıbsidised	Generic
\$	Per	✓	Manufacturer

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)12.30	200 ml OP	✓ Calogen
30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)12.30	200 ml OP	✓ Calogen
Oil30.00	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml114.92	4 OP	✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT - Special Authority	y see SA1524 above – Hospital pha	rmacy [HP3]	
Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource
		-	Beneprotein

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see	e SA1095 above -	 Hospital pharm 	nacy [HP3]
Liquid	3.75	500 ml OP	✓ Glucerna Select
·	7.50	1,000 ml OP	✓ Nutrison Advanced Diason
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA	1095 above – Ho	spital pharmacy	
Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	2.10		✓ Nutren Diabetes

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED − Special Authority see SA1525 above − Hospital pharmacy [HP3]
Powder60.48 400 g OP ✓ Monogen

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

practitioner and date contacted.			
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority s Liquid		the previous pa 500 ml OP	ge – Hospital pharmacy [HP3] ✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see Liquid		ne previous page 500 ml OP	 Hospital pharmacy [HP3] Nutrini RTH Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Spec pharmacy [HP3]	ial Authority se	ee SA1379 on th	e previous page – Hospital
Liquid	6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see S Liquid (strawberry) Liquid (vanilla)	1.60	previous page – 200 ml OP 200 ml OP 500 ml OP	Hospital pharmacy [HP3] Fortini Fortini Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.07 1.07	evious page – F 200 ml OP 200 ml OP 200 ml OP 250 ml OP	dospital pharmacy [HP3] Pediasure Pediasure Pediasure Pediasure Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special A pharmacy [HP3]	uthority see SA	A1379 on the pr	evious page – Hospital
Liquid (unflavoured) Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.60 1.60	200 ml OP 200 ml OP 200 ml OP 200 ml OP	 ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 or Powder		page – Hospital 400 g OP	pharmacy [HP3] Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/M	L – Special Authority see SA1101 above –	 Hospital ph 	armacy [HP3]
Liquid	6.08	500 ml OF	✓ Nepro HP RTH

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per 🗸	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid		20 ml OP	oharmacy [HP3] Nepro HP (strawberry) Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110		•	armacy [HP3]
Liquid, 200 ml bottle	11.52 (13.24)	4 OP	NovaSource Renal
Liquid (apricot) 125 ml Liquid (caramel) 125 ml	11.52 [′]	4 OP ✓	Renilon 7.5 Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption: or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - Special Authority see SA Liquid	1377 above – Hospital pharmacy [HP3] 00 ml OP ✓ Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see SA1377 above - Ho	spital pharmacy [HP3]
Liquid (grapefruit), 250 ml carton171.00	18 OP ✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton171.00	18 OP ✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton171.00	18 OP ✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see SA1377 above - Hosp Powder (unflavoured)4.50 8	ital pharmacy [HP3] 0 g OP ✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Authority see SA1377 about 12.04 1,000 1,00	OO ml OP Volumination Advanced Peptisorb
	✓ Peptisorb

(Peptisorb Liquid to be delisted 1 June 2023)

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML -	Special Authority	see SA1196 at	ove – Ho	spital pharmacy [HP3]
Liquid	4.00	500 ml OP	✓ Nut	rini Low Energy
			M	ulti Fibre

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price)	S	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

	Subsidy (Manufacturer's \$		Fully Brand or dised Generic Manufacturer
ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 (Liquid		lospital pharmacy 250 ml OP 1,000 ml OP	/ [HP3] ✓ Ensure Plus HN ✓ Ensure Plus RTH ✓ Nutrison Energy
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 or Liquid		spital pharmacy [250 ml OP 1,000 ml OP	HP3] Isosource Standard Nutrison Standard RTH Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authori Liquid	•	on page 260 – Ho 1,000 ml OP	ospital pharmacy [HP3] Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority s Liquid		1,000 ml OP	ital pharmacy [HP3] ✓ Jevity RTH ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority Liquid		page 260 – Hos 1,000 ml OP	pital pharmacy [HP3] ✓ Jevity Plus
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority Liquid		page 260 – Hos 1,000 ml OP	pital pharmacy [HP3] ✓ Jevity HiCal RTH ✓ Nutrison Energy Multi Fibre
ORAL FEED (POWDER) – Special Authority see SA1859 on pa Powder (chocolate)	14.00	840 g OP	✓ Sustagen Hospital Formula
Powder (vanilla)	26.00 14.00	850 g OP 840 g OP	✓ Ensure ✓ Sustagen Hospital Formula Active
	26.00	850 g OP	✓ Ensure

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	•	Manufacturer	

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 260 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with	0.70	000 I OD	
Endorsement	0.72 (1.26) (1.26)	200 ml OP	Ensure Plus Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	Ensure Plus
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with	,	000 ml OD	Liisule Flus
Endorsement	0.72 (1.26)	200 ml OP	Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with Endorsement	0.85	237 ml OP	
LINOISEMENT	(1.33)		Ensure Plus
	0.72 (1.26)	200 ml OP	Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 260 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	/	Manufacturer	

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195	on the previous p	age - Hospital p	harmacy [HP3]
Liquid	5.50	500 ml OP	✓ Nutrison
			Concentrated
	11.00	1,000 ml OP	✓ Ensure Two Cal HN
			RTH

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

SPECIAL FOODS

	Subsidy		Fully	Brand or	
	(Manufacturer's Pric	e) Sub	sidised	Generic	
	\$	Per	•	Manufacturer	
FOOD THICKENER	- Special Authority see SA1106 on the previous page - Hospi	tal pharmac	y [HP3]		
Powder	6.53	300 g OP	✓ N	lutilis	
	7.25	380 g OP	√ F	eed Thickener	
				Karicare Antamil	

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA1729 al Powder		pharmacy [HP3] 1,000 g OP	
	(5.15)	, 3 -	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729 ab	ove - Hospital	pharmacy [HP3]	
Powder	3.93	1,000 g OP	
	(7.32)	-	NZB Low Gluten Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 above -	- Hospital phari	macy [HP3]	
Powder	5.62	2,000 g OP	
	(18.10)		Horleys Flour

	Subsidy (Manufacturer's Pr		Fully sidised	Brand or Generic
	\$	Per		Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page – H	lospital pharm	nacy [HF	23]
Buckwheat Spirals	2.00	250 g OP		•
	(3.11)		0	rgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)		0	rgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)		0	rgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)		0	rgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)		0	rgran
Rice and Corn Penne	2.00	250 g OP		
	(2.92)		0	rgran
Rice and Maize Pasta Spirals	2.00	250 g OP		
	(2.92)		0	rgran
Rice and Millet Spirals	2.00	250 g OP		
	(3.11)		0	rgran
Rice and corn spaghetti noodles	2.00	375 g OP		
	(2.92)		0	rgran
Vegetable and Rice Spirals	2.00	250 g OP		
	(2.92)		0	rgran
Italian long style spaghetti	2.00	220 g OP		
	(3.11)		0	rgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic

\$ Per ✓ Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE — Special Authority see SA1108 on the previous page — Hospital pharmacy [HP3]

armacy [HP3]			
Tabs		75 OP	Phlexy 10
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex
			Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior
, ,			Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex
		•	Powder
Powder (neutral) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (orange) 28 g sachets		30	✓ PKU Lophlex
1 owder (orange) 20 g sacriets	950.00	30	Powder
Powder (orange) 36 g sachet	202.00	30	✓ PKU Anamix Junior
rowder (orange) so g sacher	393.00	30	
Decides (conflict 00 marches	000.00	00	Orange
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior
			Vanilla
Infant formula		400 g OP	✓ PKU Anamix Infant
Powder (orange)		500 g OP	XP Maxamum
Powder (unflavoured)	320.00	500 g OP	XP Maxamum
Liquid (berry)	13.10	125 ml OP	PKU Anamix Junior
			LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior
			LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior
_ 1 ()			LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex
Oral contraction (beines) roo g		00 01	Sensation 20
Liquid (juicy berries) 62.5 ml	030.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
		60 OP	✓ PKU Lophiex LQ 10
Liquid (juicy orange) 62.5 ml			•
Liquid (juicy berries) 125 ml.		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING MIX – Special Authority see SA1108 on the previous page – Hospital pharmacy [HP3]

Powder8.22 500 g OP ✓ Loprofin Mix

LOW PROTEIN PASTA - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne	5.95	250 g OP	✓ Loprofin
Low protein rice pasta	11.91	500 g OP	✓ Loprofin
Macaroni		250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals	11.91	500 g OP	✓ Loprofin

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

MINO ACID FORMULA – Special Authority see SA2092 below – Hospital phar Powder43.60	400 g OP	✓ Alfamino
Powder (unflavoured)53.00	400 g OP	✓ Alfamino Junio ✓ Elecare ✓ Elecare LCP ✓ Neocate Gold
		✓ Neocate Junior Unflavoured
Powder (vanilla)53.00	400 g OP	✓ Neocate SYNEO ✓ Elecare ✓ Neocate Junior Vanilla

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis: or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency: or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA	- Special Authority see SA1953 below	– Hospitai phar	macy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
·			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption: or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

SPECIAL FOODS

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
- 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms: or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption: or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea: or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy (Manufacturer's Price)	Full Subsidise	
\$	Per 🗸	Manufacturer

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

 Powder (unflavoured)
 35.50
 300 g OP
 KetoCal 4:1

 V Ketocal 3:1
 KetoCal 4:1

 Powder (vanilla)
 35.50
 300 g OP
 KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent......0.00

0 ✓ BCG Vaccine

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
 - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old; or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.

Ini 2 III diphtheria toxoid with 20 III tetanus toxoid, 8 mg	Ini	211	I dinhtheria	toxoid with	20 11 1	tetanus	hioxot	8 mcc
--	-----	-----	--------------	-------------	---------	---------	--------	-------

pertussis toxoid. 8 mcg pertussis filamentous

✓ Boostrix

	Subsidy (Manufacturer's Price)	Subsid Per	lised (Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – Funded for any of the following:	[Xpharm]			
 A single dose for children up to the age of 7 who have c A course of four vaccines is funded for catch up program primary immunisation; or 	nmes for children (to	the age of	10 years	,
 An additional four doses (as appropriate) are funded for pre- or post splenectomy; pre- or post solid organ transpregimens; or 	plant, renal dialysis a			
Five doses will be funded for children requiring solid org				
Note: Please refer to the Immunisation Handbook for approp	riate schedule for cat	ch up progi	ammes.	
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units				
poliomyelitis virus in 0.5ml syringe	0.00	10	✓ Infa	nrix IPV
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AI [Xpharm]	ND HAEMOPHILUS I	NFLUENZ	AE TYPE	B VACCINE -
Funded for patients meeting any of the following criteria:				
1) Up to four doses for children up to and under the age of 2) An additional four doses (as appropriate) are funded for 10 who are patients post haematopoietic stem cell trans post solid organ transplant, renal dialysis and other seven	(re-)immunisation for plantation, or chemore erely immunosuppres	r children u therapy; pre sive regime	p to and e or post ens; or	splenectomy; pre- or
3) Up to five doses for children up to and under the age of				
Note: A course of up-to four vaccines is funded for catch up to complete full primary immunisation. Please refer to the Immunisation.				
programmes.				
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous				
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,				
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10	✓ Infa	nrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]				
One dose for patients meeting any of the following:				
1) For primary vaccination in children; or				
 An additional dose (as appropriate) is funded for (re-)im transplantation, or chemotherapy; functional asplenic; pi or post cochlear implants, renal dialysis and other sever 	re or post splenector	ny; pre- or p	ost solic	
 For use in testing for primary immunodeficiency disease paediatrician. 				medicine physician or
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg prefilled syringe plus vial 0.5 ml		1	✓ Hib	erix
HEPATITIS A VACCINE - [Xpharm]				
Funded for patients meeting any of the following criteria:				
 Two vaccinations for use in transplant patients; or 				
2) Two vaccinations for use in children with chronic liver di3) One dose of vaccine for close contacts of known hepatit				
Inj 1440 ELISA units in 1 ml syringe	0.00	1	✓ Hav	rix
Inj 720 ELISA units in 0.5 ml syringe		1		rix Junior
,				

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per		Manufacturer
HEPATITIS B RECOMBINANT VACCINE - [Xpharm]				
Inj 10 mcg per 0.5 ml prefilled syringe		1	✓ E	ngerix-B
Funded for patients meeting any of the following				
 for household or sexual contacts of known 				s; or
2) for children born to mothers who are hepat				
3) for children up to and under the age of 18 y				achieved a positive
serology and require additional vaccination	or require a primary course of	or vaco	ination; or	
4) for HIV positive patients; or5) for hepatitis C positive patients; or				
6) for patients following non-consensual sexu	al intercourse: or			
7) for patients following immunosuppression;	· ·			
8) for solid organ transplant patients; or	01			
for post-haematopoietic stem cell transplan	t (HSCT) patients; or			
10) following needle stick injury.	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			
Inj 20 mcg per 1 ml prefilled syringe		1	✓ E	ngerix-B
Funded for patients meeting any of the following		4:4		
 for household or sexual contacts of known for children born to mothers who are hepat 				S, Of
3) for children up to and under the age of 18 y				achieved a nositive
serology and require additional vaccination				domoved a positive
for HIV positive patients; or	or require a primary course t			
5) for hepatitis C positive patients; or				
6) for patients following non-consensual sexu	al intercourse; or			
for patients following immunosuppression;	or			
8) for solid organ transplant patients; or				
for post-haematopoietic stem cell transplan	it (HSCT) patients; or			
10) following needle stick injury; or				
11) for dialysis patients; or				
12) for liver or kidney transplant patients.				
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52	2 AND 58) VACCINE [HPV] -	- [Xph	arm]	
Any of the following:				
 Maximum of two doses for children aged 14 year 				
Maximum of three doses for patients meeting a	ny of the following criteria:			
1) People aged 15 to 26 years inclusive; or				
2) Either:				
People aged 9 to 26 years inclusive				
Confirmed HIV infection; or Transplant (including stem cell) notice	anta. ar			
Transplant (including stem cell) patie Maximum of four doors for people again 0 to 26				
3) Maximum of four doses for people aged 9 to 26	years inclusive post chemoti	ierapy	'	

	Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic	
	\$	Per 🗸	Manufacturer	
INFLUENZA VACCINE				

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

✓ Afluria Quad Junior (2022 formulation)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness:

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment for the supply of influenza vaccine ini 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	110.00 10	•	Afluria Quad
			(2022 formulation)

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable

С

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) People 55 to 64 years of age (inclusive) and is Māori or any Pacific ethnicity; or
- c) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes: or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- d) children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- e) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or
- f) children 3 to 12 years of age (inclusive), from 1 July 2022 to 31 December 2022;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy (Manufacturer's Price	e)	Fully Subsidised	Brand or Generic	
<u> </u>	Per	1	Manufacturer	

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.
- Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Either:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant: or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients: or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Ini 10 mcg of each meningococcal polysaccharide conjugated

to a total of approximately 55 mcg of tetanus toxoid carrier

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm] Either: A) Both: 1) Child is under one year of age; and 2) Any of the following: i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant: or ii) up to three doses for close contacts of meningococcal cases of any group; or iii) up to three doses for child who has previously had meningococcal disease of any group; or iv) up to three doses for bone marrow transplant patients; or v) up to three doses for child pre- and post-immunosuppression*; or B) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients; or v) up to two doses for person pre- and post-immunosuppression*. *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 12 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV. complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression*. Note: children under 12 months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine. *Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Neisvac-C 1 PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm] 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B. 7F, 9V, 14 and 23F; 3 mcg of pneumococcal

polysaccharide serotypes 4, 18C and 19F in 0.5 ml

prefilled syringe0.00

✓ Synflorix

10

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
- Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection; or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) cochlear implants or intracranial shunts: or
 - g) cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - I) diabetes; or
 - m) Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully idised	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - Either:	[Xpharm]			
Up to three doses (as appropriate) for patients with H chemotherapy; pre- or post-splenectomy or with func complement deficiency (acquired or inherited), cochle All of the following:	tional asplenia, pre- or p	oost-solid o	organ t	ransplant, renal dialysis,
 a) Patient is a child under 18 years for (re-)immuni b) Treatment is for a maximum of two doses; and c) Any of the following: 	sation; and			
i) on immunosuppressive therapy or radiatic immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; v) who are immune-suppressed following orgor or vi) with cochlear implants or intracranial shur vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more the prednisone of 2 mg/kg per day or greater, 20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks ge xi) with cardiac disease, with cyanosis or failuxii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	or gan transplantation (incl tts; or han two weeks, and wh or children who weigh r asthma treated with hig station; or ure; or	luding haei o are on a more than	matopo n equiv 10 kg (oietic stem cell transplant); valent daily dosage of on a total daily dosage of
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)		1	√ <u>P</u>	neumovax 23
For partially vaccinated or previously unvaccinated in 2) For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for appr	dividuals; or	tch-up prod	gramm	es.
Inj 80D antigen units in 0.5 ml syringe		1 ' `	' ' <u>I</u> I	
first dose to be administered in infants aged under 14 no vaccination being administered to children aged 2	•			
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	✓ <u>F</u>	Rotarix

	Subsidy (Manufacturer's Price) \$	Sub:	Fully sidised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm] Either:				
Maximum of one dose for primary vaccination for either	:			
a) Any infant born on or after 1 April 2016; or				
 b) For previously unvaccinated children turning 11 years varicella infection (chickenpox), or 	ears old on or after 1	July 2017	, who h	ave not previously had a
Maximum of two doses for any of the following:				
 a) Any of the following for non-immune patients: 				
 i) with chronic liver disease who may in future ii) with deteriorating renal function before trans iii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, o 	plantation; or r		ion; or	
v) for post exposure prophylaxis who are immu				
b) For patients at least 2 years after bone marrow tra				
 c) For patients at least 6 months after completion of d) For HIV positive non immune to varicella with mile 				
For patients with inborn errors of metabolism at ris varicella, or				
 f) For household contacts of paediatric patients who immune compromise where the household contact 	t has no clinical histo	ry of vario	cella, or	
g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure has no clinical history of varicella.				
* immunosuppression due to steroid or other immunosuppres	ssive therapy must be	for a trea	atment p	eriod of greater than
28 days				
Inj 1350 PFU prefilled syringe	0.00	1 10		<u>arivax</u> arivax
VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] - [Xpha	arm]			
Funded for patients meeting the following criteria:				
1) Two doses for all people aged 65 years				
Inj 50 mcg per 0.5 ml vial plus vial	0.00	1	√ S	hingrix
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATE		LES VAC		•
Funded for patients meeting the following criteria:			•	
1) One dose for all people aged 65 years				
Inj 19,400 PFU prefilled syringe plus vial	0.00	1 10		ostavax ostavax
		10	• 2	USIAVAX
Diagnostic Agents				
TUBERCULIN PPD [MANTOUX] TEST - [Xpharm]				
Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	√ <u>T</u>	<u>ubersol</u>

- Symbols -	Agents Used in the Treatment of		Amphotericin B	
3TC105	Poisonings	248	Amsacrine	152
7 MED NSHA Silver/Copper	Agrylin	152	AmsaLyo	
Short	Albendazole	89	Amsidine	152
- A -	Albey	234	Amzoate	
A-Scabies66	Albustix	76	Anaesthetics	121
Abacavir sulphate105	Alchemy	161	Anagrelide hydrochloride	152
Abacavir sulphate with	Alchemy Oxybutynin	75	Analgesics	122
lamivudine105	Aldurazyme	29	Anastrozole	170
Abacavir/Lamivudine Viatris105	Alecensa	160	Anatrole	170
Abiraterone acetate167	Alectinib	160	Andriol Testocaps	80
Acarbose11	Alendronate sodium	110	Androderm	80
Accarb11	Alendronate sodium with		Anoro Ellipta	238
Accuretic 1047	colecalciferol	110	Antabuse	145
Accuretic 2047	Alfacalcidol	33	Antacids and Antiflatulents	6
Acetazolamide245	Alfamino	269	Anthelmintics	89
Acetec46	Alfamino Junior	269	Antiacne Preparations	60
Acetic acid with hydroxyquinoline and	Alginic acid	6	Antiallergy Preparations	234
ricinoleic acid74	Alglucosidase alfa	26	Antianaemics	36
Acetylcysteine248	Alkeran	149	Antiandrogen Oral	
Aci-Jel74		149	Contraceptives	74
Aciclovir	Allerfix	245	Antiarrhythmics	48
Infection100	Allerpro Syneo 1	272	Antibacterials	89
Sensory243	Allerpro Syneo 2	272	Antibacterials Topical	60
Acidex	Allersoothe	235	Anticholinergic Agents	237
Acipimox54	Allmercap	151	Anticholinesterases	109
Acitretin66		118	Antidepressants	126
Aclasta113	Alpha-Adrenoceptor Blockers	46	Antidiarrhoeals	6
Actemra221	Alpha-Keri Lotion	64	Antiepilepsy Drugs	128
Actemra S29221	Alphamox	92	Antifibrinolytics, Haemostatics and	
Actinomycin D153	Alphamox 125	92	Local Sclerosants	37
Actrapid10	Alphamox 250	92	Antifibrotics	238
Actrapid Penfill10	Alprolix	37	Antifungals	96
Acupan122	Alu-Tab	6	Antifungals Topical	61
Adalimumab (Amgevita)177	Aluminium hydroxide	6	Antihistamines	235
Adalimumab (Humira - Alternative	Alvogen	49	Antihypotensives	49
brand) 186	Amantadine hydrochloride	120	Antimalarials	98
Adapalene60	Ambrisentan	57	Antimigraine Preparations	131
ADR Cartridge 1.823	Ambrisentan Mylan	57	Antinausea and Vertigo Agents	131
Adrenaline56	Ambrisentan Viatris	57	Antipruritic Preparations	62
Advantan63	Amgevita	177	Antipsychotics	
Advate39	Amiloride hydrochloride	52	Antiretrovirals	103
Adynovate40	Amiloride hydrochloride with		Antirheumatoid Agents	110
Afinitor230	furosemide	52	Antispasmodics and Other Agents	
Aflibercept192	Amiloride hydrochloride with		Altering Gut Motility	8
Afluria Quad	hydrochlorothiazide		Antithrombotic Agents	40
(2022 formulation) 277	Aminophylline	240	Antithymocyte globulin	
Afluria Quad Junior	Amiodarone hydrochloride		(equine)	. 177
(2022 formulation)277	Amisulpride	132	Antitrichomonal Agents	98
AFT-Pyrazinamide100			Antituberculotics and	
Agents Affecting the	Amlodipine		Antileprotics	
Renin-Angiotensin System 46			Antiulcerants	
Agents for Parkinsonism and Related	Amoxicillin		Antivirals	
Disorders 120	Amoxicillin with clavulanic acid	92	Anxiolytics	136

Anzatax	157	Atropine sulphate		Beta Scalp	6
Apidra		Cardiovascular	48	Beta-Adrenoceptor Agonists	
Apidra SoloStar		Sensory		Beta-Adrenoceptor Blockers	
APO-Atomoxetine		Atropt		Betadine	6
APO-Atomoxetine S29		Atrovent		Betadine Skin Prep	
Apo-Azithromycin		AU Synacthen		Betaferon	
Apo-Diltiazem CD		Aubagio		Betahistine dihydrochloride	
				Betaine	
Apo-Temozolomide		Augmentin			
Apomorphine hydrochloride		Aurorix		Betaloc CR	
Aprepitant		AutoSoft 30	23	Betamethasone dipropionate	6
Apresoline		AutoSoft 90		Betamethasone dipropionate with	
Aqueous cream		Avallon		calcipotriol	
Aratac		Avelox		Betamethasone sodium phosphate	
Arava		Avonex		with betamethasone acetate	
Arginine		Avonex Pen		Betamethasone valerate	
Aripiprazole		Azacitidine		Betamethasone valerate with sodiu	
Aripiprazole Sandoz	132	Azacitidine Dr Reddy's		fusidate [fusidic acid]	6
Aristocort	63	Azamun	170	Betaxolol	24
Arrotex-Prazosin S29	46	Azathioprine	170	Betnovate	6
Arrow - Clopid	40	Azilect	120	Betoptic	24
Arrow - Lattim	246	Azithromycin		Betoptic S	
Arrow-Amitriptyline		Azopt		Bexsero	
Arrow-Bendrofluazide		AZT		Bezafibrate	
Arrow-Brimonidine		- B -		Bezalip	
Arrow-Diazepam		B-D Micro-Fine	15	Bezalip Retard	
Arrow-Doxorubicin		B-D Ultra Fine		Bicalutamide	
Arrow-Fluoxetine		B-D Ultra Fine II		Bicillin LA	10
Arrow-Losartan &	127	Bacillus Calmette-Guerin (BCG)	10	BiCNU	
Hydrochlorothiazide	47	vaccine	177	Bile and Liver Therapy	
•		Bacillus Calmette-Guerin	177	Biltricide	
Arrow-Norfloxacin			074		
		vaccine		Bimatoprost	
Arrow-Quinapril 10		Baclofen		Bimatoprost Multichem	
Arrow-Quinapril 20		Bactroban		Binarex	
Arrow-Quinapril 5		Balance		Binocrit	
Arrow-Roxithromycin		Barrier Creams and Emollients		Biodone	
Arrow-Timolol		BCG Vaccine		Biodone Extra Forte	
Arrow-Topiramate		Beclazone 100		Biodone Forte	
Arrow-Tramadol	126	Beclazone 250	235	Bisacodyl	
Arsenic trioxide		Beclazone 50	235	Bisacodyl Viatris	
Asacol		Beclomethasone dipropionate	235	Bisoprolol fumarate	4
Ascorbic acid	33	Bee venom allergy treatment	234	Bisoprolol Mylan	49
Aspen Adrenaline	56	Bendamustine hydrochloride	147	BK Lotion	
Aspirin		Bendrofluazide	53	Bleomycin sulphate	15
Blood	40	Bendroflumethiazide		Blood Colony-stimulating	
Nervous		[Bendrofluazide]	53	Factors	4
Asthalin	237	Benralizumab		Blood glucose diagnostic test	
Atazanavir Mylan	105	Benzathine benzylpenicillin		meter	14
Atazanavir sulphate		Benzatropine mesylate		Blood glucose diagnostic test	
Atenolol		Benzbromaron AL 100		strip	14
Atenolol AFT		Benzbromarone		Blood glucose test strips (visually	!
Atenolol AFT S29		Benztrop		impaired)	41
ATGAM		Benzydamine hydrochloride		Blood Ketone Diagnostic Test	15
			32		44
Ativan		Benzylpenicillin sodium [Penicillin	00	Strip	1
Atnahs Olsalazine		G]		Bonjela	
Atomoxetine		Beta Cream		Boostrix	
Atorvastatin	54	Beta Ointment	62	Bortezomib	15

INDEX: Generic Chemicals and Brands

Bortezomib Dr-Reddy's153	Capercit150	Chlorpheniramine maleate23
Bosentan57	Capoten46	Chlorpromazine hydrochloride 13
Bosentan Dr Reddy's57	Capsaicin	Chlortalidone [Chlorthalidone]5
Bplex33	Musculoskeletal110	Chlorthalidone5
Breo Ellipta236	Nervous122	Chlorvescent4
Brevinor 1/2873	Captopril46	Choice 380 7med Nsha Silver/copper
Bricanyl Turbuhaler237	Carafate10	Short
Brilinta40	Carbaccord148	Choice Load 3757
Brimonidine tartrate246	Carbamazepine128	Choice TT380 Short7
Brimonidine tartrate with timolol	Carbimazole82	Choice TT380 Standard7
maleate246	Carbomer247	Choline salicylate with cetalkonium
Brinzolamide245	Carboplatin148	chloride3
Brolene243	Carboplatin Ebewe148	Ciclosporin23
Brufen SR109	Carbosorb-X248	Cilazapril4
Buccastem132	Cardinol LA50	Cilicaine9
Budesonide	Cardizem CD51	Cilicaine VK9
Alimentary6	CareSens Dual14	Cinacalcet7
Respiratory235, 241	CareSens N14–15	Cipflox9
Budesonide with eformoterol236	CareSens N POP14	Ciprofloxacin
Bumetanide51	CareSens N Premier14	Infection9
Buprenorphine Naloxone BNM 144	CareSens PRO15	Sensory24
Buprenorphine with naloxone144	Carmellose sodium with gelatin and	Ciprofloxacin Teva24
Bupropion hydrochloride144	pectin32	Cisplatin14
Burinex51	Carmustine148	Cisplatin Ebewe14
Burinex S2951	Carnitor29	Citalopram hydrobromide12
Buscopan8	Carvedilol49	Cladribine15
Buspirone hydrochloride136	Carvedilol Sandoz49	Clarithromycin
Buspirone Viatris136	Casirivimab and imdevimab194	Alimentary
Busulfan148	Catapres51	Infection9
- C -	CeeNU149	Clexane4
Cabergoline87	Cefaclor monohydrate89	Clexane Forte4
Caffeine citrate242	Cefalexin89	Climara8
Calamine62	Cefazolin89	Clindamycin9
Calamine-AFT62	Ceftriaxone89	Clinicians2
Calci-Tab 50034	Ceftriaxone-AFT89	Clinicians Renal Vit3
Calcipotriol67	Cefuroxime axetil90	Clobazam12
Calcitonin77	Celapram127	Clobetasol propionate62, 6
Calcitriol33	Celebrex109	Clobetasone butyrate6
Calcitriol-AFT33	Celecoxib109	Clofazimine9
Calcium 500 mg Hexal34	Celecoxib Pfizer109	Clomazol
Calcium carbonate6, 34	Celestone Chronodose78	Dermatological6
Calcium Channel Blockers50	Cellcept170	Genito-Urinary7
Calcium Disodium Versenate249	Centrally-Acting Agents51	Clomifene citrate8
Calcium folinate150	Cephalexin ABM89	Clomipramine hydrochloride12
Calcium Folinate Ebewe150	Cetirizine hydrochloride235	Clomipramine Teva12
Calcium Folinate Sandoz150	Cetomacrogol64	Clonazepam128, 13
Calcium Folinate Sandoz S29 150	Cetomacrogol with glycerol64	Clonidine5
Calcium gluconate34	Cetomacrogol-AFT64	Clonidine hydrochloride5
Calcium Homeostasis77	Cetuximab194	Clonidine Teva5
Calcium polystyrene sulphonate45	Charcoal248	Clopidogrel4
Calcium Resonium45	Chemotherapeutic Agents147	Clopidogrel Multichem4
Calogen255	Chickenpox vaccine283	Clopine13
Candesartan cilexetil47	Chlorafast243	Clopixol134, 13
Candestar47	Chlorambucil148	Clotrimazole
Canesten61	Chloramphenicol243	Dermatological6
Capecitabine150	Chlorothiazide53	Genito-Urinary7

Clozapine	133	Cyclizine lactate	131	DBL Naloxone Hydrochloride	24
Clozaril	133	Cyclogyl	246	DBL Pethidine Hydrochloride	
Co-trimoxazole	96	Cyclonex		DBL Vincristine Sulfate	160
Coal tar	67	Cyclopentolate hydrochloride	246	Decozol	3
Coal tar with allantoin, menthol,		Cyclophosphamide		Deferasirox	
phenol and sulphur	67	Cyclorin		Deferiprone	24
Coal tar with salicylic acid and		Cycloserine		Deferoxamine Pfizer S29	24
sulphur	67	Cyproterone acetate		Denosumab	110
Coco-Scalp		Cyproterone acetate with		Deolate	9
Codeine phosphate		ethinyloestradiol	74	Deoxycoformycin	
Extemporaneous	251	Cystadane		Depo-Medrol	
Nervous		Cytarabine		Depo-Provera	
Coenzyme Q10	28	Cytotec		Depo-Testosterone	8
Colchicine		Cytoxan		Deprim	9
Colecalciferol	33	- D -		Dermol	
Colestid		D-Penamine	110	Desferrioxamine mesilate	
Colestipol hydrochloride		Dabigatran		Desmopressin	
Colgout		Dacarbazine		Desmopressin acetate	
Colifoam		Dacarbazine APP		Desmopressin-PH&T	
Colistin sulphomethate		Dactinomycin [Actinomycin D]		Desuric	
Colistin-Link		Daivobet		Detection of Substances in	
Collodion flexible		Daivonex		Urine	70
Colloidal bismuth subcitrate		Daktarin		Dexamethasone	
Colofac		Dalacin C		Hormone	7
Coloxyl		Dantrium		Sensory	
Combigan		Dantrium S29		Dexamethasone phosphate	
Compound electrolytes		Dantrolene		Dexamethasone Phosphate	
Compound electrolytes with gluc		Daonil		Panpharma	79
[Dextrose]		Dapa-Tabs		Dexamethasone with framycetin	
Compound hydroxybenzoate		Dapsone		gramicidin	
Comtan		Daraprim	95	Dexamethasone with neomycin	
Concerta		Darunavir		sulphate and polymyxin B	
Condoms		Darunavir Mylan		sulphate	24
Condyline		Darunavir Viatris		Dexamfetamine sulfate	
Contraceptives - Hormonal		Dasatinib		Dexmethsone	
Contraceptives - Non-hormonal		Daunorubicin		Dextrochlorpheniramine	
Copaxone		Daunorubicin Zentiva		maleate	23
Cordarone-X		David One Step Cassette Pregn		Dextrose	
Corticosteroids and Related Age		Test		DHC Continus	
for Systemic Use		DBL Adrenaline		Diabetes	
Corticosteroids Topical		DBL Aminophylline		Diabetes Management	
Cortifoam		DBL Bleomycin Sulfate		Diacomit	130
Cosentyx		DBL Bortezomib		Diagnostic Agents	
Cosmegen		DBL Carboplatin		Diamide Relief	
Coumadin		DBL Cisplatin		Diamox	
Country Life		DBL Dacarbazine		Diasip	
Coversyl		DBL Desferrioxamine Mesylate t		Diazepam1	
Creon 10000		BP		Diazoxide	
Creon 25000		DBL Docetaxel		Dibenzyline	
Creon Micro		DBL Ergometrine		Diclofenac Sandoz	100
Crotamiton		DBL Gemcitabine		Diclofenac Sandoz Diclofenac sodium	10
Crystaderm		DBL Gernchabine DBL Gentamicin		Musculoskeletal	100
Curam					
Curam Duo 500/125		DBL Heparin Sodium DBL Leucovorin Calcium		SensoryDifferin	24
Cvite		DBL Methotrexate Onco-Vial		Difflam	
Cyclizine hydrochloride	131	DBL Morphine Sulphate	125	Diflucan	9

INDEX: Generic Chemicals and Brands

Digestives Including Enzymes24	Drugs Affecting Bone	Enalapril maleate46
Digoxin48	Metabolism 110	Enbrel170
Dihydrocodeine tartrate124	Dual blood glucose and blood ketone	Endocrine Therapy167
Dilantin129	diagnostic test meter14	Endoxan148
Dilantin Infatab129	Dulaglutide12	Engerix-B276
Diltiazem CD Clinect51	Dulcolax SP Drop26	Enlafax XR127
Diltiazem hydrochloride51	Duocal Super Soluble Powder254	Enoxaparin sodium42
Dimethicone63, 65	Duolin237	Enstilar67
Dimethyl fumarate137	Duolin HFA237	Ensure263
Dipentum8	DuoResp Spiromax236	Ensure Plus264
Diphtheria, tetanus and pertussis	Duride56	Ensure Plus HN263
vaccine274	Durvalumab227	Ensure Plus RTH
Diphtheria, tetanus, pertussis and	- E -	Ensure Two Cal HN RTH265
polio vaccine	e-chamber La Grande241	Entacapone
Diphtheria, tetanus, pertussis, polio,	e-chamber Mask241	Entecavir
hepatitis B and haemophilus	e-chamber Turbo241	Entecavir Mylan
influenzae type B vaccine 275	E-Mycin91	Entecavir Naylari 100
Diprosone	e5 Pharma10	Entocort CIR6
		Entresto 24/26
Diprosone OV	Ear Preparations	
Dipyridamole	Ear/Eye Preparations	Entresto 49/51
Disopyramide phosphate	Easiphen Liquid	Entresto 97/103
Disulfiram145	Econazole nitrate61	Epilim
Diuretics51	Efavirenz104	Epilim Crushable
Dizole96	Efavirenz with emtricitabine and	Epilim IV
Docetaxel	tenofovir disoproxil	Epilim S/F Liquid129
Docetaxel Accord153	Eformoterol fumarate236	Epilim Syrup129
Docetaxel Sandoz153	Eformoterol fumarate dihydrate 236	Epirubicin Ebewe154
Docusate sodium25	Eftrenonacog alfa [Recombinant	Epirubicin hydrochloride154
Docusate sodium with	factor IX] 37	Eplerenone52
sennosides25	Efudix69	Epoetin alfa36
Dolutegravir106	Egopsoryl TA67	Epoprostenol59
Domperidone 131	Elaprase28	Eptacog alfa [Recombinant factor
Domperidone Viatris131	Eldepryl120	VIIa]39
Donepezil hydrochloride143	Elecare269	Erbitux194
Donepezil-Rex143	Elecare LCP269	Ergometrine maleate74
Dornase alfa240	Electral45	Erlotinib161
Dortimopt245	Elelyso31	Erythrocin IV91
Dorzolamide hydrochloride245	Elemental 028 Extra259	Erythromycin (as lactobionate)91
Dorzolamide with timolol245	Elidel67	Erythromycin ethyl succinate91
Dostinex87	Elocon63	Esbriet239
Dosulepin [Dothiepin]	Elocon Alcohol Free63	Escitalopram127
hydrochloride126	Eltrombopag37	Escitalopram (Ethics)127
Dosulepin Mylan126	Eltroxin82	Eskazole89
Dosulepin Viatris126	EMB Fatol99	Essential Ethosuximide128
Dothiepin	Emend Tri-Pack131	Essential Generics49
Doxazosin46	Emicizumab38	Essential Prednisolone8
Doxazosin Clinect46	EMLA122	Estraderm MX81
Doxine93	Empagliflozin	Estradiol TDP Mylan81
Doxorubicin Ebewe	Empagliflozin with metformin	Estradot
Doxorubicin hydrochloride153	hydrochloride13	Estradot 50 mcg81
Doxycycline93	Emtricitabine105	Estrofem
DP Lotion64	Emtricitabine with tenofovir	Etanercept
DP Lotn HC	disoproxil	Ethambutol hydrochloride99
DP-Allopurinol	Emtriva	Ethics Aspirin 122
Dr Reddy's Omeprazole9	Emulsifying ointment	Ethics Aspirin EC
Drofate50	Emulsifying Ointment ADE64	Ethics Lisinopril46

Ethinyloestradiol	82	Ferrous fumarate with folic acid.	34	Food Thickeners	265
Ethinyloestradiol with		Ferrous sulfate	35	Foods And Supplements For Inb	orn
desogestrel	72	Fexofenadine hydrochloride	235	Errors Of Metabolism	267
Ethinyloestradiol with		Fibro-vein	40	Foradil	236
levonorgestrel	73	Filgrastim	43	Forteo	112
Ethinyloestradiol with		Finasteride		Fortini	258
norethisterone	73	Fingolimod	137	Fortini Multi Fibre	258
Ethosuximide	128	Firazyr	234	Fortisip	264
Etopophos	154	Flagyl	98	Fortisip Multi Fibre	264
Etoposide	154	Flagyl-S	98	Fosamax	110
Etoposide phosphate	154	Flamazine	61	Fosamax Plus	
Etravirine	104	Flecainide acetate	48	Framycetin sulphate	243
Eumovate	62	Flecainide BNM	48	Frisium	
Evara	64	Flecainide Controlled Release		Frumil	52
Everet	129	Teva		Frusemide	52
Everolimus	230	Fleet Phosphate Enema	26	Fucicort	63
Evista		Flixonase Hayfever & Allergy	241	Fucidin	
Evusheld	221	Flixotide	236	Fucithalmic	243
Exemestane		Flixotide Accuhaler	236	Fulvestrant	167
Exjade	248	Florinef	79	Fungilin	
Extemporaneously Compound	ded	Fluanxol	134	Furosemid-Ratiopharm	
Preparations and		Flucil		Furosemide [Frusemide]	52
Galenicals		Flucloxacillin		Furosemide-Baxter	52
Eye Preparations	243	Flucloxacillin-AFT	92	fusidic acid	
Eylea		Flucloxin		Dermatological	
Ezetimibe	55	Flucon	244	Infection	
Ezetimibe Sandoz		Fluconazole		Sensory	243
Ezetimibe with simvastatin	55	Fludara Oral		- G -	
-F-		Fludarabine Ebewe		Gabapentin	128
Factor eight inhibitor bypassin	g	Fludarabine phosphate		Gacet	
fraction		Fludrocortisone acetate		Galsulfase	
Famotidine		Fluids and Electrolytes		Galvumet	
Famotidine Hovid		Flumetasone pivalate	243	Galvus	
Fasenra		Fluocortolone caproate with		Gardasil 9	
Faslodex		fluocortolone pivalate and	_	Gastrodenol	
Fatty Cream AFT		cinchocaine		Gaviscon Double Strength	
Febuxostat		Fluorometholone		Gaviscon Infant	
Febuxostat multichem	118	Fluorouracil		Gazyva	
Feed Thickener Karicare	000	Fluorouracil Accord		Gefitinib	
Aptamil		Fluorouracil sodium		GEM Aqueous Cream	
FEIBA NF		Fluox		Gemcitabine Ebewe	
Felo 10 ER		Fluoxetine hydrochloride		Gemcitabine hydrochloride	
Felo 5 ER		Flupenthixol decanoate		Gemtuzumab ozogamicin	
Felodipine		Flutamide		Genoptic	243
Femme-Tab ED		Flutamin		Gentamicin sulphate	0.4
Fenpaed 100 mg per 5 ml		Fluticasone	236	Infection	
Fentanyl		Fluticasone furoate with	000	Sensory	
Fentanyl Sandoz		vilanterol		Gilenya	
Ferinject		Fluticasone propionate		Ginet	
Ferodan		Fluticasone with salmeterol		Glatiramer acetate	
Ferriprox		Flynn		Glecaprevir with pibrentasvir	
Ferro-F-Tabs		FML		Glibenclamide	
Ferro-tab		Foban		Gliclazide	
Ferrograd		Folic acid		Glipizide	
FerrosigFerrous fumarate		Folic Acid Mulan		GlivecGlizide	162
renous iumarate	34	Folic Acid Mylan	3/	GIIZIQE	11

Glucagen Hypokit10	Horleys Bread Mix266	Idarubicin hydrochloride154
Glucagon hydrochloride10	Horleys Flour266	Idursulfase28
Glucerna Select256	Hormone Replacement Therapy -	Ifosfamide148
Glucose [Dextrose]44	Systemic81	Igroton53
Gluten Free Foods266	HPV276	Ikorel57
Glycerin with sodium saccharin251	Humalog11	Iloprost59
Glycerin with sucrose251	Humalog Mix 2511	Imatinib mesilate162
Glycerol	Humalog Mix 5011	Imatinib-Rex162
Alimentary26	Human papillomavirus (6, 11, 16, 18,	Imbruvica154
Extemporaneous251	31, 33, 45, 52 and 58) vaccine	Imfinzi227
Glyceryl trinitrate	[HPV] 276	Imigran131
Alimentary8	Humatin95	Imipramine hydrochloride126
Cardiovascular56	Humira186	Imiquimod69
Glycopyrronium238	HumiraPen186	Immune Modulators106
Glycopyrronium bromide8	Humulin 30/7011	Immunosuppressants170
Glycopyrronium with	Humulin NPH10	Incruse Ellipta238
indacaterol238	Humulin R10	Indacaterol236
Go Healthy28	Hyaluronic acid247	Indapamide53
Gold Knight71	Hydralazine56	Infanrix IPV275
Gold Knight XL71–72	Hydralazine hydrochloride56	Infanrix-hexa275
Goserelin87	Hydrocortisone	Infant Formulae269
Gutron49	Dermatological62	Infatrini273
Gynaecological Anti-infectives74	Hormone79	Infliximab195
-H-	Hydrocortisone (PSM)62	Influenza vaccine277
Habitrol145	Hydrocortisone acetate7	Inhaled Corticosteroids235
Haemophilus influenzae type B	Hydrocortisone acetate with	Inhaled Long-acting
vaccine275	pramoxine hydrochloride7	Beta-adrenoceptor Agonists 236
Haldol134	Hydrocortisone and paraffin liquid	Inspra52
Haldol Concentrate134	and lanolin 62	Instillagel Lido121
Haldol Decanoas134	Hydrocortisone butyrate63, 68	Insulin aspart11
Haloperidol133	Hydrocortisone with cinchocaine8	Insulin aspart with insulin aspart
Haloperidol decanoate134	Hydrocortisone with miconazole63	protamine10
Harvoni102	Hydrocortisone with natamycin and	Insulin glargine11
Havrix275	neomycin 63	Insulin glulisine11
Havrix Junior275	Hydrogen peroxide60	Insulin isophane10
Haylor syrup235	Hydroxocobalamin32	Insulin isophane with insulin
healthE Dimethicone 10%63	Hydroxocobalamin Panpharma32	neutral11
healthE Dimethicone 4% Lotion65	hydroxycarbamide 154	Insulin lispro11
healthE Dimethicone 5%63	Hydroxychloroquine110	Insulin lispro with insulin lispro
healthE Glycerol BP251	Hydroxyurea	protamine11
healthE Urea Cream64	[hydroxycarbamide] 154	Insulin neutral10
Healtheries Simple Baking Mix266	Hygroton53	Insulin pen needles15
Hemastix76	Hylo-Fresh247	Insulin pump16
Hemlibra38	Hymenoptera234	Insulin pump cartridge21
Heparin sodium43	Hyoscine butylbromide8	Insulin pump infusion set (steel
Heparinised saline43	Hyoscine hydrobromide132	cannula)21
Heparon Junior257	Hypam138	Insulin pump infusion set (steel
Hepatitis A vaccine275	Hyperuricaemia and Antigout 118	cannula, straight insertion) 21
Hepatitis B recombinant	Hypromellose246	Insulin pump infusion set (teflon
vaccine	Hypromellose with dextran246	cannula)22
Herceptin225	-1-	Insulin pump infusion set (teflon
Hiberix275	Ibiamox92	cannula, angle insertion with
Hiprex108	Ibrance163	insertion device)23
Histaclear235	Ibrutinib154	Insulin pump infusion set (teflon
Histafen235	Ibuprofen109	cannula, angle insertion) 23
Holoxan148	Icatibant234	Insulin pump infusion set (teflon

cannula, straight insertion with		Kaletra	106	Laxsol	25
insertion device)	23	Kalydeco	240	Ledipasvir with sofosbuvir	102
Insulin pump infusion set (teflon		Kemadrin	120	Leflunomide	110
cannula, straight insertion)	23	Kenacomb	243	Lenalidomide	155
Insulin pump reservoir	23	Kenacort-A 10	80	Letrole	170
Insulin syringes, disposable with		Kenacort-A 40	80	Letrozole	
attached needle	16	Kenalog in Orabase		Leukeran FC	
Intelence		Ketocal 3:1		Leukotriene Receptor	
Interferon beta-1-alpha		KetoCal 4:1		Antagonists	240
Interferon beta-1-beta		Ketoconazole		Leuprorelin	87
Intra-uterine device		Dermatological	68	Leustatin	
Invega Sustenna		Infection		Levetiracetam	
Invega Trinza		Ketogenic Diet		Levetiracetam-AFT	
IPCA-Frusemide		Ketoprofen		Levlen ED	
IPCA-Metoprolol		KetoSens		Levocabastine	
IPCA-Propranolol		Keytruda		Levocarnitine	
·		•		Levodopa with benserazide	
IPOL		Kindergen			
Ipratropium bromide237		Kivexa	105	Levodopa with carbidopa	
Iressa		Klacid	•	Levomepromazine	133
Irinotecan Actavis 100		Alimentary		Levomepromazine	400
Irinotecan hydrochloride		Infection		hydrochloride	133
Irinotecan-Rex		Kliogest		Levonorgestrel	
Iron (as ferric carboxymaltose)		Kliovance		Genito-Urinary	73–74
Iron polymaltose		Kogenate FS		Hormone	
Isentress		Konakion MM		Levonorgestrel BNM	
Isentress HD		Konsyl-D	25	Levothyroxine	82
Ismo 20	56	Kuvan	<mark>30</mark>	Lidocaine [Lignocaine]	121–122
Ismo 40 Retard	56	-L-		Lidocaine [Lignocaine]	
Isoniazid	99	Labetalol	49	hydrochloride	122
Isoniazid with rifampicin	99	Lacosamide	128	Lidocaine [Lignocaine] with	
Isoptin	51	Lactulose	26	chlorhexidine	122
Isoptin Retard	51	Laevolac	<mark>26</mark>	Lidocaine [Lignocaine] with	
Isoptin SR		Lagevrio	103	prilocaine	122
Isopto Carpine		Lamictal	129	Lidocaine-Baxter	122
Isosorbide mononitrate		Lamivudine	100. 105	Lidocaine-Claris	
Isosource Standard		Lamivudine Alphapharm		Life Extension	
Isotretinoin		Lamotrigine		Lignocaine	
Ispaghula (psyllium) husk		Lamprene		Lioresal Intrathecal	
Itch-Soothe		Lanoxin		Lipid-Modifying Agents	
Itraconazole		Lanoxin Paediatric Elixir		Liquigen	
Itrazole		Lanoxin PG		Lisinopril	46
Ivacaftor		Lanoxin S29		Litak	
Ivermectin		Lansoprazole		Lithium carbonate	
- J -	00	Lantus		Livostin	
Jadelle	72	Lantus SoloStar		LMX4	
				Locacorten-Viaform ED's	
Jakavi Jardiamet		Lanvis Lanzol Relief			
				Local preparations for Anal an	
Jardiance		Lapatinib ditosylate		Rectal Disorders	
Jaydess		Largactil		Locasol	
Jevity HiCal RTH		Laronidase		Locoid	
Jevity Plus		Lasix		Locoid Crelo	
Jevity RTH		Latanoprost		Locoid Lipocream	
Jinarc		Latanoprost with timolol		Locorten-Vioform	
Juno Pemetrexed	152	Lax-Suppositories		Lodoxamide	
- K -		Lax-suppositories Glycerol		Logem	
Kadcyla	226	Laxatives	25	Lomide	245

Lomustine	149	Megestrol acetate168	Metoclopramide hydrochloride	13
Loniten	56	Melatonin137	Metolazone	5
Loperamide hydrochloride	<mark>6</mark>	Melpha149	Metopirone	8
Lopinavir with ritonavir	106	Melphalan149	Metoprolol IV Mylan	4
Lopinavir/Ritonavir Mylan	106	Menactra279	Metoprolol succinate	4
Loprofin		Meningococcal (groups A, C, Y and	Metoprolol tartrate	4
Loprofin Mix	268	W-135) conjugate vaccine 279	Metrogyl	9
Lorafix		Meningococcal B multicomponent	Metronidazole	9
Loratadine		vaccine 280	Metyrapone	8
Lorazepam	136	Meningococcal C conjugate	Mexiletine hydrochloride	4
Lorstat	54	vaccine280	Miacalcic	7
Losartan Actavis	47	MenQuadfi279	Micolette	2
Losartan potassium	47	Menthol62	Micolette-S29	2
Losartan potassium with		Mepolizumab202	Miconazole	3
hydrochlorothiazide	47	Mercaptopurine151	Miconazole nitrate	
Lovir		Mercilon 2872	Dermatological	6
Loxamine	127	Mesalazine7	Genito-Urinary	7
Lucrin Depot 1-month	87	Mesna155	Micreme	
Lucrin Depot 3-month		Mestinon109	Micreme H	6
Lyderm	66	Metabolic Disorder Agents26	Microgynon 20 ED	7
Lynparza	156	Metformin hydrochloride11	Microgynon 30	7
Lyrica		Metformin Mylan11	Microlut	7
- M -		Metformin Viatris11	Midazolam	13
m-Eslon	125	Methadone BNM124	Midazolam-Baxter	13
Mabthera	206	Methadone hydrochloride	Midodrine	4
Macro Organic Psyllium Husk .	25	Extemporaneous251	Mifegyne	7
Macrobid	108	Nervous124	Mifepristone	7
Macrogol 3350 with potassium		Methatabs124	Milpharm	12
chloride, sodium bicarbonat	e and	Methenamine (hexamine)	Minerals	3
sodium chloride		hippurate108	Mini-Wright AFS Low Range	24
Madopar 125	120	Methopt246	Mini-Wright Standard	24
Madopar 250	120	Methotrexate151	Minidiab	1
Madopar 62.5	120	Methotrexate DBL Onco-Vial151	MiniMed 1.8 Reservoir	
Madopar HBS		Methotrexate Ebewe151	MMT-326A	2
Madopar Rapid	120	Methotrexate Sandoz151	MiniMed 3.0 Reservoir	
Magnesium hydroxide	35	Methyl hydroxybenzoate251	MMT-332A	2
Magnesium sulphate	35	Methylcellulose251	MiniMed 770G	1
Mantoux	283	Methylcellulose with glycerin and	MiniMed Mio MMT-921A	
Marevan	43	sodium saccharin251	MiniMed Mio MMT-923A	2
Marine Blue Lotion SPF 50+	69	Methylcellulose with glycerin and	MiniMed Mio MMT-925A	2
Martindale Pharma		sucrose251	MiniMed Mio MMT-941A	
Mask for spacer device	241	Methyldopa51	MiniMed Mio MMT-943A	
Maviret	102	Methyldopa Mylan51	MiniMed Mio MMT-945A	
Maxidex		Methyldopa Mylan S2951	MiniMed Mio MMT-965A	2
Maxitrol		Methylnaltrexone bromide25	MiniMed Mio MMT-975A	
MCT oil (Nutricia)	255	Methylphenidate ER - Teva141	MiniMed Quick-Set MMT-386A	
Measles, mumps and rubella		Methylphenidate hydrochloride 141	MiniMed Quick-Set MMT-387A	
vaccine	279	Methylphenidate hydrochloride	MiniMed Quick-Set MMT-396A	2
Mebendazole	89	extended-release142	MiniMed Quick-Set MMT-397A	
Mebeverine hydrochloride		Methylprednisolone79	MiniMed Quick-Set MMT-398A	2
Medrol	79	Methylprednisolone (as sodium	MiniMed Quick-Set MMT-399A	
Medroxyprogesterone acetate		succinate) 79	MiniMed Silhouette MMT-368A	
Genito-Urinary	73	Methylprednisolone aceponate63	MiniMed Silhouette MMT-377A	
Hormone		Methylprednisolone acetate79	MiniMed Silhouette MMT-378A	
Mefenamic acid	109	Methylxanthines240	MiniMed Silhouette MMT-381A	2
Megace	168	Metoclopramide Actavis 10132	MiniMed Silhouette MMT-382A	2

MiniMed Silhouette MMT-383A	22	Myambutol	99	Nicotine	14
MiniMed Silhouette MMT-384A	22	Mycobutin		Nifedipine	
MiniMed Sure-T MMT-864A	21	MycoNail		Nifuran	
MiniMed Sure-T MMT-866A	21	Mycophenolate mofetil		Nilotinib	16
MiniMed Sure-T MMT-874A	21	Mydriacyl		Nilstat	
MiniMed Sure-T MMT-876A	21	Mylan (12 hr release)		Alimentary	3
MiniMed Sure-T MMT-884A	21	Mylan (24 hr release)		Genito-Urinary	
MiniMed Sure-T MMT-886A	21	Mylan Atenolol		Infection	
Minims Pilocarpine	246	Mylan Clomiphen		Nintedanib	
Minims Prednisolone	245	Mylan Indapamide	53	Nipent	15
Minirin	87	Mylan Italy (24 hr release)		Nirmatrelvir with ritonavir	
Minirin Melt	87	Myleran		Nitrates	5
Mino-tabs	93	Mylotarg		Nitroderm TTS	5
Minocycline hydrochloride	93	Myometrial and Vaginal Hormo	ne	Nitrofurantoin	10
Minomycin		Preparations		Nitrolingual Pump Spray	
Minor Skin Infections		Myozyme	26	Nivestim	
Minoxidil	56	- N -		Nivolumab	22
Mirena	82	Nadolol	50	Nizoral	9
Mirtazapine	127	Nadolol BNM	50	Nodia	
Misoprostol		Naglazyme	28	Noflam 250	
Mitomycin C	156	Nalcrom		Noflam 500	
Mitozantrone		Naloxone hydrochloride		Non-Steroidal Anti-Inflammatory	
Mitozantrone Ebewe		Naltraccord		Drugs	10
Mixtard 30		Naltrexone hydrochloride		Nonacog gamma, [Recombinant	
MMR II		Naphazoline hydrochloride		Factor IX]	39
Moclobemide		Naphcon Forte		Norethisterone	
Modafinil		Naprosyn SR 1000		Genito-Urinary	74
Modavigil	143	Naprosyn SR 750		Hormone	8
Moduretic	52	Naproxen		Norflex	
Molaxole		Narcaricin mite		Norfloxacin	
Molnupiravir		Nasal Preparations		Noriday 28	
Moments		Natalizumab		Norimin	
Mometasone furoate	63	Natulan	157	Normacol Plus	
Monogen		Nausafix		Normison	
Montelukast		Nausicalm		Norpress	
Montelukast Mylan		Navelbine		Nortriptyline hydrochloride	
Montelukast Viatris		Nefopam hydrochloride		Norvir	
Moroctocog alfa [Recombinant fa		Neisvac-C		Noumed12	
VIII]		Neo-Mercazole		Noumed Paracetamol	
Morphine hydrochloride		Neocate Gold		NovaSource Renal	
Morphine sulphate		Neocate Junior Unflavoured		Novatretin	
Motetis		Neocate Junior Vanilla		NovoMix 30 FlexPen	
Mouth and Throat		Neocate SYNEO		NovoRapid	1
Movapo		Neoral		NovoRapid FlexPen	
Moxifloxacin		Neostigmine metilsulfate		NovoRapid Penfill	
MSUD Maxamum		Nepro HP (strawberry)		NovoSeven RT	
Mucilaginous laxatives with		Nepro HP (vanilla)	259	Noxafil	
stimulants	25	Nepro HP RTH	258	Nozinan	
Mucolytics		Neulactil		Nozinan (Swiss)	
Mucosoothe		Neulastim		Nozinan S29	13:
Multiple Sclerosis Treatments		Neuraxpharm		Nucala	
Multivitamin renal		NeuroTabs	34	Nuelin	
Multivitamins		Nevirapine		Nuelin-SR	
Mupirocin		Nevirapine Alphapharm		Nupentin	
Muscle Relaxants		Nevirapine Viatris		Nusinersen	
Myite		Nicorandil		Nutilis	

Nutren Diabetes	256	Omeprazole actavis 40	9	Paclitaxel Ebewe	15
Nutrient Modules	253	Omnitrope	83	Paediatric Seravit	3
Nutrini Energy Multi Fibre	258	Omnitrope S29		Palbociclib	
Nutrini Energy RTH	258	Onbrez Breezhaler	2 <mark>36</mark>	Paliperidone	13
Nutrini Low Energy Multi Fibre2		Oncaspar LYO		Paliperidone palmitate	
Nutrini Peptisorb2	271	OncoTICE	177	Palivizumab	
Nutrini Peptisorb Energy2	271	Ondansetron	132	Pamidronate disodium	
Nutrini RTH	258	Ondansetron ODT-DRLA	132	Pamisol	11°
Nutrison 800 Complete Multi		One-Alpha	33	Pamol	
Fibre2	263	One-Alpha S29	<mark>33</mark>	Pancreatic enzyme	2
Nutrison Advanced Diason2	256	Onrex	132	Pantoprazole	
Nutrison Advanced Peptisorb2	259	Opdivo		Panzop Relief	
Nutrison Concentrated	265	Ora-Blend	251	Panzytrat	2
Nutrison Energy2	263	Ora-Blend SF		Papaverine hydrochloride	5
Nutrison Energy Multi Fibre2		Ora-Plus	251	Para-amino salicylic acid	99
Nutrison Multi Fibre	263	Ora-Sweet	251	Paracare	
Nutrison Standard RTH	263	Ora-Sweet SF	251	Paracare Double Strength	
Nyefax Retard	.50	Orabase	<mark>32</mark>	Paracetamol	
Nystatin		Oral and Enteral Feeds	256	Paracetamol (Ethics)	123
Alimentary	32	Oratane	60	Paracetamol + Codeine	
Genito-Urinary	74	Ordine		(Relieve)	12
Infection	97	Orgran	267	Paracetamol with codeine	
NZB Low Gluten Bread Mix2	266	Ornidazole		Paraffin	64
- 0 -		Orphenadrine citrate		Paraffin liquid with wool fat	24
Obinutuzumab	203	Ortho-tolidine	<mark>76</mark>	Parasiticidal Preparations	6
Obstetric Preparations	76	Oruvail SR	109	Parnate	126
Ocrelizumab1	137	Osmolite RTH	263	Parnate S29	126
Ocrevus	137	Other Endocrine Agents	87	Paromomycin	9
Octocog alfa [Recombinant factor		Other Oestrogen Preparations	s82	Paroxetine	
VIII] (Advate)	39	Other Progestogen		Paser	99
Octocog alfa [Recombinant factor		Preparations	82	Paxam	136
VIII] (Kogenate FS)	40	Other Skin Preparations	69	Paxlovid	103
Octreotide1	168	Ovestin		Pazopanib	164
Octreotide Depot Teva1	168	Genito-Urinary	74	Peak flow meter	24
Octreotide GH	168	Hormone	82	Pedialyte - Bubblegum	4
Octreotide long-acting1	168	Oxaliplatin	149	Pediasure	
Oestradiol	.81	Oxaliplatin Accord	149	Pediasure Plus	
Oestradiol valerate	.81	Oxaliplatin Actavis 100	149	Pediasure RTH	
Oestradiol with norethisterone	82	Oxaliplatin Ebewe	149	Pegaspargase	157
Oestriol		Oxis Turbuhaler	2 <mark>36</mark>	Pegasys	10
Genito-Urinary	.74	Oxpentifylline	57	Pegfilgrastim	4
Hormone	82	Oxybutynin	75	Pegylated interferon alfa-2a	10
Oestrogens	.81	Oxycodone hydrochloride	125	Pembrolizumab	
Ofev	238	Oxycodone Sandoz	125	Pemetrexed	152
Oil in water emulsion		OxyNorm	125	Penicillamine	110
Olanzapine133-1	134	Oxytocin	75	Penicillin G	92
Olaparib1	156	Oxytocin BNM	75	PenMix 30	1
Olbetam	54	Oxytocin GH	75	PenMix 50	1
Olbetam S29	54	Oxytocin with ergometrine		Pentasa	
Olopatadine	247	maleate	75	Pentostatin [Deoxycoformycin]	
Olopatadine Teva		Ozurdex	244	Pentoxifylline [Oxpentifylline]	
Olsalazine	8	-P-		Peptamen Junior	258
Omalizumab	204	Pacifen	119	Pepti-Junior	
Omeprazole	9	Pacimol		Peptisorb	259
Omeprazole actavis 10		Paclitaxel	157	Perhexiline maleate	5
Omeprazole actavis 20		Paclitaxel Actavis		Pericyazine	

Perindopril	46	Podophyllotoxin	69	Prolia	110
Perjeta	206	Polaramine	235	Promethazine hydrochloride	235
Permethrin	66	Poliomyelitis vaccine	282	Propafenone hydrochloride	48
Perrigo	69	Poloxamer	25	Propamidine isethionate	
Pertuzumab		Poly-Gel	247	Propranolol	50
Peteha	99	Poly-Tears		Propylene glycol	
Pethidine hydrochloride	126	Poly-Visc		Propylthiouracil	
Pevaryl	61	Polycal		Prostacur	
Pexsig		Polyethylene glycol 400 and		Protaphane	10
Pfizer Exemestane		propylene glycol	247	Protaphane Penfill	
Pharmacy Health Sorbolene w	<i>i</i> ith	Ponstan		Protifar	
Glycerin		Posaconazole	97	Protionamide	
Pheburane		Posaconazole Juno	97	Provera	81
Phenasen		Postinor-1		Provera HD	
Phenobarbitone	129	Potassium chloride	44–45	PSM Citalopram	
Phenobarbitone sodium		Potassium citrate		Psoriasis and Eczema	
Extemporaneous	251	Potassium iodate		Preparations	66
Nervous		Povidone iodine		PTU	
Phenoxybenzamine		Pradaxa		Pulmicort Turbuhaler	
hydrochloride	46	Pramipexole hydrochloride		Pulmozyme	
Phenoxymethylpenicillin (Peni		Pravastatin		Puri-nethol	
V)		Pravastatin Mylan		Puria	
Phenytoin sodium		Praziquantel		Pyrazinamide	
Phillips Milk of Magnesia		Prazosin		Pyridostigmine bromide	
Phlexy 10		Pred Forte		Pyridoxine hydrochloride	
Phosphate Phebra		Prednisolone		Pyridoxine multichem	
Phosphorus		Prednisolone acetate		Pyrimethamine	
Phytomenadione		Prednisolone sodium		Pytazen SR	
Pilocarpine hydrochloride		Prednisolone sodium		- Q -	
Pimafucort		phosphate	245	Quetapel	134
Pimecrolimus		Prednisolone-AFT		Quetiapine	
Pine tar with trolamine laurilsul		Prednisone		Quick-Set MMT-392	
and fluorescein		Prednisone Clinect		Quick-Set MMT-393	
Pinetarsol		Pregabalin		Quinapril	
Pioglitazone		Pregabalin Pfizer		Quinapril with	
Pirfenidone		Pregnancy Tests - hCG Urine		hydrochlorothiazide	47
Pizotifen		Premarin	81	Qvar	
PKU Anamix Infant		Prevenar 13		- R -	200
PKU Anamix Junior		Priadel		RA-Morph	12/
PKU Anamix Junior Chocolate		Primaguine		Ralicrom	
PKU Anamix Junior LQ		Primidone		Raloxifene hydrochloride	
PKU Anamix Junior Orange		Primidone Clinect		Raltegravir potassium	
PKU Anamix Junior Vanilla		Primolut N		Ramipex	
PKU Lophlex LQ 10		Priorix		Ramipril	
PKU Lophlex LQ 20		Probenecid		Ranbaxy-Cefaclor	
PKU Lophlex Powder		Probenecid-AFT	118	Ranbaxy-Cefaclor S29	
PKU Lophlex Sensation 20		Procaine penicillin		Rapamune	221
'		Procarbazine hydrochloride	157	Rasagiline	
Plaquenil				Reandron 1000	
Plendil ER		Prochlorperazine Proctofoam		Recombinant factor IX	
Pneumococcal (PCV10) conju		Proctosedyl		Recombinant factor VIIa	
vaccine				Recombinant factor VIII	
Pneumococcal (PCV13) conju- vaccine	yait 201	Procyclidine hydrochloride Progesterone			
Pneumococcal (PPV23)	201			Rectogesic	
	000	Proglicem	10	RedipredRelieve	
polysaccharide vaccine Pneumovax 23		Proglycem		Relistor	
FIIEUIIIUVAX ZJ	202	F1099110Va	Ö I	TEIISIUI	25

Remicade	195	Rubifen141	Simvastatin Mylan5
Renilon 7.5		Rubifen SR141	Sinemet12
Resonium-A	45	Rugby Capsaicin Topical Cream	Sinemet CR12
Resource Beneprotein	255	Musculoskeletal110	Sirolimus23
Respigen		Nervous122	Siterone8
Respiratory Devices		Rulide D91	Slow-Lopresor4
Respiratory Stimulants		Rurioctocog alfa pegol [Recombinant	Smith BioMed Rapid Pregnancy
Retinol palmitate		factor VIII]	Test7
ReTrieve		Ruxolitinib	Sodibic4
Retrovir		Rythmodan	Sodium acid phosphate2
Revlimid			Sodium alginate
		Rytmonorm48	
Revolade		Sabril	Sodium benzoate
			Sodium bicarbonate
Riboflavin		Sacubitril with valsartan47	Blood44–4
Ribomustin		Sagent	Extemporaneous25
Ricit		SalAir237	Sodium calcium edetate24
Rifabutin		Salazopyrin8	Sodium chloride
Rifadin		Salazopyrin EN8	Blood4
Rifampicin		Salbutamol237	Respiratory24
Rifaximin	10	Salbutamol with ipratropium	Sodium citrate with sodium lauryl
Rifinah		bromide237	sulphoacetate2
Rilutek	121	Salicylic acid68	Sodium citro-tartrate7
Riluzole	121	Salmeterol236	Sodium cromoglicate
RINVOQ	232	Sandomigran 131	Alimentary
Riodine	65	Sanofi Primaquine98	Sensory24
Risedronate Sandoz	112	Sapropterin dihydrochloride30	Sodium fluoride3
Risedronate sodium	112	Scalp Preparations68	Sodium Fusidate [fusidic acid]
Risperdal Consta	135	Scopoderm TTS132	Dermatological6
Risperidone		Sebizole68	Infection9
Risperidone (Teva)		Secukinumab219	Sensory24
Risperon		Sedatives and Hypnotics137	Sodium hyaluronate [Hyaluronic
Ritalin		Seebri Breezhaler238	acid]24
Ritalin LA		Selegiline hydrochloride120	Sodium phenylbutyrate3
Ritonavir		Senna	Sodium picosulfate2
Rituximab (Mabthera)		Senokot	Sodium polystyrene sulphonate4
Rituximab (Riximyo)		SensoCard	Sodium tetradecyl sulphate4
Rivaroxaban		Serc	Sodium valproate12
Rivastigmine		Serenace	Sofradex24
		Seretide	
Rivastigmine Patch BNM 10			Soframycin24
Rivastigmine Patch BNM 5		Seretide Accuhaler237	Solgar27–29, 3
Rivotril		Serevent 236	Solifenacin Mylan
Riximyo		Serevent Accuhaler236	Solifenacin succinate7
RIXUBIS		Sertraline	Solu-Cortef7
Rizamelt		Setrona127	Solu-Medrol7
Rizatriptan	131	Setrona AU127	Solu-Medrol-Act-O-Vial7
RoActemra S29	221	Sevredol125	Somatropin (Omnitrope)8
Ronapreve		Sex Hormones Non	Sotalol5
Ropin	120	Contraceptive80	Spacer device24
Ropinirole hydrochloride		Shield XL71	Span-K4
Rosuvastatin	54	Shingles vaccine283	Spinal Muscular Atrophy13
Rosuvastatin Viatris		Shingrix283	Spinraza13
Rotarix	282	SII-Onco-BCG177	Spiolto Respimat23
Rotavirus oral vaccine	282	Sildenafil58	Spiractin5
Roxane		Silhouette MMT-37323	Spiriva23
Roxane-Propranolol		Siltuximab221	Spiriva Respimat23
Roxithromycin		Simvastatin	Spironolactone5
			- F 3

Sporanox	96	Tamsulosin-Rex	75	Tixagevimab with cilgavimab	221
Sprycel		Tandem Cartridge		TMP	95
Stemetil		Tandem t:slim X2 with Basal-IQ		Tobramycin	
Steril-Gene	80	Tap water	252	Infection	
SteroClear	241	Tarceva	161	Sensory	243
Stesolid	128	Taro	97	Tobramycin BNM	
Stimulants/ADHD Treatments	139	Taro-Testosterone	80	Tobramycin Mylan	95
Stiripentol	130	Tasigna	163	Tobrex	243
Stocrin	104	Tasmar	120	Tocilizumab	
Stomahesive		Taurine	31	Tofranil	126
Strattera	140	Tecfidera	137	Tolcapone	
Strides Shasun	97	Tegretol	128	Tolvaptan	53
Stromectol	65	Tegretol CR	128	Topamax	130
Sucralfate	10	Telfast	235	Topical Products for Joint and	
Sulfadiazine Silver	61	Teligent	94	Muscular Pain	110
Sulfadiazine sodium	95	Temaccord	157	Topiramate	130
Sulfasalazine		Temazepam	138	Topiramate Actavis	130
Sulphur	68	Temozolomide	157	Total parenteral nutrition (TPN)	44
Sulprix		Tenofovir disoproxil	100	TPN	
Sumagran		Tenofovir Disoproxil Emtricitabine	9	Tramadol hydrochloride	126
Sumatriptan	131	Mylan	102	Tramal SR 100	126
Sunitinib	165	Tenofovir Disoproxil Mylan	100	Tramal SR 150	126
Sunitinib Pfizer	165	Tenoxicam	109	Tramal SR 200	126
Sunscreens	69	Tensipine MR10	50	Trandate	49
Sunscreens, proprietary	69	Tepadina	149	Tranexamic acid	40
Sure-T MMT-863	21	Terbinafine		Tranylcypromine sulphate	126
Sure-T MMT-873	21	Terbutaline sulphate	237	Trastuzumab	
Sustagen Hospital Formula	263	Teriflunomide	137	Trastuzumab emtansine	226
Sustagen Hospital Formula		Teriparatide	112	Travatan	246
Active	263	Testosterone		Travoprost	
Sustanon Ampoules		Testosterone cipionate		Treatments for Dementia	143
Sylvant		Testosterone esters		Treatments for Substance	
Symbicort Turbuhaler 100/6		Testosterone undecanoate	80	Dependence	144
Symbicort Turbuhaler 200/6		Tetrabenazine		Trental 400	
Symbicort Turbuhaler 400/12		Tetrabromophenol		Tretinoin	
Symmetrel		Tetracosactrin		Dermatological	60
Sympathomimetics		Tetracycline		Oncology	
Synacthen		Teva Lisinopril		Trexate	
Synacthen Depot		Thalidomide		Triamcinolone acetonide	
Synacthene Retard		Thalomid		Alimentary	32
Synagis		Theophylline		Dermatological	63
Synflorix		Thiamine hydrochloride		Hormone	
Synthroid		Thiamine multichem		Triamcinolone acetonide with	
Syntometrine		THIO-TEPA		gramicidin, neomycin and nysta	atin
Syrup (pharmaceutical grade)		Thioguanine		Dermatological	
Systane Unit Dose		Thiotepa		Sensory	
-T-		Thyroid and Antithyroid Agents		Triazolam	
Tacrolimus		Ticagrelor		Trimethoprim	
Dermatological	68	Ticagrelor Sandoz		Trimethoprim with	
Oncology		Tilcotil		sulphamethoxazole	
Tacrolimus Sandoz		Timolol		[Co-trimoxazole]	96
Taliglucerase alfa		Timoptol XE		Trisequens	82
Tambocor		Tiotropium bromide		Trisul	
Tamoxifen citrate		Tiotropium bromide with		Trophic Hormones	
Tamoxifen Sandoz		olodaterol	238	Tropicamide	
Tamsulosin hydrochloride		Tivicay		Trulicity	10
ramoulooni nyuloonionu c	/ 🗸	1 1 1 1 0 ay		11 Gillotty	14

Trusopt	245	Ventolin	237	Xylocaine 2% Jelly	12
TruSteel	21	Vepesid	154	Xyntha	39
Tryzan		Verapamil hydrochloride	51	- Z -	
Tuberculin PPD [Mantoux] test	283	Vermox		Zapril	46
Tubersol		Versacloz	133	Zarontin	
Two Cal HN	265	Vesanoid	159	Zaroxolyn	52
Tykerb	162	Vexazone	12	Zavedos	
Tysabri		Vfend	98	Zeffix	100
. U -		Viaderm KC	63	Zematop	68
UK Synacthen	80	Viatris		Zetlam	
Ultibro Breezhaler		Cardiovascular	49	Ziagen	10
Ultraproct	8	Infection	95	Zidovudine [AZT]	10
Umeclidinium		Vigabatrin	130	Zidovudine [AZT] with	
Umeclidinium with vilanterol	238	Vigisom	137	lamivudine	10
Univent2	237, 241	Vildagliptin		Ziextenzo	4
Upadacitinib	232	Vildagliptin with metformin		Zimybe	5
Ural	76	hydrochloride	12	Zinc and castor oil	
Urea	64	Vimpat		Zinc sulphate	
Urex Forte	52	Vinblastine sulphate		Zincaps	3
Urinary Agents		Vincristine sulphate		Zinnat	90
Urinary Tract Infections		Vinorelbine		Ziprasidone	
Urinorm		Vinorelbine Ebewe		Zista	
Uromitexan		Viramune Suspension		Zithromax	
Ursodeoxycholic acid	24	ViruPOS		Zoledronic acid	
Ursosan		Vit.D3	33	Hormone	78
Utrogestan		VitA-POS		Musculoskeletal	113
- V -		Vitabdeck		Zoledronic acid Mylan	78
Vaccinations	274	Vital		Zoledronic acid Viatris	
Vaclovir		Vitamin B complex		Hormone	78
Valaciclovir		Vitamin B6 25		Musculoskeletal	
Valganciclovir		Vitamins		Zopiclone	
Valganciclovir Mylan		Vivonex TEN		Zopiclone Actavis	138
Vancomycin		Voltaren		Zostavax	
Vannair		Voltaren D		Zostrix	
Varenicline Pfizer		Voltaren Ophtha		Zostrix HP	
Varenicline tartrate		Voltaren SR		Zuclopenthixol decanoate	
Varicella vaccine [Chickenpox		Volumatic		Zuclopenthixol hydrochloride	
vaccine]	283	Voriconazole		Zusdone	
Varicella zoster vaccine [Shingle		Votrient		Zyban	
vaccine]		Vttack		Zypine	
Varicella zoster virus (Oka strain		- W -		Zypine ODT	130
attenuated vaccine [shingles	,	Warfarin sodium	43	Zyprexa Relprevv	
vaccine]	283	Wart Preparations		Zytiga	
Varivax		Wasp venom allergy treatmen		_,.9	
Vasodilators		Water			
Vasopressin Agonists		Blood	44		
Vasorex		Extemporaneous			
Vebulis		White Soft Liquid Paraffin AF			
Vedafil		Wool fat with mineral oil			
Veletri		- X -			
Venclexta		Xarelto	43		
Venetoclax		Xifaxan			
Venlafaxine		XMET Maxamum			
Venomil		Xolair			
VENOX		XP Maxamum			
Ventavis		Xylocaine			
· • · · · · · · · · · · · · · · · · · ·					