February 2023 Volume 30

Editors:

Kaye Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz
@Pharmaceutical Management Agency
ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

		introducing Filannac	۷
	Section A	General Rules	5
	Section B	Alimentary Tract & Metabolism	6
		Blood & Blood Forming Organs	36
		Cardiovascular System	46
		Dermatologicals	61
		Genito Urinary System	71
		Hormone Preparations – Systemic	78
		Infections – Agents For Systemic Use	90
		Musculoskeletal System	110
		Nervous System	121
		Oncology Agents & Immunosuppressants	149
		Respiratory System & Allergies	240
		Sensory Organs	249
		Various	254
	Section C	Extemporaneous Compounds (ECPs)	256
	Section D	Special Foods	259
/	Section I	National Immunisation Schedule	280

Introducing Pharmac

Index

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION BY ALIMENTARY TRACT AND METAROLISM

	SECTION B: ALIMENTARY TRACT AND ME	Subsidy (Manufacturer's Price)		Fully Subsidised	
		(Manufacturer's Frice)	Per	Jubsidised	
Α	ntacids and Antiflatulents				
Α	ntacids and Reflux Barrier Agents				
AL	GINIC ACID				
	Sodium alginate 225 mg and magnesium alginate 87.5 mg posachet		30	✓	Gaviscon Infant
SO	DIUM ALGINATE				
*	Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
*	Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m	nl	Acidex
P	hosphate Binding Agents				
	JMINIUM HYDROXIDE				
	Tab 600 mg	12.56	100	•	Alu-Tab
CA	LCIUM CARBONATE				
	Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) — Subsidy by endorsement Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed according	cium carbonate table	500 m		Roxane ium carbonate tablets are
٨	ntidiarrhoeals				
	midial mocals				
A	gents Which Reduce Motility				
*	PERAMIDE HYDROCHLORIDE – Up to 30 cap available on a Tab 2 mg	10.75	400 400	_	Nodia Diamide Relief

LOPERAMIDE HYDROCHLORIDE - Up to 30 c	ap available on a PSO		
* Tab 2 mg	10.75	400	✓ Nodia
* Cap 2 mg	7.25	400	✓ Diamide Relief

Rectal and Colonic Anti-inflammatories

BUDESONIDE

Cap 3 mg - Special Authority see SA1886 below - Retail 90 ✓ Entocort CIR pharmacy......166.50

⇒SA1886 Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	✓	

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP 21.1 g OP	✓ Cortifoam S29✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam §29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
OLSALAZINE Tab 500 mg	56.02	60	1	Atnahs Olsalazine \$29
Cap 250 mg	93.37 53.00	100 100		Dipentum Dipentum
PREDNISOLONE SODIUM Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential Prednisolone \$29
SODIUM CROMOGLICATE Cap 100 mg(Nalcrom Cap 100 mg to be delisted 1 April 2023)	92.91	100		Nalcrom Ralicrom
SULFASALAZINE * Tab 500 mg * Tab EC 500 mg		100 100		Salazopyrin Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g11.06	30 g OP	✓ Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg7.30	12	✓ Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a		_	
PSO65	.45	10	Max Health
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg6	.35 1	00	Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO6	.35	5 ✓	Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg9	.20	90 🗸	Colofac

	ALIMENTARY	IHAC	I AND	METABOLISM
	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
Antiulcerants				
Antisecretory and Cytoprotective				
MISOPROSTOL ☀ Tab 200 mcg – Up to 120 tab available on a PSO	47.73	120	√ C	ytotec
Helicobacter Pylori Eradication				
CLARITHROMYCIN Tab 500 mg – Subsidy by endorsement	ri eradication and presci		ndorsed	
H2 Antagonists				
FAMOTIDINE – Only on a prescription * Tab 20 mg	4.91	100	✓ Fa	amotidine Hovid 829
★ Tab 40 mg	8.48	100	✓ Fa	amotidine Hovid \$29
Inj 10 mg per ml, 4 ml - Subsidy by endorsement Subsidy by endorsement - Subsidised for patients red		10 t of palliat		ylan S29
Proton Pump Inhibitors	,	•		
ANSOPRAZOLE				
€ Cap 15 mg		100 100	_	anzol Relief anzol Relief
For omeprazole suspension refer Standard Formulae, page	je 256			
€ Cap 10 mg		90	√ <u>0</u>	meprazole actavis 10
≮ Cap 20 mg	1.86	90	√ <u>0</u>	meprazole actavis 20
≮ Cap 40 mg	3.11	90	√ <u>0</u>	meprazole actavis 40
Powder – Only in combination Only in extemporaneously compounded omeprazole s	42.50 suspension.	5 g	✓ M	idwest
€ Inj 40 mg ampoule with diluent	37.38	5	_	r Reddy's Omeprazole cicure S29
ANTOPRAZOLE				
★ Tab EC 20 mg ★ Tab EC 40 mg		90 90	_	anzop Relief anzop Relief
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE				
Tab 120 mg	14.51	50	✓ G	astrodenol 829

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
SUCRALFATE Tab 1 g	35.50 (48.28)	120		Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail Tab 550 mg SA1461 Special Authority for Subsidy Initial application only from a gastroenterologist, hepatological process.	625.00		mendation	
hepatologist. Approvals valid for 6 months where the patiel tolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Prahepatologist. Approvals valid without further renewal unles benefiting from treatment.	actitioner on the recomme	endatio	n of a gastr	oenterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE - Special Authority see SA1320 below - Reta Cap 25 mg Cap 100 mg Oral liq 50 mg per ml	110.00	100 100 30 ml 0	✓ OP ✓	Proglicem \$29 Proglicem \$29 Proglycem \$29 e5 Pharma \$29
➤ SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approval hypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid with appropriate and the patient is benefiting from treatment.			d for the tre	eatment of confirmed
GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	✓	Glucagen Hypokit
Insulin - Short-acting Preparations				
INSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml (Actrapid Humulin B
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓.	Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
INSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	•	NovoMix 30 FlexPen
INSULIN ISOPHANE ▲ Inj human 100 u per ml	17.68	10 ml (Humulin NPH Protaphane

▲ Inj human 100 u per ml, 3 ml......29.86

5

✓ Humulin NPH

✓ Protaphane Penfill

	Cubaidu		Fully Drand or
	Subsidy (Manufacturer's F	Price) Subs	Fully Brand or idised Generic
	\$	Per	✓ Manufacturer
ISULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
		_	✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
			✓ PenMix 30✓ PenMix 50
			▼ Penwix 50
ISULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml, 3 ml		5	✓ Humalag Miv 25
		5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml, 3 ml		5	✓ Humalog Mix 50
O III			- Hamaiog inix oo
Insulin - Long-acting Preparations			
SULIN GLARGINE			•
Inj 100 u per ml, 10 ml		1	✓ Lantus
Inj 100 u per ml, 3 ml		5 5	✓ Lantus✓ Lantus SoloStar
IIIJ 100 u per IIII, 3 IIII disposable peri	94.50	5	V Lantus SoloStar
nsulin - Rapid Acting Preparations			
SULIN ASPART			
Inj 100 u per ml, 10 ml		1	✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill ✓ NovoRapid FlorBox
Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
ISULIN GLULISINE			4
Inj 100 u per ml, 10 ml		1	✓ Apidra
Inj 100 u per ml, 3 ml		5 5	✓ Apidra✓ Apidra SoloStar
ISULIN LISPRO		3	Apidia odiootai
isolin liseno . Inj 100 u per ml, 10 ml	3/1 02	10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml		5	✓ Humalog
Alpha Glucosidase Inhibitors			J
CARBOSE			
: Tab 50 mg	8.95	90	✓ Accarb
Tab 100 mg		90	✓ Accarb
Oral Hypoglycaemic Agents			
LIBENCLAMIDE			
: Tab 5 mg	7.50	100	✓ Daonil
LICLAZIDE			
: Tab 80 mg	15.18	500	✓ Glizide
LIPIZIDE			_
: Tab 5 mg	4.58	100	✓ Minidiab
ETFORMIN HYDROCHLORIDE			
Tab immediate-release 500 mg	14.74	1,000	✓ Metformin Mylan
		.,500	✓ Metformin Viatris
Tab immediate-release 850 mg	11.28	500	Metformin Mylan
•			-

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
	\$	Per	•	Manufacturer
PIOGLITAZONE				
* Tab 15 mg	6.80	90	✓ V	/exazone
* Tab 30 mg	7.30	90	✓ V	/exazone
* Tab 45 mg		90	✓ V	/exazone
VILDAGLIPTIN				
Tab 50 mg	35.00	60	√ (alvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	√ (Salvumet
Tab 50 mg with 850 mg metformin hydrochloride		60	√ (Salvumet

GLP-1 Agonists

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Maori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*: and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

DULAGLUTIDE - Special Authority see SA2065 above - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*: or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 10 mg58.56	30	Jardiance
	Tab 25 mg58.56	00	Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

✓ CareSens Dual 1 OP

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes; or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

1 OP ✓ CareSens N ✓ CareSens N POP 20.00 ✓ CareSens N Premier

Note: Only 1 meter available per PSO

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips	26.20	50 test OP	SensoCard
---------------------------	-------	------------	-----------------------------

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

*	29 g × 12.7 mm10.9	5 100	✓ B-D Micro-Fine
	31 g × 5 mm12.2		✓ B-D Micro-Fine
	31 g × 6 mm9.5		✓ Berpu
	31 g × 8 mm10.9		✓ B-D Micro-Fine
	32 g × 4 mm10.9		✓ B-D Micro-Fine

15

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	
_		\$	Per		Manufacturer
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	- Maximum of 200	dev p	er prescrip	otion
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.36	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.36	10		
		(1.99)			B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — **(severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

1 OP

1 OP

✓ TruSteel

✓ TruSteel

	ALIMENTALLI	1117	OI AIID	METABOLISM
	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
continued than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mr 3 The patient has not had an increase in severe unexplaine				ne; and
 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within the 	ir vocational scope.			
INSULIN PUMP CARTRIDGE - Special Authority see SA1985	on page 19 – Retail ph	armac	y	
 a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded pe 	r year.			
Cartridge 300 U, t:lock × 10		1 OP	✓ Ta	andem Cartridge
 INSULIN PUMP INFUSION SET (STEEL CANNULA) – Special a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 	Authority see SA1985	on pa	ge 19 – Re	tail pharmacy
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-866A
8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-874A
8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T MMT-876A
6 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock	130.00	1 OP	√ S	ure-T MMT-863
8 mm steel needle; 29 G; manual insertion; 60 cm tubing \times 10 with 10 needles; luer lock		1 OP	_	ure-T MMT-873
INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGH' Retail pharmacy a) Maximum of 3 sets per prescription	Г INSERTION) – Spe	cial Au	thority see	SA1985 on page 19 –
b) Only on a prescriptionc) Maximum of 13 infusion sets will be funded per year.				
6 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles	130.00	1 OP	✓ T	ruSteel
8 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	130.00	1 OP	✓ Ti	ruSteel
6 mm steel cannula; straight insertion; 60 cm line × 10 with				

8 mm steel cannula; straight insertion; 60 cm line × 10 with

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing x 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing \times 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing \times 10	130.00	1 OP	✓ MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-386A

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION WITH IN	SERT	ION DEVIC	E) - Special Authority see
SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c				
line × 10 with 10 needles		1 OP		AutoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cm			_	
line × 10 with 10 needles		1 OP		AutoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION) - Speci	al Aut	thority see S	SA1985 on page 19 -
Retail pharmacy				
a) Maximum of 3 sets per prescription Only on a prescription				
b) Only on a prescriptionc) Maximum of 13 infusion sets will be funded per year.				
17 mm teflon cannula; angle insertion; 60 cm line × 10 with				
10 needles; luer lock	130.00	1 OP	1	Silhouette MMT-373
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH		INS	ERTION DE	FVICE) - Special Authority
see SA1985 on page 19 – Retail pharmacy	T INCENTION WITH	1 11 401		evice) openial realionty
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; insertion device;			_	
110 cm line × 10 with 10 needles		1 OP	•	AutoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 cr		1 00		AutoSoft 90
line x 10 with 10 needles9 mm teflon cannula; straight insertion; insertion device;	140.00	1 OP	• /	AU105011 90
110 cm line × 10 with 10 needles	140.00	1 OP	•	AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 cr		. 0.		-utocont 50
line × 10 with 10 needles		1 OP	1	AutoSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH	T INSERTION) - S	necial	Authority s	see SA1985 on page 19 -
Retail pharmacy	1 11021111011, 0	poolai	riamonty o	oo on nood on page 10
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with	1 400.00	4.00		
10 needles; luer lock		1 OP	✔ (Quick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles; luer lock		1 OP		Quick-Set MMT-392
•				Juick-Set MiM1-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 or	1 page 19 – Retail pr	narma	ıcy	
a) Maximum of 3 sets per prescriptionb) Only on a prescription				
c) Maximum of 13 packs of reservoir sets will be funded per	vear			
10 × luer lock conversion cartridges 1.8 ml for Paradigm pum		1 OP	1	ADR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 10		1 OP		MiniMed
				1.8 Reservoir
				MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP	✓ I	MiniMed
				3.0 Reservoir
				MMT-332A

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Digestives Including Enzymes

_			. ~	_	_		_	_	_		_		
Ρ	Δ	Λ	Ю	ж	н	Δ	ш	(:	ы	N	'	ΥN	ЛF

1 / WOTE/WIO ENZIME			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))	94.40	100	✓ Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)	34.93	20 g OP	✓ Creon Micro
(Panzytrat Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amy	lase, 1,250 U p	rotease)) to	be delisted 1 June 2023)
URSODEOXYCHOLIC ACID – Special Authority see SA1739 below		nacy 100	✓ <u>Ursosan</u>

⇒SA1739 Special Authority for Subsidy

Initial application — (Alaquille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Fither:
 - 1 Patient has been diagnosed with Alagille syndrome; or
 - 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults: and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- Both:
 - 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
 - 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN): and
- 2 Liver function has not improved with modifying the TPN composition.

	Subsidy	Fully		Brand or
(Man	ufacturer's Price)	Subsidi	sed	Generic
	\$	Per	✓	Manufacturer

continued...

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Laxatives

Bulk-forming Agents

* Powder for oral soin	5.00	250 g OP	Psyllium Husk
	12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS			
* Dry	6.02	500 g OP	
	(17.32)	_	Normacol Plus
	2.41	200 g OP	
	(8.72)	Ü	Normacol Plus

Faecal Softeners

* Tab 50 mg 2.3 * Tab 120 mg 3.1			✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u>
DOCUSATE SODIUM WITH SENNOSIDES Tab 50 mg with sennosides 8 mg	50 :	200	✓ <u>Laxsol</u>
POLOXAMER — Only on a prescription Not funded for use in the ear.			
* Oral drops 10%	98 30	ml OP	✓ Coloxyl

Opioid Receptor Antagonists - Peripheral

DOCUSATE SODIUM - Only on a prescription

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below - Retail p	harmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy (Manufacturer's Price \$) Sub	Fully sidised	Brand or Generic Manufacturer
Osmotic Laxatives				
GLYCEROL * Suppos 4 g - Only on a prescription	10.39	20		ax-suppositories Glycerol
LACTULOSE – Only on a prescription * Oral liq 10 g per 15 ml Laevolac to be Principal Supply on 1 April 2023	3.61	500 ml	✓ La	aevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BI Powder for oral soln 13.125 g with potassium chloride 46.6 n sodium bicarbonate 178.5 mg and sodium chloride 350.	ng,	SODIUM C		olaxole
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1		eet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,	, ,	ription		
5 ml		50		icolette icolette-S29 S29
Stimulant Laxatives				
BISACODYL – Only on a prescription * Tab 5 mg * Suppos 10 mg		200 10	_	isacodyl Viatris ax-Suppositories

* Tab, standardised......2.17

SODIUM PICOSULFATE - Special Authority see SA2053 below - Retail pharmacy

SENNA - Only on a prescription

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

continued...

100

20

30 ml OP

Senokot

Senokot

✓ Dulcolax SP Drop

(8.21)

0.43

(2.06)

Subsi	dy	Fully	Brand or
(Manufacture	er's Price) Subsi	dised	Generic
\$	Per	1	Manufacturer

continued...

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease;
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

		ARGININE – Special Authority see SA2042 below – Retail pharmacy	Α
Clinicians	90	Tab 1,000 mgCBS	
✓ Solgar	50	Cap 500 mg	
g ✓ Biomed	400 g	PowderCBS	

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE - Special Authority see SA1987 below - Retail pharmacy		
Powder for oral soln575.00	180 g OP	Cystadane

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient has a confirmed diagnosis of homocystinuria; and

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

OOLINZ I WIL Q TO - Special Authority See SA2009 belo	w – Hetali pilatiliacy		
Cap 120 mg	CBS	30	✓ Solgar
Cap 160 mg		60	✓ Go Healthy

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts: or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
 \$	Per	•	

continued...

- 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE - Special Authority see SA1695 below - Reta	ail pharmacy		
Inj 100 U per ml, 5 ml vial	1,335.16	1	✓ Aldurazyme

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARINITINE - Special Authority see SA2040 below - Re	ali pharmacy		
Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg		60	✓ Balance
Oral liq 1 g per 10 ml	CBS	118 ml	✓ Carnitor S29
Oral liq 500 mg per 10 ml		300 ml	✓ Balance
Oral liq 300 mg per 10 mi		300 1111	Dalalice

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN – Special Authority see SA2041 below – F	Retail pharmacy		
Tab 100 mg	CBS	100	 Country Life
Cap 100 mg	CBS	100	✓ Solgar

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE – Special Authority see SA1990 on the next page – Retail pharmacy
Grans 483 mg per g......2,016.00 174 g OP ✓ Pheburane

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	 Life Extension

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA2137 below - Retail pharmacy		
Inj 200 unit vial1,072.00	1	✓ Elelyso

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and

	Subsidy (Manufacturer's Price) \$	Sub: Per	Fully sidised	Brand or Generic Manufacturer
continued				
 3 Radiological (MRI) signs of bone activity performed at tw demonstrate no deterioration shown by the MRI, compar or adjusted dose; and 4 Patient has not developed another medical condition tha ERT; and 5 Patient is adherent with regular treatment and taliglucera every other week rounded to the nearest whole vial (200 	red with MRI taken imn at might reasonably be ase alfa is to be admini	nediately expected	orior to o	commencement of therapy promise a response to
Mouth and Throat				
Agents Used in Mouth Ulceration				
BENZYDAMINE HYDROCHLORIDE Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with		500 ml		

Endorsement	9.00	500 ml	
	(20.31)		Difflam
Additional subsidy by endorsement for a patient wh prescription is endorsed accordingly.	o has oral mucositis a	s a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	· ·	Orabase
	`1.52 [′]	5 g OP	
	(3.60)	Ü	Orabase
Powder	, ,	28 g OP	
	(10.95)	3 -	Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE	` ,		
* Adhesive gel 8.7% with cetalkonium chloride 0.01%		15 g OP	
* Adilesive gel 6.7 /6 with Cetalkorlium Chloride 0.01 /6	(6.00)	13 g OF	Bonjela
	(0.00)		Donjela
TRIAMCINOLONE ACETONIDE			_
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
		20	· rungiiii
MICONAZOLE			
Oral gel 20 mg per g	4.74	40 g OP	✓ <u>Decozol</u>
NYSTATIN			

Vit	am	Πî	e

Vitamin B

HY	DROXOCOBALAMIN		
*	Inj 1 mg per ml, 1 ml ampoule - Up to 6 inj available on a PSO2.46	3	✓ <u>Hydroxocobalamin</u> Panpharma

Oral liq 100,000 u per ml1.76

24 ml OP

✓ Nilstat

	Subsidy	,	Fully	
	(Manufacturer's Price \$	e) Per	Subsidised	Generic Manufacturer
YRIDOXINE HYDROCHLORIDE				
a) No more than 100 mg per dose				
b) Only on a prescription				
★ Tab 25 mg – No patient co-payment payable	2.70	90	✓	Vitamin B6 25
Tab 50 mg	23.45	500	✓	Pyridoxine
				multichem
THIAMINE HYDROCHLORIDE - Only on a prescription				
₭ Tab 50 mg		100		Thiamine multichem
	7.09		✓	Max Health
Thiamine multichem to be Principal Supply on 1 April 20	123			
Max Health Tab 50 mg to be delisted 1 April 2023)				
/ITAMIN B COMPLEX			_	
★ Tab, strong, BPC	7.15	500	•	Bplex
Vitamin C				
SCORBIC ACID				
a) No more than 100 mg per dose				
b) Only on a prescription				
k Tab 100 mg	12.50	500	1	Cvite
				<u></u>
Vitamin D				
ALFACALCIDOL				
* Cap 0.25 mcg	26.32	100	✓	One-Alpha
k Cap 1 mcg	87.98	100	✓	One-Alpha
			✓	One-Alpha S29 S29
♦ Oral drops 2 mcg per ml	60.68	20 ml C)P 🗸	One-Alpha
CALCITRIOL				
★ Cap 0.25 mcg		100	1	Calcitriol-AFT
├ Cap 0.5 mcg	13.68	100	✓	Calcitriol-AFT
COLECALCIFEROL				
 Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescript 	tion2.95	12		Vit.D3
★ Oral liq 188 mcg per ml (7,500 iu per ml)	9.00	4.8 ml (OP 🗸	Puria
Multivitamin Preparations				
·				
MULTIVITAMIN RENAL - Special Authority see SA1546 below			_	
<u></u>	6.49	30	•	Clinicians Renal Vit
⇒SA1546 Special Authority for Subsidy				
nitial application from any relevant practitioner. Approvals vali	d without further rer	newal u	nless notif	ied for applications meet
ne following criteria:				
lither:				

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

MULTI	IVITAMINS - Special Authority see SA1036 on the next page	 Retail phar 	macy	
* Po	owder	72.00	200 g OP	✓ Paediatric Seravit

	Subsidy Fully (Manufacturer's Price) Subsidised		Brand or Generic	
	\$	Per	•	Manufacturer
TO CA1026 Chaolal Authority for Cubaidy				

SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINS

- 1.000 Mvite
- * Cap (fat soluble vitamins A, D, E, K) Special Authority see 60 ✓ Vitabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

Minerals

Calcium

CA	LCIUM CARBONATE		
*	Tab 1.25 g (500 mg elemental)	250	✓ Calci-Tab 500
*	Tab eff 1.25 g (500 mg elemental) - Subsidy by endorsement260.00	100	✓ Calcium 500 mg
			Hexal S29

Subsidy by endorsement - Only when prescribed for paediatric patients (< 5 years) where calcium carbonate oral liquid is considered unsuitable.

CALCIUM GLUCONATE

*	Inj 10%, 10 ml ampoule	32.00	10	Max Health -
				Hameln S29
		64 00	20	✓ May Health \$29

Fluoride

SODIUM FLUORIDE		
* Tab 1.1 mg (0.5 mg elemental)	100	✓ PSM
(PSM Tab 1.1 mg (0.5 mg elemental) to be delisted 1 March 2023)		

lodine

DOTACCILINATODATE

ΓU	I AGGIUWI IUDA I E			
*	Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓ NeuroTabs

Iron

FERROUS FUMARATE			
* Tab 200 mg (65 mg elemental)	3.04	100	✓ Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID			

	Subsidy (Manufacturer's Price) \$	Sub:	Fully sidised	Brand or Generic Manufacturer
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental) * Oral liq 30 mg (6 mg elemental) per 1 ml		30 500 ml		<u>Ferrograd</u> <u>Ferodan</u>
IRON (AS FERRIC CARBOXYMALTOSE) — Special Authority se Inj 50 mg per ml, 10 ml vial		Retail phar 1	•	Ferinject

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE * Inj 50 mg per ml, 2 ml ampoule34.50	5	✓ Ferrosig
Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%33.60	355 ml	✓ Phillips Milk of Magnesia \$29
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule25.53	10	✓ <u>Martindale</u>
Zinc		
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)11.00	100	✓ Zincaps

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

BLOOD AND BLOOD FORMING ORGANS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA1775 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe	100.00	6	Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	✓ Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	✓ Binocrit
Inj 6,000 iu in 0.6 ml, syringe		6	✓ Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	✓ Binocrit
Inj 10,000 iu in 1 ml, syringe		6	✓ Binocrit
Inj 40,000 iu in 1 ml, syringe		1	✓ Binocrit

	Subsidy (Manufacturer's Price) \$		ully sed	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID Tab 0.8 mg	26.60	1,000		olic Acid multichem
* Tab 5 mg Oral liq 50 mcg per ml		100 5 ml OP		olic Acid Mylan iomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

reaters Group in conjunction with the National F	iaemopnilia ivianagement grou	ıр.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial		1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 be Wastage claimable	elow – Retail pharmacy		
Tab 25 mg	1,550.00	28	✓ Revolade
Tab 50 mg	3,100.00	28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy. Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	1	Manufacturer

continued...

and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

Inj 30 mg in 1 ml vial	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	7,138.00	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial		1	✓ Hemlibra
Inj 150 mg in 1 ml vial	17,846.00	1	✓ Hemlibra

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Either:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
 - 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	·	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria

Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	Xyntha
Inj 1,000 iu prefilled syringe		1	Xyntha
Inj 2,000 iu prefilled syringe		1	Xyntha
Inj 3,000 iu prefilled syringe		1	Xyntha

NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group

man are manerial macricepinna management on	, ap.		
Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial	870.00	1	✓ RIXUBIS
Inj 2,000 iu vial		1	✓ RIXUBIS
Ini 3.000 iu vial	,	1	✓ RIXUBIS

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [Xpharm]

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

managed by the naemophilia Treaters Group in conjunction	n with the national	паетнорпіна	Management G
Inj 250 iu vial	210.00	1	✓ Advate
Inj 500 iu vial	420.00	1	Advate
Inj 1,000 iu vial	840.00	1	Advate
Inj 1,500 iu vial	1,260.00	1	Advate
Inj 2,000 iu vial	1,680.00	1	Advate
Inj 3,000 iu vial	2,520.00	1	Advate

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	_	Subsidised	
	\$	Per		Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE				
For patients with haemophilia. Rare Clinical Circumstances E				
treatment is managed by the Haemophilia Treaters Group in	conjunction with the f	Natio	nal Haemo	ophilia Management Group,
subject to criteria.	007.50	_	,	V
Inj 250 iu vial		1		Kogenate FS
Inj 500 iu vialInj 1,000 iu vial		1		Kogenate FS Kogenate FS
Inj 2,000 iu vial		1		Kogenate FS
Inj 3,000 iu vial		1		Kogenate FS
• •	,		•	Rogenate i o
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] For patients with haemophilia A receiving prophylaxis treatment		d trac	tmont io n	onnaged by the Heemenhilie
Treaters Group in conjunction with the National Haemophilia		u li ea	auneni is n	nanaged by the Haemophila
Inj 250 iu vial	0 0 1	1	1	Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial		1		Adynovate
Inj 2,000 iu vial		1		Adynovate
SODIUM TETRADECYL SULPHATE	2, 100.00	•		raynorato
* Inj 3% 2 ml	28 50	5		
* III 5 /6 2 III	(73.00)	J		Fibro-vein
TRANSVALUO A OIR	(70.00)			I IDIO VOIII
TRANEXAMIC ACID	10.45	00		Manarimi Dhamaa
Tab 500 mg	10.45	60	•	Mercury Pharma
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO	8.00	5	1	Konakion MM
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO		5	1	Konakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	14.95	990	1	Ethics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	4.60	84	1	Clopidogrel
•				Multichem
	5.07		✓	Arrow - Clopid
Arrow - Clopid to be Principal Supply on 1 May 2023				•
(Clopidogrel Multichem Tab 75 mg to be delisted 1 May 2023)				
DIPYRIDAMOLE				
* Tab long-acting 150 mg	10.90	60	✓	Pytazen SR
TICAGRELOR – Special Authority see SA1955 on the next page				
* Tab 90 mg		56	1	Ticagrelor Sandoz
	90.00			Brilinta
Ticagrelor Sandoz to be Principal Supply on 1 March 202	23			
(Brilinta Tab 90 mg to be delisted 1 March 2023)				

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Fither:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Fither:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment: and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA2152 be	elow - Retail pharmacy		
Inj 20 mg in 0.2 ml syringe	31.28	10	Clexane
Inj 40 mg in 0.4 ml syringe	42.49	10	Clexane
Inj 60 mg in 0.6 ml syringe	60.67	10	Clexane
Inj 80 mg in 0.8 ml syringe	80.89	10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe	125.87	10	 Clexane Forte
Inj 150 mg in 1 ml syringe	143.86	10	Clexane Forte

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	•	Manufacturer
HEPARIN SODIUM				
Inj 1,000 iu per ml, 5 ml ampoule	86.11	50	1	Pfizer
Inj 5,000 iu per ml, 5 ml vial		10	1	Heparin Sodium
, . , ,				Panpharma
Inj 5,000 iu per ml, 1 ml	32.66	5	✓	DBL Heparin
· , -,		-		Sodium S29
	70.33		1	Hospira
Inj 5,000 iu per ml, 5 ml ampoule		50		Pfizer
Inj 25,000 iu per ml, 0.2 ml		5		Hospira
111 20,000 tu pet 1111, 0.2 1111	42.40	J		Heparin DBL S29
				•
Di	482.20	50	•	Heparin DBL \$29
Pfizer Inj 5,000 iu per ml, 5 ml ampoule to be delisted 1 July	/ 2023)			
HEPARINISED SALINE				
Inj 10 iu per ml, 5 ml	65.48	50	✓	Pfizer
Oral Anticoagulants				
A DIO ATDAN				
DABIGATRAN	70.00	00		Duadava
Cap 75 mg – No more than 2 cap per day		60		Pradaxa
Cap 110 mg		60	_	Pradaxa
Cap 150 mg	/6.36	60	•	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day		30		Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28		Xarelto
Tab 20 mg	77.56	28	✓	Xarelto
VARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
★ Tab 1 mg		50	✓	Coumadin
ŭ	6.46	100	✓	Marevan
★ Tab 2 mg	4.31	50	✓	Coumadin
₭ Tab 3 mg	10.03	100	✓	Marevan
₭ Tab 5 mg		50	✓	Coumadin
-	11.48	100	✓	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM – Special Authority see SA1259 below – Reta			_	
Inj 300 mcg per 0.5 ml prefilled syringe		10		Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	148.58	10	/	<u>Nivestim</u>
⇒SA1259 Special Authority for Subsidy				

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or

	Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic	
	\$	Per	✓	Manufacturer	
continued					
4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10 ⁹ /	L); or				

5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

(Neulastim Inj 6 mg per 0.6 ml syringe to be delisted 1 June 2023)

* Inj 50%, 10 ml ampoule - Up to 5 inj available on a PSO......30.65

Infusion......CBS

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

GLUCOSE [DEXTROSE]

Intravenous Administration

POTASSIUM CHLORIDE			
* Inj 75 mg per ml, 10 ml	65.00	50	✓ Juno
SODIUM BICARBONATE			
Inj 8.4%, 50 ml	21.40	1	✓ Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			
Inj 8.4%, 100 ml	21.95	1	✓ Biomed
 a) Up to 5 inj available on a PSO 			
b) Not in combination			
SODIUM CHLORIDE			
Not funded for use as a nasal drop. Not funded for nebuliser	use except whe	n used in conju	unction with an antibiotic intended
for nebuliser use.			
Inj 0.9%, bag – Up to 2000 ml available on a PSO	1.33	500 ml	✓ Baxter
	1.36	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, ma	ternity or post-na	ital care in the	home of the patient, or on a PSO
for emergency use. (500 ml and 1,000 ml packs)			
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standar			4
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO		50	Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	Fresenius Kabi

TOTAL PARENTERAL NUTRITION (TPN)

✓ TPN

1 OP

Biomed

Biomed

✓ Pfizer

Subsidy (Manufacturer's Price)	Subs	Fully idised	Brand or Generic	
\$	Per	/	Manufacturer	

WATER

 On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or

7 19

2) On a bulk supply order; or

Ini 10 ml amnoule - Un to 5 ini available on a PSO

- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 20 ml ampoule – Up to 5 inj available on a PSO	20	✓ Fresenius Kabi
Oral Administration		
CALCIUM POLYSTYRENE SULPHONATE Powder169.85 COMPOUND ELECTROLYTES	300 g OP	✓ Calcium Resonium
Powder for oral soln — Up to 5 sach available on a PSO9.53	50	✓ Electral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)	1,000 ml OP	✓ Pedialyte - Bubblegum
PHOSPHORUS		-
Tab eff 500 mg (16 mmol)82.50	100	Phosphate Phebra
POTASSIUM CHLORIDE		
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)5.26 (17.10)	60	Chlorvescent
* Tab long-acting 600 mg (8 mmol)	200	✓ Span-K
SODIUM BICARBONATE	100	✓ Sodibic
Cap 840 mg8.52	100	✓ Sodibic ✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE		
Powder84.65	454 g OP	✓ Resonium-A

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	Generic
		\$	Per	1	Manufacturer
Alpha-Adren	oceptor Blockers				
Alpha Adrend	oceptor Blockers				
DOXAZOSIN					
		17.35	500	1	Doxazosin Clinect
			500		Doxazosin Clinect
J		20.04	300	•	DOXUZUSIII OIIIICCI
PHENOXYBENZA	MINE HYDROCHLORIDE				
* Cap 10 mg		65.00	30	✓	BNM S29
		216.67	100	✓	Dibenzyline S29
PRAZOSIN					•
		E E0	100	./	Arrotex-Prazosin
★ Tab Ting		3.33	100	•	
				_	S29 S29
* Tab 2 mg		7.00	100	✓	Arrotex-Prazosin
					S29 S29
* Tab 5 mg		11.70	100	/	Arrotex-Prazosin
					S29 S29
					329 023
Agonto Affoo	ting the Renin-Angiotensin Systen	•			
Agents Anec	ung me nemin-Angiotensin System				
A OF 1-1-1-1-1-1					
ACE Inhibitor	rs .				
CAPTOPRIL					
		04.00	0	n /	0
	per ml	94.99 95	5 ml O	Ρ •	Capoten
Oral liquid	restricted to children under 12 years of age.				
CILAZAPRIL - Su	bsidy by endorsement				
	dorsement - Subsidised for patients who were	taking cilazapril prior	r to 1 I	May 2021	and the prescription is
	ordingly. Pharmacists may annotate the presc				
dispensing of d					
, ,		2 69	90	1	Zapril
0			90		Zapril
•			90		Zapril
ŭ		10.00	50	•	Zapin
ENALAPRIL MALE				_	
			100		Acetec
* Tab 10 mg		2.02	100	✓	Acetec
* Tab 20 mg		2.42	100	✓	Acetec
LISINOPRIL					
		11 07	90	1	Ethics Lisinopril
* Tub o mg			00		Teva Lisinopril
* Tab 10 mg		11.67	90		Ethics Lisinopril
• 1 ab 10 1119		11.01	90		
₩ Tab 00 m=		14.60	00		Teva Lisinopril
* Tab 20 mg		14.69	90		Ethics Lisinopril
				•	Teva Lisinopril
PERINDOPRIL					
				_	

30

30

30

✓ Coversyl

✓ Coversyl

✓ Coversyl

	•	טואי	IOVAGOULATI STSTEW
	Subsidy (Manufacturer's Price) \$	S Per	Fully Brand or Subsidised Generic ✓ Manufacturer
QUINAPRIL			
Tab 5 mg	5.97	90	✓ Arrow-Quinapril 5
Tab 10 mg	5.18	90	✓ Arrow-Quinapril 10
Tab 20 mg	7.95	90	Arrow-Quinapril 20
RAMIPRIL			
* Cap 1.25 mg	6.90	90	✓ Tryzan
Tryzan to be Principal Supply on 1 May 2023			•
* Cap 2.5 mg	6.60	90	✓ Tryzan
Tryzan to be Principal Supply on 1 May 2023			
* Cap 5 mg	6.75	90	✓ Tryzan
Tryzan to be Principal Supply on 1 May 2023			
* Cap 10 mg	7.05	90	✓ Tryzan
Tryzan to be Principal Supply on 1 May 2023			
ACE Inhibitors with Diuretics			
exists a record of prior dispensing of quinapril with hydrochle Tab 10 mg with hydrochlorothiazide 12.5 mg Tab 20 mg with hydrochlorothiazide 12.5 mg Angiotensin II Antagonists	4.10	30 30	✓ Accuretic 10 ✓ Accuretic 20
CANDESARTAN CILEXETIL			
* Tab 4 mg	2.00	90	✓ Candestar
* Tab 8 mg		90	✓ Candestar
* Tab 16 mg		90	✓ Candestar
* Tab 32 mg		90	✓ Candestar
LOSARTAN POTASSIUM			
* Tab 12.5 mg	1.56	84	✓ Losartan Actavis
* Tab 25 mg		84	✓ Losartan Actavis
* Tab 50 mg		84	✓ Losartan Actavis
* Tab 100 mg		84	✓ Losartan Actavis
Angiotensin II Antagonists with Diuretics			
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE			
* Tab 50 mg with hydrochlorothiazide 12.5 mg	4.00	30	✓ Arrow-Losartan &
Table 55g marry droomore and rate rate ring		00	Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see	SA1905 on the next page -	- Retail p	harmacy
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure: and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 121

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocalne hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, p	age 121	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg3.49	30	✓ Aratac
▲ Tab 200 mg4.49	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a PSO9.12	6	✓ Cordarone-X
15.22	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a		
PSO	10	✓ Martindale
DIGOXIN		<u></u>
* Tab 62.5 mcg – Up to 30 tab available on a PSO7.80	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO	240	✓ Lanoxin
* Oral lig 50 mcg per ml	60 ml	✓ Lanoxin
Total liq 30 micg per mi	00 1111	✓ Lanoxin Paediatric
		Elixir S29
		✓ Lanoxin S29 S29
		LalloxIII S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan
FLECAINIDE ACETATE		
▲ Tab 50 mg19.95	60	Flecainide BNM
▲ Cap long-acting 100 mg108.65	90	✓ Flecainide
		Controlled
		Release Teva
▲ Cap long-acting 200 mg167.92	90	✓ Flecainide
		Controlled
		Release Teva
Inj 10 mg per ml, 15 ml ampoule100.00	5	✓ Tambocor
MEXILETINE HYDROCHLORIDE		
▲ Cap 150 mg162.00	100	✓ Teva S29
▲ Cap 250 mg202.00	100	✓ Teva S29
= Oup 200 mg202.00	100	- 1640

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PROPAFENONE HYDROCHLORIDE Tab 150 mg	40.90	50	✓ F	Rytmonorm
Antihypotensives				
MIDODRINE - Special Authority see SA1474 below - Retail p	harmacy			
Tab 2.5 mg	•	100	√ 0	Gutron
Tab 5 mg	79.00	100	√ (Gutron
⇒SA1474 Special Authority for Subsidy				

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATENOLOL			
* Tab 50 mg	9.33	500	✓ Mylan Atenolol
			✓ Viatris
* Tab 100 mg		500	✓ Mylan Atenolol
* Oral liq 25 mg per 5 ml	21.25	300 ml OP	✓ Atenolol AFT
			S29 S29
	38.20		✓ Essential
			Generics S29
	49.85		✓ Atenolol AFT
Restricted to children under 12 years of age.			
BISOPROLOL FUMARATE			
* Tab 2.5 mg	1.84	90	✓ Bisoprolol Mylan
			✓ Bisoprolol Viatris
* Tab 5 mg	2.55	90	✓ Bisoprolol Mylan
•			✓ Bisoprolol Viatris
* Tab 10 mg	3.62	90	✓ Bisoprolol Mylan
-			✓ Bisoprolol Viatris
CARVEDILOL			
* Tab 6.25 mg	2.24	60	✓ Carvedilol Sandoz
* Tab 12.5 mg		60	✓ Carvedilol Sandoz
* Tab 25 mg		60	✓ Carvedilol Sandoz
LABETALOL			
* Tab 100 mg	14.50	100	✓ Trandate
* Tab 200 mg		100	✓ Trandate
* Inj 5 mg per ml, 20 ml ampoule		5	
, ,	(88.60)		Trandate
* inj 5 mg per ml, 20 ml vial	42.29	1	
	(48.20)		Alvogen S29
METOPROLOL SUCCINATE			-
* Tab long-acting 23.75 mg	1.45	30	✓ Betaloc CR
* Tab long-acting 47.5 mg		30	✓ Betaloc CR
* Tab long-acting 95 mg		30	✓ Betaloc CR
* Tab long-acting 190 mg		30	✓ Betaloc CR

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

^{*}Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
METOPROLOL TARTRATE				
* Tab 50 mg	5.66	100	✓	IPCA-Metoprolol
* Tab 100 mg		60	✓	IPCA-Metoprolol
* Tab long-acting 200 mg	23.40	28	✓	Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial	26.50	5	✓	Metoprolol IV Mylan
NADOLOL				
* Tab 40 mg	19.19	100	✓	Nadolol BNM
* Tab 80 mg		100	✓	Nadolol BNM
PROPRANOLOL				
Tab 10 mg	7.04	100	1	Drofate
* Tab 40 mg		100	✓	IPCA-Propranolol
* Cap long-acting 160 mg	18.17	100	✓	Cardinol LA
* Oral liq 4 mg per ml - Special Authority see SA1327 below	_			
Retail pharmacy	CBS	500 m		Roxane-
				Propranolol S29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

AMI ODIDINE

*	Tab 80 mg	37.50	500	Mylan
*	Tah 160 mg	14 00	100	✓ Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AIV	ILODIPINE			
*	Tab 2.5 mg1.	08	90	✓ Vasorex
*	Tab 5 mg	96	90	✓ Vasorex
	Tab 10 mg1.		90	✓ Vasorex
FΕ	LODIPINE			
*	Tab long-acting 2.5 mg	45	30	✓ Plendil ER
*	Tab long-acting 5 mg4.	07	90	Felo 5 ER
*	Tab long-acting 10 mg4.	32	90	✓ Felo 10 ER

	Subsidy (Manufacturer's Price)	D-	Fully Subsidised	Brand or Generic
	\$	Per		Manufacturer
NIFEDIPINE				
* Tab long-acting 10 mg	18.80	56	1	Tensipine MR10 S29
* Tab long-acting 20 mg	9.12	50	✓	Mylan (12 hr release) \$29
	17.72	100	1	Nyefax Retard
* Tab long-acting 30 mg	4.78	14	✓	Mylan Italy (24 hr release) \$29
	34.10	100	•	Mylan (24 hr release) \$29
* Tab long-acting 60 mg	52.81	100	✓	Mylan (24 hr release) \$29
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
Cap extended-release 120 mg	44.40	100	1	Accord S29
* Cap long-acting 120 mg		500	1	Apo-Diltiazem CD
	65.35		1	Diltiazem CD Clinect
★ Cap long-acting 180 mg	7.00	30	✓	Cardizem CD
★ Cap long-acting 240 mg		30	✓	Cardizem CD
Apo-Diltiazem CD Cap long-acting 120 mg to be delisted 1 Juni PERHEXILINE MALEATE K Tab 100 mg	,	100	•	Pexsig
VERAPAMIL HYDROCHLORIDE				
* Tab 40 mg		100		Isoptin
₭ Tab 80 mg		100	✓	Isoptin
★ Tab long-acting 120 mg	36.02	100	✓	Isoptin Retard S29
				Isoptin SR
* Tab long-acting 240 mg		30	•	Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO		5	1	Isoptin
Centrally-Acting Agents				
CLONIDINE				
★ Patch 2.5 mg, 100 mcg per day – Only on a prescription	10.34	4	1	Mylan
Patch 5 mg, 200 mcg per day — Only on a prescription		4		Mylan
Patch 7.5 mg, 300 mcg per day — Only on a prescription		4		Mylan
CLONIDINE HYDROCHLORIDE		•	-	
	20 22	112	1	Clonidine Teva
★ Tab 25 mcg ★ Tab 150 mcg		100		Catapres
k Inj 150 mcg per ml, 1 ml ampoule		100		<u>Catapres</u> Medsurge
r III) 150 Mcg per Mi, 1 Mi ampoule	29.00	10	•	<u>measurge</u>
k Tab 250 mg	15 10	100	1	Methyldopa Mylan
	52.85	500		Methyldopa Mylan
	02.00	500	•	S29 S29

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price		idised	Generic
	\$	Per		Manufacturer
Diuretics				
Didictios				
Loop Diuretics				
BUMETANIDE				
* Tab 1 mg	4.91	30	✓ B	Burinex S29 S29
G	16.36	100	✓ B	Burinex
* Inj 500 mcg per ml, 4 ml vial	7.95	5	✓ B	Burinex
FUROSEMIDE [FRUSEMIDE]				
Tab 40 mg - Up to 30 tab available on a PSO	8.00	1,000	✓	PCA-Frusemide
* Tab 500 mg		50	√ Ū	Irex Forte
	89.48		√ F	urosemid-
				Ratiopharm S29
	169.96	100	√ F	urosemid-
	. 55.55	.00	•	Ratiopharm \$29
				•
* Oral liq 10 mg per ml		30 ml OP		asix
* Inj 10 mg per ml, 25 ml ampoule		6		asix
* Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a l	PSO2.40	5	✓ F	urosemide-Baxter
Potassium Sparing Diuretics				
AMILORIDE HYDROCHLORIDE				
Oral liq 1 mg per ml	32.10	25 ml OP	✓ B	Biomed
EPLERENONE - Special Authority see SA1728 below - Retail				
Tab 25 mg		30	√ Ir	nspra
Tab 50 mg		30		nspra
■ SA1728 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Both: 1 Patient has heart failure with ejection fraction less than 40 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolad	0%; and			
2.2 Patient has experienced a clinically significant adv	erse effect while o	n optimal dos	ing of s	spironolactone.
METOLAZONE				
Tab 5 mg	CBS	1	✓ N	letolazone S29
		50	✓ Z	aroxolyn S29
SPIRONOLACTONE				
* Tab 25 mg	3.68	100	✓ S	piractin
* Tab 100 mg	10.65	100	✓ S	piractin
Oral liq 5 mg per ml	30.60	25 ml OP	✓ B	Biomed
Potassium Sparing Combination Diuretics				
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE				
* Tab 5 mg with furosemide 40 mg	8 63	28	√ □	rumil
		20	Ψ Γ	runnii
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZ		F0	./ •	laduratia
* Tab 5 mg with hydrochlorothiazide 50 mg	0.00	50	→ IV	loduretic

	Subsidy (Manufacturer's Pric \$	e) Subs Per	Fully sidised	Brand or Generic Manufacturer
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	_	rrow- Bendrofluazide
May be supplied on a PSO for reasons other than eme * Tab 5 mg	0 ,	500		rrow- Bendrofluazide
CHLOROTHIAZIDE	07.00	05I OD	4 D	lamad.
Oral liq 50 mg per ml	27.82	25 ml OP	• 6	iomed
Tab 25 mg	3.90	30	🗸 Ig	roton S29
•	6.95	50	✓ H	ygroton
Hygroton to be Principal Supply on 1 April 2023 (Igroton \$29 Tab 25 mg to be delisted 1 April 2023) INDAPAMIDE				
* Tab 2.5 mg	10.45	90	✓ D	apa-Tabs
Ü	11.61	100	✓ M	ylan Indapamide S29
Vasopressin receptor antagonists				

ГОLVAPTAN – Special Authority see SA2166 below – Re	tail pharmacy		
Tab 15 mg	873.50	28 OP	Jinarc
Tab 30 mg	873.50	28 OP	Jinarc
Tab 45 mg + 15 mg		56 OP	Jinarc
Tab 60 mg + 30 mg		56 OP	Jinarc
Tab 90 mg + 30 mg		56 OP	Jinarc

⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Fither:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

	Subsidy (Manufacturer's Price) \$	Full Subsidise Per •	•
Lipid-Modifying Agents			
Fibrates			
BEZAFIBRATE * Tab 200 mg * Tab long-acting 400 mg			Bezalip Bezalip Retard
Other Lipid-Modifying Agents			
ACIPIMOX * Cap 250 mg	21.56 25.44		Olbetam S29 S29 Olbetam
Resins			
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	32.89	30	Colestid
HMG CoA Reductase Inhibitors (Statins)			
ATORVASTATIN * Tab 10 mg * Tab 20 mg * Tab 40 mg * Tab 80 mg PRAVASTATIN * Tab 20 mg * Tab 40 mg ROSUVASTATIN - Special Authority see SA2093 below - Retai * Tab 5 mg * Tab 10 mg * Tab 20 mg * Tab 40 mg * Tab 40 mg * Tab 40 mg * Tab 5 mg * Tab 5 mg * Tab 5 mg * Tab 10 mg * Tab 10 mg * Tab 10 mg * Tab 20 mg * Tab 40 mg * Tab 40 mg * Tab 40 mg * Tab 40 mg * Tab 5 mg * Tab 5 mg * Tab 60 mg	9.24 26.54 26.54 3.61 il pharmacy 1.70 2.42 3.92 5.28	500 500 500 28 28 28 30 30 30 30 30 30 30	Lorstat Lorstat Lorstat Lorstat Pravastatin Mylan Pravastatin Mylan Rosuvastatin Viatris Rosuvastatin Viatris Rosuvastatin Viatris Rosuvastatin Viatris
1.1 Patient is considered to be at risk of cardiovascula1.2 Patient is Māori or any Pacific ethnicity; or2 Both:	r disease; and		
O.1. Deticat has a calculated vials of conditions and an elec-	of ot locat 450/ -		1

- 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
- 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

Subsidy (Manufacturer's Price)	F Subsidi	ully	Brand or Generic
 \$	Per	•	Manufacturer

continued...

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
 - 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIMVASTATIN

*	Tab 10 mg	90	Simvastatin Mylan
	Tab 20 mg	90	✓ Simvastatin Mylan
	Tab 40 mg	90	✓ Simvastatin Mylan
	Tab 80 mg7.12	90	✓ Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Special Authority see SA1045 below – Retail	pharmacy		
* Tab 10 mg	1.95	30	✓ Ezetimibe Sandoz

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 x normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 on the next page - Retail pharmacy

Tab 10 mg with simvastatin 10 mg	5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg	7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	✓ Zimybe

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per	Manufacturer
⇒SA1046 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals	valid for 2 years for applic	ations meeting t	he following criteria:

All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and

* Oral numn spray 400 mcg per dose - Unito 250 dose

Tab long-acting 60 mg.......9.25

3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

-1-	available on a PSO	6.09	250 dose OP	✓ Nitrolingual Pump Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
*	Patch 50 mg, 10 mg per day	18.62	30	✓ Nitroderm TTS
ISC	SORBIDE MONONITRATE			
*	Tab 20 mg	19.55	100	✓ Ismo 20
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard

Svm	pat	hor	nime	tics

ADF	REN	AL	INE
-----	-----	----	-----

DRENALINE		
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	Aspen Adrenaline
12.65		✓ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	Aspen Adrenaline

Vasodilators

HYDRALAZINE HYDROCHLORIDE

*	Tab 25 mg - Special Authority see SA1321 below - Retail			
	pharmacy	CBS	1	Hydralazine
			56	✓ Onelink S29
			84	✓ AMDIPHARM S29
			100	✓ Onelink S29
*	Inj 20 mg ampoule	25.90	5	✓ Apresoline

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIL

▲ Tab 10 mg	78.40	100	Loniten
-------------	-------	-----	---------------------------

90

✓ Duride

Reddy's

Reddy's

✓ Bosentan Dr

60

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	
NICORANDIL				
▲ Tab 10 mg	25.57	60	✓	Ikorel
▲ Tab 20 mg	32.28	60	•	Ikorel
PAPAVERINE HYDROCHLORIDE				
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	✓	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]				
Tab 400 mg	42.26	50	1	Trental 400
Endothelin Receptor Antagonists				
AMBRISENTAN – Special Authority see SA1702 below – Retail Tab 5 mg Tab 10 mg	1,550.00	30 30	√	Ambrisentan Mylan Ambrisentan Mylan Ambrisentan Viatris Mylan
(Ambrisentan Mylan Tab 10 mg to be delisted 1 March 2023)				•
Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertensic Notes: Application details may be obtained from Pharmac's websithe Coordinator, PAH Panel Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac BOSENTAN − Special Authority see SA1991 below − Retail pharmac	site <u>schedule.pharma</u> .govt.nz rmacy			
Tab 62.5 mg	119.00	60	V	Bosentan Dr

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and

- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and

|--|

continued...

- 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1992 below – Retail phar	macy		
Tab 25 mg	0.85	4	✓ Vedafil
Tab 50 mg	1.70	4	✓ Vedafil
Tab 100 mg		12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
 \$	Per	•	Manufacturer

continued

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV: and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dvn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL – Special Authority see SA1696 below – Retail pharmacy		
Inj 500 mcg vial	1	✓ Veletri
Inj 1.5 mg vial73.21	1	✓ Veletri
⇒SA1696 Special Authority for Subsidy		
Special Authority approved by the Pulmonary Arterial Hypertension Panel		
A P P P P P P P P P P P P P P P P P P P		/O A E

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 on the next page - Retail pharmacy

30 ✓ Vebulis 740.10 ✓ Ventavis

Vebulis to be Principal Supply on 1 March 2023

(Ventavis Nebuliser soln 10 mcg per ml, 2 ml to be delisted 1 March 2023)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA2023 below - Retail	pharmacy		
Cap 5 mg	11.26	60	Oratane
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice: and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

TRFTINOIN

Crm 0.5 mg per g − Maximum of 50 g per prescription15.57 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

HYDROGEN PEROXIDE

* Crm 1%	8.56		✓ Crystaderm✓ Crystaderm
MUPIROCIN			
Oint 2%	6.60	15 g OP	
	(11.50)	Ü	Bactroban

- a) Only on a prescription
- b) Not in combination

DERMATOLOGICALS

	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.59	5 g OP	√ F	oban
a) Maximum of 5 g per prescriptionb) Only on a prescriptionc) Not in combination		Ü	_	
Oint 2% a) Maximum of 5 g per prescription b) Only on a prescription c) Not in combination	1.59	5 g OP	√ <u>F</u>	<u>oban</u>
SULFADIAZINE SILVER Crm 1%	10.80	50 g OP	√ F	ilamazine
Antifungals Topical For systemic antifungals, refer to INFECTIONS, Antifungals, p.	age 97			
AMOROLFINE				
a) Only on a prescription b) Not in combination				
Nail soln 5%	14.93	5 ml OP	✓ <u>N</u>	<u>IycoNail</u>
CLOTRIMAZOLE * Crm 1%	1.10	20 g OP	✓ 0	Clomazol
a) Only on a prescription b) Not in combination				
c) Clomazol to be Principal Supply on 1 April 2023 * Soln 1%	4.36 (7.55)	20 ml OP	(Canesten
a) Only on a prescriptionb) Not in combination	(7.00)			variosion
ECONAZOLE NITRATE				
Crm 1%	1.00 (7.48)	20 g OP	F	evaryl
a) Only on a prescription b) Not in combination				
Foaming soln 1%, 10 ml sachets	9.89 (17.23)	3	P	'evaryl
, 0 ,	` -/		-	,

a) Only on a prescriptionb) Not in combination

	Subsidy		Fully	Brand or
	(Manufacturer's Pr \$	rice) Subs Per	sidised •	Generic Manufacturer
MICONAZOLE NITRATE				
* Crm 2%	0.81	15 g OP	✓ <u>N</u>	<u>lultichem</u>
b) Not in combination				
* Lotn 2%	4.36 (10.03)	30 ml OP	D)aktarin
a) Only on a prescriptionb) Not in combination				
* Tinct 2%	4.36 (12.10)	30 ml OP	D)aktarin
a) Only on a prescriptionb) Not in combination				
Antipruritic Preparations				
CALAMINE				

٠.			•••	•	•
	2	1)	C	nl	v

a) Only on a prescription

b) Not in combination

CROTAMITON

a) Only on a prescription

b) Not in combination

Crm 10%......3.29 20 g OP

29.60

✓ Itch-Soothe

MENTHOL - Only in combination

1) Only in combination with a dermatological base or proprietary Topical Corticosteriod - Plain

2) With or without other dermatological galenicals.

_,g.....g......g......

25 g 100 g ✓ MidWest

✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 79

Crystals......6.92

Corticosteroids - Plain

BETAMETHASONE DIPROPIONATE		
Crm 0.05%2.96	15 g OP	✓ Diprosone
36.00	50 g OP	✓ Diprosone
Oint 0.05%2.96	15 g OP	✓ Diprosone
36.00	50 g OP	✓ Diprosone
Oint 0.05% in propylene glycol base4.33	30 g OP	Diprosone OV
BETAMETHASONE VALERATE		
* Crm 0.1%4.53	50 g OP	Beta Cream
* Oint 0.1%	50 g OP	✓ Beta Ointment
* Lotn 0.1%	50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE		
* Crm 0.05%	30 g OP	✓ Dermol
* Oint 0.05%2.33	30 g OP	✓ Dermol

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's P	rico) Subc	Fully Brand idised Gener	
	(Manufacturer's P	Per Subs		acturer
LOBETASONE BUTYRATE				
Crm 0.05%	5.38	30 g OP		
	(10.00)		Eumovat	te
YDROCORTISONE				
Crm 1% - Only on a prescription	1.78	30 g OP	✓ Ethics	
, , ,	17.15	500 g	✓ Hydroco	rtisone
			(PSM)	
Ethics to be Principal Supply on 1 April 2023				
Powder – Only in combination		25 g	✓ ABM	
Up to 5% in a dermatological base (not proprietary Top	ical Corticosteriod	- Plain) with o	r without other	dermatologica
galenicals				
YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only	on			
a prescription		250 ml	DP Lotn	HC
YDROCORTISONE BUTYRATE				
Lipocream 0.1%	4.85	100 g OP	✓ Locoid I	_ipocream
Oint 0.1%		100 g OP	✓ Locoid	
Milky emul 0.1%	12.33	100 ml OP	✓ Locoid (Crelo
ETHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.46	15 g OP	✓ Advanta	ın
Oint 0.1%	4.46	15 g OP	✓ Advanta	_
OMETASONE FUROATE		Ü		_
Crm 0.1%	1 95	15 g OP	✓ Flocon	Alcohol Free
G111 G1 7/6	3.10	50 g OP		Alcohol Free
Oint 0.1%		15 g OP	✓ Elocon	
	2.90	50 g OP	✓ Elocon	
Lotn 0.1%	4.50	30 ml OP	✓ Elocon	
RIAMCINOLONE ACETONIDE				
Crm 0.02%	6.30	100 g OP	✓ Aristoco	ort
Oint 0.02%		100 g OP	✓ Aristoco	
				_
Corticosteroids - Combination				
ETAMETHASONE VALERATE WITH SODIUM FUSIDATE [F	USIDIC ACID]			
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		
	(10.45)	-	Fucicort	
a) Maximum of 15 g per prescription				
b) Only on a prescription				
YDROCORTISONE WITH MICONAZOLE - Only on a prescr	iption			
Crm 1% with miconazole nitrate 2%	•	15 g OP	✓ Micreme	H
YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN -	Only on a prescrip	tion		
Crm 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP	✓ Pimafuc	ort
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP	✓ Pimafuc	ort
Pimafucort Crm 1% with natamycin 1% and neomycin sulphate)	
RIAMCINOLONE ACETONIDE WITH GRAMICIDIN NEOMY				
RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMY(
RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYO Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 r and gramicidin 250 mcg per g — Only on a prescription	ng	15 g OP		

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

	\$	Per	_	Manufacturer
Barrier Creams and Emollients				
Barrier Creams				
DIMETHICONE				
* Crm 5% pump bottle	4.30	500 ml OP	/	healthE
* Crm 10% pump bottle	4.52	500 ml OP	/	Dimethicone 5% healthE Dimethicone 10%
ZINC AND CASTOR OIL				
* Oint	4.65	500 g	•	Boucher
Finalliante				
Emollients				
AQUEOUS CREAM				
* Crm	1.73	500 g		Evara GEM Aqueous Cream
(Evara Crm to be delisted 1 April 2023)				Oreani
CETOMACROGOL				
* Crm BP	1.99	500 g	1	Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL		· ·		
Crm 90% with glycerol 10%	2.13	500 ml OP	1	Evara
	2.35			Boucher
			/	Pharmacy Health Sorbolene with Glycerin
	3.10	1.000 ml OP	/	Boucher
	3.50	1,000 1111 01		Evara
(Boucher Crm 90% with glycerol 10% to be delisted 1 March 2023)				
(Pharmacy Health Sorbolene with Glycerin Crm 90% with glycerol 10% (Boucher Crm 90% with glycerol 10% to be delisted 1 March 2023)	to be delis	sted 1 July 2023))	
EMULSIFYING OINTMENT	0.40	500	,	
* Oint BP	3.40	500 g	•	Emulsifying Ointment ADE
OIL IN WATER EMILI SION				Ontillent ADE
OIL IN WATER EMULSION * Crm	2 04	500 g	/	Fatty Cream AFT
PARAFFIN	2.07	300 g	•	ratty Orealii Ai I
Oint liquid paraffin 50% with white soft paraffin 50%	4.94	500 g OP	1	White Soft Liquid Paraffin AFT
White Soft Liquid Paraffin AFT to be Principal Supply on 1 May	2023			
Oint liquid paraffin 50% with white soft paraffin 50%,(healthE Oint liquid paraffin 50% with white soft paraffin 50%, to be deli		500 ml OP y 2023)	•	healthE
UREA				
* Crm 10%	1.37	100 g OP	/	healthE Urea Cream

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

DERMATOLOGICALS

	Subsidy (Manufacturer's F \$	Price) Subsi	Fully idised	Brand or Generic Manufacturer
WOOL FAT WITH MINERAL OIL - Only on a prescription				
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml		
	(11.95)		D	P Lotion
	1.40	250 ml OP		
	(4.53)		D	P Lotion
	5.60	1,000 ml		
	(20.53)		Α	lpha-Keri Lotion
	(23.91)		В	K Lotion
	1.40	250 ml OP		
	(7.73)		В	K Lotion

Other Dermatological Bases

PARAFFIN

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

Minor Skin Infections

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	4.15	100 ml	✓ Riodine
Antiseptic soln 10%		15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer

Parasiticidal Preparations

DIM	ETF	IIC(ONE

		Lotion	
IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy			
Tab 3 mg - Up to 100 tab available on a PSO17.20	4	✓ Stromectol	

- PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

continued...

200 ml OP

✓ healthE

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

DERMATOLOGICALS

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	•	Manufacturer

continued...

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5%5.75	30 g OP	✓ Lyderm
Lotn 5%	30 ml OP	✓ A-Scabies

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA2024 below - Retail pl	harmacy		
Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment:
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CAI CIPOTRIOI

59.95	60 g OP	Enstilar
39.35	60 g OP	✓ Daivobet
15.90	30 g OP	 Daivobet
40.00	120 g OP	Daivonex
36.25	200 ml	✓ Midwest
	39.35 15.90 40.00	39.35 60 g OP 30 g OP 30 g OP 40.00 120 g OP

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

	Subsidy		Fully	Brand or
	(Manufacturer's Price \$	e) Subs Per	sidised	Generic Manufacturer
COAL TAR WITH ALL ANTOIN MENTION RUENOL AND OUR	*	FEI		Manuacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SUL				
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% ar		75 ~ OD		
allantoin crm 2.5%	(8.00)	75 g OP	E	gopsoryl TA
	3.43	30 g OP	Ľ	Jopsoryi TA
	(4.35)	00 g 01	Ed	gopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR	(1122)		_,	,-p,
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ C	oco-Scalp
2011 12/2 1111 2413/10 4514 2/2 4/14 241/2 11/2 21/1111111111111	7.95	40 g OP		oco-Scalp
PIMECROLIMUS - Special Authority see SA1970 below - Reta	il nharmacy	3		
a) Maximum of 15 g per prescription	iii priarriacy			
b) Note: a maximum of 15 g per prescription and no more	than one prescription	n per 12 we	eks.	
Cream 1%	28.50	15 g OP	✓ EI	idel
⇒SA1970 Special Authority for Subsidy		Ü	_	
Initial application only from a dermatologist, paediatrician, opht	halmologist or any r	elevant prad	ctitioner o	on the recommendation
of a dermatologist, paediatrician or ophthalmologist. Approvals				
meeting the following criteria:				
Both:				
1 Patient has atopic dermatitis on the eyelid; and				
2 Patient has at least one of the following contraindications				
documented epidermal atrophy, documented allergy to to	pical corticosteroids	, cataracts,	glaucom	na, or raised intraocular
pressure.				
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE				
* Soln 2.3% with trolamine laurilsulfate and fluorescein sodiur	n4.44	500 ml	✓ <u>Pi</u>	netarsol
SALICYLIC ACID				
Powder – Only in combination	18.88	250 g	✓ M	idwest
1) Only in combination with a dermatological base or	proprietary Topical	Corticostero	oid – Pla	in or collodion flexible
With or without other dermatological galenicals.				
SULPHUR				
Precipitated - Only in combination	6.35	100 g	✓ M	idwest
1) Only in combination with a dermatological base or	proprietary Topical	Corticostero	oid – Pla	in
With or without other dermatological galenicals.				
TACROLIMUS				
Oint 0.1% - Special Authority see SA2074 below - Retail				
pharmacy	33.00	30 g OP	✓ Ze	ematop
a) Maximum of 30 g per prescription				
b) Note: a maximum of 30 g per prescription and no m	ore than one prescr	iption per 12	2 weeks.	
⇒SA2074 Special Authority for Subsidy				
The state of the s				

2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist,

paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:

1 Patient has atopic dermatitis on the face; and

Both:

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Scalp Preparations

N 0 1 0 10 10 10 10 10 10 10 10 10 10 10	
★ Scalp app 0.1%	calp
* Scalp app 0.05%	<u>) l</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	<u>i</u>
KETOCONAZOLE Shampoo 2%	ole ole

a) Maximum of 100 ml per prescription

b) Only on a prescription

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Lotn,......6.50 200 g OP

✓ Marine Blue Lotion SPF 50+

Marine Blue Lotion SPF 50+ to be Principal Supply on 1 April 2023

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

IMIQUIMOD

PODOPHYLLOTOXIN

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

GENITO-URINARY SYSTEM

				_
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	•	Manufacturer	

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised <	Generic Manufacture
Contraceptives - Non-hormonal				
Condoms				
ONDOMS				
49 mm - Up to 144 dev available on a PSO	11.42	144	1	Moments
₹ 53 mm	0.95	10	1	Moments
	11.64	144	•	Moments
 a) Maximum of 60 dev per prescription 				
b) Up to 60 dev available on a PSO				
53 mm, 0.05 mm thickness		10		Moments
	11.42	144	/	Moments
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription			_	
53 mm, chocolate, brown		10		Moments
)	11.64	144	•	Moments
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription	0.05	10	./	Moments
53 mm, strawberry, red		144		Moments
a). Un to 60 day available on a DCO	11.64	144	•	woments
a) Up to 60 dev available on a PSOb) Maximum of 60 dev per prescription				
56 mm	n 97	10	1	Moments
30 11111	11.64	144		Moments
a) Maximum of 60 dev per prescription	11.01			momonto
b) Up to 60 dev available on a PSO				
56 mm, 0.05 mm thickness	1.30	12	1	Gold Knight
•	15.57	144		Gold Knight
a) Up to 60 dev available on a PSO				_
b) Maximum of 60 dev per prescription				
56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓	Gold Knight
 a) Maximum of 60 dev per prescription 				
b) Up to 60 dev available on a PSO				
56 mm, 0.08 mm thickness		10		Moments
	11.64	144	•	Moments
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
56 mm, 0.08 mm thickness, red		10		Moments
a) He to 00 days and lighter as a BOO	11.64	144	•	Moments
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription 56 mm, chocolate	1 20	12	./	Gold Knight
Jo min, Giocolate	15.57	144		Gold Knight
a) Up to 60 dev available on a PSO	10.07	177	•	acia itiligili
b) Maximum of 60 dev per prescription				
56 mm, strawberry	1.30	12	1	Gold Knight
· · · · · · · · · · · · · · · · · ·	15.57	144		Gold Knight
a). Un to 60 day available on a DCO				

a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription

12

144

14.87

17.02

✓ Gold Knight XL

✓ Gold Knight XL

✓ Shield XL

copper Short

✓ Choice TT380 Short

TT380 Standard

✓ Choice Load 375

✓ Choice

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
# 60 mm (bulk pack) a) Maximum of 60 dev per prescription b) Up to 60 dev available on a PSO	14.87	144	√	Gold Knight XL
Contraceptive Devices				
INTRA-UTERINE DEVICE a) Up to 40 dev available on a PSO b) Only on a PSO * IUD 29.1 mm length × 23.2 mm width	29.80	1		7 MED NSHA Silver/ Copper Short Choice 380 7med Nsha Silver/

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and

Choice TT380 Standard to be Principal Supply on 1 April 2023

* IUD 35.5 mm length × 19.6 mm width33.00

Choice Load 375 to be Principal Supply on 1 April 2023

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- . on a Social Welfare benefit; or
- · have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

★ Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO.......10.00 84 Mercilon 28

GENITO-URINARY SYSTEM

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	✓	Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	2.18	84	✓ I	Microgynon 20 ED
'	6.45	112		Femme-Tab ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
	(16.50)		N	Microgynon 30
 a) Higher subsidy of \$15.00 per 63 tab with Special Aut b) Up to 63 tab available on a PSO * Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets 	·	the p	orevious pa	ge
Up to 112 tab available on a PSO		84	✓ I	evlen ED
op to 112 ab available 511 a 1 00	6.45	112		Femme-Tab ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to 84 tab available on a PSO	12.25	84	✓ [Brevinor 1/28
to 84 tab available on a PSO		84	1	Norimin
	29.32	112	√ i	Norimin

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

* Tab 30 mcg - Up to 84 tab available on a PSO16.50 22.00	84 112	✓ Microlut✓ Microlut
* Subdermal implant (2 × 75 mg rods) – Up to 3 pack available on a PSO106.92	1	✓ <u>Jadelle</u>
MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO9.18	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price)		Fully	Brand or Generic	
	\$	Per		Manufacturer	
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84	✓ <u>N</u>	oriday 28	
Emergency Contraceptives					
LEVONORGESTREL					
* Tab 1.5 mg	1.75	1		evonorgestrel BNM	
	4.95		✓ P	ostinor-1	
a) Maximum of 2 tab per prescription					
b) Up to 5 tab available on a PSO					
c) Note: Direct Provision by a pharmacist permitted u	inder the provisions in F	Part I of	Section A		

(Postinor-1 Tab 1.5 mg to be delisted 1 June 2023) Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID		
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP	
(24.15)		Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators	35 g OP	✓ Clomazol
Clomazol to be Principal Supply on 1 April 2023	· ·	
* Vaginal crm 2% with applicators	20 g OP	✓ Clomazol
Clomazol to be Principal Supply on 1 April 2023	3 -	
MICONAZOLE NITRATE		
	40 ~ OD	./ Mierome
* Vaginal crm 2% with applicator6.89	40 g OP	✓ <u>Micreme</u>
NYSTATIN		
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ Nilstat
Myometrial and Vaginal Hormone Preparations		
myoniculal and vaginal normone i reparations		

EDCOMETRINE MALEATE

Lnc	OWILTHINE WALLATE			
	Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj ava	ilable on a		
	PSO	160.00	5	 DBL Ergometrine
OES	STRIOL			
*	Crm 1 mg per g with applicator	6.62	15 g OP	✓ Ovestin
*	Pessaries 500 mcg	6.86	15	✓ Ovestin

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
OXYTOCIN – Up to 5 inj available on a PSO				
Inj 5 iu per ml, 1 ml ampoule	4.98	5	✓	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	5.98	5	✓	Oxytocin BNM
			1	Oxytocin GH S29
OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj avail Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo		5	•	Syntometrine

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

Cassette	12.00	40 test OP	✓ Smith BioMed Rapid
	16.00		Pregnancy Test ✓ David One Step
			Cassette
			Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 107

5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

* Tab 5 mg4.81 100

Ricit

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Fither:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE – Special Authority see SA1032 below – Retail pharmacy

★ Cap 400 mcg22.31 100

✓ Tamsulosin-Rex

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN

GENITO-URINARY SYSTEM

GENITO-ONINANT STSTEM					
	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully sidised	Brand or Generic Manufacturer	
POTASSIUM CITRATE					
Oral liq 3 mmol per ml — Special Authority see SA1083 belo Retail pharmacy		200 ml OP	✓ B	iomed	
⇒SA1083 Special Authority for Subsidy					
Initial application from any relevant practitioner. Approvals vali Both:	d for 12 months f	or applications	meetin	g the following criteria:	
1 The patient has recurrent calcium oxalate urolithiasis; and	١				
2 The patient has had more than two renal calculi in the two		e application.			
Renewal from any relevant practitioner. Approvals valid for 2 ye benefitting from the treatment.			is appro	priate and the patient is	
SODIUM CITRO-TARTRATE					
* Grans eff 4 g sachets	2.22	28	√ <u>U</u>	<u> ral</u>	
SOLIFENACIN SUCCINATE					
Tab 5 mg	2.05	30	_	olifenacin Mylan	
Tah 10 mg	2.72	30	-	olifenacin Viatris olifenacin Mylan	
Tab 10 mg	3.72	30	_	olifenacin Viatris	
Detection of Substances in Urine					
ORTHO-TOLIDINE	7.50	50 to at 0.5			
* Compound diagnostic sticks	7.50 (8.25)	50 test OP	Н	lemastix	
TETRABROMOPHENOL	. ,				
* Blue diagnostic strips	13.92	100 test OP	✓ A	lbustix	

Obstetric Preparations

Antiprogesterones

NΛ	PRIS	חדי	NIE.
IVI	יוחי	ווו	חעו

			MILELLISIONE
✓ Mifegyne	1	- Up to 15 tab available on a PSO60.00	Tab 200 mg
✓ Mifegyne	3	180.00	

Subsidy (Manufacturer's Price)	Fully		Brand or
(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer

Calcium Homeostasis

CALCITONIN * Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ Miacalcic
CINACALCET - Special Authority see SA2170 below - Retail	pharmacy		
Tab 30 mg - Wastage claimable	42.06	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable	84.12	28	✓ Cinacalet Devatis

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

- Either:

 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
 - 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (**primary hyperparathyroidism**) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

|--|

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

⇒SA2109 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement: and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer*) from any relevant practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 3 years.

Note: Indications marked with * are unapproved indications.

Initial application — (symptomatic hypercalcaemia*) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has symptomatic hypercalcaemia .

Note: Indications marked with * are unapproved indications.

Corticosteroids and Related Agents for Systemic Use

DETAMETIMACONE CODUNA DINOCRIMATE MUTU DETAMETIMACONE ACETATE

RE	TAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE /	ACETATE		
*	Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	19.20	5	
	(3	36.96)		Celestone
	`	,		Chronodose
DE	XAMETHASONE			
*	Tab 0.5 mg - Up to 60 tab available on a PSO	.1.50	30	 Dexmethsone
*	Tab 4 mg - Up to 30 tab available on a PSO	.2.65	30	✓ Dexmethsone
	Oral lig 1 mg per ml4	18.15 2	25 ml OP	✓ Biomed

	Subsidy (Manufactureria Bria	٠ (-	Fully	
	(Manufacturer's Pric \$	e) S Per	Subsidised •	Generic Manufacturer
EXAMETHASONE PHOSPHATE	·			
Dexamethasone phosphate injection will not be funded for	oral use.			
Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a		10	1	Hameln
Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a	PSO 13.10	10		Hameln
	1 00 10.10			<u> </u>
LUDROCORTISONE ACETATE Tab 100 mcg	44.40	100		Florings
Tab 100 mcg	11.46	100	•	<u>Florinef</u>
'DROCORTISONE				
Tab 5 mg	8.10	100		Douglas
Tab 20 mg	20.32	100	•	Douglas
Inj 100 mg vial	4.38	1	•	Solu-Cortef
a) Up to 5 inj available on a PSO				
b) Only on a PSO				
THYLPREDNISOLONE				
Tab 4 mg	112 00	100	1	Medrol
Tab 100 mg		20		Medrol
· ·	220.10	20	•	Wedioi
ETHYLPREDNISOLONE (AS SODIUM SUCCINATE)			_	
Inj 40 mg vial	22.30	1	•	Solu-Medrol-Act-
				O-Vial
let 405 er er et al	04.40		,	Only Madeal Art
Inj 125 mg vial	34.10	1	•	Solu-Medrol-Act-
				O-Vial
Inj 500 mg vial	26.00	1	1	Solu-Medrol-Act-
iiij 500 iiig viai	20.00	'	•	O-Vial
				U-Viai
Inj 1 g vial	32.84	1	1	Solu-Medrol
ETHYLPREDNISOLONE ACETATE				
Inj 40 mg per ml, 1 ml vial	47.06	5	./	Depo-Medrol
	47.06	5	•	Depo-ineuroi
EDNISOLONE				
Oral liq 5 mg per ml - Up to 30 ml available on a PSO	6.00	30 ml Ol		Redipred
Restricted to children under 12 years of age.				
EDNISONE				
Tab 1 mg	18.58	500	1	Prednisone Clinect
Tab 2.5 mg		500	1	Prednisone Clinect
Tab 5 mg - Up to 30 tab available on a PSO		500		Prednisone Clinect
Tab 20 mg - Up to 30 tab available on a PSO		500	1	Prednisone Clinect
TRACOSACTRIN				
	75.00	1	./	All Synaether
Inj 250 mcg per ml, 1 ml ampoule	75.00	1		AU Synacthen
				Synacthen
Ini 1 ma nor ml. 1 ml amnoula	600.00	4		UK Synacthen
Inj 1 mg per ml, 1 ml ampoule	090.00	1		Synacthen Depot
			•	Synacthene
				Retard S29
U Synacthen Inj 250 mcg per ml, 1 ml ampoule to be deliste	d 1 March 2023)			
NAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	1	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5		Kenacort-A 40
", 10 mg por mi, 1 mi ampoulo		J	•	INDIGOUS TO

Hormones Non Contraceptive					Ī
	\$	Per	✓	Manufacturer	
	(Manufacturer's Price)		Subsidised	Generic	
	Subsidy		Fully	Brand or	

Androgen Agonists and Antagonists

CYPROTERONE ACETATE		
Tab 50 mg14.37	50	✓ <u>Siterone</u>
Tab 100 mg28.03	50	✓ Siterone
TESTOSTERONE		
Patch 5 mg per day225.00	30	✓ Androderm
TESTOSTERONE CIPIONATE		
Inj 100 mg per ml, 10 ml vial85.00	1	✓ Depo-Testosterone
393.00		✓ Taro-
		Testosterone S29
TESTOSTERONE ESTERS		
Inj 250 mg per ml, 1 ml12.98	1	✓ Sustanon Ampoules
TESTOSTERONE UNDECANOATE		
Cap 40 mg - Subsidy by endorsement21.00	60	✓ Andriol Testocaps
35.00	100	✓ Steril-Gene S29

Subsidy by endorsement - subsidised for patients who were taking testosterone undecanoate cap 40mg prior to 1 November 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of testosterone undecanoate cap 40 mg in the preceding 12 months.

✓ Reandron 1000 Inj 250 mg per ml, 4 ml vial......86.00

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer **Hormone Replacement Therapy - Systemic** Oestrogens **OESTRADIOL** 28 OP Estrofem Tab 2 mg4.12 28 OP Estrofem Patch 50 mcg per 24 hours7.04 ✓ Climara a) No more than 1 patch per week b) Only on a prescription ✓ Estradot 13.50 ✓ Fetraderm MX S29 a) No more than 2 patch per week b) Only on a prescription Patch 50 mcg per day......7.04 8 ✓ Estradot 50 mcg ✓ Estradiol TDP Mylan S29 14.50 ✓ Estraderm MX S29 a) No more than 2 patch per week b) Only on a prescription Patch 75 mcg per day......7.91 8 ✓ Estradot ✓ Estradiol TDP 10.60 Mylan S29 a) No more than 2 patch per week b) Only on a prescription Patch 100 mcg per day......7.91 ✓ Estradot ✓ Estraderm MX S29 15.50 a) No more than 2 patch per week b) Only on a prescription **OESTRADIOL VALERATE** 84 ✓ Progynova ✓ Progynova **OESTROGENS** Conjugated, equine tab 300 mcg......3.01 28 (17.50)Premarin Conjugated, equine tab 625 mcg......4.12 28 (17.50)Premarin

Progestogens

ME	DROXYPROGESTERONE ACETATE		
*	Tab 2.5 mg	30	✓ Provera
	8.75	56	Provera
*	Tab 5 mg	56	Provera
	17.50	100	Provera
*	Tab 10 mg8.94	30	Provera

	Subsidy (Manufacturer's Pric	e) Sub	Fully Brand or sidised Generic ✓ Manufacturer
Progestogen and Oestrogen Combined Prepara	ntions		
OESTRADIOL WITH NORETHISTERONE			
* Tab 1 mg with 0.5 mg norethisterone acetate		28 OP	171
W. Tab O man with 4 man manathistanana anatata	(18.10)	00.00	Kliovance
* Tab 2 mg with 1 mg norethisterone acetate	(18.10)	28 OP	Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	(10.10)		Miogest
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP	
ocolitation tab (12) and 1 mg ocolitation tab (0)	(18.10)	20 01	Trisequens
	(10110)		· · · · · · · · · · · · · · · · · · ·
Other Oestrogen Preparations			
OESTRIOL			
* Tab 2 mg	7.00	30	✓ Ovestin
-			<u> </u>
Other Progestogen Preparations			
LEVONORGESTREL			
* Intra-uterine device 52 mg	269.50	1	✓ Mirena
* Intra-uterine device 13.5 mg	215.60	1	✓ Jaydess
MEDROXYPROGESTERONE ACETATE			
Tab 100 mg	116.15	100	✓ Provera HD
NORETHISTERONE			
* Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Primolut N
PROGESTERONE			
* Cap 100 mg	14.85	30	✓ Utrogestan
Utrogestan to be Principal Supply on 1 May 2023			3
Thyroid and Antithyroid Agents			
Thyroid and Antithyroid Agents			
CARBIMAZOLE			
* Tab 5 mg	7.56	100	✓ Neo-Mercazole
LEVOTHYROXINE			
* Tab 25 mcg	5.55	90	✓ Synthroid
* Tab 50 mcg		28	✓ Mercury Pharma
	5.79	90	Synthroid
¥ Tob 100 mag	64.28	1,000	✓ Eltroxin
* Tab 100 mcg	1./8 6.01	28 90	✓ Mercury Pharma✓ Synthroid
	6.01	1,000	✓ Synthroid ✓ Eltroxin
DDODYLTHOUDAOL Coosial Authority and CA4400 halves		1,000	- LIUVAIII
PROPYLTHIOURACIL – Special Authority see SA1199 below –		400	/ DTIL
Tab 50 mg	35.00	100	✓ PTU \$29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

_			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA2032 below	- Retail pharm	acy	
*	Inj 5 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 15 mg cartridge	139.50	1	✓ Omnitrope
	, ,			✓ Omnitrope S29 S29

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

-	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
=	<u> </u>				

continued...

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred;
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and

|--|

continued...

- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Price) Subsidise	d Generic
\$ Per	Manufacturer

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GOSERELIN	
-----------	--

Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe	122.37	1	✓ Teva

I FUPRORFI IN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of

(221.60)

Lucrin Depot 1-month

Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy of \$591.68 per 1 inj with Endorsement.......177.50

(591.68)Lucrin Depot 3-month

Vasopressin Agonists

DESMOPRESSIN Wafer 120 mcg47.00	30	✓ Minirin Melt
DESMOPRESSIN ACETATE		
Tab 100 mcg25.00	30	✓ Minirin
Tab 200 mcg54.45	30	✓ Minirin
▲ Nasal spray 10 mcg per dose27.95	6 ml OP	✓ <u>Desmopressin-</u> <u>PH&T</u>
Inj 4 mcg per ml, 1 ml67.18	10	✓ Minirin

Other Endocrine Agents

CABERGOI INF

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be	
Dostinex	2	waived by Special Authority see SA2070 below4.43	
✓ Dostinex	8	17.94	

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLOMIFFNE CITRATE

✓ Mylan 10 Clomiphen S29

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
METYRAPONE Cap 250 mg	558.00	50	✓ <u>M</u>	etopirone_

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	idised	Generic	
\$	Per	1	Manufacturer	

Anthelmintics

ALBENDAZOLE - Special Authority see SA1318 below - Reta	ail pharmacy		
Tab 400 mg	469.20	60	✓ Eskazole S29

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

Tab 100 mg	7.97	6	✓ Vermox
Oral liq 100 mg per 5 ml		15 ml	
	(7.53)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	 Biltricide

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 61
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 249

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE			
Cap 250 mg	24.70	100	✓ Ranbaxy-Cefactor S29 S29
	25.85		✓ Ranbaxy-Cefaclor
Ranbaxy-Cefaclor to be Principal Supply on 1 April 2023			
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.53	100 ml	✓ Ranbaxy-Cefactor S29 S29
	3.75		✓ Ranbaxy-Cefactor
Ranbaxy-Cefaclor to be Principal Supply on 1 April 2023			
(Ranbaxy-Cefaclor S29 S29 Cap 250 mg to be delisted 1 April 20	23)		
(Ranbaxy-Cefaclor S29 S29 Grans for oral liq 125 mg per 5 ml to	be delisted 1 A	pril 2023)	
CEFALEXIN			
Cap 250 mg	3.85	20	Cephalexin ABM
Cephalexin ABM to be Principal Supply on 1 April 2023			
Cap 500 mg	5.85	20	Cephalexin ABM
Cephalexin ABM to be Principal Supply on 1 April 2023			
Grans for oral liq 25 mg per ml - Wastage claimable		100 ml	✓ <u>Flynn</u>
Grans for oral liq 50 mg per ml – Wastage claimable	10.38	100 ml	✓ <u>Flynn</u>
CEFAZOLIN - Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance with a endorsed accordingly.	Health NZ Hos	oital approved	protocol and the prescription is
Inj 500 mg vial	3.39	5	✓ AFT
lnj 1 g vial		5	✓ AFT

	Subsidy (Manufacturer's Price) \$		Fully dised	Brand or Generic Manufacturer
CEFTRIAXONE – Subsidy by endorsement				
a) Up to 10 inj available on a PSO				
 Subsidised only if prescribed for a dialysis or cystic fibros pelvic inflammatory disease, or the treatment of suspecte endorsed accordingly. 				
Inj 500 mg vial	0.79	1	✓ C	eftriaxone-AFT
Inj 1 g vial	3.59	5	√ C	eftriaxone-AFT
Ceftriaxone-AFT to be Principal Supply on 1 April 2023		Ü		oranazono ya r
CEFUROXIME AXETIL - Subsidy by endorsement				
Only if prescribed for prophylaxis of endocarditis and the pre	scription is endorsed	accordingly	' .	
Tab 250 mg	45.93	50	✓ Z	innat

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg8.	19 30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO2.		✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage		
claimable16.	97 15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

(Zinnat Tab 250 mg to be delisted 1 March 2024)

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and

3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see \$A1857 below

Tab 250 mg	8.53	14	✓ Klacid
Grans for oral lig 250 mg per 5 ml - Wastage claimable	192.00	50 ml	Klacid

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Atypical mycobacterial infection; or

ERYTHROMYCIN (AS LACTOBIONATE)

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated.

Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician.

Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

lnj 1 g vial	10.00	1	 Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE Tab 400 mg	16 95	100	✓ E-Mvcin
a) Up to 20 tab available on a PSOb) Up to 2 x the maximum PSO quantity for RFPP			,
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	✓ E-Mycin
c) Wastage claimable Grans for oral liq 400 mg per 5 ml	6.77	100 ml	✓ E-Mycin

0.//	100 1111	♥ E-WyCifi
8.29	10	Rulide D
29.50	50	✓ Arrow-
		Roxithromycin
35.50	50	✓ Arrow-
		Roxithromycin
		8.29 10 29.50 50

(Rulide D Tab disp 50 mg to be delisted 1 March 2023)

	Subsidy		Fully	Brand or
	(Manufacturer's Pi	rice) Subs Per	idised	Generic Manufacturer
	\$	rei	<u> </u>	Manuacturei
Penicillins				
AMOXICILLIN				
Cap 250 mg	43.45	500	1	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	66.44	500	1	Alphamox
 a) Up to 30 cap available on a PSO 				
b) Up to 10 x the maximum PSO quantity for RFPP			_	
Grans for oral liq 125 mg per 5 ml	1.40	100 ml		Alphamox 125
 a) Up to 200 ml available on a PSO 				
b) Wastage claimable			_	
Grans for oral liq 250 mg per 5 ml	1.73	100 ml	/	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable	15.07	10	./	lhiamay
Inj 250 mg vialInj 500 mg vial		10 10		Ibiamox Ibiamox
Inj 1 g vial — Up to 5 inj available on a PSO		10		Ibiamox
	21.04	10	٠	IDIAIIIOX
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab	2.22	40		0 D 500//05
available on a PSO		10	•	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 i	•	100		A
per ml	0.50	100 ml	٧	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r per ml – Up to 200 ml available on a PSO	•	100 ml OP	./	Curam
·	2.20	100 IIII OF	•	Curain
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj	075 07	40	,	District A
available on a PSO	3/5.9/	10	•	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]			_	
Inj 600 mg (1 million units) vial – Up to 5 inj available on a P	SO 11.09	10		Sandoz
FLUCLOXACILLIN				
Cap 250 mg - Up to 30 cap available on a PSO		250		Flucloxacillin-AFT
Cap 500 mg - Up to 30 cap available on a PSO		500		Flucloxacillin-AFT
Grans for oral liq 25 mg per ml	3.29	100 ml	/	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable	0.00	4001	,	A F-T
Grans for oral liq 50 mg per ml	3.68	100 ml	•	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable Inj 250 mg vial	17.56	10		Flucioxin
Inj 500 mg vial		10		Flucioxin
Inj 1 g vial — Up to 5 inj available on a PSO		5	_	Flucil
,		•	-	

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer	
PHENOXYMETHYLPENICILLIN (PENICILLIN V) Cap 250 mg — Up to 30 cap available on a PSO Cap 500 mg a) Up to 20 cap available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP		50 50	_	ilicaine VK ilicaine VK	
Grans for oral liq 125 mg per 5 ml	3.40	100 ml	✓ <u>A</u>	<u>IFT</u>	
Grans for oral liq 250 mg per 5 ml	4.24 1	100 ml	✓ <u>A</u>	<u>.FT</u>	

Tetracyclines

DO	XYCYCLINE			
*	Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	✓ Doxine
MIN	NOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	19.32	100	
		(52.04)		Minomycin

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

TETRACYCLINE - Special Authority see SA1332 below - Retail pharmacy

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 61

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO 2.42 Tab 500 mg - Up to 5 tab available on a PSO 3.40 Tab 750 mg 5.95	28 28 28	✓ Cipflox ✓ Cipflox ✓ Cipflox
CLINDAMYCIN		
Cap hydrochloride 150 mg5.30	24	Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule39.00	10	Dalacin C

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Generic
	Ψ	rei		Manufacturer
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S	ubsidy by endorseme	nt		
Only if prescribed for dialysis or cystic fibrosis patient and the	prescription is endor	sed a	accordingly	y.
Inj 150 mg	65.00	1	1	Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement	95.00	5	/	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient of				
endorsed accordingly.	n complicated unitary	ιιασι	IIIICCIIOII	and the prescription is
3,	01.00	5	./	Wockhardt S29
Inj 10 mg per ml, 2 ml ampoule – Subsidy by endorsement				
	182.00	10		Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.	or complicated urinary	tract	infection	and the prescription is
Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement	18.38	10	1	Pfizer
ing to mg por mi, a m ampould — case by shacked memoria	87.50	50		Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient of	or complicated urinary			
endorsed accordingly.	or complicated unitary	liuot	moodon	and the presemption is
MOXIFLOXACIN - Special Authority see SA1740 below - Retail	pharmacy			
No patient co-payment payable				
Tab 400 mg	42.00	5	1	Avelox
⇒SA1740 Special Authority for Subsidy				

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications: or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 on the next page - Retail pharmacy

Cap 250 mg......126.00 16 ✓ Humatin S29

INFECTIONS - AGENTS FOR SYSTEMIC USE	Ē			
	Subsidy (Manufacturer's Price \$) Sub	Fully sidised	Brand or Generic Manufacturer
■ SA1689 Special Authority for Subsidy Initial application only from an infectious disease specialist, clinic month for applications meeting the following criteria: Either:	cal microbiologist o	r gastroent	erologis	t. Approvals valid for 1
Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage.				
Renewal only from an infectious disease specialist, clinical microb applications meeting the following criteria: Either: 1 Patient has confirmed cryptosporidium infection; or	oiologist or gastroer	nterologist.	Approv	als valid for 1 month for
2 For the eradication of Entamoeba histolyica carriage.				
PYRIMETHAMINE – Special Authority see SA1328 below – Reta Tab 25 mg		30	✓ D	araprim S29
■ SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following:	without further ren	ewal unles	s notifie	d for applications meeting
For the treatment of toxoplasmosis in patients with HIV for For pregnant patients for the term of the pregnancy; or For infants with congenital toxoplasmosis until 12 months of the pregnancy.	•	ns; or		
SODIUM FUSIDATE [FUSIDIC ACID]				
Tab 250 mg		36	√ F	ucidin
SULFADIAZINE SODIUM – Special Authority see SA1331 below Tab 500 mg	, ,	56	✓ W	ockhardt S29
■ SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of	a period of 3 montl		s notifie	d for applications meeting
TOBRAMYCIN Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement	18.50	5		obramycin Mylan iatris
Only if prescribed for dialysis or cystic fibrosis patient and	the prescription is	endorsed		
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by endorsement	395.00	56 dose	✓ <u>T</u>	obramycin BNM
a) Wastage claimableb) Only if prescribed for a cystic fibrosis patient and the p	orescription is endo	rsed acco	dingly.	
TRIMETHOPRIM			. –	
* Tab 300 mg - Up to 30 tab available on a PSO		50	✓ Ī	<u>MP</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXA * Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - U				
to 20 tob available on a PSO	64 8U	500	∠ ⊤	rieul

to 30 tab available on a PSO......64.80

available on a PSO......2.97

Oral liq 8 mg sulphamethoxazole 40 mg per ml - Up to 200 ml

✓ Trisul

✓ Deprim

500

100 ml

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

VANCOMYCIN - Subsidy by endorsement

Only if prescribed for a dialysis or cystic fibrosis patient or for prophylaxis of endocarditis or for treatment of Clostridium difficile following metronidazole failure and the prescription is endorsed accordingly.

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 62
- b) For topical antifungals refer to GENITO URINARY, page 75

FLUCONAZOLE

Cap 50 mg	2.75	28	Dizole
			Mylan
Cap 150 mg	0.65	1	Mylan
Cap 200 mg	12.89	28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority			
see SA1359 below - Retail pharmacy	129.02	35 ml	Diflucan
Wastage claimable			

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg	.4.27	15	✓ Itrazole
Oral lig 10 mg per ml - Special Authority see SA1322 below -			
Retail pharmacy1	41.80	150 ml OP	✓ Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

	Subsidy (Manufacturer's Pric \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
KETOCONAZOLE				
Tab 200 mg - PCT	CBS	30	✓	Burel S29
			1	Link Healthcare S29
			✓	Nizoral S29
		100	✓	Strides Shasun S29
			1	Taro S29
(Link Healthcare S29 Tab 200 mg to be delisted 1 July 2023)				
(Nizoral S29 Tab 200 mg to be delisted 1 July 2023)				
(Strides Shasun S29 Tab 200 mg to be delisted 1 July 2023)				
NYSTATIN				
Tab 500,000 u	14 16	50		
745 000,000 u	(17.09)	00		Nilstat
Cap 500,000 u	` ,	50		
•	(15.47)			Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Reta	il pharmacy			
Tab modified-release 100 mg		24	1	Posaconazole Juno
•	869.86		1	Noxafil
Posaconazole Juno to be Principal Supply on 1 April 202				
Oral liq 40 mg per ml		105 ml (Devatis
	761.13		•	Noxafil
Devatis to be Principal Supply on 1 May 2023				

Devatis to be Principal Supply on 1 May 2023

(Noxafil Tab modified-release 100 mg to be delisted 1 April 2023) (Noxafil Oral lig 40 mg per ml to be delisted 1 May 2023)

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERRINAFINE

* Tab 250 mg	8.15	84	✓ <u>Deolate</u>	
VORICONAZOLE - Special Authority see SA1273 on the ne	xt page - Retail phar	macy		
Tab 50 mg	91.00	56	✓ Vttack	
Tab 200 mg	350.00	56	✓ Vttack	
Powder for oral suspension 40 mg per ml - Wastage				
claimable	1,523.22	70 ml	✓ Vfend	

5	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE - Special Authority see SA1684 belo	w – Retail pharmacy			
Tab 15 mg	400.00	100	✓ Sanofi	
· ·			Primaquine S29	

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.16	10	✓ Arrow-Ornidazole

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
Antituberculotics and Antileprotics				
Note: There is no co-payment charge for all pharmaceuticals lis immigration status.	ted in the Antitubercul	lotics a	nd Antilepi	rotics group regardless of
CLOFAZIMINE - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda dermatologist.	tion of, an infectious d	lisease	physician,	clinical microbiologist or
* Cap 50 mg	442.00	100	✓ L	amprene S29
CYCLOSERINE - Retail pharmacy-Specialist				•
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician. 				-
Cap 250 mg	344.00	60	√ (Cyclorin S29
DAPSONE - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda dermatologist				
Tab 25 mg		100		Dapsone
Tab 100 mg		100	V L	Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Special	st			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician 	tion of, an infectious d	lisease	physician,	clinical microbiologist or
Tab 100 mg	85.73	100	✓ E	MB Fatol S29
Tab 400 mg	49.34	56	✓ N	Myambutol S29
ISONIAZID - Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda microbiologist, dermatologist or public health physician 	tion of, an internal me	dicine ¡	ohysician,	paediatrician, clinical
* Tab 100 mg	23.00	100	√ F	PSM
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist			_	
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda microbiologist, dermatologist or public health physician	tion of, an internal me	dicine ¡	ohysician,	paediatrician, clinical
* Tab 100 mg with rifampicin 150 mg	89.82	100	√ F	Rifinah
* Tab 150 mg with rifampicin 300 mg		100	✓ <u>F</u>	Rifinah
PARA-AMINO SALICYLIC ACID - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician	tion of, an infectious d	lisease	specialist,	clinical microbiologist or
Grans for oral liq 4 g sachet	280.00	30	√ F	Paser \$29
PROTIONAMIDE – Retail pharmacy-Specialist		00		4001
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician	tion of, an infectious d	lisease	specialist,	clinical microbiologist or
Tab 250 mg	305.00	100	✓ F	Peteha S29
✓ fully subsidised	\$29 Unapprove	d modici	ing gunnlind	under Coation 20

25

35

✓ Lovir

✓ Lovir

✓ Lovir

	INFECTIONS -	AGENTS	FOR SYSTEMIC U	ISE
	Subsidy (Manufacturer's Pric	ce) Sub	Fully Brand or sidised Generic Manufacturer	
PYRAZINAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommen- respiratory physician	dation of, an infectious	s disease ph	ysician, clinical microbic	ologist or
* Tab 500 mg	64.95	100	✓ AFT-Pyrazinami	de
RIFABUTIN - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommended gastroenterologist		·		sician or
* Cap 150 mg	353.71	30	✓ Mycobutin	
RIFAMPICIN – Subsidy by endorsement				
 a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus infect antimicrobial based on susceptibilities and the prescrip Retail pharmacy - Specialist. Specialist must be an in paediatrician, or public health physician. 	otion is endorsed acco	ordingly; can	be waived by endorsen	nent -
* Cap 150 mg		100	✓ <u>Rifadin</u>	
* Cap 300 mg		100	Rifadin	
* Oral liq 100 mg per 5 ml	12.60	60 ml	✓ <u>Rifadin</u>	
Antivirals				
For eye preparations refer to Eye Preparations, Anti-Infective	Preparations, page 24	19		
Hepatitis B Treatment				
* Tab 0.5 mg	52.00	30	✓ Entecavir Mylan✓ Entecavir Sando	
LAMIVUDINE - Special Authority see SA1685 below - Retail	pharmacy			
Tab 100 mg		28	✓ <u>Zetlam</u>	
Oral liq 5 mg per ml	270.00	240 ml OP	✓ Zeffix	
⇒SA1685 Special Authority for Subsidy				
Initial application only from a relevant specialist or medical p Approvals valid for 1 year where used for the treatment or pre			n of a relevant specialist	
Renewal from any relevant practitioner. Approvals valid for 2	years where used for	the treatme	nt or prevention of hepa	titis B.
TENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for the	e treatment of HIV is in	ncluded in th	ne count of up to 4 subsi	dised
antiretrovirals for the purposes of Special Authority SA213				
* Tab 245 mg (300 mg as a maleate)		30	✓ <u>Tenofovir Disop</u> <u>Mylan</u>	<u>roxil</u>
Herpesvirus Treatments				

▲Three months supply ma	be dispensed at one time if endorsed	"certified exemption" by the prescriber or pharmacist.

Lovir to be Principal Supply on 1 March 2023

Lovir to be Principal Supply on 1 April 2023

Lovir to be Principal Supply on 1 April 2023

* Tab dispersible 800 mg6.46

ACICLOVIR

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
VALACICLOVIR Tab 500 mg Tab 1,000 mg		30 30		aclovir aclovir	
VALGANCICLOVIR - Special Authority see SA1993 below - Ret Tab 450 mg	, ,	60	✓ <u>v</u>	alganciclovir Mylan	

⇒SA1993 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

continued...

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – Subsidy by endorsement; can be waived by Special Authority see SA2138 on the next page

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

* Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a

> ✓ Tenofovir Disoproxil Emtricitabine Viatr

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully Brand or idised Generic ✓ Manufacturer
Non-nucleosides Reverse Transcriptase Inhibit	ors		
EFAVIRENZ – Special Authority see SA2139 on page 104 – Re Tab 200 mg Tab 600 mg	190.15	90 30	✓ Stocrin ✓ Stocrin
ETRAVIRINE – Special Authority see SA2139 on page 104 – R. Tab 200 mg		60	✓ Intelence
NEVIRAPINE – Special Authority see SA2139 on page 104 – R Tab 200 mg		60	✓ <u>Nevirapine</u> Alphapharm ✓ Nevirapine Viatris
Oral suspension 10 mg per ml	203.55	240 ml OP	✓ Viramune Suspension
Nucleosides Reverse Transcriptase Inhibitors			
ABACAVIR SULPHATE – Special Authority see SA2139 on pag Tab 300 mg Oral liq 20 mg per ml	180.00	60 240 ml OP	✓ Ziagen ✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE — Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	29.50	30	Abacavir/ Lamivudine Viatris
Abacavir/Lamivudine Viatris to be Principal Supply on 1 (Kivexa Tab 600 mg with lamivudine 300 mg to be delisted 1 Ma	•		✓ Kivexa
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOP pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil canti-retroviral Special Authority	ounts as three an	•	
Tab 600 mg with emtricitabine 200 mg and tenofovir disopro 245 mg (300 mg as a maleate)		30	✓ Mylan✓ Viatris
EMTRICITABINE – Special Authority see SA2139 on page 104 Cap 200 mg	307.20	30	✓ Emtriva
LAMIVUDINE - Special Authority see SA2139 on page 104 - R Tab 150 mg		60	✓ <u>Lamivudine</u> <u>Alphapharm</u> ✓ Lamivudine Viatris
Oral liq 10 mg per ml		240 ml OP	✓ 3TC
ZIDOVUDINE [AZT] – Special Authority see SA2139 on page 10 Cap 100 mg Oral liq 10 mg per ml	152.25	100 200 ml OP	✓ Retrovir ✓ Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE — Special Authority se Note: zidovudine [AZT] with lamivudine (combination tablet the anti-retroviral Special Authority.			
Tab 300 mg with lamivudine 150 mg	33.00	60	✓ Alphapharm

Fully

Subsidised

Brand or

Generic

Subsidy

(Manufacturer's Price)

	(Manufacturer's Price)	Per	✓ Manufacturer
Protease Inhibitors			
ATAZANAVIR SULPHATE – Special Authority see SA2139 on p Cap 150 mg	•	armacy 60	✓ Atazanavir Mylan✓ Teva
Atazanavir Mylan to be Principal Supply on 1 May 2023 Cap 200 mg Atazanavir Mylan to be Principal Supply on 1 May 2023 (Teva Cap 150 mg to be delisted 1 May 2023) (Teva Cap 200 mg to be delisted 1 May 2023)	188.91	60	✓ Atazanavir Mylan✓ Teva
DARUNAVIR – Special Authority see SA2139 on page 104 – Re Tab 400 mg Tab 600 mg	132.00	60 60	✓ Darunavir Mylan ✓ Darunavir Mylan ✓ Darunavir Viatris
LOPINAVIR WITH RITONAVIR – Special Authority see SA2139 Tab 100 mg with ritonavir 25 mg	150.00	60	✓ <u>Lopinavir/Ritonavir</u> <u>Mylan</u>
Tab 200 mg with ritonavir 50 mg Oral lig 80 mg with ritonavir 20 mg per ml		120 00 ml OP	✓ <u>Lopinavir/Ritonavir</u> <u>Mylan</u> ✓ Kaletra
RITONAVIR – Special Authority see SA2139 on page 104 – Ret Tab 100 mg	ail pharmacy	30	✓ Norvir
Strand Transfer Inhibitors			
DOLUTEGRAVIR – Special Authority see SA2139 on page 104 Tab 50 mg	1,090.00 on page 104 – Retail	30 pharmacy 60	✓ Tivicay ✓ Isentress
Tab 600 mg	,	60	✓ Isentress HD
PEGYLATED INTERFERON ALFA-2A — Special Authority see S Note: Pharmac will consider funding ribavirin for the small g Special Authority criteria. Please contact the Hepatitis C Co	roup of patients who	have a clinic	cal need for ribavirin and meet

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or

Inj 180 mcg prefilled syringe......500.00

- 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general

continued...

Pegasys

((Subsidy Manufacturer's Price)	Sub	Fully	Brand or Generic
	\$	Per	1	Manufacturer

continued...

physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has chronic hepatitis C. genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Fither:
 - 5.1 HBeAg positive: or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis): and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

1 Patient has a cutaneous T cell lymphoma*: or

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
 - 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate: or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	19.95	100	✓ <u>Hiprex</u>
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ <u>Nifuran</u>
* Tab 100 mg	37.50	100	✓ <u>Nifuran</u>
* Cap modified-release 100 mg - Up to 15 cap available on a			
PSO	86.40	100	✓ <u>Macrobid</u>
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	245.00	100	✓ Arrow-Norfloxacin
Only if prescribed for a patient with an uncomplicated urin	ary tract infection		

٥r with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)) Per	Subsidised	Generic Manufacturer
	φ	r ei		Manufacturer
Anticholinesterases				
OSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	✓	Max Health
RIDOSTIGMINE BROMIDE				
Tab 60 mg	45.79	100	1	Mestinon
Ion-Steroidal Anti-Inflammatory Drugs				
CLOFENAC SODIUM				
Tab EC 25 mg	1 00	50	1	Diclofenac Sandoz
Tab 50 mg dispersible		20		Voltaren D
		50		Diclofenac Sandoz
Tab EC 50 mg Tab long-acting 75 mg				
		100		Voltaren SR
Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a		5		Voltaren
Suppos 12.5 mg		10		Voltaren
Suppos 25 mg		10		Voltaren
Suppos 50 mg – Up to 10 supp available on a PSO		10		Voltaren
Suppos 100 mg	7.00	10	•	Voltaren
JPROFEN				
Tab 200 mg	21.40	1,000	/	Relieve
Tab long-acting 800 mg	3.05	30	✓	Brufen SR
Oral liq 20 mg per ml		200 m	ı 🗸	Ethics
	11.29		✓	Fenpaed 100 mg per
				5 ml
TOPROFEN	10.07	00	,	O!! OD
Cap long-acting 200 mg	12.07	28	•	Oruvail SR
FENAMIC ACID				
Cap 250 mg	1.25	50		
	(10.82)			Ponstan
	0.50	20		
	(7.50)			Ponstan
PROXEN				
Tab 250 mg	32 69	500	1	Noflam 250
Tab 500 mg		250		Noflam 500
Tab long-acting 750 mg		28		Naprosyn SR 750
Tab long-acting 1 g		28		Naprosyn SR 1000
	0.02	20	•	ivapiusyii on 1000
NOXICAM			_	
Tab 20 mg		100		Tilcotil
Inj 20 mg vial	9.95	1	•	AFT
ISAIDs Other				
ELECOXIB				
Cap 100 mg	3.45	60		Celebrex
				Celecoxib Pfizer
Cap 200 mg	3.20	30	/	Celebrex
				Celecoxib Pfizer

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

Topical Products for Joint and Muscular Pain

CAPSAICIN

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary)*, and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroquine. Note: Indication marked with a * is an unapproved indication.

* Tab 200 mg	8.78	100	✓ Plaquenil
LEFLUNOMIDE			
Tab 10 mg	6.00	30	✓ Arava
Tab 20 mg	6.00	30	✓ Arava
PENICILLAMINE			
Tab 125 mg	67.23	100	D-Penamine
Tab 250 mg	110.12	100	D-Penamine

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODIUM			
* Tab 70 mg	2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL			
* Tab 70 mg with colecalciferol 5,600 iu	1.51	4	✓ Fosamax Plus

Other Treatments

DENOSUMAB - Special Authority see SA1777 below - Retail	pharmacy		
Inj 60 mg prefilled syringe	326.00	1	✓ Prolia

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither:
 - 2.1 The patient is female and postmenopausal; or

Subsidy (Manufacturer's Price)	F Subsidis	ully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	Pamisol
Inj 6 mg per ml, 10 ml vial	88.11	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	94.34	1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see	e SA1779 below – Retail p	harmacy	
* Tab 60 mg	53.76	28	Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

(Ma	Subsidy nufacturer's Price)	F Subsid	ully	Brand or Generic
	\$	Per	1	Manufacturer

continued...

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM			
Tab 35 mg	2.50	4	✓ Risedronate Sandoz
TERIPARATIDE - Special Authority see SA1139 on the next page	- Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml	490.00	1	✓ Forteo

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, bag – Special Authority see SA2110 on page 116 – Retail pharmacy22.	.53 100	-	Zoledronic Acid Viatris Zoledronic-US \$29
Inj 0.05 mg per ml, 100 ml, vial — Special Authority see SA2110 below — Retail pharmacy	.00 100) ml OP 💌	Aclasta

⇒SA2110 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain: or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or

Subsidy (Manufacturer's Price)	Subs	Fully idised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
 - 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or

- 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (spinal cord injury*) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has experienced an acute traumatic spinal cord injury in the last six months; and
- 2 Patient is being managed by a specialist spinal acute care and rehabilitation unit; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Note: Indications marked with * are unapproved indications.

Renewal — (spinal cord injury*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period; and
- 2 The patient has not received more than two doses of zoledronic acid for this indication.

The patient must not have had more than 1 prior approval.

Notes: No further renewals will be subsidised. A maximum of 2 vials of zoledronic acid treatment for spinal cord injury will be subsidised. Indications marked with * are unapproved indications.

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

⇒SA2110 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain: or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or
 (Manufacturer's Price) \$	Per 🗸	Generic Manufacturer

continued...

- 2.4 Asymptomatic disease, but risk of complications; or
- 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note): or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (spinal cord injury*) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has experienced an acute traumatic spinal cord injury in the last six months; and
- 2 Patient is being managed by a specialist spinal acute care and rehabilitation unit; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Note: Indications marked with * are unapproved indications.

Renewal — (spinal cord injury*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period; and
- 2 The patient has not received more than two doses of zoledronic acid for this indication.

The patient must not have had more than 1 prior approval.

Notes: No further renewals will be subsidised. A maximum of 2 vials of zoledronic acid treatment for spinal cord injury will be subsidised. Indications marked with * are unapproved indications.

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Brand or

Fully

	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
Hyperuricaemia and Antigout				
ALLOPURINOL				
* Tab 100 mg		500	✓ [P-Allopurinol
* Tab 300 mg	28.57	500	✓ [P-Allopurinol
BENZBROMARONE - Special Authority see SA1963 below - I	Retail pharmacy			
Tab 50 mg	22.50	100	✓ N	larcaricin mite S29
Tab 100 mg	13.50	30	✓ [esuric \$29
			√ (Irinorm \$29
	45.00	100	√ E	Senzbromaron AL
				100 \$29

Subsidy

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg6.00 100	✓ Colgout
FEBUXOSTAT - Special Authority see SA2054 below - Retail pharmacy	
Tab 80 mg20.00 28	✓ Febuxostat
	multichem
Tab 120 mg20.00 28	✓ <u>Febuxostat</u> multichem

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose: or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

PROBENECID

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer	
Muscle Belayants					

masore relaxants			
BACLOFEN	<u></u>		
* Tab 10 mg	4.20	100	✓ Pacifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	✓ Lioresal Intrathecal
Subsidised only for use in a programmable pump in patient caused intolerable side effects and the prescription is endo	s where oral antis		ents have been ineffective or have
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement	306.82	5	✓ <u>Medsurge</u>
Subsidised only for use in a programmable pump in patient caused intolerable side effects and the prescription is endo			ents have been ineffective or have
DANTROLENE			
Cap 25 mg	112.13	100	✓ Dantrium
			✓ Dantrium S29 S29
Cap 50 mg	77.00	100	✓ Dantrium
ORPHENADRINE CITRATE			
Tab 100 mg	20.76	100	✓ <u>Norflex</u>

Subsidy (Manufacturer's Price) S

Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists a	and Related Agents
----------------------------	--------------------

AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg	38.24	60	 Symmetrel
	63.73	100	Symmetrel
APOMORPHINE HYDROCHLORIDE			
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	121.84	5	✓ Movapo
ENTACAPONE			
▲ Tab 200 mg	18.04	100	✓ Comtan
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg	13.75	100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg		100	Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg	22.85	100	Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
* Tab 100 mg with carbidopa 25 mg	21.11	100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg		100	✓ Sinemet CR
* Tab 250 mg with carbidopa 25 mg	38.39	100	✓ Sinemet
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	5.51	100	✓ Ramipex
▲ Tab 1 mg	18.66	100	✓ Ramipex
RASAGILINE			
* Tab 1 mg	53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	4.05	84	✓ Ropin
▲ Tab 1 mg	4.95	84	✓ Ropin
▲ Tab 2 mg	6.48	84	✓ Ropin
▲ Tab 5 mg	14.50	84	✓ Ropin

SELEGILINE HYDROCHLORIDE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of selegiline hydrochloride.

*	Tab 5 mg48.00	100	✓ Eldepryl S29
TO	LCAPONE		
	Tab 100 mg152.38	100	✓ Tasmar

Anticholinergics

BENZATROPINE MESYLATE			
Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
a) Up to 10 inj available on a PSO			
b) Only on a PSO			
PROCYCLIDINE HYDROCHLORIDE			

Tab 5 mg7.40 100

✓ Kemadrin



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE - Special Authority see SA1403 below - Retail pharmacy

Wastage claimable

⇒SA1403 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Anaesthetics

Local

LIDOCAINE [LIGNOCAINE]

Gel 2%, tube − Subsidy by endorsement14.50 30 ml ✓ Xylocaine 2% Jelly

- a) Up to 150 ml available on a PSO
- b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Gel 2%, 11 ml urethral syringe − Subsidy by endorsement...............59.50 10 ✓ Instillagel Lido

- a) Up to 5 each available on a PSO
- Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e)	Subsidised	I Generic
	\$	Per	1	Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 m	/	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO		25	1	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	9.00 [°]	25	1	Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20	5	1	Lidocaine-Claris
	6.85		1	Lidocaine-Baxter
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	7.15	5	1	Lidocaine-Baxter
(Lidocaine-Claris Inj 1%, 20 ml vial to be delisted 1 June 2023)				
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –				
Subsidy by endorsement	103 32	10	1	Pfizer
	100.02	10	•	1 11201
a) Up to 5 each available on a PSO				

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDUCAINE [LIGNOCAINE] - Special Authority see SA0906 and	ve – Retali pnar	macy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Author	ority see SA0906	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓ EMLA

Analgesics

ASPIRIN

Non-opioid Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

*	Tab dispersible 300 mg - Up to 30 tab available on a PSO4.50	100	Ethics Aspirin
CA	PSAICIN – Subsidy by endorsement		
	Subsidised only if prescribed for post-herpetic neuralgia or diabetic peripheral neuralgia	uropathy	and the prescription is endorsed
	accordingly.		

Subsidised only if prescribed for post-fierpetic fiedralgia of	ulabelic peripriera	ii iicuiopaiiiy a	ila tile prescription is endorse
accordingly.			
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.14	57 g OP	✓ Rugby Capsaicin
		•	Topical
			Cream S29

NEFOPAM HYDROCHLORIDE			
Tab 30 mg	23.40	90	Acupan

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturaria Brica	.\ 0	Fully Brand	
	(Manufacturer's Price \$	Per Subs	idised Gener Manuf	acturer
RACETAMOL				
Tab 500 mg - blister pack	19.75	1,000	✓ Pacimol	
 a) Maximum of 300 tab per prescription; can be wai b) Up to 30 tab available on a PSO c) 	ved by endorsement			
Subsidy by endorsement for higher quantities	es is available for patier	nts with long	term condition	ns who require
regular daily dosing for one month or greate				
annotate the prescription as endorsed wher	e dispensing history su	pports a lon	g-term condition	n.
Maximum of 100 tab per dispensing for non				
(for non-endorsed patients), then dispense	in repeat dispensings n	ot exceeding	g 100 tab per d	lispensing.
Tab 500 mg - bottle pack - Maximum of 300 tab per			.	
prescription; can be waived by endorsement	17.92	1,000	✓ <u>Noumed</u> Parace	<u>[</u> etamol
1) Subsidy by endorsement for higher quantities is	s available for natients	with long ter		
daily dosing for one month or greater, and the				
prescription as endorsed where dispensing his				,
2) Maximum of 100 tab per dispensing for non-en				than 100 tabs (f
non-endorsed patients), then dispense in repea				
0.111.42			4.5	
Oral liq 120 mg per 5 ml	3.98	200 ml	✓ Paraceta	
	5 45	4 000 1	(Ethics	
		1,000 ml 200 ml OP	✓ Paracare✓ Avallon	9
a) Manimum of COO and non-manimizing combined		200 mi OP	Availon	
a) Maximum of 600 ml per prescription; can be waivb) Up to 200 ml available on a PSO	ed by endorsement			
c) Not in combination				
d)				
Maximum of 200 ml per dispensing for non-	endorsed patients. If o	uantities pre	escribed excee	d 200 ml (for
non-endorsed patients), then dispense in re				,
2) Subsidy by endorsement for higher quantitie				
regular daily dosing for one month or greate				
Pharmacists may annotate the prescription	as endorsed where dis	pensing hist	ory supports a	long-term
condition.				
Oral liq 250 mg per 5 ml		200 ml	✓ Pamol	
	6.25	1,000 ml	✓ Paracare Streng	
a) Maximum of 600 ml per prescription; can be waiv	ed by endorsement		·	,
b) Up to 200 ml available on a PSO				
b) Up to 200 ml available on a PSO c) Not in combination d)				
 b) Up to 200 ml available on a PSO c) Not in combination d) 1) Maximum of 200 ml per dispensing for non- 				
b) Up to 200 ml available on a PSO c) Not in combination d) 1) Maximum of 200 ml per dispensing for nonnon-endorsed patients), then dispense in re	peat dispensing not ex	ceeding 200	ml per dispens	sing.
 b) Up to 200 ml available on a PSO c) Not in combination d) 1) Maximum of 200 ml per dispensing for non-non-endorsed patients), then dispense in re 2) Subsidy by endorsement for higher quantitie 	epeat dispensing not ex es is available for patier	ceeding 200 nts with long	ml per dispens term condition	sing. ns who require
 b) Up to 200 ml available on a PSO c) Not in combination d) 1) Maximum of 200 ml per dispensing for non-non-endorsed patients), then dispense in re 2) Subsidy by endorsement for higher quantitic regular daily dosing for one month or greate 	epeat dispensing not exe es is available for patier er and the prescription is	ceeding 200 nts with long s endorsed	ml per dispens term condition or annotated ac	sing. ns who require ccordingly.
 b) Up to 200 ml available on a PSO c) Not in combination d) 1) Maximum of 200 ml per dispensing for non-non-endorsed patients), then dispense in re 2) Subsidy by endorsement for higher quantitic regular daily dosing for one month or greate Pharmacists may annotate the prescription 	epeat dispensing not exe es is available for patier er and the prescription is	ceeding 200 nts with long s endorsed	ml per dispens term condition or annotated ac	sing. ns who require ccordingly.
 b) Up to 200 ml available on a PSO c) Not in combination d) 1) Maximum of 200 ml per dispensing for non-non-endorsed patients), then dispense in re 2) Subsidy by endorsement for higher quantitic regular daily dosing for one month or greate Pharmacists may annotate the prescription condition. 	epeat dispensing not exe es is available for patier er and the prescription is	ceeding 200 nts with long s endorsed	ml per dispens term condition or annotated ac	sing. ns who require ccordingly.
 b) Up to 200 ml available on a PSO c) Not in combination d) 1) Maximum of 200 ml per dispensing for non-non-endorsed patients), then dispense in re 2) Subsidy by endorsement for higher quantitic regular daily dosing for one month or greate Pharmacists may annotate the prescription 	epeat dispensing not ex es is available for patier er and the prescription i as endorsed where dis	ceeding 200 nts with long s endorsed	ml per dispens term condition or annotated ac	sing. ns who require ccordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Per	Subsidised	Generic
₭ Suppos 500 mg	12.40	50	1	Gacet
Paracare Oral liq 120 mg per 5 ml to be delisted 1 June 2023 Paracare Double Strength Oral liq 250 mg per 5 ml to be del				
Opioid Analgesics				
ODEINE PHOSPHATE - Safety medicine; prescriber may	determine dispensing fre	quen	су	
Tab 15 mg	, ,	100	•	Noumed
Č	6.25		✓	PSM
Noumed to be Principal Supply on 1 May 2023				
Tab 30 mg	6.98	100	✓	Aspen
			✓	Noumed
	7.45		✓	PSM
Noumed to be Principal Supply on 1 April 2023				
Tab 60 mg		100		Noumed
	14.25		/	PSM
Noumed to be Principal Supply on 1 April 2023				
PSM Tab 15 mg to be delisted 1 May 2023)				
PSM Tab 30 mg to be delisted 1 April 2023)				
PSM Tab 60 mg to be delisted 1 April 2023)				
DIHYDROCODEINE TARTRATE				
Tab long-acting 60 mg	8.60	60	•	DHC Continus
ENTANYL				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing	g frequency			
Inj 50 mcg per ml, 2 ml ampoule		10		Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10		Boucher and Muir
Patch 12.5 mcg per hour		5		Fentanyl Sandoz
Patch 25 mcg per hour		5		Fentanyl Sandoz
Patch 50 mcg per hour		5		Fentanyl Sandoz
Patch 75 mcg per hour		5		Fentanyl Sandoz
Patch 100 mcg per hour	18.59	5	•	Fentanyl Sandoz
ETHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing				
d) Extemporaneously compounded methadone will only	be reimbursed at the rat	e of th	ne cheape	st form available
(methadone powder, not methadone tablets).				
e) For methadone hydrochloride oral liquid refer Standar				
Tab 5 mg		10	_	Methadone BNM
Oral liq 2 mg per ml		200 m		Biodone
Oral liq 5 mg per ml		200 m		Biodone Forte
Oral lig 10 mg par ml	7 50	200 ~	al .//	Diadona Eytra Ear

200 ml

10

✓ Biodone Extra Forte

✓ AFT

	Subsidy		Fully Brand or
(Manufacturer's P	rice) Sub Per	sidised Generic Manufacturer
ORPHINE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing frequency	lliency		
Oral lig 1 mg per ml		200 ml	✓ RA-Morph
Oral lig 2 mg per ml		200 ml	✓ RA-Morph
Oral lig 5 mg per ml		200 ml	✓ Ordine S29
Oral liq 5 mg per mil	13.44	200 1111	✓ RA-Morph
Ovel lie 10 men men mi	07.74	0001	•
Oral liq 10 mg per ml	27.74	200 ml	✓ Ordine \$29
			✓ RA-Morph
ORPHINE SULPHATE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing freq			
Tab immediate-release 10 mg		10	✓ <u>Sevredol</u>
Tab immediate-release 20 mg		10	✓ <u>Sevredol</u>
Cap long-acting 10 mg	3.00	10	✓ m-Eslon
m-Eslon to be Principal Supply on 1 April 2023			
Cap long-acting 30 mg	4.30	10	✓ m-Eslon
m-Eslon to be Principal Supply on 1 April 2023			
Cap long-acting 60 mg	9.00	10	✓ m-Eslon
m-Eslon to be Principal Supply on 1 April 2023			
Cap long-acting 100 mg	10.50	10	✓ m-Eslon
m-Eslon to be Principal Supply on 1 April 2023			
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSC		5	Medsurge
	6.99		DBL Morphine
			Sulphate
Medsurge to be Principal Supply on 1 March 2023			
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	SO 4.68	5	Medsurge
	5.61		DBL Morphine
			Sulphate
Medsurge to be Principal Supply on 1 March 2023			
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	SO5.53	5	✓ Medsurge
	7.08		✓ DBL Morphine
			Sulphate
Medsurge to be Principal Supply on 1 March 2023			
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	SO6.28	5	✓ Medsurge
	7.28		✓ DBL Morphine
			Sulphate
Medsurge to be Principal Supply on 1 March 2023			-
DBL Morphine Sulphate Inj 5 mg per ml, 1 ml ampoule to be delisi	ted 1 March 202	23)	
DBL Morphine Sulphate Inj 10 mg per ml, 1 ml ampoule to be deli			
DBL Morphine Sulphate Inj 15 mg per ml, 1 ml ampoule to be deli			
OBL Morphine Sulphate Inj 30 mg per ml, 1 ml ampoule to be deli			

(DBL Morphine Sulphate Inj 30 mg per ml, 1 ml ampoule to be delisted 1 March 2023)

	Subsidy		Fully Brand or	
	(Manufacturer's Price)		Subsidised Generic Manufacturer	
	\$	Per	✓ Manufacturer	
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				
Tab controlled-release 5 mg		20	✓ Oxycodone Sandoz	
Tab controlled-release 10 mg		20	✓ Oxycodone Sandoz	
Tab controlled-release 20 mg		20	✓ Oxycodone Sandoz	
Tab controlled-release 40 mg		20	✓ Oxycodone Sandoz	
Tab controlled-release 80 mg		20	✓ Oxycodone Sandoz	
Cap immediate-release 5 mg		20	OxyNorm OxyNorm	
Cap immediate-release 10 mg		20	OxyNorm OxyNorm	
Cap immediate-release 20 mg		20 250 m	OxyNorm OxyNorm	
Oral liq 5 mg per 5 ml			<u></u>	
Inj 10 mg per ml, 1 ml ampoule		5	✓ <u>Hameln</u>	
Inj 10 mg per ml, 2 ml ampoule		5 5	✓ <u>Hameln</u> ✓ Hameln	
Inj 50 mg per ml, 1 ml ampoule		-		
PARACETAMOL WITH CODEINE - Safety medicine; prescriber				
* Tab paracetamol 500 mg with codeine phosphate 8 mg	27.50	1,000		
			Codeine (Relieve)	
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 Safety medicine; prescriber may determine dispensing free 	quency			
Tab 50 mg	4.70	10	✓ PSM	
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	SO29.88	5	DBL Pethidine	
			Hydrochloride	
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a P	SO30.72	5	DBL Pethidine	
			Hydrochloride	
TRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg	1.52	20	✓ Tramal SR 100	
Tab sustained-release 150 mg		20	✓ Tramal SR 150	
Tab sustained-release 200 mg		20	✓ Tramal SR 200	
Cap 50 mg		100	✓ Arrow-Tramadol	
Antidepressants				
·				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine d	ispensing frequency			
Tab 10 mg		100	✓ Arrow-Amitriptyline	ı
Tab 25 mg		100	✓ Arrow-Amitriptyline	
Tab 50 mg		100	✓ Arrow-Amitriptyline	
CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescri		ienan		
Tab 10 mg	•	30	✓ Clomipramine Teva	
Tab 25 mg		30	✓ Clomipramine Teva	
- 4.0 ±0 mg	11.00	00	- Otompramme reva	

	Subsidy		ully	Brand or
(I	Manufacturer's Price)	Subsid Per	ised •	Generic Manufacturer
DOOLII EDINI DOTLIIEDINI LIVDEQQUI ORIDE Q. b. c. b.	<u> </u>	1 51	_	manuacture!
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endo				
Safety medicine; prescriber may determine dispensing frequency by Subsidy by endorsement – Subsidised for patients who wer 2019 and the prescription is endorsed accordingly. Pharma exists a record of prior dispensing of dosulepin [dothiepin] has been supported by the prescription of the prior dispensing of dosulepin [dothiepin] has been supported by the prescription of the provided by the p	e taking dosulepin acists may annotate lydrochloride.	the prescri	ption	as endorsed where there
Tab 75 mg	3.85	30		Dosulepin Mylan Dosulepin Viatris
Cap 25 mg	7.83	50		Oosulepin Mylan 829 Oosulepin
(D			-	Viatris S29
(Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023)				
IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber m		nsing freque		ofranil
Tab 10 mg	10.96	100		ofranii Tofranii
Tab 25 mg		50		ofranil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; prescrib	er may determine o	dispensing f	reque	ency
Tab 10 mg	2.46	100	✓ N	lorpress
Norpress to be Principal Supply on 1 May 2023 Tab 25 mg	6.20	180	./ \	larnraaa
Norpress to be Principal Supply on 1 May 2023	0.29	100	• 1	lorpress
, , , , , ,				
Monoamine-Oxidase Inhibitors (MAOIs) - Non Sel	ective			
TRANYLCYPROMINE SULPHATE				
Tab 10 mg		28		Parnate S29 S29
	22.94	50	-	Parnate
	45.88	100		Parnate S29 S29
	96.00		V F	Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg		60	_	Aurorix
* Tab 300 mg	19.25	60	• •	<u>Aurorix</u>
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE				
* Tab 20 mg		84		PSM Citalopram
Celapram to be Principal Supply on 1 March 2023	2.86		• (Celapram
(PSM Citalopram Tab 20 mg to be delisted 1 March 2023)				
ESCITALOPRAM				
* Tab 10 mg	1.07	28	✓ <u>E</u>	<u>scitalopram</u>
			_	(Ethics)
* Tab 20 mg	1.92	28	✓ E	Escitalopram (Ethica)
				(Ethics)

_					TIVOOO OTOTEM
		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
	JOXETINE HYDROCHLORIDE Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement	2.50	28	✓	Fluox
	 When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multip endorsed. Note: Tablets should be combined with 	le of 20 mg in which	case	the prescr	iption is deemed to be
	Cap 20 mg	2.91	84	1	Fluox
		3.13	90	✓	Arrow-Fluoxetine
(FI	lox Cap 20 mg to be delisted 1 June 2023)				
PΑ	ROXETINE				
*	Tab 20 mg	4.11	90	•	Loxamine
SE	RTRALINE				
*	Tab 50 mg	0.99	30		Setrona
				•	Setrona AU
	Setrona to be Principal Supply on 1 April 2023	4.74	00		Catuana
木	Tab 100 mg	1.74	30		Setrona Setrona AU
	Setrona to be Principal Supply on 1 April 2023			•	octiona Ao
(Se	etrona AU Tab 50 mg to be delisted 1 April 2023)				
•	etrona AU Tab 100 mg to be delisted 1 April 2023)				
0	ther Antidepressants				
NAID	RTAZAPINE				
IVIII	Tab 30 mg	2 60	28	/	Noumed
	Tab 45 mg		28		Noumed
۷F	NLAFAXINE				
	Cap 37.5 mg	6.38	84	/	Enlafax XR
	Cap 75 mg		84	/	Enlafax XR
*	Cap 150 mg	11.16	84	✓	Enlafax XR
A	ntiepilepsy Drugs				
Α	gents for Control of Status Epilepticus				
		ing fraguancy			
IJŀ	ZEPAM – Safety medicine; prescriber may determine dispens Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement	. ,	5	1	Hospira
	a) Up to 5 inj available on a PSO		Ü	•	Поорни
	b) Only on a PSO				
	c) PSO must be endorsed "not for anaesthetic procedure	es".			
	Rectal tubes 5 mg - Up to 5 tube available on a PSO	54.58	5	/	Stesolid
РΗ	ENYTOIN SODIUM				
*	Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a				
	PSO	104.58	5	✓	Hospira
*	Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a				
	PSO	154.01	5	/	Hospira

	Subsidy		Fully	Brand or
	(Manufacturer's Pric		sidised	Generic
	\$	Per		Manufacturer
Control of Epilepsy				
Control of Ephicpsy				
CARBAMAZEPINE				
* Tab 200 mg		100		egretol
* Tab long-acting 200 mg		100		egretol CR
	33.96	200		egretol CR
* Tab 400 mg	34.58	100	✓ Te	egretol
* Tab long-acting 400 mg	39.17	100	✓ Te	egretol CR
* Oral liq 20 mg per ml	26.37	250 ml	✓ Te	egretol
CLOBAZAM - Safety medicine; prescriber may determine dispe	ensing frequency			
Tab 10 mg		50	✓ F	risium
•			•	IOIUIII
CLONAZEPAM - Safety medicine; prescriber may determine d			4.5	
Oral drops 2.5 mg per ml	7.38	10 ml OP	✓ R	ivotril
ETHOSUXIMIDE				
Cap 250 mg	78.89	56	✓ E	ssential
				Ethosuximide \$29
	140.88	100		arontin
Oral lig 250 mg per 5 ml		200 ml		arontin
1 01		200 1111	• 20	aronun
GABAPENTIN				
Note: Not subsidised in combination with subsidised prega				
* Cap 100 mg		100	_	upentin
* Cap 300 mg	8.45	100	_	upentin
* Cap 400 mg	10.26	100	✓ N	<u>upentin</u>
LACOSAMIDE - Special Authority see SA1125 below - Retail	oharmacy			
▲ Tab 50 mg		14	✓ Vi	mpat
▲ Tab 100 mg		14		mpat
_ 140 100 mg	200.24	56		mpat
▲ Tab 150 mg		14		mpat
_ 100 100 mg	300.40	56		mpat
▲ Tab 200 mg		56		mpat
	400.55	90	▼ VI	ιιιμαι

Cubaidy

Eully

Drand or

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see note).

Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

LAMOTRIGINE

55.00	30	✓ Lamictal
	30	✓ Lamictal
	56	✓ Logem
	56	✓ Logem
	56	✓ Logem
	55.00 50.00 2.76 3.31 4.40	

	Subsidy		Fully Brand or
	(Manufacturer's P \$	rice) Subs Per	idised Generic Manufacturer
EVETIRACETAM	<u> </u>		
Tab 250 mg	4.99	60	✓ Everet
Tab 500 mg		60	✓ Everet
Tab 750 mg		60	✓ Everet
Tab 1,000 mg	18.59	60	✓ Everet
Oral liq 100 mg per ml	44.78	300 ml OP	✓ Levetiracetam-AFT
HENOBARBITONE			
For phenobarbitone oral liquid refer Standard For	mulae, page 256		
Tab 15 mg		500	✓ PSM
Tab 30 mg		500	✓ PSM
HENYTOIN SODIUM			
Tab 50 mg	75.00	200	✓ Dilantin Infatab
Cap 30 mg		200	✓ Dilantin illiatab
Cap 100 mg		200	✓ Dilantin
Oral lig 30 mg per 5 ml		500 ml	✓ Dilantin
REGABALIN			
Note: Not subsidised in combination with subsidiation	and anhanontin		
Cap 25 mg		56	✓ Pregabalin Pfizer
Cap 25 mg		50	•
· Cap 75 mg	7.80	FC	✓ Milpharm \$29
Cap 75 mg		56	✓ Pregabalin Pfizer
0 150	8.10	F.C	✓ Milpharm S29
Cap 150 mg	4.01	56	✓ Lyrica
	40.44		✓ Pregabalin Pfizer
0 000	12.44		✓ Milpharm S29
Cap 300 mg		56	Pregabalin Pfizer
RIMIDONE			
Tab 250 mg	37.35	100	Primidone Clinect
DDIUM VALPROATE			
Tab 100 mg	13.65	100	✓ Epilim Crushable
Tab 200 mg EC	27.44	100	✓ Epilim
Tab 500 mg EC	52.24	100	✓ Epilim
Oral liq 200 mg per 5 ml	20.48	300 ml	✓ Epilim S/F Liquid
			✓ Epilim Syrup
Inj 100 mg per ml, 4 ml	41.50	1	✓ Epilim IV
TIRIPENTOL - Special Authority see SA1330 below	/ - Retail pharmacy		
Cap 250 mg		60	✓ Diacomit S29
		60	✓ Diacomit \$29
Powder for oral liq 250 mg sachet		OU	▼ Diacollile 329

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
	\$	Per	✓	Manufacturer
TOPIRAMATE				
▲ Tab 25 mg	11.07	60	1	Arrow-Topiramate
				Topiramate Actavis
	26.04		1	Topamax
▲ Tab 50 mg	18.81	60	1	Arrow-Topiramate
•			1	Topiramate Actavis
	44.26		1	Topamax
▲ Tab 100 mg	31.99	60	1	Arrow-Topiramate
•			1	Topiramate Actavis
	75.25		1	Topamax
▲ Tab 200 mg	55.19	60	1	Arrow-Topiramate
·			1	Topiramate Actavis
	129.85		1	Topamax
▲ Sprinkle cap 15 mg	20.84	60	1	Topamax
▲ Sprinkle cap 25 mg		60	1	Topamax
VIGABATRIN - Special Authority see SA2088 below - Retail pha				-
▲ Tab 500 mg		100	1	Sabril

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents: or
 - 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:

2 Either:

- 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
- 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Acute Migraine Treatment

DI	7A	TI	DΙ	דם	٦Λ.	NI
к	/ A	ш	ĸп	РΙ	А	N

✓ Rizamelt Tab orodispersible 10 mg......3.65 30

	Subsidy (Manufacturer's Price) \$) Per	Fully Subsidised	
SUMATRIPTAN Tab 50 mg Tab 100 mg Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj	22.68	90 90		Sumagran Sumagran
prescription	34.00	2 OP	•	Imigran
Prophylaxis of Migraine				
For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR PIZOTIFEN	SYSTEM, page 49			
* Tab 500 mcg	23.21	100	1	Sandomigran
Antinausea and Vertigo Agents				
For Antispasmodics refer to ALIMENTARY TRACT, page 8				
APREPITANT – Special Authority see SA0987 below – Retail	nharmacy			
Cap 2 × 80 mg and 1 × 125 mg		3 OP	1	Emend Tri-Pack
■ SA0987 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals viewedgenic chemotherapy and/or anthracycline-based chemot Renewal from any relevant practitioner. Approvals valid for 12 chemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE	herapy for the treatme months where the pat	nt of m	nalignancy	
* Tab 16 mg	4.62	100	1	Serc
CYCLIZINE HYDROCHLORIDE				
Tab 50 mg	0.49	10	1	Nausicalm
CYCLIZINE LACTATE				
Inj 50 mg per ml, 1 ml ampoule - Up to 10 inj available on	a			
PSO	16.36	10	1	<u>Hameln</u>
DOMPERIDONE				
* Tab 10 mg	2.85 4.00	100		Pharmacy Health Domperidone Viatris
(Pharmacy Health Tab 10 mg to be delisted 1 June 2023)				
HYOSCINE HYDROBROMIDE				
* Inj 400 mcg per ml, 1 ml ampoule		10	/	Martindale S29
B				

⇒SA1998 Special Authority for Subsidy

Patch 1.5 mg - Special Authority see SA1998 below - Retail

pharmacy......17.70

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

✓ Scopoderm TTS

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Per	Subsidised Generic Manufacturer
ETOCI OPPAMINE HYDROCHI OPINE	Ψ	1 01	- Manadataron
ETOCLOPRAMIDE HYDROCHLORIDE Tab 10 mg - Up to 30 tab available on a PSO	1 20	100	✓ Metoclopramide
Tab 10 mg - Up to 30 tab available on a PSO	1.30	100	Actavis 10
: Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a	PSO 7.00	10	✓ Baxter
	1 00	10	Daxiei
NDANSETRON	0.60	E0	✓ Onrex
Tab 4 mg		50	
Tab disp 4 mg - Up to 10 tab available on a PSO	0.76	10	✓ Ondansetron ODT-DRLA
- Toh 9 mg	4.57	50	✓ Onrex
Tab 8 mg			✓ Ondansetron
Tab disp 8 mg - Up to 10 tab available on a PSO	1.13	10	Ondanseiron ODT-DRLA
DOCUM ORDER AZIME			ODI-DREA
ROCHLORPERAZINE	F 07		
Tab 3 mg buccal		50	Duessatana
Tab E ma . Un to 20 tab available on a DCO	(30.00)	OFO	Buccastem
Tab 5 mg – Up to 30 tab available on a PSO		250	✓ <u>Nausafix</u>
Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO	∠5.81	10	✓ Stemetil
Antipsychotics			
Antipayenotica			
General			
MISULPRIDE – Safety medicine; prescriber may determine o			
Tab 100 mg		30	✓ Sulprix
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg	29.78	60	Sulprix
RIPIPRAZOLE - Safety medicine; prescriber may determine	dispensing frequency		
Tab 5 mg	10.50	30	✓ Aripiprazole San
Tab 10 mg	10.50	30	 Aripiprazole Sand
Tab 15 mg	10.50	30	 Aripiprazole Sand
Tab 20 mg	10.50	30	 Aripiprazole Sand
Tab 30 mg	10.50	30	Aripiprazole San
HLORPROMAZINE HYDROCHLORIDE - Safety medicine; ¡	prescriber may determi	ne dis	spensing frequency
Tab 10 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Tab 25 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO	36.73	100	✓ Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO		10	✓ Largactil
OZAPINE - Hospital pharmacy [HP4]			-
Safety medicine; prescriber may determine dispensing free	quency		
Tab 25 mg		50	✓ Clopine
······································			✓ Clozaril
	13.37	100	✓ Clopine
			✓ Clozaril
			✓ Clopine
Tab 50 mg	8.67	50	• Clopine
Tab 50 mg	8.67 17.33	50 100	✓ Clopine
Tab 50 mg Tab 100 mg	17.33		
·	17.33	100	✓ Clopine
·	17.33	100	✓ Clopine✓ Clopine
·	17.33 17.33	100 50	✓ Clopine ✓ Clopine ✓ Clozaril
·	17.33 17.33 34.65	100 50	✓ Clopine✓ Clopine✓ Clozaril✓ Clopine

✓ Versacloz

100 ml

	Subsidy		Fully	Brand or
	(Manufacturer's Price) Per	Subsidised	Generic Manufacturer
	Ψ	rei		Ivialiulaciulei
HALOPERIDOL – Safety medicine; prescriber may determine d				_
Tab 500 mcg - Up to 30 tab available on a PSO		100		Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg - Up to 30 tab available on a PSO	14.86	50	_	Serenace
	29.72	100		Serenace
Oral liq 2 mg per ml - Up to 200 ml available on a PSO		100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	PSO21.55	10	✓ :	Serenace
LEVOMEPROMAZINE - Safety medicine; prescriber may dete	ermine dispensing free	quency	,	
Tab 25 mg (33.8 mg as a maleate)		100		Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan
Tab 100 mg (135 mg as a maleate)		100		Nozinan (Swiss)
Tab 100 mg as a maleate		100		Nozinan (Omeo)
•				
_EVOMEPROMAZINE HYDROCHLORIDE - Safety medicine;				
Inj 25 mg per ml, 1 ml ampoule	16.75	5	•	Neuraxpharm S29
			✓	Nozinan S29 S29
	24.48	10		Wockhardt
	33.50		✓	Nozinan
Wockhardt to be Principal Supply on 1 April 2023				
Nozinan Inj 25 mg per ml, 1 ml ampoule to be delisted 1 April 2	2023)			
LITHIUM CARBONATE - Safety medicine; prescriber may dete	ermine dispensing fre	auency	ı	
Tab long-acting 400 mg		100		Priadel
Cap 250 mg		100		Douglas
		100	•	Douglas
DLANZAPINE – Safety medicine; prescriber may determine dis				
Tab 2.5 mg		28		Zypine
Tab 5 mg		28		<u>Zypine</u>
Tab orodispersible 5 mg	1.81	28		Zypine ODT
Tab 10 mg	2.01	28	✓ ;	<u>Zypine</u>
Tab orodispersible 10 mg	2.38	28	✓ ;	Zypine ODT
PERICYAZINE - Safety medicine; prescriber may determine di	spensina frequency			
Tab 2.5 mg		84	✓	Neulactil
	12.49	100		Neulactil
Tab 10 mg		84		Neulactil
145 17 mg	44.45	100		Neulactil
OLIETIA DINIE				
QUETIAPINE – Safety medicine; prescriber may determine disp		00		0
Tab 25 mg		90		Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg	12.86	90	•	<u>Quetapel</u>
RISPERIDONE - Safety medicine; prescriber may determine d	ispensing frequency			
Tab 0.5 mg	1.86	60	✓	Risperidone (Teva)
Tab 1 mg	2.06	60	✓	Risperidone (Teva)
Tab 2 mg	2.29	60	✓	Risperidone (Teva)
Tab 3 mg		60		Risperidone (Teva)
Tab 4 mg		60		Risperidone (Teva)
Oral lig 1 mg per ml		30 ml		Risperon
, •,		55 1111	- !	<u></u>
ZIPRASIDONE – Safety medicine; prescriber may determine di		00		7d.aa
Cap 20 mg		60		Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg	40	60		Zusdone

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once



	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; po	rescriber may determine			equency
Tab 10 mg	31.45	100		Clopixol
Depot Injections				
FLUPENTHIXOL DECANOATE - Safety medicine; prescriber	may determine dispens	ing fı	requency	
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO		5		Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO		5		Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	•	Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber n	nay determine dispensir	ng fre	equency	
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO		5	✓	Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	55.90	5		Haldol Concentrate
			•	Haldol
				Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - Retail	pharmacy			
Safety medicine; prescriber may determine dispensing freq	quency			
Inj 210 mg vial		1		Zyprexa Relprevv
Inj 300 mg vial		1		Zyprexa Relprevv
Inj 405 mg vial	504.00	1	•	Zyprexa Relprevv
⇒SA1428 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals va	alid for 12 months for ap	plica	tions meet	ting the following criteria:
Either:				
1 The patient has had an initial Special Authority approval	for paliperidone depot	njec	tion or risp	eridone depot injection; or
2 All of the following:				
2.1 The patient has schizophrenia; and				
2.2 The patient has tried but failed to comply with tre				
2.3 The patient has been admitted to hospital or trea		tens	ive outpati	ent or nome-based
treatment for 30 days or more in the last 12 month	ins.			

of an atypical antipsychotic depot injection. PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine: prescriber may determine dispensing frequency

dately medicine, prescriber may determine dispensing frequency		
Inj 25 mg syringe194.25	1	✓ Invega Sustenna
Inj 50 mg syringe271.95	1	✓ Invega Sustenna
Inj 75 mg syringe357.42	1	✓ Invega Sustenna
Inj 100 mg syringe435.12	1	✓ Invega Sustenna
Inj 150 mg syringe435.12	1	✓ Invega Sustenna

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic r ✓ Manufacturer	
PALIPERIDONE PALMITATE - Special Authority see SA2167 b	elow – Retail pharma	су		
Inj 175 mg syringe	815.85	1	✓ Invega Trinza	
Inj 263 mg syringe	1,072.26	1	✓ Invega Trinza	
Inj 350 mg syringe		1	✓ Invega Trinza	
Inj 525 mg syringe		1	✓ Invega Trinza	
⇒SA2167 Special Authority for Subsidy			-	

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy Safety medicine; prescriber may determine dispensing frequency ✓ Risperdal Consta ✓ Risperdal Consta ✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or clanzapine depot injection; or
- 2 All of the following:

Anxiolytics

- 2.1 The patient has schizophrenia or other psychotic disorder; and
- 2.2 Has tried but failed to comply with treatment using oral atvoical antipsychotic agents: and
- 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency ✓ Clopixol

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	18.50	100	✓ Buspirone Viatris
* Tab 10 mg	12.50	100	 Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may determine	ne dispensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine d	ispensing frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine	e dispensing frequency		
Tab 1 mg	9.72	250	✓ Ativan
Tab 2.5 mg	12.50	100	✓ Ativan

Subsidy (Manufacturer's Price) Per

Fully Subsidised Brand or Generic Manufacturer

Multiple Sclerosis Treatments

⇒SA2176 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 2 Patients has an EDSS score between 0 6.0; and
- 3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months: and
- 4 All of the following:
 - 4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 4.5 Either:
 - 4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 6 Any of the following:
 - 6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion: or
 - 6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2176 above - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 120 mg	520.00	14	✓ Tecfidera
Cap 240 mg	2,000.00	56	Tecfidera

FINGOLIMOD - Special Authority see SA2176 above - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.
- Cap 0.5 mg......2,200.00 ✓ Gilenya

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price)	9	Fully Subsidised	Brand or Generic
	\$	Per	✓	Manufacturer
GLATIRAMER ACETATE – Special Authority see SA2176 on th Note: Treatment on two or more funded multiple sclerosis tr Inj 40 mg prefilled syringe	reatments simultaneou		not permit	tted. Copaxone
INTERFERON BETA-1-ALPHA — Special Authority see SA2176 Note: Treatment on two or more funded multiple sclerosis tr Inj 6 million iu prefilled syringe	reatments simultaneou 1,170.00		not permit	
INTERFERON BETA-1-BETA – Special Authority see \$A2176 c Note: Treatment on two or more funded multiple sclerosis tr Inj 8 million iu per 1 ml	reatments simultaneou		not permit	•
NATALIZUMAB — Special Authority see SA2176 on the previous Note: Treatment on two or more funded multiple sclerosis tr Inj 20 mg per ml, 15 ml vial	reatments simultaneou	•		tted. Tysabri
OCRELIZUMAB – Special Authority see SA2176 on the previou Note: Treatment on two or more funded multiple sclerosis tr Inj 30 mg per ml, 10 ml vial	reatments simultaneou	•		tted. Ocrevus
TERIFLUNOMIDE – Special Authority see SA2176 on the previous) a) Wastage claimable b) Note: Treatment on two or more funded multiple scleros Tab 14 mg	is treatments simultar	•	y is not pe	rmitted. Aubagio

Sedatives and Hypnotics

0 ✓ Vigisom

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	I Generic
MIDAZOLAM Cofety medicines procesibes may determine diene	<u> </u>	rei		Manuacturei
MIDAZOLAM – Safety medicine; prescriber may determine disper Inj 1 mg per ml, 5 ml ampoule		10		Midazolam Mylan Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available on a PSO	17.28	10		Pfizer
Inj 5 mg per ml, 3 ml ampoule	5.00	5 5		Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule — Up to 5 inj available of a PSO On a PSO for status epilepticus use only. PSO must be e	13.09	5 epiler		Pfizer only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 b	elow – Retail pharma	acy		•
Inj 200 mg per ml, 1 ml ampoule	103.30	10	1	Max Health S29
⇒SA1386 Special Authority for Subsidy				

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM – Safety medicine; prescriber may determine di Tab 10 mg		25	✓ <u>Normison</u>
TRIAZOLAM - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 125 mcg	5.10	100	
	(9.85)		Hypam
Tab 250 mcg	4.10	100	
	(11.20)		Hypam
ZOPICLONE - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 7.5 mg	10.80	500	✓ Zopiclone Actavis

Spinal Muscular Atrophy

NUSINERSEN - PCT only - Special Authority see SA2174 below Inj 12 mg per 5 ml vial120,000.00 ✓ Spinraza

⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

Stimulants/ADHD Treatments

ATOMOXETINE			
Cap 10 mg	18.41	28	✓ APO-Atomoxetine ✓ APO-Atomoxetine S29 S29
			✓ Generic Partners
0 40	107.03	20	✓ Strattera
Cap 18 mg	27.06	28	✓ APO-Atomoxetine
	407.00		✓ Generic Partners
0 05	107.03		✓ Strattera
Cap 25 mg	29.22	28	✓ APO-Atomoxetine
0 40	00.00		✓ Generic Partners
Cap 40 mg	29.22	28	✓ APO-Atomoxetine
			✓ Generic Partners
• ••	107.03		✓ Strattera
Cap 60 mg	46.51	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine S29 S29
			Generic Partners
Cap 80 mg	56.45	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine S29 S29
			Generic Partners
Cap 100 mg	58.48	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine S29 S29
			✓ Generic Partners
(Strattera Cap 10 mg to be delisted 1 November 2023)			
(Strattera Cap 18 mg to be delisted 1 November 2023)			
(Strattera Cap 40 mg to be delisted 1 November 2023)			
DEXAMFETAMINE SULFATE – Special Authority see SA1149 below	w Dotoil ph	ormoov	
·	w – netali pri	amacy	
a) Only on a controlled drug form			
b) Safety medicine; prescriber may determine dispensing freque	•	100	/ DOM
Tab 5 mg		100	✓ <u>PSM</u>
	28.50		✓ Aspen

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:



Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

a) Only on a controlled drug form

frequency		
3.20	30	✓ Rubifen
	30	✓ Ritalin
		✓ Rubifen
7.75	30	 Methylphenidate ER
		- Teva
7.85	30	✓ Rubifen
	30	Rubifen SR
	30	 Methylphenidate ER
		- Teva
15.50	30	✓ Methylphenidate ER
		- Teva
22.25	30	✓ Methylphenidate ER
		- Teva
	7.75	3.20 30 30 30 307.75 307.85 3010.95 3011.45 3015.50 30

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	(Fully Subsidised	Brand or Generic	
	Per	✓	Manufacturer	

continued...

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency ✓ Concerta Tab extended-release 18 mg.......58.96 30 ✓ Concerta Tab extended-release 27 mg......65.44 30 ✓ Concerta Tab extended-release 36 mg......71.93 ✓ Concerta Tab extended-release 54 mg......86.24 30 Cap modified-release 10 mg15.60 30 ✓ Ritalin LA ✓ Ritalin LA 30 Cap modified-release 30 mg25.52 30 ✓ Ritalin LA ✓ Ritalin LA Cap modified-release 40 mg30.60 30

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:

NERVOUS SYSTEM

Subsidy (Manufacturer's	Ful Price) Subsidise	·
	Per •	Manufacturer

continued...

- 3.1 Applicant is a paediatrician or psychiatrist; or
- 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see SA1999 below – Retail pharmacy
Tab 100 mg29.13 60 ✓ Modavigil

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 0.1
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia			
DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 on the next p	oage – Retail pharma	acy	
Patch 4.6 mg per 24 hour	38.00	30	 Rivastigmine Patch
			<u>BNM 5</u>
Patch 9.5 mg per 24 hour	38.00	30	 Rivastigmine Patch
			BNM 10

Treatments for Demontia

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate: and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg	. ,	28	✓ <u>Buprenorphine</u> Naloxone BNM
Tab sublingual 8 mg with naloxone 2 mg	34.00	28	✓ <u>Buprenorphine</u> Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and



Subsidy (Manufacturer's Price)	S	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE Tab modified-release 150 mg	11.00	30	✓ Zyban
DISULFIRAM			
Tab 200 mg	236.40	100	✓ Antabuse S29
NALTREXONE HYDROCHLORIDE - Special Authority	see SA1408 below – Retai	I pharmacy	

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or

- 2.2 Patient achieved significant improvement but requires further treatment; or
- 2.3 Patient is well controlled but requires maintenance therapy.

✓ Naltraccord

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. Patch 7 mg - Up to 28 patch available on a PSO19.14 ✓ Habitrol 28 Patch 7 mg for direct distribution only - [Xpharm]......4.13 7 ✓ Habitrol Patch 14 mg - Up to 28 patch available on a PSO21.05 28 ✓ Habitrol Patch 14 mg for direct distribution only - [Xpharm]......6.48 7 ✓ Habitrol Patch 21 mg - Up to 28 patch available on a PSO24.12 28 ✓ Habitrol Patch 21 mg for direct distribution only - [Xpharm]......10.93 7 ✓ Habitrol Lozenge 1 mg - Up to 216 loz available on a PSO......19.76 216 ✓ Habitrol 36 ✓ Habitrol Lozenge 2 mg - Up to 216 loz available on a PSO......21.65 216 ✓ Habitrol Lozenge 2 mg for direct distribution only - [Xpharm]3.40 36 ✓ Habitrol ✓ Habitrol Gum 2 mg (Fruit) - Up to 384 piece available on a PSO38.21 384 Gum 2 mg (Fruit) for direct distribution only - [Xpharm]................9.04 96 ✓ Habitrol ✓ Habitrol Gum 2 mg (Mint) - Up to 384 piece available on a PSO......21.42 204 384 ✓ Habitrol Gum 2 mg (Mint) for direct distribution only - [Xpharm]......9.04 ✓ Habitrol 96 Gum 4 mg (Fruit) - Up to 384 piece available on a PSO24.17 204 ✓ Habitrol ✓ Habitrol 384 ✓ Habitrol Gum 4 mg (Fruit) for direct distribution only - [Xpharm].................10.47 96 ✓ Habitrol Gum 4 mg (Mint) - Up to 384 piece available on a PSO......44.17 384 Gum 4 mg (Mint) for direct distribution only - [Xpharm]......10.47 96 ✓ Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 4216.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg17.62	56	✓ Varenicline Pfizer

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

NERVOUS SYSTEM

Subs (Manufactur		
\$	Per	Manufacturer

continued...

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA2153 below

Inj 25 mg vial77.00	•	✓ Ribomustin
Inj 100 mg vial308.00) 1	✓ Ribomustin
Inj 1 mg for ECP	3 1 mc	✓ Baxter

⇒SA2153 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+): and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 ` \$	Per	✓	

continued...

- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

booder Air — i o i — netaii phannacy-opecialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
, ,	45.20		 Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	710.00	1	✓ BiCNU
Inj 100 mg for ECP		100 mg OP	✓ Baxter
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist		-	
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	✓ Cisplatin Ebewe
.,	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP	0.31	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE		· ·	
Tab 50 mg - PCT - Retail pharmacy-Specialist	145.00	50	✓ Cyclonex
Inj 1 g vial – PCT – Retail pharmacy-Specialist		1	✓ Endoxan
, 3 , ,	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist	71.25	1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist		1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
Inj 1 g	96.00	1	✓ Holoxan
Inj 2 g		1	✓ Holoxan
Inj 1 mg for ECP		1 mg	✓ Baxter
<i>,</i>		Ū	

1 ma

✓ Baxter

	Subsidy		Fully	
((Manufacturer's Price)	Per	Subsidised	
	Ф	Per		Manufacturer
LOMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg		20		CeeNU
Cap 40 mg	399.15	20	/	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist		1	✓	Melpha
	67.80		1	Alkeran
			1	Alkeran S29 S29
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
,g	20.0	•		100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial		1		Oxaliplatin Accord
Inj 1 mg for ECP		1 mg	_	Baxter
THIOTEPA - PCT only - Specialist				
• •	CDC	1	./	Bedford \$29
Inj 15 mg vial		1		
				Max Health S29
				THIO-TEPA \$29
			•	Tepadina S29
Inj 100 mg vial	CBS	1	✓	Max Health S29
			•	Tepadina S29
Antimetabolites				
Anumetabonies				
AZACITIDINE - PCT only - Specialist - Special Authority see SAZ	2141 below			
Inj 100 mg vial		1	1	Azacitidine Dr
				Reddy's

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome: or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully Brand or	
	(Manufacturer's Pric		bsidised Generic	
	\$	Per	✓ Manufacturer	
CALCIUM FOLINATE				
Tab 15 mg - PCT - Retail pharmacy-Specialist	135.33	10	DBL Leucovorin	
			Calcium	
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist		5	Hospira	
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Special	ist7.28	1	✓ Calcium Folinate	•
			Sandoz	
			✓ Calcium Folinate	
			Sandoz S29 S2	9
Inj 50 mg - PCT - Retail pharmacy-Specialist	72.80	10	Leucovorin	
			Pharmacia S29	
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	 Calcium Folinate 	•
			Sandoz	
Inj 100 mg - PCT only - Specialist	7.33	1	 Calcium Folinate 	•
			Ebewe	
	94.90	10	Leucovorin	
			Pharmacia S29	
Inj 300 mg - PCT only - Specialist	22.51	1	 Calcium Folinate 	•
			Ebewe	
	25.14		 Leucovorin DBL 	S29
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25 14	1	✓ Calcium Folinate	
ing forming per finit, 65 this vital in Ort Only Openialist	20.17	'	Sandoz	•
			✓ Calcium Folinate	•
			Sandoz S29 S2	
Inj 1 g - PCT only - Specialist	67 51	1	✓ Calcium Folinate	
injing nondring openialisticismississississississississississississi			Ebewe	
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	✓ Calcium Folinate	9
,		•	Sandoz	
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓ Baxter	
CAPECITABINE – Retail pharmacy-Specialist		ŭ		
Tab 150 mg	10.00	60	✓ Capercit	
Tab 500 mg		120	✓ Capercit	
CLADRIBINE – PCT only – Specialist			•	
Inj 2 mg per ml, 5 ml	749 96	1	✓ Litak S29	
Inj 1 mg per ml, 10 ml		1	✓ Leustatin	
Inj 10 mg for ECP		10 mg OP	✓ Baxter	
CYTARABINE				
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special	ist472.00	5	✓ Pfizer	
Inj 100 mg per ml, 20 ml vial – PCT – Retail		· ·		
pharmacy-Specialist	48.80	1	✓ Pfizer	
Inj 1 mg for ECP - PCT only - Specialist		10 mg	✓ Baxter	
Inj 100 mg intrathecal syringe for ECP - PCT only - Special	ist94.40	I00 mg OP	✓ Baxter	
FLUDARABINE PHOSPHATE		-		
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	✓ Fludara Oral	
Inj 50 mg vial - PCT only - Specialist		5	✓ Fludarabine Ebe	we
Inj 50 mg for ECP - PCT only - Specialist	126.80	50 mg OP	✓ Baxter	

	Subsidy (Manufacturer's Price		Fully bsidised	Brand or Generic
	\$	Per		Manufacturer
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	1	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	29.44	1	1	Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist		100 mg	1	Baxter
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist		ŭ		
Inj 1 g, 26.3 ml vial	62.50	1	1	DBL Gemcitabine
Inj 1 g		1	_ '	Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg		Baxter
		ing		Daxio
IRINOTECAN HYDROCHLORIDE – PCT only – Specialist	50 FT		,	
Inj 20 mg per ml, 5 ml vial		1		Accord
	71.44			Irinotecan Actavis 100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	1	Baxter
MERCAPTOPURINE		Ü		
Tab 50 mg - PCT - Retail pharmacy-Specialist	25.00	25	1	Puri-nethol
. , ,		23	•	<u>r un incunoi</u>
Oral suspension 20 mg per ml – Retail pharmacy-Specialist		100I OD	,	A.U
Special Authority see SA1725 below	428.00	100 ml OP		Allmercap

■ SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist9.98	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist33.71	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist56.05	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	Methotrexate Sandoz
*	Inj 10 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 15 mg prefilled syringe14.77	1	Methotrexate Sandoz
*	Inj 20 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 25 mg prefilled syringe14.99	1	Methotrexate Sandoz
*	Inj 30 mg prefilled syringe15.09	1	Methotrexate Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	Methotrexate DBL Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist79.99	1	✓ Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist0.06	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PEMETREXED - PCT only - Specialist - Special Authority see	SA1679 below			
Inj 100 mg vial	60.89	1	✓	Juno Pemetrexed
Inj 500 mg vial	217.77	1	✓	Juno Pemetrexed
Inj 1 mg for ECP	0.55	1 mg	✓	Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:

Tah 40 mg

- 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
- 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

126.31

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

140 40 mg	20	Lanvio
Other Cytotoxic Agents		
AMSACRINE - PCT only - Specialist		
Inj 50 mg per ml, 1.5 ml ampoule1,500.00	6	✓ Amsidine S29
4,736.00		✓ Amsidine S29
Inj 75 mg1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist		
Cap 0.5 mg1,175.87	100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist		
Inj 1 mg per ml, 10 ml vial	10	✓ Phenasen
Inj 10 mg for ECP481.70	10 mg OP	✓ Baxter

25

I anvis

	Subsidy (Manufacturer's Pr	,	Fully sidised	Brand or Generic
	\$	Per		Manufacturer
BLEOMYCIN SULPHATE - PCT only - Specialist				
Inj 15,000 iu, vial	185.16	1	✓ 0	BL Bleomycin Sulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	√ E	Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see S	A1889 below			
Inj 3.5 mg vial		1		DBL Bortezomib Bortezomib Dr-Reddy's
Inj 1 mg for ECP(Bortezomib Dr-Reddy's Inj 3.5 mg vial to be delisted 1 May 2023		1 mg	✓ B	Baxter

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	72.11	1	DBL Dacarbazine
, 3	580.60	10	✓ Dacarbazine
			APP S29
Inj 200 mg for ECP	72.11	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist			
Inj 2 mg per ml, 10 ml	171.93	1	✓ Pfizer
Inj 20 mg vial		10	✓ Daunorubicin
	•		Zentiva S29
Inj 20 mg for ECP	171.93	20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist			
Inj 20 mg	48.75	1	✓ Docetaxel Sandoz
lnj 10 mg per ml, 8 ml vial	46.89	1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	✓ Docetaxel
			Accord S29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP	0.65	1 mg	✓ Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	10.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1	Doxorubicin Ebewe
	17.00		✓ Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	65.00	1	✓ Arrow-Doxorubicin
	69.99		Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	✓ Baxter

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	I Generic Manufacturer
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml vial	25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial	30.00	1	✓	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓	Epirubicin Ebewe
Inj 1 mg for ECP		1 mg	/	Baxter
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	1	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist	340.73	10	1	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Special	list7.90	1	1	Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	/	Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist				
Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	/	Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pha	rmacy-Specialist			
Cap 500 mg	, ,	100	1	Devatis
IBRUTINIB - Special Authority see SA2168 below - Retail phan	macv			
Tab 140 mg	•	30	1	Imbruvica
Tab 420 mg		30	1	Imbruvica

⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
 - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
 - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
 - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

Inj 5 mg vial - PCT only - Specialist	109.74	1	✓ Zavedos
Inj 10 mg vial - PCT only - Specialist	233.64	1	✓ Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
LENALIDOMIDE - Retail pharmacy-Specialist - Special Autho	rity see SA2047 below			
Wastage claimable				
Cap 5 mg	5,122.76	28	✓	Revlimid
Cap 10 mg	4,655.25	21	✓	Revlimid
•	6,207.00	28	✓	Revlimid
Cap 15 mg	5,429.39	21	✓	Revlimid
	7,239.18	28	✓	Revlimid
Cap 25 mg	7,627.00	21	✓	Revlimid

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Fither
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MFSNA

Tab 400 mg - PCT - Retail pharmacy-Specialist314.00	50	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist448.50	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.45	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	15	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	100 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
MITOMYCIN C - PCT only - Specialist				
Inj 5 mg vial	641.70	1	1	Accord S29
Inj 20 mg vial		1	✓.	Teva
Inj 1 mg for ECP	269.85	1 mg	✓	Baxter
MITOZANTRONE - PCT only - Specialist				
Inj 2 mg per ml, 10 ml vial	97.50	1	✓	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	✓	Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority se	ee SA2163 below			
Tab 100 mg	3,701.00	56	✓	Lynparza
Tab 150 mg		56	✓	Lynparza

⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Fither:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment

Subsidy		Fully	Brand or	Τ
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or

5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg	24.00	1	✓ Paclitaxel Ebewe
, ,	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
, ,	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	Paclitaxel Ebewe
	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see	SA1979 below		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMII F)

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 The patient has relapsed acute lymphoblastic leukaemia; and

DENITORIATINI (DEOVVCOEODMVCINI) DOT only Charlette

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist			
Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharmacy-Sp	ecialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 on the next page	– Retail pha	rmacy	
Cap 5 mg	9.13	5	✓ Temaccord
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
Cap 100 mg	35.98	5	✓ Temaccord
	40.20		✓ Apo-Temozolomide
Cap 140 mg	50.12	5	✓ Temaccord
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ Temaccord

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Sp	ecial Authority see SA1124 o	n the next page	
Cap 50 mg	378.00	28	Thalomid
Can 100 mg	756.00	28	✓ Thalomid

Subs	sidy Full	/ Brand or
(Manufactur	rer's Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

100 ✓ Vesanoid	100	Cap 10 mg - PCT - Retail pharmacy-Specialist
1	pelow	ENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA1868 be
42 OP ✓ Venclexta	42 OP	Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg
14 OP ✓ Venclexta	14 OP	Tab 10 mg95.78
7 OP ✓ Venclexta	7 OP	Tab 50 mg239.44
120 ✓ Venclexta	120	Tab 100 mg - Wastage claimable

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax: and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

Subsidy		Fully	
(Manufacturer's Price		Subsidised	
<u></u>	Per		Manufacturer
VINBLASTINE SULPHATE			
Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37	5	✓	Hospira
Inj 1 mg for ECP - PCT only - Specialist6.00	1 mg	· ·	Baxter
VINCRISTINE SULPHATE			
Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Specialist74.52	5	•	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist102.73	5	•	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	·	Baxter
VINORELBINE - PCT only - Specialist			
Inj 10 mg per ml, 1 ml vial12.00	1	✓	Navelbine
42.00		✓	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00	1	✓	Navelbine
210.00		✓	Vinorelbine Ebewe
328.65		✓	Sagent S29
Inj 1 mg for ECP1.25	1 mg	·	Baxter
	50 mg	OP 🗸	Baxter (Sagent)

Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable			
Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel
•			

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and

bsidy Fu turer's Price) Subsidis	Illy Brand or ed Generic	
 \$ Per	✓ Manufacturer	

continued...

- 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day: or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB — Retail pharmacy-Specialist — Special Authority see SA2115 below Brand switch fee payable (Pharmacode 2651564) - see page 254 for details

Tab 100 mg	329.70	30	✓ Alchemy
Tab 150 mg	569.70	30	✓ Alchemy

⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Tab 250 mg	 	 918.00	30	✓ Iressa

	Subsidy	Fully	Brand or
(Manufa	,	sidised	Generic
	\$ Per	·	Manutacturer

⇒SA2116 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Fither:
 - 2.1 Patient is treatment naive: or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
 - 2 Gefitinib to be discontinued at progression; and
 - 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

Authority see SA1460	
2,400.00	60 ✓ Glivec
58.23	60 ✓ Imatinib-Rex
	30 ✓ Imatinib-Rex
84.79	

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website <u>schedule.pharmac.govt.nz/SAForms</u>, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST - access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 on the next page - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6,532.00	120	Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day: and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

W	as	tage	cla	iima	ble	
---	----	------	-----	------	-----	--

Tab 75 mg4,000.00	21	Ibrance
Tab 100 mg4,000.00	21	Ibrance
Tab 125 mg4,000.00	21	✓ Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state: and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
- 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal: or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
RUXOLITINIB – Special Authority see SA1890 below – Retail p Wastage claimable	pharmacy			
Tab 5 mg	2,500.00	56	1	Jakavi
Tab 10mg	5,000.00	56	✓	Jakavi
Tab 15 mg	5,000.00	56	✓	Jakavi
Tab 20 mg	5,000.00	56	/	Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
- 5.2 Haemoglobin level < lower limit of normal; or
- 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
- 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
- 5.5 Karnofsky performance score of less than or equal to 70; or
- 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

1 Any of the following:

Both:

- 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
- 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✓ Manufacturer

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 84

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

Tab 250 mg4,276.19 120 **✓ Zytiga**

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

DIGALLITANIDE

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Tab 50 mg	4.21	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
· ·	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Authority	see SA1895 on	the next page	9
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex

Subsidy	0.1	Fully	Brand or	
(Manufacturer's Price) \$	Per	sidised •	Generic Manufacturer	

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

OCTRECTIDE

OUTILOTIDE	27.50	_	4 88 11 111
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓ Max Health
			✓ Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓ Max Health
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓ Max Health
OCTREOTIDE LONG-ACTING - Special Authority see	SA2119 below – Retail pha	armacy	
Inj depot 10 mg prefilled syringe	439.97	1	 Octreotide Depot
			<u>Teva</u>
Inj depot 20 mg prefilled syringe	647.03	1	 Octreotide Depot
			Teva
Inj depot 30 mg prefilled syringe	718.55	1	 Octreotide Depot
,			Teva

⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		idised	Generic	
 \$	Per	✓	Manufacturer	

continued...

2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

TAMOXIFEN CITRATE

*	Tab 10 mg	60	✓ Tamoxifen Sandoz
*	Tab 20 mg6.65	60	✓ <u>Tamoxifen Sandoz</u>

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer	
Aromatase Inhibitors					
ANASTROZOLE * Tab 1 mg EXEMESTANE * Tab 25 mg		30		natrole fizer Exemestane	
LETROZOLE * Tab 2.5 mg		30	_	etrole	

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE		
* Tab 25 mg	60	Azamun
Azamun to be Principal Supply on 1 April 2023		
* Tab 50 mg8.10	100	Azamun
Azamun to be Principal Supply on 1 March 2023		
MYCOPHENOLATE MOFETIL		
Tab 500 mg35.90	50	✓ Cellcept
Cap 250 mg35.90	100	✓ Cellcept
Powder for oral lig 1 g per 5 ml - Subsidy by endorsement	165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT - Special Authority see SA2103 below -	Retail pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Inj 50 mg prefilled syringe		4	✓ Enbrel

⇒SA2103 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg,

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and

2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs. CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less;
 - 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
 - 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
- 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and

\$ Per ✔ Manufacturer		Subsidy (Manufacturer's Price) \$		Fully Subsidised	Brand or Generic Manufacturer
-----------------------	--	---	--	---------------------	-------------------------------------

continued...

- 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 212 Fither
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Fither:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Sp	ecialist		
Inj 50 mg per ml, 5 ml	2,774.48	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT	only – Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29

Monoclonal Antibodies

	'	8 below – Retail pharm	ADALIMUMAB (AMGEVITA) - Special Authority see SAZI
Amgevita	1	190.00	Inj 20 mg per 0.4 ml prefilled syringe
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled pen
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled syringe

⇒SA2178 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
 - 2 Fither:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	√	

continued...

4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g., inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline: and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plague psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis: or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate. ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

1 Both:

2 Both:

- 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.2 Fither:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

	Subsidy	F	ully	Brand or
(Ma	nufacturer's Price)	Subsidi	sed	Generic
·	\$	Per	✓	Manufacturer

continued...

Both:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose: or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < 1/2+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or 2 Both:
- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

1 The patient has had a good clinical response following 3 initial doses: or

Subsidy	Fully	/ Brand or	
(Manufacturer's Price)	Subsidised	d Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or

Subsidy	,	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

zitner:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
- 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD: and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate: and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Fither:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated): and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		ubsidised	Generic	
\$	Per		Manufacturer	

continued...

- 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or
- 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroillitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

significant response to treatment in the opinion of the physician; or

2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see \$A2157 below - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe	1,599.96	2	Humira
Inj 40 mg per 0.4 ml prefilled syringe	1,599.96	2	Humira
Inj 40 mg per 0.8 ml prefilled pen		2	✓ HumiraPen
Ini 40 mg per 0.8 ml prefilled syringe		2	✓ Humira

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Any of the following:

- 1.1 The patient has had a good clinical response following 12 weeks' initial treatment: or
- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or

Subsidy		Fully	Brand or	
Gabblay		. ,	Diana oi	
(Manufacturer's Price)		Subsidised	Generic	
(Wandacturer 3 Frice)		Oubsidisca	acricio	
\$	Per	_	Manufacturer	
Ψ	1 01	•	Manadatarer	

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per ✔ Manufacturer

continued...

least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Fither:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacy

1 ✓ Eylea

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

- Fither:
 - 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Fither:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
 - 2 Fither:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and

ONCOLOGY AGENTS AND IMMUNOSU	PPRESSANTS		
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
continued			
 8 Patient has an Asthma Control Test (ACT) score of using the ACT and oral corticosteroid dose must be the first dose to assess response to treatment; and 9 Either: 			
9.1 Patient has not previously received an anti-IL9.2 Both:	.5 biological therapy for the	ir severe eosinop	philic asthma; or
9.2.1 Patient was refractory or intolerant to 9.2.2 Patient was not eligible to continue tre within 12 months of commencing trea	eatment with previous anti-litment.	L5 biological the	
Renewal — (Severe eosinophilic asthma) only from a re years for applications meeting the following criteria: Both:	spiratory physician or clinic	al immunologist.	Approvals valid for 2
1 An increase in the Asthma Control Test (ACT) score2 Either:	of at least 5 from baseline;	and	
2.1 Exacerbations have been reduced from base2.2 Reduction in continuous oral corticosteroid u control.	•		·
CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Au	thority see SA2096 below		
Inj 120 mg per ml casirivimab, 11.1 ml vial (1) and inj 1 per ml imdevimab, 11.1 ml vial (1)	•	1 OP ✓ F	Ronapreve
SA2096 Special Authority for Subsidy Initial application — (Treatment of profoundly immunod valid for 2 weeks for applications meeting the following crite All of the following:		om any relevant	practitioner. Approvals
 Patient has confirmed (or probable) COVID-19; and The patient is in the community with mild to moderat Patient is profoundly immunocompromised** and is against COVID-19 or is unvaccinated; and 		ed an adequate r	esponse to vaccination
 4 Patient's symptoms started within the last 10 days; a 5 Patient is not receiving high flow oxygen or assisted 6 Casirivimab and imdevimab is to be administered at 	/mechanical ventilation; and a maximum dose of no gre	ater than 2,400 i	mg.
Notes: * Mild to moderate disease severity as described or ** Examples include B-cell depletive illnesses or patients re			
CETUXIMAB – PCT only – Specialist – Special Authority s Inj 5 mg per ml, 20 ml vial Inj 5 mg per ml, 100 ml vial Inj 1 mg for ECP	364.00 1,820.00	1 🗸 E	Erbitux Erbitux Baxter
⇒SA1697 Special Authority for Subsidy			
Initial application only from a medical oncologist or medical Approvals valid for 6 months for applications meeting the for All of the following:		mendation of a n	nedical oncologist.
Patient has locally advanced, non-metastatic, squan Patient is contraindicated to, or is intolerant of, cisple Patient has good performance status; and To be administered in combination with radiation the	atin; and	d and neck; and	

GEMTUZUMAB OZOGAMICIN - PCT only - Specialist - Special Authority see SA2158 on the next page

✓ Mylotarg

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA2158 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2179 below

Inj 100 mg	428.00	1	Remicade
Inj 1 mg for ECP	4.40	1 mg	Baxter

⇒SA2179 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Crohn's disease (children))** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 1 Paediatric patient has active Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (**Graft vs host disease**) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — **(Pulmonary sarcoidosis)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less: and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist.

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement: and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

- Fither:
 - 1 A withdrawal period has been tried and the patient has relapsed; or
 - 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances: or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plague psoriasis; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Patient has "whole body" severe chronic plague psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation; or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis: or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis; or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:
All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Fither:

Subsidy (Manufactured a Drice)		Fully	Brand or
 (Manufacturer's Price) \$	Subside Per	JISEO 🗸	Generic Manufacturer

continued...

- 2.1 Patients SCCAI is greater than or equal to 4; or
- 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs: and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis - peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacture's	Price)	Fully Subsidised	Brand or Generic	
\$	Pei	•	Manufacturer	

continued...

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
 - 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
 - 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
 - 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
 - 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

MEPOLIZUMAB - Special Authority see SA2154 below	v – Retail pharmacy		
Inj 100 mg prefilled pen	1,638.00	1	Nucala
Inj 100 mg vial	1,638.00	1	Nucala

⇒SA2154 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA2155 below

Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive: and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe	450.00	1	Xolair
Ini 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months. unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
- 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Specialist - Special Authority see SA2143 below

⇒SA2143 Special Authority for Subsidy

Initial application — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Infant was born in the last 2 years and has severe lung, airway, neurological or neuromuscular disease that requires ongoing, life-sustaining community ventilation; or
- 2 Both:
 - 2.1 Infant was born in the last 12 months; and
 - 2.2 Any of the following:
 - 2.2.1 Patient was born at less than 28 weeks gestation; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient was born at less than 32 weeks gestation; and
 - 2.2.2.2 Either:
 - 2.2.2.2.1 Patient has chronic lung disease; or
 - 2.2.2.2. Patient is Māori or any Pacific ethnicity; or
 - 2.2.3 Both:
 - 2.2.3.1 Patient has haemodynamically significant heart disease; and
 - 2.2.3.2 Any of the following:
 - 2.2.3.2.1 Patient has unoperated simple congenital heart disease with significant left to right shunt (see note a); or
 - 2.2.3.2.2 Patient has unoperated or surgically palliated complex congenital heart disease; or
 - 2.2.3.2.3 Patient has severe pulmonary hypertension (see note b); or
 - 2.2.3.2.4 Patient has moderate or severe LV failure (see note c).

Notes:

a) Patient requires/will require heart failure medication, and/or patient has significant pulmonary hypertension, and/or patient

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

will require surgical palliation/definitive repair within the next 3 months.

- b) Mean pulmonary artery pressure more than 25 mmHg.
- c) LV Ejection Fraction less than 40%.

Renewal — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months where patient still meets initial criteria.

PERTUZUMAB - PCT only - Specialist - Special Authori	ity see SA1606 below		
Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	✓ Baxter

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3 The patient has good performance status (ECOG grade 0-1); and
 - 4 Pertuzumab to be administered in combination with trastuzumab; and
 - 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
 - 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Spe	ecial Authority see SA197	'6 below	
Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin: or

-			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per ✓	Manufacturer

continued...

- 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept: or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or

Subsidy		Fully	Brand or
(Manufacturer's	Price) Subs	sidised	Generic
<u> </u>	Per	1	Manufacturer

continued...

- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

HITUXIMAB (HIXIMYO) - PCT only - Specialist -	- Special Authority see SA2114 below
Ini 100 ma nor 10 ml vial	275 22

inj 100 mg per 10 mi viai	2/5.33	2	▼ <u>RIXIMYO</u>
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2114 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Fither:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD: and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1.000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and

Subsidy (Manufacturer's I	Price) Sub	Fully	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1.000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	1	Manufacturer	

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy): and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- - 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
 - 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia*

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and

Subsidy	Full		
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2 The patient has not received rituximab in the previous 6 months; and
 - 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's	Price) Su	ıbsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Roth:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment: or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

experienced intolerable side effects.

d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient has "whole body" severe chronic plague psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plagues have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plague psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 on the next page - Retail pharmacy

Note: Siltuximab is to be administered at doses no g		y 3 weeks.	
Inj 100 mg vial	770.57	1	✓ Sylvant
Inj 400 mg vial	3,082.33	1	✓ Sylvant

Subsidy	Fı	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	/	Manufacturer

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Inj 100 mg per ml, 1.5 ml vial with cilgavimab 100 mg per ml,1.5 ml vial	0.00	1	✓ Evusheld
TOCILIZUMAB - PCT only - Special Authority see SA2159 belo			
Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
	880.00	4	✓ RoActemra S29 S29
Inj 20 mg per ml, 10 ml vial	550.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
Inj 20 mg per ml, 20 ml vial	1,100.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
	4,400.00	4	✓ RoActemra S29 S29
Inj 1 mg for ECP	2.85	1 mg	✓ Baxter

⇒SA2159 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Roth
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Fither:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and

- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

✓ Herceptin

 Inj 440 mg vial
 3,875.00
 1
 ✓ Herceptin

 Inj 1 mg for ECP
 9.36
 1 mg
 ✓ Baxter

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib: and
- 3 Fither:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria: All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and

Subsidy (Manufacturer's Pric	ce) Su	Fully bsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Fither:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2144 below

Inj 100 mg vial	2,320.00	1	Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	24.52	1 mg	✓ Baxter

⇒SA2144 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 5.1 Patient does not have symptomatic brain metastases; or
- 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe.......4,162.00 1 ✓ Stelara

⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

insufficient benefit to meet renewal criteria: or

2.2.2 Both:

2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and

2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active ulcerative colitis: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

⇒SA2183 Special Authority for Subsidy

Initial application — (**Crohn's disease - adults**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated): or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
- 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
- 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids: or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has active ulcerative colitis; and

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Programmed Cell Death-1 (PD-1) Inhibitors

DURVALUMAB - PCT only - Specialist - Special Authority see	SA2164 below		
Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	Imfinzi
Inj 1 mg for ECP	9.59	1 mg	Baxter

⇒SA2164 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
 - 3 Treatment with durvalumab to cease upon signs of disease progression; and
 - 4 Total continuous treatment duration must not exceed 12 months.

NIVOLUMAB - PCT only - Specialist - Special Authority see SA2120 below

Opdivo	1	1,051.98	Inj 10 mg per ml, 4 ml vial
✓ Opdivo	1	2,629.96	Inj 10 mg per ml, 10 ml vial
✓ Baxter	1 mg	27.62	Inj 1 mg for ECP

⇒SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV: and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab: or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging

Subsidy	Full		
 (Manufacturer's Price) \$	Subsidise Per 🗸	d Generic Manufacturer	

continued...

or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special A	Authority see SA2121 below	
Inj 25 mg per ml, 4 ml vial	4,680.00 1	✓ Keytruda
Inj 1 mg for ECP	49.14 1 mg	✓ Baxter

⇒SA2121 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV: and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and

	Subsidy	Fully	Brand or
(Manu	acturer's Price)	Subsidised	Generic
	\$ Per		Manufacturer

continued...

- 2.2 Patient has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml		50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail	ail pharmacy		
Wastage claimable			
Tab 10 mg	6,512.29	30	Afinitor
Tab 5 mg	4,555.76	30	✓ Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

SIROLIMLIS	 Special Authority see 	SA2005 on the	next nage -	Retail nharmacy

Tab 1 mg	 749.99	100	Rapamune
Tab 2 mg	 1,499.99	100	✓ Rapamune
•	449.99		✓ Rapamune

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✔ Manufacturer

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP: or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

 $\textbf{Renewal--(renal angiomyolipoma(s) associated with tuberous sclerosis complex*)} \ \ \text{from any relevant practitioner}.$

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufactu 	rer

continued...

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg		100	✓ Tacrolimus Sandoz
Cap 0.75 mg		100	✓ Tacrolimus Sandoz
Cap 1 mg		100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Antiallergy Preparations

Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 ini per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

Inj 0.15 mg per 0.3 ml auto-injector	90.00	1 OP	Epipen Jr
Inj 0.3 mg per 0.3 ml auto-injector	90.00	1 OP	Epipen

⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
 - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA13	367 above –	Retail pharma	су
Initiation kit - 5 vials freeze dried venom with diluent	305.00	1 OP	✓ VENOX \$29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX \$29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with			
diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)		sidised	Generic	
	\$	Per	•	Manufacturer	
WASP VENOM ALLERGY TREATMENT - Special Authority see	e SA1367 on the pre	vious page	- Reta	l pharmacy	_
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze	•			,	
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ A	lbey	
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		-		•	
dried venom, with diluent	305.00	1 OP	✓ H	ymenoptera S29	
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze		-		,	
dried venom, with diluent	305.00	1 OP	✓ ∨	enomil \$29	
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze		-			
dried venom, with diluent	305.00	1 OP	✓ H	ymenoptera S29	
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze			-	,	
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305 00	1 OP	✓ A	lhev	
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze		. 0.		,	
dried venom, with diluent		1 OP	✓ V	enomil S29	
and vonom, was and on		1 01		011011111	
Antihistamines					
7 Hillington Hilling					
CETIRIZINE HYDROCHLORIDE					
* Tab 10 mg		100	✓ Z		
* Oral liq 1 mg per ml	2.84	200 ml	✓ H	<u>istaclear</u>	
CHLORPHENIRAMINE MALEATE					
* Oral liq 2 mg per 5 ml	9.37	500 ml	✓ H	istafen	
DEXTROCHLORPHENIRAMINE MALEATE					
* Tab 2 mg	2.02	40			
, 3	(8.40)		Р	olaramine	
	`1.01 [′]	20			
	(5.99)		Р	olaramine	
* Oral liq 2 mg per 5 ml	1.77	100 ml			
	(10.29)		Р	olaramine	
FEXOFENADINE HYDROCHLORIDE					
* Tab 60 mg	4.34	20			
· ·	(8.23)		Т	elfast	
* Tab 120 mg	4.74	10			
	(8.23)		Т	elfast	
	14.22	30			
	(26.44)		Т	elfast	
LORATADINE					
* Tab 10 mg		100	√ <u>L</u>	orafix	
* Oral liq 1 mg per ml	1.43	100 ml	✓ H	aylor syrup	
PROMETHAZINE HYDROCHLORIDE					
* Tab 10 mg	1.39	50	✓ A	llersoothe	
* Tab 25 mg		50	_	llersoothe	
* Oral liq 1 mg per 1 ml		100 ml	_	llersoothe	
* Inj 25 mg per ml, 2 ml ampoule - Up to 5 inj available on a F	PSO21.09	5	✓ H	ospira	
·					

	Subsidy	D.:) O.:I:	Fully Brand or
	(Manufacturer's \$	Price) Subsi Per	dised Generic ✓ Manufacturer
	Ψ	Геі	- Manuacturei
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose	14.01	200 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free	8.54	200 dose OP	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose	17.52	200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free	12.50	200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17 00	200 dose OP	✓ Pulmicort
1 owder for initialization, 100 mag per doce		200 0000 01	Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	✓ Pulmicort
1 owder for initialiation, 200 mag per dose	10.00	200 0030 01	Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	✓ Pulmicort
1 owder for illinatation, 400 micg per dose	52.00	200 dose Oi	Turbuhaler
FLUTIOACONE			Turbunaici
FLUTICASONE	7.40	100 00	
Aerosol inhaler, 50 mcg per dose		120 dose OP	Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose OP	 ✓ <u>Flixotide</u> ✓ Flixotide Accuhaler
Powder for inhalation, 250 mcg per dose	11.93	60 dose OP	Flixotide Accunaier
Inhaled Long-acting Beta-adrenoceptor Agonis	ts		
EFORMOTEROL FUMARATE			
Powder for inhalation, 12 mcg per dose, and monodose dev		60 dose	E 19
(F	(35.80)	"	Foradil
(Foradil Powder for inhalation, 12 mcg per dose, and monodose	device to be dei	listea 1 July 2023	3)
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dose	e)10.32	60 dose OP	
	(16.90)		Oxis Turbuhaler
INDACATEROL			
Powder for inhalation 150 mcg	61.00	30 dose OP	✓ Onbrez Breezhaler
Powder for inhalation 300 mcg		30 dose OP	✓ Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose	26.25	120 dose OP	✓ Serevent
Acrosor initialer of O-free, 25 may per dose	20.23	120 dose OF	- Ocieveni

Subsidy

Fully

Brand or

Powder for inhalation, 50 mcg per dose, breath activated26.25

60 dose OP ✓ Serevent Accuhaler

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

Inhaled Corticosteroids with Long-Acting Beta-Adrenocep	tor Agonists	
BUDESONIDE WITH EFORMOTEROL Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide with 6 mcg eformoterol fumarate metered dose)	120 dose OP	✓ DuoResp Spiromax
eformoterol fumarate metered dose) – No more than 2		
dose per day82.50	120 dose OP	✓ DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg	120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg 33.74	120 dose OP	✓ Symbicort Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg21.40	120 dose OP	✓ Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg 33.74	120 dose OP	✓ Symbicort Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate		
12 mcg - No more than 2 dose per day33.74	60 dose OP	✓ Symbicort Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL		
Powder for inhalation 100 mcg with vilanterol 25 mcg44.08	30 dose OP	✓ Breo Ellipta
FLUTICASONE WITH SALMETEROL		
Aerosol inhaler 50 mcg with salmeterol 25 mcg	120 dose OP	Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg	120 dose OP	✓ <u>Seretide</u>
more than 2 dose per day33.74 Powder for inhalation 250 mcg with salmeterol 50 mcg – No	60 dose OP	✓ Seretide Accuhaler
more than 2 dose per day44.08	60 dose OP	✓ Seretide Accuhaler
Beta-Adrenoceptor Agonists		
SALBUTAMOL		
Oral liq 400 mcg per ml40.00	150 ml	✓ <u>Ventolin</u>
Infusion 1 mg per ml, 5 ml	10 5	✓ Ventolin✓ Ventolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO53.00	5	ventolin
Inhaled Beta-Adrenoceptor Agonists		
SALBUTAMOL		
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO3.80	200 dose OP	✓ Respigen
(6.20)		✓ SalAir Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb		VOITIOIIII
available on a PSO8.96	20	✓ <u>Asthalin</u>
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO9.43	20	✓ <u>Asthalin</u>
TERBUTALINE SULPHATE		
Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated22.20	120 dose OP	✓ Bricanyl Turbuhaler

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Anticholinergic Agents

IPRATROPIUM BROMIDE

Aerosol inhaler, 20 mcg per dose CFC-free - Up to 400 dose			
available on a PSO16.	.20 200 do	ose OP 🗸 🗸	trovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 neb			
available on a PSO11.	.73 2	20 🗸 U	Jnivent
28	20	✓ A	ccord \$29

Inhaled Beta-Adrenoceptor Agonists with Anticholinergic Agents

SALBUTAMOL WITH IPRATROPIUM BROMIDE

Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg per		
dose CFC-free12.19	200 dose OP	Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per		
vial 2.5 ml ampoule – Un to 20 neb available on a PSO 11.04	20	✓ Duolin

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	

continued...

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
 - 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL – Special Authority see SA1584 Powder for Inhalation 50 mcg with indacaterol 110 mcg81.00	on the previous page – Retail pharmacy 30 dose OP ✓ Ultibro Breezhaler
TIOTROPIUM BROMIDE WITH OLODATEROL – Special Authority see SA15i Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00	84 on the previous page − Retail pharmacy 60 dose OP ✓ Spiolto Respimat
UMECLIDINIUM WITH VILANTEROL – Special Authority see SA1584 on the Powder for inhalation 62.5 mcg with vilanterol 25 mcg77.00	orevious page – Retail pharmacy 30 dose OP ✓ Anoro Ellipta

Antifibrotics

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg	2,554.00	60 OP	Ofev
Cap 150 mg	3,870.00	60 OP	Ofev

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 on the next page

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90	Esbriet
Tab 267 mg	1,215.00	90	Esbriet

Subsidy		Fully	Brand or
(Manufacturer's Price)	_	Subsidised	Generic
<u> </u>	Per		Manutacturer

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists

МО	NTELUKAST			
*	Tab 4 mg	3.10	28	✓ Montelukast Mylan
*	Tab 5 mg	3.10	28	✓ Montelukast Mylan
	•			✓ Montelukast Viatris
*	Tab 10 mg	2.90	28	✓ Montelukast Mylan
	·			✓ Montelukast Viatris

Methylxanthines

AM	IN	0	PHYLLINE	Ξ
			-	

* Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available	e on a		
PSO	180.00	5	✓ DBL Aminophylline
THEOPHYLLINE			
* Tab long-acting 250 mg	23.02	100	✓ Nuelin-SR
* Oral liq 80 mg per 15 ml		500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA - Special Authority see SA1978 below - Reta	ail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

continued...

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

IVACAFTOR - PCT only - Specialist - Special Authority see \$A2017 below

Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 9 Fither
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele: or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele: and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

Not funded for use as a nasal drop.

Nasal Preparations

Allergy Prophylactics

BUDESONIDE

Metered aqueous nasal spray, 50 mcg per dose		✓ <u>SteroClear</u> ✓ SteroClear
FLUTICASONE PROPIONATE	2.01	<u> </u>
Metered aqueous nasal spray, 50 mcg per dose	1.98 120 dose OP	✓ Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%	5.23 15 ml OP	✓ Univent

Oral liq 20 mg per ml (10 mg base per ml)......15.10

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Respiratory Devices				
MASK FOR SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO c) Only for children aged six years and under Small	2.20	1	√ e-	chamber Mask
PEAK FLOW METER a) Up to 25 dev available on a PSO b) Only on a PSO				
Low range	9.54	1		ini-Wright AFS Low Range
Normal range	9.54	1		ini-Wright Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSOb) Only on a PSO				
220 ml (single patient)	2.95	1	√ e-	chamber Turbo
510 ml (single patient)	5.12	1	•	chamber La Grande
800 ml	6.50	1	√ Vo	olumatic
Respiratory Stimulants				
CAFFEINE CITRATE				

✓ Biomed

25 ml OP

	Subsidy (Manufacturer's F	Price) Sub	Fully	Brand or Generic
	\$	Per	✓	Manufacturer
Ear Preparations				
FLUMETASONE PIVALATE				
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP		ocacorten-Viaform
			_	ED's
TRIANCINOLONE ACTONIDE WITH CRANICIDIA NEOMYC	INI ANID NIVOTAT	TINI	▼ L(ocorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate	IN AND NYSTAT	I IIN		
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ K	enacomb
0 0 0.0				
Ear/Eye Preparations				
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN				
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and				
gramicidin 50 mcg per ml	4.50	8 ml OP		
	(9.27)		So	ofradex
FRAMYCETIN SULPHATE	4.40	0 OD		
Ear/Eye drops 0.5%	4.13 (8.65)	8 ml OP	Sc	oframycin
	(0.00)		00	Siramyon
Eye Preparations				
Eye preparations are only funded for use in the eye, unless expli-	citly stated other	wise.		
Anti-Infective Preparations				
ACICLOVIR				
* Eye oint 3%	14.88	4.5 g OP	✓ <u>Vi</u>	<u>ruPOS</u>
CHLORAMPHENICOL	4.00	5 × OD	4 D	
Eye oint 1% Eye drops 0.5%		5 g OP 10 ml OP	_	<u>evatis</u> hlorafast
Funded for use in the ear*. Indications marked with * ar			• 01	moraidat
CIPROFLOXACIN				
Eye drops 0.3% - Subsidy by endorsement	9.73	5 ml OP	✓ <u>C</u> i	profloxacin Teva
When prescribed for the treatment of bacterial keratitis of				
for the second line treatment of chronic suppurative otitic Note: Indication marked with a * is an unapproved indic		*; and the pres	scription i	is endorsed accordingly.
GENTAMICIN SULPHATE	alion.			
Eye drops 0.3%	11 40	5 ml OP	√ G	enoptic
(Genoptic Eye drops 0.3% to be delisted 1 August 2023)		0 1111 01	- 0	0.10040
PROPAMIDINE ISETHIONATE				
* Eye drops 0.1%	2.97	10 ml OP		
	(14.55)		Br	rolene

5 g OP

3.5 g OP 5 ml OP ✓ Fucithalmic

✓ Tobrex

✓ Tobrex

SODIUM FUSIDATE [FUSIDIC ACID]

TOBRAMYCIN



Subsidy	Fι	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
 \$	Per	•	Manufacturer

Corticosteroids and Other Anti-Inflammatory Preparations

DE	XAMETHASONE			
*	Eye oint 0.1%	5.86	3.5 g OP	✓ Maxidex
*	Eye drops 0.1%	4.50	5 ml OP	✓ Maxidex
	Ocular implant 700 mcg - Special Authority see SA1680 below			
	- Retail pharmacy1,4	144.50	1	Ozurdex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b		
sulphate 6,000 u per g5.39	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin	-	
b sulphate 6,000 u per ml4.50	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM		
Eye drops 0.1%	5 ml OP	✓ Voltaren Ophtha
FLUOROMETHOLONE		· · · · · · · · · · · · · · · · · · ·
* Eye drops 0.1%	5 ml OP	✓ FML
5.20		✓ Flucon

	Subsidy		Fully	Brand or
	(Manufacturer's P	rice) Subs	idised	Generic
	\$	Per	1	Manufacturer
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
, , ,	(10.34)		L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	√ L	omide
PREDNISOLONE ACETATE				
Eye drops 1%	6.92	10 ml OP	✓ P	rednisolone-AFT
, .	7.00	5 ml OP	✓ P	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority se	ee SA1715 below	– Retail pharn	nacy	
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	✓ N	linims Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE

Eye drops 2%1.79	5 ml OP	✓ Rexacrom
2.62	10 ml OP	✓ Allerfix

Allerfix to be Principal Supply on 1 March 2023 (Rexacrom Eye drops 2% to be delisted 1 March 2023)

Glaucoma Preparations - Beta Blockers

# Eye drops 0.25% * Eye drops 0.5%		5 ml OP 5 ml OP	✓ Betoptic S✓ Betoptic
TIMOLOL # Eye drops 0.25% # Eye drops 0.5% # Eye drops 0.5%, gel forming	2.04	5 ml OP 5 ml OP 2.5 ml OP	✓ Arrow-Timolol ✓ Arrow-Timolol ✓ Timoptol XE
Clausema Prenerations Carbonia Anhyd	roce Inhihiters		

Giauconia Preparations - Carbonic Annyurase inilibitors		
ACETAZOLAMIDE * Tab 250 mg17.03	100	✓ Diamox
BRINZOLAMIDE	5 ml OP	✓ <u>Azopt</u>
DORZOLAMIDE HYDROCHLORIDE * Eye drops 2%	5 ml OP	
(17.44)		Trusopt
DORZOLAMIDE WITH TIMOLOL * Eye drops 2% with timolol 0.5%	5 ml OP	✓ Dortimopt

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's F \$	Price) Subs Per	idised C	Brand or Generic Manufacturer
Glaucoma Preparations - Prostaglandin Analog	jues			
BIMATOPROST * Eye drops 0.03%	5.95	3 ml OP		atoprost ultichem
LATANOPROST * Eye drops 0.005%	1.82	2.5 ml OP	✓ <u>Teva</u>	<u>a</u>
TRAVOPROST * Eye drops 0.004%	9.75	2.5 ml OP	✓ <u>Trav</u>	<u>ratan</u>
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE * Eye drops 0.2%	4.29	5 ml OP	✓ Arro	ow-Brimonidine
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Con	nbigan
LATANOPROST WITH TIMOLOL Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	✓ Arro	ow - Lattim
PILOCARPINE HYDROCHLORIDE * Eye drops 1% * Eye drops 2%		15 ml OP 15 ml OP	✓ Isop	oto Carpine oto Carpine
Eye drops 4% Subsidised for oral use pursuant to the Standard Formu Eye drops 2% single dose – Special Authority see SA0895	lae.	15 ml OP	·	to Carpine
below – Retail pharmacy	d for 2 years for			ims Pilocarpine following criteria:
2 Patient wears soft contact lenses. Note: Minims for a general practice are considered to be "tools or Renewal from any relevant practitioner. Approvals valid for 2 yes benefiting from treatment.				
Mydriatics and Cycloplegics				
ATROPINE SULPHATE * Eye drops 1%	17.36	15 ml OP	✓ Atro	p <u>t</u>
* Eye drops 1%TROPICAMIDE	8.76	15 ml OP	✓ Cyc	logyl
* Eye drops 0.5%* * Eye drops 1%		15 ml OP 15 ml OP	✓ Myd	,

Preparations for Tear Deficiency

For acetylcysteine eye drops refer Standard Formulae, page 256

15 ml OP

✓ Methopt

HYPROMELLOSE

	Subsidy (Manufacturer's P	rice) Sub	Fully sidised	Brand or Generic
	\$	Per	•	Manufacturer
HYPROMELLOSE WITH DEXTRAN * Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	√ P	oly-Tears

Preservative Free Ocular Lubricants

⇒SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA2134 above – Retail pha	armacy			
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel	
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL -	Special Authority	see SA2134	above – Retail pharmacy	
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	Systane Unit Dose	
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml	10.78	30	✓ Systane Unit Dose	
(Systane Unit Dose Eye drops 0.4% and propylene glycol 0.3%,	0.4 ml to be deliste	ed 1 June 20	023)	
SODIUM HYALURONATE [HYALURONIC ACID] - Special Auth	nority see SA2134	above – Re	tail pharmacy	
Eye drops 1 mg per ml	13.85	10 ml OP	✓ <u>Hylo-Fresh</u>	
Hylo-Fresh has a 6 month expiry after opening. The Ph	armacy Procedure	s Manual re	striction allowing one bottle per	
month is not relevant and therefore only the prescribed of	dosage to the near	est OP may	be claimed.	

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%2.17	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g3.80	5 g OP	✓ VitA-POS



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Various

PHARMACY SERVICES

May only be claimed once per patient.

* Brand switch fee......4.50 1 fee ✓ BSF Alchemy

The Pharmacode for BSF Alchemy is 2651564 - see also page 163

(BSF Alchemy Brand switch fee to be delisted 1 May 2023)

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE

Inj 200 mg per ml, 10 ml ampoule52.88 10 ✓ Martindale Pharma

NAI OXONE HYDROCHI ORIDE

a) Up to 10 ini available on a PSO

b) Only on a PSO

10 ✓ Hameln

Removal and Elimination

CHARCOAL

✓ Carbosorb-X 250 ml OP

a) Up to 250 ml available on a PSO

b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	Exjade
Tab 500 mg dispersible	1.105.00	28	✓ Exiade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per µL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per uL).

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

Versenate

	Subsidy		Fully	Brand or
	(Manufacturer's P		idised	Generic
	\$	Per		Manufacturer
DEFERIPRONE - Special Authority see SA1480 below - Retail	pharmacy			
Tab 500 mg		100	√ F	erriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	√ F	erriprox
⇒SA1480 Special Authority for Subsidy				
Initial application only from a haematologist. Approvals valid wi	thout further ren	ewal unless no	tified fo	or applications meeting the
following criteria:				
Either:				
1 The patient has been diagnosed with chronic iron overload				or
2 The patient has been diagnosed with chronic iron overload	d due to acquired	l red cell aplas	ia.	
DESFERRIOXAMINE MESILATE				
* Inj 500 mg vial	151.31	10	✓ D	BL
•				Desferrioxamine
				Mesylate for Inj
				BP
			✓ [eferoxamine Pfizer
				S29 S29
SODIUM CALCIUM EDETATE				
* Inj 200 mg per ml, 5 ml	53.31	6		
	(156.71)		C	Calcium Disodium



Standard Formulae

Standard Formulae			
ACETYLCYSTEINE EYE DROPS		PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Acetylcysteine inj 200 mg per ml, 10 ml	qs	mg per ml)	
Suitable eye drop base	qs	Phenobarbitone Sodium	400 mg
00050151100710070		Glycerol BP	4 ml
CODEINE LINCTUS (3 mg per 5 ml)	00	Water	to 40 ml
Codeine phosphate	60 mg	DIL COADDINE ODAL LICHID	
Glycerol	40 ml	PILOCARPINE ORAL LIQUID	
Preservative	qs	Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative Water	qs to 500 ml
CODEINE LINCTUS (15 mg per 5 ml)		(Preservative should be used if quantity supplied is	
Codeine phosphate	300 mg	than 5 days.)	ioi illore
Glycerol	40 ml	than 5 days.)	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	
Water	to 100 ml	Methylcellulose	5 g
501 W 10 M 01 T 1 W 10 M		Preservative	qs
FOLINIC MOUTHWASH	4.1	Water	to 500 ml
Calcium folinate 15 mg tab	1 tab	(Preservative should be used if quantity supplied is	for more
Preservative	qs to 500 ml	than 5 days. Maximum 500 ml per prescription.)	
Water (Preservative should be used if quantity supplied is		SODIUM CHLORIDE ORAL LIQUID	
than 5 days. Maximum 500 ml per prescription.)	, ioi illole	Sodium chloride inj 23.4%, 20 ml	qs
than 5 days. Maximum 500 mi per prescription.		Water	qs
METHADONE MIXTURE		(Only funded if prescribed for treatment of hyponatr	aemia)
Methadone powder	qs	VANCOMATON ORAL COLUTION (OF THE TOTAL)	
Glycerol	qs	VANCOMYCIN ORAL SOLUTION (25 mg per ml)	5 vials
Water	to 100 ml	Vancomycin 500 mg injection Glycerin with sucrose suspension	37.5 ml
METHYL HYDROXYBENZOATE 10% SOLUTION	ı	Water	to 100 ml
Methyl hydroxybenzoate	10 g	(Only funded if prescribed for treatment of Clostridia	
Propylene glycol	to 100 ml		ann dimene
(Use 1 ml of the 10% solution per 100 ml of oral lig		· · · · · · · · · · · · · · · · · · ·	
,	,	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	
OMEPRAZOLE SUSPENSION		Vancomycin 500 mg injection	10 vials
Omeprazole capsules or powder	qs	Glycerol BP	40 ml
Sodium bicarbonate powder BP Water	8.4 g to 100 ml	Water	to 100 ml
vvalei	10 100 1111	(Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	am dillicile
PHENOBARBITONE ORAL LIQUID		ioliowing metroriluazore failure)	
Phenobarbitone Sodium	1 g		
Glycerol BP	70 ml		
Water	to 100 ml		

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per Manufacturer

Extemporaneously Compounded Preparations a	nd Galenica	ıls	
CODEINE PHOSPHATE - Safety medicine; prescriber may deter	mine dispensino	g frequency	
Powder – Only in combination		25 g	Dauglas
Only in extemporaneously compounded codeine linctus.	(90.09)		Douglas
COLLODION FLEXIBLE			
Note: This product is no longer being manufactured by the su	pplier and will b	e delisted fror	n the Schedule at a date to be
determined.			4
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE – Only in combination Only in extemporaneously compounded oral mixtures.			
Soln	30.00	100 ml	✓ Midwest
GLYCERIN WITH SODIUM SACCHARIN - Only in combination			
Only in combination with Ora-Plus or when used in the vancor	nycin oral Iquuid	d Standard Fo	rmulae.
Suspension	30.95	473 ml	Ora-Sweet SF
GLYCERIN WITH SUCROSE - Only in combination			
Only in combination with Ora-Plus or when used in the vancor Suspension		d Standard Fo 473 ml	rmulae. ✓ Ora-Sweet
GLYCEROL	30.33	4731111	• Ola-Sweet
Liquid – Only in combination	3.23	500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid prepara			
METHADONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
 c) Safety medicine; prescriber may determine dispensing free d) Extemporaneously compounded methadone will only be re 		rate of the ch	neanest form available
(methadone powder, not methadone tablets).	imbaroca at tric	rate or the or	icapeot form available
Powder	7.84	1 g	✓ AFT
METHYL HYDROXYBENZOATE			_
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE	00.05	400	A MILINAY 4
Powder Suspension – Only in combination		100 g 473 ml	✓ MidWest ✓ Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHA			• Old-Flus
Suspension		473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only			
Suspension		473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM			
Powder - Only in combination		10 g	✓ MidWest
Only in children up to 12 years	325.00	100 g	✓ MidWest
Only in children up to 12 years PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxybenzo	ate 10% solutio	n.	
Liq		500 ml	✓ Midwest
SODIUM BICARBONATE			
Powder BP - Only in combination		500 g	✓ Midwest

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price)	S Per	Fully Subsidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatio		500 ml	✓ M	idwest	_
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water	

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)	12.30	200 ml OP	✓ Calogen
	30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)	12.30	200 ml OP	✓ Calogen
Oil	30.00	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml	114.92	4 OP	✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	rmacy [HP3]	PROTEIN SUPPLEMENT — Special Authority see SA1524 above — Hospital pha
✓ Protifar	225 g OP	Powder
✓ Resource	227 g OP	8.95
Beneprotein		

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see	SA1095 above -	 Hospital pharm 	nacy [HP3]
Liquid	3.75	500 ml OP	✓ Glucerna Select
·	7.50	1,000 ml OP	✓ Nutrison Advanced
			Diason
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1	095 above – Ho	spital pharmacy	[HP3]
Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
,	2.10		✓ Nutren Diabetes

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED − Special Authority see SA1525 above − Hospital pharmacy [HP3]
Powder60.48 400 g OP ✓ Monogen

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child: or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

SPECIAL FOODS

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see Liquid		the previous page 500 ml OP	ge – Hospital pharmacy [HP3] Nutrini Energy RTH Frebini Energy
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see S Liquid		ne previous page 500 ml OP	✓ Nutrini RTH✓ Pediasure RTH
	6.50		✓ Frebini Original
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special pharmacy [HP3]	Authority se	ee SA1379 on the	e previous page – Hospital
Liquid	6.00	500 ml OP	Nutrini Energy Multi Fibre
	7.00		✓ Frebini Energy Fibre
PAEDIATRIC ENTERAL FEED WITH FIBRE 1KCAL/ML - Special A pharmacy [HP3]	uthority see	SA1379 on the	previous page – Hospital
Liquid	7.00	500 ml OP	✓ Frebini Original Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA1	1379 on the	previous page -	
Liquid (strawberry)		200 ml OP	✓ Fortini
Liquid (vanilla)		200 ml OP	✓ Fortini
	6.99	500 ml OP	✓ Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA13	79 on the pr		
Liquid (chocolate)		200 ml OP	✓ Pediasure
Liquid (strawberry)		200 ml OP	✓ Pediasure
Liquid (vanilla)		200 ml OP	✓ Pediasure
	1.34	250 ml OP	✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authorization pharmacy [HP3]	nority see S	A1379 on the pre	evious page – Hospital
Liquid (unflavoured)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)	1.60	200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on to	he previous	page - Hospital	pharmacy [HP3]
Powder		400 g OP	✓ Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority s Liquid			Hospital pharmacy [HP3] Nepro HP RTH
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see S Liquid			
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA	1101 on the previou	s page – Hosp	ital pharmacy [HP3]
Liquid, 200 ml bottle	11.52	4 OP	
	(13.24)		NovaSource Renal
Liquid (apricot) 125 ml	11.52	4 OP	✓ Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	✓ Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 vear for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome: or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Liquid (caramel) 125 ml.......11.52

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - S	Special Authority see	SA1377 above	- Hospital pharmacy [HP3]
Liquid	18.06	1,000 ml OP	✓ Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority	see SA1377 above -	- Hospital pharm	acy [HP3]
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully sidised	Brand or Generic Manufacturer
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see S Powder (unflavoured)		revious page – 80 g OP		ıl pharmacy [HP3] 'ivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Auth [HP3]	ority see SA137	77 on the previo	us page	- Hospital pharmacy
Liquid	9.60	500 ml OP	✓ S	Survimed OPD
·	12.04	1,000 ml OP	-	lutrison Advanced Peptisorb Peptisorb
(Pantisorh Liquid to be delisted 1. June 2023)				

(Peptisorb Liquia to be delisted 1 June 2023)

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Child aged one to eight years: and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Liquid	4.00	500 ml OP	1	Nutrini Low Energy	
				Multi Eibre	

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and

Subsidy (Manufacturer's P	rice) Sı	Fully ubsidised	Brand or Generic	
\$	Per	1	Manufacturer	

3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (**Long-term medical condition**) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia: or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	•	Manufacturer	

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 266 - H Liquid	ospital pharmac 250 ml OP 1,000 ml OP	y [HP3] ✓ Ensure Plus HN ✓ Ensure Plus RTH ✓ Nutrison Energy ✓ Fresubin HP Energy
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on page 266 - Hos Liquid1.24 5.29	spital pharmacy 250 ml OP 1,000 ml OP	 ✓ Isosource Standard ✓ Nutrison Standard RTH ✓ Osmolite RTH
6.50 ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority see SA1859 o Liquid	n page 266 – Ho 1,000 ml OP	✓ Fresubin Original ospital pharmacy [HP3] ✓ Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority see SA1859 on p Liquid	age 266 - Hosp 1,000 ml OP	
ENTERAL FEED WITH FIBRE 1.2KCAL/ML – Special Authority see SA1859 on Liquid	page 266 – Hos 1,000 ml OP	pital pharmacy [HP3] ✓ Jevity Plus
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see SA1859 on Liquid	page 266 - Hos 1,000 ml OP	pital pharmacy [HP3] ✓ Jevity HiCal RTH ✓ Nutrison Energy Multi Fibre
9.80		✓ Fresubin HP Energy Fibre
ENTERAL FEED WITH PROTEIN 1.2KCAL/ML - Special Authority see SA1859 Liquid9.60	on page 266 – H 500 ml OP	Hospital pharmacy [HP3] Fresubin Intensive
ORAL FEED (POWDER) - Special Authority see SA1859 on page 266 - Hospita Powder (chocolate)	al pharmacy [HP 840 g OP	3] ✓ Sustagen Hospital Formula
Powder (vanilla)	850 g OP 840 g OP	✓ Ensure ✓ Sustagen Hospital Formula Active
26.00	850 g OP	✓ Ensure

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	1	Manufacturer	

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 266 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
LINGISCHICIL	(1.26) (1.26)	200 1111 01	Ensure Plus Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 ml with Endorsement		200 ml OP	France Dive
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with	(1.26)		Ensure Plus
Endorsement	0.72 (1.26)	200 ml OP	Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with Endorsement	0.85	237 ml OP	
	(1.33) 0.72	200 ml OP	Ensure Plus
	(1.26) (1.26)	200 1111 01	Ensure Plus Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 266 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisin Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML	 Special Authority see SA1195 on the previous p 	age – Hospital p	harmacy [HP3]
Liquid	5.50	500 ml OP	✓ Nutrison
			Concentrated
	6.50		✓ Fresubin 2kcal HP
	11.00	1,000 ml OP	Ensure Two Cal HN
			RTH

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

continued...

SPECIAL FOODS

	Subsidy		Fully	Brand or
(N	fanufacturer's Price)	5	Subsidised	Generic
	\$	Per	•	Manufacturer

continued...

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX - Special Authority see SA17			
Powder	2.81	1,000 g OP	
	(5.15)	-	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA172	9 above – Hospital p	oharmacy [HP3]	
Powder	3.93 ·	1,000 g OP	
	(7.32)		NZB Low Gluten
	, ,		Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 ab	ove – Hospital pharr	nacy [HP3]	
Powder	5.62	2,000 g OP	
	(18.10)	-	Horlevs Flour

	Subsidy		Fully	Brand or
	(Manufacturer's Pri		sidised	Generic
	\$	Per	1	Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	orevious page – H	lospital pharr	nacy [HF	23]
Buckwheat Spirals	2.00	250 g OP		•
	(3.11)		0	rgran
Corn and Vegetable Shells	2.00	250 g OP		•
•	(2.92)		0	rgran
Corn and Vegetable Spirals	2.00	250 g OP		•
•	(2.92)		0	rgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		•
•	(3.82)		0	rgran
Rice and Corn Macaroni	2.00 [°]	250 g OP		
	(2.92)	· ·	0	rgran
Rice and Corn Penne	2.00 [°]	250 g OP		
	(2.92)		0	rgran
Rice and Maize Pasta Spirals	2.00	250 g OP		•
·	(2.92)		0	rgran
Rice and Millet Spirals	2.00	250 g OP		•
	(3.11)		0	rgran
Rice and corn spaghetti noodles	2.00	375 g OP		•
	(2.92)		0	rgran
Vegetable and Rice Spirals	2.00	250 g OP		•
	(2.92)	_	0	rgran
Italian long style spaghetti	2.00	220 g OP		-
- · · · ·	(3.11)	-	0	rgran
				=

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic S Per ✓ Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE – Special Authority see SA1108 on the previous page – Hospital pharmacy [HP3]

annacy [rif5]			
Tabs	99.00	75 OP	✓ Phlexy 10
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex
			Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior
			Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex
			Powder
Powder (neutral) 36 g sachets	393.00	30	PKU Anamix Junior
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex
			Powder
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior
(0 / 0			Orange
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior
· · · · · · · · · · · · · · · · · · ·			Vanilla
Infant formula	174 72	400 g OP	✓ PKU Anamix Infant
Powder (orange)		500 g OP	✓ XP Maxamum
Powder (unflavoured)		500 g OP	✓ XP Maxamum
Liquid (berry)		125 ml OP	✓ PKU Anamix Junior
Elquid (5511y)		120 1111 01	LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior
1 (3-)			LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior
,			LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex
1 1 (11 11)	,		Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml		30 OP	✓ PKU Lophlex LQ 20
qa.a (ja.a) a.aga,a		00 01	zepinok za zo

Foods

LOW PROTEIN PASTA - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne		250 g OP	✓ Loprofin
Low protein rice pasta	11.91	500 g OP	✓ Loprofin
Macaroni		250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals	11.91	500 a OP	✓ Loprofin

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA - Special Authority see SA2092 below - Hospital phar	macy [HP3]	
Powder	400 g OP	✓ Alfamino✓ Alfamino Junior
Powder (unflavoured)53.00	400 g OP	✓ Elecare ✓ Elecare LCP ✓ Neocate Gold ✓ Neocate Junior Unflavoured
Powder (vanilla)53.00	400 g OP	✓ Neocate SYNEO✓ Elecare✓ Neocate Junior✓ Vanilla

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis: or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency: or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA - Special Auth	nority see SA1953 below -	Hospital pharr	nacy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption: or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

SPECIAL FOODS

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	✓	

continued...

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
- 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms: or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption: or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea: or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

 Powder (unflavoured)
 35.50
 300 g OP

 KetoCal 4:1

 Powder (vanilla)
 35.50
 300 g OP

 KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent.................0.00 10 ✓ BCG Vaccine

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
 - 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old; or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 - 9 above.

pertussis toxoid. 8 mcg pertussis filamentous

10 **Boostrix Boostrix**

✓ fully subsidised

Principal Supply

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	Fully Brand or ised Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE	- [Xpharm]		
Funded for any of the following:	., .		
 A single dose for children up to the age of 7 who have A course of four vaccines is funded for catch up progra primary immunisation; or 			
 An additional four doses (as appropriate) are funded fo pre- or post splenectomy; pre- or post solid organ trans regimens; or 			
4) Five doses will be funded for children requiring solid or	gan transplantation.		
Note: Please refer to the Immunisation Handbook for approp	oriate schedule for cat	ch up progr	ammes.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units	0.00	40	A la familia IDV
poliomyelitis virus in 0.5ml syringe		10	✓ <u>Infanrix IPV</u>
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B A	IND HAEMOPHILUS I	NFLUENZA	LE TYPE B VACCINE -
[Xpharm]			
Funded for patients meeting any of the following criteria: 1) Up to four doses for children up to and under the age of 2) An additional four doses (as appropriate) are funded for 10 who are patients post haematopoietic stem cell tran post solid organ transplant, renal dialysis and other set 3) Up to five doses for children up to and under the age of Note: A course of up-to four vaccines is funded for catch up to complete full primary immunisation. Please refer to the Improgrammes. Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus, 10 mcg hepatitis B surface antigen in 0.5 ml syringe HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm] One dose for patients meeting any of the following: 1) For primary vaccination in children; or 2) An additional dose (as appropriate) is funded for (re-)in transplantation, or chemotherapy; functional asplenic; por post cochlear implants, renal dialysis and other seven	r (re-)immunisation for splantation, or chemoty splantation, or chemoty rerely immunosuppres of 10 receiving solid organization Handbook munisation Handbook	r children up therapy; pre sive regime gan transpla ren (up to a c for the app 10 10 ts post haer ny; pre- or p	e or post splenectomy; pre- or ens; or antation. Ind under the age of 10 years) propriate schedule for catch up Infanrix-hexa matopoietic stem cell lost solid organ transplant, pre-
 For use in testing for primary immunodeficiency diseas paediatrician. 	es, on the recommend	lation of an	internal medicine physician or
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mc prefilled syringe plus vial 0.5 ml		1	✓ Hiberix
HEPATITIS A VACCINE – [Xpharm] Funded for patients meeting any of the following criteria: 1) Two vaccinations for use in transplant patients; or 2) Two vaccinations for use in children with chronic liver of 3) One dose of vaccine for close contacts of known hepat			
Inj 1440 ELISA units in 1 ml syringe Inj 720 ELISA units in 0.5 ml syringe		1	✓ <u>Havrix</u> ✓ <u>Havrix Junior</u>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B RECOMBINANT VACCINE – [Xpharm] Inj 10 mcg per 0.5 ml prefilled syringe Funded for patients meeting any of the following crite		1	√ E	ngerix-B
 for household or sexual contacts of known acut for children born to mothers who are hepatitis B for children up to and under the age of 18 years serology and require additional vaccination or reference for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual int for patients following immunosuppression; or for solid organ transplant patients; or for post-haematopoietic stem cell transplant (HS) following needle stick injury. 	s surface antigen (HBsAgs s inclusive who are consic equire a primary course of ercourse; or) pos derec	itive; or I not to have	
Inj 20 mcg per 1 ml prefilled syringe Funded for patients meeting any of the following crite		1	✓ <u>E</u>	ngerix-B
 for household or sexual contacts of known acut for children born to mothers who are hepatitis B for children up to and under the age of 18 years serology and require additional vaccination or reference of the household of	s surface antigen (HBsAgs s inclusive who are consic equire a primary course of tercourse; or) pos derec	itive; or I not to have	
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AN Any of the following:	,	- [Xpl	narm]	
1) Maximum of two doses for children aged 14 years a 2) Maximum of three doses for patients meeting any o 1) People aged 15 to 26 years inclusive; or 2) Either: People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or 2) Transplant (including stem cell) patients: 3) Maximum of four doses for people aged 9 to 26 years.	f the following criteria: or	nerap	у	
Inj 270 mcg in 0.5 ml syringe	0.00	10	√ <u>G</u>	ardasil 9

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer
INELLIENZA VACCINE			

INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

✓ Afluria Quad Junior - [Xpharm]......11.00 (2022 formulation)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease; or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness:

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	110.00 10	•	Afluria Quad
			(2022 formulation)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- a) Only on a prescription
- b) No patient co-payment payable

С

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) People 55 to 64 years of age (inclusive) and is Māori or any Pacific ethnicity; or
- c) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- d) children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- e) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or
- f) children 3 to 12 years of age (inclusive), from 1 July 2022 to 31 December 2022;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	•	Manufacturer

MEASUES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression: or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Either:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients: or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Ini 10 mcg of each meningococcal polysaccharide conjugated

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm] Either: A) Both: 1) Child is under one year of age; and 2) Any of the following: i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant: or ii) up to three doses for close contacts of meningococcal cases of any group; or iii) up to three doses for child who has previously had meningococcal disease of any group; or iv) up to three doses for bone marrow transplant patients; or v) up to three doses for child pre- and post-immunosuppression*; or B) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients; or v) up to two doses for person pre- and post-immunosuppression*. *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 12 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV. complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression*. Note: children under 12 months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine. *Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Neisvac-C 1 PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm] 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B.

7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

prefilled syringe0.00

✓ Synflorix

10

			_
Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
- 2) Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection; or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) cochlear implants or intracranial shunts: or
 - g) cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - I) diabetes; or
 - m) Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note:	please refer to the	Immunisation	Handbook for	the appropri	iate schedule	for catch u	up prod	irammes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

0, 0. 1, 0.2,, 0.1,, 100, 10. 1, 10. 11. 11. 11. 11. 11. 11. 11. 11. 11.		
syringe0.00	10	Prevenar 13
	1	✓ Prevenar 13

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - Either:	- [Xpharm]			
 Up to three doses (as appropriate) for patients with H chemotherapy; pre- or post-splenectomy or with func complement deficiency (acquired or inherited), cochle All of the following: 	tional asplenia, pre- or p	oost-s	solid organ t	ransplant, renal dialysis,
a) Patient is a child under 18 years for (re-)immunib) Treatment is for a maximum of two doses; andc) Any of the following:	isation; and			
 i) on immunosuppressive therapy or radiatic immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or 	n therapy, vaccinate wh	nen th	nere is expe	cted to be a sufficient
iv) with renal failure, or nephrotic syndrome;v) who are immune-suppressed following orgor	gan transplantation (incl	luding	j haematopo	pietic stem cell transplant);
 vi) with cochlear implants or intracranial shur vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more to prednisone of 2 mg/kg per day or greater, 20 mg or greater; or 	han two weeks, and wh			, ,
ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks ge xi) with cardiac disease, with cyanosis or failu xii) with diabetes; or xiii) with Down syndrome; or	station; or	gh-do	se corticost	eroid therapy); or
xiv) who are pre-or post-splenectomy, or with	functional asplenia.			
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	0.00	1	√ <u>P</u>	neumovax 23
POLIOMYELITIS VACCINE – [Xpharm] Up to three doses for patients meeting either of the followir 1) For partially vaccinated or previously unvaccinated in	ŭ			
For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for appr Inj 80D antigen units in 0.5 ml syringe	•	tch-up	o programm	
ROTAVIRUS ORAL VACCINE – [Xpharm] Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 14	weeks of age; and			
2) no vaccination being administered to children aged 2	4 weeks or over.			
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	√ F	Rotarix

NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price) \$	Subsid Per	lised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm] Either:				
1) Maximum of one dose for primary vaccination for eith	ner:			
 a) Any infant born on or after 1 April 2016; or b) For previously unvaccinated children turning 11 varicella infection (chickenpox), or 	years old on or after 1	July 2017, v	who hav	re not previously had a
2) Maximum of two doses for any of the following:				
a) Any of the following for non-immune patients:				
 i) with chronic liver disease who may in futu ii) with deteriorating renal function before tra iii) prior to solid organ transplant; or iv) prior to any elective immunosuppression* v) for post exposure prophylaxis who are im 	nsplantation; or , or mune competent inpatic	ents.; or		
b) For patients at least 2 years after bone marrow				
c) For patients at least 6 months after completion				
d) For HIV positive non immune to varicella with n				
 e) For patients with inborn errors of metabolism at varicella, or 	risk of major metabolic	aecompen	sation, v	vitn no clinical history of
f) For household contacts of paediatric patients w immune compromise where the household con	tact has no clinical histo	ory of varice	lla, or	
g) For household contacts of adult patients who he immunocompromised, or undergoing a procedule has no clinical history of varicella.				
* immunosuppression due to steroid or other immunosupp	ressive therapy must be	e for a treatr	nent pe	riod of greater than
28 days			·	•
Inj 1350 PFU prefilled syringe	0.00	1 10	✓ <u>Val</u> ✓ <u>Val</u>	
VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] - [X	oharm]			
Funded for patients meeting the following criteria:				
 Two doses for all people aged 65 years 				
Inj 50 mcg per 0.5 ml vial plus vial	0.00	1	✓ Shi	ingrix
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUA Funded for patients meeting the following criteria:	TED VACCINE [SHING	LES VACCI	INE] –[Xpharm]
1) One dose for all people aged 65 years				
Inj 19,400 PFU prefilled syringe plus vial	0.00	1 10		stavax stavax
Diagnostic Agents				
TUDEDCUI IN DDD (MANITOLIV) TECT (Vek)				
TUBERCULIN PPD [MANTOUX] TEST - [Xpharm] Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	✓ <u>Tul</u>	<u>persol</u>

- Symbols -	Agents for Parkinsonism and Related	Amoxicillin93
3TC106	Disorders 121	Amoxicillin with clavulanic acid93
7 MED NSHA Silver/Copper	Agents Used in the Treatment of	Amphotericin B32
Short73	Poisonings254	Amsacrine154
- A -	Agrylin154	AmsaLyo154
A-Scabies68	Albendazole90	Amsidine154
Abacavir sulphate106	Albey240–241	Amzoate30
Abacavir sulphate with	Albustix77	Anaesthetics122
lamivudine 106	Alchemy163	Anagrelide hydrochloride154
Abacavir/Lamivudine Viatris106	Alchemy Oxybutynin76	Analgesics 123
Abiraterone acetate169	Aldurazyme29	Anastrozole172
Acarbose11	Alecensa162	Anatrole172
Accarb11	Alectinib162	Andriol Testocaps81
Accuretic 1047	Alendronate sodium111	Androderm81
Accuretic 2047	Alendronate sodium with	Anoro Ellipta245
Acetazolamide251	colecalciferol111	Antabuse146
Acetec46	Alfacalcidol33	Antacids and Antiflatulents6
Acetic acid with hydroxyquinoline and	Alfamino275	Anthelmintics90
ricinoleic acid75	Alfamino Junior275	Antiacne Preparations61
Acetylcysteine254	Alginic acid6	Antiallergy Preparations240
Aci-Jel75	Alglucosidase alfa26	Antianaemics36
Aciclovir	Alkeran 151	Antiandrogen Oral
Infection101	Alkeran S29151	Contraceptives75
Sensory249	Allerfix251	Antiarrhythmics48
Acidex6	Allerpro Syneo 1278	Antibacterials90
Acipimox54	Allerpro Syneo 2278	Antibacterials Topical61
Acitretin68	Allersoothe241	Anticholinergic Agents244
Aclasta114	Allmercap	Anticholinesterases110
Actemra224	Allopurinol119	Antidepressants127
Actemra S29224	Alpha-Adrenoceptor Blockers46	Antidiarrhoeals6
Actinomycin D155	Alpha-Keri Lotion66	Antiepilepsy Drugs129
Actrapid10	Alphamox93	Antifibrinolytics, Haemostatics and
Actrapid Penfill10	Alphamox 12593	Local Sclerosants37
Acupan 123	Alphamox 25093	Antifibrotics245
Adalimumab (Amgevita)179	Alprolix37	Antifungals97
Adalimumab (Humira - Alternative	Alu-Tab6	Antifungals Topical62
brand) 188	Aluminium hydroxide6	Antihistamines241
Adapalene61	Alvogen49	Antihypotensives49
ADR Cartridge 1.823	Amantadine hydrochloride121	Antimalarials99
Adrenaline	Ambrisentan57	Antimigraine Preparations132
Cardiovascular56	Ambrisentan Mylan57	Antinausea and Vertigo Agents 133
Respiratory240	Ambrisentan Viatris57	Antipruritic Preparations63
Advantan64	Amgevita179	Antipsychotics134
Advate39	Amiloride hydrochloride52	Antiretrovirals104
Adynovate40	Amiloride hydrochloride with	Antirheumatoid Agents111
Afinitor236	furosemide52	Antispasmodics and Other Agents
Aflibercept194	Amiloride hydrochloride with	Altering Gut Motility8
Afluria Quad	hydrochlorothiazide52	Antithrombotic Agents40
(2022 formulation)	Aminophylline246	Antithymocyte globulin
Afluria Quad Junior	Amiodarone hydrochloride48	(equine) 179
(2022 formulation) 283	Amisulpride134	Antitrichomonal Agents99
AFT-Pyrazinamide101	Amitriptyline127	Antituberculotics and
Agents Affecting the	Amlodipine50	Antileprotics100
Renin-Angiotensin System 46	Amorolfine62	Antiulcerants9

Antivirals	101	Atomoxetine	141	Beta Cream	6
Anxiolytics		Atorvastatin		Beta Ointment	
Anzatax		Atropine sulphate		Beta Scalp	
Apidra		Cardiovascular	48	Beta-Adrenoceptor Agonists	
Apidra SoloStar		Sensory		Beta-Adrenoceptor Blockers	
APO-Atomoxetine		Atropt		Betadine	
APO-Atomoxetine S29		Atrovent		Betadine Skin Prep	
Apo-Azithromycin		AU Synacthen		Betaferon	
Apo-Diltiazem CD		Aubagio		Betahistine dihydrochloride	
Apo-Temozolomide	159	Augmentin		Betaine	. 2
Apomorphine hydrochloride		Aurorix		Betaloc CR	
Aprepitant		AutoSoft 30		Betamethasone dipropionate	
Apresoline		AutoSoft 90		Betamethasone dipropionate with	
Aqueous cream		Avallon		calcipotriol	6
Aratac		Avelox		Betamethasone sodium phosphate	
Arava		Avonex		with betamethasone acetate	79
Arginine		Avonex Pen		Betamethasone valerate	
Aripiprazole		Azacitidine		Betamethasone valerate with sodiur	,
Aripiprazole Sandoz		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Aristocort		Azamun		Betaxolol	
Arrotex-Prazosin S29		Azathioprine		Betnovate	
Arrow - Clopid		Azilect		Betoptic	
Arrow - Lattim		Azithromycin		Betoptic S	
Arrow-Amitriptyline		Azopt		Bexsero	
Arrow-Bendrofluazide		AZT		Bezafibrate	
Arrow-Brimonidine		- B -	100	Bezalip	
Arrow-Diazepam		B-D Micro-Fine	15	Bezalip Retard	
Arrow-Doxorubicin		B-D Ultra Fine		Bicalutamide	
Arrow-Fluoxetine		B-D Ultra Fine II		Bicillin LA	
Arrow-Losartan &	123	Bacillus Calmette-Guerin (BCG)	10	BiCNU	
Hydrochlorothiazide	47	vaccine	170	Bile and Liver Therapy	
Arrow-Norfloxacin		Bacillus Calmette-Guerin	173	Biltricide	
Arrow-Ornidazole		vaccine	200	Bimatoprost	
Arrow-Quinapril 10		Baclofen		Bimatoprost Multichem	
		Bactroban		Binarex	
Arrow Quinapril 5				Binocrit	
Arrow Povithromyoin		Barrier Creams and Emollients		Biodone	
Arrow-Roxithromycin					
Arrow-Timolol		BCG Vaccine		Biodone Extra Forte	
Arrow-Topiramate		Beclazone 100		Biodone Forte	
Arrow-Tramadol		Beclazone 250		Bisacodyl	21
Arsenic trioxide		Beclazone 50		Bisacodyl Viatris	
Asacol		Beclomethasone dipropionate		Bisoprolol fumarate	
Ascorbic acid		Bee venom allergy treatment		Bisoprolol Mylan	
Aspen Adrenaline	50	Bendamustine hydrochloride		Bisoprolol Viatris	
Aspirin	40	Bendrofluazide	53	BK Lotion	
Blood		Bendroflumethiazide		Bleomycin sulphate	. 15
Nervous		[Bendrofluazide]		Blood Colony-stimulating	
Asthalin		Benralizumab		Factors	4
Atazanavir Mylan		Benzathine benzylpenicillin		Blood glucose diagnostic test	
Atazanavir sulphate		Benzatropine mesylate	121	meter	14
Atenolol		Benzbromaron AL 100		Blood glucose diagnostic test	
Atenolol AFT		Benzbromarone		strip	1
Atenolol AFT S29		Benztrop		Blood glucose test strips (visually	٠.
ATGAM		Benzydamine hydrochloride		impaired)	1
Ativan		Benzylpenicillin sodium [Penicillin		Blood Ketone Diagnostic Test	
Atnahs Olsalazine	8	G]	93	Strip	1

Bonjela32	Calogen	261	Chickenpox vaccine	28
Boostrix280	Candesartan cilexetil	47	Chlorafast	24
Bortezomib 155	Candestar	47	Chlorambucil	15
Bortezomib Dr-Reddy's155	Canesten	62	Chloramphenicol	24
Bosentan57	Capecitabine	152	Chlorothiazide	
Bosentan Dr Reddy's57	Capercit	152	Chlorpheniramine maleate	24
Bplex33	Capoten		Chlorpromazine hydrochloride	13
Breo Ellipta243	Capsaicin		Chlortalidone [Chlorthalidone]	5
Brevinor 1/2874	Musculoskeletal	111	Chlorthalidone	5
Bricanyl Turbuhaler243	Nervous	123	Chlorvescent	4
Brilinta40	Captopril	46	Choice 380 7med Nsha Silver/cop	
Brimonidine tartrate252	Carafate		Short	7
Brimonidine tartrate with timolol	Carbaccord		Choice Load 375	7
maleate252	Carbamazepine	130	Choice TT380 Short	7
Brinzolamide251	Carbimazole	83	Choice TT380 Standard	7
Brolene249	Carbomer		Choline salicylate with cetalkoniur	m
Brufen SR110	Carboplatin		chloride	
BSF Alchemy254	Carboplatin Ebewe		Ciclosporin	
Buccastem134	Carbosorb-X		Cilazapril	
Budesonide	Cardinol LA		Cilicaine VK	
Alimentary6	Cardizem CD	51	Cinacalcet	
Respiratory242, 247	CareSens Dual	14	Cipflox	9
Budesonide with eformoterol243	CareSens N		Ciprofloxacin	
Bumetanide52	CareSens N POP		Infection	9
Buprenorphine Naloxone BNM145	CareSens N Premier	14	Sensory	
Buprenorphine with naloxone145	CareSens PRO	15	Ciprofloxacin Teva	24
Bupropion hydrochloride146	Carmellose sodium with gelati		Cisplatin	15
Burel98	pectin		Cisplatin Ebewe	
Burinex52	Carmustine		Citalopram hydrobromide	
Burinex S2952	Carnitor	29	Cladribine	
Buscopan8	Carvedilol	49	Clarithromycin	
Buspirone hydrochloride137	Carvedilol Sandoz	49	Alimentary	
Buspirone Viatris137	Casirivimab and imdevimab	196	Infection	9
Busulfan150	Catapres	51	Clexane	4
- C -	CeeNU		Clexane Forte	4
Cabergoline88	Cefaclor monohydrate		Climara	
Caffeine citrate248	Cefalexin		Clindamycin	
Calamine63	Cefazolin	90	Clinicians	2
Calamine-AFT63	Ceftriaxone		Clinicians Renal Vit	3
Calci-Tab 50034	Ceftriaxone-AFT	91	Clobazam	
Calcipotriol68	Cefuroxime axetil	91	Clobetasol propionate	.63, 7
Calcitonin78	Celapram	128	Clobetasone butyrate	
Calcitriol33	Celebrex		Clofazimine	
Calcitriol-AFT33	Celecoxib	110	Clomazol	
Calcium 500 mg Hexal34	Celecoxib Pfizer	110	Dermatological	6
Calcium carbonate6, 34	Celestone Chronodose	79	Genito-Urinary	<mark>7</mark>
Calcium Channel Blockers50	Cellcept		Clomifene citrate	
Calcium Disodium Versenate255	Centrally-Acting Agents		Clomipramine hydrochloride	12
Calcium folinate152	Cephalexin ABM		Clomipramine Teva	
Calcium Folinate Ebewe152	Cetirizine hydrochloride		Clonazepam13	
Calcium Folinate Sandoz152	Cetomacrogol		Clonidine	
Calcium Folinate Sandoz S29 152	Cetomacrogol with glycerol		Clonidine hydrochloride	
Calcium gluconate34	Cetomacrogol-AFT	65	Clonidine Teva	
Calcium Homeostasis	Cetuximab		Clopidogrel	
Calcium polystyrene sulphonate45	Charcoal		Clopidogrel Multichem	
Calcium Resonium45	Chemotherapeutic Agents		Clopine	

Clopixol13	6-137	Curam	93	DBL Heparin Sodium	43
Clotrimazole		Curam Duo 500/125	93	DBL Leucovorin Calcium	152
Dermatological	62	Cvite	33	DBL Methotrexate Onco-Vial	153
Genito-Urinary		Cyclizine hydrochloride	133	DBL Morphine Sulphate	126
Clozapine		Cyclizine lactate		DBL Pethidine Hydrochloride	
Clozaril		Cyclogyl	252	DBL Vincristine Sulfate	
Co-trimoxazole		Cyclonex		Decozol	
Coal tar		Cyclopentolate hydrochloride		Deferasirox	
Coal tar with allantoin, menthol,		Cyclophosphamide		Deferiprone	
phenol and sulphur	69	Cyclorin		Deferoxamine Pfizer S29	
Coal tar with salicylic acid and		Cycloserine		Denosumab	
sulphur	69	Cyproterone acetate		Deolate	
Coco-Scalp		Cyproterone acetate with		Deoxycoformycin	
Codeine phosphate	09	,,	75		
Extemporaneous	257	ethinyloestradiol		Depo-Medrol	7،
		Cystadane		Depo-Provera	/4
Nervous		Cytarabine		Depo-Testosterone	0
Coenzyme Q10		Cytotec		Deprim	
Colchicine		Cytoxan	150	Dermol	
Colecalciferol		-D-		Desferrioxamine mesilate	
Colestid		D-Penamine		Desmopressin	
Colestipol hydrochloride		Dabigatran		Desmopressin acetate	
Colgout		Dacarbazine		Desmopressin-PH&T	
Colifoam		Dacarbazine APP		Desuric	119
Colistin sulphomethate	95	Dactinomycin [Actinomycin D].		Detection of Substances in	
Colistin-Link		Daivobet	68	Urine	77
Collodion flexible	257	Daivonex	68	Dexamethasone	
Colloidal bismuth subcitrate	9	Daktarin	63	Hormone	79
Colofac	8	Dalacin C	94	Sensory	250
Coloxyl	25	Dantrium	120	Dexamethasone phosphate	80
Combigan	252	Dantrium S29	120	Dexamethasone with framycetin	and
Compound electrolytes	45	Dantrolene	120	gramicidin	249
Compound electrolytes with glucos	se	Daonil	11	Dexamethasone with neomycin	
[Dextrose]		Dapa-Tabs	53	sulphate and polymyxin B	
Compound hydroxybenzoate		Dapsone		sulphate	250
Comtan		Daraprim		Dexamfetamine sulfate	
Concerta		Darunavir		Dexmethsone	
Condoms		Darunavir Mylan		Dextrochlorpheniramine	
Condyline		Darunavir Viatris		maleate	241
Contraceptives - Hormonal		Dasatinib		Dextrose	
Contraceptives - Non-hormonal		Daunorubicin		DHC Continus	
Copaxone		Daunorubicin Zentiva		Diabetes	
Cordarone-X		David One Step Cassette Preg		Diabetes Management	
Corticosteroids and Related Agent		Test	•	Diacomit	
•		DBL Adrenaline		Diagnostic Agents	
for Systemic Use					
Corticosteroids Topical		DBL Aminophylline		Diamide Relief	
Cortifoam		DBL Bleomycin Sulfate		Diamox	
Cosentyx		DBL Bortezomib		Diasip	262
Cosmegen		DBL Carboplatin		Diazepam1	29, 137
Coumadin		DBL Cisplatin		Diazoxide	
Country Life		DBL Dacarbazine		Dibenzyline	46
Coversyl	46	DBL Desferrioxamine Mesylate		Diclofenac Sandoz	110
Creon 10000		BP		Diclofenac sodium	
Creon 25000		DBL Docetaxel		Musculoskeletal	
Creon Micro		DBL Ergometrine	75	Sensory	250
Crotamiton	63	DBL Gemcitabine		Differin	
Crystaderm	61	DBL Gentamicin	95	Difflam	32

Diflucan97	Drofate	50	Emulsifying Ointment ADE	6
Digestives Including Enzymes24	Drugs Affecting Bone		Enalapril maleate	4
Digoxin48	Metabolism	111	Enbrel	17
Dihydrocodeine tartrate125		one	Endocrine Therapy	16
Dilantin131	diagnostic test meter	14	Endoxan	
Dilantin Infatab131	Dulaglutide	12	Engerix-B	28
Diltiazem CD Clinect51	Dulcolax SP Drop	26	Enlafax XR	12
Diltiazem hydrochloride51	Duocal Super Soluble Powder		Enoxaparin sodium	
Dimethicone65–66			Enstilar	
Dimethyl fumarate138	Duolin HFA	244	Ensure	
Dipentum8		243	Ensure Plus	27
Diphtheria, tetanus and pertussis	Duride		Ensure Plus HN	
vaccine280			Ensure Plus RTH	26
Diphtheria, tetanus, pertussis and	- E -		Ensure Two Cal HN RTH	
polio vaccine	e-chamber La Grande	248	Entacapone	
Diphtheria, tetanus, pertussis, polio,	e-chamber Mask		Entecavir	
hepatitis B and haemophilus	e-chamber Turbo		Entecavir Mylan	
influenzae type B vaccine 281			Entecavir Sandoz	
Diprosone63			Entocort CIR	
Diprosone OV63			Entresto 24/26	
Dipyridamole40			Entresto 49/51	4
			Entresto 97/103	
Disopyramide phosphate48 Disulfiram146			Entyvio	
Diuretics			•	
Dizole		100	Epilim Epilim Crushable	
		100		
Docetaxel 155	•		Epilim IV	
Docetaxel Accord			Epilim S/F Liquid	
Docetaxel Sandoz	•	242	Epilim Syrup	
Docusate sodium25		07	Epipen	
Docusate sodium with	factor IX]		Epipen Jr	
sennosides25			Epirubicin Ebewe	15
Dolutegravir107			Epirubicin hydrochloride	15
Domperidone			Eplerenone	5
Domperidone Viatris			Epoetin alfa	
Donepezil hydrochloride144			Epoprostenol	5
Donepezil-Rex144			Eptacog alfa [Recombinant factor	
Dornase alfa246			VIIa]	
Dortimopt251	Elelyso		Erbitux	
Dorzolamide hydrochloride251	Elemental 028 Extra		Ergometrine maleate	
Dorzolamide with timolol251	Elidel		Erlotinib	
Dostinex88			Erythrocin IV	
Dosulepin [Dothiepin]	Elocon Alcohol Free		Erythromycin (as lactobionate)	
hydrochloride 128			Erythromycin ethyl succinate	9
Dosulepin Mylan128	Eltroxin	83	Esbriet	
Dosulepin Viatris128	EMB Fatol		Escitalopram	
Dothiepin128	Emend Tri-Pack	133	Escitalopram (Ethics)	12
Doxazosin46	Emicizumab	38	Eskazole	
Doxazosin Clinect46	EMLA	123	Essential Ethosuximide	
Doxine94	Empagliflozin	13	Essential Generics	4
Doxorubicin Ebewe155			Essential Prednisolone	
Doxorubicin hydrochloride 155		13	Estraderm MX	8
Doxycycline94			Estradiol TDP Mylan	
DP Lotion66			Estradot	
DP Lotn HC64		103	Estradot 50 mcg	
DP-Allopurinol119			Estrofem	
Dr Reddy's Omeprazole9		65	Etanercept	
·				

Ethambutol hydrochloride		Ferrograd		Folic acid	
Ethics Aspirin		Ferrosig		Folic Acid multichem	
Ethics Aspirin EC		Ferrous fumarate		Folic Acid Mylan	
Ethics Lisinopril	46	Ferrous fumarate with folic acid		Food Thickeners	
Ethinyloestradiol with		Ferrous sulfate		Foods And Supplements For I	
desogestrel	73	Fexofenadine hydrochloride	241	Errors Of Metabolism	
Ethinyloestradiol with		Fibro-vein	40	Foradil	
levonorgestrel	74	Filgrastim	43	Forteo	113
Ethinyloestradiol with		Finasteride		Fortini	26 ²
norethisterone	74	Fingolimod	138	Fortini Multi Fibre	
Ethosuximide	130	Firazyr		Fortisip	
Etopophos	156	Flagyl	99	Fortisip Multi Fibre	
Etoposide	156	Flagyl-S		Fosamax	
Etoposide phosphate	156	Flamazine	62	Fosamax Plus	
Etravirine	106	Flecainide acetate	48	Framycetin sulphate	249
Eumovate	64	Flecainide BNM	48	Frebini Energy	26 ²
Evara	65	Flecainide Controlled Release		Frebini Energy Fibre	26 ²
Everet	131	Teva	48	Frebini Original	26 ²
Everolimus	236	Fleet Phosphate Enema	26	Frebini Original Fibre	
Evista	112	Flixonase Hayfever & Allergy	247	Fresubin 2kcal HP	
Evusheld	224	Flixotide		Fresubin HP Energy	269
Exemestane	172	Flixotide Accuhaler	242	Fresubin HP Energy Fibre	
Exjade		Florinef		Fresubin Intensive	
Extemporaneously Compound		Fluanxol		Fresubin Original	
Preparations and		Flucil	93	Fresubin Original Fibre	
Galenicals	257	Flucloxacillin	93	Frisium	
Eye Preparations		Flucloxacillin-AFT		Frumil	
Eylea		Flucloxin		Frusemide	
Ezetimibe		Flucon		Fucicort	
Ezetimibe Sandoz		Fluconazole		Fucidin	96
Ezetimibe with simvastatin		Fludara Oral	152	Fucithalmic	249
-F-		Fludarabine Ebewe	152	Fulvestrant	
Factor eight inhibitor bypassin	a	Fludarabine phosphate	152	Fungilin	
fraction		Fludrocortisone acetate		Furosemid-Ratiopharm	
Famotidine		Fluids and Electrolytes		Furosemide [Frusemide]	
Famotidine Hovid		Flumetasone pivalate		Furosemide-Baxter	
Fasenra		Fluocortolone caproate with		fusidic acid	
Faslodex		fluocortolone pivalate and		Dermatological	62.64
Fatty Cream AFT		cinchocaine	8	Infection	
Febuxostat		Fluorometholone		Sensory	
Febuxostat multichem		Fluorouracil		- G -	
Feed Thickener Karicare		Fluorouracil Accord		Gabapentin	130
Aptamil	272	Fluorouracil sodium		Gacet	
FEIBA NF		Fluox		Galsulfase	
Felo 10 ER		Fluoxetine hydrochloride		Galvumet	
Felo 5 ER		Flupenthixol decanoate		Galvus	
Felodipine		Flutamide		Gardasil 9	
Femme-Tab ED		Flutamin		Gastrodenol	
Fenpaed 100 mg per 5 ml		Fluticasone		Gaviscon Double Strength	
Fentanyl		Fluticasone furoate with	٢٦٢	Gaviscon Infant	
Fentanyl Sandoz		vilanterol	243	Gazyva	
Ferinject		Fluticasone propionate		Gefitinib	
Ferodan		Fluticasone with salmeterol		GEM Aqueous Cream	
Ferriprox		Flynn		Gemcitabine Ebewe	
Ferro-F-Tabs		FML		Gemcitabine hydrochloride	
Ferro-tab		Foban		Gemtuzumab ozogamicin	
ι στιστασ		ı uvalı	∪∠	Gorniuzumas Uzugamicili	130

Genoptic	249	Hemlibra	38	Hylo-Fresh	253
Gentamicin sulphate		Heparin sodium	43	Hymenoptera	240-241
Infection	95	Heparin Sodium Panpharma	43	Hyoscine butylbromide	8
Sensory	249	Heparinised saline		Hyoscine hydrobromide	
Gilenya		Heparon Junior		Hypam	
Ginet		Hepatitis A vaccine		Hyperuricaemia and Antigout	
Glatiramer acetate		Hepatitis B recombinant		Hypromellose	
Glecaprevir with pibrentasvir		vaccine	282	Hypromellose with dextran	
Glibenclamide		Herceptin		-1-	
Gliclazide		Hiberix		lbiamox	93
Glipizide		Hiprex		Ibrance	
Glivec		Histaclear		Ibrutinib	
Glizide		Histafen		Ibuprofen	
Glucagen Hypokit		Holoxan		Icatibant	
Glucagon hydrochloride		Horleys Bread Mix		Idarubicin hydrochloride	
Glucerna Select		Horleys Flour		Idursulfase	
Glucose [Dextrose]				Ifosfamide	
		Hormone Replacement Therapy			
Gluten Free Foods		Systemic		Igroton	
Glycerin with sodium saccharia		HPV		lkorel	
Glycerin with sucrose	207	Humalog		lloprost	
Glycerol	00	Humalog Mix 25		Imatinib mesilate	
Alimentary		Humalog Mix 50		Imatinib-Rex	
Extemporaneous	257	Human papillomavirus (6, 11, 16		Imbruvica	
Glyceryl trinitrate	•	31, 33, 45, 52 and 58) vaccin		Imfinzi	
Alimentary		[HPV]		Imigran	
Cardiovascular		Humatin		Imipramine hydrochloride	
Glycopyrronium		Humira		Imiquimod	
Glycopyrronium bromide	8	HumiraPen		Immune Modulators	
Glycopyrronium with		Humulin 30/70		Immunosuppressants	
indacaterol		Humulin NPH		Incruse Ellipta	
Go Healthy		Humulin R		Indacaterol	
Gold Knight	72	Hyaluronic acid	253	Indapamide	53
Gold Knight XL	72–73	Hydralazine	56	Infanrix IPV	281
Goserelin		Hydralazine hydrochloride	56	Infanrix-hexa	281
Gutron	49	Hydrocortisone		Infant Formulae	275
Gynaecological Anti-infectives	75	Dermatological	64	Infatrini	279
- H -		Hormone	80	Infliximab	197
Habitrol		Hydrocortisone (PSM)	64	Influenza vaccine	283
Haemophilus influenzae type E	3	Hydrocortisone acetate	7	Inhaled Corticosteroids	242
vaccine		Hydrocortisone acetate with		Inhaled Long-acting	
Haldol	136	pramoxine hydrochloride	7	Beta-adrenoceptor Agonists.	242
Haldol Concentrate	136	Hydrocortisone and paraffin liqu	iid	Inspra	52
Haldol Decanoas	136	and lanolin	64	Instillagel Lido	122
Haloperidol	135	Hydrocortisone butyrate		Insulin aspart	
Haloperidol decanoate		Hydrocortisone with cinchocaine		Insulin aspart with insulin aspar	
Harvoni		Hydrocortisone with miconazole		protamine	
Havrix	281	Hydrocortisone with natamycin a		Insulin glargine	
Havrix Junior	281	neomycin		Insulin glulisine	
Haylor syrup		Hydrogen peroxide		Insulin isophane	
healthE Dimethicone 10%		Hydroxocobalamin		Insulin isophane with insulin	
healthE Dimethicone 4% Lotio		Hydroxocobalamin Panpharma.		neutral	11
healthE Dimethicone 5%		hydroxycarbamide		Insulin lispro	
healthE Glycerol BP		Hydroxychloroguine		Insulin lispro with insulin lispro	
healthE Urea Cream		Hydroxyurea	111	protamine	11
Healtheries Simple Baking Mix		[hydroxycarbamide]	156	Insulin neutral	11 10
Hemastix		Hygroton		Insulin pen needles	
1 IO11103UA	/ /	1 1yg101011		modili pen needles	13

Insulin pump	16	Ivermectin	66	Lanoxin S29	48
Insulin pump cartridge	21	- J -		Lansoprazole	9
Insulin pump infusion set (steel		Jadelle	74	Lantus	11
cannula)	21	Jakavi		Lantus SoloStar	
Insulin pump infusion set (steel		Jardiamet		Lanvis	
cannula, straight insertion)	21	Jardiance		Lanzol Relief	
Insulin pump infusion set (teflon	21	Jaydess		Lapatinib ditosylate	
cannula)	22	Jevity HiCal RTH		Largactil	13/
Insulin pump infusion set (teflon	22			Laronidase	
		Jevity Plus Jevity RTH	209	Lasix	
cannula, angle insertion with	00				
insertion device)	23	Jinarc		Latanoprost	
Insulin pump infusion set (teflon	00	Juno Pemetrexed	154	Latanoprost with timolol	
cannula, angle insertion)	23	- K -	000	Lax-Suppositories	
Insulin pump infusion set (teflon		Kadcyla		Lax-suppositories Glycerol	
cannula, straight insertion with		Kaletra		Laxatives	
insertion device)	23	Kalydeco		Laxsol	
Insulin pump infusion set (teflon		Kemadrin		Ledipasvir with sofosbuvir	
cannula, straight insertion)	23	Kenacomb	249	Leflunomide	
Insulin pump reservoir	23	Kenacort-A 10	80	Lenalidomide	157
Insulin syringes, disposable with		Kenacort-A 40	80	Letrole	172
attached needle	16	Kenalog in Orabase	32	Letrozole	172
Intelence	106	Ketocal 3:1	279	Leukeran FC	150
Interferon beta-1-alpha	139	KetoCal 4:1	279	Leukotriene Receptor	
Interferon beta-1-beta	139	Ketoconazole		Antagonists	246
Intra-uterine device	73	Dermatological	70	Leuprorelin	
Invega Sustenna		Infection		Leustatin	
Invega Trinza		Ketogenic Diet		Levetiracetam	
IPCA-Frusemide		Ketoprofen		Levetiracetam-AFT	
IPCA-Metoprolol		KetoSens		Levlen ED	
IPCA-Propranolol		Keytruda		Levocabastine	
IPOL		Kindergen		Levocarnitine	
		Kivexa			
Ipratropium bromide244 Iressa		Klacid	100	Levodopa with benserazide	
			0	Levodopa with carbidopa	
Irinotecan Actavis 100		Alimentary		Levomepromazine	130
Irinotecan hydrochloride		Infection		Levomepromazine	401
Irinotecan-Rex		Kliogest		hydrochloride	138
Iron (as ferric carboxymaltose)		Kliovance		Levonorgestrel	
Iron polymaltose		Kogenate FS		Genito-Urinary	
Isentress		Konakion MM		Hormone	
Isentress HD		Konsyl-D		Levonorgestrel BNM	
Ismo 20	56	Kuvan	30	Levothyroxine	
Ismo 40 Retard	56	-L-		Lidocaine [Lignocaine]	. 122–123
Isoniazid		Labetalol		Lidocaine [Lignocaine]	
Isoniazid with rifampicin	100	Lacosamide	130	hydrochloride	123
Isoptin	51	Lactulose	<mark>26</mark>	Lidocaine [Lignocaine] with	
Isoptin Retard	51	Laevolac	<mark>26</mark>	chlorhexidine	123
Isoptin SR	51	Lagevrio	104	Lidocaine [Lignocaine] with	
Isopto Carpine	252	Lamictal	130	prilocaine	123
Isosorbide mononitrate		Lamivudine		Lidocaine-Baxter	123
Isosource Standard		Lamivudine Alphapharm		Lidocaine-Claris	123
Isotretinoin		Lamivudine Viatris		Life Extension	
Ispaghula (psyllium) husk		Lamotrigine		Lignocaine	
Itch-Soothe		Lamprene		Lioresal Intrathecal	
Itraconazole		Lanoxin		Lipid-Modifying Agents	
Itrazole		Lanoxin Paediatric Elixir		Liquigen	
Ivacaftor		Lanoxin PG		Lisinopril	
IVavaIIVI	41	Landain Fu	40	∟ιοιιιομπι	40

Litak	152	Mask for spacer device	248	Methylnaltrexone bromide	2
Lithium carbonate	135	Maviret	103	Methylphenidate ER - Teva	14
Livostin	251	Maxidex	250	Methylphenidate hydrochloride	14
LMX4	123	Maxitrol	250	Methylphenidate hydrochloride	
Locacorten-Viaform ED's	249	MCT oil (Nutricia)	261	extended-release	14
Local preparations for Anal and		Measles, mumps and rubella		Methylprednisolone	8
Rectal Disorders	8	vaccine	285	Methylprednisolone (as sodium	
Locasol	275	Mebendazole	90	succinate)	8
Locoid	.64, 70	Mebeverine hydrochloride	8	Methylprednisolone aceponate	6
Locoid Crelo	64	Medrol		Methylprednisolone acetate	8
Locoid Lipocream	64	Medroxyprogesterone acetate		Methylxanthines	24
Locorten-Vioform	249	Genito-Urinary	74	Metoclopramide Actavis 10	13
Lodoxamide	251	Hormone	82-83	Metoclopramide hydrochloride	13
Logem	130	Mefenamic acid	110	Metolazone	5
Lomide	251	Melatonin	139	Metopirone	8
Lomustine	151	Melpha	151	Metoprolol IV Mylan	5
Loniten	56	Melphalan	151	Metoprolol succinate	
Loperamide hydrochloride	6	Menactra		Metoprolol tartrate	
Lopinavir with ritonavir		Meningococcal (groups A, C, Y an	d	Metrogyl	9
Lopinavir/Ritonavir Mylan		W-135) conjugate vaccine		Metronidazole	9
Loprofin		Meningococcal B multicomponent		Metyrapone	
Loprofin Mix	274	vaccine	286	Mexiletine hydrochloride	
Lorafix	241	Meningococcal C conjugate		Miacalcic	7
Loratadine	241	vaccine	286	Micolette	2
Lorazepam	137	MenQuadfi		Micolette-S29	
Lorstat	54	Menthol	63	Miconazole	3
Losartan Actavis	47	Mepolizumab	205	Miconazole nitrate	
Losartan potassium	47	Mercaptopurine		Dermatological	6
Losartan potassium with		Mercilon 28		Genito-Urinary	7
hydrochlorothiazide	47	Mesalazine	7	Micreme	
Lovir		Mesna	157	Micreme H	
Loxamine	129	Mestinon	110	Microgynon 20 ED	
Lucrin Depot 1-month	88	Metabolic Disorder Agents	26	Microgynon 30	
Lucrin Depot 3-month		Metformin hydrochloride		Microlut	
Lyderm		Metformin Mylan		Midazolam	14
Lynparza		Metformin Viatris		Midazolam Mylan	
Lyrica		Methadone BNM		Midazolam-Baxter	14
- M -		Methadone hydrochloride		Midodrine	
m-Eslon	126	Extemporaneous	257	Mifegyne	
Mabthera		Nervous		Mifepristone	7
Macro Organic Psyllium Husk		Methenamine (hexamine)		Milpharm	
Macrobid		hippurate	109	Minerals	
Macrogol 3350 with potassium		Methopt		Mini-Wright AFS Low Range	
chloride, sodium bicarbonate a	ınd	Methotrexate		Mini-Wright Standard	
sodium chloride		Methotrexate DBL Onco-Vial		Minidiab	
Madopar 125		Methotrexate Ebewe		MiniMed 1.8 Reservoir	
Madopar 250		Methotrexate Sandoz		MMT-326A	2
Madopar 62.5		Methyl hydroxybenzoate		MiniMed 3.0 Reservoir	
Madopar HBS		Methylcellulose		MMT-332A	2
Madopar Rapid		Methylcellulose with glycerin and		MiniMed 770G	
Magnesium hydroxide		sodium saccharin	257	MiniMed Mio MMT-921A	
Magnesium sulphate	35	Methylcellulose with glycerin and		MiniMed Mio MMT-923A	2
Mantoux		sucrose	257	MiniMed Mio MMT-925A	2
Marevan		Methyldopa		MiniMed Mio MMT-941A	2
Marine Blue Lotion SPF 50+		Methyldopa Mylan		MiniMed Mio MMT-943A	
Martindale Pharma		Methyldopa Mylan S29	51	MiniMed Mio MMT-945A	
			· · · · · · · · · · · · · · · · · · ·		

MiniMed Mio MMT-965A		Movapo		Neoral	
MiniMed Mio MMT-975A		Moxifloxacin		Neostigmine metilsulfate	
MiniMed Quick-Set MMT-386A.		MSUD Maxamum	273	Nepro HP (strawberry)	
MiniMed Quick-Set MMT-387A.		Mucilaginous laxatives with		Nepro HP (vanilla)	
MiniMed Quick-Set MMT-396A.		stimulants	25	Nepro HP RTH	
MiniMed Quick-Set MMT-397A.	22	Mucolytics	246	Neulactil	
MiniMed Quick-Set MMT-398A.	<mark>22</mark>	Mucosoothe		Neulastim	
MiniMed Quick-Set MMT-399A.	<mark>22</mark>	Multiple Sclerosis Treatments	138	Neuraxpharm	13
MiniMed Silhouette MMT-368A.	<mark>22</mark>	Multivitamin renal		NeuroTabs	34
MiniMed Silhouette MMT-377A.	22	Multivitamins	33	Nevirapine	10
MiniMed Silhouette MMT-378A.	22	Mupirocin		Nevirapine Alphapharm	
MiniMed Silhouette MMT-381A.	22	Muscle Relaxants		Nevirapine Viatris	
MiniMed Silhouette MMT-382A.	22	Mvite	34	Nicorandil	
MiniMed Silhouette MMT-383A.		Myambutol	100	Nicotine	
MiniMed Silhouette MMT-384A.		Mycobutin		Nifedipine	
MiniMed Sure-T MMT-864A		MycoNail		Nifuran	
MiniMed Sure-T MMT-866A		Mycophenolate mofetil		Nilotinib	
MiniMed Sure-T MMT-874A		Mydriacyl		Nilstat	
MiniMed Sure-T MMT-876A		Mylan (12 hr release)		Alimentary	2
MiniMed Sure-T MMT-884A		Mylan (24 hr release)		Genito-Urinary	
MiniMed Sure-T MMT-886A				Infection	
		Mylan Atenolol		Nintedanib	
Minims Produissions		Mylan Clomiphen			
Minims Prednisolone		Mylan Indapamide		Nipent	
Minirin		Mylan Italy (24 hr release)	51	Nirmatrelvir with ritonavir	
Minirin Melt		Myleran		Nitrates	
Mino-tabs		Mylotarg		Nitroderm TTS	
Minocycline hydrochloride		Myometrial and Vaginal Hormo		Nitrofurantoin	
Minomycin	94	Preparations	75	Nitrolingual Pump Spray	5
Minor Skin Infections	66	Myozyme		Nivestim	4
Minor Skin Infections Minoxidil	66 56	Myozyme N -	26	Nivestim	43
Minor Skin Infections	66 56 83	Myozyme N - Nadolol	26	Nivolumab Nizoral	4; 23; 9
Minor Skin Infections	66 56 83 129	Myozyme		Nivestim Nivolumab Nizoral Nodia	234 98
Minor Skin Infections	66 56 83 129	Myozyme N - Nadolol		Nivolumab Nizoral	234 98
Minor Skin Infections	66 56 83 129	Myozyme	26 50 50 28	Nivestim Nivolumab Nizoral Nodia	4; 9; 110
Minor Skin Infections	66 56 83 129 9 158	Myozyme - N - Nadolol Nadolol BNM Naglazyme		Nivestim Nivolumab Nizoral Nodia Noflam 250	4; 9; 110
Minor Skin Infections	66 56 83 129 9 158	Myozyme	26 50 28 8	Nivestim Nivolumab Nizoral Nodia Noflam 250 Noflam 500	4; 99 110
Minor Skin Infections	66 56 129 9 158 158	Nadolol		Nivestim Nivolumab Nizoral Nodia Noflam 250 Noflam 500 Non-Steroidal Anti-Inflammatory	4; 98 110 110
Minor Skin Infections		Nadolol	26 50 28 8 254 146	Nivestim Nivolumab Nizoral Nodia Noflam 250 Noflam 500 Non-Steroidal Anti-Inflammatory Drugs	4; 23 0 110 110
Minor Skin Infections		Nadolol	26 50 28 8 254 146 146 253	Nivestim Nivolumab Nizoral Nodia Noflam 250 Noflam 500 Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant	4; 23 0 110 110
Minor Skin Infections		Nyozyme - N - Nadolol	2650288254146146253	Nivestim	
Minor Skin Infections		Nyozyme - N - Nadolol - N - Nadolol BNM - Naglazyme - Nalcrom - Naloxone hydrochloride - Naltraccord - Naltraccord - Natrexone hydrochloride - Naphazoline hydrochloride - Naphcon Forte - Naprosyn SR 1000 - Naprosyn SR 1000 - Nadolo - Naphosyn SR 1000 - Naprosyn SR 1000 - Napro	26505088254146146253253	Nivestim	
Minor Skin Infections Minoxidil Mirena Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil		Nyozyme - N - Nadolol	265050288254146146253253110110	Nivestim	
Minor Skin Infections Minoxidil		Nyozyme - N - Nadolol	2650288254146253253110110	Nivestim	
Minor Skin Infections Minoxidil		Nyozyme - N - Nadolol	26	Nivestim	
Minor Skin Infections Minoxidil		Nyozyme - N - Nadolol		Nivestim	
Minor Skin Infections Minoxidil		Nyozyme - N - Nadolol		Nivestim	
Minor Skin Infections Minoxidil		Nyozyme - N - Nadolol		Nivestim	
Minor Skin Infections Minoxidil		Nadolol		Nivestim	
Minor Skin Infections Minoxidil. Mirena. Mirtazapine. Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Molnupiravir Moments Mometasone furoate Monogen Montelukast		Nyozyme - N - Nadolol	26	Nivestim	44:44:23-44:44:44:44:44:44:44:44:44:44:44:44:44:
Minor Skin Infections Minoxidil. Mirena. Mirtazapine. Misoprostol. Mitomycin C. Mitozantrone Ebewe Mitozantrone Ebewe Mitozantrone Ebewe Mixtard 30. MMR II. Moclobemide Modafinil Modavigil Moduretic Molaxole Molnupiravir Moments Moments Mometasone furoate Monogen Montelukast Montelukast Montelukast Montelukast Minimozionic Minoxidil Montelukast Montelukast Montelukast Montelukast Minimozionic Mirena. Mi		Nyozyme - N - Nadolol		Nivestim	444 444 444 444 444 444 444 444 444 44
Minor Skin Infections Minoxidil. Mirena. Mirtazapine. Misoprostol. Mitomycin C. Mitozantrone Ebewe. Mitozantrone Ebewe. Mitozantrone Ebewe. Mitozantrone Mitozantrone Ebewe. Modafinil Modavigil Modavigil Moduretic. Molaxole Molnupiravir Moments. Moments. Mometasone furoate. Monogen. Montelukast Montelukast Mylan. Montelukast Viatris		Nyozyme - N - Nadolol BNM Nadolol BNM Naglazyme Nalcrom Naloxone hydrochloride Naphazoline hydrochloride Naphazoline hydrochloride Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Navelonide Navelbine Nefopam hydrochloride Navelbine Nefopam hydrochloride Nadologia PN - Navelbine Nefopam hydrochloride Nadologia BNM - Navelbine Nefopam hydrochloride Nadologia BNM - Navelbine Nefopam hydrochloride - Nadologia BNM - Navelbine Nefopam hydrochloride - Nadologia BNM - Navelbine Nefopam hydrochloride - Nadologia BNM - Navelbine - Nefopam hydrochloride - Nadologia BNM - Nad		Nivestim	44:44:23-44:44:44:44:44:44:44:44:44:44:44:44:44:
Minor Skin Infections		Nyozyme - N - Nadolol BNM Naglazyme Nalcrom Naloxone hydrochloride Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphazoline hydrochloride Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Nefopam hydrochloride Neisvac-C		Nivestim	4:4:4:23-4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:
Minor Skin Infections		Nadolol Nadolol BNM Naglazyme Nalcrom Naphazoline hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-Mercazole		Nivestim	4:4:4:23-4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:4:
Minor Skin Infections		Nadolol Nadolol BNM Naglazyme Nalcrom Nalcrom Nalcrom Nalcrom Nalcrom Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphazoline hydrochloride Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-Mercazole Neocate Gold Nalcrom Nadolo Neisvac-C Neo-Mercazole Neocate Gold Neisvac-C Neo-Mercazole Neisvac-C Neisva		Nivestim	44: 44: 44: 42: 42: 42: 42: 42: 42: 42:
Minor Skin Infections		Nadolol		Nivestim	44: 44: 44: 42: 42: 42: 42: 42: 42: 42:
Minor Skin Infections		Nadolol Nadolol BNM Naglazyme Nalcrom Nalcrom Nalcrom Nalcrom Nalcrom Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphazoline hydrochloride Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-Mercazole Neocate Gold Nalcrom Nadolo Neisvac-C Neo-Mercazole Neocate Gold Neisvac-C Neo-Mercazole Neisvac-C Neisva		Nivestim	44:44:23-44:44:44:44:44:44:44:44:44:44:44:44:44:

NovoRapid FlexPen	11	Oestrogens	82	Oxycodone hydrochloride	12
NovoRapid Penfill	11	Ofev	245	Oxycodone Sandoz	12
NovoSeven RT	39	Oil in water emulsion	65	OxyNorm	12
Noxafil	98	Olanzapine	135–136	Oxytocin	
Nozinan	135	Olaparib		Oxytocin BNM	
Nozinan (Swiss)	135	Olbetam		Oxytocin GH	70
Nozinan S29	135	Olbetam S29	54	Oxytocin with ergometrine	
Nucala	205	Olopatadine	253	maleate	70
Nuelin	246	Olopatadine Teva	253	Ozurdex	250
Nuelin-SR		Olsalazine		- P -	
Nupentin	130	Omalizumab	207	Pacifen	120
Nusinersen	140	Omeprazole	9	Pacimol	124
Nutilis	272	Omeprazole actavis 10	9	Paclitaxel	159
Nutren Diabetes	262	Omeprazole actavis 20	9	Paclitaxel Actavis	
Nutrient Modules	259	Omeprazole actavis 40	9	Paclitaxel Ebewe	
Nutrini Energy Multi Fibre		Omnitrope	84	Paediatric Seravit	
Nutrini Energy RTH	264	Omnitrope S29	84	Palbociclib	16
Nutrini Low Energy Multi Fibre		Onbrez Breezhaler	242	Paliperidone	136
Nutrini Peptisorb	277	Oncaspar LYO	159	Paliperidone palmitate	137
Nutrini Peptisorb Energy	277	OncoTICE	179	Palivizumab	
Nutrini RTH		Ondansetron	134	Pamidronate disodium	112
Nutrison 800 Complete Multi		Ondansetron ODT-DRLA.	134	Pamisol	112
Fibre	269	One-Alpha	33	Pamol	124
Nutrison Advanced Diason	262	One-Alpha S29		Pancreatic enzyme	24
Nutrison Advanced Peptisorb	266	Onrex	134	Pantoprazole	
Nutrison Concentrated	271	Opdivo	234	Panzop Relief	
Nutrison Energy	269	Ora-Blend	257	Panzytrat	24
Nutrison Energy Multi Fibre	269	Ora-Blend SF	257	Papaverine hydrochloride	5
Nutrison Multi Fibre		Ora-Plus	257	Para-amino salicylic acid	
Nutrison Standard RTH	269	Ora-Sweet	257	Paracare	
Nyefax Retard	51	Ora-Sweet SF	257	Paracare Double Strength	124
Nystatin		Orabase	32	Paracetamol	124
Alimentary	32	Oral and Enteral Feeds	262	Paracetamol (Ethics)	124
Genito-Urinary		Oratane	61	Paracetamol + Codeine	
Infection	98	Ordine	126	(Relieve)	12
NZB Low Gluten Bread Mix	272	Orgran	273	Paracetamol with codeine	
-0-		Ornidazole	99	Paraffin	65–66
Obinutuzumab	206	Orphenadrine citrate	120	Paraffin liquid with wool fat	25
Obstetric Preparations	77	Ortho-tolidine	77	Parasiticidal Preparations	6
Ocicure	9	Oruvail SR	110	Parnate	128
Ocrelizumab	139	Osmolite RTH	269	Parnate S29	
Ocrevus	139	Other Endocrine Agents	88	Paromomycin	9
Octocog alfa [Recombinant factor	or	Other Oestrogen Preparat	ons83	Paroxetine	
VIII] (Advate)	39	Other Progestogen		Paser	100
Octocog alfa [Recombinant factor	or	Preparations	83	Paxam	13
VIII] (Kogenate FS)		Other Skin Preparations	70	Paxlovid	104
Octreotide		Ovestin		Pazopanib	160
Octreotide Depot Teva	170	Genito-Urinary	75	Peak flow meter	248
Octreotide GH		Hormone		Pedialyte - Bubblegum	
Octreotide long-acting	170	Oxaliplatin	151	Pediasure	
Oestradiol		Oxaliplatin Accord	151	Pediasure Plus	264
Oestradiol valerate		Oxaliplatin Actavis 100		Pediasure RTH	264
Oestradiol with norethisterone	83	Oxaliplatin Ebewe		Pegaspargase	
Oestriol		Oxis Turbuhaler		Pegasys	
Genito-Urinary	75	Oxpentifylline	57	Pegfilgrastim	4
Hormone		Oxybutynin	76	Pegylated interferon alfa-2a	107

Pembrolizumab	235	PKU Anamix Junior Orange	274	Primidone Clinect	131
Pemetrexed	154	PKU Anamix Junior Vanilla	274	Primolut N	83
Penicillamine	111	PKU Lophlex LQ 10	274	Priorix	285
Penicillin G	93	PKU Lophlex LQ 20	274	Probenecid	119
PenMix 30	11	PKU Lophlex Powder	274	Probenecid-AFT	119
PenMix 50	11	PKU Lophlex Sensation 20	274	Procarbazine hydrochloride	159
Pentasa	7	Plaquenil		Prochlorperazine	134
Pentostatin [Deoxycoformycir	n] 159	Plendil ER		Proctofoam	
Pentoxifylline [Oxpentifylline].	57	Pneumococcal (PCV10) conjuga	ate	Proctosedyl	8
Peptamen Junior		vaccine	286	Procyclidine hydrochloride	121
Pepti-Junior	278	Pneumococcal (PCV13) conjuga	ate	Progesterone	83
Peptisorb	266	vaccine	287	Proglicem	10
Perhexiline maleate	51	Pneumococcal (PPV23)		Proglycem	10
Pericyazine	135	polysaccharide vaccine	288	Progynova	
Perindopril	46	Pneumovax 23	288	Prolia	
Perjeta	209	Podophyllotoxin	70	Promethazine hydrochloride	241
Permethrin	68	Polaramine	241	Propafenone hydrochloride	49
Perrigo	70	Poliomyelitis vaccine	288	Propamidine isethionate	
Pertuzumab		Poloxamer		Propranolol	
Peteha	100	Poly-Gel	253	Propylene glycol	
Pethidine hydrochloride	127	Poly-Tears		Propylthiouracil	
Pevaryl		Poly-Visc		Prostacur	
Pexsig		Polycal		Protaphane	10
Pfizer Exemestane	172	Polyethylene glycol 400 and		Protaphane Penfill	
Pharmacy Health Sorbolene	with	propylene glycol	253	Protifar	
Glycerin		Ponstan		Protionamide	100
Pharmacy Services		Posaconazole		Provera	82
Pheburane		Posaconazole Juno	98	Provera HD	
Phenasen		Postinor-1	75	PSM Citalopram	128
Phenobarbitone		Potassium chloride		Psoriasis and Eczema	
Phenobarbitone sodium		Potassium citrate	77	Preparations	68
Extemporaneous	257	Potassium iodate		PTU	
Nervous		Povidone iodine	66	Pulmicort Turbuhaler	242
Phenoxybenzamine		Pradaxa		Pulmozyme	
hydrochloride	46	Pramipexole hydrochloride		Puri-nethol	
Phenoxymethylpenicillin (Pen		Pravastatin		Puria	
V)		Pravastatin Mylan		Pyrazinamide	
Phenytoin sodium		Praziquantel		Pyridostigmine bromide	
Phillips Milk of Magnesia	35	Prazosin		Pyridoxine hydrochloride	
Phlexy 10		Pred Forte		Pyridoxine multichem	
Phosphate Phebra		Prednisolone		Pyrimethamine	
Phosphorus		Prednisolone acetate		Pytazen SR	
Phytomenadione		Prednisolone sodium		- Q -	
Pilocarpine hydrochloride		Prednisolone sodium		Quetapel	135
Pimafucort		phosphate	251	Quetiapine	
Pimecrolimus		Prednisolone-AFT		Quick-Set MMT-392	
Pine tar with trolamine laurilsu		Prednisone		Quick-Set MMT-393	
and fluorescein		Prednisone Clinect		Quinapril	
Pinetarsol		Pregabalin		Quinapril with	
Pioglitazone		Pregabalin Pfizer		hydrochlorothiazide	47
Pirfenidone		Pregnancy Tests - hCG Urine		Qvar	
Pizotifen		Premarin		- R -	
PKU Anamix Infant		Prevenar 13		RA-Morph	126
PKU Anamix Junior		Priadel		Ralicrom	
PKU Anamix Junior Chocolate		Primaquine		Raloxifene hydrochloride	
PKU Anamix Junior LQ		Primidone		Raltegravir potassium	
				y p p	

Ramipex	121	RIXUBIS	39	Sertraline	129
Ramipril	47	Rizamelt	132	Setrona	129
Ranbaxy-Cefaclor	90	Rizatriptan	132	Setrona AU	129
Ranbaxy-Cefaclor S29	90	RoActemra S29	224	Sevredol	126
Rapamune	236	Ronapreve	196	Sex Hormones Non	
Rasagiline	121	Ropin		Contraceptive	8 ⁻
Reandron 1000		Ropinirole hydrochloride	121	Shield XL	
Recombinant factor IX	37, 39	Rosuvastatin	54	Shingles vaccine	289
Recombinant factor VIIa	39	Rosuvastatin Viatris	54	Shingrix	289
Recombinant factor VIII	39–40	Rotarix	288	SII-Onco-BCG	179
Rectogesic	8	Rotavirus oral vaccine	288	Sildenafil	58
Redipred	80	Roxane	6	Silhouette MMT-373	2
Relieve	110	Roxane-Propranolol	50	Siltuximab	223
Relistor		Roxithromycin	92	Simvastatin	
Remicade	197	Rubifen	142	Simvastatin Mylan	5
Renilon 7.5	265	Rubifen SR	142	Sinemet	12°
Resonium-A	45	Rugby Capsaicin Topical Cre	am	Sinemet CR	12°
Resource Beneprotein	261	Musculoskeletal	111	Sirolimus	236
Respigen	243	Nervous	123	Siterone	8 [.]
Respiratory Devices	248	Rulide D	92	Slow-Lopresor	50
Respiratory Stimulants		Rurioctocog alfa pegol [Reco	mbinant	Smith BioMed Rapid Pregnancy	
Retinol palmitate		factor VIII]	40	Test	70
ReTrieve		Ruxolitinib	167	Sodibic	4
Retrovir	106	Rythmodan	48	Sodium acid phosphate	20
Revlimid		Rytmonorm		Sodium alginate	(
Revolade	37	- S -		Sodium benzoate	
Rexacrom	251	Sabril	132	Sodium bicarbonate	
Riboflavin	29	Sacubitril with valsartan	47	Blood	. 44-4
Ribomustin	149	Sagent	162	Extemporaneous	25
Ricit	76	SalAir	243	Sodium calcium edetate	25
Rifabutin	101	Salazopyrin	8	Sodium chloride	
Rifadin	101	Salazopyrin EN		Blood	4
Rifampicin	101	Salbutamol		Respiratory	24
Rifaximin		Salbutamol with ipratropium		Sodium citrate with sodium lauryl	
Rifinah	100	bromide	244	sulphoacetate	20
Rilutek	122	Salicylic acid	69	Sodium citro-tartrate	7
Riluzole	122	Salmeterol	242	Sodium cromoglicate	
RINVOQ	238	Sandomigran	133	Alimentary	8
Riodine	66	Sanofi Primaquine	99	Sensory	25
Risedronate Sandoz	113	Sapropterin dihydrochloride	30	Sodium fluoride	34
Risedronate sodium	113	Scalp Preparations	70	Sodium Fusidate [fusidic acid]	
Risperdal Consta	137	Scopoderm TTS	133	Dermatological	62
Risperidone		Sebizole	70	Infection	
Risperidone (Teva)	135	Secukinumab	221	Sensory	249
Risperon	135	Sedatives and Hypnotics	139	Sodium hyaluronate [Hyaluronic	
Ritalin	142	Seebri Breezhaler	244	acid]	250
Ritalin LA	143	Selegiline hydrochloride	121	Sodium phenylbutyrate	30
Ritonavir	107	Senna	26	Sodium picosulfate	20
Rituximab (Mabthera)	209	Senokot	26	Sodium polystyrene sulphonate	4
Rituximab (Riximyo)	211	SensoCard	15	Sodium tetradecyl sulphate	
Rivaroxaban		Serc	133	Sodium valproate	
Rivastigmine	144	Serenace	135	Sofradex	
Rivastigmine Patch BNM 10		Seretide		Soframycin	249
Rivastigmine Patch BNM 5	144	Seretide Accuhaler	243	Solgar27-	-29, 3 ⁻
Rivotril	130	Serevent	242	Solifenacin Mylan	
Riximyo		Serevent Accuhaler	242	Solifenacin succinate	

Solifenacin Viatris	77	Synacthene Retard	80	Thalidomide	160
Solu-Cortef		Synagis	208	Thalomid	
Solu-Medrol	80	Synflorix		Theophylline	
Solu-Medrol-Act-O-Vial	80	Synthroid	83	Thiamine hydrochloride	33
Somatropin (Omnitrope)	84	Syntometrine	76	Thiamine multichem	
Sotalol	50	Syrup (pharmaceutical grade)	258	THIO-TEPA	151
Spacer device	248	Systane Unit Dose	253	Thioguanine	154
Span-K	45	- T -		Thiotepa	15 1
Spinal Muscular Atrophy	140	Tacrolimus		Thyroid and Antithyroid Agents	83
Spinraza	140	Dermatological	69	Ticagrelor	40
Spiolto Respimat	245	Oncology	238	Ticagrelor Sandoz	40
Spiractin	52	Tacrolimus Sandoz	238	Tilcotil	110
Spiriva	244	Taliglucerase alfa	31	Timolol	251
Spiriva Respimat	244	Tambocor	48	Timoptol XE	251
Spironolactone		Tamoxifen citrate	171	Tiotropium bromide	244
Sporanox	97	Tamoxifen Sandoz	171	Tiotropium bromide with	
Sprycel	162	Tamsulosin hydrochloride	76	olodaterol	245
Stelara	230	Tamsulosin-Rex		Tivicay	
Stemetil	134	Tandem Cartridge	21	Tixagevimab with cilgavimab	224
Steril-Gene	81	Tandem t:slim X2 with Basal-IQ		TMP	
SteroClear	247	Tap water		Tobramycin	
Stesolid	129	Taro		Infection	96
Stimulants/ADHD Treatments .	141	Taro-Testosterone	81	Sensory	249
Stiripentol	131	Tasigna	165	Tobramycin BNM	
Stocrin		Tasmar		Tobramycin Mylan	
Stomahesive		Taurine	31	Tobrex	
Strattera	141	Tecfidera	138	Tocilizumab	
Strides Shasun	98	Tegretol	130	Tofranil	128
Stromectol	66	Tegretol CR		Tolcapone	
Sucralfate		Telfast		Tolvaptan	
Sulfadiazine Silver		Teligent		Topamax	
Sulfadiazine sodium		Temaccord		Topical Products for Joint and	
Sulfasalazine	8	Temazepam	140	Muscular Pain	111
Sulphur		Temozolomide		Topiramate	
Sulprix		Tenofovir disoproxil		Topiramate Actavis	
Sumagran		Tenofovir Disoproxil Emtricitabine		Total parenteral nutrition (TPN)	
Sumatriptan		Mylan	103	TPN	
Sunitinib		Tenofovir Disoproxil Emtricitabine		Tramadol hydrochloride	
Sunitinib Pfizer		Viatr		Tramal SR 100	
Sunscreens		Tenofovir Disoproxil Mylan		Tramal SR 150	
Sunscreens, proprietary		Tenoxicam		Tramal SR 200	127
Sure-T MMT-863		Tensipine MR10		Trandate	
Sure-T MMT-873		Tepadina		Tranexamic acid	
Survimed OPD	266	Terbinafine		Tranylcypromine sulphate	
Sustagen Hospital Formula		Terbutaline sulphate		Trastuzumab	
Sustagen Hospital Formula		Teriflunomide		Trastuzumab emtansine	
Active	269	Teriparatide		Travatan	
Sustanon Ampoules		Testosterone		Travoprost	
Sylvant		Testosterone cipionate		Treatments for Dementia	
Symbicort Turbuhaler 100/6		Testosterone esters		Treatments for Substance	
Symbicort Turbuhaler 200/6		Testosterone undecanoate		Dependence	149
Symbicort Turbuhaler 400/12		Tetrabenazine		Trental 400	
Symmetrel		Tetrabromophenol		Tretinoin	
Sympathomimetics		Tetracosactrin		Dermatological	61
Synacthen		Tetracycline		Oncology	
Synacthen Depot		Teva Lisinopril		Trexate	
-,		·			

Triamcinolone acetonide	vaccine]289	Voltaren SR	11
Alimentary32	Varicella zoster vaccine [Shingles	Volumatic	24
Dermatological64	vaccine] 289	Voriconazole	
Hormone80	Varicella zoster virus (Oka strain) live	Votrient	16
Triamcinolone acetonide with	attenuated vaccine shingles	Vttack	9
gramicidin, neomycin and nystatin	vaccine] 289	- W -	
Dermatological64	Various254	Warfarin sodium	4
Sensory249	Varivax289	Wart Preparations	
Triazolam140	Vasodilators56	Wasp venom allergy treatment	
Trimethoprim96	Vasopressin Agonists88	Water	
Trimethoprim with	Vasorex50	Blood	4
sulphamethoxazole	Vebulis59	Extemporaneous	25
[Co-trimoxazole]96	Vedafil58	White Soft Liquid Paraffin AFT	6
Trisequens83	Vedolizumab231	Wool fat with mineral oil	
Trisul96	Veletri59	- X -	
Trophic Hormones84	Venclexta161	Xarelto	4
Tropicamide252	Venetoclax161	Xifaxan	
Trulicity12	Venlafaxine129	XMET Maxamum	27
Trusopt251	Venomil240-241	Xolair	
TruSteel21	VENOX240	XP Maxamum	27
Tryzan47	Ventavis59	Xylocaine	
Tuberculin PPD [Mantoux] test289	Ventolin243	Xylocaine 2% Jelly	
Tubersol289	Vepesid	Xyntha	
Two Cal HN271	Verapamil hydrochloride51	- Z -	
Tykerb164	Vermox90	Zapril	4
Tysabri139	Versacloz134	Zarontin	
- U -	Vesanoid161	Zaroxolyn	
UK Synacthen80	Vexazone12	Zavedos	
Ultibro Breezhaler245	Vfend98	Zeffix	
Ultraproct8	Viaderm KC64	Zematop	
Umeclidinium244	Viatris	Zetlam	
Umeclidinium with vilanterol245	Cardiovascular49	Ziagen	
Univent244, 247	Infection96, 106	Zidovudine [AZT]	
Upadacitinib238	Vigabatrin132	Zidovudine [AZT] with	
Ural	Vigisom139	lamivudine	10
Urea	Vildagliptin12	Ziextenzo	
Urex Forte	Vildagliptin with metformin	Zimybe	
Urinary Agents76	hydrochloride	Zinc and castor oil	
Urinary Tract Infections109	Vimpat130	Zinc sulphate	
Urinorm119	Vinblastine sulphate162	Zincaps	3
Uromitexan 157	Vincristine sulphate162	Zinnat	
Ursodeoxycholic acid24	Vinorelbine162	Ziprasidone	
Ursosan	Vinorelbine Ebewe	Zista	
Ustekinumab230	Viramune Suspension	Zithromax	
Utrogestan83	ViruPOS249	Zoledronic acid	
- V -	Vit.D3	Hormone	7
Vaccinations280	VitA-POS253	Musculoskeletal	
Vaclovir	Vitabdeck34	Zoledronic acid Mylan	
Valaciclovir 102	Vital	Zoledronic acid Viatris	
Valganciclovir	Vitamin B complex33	Hormone	7
Valganciclovir Mylan102	Vitamin B complex 33	Musculoskeletal	
Vancomycin97	Vitamins	Zoledronic-US	
Vannair	Vivonex TEN	Zopiclone	
Varenicline Pfizer147	Voltaren	Zopiclone Actavis	1/1
Varenicline Frizer147 Varenicline tartrate147	Voltaren D110	Zostavax	
Varicella vaccine [Chickenpox	Voltaren Ophtha250	Zostrix	
vancena vaccine (chickenpox	voltaren Oprilia230	ZUSUIX	11

Zostrix HP	123
Zuclopenthixol decanoate	137
Zuclopenthixol hydrochloride	136
Zusdone	135
Zyban	146
Zypine	135
Zypine ODT	135
Zyprexa Relprevv	136
Zutigo	160