		•	
March 2023 Volume 30 Editors: Kaye Wilson, Doris Chong & Ayeshah Khan	Section A	General Rules	5
email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street	Section B	Alimentary Tract & Metabolism	6
PO Box 10 254 Wellington		Blood & Blood Forming Organs	37
Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)		Cardiovascular System	47
Circulation		Dermatologicals	62
You can register to have an electronic version of the Pharmaceutical Schedule (link		Genito Urinary System	72
to PDF copy) emailed to your nominated		Hormone Preparations – Systemic	79
email address each month by subscribing at pharmac.govt.nz/subscribe.		Infections – Agents For Systemic Use	90
Production		Musculoskeletal System	111
Typeset automatically from XML and TEX. XML version of the Schedule available from		Nervous System	117
schedule.pharmac.govt.nz/pub/schedule		Oncology Agents & Immunosuppressants	144
Programmers Anrik Drenth & John Geering		Respiratory System & Allergies	235
email: texschedule@pharmac.govt.nz			
©Pharmaceutical Management Agency		Sensory Organs	244
ISSN 1179-3686 BY This work is licensed under the Creative		Various	249
Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence	Section C	Extemporaneous Compounds (ECPs)	251
terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written	Section D	Special Foods	254
form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility	Section I	National Immunisation Schedule	275
for any errors or omissions, and shall not be liable for any consequences arising there from.		Index	285

Introducing Pharmac

2

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gram g	
kilogram kg	
international unit iu	

Abbreviations

Capsule Cream Device Dispersible Effervescent Emulsion	Amp Cap Crm Dev Disp Eff Emul EC
Enteric Coated	EC

microgram me	cg
milligramn	ng
millilitreı	ml

millimole	mmol
unit	u

Gelatinous	Gel	SolutionSoln
Granules	Gran	SuppositorySupp
Infusion	Inf	TabletTab
Injection	Inj	Tincture Tinc
Liquid	Liq	Trans Dermal Delivery
Long Acting	LA	SystemTDDS
Ointment	Oint	-
Sachet	Sach	

Read the General Rules : https://www.pharmac.govt.nz/section-a.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	-	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
A	ntacids and Antiflatulents				
A	ntacids and Reflux Barrier Agents				
	INIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg pe sachet DIUM ALGINATE		30	J	Gaviscon Infant
*	Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (13.61)	60		Gaviscon Extra Strength
*	Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m	ıl	Acidex
P	nosphate Binding Agents				
*	MINIUM HYDROXIDE Tab 600 mg CIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement		100 500 m	_	Alu-Tab Roxane
A	Only when prescribed for patients unable to swallow calci inappropriate and the prescription is endorsed accordingly nticliarrhoeals		ts or v	vhere calci	PAI um carbonate tablets are
A	gents Which Reduce Motility				
*	PERAMIDE HYDROCHLORIDE – Up to 30 cap available on a Tab 2 mg Cap 2 mg		400 400		Nodia <u>Diamide Relief</u>
R	ectal and Colonic Anti-inflammatories				
	DESONIDE Cap 3 mg – Special Authority see SA1886 below – Retail pharmacy A1886 Special Authority for Subsidy		90	1	Entocort CIR
Initi	al application — (Crohn's disease) from any relevant practi following criteria:	tioner. Approvals v	alid fo	r 6 months	for applications meeting
	1 Mild to moderate ileal, ileocaecal or proximal Crohn's disea 2 Any of the following:	ase; and			
	2.1 Diabetes; or				
					continued.

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🖌	Manufacturer	

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)	15 g OP	 ✓ Colifoam ✓ Cortifoam \$29
	21.1 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%	10 g OP	Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	Asacol
Tab long-acting 500 mg56.10	100	 Pentasa
Tab 800 mg	90	Asacol
Modified release granules, 1 g118.10	100 OP	 Pentasa
Enema 1 g per 100 ml41.30	7	 Pentasa
Suppos 500 mg	20	Asacol
Suppos 1 g	28	 Pentasa

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully		
	(Manufacturer's Price) \$) Per	Subsidised	d Generic Manufacturer	
DLSALAZINE	•				
Tab 500 mg		60	1	Atnahs	
				Olsalazine S29	
	93.37	100	1	Dipentum	
Cap 250 mg	53.00	100	1	Dipentum	
PREDNISOLONE SODIUM					
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	1	Essential	
				Prednisolone S29	
ODIUM CROMOGLICATE					
Cap 100 mg	92.91	100		Nalcrom	
Nalawara Cara 100 marta ha daliatad 1 April 2000)			1	Ralicrom	
Nalcrom Cap 100 mg to be delisted 1 April 2023)					
:ULFASALAZINE ₭ Tab 500 mg	16 50	100		Salazopyrin	
k Tab 500 mg		100		Salazopyrin EN	
		100	-		
Local preparations for Anal and Rectal Disorde	ers				
Antihaemorrhoidal Preparations					
•					
LUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PI	VALATE AND CINCF	IOCAI	NE		
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g	11.06	30 g O	P 🗸	Ultraproct	
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		50 y O		onaproor	
cinchocaine hydrochloride 1 mg	7.30	12	1	Ultraproct	
YDROCORTISONE WITH CINCHOCAINE				•	
Oint 5 mg with cinchocaine hydrochloride 5 mg per g		30 g O	P 🗸	Proctosedyl	
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g		12		Proctosedyl	
Management of Anal Fissures					
LYCERYL TRINITRATE – Special Authority see SA1329 belo	w – Retail pharmacy				
♦ Oint 0.2%		30 g O	P 🗸	Rectogesic	
SA1329 Special Authority for Subsidy		5			
itial application from any relevant practitioner. Approvals va	lid without further ren	ewal u	nless noti	fied where the patient has	
hronic anal fissure that has persisted for longer than three wee	ks.				
Antioneomodice and Other Agente Altering Cu	t Motility				
Antispasmodics and Other Agents Altering Gu	it would y				
SLYCOPYRRONIUM BROMIDE					
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available o					
PSO	65.45	10	-	Max Health	
YOSCINE BUTYLBROMIDE			-	_	
₭ Tab 10 mg		100		Buscopan	
 Inj 20 mg, 1 ml – Up to 5 inj available on a PSO 	6.35	5		Buscopan Buscopan COO	
			~	Buscopan S29 S29	
	0.00	~~		0.1.4.	
🖌 Tab 135 mg	9.20	90	~	Colofac	

8

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Antiulcerants				
Antisecretory and Cytoprotective				
MISOPROSTOL * Tab 200 mcg – Up to 120 tab available on a PSO	47.73	120	•	Cytotec
Helicobacter Pylori Eradication				
 CLARITHROMYCIN Tab 500 mg – Subsidy by endorsement a) Maximum of 28 tab per prescription b) Subsidised only if prescribed for helicobacter pylori er Note: the prescription is considered endorsed if clarit inhibitor and either amoxicillin or metronidazole. 	adication and prescr		is endorse	
H2 Antagonists				
FAMOTIDINE – Only on a prescription * Tab 20 mg	4.91	100	1	Famotidine Hovid S29
₭ Tab 40 mg	8.48	100	1	Famotidine Hovid S29
Inj 10 mg per ml, 4 ml – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients received		10 t of pa		Mylan S29
Proton Pump Inhibitors				
ANSOPRAZOLE * Cap 15 mg * Cap 30 mg DMEPRAZOLE For omeprazole suspension refer Standard Formulae, page 2	5.26	100 100		Lanzol Relief Lanzol Relief
* Cap 10 mg		90	1	Omeprazole actavis 10
₭ Cap 20 mg	1.86	90	1	Omeprazole actavis 20
₭ Cap 40 mg	3.11	90	1	Omeprazole actavis 40
Powder – Only in combination Only in extemporaneously compounded omeprazole susp		5 g	1	Midwest
 Inj 40 mg ampoule with diluent 		5		<u>Dr Reddy's</u> <u>Omeprazole</u> Ocicure ^{©29}
PANTOPRAZOLE 券 Tab EC 20 mg 券 Tab EC 40 mg		90 90		Panzop Relief Panzop Relief
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mg	14.51	50	1	Gastrodenol 529

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
SUCRALFATE Tab 1 g		120		
	(48.28)		(Carafate

Bile and Liver Therapy

⇒SA1461 Special Authority for Subsidy

Initial application only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

56

✓ Xifaxan

Renewal only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Diabetes

Hyperglycaemic Agents

DIAZOXIDE - Special Authority see SA1320 below - Retail pharm	nacy		
Cap 25 mg	110.00	100	Proglicem S29
Cap 100 mg	280.00	100	Proglicem S29
Oral liq 50 mg per ml	620.00	30 ml OP	Proglycem S29
			🖌 e5 Pharma S29

➡SA1320 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

GLUCAGON HYDROCHLORIDE

GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSO	1	✓ Glucagen Hypokit
Insulin - Short-acting Preparations		
INSULIN NEUTRAL ▲ Inj human 100 u per ml25.26	10 ml OP	 ✓ Actrapid ✓ Humulin R
▲ Inj human 100 u per ml, 3 ml42.66	5	 ✓ Actrapid Penfill ✓ Humulin R
Insulin - Intermediate-acting Preparations		
INSULIN ASPART WITH INSULIN ASPART PROTAMINE	5	✓ NovoMix 30 FlexPen

	r per mi, o mi premied pen		0	
INSULIN ISO	PHANE			
🔺 Inj humai	n 100 u per ml	17.68	10 ml OP	 Humulin NPH
				 Protaphane
🔺 Inj humar	n 100 u per ml, 3 ml	29.86	5	 Humulin NPH
				 Protaphane Penfill

	Subsidy		Fully	Brand or
	(Manufacturer's P	rice) Subs	idised	
	\$	Per	1	Manufacturer
NSULIN ISOPHANE WITH INSULIN NEUTRAL				
 Inj human with neutral insulin 100 u per ml 	25.26	10 ml OP	1	Humulin 30/70
				Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	-	Humulin 30/70
		Ũ		PenMix 30
				PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE				
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,		F		Humalan Mix 05
3 ml		5	v	Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,		-		Harris Min 50
3 ml		5	•	Humalog Mix 50
Insulin - Long-acting Preparations				
NSULIN GLARGINE				
Inj 100 u per ml, 10 ml	63.00	1	1	Lantus
Inj 100 u per mi, 3 mi		5		Lantus
Inj 100 u per ml, 3 ml disposable pen		5 5		Lantus SoloStar
	94.50	5	•	Lantus SoloStar
Insulin - Rapid Acting Preparations				
NSULIN ASPART				
Inj 100 u per ml, 10 ml	30.03	1	1	NovoRapid
Inj 100 u per ml, 3 ml		5		NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5		NovoRapid FlexPen
		U U		
Inj 100 u per ml, 10 ml	27.02	1	1	Apidra
Inj 100 u per ml, 3 ml		5		Apidra
Inj 100 u per ml, 3 ml disposable pen		5		Apidra SoloStar
		5	•	Apiula SoloSiai
NSULIN LISPRO				
Inj 100 u per ml, 10 ml		10 ml OP	-	Humalog
Inj 100 u per ml, 3 ml	59.52	5	~	Humalog
Alpha Glucosidase Inhibitors				
CARBOSE				
₭ Tab 50 mg	8.95	90	1	Accarb
K Tab 100 mg		90	-	Accarb
Oral Hunaghuasamia Aganta				
Oral Hypoglycaemic Agents				
GLIBENCLAMIDE			-	
🖌 Tab 5 mg	7.50	100	~	Daonil
BLICLAZIDE				
 Tab 80 mg 	15.18	500	✓	Glizide
BLIPIZIDE				
K Tab 5 mg	4.58	100	1	Minidiab
IETFORMIN HYDROCHLORIDE				
	1/ 7/	1 000	1	Metformin Mulan
Tab immediate-release 500 mg	14./4	1,000		Metformin Mylan Metformin Viatris
k Tab immediate-release 850 mg	11.00	500		Metformin Mylan
		500	•	
Metformin Mylan Tab immediate-release 500 mg to be delisted 1	August 2023)			

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

(\	Subsidy /lanufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PIOGLITAZONE				
* Tab 15 mg	6.80	90	 Image: A second s	Vexazone
* Tab 30 mg	7.30	90	 Image: A second s	Vexazone
* Tab 45 mg	12.25	90	 Image: A second s	Vexazone
VILDAGLIPTIN				
Tab 50 mg	35.00	60	v	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	1	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	~ (Galvumet
GLP-1 Agonists				
DULAGLUTIDE - Special Authority see SA2065 below - Retail pha Note: Not to be given in combination with a funded SGLT-2 inh				
* Inj 1.5mg per 0.5 ml prefilled pen		4	1	Trulicity
>SA2065 Special Authority for Subsidy				•

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

LIRAGLUTIDE - Special Authority see SA2187 below - Retail pharmacy

- a) Maximum of 9 inj per prescription
- b)
- a) Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

⇒SA2187 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

the following criteria:

Either:

- 1 Patient has previously received an initial Special Authority approval for either an SGLT-2 inhibitor or GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

		Subsidy		Fully	Brand or
		(Manufacturer's Price \$	e) Per	Subsidised	Generic Manufacturer
EM	PAGLIFLOZIN – Special Authority see SA2068 on the previo Note: Not to be given in combination with a funded GLP-1 a	ous page – Retail ph	armacy	/	
	Tab 10 mg Tab 25 mg		30 30		Jardiance Jardiance
EM	PAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - S				
pna	rmacy Note: Not to be given in combination with a funded GLP-1 a	agonist.			
	Tab 5 mg with 1,000 mg metformin hydrochloride		60		Jardiamet
	Tab 5 mg with 500 mg metformin hydrochloride		60		Jardiamet
	Tab 12.5 mg with 1,000 mg metformin hydrochloride Tab 12.5 mg with 500 mg metformin hydrochloride		60 60		Jardiamet Jardiamet
D	iabetes Management				
Κ	etone Testing				
BL(DOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by end	dorsement			
	a) Not on a BSO				
	b) Maximum of 20 strip per prescription				
	c) Up to 10 strip available on a PSO				
	d) Patient has any of the following:				
	1) type 1 diabetes; or				
	 permanent neonatal diabetes; or undergone a pancreatectomy; or 				
	4) cystic fibrosis-related diabetes; or				
	5) metabolic disease or epilepsy under the care of a p	aediatrician neurolo	naist or	metabolic	snecialist
	The prescription must be endorsed accordingly.		giot of	metabolic	opeoialiot.
	Test strips	15.50 1	0 strip	OP 🗸	KetoSens
D	ual Blood Glucose and Blood Ketone Testing				
DU	AL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC	C TEST METER – S	ubsidy	by endors	ement
	a) Maximum of 1 pack per prescription				
	b) Up to 1 pack available on a PSO				
	c) A dual blood glucose and blood ketone diagnostic test m	neter is subsidised fo	r a pati	ent who h	as:
	 type 1 diabetes; or 				
	permanent neonatal diabetes; or				
	 undergone a pancreatectomy; or 				
	 cystic fibrosis-related diabetes; or 				
	5) metabolic disease or epilepsy under the care of a p				
	The prescription must be endorsed accordingly. Only 1				
	the avoidance of doubt patients who have previously rec funded CareSens meter.	eiveu a iunueu mele	i, ouie		esens, are engible for a
	Meter with 50 lancets, a lancing device and 10 blood glucos	e			
	diagnostic test strips		1 OP	1	CareSens Dual
	č				

Subsidy (Manufacturer's Price)		Fully bsidised	Brand or Generic	
\$	Per	1	Manufacturer	

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes: or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cvstic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

strips					1 OP	 CareSens N CareSens N POP
				20.00		✓ CareSens N Premier
Note: Only	1 meter avai	ilable per F	SO			

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed: or
- Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed: or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

50 test OP SensoCard

50 test OP

✓ CareSens N

CareSens PRO

	Subsidy (Manufacturer's Price)	Si	Fully bsidised	
	(Manalatator o F 166) \$	Per	A 100 A 1	Manufacturer
Insulin Syringes and Needles				
Subsidy is available for disposable insulin syringes, needles, and				
he supply of insulin or liraglutide or when prescribed for a patier annotate the prescription as endorsed where there exists a reco				
NSULIN PEN NEEDLES - Maximum of 200 dev per prescription				
₩ 29 g × 12.7 mm		100		B-D Micro-Fine
₩ 31 g × 5 mm		100		B-D Micro-Fine
₩ 31 g × 6 mm		100		Berpu
₩ 31 g × 8 mm		100		B-D Micro-Fine
卷 32 g × 4 mm	10.95	100	~	B-D Micro-Fine
NSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDL	E – Maximum of 200	dev per	prescri	ption
Syringe 0.3 ml with 29 g × 12.7 mm needle	13.56	100	 ✓ 	B-D Ultra Fine
	1.36	10		
	(1.99)			B-D Ultra Fine
₭ Syringe 0.3 ml with 31 g × 8 mm needle		100	1	B-D Ultra Fine II
, , , ,	1.30	10		
	(1.99)			B-D Ultra Fine II
Syringe 0.5 ml with 29 g × 12.7 mm needle		100	1	B-D Ultra Fine
, , , , , , , , , , , , , , , , , , , ,	1.36	10		
	(1.99)			B-D Ultra Fine
Syringe 0.5 ml with 31 g × 8 mm needle		100	1	B-D Ultra Fine II
, , , , , , , , , , , , , , , , , , , ,	1.36	10		
	(1.99)			B-D Ultra Fine II
Syringe 1 ml with 29 g × 12.7 mm needle		100	1	B-D Ultra Fine
	1.36	10		
	(1.99)			B-D Ultra Fine
₭ Syringe 1 ml with 31 g × 8 mm needle	()	100	1	B-D Ultra Fine II
	1.36	10		
	(1.99)			B-D Ultra Fine II
Insulin Pumps				
NSULIN PUMP – Special Authority see SA1603 below – Retail	pharmacy			

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

C)	Maxi	mun	۱ of	1	insulin	pump	per	patient	each	four	year	period	

Min basal rate 0.025 U/h	8,800.00	1	 MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	 Tandem t:slim
			X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

All of the following.

16

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and

8 Either:

- 8.1 Applicant is a relevant specialist; or
- 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and

3 Either:

- 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
- 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	bsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and

3 Either:

- 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
- 3.2 The pump is due for replacement; and

4 Either:

- 4.1 Applicant is a relevant specialist; or
- 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and

8 Either:

- 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
- 8.2 The pump is due for replacement; and

9 Either:

- 9.1 Applicant is a relevant specialist; or
- 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's	Price) Sub:	sidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

➡SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and

8 Either:

8.1 Applicant is a relevant specialist; or

8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and

3 Either:

- 3.1 Applicant is a relevant specialist; or
- 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and

3 Either:

- 3.1 Applicant is a relevant specialist; or
- 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

	Subsidy (Manufacturer's Price \$) Sub Per	Fully sidised	Brand or Generic Manufacturer
 continued pump therapy; and 4 The patient is continuing to derive benefit from pump thera 5 The patient had achieved and is maintaining a HbA1c of e 6 The patient has had no increase in severe unexplained hy 7 The patient's HbA1c has not deteriorated more than 5 mm 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within thei Renewal — (Previous use before 1 September 2012) only from years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mm 	apy; and qual to or less than poglycaemic episod ol/mol from baselind r vocational scope. n a relevant speciali treatment plan and ol/mol from initial ap	80 mmol/r les from ba e; and st or nurse has maint oplication;	aseline; a e practitio ained a and	ump therapy; and and oner. Approvals valid for 2 HbA1c of equal to or less
 3 The patient has not had an increase in severe unexplained 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within thei INSULIN PUMP CARTRIDGE – Special Authority see SA1985 or 	r vocational scope.		n daselir	ie; and
 a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per Cartridge 300 U, t:lock × 10 	year.	1 OP	✔ Т	andem Cartridge
 INSULIN PUMP INFUSION SET (STEEL CANNULA) – Special A a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10 	·	5 on page 1 OP		liniMed Sure-T
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	🗸 N	MMT-884A liniMed Sure-T MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	🗸 N	IiniMed Sure-T MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	🗸 N	liniMed Sure-T MMT-866A
8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	🗸 N	liniMed Sure-T MMT-874A
8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T MMT-876A
6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	✓ s	ure-T MMT-863
8 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock	130.00	1 OP	√ S	ure-T MMT-873

	Subsidy (Manufacturer's \$	Price)	Subsic Per	Fully lised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	(INSERTION)	– Speci	al Authori	ty see	SA1985 on page 19 -
6 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles		1	OP	🗸 Ti	ruSteel
8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles		1	OP	🗸 Ti	ruSteel
6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles		1	OP	🗸 Ti	ruSteel
8 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles		1	OP	🗸 Ti	ruSteel

	Subsidy		Fully	Brand or
	(Manufacturer's Prie	ce) Si Per	ubsidised	Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA) – Specia	al Authority see SA	1985 on p	age 19 –	Retail pharmacy
 a) Maximum of 3 set per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 	,, ,		-9	· · · · · · · · · · · · · · · · · · ·
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	1	MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	1	MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing x 10	130.00	1 OP	1	MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing x 10		1 OP	1	MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10		1 OP	1	MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing x 10		1 OP	1	MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing x 10	130.00	1 OP	1	MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing x 10	130.00	1 OP	1	MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	1	MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing x 10		1 OP	1	MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing × 10	130.00	1 OP	1	MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing × 10		1 OP	1	MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing		1 OP	1	MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10		1 OP	1	MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing x 10		1 OP	1	MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10		1 OP	1	MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing x 10	130.00	1 OP	1	MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	1	MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	1	MiniMed Quick-Set MMT-386A

	Subsidy (Manufacturer's Pri		Fully osidised	Brand or Generic
	\$	Per	-	Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription	SERTION WITH	INSERTION	N DEVIC	 E) – Special Authority see
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c line × 10 with 10 needles	140.00	1 OP	✓ A	utoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cm line × 10 with 10 needles		1 OP	✓ A	utoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	SEBTION) - Sp	ecial Author	itv see S	A1985 on page 19 -
 Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 17 mm teflon cannula; angle insertion; 60 cm line × 10 with 	, -p		.,	
10 needles; luer lock	130.00	1 OP	1 9	ilhouette MMT-373
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH see SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device;	T INSERTION W	ITH INSERT	TION DE	VICE) – Special Authority
110 cm line × 10 with 10 needles 6 mm teflon cannula; straight insertion; insertion device; 60 cr		1 OP	✓ A	utoSoft 90
line × 10 with 10 needles		1 OP	✓ A	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 110 cm line × 10 with 10 needles	140.00	1 OP	✓ A	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 cr line × 10 with 10 needles		1 OP	✓ A	utoSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH Retail pharmacy a) Maximum of 3 sets per prescription	T INSERTION) -	- Special Au	thority se	ee SA1985 on page 19 –
 b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 				
10 needles; luer lock 9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with		1 OP	✓ (Quick-Set MMT-393
10 needles; luer lock		1 OP	✓ 0	uick-Set MMT-392
INSULIN PUMP RESERVOIR - Special Authority see SA1985 or	page 19 – Retai	il pharmacy		
 a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded per sets 				
$10 \times luer lock conversion cartridges 1.8 ml for Paradigm pump$		1 OP	✓ Δ	DR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 10		1 OP		liniMed 1.8 Reservoir MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP	✓ N	liniMed 3.0 Reservoir MMT-332A

24

	Subsidy (Manufacturer's Price) \$) Per	Fully Subsidised	Brand or Generic Manufacturer
Digestives Including Enzymes				
PANCREATIC ENZYME				
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)		100	1	Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amyla 1,250 U protease))		100	✓	Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)		100	✓	Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200				
Eur U) (Panzytrat Cap pancreatin (175 mg (25,000 U lipase, 22,500 U	l amylase, 1,250 U pro			Creon Micro isted 1 June 2023)
URSODEOXYCHOLIC ACID – Special Authority see SA1739 Cap 250 mg		100 ncy	~	Ursosan
➡SA1739 Special Authority for Subsidy				
Initial application — (Alagille syndrome or progressive fan Approvals valid without further renewal unless notified for appli				ny relevant practitioner.
Either:	5	Ŭ		
 Patient has been diagnosed with Alagille syndrome; or Patient has progressive familial intrahepatic cholestasis 				
Initial application — (Chronic severe drug induced cholest for 3 months for applications meeting the following criteria: All of the following:	atic liver injury) from	any re	elevant pra	ctitioner. Approvals valid
 Patient has chronic severe drug induced cholestatic live Cholestatic liver injury not due to Total Parenteral Nutrit Treatment with ursodeoxycholic acid may prevent hosp 	ion (TPN) use in adults		tion of stay	
Initial application — (Primary biliary cholangitis) from any meeting the following criteria: Both:				
 Primary biliary cholangitis confirmed by antimitochondri with or without raised serum IgM or, if AMA is negative, Patient not requiring a liver transplant (bilirubin > 100 ui 	by liver biopsy; and			d cholestatic liver enzymes
Initial application - (Pregnancy) from any relevant practitio				e the patient diagnosed with
cholestasis of pregnancy. Initial application — (Haematological Transplant) from any meeting the following criteria:	relevant practitioner.	Approv	vals valid fo	or 6 months for applications
Both: 1 Patient at risk of veno-occlusive disease or has hepatic allogenic stem cell or bone marrow transplantation; and 2 Treatment for up to 13 weeks.		lergoin	g condition	ing treatment prior to
Initial application — (Total parenteral nutrition induced ch	olestasis) from any re	elevant	practitione	r. Approvals valid for 6

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

months for applications meeting the following criteria:

Both:

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

continued...

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Laxatives

Bulk-forming Agents

ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln	6.00	250 g OP	 Macro Organic Psyllium Husk
	12.20	500 g OP	✓ <u>Konsyl-D</u>
MUCILAGINOUS LAXATIVES WITH STIMULANTS			
* Dry	6.02	500 g OP	
	(17.32)	-	Normacol Plus
	2.41	200 g OP	
	(8.72)	Ū	Normacol Plus
Faecal Softeners			
DOCUSATE SODIUM - Only on a prescription			
* Tab 50 mg	2.31	100	 Coloxyl
		400	

* Tab 120 mg3.13	100	 <u>Coloxyl</u> 	
DOCUSATE SODIUM WITH SENNOSIDES Tab 50 mg with sennosides 8 mg	200	✓ Laxsol	
POLOXAMER – Only on a prescription Not funded for use in the ear.			
* Oral drops 10%	30 ml OP	✓ Coloxyl	

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE - Special Authori	ty see SA1691 below – Retail ph	armacy	,
Inj 12 mg per 0.6 ml vial		1	 Relistor
	246.00	7	 Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

26

1 The patient is receiving palliative care; and

2 Either:

- 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
- 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully Brand or osidised Generic ✓ Manufacturer
Osmotic Laxatives			
GLYCEROL * Suppos 2.8/4.0 g – Only on a prescription		20	✓ <u>Lax-suppositories</u> Glycerol
LACTULOSE – Only on a prescription * Oral liq 10 g per 15 ml Laevolac to be Principal Supply on 1 April 2023	3.61	500 ml	✓ Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM B Powder for oral soln 13.125 g with potassium chloride 46.6		SODIUM	CHLORIDE
sodium bicarbonate 178.5 mg and sodium chloride 350		30	✓ <u>Molaxole</u>
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	✓ Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETAT Enema 90 mg with sodium lauryl sulphoacetate 9 mg per m 5 ml Micolette to be Principal Supply on 1 June 2023	l,	ription 50	✓ Micolette ✓ Micolette-S29 S29
Stimulant Laxatives			
BISACODYL – Only on a prescription * Tab 5 mg * Suppos 10 mg SENNA – Only on a prescription * Tab, standardised	3.69 2.17 (8.21)	200 10 100	 Bisacodyl Viatris Lax-Suppositories Senokot
	0.43 (2.06)	20	Senokot
SODIUM PICOSULFATE – Special Authority see SA2053 below Oral soln 7.5 mg per ml		30 ml OP	 Dulcolax SP Drop

1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and

2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

✓ Myozyme

(Ma	Subsidy nufacturer's Price)	Sut	Fully	Brand or Generic
	\$	Per	1	Manufacturer

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 below - Retail pharmacy

Tab 1,000 mg	CBS	90	 Clinicians
Cap 500 mg	CBS	50	 Solgar
Powder		400 g	 Biomed

➡SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE - Special Authority see SA1987 on the next page - Retail pharmac	су	
Powder for oral soln575.00	180 g OP	 Cystadane

28

➡SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the followina:

- 1 The patient has a confirmed diagnosis of homocystinuria: and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5.10-methylene-tetrahydrofolate reductase (MTHFR) deficiency: or
 - 2.3 A disorder of intracellular cobalamin metabolism; and

3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation. Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below - Retail p	pharmacy		
Cap 120 mg	CBS	30	🗸 Solgar
Cap 160 mg	CBS	60	🗸 Go Hea

υaμ	120 mg		30	 Solyal
Сар	160 mg	CBS	60	 Go Healthy

► SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE - Special Authority see SA1988 below - Retail pharmacy

Inj 1 mg per ml, 5 ml vial.....2,234.00 1 ✓ Naglazyme

■ SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

IDURSULFASE	- Special Authority	see SA1623 on	the next page	- Retail pharmacy

Inj 2 mg per ml, 3 ml vial	4,608.30	1	🗸 Elaprase
----------------------------	----------	---	------------

	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully osidised	Brand or Generic Manufacturer
SA1623 Special Authority for Subsidy				
nitial application only from a metabolic physician. Approvals	s valid for 24 weeks for	r applicatio	ns meet	ing the following criteria:
Il of the following:				
 The patient has been diagnosed with Hunter Syndrome Either: 		,,		
 2.1 Diagnosis confirmed by demonstration of iduron assay in cultured skin fibroblasts; or 2.2 Dataglian of a diagnost population in the in 			e blood	cells by either enzyme
2.2 Detection of a disease causing mutation in the id3 Patient is going to proceed with a haematopoietic stem idursulfase would be bridging treatment to transplant; a	cell transplant (HSCT		e next 3 r	nonths and treatment wit
 Patient has not required long-term invasive ventilation f (ERT); and 	or respiratory failure p		•	
5 Idursulfase to be administered for a total of 24 weeks (e greater than 0.5 mg/kg every week.	equivalent to 12 weeks	pre- and 1	12 weeks	s post-HSCT) at doses n
ARONIDASE – Special Authority see SA1695 below – Retail Inj 100 U per ml, 5 ml vial		1	✓ A	Idurazyme
SA1695 Special Authority for Subsidy nitial application only from a metabolic physician. Approvals Il of the following:	s valid for 24 weeks for	applicatio	ns meet	ing the following criteria:
1 The patient has been diagnosed with Hurler Syndrome 2 Either:	(mucopolysacchardos	is I-H); and	b	
 Diagnosis confirmed by demonstration of alpha- assay in cultured skin fibroblasts; or 				
2.2 Detection of two disease causing mutations in the to have Hurler syndrome; and	·	0		Ū
 Patient is going to proceed with a haematopoietic stem laronidase would be bridging treatment to transplant; ar Patient has not required long-term invasive ventilation f 	nd			
(ERT); and	or respiratory railure p			and heplacement mera
 5 Laronidase to be administered for a total of 24 weeks (e than 100 units/kg every week. 	equivalent to 12 weeks	pre- and '	12 post-l	HSCT) at doses no great
EVOCARNITINE - Special Authority see SA2040 below - Re	etail pharmacy			
Tab 500 mg	CBS	30		olgar
Cap 250 mg		30		olgar
Cap 500 mg		60		Balance
Oral liq 1 g per 10 ml		118 ml		arnitor S29
Oral liq 500 mg per 10 ml	CBS	300 ml	✓ E	Balance
»SA2040 Special Authority for Subsidy				
itial application only from a metabolic physician. Approvals	s valid for 6 months wh	iere patien	t has a s	suspected inborn error of
tabolism that may respond to carnitine supplementation.				

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN - Special Authority see SA2041 on the next page	e – Retail pharmacy		
Tab 100 mg	CBS	100	 Country Life
Cap 100 mg	CBS	100	 Solgar

30

Subsic (Manufacture	,	ully	Brand or Conorio	
\$	Per	seu ✓	Generic Manufacturer	

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- $\ensuremath{\mathbf{2}}$ The treatment remains appropriate and the patient is benefiting from treatment.

➡SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
 - 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
 - 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
 - 4 Sapropterin to be used alone or in combination with PKU dietary management; and
 - 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy

Soln 100 mg per mlCBS 100 ml 🗸 Amzoate 🐲

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 on the next page -	· Retail pharmacy	/
Grans 483 mg per g2,016.00	174 g OP 🖌	Pheburane

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price)	ę	Fully Subsidised	Brand or Generic
 (.inanalastalei e r 1166) \$	Per	 ✓ 	Manufacturer

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	Solgar
Cap 1,000 mg	CBS	90	 Life Extension
Powder		300 g	 Life Extension

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

Elelyso

1

➡SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

32

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Mouth and Throat

Agents Used in Mouth Ulceration

Agents used in mouth diceration			
BENZYDAMINE HYDROCHLORIDE Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with Endorsement	9.00	500 ml	
	(20.31)	500 m	Difflam
Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly.	· · · ·	as a result of tr	
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	 Stomahesive
	4.55	15 g OP	
	(7.90)		Orabase
	1.52	5 g OP	
	(3.60)		Orabase
Powder		28 g OP	
	(10.95)		Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
	(6.00)	-	Bonjela
TRIAMCINOLONE ACETONIDE			
Paste 0.1%		5 g OP	 Kenalog in Orabase
		- 3 -	
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	🗸 Fungilin
MICONAZOLE			
Oral gel 20 mg per g	4.74	40 g OP	 Decozol
NYSTATIN		5	
Oral liq 100,000 u per ml	1 76	24 ml OP	 Nilstat
		24 111 01	<u> Miotar</u>
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN			

* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PSO2.46

3

<u>Hydroxocobalamin</u>
 <u>Panpharma</u>

	Subsidy		Fully	Brand or
	(Manufacturer's Pr		Subsidised	I Generic
	\$	Per	/	Manufacturer
PYRIDOXINE HYDROCHLORIDE				
a) No more than 100 mg per dose				
b) Only on a prescription				
* Tab 25 mg – No patient co-payment payable		90		Vitamin B6 25
Tab 50 mg	23.45	500	•	Pyridoxine
				multichem
THIAMINE HYDROCHLORIDE – Only on a prescription				
* Tab 50 mg		100		Thiamine multichem
This wire any litch we take Driveland Operators of April 200	7.09		v	Max Health
Thiamine multichem to be Principal Supply on 1 April 202	23			
(Max Health Tab 50 mg to be delisted 1 April 2023)				
VITAMIN B COMPLEX				_ .
* Tab, strong, BPC	7.15	500	v	Bplex
Vitamin C				
ASCORBIC ACID				
a) No more than 100 mg per dose				
 b) Only on a prescription * Tab 100 mg 	10 50	500		Cvite
* Tab 100 Hig	12.50	500	•	CVILE
Vitamin D				
ALFACALCIDOL				
* Cap 0.25 mcg		100	1	One-Alpha
* Cap 1 mcg		100		One-Alpha
				One-Alpha S29 S29
* Oral drops 2 mcg per ml	60.68	20 ml O		One-Alpha
CALCITRIOL				•
* Cap 0.25 mcg	7 89	100	1	Calcitriol-AFT
* Cap 0.5 mcg		100		Calcitriol-AFT
COLECALCIFEROL				
Correction 2002 (Solution) – Maximum of 12 cap per prescription (Solution) – Maximum of 12 cap per prescription (Solution) – Maximum of 12 cap per prescription)	on 2.05	12	1	Vit.D3
* Oral liq 188 mcg per ml (7,500 iu per ml)		4.8 ml C		Puria
				i unu
Multivitamin Preparations				
MULTIVITAMIN RENAL – Special Authority see SA1546 below –	Retail pharmacy	/		
ж Сар		30	1	Clinicians Renal Vit
SA1546 Special Authority for Subsidy				
initial application from any relevant practitioner. Approvals valid	l without further r	enewal ur	less noti	fied for applications meeting
the following criteria:				
Either:				
1 The patient has chronic kidney disease and is receiving eit	ther peritoneal di	alysis or h	aemodia	ysis; or
2 The patient has chronic kidney disease grade 5, defined a 15 ml/min/1.73 m ² body surface area (BSA).				
MULTIVITAMINS – Special Authority see SA1036 on the next pa	ge – Retail pharr	nacy		
* Powder	•	200 g O	Р 🗸	Paediatric Seravit

	Subsidy (Manufacturer's Price)	Suk	Fully sidised	Brand or Generic
	(Manulacturers Frice) \$	Per		Manufacturer
► SA1036 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid	d without further rene	ewal unles	ss notified	d where the patient has
inborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid without	further renewal unle	ee notified	l whoro n	ationt has had a provinus
approval for multivitamins.		ss nouneu	i wilele p	allent has had a previous
VITAMINS				
* Tab (BPC cap strength)		1,000	✓ <u>M</u>	vite
* Cap (fat soluble vitamins A, D, E, K) – Special Authority see SA1720 below – Retail pharmacy		60	🖌 V	itabdeck
SA1720 Special Authority for Subsidy	20.40	00	•••	habacon
Initial application from any relevant practitioner. Approvals valid	d without further rene	ewal unles	ss notified	d for applications meeting
the following criteria:				
Any of the following:				
 Patient has cystic fibrosis with pancreatic insufficiency; or Patient is an infant or child with liver disease or short gut s 	wndrome: or			
3 Patient has severe malabsorption syndrome.	syndrome, or			
Minerals				
Calcium				
CALCIUM CARBONATE				
* Tab 1.25 g (500 mg elemental)		250		alci-Tab 500
* Tab eff 1.25 g (500 mg elemental) – Subsidy by endorsement	nt260.00	100	✓ C	alcium 500 mg Hexal S29
Subsidy by endorsement – Only when prescribed for pae	ediatric patients (< 5	vears) wh	nere calci	
considered unsuitable.		, ,		
CALCIUM GLUCONATE				
* Inj 10%, 10 ml ampoule		10	🗸 M	ax Health -
	64.00	20	. / M	Hameln S29 ax Health S29
	64.00	20	• IV	ax Health 529
lodine				
POTASSIUM IODATE				
* Tab 253 mcg (150 mcg elemental iodine)		90	✓ <u>N</u>	euroTabs
Iron				
FERROUS FUMARATE				
* Tab 200 mg (65 mg elemental)		100	✓ <u>F</u>	erro-tab
FERROUS FUMARATE WITH FOLIC ACID	5.00	400	<i>.</i> -	
* Tab 310 mg (100 mg elemental) with folic acid 350 mcg	5.98	100	✓ <u>F</u>	erro-F-Tabs
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental)	2 55	30	✓ F	errograd
 * Oral lig 30 mg (6 mg elemental) per 1 ml 		500 ml		erodan
IRON (AS FERRIC CARBOXYMALTOSE) - Special Authority se		ext page -	_	
Inj 50 mg per ml, 10 ml vial		1		erinject

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$ P	Ver ✓	Manufacturer

⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE

* Inj 50 mg per ml, 2 ml ampoule	5	✓ Ferrosig
Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%	355 ml	 Phillips Milk of Magnesia 529
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule25.53	10	✓ <u>Martindale</u>
Zinc		
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)11.00	100	✓ Zincaps

Subsidised

Per

Fully

Subsidy (Manufacturer's Price) \$ Brand or Generic

Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

► SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.
- Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA1775 above - Retail pharmacy

Wastage claimable

Inj 1,000 iu in 0.5 ml, syringe	250.00	6	 Binocrit
Inj 2,000 iu in 1 ml, syringe		6	 Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	 Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	 Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	 Binocrit
Inj 6,000 iu in 0.6 ml, syringe		6	 Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	 Binocrit
Inj 10,000 iu in 1 ml, syringe		6	 Binocrit
Inj 40,000 iu in 1 ml, syringe		1	 Binocrit

		Subsidy (Manufacturer's Price \$	e) Subsi Per	Fully idised	Brand or Generic Manufacturer
Μ	egaloblastic				
FO	LIC ACID				
	Tab 0.8 mg		1,000		olic Acid multichem
*	Tab 5 mg	5.82	100		<u>olic Acid Mylan</u> olic Acid Viatris
	Oral liq 50 mcg per ml		25 ml OP		iomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial		1	 Alprolix
Inj 500 iu vial	1,225.00	1	 Alprolix
Inj 1,000 iu vial	2,450.00	1	Alprolix
Inj 2,000 iu vial	4,900.00	1	Alprolix
Inj 3,000 iu vial	7,350.00	1	Alprolix
Inj 4,000 iu vial		1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 below Wastage claimable	– Retail pharmacy		
Tab 25 mg	1,550.00	28	Revolade
Tab 50 mg	3,100.00	28	 Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:

38

3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

Inj 30 mg in 1 ml vial3,570.00	1	 Hemlibra
Inj 60 mg in 0.4 ml vial7,138.00	1	 Hemlibra
Inj 105 mg in 0.7 ml vial	1	 Hemlibra
Inj 150 mg in 1 ml vial17,846.00	1	 Hemlibra

► SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Either:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
- 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.
- Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:
 - 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose
 - period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
 - 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	NovoSeven RT
Inj 2 mg syringe	2,356.60	1	NovoSeven RT
Inj 5 mg syringe		1	NovoSeven RT
Inj 8 mg syringe		1	NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	🗸 FEIBA NF
Inj 1,000 U	2,630.00	1	🖌 FEIBA NF
lnj 2,500 U	6,575.00	1	🖌 FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

inj 250 lu pretilied syringe	 1	🗸 Xyntha
Inj 500 iu prefilled syringe	 1	 Xyntha
Inj 1,000 iu prefilled syringe	1	🗸 Xyntha
Inj 2,000 iu prefilled syringe	1	🗸 Xyntha
Inj 3,000 iu prefilled syringe	1	🗸 Xyntha

NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	 1	RIXUBIS
Inj 1,000 iu vial	 1	RIXUBIS
	 1	RIXUBIS
Inj 3,000 iu vial	 1	RIXUBIS

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [Xpharm]

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. 1 ✓ Advate 1 ✓ Advate 1 Advate ✓ Advate 1 ✓ Advate 1 1 ✓ Advate

	0.1.11			
(N)	Subsidy lanufacturer's Price)	S	Fully bsidised	Brand or Generic
(1)	\$	Per		Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) – [Xnharm]			
For patients with haemophilia. Rare Clinical Circumstances Bra		e recom	binant fa	ctor VIII. Access to funded
treatment is managed by the Haemophilia Treaters Group in co				
subject to criteria.	ijunouon mar aro i	lationa	rnaomoj	onnia managomont oroup,
Inj 250 iu vial	237.50	1	1	Kogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Ini 2.000 iu vial		1		Kogenate FS
Inj 3,000 iu vial	.2,850.00	1		Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] -	Xnharml			Ū
For patients with haemophilia A receiving prophylaxis treatment		d treatn	nent is m	anaged by the Haemophilia
Treaters Group in conjunction with the National Haemophilia Ma		anoun		anagoa by the Haemophina
Inj 250 iu vial	0 0 1	1	1	Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial		1		Adynovate
Inj 2,000 iu vial	,	1		Adynovate
SODIUM TETRADECYL SULPHATE	,	-	-	,,
* Inj 3% 2 ml	28 50	5		
* IIJ 5 /6 2 III	(73.00)	5		Fibro-vein
	(73.00)			
TRANEXAMIC ACID				
Tab 500 mg	10.45	60	✓	Mercury Pharma
Mercury Pharma to be Principal Supply on 1 June 2023				
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO	8.00	5	 Image: A second s	Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO	9.21	5	 Image: A second s	Konakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	14.05	990		Ethics Aspirin EC
5	14.95	990	•	Ethics Aspirin EC
CLOPIDOGREL				.
* Tab 75 mg	4.60	84	•	Clopidogrel
			_	Multichem
	5.07			Arrow - Clopid
Arrow - Clopid to be Principal Supply on 1 May 2023				
(Clopidogrel Multichem Tab 75 mg to be delisted 1 May 2023)				
DIPYRIDAMOLE				
* Tab long-acting 150 mg	13.93	60	 Image: A second s	Pytazen SR
TICAGRELOR - Special Authority see SA1955 on the next page -	Retail pharmacy			
Brand switch fee payable (Pharmacode 2653206) - see page 24				
* Tab 90 mg		56	1	Ticagrelor Sandoz
			-	

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	<u> </u>	Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	1	Manufacturer

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer
Heparin and Antagonist Preparations				
ENOXAPARIN SODIUM - Special Authority see SA2152 below -	 Retail pharmacy 			
Inj 20 mg in 0.2 ml syringe		10	🗸 C	lexane
Inj 40 mg in 0.4 ml syringe		10	🗸 C	lexane
Inj 60 mg in 0.6 ml syringe	60.67	10	🗸 C	lexane
Inj 80 mg in 0.8 ml syringe		10	🗸 C	lexane
Inj 100 mg in 1 ml syringe		10	🗸 C	lexane
Inj 120 mg in 0.8 ml syringe		10	🗸 C	lexane Forte
Inj 150 mg in 1 ml syringe		10	✓ C	lexane Forte

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and

3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		bsidised	
	\$	Per	1	Manufacturer
HEPARIN SODIUM				
Inj 1,000 iu per ml, 5 ml ampoule		50		Pfizer
Inj 5,000 iu per ml, 5 ml vial		10	-	Heparin Sodium
				Panpharma
Inj 5,000 iu per ml, 1 ml		5	1	DBL Heparin
				Sodium S29
	70.33		✓	Hospira
Inj 5,000 iu per ml, 5 ml ampoule		50	✓	Pfizer
Inj 25,000 iu per ml, 0.2 ml	22.42	5	✓	Hospira
	42.40		1	Heparin DBL S29
	482.20	50		Heparin DBL \$29
(Pfizer Inj 5,000 iu per ml, 5 ml ampoule to be delisted 1 July 202				· · · · ·
HEPARINISED SALINE	-)			
Inj 10 iu per ml, 5 ml	65.48	50	1	Pfizer
	00.40	50	•	F 11261
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg – No more than 2 cap per day		60	1	Pradaxa
Cap 110 mg		60	1	Pradaxa
Cap 150 mg		60	✓	Pradaxa
RIVAROXABAN				
Tab 10 mg – No more than 1 tab per day	83 10	30	1	Xarelto
Tab 15 mg – Up to 14 tab available on a PSO		28		Xarelto
Tab 20 mg		28		Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	1	Coumadin
ጥ Tau Tiny		100	-	Marevan
* Tab 2 mg	••••	50		Coumadin
* Tab 2 mg		100		Marevan
* Tab 5 mg		50	-	Coumadin
1 uo o mg	11.48	100	-	Marevan
	0.11	100		marevan

Blood Colony-stimulating Factors

FILGRASTIM - Special Authority see SA1259 below - Retail ph	armacy		
Inj 300 mcg per 0.5 ml prefilled syringe		10	 <u>Nivestim</u>
Inj 480 mcg per 0.5 ml prefilled syringe	148.58	10	✓ Nivestim

➡SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

44

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or

continued...

	Subsidy (Manufacturer's Price))	Fully Subsidised	Brand or Generic
	\$	Per	1	Manufacturer
continued				
4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10				
5 Treatment of drug-induced prolonged neutropenia (ANC	,			
Note: *Febrile neutropenia risk greater than or equal to 20% after European Organisation for Research and Treatment of Cancer (tother	risk factors	s as defined by the
PEGFILGRASTIM - Special Authority see SA1912 below - Retain			-	
Inj 6 mg per 0.6 ml syringe		1		Ziextenzo
Ziextenzo to be Principal Supply on 1 June 2023	1,080.00		v	Neulastim
(Neulastim Inj 6 mg per 0.6 ml syringe to be delisted 1 June 2020	3)			
► SA1912 Special Authority for Subsidy	,			
Initial application only from a relevant specialist, vocationally re	gistered general pra	ctitione	r or medic	al practitioner on the
recommendation of a relevant specialist. Approvals valid without	t further renewal unle	ess not	ified where	e used for prevention of
neutropenia in patients undergoing high risk chemotherapy for ca				
Note: *Febrile neutropenia risk greater than or equal to 5% after		other ri	sk factors	as defined by the European
Organisation for Research and Treatment of Cancer (EORTC) g	uidelines.			
Fluids and Electrolytes				
Intravenous Administration				
GLUCOSE [DEXTROSE]				
* Inj 50%, 10 ml ampoule - Up to 5 inj available on a PSO		5	✓	Biomed
* Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO	15.00	1	✓	Biomed
POTASSIUM CHLORIDE				
* Inj 75 mg per ml, 10 ml	65.00	50	v	Juno
SODIUM BICARBONATE			_	
Inj 8.4%, 50 ml	21.40	1	-	Biomed
a) Up to 5 inj available on a PSO				
b) Not in combinationInj 8.4%, 100 ml	21.95	1	1	Biomed
a) Up to 5 inj available on a PSO	21.95	1	•	Diomed
b) Not in combination				
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Not funded for nebulise	r use except when u	sed in o	conjunctior	n with an antibiotic intended
for nebuliser use.				
Inj 0.9%, bag – Up to 2000 ml available on a PSO		500 m		Baxter
		1,000 n		Baxter
Only if prescribed on a prescription for renal dialysis, ma for emergency use. (500 ml and 1,000 ml packs)	aternity or post-natal	care in	une nome	or the patient, or on a PSO
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	1	Biomed
For Sodium chloride oral liquid formulation refer Standa				
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20		Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO		50		Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	v	Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)	000	4.05		-
Infusion	CBS	1 OP	v 1	TPN

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully sidised	Brand or Generic Manufacturer
WATER				
 On a prescription or Practitioner's Supply Order only we Schedule requiring a solvent or diluent; or On a bulk supply order; or When used in the extemporaneous compounding of ey When used for the dilution of sodium chloride soln 7% to 	e drops; or		ection lis	sted in the Pharmaceutical
Inj 10 ml ampoule – Up to 5 inj available on a PSO Inj 20 ml ampoule – Up to 5 inj available on a PSO		50 20	✓ Pi ✓ <u>Fi</u>	fizer resenius Kabi
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder		300 g OP	✔ Ca	alcium Resonium
COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO	9.53	50	✓ <u>E</u>	lectral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)	8.55	1,000 ml OP		edialyte - Bubblegum
PHOSPHORUS Tab eff 500 mg (16 mmol)		100	🗸 Pi	hosphate Phebra
 POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq) 	5.26 (17.10)	60	C	hlorvescent
* Tab long-acting 600 mg (8 mmol) SODIUM BICARBONATE		200	✓ SI	pan-K
Cap 840 mg	8.52	100	-	odibic odibic
SODIUM POLYSTYRENE SULPHONATE Powder		454 g OP	✓ R	esonium-A

	Subsidy		Fully	Brand or
	(Manufacturer's Price \$	Per	Subsidised	Generic Manufacturer
Alpha-Adrenoceptor Blockers				
Alpha Adrenoceptor Blockers				
00XAZOSIN ₭ Tab 2 mg		500	 Image: A second s	Doxazosin Clinect
₭ Tab 4 mg		500	✓ 1	Doxazosin Clinect
HENOXYBENZAMINE HYDROCHLORIDE				
← Cap 10 mg	65.00	30	 Image: A second s	BNM S29
	216.67	100	✓	Dibenzyline S29
RAZOSIN				
• Tab 1 mg	5.53	100	v	Arrotex-Prazosin S29 S29
• Tab 2 mg	7.00	100	•	Arrotex-Prazosin S29 S29
- Tab 5 mg	11.70	100	•	Arrotex-Prazosin S29 S29
ACE Inhibitors				
APTOPRIL				-
APTOPRIL		95 ml Ol	P 🗸 (Capoten
APTOPRIL ← Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement	Je.			
APTOPRIL ← Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr	je. vere taking cilazapril pri	or to 1 N	<i>l</i> lay 2021 a	and the prescription is
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril.	le. vere taking cilazapril pri escription as endorsed	or to 1 N where ti	lay 2021 a nere exists	and the prescription is a record of prior
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg	ie. vere taking cilazapril priv rescription as endorsed	or to 1 N	/lay 2021 a nere exists ✔ :	and the prescription is a record of prior Zapril
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg	ie. vere taking cilazapril privescription as endorsed 2.69 5.79	or to 1 N where th 90	/lay 2021 a nere exists ✓ : ✓ :	and the prescription is a record of prior
 APTOPRIL Oral liq 5 mg per ml	ie. vere taking cilazapril privescription as endorsed 2.69 5.79	or to 1 M where th 90 90	/lay 2021 a nere exists ✓ : ✓ :	and the prescription is a record of prior Zapril Zapril
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg Tab 5 mg NALAPRIL MALEATE	yere taking cilazapril privescription as endorsed 2.69 5.79 	or to 1 M where th 90 90	May 2021 a here exists	and the prescription is a record of prior Zapril Zapril Zapril Acetec
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg Tab 2.5 mg Tab 5 mg VALAPRIL MALEATE Tab 5 mg Tab 10 mg	ue. vere taking cilazapril privescription as endorsed 2.69 5.79 10.05 	or to 1 M where th 90 90 90 100 100	May 2021 a nere exists v : v : v : v : v : v :	and the prescription is a record of prior Zapril Zapril Zapril Acetec Acetec
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg Tab 5 mg NALAPRIL MALEATE Tab 5 mg Tab 10 mg Tab 20 mg	ue. vere taking cilazapril privescription as endorsed 2.69 5.79 10.05 	or to 1 M where th 90 90 90 100	May 2021 a nere exists v : v : v : v : v : v :	and the prescription is a record of prior Zapril Zapril Zapril Acetec
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag LAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg Tab 2.5 mg Tab 5 mg NALAPRIL MALEATE Tab 5 mg Tab 10 mg Tab 20 mg	ue. vere taking cilazapril privescription as endorsed 2.69 5.79 10.05 1.82 2.02 2.42	or to 1 M where th 90 90 90 100 100 100	Aay 2021 a nere exists 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	and the prescription is a record of prior Zapril Zapril Zapril Acetec Acetec Acetec
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg Tab 2.5 mg Tab 5 mg NALAPRIL MALEATE Tab 5 mg Tab 10 mg Tab 20 mg SINOPRIL	ue. vere taking cilazapril privescription as endorsed 2.69 5.79 10.05 1.82 2.02 2.42	or to 1 M where th 90 90 90 100 100	Aay 2021 a nere exists 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	and the prescription is a record of prior Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg Tab 5.5 mg Tab 5 mg NALAPRIL MALEATE Tab 5 mg SINOPRIL Tab 20 mg SINOPRIL Tab 5 mg	yere taking cilazapril privescription as endorsed 2.69 5.79 10.05 1.82 2.02 2.42 	or to 1 M where th 90 90 90 100 100 100	Aay 2021 a here exists 7 2 7 2 7 2 7 2 7 2 7 2 7 2 7 2 7 2 7 2	and the prescription is a record of prior Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril Feva Lisinopril Ethics Lisinopril
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg Tab 5 mg NALAPRIL MALEATE Tab 5 mg Tab 20 mg SINOPRIL Tab 20 mg Tab 10 mg Tab 10 mg	yere taking cilazapril privescription as endorsed 	or to 1 N 90 90 90 100 100 100 90	Aay 2021 a nere exists 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	and the prescription is a record of prior Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril Feva Lisinopril
APTOPRIL € Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. € Tab 0.5 mg E Tab 5 mg E Tab 5 mg E Tab 5 mg E Tab 20 mg SINOPRIL € Tab 10 mg € Tab 10 mg € Tab 20 mg	yere taking cilazapril privescription as endorsed 	or to 1 N where th 90 90 90 100 100 100 90 90	Aay 2021 a nere exists 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	and the prescription is a record of prior Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril
APTOPRIL Coral liq 5 mg per ml Oral liquid restricted to children under 12 years of ag ILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril. Tab 0.5 mg Tab 5 mg Tab 5 mg NALAPRIL MALEATE Tab 5 mg Tab 5 mg Tab 20 mg SINOPRIL Tab 10 mg Tab 20 mg ERINDOPRIL ERINDOPRIL Tab 20 mg	yere taking cilazapril privescription as endorsed 2.69 5.79 10.05 1.82 2.42 2.42 11.07 11.67 1.67 1.58	or to 1 N where th 90 90 90 100 100 100 90 90 90 90 30	Aay 2021 a nere exists 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	and the prescription is a record of prior Zapril Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Coversyl
CILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who w endorsed accordingly. Pharmacists may annotate the pr dispensing of cilazapril.	yere taking cilazapril privescription as endorsed 2.69 5.79 10.05 1.82 2.02 2.42 11.07 11.67 11.67 14.69 1.58 2.95	or to 1 N where th 90 90 90 100 100 100 90 90 90	Aay 2021 a here exists 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	and the prescription is a record of prior Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril Ethics Lisinopril

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
QUINAPRIL	Ŷ			Manufacturer
* Tab 5 mg		90	1	Arrow-Quinapril 5
* Tab 10 mg		90		Arrow-Quinapril 10
* Tab 20 mg		90		Arrow-Quinapril 20
RAMIPRIL				
* Cap 1.25 mg	6.90	90	1	Tryzan
Tryzan to be Principal Supply on 1 May 2023				
* Cap 2.5 mg	6.60	90	✓	Tryzan
Tryzan to be Principal Supply on 1 May 2023				•
* Cap 5 mg	6.75	90	✓	Tryzan
Tryzan to be Principal Supply on 1 May 2023				
* Cap 10 mg	7.05	90	✓	Tryzan
Tryzan to be Principal Supply on 1 May 2023				
ACE Inhibitors with Diuretics				
QUINAPRIL WITH HYDROCHLOROTHIAZIDE – Subsidy by e Subsidy by endorsement – Subsidised for patients who wer 2022 and the prescription is endorsed accordingly. Pharma exists a record of prior dispensing of quinapril with hydroch	e taking quinapril with lacists may annotate the orothiazide.	pres	scription as	endorsed where there
Tab 10 mg with hydrochlorothiazide 12.5 mg		30		Accuretic 10
Tab 20 mg with hydrochlorothiazide 12.5 mg	5.25	30	~	Accuretic 20
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
* Tab 4 mg	2.00	90	1	Candestar
* Tab 8 mg	2.28	90	✓	Candestar

* Tab 16 mg	3.31	90	 Candestar
* Tab 32 mg		90	✓ Candestar
LOSARTAN POTASSIUM			
* Tab 12.5 mg	1.56	84	 Losartan Actavis
* Tab 25 mg		84	 Losartan Actavis
* Tab 50 mg		84	 Losartan Actavis
* Tab 100 mg		84	 Losartan Actavis
5			

Angiotensin II Antagonists with Diuretics

LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE		
* Tab 50 mg with hydrochlorothiazide 12.5 mg4.00	30	Arrow-Losartan &
		Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1905 on	the next page -	Retail phar	macy
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	 Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	 Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	 Entresto 97/103

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) 🗧	Subsidised	Generic	
\$	Per	1	Manufacturer	

➡SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and

3 Either:

- 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
- 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 117	For lignocaine	hydrochloride refer to	NERVOUS SYSTEM,	Anaesthetics, Local, page 117
---	----------------	------------------------	-----------------	-------------------------------

AMIODARONE HYDROCHLORIDE

AMIODATIONE TIT DITOCITEOTIDE		
▲ Tab 100 mg	30	 <u>Aratac</u>
▲ Tab 200 mg	30	 Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a PSO 9.12	6	 Cordarone-X
15.22	10	Max Health
ATROPINE SULPHATE		
ing ooo meg per mi, i m ampeare op to o mjaranabio en a	10	. Martindala
PSO15.09	10	✓ <u>Martindale</u>
DIGOXIN		
* Tab 62.5 mcg – Up to 30 tab available on a PSO	240	Lanoxin PG
* Tab 250 mcg – Up to 30 tab available on a PSO 16.90	240	 Lanoxin
* Oral liq 50 mcg per ml 16.60	60 ml	 Lanoxin
		Lanoxin Paediatric
		Elixir S29
		Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE	100	
▲ Cap 100 mg23.87	100	 Rythmodan
FLECAINIDE ACETATE		
▲ Tab 50 mg 19.95	60	 Flecainide BNM
▲ Cap long-acting 100 mg	90	 Flecainide
		Controlled
		Release Teva
▲ Cap long-acting 200 mg	90	Flecainide
		Controlled
		Release Teva
Inj 10 mg per ml, 15 ml ampoule	5	✓ Tambocor
	5	
MEXILETINE HYDROCHLORIDE		
▲ Cap 150 mg162.00	100	Teva S29
▲ Cap 250 mg	100	Teva S29

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PROPAFENONE HYDROCHLORIDE Tab 150 mg	40.90	50	1	Rytmonorm
Antihypotensives				
MIDODRINE - Special Authority see SA1474 below - Retail phar	macy			
Tab 2.5 mg		100	~	Midodrine Medsurge
	53.00		✓	Gutron
Tab 5 mg	59.98	100	~	Midodrine Medsurge
	79.00		✓	Gutron
(Cutron Tab 0.5 mg to be delicted 1 August 2022)				

(Gutron Tab 2.5 mg to be delisted 1 August 2023)

(Gutron Tab 5 mg to be delisted 1 August 2023)

➡SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATE	ENOLOL			
*	Tab 50 mg	9.33	500	 ✓ <u>Mylan Atenolol</u> ✓ Viatris
*	Tab 100 mg		500	 Mylan Atenolol
*	Oral liq 25 mg per 5 ml	21.25	300 ml OP	✓ Atenolol AFT
		38.20		S29 S29 Sessential Generics S29
		49.85		 Atenolol AFT
	Restricted to children under 12 years of age.			
BIS	OPROLOL FUMARATE			
*	Tab 2.5 mg	1.84	90	 Bisoprolol Mylan Bisoprolol Viatris
*	Tab 5 mg	2.55	90	 Bisoprolol Mylan Bisoprolol Viatris
*	Tab 10 mg		90	 Bisoprolol Mylan Bisoprolol Viatris
CA	RVEDILOL			
*	Tab 6.25 mg	2.24	60	 Carvedilol Sandoz
*	Tab 12.5 mg		60	 Carvedilol Sandoz
*	Tab 25 mg		60	 Carvedilol Sandoz

_					
		Subsidy		Fully	
		(Manufacturer's Price)		Subsidised	
		\$	Per		Manufacturer
LA	BETALOL				
*	Tab 100 mg		100	✓	Trandate
*	Tab 200 mg		100	✓	Trandate
*	Inj 5 mg per ml, 20 ml ampoule		5		
		(88.60)			Trandate
*	inj 5 mg per ml, 20 ml vial		1		
		(48.20)			Alvogen S29
ME	TOPROLOL SUCCINATE				
*	Tab long-acting 23.75 mg	1.45	30	1	Betaloc CR
*	Tab long-acting 47.5 mg.	1.43	30	1	Betaloc CR
*	Tab long-acting 95 mg	2.15	30	1	Betaloc CR
*	Tab long-acting 190 mg		30	✓	Betaloc CR
ME	TOPROLOL TARTRATE				
*	Tab 50 mg		100	1	IPCA-Metoprolol
*	Tab 100 mg		60		IPCA-Metoprolol
*	Tab long-acting 200 mg		28		Slow-Lopresor
*	Inj 1 mg per ml, 5 ml vial		5		Metoprolol IV Mylan
					Metoprolol IV Viatris
NA	DOLOL				
*	Tab 40 mg		100	1	Nadolol BNM
*	Tab 80 mg		100		Nadolol BNM
PR	OPBANOLOL				
*	Tab 10 mg	7.04	100	1	Drofate
*	Tab 40 mg		100		IPCA-Propranolol
*	Cap long-acting 160 mg		100		Cardinol LA
*	Oral lig 4 mg per ml – Special Authority see SA1327 below				
	Retail pharmacy		500 m	nl 🗸	Roxane-
					Propranolol S29
					· · · · · · · · · · · · · · · · · · ·

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

*	Tab 80 mg	7.50	500	🗸 Mylan
	Tab 160 mg14		100	 Mylan

(N	Subsidy lanufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Calcium Channel Blockers				
Dihydropyridine Calcium Channel Blockers				
MLODIPINE	4.00			
₭ Tab 2.5 mg ₭ Tab 5 mg		90 90	-	/ <u>asorex</u> /asorex
k Tab 10 mg		90		asorex
ELODIPINE			-	
k Tab long-acting 2.5 mg		30	✓ F	Plendil ER
★ Tab long-acting 5 mg		90	-	elo 5 ER
Fab long-acting 10 mg	4.32	90	✓ <u>F</u>	elo 10 ER
IIFEDIPINE				
€ Tab long-acting 10 mg	18.80	56	ا 🗸	ensipine MR10 S29
 Tab long-acting 20 mg 	9.12	50	🗸 N	lylan (12 hr
				release) S29
	17.72	100	N	lyefax Retard
Tab long-acting 30 mg	4.78	14	🗸 N	Íylan Italy (24 hr
				release) S29
	34.10	100	🗸 N	lylan (24 hr
				release) S29
 Tab long-acting 60 mg 	52.81	100	🗸 N	lylan (24 hr
				release) S29
Other Calcium Channel Blockers				
ILTIAZEM HYDROCHLORIDE				
Cap extended-release 120 mg	44.40	100	✓ ↓	ccord S29
Cap long-acting 120 mg	33.42	500	✓ ↓	po-Diltiazem CD
	65.35		✓ [Diltiazem CD Clinect
Diltiazem CD Clinect to be Principal Supply on 1 June 2023				
Cap long-acting 180 mg		30	_	Cardizem CD
Cap long-acting 240 mg		30	✓ <u>(</u>	Cardizem CD
Accord S29 Cap extended-release 120 mg to be delisted 1 June 2	,			
Apo-Diltiazem CD Cap long-acting 120 mg to be delisted 1 June 20	123)			
ERHEXILINE MALEATE	60.00	100		lovoia
• Tab 100 mg		100	✓ F	Pexsig
	7.01	100		- i -
K Tab 40 mg		100		soptin
₭ Tab 80 mg		100 100		soptin
Tab long-acting 120 mg	30.02	100		soptin Retard S29 soptin SR
Tab long-acting 240 mg	15.12	30		soptin SR
 Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a 		00		
PSO	25.00	5	🗸 I:	soptin
		5		

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Per	Subsidised	Generic
	\$	rei		Manufacturer
Centrally-Acting Agents				
CLONIDINE				
* Patch 2.5 mg, 100 mcg per day – Only on a prescription		4		Mylan
* Patch 5 mg, 200 mcg per day – Only on a prescription		4		<u>Mylan</u>
* Patch 7.5 mg, 300 mcg per day – Only on a prescription		4	~	Mylan
CLONIDINE HYDROCHLORIDE	00.00	440		
* Tab 25 mcg		112		Clonidine Teva
 * Tab 150 mcg * Inj 150 mcg per ml, 1 ml ampoule 		100 10		<u>Catapres</u> Medsurge
METHYLDOPA		10	•	meusurge
* Tab 250 mg	15 10	100	1	Methyldopa Mylan
* Tab 200 mg	52.85	500		Methyldopa Mylan
	02.00			S29 S29
Diuretics				
Less Dissetter				
Loop Diuretics				
BUMETANIDE				
* Tab 1 mg	4.91	30	✓	Burinex S29 S29
5	16.36	100	1	Burinex
* Inj 500 mcg per ml, 4 ml vial	7.95	5	1	Burinex
FUROSEMIDE [FRUSEMIDE]				
Tab 40 mg – Up to 30 tab available on a PSO	8.00	1,000		IPCA-Frusemide
* Tab 500 mg		50		Urex Forte
	89.48		~	Furosemid-
				Ratiopharm S29
	169.96	100	1	Furosemid-
	100.00	100		Ratiopharm S29
* Oral liq 10 mg per ml		0 ml C		Lasix
* Inj 10 mg per ml, 25 ml ampoule		6		Lasix
* Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a P	502.40	5	•	Furosemide-Baxter
Potassium Sparing Diuretics				
AMILORIDE HYDROCHLORIDE				
Oral liq 1 mg per ml		5 ml C	P 🗸	Biomed
EPLERENONE - Special Authority see SA1728 below - Retail p	harmacy			
Tab 25 mg		30		Inspra
Tab 50 mg	25.00	30	✓	Inspra
► SA1728 Special Authority for Subsidy	I will and finally an orange			

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Patient has heart failure with ejection fraction less than 40%; and

2 Either:

- 2.1 Patient is intolerant to optimal dosing of spironolactone; or
- 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

(Ma	Subsidy Inufacturer's Pr \$	rice) Subs Per	Fully Brand or sidised Generic ✓ Manufacturer
/ETOLAZONE			
Tab 5 mg	CBS	1 50	 Metolazone S29 Zaroxolyn S29
PIRONOLACTONE		50	
₭ Tab 25 mg		100	✓ <u>Spiractin</u>
₭ Tab 100 mg Oral lig 5 mg per ml		100 25 ml OP	 ✓ <u>Spiractin</u> ✓ Biomed
		20111101	· Biolinea
Potassium Sparing Combination Diuretics			
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE			6 –
★ Tab 5 mg with furosemide 40 mg	8.63	28	 Frumil
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE ₭ Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	✓ Moduretic
Thiazide and Related Diuretics			
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]			
K Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	 <u>Arrow-</u> Bendrofluazide
			Bonaronaulao
May be supplied on a PSO for reasons other than emergenc		500	✓ Arrow-
	04.00	500	Bendrofluazide
CHLOROTHIAZIDE			
Oral liq 50 mg per ml	27.82	25 ml OP	✓ Biomed
Tab 25 mg	3.90	30	 Igroton S29
I have been been been been been a familie of the	6.95	50	 Hygroton
Hygroton to be Principal Supply on 1 April 2023 Igroton S29 Tab 25 mg to be delisted 1 April 2023)			
NDAPAMIDE			
K Tab 2.5 mg	10.45	90	Dapa-Tabs
-	11.61	100	✓ Mylan
			Indapamide S29

(Mylan Indapamide S29 Tab 2.5 mg to be delisted 1 August 2023)

Vasopressin receptor antagonists

TOLVAPTAN - Special Authority see SA2166 on the next page - Retail phar	macy	
Tab 15 mg	28 OP	 Jinarc
Tab 30 mg	28 OP	 Jinarc
Tab 45 mg + 15 mg1,747.00	56 OP	 Jinarc
Tab 60 mg + 30 mg1,747.00		 Jinarc
Tab 90 mg + 30 mg1,747.00		 Jinarc

Subsidy		Fully	Brand or	
(Manufacturer's Price)		ubsidised	Generic	
\$	Per	-	Manufacturer	

⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Either:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents

Fibrates

BEZAFIBRATE		
* Tab 200 mg	90	Bezalip
* Tab long-acting 400 mg21.21	30	 Bezalip Retard
Other Lipid-Modifying Agents		
ACIPIMOX		
* Cap 250 mg21.56	30	 Olbetam S29 S29
25.44		 Olbetam
Resins		
COLESTIPOL HYDROCHLORIDE		
Grans for oral liq 5 g	30	 Colestid
HMG CoA Reductase Inhibitors (Statins)		
ATORVASTATIN		
* Tab 10 mg6.16	500	✓ Lorstat
* Tab 20 mg9.24	500	✓ Lorstat
* Tab 40 mg14.92	500	 Lorstat
* Tab 80 mg26.54	500	Lorstat
PRAVASTATIN		
* Tab 20 mg2.11	28	 Pravastatin Mylan
		 Pravastatin Viatris
* Tab 40 mg3.61	28	Pravastatin Mylan

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

		Subsidy (Manufacturer's Price) \$		Subsidised	Brand or Generic Manufacturer
RO	SUVASTATIN – Special Authority see SA2093 below – Retail	pharmacy			
*	Tab 5 mg		30	✓ <u>F</u>	losuvastatin Viatris
*	Tab 10 mg	2.42	30	🗸 F	losuvastatin Viatris
*	Tab 20 mg	3.92	30	✓ F	losuvastatin Viatris
*	Tab 40 mg	5.28	30	✓ F	losuvastatin Viatris

⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient is considered to be at risk of cardiovascular disease; and
- 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:
 - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
 - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIMVASTATIN

56

*	Tab 10 mg	90	 Simvastatin Mylan
*	Tab 20 mg	90	 Simvastatin Mylan
*	Tab 40 mg	90	 Simvastatin Mylan
			 Simvastatin Viatris
*	Tab 80 mg7.12	90	 Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Special Authority see SA1045 on the next page – Retail pharmacy		
* Tab 10 mg1.95	30	 Ezetimibe Sandoz

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than $10 \times$ normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy					
Tab 10 mg with simvastatin 10 mg	5.15	30	 Zimybe 		
Tab 10 mg with simvastatin 20 mg	6.15	30	 Zimybe 		
Tab 10 mg with simvastatin 40 mg		30	 Zimybe 		
Tab 10 mg with simvastatin 80 mg		30	 Zimybe 		

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

*	Oral pump spray, 400 mcg per dose - Up to 250 dose			
	available on a PSO	7.48	250 dose OP	 Nitrolingual Pump Spray
*	Patch 25 mg, 5 mg per day		30	 Nitroderm TTS
*	Patch 50 mg, 10 mg per day		30	 Nitroderm TTS
	SORBIDE MONONITRATE			
*	Tab 20 mg		100	🗸 Ismo 20
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
*	Tab long-acting 60 mg		90	✓ Duride
S	ympathomimetics			
AD	RENALINE			
	Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a P	SO4.98 12.65	5	 Aspen Adrenaline DBL Adrenaline

Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a PSO27.00	5	 Hospira
49.00	10	Aspen Adrenaline

	Subsidy) 0	Fully Brand or
	(Manufacturer's Price \$	Per	ubsidised Generic Manufacturer
	Ŧ		
Vasodilators			
HYDRALAZINE HYDROCHLORIDE			
	Detail		
* Tab 25 mg – Special Authority see SA1321 below - pharmacy		1	 Hydralazine
phamady		56	✓ Onelink \$29
		84	✓ AMDIPHARM \$29
		100	✓ Onelink \$29
* Inj 20 mg ampoule	25.90	5	 ✓ Apresoline
SA1321 Special Authority for Subsidy		0	Apresonne
Initial application from any relevant practitioner. Appro	wals valid without further ren		ass notified for applications meeting
he following criteria:		ewar unit	cos notified for applications meeting
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure in combination	with a nitrate, in patients who	are intol	erant or have not responded to A
inhibitors and/or angiotensin receptor blockers.			
MINOXIDIL			
Tab 10 mg		100	 Loniten
NICORANDIL			
▲ Tab 10 mg		60	✓ Ikorel
▲ Tab 20 mg		60	✓ Ikorel
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	 Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg		50	 Trental 400
•			
Endothelin Receptor Antagonists			
AMBRISENTAN – Special Authority see SA1702 below	Potail pharmaoy		
Tab 5 mg	, ,	30	 Ambrisentan Mylan
Tab 10 mg		30	✓ Ambrisentan Viatris
	,		 Mylan
SA1702 Special Authority for Subsidy			
Special Authority approved by the Pulmonary Arterial Hy	pertension Panel		
Notes: Application details may be obtained from Pharm	, ac's website <u>schedule.pharm</u>	nac.govt.r	nz/SAForms or:
The Coordinator, PAH Panel			
Pharmac, PO Box 10-254, WELLINGTON			
Tel: (04) 916 7561, Fax: (04) 974 4858, Email: <u>PAH@</u>			
BOSENTAN – Special Authority see SA1991 below – F			/ - · -
Tab 62.5 mg	119.85	60	Bosentan Dr
	110.05	00	Reddy's
Tab 125 mg	119.85	60	 Bosentan Dr Boddula
			Reddy's
SA1991 Special Authority for Subsidy			

► SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	1	Manufacturer	

continued...

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- of the following
- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

	Subsidy (Manufacturer's Price)		Fully sidised	Brand or Generic Manufacturer	
Phosphodiesterase Type 5 Inhibitors	\$	Per		Manufacturer	
SILDENAFIL – Special Authority see SA1992 below – Retail ph	armacy				
Tab 25 mg		4	✓ V	edafil	
Tab 50 mg	1.70	4	✓ V	edafil	
Tab 100 mg	10.20	12	✓ V	edafil	
► SA1992 Special Authority for Subsidy	avent prostitionar Ar		olid with	aut furthar rangur	

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III; or
 - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- Both:
 - 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
 - 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Prostacyclin Analogues				
 EPOPROSTENOL – Special Authority see SA1696 below – Ret Inj 500 mcg vial Inj 1.5 mg vial ⇒SA1696 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertensi Notes: Application details may be obtained from Pharmac's web The Coordinator, PAH Panel Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac 		1 1 <u>c.gov</u>	✓ V	eletri eletri 1 <u>s</u> or:
 ILOPROST – Special Authority see SA1705 below – Retail phar Brand switch fee payable (Pharmacode 2653214) - see pag Nebuliser soln 10 mcg per ml, 2 ml ⇒SA1705 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertensis Notes: Application details may be obtained from Pharmac's web The Coordinator, PAH Panel Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac 	macy a 249 for details 	30 <u>c.gov</u>	_	ebulis 1 <u>s</u> or:

	Subsidy (Manufacturer's Price \$) Sub Per	Fully sidised	Brand or Generic Manufacturer
Antiacne Preparations				
For systemic antibacterials, refer to INFECTIONS, Antibacterials	s, page 90			
ADAPALENE				
a) Maximum of 30 g per prescription				
b) Only on a prescription				
Crm 0.1%		30 g OP	_	Differin
Gel 0.1%		30 g OP	✓ □	Differin
ISOTRETINOIN - Special Authority see SA2023 below - Retail	pharmacy			
Cap 5 mg		60	✓ 0	Dratane
Cap 10 mg		120	✓ 0	Dratane
Cap 20 mg		120	✓ <u>c</u>	Dratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
- 2 Patient is not of child bearing potential.

TRETINOIN

Crm 0.5 mg per g – Maximum of 50 g per prescription15.57	50 g OP	✓ <u>ReTrieve</u>
Antibacterials Topical		
For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90		
HYDROGEN PEROXIDE		
* Crm 1%	10 g OP	 Crystaderm
MUPIROCIN		
Oint 2%	15 g OP	
(11.50)		Bactroban
a) Only on a prescription		
 b) Not in combination 		

	Subsidy (Manufacturer's F		Fully Brand or sidised Generic	
	\$	Per	 Manufacturer 	
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1 59	5 g OP	✓ Foban	
a) Maximum of 5 g per prescription	1.03	5 y 01		
b) Only on a prescription				
c) Not in combination				
Oint 2%	1.59	5 g OP	 Foban 	
 a) Maximum of 5 g per prescription b) Only on a prescription 				
b) Only on a prescriptionc) Not in combination				
SULFADIAZINE SILVER				
Crm 1%		50 g OP	 Flamazine 	
a) Up to 250 g available on a PSO				
b) Not in combination				
Antifungela Tanical				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals, pa	age 97			
AMOROLFINE	-			
a) Only on a prescription				
b) Not in combination			_	
Nail soln 5%	14.93	5 ml OP	✓ MycoNail	
	4.40	00 - 00		
 Crm 1% a) Only on a prescription 	1.10	20 g OP	 Clomazol 	
b) Not in combination				
c) Clomazol to be Principal Supply on 1 April 2023				
* Soln 1%	4.36	20 ml OP		
	(7.55)		Canesten	
 a) Only on a prescription b) Not in combination 				
b) Not in combination				
ECONAZOLE NITRATE Crm 1%	1.00	20 g OP		
Unit 170	(7.78)	20 9 01	Pevaryl	
a) Only on a prescription	(
b) Not in combination				
Foaming soln 1%, 10 ml sachets		3	5	
a). Only an a pressription	(17.92)		Pevaryl	
a) Only on a prescriptionb) Not in combination				

	Subsidy		Fully Brand or
	(Manufacturer's F		idised Generic
	\$	Per	 Manufacturer
MICONAZOLE NITRATE			
* Crm 2%	0.81	15 g OP	 <u>Multichem</u>
a) Only on a prescription			
 b) Not in combination * Lotn 2% 	4 36	30 ml OP	
	(10.03)	00 111 01	Daktarin
a) Only on a prescription	(
b) Not in combination			
* Tinct 2%		30 ml OP	D. I. I.
	(12.10)		Daktarin
a) Only on a prescriptionb) Not in combination			
b) Not in combination			
Antipruritic Preparations			
CALAMINE a) Only on a prescription			
b) Not in combination			
Crm, aqueous, BP	1.08	100 g	 Calamine-AFT
CROTAMITON		-	
a) Only on a prescription			
b) Not in combination			
Crm 10%	3.29	20 g OP	✓ Itch-Soothe
MENTHOL – Only in combination			
1) Only in combination with a dermatological base		orticosteriod -	Plain
With or without other dermatological galenicals	3.		
Crystals	6.02	25 g	✓ MidWest
Ciysials	29.60	20 g	✓ MidWest
	20100		
Corticosteroids Topical			
For systemic corticosteroids, refer to CORTICOSTEROI	DS AND RELATED AGE	NTS, page 80	
Corticosteroids - Plain			
BETAMETHASONE DIPROPIONATE			
Crm 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	 Diprosone

	CIII 0.05%	15 g OP	Diprosone
	36.00	50 g OP	 Diprosone
	Oint 0.05%2.96	15 g OP	 Diprosone
	36.00	50 g OP	 Diprosone
	Oint 0.05% in propylene glycol base4.33	30 g OP	 Diprosone OV
BE	TAMETHASONE VALERATE		
*	Crm 0.1%	50 g OP	 Beta Cream
*	Oint 0.1%	50 g OP	 Beta Ointment
*	Lotn 0.1%	50 ml OP	 Betnovate
CI	OBETASOL PROPIONATE		
	Crm 0.05%	30 g OP	 Dermol
	Oint 0.05%	30 g OP	✓ Dermol

	Subsidy		Fully	Brand or
	(Manufacturer's I	Price) Subs	idised	Generic
	\$	Per	1	Manufacturer
CLOBETASONE BUTYRATE				
Crm 0.05%	5.38	30 g OP		
	(10.00)	0		Eumovate
HYDROCORTISONE				
* Crm 1% – Only on a prescription	1.78	30 g OP	1	Ethics
	17.15	500 g		Hydrocortisone
		y		(PSM)
	20.40		1	Noumed
Ethics to be Principal Supply on 1 April 2023				
* Powder – Only in combination		25 g	1	ABM
Up to 5% in a dermatological base (not proprietary Topica	al Corticosterio		or with	out other dermatological
galenicals		,		0
(Hydrocortisone (PSM) Crm 1% to be delisted 1 August 2023)				
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only o	n			
a prescription		250 ml	1	DP Lotn HC
HYDROCORTISONE BUTYRATE		200 111		DI LOUITIO
Lipocream 0.1%	4.05	100 ~ 00		
		100 g OP 100 g OP		Locoid Lipocream Locoid
Oint 0.1% Milky emul 0.1%		100 g OP 100 ml OP		Locoid Crelo
	12.00	100 IIII OF	•	
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%		15 g OP		Advantan
Oint 0.1%	4.46	15 g OP		<u>Advantan</u>
MOMETASONE FUROATE			_	
Crm 0.1%		15 g OP		Elocon Alcohol Free
0' 10.40	3.10	50 g OP		Elocon Alcohol Free
Oint 0.1%		15 g OP		Elocon
Late 0.40/	2.90	50 g OP	- 1	Elocon
Lotn 0.1%	4.50	30 ml OP	•	<u>Elocon</u>
TRIAMCINOLONE ACETONIDE			_	
Crm 0.02%		100 g OP		Aristocort
Oint 0.02%	6.35	100 g OP	 Image: A second s	Aristocort
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUS Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		
		15 y OP		Fucicort
a) Maximum of 15 g per prescription	(10.43)			
b) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE - Only on a prescript		45 00		
* Crm 1% with miconazole nitrate 2%		15 g OP		Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - Or			_	
Crm 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP		Pimafucort
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP		Pimafucort
(Pimafucort Crm 1% with natamycin 1% and neomycin sulphate 0			9)	
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCII	N AND NYSTA	TIN		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg				
and gramicidin 250 mcg per g – Only on a prescription	3.49	15 g OP		
	(9.28)			Viaderm KC

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully Brand or
	(Manufacturer's	Price) Subsid Per	dised Generic ✓ Manufacturer
	ð	Per	Manufacturer
Barrier Creams and Emollients			
Barrier Greating and Emoments			
Barrier Creams			
DIMETHICONE			
* Crm 5% pump bottle	4.30	500 ml OP	✓ <u>healthE</u>
			Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ healthE
			Dimethicone 10%
ZINC AND CASTOR OIL			
* Oint	4.65	500 g	 Boucher
Emollients			
AQUEOUS CREAM			
* Cm	1 73	500 g	✓ Evara
		500 g	✓ <u>GEM Aqueous</u>
			Cream
(Evara Crm to be delisted 1 April 2023)			<u></u>
CETOMACROGOL			
* Cm BP	1 99	500 g	Cetomacrogol-AFT
		500 g	
CETOMACROGOL WITH GLYCEROL	0.10	500 ml OD	
Crm 90% with glycerol 10%	2.13 2.35	500 ml OP	 ✓ Evara ✓ Pharmacy Health
	2.00		Sorbolene with
			Glycerin
	3.50	1.000 ml OP	✓ Evara
(Pharmacy Health Sorbolene with Glycerin Crm 90% with glycer		.,	
EMULSIFYING OINTMENT		10100 1 0019 2020	7
* Oint BP	3.40	500 g	 Emulsifying
		500 g	Ointment ADE
OIL IN WATER EMULSION * Crm	2.04	500 a	- Eatty Croom AET
-	2.04	500 g	 Fatty Cream AFT
PARAFFIN	4.04	500 × 05	
Oint liquid paraffin 50% with white soft paraffin 50%	4.94	500 g OP	✓ White Soft Liquid
Milling Optimized Description AFT to be Definite to Complete	4 14 0000		Paraffin AFT
White Soft Liquid Paraffin AFT to be Principal Supply or			√ haalihE
Oint liquid paraffin 50% with white soft paraffin 50%,		500 ml OP	 healthE
(healthE Oint liquid paraffin 50% with white soft paraffin 50%, to	be delisted T Ma	ay 2023)	
UREA		100 00	
* Crm 10%	1.37	100 g OP	 healthE Urea Cream

	Subsidy (Manufacturer's P	rico) Subo	Fully Brand or idised Generic
	(Manulaciulei S F	Per	Manufacturer
	*	-	
WOOL FAT WITH MINERAL OIL – Only on a prescription * Lotn hydrous 3% with mineral oil	5 60	1,000 ml	
	(14.96)	1,000 mi	DP Lotion
	()		
	(20.53) 1.40	250 ml OP	Alpha-Keri Lotion
		250 III OF	DP Lotion
	(5.87) 5.60	1,000 ml	DF LOUOT
		1,000 mi	BK Lotion
	(23.91)		BK LOUON
	1.40	250 ml OP	DKLation
	(7.73)		BK Lotion
Other Dermatological Bases			
PARAFFIN			
White soft – Only in combination		450 g	✓ healthE
	19.99	2,500 g	✓ healthE
Only in combination with a dermatological galenical or a			
,			
Minor Skin Infections			
POVIDONE IODINE			
Oint 10%	7.40	65 g OP	 Betadine
	7.40	05 y OF	• Detaume
a) Maximum of 130 g per prescription			
b) Only on a prescription	4.45	100 1	Diadina
Antiseptic Solution 10%		100 ml 15 ml	 ✓ <u>Riodine</u> ✓ Riodine
Antiseptic soln 10%			
	5.40	500 ml	 Riodine
Skin preparation, povidone iodine 10% with 30% alcohol		100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol		100 ml	24
	(7.78)		Pfizer
Parasiticidal Preparations			
DIMETHICONE			
* Lotn 4%		200 ml OP	✓ healthE
			Dimethicone 4%
			Lotion
VERMECTIN – Special Authority see SA1225 below – Retail p	harmaov		
Tab 3 mg – Up to 100 tab available on a PSO		4	✓ Stromectol
1) PSO for institutional use only. Must be endorsed		the institution f	
a valid Special Authority for patient of that institution			or which the 1 SO is required all
2) Ivermectin available on BSO provided the BSO in		aial Authority f	or a patient of the institution
 For the purposes of subsidy of ivermectin, instituti 			
facilities or prisons.	on means age lei	aleu residerilla	i care raciilles, disability care
actitutes of prisons.			

➡SA1225 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

continued...

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

	Subsidy (Manufacturer's Pri \$	ce) Si Per	Fully ubsidised	Brand or Generic Manufacturer
 continued Renewal — (Other parasitic infections) only from an infectiou approvals valid for 1 month for applications meeting the following of the following: Filaricides; or Cutaneous larva migrans (creeping eruption); or 		st, clinical n	nicrobiolog	gist or dermatologist.
3 Strongyloidiasis.				
PERMETHRIN Crm 5% Lotn 5%		30 g OP 30 ml OP		<u>yderm</u> I-Scabies
Psoriasis and Eczema Preparations				
CITRETIN - Special Authority see SA2024 below - Retail pha	rmacv			
Cap 10 mg.		60		lovatretin
Cap 25 mg	41.36	60	✓ <u>N</u>	lovatretin
 SA2024 Special Authority for Subsidy SA2024 Special Authority for Subsidy initial application from any relevant practitioner. Approvals valial of the following: Applicant is a vocationally registered dermatologist, vocative working in a relevant scope of practice; and Applicant has an up to date knowledge of the safety issued Either: Patient is of child bearing potential and the possible treatment and patient has been counselled and ur pregnancy and that they must not become pregna completion of treatment; or Patient is not of child bearing potential. Patient is of child bearing potential. Patient is of child bearing potential and the possibility of p treatment and patient has been counselled and ur or become pregna completion of treatment; or Patient is not of child bearing potential. 	ionally registered as around acitretin lity of pregnancy h iderstands the risk nt during treatmen ar for applications regnancy has bee	general pra and is com nas been e: of teratoge t and for a meeting th n excluded	actitioner, a petent to kcluded pr enicity if a period of the e followin	or nurse practitioner prescribe acitretin; and rior to commencement of citretin is used during three years after the g criteria: ommencement of
and that they must not become pregnant during treatmen or 2 Patient is not of child bearing potential.	t and for a period of	of three yea	ars after th	e completion of treatmen
Foam spray 500 mcg with calcipotriol 50 mcg per g Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g		60 g OP 60 g OP 30 g OP	✓ <u>□</u>	instilar <u>Jaivobet</u> Jaivobe <u>t</u>
ALCIPOTRIOL Oint 50 mcg per g	40.00	120 g OP	. –	aivonex
COAL TAR		120 y UF	• 0	
Soln BP – Only in combination		200 ml	🗸 N	lidwest
 Up to 10% only in combination with a dermatologic With or without other dermatological galenicals. 	al base or proprie	tary Topica	I Corticos	teriod – Plain

/\./	Subsidy	ioo) C	Fully Brand or sidised Generic
(Mar	nufacturer's Pri \$	Per Sub	Manufacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUF			
	١		
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5%	6 50	75 g OP	
	(8.00)	75 y OF	Egopsoryl TA
	3.43	30 g OP	
	(4.35)	00 9 01	Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR	(100)		
Soln 12% with salicylic acid 2% and sulphur 4% oint	4 97	25 g OP	Coco-Scalp
	7.95	40 g OP	✓ Coco-Scalp
DIMECROLIMUS Chastiel Authority and SA1070 below. Betail abo		10 9 01	
PIMECROLIMUS – Special Authority see SA1970 below – Retail pha	inacy		
 a) Maximum of 15 g per prescription b) Note: a maximum of 15 g per prescription and no more than c 	no procorinti	ion nor 12 wa	ooko
Cream 1%		15 g OP	✓ <u>Elidel</u>
SA1970 Special Authority for Subsidy		10 9 01	
Initial application only from a dermatologist, paediatrician, ophthalmo	logict or any	rolovant pra	atitionar on the recommandation
of a dermatologist, paediatrician or ophthalmologist. Approvals valid v			
meeting the following criteria:		a renewar un	less notified for applications
Both:			
1 Patient has atopic dermatitis on the eyelid; and			
2 Patient has at least one of the following contraindications to top	vical corticos	teroids: peri	orificial dermatitis. rosacea.
documented epidermal atrophy, documented allergy to topical			
pressure.			
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEI	N – Onlv on	a prescriptio	n
* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium		500 ml	 Pinetarsol
SALICYLIC ACID			
Powder – Only in combination	18.88	250 g	✓ Midwest
 Only in combination with a dermatological base or propr 		0	oid – Plain or collodion flexible
 With or without other dermatological galenicals. 	lotary ropioc		
SULPHUR			
Precipitated – Only in combination	6 35	100 g	✓ Midwest
		0	
 Only in combination with a dermatological base or propr With or without other dermatological galenicals. 	letary Topica	a Conicoster	olu – Plain
TACROLIMUS			
Oint 0.1% – Special Authority see SA2074 below – Retail	22.00	00 ~ OD	. Tomatan
pharmacy	33.00	30 g OP	Zematop
 a) Maximum of 30 g per prescription b) Note: a maximum of 20 g per prescription and no more the 		orintion nor 1	0 wooko
b) Note: a maximum of 30 g per prescription and no more th	an one preso	copuon per 1	Z WEEKS.
► SA2074 Special Authority for Subsidy			
Initial application only from a dermatologist, paediatrician or any rele			
paediatrician, . Approvals valid without further renewal unless notified Both:	ior application	ons meeting	the following chiefta:
 Patient has atopic dermatitis on the face; and 			

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
Scalp Preparations			
BETAMETHASONE VALERATE			
* Scalp app 0.1%	9.84	100 ml OP	 Beta Scalp
CLOBETASOL PROPIONATE			
* Scalp app 0.05%	6.26	30 ml OP	✓ Dermol
HYDROCORTISONE BUTYRATE			4
Scalp lotn 0.1%	6.57	100 ml OP	 Locoid
KETOCONAZOLE	0.00	100	
Shampoo 2%	3.23	100 ml OP	 ✓ <u>Sebizole</u> ✓ Sebizole
a) Maximum of 100 ml per prescriptionb) Only on a prescription			
Sunscreens			
SUNSCREENS, PROPRIETARY – Subsidy by endorsement Only if prescribed for a patient with severe photosensitivity s endorsed accordingly. Lotn,	-	efined clinical co 200 g OP	✓ Marine Blue Lotion
Marine Blue Lotion SPF 50+ to be Principal Supply on 1	April 2023		SPF 50+
Wart Preparations			
For salicylic acid preparations refer to PSORIASIS AND ECZEM IMIQUIMOD	A PREPARATIO	ONS, page 69	
Crm 5%, 250 mg sachet	21.72	24	✓ Perrigo
PODOPHYLLOTOXIN Soln 0.5% a) Maximum of 3.5 ml per prescription b) Only on a prescription	33.60	3.5 ml OP	✓ Condyline
Other Skin Preparations			
Antineoplastics			
FLUOROURACIL SODIUM Crm 5%	6.95	20 g OP	✓ <u>Efudix</u>

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	Manufacturer

(Manufacturer's Price) Subsidised Generic \$ Per Manufacturer **Contraceptives - Non-hormonal** Condoms CONDOMS * 49 mm - Up to 144 dev available on a PSO 11.42 Moments 144 Moments 10 11.64 Moments 144 a) Maximum of 60 dev per prescription b) Up to 60 dev available on a PSO * 53 mm, 0.05 mm thickness......0.95 10 ✓ Moments ✓ Moments 144 11.42 a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription * 53 mm. chocolate. brown0.95 ✓ Moments 10 Moments 11.64 144 a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription 53 mm, strawberry, red.....0.95 ✓ Moments * 10 11.64 144 ✓ Moments a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription Moments * 56 mm......0.97 10 11.64 144 ✓ Moments a) Maximum of 60 dev per prescription b) Up to 60 dev available on a PSO * 56 mm. 0.05 mm thickness......1.30 Gold Knight 12 Gold Knight 15.57 144 a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription 144 Gold Knight a) Maximum of 60 dev per prescription b) Up to 60 dev available on a PSO 56 mm, 0.08 mm thickness......0.97 ✓ Moments 10 * ✓ Moments 11.64 144 a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription * 56 mm. 0.08 mm thickness. red0.97 ✓ Moments 10 11.64 144 Moments a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription Gold Knight 12 15.57 Gold Knight 144 a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription Gold Knight 12 15.57 144 Gold Knight a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription 12 Gold Knight XL 14.87 144 Shield XL 17.02 Gold Knight XL

▲Three おい州を対理して 命気のとないないないないでは、 if endorsed "certified exemption" by the prescriber or pharmacist. *Three おい州なら 気 保めいれるはないないないないで、

73

GENITO-URINARY SYSTEM

Brand or

Fully

Subsidy

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
 # 60 mm (bulk pack) a) Maximum of 60 dev per prescription b) Up to 60 dev available on a PSO 	14.87	144	v	Gold Knight XL
Contraceptive Devices				
INTRA-UTERINE DEVICE a) Up to 40 dev available on a PSO b) Only on a PSO # IUD 29.1 mm length × 23.2 mm width Choice TT380 Short to be Principal Supply on 1 April 2		1	v	7 MED NSHA Silver/ Copper Short Choice 380 7med Nsha Silver/ copper Short Choice TT380 Short
 * IUD 33.6 mm length × 29.9 mm width 		1	1	Choice TT380 Standard
Choice TT380 Standard to be Principal Supply on 1 Ap # IUD 35.5 mm length × 19.6 mm width Choice Load 375 to be Principal Supply on 1 April 2023		1	1	Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab – Up to

84 tab available on a PSO...... 10.00 84 ✓ Mercilon 28

S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL	Ŷ			manalaotaron
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets				
Up to 112 tab available on a PSO		84	1	Lo-Oralcon 20 ED
	2.18	04		Microgynon 20 ED
	6.45	112		Femme-Tab ED
* Tab 30 mcg with levonorgestrel 150 mcg	•••••	63	•	
	(16.50)	00		Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Aut	(/	the i	orovious n	0,
b) Up to 63 tab available on a PSO				aye
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO		84	1	Oralcon 30 ED
	1.77	04		Levien ED
	6.45	112		Femme-Tab ED
(Microgynon 20 ED Tab 20 mcg with levonorgestrel 100 mcg and				
(Femme-Tab ED Tab 20 mcg with levonorgestrel 100 mcg and 7			0	,
(Levlen ED Tab 30 mcg with levonorgestrel 150 mcg and 7 inert			0	,
(Femme-Tab ED Tab 30 mcg with levonorgestrel 150 mcg and 7				
ETHINYLOESTRADIOL WITH NORETHISTERONE			Juli)
Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to 84 tab available on a PSO		84		Brevinor 1/28
		64	v	Drevinor 1/20
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab – L		04		No vive in
to 84 tab available on a PSO		84		Norimin
	29.32	112	~	Norimin

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

(Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	
LEVONORGESTREL				
* Tab 30 mcg – Up to 84 tab available on a PSO	16.50 22.00	84 112		Microlut Microlut
Subdermal implant (2 × 75 mg rods) – Up to 3 pack available on a PSO	106.92	1	1	Jadelle
MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PS	O9.18	1	1	Depo-Provera
NORETHISTERONE Tab 350 mcg – Up to 84 tab available on a PSO	12.25	84	1	Noriday 28
Emergency Contraceptives				
LEVONORGESTREL				
* Tab 1.5 mg	1.75	1	~	Levonorgestrel BNM
	4.95		1	Postinor-1
 a) Maximum of 2 tab per prescription b) Up to 5 tab available on a PSO c) Note: Direct Provision by a pharmacist permitted under 	r the provisions in F	Part I	of Section	A.

- c) Note. Direct Provision by a pharmacist permitted under the provisions in
- d) Levonorgestrel BNM to be Principal Supply on 1 June 2023

(Postinor-1 Tab 1.5 mg to be delisted 1 June 2023)

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator		100 g OP	
	(24.15)	100 g OF	Aci-Jel
CLOTRIMAZOLE			
* Vaginal crm 1% with applicators	3.50	35 g OP	 Clomazol
Clomazol to be Principal Supply on 1 April 2023			
 Vaginal crm 2% with applicators 	3.85	20 g OP	 Clomazol
Clomazol to be Principal Supply on 1 April 2023			
MICONAZOLE NITRATE			
* Vaginal crm 2% with applicator	6.89	40 g OP	✓ Micreme
NYSTATIN			
Vaginal crm 100,000 u per 5 g with applicator(s)	4.00	75 g OP	✓ Nilstat
		-	

76

	Subsidy (Manufacturer's Pric \$		Fully Brand or dised Generic ✓ Manufacturer
Myometrial and Vaginal Hormone Preparations			
ERGOMETRINE MALEATE Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO		5	✓ DBL Ergometrine
* Crm 1 mg per g with applicator * Pessaries 500 mcg OXYTOCIN – Up to 5 inj available on a PSO		15 g OP 15	✓ <u>Ovestin</u> ✓ <u>Ovestin</u>
Inj 5 iu per ml, 1 ml ampoule Oxytocin BNM to be Principal Supply on 1 June 2023	4.98	5	 Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	5.98	5	 ✓ Oxytocin BNM ✓ Oxytocin GH \$29
Oxytocin BNM to be Principal Supply on 1 June 2023 OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj avai Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo		5	✓ <u>Syntometrine</u>
Pregnancy Tests - hCG Urine PREGNANCY TESTS - HCG URINE a) Up to 200 test available on a PSO b) Only on a PSO Cassette	12.00 16.00	40 test OP	 Smith BioMed Rapid Pregnancy Test David One Step Cassette Pregnancy Test
Urinary Agents			
For urinary tract Infections refer to INFECTIONS, Antibacterials,	page 108		
5-Alpha Reductase Inhibitors			
FINASTERIDE – Special Authority see SA0928 below – Retail p * Tab 5 mg SA0928 Special Authority for Subsidy	4.81	100	✓ <u>Ricit</u>
Initial application from any relevant practitioner. Approvals value the following criteria: Both: 1 Patient has symptomatic benign prostatic hyperplasia; and 2 Either: 2.1 The patient is intolerant of non-selective alpha bloo 2.2 Symptoms are not adequately controlled with non-	d ckers or these are	contraindicate	
Alpha-1A Adrenoreceptor Blockers			
TAMSULOSIN HYDROCHLORIDE – Special Authority see SA1 * Cap 400 mcg		ige – Retail ph 100	armacy ✓ <u>Tamsulosin-Rex</u>

	Subsidy (Manufacturer's Pr \$		Fully dised	Brand or Generic Manufacturer
 SA1032 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valithe following criteria: Both: 1 Patient has symptomatic benign prostatic hyperplasia; and 2 The patient is intolerant of non-selective alpha blockers of 	d		notifie	ed for applications meeting
Other Urinary Agents				
OXYBUTYNIN * Tab 5 mg	5.42	100	✓ J	Alchemy Oxybutynin ⁶²⁹
POTASSIUM CITRATE Oral liq 3 mmol per ml – Special Authority see SA1083 belo Retail pharmacy SA1083 Special Authority for Subsidy		200 ml OP	✓ E	Biomed
Both: 1 The patient has recurrent calcium oxalate urolithiasis; and 2 The patient has had more than two renal calculi in the two Renewal from any relevant practitioner. Approvals valid for 2 ye benefitting from the treatment. SODIUM CITRO-TARTRATE	years prior to the ars where the trea		appro	opriate and the patient is
* Grans eff 4 g sachets	2.22	28	✓ <u>l</u>	Jral
SOLIFENACIN SUCCINATE Tab 5 mg	2.05	30		Solifenacin Mylan
Tab 10 mg	3.72	30	✓ <u>s</u>	Solifenacin Viatris Solifenacin Mylan Solifenacin Viatris
Detection of Substances in Urine				
ORTHO-TOLIDINE Compound diagnostic sticks 	7.50 (8.25)	50 test OP	ŀ	Hemastix
TETRABROMOPHENOL Blue diagnostic strips 		100 test OP	~	Albustix
Obstetric Preparations				
Antiprogesterones				
MIFEPRISTONE Tab 200 mg – Up to 15 tab available on a PSO	60.00 180.00	1 3	-	Aifegyne Aifegyne

78

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
Calcium Homeostasis				
CALCITONIN * Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ M	liacalcic
CINACALCET – Special Authority see SA2170 below – Retail ph Tab 30 mg – Wastage claimable Tab 60 mg – Wastage claimable		28 28	_	inacalet Devatis inacalet Devatis
⇒SA2170 Special Authority for Subsidy				

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 All of the following:

- 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
- 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
- 1.3 The patient is symptomatic; or

2 All of the following:

- 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
- 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
- 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.
- Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
- 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	acturer's Price) Per	Subsidised	
ontinued		-		
3.2 Parathyroid tissue is surgically inaccessible; or				
3.3 Parathyroid surgery is not feasible.				
Renewal — (secondary or tertiary hyperparathyroidism) from any re	levant practi	tioner.	Approval	s valid for 12 months for
applications meeting the following criteria: Either:				
 The patient has had a kidney transplant, and following a treatmer parathyroid hormone (PTH) level to support ongoing cessation of The patient has not received a kidney transplant and trial of withd 	treatment ha	as not b	een reac	hed; or
ZOLEDRONIC ACID				
Inj 4 mg per 5 ml, vial	8.00	1	1	Zoledronic acid
				<u>Mylan</u>
			1	Zoledronic acid
				Viatris
Corticosteroids and Related Agents for Systemic Use	1			
BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE	ACETATE			
Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml		5		
	36.96)			Celestone
·	,			Chronodose
DEXAMETHASONE				
K Tab 0.5 mg – Up to 60 tab available on a PSO	.1.50	30		Dexmethsone
₭ Tab 4 mg – Up to 30 tab available on a PSO		30	-	Dexmethsone
Oral liq 1 mg per ml	18.15 2	25 ml O	P 🗸	Biomed
DEXAMETHASONE PHOSPHATE				
Dexamethasone phosphate injection will not be funded for oral use.				
k Inj 4 mg per ml, 1 ml ampoule − Up to 5 inj available on a PSO		10		Hameln
Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	13.10	10	•	Hameln
LUDROCORTISONE ACETATE				
₭ Tab 100 mcg	1.46	100	~	Florinef
IYDROCORTISONE				
₭ Tab 5 mg		100		Douglas
K Tab 20 mg		100		Douglas
Inj 100 mg vial	.4.38	1	•	Solu-Cortef
a) Up to 5 inj available on a PSOb) Only on a PSO				
/ETHYLPREDNISOLONE				
# Tab 4 mg1	12.00	100	1	Medrol
* Tab 4 mg		20	-	Medrol
C C	-0.10	20	·	Medior
IETHYLPREDNISOLONE (AS SODIUM SUCCINATE) Inj 40 mg vial	02 30	1	1	Solu-Medrol-Act-
iiij +0 iiig viai	2.00	,	•	O-Vial
				
Inj 125 mg vial	34.10	1	1	Solu-Medrol-Act- O-Vial
Inj 500 mg vial	26.88	1	~	Solu-Medrol-Act- O-Vial
loi 1 a vial	0.04	4		Solu-Medrol
Inj 1 g vial	0∠.04	1	•	Solu-mearol

 fully subsidised Principal Supply Sole Subsidised Supply

	Subsidy (Manufacturer's Pric \$	e) Per	Fully Subsidised	
METHYLPREDNISOLONE ACETATE	Ŷ	1.01		Manufacturor
Inj 40 mg per ml, 1 ml vial	47.06	5	1	Depo-Medrol
PREDNISOLONE		0	-	Dopo mouror
 * Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age. 	6.00	30 ml O	P 🗸	Redipred
PREDNISONE				
* Tab 1 mg		500		Prednisone Clinect
* Tab 2.5 mg		500		Prednisone Clinect
* Tab 5 mg – Up to 30 tab available on a PSO		500		Prednisone Clinect
* Tab 20 mg – Up to 30 tab available on a PSO	50.51	500	1	Prednisone Clinect
TETRACOSACTRIN				
 Inj 250 mcg per ml, 1 ml ampoule 	75.00	1		Synacthen
				UK Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1		Synacthen Depot Synacthene Retard S29
TRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule		5		Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule	51.10	5	/	Kenacort-A 40
Sex Hormones Non Contraceptive Androgen Agonists and Antagonists				
CYPROTERONE ACETATE				
Tab 50 mg		50		Siterone
Tab 100 mg		50	-	Siterone
TESTOSTERONE				
Patch 5 mg per day		30	1	Androderm
TESTOSTERONE CIPIONATE				
Inj 100 mg per ml, 10 ml vial		1	1	Depo-Testosterone
	393.00			Taro-
				Testosterone S29
TESTOSTERONE ESTERS				• · · · ·
Inj 250 mg per ml, 1 ml		1	-	Sustanon Ampoules
TESTOSTERONE UNDECANOATE				
Cap 40 mg – Subsidy by endorsement	21.00	60	1	Andriol Testocaps
	35.00	100		Steril-Gene S29
Subsidy by endorsement – subsidised for patients who w				
 November 2021 and the prescription is endorsed according where there exists a record of prior dispensing of testostic 				

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	(Manufacturer's Price) Su	bsidised	Generic
	\$	Per	1	Manufacturer
	Ψ	1.01	-	Manulastaron
Hormone Replacement Therapy - Systemic				
normone neplacement merapy - Systemic				
Oestrogens				
DESTRADIOL				
🗧 Tab 1 mg	4.12	28 OP		
5	(11.10)		I	Estrofem
₭ Tab 2 mg	(-)	28 OP		
1 ab 2 mg		20 01	,	Estrofem
	(11.10)			
Patch 50 mcg per 24 hours	7.04	4	v (Climara
 a) No more than 1 patch per week 				
b) Only on a prescription				
Patch 25 mcg per day	6 1 2	8	1	Estradot
r alon 20 mog por day		0		
	13.50		v	Estraderm MX S29
 a) No more than 2 patch per week 				
b) Only on a prescription				
Patch 50 mcg per day		8	 I 	Estradot 50 mcg
r alon oo mog por aay	9.22	Ũ		Estradiol TDP
	5.22		• •	
				Mylan S29
	14.50		✓	Estraderm MX S29
a) No more than 2 patch per week				
, , ,				
b) Only on a prescription	- 04	•		-
Patch 75 mcg per day		8		Estradot
	10.60		✓	Estradiol TDP
				Mylan S29
a) No more than 2 patch per week				
, , ,				
b) Only on a prescription				
Patch 100 mcg per day	7.91	8	✓ 1	Estradot
	15.50		1	Estraderm MX S29
a) No more than 2 patch per week				
, , ,				
b) Only on a prescription				
DESTRADIOL VALERATE				
Fab 1 mg	12.36	84	1	Progynova
 Fab 2 mg 		84		Progynova
-	12.00	04	• 1	iogynova
DESTROGENS				
Conjugated, equine tab 300 mcg	3.01	28		
	(17.50)		I	Premarin
 Conjugated, equine tab 625 mcg 		28		
· oonjugatou, oquino tab 020 mog		20	,	Premarin
	(17.50)		1	
Progestogens				
IEDROXYPROGESTERONE ACETATE				
 Tab 2.5 mg 	4.69	30	✓	Provera
,	8.75	56	 Image: A second s	Provera
₭ Tab 5 mg	••	56	-	Provera
• 100 0 Hig				
₭ Tab 10 mg	17.50	100 30	-	Provera
🗧 Tab 10 mg				Provera

	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully Brand or osidised Generic Manufacturer
Progestogen and Oestrogen Combined Prepara	ations		
OESTRADIOL WITH NORETHISTERONE * Tab 1 mg with 0.5 mg norethisterone acetate		28 OP	Kliavanaa
* Tab 2 mg with 1 mg norethisterone acetate	(18.10) 5.40 (18.10)	28 OP	Kliovance Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40 (18.10)	28 OP	Trisequens
Other Oestrogen Preparations			
OESTRIOL * Tab 2 mg	7.00	30	✓ <u>Ovestin</u>
Other Progestogen Preparations			
LEVONORGESTREL * Intra-uterine device 52 mg * Intra-uterine device 13.5 mg		1 1	✓ Mirena ✓ Jaydess
MEDROXYPROGESTERONE ACETATE Tab 100 mg NORETHISTERONE	116.15	100	✓ Provera HD
K Tab 5 mg − Up to 30 tab available on a PSO PROGESTERONE	5.49	30	✓ Primolut N
Cap 100 mg Utrogestan to be Principal Supply on 1 May 2023	14.85	30	 Utrogestan
Thyroid and Antithyroid Agents			
CARBIMAZOLE ₭ Tab 5 mg EVOTHYROXINE	7.56	100	✓ <u>Neo-Mercazole</u>
✤ Tab 25 mcg	5.55	90	 Synthroid
k Tab 50 mcg	1.71	28	 Mercury Pharma
	5.79	90	 Synthroid
K T-h 100 men	64.28	1,000	 Eltroxin Management Pharmace
₭ Tab 100 mcg	1.78 6.01	28 90	 Mercury Pharma Synthroid
	66.78	90 1,000	 ✓ Synthroid ✓ Eltroxin
PROPYLTHIOURACIL – Special Authority see SA1199 below –		.,	
Tab 50 mg		100	✓ PTU \$29
■SA1199 Special Authority for Subsidy		100	• FIU

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Trophic Hormones				
Growth Hormones				
SOMATROPIN (OMNITROPE) - Special Authority see SA2032 t	<mark>below – R</mark> etail pharm	асу		
* Inj 5 mg cartridge	69.75	1	-	0 <u>mnitrope</u> 0mnitrope S29 S29
* Inj 10 mg cartridge	69.75	1	✓ <u>c</u>	mnitrope
* Inj 15 mg cartridge	139.50	1	✓ <u>C</u>	Omnitrope S29 S29 Omnitrope Omnitrope S29 S29

➡SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
 - Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under ; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

86

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
- 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

		Subsidy		Fully	Brand or
		(Manufacturer's Price \$	e) Sub Per	sidised ✓	Generic Manufacturer
ntinu	ed				
ose o	f somatropin not to exceed 0.7 mg per day for male patier				
	commencement of treatment for hypopituitarism, patients in of corticosteroid and levothyroxine.	nust be monitored f	or any requ	iired adj	ustment in replacement
GnR	H Analogues				
OSEI	RELIN				
	plant 3.6 mg, syringe		1	✓ <u>⊺</u>	
	plant 10.8 mg, syringe		1	✓ <u>⊺</u>	eva
Ad	ditional subsidy by endorsement where the patient is a ch serelin and the prescription is endorsed accordingly.	ild or adolescent an	d is unable	to tolera	ate administration of
-	3.75 mg prefilled dual chamber syringe – Higher subsidy	of			
	\$221.60 per 1 inj with Endorsement		1		
		(221.60)		L	ucrin Depot 1-month
Inj	11.25 mg prefilled dual chamber syringe – Higher subsid				
	of \$591.68 per 1 inj with Endorsement		1		uarin Danat 2 month
		(591.68)		L	ucrin Depot 3-month
Vaso	ppressin Agonists				
	DPRESSIN				
Wa	afer 120 mcg	47.00	30	🗸 N	linirin Melt
ESM	OPRESSIN ACETATE				
	b 100 mcg		30		linirin
	b 200 mcg		30		linirin
⊾ Na	sal spray 10 mcg per dose	27.95	6 ml OP	✓ [<u>esmopressin-</u> <u>PH&T</u>
Inj	4 mcg per ml, 1 ml	67.18	10	🗸 N	linirin
Othe	er Endocrine Agents				
	RGOLINE				
	b 0.5 mg – Maximum of 2 tab per prescription; can be				
, in	waived by Special Authority see SA2070 below	4.43	2	✓ C	Oostinex
		17.94	8	-	ostinex
⇒SA2	070 Special Authority for Waiver of Rule				
nitial a	application from any relevant practitioner. Approvals vali	d without further rer	newal unles	s notifie	d for applications meetir
ne foll	owing criteria:				
	the following:				
	Hyperprolactinemia; or				
	Acromegaly*; or				
	Inhibition of lactation.				\ (
ractiti /hich l	al — (for patients who have previously been funded u oner. Approvals valid without further renewal unless notifi has expired and the treatment remains appropriate and the	ed where the patier	nt has previo	ously he	
	Indication marked with * is an unapproved indication.				
	FENE CITRATE	20 94	10	. A	lylan
ıa	b 50 mg	29.04	10	• N	Clomiphen S29
	fully subsidized	C20 man	od modiair -	ounelie -	under Section 00
38	✓ fully subsidised Principal Supply	Sole Subsidise		supplied	under Section 29

S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
METYRAPONE Cap 250 mg	558.00	50	✓ <u>N</u>	letopirone

	Subsidy (Manufacturer's Price) Sub	Fully sidised	Brand or Generic
	`\$	Per	1	Manufacturer
Anthelmintics				
ALBENDAZOLE - Special Authority see SA1318 below - Retail	pharmacy			
Tab 400 mg		60	✓ E	skazole S29
➡SA1318 Special Authority for Subsidy				
Initial application only from an infectious disease specialist or c patient has hydatids.	linical microbiologist	. Approva	s valid f	or 6 months where the
Renewal only from an infectious disease specialist or clinical mic remains appropriate and the patient is benefitting from the treatm		als valid fo	r 6 moni	ths where the treatment
MEBENDAZOLE – Only on a prescription				
Tab 100 mg		6	✓ <u>v</u>	/ermox
Oral liq 100 mg per 5 ml		15 ml		,
	(7.53)		V	/ermox
PRAZIQUANTEL	<u> </u>	8		Biltricide
Tab 600 mg		ð	• •	Sillricide
Antibacterials				
a) For topical antibacterials, refer to DERMATOLOGICALS, pag				
b) For anti-infective eye preparations, refer to SENSORY ORGA	NS, page 244			
Cephalosporins and Cephamycins				
CEFACLOR MONOHYDRATE				
Cap 250 mg	24.70	100	✔ F	Ranbaxy-Cefaclor S29 S29
	25.85		🗸 F	anbaxy-Cefaclor
Ranbaxy-Cefaclor to be Principal Supply on 1 April 2023		100		
Grans for oral liq 125 mg per 5 ml – Wastage claimable		100 ml	✓ H	Ranbaxy-Cefaclor
	3.75		. -	S29 ^(S29) Ranbaxy-Cefaclor
Ranbaxy-Cefaclor to be Principal Supply on 1 April 2023			• 1	andaxy-deración
(Ranbaxy-Cefaclor S29 S29) Cap 250 mg to be delisted 1 April 2				
(Ranbaxy-Cefaclor S29 S29 Grans for oral liq 125 mg per 5 ml i	,	2023)		
CEFALEXIN	r i i i i i i i i i i i i i i i i i i i	/		
Cap 250 mg		20	✓ 0	Cephalexin ABM
Cephalexin ABM to be Principal Supply on 1 April 2023				•
Cap 500 mg	5.85	20	✓ 0	Cephalexin ABM
Cephalexin ABM to be Principal Supply on 1 April 2023	7.00	100		1
Grans for oral liq 25 mg per ml – Wastage claimable Grans for oral lig 50 mg per ml – Wastage claimable		100 ml 100 ml	_	<u>lynn</u> lynn
CEFAZOLIN – Subsidy by endorsement		100 111	• [<u>17111</u>
Only if prescribed for dialysis or cellulitis in accordance with	a Health NZ Hospita	l approved	protoco	ol and the prescription is
endorsed accordingly. Inj 500 mg vial	3 30	5	~ ^	FT
Inj 1 g vial		5 5	✓ A	NFT
······································		÷	· <u>-</u>	<u></u>

90

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CEFTRIAXONE – Subsidy by endorsement				
a) Up to 10 inj available on a PSO				
b) Subsidised only if prescribed for a dialysis or cystic fibros pelvic inflammatory disease, or the treatment of suspecte endorsed accordingly.				
Inj 500 mg vial	0.79	1	✓ (Ceftriaxone-AFT
Ceftriaxone-AFT to be Principal Supply on 1 April 2023 Inj 1 g vial	3 59	5	10	Ceftriaxone-AFT
Ceftriaxone-AFT to be Principal Supply on 1 April 2023		Ũ		
CEFUROXIME AXETIL – Subsidy by endorsement Only if prescribed for prophylaxis of endocarditis and the pre	scription is endorsed	accol	rdinalv.	
Tab 250 mg (Zinnat Tab 250 mg to be delisted 1 March 2024)		50		linnat
Macrolides				

Tab 500 mg – Up to 8 tab available on a PSO	2.57	2	 Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) – Wastage			
claimable	16.97	15 ml	 Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

Tab 250 mg	8.53	14	 Klacid
Grans for oral liq 250 mg per 5 ml - Wastage claimable	2.00	50 ml	 Klacid

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Atypical mycobacterial infection; or

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and

2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE)

Inj 1 g vial		1	 Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg		100	 E-Mycin
 a) Up to 20 tab available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 200 mg per 5 ml 	5.00	100 ml	✔ E-Mycin
 a) Up to 300 ml available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP c) Wastage claimable 			
Grans for oral liq 400 mg per 5 ml a) Up to 200 ml available on a PSO b) Wastage claimable	6.77	100 ml	 E-Mycin
ROXITHROMYCIN			
Tab 150 mg	13.19	50	 Arrow- Roxithromycin
Tab 300 mg	25.00	50	✓ Arrow- Roxithromycin

92

	Subsidy	、	Fully	
	(Manufacturer's Price \$	Per	Subsidised	Generic Manufacturer
Penicillins				
AMOXICILLIN				
Cap 250 mg	43.45	500	✓	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP	CC 44	500		Alubaman
Cap 500 mg		500	•	Alphamox
 a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP 				
Grans for oral lig 125 mg per 5 ml	1.40	100 ml	1	Alphamox 125
a) Up to 200 ml available on a PSO	1.40	100 111	•	Alphaniox 125
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	1 73	100 ml	1	Alphamox 250
a) Up to 300 ml available on a PSO		100 111	-	- upitalitex 200
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Inj 250 mg vial		10	1	Ibiamox
Inj 500 mg vial		10	1	Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO	21.64	10	1	Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab				
available on a PSO	0.89	10	✓	Curam Duo 500/125
Grans for oral lig amoxicillin 25 mg with clavulanic acid 6.25 r	ng			
per ml		100 ml	1	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r	ng			
per ml – Up to 200 ml available on a PSO		00 ml C	P 🗸	Curam
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj				
available on a PSO		10	1	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS	SO 11.09	10	1	Sandoz
FLUCLOXACILLIN				
Cap 250 mg – Up to 30 cap available on a PSO	15.79	250	1	Flucloxacillin-AFT
Cap 500 mg – Up to 30 cap available on a PSO		500		Flucloxacillin-AFT
Grans for oral lig 25 mg per ml		100 ml		AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 50 mg per ml	3.68	100 ml	~	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Inj 250 mg vial		10		Flucloxin
Inj 500 mg vial		10		Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.70	5	~	Flucil

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	Generic
PHENOXYMETHYLPENICILLIN (PENICILLIN V) Cap 250 mg – Up to 30 cap available on a PSO Cap 500 mg a) Up to 20 cap available on a PSO		50 50		<u>Cilicaine VK</u> <u>Cilicaine VK</u>
 b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml a) Up to 200 ml available on a PSO 	3.40	100 m	I 🗸	<u>AFT</u>
 b) Wastage claimable Grans for oral liq 250 mg per 5 ml a) Up to 300 ml available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP c) Wastage claimable 	4.24	100 m	🗸	<u>AFT</u>
Tetracyclines				
DOXYCYCLINE * Tab 100 mg – Up to 30 tab available on a PSO MINOCYCLINE HYDROCHLORIDE	64.43	500	r	Doxine
* Tab 50 mg – Additional subsidy by Special Authority see SA1355 below – Retail pharmacy	5.79 (12.05)	60		Mino-tabs
* Cap 100 mg		100		Minomycin
► SA1355 Special Authority for Manufacturers Price Initial application from any relevant practitioner. Approvals vali rosacea. TETRACYCLINE – Special Authority see SA1332 below – Retai		ewal u	nless notif	ied where the patient has
Tab 250 mg		28	1	Accord S29
► SA1332 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vali Both: 1 For the eradication of helicobacter pylori following unsucc	d for 3 months for ap			0

1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and

2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 62

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg – Up to 5 tab available on a PSO	.2.42	28	 Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	.3.40	28	 Cipflox
Tab 750 mg	.5.95	28	 Cipflox

94

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
CLINDAMYCIN				
Cap hydrochloride 150 mg	5.30	24	✓	Dalacin C
Inj 150 mg per ml, 4 ml ampoule	35.10	10	✓	Hameln
	39.00		✓	Dalacin C
Dalacin C Inj 150 mg per ml, 4 ml ampoule to be delisted 1 Augu	ıst 2023)			
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – S	ubsidy by endorseme	ent		
Only if prescribed for dialysis or cystic fibrosis patient and the	e prescription is endor	rsed	accordingly	y.
Inj 150 mg		1		Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement		5	1	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.			t infection a	and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	1	Wockhardt S29
J - 34: , ,	182.00	10	1	Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient o endorsed accordingly.				•
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement		10	1	Pfizer
1 - 34: ,	87.50	50	1	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	t infection a	and the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Retail No patient co-payment payable	pharmacy			
Tab 400 mg		5	✓	Avelox

► SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

- Any of the following:
 - 1 Both:
 - 1.1 Active tuberculosis*; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
 - 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
 - 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:

	Subsidy (Manufacturer's Price \$) Sub Per	Fully sidised	Brand or Generic Manufacturer
continued				
2.1 Has tried and failed to clear infection using azithr 2.2 Has laboratory confirmed azithromycin resistance				
3 Treatment is only for 7 days. nitial application — (Penetrating eye injury) only from an or	abthalmologist Appr	wale valid	for 1 mc	onth where the nationt
equires prophylaxis following a penetrating eye injury only non-an-op- Note: Indications marked with * are unapproved indications.				man where the patient
PAROMOMYCIN – Special Authority see SA1689 below – Retain	ail pharmacy			
Cap 250 mg	126.00	16	✓ Н	lumatin S29
SA1689 Special Authority for Subsidy nitial application only from an infectious disease specialist, cli nonth for applications meeting the following criteria: Either:	inical microbiologist o	r gastroent	erologis	t. Approvals valid for 1
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 				
Renewal only from an infectious disease specialist, clinical mice applications meeting the following criteria: Either:	robiologist or gastroer	nterologist.	Approv	vals valid for 1 month for
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 				
PYRIMETHAMINE – Special Authority see SA1328 below – Re	etail pharmacy			
Tab 25 mg		30	🗸 D	araprim S29
 nitial application from any relevant practitioner. Approvals value following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 month 	or a period of 3 month		s notifie	d for applications meetin
SODIUM FUSIDATE [FUSIDIC ACID]				
Tab 250 mg	67.85	36	✔ F	ucidin
ULFADIAZINE SODIUM - Special Authority see SA1331 belo	w – Retail pharmacy			
Tab 500 mg Salassi Special Authority for Subsidy	543.20	56	✓ V	Vockhardt S29
itial application from any relevant practitioner. Approvals va e following criteria: ny of the following:	lid without further ren	ewal unles	s notifie	d for applications meetin
 For the treatment of toxoplasmosis in patients with HIV f For pregnant patients for the term of the pregnancy; or For infants with congenital toxoplasmosis until 12 month 		ns; or		
OBRAMYCIN Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement		5		<u>obramycin Mylan</u> /iatris
Only if prescribed for dialysis or cystic fibrosis patient a	and the prescription is	endorsed		
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by	005.00	50 de		
endorsement		56 dose	✓ Ī	obramycin BNM
a) Wastage claimableb) Only if prescribed for a cystic fibrosis patient and th	e prescription is endo	rsed accor	dingly.	

96

	Subsidy		Fully	
	(Manufacturer's Price \$) Per	Subsidised	I Generic Manufacturer
TRIMETHOPRIM	· · · · · · · · · · · · · · · · · · ·			
* Tab 300 mg – Up to 30 tab available on a PSO		50	1	TMP
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXA				
 * Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – U 				
to 30 tab available on a PSO	•	500	1	Trisul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 m				
available on a PSO		100 m	l 🗸	Deprim
VANCOMYCIN – Subsidy by endorsement				- F
Only if prescribed for a dialysis or cystic fibrosis patient or for	prophylaxis of end	ocarditi	s or for tre	eatment of Clostridium
difficile following metronidazole failure and the prescription is	endorsed according	alv.		
Ini 500 mg vial		1	1	Mylan
, .				<i></i>
Antifungals				
a) For topical antifungals refer to DERMATOLOGICALS, page 63				
b) For topical antifungals refer to GENITO URINARY, page 76				
FLUCONAZOLE				
Cap 50 mg	2.75	28		Dizole
Cap 150 mg	0.65	1		<u>Mylan</u> Mylan
Cap 150 mg		28		<u>Mylan</u> Mylan
Powder for oral suspension 10 mg per ml – Special Authority		20	•	wyian
see SA1359 below – Retail pharmacy		35 ml	1	Diflucan
Wastage claimable		00 111		Dinadan
(Dizole Cap 50 mg to be delisted 1 August 2023)				
SA1359 Special Authority for Subsidy				
Initial application — (Systemic candidiasis) from any relevant	practitioner. Appro	vals va	lid for 6 v	eeks for applications
meeting the following criteria:				
Both:				
1 Patient requires prophylaxis for, or treatment of systemic ca	andidiasis; and			
2 Patient is unable to swallow capsules.				
Initial application — (Immunocompromised) from any relevant	practitioner. Appr	ovals v	alid for 6	months for applications
meeting the following criteria:				
All of the following:				
1 Patient is immunocompromised; and				
 Patient is at moderate to high risk of invasive fungal infection Patient is unable to smaller 	on; and			
3 Patient is unable to swallow capsules.	ar Anneciale	for 0	upplys for	onnligations manthematic
Renewal — (Systemic candidiasis) from any relevant practitione	er. Approvais valio	1 10r 6 V	VEEKS TOP	applications meeting the
following criteria: Both:				
1 Patient requires prophylaxis for, or treatment of systemic ca	andidiasis: and			
2 Patient is unable to swallow capsules.	anunuasis, anu			
Renewal — (Immunocompromised) from any relevant practition	er Annrovale vali	d for 6	monthe fo	r applications meeting the
following criteria:	ion. Approvais vali			applications meeting the
All of the following:				

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pr \$	ice) Sub Per	Fully sidised	
ITRACONAZOLE	÷			
Cap 100 mg	4.27	15	1	Itrazole
Oral liq 10 mg per ml - Special Authority see SA1322 below	_			
Retail pharmacy	141.80	150 ml OP	1	Sporanox
SA1322 Special Authority for Subsidy				
nitial application only from an infectious disease specialist, clini	0	,		, ,
practitioner on the recommendation of a infectious disease physic		obiologist or o	clinical	immunologist. Approvals
valid for 6 months where the patient has a congenital immune def Renewal from any relevant practitioner. Approvals valid for 6 mo		cotmont rom	oino or	proprieto and the patient
benefitting from the treatment.		eaunentien	ains af	propriate and the patient
KETOCONAZOLE				
Tab 200 mg – PCT	CBS	30	1	Burel S29
100 200 mg 1 01		00		Link Healthcare S29
				Nizoral S29
		100		Strides Shasun S29
		100		Taro S29
(Link Healthcare see Tab 200 mg to be delisted 1 July 2023)				
(Nizoral sea) Tab 200 mg to be delisted 1 July 2023)				
NYSTATIN				
Tab 500.000 u	14.16	50		
	(17.09)			Nilstat
Cap 500,000 u	12.81	50		
	(15.47)			Nilstat
POSACONAZOLE – Special Authority see SA1285 below – Reta				
Tab modified-release 100 mg		24		Posaconazole Juno
December 1, has to be Discipled and the Horse	869.86		1	Noxafil
Posaconazole Juno to be Principal Supply on 1 April 202		105 ml OP		Devatis
Oral liq 40 mg per ml		100 III OP		Noxafil
	701.10		•	ItoAuiii

Devatis to be Principal Supply on 1 May 2023

(Noxafil Tab modified-release 100 mg to be delisted 1 April 2023)

(Noxafil Oral liq 40 mg per ml to be delisted 1 May 2023)

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

98

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Subsidy (Manufacturer's Price)	Fu Subsidis	,	
 \$	Per	 Manufacturer 	

continued...

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

* Tab 250 mg	8.15	84	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 below - Retail pl	narmacy		
Tab 50 mg	91.00	56	 Vttack
Tab 200 mg	350.00	56	 Vttack
Powder for oral suspension 40 mg per ml - Wastage			
claimable	1,523.22	70 ml	 Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4~ Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE	- Special Authority see SA1684 below - Retail pharmacy	
Tab 15 mg		

✓ Sanofi Primaguine S29

100

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
ontinued ne following criteria: loth:			
 The patient has relapsed vivax or ovale malaria; and Primaquine is to be given for a maximum of 21 days. 			
Antitrichomonal Agents			
IETRONIDAZOLE			
Tab 200 mg – Up to 30 tab available on a PSO		250	 Metrogyl
Tab 400 mg – Up to 15 tab available on a PSO		21	Metrogyl
Oral liq benzoate 200 mg per 5 ml		100 ml 10	 ✓ FlagyI-S ✓ FlagyI
Suppos 500 mg	24.40	10	• Flagyl
NRIDAZOLE Tab 500 mg		10	✓ <u>Arrow-Ornidazole</u>
Antituberculotics and Antileprotics			
lote: There is no co-payment charge for all pharmaceuticals lis	sted in the Antitubercu	lotics a	nd Antileprotics group regardless
nmigration status.			
LOFAZIMINE – Retail pharmacy-Specialist			
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendation	ation of, an infectious	disease	physician, clinical microbiologist
b) Prescriptions must be written by, or on the recommendated dermatologist.		disease 100	
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg 			physician, clinical microbiologist
 b) Prescriptions must be written by, or on the recommendate dermatologist. ♦ Cap 50 mg YCLOSERINE – Retail pharmacy-Specialist 			
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg YCLOSERINE – Retail pharmacy-Specialist a) No patient co-payment payable 	442.00	100	✓ Lamprene \$29
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg	442.00	100	✓ Lamprene \$29
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100	✓ Lamprene \$29
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg YCLOSERINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendate respiratory physician. Cap 250 mg 		100 disease	 Lamprene S29 physician, clinical microbiologist
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg YCLOSERINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendate respiratory physician. Cap 250 mg 		100 disease	 Lamprene S29 physician, clinical microbiologist
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg YCLOSERINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendate respiratory physician. Cap 250 mg APSONE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendate respiratory physician. 	442.00 ation of, an infectious of	100 disease 60	 Lamprene \$29 physician, clinical microbiologist Cyclorin \$29
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg	442.00 ation of, an infectious of 	100 disease 60	 Lamprene \$29 physician, clinical microbiologist Cyclorin \$29 physician, clinical microbiologist
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease	 Lamprene \$29 physician, clinical microbiologist Cyclorin \$29
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease 100	 Lamprene \$29 physician, clinical microbiologist Cyclorin \$29 physician, clinical microbiologist Dapsone
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease 100	 Lamprene \$29 physician, clinical microbiologist Cyclorin \$29 physician, clinical microbiologist Dapsone
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease 100 100	 Lamprene \$29 physician, clinical microbiologist Cyclorin \$29 physician, clinical microbiologist Dapsone Dapsone
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease 100 100	 Lamprene \$29 physician, clinical microbiologist Cyclorin \$29 physician, clinical microbiologist Dapsone Dapsone
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease 100 100 disease	 Lamprene S29 physician, clinical microbiologist Cyclorin S29 physician, clinical microbiologist Dapsone Dapsone physician, clinical microbiologist
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease 100 100 disease 100	 Lamprene S29 physician, clinical microbiologist Cyclorin S29 physician, clinical microbiologist Dapsone Dapsone physician, clinical microbiologist EMB Fatol S29
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease 100 100 disease 100	 Lamprene S29 physician, clinical microbiologist Cyclorin S29 physician, clinical microbiologist Dapsone Dapsone physician, clinical microbiologist EMB Fatol S29
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease 100 100 disease 100 56	 Lamprene S29 physician, clinical microbiologist Cyclorin S29 physician, clinical microbiologist Dapsone Dapsone physician, clinical microbiologist EMB Fatol S29 Myambutol S29
 b) Prescriptions must be written by, or on the recommendate dermatologist. Cap 50 mg		100 disease 60 disease 100 100 disease 100 56	 Lamprene S29 physician, clinical microbiologist Cyclorin S29 physician, clinical microbiologist Dapsone Dapsone physician, clinical microbiologist EMB Fatol S29 Myambutol S29

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per	1	Manufacturer
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation microbiologist, dermatologist or public health physician 	on of, an internal me	dicine	physician	, paediatrician, clinical
* Tab 100 mg with rifampicin 150 mg.		100	1	Rifinah
* Tab 150 mg with rifampicin 300 mg		100		Rifinah
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation respiratory physician Grans for oral lig 4 g sachet 		isease 30		t, clinical microbiologist or
PROTIONAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
 b) Prescriptions must be written by, or on the recommendation respiratory physician 	on of, an infectious d	isease	specialis	t, clinical microbiologist or
Tab 250 mg		100	~	Peteha S29
PYRAZINAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation respiratory physician				-
* Tab 500 mg	64.95	100	1	AFT-Pyrazinamide
RIFABUTIN – Retail pharmacy-Specialist				
 a) No patient co-payment payable 				
b) Prescriptions must be written by, or on the recommendation gastroenterologist				
* Cap 150 mg	353.71	30	1	Mycobutin
RIFAMPICIN – Subsidy by endorsement				
a) No patient co-payment payable				
 b) For confirmed recurrent Staphylococcus aureus infection i antimicrobial based on susceptibilities and the prescription Retail pharmacy - Specialist. Specialist must be an intern paediatrician, or public health physician. 	n is endorsed accord	ingly; (can be wa	ived by endorsement -
* Cap 150 mg		100		<u>Rifadin</u>
* Cap 300 mg		100		Rifadin
* Oral liq 100 mg per 5 ml	12.60	60 ml	1	Rifadin
Antivirals				
For eye preparations refer to Eye Preparations, Anti-Infective Pre	parations, page 244			
Hepatitis B Treatment				
ENTECAVIR				
* Tab 0.5 mg		30		Entecavir Mylan Entecavir Sandoz

LAMIVUDINE - Special Authority see SA1685 on the next page -	- Retail pharma	су	
Tab 100 mg	6.95	28	 Zetlam
Oral lig 5 mg per ml		240 ml OP	✓ Zeffix

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

(Subsidy Manufacturer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer
SA1685 Special Authority for Subsidy Initial application only from a relevant specialist or medical practit Approvals valid for 1 year where used for the treatment or prevention Renewal from any relevant practitioner. Approvals valid for 2 years TENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for the treat antiretrovirals for the purposes of Special Authority SA2139., p	on of hepatitis B. s where used for the tment of HIV is inclu	e treatmen	t or pre	evention of hepatitis B.
 Tab 245 mg (300 mg as a maleate) 		30	✓ <u>T</u>	enofovir Disoproxil <u>Mylan</u>
Herpesvirus Treatments				
ACICLOVIR				
K Tab dispersible 200 mg		25	✓ <u>L</u>	
Tab dispersible 400 mg Lovir to be Principal Supply on 1 April 2023	5.81	56	✓ L	ovir
Tab dispersible 800 mg Lovir to be Principal Supply on 1 April 2023	6.46	35	✓ L	ovir
/ALACICLOVIR				
Tab 500 mg		30	✓ V	aclovir
Tab 1,000 mg	13.76	30	✓ <u>v</u>	aclovir
ALGANCICLOVIR – Special Authority see SA1993 below – Reta				
Tab 450 mg	132.00	60	✓ <u>v</u>	<u>alganciclovir</u> Mylan

⇒SA1993 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or

2 Both:

- 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
- 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
 - 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12

Subsidy	y Fu	lly Brand or	
(Manufacturer	/	ed Generic	
\$	Per	 Manufacturer 	

continued...

months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR – [Xpharm] Note the supply of treatment is via Pharmac's approved dire website https://pharmac.govt.nz/maviret	ct distribution sup	ply. Further c	details can be found on Pharmac's
Tab 100 mg with pibrentasvir 40 mg	24,750.00	84 OP	 Maviret
LEDIPASVIR WITH SOFOSBUVIR – [Xpharm] – Special Autho No patient co-payment payable		elow	
Tab 90 mg with sofosbuvir 400 mg	24,363.46	28	 Harvoni
▶ SA1605 Special Authority for Subsidy Special Authority approved by the Hepatitis C Treatment Panel Notes: By application to the Hepatitis C Treatment Panel (HepC Applications will be considered by HepCTP and approved subje Application details may be obtained from Pharmac's website <u>http</u> The Coordinator, Hepatitis C Treatment Panel	CTP). ct to confirmation of	0,	<u>et</u> or:
Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,			

Email: hepcpanel@pharmac.govt.nz

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

30

 Tenofovir Disoproxil Emtricitabine Mvlan Tenofovir Disoproxil **Emtricitabine Viatr**

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement: can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV. prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 105 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

* Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a

-CA0120	Special Authority	for Subsidy
	Special Authority	V IOI SUDSIUV

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion: and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.
- Notes: Refer to local health pathways or the Australasian Society for HIV. Viral Hepatitis and Sexual Health Medicine clinical auidelines:

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion: and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on Pharmac's website) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability. 40 Lagevrio

Cap 200 mg......0.00

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer	
NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by er	ndorsement				

- a) No patient co-payment payable

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA2139 on the previous page - Retail phan	macy	
Tab 200 mg	90	 Stocrin
Tab 600 mg63.38	30	 Stocrin
ETRAVIRINE - Special Authority see SA2139 on the previous page - Retail pha	irmacy	
Tab 200 mg770.00	60	 Intelence
NEVIRAPINE - Special Authority see SA2139 on the previous page - Retail pha	armacy	
Tab 200 mg	60	 Nevirapine
		<u>Alphapharm</u>
		 Nevirapine Viatris
Oral suspension 10 mg per ml203.55	240 ml OP	 Viramune
		Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA2139 on the Tab 300 mg Oral liq 20 mg per ml		Retail pharmad 60 240 ml OP	∠y ✓ Ziagen ✓ Ziagen
NBACAVIR SULPHATE WITH LAMIVUDINE – Special Authori Note: abacavir with lamivudine (combination tablets) count anti-retroviral Special Authority.	,		v 1 7
Tab 600 mg with lamivudine 300 mg	29.50	30	✓ Abacavir/ Lamivudine Viatris
	75.00		 Kivexa
Abacavir/Lamivudine Viatris to be Principal Supply on	May 2023		

(Kivexa Tab 600 mg with lamivudine 300 mg to be delisted 1 May 2023)

	Subsidy		Fully Brand or	
	(Manufacturer's P			
	\$	Per	 Manufactur 	· · · · · · · · · · · · · · · · · · ·
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPF	IOXIL – Special	Authority see S	A2139 on page 10	05 – Retail
pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil co anti-retroviral Special Authority	unts as three ar	nti-retroviral mec	lications for the pu	irposes of the
Tab 600 mg with emtricitabine 200 mg and tenofovir disoprox	il			
245 mg (300 mg as a maleate)		30	✓ Mylan✓ Viatris	
EMTRICITABINE – Special Authority see SA2139 on page 105 – Cap 200 mg		зу 30	✓ Emtriva	
LAMIVUDINE - Special Authority see SA2139 on page 105 - Re				
Tab 150 mg		60	✓ <u>Lamivudine</u> <u>Alphaphar</u>	
Oral liq 10 mg per ml	102.50	240 ml OP	 Lamivudine 3TC 	Viatris
ZIDOVUDINE [AZT] - Special Authority see SA2139 on page 10	5 – Retail pharm	nacy		
Cap 100 mg Oral liq 10 mg per ml		100 200 ml OP	RetrovirRetrovir	
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets)				ourposes of
the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg		60	✓ Alphapharm	
Protease Inhibitors				
ATAZANAVIR SULPHATE - Special Authority see SA2139 on pa	age 105 – Retail	pharmacy		
Cap 150 mg		60	 ✓ Atazanavir M ✓ Teva 	lylan
Atazanavir Mylan to be Principal Supply on 1 May 2023				
Cap 200 mg		60	✓ Atazanavir N	lylan
Atazanavir Mylan to be Principal Supply on 1 May 2023	188.91		 Teva 	
(Teva Cap 150 mg to be delisted 1 May 2023) (Teva Cap 200 mg to be delisted 1 May 2023)				
DARUNAVIR - Special Authority see SA2139 on page 105 - Rei	ail pharmacy			
Tab 400 mg		60	 Darunavir M 	ylan
Tab 600 mg	196.65	60	✓ <u>Darunavir M</u>	
(Darunavir Mylan Tab 600 mg to be delisted 1 August 2023)			✓ Darunavir Vi	latris
LOPINAVIR WITH RITONAVIR – Special Authority see SA2139	on nago 105 _ F	Rotail nharmaou		
Tab 100 mg with ritonavir 25 mg		60	✓ Lopinavir/Ri	tonavir
Tab 200 mg with ritonavir 50 mg	295.00	120	<u>Mylan</u> ✓ <u>Lopinavir/Ri</u>	tonavir
Oral liq 80 mg with ritonavir 20 mg per ml	735.00	300 ml OP	<u>Mylan</u> ✓ Kaletra	
RITONAVIR – Special Authority see SA2139 on page 105 – Reta			- Nulstia	
Tab 100 mg		30	 Norvir 	
Strand Transfer Inhibitors				
DOLUTEGRAVIR – Special Authority see SA2139 on page 105 - Tab 50 mg		cy 30	🗸 Tivicay	

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		ully	Brand or	
	(Manufacturer's Price)	Subsidis		Generic	
	\$	Per	1	Manufacturer	
RALTEGRAVIR POTASSIUM – Special Authority see SA2139 on page 105 – Retail pharmacy					
Tab 400 mg	1,090.00	60	🗸 İs	entress	
Tab 600 mg	1,090.00	60	🗸 İs	entress HD	
Tab 600 mg	1,090.00	60	✓ Is	entress HD	

Immune Modulators

PEGYLATED INTERFERON ALFA-2A – Special Authority see SA2034 below – Retail pharmacy

► SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and

3.2.2 Either:

- 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
- 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

INFECTIONS - AGENTS FOR SYSTEMIC USE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
NITROFURANTOIN				
* Tab 50 mg – Up to 30 tab available on a PSO		100	A M	<u>Nifuran</u>
* Tab 100 mg		100	A M A	<u>Nifuran</u>
* Cap modified-release 100 mg – Up to 15 cap available on a PSO		100	✓ <u>N</u>	<i>Macrobid</i>
NORFLOXACIN Tab 400 mg – Subsidy by endorsement		100	√	Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

MUSCULOSKELETAL SYSTEM

	Subsidy		Fully Brand or
	(Manufacturer's Price)		ubsidised Generic
	\$	Per	 Manufacturer
Anticholinesterases			
NEOSTIGMINE METILSULFATE			
Inj 2.5 mg per ml, 1 ml ampoule		10	Max Health
PYRIDOSTIGMINE BROMIDE			
▲ Tab 60 mg	50.28	100	 Mestinon
		100	· mootmon
Non-Steroidal Anti-Inflammatory Drugs			
· · ·			
	1.00	50	
* Tab EC 25 mg		50 20	 ✓ <u>Diclofenac Sandoz</u> ✓ Voltaren D
 * Tab 50 mg dispersible * Tab EC 50 mg 		20 50	 Vonaren D ✓ Diclofenac Sandoz
 Tab EC 50 mg Tab long-acting 75 mg 		100	✓ Voltaren SR
 Tab long-acting 75 mg. Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a l 		5	✓ Voltaren
* Suppos 12.5 mg		10	✓ Voltaren
* Suppos 25 mg		10	✓ Voltaren
 Suppos 50 mg – Up to 10 supp available on a PSO 		10	✓ Voltaren
* Suppos 100 mg		10	 Voltaren
BUPROFEN			
* Tab 200 mg	21 40	1,000	 Relieve
* Tab long-acting 800 mg		30	✓ Brufen SR
* Oral liq 20 mg per ml		200 ml	✓ Ethics
	11.29		 Fenpaed 100 mg per
			5 ml
KETOPROFEN			
* Cap long-acting 200 mg		28	✓ Oruvail SR
MEFENAMIC ACID			
* Cap 250 mg	1 25	50	
* Oap 200 mg	(10.82)	50	Ponstan
	0.50	20	ronotan
	(7.50)		Ponstan
NAPROXEN	、		
* Tab 250 mg	32.69	500	 Noflam 250
* Tab 500 mg		250	✓ Noflam 500
* Tab long-acting 750 mg		28	✓ Naprosyn SR 750
* Tab long-acting 1 g		28	 Naprosyn SR 1000
TENOXICAM			
* Tab 20 mg		100	 Tilcotil
* Inj 20 mg vial		1	✓ AFT
NSAIDs Other			
CELECOXIB			
Cap 100 mg	3.45	60	 Celebrex
		00	 Celecoxib Pfizer
Cap 200 mg		30	✓ Celebrex
- · · · · · · · · · · · · · · · · · · ·			 Celecoxib Pfizer
			<u></u>

	(Manufacturer's Pri \$	ce) Sub Per	osidised Generic Manufacturer
Topical Products for Joint and Muscular Pain			
CAPSAICIN			
Crm 0.025% – Special Authority see SA1289 below – Retail pharmacy	9.75 13.00	45 g OP 60 g OP	 ✓ <u>Zostrix</u> ✓ Rugby Capsaicin Topical Cream ^{S29}
SA1289 Special Authority for Subsidy itial application from any relevant practitioner. Approvals valid steoarthritis that is not responsive to paracetamol and oral non-si	without further re eroidal anti-inflar	enewal unles mmatories a	ss notified where the patient has re contraindicated.
Antirheumatoid Agents			
YDROXYCHLOROQUINE – Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systemic suppression, relevant dermatological conditions (cutaneous for mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmor Pharmacists may annotate the prescription as endorsed when hydroxychloroquine. Note: Indication marked with a * is an u	rms of lupus and ary)*, and the pr e there exists a r napproved indica	l lichen plan escription is ecord of pric ation.	us, cutaneous vasculitides and endorsed accordingly. or dispensing of
÷ Tab 200 mg EFLUNOMIDE	8.78	100	Plaquenil
Tab 10 mg Tab 20 mg		30 30	✓ <u>Arava</u> ✓ Arava
ENICILLAMINE Tab 125 mg Tab 250 mg		100 100	✓ D-Penamine✓ D-Penamine
Drugs Affecting Bone Metabolism			
Alendronate for Osteoporosis			
LENDRONATE SODIUM Tab 70 mg LENDRONATE SODIUM WITH COLECALCIFEROL	2.44	4	✓ Fosamax
Tab 70 mg with colecalciferol 5,600 iu	1.51	4	 Fosamax Plus
Other Treatments			
ENOSUMAB – Special Authority see SA1777 below – Retail pha Inj 60 mg prefilled syringe SA1777 Special Authority for Subsidy itial application from any relevant practitioner. Approvals valid e following criteria:	326.00	1 enewal unles	Prolia ss notified for applications meeting
 of the following: The patient has severe, established osteoporosis; and Fither: 			

- 2 Either:
 - 2.1 The patient is female and postmenopausal; or

MUSCULOSKELETAL SYSTEM

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

- 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	Pamisol
Inj 6 mg per ml, 10 ml vial		1	 Pamisol
Inj 9 mg per ml, 10 ml vial	94.34	1	 Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA1779	below – Retail ph	armacy	
* Tab 60 mg	53.76	28	 Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

continued...

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

4	 Risedronate Sandoz
1	 Forteo
	4

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

	Subsidy (Manufacturer's F \$	Price) Sub Per	Fully sidised	Brand or Generic Manufacturer
ntinued				
 c) A vertebral fracture is defined as a 20% or greater reducer relative to the posterior height of that body, or a 20% or body above or below the affected vertebral body. d) A maximum of 18 months of treatment (18 cartridges) w 	greater reduction			
DLEDRONIC ACID Inj 0.05 mg per ml, 100 ml, bag	22.53	100 ml OP	✓ Z	oledronic Acid Viatris
			✓ Z	oledronic-US S29
Zoledronic Acid Viatris to be Principal Supply on 1 Jun Inj 0.05 mg per ml, 100 ml, vial clasta Inj 0.05 mg per ml, 100 ml, vial to be delisted 1 June 2	60.00	100 ml OP	✓ A	clasta
Hyperuricaemia and Antigout				
LOPURINOL				
Tab 100 mg		500	✓ D	P-Allopurinol
Tab 300 mg		500	✓ □	P-Allopurinol
ENZBROMARONE - Special Authority see SA1963 below -	Retail pharmacy			
Tab 50 mg	22.50	100	🗸 N	larcaricin mite S29
		30	🗸 D	esuric S29
Tab 100 mg			1 1	Irinorm S29
Tab 100 mg	45.00	100	-	Senzbromaron AL

MUSCULOSKELETAL SYSTEM

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

00100		
* Tab 500 mcg6.00	100	✓ Colgout
FEBUXOSTAT – Special Authority see SA2054 below – Retail pharmacy		
Tab 80 mg	28	 Febuxostat
		multichem
Tab 120 mg20.00	28	 Febuxostat
		multichem

➡SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required

	Subsidy (Manufacturer's Price) \$	Sub: Per	Fully sidised	Brand or Generic Manufacturer
continued				
and serum urate remains greater than 0.36 mmol maximum tolerated dose; or				
2.3 The patient has renal impairment such that probe remains greater than 0.36 mmol/l despite optimal	treatment with allopur	rinol (see l	Note); o	r
2.4 The patient has previously had an initial Special A nitial application — (Tumour lysis syndrome) only from a ha				
applications meeting the following criteria: Both:		ууы. Арр	rovais v	and for 6 weeks for
 Patient is scheduled to receive cancer therapy carrying a Patient has a documented history of allopurinol intolerand 		n risk of tur	mour lys	sis syndrome; and
Renewal — (Gout) from any relevant practitioner. Approvals v batient is benefitting from treatment.	alid for 2 years where	the treatm	nent ren	nains appropriate and the
Renewal — (Tumour lysis syndrome) only from a haematolog reatment remains appropriate and the patient is benefitting from		orovals va	lid for 6	weeks where the
PROBENECID 券 Tab 500 mg		100	✓ P	Probenecid-AFT
Muscle Relaxants				
3ACLOFEN * Tab 10 mg	4.20	100	✓ P	acifen
Inj 0.05 mg per ml, 1 ml ampoule – Subsidy by endorseme		1	-	ioresal Intrathecal
Subsidised only for use in a programmable pump in pat caused intolerable side effects and the prescription is e		pastic age	nts hav	e been ineffective or have
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement		5		ledsurge
Subsidised only for use in a programmable pump in pat caused intolerable side effects and the prescription is e		pastic age	nts hav	e been ineffective or have
DANTROLENE			_	
Cap 25 mg	112.13	100	-	antrium
Cap 50 mg	77.00	100		Dantrium S29 S29 Dantrium
		100	• 1	
ORPHENADRINE CITRATE Tab 100 mg	20.76	100	🖌 N	lorflex
		100	• 1	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Agents for Parkinsonism and Related Disorder	'S			
Dopamine Agonists and Related Agents				
AMANTADINE HYDROCHLORIDE				
▲ Cap 100 mg		60	✓	Symmetrel
	63.73	100	✓	Symmetrel
APOMORPHINE HYDROCHLORIDE				
Inj 10 mg per ml, 2 ml ampoule		5	1	Movapo
Inj 10 mg per ml, 5 ml ampoule	121.84	5	1	<u>Movapo</u>
ENTACAPONE				
▲ Tab 200 mg		100	1	Comtan
LEVODOPA WITH BENSERAZIDE				
* Tab dispersible 50 mg with benserazide 12.5 mg		100	1	Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	-	Madopar 62.5
* Cap 100 mg with benserazide 25 mg		100	1	Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100	✓	Madopar HBS
* Cap 200 mg with benserazide 50 mg		100	1	Madopar 250
EVODOPA WITH CARBIDOPA				
* Tab 100 mg with carbidopa 25 mg		100	1	Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg		100	1	Sinemet CR
* Tab 250 mg with carbidopa 25 mg		100	✓	Sinemet
PRAMIPEXOLE HYDROCHLORIDE				
▲ Tab 0.25 mg	5.51	100	1	Ramipex
▲ Tab 1 mg		100		Ramipex
RASAGILINE				
* Tab 1 mg	53 50	30	1	Azilect S29
ROPINIROLE HYDROCHLORIDE		00	-	
▲ Tab 0.25 mg	4.05	84	1	Ropin
Tab 1 mg		84		Ropin
Tab 2 mg		84	-	Ropin
▲ Tab 5 mg		84	-	Ropin
SELEGILINE HYDROCHLORIDE – Subsidy by endorsement		•		
Subsidy by endorsement – Subsidised for patients who wer prescription is endorsed accordingly. Pharmacists may anr prior dispensing of selegiline hydrochloride.	notate the prescription	as er 100	ndorsed wh	ere there exists a record of
▲ Tab 100 mg	152.38	100	1	Tasmar
Anticholinergics				
BENZATROPINE MESYLATE				
Tab 2 mg		60		Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	1	Phebra
a) Up to 10 inj available on a PSO				
b) Only on a PSO				
PROCYCLIDINE HYDROCHLORIDE				
Tab 5 mg	7.40	100	1	Kemadrin

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Agents for Essential Tremor, Chorea and Relat	ed Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail phar Wastage claimable	macy			
Tab 50 mg		56	✓ <u>R</u>	ilutek
 SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory speciali following criteria: All of the following: The patient has amyotrophic lateral sclerosis with diseas: The patient has at least 60 percent of predicted forced viti 	e duration of 5 years c	or less; a	nd	-
 The patient has an easi to percent of predicted forced with 3 The patient has not undergone a tracheostomy; and The patient has not experienced respiratory failure; and Any of the following: The patient is ambulatory; or The patient is able to use upper limbs; or The patient is able to swallow. 	ai capacity within 2 m	onins pri	or to the	initian application, and
 Renewal from any relevant practitioner. Approvals valid for 18 r All of the following: The patient has not undergone a tracheostomy; and The patient has not experienced respiratory failure; and Any of the following: The patient is ambulatory; or The patient is able to use upper limbs; or The patient is able to swallow. 	nonths for applications	s meetin	g the follo	wing criteria:
TETRABENAZINE Tab 25 mg Motetis to be Principal Supply on 1 April 2023	106.59	112	✓ M	lotetis
Anaesthetics Local				
 LIDOCAINE [LIGNOCAINE] Gel 2%, tube - Subsidy by endorsement a) Up to 150 ml available on a PSO b) Subsidised only if prescribed for urethral or cervical Gel 2%, 11 ml urethral syringe - Subsidy by endorsement a) Up to 5 each available on a PSO b) Subsidised only if prescribed for urethral, cervical or accordingly. 	administration and the	10	otion is er ✔ <u>Ir</u>	stillagel Lido

	Subsidy		Fully Brand or
	(Manufacturer's Pr	rice) Subs	sidised Generic
	\$	Per	 Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Oral (gel) soln 2%		200 ml	 Mucosoothe
Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO		25	✓ Lidocaine-Baxter
,	17.50	50	
	(35.00)		Xylocaine
Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO	(/	25	 Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO		5	
,,	(20.00)		Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO	()	5	 Lidocaine-Claris
j ,	6.85		 Lidocaine-Baxter
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO		5	 Lidocaine-Baxter
(Lidocaine-Claris Inj 1%, 20 ml vial to be delisted 1 June 2023)		Ū.	
, , , , , , , , , , , , , , , , , , , ,			
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE			
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes -			4 - 11
Subsidy by endorsement		10	 Pfizer
 a) Up to 5 each available on a PSO 			
b) Subsidised only if prescribed for urethral or cervical	administration and	the prescripti	on is endorsed accordingly.
Topical Local Anaesthetics			
benefiting from treatment. LIDOCAINE [LIGNOCAINE] – Special Authority see SA0906 ab Crm 4%	5.40 27.00	5 g OP 30 g OP	✓ LMX4 ✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Auth	,		
Crm 2.5% with prilocaine 2.5%		30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓ EMLA
Analgesics			
For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, p	age 111		
Non-opioid Analgesics			
ASPIRIN			
	4.50	100	Cethian Acrisia
* Tab dispersible 300 mg – Up to 30 tab available on a PSO.	4.50	100	✓ Ethics Aspirin
 Tab dispersible 300 mg - Up to 30 tab available on a PSO. CAPSAICIN - Subsidy by endorsement 			·
* Tab dispersible 300 mg – Up to 30 tab available on a PSO.			·
 Tab dispersible 300 mg - Up to 30 tab available on a PSO. CAPSAICIN - Subsidy by endorsement 			·
Tab dispersible 300 mg – Up to 30 tab available on a PSO. CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or of	liabetic peripheral		·
Tab dispersible 300 mg – Up to 30 tab available on a PSO. CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or o accordingly.	liabetic peripheral	neuropathy a	nd the prescription is endorsed
Tab dispersible 300 mg – Up to 30 tab available on a PSO. CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or o accordingly.	liabetic peripheral	neuropathy a 45 g OP	nd the prescription is endorsed ✓ <u>Zostrix HP</u> ✓ Rugby Capsaicin Topical
* Tab dispersible 300 mg – Up to 30 tab available on a PSO. CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or of accordingly. Crm 0.075% NEFOPAM HYDROCHLORIDE	liabetic peripheral 11.95 15.14	neuropathy a 45 g OP	nd the prescription is endorsed ✓ <u>Zostrix HP</u> ✓ Rugby Capsaicin Topical Cream 529
Tab dispersible 300 mg – Up to 30 tab available on a PSO. CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or or accordingly. Crm 0.075%	liabetic peripheral 11.95 15.14	neuropathy a 45 g OP 57 g OP	nd the prescription is endorsed ✓ <u>Zostrix HP</u> ✓ Rugby Capsaicin Topical

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

nent ription is story supp nts. If qu sings not atients wi nnotated ong-term If quant ot exceed a 2 5 1, 0 20 nent	Per 1,000 as with long annotate ports a lor uantities p t exceedir 1,000 ith long te according n condition tities pres- ding 100 t 200 ml 000 ml OP perturbation of the condition of the port of the	g term d acco ng-tern rescril ng 100 v rm co logly. Pl cribed rab per ab per v v v	 Manufacturer Pacimol r conditions who require ordingly. Pharmacists m m condition. ibed for more than 100 ta 0 tab per dispensing. Y Noumed Paracetamol orditions who require reg Pharmacists may annotate d for more than 100 tabs er dispensing. Y Paracetamol (Ethics) Y Paracare Avallon Avallon
nent ription is story supp nts. If qu sings not atients wi notated ong-term If quant ot exceed 3 2 5 1, 0 20 nent ats. If qua not excee or patients	is with long annotate ports a lor uantities p t exceedir 1,000 ith long te according o condition tities press ding 100 t 200 ml 00 ml OP	g term d acco ng-tern rescril ng 100 v rm co logly. Pl cribed rab per ab per v v v	n conditions who require ordingly. Pharmacists m m condition. ibed for more than 100 ta 0 tab per dispensing. 7 <u>Noumed Paracetamol</u> onditions who require reg Pharmacists may annotate d for more than 100 tabs er dispensing. 7 Paracetamol (Ethics) 7 Paracare 7 Avallon bed exceed 200 ml (for per dispensing.
nent ription is story supp nts. If qu sings not atients wi notated ong-term If quant ot exceed 3 2 5 1, 0 20 nent ats. If qua not excee or patients	is with long annotate ports a lor uantities p t exceedir 1,000 ith long te according o condition tities press ding 100 t 200 ml 00 ml OP	g term d acco ng-tern rescril ng 100 v rm co logly. Pl cribed rab per ab per v v v	n conditions who require ordingly. Pharmacists m m condition. ibed for more than 100 ta 0 tab per dispensing. 7 <u>Noumed Paracetamol</u> onditions who require reg Pharmacists may annotate d for more than 100 tabs er dispensing. 7 Paracetamol (Ethics) 7 Paracare 7 Avallon bed exceed 200 ml (for per dispensing.
er patients ription is story supp nts. If qu sings not atients wi notated ong-term If quant ot exceece atients atients. If qua not excee or patients	annotate ports a lor uantities p t exceedir 1,000 ith long te according o condition tities pres ding 100 t 200 ml ,000 ml 00 ml OP lantities pr eeding 200	d accomp-tern rescril and 100 arm coordination and period and	ordingly. Pharmacists m m condition. ibed for more than 100 ta 0 tab per dispensing. 7 <u>Noumed Paracetamol</u> onditions who require reg Pharmacists may annotate d for more than 100 tabs or dispensing. 7 Paracetamol (Ethics) 7 Paracare 7 Avallon bed exceed 200 ml (for per dispensing.
ription is story supp nts. If qu sings not atients wi notated org-term If quant ot exceed 3 2 5 1, 0 20 nent ats. If quant not exceed or patients	annotate ports a lor uantities p t exceedir 1,000 ith long te according o condition tities pres ding 100 t 200 ml ,000 ml 00 ml OP lantities pr eeding 200	d accomp-tern rescril and 100 arm coordination and period and	ordingly. Pharmacists m m condition. ibed for more than 100 ta 0 tab per dispensing. 7 <u>Noumed Paracetamol</u> onditions who require reg Pharmacists may annotate d for more than 100 tabs or dispensing. 7 Paracetamol (Ethics) 7 Paracare 7 Avallon bed exceed 200 ml (for per dispensing.
atients wi notated ong-term If quant ot exceed 3 2 5 1, 0 20 nent ats. If qua not exceed or patients	ith long te according n condition tities prese ding 100 t 200 ml ,000 ml 00 ml OP nantities pr eeding 200	erm co gly. Pl 1. cribed ab per 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	Paracetamol proditions who require reg pharmacists may annotate of or more than 100 tabs or dispensing. Paracetamol (Ethics) Paracare Avallon bed exceed 200 ml (for per dispensing.
nnotated a ong-term If quant ot exceed 3 2 5 1, 0 20 nent ats. If qua not exceed or patients	according o condition tities prese ding 100 t 200 ml ,000 ml 00 ml OP eantities pr eeding 200	gly. Pl n. cribed ab per v v v v rescrib 0 ml p	Pharmacists may annotate d for more than 100 tabs er dispensing. ^ Paracetamol (Ethics) ^ Paracare ^ Avallon bed exceed 200 ml (for ber dispensing.
5 1,) 20 nent nts. If qua not exce or patients	,000 ml 00 ml OP antities pr eeding 200	rescrib 0 ml p	(Ethics) Yearacare Yavallon bed exceed 200 ml (for ber dispensing.
) 20 nent nts. If qua not exce or patients	0 ml OP antities pr eeding 20	rescrib 0 ml p	Availon bed exceed 200 ml (for ber dispensing.
nts. If qua not exce	eeding 20	0 ml p	per dispensing.
not exce pr patients	eeding 20	0 ml p	per dispensing.
	endorsed	or an	n conditions who require motated accordingly. supports a long-term
			Pamol Paracare Double Strength
nent			u u u u
not exce or patients ription is o	eeding 20 s with lon endorsed	0 ml p g term or an	per dispensing. n conditions who require notated accordingly.
)	10		Gacet
r g	ment ents. If qu g not exc ior patient cription is	1,000 ml ment ints. If quantities pi g not exceeding 20 for patients with lon cription is endorsed	5 1,000 ml 🗸

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	I Generic Manufacturer
 Suppos 500 mg 		50	1	Gacet
Paracare Oral lig 120 mg per 5 ml to be delisted 1 June 2023				
Paracare Double Strength Oral liq 250 mg per 5 ml to be deli	sted 1 April 2023)			
Opioid Analgesics				
CODEINE PHOSPHATE – Safety medicine; prescriber may c				N
Tab 15 mg	5.92 6.25	100		Noumed PSM
Noumed to be Principal Supply on 1 May 2023	0.20			r JWI
Tab 30 mg	6.98	100	1	Aspen
				Noumed
	7.45			PSM
Noumed to be Principal Supply on 1 April 2023				
Tab 60 mg	13.89	100	1	Noumed
	14.25		1	PSM
Noumed to be Principal Supply on 1 April 2023				
PSM Tab 15 mg to be delisted 1 May 2023)				
PSM Tab 30 mg to be delisted 1 April 2023)				
PSM Tab 60 mg to be delisted 1 April 2023)				
DIHYDROCODEINE TARTRATE				
Tab long-acting 60 mg	8.60	60	v	DHC Continus
ENTANYL				
a) Only on a controlled drug form				
 b) No patient co-payment payable c) Sofate modulation group data mission disconsistent data mission data mission disconsistent data mission disconsistent data mission disconsistent data mission disconsistent data mission data miss	. .			
c) Safety medicine; prescriber may determine dispensing Inj 50 mcg per ml, 2 ml ampoule		10		Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10		Boucher and Muir
Patch 12.5 mcg per hour		5		Fentanyl Sandoz
Patch 25 mcg per hour		5		Fentanyl Sandoz
Patch 50 mcg per hour		5		Fentanyl Sandoz
Patch 75 mcg per hour		5		Fentanyl Sandoz
Patch 100 mcg per hour		5	1	Fentanyl Sandoz
IETHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing	frequency			
d) Extemporaneously compounded methadone will only b		te of th	le cheape	st form available
(methadone powder, not methadone tablets).				
e) For methadone hydrochloride oral liquid refer Standard				
Tab 5 mg		10		Methadone BNM
Oral liq 2 mg per ml		200 m		Biodone Biodone Forte
Oral liq 5 mg per ml	••••	200 m		Biodone Forte
Oral liq 10 mg per ml		200 m 10		Biodone Extra Forte
Inj 10 mg per ml, 1 ml		10	v	AFI

	Subsidy		Fully Brand or
(Manufacturer's Pri		sidised Generic
	\$	Per	 Manufacturer
IORPHINE HYDROCHLORIDE			
 a) Only on a controlled drug form 			
 b) No patient co-payment payable 			
c) Safety medicine; prescriber may determine dispensing freq	uency		
Oral liq 1 mg per ml	11.98	200 ml	 RA-Morph
Oral liq 2 mg per ml		200 ml	 RA-Morph
Oral liq 5 mg per ml		200 ml	 Ordine S29
			RA-Morph
Oral liq 10 mg per ml		200 ml	✓ Ordine S29
		200	✓ RA-Morph
IORPHINE SULPHATE			· F
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing freq	uency		
Tab immediate-release 10 mg		10	Sevredol
Tab immediate-release 20 mg		10	✓ Sevredol
Cap long-acting 10 mg		10	✓ m-Eslon
m-Eslon to be Principal Supply on 1 April 2023		10	
Cap long-acting 30 mg	4.30	10	✓ m-Eslon
m-Eslon to be Principal Supply on 1 April 2023	4.00	10	
Cap long-acting 60 mg	9.00	10	✓ m-Eslon
m-Eslon to be Principal Supply on 1 April 2023		10	
Cap long-acting 100 mg	10 50	10	✓ m-Eslon
m-Eslon to be Principal Supply on 1 April 2023		10	
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSC	5 38	5	 Medsurge
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5	✓ Medsurge
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5	✓ Medsurge
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5	✓ Medsurge
XYCODONE HYDROCHLORIDE	0	Ŭ	<u></u>
a) Only on a controlled drug form			
 b) No patient co-payment payable c) Softward distance of the payable 			
 c) Safety medicine; prescriber may determine dispensing freq Tab controlled-release 5 mg 		20	 Oxycodone Sandoz
Tab controlled-release 5 mg		20	✓ Oxycodone Sandoz
Tab controlled-release 20 mg		20	 Oxycodone Sandoz Oxycodone Sandoz
5		20 20	 Oxycodone Sandoz Oxycodone Sandoz
Tab controlled-release 40 mg		20 20	
Tab controlled-release 80 mg Cap immediate-release 5 mg		20 20	 ✓ <u>Oxycodone Sandoz</u> ✓ <u>OxyNorm</u>
· · ·		20 20	✓ <u>OxyNorm</u> ✓ OxyNorm
Cap immediate-release 10 mg Cap immediate-release 20 mg		20 20	✓ OxyNorm ✓ OxyNorm
Oral lig 5 mg per 5 ml		20 250 ml	✓ OxyNorm ✓ OxyNorm
		250 mi 5	✓ <u>OxyNorm</u> ✓ Hameln
Inj 10 mg per ml, 1 ml ampoule		5 5	✓ Hamein
Inj 10 mg per ml, 2 ml ampoule Inj 50 mg per ml, 1 ml ampoule		э 5	✓ <u>Hameln</u>
		°,	
PARACETAMOL WITH CODEINE – Safety medicine; prescriber m			
K Tab paracetamol 500 mg with codeine phosphate 8 mg	27.50	1,000	Paracetamol +
			Codeine (Relieve)

	0 1 1			
	Subsidy		Fully Subsidised	
	(Manufacturer's Price) \$	Per		Manufacturer
	Ψ	1 61		Manulaciulei
ETHIDINE HYDROCHLORIDE				
 a) Only on a controlled drug form 				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fro	equency			
Tab 50 mg		10	✓	PSM
.	8.68		1	Noumed Pethidine
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	PSO 29.88	5	1	DBL Pethidine
Jee Ster (a feet of the Jack and the				Hydrochloride
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a F	250 30.72	5	1	DBL Pethidine
	0000.72	Ū	-	Hydrochloride
SM Tab 50 mg to be delisted 1 August 2023)				nyaroomonae
c c ,				
RAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg		20		Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg	2.80	100	✓	Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
MITRIPTYLINE – Safety medicine; prescriber may determine of				
Tab 10 mg		100		Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg	2.51	100	1	Arrow-Amitriptyline
LOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescr	riber mav determine c	lispens	ina freau	encv
Tab 10 mg		30		Clomipramine Teva
5		00		Clomipramine Teva
Tab 25 mg		30	 ✓ 	
Tab 25 mg		30	1	
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er	ndorsement	30	-	<u>elemplamie reva</u>
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fro	ndorsement equency			
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fro b) Subsidy by endorsement – Subsidised for patients who w	ndorsement equency vere taking dosulepin	[dothie	pin] hydi	rochloride prior to 1 June
 OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fra b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar 	ndorsement equency vere taking dosulepin macists may annotate	[dothie	pin] hydi	rochloride prior to 1 June
 OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin 	ndorsement equency vere taking dosulepin macists may annotate ı] hydrochloride.	[dothie e the p	pin] hydi rescriptio	rochloride prior to 1 June n as endorsed where the
 OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fra b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar 	ndorsement equency vere taking dosulepin macists may annotate ı] hydrochloride.	[dothie	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg	ndorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris
 OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin 	ndorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg	ndorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg	ndorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg	ndorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg.	ndorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529 Dosulepin
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg	ndorsement equency vere taking dosulepin macists may annotate n] hydrochloride. 	[dothie e the p 30 50	epin] hydr rescriptio v	rochloride prior to 1 June n as endorsed where the Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529 Dosulepin Viatris 529
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023) IIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber	ndorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30 50	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529 Dosulepin Viatris 529
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg	ndorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30 50 ensing f	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529 Dosulepin Viatris 529
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023) IIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber Tab 10 mg	ndorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30 50 ensing f 50 100	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Mylan 529 Dosulepin Viatris 529 Y
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023) IIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber Tab 10 mg Tab 25 mg	hdorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30 50 50 ensing f 50 100 50	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529 Dosulepin Viatris 529 Y Tofranil Tofranil Tofranil
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023) IIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber Tab 10 mg	hdorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30 50 50 ensing f 50 100 50	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529 Dosulepin Viatris 529 Y Tofranil Tofranil Tofranil
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023) IIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber Tab 10 mg Tab 25 mg	hdorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30 50 50 ensing f 50 100 50	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529 Dosulepin Viatris 529 Y Tofranil Tofranil Tofranil
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023) IIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber Tab 25 mg Tab 25 mg ORTRIPTYLINE HYDROCHLORIDE – Safety medicine; presc	hdorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30 50 50 100 50 dispens	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529 Dosulepin Viatris 529 Y Tofranil Tofranil Tofranil Jency
OSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er a) Safety medicine; prescriber may determine dispensing fr b) Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin Tab 75 mg Cap 25 mg Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023) IIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber Tab 10 mg Tab 25 mg ORTRIPTYLINE HYDROCHLORIDE – Safety medicine; prescriber Tab 10 mg	hdorsement equency vere taking dosulepin macists may annotate 1) hydrochloride. 	[dothie e the p 30 50 50 100 50 dispens	epin] hydr rescriptio	rochloride prior to 1 June n as endorsed where th Dosulepin Mylan Dosulepin Viatris Dosulepin Mylan 529 Dosulepin Viatris 529 Y Tofranil Tofranil Tofranil Jency

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic ✓ Manufacturer
Monoamine-Oxidase Inhibitors (MAOIs) - N	Ion Selective		
TRANYLCYPROMINE SULPHATE			
Tab 10 mg		28	 ✓ Parnate S29 S29 ✓ Parnate
	22.94 45.88	50 100	
	45.88 96.00	100	✓ Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE			
* Tab 150 mg	11.80	60	✓ <u>Aurorix</u>
* Tab 300 mg		60	✓ <u>Aurorix</u>
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE – Brand switch fee pay	,	<i>,</i>	
* Tab 20 mg	2.86	84	 <u>Celapram</u>
	1.07	00	
* Tab 10 mg	1.07	28	 Escitalopram (Ethics)
券 Tab 20 mg	1.92	28	✓ Escitalopram
FLUOXETINE HYDROCHLORIDE			(Ethics)
 Tab dispersible 20 mg, scored – Subsidy by endorser Subsidised by endorsement 	nent2.50	28	✓ <u>Fluox</u>
 When prescribed for a patient who cannot so accordingly; or 	wallow whole tablets or cape	sules	and the prescription is endorsed
2) When prescribed in a daily dose that is not a endorsed. Note: Tablets should be combin			
Cap 20 mg		30	Brown & Burk S29
	2.91	84	✓ Fluox
	3.13	90	Arrow-Fluoxetine
Arrow-Fluoxetine to be Principal Supply on 1 June	2023		
(Fluox Cap 20 mg to be delisted 1 June 2023)			
PAROXETINE * Tab 20 mg	/ 11	90	 Loxamine
SERTRALINE		30	
* Tab 50 mg	0.99	30	 Setrona
· · · · · · · · · · · · · · · · · · ·			✓ Setrona AU
Setrona to be Principal Supply on 1 April 2023			
* Tab 100 mg	1.74	30	 ✓ Setrona ✓ Setrona AU
Setrona to be Principal Supply on 1 April 2023			
(Setrona AU Tab 50 mg to be delisted 1 April 2023)			
(Setrona AU Tab 100 mg to be delisted 1 April 2023)			

	Subsidy		Fully	Brand or
	(Manufacturer's Price		sidised	Generic
	\$	Per	/	Manufacturer
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2.60	28	✓ No	umed
Tab 45 mg	3.45	28	✓ No	umed
VENLAFAXINE				
* Cap 37.5 mg		84		lafax XR
* Cap 75 mg		84		lafax XR lafax XR
* Cap 150 mg	11.10	84	V EN	
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
DIAZEPAM – Safety medicine; prescriber may determine dispen	sing frequency			
Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorsement	27.92	5	🗸 Ho	spira
a) Up to 5 inj available on a PSO				
b) Only on a PSO				
c) PSO must be endorsed "not for anaesthetic procedur Rectal tubes 5 mg – Up to 5 tube available on a PSO		5	✓ Ste	eolid
5		5	• <u>51</u>	-50110
PHENYTOIN SODIUM				
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	104 58	5	🖌 Ho	enira
* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a		0	• 110	эрпа
PSO	154.01	5	🖌 Ho	spira
Control of Epilepsy				
CARBAMAZEPINE * Tab 200 mg	1/ 53	100	🖌 To	gretol
* Tab long-acting 200 mg		100		gretol CR
	33.96	200		gretol CR
* Tab 400 mg		100		gretol
* Tab long-acting 400 mg		100	🗸 Te	gretol CR
* Oral liq 20 mg per ml		250 ml	🗸 Te	gretol
CLOBAZAM - Safety medicine; prescriber may determine disper				
Tab 10 mg	9.12	50	🗸 Fri	sium
CLONAZEPAM - Safety medicine; prescriber may determine dis				
Oral drops 2.5 mg per ml	7.38	10 ml OP	🗸 Riv	/otril
ETHOSUXIMIDE				
Cap 250 mg	78.89	56		sential
	1 10 22	100	· ·	Ethosuximide S29
Oral lig 250 mg par 5 ml	140.88	100 200 ml		rontin
Oral liq 250 mg per 5 ml		200 ml	▼ Za	rontin
GABAPENTIN Note: Not subsidized in combination with subsidized progab	alin			
Note: Not subsidised in combination with subsidised pregab. * Cap 100 mg		100	🖌 Nu	pentin
* Cap 100 mg		100		pentin
* Cap 400 mg		100		pentin

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
LACOSAMIDE – Special Authority see SA1125 below – Retail p	harmacy				
▲ Tab 50 mg	25.04	14	 	/impat	
▲ Tab 100 mg		14	 	/impat	
-	200.24	56	 	/impat	
▲ Tab 150 mg	75.10	14	 	/impat	
J. J. J. J. J. J. J. J. J. J. J. J. J. J	300.40	56	 	/impat	
▲ Tab 200 mg		56	 	/impat	

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see note).

Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment. LAMOTRIGINE

LA	MOTRIGINE			
	Tab dispersible 2 mg55	5.00	30 🗸	Lamictal
	Tab dispersible 5 mg50	0.00	30 🗸	Lamictal
*	Tab dispersible 25 mg2	2.76	56 🖌	Logem
*	Tab dispersible 50 mg	3.31	56 🗸	Logem
*	Tab dispersible 100 mg	1.40	56 🖌	Logem
LE'	VETIRACETAM			
	Tab 250 mg	1.99	60 🗸	Everet
	Tab 500 mg	3.79	60 🗸	Everet
	Tab 750 mg14	1.39	60 🗸	Everet
	Tab 1,000 mg	3.59	60 🗸	Everet
	Oral liq 100 mg per ml44	1.78 300	ml OP 🛛 🗸	Levetiracetam-AFT
PH	ENOBARBITONE			
	For phenobarbitone oral liquid refer Standard Formulae, page 251			
*	Tab 15 mg).00 5	00 🗸	PSM
*	Tab 30 mg40		00 🗸	PSM
РН				
*	Tab 50 mg	5 00 2	00 🗸	Dilantin Infatab
	Cap 30 mg			Dilantin
	Cap 100 mg			Dilantin
*	Oral lig 30 mg per 5 ml			Dilantin
	EGABALIN			
гn	Note: Not subsidised in combination with subsidised gabapentin			
	Cap 25 mg	25	56 🗸	Pregabalin Pfizer
		7.80		Milpharm S29
*	Cap 75 mg			Pregabalin Pfizer
ጥ		3.10		•
				Milpharm S29
	Cap 150 mg4	.01 ;		Lyrica
	47			Pregabalin Pfizer
		2.44		Milpharm S29
	Cap 300 mg7	.38	56 🗸	Pregabalin Pfizer
PR	IMIDONE			
*	Tab 250 mg	7.35 1	00 🗸	Primidone Clinect

126 fully subsidised Principal Supply

	Subsidy (Manufacturer's Price)	Fully Subsidised	
	\$	Per	1	Manufacturer
SODIUM VALPROATE				
Tab 100 mg		100	1	Epilim Crushable
Tab 200 mg EC	27.44	100	1	Epilim
Tab 500 mg EC		100	1	Epilim
* Oral liq 200 mg per 5 ml		300 m	nl 🗸	Epilim S/F Liquid
			1	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	✓	Epilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retail pl	harmacy			
Cap 250 mg		60	1	Diacomit S29
Powder for oral liq 250 mg sachet		60	1	Diacomit S29

⇒SA1330 Special Authority for Subsidy

TOPIRAMATE

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

▲ Tab 25 mg	11.07	60	Arrow-Topiramate
ů –			 Topiramate Actavis
	26.04		 Topamax
Tab 50 mg	18.81	60	Arrow-Topiramate
C C C C C C C C C C C C C C C C C C C			 Topiramate Actavis
	44.26		 Topamax
Tab 100 mg	31.99	60	Arrow-Topiramate
ů.			 Topiramate Actavis
	75.25		 Topamax
Tab 200 mg	55.19	60	Arrow-Topiramate
•			 Topiramate Actavis
	129.85		 Topamax
Sprinkle cap 15 mg	20.84	60	 Topamax
Sprinkle cap 25 mg	26.04	60	 Topamax
IGABATRIN – Special Authority see SA2088 below – Retail pharr	nacy		-
Tab 500 mg		100	✓ Sabril

➡SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

1.3 Patient has tuberous sclerosis complex; and

2 Either:

- 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
- 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields..

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and 2 Either:
- 2 Eithe
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 111

Acute Migraine Treatment

RIZATRIPTAN Tab orodispersible 10 mg	3 65	30	✓ Rizamelt
SUMATRIPTAN	.0.00	50	
Tab 50 mg	14.41	90	✓ Sumagran
Tab 100 mg2	22.68	90	 Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj per prescription	34.00	2 OP	✓ Imigran
Prophylaxis of Migraine			
For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM,	page 50		
PIZOTIFEN * Tab 500 mcg2	23.21	100	 Sandomigran
Antinausea and Vertigo Agents			
For Antispasmodics refer to ALIMENTARY TRACT, page 8			
APREPITANT – Special Authority see SA0987 below – Retail pharmacy			
Cap 2 × 80 mg and 1 × 125 mg	30.00	3 OP	 Emend Tri-Pack
SA0987 Special Authority for Subsidy			is underseine biebb
Initial application from any relevant practitioner. Approvals valid for 12 emetogenic chemotherapy and/or anthracycline-based chemotherapy for			0 0 0 7
Renewal from any relevant practitioner. Approvals valid for 12 months w	vhere the patie	ent is underg	
chemotherapy and/or anthracycline-based chemotherapy for the treatme	nt of malignar	ncy.	
BETAHISTINE DIHYDROCHLORIDE * Tab 16 mg	4.00	100	1 Carro
★ lab 16 mg	.4.62	100	✓ <u>Serc</u>
CYCLIZINE HYDROCHLORIDE			

	Subsidy		Fully	
	(Manufacturer's Price) \$	Su Per	ubsidisec	I Generic Manufacturer
YCLIZINE LACTATE	.			
Inj 50 mg per ml, 1 ml ampoule – Up to 10 inj available on	9			
PSO		10	1	HameIn
OMPERIDONE				
k Tab 10 mg	2.85	100	1	Pharmacy Health
	4.00	100		Domperidone Viatris
Pharmacy Health Tab 10 mg to be delisted 1 June 2023)				
 Inj 400 mcg per ml, 1 ml ampoule 	93.00	10	1	Martindale S29
Patch 1.5 mg – Special Authority see SA1998 below – Re		10		
pharmacy		2	1	Scopoderm TTS
SA1998 Special Authority for Subsidy				•
nitial application from any relevant practitioner. Approvals va	alid for 1 year for applic	ations m	neetina	the following criteria.
	y respond to oral anti-n s of at least two other a	ausea a Iternativ	igents; (ve treatr	nents have proven
where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 y enefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE	y respond to oral anti-n s of at least two other a year where the treatment	ausea a Iternativ nt remai	igents; o ve treatr	nents have proven opriate and the patient
where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 y enefiting from treatment.	y respond to oral anti-n s of at least two other a year where the treatment	ausea a Iternativ	igents; o ve treatr	nents have proven opriate and the patient Metoclopramide
where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 y enefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE	y respond to oral anti-n s of at least two other a year where the treatment 	ausea a Iternativ nt remai	igents; o ve treatr	nents have proven opriate and the patient
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE K Tab 10 mg – Up to 30 tab available on a PSO k Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a 	y respond to oral anti-n s of at least two other a year where the treatment 	ausea a Iternativ nt remai 100	igents; o ve treatr	nents have proven opriate and the patient Metoclopramide Actavis 10
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE K Tab 10 mg – Up to 30 tab available on a PSO 	y respond to oral anti-n s of at least two other a vear where the treatmen 1.30 PSO7.00	ausea a Iternativ nt remai 100	igents; (ve treatr	nents have proven opriate and the patient Metoclopramide Actavis 10
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE K Tab 10 mg – Up to 30 tab available on a PSO k Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON K Tab 4 mg 	y respond to oral anti-n s of at least two other a vear where the treatmen 	ausea a Iternativ nt remai 100 10	igents; (re treatr ns appr	nents have proven opriate and the patient Metoclopramide Actavis 10 Baxter
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE K Tab 10 mg – Up to 30 tab available on a PSO k Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON 	y respond to oral anti-n s of at least two other a vear where the treatmen 	ausea a Iternativ nt remai 100 10	igents; (re treatr ns appr	nents have proven opriate and the patient Metoclopramide Actavis 10 Baxter Periset Onrex Ondansetron
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE K Tab 10 mg – Up to 30 tab available on a PSO Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON K Tab 4 mg Tab disp 4 mg – Up to 10 tab available on a PSO 	y respond to oral anti-n s of at least two other a rear where the treatmen 	ausea a Iternativ nt remai 100 10 50 10	igents; (/e treatr ns appr	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u>
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE K Tab 10 mg – Up to 30 tab available on a PSO k Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON K Tab 4 mg 	y respond to oral anti-n s of at least two other a rear where the treatmen 	ausea a Iternativ nt remai 100 10 50	igents; (ve treatr ins appr v	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u> Periset
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE Tab 10 mg – Up to 30 tab available on a PSO Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON Tab 4 mg Tab disp 4 mg – Up to 10 tab available on a PSO Kab 8 mg 	y respond to oral anti-n s of at least two other a vear where the treatmen 	ausea a Iternativ nt remai 100 10 50 10 50	igents; (ve treatr ins appr v v v v v	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u> Periset Onrex
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE K Tab 10 mg – Up to 30 tab available on a PSO Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON K Tab 4 mg Tab disp 4 mg – Up to 10 tab available on a PSO 	y respond to oral anti-n s of at least two other a vear where the treatmen 	ausea a Iternativ nt remai 100 10 50 10	igents; (ve treatr ins appr v v v v v	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u> Periset Onrex <u>Ondansetron</u>
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE K Tab 10 mg – Up to 30 tab available on a PSO k Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON K Tab 4 mg Tab disp 4 mg – Up to 10 tab available on a PSO K Tab 8 mg Tab disp 8 mg – Up to 10 tab available on a PSO 	y respond to oral anti-n s of at least two other a vear where the treatmen 	ausea a Iternativ nt remai 100 10 50 10 50	igents; (ve treatr ins appr v v v v v	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u> Periset Onrex
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE Tab 10 mg – Up to 30 tab available on a PSO k Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON Tab disp 4 mg – Up to 10 tab available on a PSO K Tab 8 mg Tab disp 8 mg – Up to 10 tab available on a PSO Conrex Tab 4 mg to be delisted 1 August 2023) 	y respond to oral anti-n s of at least two other a vear where the treatmen 	ausea a Iternativ nt remai 100 10 50 10 50	igents; (ve treatr ins appr v v v v v	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u> Periset Onrex <u>Ondansetron</u>
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE Tab 10 mg – Up to 30 tab available on a PSO Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON Tab 4 mg Tab disp 4 mg – Up to 10 tab available on a PSO Tab disp 8 mg – Up to 10 tab available on a PSO Onrex Tab 4 mg to be delisted 1 August 2023) 	y respond to oral anti-n s of at least two other a vear where the treatmen 	ausea a Iternativ nt remai 100 10 50 10 50	igents; (ve treatr ins appr v v v v v	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u> Periset Onrex <u>Ondansetron</u>
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE k Tab 10 mg - Up to 30 tab available on a PSO k Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a DNDANSETRON k Tab 4 mg Tab disp 4 mg - Up to 10 tab available on a PSO k Tab 8 mg Tab disp 8 mg - Up to 10 tab available on a PSO Mark Tab 4 mg to be delisted 1 August 2023) Onrex Tab 8 mg to be delisted 1 August 2023) PROCHLORPERAZINE 	y respond to oral anti-n s of at least two other a rear where the treatmen 	ausea a Iternativ nt remai 100 10 50 10 50	igents; (ve treatr ins appr v v v v v	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u> Periset Onrex <u>Ondansetron</u>
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE Tab 10 mg – Up to 30 tab available on a PSO Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a DNDANSETRON Tab 4 mg Tab disp 4 mg – Up to 10 tab available on a PSO Tab disp 8 mg – Up to 10 tab available on a PSO Onrex Tab 4 mg to be delisted 1 August 2023) PROCHLORPERAZINE 	y respond to oral anti-n s of at least two other a rear where the treatmen 	ausea a Iternativ 100 10 50 10 50 10	igents; (ve treatr ins appr v v v v v	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u> Periset Onrex <u>Ondansetron</u>
 where the patient cannot tolerate or does not adequatel 2 Control of clozapine-induced hypersalivation where trial ineffective. Renewal from any relevant practitioner. Approvals valid for 1 yenefiting from treatment. METOCLOPRAMIDE HYDROCHLORIDE k Tab 10 mg - Up to 30 tab available on a PSO k Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a DNDANSETRON k Tab 4 mg Tab disp 4 mg - Up to 10 tab available on a PSO k Tab 8 mg Tab disp 8 mg - Up to 10 tab available on a PSO Mark Tab 4 mg to be delisted 1 August 2023) Onrex Tab 8 mg to be delisted 1 August 2023) PROCHLORPERAZINE 	y respond to oral anti-n s of at least two other a rear where the treatmen 	ausea a Iternativ 100 10 50 10 50 10	igents; (re treatr ns appr	nents have proven opriate and the patient <u>Metoclopramide</u> <u>Actavis 10</u> <u>Baxter</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u> Periset Onrex <u>Ondansetron</u> <u>ODT-DRLA</u>

General

AMISULPRIDE - Safety medicine; prescriber may deterr	nine dispensing frequency		
Tab 100 mg		30	 Sulprix
Tab 200 mg	14.96	60	 Sulprix
Tab 400 mg		60	 Sulprix

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	(Manufacturer's Price		Subsidised	
	\$	Per		Manufacturer
RIPIPRAZOLE - Safety medicine; prescriber may determine c	lispensing frequency	/		
Tab 5 mg		30	✓	Aripiprazole Sandoz
Tab 10 mg		30	✓	Aripiprazole Sandoz
Tab 15 mg		30	✓	Aripiprazole Sandoz
Tab 20 mg		30	✓	Aripiprazole Sandoz
Tab 30 mg		30	1	Aripiprazole Sandoz
ILORPROMAZINE HYDROCHLORIDE – Safety medicine; pr	rescriber may deterr	nine disr	oensina fr	equency
Tab 10 mg – Up to 30 tab available on a PSO		100		Largactil
Tab 25 mg – Up to 30 tab available on a PSO		100		Largactil
Tab 100 mg – Up to 30 tab available on a PSO		100		Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO		10		Largactil
OZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing frequ	IODOV			
Tab 25 mg		50	1	Clopine
1 ab 20 mg	0.09	50		Clozaril
	13.37	100		Clopine
	10.07	100		Clozaril
Tab 50 mg	8 67	50		Clopine
Tab 50 mg	17.33	100		Clopine
Tab 100 mg		50		Clopine
Tab 100 mg		50		Clozaril
	34.65	100		Clopine
	04.00	100		Clozaril
Tab 200 mg	34 65	50		Clopine
Tab 200 mg	69.30	100		Clopine
Suspension 50 mg per ml		100 ml		Versacloz
			•	VCIGUOIOZ
LOPERIDOL – Safety medicine; prescriber may determine d				C
Tab 500 mcg – Up to 30 tab available on a PSO		100		Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg – Up to 30 tab available on a PSO		50		Serenace
Oral lig 0 mg nor ml Un to 000 ml available on a PCO	29.72	100		Serenace Serenace
Oral liq 2 mg per ml – Up to 200 ml available on a PSO		100 ml 10		Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a P			•	Selellace
/OMEPROMAZINE - Safety medicine; prescriber may dete				
Tab 25 mg (33.8 mg as a maleate)		100		Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan
Tab 100 mg (135 mg as a maleate)		100		Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	~	Nozinan
/OMEPROMAZINE HYDROCHLORIDE – Safety medicine;	prescriber may dete	rmine di	spensing	frequency
Inj 25 mg per ml, 1 ml ampoule		5	✓	Neuraxpharm S29
				Nozinan S29 S29
	24.48	10		Wockhardt
	33.50			Nozinan
Wockhardt to be Principal Supply on 1 April 2023 bzinan Inj 25 mg per ml, 1 ml ampoule to be delisted 1 April 2				
	,			
THIUM CARBONATE – Safety medicine; prescriber may dete				Driedel
Tab long-acting 400 mg		100	-	Priadel Develop
Cap 250 mg	9.42	100	•	Douglas

	Cubaidu		Fully	Drand ar
	Subsidy (Manufacturer's Price)		Fully Subsidised	
	(Manulaciulei S Flice)	Per		Manufacturer
	•			
OLANZAPINE – Safety medicine; prescriber may determine disp	• • •	~~		_ .
Tab 2.5 mg		28		Zypine
Tab 5 mg		28		Zypine
Tab orodispersible 5 mg		28	-	Zypine ODT
Tab 10 mg	2.01	28		Zypine
Tab orodispersible 10 mg	2.38	28	✓	Zypine ODT
PERICYAZINE - Safety medicine; prescriber may determine dis	pensina frequency			
Tab 2.5 mg		84	1	Neulactil
	12.49	100		Neulactil
Tab 10 mg		84		Neulactil
	44.45	100		Neulactil
		100	•	neulaciii
QUETIAPINE – Safety medicine; prescriber may determine disp	ensing frequency			
Tab 25 mg	2.15	90	-	Quetapel
Tab 100 mg	5.06	90	✓	Quetapel
Tab 200 mg	8.90	90	✓	Quetapel
Tab 300 mg		90	✓	Quetapel
RISPERIDONE - Safety medicine; prescriber may determine dis	snensing frequency			
Tab 0.5 mg		60	1	Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
		60		Risperidone (Teva)
Tab 3 mg		60 60		
Tab 4 mg				Risperidone (Teva)
Oral liq 1 mg per ml	8.90	30 m	✓	Risperon
ZIPRASIDONE – Safety medicine; prescriber may determine dis	pensing frequency			
Cap 20 mg	17.90	60	✓	Zusdone
Cap 40 mg	27.41	60	✓	Zusdone
Cap 60 mg		60	✓	Zusdone
Cap 80 mg		60	✓	Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre	scriber may determin	o dier	oonsina fra	
Tab 10 mg	•	100		Clopixol
Tab T0 Hig		100	•	Сюріхої
Depot Injections				
Depot injections				
FLUPENTHIXOL DECANOATE - Safety medicine; prescriber m	av determine dispens	sina fi	requency	
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Fluanxol
Inj 20 mg per ml, 2 ml $-$ Up to 5 inj available on a PSO		5		Fluanxol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Fluanxol
		-		TIUATIAO
HALOPERIDOL DECANOATE – Safety medicine; prescriber ma		•		
Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Haldol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Haldol Concentrate
			✓	Haldol
				Decanoas S29
OLANZAPINE - Special Authority see SA1428 on the next page	- Retail nharmaou			
Safety medicine; prescriber may determine dispensing frequ				
		1		Zuprova Polarova
Inj 210 mg vial				Zyprexa Relprevv
Inj 300 mg vial		1		Zyprexa Relprevv
Inj 405 mg vial		1	•	Zyprexa Relprevv

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy	Ful	ly Brand or
(Manufacturer's Pr	rice) Subsidise	d Generic
\$	Per •	Manufacturer

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or

- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency
--

Inj 25 mg syringe	 1	Invega Sustenna
Inj 50 mg syringe	 1	Invega Sustenna
Inj 75 mg syringe	 1	Invega Sustenna
Inj 100 mg syringe	 1	Invega Sustenna
Inj 150 mg syringe	 1	 Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or 2 All of the following:

- 2.1 The patient has schizophrenia or other psychotic disorder; and
- 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
- 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE - Special	Authority see SA2167 below – Retail pharm	пасу	
Inj 175 mg syringe		1	🗸 Invega Trinza
Inj 263 mg syringe		1	🗸 Invega Trinza
Inj 350 mg syringe		1	🗸 Invega Trinza
Inj 525 mg syringe		1	🗸 Invega Trinza
			-

⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The patient has schizophrenia; and

2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA1427 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing	frequency		
Inj 25 mg vial		1	Risperdal Consta
Inj 37.5 mg vial		1	Risperdal Consta
Inj 50 mg vial		1	 Risperdal Consta

Subsidy (Manufacturer's Price)	Subsid	Fully lised	Brand or Generic
`\$	Per	✓	Manufacturer

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml – Up to 5 inj available on a PSO19.80	5	 Clopixol
---	---	------------------------------

Anxiolytics

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg		100	 Buspirone Viatris
* Tab 10 mg	12.50	100	 Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may determ	ne dispensing frequency		
Tab 500 mcg	5.64	100	🗸 Paxam
Tab 2 mg	10.78	100	 Paxam
DIAZEPAM - Safety medicine; prescriber may determine of	dispensing frequency		
Tab 2 mg	61.07	500	 Arrow-Diazepam
Tab 5 mg	73.60	500	 Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determin	e dispensing frequency		
Tab 1 mg	9.72	250	 Ativan
Tab 2.5 mg		100	✓ Ativan

Multiple Sclerosis Treatments

⇒SA2176 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 2 Patients has an EDSS score between 0 6.0; and
- 3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 4 All of the following:
 - 4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and

Subsidy (Manufacturer's Price)	Subs	Fully sidised	Brand or Generic	
\$	Per	✓	Manufacturer	

- 4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
- 4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and

4.5 Either:

- 4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
- 4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and

6 Any of the following:

- 6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
- 6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
- 6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
- 6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
- 6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2176 on the previous page - Retail pharmacy

a) Wastage claimable		
b) Note: Treatment on two or more funded multiple sclerosis treatments sim	ultaneously is	not permitted.
Cap 120 mg520.00	14	 Tecfidera
Cap 240 mg2,000.00	56	 Tecfidera
FINGOLIMOD - Special Authority see SA2176 on the previous page - Retail pha	armacy	
a) Wastage claimable		
b) Note: Treatment on two or more funded multiple sclerosis treatments sim	ultaneously is	not permitted.
Cap 0.5 mg2,200.00	28	Gilenya
GLATIRAMER ACETATE - Special Authority see SA2176 on the previous page	– Retail pharm	nacv
Note: Treatment on two or more funded multiple sclerosis treatments simulta		
Inj 40 mg prefilled syringe1,137.48	12	Copaxone
INTERFERON BETA-1-ALPHA - Special Authority see SA2176 on the previous	page – Retail	pharmacv
Note: Treatment on two or more funded multiple sclerosis treatments simulta		
Inj 6 million iu prefilled syringe1,170.00	4	Avonex
Injection 6 million iu per 0.5 ml pen injector1,170.00	4	 Avonex Pen
INTERFERON BETA-1-BETA - Special Authority see SA2176 on the previous p	age – Retail pl	narmacy
Note: Treatment on two or more funded multiple sclerosis treatments simulta	neously is not	permitted.
Inj 8 million iu per 1 ml1,322.89	15	 Betaferon
NATALIZUMAB - Special Authority see SA2176 on the previous page - Retail p	narmacy	
Note: Treatment on two or more funded multiple sclerosis treatments simulta	neously is not	permitted.
Inj 20 mg per ml, 15 ml vial1,750.00	1	 Tysabri
OCRELIZUMAB - Special Authority see SA2176 on the previous page - Retail p	harmacy	
Note: Treatment on two or more funded multiple sclerosis treatments simulta		permitted.
Inj 30 mg per ml, 10 ml vial9,346.00	1	 Ocrevus

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	ully sed	Brand or Generic Manufacturer
TERIFLUNOMIDE - Special Authority see SA2176 on page 133 -	- Retail pharmacy			
a) Wastage claimable	notali phannaoy			
 b) Note: Treatment on two or more funded multiple sclerosis 	treatments simultan	eouslv is no	t perr	nitted.
Tab 14 mg		28	•	ubagio
			_	
Sedatives and Hypnotics				
MELATONIN - Special Authority see SA1666 below - Retail pha	rmacy			
Tab modified-release 2 mg – No more than 5 tab per day		30	✓ V	igisom
➡SA1666 Special Authority for Subsidy				
Initial application only from a psychiatrist, paediatrician, neurolog	gist, respiratory speci	alist or med	ical p	ractitioner on the
recommendation of a psychiatrist, paediatrician, neurologist or res	piratory specialist. A	pprovals va	lid fo	r 12 months for
applications meeting the following criteria:				
All of the following:				
1 Patient has been diagnosed with persistent and distressing				
(including, but not limited to, autism spectrum disorder or a	21	,		,
2 Behavioural and environmental approaches have been trie				propriate; and
 Funded modified-release melatonin is to be given at doses Patient is aged 18 years or under*. 	no greater than 10 h	ig per day;	anu	
Renewal only from a psychiatrist, paediatrician, neurologist, respi	ratory energialist or m	odical pract	tiona	r on the recommendation
of a psychiatrist, paediatrician, neurologist or respiratory specialist				
following criteria:			ioi up	photono mooting the
All of the following:				
1 Patient is aged 18 years or under*; and				
2 Patient has demonstrated clinically meaningful benefit from	n funded modified-rel	ease melato	onin (d	clinician determined); and
3 Patient has had a trial of funded modified-release melatoni	n discontinuation with	nin the past	12 m	onths and has had a
recurrence of persistent and distressing insomnia; and				
4 Funded modified-release melatonin is to be given at doses	no greater than 10 n	ng per day.		
Note: Indications marked with * are unapproved indications.				
MIDAZOLAM - Safety medicine; prescriber may determine dispe	nsing frequency			
Inj 1 mg per ml, 5 ml ampoule	3.95	10	🗸 W	lidazolam Mylan
	6.10		🗸 W	lidazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available				
on a PSO		10	✓ P	
On a PSO for status epilepticus use only. PSO must be				
Inj 5 mg per ml, 3 ml ampoule		5	✓ IVI	lidazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule – Up to 5 inj available o		5	✓ P	finar
a PSO On a PSO for status epilepticus use only. PSO must be e		-		
			Se Uli	iy.
PHENOBARBITONE SODIUM – Special Authority see SA1386 b			<i>.</i>	11 kili 🚳
Inj 200 mg per ml, 1 ml ampoule		10	✓ IVI	ax Health S29
SA1386 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid	without further renew	val unless n	otified	d for applications meeting
the following criteria:				
Both:	to other ensures and			
1 For the treatment of terminal agitation that is unresponsive	U .			
2 The applicant is part of a multidisciplinary team working in				
TEMAZEPAM – Safety medicine; prescriber may determine dispe				
Tab 10 mg	1.33	25	✓ N	ormison

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
TRIAZOLAM – Safety medicine; prescriber may determine disp	ensing frequency			
Tab 125 mcg	5.10	100		
	(9.85)		ł	Hypam
Tab 250 mcg		100		
	(11.20)		I	Hypam
ZOPICLONE - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 7.5 mg		500	 Image: A second s	Zopiclone Actavis
Spinal Muscular Atrophy				
NUSINERSEN – PCT only – Special Authority see SA2174 belo	W			
Inj 12 mg per 5 ml vial	120,000.00	1	√ 9	Spinraza
► SA2174 Special Authority for Subsidy				

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

NERVOUS	SYSTEM
----------------	--------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
timulants/ADHD Treatments				
OMOXETINE				
Cap 10 mg	18.41	28	-	APO-Atomoxetine APO-Atomoxetine S29 S29
			✓ (Generic Partners
	107.03		✓ :	Strattera
Cap 18 mg		28	✓	APO-Atomoxetine
			✓ (Generic Partners
	107.03		✓ :	Strattera
Cap 25 mg		28	✓	APO-Atomoxetine
			✓ (Generic Partners
Cap 40 mg		28	 Image: A second s	APO-Atomoxetine
			✓ (Generic Partners
	107.03		✓ :	Strattera
Cap 60 mg		28	 Image: A second s	APO-Atomoxetine
			v 1	APO-Atomoxetine S29 S29
			✓ (Generic Partners
Cap 80 mg		28	1	APO-Atomoxetine
			•	APO-Atomoxetine S29 S29
			1	Generic Partners
Cap 100 mg	58.48	28		APO-Atomoxetine
			_	APO-Atomoxetine S29 S29
			✓ (Generic Partners
rattera Cap 10 mg to be delisted 1 November 2023) rattera Cap 18 mg to be delisted 1 November 2023) rattera Cap 40 mg to be delisted 1 November 2023)				
XAMFETAMINE SULFATE – Special Authority see SA1149 t	pelow – Retail pharma	cv		
 a) Only on a controlled drug form b) Safety medicine; prescriber may determine dispensing free 		.,		
Tab 5 mg		100	✓	PSM
	28.50			Aspen

SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
(Man	ufacturer's Price)	Subsid	dised	Generic
	\$	Per	1	Manufacturer

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and

2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

1 The treatment remains appropriate and the patient is benefiting from treatment; and

2 Either:

- 2.1 Applicant is a paediatrician or psychiatrist; or
- 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

a)	Only	y on a	controlled	drug form	
----	------	--------	------------	-----------	--

b)	Safety medicine;	prescriber ma	y determine	dispensing	frequency
----	------------------	---------------	-------------	------------	-----------

b) Salety medicine; prescriber may determine disper	ising frequency		
Tab immediate-release 5 mg		30	 Rubifen
Tab immediate-release 10 mg		30	 Ritalin
0			 Rubifen
Tab extended-release 18 mg	7.75	30	 Methylphenidate ER Teva
Tab immediate-release 20 mg	7.85	30	 Rubifen
Tab sustained-release 20 mg		30	 Rubifen SR
Tab extended-release 27 mg		30	 Methylphenidate ER Teva
Tab extended-release 36 mg		30	 Methylphenidate ER Teva
Tab extended-release 54 mg	22.25	30	 Methylphenidate ER Teva

SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or	
(Manuacturer's Frice)		Subsidised	Generic	
\$	Per	1	Manufacturer	

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER - Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER - Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	 Concerta
Tab extended-release 27 mg		30	 Concerta
Tab extended-release 36 mg		30	 Concerta
Tab extended-release 54 mg		30	 Concerta
Cap modified-release 10 mg		30	 Ritalin LA
Cap modified-release 20 mg		30	 Ritalin LA
Cap modified-release 30 mg	25.52	30	 Ritalin LA
Cap modified-release 40 mg	30.60	30	 Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and

4 Either:

4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Si	ubsidised	Generic	
\$	Per	✓	Manufacturer	

4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Retail pharmacy

Tab 100 mg29.13 60

➡SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and

3 Either:

- 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
- 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE

* Tab 5 mg * Tab 10 mg	90 90	 ✓ Donepezil-Rex ✓ Donepezil-Rex
RIVASTIGMINE – Special Authority see SA1488 below – Re Patch 4.6 mg per 24 hour	 30	✓ <u>Rivastigmine Patch</u> BNM 5
Patch 9.5 mg per 24 hour	 30	✓ <u>Rivastigmine Patch</u> BNM 10

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Modavigil

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	ully Brand or sed Generic Manufacturer
Treatments for Substance Dependence			
BUPRENORPHINE WITH NALOXONE – Special Authority see 5 a) No patient co-payment payable b) Safety medicine; prescriber may determine dispensing fre		il pharmacy	
Tab sublingual 2 mg with naloxone 0.5 mg	11.76	28	 <u>Buprenorphine</u> Naloxone BNM
Tab sublingual 8 mg with naloxone 2 mg	34.00	28	✓ <u>Buprenorphine</u> Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health..

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release	150 mg	11.00	30	🗸 Zyban
----------------------	--------	-------	----	---------

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per		Brand or Generic Manufacturer
DISULFIRAM				
Tab 200 mg		100	✓ <u>I</u>	Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see SA1	408 below - Retail p	harma	асу	
Tab 50 mg		30	A Market Ma Market Market Mar	Valtraccord

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A

b) Note. Direct i tovision by a phannacist pennitted under the provision		
Patch 7 mg - Up to 28 patch available on a PSO 19.14	- 28	 Habitrol
Patch 7 mg for direct distribution only - [Xpharm]4.13	7	 Habitrol
Patch 14 mg - Up to 28 patch available on a PSO21.05	28	 Habitrol
Patch 14 mg for direct distribution only - [Xpharm]6.48	7	 Habitrol
Patch 21 mg - Up to 28 patch available on a PSO24.12	28	 Habitrol
Patch 21 mg for direct distribution only - [Xpharm]10.93	7	 Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO 19.76	216	 Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	36	 Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO21.65	216	 Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	36	 Habitrol
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO	384	 Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]9.04	96	 Habitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO21.42	204	 Habitrol
38.21	384	 Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]9.04	96	 Habitrol
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO24.17	204	 Habitrol
44.17	384	 Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]10.47	96	 Habitrol
Gum 4 mg (Mint) - Up to 384 piece available on a PSO	384	 Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm] 10.47	96	 Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	53 OP	 Varenicline Pfizer
Tab 1 mg17.62	56	 Varenicline Pfizer

Subsidy (Manufacturer's Price)		Fully ubsidised	Brand or Generic	
\$	Per	1	Manufacturer	

► SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Sul	osidised	Generic
	\$	Per	 ✓ 	Manufacturer
Chemotherapeutic Agents				
Alkylating Agents				
BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist -	Special Authority see	SA215	3 below	
Inj 25 mg vial		1	🗸 R	libomustin
Ini 100 mg vial		1	🗸 R	libomustin
Inj 1 mg for ECP		1 mg	🗸 В	axter
► SA2153 Special Authority for Subsidy				
Initial application - (treatment naive CLL) only from a relevant	nt specialist or medica	al practiti	oner on t	the recommendation of a
relevant specialist. Approvals valid for 12 months for applications	meeting the followin	g criteria	l:	

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2 Both:

- 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2.2 Either:

2.2.1 Both:

- 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
- 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application ---- (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2: and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and

~

5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN – PCT – Retail pharmacy-Specialist	05 400	/ .
Tab 2 mg	.25 100	 Myleran
CARBOPLATIN – PCT only – Specialist		
Inj 10 mg per ml, 45 ml vial32	.59 1	 DBL Carboplatin
45	.20	 Carboplatin Ebewe
48	.50	 Carbaccord
Inj 1 mg for ECP0	.10 1 mg	 Baxter
CARMUSTINE – PCT only – Specialist		
Inj 100 mg vial710	.00 1	✓ BICNU
Inj 100 mg for ECP710	.00 100 mg OP	✓ Baxter
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist	-	
Tab 2 mg	.06 25	 Leukeran FC
CISPLATIN – PCT only – Specialist		
	.00 1	 Cisplatin Ebewe
Inj 1 mg per ml, 50 ml vial15 Inj 1 mg per ml, 100 ml vial21		 Cisplatin Ebewe Cisplatin Ebewe
nij i nig per nii, roo nii vial		✓ DBL Cisplatin
Inj 1 mg for ECP0		✓ Baxter
, ,	.or ring	
CYCLOPHOSPHAMIDE	00 F0	(O)
Tab 50 mg – PCT – Retail pharmacy-Specialist		 <u>Cyclonex</u>
Inj 1 g vial – PCT – Retail pharmacy-Specialist		 Endoxan
127.		 Cytoxan
Inj 2 g vial – PCT only – Specialist		 Endoxan
Inj 1 mg for ECP – PCT only – Specialist0	.04 1 mg	 Baxter
IFOSFAMIDE – PCT only – Specialist		
Inj 1 g96		 Holoxan
lnj 2 g180		 Holoxan
Inj 1 mg for ECP0	.10 1 mg	 Baxter

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

(/	Subsidy /anufacturer's Price) \$	Per	Fully Subsidised	Generic
LOMUSTINE – PCT – Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	✓	CeeNU
Cap 40 mg	399.15	20	~	CeeNU
MELPHALAN				
Tab 2 mg – PCT – Retail pharmacy-Specialist	40.70	25	✓	Alkeran
Inj 50 mg – PCT only – Specialist	65.00	1	✓	Melpha
	67.80		✓	Alkeran
			1	Alkeran S29 S29
XALIPLATIN – PCT only – Specialist				
Inj 100 mg vial	25.01	1	~	Oxaliplatin Actavis 100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial		1	✓	Oxaliplatin Accord
Inj 1 mg for ECP	0.48	1 mg	, 🗸	Baxter
HIOTEPA – PCT only – Specialist				
Inj 15 mg vial	CBS	1	1	Bedford S29
, ,			1	Max Health S29
			1	THIO-TEPA S29
			1	Tepadina S29
Inj 100 mg vial	CBS	1		Max Health S29
		•		Tepadina S29
			•	i opudinu dav
Antimetabolites				
ZACITIDINE – PCT only – Specialist – Special Authority see SA2 Inj 100 mg vial		1	1	Azacitidine Dr

Inj 100 mg vial		1	 <u>Azacitidine Dr</u> Reddv's
Inj 1 mg for ECP	0.83	1 mg	✓ Baxter

⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

- Both:
 - 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully Brand or
	(Manufacturer's Price	ce) Subs	sidised Generic
	\$	Per	 Manufacturer
CALCIUM FOLINATE			
Tab 15 mg - PCT - Retail pharmacy-Specialist	135.33	10	 DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist	17.10	5	🗸 Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia	alist7.28	1	 Calcium Folinate Sandoz
			 Calcium Folinate Candez CO0 con
	70.00	10	Sandoz S29 S29
Inj 50 mg – PCT – Retail pharmacy-Specialist		10	 Leucovorin Pharmacia S29
Inj 10 mg per ml, 10 ml vial – PCT only – Specialist	9.49	1	 Calcium Folinate Sandoz
Inj 100 mg – PCT only – Specialist	7.33	1	 Calcium Folinate Ebewe
	94.90	10	✓ Leucovorin
			Pharmacia S29
Inj 300 mg – PCT only – Specialist	22.51	1	 Calcium Folinate Ebewe
	25.14		 Leucovorin DBL S29
Inj 10 mg per ml, 35 ml vial – PCT only – Specialist	25.14	1	 Calcium Folinate Sandoz
			 Calcium Folinate
			Sandoz S29 S29
Inj 1 g – PCT only – Specialist	67.51	1	 Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial – PCT only – Specialist	72.00	1	 Calcium Folinate Sandoz
Inj 1 mg for ECP – PCT only – Specialist	0.06	1 mg	✓ Baxter
CAPECITABINE – Retail pharmacy-Specialist			
Tab 150 mg	10.00	60	 Capercit
Tab 500 mg		120	✓ Capecitabine- DRLA S29
			✓ Capercit
CLADRIBINE – PCT only – Specialist			
Inj 2 mg per ml, 5 ml		1	 Litak S29
Inj 1 mg per ml, 10 ml		1	 Leustatin
Inj 10 mg for ECP	749.96	10 mg OP	 Baxter
CYTARABINE			
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia Inj 100 mg per ml, 20 ml vial – PCT – Retail	alist472.00	5	✓ Pfizer
pharmacy-Specialist		1	 Pfizer
Inj 1 mg for ECP – PCT only – Specialist Inj 100 mg intrathecal syringe for ECP – PCT only – Specia		10 mg 100 mg OP	✓ Baxter✓ Baxter
FLUDARABINE PHOSPHATE			
Tab 10 mg - PCT - Retail pharmacy-Specialist		20	 Fludara Oral
Inj 50 mg vial – PCT only – Specialist Inj 50 mg for ECP – PCT only – Specialist	634.00	5	 Fludarabine Ebewe

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pric	e) Sub	Fully sidised	Brand or Generic
	(Manulacialers i fic \$	Per		Manufacturer
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist		1	✓	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist		1	✓	Fluorouracil Accord
Inj 1 mg for ECP – PCT only – Specialist	0.62	100 mg	✓	Baxter
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	✓	DBL Gemcitabine
lnj 1 g		1		Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg		Baxter
IRINOTECAN HYDROCHLORIDE – PCT only – Specialist		U U		
Inj 20 mg per ml, 5 ml vial		1	1	Accord
j - j	71.44		✓	rinotecan Actavis
				100
	100.00		✓	rinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	✓	Baxter
MERCAPTOPURINE		0		
Tab 50 mg – PCT – Retail pharmacy-Specialist		25	✓	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialis			-	
Special Authority see SA1725 below		100 ml OP	1	Allmercap

➡SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

Tab 2.5 mg – PCT – Retail pharmacy-Specialist	90	✓ <u>Trexate</u>
	••	✓ <u>Trexate</u>
	5	 Methotrexate DBL
Inj 7.5 mg prefilled syringe14.61	1	 Methotrexate Sandoz
Inj 10 mg prefilled syringe14.66	1	 Methotrexate Sandoz
Inj 15 mg prefilled syringe14.77	1	 Methotrexate Sandoz
Inj 20 mg prefilled syringe14.88	1	 Methotrexate Sandoz
Inj 25 mg prefilled syringe14.99	1	 Methotrexate Sandoz
Inj 30 mg prefilled syringe15.09	1	 Methotrexate Sandoz
Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist30.00	5	 Methotrexate DBL Onco-Vial
Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Specialist45.00	1	 DBL Methotrexate Onco-Vial
Ini 100 mg per ml. 10 ml – PCT – Retail pharmacy-Specialist25.00	1	Methotrexate Ebewe
	-	
3 01 7	1	✓ Methotrexate Ebewe
	-	
	•	 Baxter
Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist	5 mg OP	 Baxter
	Tab 10 mg – PCT – Retail pharmacy-Specialist	Tab 10 mg - PCT - Retail pharmacy-Specialist

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PEMETREXED – PCT only – Specialist – Special Authority see	SA1679 below			
Inj 100 mg vial	60.89	1	✓,	Juno Pemetrexed
Inj 500 mg vial		1	✓.	Juno Pemetrexed
Inj 1 mg for ECP	0.55	1 mg	✓	Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

THIOGUANINE - PCT - Retail pharmacy-Specialist

Tab 40 mg	25	 Lanvis
Other Cytotoxic Agents		
AMSACRINE – PCT only – Specialist		
Inj 50 mg per ml, 1.5 ml ampoule1,500.00	6	 Amsidine S29
4,736.00		 Amsidine S29
Inj 75 mg1,250.00	5	AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist		
Cap 0.5 mg1,175.87	100	🗸 Agrylin
ARSENIC TRIOXIDE – PCT only – Specialist		
Inj 1 mg per ml, 10 ml vial4,817.00	10	 Phenasen
Inj 10 mg for ECP	10 mg OP	 Baxter

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	(Manufacturer's Pr \$	rice) Subs Per	sidised	Generic Manufacturer
	φ	FEI		Wallulaciulei
BLEOMYCIN SULPHATE – PCT only – Specialist				
Inj 15,000 iu, vial		1	✓ [DBL Bleomycin Sulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	🗸 E	Baxter
BORTEZOMIB – PCT only – Specialist – Special Authority see	SA1889 below			
Inj 3.5 mg vial	74.93	1	✓ [OBL Bortezomib
	105.00		✓ E	Bortezomib Dr-Reddy's
Inj 1 mg for ECP		1 mg	🗸 E	Baxter
Bortezomib Dr-Reddy's Inj 3.5 mg vial to be delisted 1 May 202	3)	-		
ollowing criteria:	it luttier tenewart	uniess notified	for app	blications meeting tr
ollowing criteria: Either: 1 The patient has symptomatic multiple myeloma; or 2 The patient has symptomatic systemic AL amyloidosis *.	in luriner reneward	uniess notined	for app	plications meeting in
bilowing criteria: ither: 1 The patient has symptomatic multiple myeloma; or 2 The patient has symptomatic systemic AL amyloidosis *. lote: Indications marked with * are unapproved indications. MACARBAZINE – PCT only – Specialist				, , , , , , , , , , , , , , , , , , ,
ollowing criteria: ither: 1 The patient has symptomatic multiple myeloma; or 2 The patient has symptomatic systemic AL amyloidosis *. lote: Indications marked with * are unapproved indications.	72.11	1	✓ [DBL Dacarbazine
ollowing criteria: Either: 1 The patient has symptomatic multiple myeloma; or 2 The patient has symptomatic systemic AL amyloidosis *. Iote: Indications marked with * are unapproved indications. DACARBAZINE – PCT only – Specialist			✓ [, , , , , , , , , , , , , , , , , , ,
bilowing criteria: ither: 1 The patient has symptomatic multiple myeloma; or 2 The patient has symptomatic systemic AL amyloidosis *. lote: Indications marked with * are unapproved indications. MACARBAZINE – PCT only – Specialist		1	✓ [✓ [DBL Dacarbazine Dacarbazine
bilowing criteria: ither: 1 The patient has symptomatic multiple myeloma; or 2 The patient has symptomatic systemic AL amyloidosis *. lote: Indications marked with * are unapproved indications. NACARBAZINE – PCT only – Specialist Inj 200 mg vial Inj 200 mg for ECP		1 10	✓ [✓ [DBL Dacarbazine Dacarbazine APP 529
bilowing criteria: ither: 1 The patient has symptomatic multiple myeloma; or 2 The patient has symptomatic systemic AL amyloidosis *. lote: Indications marked with * are unapproved indications. ACARBAZINE – PCT only – Specialist Inj 200 mg vial Inj 200 mg for ECP ACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist Inj 0.5 mg vial		1 10	✓ [✓ [✓ [DBL Dacarbazine Dacarbazine APP 529 Baxter Cosmegen
billowing criteria: 1 The patient has symptomatic multiple myeloma; or 2 The patient has symptomatic systemic AL amyloidosis *. Iote: Indications marked with * are unapproved indications. DACARBAZINE – PCT only – Specialist Inj 200 mg vial Inj 200 mg for ECP DACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist		1 10 200 mg OP	✓ [✓ [✓ [DBL Dacarbazine Dacarbazine APP 529 Baxter
billowing criteria: ither: 1 The patient has symptomatic multiple myeloma; or 2 The patient has symptomatic systemic AL amyloidosis *. lote: Indications marked with * are unapproved indications. DACARBAZINE – PCT only – Specialist Inj 200 mg for ECP DACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist Inj 0.5 mg vial Inj 0.5 mg for ECP DAUNORUBICIN – PCT only – Specialist		1 10 200 mg OP 1	✓ [✓ [✓ [✓ [✓ [DBL Dacarbazine Dacarbazine APP 529 Baxter Cosmegen Baxter
2 The patient has symptomatic systemic ÅL amyloidosis *. Note: Indications marked with * are unapproved indications. DACARBAZINE – PCT only – Specialist Inj 200 mg vial Inj 200 mg for ECP DACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist Inj 0.5 mg vial		1 10 200 mg OP 1		DBL Dacarbazine Dacarbazine APP 529 Baxter Cosmegen

BAONONOBION TOTONY Opecialist		
Inj 2 mg per ml, 10 ml171.93	1	 Pfizer
Inj 20 mg vial1,495.00	10	 Daunorubicin
		Zentiva S29
Inj 20 mg for ECP171.93	20 mg OP	 Baxter
DOCETAXEL – PCT only – Specialist		
lnj 20 mg48.75	1	 Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial	1	 DBL Docetaxel
Inj 20 mg per ml, 4 ml vial26.95	1	 Docetaxel
		Accord S29
Inj 80 mg195.00	1	 Docetaxel Sandoz
Inj 1 mg for ECP0.65	1 mg	 Baxter
DOXORUBICIN HYDROCHLORIDE – PCT only – Specialist		
Inj 2 mg per ml, 5 ml vial10.00	1	 Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial11.50	1	 Doxorubicin Ebewe
17.00		Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial23.00	1	 Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial65.00	1	Arrow-Doxorubicin
69.99		 Doxorubicin Ebewe
Inj 1 mg for ECP0.35	1 mg	✓ Baxter

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
EPIRUBICIN HYDROCHLORIDE – PCT only – Specialist				
Inj 2 mg per ml, 5 ml vial	25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓	Epirubicin Ebewe
Inj 1 mg for ECP		1 mg	1 🗸	Baxter
ETOPOSIDE				
Cap 50 mg – PCT – Retail pharmacy-Specialist		20	1	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	1	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specia	alist7.90	1	✓	Rex Medical
Inj 1 mg for ECP – PCT only – Specialist	0.09	1 mg	1 🗸	Baxter
ETOPOSIDE PHOSPHATE – PCT only – Specialist				
Inj 100 mg (of etoposide base)		1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg		Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pha	armacy-Specialist	-		
Cap 500 mg		100	1	Devatis
IBRUTINIB – Special Authority see SA2168 below – Retail pha				
Tab 140 mg		30	1	Imbruvica
Tab 420 mg		30	-	Imbruvica
		00	•	

⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and

- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
 - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and

4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or

4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

Inj 5 mg vial – PCT only – Specialist	109.74	1	 Zavedos
Inj 10 mg vial - PCT only - Specialist		1	 Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	 Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
LENALIDOMIDE - Retail pharmacy-Specialist - Special Au	thority see SA2047 below			
Wastage claimable				
Cap 5 mg	5,122.76	28	✓	Revlimid
Cap 10 mg		21	✓	Revlimid
	6,207.00	28	1	Revlimid
Cap 15 mg		21	1	Revlimid
	7.239.18	28	✓	Revlimid
Cap 25 mg	7 607 00	21		Revlimid

SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and

3 Either:

- 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
- 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

Tab 400 mg – PCT – Retail pharmacy-Specialist	50	 Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	50	 Uromitexan
Inj 100 mg per ml, 4 ml ampoule – PCT only – Specialist 177.45	15	 Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	15	 Uromitexan
Inj 1 mg for ECP – PCT only – Specialist2.96	100 mg	 Baxter

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	1	Manufacturer
MITOMYCIN C – PCT only – Specialist				
Inj 5 mg vial	641.70	1	1	Accord S29
Inj 20 mg vial		1	1	Teva
Inj 1 mg for ECP		1 mg	· ·	Baxter
MITOZANTRONE – PCT only – Specialist				
Inj 2 mg per ml, 10 ml vial		1	1	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg		Baxter
DLAPARIB – Retail pharmacy-Specialist – Special Authority s		Ŭ		
Tab 100 mg		56	1	Lynparza
Tab 150 mg		56	1	Lynparza

⇒SA2163 Special Authority for Subsidy

Initial application - (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen: and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment: and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal - (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the followina:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Fither:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment: and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Fither:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or

5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component. **A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL - PCT only - Specialist

Inj 30 mg		5	Paclitaxel Ebewe
Inj 100 mg		1	Paclitaxel Ebewe
	91.67		Paclitaxel Actavis
Inj 150 mg		1	Paclitaxel Ebewe
	137.50		 Anzatax
			Paclitaxel Actavis
Inj 300 mg		1	Paclitaxel Ebewe
	275.00		Anzatax
			Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	 Baxter
PEGASPARGASE - PCT only - Special Authority s	ee SA1979 below		
Inj 750 iu per ml, 5 ml vial		1	 Oncaspar LYO \$29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The patient has newly diagnosed acute lymphoblastic leukaemia; and

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The patient has relapsed acute lymphoblastic leukaemia; and

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist

Inj 10 mg	CBS	1	 Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharr	nacy-Specialist		
Cap 50 mg		50	 Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 on the n	ext page – Retail phar	macy	
Cap 5 mg	9.13	5	 Temaccord
Cap 20 mg		5	 Temaccord
	18.30		Apo-Temozolomide
Cap 100 mg		5	 Temaccord
	40.20		Apo-Temozolomide
Cap 140 mg		5	 Temaccord
Cap 180 mg	620.00	14	Accord S29
Cap 250 mg		5	 Temaccord

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDO	MIDE – Retail	pharmacy-Special	ist – Special	Authority see	: SA1124 (on the next page	Э
•						~~	

Cap 50 mg	 28	 Thalomid
Cap 100 mg	 28	 Thalomid

155

Subsidy	F	ully	Brand or
(Manufacturer's	s Price) Subsidi	sed	Generic
\$	Per	✓	Manufacturer

► SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Fither

1 The notiont has

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

Cap 10 mg – PCT – Retail pharmacy-Specialist	479.50	100	 Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority	see SA1868 belo	w	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OP	 Venclexta
Tab 10 mg		14 OP	 Venclexta
Tab 50 mg	239.44	7 OP	 Venclexta
Tab 100 mg - Wastage claimable	8,209.41	120	 Venclexta

➡SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

	Subsidy		Fully Brand or
	(Manufacturer's P \$	rice) Subs Per	sidised Generic Manufacturer
	Ψ	10	• Manalactarci
INBLASTINE SULPHATE		-	. Haamina
Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-S		5	 ✓ Hospira ✓ Baxter
Inj 1 mg for ECP – PCT only – Specialist		1 mg	• Daxler
INCRISTINE SULPHATE		_	4 - - · · · · · ·
Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Sp	ecialist74.52	5	 DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Sp	ecialist102.73	5	 DBL Vincristine Sulfate
Inj 1 mg for ECP – PCT only – Specialist		1 mg	 Baxter
INORELBINE – PCT only – Specialist		Ũ	
Inj 10 mg per ml, 1 ml vial	12.00	1	Navelbine
	42.00	•	✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial		1	✓ Navelbine
	210.00		 Vinorelbine Ebewe
	328.65		 Sagent S29
Inj 1 mg for ECP		1 mg	✓ Baxter
, , , , , , , , , , , , , , , , , , , ,		•	 Baxter (Sagent)
Inj 50 mg for ECP Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octor	ber 2024)	50 mg OP	
	ber 2024)	50 mg OP	• Barter (Sagent)
Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octol Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octol Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable	ber 2024) ber 2024) ity see SA1870 below	SU mg OP	• Bartel (Sagent)
Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octor Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octor Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable Cap 150 mg	ber 2024) ber 2024) ity see SA1870 below	224	 Alecensa
Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octor Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octor Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable Cap 150 mg	ber 2024) ber 2024) ity see SA1870 below 	224 ecommendation	✓ Alecensa In of a relevant specialist.
 Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octoo Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octoo Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable Cap 150 mg	ber 2024) ber 2024) ity see SA1870 below 7,935.00 al practitioner on the re illowing criteria: table, non-small cell lu	224 ecommendation	 Alecensa n of a relevant specialist. d
 Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octor Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octor Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable Cap 150 mg	ber 2024) ber 2024) ity see SA1870 below 7,935.00 al practitioner on the re illowing criteria: table, non-small cell lu	224 ecommendation	 Alecensa n of a relevant specialist. d
 Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octob Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octob Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable Cap 150 mg	ber 2024) ber 2024) ity see SA1870 below 7,935.00 al practitioner on the re illowing criteria: table, non-small cell lu	224 ecommendation ing cancer; and hase gene rear	 Alecensa n of a relevant specialist. d rrangement using an appropriation
 Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octoo Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octoo Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable Cap 150 mg	ber 2024) ber 2024) ity see SA1870 below 	224 ecommendation ing cancer; and hase gene rear	 Alecensa n of a relevant specialist. d rrangement using an appropriation
 Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octoo Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octoo Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable Cap 150 mg	ber 2024) ber 2024) ity see SA1870 below 7,935.00 al practitioner on the re illowing criteria: table, non-small cell lu has an ALK tyrosine kir oner on the recommen ECIST criteria; and ent.	224 ecommendation ing cancer; and hase gene rear	 Alecensa In of a relevant specialist. d drangement using an appropriation
 Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octoo Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octoo Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable Cap 150 mg	ber 2024) ber 2024) ity see SA1870 below 	224 ecommendation ing cancer; and hase gene rear	 Alecensa n of a relevant specialist. d rrangement using an appropria evant specialist. Approvals va
 Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octoo Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octoo Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Author Wastage claimable Cap 150 mg	ber 2024) ber 2024) ity see SA1870 below 7,935.00 al practitioner on the re- illowing criteria: stable, non-small cell lu- table, non-small cell lu- stable, non-small cell lu- stable, non-small cell lu- table, non-small cell lu- stable, non-small cell lu- stable, non-small cell lu- table, non-small cell lu- stable, non-small cell lu- sta	224 ecommendation ing cancer; and nase gene rear dation of a rele	 Alecensa In of a relevant specialist. d drangement using an appropriation

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Any of the following:

				Subsidy		Fully	Brand or
				(Manufacturer's Pri \$	ce) Pe	Subsidised er ✓	Generic Manufacturer
continued							
1 Both:							
		itient has a dia um dose of 14	• •	eloid leukaemia (CML) in b	last cri	isis or accele	rated phase; and
2 Both:							
		tient has a dia um dose of 14		a chromosome-positive ac	ute lyn	nphoid leukae	emia (Ph+ ALL); and
3 All of	the follo	wing:					
3.2	2 Maxim	tient has a dia um dose of 10 the following:	gnosis of CML in chr 0 mg/day; and	onic phase; and			
	3.3.2	Patient has ex	perienced treatment	• •		•	r treatment with imatinib; or
	3.3.4	Patients is en	olled in the KISS stu	CML defined by the Soka dy** and requires dasatinit	treatr	nent accordir	ng to the study protocol.
	meeting	haematologist the following c		e recommendation of a had	ematol	ogist. Appro	vals valid for 6 months for
	•	nent failure whi	le on dasatinib*; and				
			,	patient is benefiting from ti	eatme	nt; and	
	num das e CML.	atinib dose of	140 mg/day for accel	erated or blast phase CML	and P	h+ ALL, and	100 mg/day for chronic
		ure for CML as alsnz.ac.nz/kis		ia Net Guidelines. **Kinas	e-Inhil	bition Study v	vith Sprycel Start-up
ERLOTINIB	– Retail	pharmacy-Spe	ecialist – Special Auth	nority see SA2115 below			
https://www.o	cancertri	alsnz.ac.nz/kis	s/			,	

Brand switch fee payable (Pharmacode 2651564) - see	e page 249 for details		
Tab 100 mg		30	 Alchemy
Tab 150 mg	569.70	30	 Alchemy

⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

		Subsidy	Fu	
		(Manufacturer's Price) \$	Subsidise Per	ed Generic Manufacturer
		Ŧ	1.01	manulaotaron
GEFITINIB – Retail pharmacy-Sp Tab 250 mg	ecialist – Special Authority see		30 •	Iressa
⇒SA2116 Special Authority fo	r Subsidy			
Initial application only from a rel			nendation of a	a relevant specialist.
Approvals valid for 4 months for a All of the following:	pplications meeting the followir	ng criteria:		
Ũ	ed, or metastatic, unresectable	, non-squamous Non S	Small Cell Lur	ng Cancer (NSCLC); and
2.1 Patient is treatment2.2 Both:	naive; or			
	has discontinued erlotinib due t did not progress whilst on erlot			
3 There is documentation co4 Gefitinib is to be given for a	nfirming that disease expresse a maximum of 3 months.	s activating mutations	of EGFR tyro	sine kinase; and
Renewal only from a relevant spe for 6 months where radiological as Renewal — (pandemic circums) the following criteria:	ssessment (preferably including	g CT scan) indicates N	ISCLC has no	ot progressed.
All of the following:				
2 Gefitinib to be discontinue	1 0			
3 The regular Special Autho	rity renewal requirements cann	ot be met due to COV	ID-19 constra	ints on the health sector.
	atinib mesilate (supplied by No d/or metastatic malignant GIST			
Tab 100 mg [Vabarm] Cr	anial Authority and CA1460			
Tab 100 mg – [Xpharm] – Sp		2 400 00	60 •	Glivec
* Cap 100 mg				Imatinib-Rex
* Cap 400 mg				/ Imatinib-Rex
SA1460 Special Authority fo Special Authority approved by the Notes: Application details may be should be sent to:	r Subsidy CML/GIST Co-ordinator			
The CML/GIST Co-ordinator	Phone: (04) 460 4990			
Pharmac	Facsimile: (04) 916 7571			
PO Box 10 254	Email: cmlgistcoordinator@pl	harmac.govt.nz		
Wellington	······································			
Special Authority criteria for GI Funded for patients:	ST – access by application			
	d by an oncologist) of unresect	table and/or metastation	c malignant ga	astrointestinal stromal tumour
b) Maximum dose of 400 mg/	dav			
	nd subsequent prescriptions ca	n be written by an onc	ologist.	

 d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 on the next page - Retail pharmacy

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
SA2035 Special Authority for Subsidy lenewal — (metastatic breast cancer) only from a relevant spelevant specialist. Approvals valid for 12 months for application Il of the following:				ecommendation of a
 The patient has metastatic breast cancer expressing HEF and The cancer has not progressed at any time point during t Lapatinib not to be given in combination with trastuzumat Lapatinib to be discontinued at disease progression. 	he previous 12 month		0	
ILOTINIB – Special Authority see SA1489 below – Retail phar Wastage claimable				
Cap 150 mg Cap 200 mg		120 120		asigna asigna
 SA1489 Special Authority for Subsidy initial application only from a haematologist. Approvals valid for Il of the following: Patient has a diagnosis of chronic myeloid leukaemia (CI 2 Either: Patient has documented CML treatment failure* w 2.2 Patient has experienced treatment limiting toxicity 	ML) in blast crisis, acc	celerat	ted phase, o	r in chronic phase; and
3 Maximum nilotinib dose of 800 mg/day; and4 Subsidised for use as monotherapy only.		ing iu		ent with inidinity, and
ote: *treatment failure as defined by Leukaemia Net Guideline enewal only from a haematologist. Approvals valid for 6 mont I of the following:		eeting	the following	g criteria:
 Lack of treatment failure while on nilotinib as defined by I Nilotinib treatment remains appropriate and the patient is Maximum nilotinib dose of 800 mg/day; and Subsidised for use as monotherapy only. 				
ALBOCICLIB – Retail pharmacy-Specialist – Special Authority Wastage claimable	v see SA1894 below			
Tab 75 mg	,	21		brance
Tab 100 mg	,	21		brance
Tab 125 mg	4,000.00	21	✓	brance
SA1894 Special Authority for Subsidy tial application only from a medical oncologist or medical pra provals valid for 6 months for applications meeting the followin of the following: 1 Patient has unresectable locally advanced or metastatic I 2 There is documentation confirming disease is hormone-re-	ng criteria: breast cancer; and			Ū
 3 Patient has an ECOG performance score of 0-2; and 4 Either: 				
second or subsequent line setting 4.1 Disease has relapsed or progressed during prior e 4.2 Both:	endocrine therapy; or			

- 4.2 Both:
 - first line setting
 - 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal

|--|

continued...

- state: and
- 4.2.2 Either:

4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or

- 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
- 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	1,334.70	30	 Votrient
Tab 400 mg	2,669.40	30	 Votrient

■SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the followina:

- - 1 The patient has metastatic renal cell carcinoma; and
 - 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
 - 3 The patient has good performance status (WHO/ECOG grade 0-2); and
 - 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
 - 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal: or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
 - 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression: and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

(Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	Generic
RUXOLITINIB – Special Authority see SA1890 below – Retail pha	rmacy			
Wastage claimable				
Tab 5 mg	2,500.00	56	✓	Jakavi
Tab 10mg	5,000.00	56	✓	Jakavi
Tab 15 mg	5,000.00	56	✓	Jakavi
Tab 20 mg	5,000.00	56	1	Jakavi
⇒SA1890 Special Authority for Subsidy				

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the followina:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia mvelofibrosis: and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2117 below - Retail pharmacy

Cap 12.5 mg		28	 Sunitinib Pfizer
Cap 25 mg		28	 Sunitinib Pfizer
Cap 50 mg	694.62	28	 Sunitinib Pfizer

■ SA2117 Special Authority for Subsidy

Initial application - (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval: or

2.4 Both:

2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and

- 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
- 5.2 Haemoglobin level < lower limit of normal; or
- 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
- 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
- 5.5 Karnofsky performance score of less than or equal to 70; or
- 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

	Subsidy (Manufacturer's Price) \$	Subsic Per	Fully dised	Brand or Generic Manufacturer
Endocrine Therapy				
For GnRH ANALOGUES – refer to HORMONE PREPARATIONS	6, Trophic Hormones,	page 84		
ABIRATERONE ACETATE – Retail pharmacy-Specialist – Spec Wastage claimable	ial Authority see SA2	118 below		
Tab 250 mg	4,276.19	120	✓ Zy	/tiga
⇒SA2118 Special Authority for Subsidy				
Initial application only from a medical oncologist, radiation onco a medical oncologist, radiation oncologist or urologist. Approvals All of the following:				
1 Patient has prostate cancer; and				
2 Patient has metastases; and				
3 Patient's disease is castration resistant; and4 Either:				
4.1 All of the following:				
4.1.1 Patient is symptomatic; and				
4.1.2 Patient has disease progression (rising ser 4.1.3 Patient has ECOG performance score of 0-	1; and		ndroge	n therapy; and
4.1.4 Patient has not had prior treatment with tax	ane chemotherapy; o	r		
4.2 All of the following:				
4.2.1 Patient's disease has progressed following4.2.2 Patient has ECOG performance score of 0-4.2.3 Patient has not had prior treatment with abi	2; and	ontaining a	taxane	e; and
Renewal — (abiraterone acetate) only from a medical oncologi		st. urologist	or me	dical practitioner on the
recommendation of a medical oncologist, radiation oncologist or				
the following criteria:	0 11			
All of the following:				
1 Significant decrease in serum PSA from baseline; and				
2 No evidence of clinical disease progression; and				

- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE

Tab 50 mg	4.21	28	 Binarex
FLUTAMIDE			
Tab 250 mg		90	Prostacur S29
·	119.50	100	 Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Authority	y see SA1895 on	the next pag	е
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	 Faslodex

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
 \$	Per	 Manufacturer 	

➡SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

OCTREOTIDE

Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓ Max Health
			✓ Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	 Max Health
			 Octreotide GH S29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	Max Health
OCTREOTIDE LONG-ACTING - Special Authority see SA2119	below - Retail pha	armacy	
Inj depot 10 mg prefilled syringe		1	 Octreotide Depot
			Teva
Inj depot 20 mg prefilled syringe	647.03	1	 Octreotide Depot
			Teva
Inj depot 30 mg prefilled syringe	718.55	1	 Octreotide Depot
			Teva

➡SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

has failed; or

2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or

5 Both:

- 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
- 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

TAMOXIFEN CITRATE

*	Tab 10 mg	5.00	50	 <u>Tamoxifen Sandoz</u>
*	Tab 20 mg6	6.65	60	 <u>Tamoxifen Sandoz</u>

(Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	Generic
Aromatase Inhibitors				
ANASTROZOLE * Tab 1 mg	4.55	30	1	Anatrole
EXEMESTANE * Tab 25 mg	14.50	30	1	Pfizer Exemestane
LETROZOLE * Tab 2.5 mg	5.84	30	1	Letrole
Immunosuppressants				
Cytotoxic Immunosuppressants				
AZATHIOPRINE				
* Tab 25 mg Azamun to be Principal Supply on 1 April 2023	7.36	60	1	Azamun
* Tab 50 mg	8.10	100	1	Azamun
MYCOPHENOLATE MOFETIL				• • •
Tab 500 mg		50		Cellcept
Cap 250 mg Powder for oral lig 1 g per 5 ml – Subsidy by endorsement		100 35 ml C		Cellcept Cellcept
Mycophenolate powder for oral liquid is subsidised only fo the prescription is endorsed accordingly.				
Fusion Proteins				
ETANERCEPT – Special Authority see SA2103 below – Retail pha Inj 25 mg Inj 25 mg autoinjector Inj 50 mg autoinjector Inj 50 mg prefilled syringe	690.00 690.00 1,050.00	4 4 4 4	√ √	Enbrei Enbrei Enbrei Enbrei
SA2103 Special Authority for Subsidy Initial application — (adult-onset Still's disease) only from a rhe	eumatologist App	rovals	valid for 6	months for applications

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either: 1 Both:

- - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

- 18-24 years Male: 7.0 cm; Female: 5.5 cm
- 25-34 years Male: 7.5 cm; Female: 5.5 cm
- 35-44 years Male: 6.5 cm; Female: 4.5 cm
- 45-54 years Male: 6.0 cm; Female: 5.0 cm
- 55-64 years Male: 5.5 cm; Female: 4.0 cm
- 65-74 years Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

	Subsidy	I	ully	Brand or
(N	lanufacturer's Price)	Subsid	ised	Generic
	\$	Per	✓	Manufacturer

continued...

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

1 Both:

- The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Sub	sidy F	ully	Brand or
(Manufactu	irer's Price) Subsid	ised	Generic
\$	B Per	1	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and

1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects; or
- 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Either:

	Subsidy	Fully	Brand or
(Manufa	acturer's Price) S	Subsidised	Generic
	\$ Per	~	Manufacturer

continued...

- 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

- Either:
 - 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
 - 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.
- Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only – Specialist Inj 50 mg per ml, 5 ml	8 5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE – PCT only – Specialist Subsidised only for bladder cancer.		
Inj 2-8 × 100 million CFU	7 1	 OncoTICE
Inj 40 mg per ml, vial176.9		✓ SII-Onco-BCG S29

Monoclonal Antibodies

ADALIMUMAB (AMGEVITA) - Special Authority see SA2178	B below – Retail pharm	nacy	
Inj 20 mg per 0.4 ml prefilled syringe		1	 Amgevita
Inj 40 mg per 0.8 ml prefilled pen		2	 Amgevita
Inj 40 mg per 0.8 ml prefilled syringe		2	 Amgevita

⇒SA2178 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and

2 Either:

- 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
- 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal - (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Patient has pyoderma gangrenosum*; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	~	Manufacturer	

continued...

2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🖌	Manufacturer	

continued...

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of

Subsidy		Fully	Brand or	
(Manufacturer's Price	ce) Si	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and

1.2 Either:

- 1.2.1 Patient has experienced intolerable side effects; or
- 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or

1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or

- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	lbsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per	 Manufactu 	irer

continued...

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Fither:

1 Both:

1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and

1.2 Either:

- 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
- 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
continued treating physician.				
ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - S	pecial Authority see SA2157	below – F	Retail ph	armacy
Inj 20 mg per 0.2 ml prefilled syringe	1,599.96	2	🖌 И	lumira
Inj 40 mg per 0.4 ml prefilled syringe	1,599.96	2	🗸 Н	lumira
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	🗸 Н	lumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	-	2	🗸 Н	lumira

➡SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- treatment; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
 - 2 Patient has received a maximum of 6 months treatment with Amgevita; and
 - 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- $3\;$ Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
 - 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- Both:
 - 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
 - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsic	dised	Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT – Special Authority see SA1772 below – Retail pharmacy

Inj 40 mg per ml, 0.1 ml vial...... 1,250.00 1 🗸 Eylea

■SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application - (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

Fasenra

1

► SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10°9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
 \$	Per	1	Manufacturer

continued...

the first dose to assess response to treatment; and

9 Either:

9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or

9.2 Both:

- 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
- 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and

2 Either:

- 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
- 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

Inj 120 mg per ml casirivimab, 11.1 ml vial (1) and inj 120 mg

per ml imdevimab, 11.1 ml vial (1).....0.00 1 OP 🗸 Ronapreve

⇒SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.
- Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 below

Inj 5 mg per ml, 20 ml vial		1	🖌 Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	🖌 Erbitux
Inj 1 mg for ECP		1 mg	 Baxter

► SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

GEMTUZUMAB OZOGAMICIN - PO	CT only - Specialist - Special Authority s	ee SA2158 on the next page
----------------------------	--	----------------------------

Inj 5 mg vial	12,973.00	1	 Mylotarg
---------------	-----------	---	------------------------------

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2158 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2179 below

Inj 100 mg		 Remicade
Inj 1 mg for ECP	4.40 1 mg	 Baxter

⇒SA2179 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and 2 Fither:

- 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
- 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and

Subsidy	Fully	y Brand or	
(Manufacturer's Price)	Subsidised	d Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application - (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🖌	Manufacturer

continued...

Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and

2.3 Either:

- 2.3.1 There has been an improvement in MRI appearances; or
- 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or

1.2 Both:

- 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation; or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis; or
 - 2.12 Neurosarcoidosis; or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal - (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy	Ful	y Brand or
(Manufacturer's Price)	Subsidise	d Generic
 \$	Per 🖌	Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application - (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician: and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application - (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 The patient has severe Behcet's disease which is significantly impacting the patient's guality of life (see Notes); and

2 Fither

- 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes): or
- 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004:31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.1 Patients SCCAI is greater than or equal to 4; or
- 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 Either:

- 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
- 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment .

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.
- MEPOLIZUMAB Special Authority see SA2154 below Retail pharmacy

Inj 100 mg prefilled pen		1	 Nucala
Inj 100 mg vial	1,638.00	1	 Nucala

⇒SA2154 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5×10^{9} cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and

6 Either:

- 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
- 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

9 Either:

9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or 9.2 Both:

- 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
- 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and

2 Either:

- 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
- 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA2155 below

Inj 25 mg per ml, 40 ml vial	5,910.00	1	🗸 Gazyva
Inj 1 mg for ECP	6.21	1 mg	 Baxter

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 Patient has follicular lymphoma; or
- 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Subsidy		Fully	Brand or	
(Manufacturer's P	rice) Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe		1	🗸 Xolair
Inj 150 mg vial	450.00	1	🗸 Xolair

➡SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and

4 Either:

Subsidy	Ful	ly Brand or
(Manufacturer's	Price) Subsidise	d Generic
\$	Per	Manufacturer

continued...

- 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
- 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient has previously adequately responded* to 6 doses of omalizumab; or

2 Both:

- 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
- 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB – PCT only – Specialist – Special Authority see SA2143 below

Inj 100 mg per ml, 1 ml vial...... 1,700.00 1 🖌 Synagis

(Synagis Inj 100 mg per ml, 1 ml vial to be delisted 1 January 2024)

⇒SA2143 Special Authority for Subsidy

Initial application — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Infant was born in the last 2 years and has severe lung, airway, neurological or neuromuscular disease that requires ongoing, life-sustaining community ventilation; or
- 2 Both:
 - 2.1 Infant was born in the last 12 months; and
 - 2.2 Any of the following:
 - 2.2.1 Patient was born at less than 28 weeks gestation; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient was born at less than 32 weeks gestation; and
 - 2.2.2.2 Either:
 - 2.2.2.2.1 Patient has chronic lung disease; or
 - 2.2.2.2.2 Patient is Māori or any Pacific ethnicity; or
 - 2.2.3 Both:
 - 2.2.3.1 Patient has haemodynamically significant heart disease; and
 - 2.2.3.2 Any of the following:
 - 2.2.3.2.1 Patient has unoperated simple congenital heart disease with significant left to right shunt (see note a); or
 - 2.2.3.2.2 Patient has unoperated or surgically palliated complex congenital heart disease; or
 - 2.2.3.2.3 Patient has severe pulmonary hypertension (see note b); or
 - 2.2.3.2.4 Patient has moderate or severe LV failure (see note c).

Notes:

a) Patient requires/will require heart failure medication, and/or patient has significant pulmonary hypertension, and/or patient

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pric \$	e) Subs Per	Fully sidised	Brand or Generic Manufacturer
ontinued will require surgical palliation/definitive repair within th b) Mean pulmonary artery pressure more than 25 mmH c) LV Ejection Fraction less than 40%.				
Renewal — (RSV prophylaxis for the 2022/2023 RSV sea Approvals valid for 6 months where patient still meets initial of		f COVID-19)	only fro	om a paediatrician.
PERTUZUMAB – PCT only – Specialist – Special Authority Inj 30 mg per ml, 14 ml vial Inj 420 mg for ECP		1 120 mg OP	✓ Pe ✓ Ba	•
SA1606 Special Authority for Subsidy nitial application — (metastatic breast cancer) only from if a relevant specialist. Approvals valid for 12 months for ap II of the following:				on the recommendation
 The patient has metastatic breast cancer expressing and Either: 2.1 Patient is chemotherapy treatment naïve; or 	HER-2 IHC 3+ or ISH+	(including Fl	SH or ot	her current technology
2.2 Patient has not received prior treatment for the 12 months between prior (neo)adjuvant chemo	otherapy treatment and			
 3 The patient has good performance status (ECOG gra 4 Pertuzumab to be administered in combination with tr 5 Pertuzumab maximum first dose of 840 mg, followed 6 Pertuzumab to be discontinued at disease progressic 	astuzumab; and by maximum of 420 mg	gevery 3 we	eks; and	
Renewal — (metastatic breast cancer) only from a relevant specialist. Approvals valid for 12 months for application.			on the re	commendation of a
1 The patient has metastatic breast cancer expressing and		, U		
2 The cancer has not progressed at any time point duri	o 1		i periuzu	map and trastuzumap.
ITUXIMAB (MABTHERA) – PCT only – Specialist – Specia	al Authority see SA1976	2 Delow	🗸 Ma	abthera abthera

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer, and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

6 Either:

- 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
- 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following:
- wrist, elbow, knee, ankle, and either shoulder or hip; and

7 Either:

- 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and

8 Either:

- 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Both:

- 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and

3 Either:

3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Si	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2114 below

Inj 100 mg per 10 ml vial	275.33 2	Riximyo
Inj 500 mg per 50 ml vial	688.20 1	 Riximyo
Inj 1 mg for ECP	1.38 1 mg	 Baxter (Riximyo)

SA2114 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*. Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and

- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and

4 Either:

- 4.1 The patient does not have chromosome 17p deletion CLL; or
- 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2. **Renewal — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	1	Manufacturer

continued...

maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and

2 Either:

2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or

2.2 Both:

2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

- All of the following:
 - 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
 - 2 An initial response lasting at least 12 months was demonstrated; and
 - 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS))

only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only

from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Subsidy	Fu	ly Bra	nd or
(Manufacturer's Price)	Subsidise	d Gei	neric
\$	Per	/ Mai	nufacturer

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or

2 Both:

- 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
- 2.2 To be used for a maximum of 6 treatment cycles.
- Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.
- Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant muccoutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.
- Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia*

Subs (Manufactur		
\$	Per	 Manufacturer

continued...

associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and

Subsidy	Fully	/ Brand or	
(Manufacturer's Price)	Subsidised	d Generic	
\$	Per 🗸	Manufacturer	

continued...

- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.
- Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.
- Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.
- Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🖌	Manufacturer	

continued...

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
- 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

1 Patient was previously treated with rituximab for membranous nephropathy*; and

2 Either:

- 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
- 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has

Subsidy		Fully	Brand or	
(Manufacturer's Price)	:	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

experienced intolerable side effects.

d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.
- Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2084 below - Retail pharmacy

 Cosentyx 	1		1 ml prefilled syringe	Inj 150 mg per ml
 Cosentyx 	2	1,599.00		

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy	Ful	ly Brand or
(Manufacturer's Price)	Subsidise	d Generic
 \$	Per •	Manufacturer

continued...

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

 Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
`\$	Per	1	Manufacturer

continued...

2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 on the next page - Retail pharmacy

Note:	: Siltuximab is to be administered at doses no greater than 1	1 mg/kg every 3 weeks.

Inj 100 mg vial	770.57	1	 Sylvant
Inj 400 mg vial		1	 Sylvant

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.
- Ini 100 mg per ml. 1.5 ml vial with cilgavimab 100 mg per ml,1.5 ml vial.....0.00 1 Evusheld TOCILIZUMAB - PCT only - Special Authority see SA2159 below 1 ✓ Actemra ✓ Actemra S29 S29 ✓ RoActemra S29 S29 ✓ RoActemra S29 S29 880.00 4 Inj 20 mg per ml, 10 ml vial......550.00 ✓ Actemra 1 Actemra S29 S29 ✓ BoActemra S29 S29 Inj 20 mg per ml, 20 ml vial......1,100.00 1 Actemra Actemra S29 S29 ✓ RoActemra S29 S29 4,400.00 ✓ RoActemra S29 S29 1 mg ✓ Baxter

⇒SA2159 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the

(Ma	Subsidy anufacturer's Price)	Subs	Fully sidised	Brand or Generic
·	\$	Per	1	Manufacturer

continued...

following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2 Tocilizumab is to be used as monotherapy; and
 - 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
 - 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
 - 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 6 Either:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
 \$	Per	✓	

continued...

- 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both

- 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

(Manu	Subsidy Ifacturer's Price)	Full Subsidise	
· · · · · · · · · · · · · · · · · · ·	\$ [´] F	Per 🗸	Manufacturer

continued...

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB – PCT	only – Specialist – Special Authority see SA1632 on t	he next page	
Inj 150 mg vial		1	 Herceptin
Inj 440 mg vial		1	 Herceptin
Inj 1 mg for ECP		1 mg	 Baxter

Subs	sidy F	ully B	rand or
(Manufactur	rer's Price) Subsidi	sed G	eneric
\$	Per	✓ M	anufacturer

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria: All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2144 below

Inj 100 mg vial	2,320.00	1	🗸 Kadcyla
Inj 160 mg vial		1	🗸 Kadcyla
Inj 1 mg for ECP		1 mg	 Baxter

⇒SA2144 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and

3 Either:

- 3.1 The patient has received prior therapy for metastatic disease*; or
- 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and

5 Either:

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer	

continued...

- 5.1 Patient does not have symptomatic brain metastases; or
- 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB – Special Authority see SA2182 below – Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe......4,162.00 1 🖌 Stelara

⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

insufficient benefit to meet renewal criteria; or

2.2.2 Both:

2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and

2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or

2 Both:

- 2.1 Patient has active ulcerative colitis; and
- 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or

1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and

2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

⇒SA2183 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
- 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
- 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has active ulcerative colitis; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	:	Subsidised	Generic
\$	Per	~	Manufacturer

continued...

- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20^* ; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Programmed Cell Death-1 (PD-1) Inhibitors

DURVALUMAB – PCT only – Specialist – Special Authority see SA2164 below	DURVALUMAB	- PCT only -	Specialist - Spec	ial Authority se	e SA2164 below
---	------------	--------------	-------------------	------------------	----------------

Inj 50 mg per ml, 10 ml vial	4,700.00	1	🗸 Imfinzi
Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	🗸 Imfinzi
Inj 1 mg for ECP	9.59	1 mg	 Baxter

⇒SA2164 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and

	Subsidy (Manufacturer's Price) \$	Sut Per	Fully osidised	Brand or Generic Manufacturer
continued				
2 Either:				
2.1 Durvalumab is to be used at a maximum dose of n	o greater than 10 mg/	kg every	2 weeks	s; or
2.2 Durvalumab is to be used at a flat dose of 1500 mg	every 4 weeks; and			
3 Treatment with durvalumab to cease upon signs of diseas	e progression; and			
4 Total continuous treatment duration must not exceed 12 m	nonths.			
NIVOLUMAB - PCT only - Specialist - Special Authority see SA	2120 below			
Inj 10 mg per ml, 4 ml vial	1,051.98	1	✓ 0	pdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	√ 0	pdivo
Inj 1 mg for ECP		1 mg	✔ В	axter

⇒SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

- Either:
 - 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB – PCT only – Specialist – Special Authority see SA2121 below

Inj 25 mg per ml, 4 ml vial	4,680.00	1	 Keytruda
Inj 1 mg for ECP		1 mg	 Baxter

⇒SA2121 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and

Subsidy		Fully	Brand or	
(Manufacturer's P	rice) S	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

2.2 Patient has signs of disease progression; and

2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN

Cap 25 mg Cap 50 mg Cap 100 mg		50 50 50	 ✓ Neoral ✓ Neoral ✓ Neoral
Oral liq 100 mg per ml		50 ml OP	 Neoral
EVEROLIMUS – Special Authority see SA2008 below – Retail Wastage claimable	pharmacy		
Tab 10 mg		30	✓ Afinitor
Tab 5 mg	4,555.76	30	 Afinitor

SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

SIROLIMUS - S	pecial Authority see	SA2005 on the	e next page – Retai	l pharmacy
---------------	----------------------	---------------	---------------------	------------

Tab 1 mg		100	 Rapamune
Tab 2 mg	1,499.99	100	 Rapamune
Oral liq 1 mg per ml		60 ml OP	 Rapamune

*Three months or six months, as applicable, dispensed all-at-once

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	
--	--

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- · Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- · Leukoencepthalopathy; or
- Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal --- (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Sut	bsidy	Fully	Brand or
(Manufacti	turer's Price) Subsi	dised	Generic
	\$ Per	~	Manufacturer

continued...

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.
- Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg		 Tacrolimus Sandoz
Cap 0.75 mg	99.30 100	 Tacrolimus Sandoz
Cap 1 mg		 Tacrolimus Sandoz
Cap 5 mg	248.20 50	 Tacrolimus Sandoz

➡SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – Special Authority see SA2079 on the next page – Retail pharmacy Tab 15 mg1,271.00 28

✓ RINVOQ

Subsidy (Manufacturer's Price) \$) Subsi Per	Fully idised	Brand or Generic Manufacturer	
---	----------------	-----------------	-------------------------------------	--

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and

3 Either:

- 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

	Subsidy (Manufacturer's Price) \$		Fully Brand or lised Generic Manufacturer	
Antiallergy Preparations				
Allergic Emergencies				
 ADRENALINE – Special Authority see SA2185 below – Retail pl a) Maximum of 2 inj per prescription b) Additional prescriptions limited to replacement of up to tw treatment of anaphylaxis. 	vo devices prior to exp	biry, or repla	cement of used device for	r
Inj 0.15 mg per 0.3 ml auto-injector Inj 0.3 mg per 0.3 ml auto-injector		1 OP 1 OP	 Epipen Jr Epipen 	
Initial application — (anaphylaxis) from any relevant practitior applications meeting the following criteria: Both:	ner. Approvals valid v	vithout furth	er renewal unless notified	for
 Either: 1 Either: 1.1 Patient has experienced an anaphylactic reaction department; or 1.2 Patient has been assessed to be at significant risk 2 Patient is not to be prescribed more than two devices in ir 	of anaphylaxis by a r		, ,	су
ICATIBANT – Special Authority see SA1558 below – Retail phar Inj 10 mg per ml, 3 ml prefilled syringe	rmacy	1	✓ Firazyr	
SA1558 Special Authority for Subsidy Initial application only from a clinical immunologist or relevant s the following criteria: Both:	pecialist. Approvals	valid for 12	months for applications m	eeting
1 Supply for anticipated emergency treatment of laryngeal/	pro-pharyngeal or sev	ere abdomi	nal attacks of acute hered	litary

- angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

■ SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy

Initiation kit - 5 vials freeze dried venom with diluent	5.00	1 OP 🗸	VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	5.00	1 OP 🖌	VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with			
diluent28	35.00	1 OP 🖌 🖌	Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml30	5.00	1 OP 🖌 🖌	Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent 30	5.00	1 OP 🖌	Hymenoptera S29

	Subsidy		Fully Brand or
	(Manufacturer's Pric	e) Subs	sidised Generic
	\$	Per	 Manufacturer
NASP VENOM ALLERGY TREATMENT – Special Authority se	e SA1367 on the p	revious page	- Retail pharmacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			1 5
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml		1 OP	 Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent		1 OP	 Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent		1 OP	 Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze	9		
dried venom, with diluent		1 OP	 Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze			
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml		1 OP	 Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freez	e		
dried venom, with diluent		1 OP	 Venomil S29
		_	
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	7 50	100	 Zista
* Oral lig 1 mg per ml		200 ml	✓ Histaclear
		200	
Oral liq 2 mg per 5 ml	9.37	500 ml	 Histafen
		500 m	• mataien
	0.00	40	
* Tab 2 mg		40	Polaramine
	(8.40) 1.01	20	Foldidillille
	(5.99)	20	Polaramine
* Oral liq 2 mg per 5 ml		100 ml	rolaramine
	(10.29)	100 111	Polaramine
EXOFENADINE HYDROCHLORIDE	()		
* Tab 60 mg	4 34	20	
	(8.23)	20	Telfast
* Tab 120 mg	()	10	· ondot
	(8.23)		Telfast
	14.22	30	
	(26.44)		Telfast
ORATADINE			
* Tab 10 mg	1.78	100	 Lorafix
* Oral liq 1 mg per ml		100 ml	✓ Haylor syrup
* Tab 10 mg		50	 Allersoothe
* Tab 25 mg		50	✓ Allersoothe
* Oral liq 1 mg per 1 ml		100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a		5	 Hospira

	Subsidy		Fully	Brand or
	(Manufacturer's I	Price) S	ubsidised	Generic
	\$	Per	1	Manufacturer
Inhaled Corticosteroids				
Innaled Controosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose		200 dose C)P 🗸	Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose C	P 🗸	Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose C	P 🗸	Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose C	P 🗸	Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free		200 dose C	P 🗸	Beclazone 250
BUDESONIDE				
	17.00	000 dasa (Pulmicort
Powder for inhalation, 100 mcg per dose	17.00	200 dose C	•	
				Turbuhaler
Powder for inhalation, 200 mcg per dose		200 dose C	P ✓	Pulmicort
				Turbuhaler
Powder for inhalation, 400 mcg per dose		200 dose C	P 🗸	Pulmicort
				Turbuhaler
FLUTICASONE				
Aerosol inhaler, 50 mcg per dose	7 19	120 dose C	P 🗸	Flixotide
Powder for inhalation, 50 mcg per dose		60 dose O		Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose O		Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose C		Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose C		Flixotide
Powder for inhalation, 250 mcg per dose		60 dose O		Flixotide Accuhaler
Powder for initialation, 250 mcg per dose		00 00se 0	г •	Flixolide Accultater
Inhaled Long-acting Beta-adrenoceptor Agonis	ts			
EFORMOTEROL FUMARATE				
Powder for inhalation, 12 mcg per dose, and monodose devi		60 dose		
	(35.80)			Foradil
(Foradil Powder for inhalation, 12 mcg per dose, and monodose	device to be del	isted 1 July 2	2023)	
EFORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated				
(equivalent to eformoterol fumarate 6 mcg metered dose) 10.32	60 dose O	P	
(equivalent to clothioteron famarate of meg meterod dost	(16.90)	00 0030 0		Oxis Turbuhaler
	(10.50)			
INDACATEROL				.
Powder for inhalation 150 mcg		30 dose O		Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose O	Р 🗸	Onbrez Breezhaler
SALMETEROL				
Aerosol inhaler CFC-free, 25 mcg per dose		120 dose C	P 🗸	Serevent
Powder for inhalation, 50 mcg per dose, breath activated		60 dose O	Р 🗸	Serevent Accuhaler

	Subsidy		Fully	Brand or
	(Manufacturer's \$	Price) Subsic Per		Generic Manufacturer
Inhaled Corticosteroids with Long-Acting Beta-A	drenocept	or Agonists		
BUDESONIDE WITH EFORMOTEROL				
Powder for inhalation 160 mcg with 4.5 mcg eformoterol				
fumarate per dose (equivalent to 200 mcg budesonide wit 6 mcg eformoterol fumarate metered dose)		120 dose OP	1	DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumara		120 0036 01	•	Duonesp Spirollax
per dose (equivalent to 400 mcg budesonide with 12 mcg				
eformoterol fumarate metered dose) - No more than 2	00.50	100 days 00		D
dose per day Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP 120 dose OP		DuoResp Spiromax Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg		120 dose OP		Symbicort
-	-			Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP		Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 mc	cg33.74	120 dose OP	~	Symbicort Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate				
12 mcg – No more than 2 dose per day		60 dose OP	✓	Symbicort
				Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL				
Powder for inhalation 100 mcg with vilanterol 25 mcg		30 dose OP	~	Breo Ellipta
FLUTICASONE WITH SALMETEROL Aerosol inhaler 50 mcg with salmeterol 25 mcg	25 70	120 dose OP	1	Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP		Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg – No				
more than 2 dose per day		60 dose OP	~	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No	44.00	60 dose OP		Seretide Accuhaler
more than 2 dose per day		00 dose OF	•	Serelide Accurater
Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Oral liq 400 mcg per ml Infusion 1 mg per ml, 5 ml		150 ml 10		Ventolin Ventolin
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO		5		Ventolin
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO	2 20	200 dose OP	1	Respigen
		200 0036 01		SalAir
	(6.20)			Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb		00		A
available on a PSO Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb	8.96	20	•	<u>Asthalin</u>
available on a PSO	9.43	20	✓	Asthalin
TERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to				
250 mcg metered dose), breath activated		120 dose OP	✓	Bricanyl Turbuhaler

	Subsidy (Manufacturer's	Price)	Fully Subsidised	Brand or Generic
	\$	Per	1	Manufacturer
Anticholinergic Agents				
PRATROPIUM BROMIDE Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose available on a PSO Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne	16.20	200 dose	OP 🗸 A	Atrovent
available on a PSO		20		Inivent Accord 529
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	Agents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p dose CFC-free Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO	12.19	200 dose 20		Duolin HFA Duolin
Long-Acting Muscarinic Antagonists				
 GLYCOPYRRONIUM – Subsidy by endorsement a) Inhaled glycopyrronium treatment will not be subsidised if umeclidinium. b) Glycopyrronium powder for inhalation 50 mcg per dose is having COPD using spirometry if spirometry is possible, a Powder for inhalation 50 mcg per dose 	subsidised only	y for patients	s who have rsed accore	been diagnosed as
 TIOTROPIUM BROMIDE – Subsidy by endorsement a) Tiotropium treatment will not be subsidised if patient is als umeclidinium. b) Tiotropium bromide is subsidised only for patients who ha spirometry is possible, and the prescription is endorsed at 1 October 2018 with a valid Special Authority are deemed Powder for inhalation, 18 mcg per dose	ve been diagno ccordingly. Pat l endorsed. 	osed as havi	ng COPD (ad tiotropiu	using spirometry if
 JMECLIDINIUM – Subsidy by endorsement a) Umeclidinium will not be subsidised if patient is also recei tiotropium bromide. b) Umeclidinium powder for inhalation 62.5 mcg per dose is COPD using spirometry if spirometry is possible, and the Powder for inhalation 62.5 mcg per dose. 	ving treatment subsidised only prescription is e	with subsidis	sed inhaled who have cordingly.	l glycopyrronium or
Long-Acting Muscarinic Antagonists with Long-	Acting Beta	a-Adreno	ceptor A	Igonists
Combination long acting muscarinic antagonist and long acting be reatment with a combination inhaled corticosteroid and long actir			osidised if p	patient is also receiving

► SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

Sub		-ully ised	Brand or Generic
	\$ Per	✓	Manufacturer

continued...

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL – Special Authority s Powder for Inhalation 50 mcg with indacaterol 110 mcg		ge – Retail pharmacy ✓ Ultibro Breezhaler
TIOTROPIUM BROMIDE WITH OLODATEROL – Special Authorit Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg		
UMECLIDINIUM WITH VILANTEROL – Special Authority see SA1		etail pharmacy

Antifibrotics

NINTEDANIB - Special Authority see SA2012 below -	- Retail pharmacy		
Note: Nintedanib not subsidised in combination w	ith subsidised pirfenidone.		
Cap 100 mg	2,554.00	60 OP	 Ofev
Cap 150 mg		60 OP	 Ofev

SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 on the next page

Note: Pirfenidone is not subsidised in combination w	vith subsidised nintedanib.		
Tab 801 mg		90	 Esbriet
Tab 267 mg	1,215.00	90	 Esbriet

28

Montelukast Mylan

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	1	Manufacturer	

► SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists		
MONTELUKAST * Tab 4 mg * Tab 5 mg	28 28	 <u>Montelukast Mylan</u> <u>Montelukast Mylan</u> Montelukast Viatris

			 Montelukast Viatris
Methylxanthines			
AMINOPHYLLINE			
* Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a			
PSO	180.00	5	 DBL Aminophylline
THEOPHYLLINE			
* Tab long-acting 250 mg	23.94	100	 Nuelin-SR
* Oral liq 80 mg per 15 ml	17.62	500 ml	 Nuelin

Mucolytics

DORNASE ALFA - Special Authority see SA1978 below - Retail p			
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	 Pulmozyme
SA1978 Special Authority for Subsidy			

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

IVACAFTOR - PCT only - Specialist - Special Authority	see SA2017 below		
Tab 150 mg		56	 Kalydeco
Oral granules 50 mg, sachet		56	 Kalydeco
Oral granules 75 mg, sachet		56	 Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

Not funded for use as a nasal drop. Soln 7%24.50	90 ml OP	✓ Biomed
Nasal Preparations		
Allergy Prophylactics		
BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose	200 dose OP 200 dose OP	✓ <u>SteroClear</u> ✓ <u>SteroClear</u>
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose1.98	120 dose OP	✓ <u>Flixonase Hayfever</u> <u>& Allergy</u>
IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%5.23	15 ml OP	✓ <u>Univent</u>

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	Brand or Generic Manufacturer
Respiratory Devices				
MASK FOR SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
 c) Only for children aged six years and under Small 		1	√ e	e-chamber Mask
PEAK FLOW METER				
a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range	9.54	1	✓ I	Mini-Wright AFS Low Range
Normal range	9.54	1	✓ N	Mini-Wright Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)		1	-	e-chamber Turbo
510 ml (single patient)	5.95	1		e-chamber La Grande
800 ml	6.50	1	۷ ۱	/olumatic
Respiratory Stimulants				
Hoophatory otimulanto				
CAFFEINE CITRATE				
Oral liq 20 mg per ml (10 mg base per ml)		25 ml (OP 🖌 E	Biomed

	Subsidy		Fully Brand or	
	(Manufacturer's P		sidised Generic	
	\$	Per	 Manufacturer 	
Ear Preparations				
FLUMETASONE PIVALATE				
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	 Locacorten-Viaform ED's 	
			✓ Locorten-Vioform	
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	IN		
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			.	
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	 Kenacomb 	
Ear/Eye Preparations				
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN				
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and				
gramicidin 50 mcg per ml	4 50	8 ml OP		
		0 IIII OF	Sofradex	
	(9.27)		Sunduex	
FRAMYCETIN SULPHATE	4.40			
Ear/Eye drops 0.5%		8 ml OP	0	
	(8.65)		Soframycin	
Eye Preparations				
Eye preparations are only funded for use in the eye, unless explicit	citly stated otherv	vise.		
Anti-Infective Preparations				
ACICLOVIR				
* Eye oint 3%	14.88	4.5 g OP	 ViruPOS 	
CHLORAMPHENICOL		•		
Eve oint 1%	1.09	5 g OP	 Devatis 	
Eye drops 0.5%		10 ml OP	✓ Chlorafast	
Funded for use in the ear*. Indications marked with * ar			••••••	
CIPROFLOXACIN	e anappierea ne			
Eye drops 0.3% – Subsidy by endorsement	0.72	5 ml OP	 Ciprofloxacin Teva 	
When prescribed for the treatment of bacterial keratitis of				· or
for the second line treatment of chronic suppurative otitis		,		·
Note: Indication marked with a * is an unapproved indic		, and the pies		gıy.
GENTAMICIN SULPHATE	11 40		- Conontin	
Eye drops 0.3%	11.40	5 ml OP	 Genoptic 	
(Genoptic Eye drops 0.3% to be delisted 1 August 2023)				
PROPAMIDINE ISETHIONATE				
* Eye drops 0.1%		10 ml OP		
	(14.55)		Brolene	
SODIUM FUSIDATE [FUSIDIC ACID]				
Eye drops 1%	5.29	5 g OP	 Fucithalmic 	
TOBRAMYCIN				
Eye oint 0.3%		3.5 g OP	 Tobrex 	
Eye drops 0.3%		5 ml OP	✓ Tobrex	
	-			

	Subsidy (Manufacturer's Pr		Fully Brand or sidised Generic
	(Manulacturer's Pr	Per Sub	Manufacturer
Corticosteroids and Other Anti-Inflammatory	Preparations		
DEXAMETHASONE			
* Eye oint 0.1%		3.5 g OP	 Maxidex
* Eye drops 0.1%		5 ml OP	 Maxidex
Ocular implant 700 mcg – Special Authority see SA1680 – Retail pharmacy		1	✓ Ozurdex
SA1680 Special Authority for Subsidy			
Initial application — (Diabetic macular oedema) only from	an ophthalmologist.	Approvals va	alid for 12 months for applications
meeting the following criteria:			
All of the following:			
1 Patient has diabetic macular oedema with pseudopha			and the state of the second
2 Patient has reduced visual acuity of between 6/9 - 6/43 Either:			uction in vision; and
3.1 Patient's disease has progressed despite 3 inje3.2 Patient is unsuitable or contraindicated to treat			
4 Dexamethasone implants are to be administered not r maximum of 3 implants per eye per year.	nore frequently than o	once every 4	months into each eye, and up to a
Renewal - (Diabetic macular oedema) only from an ophth	almologist. Approva	s valid for 12	months for applications meeting
the following criteria: Both:			
1 Patient's vision is stable or has improved (prescriber d	etermined); and		
2 Dexamethasone implants are to be administered not r	nore frequently than o	once every 4	months into each eye, and up to a
maximum of 3 implants per eye per year.			
Initial application - (Women of child bearing age with dia	abetic macular oede	ma) only from	m an ophthalmologist. Approvals
valid for 12 months for applications meeting the following crite	eria:		
All of the following:			
1 Patient has diabetic macular oedema; and			
2 Patient has reduced visual acuity of between 6/9 - 6/4			uction in vision; and
3 Patient is of child bearing potential and has not yet co			
4 Dexamethasone implants are to be administered not r maximum of 2 implants par are per year	nore frequently than o	once every 4	months into each eye, and up to a
maximum of 3 implants per eye per year.		furne en enki	
Renewal — (Women of child bearing age with diabetic mathematications meeting the following criteria:	acular oedema) only	from an opn	thaimologist. Approvals valid for
All of the following:			
1 Patient's vision is stable or has improved (prescriber d	atorminad); and		
2 Patient is of child bearing potential and has not yet co	<i>,</i> .		
3 Dexamethasone implants are to be administered not r			months into each eve, and up to a
maximum of 3 implants per eye per year.			
DEXAMETHASONE WITH NEOMYCIN SULPHATE AND PC		ATE	
* Eye oint 0.1% with neomycin sulphate 0.35% and polymy			
sulphate 6,000 u per g		3.5 g OP	 Maxitrol
 Eye drops 0.1% with neomycin sulphate 0.35% and poly. 		0.0 9 01	
b sulphate 6,000 u per ml		5 ml OP	 Maxitrol
DICLOFENAC SODIUM			
Eye drops 0.1%	8 80	5 ml OP	 Voltaren Ophtha
	0.00		
FLUOROMETHOLONE * Eye drops 0.1%	2 00	5 ml OP	✓ FML
- Eye ulups 0.1%	3.U9 5.00	5 III OP	✓ FML

*Three months or six months, as applicable, dispensed all-at-once

5.20

✓ Flucon

SENSORY ORGANS

SENSORY ORGANS

(M	Subsidy anufacturer's P \$	Price) Subs Per	Fully sidised	Brand or Generic Manufacturer
EVOCABASTINE	Ŷ	101		Manufacturor
Eve drops 0.5 mg per ml	8 71	4 ml OP		
	(10.34)	4 111 01	L	_ivostin
ODOXAMIDE	()			
Eve drops 0.1%	8.71	10 ml OP	✓ 1	Lomide
PREDNISOLONE ACETATE				
Eve drops 1%	6.92	10 ml OP	1	Prednisolone-AFT
	7.00	5 ml OP	 I 	Pred Forte
REDNISOLONE SODIUM PHOSPHATE – Special Authority see S	A1715 below	– Retail pharr	nacv	
Eye drops 0.5%, single dose (preservative free)		20 dose		Vinims
, , , , , , , , , , , , , , , , , , ,				Prednisolone
itial application only from an ophthalmologist or optometrist. App ollowing criteria:	rovals valid f	or 6 months for	r applic	ations meeting the
 SA1715 Special Authority for Subsidy nitial application only from an ophthalmologist or optometrist. App ollowing criteria: Both: Patient has severe inflammation; and Patient has a confirmed allergic reaction to preservative in ey Renewal from any relevant practitioner. Approvals valid for 6 month benefiting from treatment. SODIUM CROMOGLICATE Eye drops 2% 	e drops. s where the t		ins ap	, , , , , , , , , , , , , , , , , , ,
nitial application only from an ophthalmologist or optometrist. App ollowing criteria: Both: 1 Patient has severe inflammation; and 2 Patient has a confirmed allergic reaction to preservative in ey Renewal from any relevant practitioner. Approvals valid for 6 month benefiting from treatment. SODIUM CROMOGLICATE	e drops. s where the t	treatment rema	ins ap	propriate and the patient
nitial application only from an ophthalmologist or optometrist. Applolowing criteria: Both: 1 Patient has severe inflammation; and 2 Patient has a confirmed allergic reaction to preservative in ey Renewal from any relevant practitioner. Approvals valid for 6 month SODIUM CROMOGLICATE Eye drops 2%	e drops. s where the t	treatment rema	ins ap	propriate and the patient
nitial application only from an ophthalmologist or optometrist. App ollowing criteria: Both: 1 Patient has severe inflammation; and 2 Patient has a confirmed allergic reaction to preservative in ey Renewal from any relevant practitioner. Approvals valid for 6 month enefiting from treatment. SODIUM CROMOGLICATE Eye drops 2% Glaucoma Preparations - Beta Blockers BETAXOLOL k Eye drops 0.25%	e drops. s where the t 2.62	treatment rema 10 ml OP 5 ml OP	iins apj ✓ <u>j</u>	propriate and the patient <u>Allerfix</u> Betoptic S
nitial application only from an ophthalmologist or optometrist. App ollowing criteria: Both: 1 Patient has severe inflammation; and 2 Patient has a confirmed allergic reaction to preservative in ey Renewal from any relevant practitioner. Approvals valid for 6 month enefiting from treatment. SODIUM CROMOGLICATE Eye drops 2% Glaucoma Preparations - Beta Blockers BETAXOLOL k Eye drops 0.25%	e drops. s where the t 2.62	treatment rema 10 ml OP	iins apj ✓ <u>j</u>	propriate and the patient Allerfix
nitial application only from an ophthalmologist or optometrist. App ollowing criteria: Both: 1 Patient has severe inflammation; and 2 Patient has a confirmed allergic reaction to preservative in ey Renewal from any relevant practitioner. Approvals valid for 6 month benefiting from treatment. SODIUM CROMOGLICATE Eye drops 2% BETAXOLOL k Eye drops 0.25% k Eye drops 0.5%	e drops. s where the t 2.62	treatment rema 10 ml OP 5 ml OP	iins apj ✓ <u>j</u>	propriate and the patient <u>Allerfix</u> Betoptic S
nitial application only from an ophthalmologist or optometrist. App ollowing criteria: Both: 1 Patient has severe inflammation; and 2 Patient has a confirmed allergic reaction to preservative in ey Renewal from any relevant practitioner. Approvals valid for 6 month enefiting from treatment. SODIUM CROMOGLICATE Eye drops 2% Glaucoma Preparations - Beta Blockers BETAXOLOL k Eye drops 0.25% MOLOL k Eye drops 0.25%	e drops. s where the t 2.62 11.80 7.50 1.81	treatment rema 10 ml OP 5 ml OP 5 ml OP 5 ml OP 5 ml OP	iins apj ✓ <u>j</u> ✓ I	propriate and the patient <u>Allerfix</u> Betoptic S Betoptic Arrow-Timolol
nitial application only from an ophthalmologist or optometrist. Application only from an ophthalmologist or optometrist. Application of the second structure is a confirmed allergic reaction to preservative in ey Renewal from any relevant practitioner. Approvals valid for 6 monthologist from treatment. SODIUM CROMOGLICATE Eye drops 2% Glaucoma Preparations - Beta Blockers BETAXOLOL	e drops. s where the t 2.62 11.80 7.50 1.81 2.04	treatment rema 10 ml OP 5 ml OP 5 ml OP 5 ml OP	iins app	propriate and the patient <u>Allerfix</u> Betoptic S Betoptic

(Timoptol XE Eye drops 0.5%, gel forming to be delisted 1 March 2024)

Glaucoma Preparations - Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE * Tab 250 mg	100	 Diamox
BRINZOLAMIDE * Eye drops 1%	5 ml OP	✓ <u>Azopt</u>
DORZOLAMIDE HYDROCHLORIDE * Eye drops 2%	5 ml OP	Trusopt
(Trusopt Eye drops 2% to be delisted 1 March 2024) DORZOLAMIDE WITH TIMOLOL		
 * Eye drops 2% with timolol 0.5%2.73 Glaucoma Preparations - Prostaglandin Analogues 	5 ml OP	✓ Dortimopt
and contain operations in rootagianam / indioguos		
BIMATOPROST * Eye drops 0.03%5.95	3 ml OP	✓ <u>Bimatoprost</u> <u>Multichem</u>

246 ✓ fully subsidised Principal Supply Sole Subsidised Supply

SENSORY ORGANS

	Subsidy		Fully	Brand or
(Ν	/lanufacturer's F		ubsidised	Generic
	\$	Per		Manufacturer
ATANOPROST				
* Eye drops 0.005%		2.5 ml OF	, √ ⊺	eva
TRAVOPROST			_	
✤ Eye drops 0.004%	9 75	2.5 ml OF	, 🖌 т	ravatan
		2.0 111 01	• 1	Tavatan
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE				
Eye drops 0.2%	4.29	5 ml OP	✓ <u>A</u>	rrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE				
Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ C	ombigan
ATANOPROST WITH TIMOLOL				-
Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OF	> ✓ A	rrow - Lattim
PILOCARPINE HYDROCHLORIDE			-	
	4.26	15 ml OP	1	opto Carpine
 ₭ Eye drops 1% ₭ Eye drops 2% 		15 ml OP		sopto Carpine
 ▲ Eye drops 2 %		15 ml OP		sopto Carpine
Subsidised for oral use pursuant to the Standard Formulae		13 111 01	• 18	sopto carpine
✤ Eye drops 2% single dose – Special Authority see SA0895				
below – Retail pharmacy	31.05	20 dose	🗸 N	linims Pilocarpine
		20 0050	• 10	ininina Filocalpine
SA0895 Special Authority for Subsidy				

SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

A I ROPINE SULPHATE * Eye drops 1%	.17.36	15 ml OP	✓ <u>Atropt</u>
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%	8.76	15 ml OP	 Cyclogyl
TROPICAMIDE * Eye drops 0.5% * Eye drops 1%		15 ml OP 15 ml OP	 ✓ Mydriacyl ✓ Mydriacyl

Preparations for Tear Deficiency

For acetylcysteine eye drops refer Standard Formulae, page 251

HYPROMELLOSE			
* Eye drops 0.5%	19.50	15 ml OP	 Methopt
HYPROMELLOSE WITH DEXTRAN			
* Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	 Poly-Tears

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Preservative Free Ocular Lubricants				
SA2134 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals va toth:	alid for 12 months for	applica	itions meetin	g the following criteria:
 Confirmed diagnosis by slit lamp or Schirmer test of sev Either: 	vere secretory dry eye	; and		
2.1 Patient is using eye drops more than four times of2.2 Patient has had a confirmed allergic reaction to patient of				
tenewal from any relevant practitioner. Approvals valid for 24 rops and has benefited from treatment.	months where the pa	atient c	ontinues to r	equire lubricating eye
ARBOMER – Special Authority see SA2134 above – Retail p Ophthalmic gel 0.3%, 0.5 g		30	√ P	oly-Gel
OLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml Systane Unit Dose Eye drops 0.4% and propylene glycol 0.3%	4.30 10.78	24 30	✓ S ✓ S	Retail pharmacy ystane Unit Dose ystane Unit Dose
ODIUM HYALURONATE [HYALURONIC ACID] – Special Au Eye drops 1 mg per ml Hylo-Fresh has a 6 month expiry after opening. The F month is not relevant and therefore only the prescribed		10 ml (Manua	DP	ylo-Fresh allowing one bottle per
Other Eye Preparations				
IAPHAZOLINE HYDROCHLORIDE ≰ Eye drops 0.1%		15 ml (OP 🗸 N	aphcon Forte
Eye drops 0.1% Eye drops 0.1% ARAFFIN LIQUID WITH WOOL FAT	2.17	5 ml C	P ✓ <u>C</u>	lopatadine Teva
Eye oint 3% with wool fat 3% ETINOL PALMITATE	3.63	3.5 g C)P 🖌 P	oly-Visc
ETINOL PALMITATE Eye oint 138 mcg per g	3.80	5 g O	P 🗸 V	itA-POS

VARIOUS

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Various	Ψ	1 61		Manufacturer
PHARMACY SERVICES May only be claimed once per patient.	4.50	1 fee	 Image: A second s	BSF Alchemy BSF Celapram BSF Ticagrelor
a) The Dharmanado for DSE Alabamy is 2651564 and				Sandoz BSF Vebulis
 a) The Pharmacode for BSF Alchemy is 2651564 - see is b) The Pharmacode for BSF Ticagrelor Sandoz is 26532 c) The Pharmacode for BSF Vebulis is 2653214 - see aid) The Pharmacode for BSF Celapram is 2653222 - see (BSF Alchemy Brand switch fee to be delisted 1 May 2023) (BSF Celapram Brand switch fee to be delisted 1 June 2023) (BSF Ticagrelor Sandoz Brand switch fee to be delisted 1 June 2023) (BSF Vebulis Brand switch fee to be delisted 1 June 2023) 	206 - see also page lso page 61 e also page 124	41		
Agents Used in the Treatment of Poisonings				
Antidotes				
ACETYLCYSTEINE Inj 200 mg per ml, 10 ml ampoule NALOXONE HYDROCHLORIDE a) Up to 10 inj available on a PSO	52.88	10	√ į	Martindale Pharma
 b) Only on a PSO * Inj 400 mcg per ml, 1 ml ampoule 	35.26	10	√ !	Hameln
Removal and Elimination				
CHARCOAL * Oral liq 50 g per 250 ml a) Up to 250 ml available on a PSO b) Only on a PSO	43.50 2	50 ml (OP 🗸	Carbosorb-X
DEFERASIROX – Special Authority see SA1492 below – Retail p Wastage claimable	oharmacy			
Tab 125 mg dispersible Tab 250 mg dispersible Tab 500 mg dispersible	552.00	28 28 28	 Image: A second s	Exjade Exjade Exjade
SA1492 Special Authority for Subsidy Initial application only from a haematologist. Approvals valid for All of the following:	r 2 years for applica	tions m	neeting the	following criteria:

1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and

- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or

Subsidy	Fi	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
- 3.3 Treatment with deferiprone has resulted in arthritis; or
- 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

		DEFERIPRONE – Special Authority see SA1480 below – Retail pharmacy
 Ferriprox 	100	Tab 500 mg533.17
 Ferriprox 	250 ml OP	Oral lig 100 mg per 1 ml

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

I

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

*	Inj 500 mg vial	151.31	10	✓ DBL Desferrioxamine Mesylate for Inj BP
				 Deferoxamine Pfizer S29 S29
SOD	DIUM CALCIUM EDETATE			
*	Inj 200 mg per ml, 5 ml	53.31	6	
		(156.71)		Calcium Disodium Versenate

Standard Formulae

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
Suitable eye drop base	qs	Phenobarbitone Sodium Glycerol BP	400 mg 4 ml
CODEINE LINCTUS (3 mg per 5 ml)		Water	to 40 ml
Codeine phosphate	60 mg		
Glycerol	40 ml	PILOCARPINE ORAL LIQUID	
Preservative	qs	Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative Water	qs to 500 ml
CODEINE LINCTUS (15 mg per 5 ml)		(Preservative should be used if quantity supplied is	
Codeine phosphate	300 mg	than 5 days.)	
Glycerol	40 ml	. ,	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	_
Water	to 100 ml	Methylcellulose	5 g
FOLINIC MOUTHWASH		Preservative Water	qs to 500 ml
Calcium folinate 15 mg tab	1 tab	(Preservative should be used if quantity supplied is	
Preservative	qs	than 5 days. Maximum 500 ml per prescription.)	
Water	to 500 ml	, , ,	
(Preservative should be used if quantity supplied is	for more	SODIUM CHLORIDE ORAL LIQUID	
than 5 days. Maximum 500 ml per prescription.)		Sodium chloride inj 23.4%, 20 ml Water	qs
METHADONE MIXTURE		(Only funded if prescribed for treatment of hyponatr	qs aemia)
Methadone powder	qs		aonnay
Glycerol	qs	VANCOMYCIN ORAL SOLUTION (25 mg per ml)	
Water	to 100 ml	Vancomycin 500 mg injection	5 vials
METHYL HYDROXYBENZOATE 10% SOLUTION		Glycerin with sucrose suspension Water	37.5 ml to 100 ml
Methyl hydroxybenzoate	10 g	(Only funded if prescribed for treatment of Clostridiu	
Propylene glycol	to 100 ml	following metronidazole failure)	
(Use 1 ml of the 10% solution per 100 ml of oral liqu		-	
	,	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	40.11
OMEPRAZOLE SUSPENSION	~~	Vancomycin 500 mg injection	10 vials 40 ml
Omeprazole capsules or powder Sodium bicarbonate powder BP	qs 8.4 q	Glycerol BP Water	40 mi to 100 ml
Water	to 100 ml	(Only funded if prescribed for treatment of Clostridiu	
		following metronidazole failure)	
PHENOBARBITONE ORAL LIQUID		,	
Phenobarbitone Sodium	1 g		
Glycerol BP Water	70 ml to 100 ml		
Walei			

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy		Fully Brand or	
	(Manufacturer's P		sidised Generic	
	\$	Per	 Manufacturer 	
Extemporaneously Compounded Preparations	and Galenica	lls		
CODEINE PHOSPHATE – Safety medicine; prescriber may det	ermine dispensin	a frequency		
Powder – Only in combination		25 g		
,	(90.09)	5	Douglas	
Only in extemporaneously compounded codeine linctus	•		-	
COLLODION FLEXIBLE				
Note: This product is no longer being manufactured by the	supplier and will b	e delisted fror	n the Schedule at a dat	e to be
determined.				
Collodion flexible	19.30	100 ml	PSM	
COMPOUND HYDROXYBENZOATE - Only in combination				
Only in extemporaneously compounded oral mixtures.				
Soln		100 ml	 Midwest 	
GLYCERIN WITH SODIUM SACCHARIN - Only in combination	ı			
Only in combination with Ora-Plus or when used in the vanc	omycin oral Iquui	d Standard Fo	rmulae.	
Suspension		473 ml	 Ora-Sweet SF 	
GLYCERIN WITH SUCROSE – Only in combination				
Only in combination with Ora-Plus or when used in the vanc	omycin oral Iquui	d Standard Fo	rmulae.	
Suspension		473 ml	 Ora-Sweet 	
GLYCEROL				
* Liquid – Only in combination	3.23	500 ml	healthE Glycero	I BP
Only in extemporaneously compounded oral liquid prep	arations.			
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 c) Safety medicine; prescriber may determine dispensing fr 				
d) Extemporaneously compounded methadone will only be	reimbursed at the	e rate of the ch	eapest form available	
(methadone powder, not methadone tablets).	7.04	4		
Powder	7.84	1 g	🗸 AFT	
METHYL HYDROXYBENZOATE			/ .	
Powder	8.98	25 g	 Midwest 	
METHYLCELLULOSE				
Powder		100 g	✓ MidWest	
Suspension – Only in combination		473 ml	 Ora-Plus 	
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH				
Suspension		473 ml	 Ora-Blend SF 	
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - On				
Suspension		473 ml	 Ora-Blend 	
PHENOBARBITONE SODIUM				
Powder – Only in combination	52.50	10 g	 MidWest 	
	325.00	100 g	 MidWest 	
Only in children up to 12 years				
PROPYLENE GLYCOL				
Only in extemporaneously compounded methyl hydroxyben:			/ .	
Liq	11.25	500 ml	 Midwest 	
SODIUM BICARBONATE				
Powder BP – Only in combination		500 g	 Midwest 	
Only in extemporaneously compounded omeprazole an	u lansoprazolé su	ispension.		

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
SYRUP (PHARMACEUTICAL GRADE) – Only in combination				
Only in extemporaneously compounded oral liquid preparation Liq		500 ml	✓ M	lidwest
WATER Tap – Only in combination	0.00	1 ml	✓ Та	ap water

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Either:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children; or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Initial application — (Inborn errors of metabolism)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. **Renewal — (Cystic fibrosis or renal failure)** only from a dietitian, relevant specialist, vocationally registered general

practitioner or general practitioner on the recommendation of a dietitian, relevant specialist, vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

. Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT	- Special Authority see SA1930 above -	Hospital pharmacy	[HP3]
Powder		400 g OP	Polycal

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE AND FAT SU	JPPLEMENT – Special Author	ity see SA1376 on t	he previous pag	e – Hosp	oital pharmacy [HP3]
Powder (neutral)	-		400 g OP	Duod	al Super
			-	So	uble Powder

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
 \$	Per	1	Manufacturer

- 10 ascites; or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Patho

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Autho	ity see SA1523 on the previous	page – Hospital pharmacy [HP3]
--------------------------------	--------------------------------	--------------------------------

Emulsion (neutral)		200 ml OP	✓ Calogen
	30.75	500 ml OP	 Calogen
Emulsion (strawberry)		200 ml OP	 Calogen
Oil		500 ml OP	 MCT oil (Nutricia)
Oil, 250 ml		4 OP	 Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Resource Beneprotein

	armacy [HP3]	PROTEIN SUPPLEMENT – Special Authority see SA1524 above – Hospital ph	PR
 Protifar 	225 g OP	Powder	
 Resource 	227 g OP	8.95	
Donon	•		

Subsidy (Manufacturer's Price)

\$

Per

Fully Subsidised Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML – Special Authority s Liquid		500 ml OP	
DIABETIC ORAL FEED 1KCAL/ML – Special Authority see S Liquid (strawberry) Liquid (vanilla)	1.50	spital pharmacy 200 ml OP 200 ml OP	[HP3] ✓ Diasip ✓ Diasip ✓ Nutren Diabetes

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED – Special Authority see SA1525 above – Hospital pharmacy [HP3]

Powder	400 g OP	🗸 Monogen
--------	----------	-----------

(Ma	Subsidy	Fu	illy	Brand or
	nufacturer's Price)	Subsidis	ed	Generic
·	\$	Per	~	Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.
- ENTERAL/ORAL FEED 1KCAL/ML Special Authority see SA1098 above Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see	SA1099 above – Hos	pital pharmacy	[HP3]
Powder		400 g OP	 Kindergen

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

	Subsidy (Manufacturer's Pric \$		dised	Brand or Generic Manufacturer
continued applications meeting the following criteria: Both:				
 The treatment remains appropriate and the patient is ber General Practitioners must include the name of the dietit practitioner and date contacted. 			nally regi	stered general
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authorit Liquid		e previous pa 500 ml OP	🗸 Nut	spital pharmacy [HP3] t rini Energy RTH bini Energy
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority Liquid		previous page 500 ml OP	✓ Nut ✓ Pec	rini RTH liasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Sp pharmacy [HP3]	6.50 becial Authority see	SA1379 on th		bini Original us page – Hospital
Liquid	6.00	500 ml OP		rini Energy Multi ibre
	7.00		✓ Fre	bini Energy Fibre
PAEDIATRIC ENTERAL FEED WITH FIBRE 1KCAL/ML – Spe pharmacy [HP3]	cial Authority see S	A1379 on the	previous	page – Hospital
Liquid	7.00	500 ml OP		bini Original ibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority se	e SA1379 on the pr	evious page -	- Hospita	l pharmacy [HP3]
Liquid (strawberry)		200 ml OP	🗸 For	
Liquid (vanilla)		200 ml OP	 For 	
	6.99	500 ml OP	Pec	liasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML – Special Authority see				
Liquid (chocolate)		200 ml OP		liasure
Liquid (strawberry)		200 ml OP		liasure
Liquid (vanilla)		200 ml OP		liasure
		250 ml OP		liasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML – Specia pharmacy [HP3]	al Authority see SA1	379 on the pr	evious p	age – Hospital
Liquid (unflavoured)		200 ml OP		tini Multi Fibre
Liquid (chocolate)		200 ml OP		tini Multi Fibre
Liquid (strawberry)		200 ml OP		tini Multi Fibre
Liquid (vanilla)		200 ml OP		tini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA137 Powder		a <mark>ge</mark> – Hospital 400 g OP		cy [HP3] otamen Junior
Denal Dreducto				

Renal Products

► SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...

SPECIAL FOODS

 Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
 (manulastalor o r noc) \$	Per	✓	Manufacturer

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see Liquid		o <mark>revious page</mark> – 500 ml OP	
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA Liquid		ous page – Hos 220 ml OP	pital pharmacy [HP3] ✓ Nepro HP (strawberry) ✓ Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA11	01 on the previou	<mark>is page</mark> – Hospi	tal pharmacy [HP3]
Liquid, 200 ml bottle	11.52	4 OP	
	(13.24)		NovaSource Renal
Liquid (apricot) 125 ml		4 OP	Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	 Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - Special Aut	uthority see SA1377 above – Hospital pharmacy [HP3]
Liquid	18.06 1,000 ml OP 🖌 Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see SA137	377 above – Hospital pharmacy [HP3]
Liquid (grapefruit), 250 ml carton1	
Liquid (pineapple & orange), 250 ml carton1	171.00 18 OP
Liquid (summer fruits), 250 ml carton1	171.00 18 OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer \$ ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see SA1377 on the previous page - Hospital pharmacy [HP3] Vivonex TEN Powder (unflavoured)4.50 80 g OP SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Authority see SA1377 on the previous page - Hospital pharmacy [HP3] 500 ml OP Survimed OPD 12.04 1.000 ml OP Nutrison Advanced Peptisorb Peptisorb

(Peptisorb Liquid to be delisted 1 June 2023)

Paediatric Products For Children With Low Energy Requirements

➡SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Standard Supplements

■ SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and

continued...

SPECIAL FOODS

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
\$	Per	1	Manufacturer

3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - Patient is Malnourished
 - 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
 - 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	1	Manufacturer	

- 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority

forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

continued...

SPECIAL FOODS

	Subsidy (Manufacturer's \$	Price) Per	Fully Subsidised	Brand or Generic Manufacturer
ontinued				
 Is being fed via a tube or a tube is to be inserted for the pucondition criteria); or Cystic Fibrosis; or Liver disease; or Chronic Renal failure; or Inflammatory bowel disease; or Chronic obstructive pulmonary disease with hypercapnia; or Short bowel syndrome; or Bowel fistula; or Severe chronic neurological conditions. 		ng (not naso	ogastric tub	e - refer to specific medica
INTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 o	n nage 261 – H	-losnital nha	armacy [HP	3]
Liquid		250 ml (Ensure Plus HN
	7.00	1,000 ml	OP 🗸	Ensure Plus RTH Nutrison Energy
	9.60		✓ I	Fresubin HP Energy
NTERAL FEED 1KCAL/ML - Special Authority see SA1859 on				
Liquid	1.24 5.29	250 ml (1,000 ml	OP 🗸 I	sosource Standard Nutrison Standard RTH
	6.50			Osmolite RTH Fresubin Original
NTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority Liquid		on page 26 1,000 ml		l pharmacy [HP3] Nutrison 800 Complete Multi Fibre
NTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority set				
Liquid	5.29	1,000 ml	-	Jevity RTH
	7.00			Nutrison Multi Fibre Fresubin Original Fibre
NTERAL FEED WITH FIBRE 1.2KCAL/ML – Special Authority s		n <mark>page 261</mark> 1,000 ml		bharmacy [HP3] Jevity Plus
NTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority s Liquid		1,000 ml	OP 🗸	bharmacy [HP3] Jevity HiCal RTH Nutrison Energy Multi Fibre
	9.80		√	Fresubin HP Energy Fibre
NTERAL FEED WITH PROTEIN 1.2KCAL/ML – Special Author Liquid		on page 2 500 ml (tal pharmacy [HP3] F resubin Intensive
RAL FEED (POWDER) – Special Authority see SA1859 on page	e 261 – Hospi			
Powder (chocolate)		840 g C		Sustagen Hospital Formula
	26.00	850 g C		Ensure
Powder (vanilla)		840 g C		Sustagen Hospital Formula Active -
	26.00	850 g C	א או	Ensure

			SPECIAL FOODS
	Subsidy Manufacturer's		ully Brand or ed Generic
(1	\$	Price) Subsidis Per	Manufacturer
RAL FEED 1.5KCAL/ML - Special Authority see SA1859 on pag	e 261 – Hosp	ital pharmacy [HP3	3]
Additional subsidy by endorsement is available for patients bein			
epidermolysis bullosa, or as exclusive enteral nutrition in childre			
disease, or for patients with COPD and hypercapnia, defined a endorsed accordingly.	s CO2 value (exceeding 55mmH	g. The prescription must be
Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml	. ,		i olusip
with Endorsement		200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with Endorsement		237 ml OP	
Endorsement		237 MI OP	Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Si	ubsidised	Generic	
\$	Per	✓	Manufacturer	

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous p	<mark>age</mark> – Hospital p	harmacy [HP3]
Liquid5.50	500 ml OP	 Nutrison Concentrated
6.50		 Fresubin 2kcal HP
11.00	1,000 ml OP	 Ensure Two Cal HN RTH
ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the previous page Additional subsidy by endorsement is available for patients being bolus fed t epidermolysis bullosa. The prescription must be endorsed accordingly. Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with		, , ,
Endorsement0.96 (1.90)	200 ml OP	Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER - Special Authority see SA1	106 on the previous page - Hos	pital pharmacy	[HP3]
Powder		300 g OP	
	7.25	380 g OP	 Feed Thickener Karicare Aptamil

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Fither

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA1729 abo Powder		pharmacy [HP3] 1,000 g OP	
	(5.15)	1,000 g Ol	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729 abo	/e – Hospital	pharmacy [HP3]	
Powder		1,000 g OP	
	(7.32)	-	NZB Low Gluten Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR – Special Authority see SA1729 above – Powder		macy [HP3] 2,000 g OP	
	(18.10)	_,	Horleys Flour

	Subsidy (Manufacturer's Pric \$	· _	
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page - H	ospital pharmacy	[HP3]
Buckwheat Spirals		250 g OP	
	(3.11)		Orgran
Corn and Vegetable Shells	2.00	250 g OP	
	(2.92)		Orgran
Corn and Vegetable Spirals	2.00	250 g OP	
	(2.92)		Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP	
	(3.82)		Orgran
Rice and Corn Macaroni	2.00	250 g OP	
	(2.92)		Orgran
Rice and Corn Penne	2.00	250 g OP	
	(2.92)		Orgran
Rice and Maize Pasta Spirals	2.00	250 g OP	
	(2.92)		Orgran
Rice and Millet Spirals	2.00	250 g OP	
	(3.11)		Orgran
Rice and corn spaghetti noodles	2.00	375 g OP	
	(2.92)		Orgran
Vegetable and Rice Spirals	2.00	250 g OP	
	(2.92)	-	Orgran
Italian long style spaghetti	2.00	220 g OP	-
	(3.11)		Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

AMINOACID FORMULA WITHOUT METHIONINE - S	Special Authority see SA1108	<mark>3 above</mark> – Hospital	pharmacy [HP3]
Powder		500 g OP 🛛 🗸	XMET Maxamum

Supplements For MSUD

Powder 437.22	500 a OP	MSUD Maxamum
pharmacy [HP3]		
AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - S	pecial Authority se	ee SA1108 above – Hospital

	Subsidy (Manufacturer's Pr \$	ice) Subs Per	Fully sidised	Brand or Generic Manufacturer
upplements For PKU				
IINOACID FORMULA WITHOUT PHENYLALANINE armacy [HP3]	- Special Authority see SA	A1108 on the	previous	page – Hospital
Tabs		75 OP		lexy 10
Powder (berry) 28 g sachets	936.00	30		(U Lophlex Powder
Powder (chocolate) 36 g sachet		30		(U Anamix Junio) Chocolate
Powder (neutral) 28 g sachets	936.00	30		(U Lophlex Powder
Powder (neutral) 36 g sachets		30	🗸 Pł	(U Anamix Junio
Powder (orange) 28 g sachets	936.00	30		(U Lophlex Powder
Powder (orange) 36 g sachet		30		(U Anamix Junio) Orange
Powder (vanilla) 36 g sachet		30		(U Anamix Junio) Vanilla
Infant formula		400 g OP	🗸 Pł	U Anamix Infant
Powder (orange)		500 g OP	🖌 XF	9 Maxamum
Powder (unflavoured)		500 g OP	🖌 XF	9 Maxamum
Liquid (berry)	13.10	125 ml OP		(U Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP		(U Anamix Junio) LQ
Liquid (unflavoured)	13.10	125 ml OP		(U Anamix Junior LQ
Liquid (forest berries), 250 ml carton		18 OP	🖌 Ea	siphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	🗸 Pł	U Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP		U Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	🗸 Pł	(U Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	🗸 Pł	(U Lophlex LQ 10
Liquid (juicy orange) 62.5 ml	939.00	60 OP	🗸 Pł	(U Lophlex LQ 10
Liquid (juicy berries) 125 ml	936.00	30 OP	🗸 Pł	(U Lophlex LQ 20
Liquid (juicy orange) 125 ml		30 OP	🗸 Pł	U Lophlex LQ 20

Foods

LOW PROTEIN BAKING MIX – Special Authority see SA1108 on the previous Powder		pharmacy [HP3] ✓ Loprofin Mix
LOW PROTEIN PASTA - Special Authority see SA1108 on the previous page	- Hospital pharm	nacy [HP3]
Animal shapes	500 g OP	 Loprofin
Lasagne	250 g OP	 Loprofin
Low protein rice pasta11.91	500 g OP	 Loprofin
Macaroni	250 g OP	 Loprofin
Penne	500 g OP	 Loprofin
Spaghetti	500 g OP	 Loprofin
Spirals	500 g OP	 Loprofin

SPECIAL FOODS

	Subsidy (Manufacturer's Pric \$	e) Subs Per	Fully idised	Brand or Generic Manufacturer
Infant Formulae				
For Williams Syndrome				
 SA1110 Special Authority for Subsidy hitial application only from a dietitian, relevant specialist or vo ear where the patient is an infant suffering from Williams Syndi tenewal only from a dietitian, relevant specialist, vocationally re ecommendation of a dietitian, relevant specialist or vocationally pplications meeting the following criteria: soth: 1 The treatment remains appropriate and the patient is ber 2 General Practitioners must include the name of the dietit practitioner and date contacted. WW CALCIUM INFANT FORMULA – Special Authority see S/ Powder	rome and associated egistered general pra registered general pra nefiting from treatme ian, relevant special A1110 above – Hosp	hypercalca actitioner or practitioner. nt; and st or vocatic	emia. general Approv onally re cy [HP3	practitioner on the vals valid for 1 year for egistered general
Gastrointestinal and Other Malabsorptive Prob	olems			
MINO ACID FORMULA – Special Authority see SA2092 below Powder		cy [HP3] 400 g OP	-	Alfamino Alfamino Junior
Powder (unflavoured)	53.00	400 g OP	✓ E ✓ E ✓ N ✓ N	ilecare Elecare LCP Ieocate Gold Ieocate Junior Unflavoured Ieocate SYNEO
Powder (vanilla)	53.00	400 g OP	✓ E	lecare leocate Junior Vanilla

➡SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 6.2.2 Patient has IgE mediated allergy.

Initial application - (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	~	Manufacturer	

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

Liquid 1 kcal/ml	10.45	500 ml OP	Nutrini Peptisorb
Liquid 1.5 kcal/ml		500 ml OP	 Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure; or

2.10 Both:

2.10.1 The patient is currently receiving funded amino acid formula; and

Subsidy (Manufacturer's Price)	ç	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA	 Special Authority see SA1557 bel 	ow – Hospital pł	narmacy [HP3]
Powder		450 g OP	 Pepti-Junior
	30.42	900 g OP	 Allerpro Syneo 1
		-	Allerpro Syneo 2

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy (Manufacturer's Price)	Si	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 6 months for applications meeting the following criteria:

Approvals valid for 6 months for applications meeting the folio

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority se	e SA1197	above – Retail p	pharmacy
Powder (unflavoured)	35.50	300 g OP	 KetoCal 4:1
			 Ketocal 3:1
Powder (vanilla)	35.50	300 g OP	 KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

10

10

1

Boostrix Boostrix

Fully

Brand or

BCG Vaccine

(Manufacturer's Price) Subsidised Generic Per Manufacturer \$ Vaccinations BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm] For infants at increased risk of tuberculosis. Increased risk is defined as: 1) living in a house or family with a person with current or past history of TB; or 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php. Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Subsidy

Danish strain 1331, live attenuated, vial with diluent......0.00

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
 - 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old; or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.
 - Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.
- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 - 9 above.
- Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$) Si Per	Fully ubsidised	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE Funded for any of the following:	– [Xpharm]			
 A single dose for children up to the age of 7 who have A course of four vaccines is funded for catch up progra primary immunisation; or 				ars) to complete full
 An additional four doses (as appropriate) are funded for pre- or post splenectomy; pre- or post solid organ trans regimens; or 	splant, renal dialysis			
4) Five doses will be funded for children requiring solid or	•			
Note: Please refer to the Immunisation Handbook for appro	priate schedule for c	atch up p	orogramm	es.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous				
haemagglutinin, 8 mcg pertactin and 80 D-antigen units				
poliomyelitis virus in 0.5ml syringe		10	🗸 li	nfanrix IPV
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B A				
[Xpharm]				
Funded for patients meeting any of the following criteria:				
1) Up to four doses for children up to and under the age of	of 10 for primary imm	unisatio	n; or	
2) An additional four doses (as appropriate) are funded for	or (re-)immunisation f	or childr	en up to a	nd under the age of
10 who are patients post haematopoietic stem cell tran				
post solid organ transplant, renal dialysis and other se				
Up to five doses for children up to and under the age of	-	-		
Note: A course of up-to four vaccines is funded for catch up				
to complete full primary immunisation. Please refer to the In	nmunisation Handbo	ok for the	e appropri	ate schedule for catch up
programmes.				
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous				
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,				
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10	🖌 li	nfanrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]	0.00			
One dose for patients meeting any of the following:				
1) For primary vaccination in children; or				
 An additional dose (as appropriate) is funded for (re-)ir transplantation, or chemotherapy; functional asplenic; or post cochlear implants, renal dialysis and other seve 	pre or post splenecto	my; pre-	or post s	
3) For use in testing for primary immunodeficiency diseas paediatrician.				nal medicine physician or
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mc			-	
prefilled syringe plus vial 0.5 ml	0.00	1	✓ H	liberix
HEPATITIS A VACCINE – [Xpharm]				
Funded for patients meeting any of the following criteria:				
1) Two vaccinations for use in transplant patients; or	dicasco: ar			
 Two vaccinations for use in children with chronic liver of 3). One dose of vaccine for close contacts of known bena 				
a) One dose of vaccine for close contacts of known hepa				
3) One dose of vaccine for close contacts of known hepa		1	✓ Н	lavrix
	0.00	1 1		l <u>avrix</u> lavrix Junio <u>r</u>

		Subsidy		Fully	Brand or
		(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
		Ψ	rei	•	Manulaclurei
	COMBINANT VACCINE – [Xpharm]	0.00			Ta manine D
	er 0.5 ml prefilled syringe or patients meeting any of the following criteria		1	•	Engerix-B
	household or sexual contacts of known acute		0000	itio D corrio	ro: or
	children born to mothers who are hepatitis B s				15, 01
	children up to and under the age of 18 years in				a achieved a nositive
	ology and require additional vaccination or rec				
	HIV positive patients; or				
	hepatitis C positive patients; or				
	patients following non-consensual sexual inter	rcourse; or			
7) for	patients following immunosuppression; or				
	solid organ transplant patients; or				
,	post-haematopoietic stem cell transplant (HSC	CT) patients; or			
10) follo	owing needle stick injury.				
lni 20 mca ne	er 1 ml prefilled syringe	0.00	1		Engerix-B
	for patients meeting any of the following criteria		'	• !	
	household or sexual contacts of known acute		enat	titis R carrie	rs: or
	children born to mothers who are hepatitis B s				10, 01
	children up to and under the age of 18 years i				e achieved a positive
	ology and require additional vaccination or rec				
	HIV positive patients; or				
	hepatitis C positive patients; or				
	patients following non-consensual sexual inter	rcourse; or			
	patients following immunosuppression; or				
	solid organ transplant patients; or				
	post-haematopoietic stem cell transplant (HSC	() patients; or			
	owing needle stick injury; or dialysis patients; or				
	liver or kidney transplant patients.				
12) 101					
	DMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND	58) VACCINE [HPV] -	[Xn	harml	
Any of the fol			[/ P	lannj	
	im of two doses for children aged 14 years and	d under: or			
	im of three doses for patients meeting any of t				
,	eople aged 15 to 26 years inclusive; or	Ū			
2) Ei					
Pe	eople aged 9 to 26 years inclusive				
	1) Confirmed HIV infection; or				
	2) Transplant (including stem cell) patients:	or			
3) Maximu	Im of four doses for people aged 9 to 26 years	inclusive post chemoth	nerap	у	

Inj 270 mcg in 0.5 ml syringe0.00 10 🖌 Gardasil

 INFLUENZA VACCINE Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) [Xpharm] 11 A) INFLUENZA VACCINE - child aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis is available each year for patients aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis (a) have any of the following cardiovascular diseases (a) is chaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or have either of the following chronic respiratory disease (a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lution have diabetes; or iv) have diabetes; or iv) have diabetes; or iv) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or 	ths hths who meet th es: ung function; or	Lefollowing	Manufacturer Afluria Quad Junior (2023 formulation)
 Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) [Xpharm] 11 A) INFLUENZA VACCINE – child aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis (a) have any of the following cardiovascular diseases a) ischaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired let iii) have diabetes; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ths hths who meet th es: ung function; or	he following	(2023 formulation)
 [Xpharm]11 A) INFLUENZA VACCINE - child aged 6 months to 35 monis available each year for patients aged 6 months to 35 monis available each year for patients aged 6 months to 35 moni) have any of the following cardiovascular diseases a) ischaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired luii) have diabetes; or iv) have chronic renal disease; or v) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ths hths who meet th es: ung function; or	he following	(2023 formulation)
 A) INFLUENZA VACCINE - child aged 6 months to 35 morn is available each year for patients aged 6 months to 35 morn i) have any of the following cardiovascular diseases a) ischaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired to iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ths hths who meet th es: ung function; or	he following	(2023 formulation)
 is available each year for patients aged 6 months to 35 mor i) have any of the following cardiovascular diseases a) ischaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lu iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	nths who meet the set of the set		, , , , , , , , , , , , , , , , , , ,
 is available each year for patients aged 6 months to 35 mor i) have any of the following cardiovascular diseases a) ischaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired to iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	nths who meet the set of the set		criteria, as set by Pharma
 i) have any of the following cardiovascular diseases a) ischaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired luit iii) have diabetes; or iv) have cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	es: ung function; or		criteria, as set by Pharma
 a) ischaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lu iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ung function; or		
 b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lu iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ung function; or		
 c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lu iii) have diabetes; or iv) have chronic renal disease; or v) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ung function; or		
 d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired le iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ung function; or		
 e) cerebo-vascular disease; or ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired le iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ung function; or		
 ii) have either of the following chronic respiratory disease a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired le iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ung function; or		
 a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lu iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or 	ung function; or		
 b) other chronic respiratory disease with impaired living initial have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or 	-		
 iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or 	-		
 iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or 	cancers if not ir	nvasive; or	
 v) have any cancer, excluding basal and squamous skin vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or 	cancers if not ir	nvasive; or	
 vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or 	cancers if not ir	nvasive; or	
 a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or 			
 b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or 			
 c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or 			
 d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or 			
e) neuromuscular and CNS diseases/disorders, orf) haemoglobinopathies, or			
f) haemoglobinopathies, or			
, ,			
h) have a cochlear implant, or			
i) errors of metabolism at risk of major metabolic d	lecompensation	or	
j) pre and post splenectomy, or	ee en periodaien	,	
k) down syndrome, or			
vii) have been hospitalised for respiratory illness or have a	a historv of sign	ificant respir	atory illness:
Unless meeting the criteria set out above, the following cond	, ,		
a) asthma not requiring regular preventative therapy,			5
b) hypertension and/or dyslipidaemia without evidence of	f end-organ dise	ease.	
B) Doctors are the only Contractors entitled to claim payment f	-		accine ini 30 mca in 0.25 i
syringe (paediatric quadrivalent vaccine) to patients eligible			
and they may only do so in respect of the influenza vaccine	listed in the Pha	armaceutica	l Schedule.

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)......110.00

10

 Afluria Quad (2023 formulation)

(Man	Subsidy Jacturer's Price)	Full Subsidise		
	\$ 1	Per 🖌	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable
- C)

A) INFLUENZA VACCINE - people 3 years and over

- is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:
 - a) all people 65 years of age and over; or
 - b) People 55 to 64 years of age (inclusive) and is Māori or any Pacific ethnicity; or
 - c) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
 - children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
 - e) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or
 - f) children 3 to 12 years of age (inclusive), from 1 July 2022 to 31 December 2022;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- C)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

diluent 0.5 ml	5	🖌 MMR II
250.00	10	 Priorix

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Either:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 10 mcg of each meningococcal polysaccharide conjugated to a total of approximately 55 mcg of tetanus toxoid carrier			
per 0.5 ml vial	0.00	1	 MenQuadfi
Inj 4 mcg of each meningococcal polysaccharide conjugated to a total of approximately 48 mcg of diphtheria toxoid carrier			
per 0.5 ml vial	0.00	1	 Menactra
		5	✓ Menactra

Subsidy	F	ully Bran	id or
(Manufacturer's	Price) Subsid	sed Gen	eric
\$	Per	 Man 	ufacturer

MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm]

Any of the following:

- A) Three doses for children up to 12 months of age (inclusive) for primary immunisation; or
- B) Up to three doses (dependent on age at first dose) for a catch-up programme for children from 13 months to 59 months of age (inclusive) for primary immunisation, from 1 March 2023 to 31 August 2025; or
- C) Both:
 - 1) Person is one year of age or over; and
 - 2) Any of the following:
 - i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
 - ii) up to two doses for close contacts of meningococcal cases of any group; or
 - iii) up to two doses for person who has previously had meningococcal disease of any group; or
 - iv) up to two doses for bone marrow transplant patients; or
 - v) up to two doses for person pre- and post-immunosuppression*; or

D) Both:

- 1) Person is aged between 13 and 25 years (inclusive); and
- 2) Either:
 - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) Two doses for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 March 2023 to 28 February 2024.

*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 175 mcg per 0.5 ml prefilled syringe......0.00 1 🖌 Bexsero

MENINGOCOCCAL C CONJUGATE VACCINE – [Xpharm] Both:

- 1) The child is under 12 months of age; and
- 2) Any of the following:
 - Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) Two doses for close contacts of meningococcal cases of any group; or
 - 3) Two doses for child who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for child pre- and post-immunosuppression*.

Note: children under 12 months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 10 mcg in 0.5 ml syringe......0.00 1 Veisvac-C PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE – [Xpharm]

1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes In 1 mg of pneumococcal polysaccharide services 1, 5, 6B.

7F, 9V, 14 and 23F; 3 mcg of pneumococcal		
· · · · · · · · · · · · · · · · · · ·		
polysaccharide serotypes 4, 18C and 19F in 0.5 ml		
prefilled syringe0.00	10	Synflorix

*Three months or six months, as applicable, dispensed all-at-once

Subsidy	S	Fully	Brand or
(Manufacturer's Price)		ubsidised	Generic
\$	Per	1	Manufacturer

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
- Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously
 received two doses of the primary course of PCV10; or
- 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection; or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) cochlear implants or intracranial shunts; or
 - g) cerebrospinal fluid leaks; or
 - receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - diabetes; or
 - m) Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes
- Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml

syringe).00

10

1

Prevenar 13

Prevenar 13

Subsidy (Manufacturer's Price) \$	Su Per	Fully osidised	Brand or Generic Manufacturer
- [Xpharm]			
tional asplenia, pre- or p	oost-soli	d organ t	ransplant, renal dialysis,
	(Manufacturer's Price) \$ - [Xpharm] IIV, for patients post had tional asplenia, pre- or p	(Manufacturer's Price) Sul Per - [Xpharm] IIV, for patients post haematopo tional asplenia, pre- or post-solic	(Manufacturer's Price) Subsidised \$ Per ✔

- a) Patient is a child under 18 years for (re-)immunisation; and
- b) Treatment is for a maximum of two doses; and
- c) Any of the following:
 - i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - ii) with primary immune deficiencies; or
 - iii) with HIV infection; or
 - iv) with renal failure, or nephrotic syndrome; or
 - v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - vi) with cochlear implants or intracranial shunts; or
 - vii) with cerebrospinal fluid leaks; or
 - viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - x) pre term infants, born before 28 weeks gestation; or
 - xi) with cardiac disease, with cyanosis or failure; or
 - xii) with diabetes; or
 - xiii) with Down syndrome; or
 - xiv) who are pre-or post-splenectomy, or with functional asplenia.

Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each

23 pneumococcal serotype)0.00	1	Pneumovax	23
POLIOMYELITIS VACCINE – [Xpharm]			
Up to three doses for patients meeting either of the following:			
 For partially vaccinated or previously unvaccinated individuals; or For revaccination following immunosuppression. 			
Note: Please refer to the Immunisation Handbook for appropriate schedule for	catch-up p	rogrammes.	
Inj 80D antigen units in 0.5 ml syringe0.00	1	✓ IPOL	
ROTAVIRUS ORAL VACCINE – [Xpharm]			
Maximum of two doses for patients meeting the following:			
1) first dose to be administered in infants aged under 14 weeks of age; and			
2) no vaccination being administered to children aged 24 weeks or over.			
Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, squeezable tube	10	Rotarix	
Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, prefilled oral applicator0.00	10	✓ <u>Rotarix</u>	

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm]	_			
Either:				
1) Maximum of one dose for primary vaccination for either				
a) Any infant born on or after 1 April 2016; or				
b) For previously unvaccinated children turning 11 ye varicella infection (chickenpox), or	ears old on or after 1	July 201	7, who h	ave not previously had a
Maximum of two doses for any of the following:				
 Any of the following for non-immune patients: 				
 i) with chronic liver disease who may in future ii) with deteriorating renal function before trans iii) prior to solid organ transplant; or 		nsplantat	ion; or	
iv) prior to any elective immunosuppression*, o	r			
v) for post exposure prophylaxis who are immu	une competent inpatie	ents.; or		
b) For patients at least 2 years after bone marrow transmission of the patients at least 2 years after bone marrow transmission of the patients.				
c) For patients at least 6 months after completion of				
 d) For HIV positive non immune to varicella with mile a) For activate with inhore arrays of matched line at a 				
 For patients with inborn errors of metabolism at ris varicella, or 	sk of major metabolic	aecomp	ensation	, with no clinical history of
f) For household contacts of paediatric patients who	are immunocompron	nised or	underac	ning a procedure leading to
immune compromise where the household contact				
g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure has no clinical history of varicella.	e no clinical history of	varicella	and wh	o are severely
* immunosuppression due to steroid or other immunosuppres	ssive therapy must be	for a tre	atment c	period of greater than
28 days				3
Inj 1350 PFU prefilled syringe	0.00	1 10		'arivax 'arivax
VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] - [Xph	arm]			
Funded for patients meeting the following criteria:	•			
1) Two doses for all people aged 65 years				
Inj 50 mcg per 0.5 ml vial plus vial	0.00	1	✓ s	hingrix
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATE	D VACCINE ISHING	I ES VAC	CINF1 -	- [Xpharm]
Funded for patients meeting the following criteria:]	[, [, [, [, []]]]
1) One dose for all people aged 65 years				
Inj 19,400 PFU prefilled syringe plus vial	0.00	1	17	ostavax
ing 10,400 FT O premieu synnige plus viai		10	_	lostavax
Diagnostic Agents				
TUBERCULIN PPD [MANTOUX] TEST – [Xpharm]				
Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	✓ <u>⊺</u>	ubersol

.

- Symbols -
3TC
7 MED NSHA Silver/Copper
Short
- A -
A-Scabies69
Abacavir sulphate 106
Abacavir sulphate with
lamivudine 106
Abacavir/Lamivudine Viatris 106
Abiraterone acetate 164
Acarbose11
Accarb 11
Accuretic 1048
Accuretic 2048
Acetazolamide246
Acetec
Acetic acid with hydroxyquinoline and
ricinoleic acid
Acetylcysteine249
Aci-Jel
Aciclovir
Infection102
Sensory244
Acidex6
Acipimox 55
Acitretin69
Aclasta115
Actemra219
Actemra S29219
Actinomycin D150
Actrapid 10
Actrapid Penfill10
Acupan 119
Adalimumab (Amgevita) 174
Adalimumab (Humira - Alternative
brand) 183
Adapalene
ADR Cartridge 1.8
Adrenaline
Cardiovascular
Respiratory
Advantan
Advate
Adynovate
Afinitor231 Aflibercept
Allibercept
(2023 formulation) 278
Afluria Quad Junior
(2023 formulation) 278
AFT-Pyrazinamide101
Agents Affecting the
Renin-Angiotensin System

Agents for Parkinsonism and Related
Disorders 117
Agents Used in the Treatment of
Poisonings 249
Agrylin149
Albendazole90
Albey
Albustix
Alchemy158
Alchemy Oxybutynin78
Aldurazyme
Alecensa157
Alectinib157
Alendronate sodium112
Alendronate sodium with
colecalciferol112
Alfacalcidol
Alfamino270
Alfamino Junior270
Alginic acid6
Alglucosidase alfa27
Alkeran146
Alkeran S29146
Allerfix246
Allerpro Syneo 1
Allerpro Syneo 2273
Allersoothe
Allmercap148
Allopurinol 115
Alpha-Adrenoceptor Blockers
Alpha-Keri Lotion
Alphamox
Alphamox 125
Alphamox 250
Alprolix
Alu-Tab6
Aluminium hydroxide6
Alvogen
Amantadine hydrochloride117
Ambrisentan
Ambrisentan Mylan58
Ambrisentan Viatris
Amgevita174
Amiloride hydrochloride53
Amiloride hydrochloride with
furosemide
Amiloride hydrochloride with
hydrochlorothiazide54
Aminophylline241
Amiodarone hydrochloride
Amisulpride129
Amitriptyline 123
Amlodipine
Amorolfine

Amoxicillin	93
Amoxicillin with clavulanic acid	93
Amphotericin B	33
Amsacrine	.149
AmsaLyo	. 149
Amsidine	. 149
Amzoate	
Anaesthetics	.118
Anagrelide hydrochloride	. 149
Analgesics	. 119
Anastrozole	. 167
Anatrole	. 167
Andriol Testocaps	
Androderm	81
Anoro Ellipta	.240
Antabuse	.142
Antacids and Antiflatulents	6
Anthelmintics	
Antiacne Preparations	62
Antiallergy Preparations	.235
Antianaemics	
Antiandrogen Oral	
Contraceptives	76
Antiarrhythmics	
Antibacterials	
Antibacterials Topical	
Anticholinergic Agents	239
Anticholinesterases	.111
Antidepressants	123
Antidiarrhoeals	
Antiepilepsy Drugs	
Antifibrinolytics, Haemostatics and	
Local Sclerosants	38
Antifibrotics	
Antifungals	
Antifungals Topical	63
Antihistamines	.236
Antihypotensives	
Antimalarials	
Antimigraine Preparations	. 128
Antinausea and Vertigo Agents	. 128
Antipruritic Preparations	64
Antipsychotics	. 129
Antiretrovirals	
Antirheumatoid Agents	.112
Antispasmodics and Other Agents	
Altering Gut Motility	8
Antithrombotic Agents	41
Antithymocyte globulin	
(equine)	. 174
Antitrichomonal Agents	. 100
Antituberculotics and	
Antileprotics	. 100
Antiulcerants	9

INDEX: Generic Chemicals and Brands

Antivirals 101
Anxiolytics133
Anzatax154
Apidra11
Apidra SoloStar11
APO-Atomoxetine
APO-Atomoxetine S29
Apo-Azithromycin
Apo-Diltiazem CD52
Apo-Temozolomide 154
Apomorphine hydrochloride117
Aprepitant 128
Apresoline
Aqueous cream
Aratac
Arava
Arginine
Arginine
Aripiprazole130
Aripiprazole Sandoz 130
Aristocort65
Arrotex-Prazosin S2947
Arrow - Clopid41
Arrow - Lattim
Arrow-Amitriptyline123
Arrow-Bendrofluazide
Arrow-Brimonidine
Arrow-Diazepam
Arrow Devery hisin
Arrow-Doxorubicin
Arrow-Fluoxetine 124
Arrow-Losartan &
Hydrochlorothiazide
Arrow-Norfloxacin 110
Arrow-Ornidazole100
Arrow-Quinapril 1048
Arrow-Quinapril 1048 Arrow-Quinapril 20 48
Arrow-Quinapril 2048
Arrow-Quinapril 20

Atomoxetine1	
Atorvastatin	55
Atropine sulphate	
Cardiovascular	49
Sensory2	
Atropt	
Atrovent	
Aubagio1	
Augmentin	
Aurorix 1	
AutoSoft 30	
AutoSoft 90	24
Avallon1	
Avelox	
Avonex1	
Avonex Pen	
Azacitidine 1	
Azacitidine Dr Reddy's 1	40
Azamun 1	40
Azathioprine1	
Azilect1	17
Azithromycin	91
Azopt	
AZT1	07
- в - B-D Micro-Fine	
B-D Ultra Fine	16
B-D Ultra Fine II	16
Bacillus Calmette-Guerin (BCG)	
vaccine 1	74
Bacillus Calmette-Guerin	
Bacillus Calmette-Guerin vaccine	275
Bacillus Calmette-Guerin vaccine	275 16
Bacillus Calmette-Guerin vaccine	275 16 62
Bacillus Calmette-Guerin vaccine	275 16 62 30
Bacillus Calmette-Guerin vaccine	275 16 62 30 66
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 235 44
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 235 44
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237 237 235 44 54
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237 237 237 237 235 44 54
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 230 66 275 237 237 237 237 237 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 230 66 275 237 237 237 237 237 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 37 237 237 237 237 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237 237 237 237 237 237
Bacillus Calmette-Guerin vaccine	275 16 62 30 66 275 237 237 237 237 237 237 237 237 237 237

Beta Ointment64
Beta Scalp71
Beta-Adrenoceptor Agonists238
Beta-Adrenoceptor Blockers50
Betadine
Betadine Skin Prep67
Betaferon
Betahistine dihydrochloride
Betaine
Betaloc CR
Betanethasone dipropionate
Betamethasone dipropionate with
calcipotriol
Betamethasone sodium phosphate
with betamethasone acetate 80
Betamethasone valerate64, 71
Betamethasone valerate with sodium
fusidate [fusidic acid] 65
Betaxolol246
Betnovate64
Betoptic246
Betoptic S246
Bexsero
Bezafibrate
Bezalip
Bezalip Retard
Bicalutamide
Bicillin LA
BiCNU145 Bile and Liver Therapy10
Bile and Liver Therapy10
Biltricide
Bimatoprost
Bimatoprost Multichem
Binarex
Binocrit
Biodone 121
Biodone Extra Forte121
Biodone Forte 121
Bisacodyl27
Bisacodyl Viatris
Bisoprolol fumarate
Bisoprolol Mylan 50
Bisoprolol Viatris
BK Lotion
Bleomycin sulphate
Blood Colony-stimulating
Factors
Blood glucose diagnostic test
meter
Blood glucose diagnostic test
strip
Blood glucose test strips (visually
impaired)15
Blood Ketone Diagnostic Test
Strip 14
Bonjela

D 11 075
Boostrix
Bortezomib150
Bortezomib Dr-Reddy's 150
Bosentan
Bosentan Dr Reddy's58
Bplex
Breo Ellipta
Brevinor 1/28
Bricanyl Turbuhaler
Brimonidine tartrate
Brimonidine tartrate with timolol
maleate
Illaleale
Brinzolamide
Brown & Burk 124
Brufen SR 111
BSF Alchemy249
BSF Celapram249
BSF Ticagrelor Sandoz249
BSF Vebulis249
Buccastem129
Budesonide
Alimentary6
Respiratory
Budesonide with eformoterol
Bumetanide
Buprenorphine Naloxone BNM
Buprenorphine with naloxone
Bupropion hydrochloride
Burel
Burinex
Burinex S2953
Buscopan
Buscopan S298
Buspirone hydrochloride133
Buspirone Viatris 133
Busulfan145
- C -
Cabergoline
Caffeine citrate243
Calamine64
Calamine-AFT 64
Calci-Tab 50035
Calcipotriol69
Calcitonin79
Calcitriol34
Calcitriol-AFT34
Calcium 500 mg Hexal35
Calcium carbonate
Calcium carbonate PAI6
Calcium Channel Blockers52
Calcium Disodium Versenate
Calcium folinate 147
Calcium Folinate Ebewe
Calcium Folinate Sandoz
Calcium Folinate Sandoz S29

Calcium gluconate	35
Calcium Homeostasis	79
Calcium polystyrene sulphonate	46
Calcium Resonium	46
Calogen	.256
Candesartan cilexetil	
Candestar	
Canesten	
Capecitabine	
Capecitabine-DRLA	147
Capercit	1/7
Capoten	. 147
Capsaicin	47
Musculoskeletal	110
Nervous	
Captopril	
Carafate	
Carbaccord	.145
Carbamazepine	
Carbimazole	83
Carbomer	
Carboplatin	.145
Carboplatin Ebewe	.145
Carbosorb-X	.249
Cardinol LA	51
Cardizem CD	
CareSens Dual	14
CareSens N	15
CareSens N POP	15
CareSens N Premier	
CareSens PRO	15
Carmellose sodium with gelatin and	
pectin	
Carmustine	
Carnitor	
Carvedilol	
Carvedilol Sandoz	
Casirivimab and imdevimab	101
Catapres	
CeeNU	00
Cefaclor monohydrate	. 140
Cetacior mononydrate	90
Cefalexin	
Cefazolin	
Ceftriaxone	
Ceftriaxone-AFT	91
Cefuroxime axetil	
Celapram	
Celebrex	
Celecoxib	
Celecoxib Pfizer	
Celestone Chronodose	
Cellcept	. 167
Centrally-Acting Agents	53
Cephalexin ABM	90
Cetirizine hydrochloride	.236
Cetomacrogol	

INDEX: Generic Chemicals and Brands

Cetomacrogol with glycerol6	6
Cetomacrogol-AFT6	6
Cetuximab19	
Charcoal24	9
Chemotherapeutic Agents14	4
Chickenpox vaccine	
Chlorafast24	
Chlorambucil14	
Chloramphenicol24	
Chlorothiazide5	
Chlorpheniramine maleate23	
Chlorpromazine hydrochloride13	
Chlortalidone [Chlorthalidone]5	54
Chlorthalidone	54
Chlorvescent4	6
Choice 380 7med Nsha Silver/copper	
Short	'4
Choice Load 3757	1
Choice TT380 Short7	24
Choice TT380 Standard7	
Choline salicylate with cetalkonium	1
chloride	13
Ciclosporin	
Cilazapril	17
Cilicaine VK	14
Cinacalcet	'n
Cipflox	
Ciprofloxacin	-
Infection9	14
Sensory24	л 1
Ciprofloxacin Teva	И
Cisplatin	15
Cisplatin Ebewe	
Citalopram hydrobromide 12	ν Δ
Cladribine	
Clarithromycin	
Alimentary	۵
Infection9	
Clexane	
Clexane Forte	2
Climara	
Clindamycin	15
Clinicians	
Clinicians Renal Vit	2.U
Clobazam	
Clobetasol propionate	
Clobetasone butyrate	
Clofazimine10	
Clomazol	0
Dermatological6	3
Genito-Urinary7	20
Clomifene citrate	
Clomipramine hydrochloride	
Clomipramine Teva	ט. מו
Clonazepam 125, 13	
Clonidine	03 0
	5

INDEX: Generic Chemicals and Brands

Clonidine hydrochloride53
Clonidine Teva53
Clopidogrel41
Clopidogrel Multichem
Clopine
Clopixol131, 133
Clotrimazole
Dermatological
Genito-Urinary
Clozapine
Clozaril
Co-trimoxazole
Coal tar
Coal tar with allantoin, menthol,
phenol and sulphur
Coal tar with salicylic acid and
sulphur
Coco-Scalp
Codeine phosphate
Extemporaneous
Nervous
Coenzyme Q10
Colchicine
Colecalciferol
Colestid
Colestipol hydrochloride
Colgout
Colifoam
Colistin sulphomethate
Colistin-Link
Collodion flexible
Colloidal bismuth subcitrate
Colofac
Coloxyl
Combigan
Compound electrolytes
Compound electrolytes with glucose
[Dextrose]
Compound hydroxybenzoate
Comtan
Concerta
Condoms
Condyline71
Contraceptives - Hormonal
Contraceptives - Non-hormonal72
Copaxone
Cordarone-X
Corticosteroids and Related Agents
for Systemic Use
Corticosteroids Topical
Cortifoam7
Cosentyx216
Cosmegen 150
Coumadin44
Country Life
Coversyl47

Creon 10000	25
Creon 25000	25
Creon Micro	
Crotamiton	
Crystaderm	62
Curam Curam Duo 500/125	90
Cvite	34
Cyclizine hydrochloride	
Cyclizine lactate	. 129
Cyclogyl	.247
Cyclonex	
Cyclopentolate hydrochloride	.247
Cyclophosphamide	.145
Cyclorin	. 100
Cycloserine	
Cyproterone acetate	
Cyproterone acetate with	
ethinyloestradiol	76
Cystadane	
Cytarabine	1/7
Cytotec	
Cytoxan	9
- D -	. 140
D-Penamine	110
D-Penamine	. 112
Dabigatran	
Dacarbazine	. 150
Dacarbazine APP	
Dactinomycin [Actinomycin D]	
Daivobet	69
Daivobet Daivonex	69 69
Daivobet Daivonex Daktarin	69 69 64
Daivobet Daivonex Daktarin Dalacin C	69 69 64 95
Daivobet Daivonex Daktarin	69 69 64 95
Daivobet Daivonex Daktarin Dalacin C	69 69 64 95 .116
Daivobet Daivonex Daktarin Dalacin C Dantrium	69 64 95 .116 .116
Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29	69 64 95 .116 .116 .116
Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29 Dantrolene Daonil	69 64 95 .116 .116 .116 11
Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29 Dantrolene Daonil Dapa-Tabs	69 64 95 .116 .116 .116 11 54
Daivobet Daivonex	69 69 64 95 . 116 . 116 11 11 54 . 100
Daivobet Daivonex	69 64 95 . 116 . 116 . 116 11 54 . 100 96
Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29 Dantrolene Daonil Dapa-Tabs Dapsone Daraprim Darunavir	69 64 95 . 116 . 116 . 116 11 54 . 100 96 . 107
Daivobet Daivonex Daktarin Dalacin C Dantrium S29 Dantrolene Daonil Dapa-Tabs Dapsone Daraprim Darunavir Darunavir Mylan	69 64 95 . 116 . 116 . 116 . 116 . 117 96 . 107 . 107
Daivobet	69 69 95 . 116 . 116 . 116 . 116 . 100 96 . 107 . 107 . 107
Daivobet	69 69 95 . 116 . 116 . 116 . 116 . 117 . 107 . 107 . 107 . 157
Daivobet	69 69 64 95 . 116 . 116 11 54 . 100 96 . 107 . 107 . 157 . 150
Daivobet	69 69 64 95 . 116 . 116 11 54 . 100 . 107 . 107 . 157 . 150 . 150
Daivobet	69 69 64 95 . 116 . 116 11 54 . 107 . 107 . 107 . 107 . 150 . 150 y
Daivobet	69 64 95 .116 .116 .116 11 .107 .107 .107 .150 .150 .150 y 77
Daivobet	69 64 95 . 116 . 116 . 116 11 54 . 107 . 107 . 107 . 107 . 150 . 150 y 77 57
Daivobet	69 69 95 . 116 . 116 . 116 . 117 . 100 . 107 . 107 . 107 . 150 y 77 57 . 241
Daivobet	69 69 95 . 116 . 116 . 116 . 117 . 100 . 107 . 107 . 107 . 150 y 77 57 . 241 . 150
Daivobet	69 69 64 95 . 116 . 116 11 54 . 100 96 . 107 . 107 . 157 . 150 . 150 y 77 57 . 241 . 150 . 150
Daivobet	69 69 64 95 . 116 . 116 11 54 . 107 . 107 . 107 . 107 . 150 . 150 y 77 57 . 241 . 150 . 150 . 150
Daivobet	69 69 64 95 . 116 . 116 11 54 . 100 96 . 107 . 107 . 107 . 107 . 150 . 150 57 241 . 150 . 150 145 145
Daivobet	69 69 64 95 . 116 . 116 11 54 . 100 96 . 107 . 107 . 107 . 107 . 150 . 150 y 77 57 . 241 . 150 . 150 . 145 . 145 . 150

BP	250
DBL Docetaxel	150
DBL Ergometrine	77
DBL Gemcitabine	148
DBL Gentamicin	95
DBL Heparin Sodium	44
DBL Leucovorin Calcium	
DBL Methotrexate Onco-Vial	148
DBL Pethidine Hydrochloride	123
DBL Vincristine Sulfate	157
Decozol	
Deferasirox	249
Deferiprone	250
Deferiprone Deferoxamine Pfizer S29	250
Denosumab	112
Deolate	
Deoxycoformycin	
Depo-Medrol	
Depo-Provera	01
Depo-Provera	/0
Depo-Testosterone	
Deprim	
Dermol	b4, 71
Desferrioxamine mesilate	
Desmopressin	88
Desmopressin acetate	
Desmopressin-PH&T	88
Desuric	115
Detection of Substances in	
Urine	78
Dexamethasone	
Hormone	
Sensory	
Dexamethasone phosphate	80
Dexamethasone with framycetin ar	nd
gramicidin	244
Dexamethasone with neomycin	
sulphate and polymyxin B	
sulphate	245
Dexamfetamine sulfate	137
Dexmethsone	<mark>80</mark>
Dextrochlorpheniramine	
maleate	236
Dextrose	45–46
DHC Continus	121
Diabetes	10
Diabetes Management	14
Diacomit	127
Diagnostic Agents	284
Diamide Relief	6
Diamox	
Diasip	257
Diazepam12	5. 133
Diazoxide	
Dibenzyline	
Diclofenac Sandoz	
· ··· · · · · ·	

INDEX: Generic	Chemicals	and	Brands
----------------	-----------	-----	--------

Diclofenac sodium
Musculoskeletal 111
Sensory245
Differin
Difflam
Diflucan
Digestives Including Enzymes25
Digoxin
Dihydrocodeine tartrate
Dilantin
Dilantin Infatab126
Diltiazem CD Clinect
Diltiazem hydrochloride
Dimethicone
Dimethyl fumarate
Dipentum
Diphtheria, tetanus and pertussis
vaccine
Diphtheria, tetanus, pertussis and
polio vaccine
Diabthoria tatonua nartuosia nalia
Diphtheria, tetanus, pertussis, polio,
hepatitis B and haemophilus
influenzae type B vaccine 276
Diprosone
Diprosone OV
Dipyridamole
Disopyramide phosphate
Disulfiram
Diuretics
Dizole
Docetaxel
Docetaxel Sandoz
Docusate sodium
Docusate sodium with
sennosides
Dolutegravir
Domperidone
Domperidone Viatris
Donepezil hydrochloride
Donepezil-Rex
Dornase alfa
Dortimopt
Dorzolamide hydrochloride246
Dorzolamide with timolol246
Dostinex
Dosulepin [Dothiepin]
hydrochloride
Dosulepin Mylan
Dosulepin Viatris
Dothiepin
Doxazosin
Doxazosin Clinect
Doxine
Doxorubicin Ebewe
Doxorubicin hydrochloride150

=

Doxycycline	
DP Lotion	<mark>67</mark>
DP Lotn HC	65
DP-Allopurinol	115
Dr Reddy's Omeprazole	9
Drofate	51
Drugs Affecting Bone	
Metabolism	. 112
Dual blood glucose and blood keto	
diagnostic test meter	14
Dulaglutide	12
Dulcolax SP Drop	27
Duocal Super Soluble Powder	255
Duolin	239
Duolin HFA	239
DuoResp Spiromax	
Duride	
Durvalumab	
- F -	220
e-chamber La Grande	243
e-chamber Mask	
e-chamber Turbo	
E-Mycin	
e5 Pharma	
Ear Preparations	
Ear/Eye Preparations	
Easiphen Liquid	269
Econazole nitrate	63
Efavirenz	
Efavirenz with emtricitabine and	
tenofovir disoproxil	107
Eformoterol fumarate	
Eformoterol fumarate dihydrate	
Eftrenonacog alfa [Recombinant	201
factor IX]	20
Efudix	50
Egopsoryl TA	71 70
Elaprase	70 20
Eldepryl	
Elecare	
Elecare LCP	270 270
Electral	
Elelyso	
Elemental 028 Extra	260
Elidel	
Elocon	
Elocon Alcohol Free	05 65
Eltrombopag	
Eltroxin	00 מע
EMB Fatol	
Emend Tri-Pack	
Emicizumab	
EMLA	
Empagliflozin	
Empagliflozin with metformin	14
hydrochloride	14

Emtricitabine	107
Emtricitabine with tenofovir	
disoproxil	104
Emtriva	107
Emulsifying ointment	. 66
Emulsifying Ointment ADE	. 66
Enalapril maleate	.47
Enbrel	167
Endocrine Therapy	164
Endoxan	145
Engerix-B	277
Enlafax XR	125
Enoxaparin sodium	43
Enstilar	69
Ensure	264
Ensure Plus	265
Ensure Plus HN	264
Ensure Plue RTH	264
Ensure Plus RTH Ensure Two Cal HN RTH	204
Entacapone	200
Entecavir	101
Entecavir Mylan	
Entecavir Sandoz	101
Entocort CIR	101
Entresto 24/26	
Entresto 49/51	
Entresto 97/103	40 10
Entyvio	40 006
Epilim	107
Epilim Crushable	107
Epilim IV	107
Epilim S/F Liquid	107
Epilim Syrup	107
Epillin Syrup	127
Epipen	230
Epipen Jr	235
Epirubicin Ebewe	151
Epirubicin hydrochloride	151
Eplerenone	
Epoetin alfa	
Epoprostenol Eptacog alfa [Recombinant factor	61
Eptacog alta [Recombinant factor	
VIIa]	. 40
Erbitux	191
Ergometrine maleate	
Erlotinib	158
Erythrocin IV	. 92
Erythromycin (as lactobionate)	. 92
Erythromycin ethyl succinate	. 92
Esbriet	240
Escitalopram Escitalopram (Ethics)	124
Escitalopram (Ethics)	124
Eskazole	. 90
Essential Ethosuximide	125
Essential Generics	50
Essential Prednisolone	
Estraderm MX	. 82

Estradiol TDP Mylan82
Estradot82
Estradot 50 mcg82
Estrofem82
Etanercept 167
Ethambutol hydrochloride100
Ethics Aspirin119
Ethics Aspirin EC41
Ethics Lisinopril47
Ethinyloestradiol with
desogestrel
Ethinyloestradiol with
levonorgestrel
Ethinyloestradiol with
norethisterone
Ethosuximide
Etopophos 151
Etoposide
Etoposide phosphate
Etravirine
Eumovate65
Evara
Everet
Everolimus231
Evista
Evusheld219
Exemestane167
Exjade249
Extemporaneously Compounded
Preparations and
Preparations and Galenicals
Preparations and Galenicals252 Eye Preparations244
Preparations and Galenicals
Preparations and Galenicals
Preparations and Galenicals
Preparations and Galenicals
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe with simvastatin 57 - F - -
Preparations and Galenicals
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe with simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40
Preparations and Galenicals
Preparations and Galenicals
Preparations and Galenicals
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe with simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Famotidine 9 Fasenra 190 Faslodex 164
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe with simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Famotidine 9 Fasenra 190 Faslodex 164 Fatty Cream AFT 66
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe with simvastatin 57 - F - Factor eight inhibitor bypassing fraction fraction 9 Famotidine 9 Fasenra 190 Fastodex 164 Fatty Cream AFT 66 Febuxostat 115
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe with simvastatin 57 - F - Factor eight inhibitor bypassing fraction fraction 40 Famotidine 9 Fasenra 190 Faslodex 164 Fatty Cream AFT 66 Febuxostat 115
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe with simvastatin 57 - F - Factor eight inhibitor bypassing fraction fraction 9 Famotidine 9 Fasenra 190 Fastodex 164 Fatty Cream AFT 66 Febuxostat 115
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe With simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Fasenra 190 Faslodex 164 Fatty Cream AFT 66 Febuxostat multichem 115 Feed Thickener Karicare Aptamil
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe With simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Fasenra 190 Faslodex 164 Febuxostat 115 Febuxostat multichem 115 Fede Thickener Karicare Aptamil Aptamil 267 FEIBA NF 40
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe Sandoz 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Fasenra 190 Faslodex 164 Fatty Cream AFT 66 Febuxostat 115 Feed Thickener Karicare 40 Aptamil 267 FEIBA NF 40 Felo 10 ER 52
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe With simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Fasenra 190 Faslodex 164 Febuxostat 115 Febuxostat multichem 115 Fede Thickener Karicare Aptamil Aptamil 267 FEIBA NF 40
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe With simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Fasenra 190 Faslodex 164 Fatty Cream AFT 66 Febuxostat multichem 115 Feed Thickener Karicare Aptamil. Aptamil. 267 FEIBA NF 40 Felo 10 ER 52 Felo 5 ER 52
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe with simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Fasenra 190 Faslodex 164 Fetty Cream AFT 66 Febuxostat multichem 115 Feed Thickener Karicare 40 Felba NF 40 Felb NF 40 Felb NF 52 Felo 5 ER 52 Felo 5 ER 52 Felodipine 52
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe With simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Fasenra 190 Faslodex 164 Fatty Cream AFT 66 Febuxostat multichem 115 Feed Thickener Karicare Aptamil. Aptamil. 267 FEIBA NF 40 Felo 10 ER 52 Felo 5 ER 52
Preparations and Galenicals 252 Eye Preparations 244 Eylea 189 Ezetimibe 56 Ezetimibe Sandoz 56 Ezetimibe with simvastatin 57 - F - Factor eight inhibitor bypassing fraction 40 Famotidine 9 Fasenra 190 Faslodex 164 Fetty Cream AFT 66 Febuxostat multichem 115 Febuxostat multichem 115 Felb XNF 40 Felb NF 40 Felo 10 ER 52 Felo 5 ER 52 Felodipine 52 Felodipine 52 Felodipine 52 Felodipine 52 Felodipine 52

Ferinject	35
Ferodan	35
Ferriprox	250
Ferro-F-Tabs	35
Ferro-tab	35
Ferrograd	
Ferrosig	
Ferrous fumarate	35
Ferrous fumarate with folic acid	
Ferrous sulfate	
Fexofenadine hydrochloride	236
Fibro-vein	41
Filgrastim	44
Finasteride	
Fingolimod	
Firazyr	
Flagyl	100
Flagyl-S	
Flamazine	
Flecainide acetate	
Flecainide BNM Flecainide Controlled Release	49
Teva	
Fleet Phosphate Enema	
Flixonase Hayfever & Allergy	242
Flixotide	237
Flixotide Accuhaler	
Florinef	
Fluanxol	
Flucil	
Flucloxacillin	93
Flucloxacillin-AFT	
Flucloxin	
Flucon	
Fluconazole	
Fludara Oral	
Fludarabine Ebewe	147
Fludarabine phosphate	
Fludrocortisone acetate	80
Fluids and Electrolytes	45
Flumetasone pivalate	244
Fluocortolone caproate with	
fluocortolone pivalate and	
cinchocaine	8
Fluorometholone	
Fluorouracil	
Fluorouracil Accord	
Fluorouracil sodium	71
Fluox	
Fluoxetine hydrochloride	
Flupenthixol decanoate	
Flutamide	
Flutamin	
Fluticasone	
Fluticasone furoate with	
vilanterol	238
	200

Fluticasone propionate242
Fluticasone with salmeterol238
Flynn
FML
Foban
Folic acid
Folic Acid multichem
Folic Acid Mylan
Folic Acid Viatris
Food Thickeners
Foods And Supplements For Inborn
Errors Of Metabolism
Foradil
Forteo
Fortini
Fortini Multi Fibre
Fortisip
Fortisip
Fortisip Multi Fibre
Fosamax
Fosamax Plus
Framycetin sulphate
Frebini Energy
Frebini Energy Fibre
Frebini Original
Frebini Original Fibre
Fresubin 2kcal HP 266
Fresubin HP Energy
Fresubin Intensive264
Fresubin Original 264
Fresubin Original Fibre
Frisium125
Frumil54
Frusemide53
Fucicort
Fucidin
Fucithalmic
Fulvestrant164
Fungilin
Furosemid-Ratiopharm53
Furosemide [Frusemide]53
Furosemide-Baxter
fusidic acid
Dermatological
Infection
Sensory
- G -
Gabapentin
Gacet
Galsulfase
Galvumet
Galvus
Gardasil 9277
Gastrodenol
Gaviscon Extra Strength
Gaviscon Infant6

Gazyva201
Gefitinib159
GEM Aqueous Cream
Gemcitabine Ebewe
Gemcitabine hydrochloride
Gemtuzumab ozogamicin
Genoptic
Gentamicin sulphate
Infection95
Sensory244
Gilenya
Ginet
Glatiramer acetate 134
Glecaprevir with pibrentasvir 103
Glibenclamide11
Gliclazide11
Glipizide11
Glivec
Glizide11
Glucagen Hypokit 10
Glucagon hydrochloride10
Glucerna Select257
Glucose [Dextrose]45
Gluten Free Foods
Glycerin with sodium saccharin 252
Glycerin with sucrose
Glycerol
Alimentary27
Extemporaneous
Glyceryl trinitrate
Alimentary8
Cardiovascular
Glycopyrronium239
Glycopyrronium bromide8
Glycopyrronium with
indacaterol240
Go Healthy29
Gold Knight73
Gold Knight XL
Goserelin
Gutron
Gynaecological Anti-infectives
- H -
Habitrol
Haemophilus influenzae type B
vaccine
Haldol 131
Haldol Concentrate131
Haldol Decanoas 131
Haloperidol130
Haloperidol decanoate
Harvoni 103
Havrix
Havrix Junior
Havior syrup
healthE Dimethicone 10%

healthE Dimethicone 4% Lotion	
healthE Dimethicone 5%	
healthE Glycerol BP29	
healthE Urea Cream	66
Healtheries Simple Baking Mix2	67
Hemastix	
Hemlibra	
Heparin sodium	44 1 /
Heparinised saline	
Heparon Junior	
Hepatitis A vaccine	
Hepatitis B recombinant	
vaccine	77
Herceptin	22
Hiberix	
Hiprex 10	
Histaclear2	
Histafen23	
Holoxan14	
Horleys Bread Mix	67
Horleys Flour	<u>57</u>
Hormone Replacement Therapy -	
Systemic	
HPV21 Humalog	
Humalog Mix 25	11
Humalog Mix 50	
rianialog mix oo	
Human papillomavirus (6, 11, 16, 18,	
Human papillomavirus (6, 11, 16, 18,	
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV]	
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV]	77 96
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV]	77 96
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 2 Humatin 2 Humira 18 Humira 18	77 96 33
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 2 Humatin 2 Humira 18 Humulin 30/70 18	77 96 33 33
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 2 Humatin 2 Humira 18 HumiraPen 18 Humulin 30/70 18	77 96 33 33 11
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 2 Humatin 2 Humira 18 Humira 18 Humira 18 Humira 18 Humulin 30/70 18 Humulin R 18	77 96 33 11 10
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 2 Humatin 2 Humira 18 HumiraPen 18 Humulin 30/70 18 Humulin R 14 Humulin R 14	77 96 33 11 10 10
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humira 18 Humira 18 HumiraPen 18 Humulin 30/70 18 Humulin R 18 Hyaluronic acid 22	77 96 33 11 10 10 48 58
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humatin 27 Humira 18 HumiraPen 18 Humulin 30/70 18 Humulin R 19 Hydralazine 24 Hydralazine 24	77 96 33 11 10 10 48 58
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humira 18 Humira 18 HumiraPen 18 Humulin 30/70 19 Humulin NPH 19 Humulin R 24 Hydralazine 14 Hydralazine 14 Hydrocortisone 14	77 96 33 11 10 10 48 58
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humira 18 HumiraPen 18 Humulin 30/70 18 Humulin R 18 Hydralazine 24 Hydralazine hydrochloride 24 Hydrocortisone 19 Dermatological 6	77 96 33 11 10 10 48 58 58 58
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] IHPV] 2 Humatin 2 Humira 1 HumiraPen 1 Humulin 30/70 1 Humulin ccid 2 Hydralazine 2 Hydralazine 2 Hydralazine 3 Hydrocortisone 2 Dermatological 6 Hormone 6	77 96 33 11 10 10 48 58 58 55 30
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 2 Humatin 2 Humira 1 Humira 1 Humira 1 HumiraPen 1 Humulin 30/70 1 Humulin R 2 Hydralazine 4 Hydralazine 4 Hydrocortisone 1 Dermatological 6 Hydrocortisone (PSM) 6	77 96 33 11 10 10 48 58 58 55 30 55
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 2 Humatin 2 Humira 1 HumiraPen 1 Humulin 30/70 1 Humulin caid 2 Hydralazine 2 Hydralazine 3 Hydrocortisone 2 Dermatological 4 Hydrocortisone (PSM) 6 Hydrocortisone acetate 6	77 96 33 11 10 10 48 58 58 55 30 55
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 2 Humatin 2 Humatin 2 Humira 18 Humira 18 Humira Pen 18 Humulin 30/70 18 Humulin NPH 18 Hydralazine 24 Hydralazine 24 Hydrocortisone 25 Hydrocortisone 24 Hydrocortisone 24 Hydrocortisone	77 93 33 11 10 48 58 53 53 53 53 53 53 53 53 53 53 53 53 53
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humatin 27 Humira 18 Humira 18 Humira 18 Humulin 30/70 18 Humulin 30/70 18 Humulin NPH 18 Hydralazine 18 Hydralazine 19 Hydrocortisone 10 Dermatological 10 Hydrocortisone (PSM) 10 Hydrocortisone acetate 10 Hydrocortisone hydrochloride 10	77 93 33 11 10 48 58 53 53 53 53 53 53 53 53 53 53 53 53 53
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 2 Humatin 2 Humatin 2 Humira 18 Humira 18 Humira Pen 18 Humulin 30/70 18 Humulin NPH 18 Hydralazine 24 Hydralazine 24 Hydrocortisone 25 Hydrocortisone 24 Hydrocortisone 24 Hydrocortisone	77 96 33 11 10 10 48 58 55 56 55 57 7 7
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humatin 27 Humatin 27 Humira 18 HumiraPen 18 Humulin 30/70 18 Humulin NPH 18 Hydralazine 24 Hydralazine 24 Hydrocortisone 27 Dermatological 26 Hydrocortisone acetate 26 Hydrocortisone acetate 26 Hydrocortisone acetate 26 Hydrocortisone acetate 27 Hydrocortisone acetate 26 Hydrocortisone acetate 27 Hydrocortisone butyrate 26	77933311 1014855 530557 71
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humatin 27 Humira 18 HumiraPen 18 Humulin 30/70 18 Humulin R 24 Hydralazine 24 Hydralazine 24 Hydrocortisone 26 Dermatological 26 Hydrocortisone acetate 26 Hydrocortisone acetate 27 Hydrocortisone acetate 28 Hydrocortisone acetate 29 Hydrocortisone butyrate 26 Hydrocortisone with cinchocaine 29	77 96 33 33 31 10 10 10 10 10 10 10 10 10 10 10 10 10
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humatin 27 Humira 18 HumiraPen 18 Humulin 30/70 18 Humulin 70 19 Humulin 8 24 Hydralazine 26 Hydralazine 27 Hydrocortisone 26 Dermatological 26 Hydrocortisone acetate 26 Hydrocortisone acetate with 27 Hydrocortisone acetate with 26 Hydrocortisone acetate with 26 Hydrocortisone butyrate 65 Hydrocortisone with cinchocaine 26 Hydrocortisone with miconazole 27	77 96 33 33 31 10 10 10 10 10 10 10 10 10 10 10 10 10
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humira 18 Humira 18 Humira 18 Humira 18 Humira 18 Humulin 30/70 19 Humulin 8 18 Hydronic acid 24 Hydralazine 18 Hydralazine 19 Hydrocortisone 10 Dermatological 6 Hydrocortisone acetate 10 Hydrocortisone acetate with 10 pramoxine hydrochloride 10 Hydrocortisone acetate with 10 pramoxine hydrochloride 10 Hydrocortisone butyrate 65, 7 Hydrocortisone with cinchocaine 10 Hydrocortisone with miconazole 10 Hydrocortisone with matamycin and 10	77 96 33 33 31 11 10 10 85 85 85 85 85 77 7 7 7 55 71 .85 5
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humira 18 Humira 18 Humira 18 Humira 18 Humulin R 18 Humulin NPH 18 Hydroncic acid 24 Hydralazine 18 Hydrocortisone 26 Dermatological 6 Hydrocortisone acetate 19 Hydrocortisone acetate with 6 Hydrocortisone and paraffin liquid 6 Hydrocortisone butyrate 65 Hydrocortisone with cinchocaine 6 Hydrocortisone with cinchocaine 6 Hydrocortisone with natamycin and 6	77 96 33 33 31 11 10 10 85 85 85 85 85 85 85 85 85
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] 27 Humatin 27 Humira 18 Humira 18 Humira 18 Humira 18 Humira 18 Humulin 30/70 19 Humulin 8 18 Hydronic acid 24 Hydralazine 18 Hydralazine 19 Hydrocortisone 10 Dermatological 6 Hydrocortisone acetate 10 Hydrocortisone acetate with 10 pramoxine hydrochloride 10 Hydrocortisone acetate with 10 pramoxine hydrochloride 10 Hydrocortisone butyrate 65, 7 Hydrocortisone with cinchocaine 10 Hydrocortisone with miconazole 10 Hydrocortisone with matamycin and 10	77 96 33 33 11 10 10 48 58 55 50 55 52 52 52 55 55 52 55 55 55

Hydroxocobalamin Panpharma 33
hydroxycarbamide 151
Hydroxychloroquine112
Hydroxyurea
[hydroxycarbamide]
Hygroton54
Hylo-Fresh248
Hymenoptera235–236
Hyoscine butylbromide8
Hyoscine hydrobromide129
Hypam 136
Hyperuricaemia and Antigout 115
Hypromellose247
Hypromellose with dextran 247
-1-
lbiamox93
Ibrance160
Ibrutinib151
Ibuprofen 111
Icatibant235
Idarubicin hydrochloride151
Idursulfase
Ifosfamide145
Igroton54
Ikorel58
Iloprost61
Imatinib mesilate159
Imatinib-Rex 159
Imbruvica 151
Imfinzi 228
Imigran128
Imipramine hydrochloride 123
Imiquimod71
Immune Modulators108
Immunosuppressants 167
Incruse Ellipta 239
Indacaterol237
Indapamide54
Infanrix IPV
Infanrix-hexa276
Infant Formulae270
Infatrini274
Infliximab 192
Influenza vaccine278
Inhaled Corticosteroids237
Inhaled Long-acting
Beta-adrenoceptor Agonists 237
Inspra53
Instillagel Lido118
Insulin aspart11
Insulin aspart with insulin aspart
protamine10
Insulin glargine11
Insulin glulisine 11
Insulin isophane10
Insulin isophane with insulin

neutral11
Insulin lispro11
Insulin lispro with insulin lispro
protamine11
Insulin neutral 10
Insulin pen needles16
Insulin pump16
Insulin pump cartridge21
Insulin pump infusion set (steel
cannula)21
Insulin pump infusion set (steel
cannula, straight insertion) 22
Insulin pump infusion set (teflon
cannula)23
Insulin pump infusion set (teflon
cannula, angle insertion with
insertion device) 24
Insulin pump infusion set (teflon
cannula, angle insertion) 24
Insulin pump infusion set (teflon
cannula, straight insertion with
insertion device) 24
Insulin pump infusion set (teflon
cannula, straight insertion)24
Insulin pump reservoir24
Insulin syringes, disposable with
attached needle 16
Intelence 106
Interferon beta-1-alpha134
Interferon beta-1-beta134
Intra-uterine device74
Invega Sustenna132
Invega Trinza132
IPCA-Frusemide53
IPCA-Metoprolol51
IPCA-Propranolol51
IPOL
Ipratropium bromide239, 242
Iressa159
Irinotecan Actavis 100 148
Irinotecan hydrochloride 148
Irinotecan-Rex
Iron (as ferric carboxymaltose)
Iron polymaltose
Isentress
Isentress HD108
Ismo 20
Ismo 40 Retard
Isoniazid
Isoniazid with rifampicin
Isoptin
Isoptin Retard
Isoptin SR
Isopto Carpine
Isosorbide mononitrate
Isosource Standard 264

Isotretinoin
Ispaghula (psyllium) husk26
Itch-Soothe64
Itraconazole
Itrazole
Ivacaftor
Ivermectin
- J -
Jadelle
Jakavi
Jardiamet14
Jardiance
Jaydess
Jevity HiCal RTH
Jevity Plus
Jevity RTH264
Jinarc
Juno Pemetrexed149
- K -
Kadcyla224
Kaletra 107
Kalydeco242
Kemadrin 117
Kenacomb244
Kenacort-A 1081
Kenacort-A 4081
Kenalog in Orabase
Ketocal 3:1274
KetoCal 4:1274
Ketoconazole
Dermatological
Infection
Ketogenic Diet274
Ketoprofen111
KetoSens14
Keytruda230
Kindergen258
Kivexa
Klacid
Alimentary9
Infection
Kliogest83
Kliovance
Kogenate FS41
Konakion MM
Konsyl-D
Kuvan
-1 -
Labetalol51
Lacosamide
Lactulose
Laevolac
Lagevrio 104
Lagevrio104
Lamictal

Lamivudine Viatris	
Lamotrigine	.126
Lamprene	. 100
Lanoxin	. 49
Lanoxin Paediatric Elixir	40
Lanoxin PG	
Lanoxin S29	
Lansoprazole	
Lantus	
Lantus SoloStar	
Lanua Solosiai	140
Lanvis	. 149
Lanzol Relief	9
Lapatinib ditosylate	. 159
Largactil	
Laronidase	
Lasix	
Latanoprost	
Latanoprost with timolol	.247
Lax-Suppositories	27
Lax-suppositories Glycerol	27
Laxatives	26
Laxsol	
Ledipasvir with sofosbuvir	.103
Leflunomide	.112
Lenalidomide	. 152
Letrole	. 167
Letrozole	. 167
Leukeran FC	. 145
Leukeran FC	
Leukotriene Receptor	
Leukotriene Receptor Antagonists	241
Leukotriene Receptor Antagonists Leuprorelin	241 88
Leukotriene Receptor Antagonists Leuprorelin Leustatin	241 88 .147
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam	241 88 .147 .126
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam-AFT	241 88 .147 .126 .126
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam-AFT Levlen ED	241 88 .147 .126 .126 75
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam.AFT Levlen ED Levocabastine	241 88 .147 .126 .126 75 .246
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam-AFT Levlen ED Levocabastine Levocamitine	241 88 .147 .126 .126 75 .246 30
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam-AFT Levlen ED Levocabastine Levocamitine Levodopa with benserazide	241 88 .147 .126 .126 75 .246 30 .117
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam-AFT Levlen ED Levocabastine Levocapa with benserazide Levodopa with carbidopa	241 88 .147 .126 .126 75 .246 30 .117 .117
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam-AFT Levlen ED Levocabastine Levocapa with benserazide Levodopa with carbidopa Levomepromazine	241 88 .147 .126 .126 75 .246 30 .117 .117
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam-AFT Levlen ED Levocabastine Levocarnitine Levodopa with benserazide Levodopa with benserazide Levomepromazine Levomepromazine	241 88 .147 .126 .126 75 .246 30 .117 .117 .130
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam-AFT Levlen ED Levocabastine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine hydrochloride	241 88 .147 .126 .126 75 .246 30 .117 .117 .130
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam.AFT Levlen ED Levocabastine Levocarnitine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine hydrochloride Levonrgestrel	241 88 .147 .126 .126 30 .117 .117 .130
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam.AFT Levlen ED Levocabastine Levocarnitine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine hydrochloride Levonorgestrel Genito-Urinary	241 88 .147 .126 .126 75 .246 30 .117 .117 .130 .130
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam.AFT Levlen ED Levocabastine Levocanitine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levodopa with carbidopa Levomepromazine hydrochloride Levonorgestrel Genito-Urinary Hormone	241 88 .147 .126 .126 75 .246 30 .117 .117 .130 .130 76 83
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam.AFT Levlen ED Levocabastine Levocabastine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine hydrochloride Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM	241 88 .147 .126 .126 75 .246 30 .117 .130 .130 76 83 76
Leukotriene Receptor Antagonists Leuprorelin Levestatin Levetiracetam Levetiracetam.AFT Levine ED Levocabastine Levocabastine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine hydrochloride Levomepromazine hydrochloride Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM	241 88 .147 .126 .126 75 .246 75 .117 .117 .130 76 83 76 83
Leukotriene Receptor Antagonists Leuprorelin Levestatin Levetiracetam Levetiracetam.AFT Levine ED Levocabastine Levocabastine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine hydrochloride Levomepromazine hydrochloride Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM Levonorgestrel BNM Levothyroxine	241 88 .147 .126 .126 75 .246 75 .117 .117 .130 76 83 76 83
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam.AFT Leven ED Levocabastine Levocabastine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine hydrochloride Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levothyroxine Lidocaine [Lignocaine]	241 88 .147 .126 .126 75 .246 30 .117 .130 .130 76 83 76 83 76
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam.AFT Levien ED Levocanitine Levocarnitine Levodopa with benserazide Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine hydrochloride Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM	241 88 .147 .126 .126 75 .246 30 .117 .130 .130 76 83 76 83 76
Leukotriene Receptor Antagonists Leuprorelin Levetiracetam Levetiracetam.AFT Levien ED Levocabastine Levocarnitine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine Levomepromazine Levomepromazine Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Lidocaine [Lignocaine]	241 88 .147 .126 .126 75 .246 30 .117 .130 .130 76 83 76 83 76 83 76 83 83
Leukotriene Receptor Antagonists Leuprorelin Leustatin Levetiracetam Levetiracetam Levelin ED Levocabastine Levocapa with benserazide Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine hydrochloride Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM Levonorgestrel BNM Levothyroxine Lidocaine [Lignocaine] hydrochloride Lidocaine [Lignocaine] with chlorhexidine	241 88 .147 .126 .126 75 .246 30 .117 .130 .130 76 83 76 83 76 83 76 83 83
Leukotriene Receptor Antagonists Leuprorelin Levetiracetam Levetiracetam.AFT Leveln ED Levocabastine Levocabastine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine Levomepromazine Levomepromazine Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levontyroxine Lidocaine [Lignocaine]	241 88 .147 .126 75 .246 30 .117 .117 .130 76 83 76 83 -119 .119
Leukotriene Receptor Antagonists Leuprorelin Levetiracetam Levetiracetam AFT Leveln ED Levocabastine Levocabastine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine Levomepromazine Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM Levothyroxine Lidocaine [Lignocaine]	241 88 .147 .126 75 .246 30 .117 .130 .117 .130 76 83 76 83 76 83 76 83 719
Leukotriene Receptor Antagonists Leuprorelin Levetiracetam Levetiracetam.AFT Leveln ED Levocabastine Levocabastine Levodopa with benserazide Levodopa with benserazide Levodopa with carbidopa Levomepromazine Levomepromazine Levomepromazine Levonorgestrel Genito-Urinary Hormone Levonorgestrel BNM Levonorgestrel BNM Levonorgestrel BNM Levontyroxine Lidocaine [Lignocaine]	. 241 88 . 147 . 126 75 . 246 75 . 130 . 117 . 130 . 130 76 83 76 83 76 83 76 83 76 83 75 119 . 119 . 119 . 119 . 119 . 119

Life Extension32
Lignocaine 118–119
Lioresal Intrathecal 116
Lipid-Modifying Agents55
Liquigen256
Liraglutide12
Lisinopril
Litak
Lithium carbonate 130
Livostin
LMX4 119
Lo-Oralcon 20 ED75
Locacorten-Viaform ED's244
Local preparations for Anal and
Rectal Disorders 8
Locasol
Locoid
Locoid Crelo65
Locoid Lipocream65
Locorten-Vioform
Lodoxamide246
Logem126
Lomide246
Lomustine146
Loniten58
Loperamide hydrochloride6
Lopinavir with ritonavir107
Lopinavir/Ritonavir Mylan 107
Loprofin
Loprofin Mix269
Lorafix236
Loratadine236
Lorazepam133
Lorstat55
Losartan Actavis
Losartan potassium
Losartan potassium with
hydrochlorothiazide 48
Lovir
Loxamine
Lucrin Depot 1-month
Lucrin Depot 3-month
Lyderm
Lynparza153
Lyrica 126
- M - m-Eslon
Mabthera
Macro Organic Psyllium Husk
Macrobid110 Macrogol 3350 with potassium
chloride, sodium bicarbonate and sodium chloride27
Madopar 125 117
Madopar 250 117
Madopar 250
maaopa 02.0 117

Madopar HBS	
Madopar Rapid	. 117
Magnesium hydroxide	<mark>36</mark>
Magnesium sulphate	36
Mantoux	
Marevan	
Marine Blue Lotion SPF 50+	71
Martindale Pharma	
Mask for spacer device	243
Maviret	103
Maxidex	245
Maxitrol	245
MCT oil (Nutricia)	240
Measles, mumps and rubella	200
vaccine	000
Mebendazole	
Mebeverine hydrochloride	90
webeverine hydrochioride	8
Medrol	80
Medroxyprogesterone acetate	
Genito-Urinary	
Hormone8	
Mefenamic acid	
Melatonin	
Melpha	
Melphalan	
Menactra	
Meningococcal (groups A, C, Y and	
W-135) conjugate vaccine	280
	. 200
Meningococcal B multicomponent	. 200
Meningococcal B multicomponent	
Meningococcal B multicomponent vaccine	
Meningococcal B multicomponent vaccine Meningococcal C conjugate	. 281
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine	. 281 . 281
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi	. 281 . 281 . 281
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol	. 281 . 281 280 64
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Mepolizumab	. 281 . 281 280 64 200
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Mepolizumab Mercaptopurine	. 281 . 281 280 64 200 148
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Mepolizumab Mercaptopurine Mercilon 28	. 281 . 281 280 64 200 148 74
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Mercaptopurine Mercaptopurine Mercalazine	. 281 . 281 280 64 200 148 74 7
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Mercaptopurine Mercaptopurine Mercalazine Mesalazine	. 281 . 281 280 64 200 148 74 74 72
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Meroptopurine Mercaptopurine Mesalazine Mesna Mestinon	. 281 . 281 280 64 200 148 74 74 72 152 111
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Meraptopurine Mercalptopurine Mesalazine Mestanon Mestanon Metabolic Disorder Agents	. 281 . 281 280 64 200 148 74 74 7 152 111 27
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Mercaptopurine Mercaptopurine Mesalazine Mesna Mestinon Metabolic Disorder Agents Metformin hydrochloride	. 281 . 281 280 64 200 148 74 7 152 111 27 11
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Meraptopurine Mercalptopurine Mercalptopurine Mestalpto 28 Mestalpto 28 Mestanon Mestanon Mestanon Metabolic Disorder Agents Metformin hydrochloride Metformin Mylan	. 281 . 281 280 64 74 74 152 111 27 11
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Meraptopurine Mercaptopurine Mercaptopurine Mesalazine Mesalazine Mestionon Metabolic Disorder Agents Metformin hydrochloride Metformin Mylan Metformin Viatris	. 281 . 281 280 64 200 148 74 152 111 27 11 11
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Merabolizumab Mercaptopurine Mercaptopurine Mesalazine Mesalazine Mestanon Mestonon Metabolic Disorder Agents Metformin hydrochloride Metformin Mylan Metformin Viatris Methadone BNM	. 281 . 281 280 64 200 148 74 152 111 27 11 11
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Merolizumab Merolizumab Mercaptopurine Mercaptopurine Mesalazine Mestalolic Disorder Agents Metformin hydrochloride Metformin Mylan Metformin Viatris Methadone BNM	. 281 . 281 280 64 200 148 74 152 111 27 11 11 121
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Menthol Merthol Mercaptopurine Mercaptopurine Mercaptopurine Mesalazine Mesalazine Mestalolic Disorder Agents Metformin hydrochloride Metformin Viatris Metformin Viatris Methadone BNM Methadone hydrochloride Extemporaneous	. 281 . 281 280
Meningococcal B multicomponent vaccine Meningococcal C conjugate vaccine MenQuadfi Merolizumab Merolizumab Merolizumab Merolizumab Merolizumab Merolizumab Mesalazine Mesalazine Mestabolic Disorder Agents Metformin hydrochloride Metformin Mylan Metformin Mylan Methadone BNM Methadone BNM Methadone hydrochloride Extemporaneous Nervous	. 281 . 281 280
Meningococcal B multicomponent vaccine	. 281 . 281 280 64 7 152 111 7 152 111 11 27 11 21
Meningococcal B multicomponent vaccine	. 281 . 281 280 64 7 152 111 27 11 11 252 121 109
Meningococcal B multicomponent vaccine	. 281 . 281 280 64 200 148 74 152 111 11 121 27 121 222 121 109 247
Meningococcal B multicomponent vaccine	. 281 . 281 280 64 200 148 74 152 111 11 121 27 121 212 121 129 247 148
Meningococcal B multicomponent vaccine	. 281 . 281 280 64 200 148 74 152 111 27 11 121 252 121 109 247 148 148
Meningococcal B multicomponent vaccine	. 281 . 281 280 64 200 148 74 152 111 27 11 121 252 121 129 247 148 148
Meningococcal B multicomponent vaccine	. 281 . 281 280 64 74 77 152 111 77 11 121 252 121 252 121 247 148 148 148
Meningococcal B multicomponent vaccine	. 281 . 281 280 64 7 7 7 152 111 7 7 11

Methylcellulose 252
Methylcellulose with glycerin and
sodium saccharin
Methylcellulose with glycerin and
sucrose
Methyldopa53
Methyldopa Mylan53
Methyldopa Mylan S2953
Methylnaltrexone bromide
Methylphenidate ER - Teva
Methylphenidate hydrochloride 138
Methylphenidate hydrochloride
inethylphenidate hydrochlonde
extended-release
Methylprednisolone80
Methylprednisolone (as sodium
succinate) 80
Methylprednisolone aceponate65
Methylprednisolone acetate81
Methylxanthines241
Metoclopramide Actavis 10 129
Metoclopramide hydrochloride 129
Metolazone54
Metopirone89
Metoprolol IV Mylan51
Metoprolol IV Viatris51
Metoprolol succinate
Metoprolol tartrate
Metrogyl
Metronidazole
Metyrapone
Mexiletine hydrochloride
Miacalcic
Micolette
Micolette-S2927
Miconazole
Miconazole nitrate
Dermatological64
Genito-Urinary76
Micreme76
Micreme H 65
Microgynon 20 ED75
Microgynon 3075
Microlut
Midazolam135
Midazolam Mylan 135
Midazolam-Baxter
Midodrine
Mifegyne
Mifepristone
Milpharm
Minerals
Mini-Wright AFS Low Range
Mini-Wright Standard243
Winit-Wright Standard243
Minidiab11
MiniMed 1.8 Reservoir
MMT-326A24

MiniMed 3.0 Reservoir
MMT-332A 24
MiniMed 770G 16
MiniMed Mio MMT-921A23
MiniMed Mio MMT-923A23
MiniMed Mio MMT-925A23
MiniMed Mio MMT-941A23
MiniMed Mio MMT-943A23
MiniMed Mio MMT-945A23
MiniMed Mio MMT-965A23
MiniMed Mio MMT-975A23
MiniMed Quick-Set MMT-386A23
MiniMed Quick-Set MMT-387A23
MiniMed Quick-Set MMT-396A23
MiniMed Quick-Set MMT-397A23
MiniMed Quick-Set MMT-398A23
MiniMed Quick-Set MMT-399A23
MiniMed Silhouette MMT-368A23
MiniMed Silhouette MMT-377A23
MiniMed Silhouette MMT-378A23
MiniMed Silhouette MMT-381A23
MiniMed Silhouette MMT-382A23
MiniMed Silhouette MMT-383A23
MiniMed Silhouette MMT-384A23
MiniMed Sure-T MMT-864A21
MiniMed Sure-T MMT-866A21
MiniMed Sure-T MMT-874A21
MiniMed Sure-T MMT-876A21
MiniMed Sure-T MMT-884A21
MiniMed Sure-T MMT-886A21
Minims Pilocarpine
Minims Prednisolone
Minirin
Minirin Melt88
Mino-tabs94
Minocycline hydrochloride94
Minomycin
Minor Skin Infections67
Minoxidil58
Mirena83
Mirtazapine 125
Misoprostol9
Mitomycin C 153
Mitozantrone
Mitozantrone Ebewe153
Mixtard 3011
MMR II
Moclobemide 124
Modafinil
Modavigil 140
Moduretic
Molaxole
Molnupiravir104
Moments
Mometasone furoate

Montelukast 241
Montelukast Mylan241
Montelukast Viatris
Moroctocog alfa [Recombinant factor
VIII]
Morphine hydrochloride 122
Morphine sulphate 122
Motetis118
Mouth and Throat
Movapo117
Moxifloxacin
MSUD Maxamum
Mucilaginous laxatives with
stimulants 26
Mucolytics
Mucosoothe 119
Multiple Sclerosis Treatments 133
Multivitamin renal
Multivitamins
Mupirocin
Muscle Relaxants 116
Mvite
Myambutol 100
Mycobutin 101
MycoNail
Mycophenolate mofetil
Mydriacyl
Mylan (12 hr release)
Mylan (24 hr release)
Mylan Atenolol
Mylan Clomiphen
Mylan Indapamide
Mylan Italy (24 hr release)
Myleran
Mylotarg191
Myometrial and Vaginal Hormone
Preparations
Myozyme
- N -
Nadolol
Nadolol BNM
Naglazyme
Nalcrom
Naloxone hydrochloride
Naltraccord
Naltrexone hydrochloride
Naphazoline hydrochloride
Naphcon Forte
Naprosyn SR 1000
Naprosyn SR 750 111
Naproxen111
Narcaricin mite
Nasal Preparations
Natalizumab134
Natalizumab
Nausafix
11000011/

Nausicalm	128
Navelbine	157
Nefopam hydrochloride	119
Neisvac-C	281
Neo-Mercazole	83
Neocate Gold	270
Neocate Junior Unflavoured	270
Neocate Junior Vanilla	270
Neocate SYNEO	270
Neoral	
Neostigmine metilsulfate	111
Nepro HP (strawberry)	260
Nepro HP (vanilla)	260
Nepro HP RTH	260
Neulactil	
Neulastim	45
Neuraxpharm	130
NeuroTabs	
Nevirapine	106
Nevirapine Alphapharm	106
Nevirapine Viatris	106
Nicorandil	58
Nicotine	142
Nifedipine	52
Nifuran	110
Nilotinib	160
Nilstat	
Alimentary	
Genito-Urinary	
Infection	98
Nintedanib	240
Nipent	154
Nirmatrelvir with ritonavir	105
Nitrates	57
Nitroderm TTS	
Nitrofurantoin	110
Nitrolingual Pump Spray	57
Nivestim	44
Nivolumab	229
Nizoral	
Nodia	
Noflam 250	
Noflam 500	111
Non-Steroidal Anti-Inflammatory	
Drugs	111
Nonacog gamma, [Recombinant	
Factor IX]	40
Norethisterone	
Genito-Urinary	
Hormone	
Norflex	116
Norfloxacin	110
Noriday 28	76
Norimin	
Normacol Plus	
Normison	135

Norpress 123
Nortriptyline hydrochloride
Norvir
Noumed
Dermatological65
Nervous121, 125
Noumed Paracetamol120
Noumed Pethidine 123
NovaSource Renal260
Novatretin69
NovoMix 30 FlexPen 10
NovoRapid11
NovoRapid FlexPen11
NovoRapid Penfill
NovoSeven RT40
Noxafil
Nozinan130
Nozinan (Swiss)130
Nozinan S29130
Nucala
Nuelin
Nuelin-SR
Nupentin
Nusinersen
Nutilis
Nutren Diabetes
Nutrient Modules
Nutrini Energy Multi Fibre259
Nutrini Energy Multi Fibre
Nutrini Energy RTH
Nutrini Low Energy Multi Fibre
Nutrini Peptisorb
Nutrini Peptisorb Energy272
Nutrini RTH
Nutrison 800 Complete Multi
Fibre
Nutrison Advanced Diason 257
Nutrison Advanced Peptisorb
Nutrison Concentrated266
Nutrison Energy264
Nutrison Energy Multi Fibre 264
Nutrison Multi Fibre264
Nutrison Standard RTH
Nyefax Retard52
Nystatin
Alimentary33
Genito-Urinary76
Infection
NZB Low Gluten Bread Mix 267
-0-
Obinutuzumab
Obstetric Preparations
Ocicure
Ocrelizumab134
Ocrevus
Octocog alfa [Recombinant factor
VIII] (Advate)
•] (/

Octocog alfa [Recombinant factor VIII] (Kogenate FS)41
Optro atida
Octreotide
Octreotide Depot Teva 165
Octreotide GH165
Octreotide long-acting 165
Oestradiol
Oestradiol valerate82
Oestradiol with norethisterone83
Oestriol
Genito-Urinary77
Hormone
Oestrogens82
Ofev
Oil in water emulsion
Olanzapine
Olaparib
Olbetam
Olbetam S29
Olopatadine
Olopatadine Teva
Olsalazine
Omalizumab202
Omeprazole9
Omeprazole actavis 109
Omeprazole actavis 209
Omeprazole actavis 409
Omnitrope84
Omnitrope S2984
Onbrez Breezhaler
Oncaspar LYO154
OncoTICE174
Ondansetron129
Ondansetron ODT-DRLA 129
One-Alpha
One-Alpha S29
Onrex
Opdivo
Ora-Blend
Ora-Blend SF252
Ora-Plus
Ora-Sweet
Ora-Sweet SF252
Orabase
Oral and Enteral Feeds
Oralcon 30 ED75
Oratane62
Ordine122
Orgran
Ornidazole 100
Orphenadrine citrate116
Ortho-tolidine
Oruvail SR 111
Osmolite RTH264
Other Endocrine Agents
Other Oestrogen Preparations83

Other Progestogen
Preparations 83
Other Skin Preparations71
Ovestin
Genito-Urinary77
Hormone
Oxaliplatin146
Oxaliplatin Accord146
Oxaliplatin Actavis 100 146
Oxaliplatin Ebewe146
Oxis Turbuhaler237
Oxpentifylline
Oxybutynin78
Oxycodone hydrochloride122
Oxycodone Sandoz 122
OxyNorm
Oxytocin77
Oxytocin BNM77
Oxytocin GH77
Oxytocin with ergometrine
maleate77
Ozurdex245
- P -
Pacifen116
Pacimol120
Paclitaxel
Paclitaxel Actavis154
Paclitaxel Ebewe 154
Paediatric Seravit
Palbociclib 160
Paliperidone132
Paliperidone palmitate 132
Palivizumab
Pamidronate disodium113
Pamisol113
Pamol
Pancreatic enzyme25
Pantoprazole
Panzop Relief
Panzytrat
Papaverine hydrochloride58
Para-amino salicylic acid 101
Paracare 120
Paracare Double Strength 120
Paracetamol 120
Paracetamol (Ethics) 120
Paracetamol + Codeine
(Relieve)
Paracetamol with codeine
Paraffin
Paraffin liquid with wool fat
Parasiticidal Preparations
Parnate
Parnate S29124
Paromomycin
Paroxetine

Paser 101
Paxam 133
Paxlovid 105
Pazopanib161
Peak flow meter243
Pedialyte - Bubblegum46
Pediasure259
Pediasure Plus259
Pediasure RTH259
Pegaspargase 154
Pegasys108
Pegfilgrastim45
Pegylated interferon alfa-2a 108
Pembrolizumab230
Pemetrexed149
Penicillamine 112
Penicillin G93
PenMix 3011
PenMix 5011
Pentasa7
Pentostatin [Deoxycoformycin] 154
Pentoxifylline [Oxpentifylline]58
Peptamen Junior
Pepti-Junior
Peptisorb
Perhexiline maleate
Pericyazine 131
Perindopril
Periset
Perjeta
Permethrin
Perrigo
Pertuzumab
Peteha 101
Pethidine hydrochloride 123
Pevaryl
Pexsig
Pfizer Exemestane 167
Pharmacy Health Sorbolene with
Glycerin 66
Pharmacy Services
Pheburane
Phenasen149
Phenobarbitone 126
Phenobarbitone sodium
Extemporaneous252
Nervous
Phenoxybenzamine
hydrochloride 47
Phenoxymethylpenicillin (Penicillin
V)
Phenytoin sodium125–126
Phillips Milk of Magnesia
Phlexy 10
Phosphate Phebra46
Phosphorus46

Phytomenadione	44
Pilocarpine hydrochloride	247
Pimafucort	65
Pimecrolimus	
Pine tar with trolamine laurilsulfate	
	70
and fluorescein	
Pinetarsol	
Pioglitazone	12
Pirfenidone	240
Pizotifen	
PKU Anamix Infant	
PKU Anamix Junior	
PKU Anamix Junior Chocolate	269
PKU Anamix Junior LQ	269
PKU Anamix Junior Orange	269
PKU Anamix Junior Vanilla	
PKU Lophlex LQ 10	
	209
PKU Lophlex LQ 20	
PKU Lophlex Powder	
PKU Lophlex Sensation 20	269
Plaquenil	. 112
Plendil ER	
Pneumococcal (PCV10) conjugate	52
vaccine	. 281
Pneumococcal (PCV13) conjugate	
vaccine	. 282
Pneumococcal (PPV23)	
polysaccharide vaccine	202
	. 200
Pneumovax 23	
Podophyllotoxin	
Polaramine	
Poliomyelitis vaccine	283
Poloxamer	
Poly-Gel	
Poly-Tears	
Poly-Visc	
Polycal	254
Polyethylene glycol 400 and	
propylene glycol	248
Ponstan	
Posaconazole	
Posaconazole Juno	
Postinor-1	
Potassium chloride4	5-46
Potassium citrate	
Potassium iodate	35
Povidone iodine	
Pradaxa	
Pramipexole hydrochloride	117
Pravastatin	55
Pravastatin Mylan	
Pravastatin Viatris	
Praziguantel	
Prazosin	
Pred Forte	
Prednisolone	<mark>81</mark>

Prednisolone acetate	246
Prednisolone sodium	8
Prednisolone sodium	
phosphate	246
Prednisolone-AFT	246
Prednisone	
Prednisone Clinect	81
Pregabalin	
Pregabalin Pfizer	126
Pregnancy Tests - hCG Urine	77
Premarin	
Prevenar 13	
Priadel	
Primaquine	100
Primidone	126
Primidone Clinect	126
Primolut N	120
Priorix	
Probenecid	
Probenecid-AFT	
Procarbazine hydrochloride	154
Prochlorperazine	104
Proctofoam	129
Proctosedyl	
Procyclidine hydrochloride Progesterone	
Proglicem Proglycem	
Progynova	10
Prolia	
Promethazine hydrochloride	
Propafenone hydrochloride	
Propamidine isethionate	
Propranolol Propylene glycol	
Propylthiouracil	
Prostacur	
Protaphane	10
Protaphane Penfill	
Protifar	
Protionamide Provera	101
Provera HD	03
Psoriasis and Eczema	~~~
Preparations	69
PTU	83
Pulmicort Turbuhaler	237
Pulmozyme	241
Puri-nethol	
Puria	
Pyrazinamide	
Pyridostigmine bromide	
Pyridoxine hydrochloride	
Pyridoxine multichem	
Pyrimethamine	
Pytazen SR	41

- Q -
Quetapel131
Quetiapine131
Quick-Set MMT-39224
Quick-Set MMT-39324
Quinapril
Quinapril with
hydrochlorothiazide
Qvar
- B -
RA-Morph
Ralicrom
Raloxifene hydrochloride 113
Raltegravir potassium
Ramipex 117
Ramipril
Ranbaxy-Cefaclor
Ranbaxy-Cefaclor S29
Rapamune
Rasagiline117
Reandron 1000
Recombinant factor IX
Recombinant factor VIIa40 Recombinant factor VIII40–41
Rectogesic
Redipred81 Relieve111
Relieve
Remicade
Renilon 7.5
Resonium-A
Respigen
Respiratory Devices
Respiratory Stimulants
Retinol palmitate
ReTrieve
Revlimid
Revolade
Ribomustin
Ribornusuri
Rifabutin
Rifadin
Rifampicin101 Rifaximin10
Rifinah
Rilutek
Riluzole
RINVOQ
Riodine
Risedronate Sandoz
Risedronate sodium
Risperdal Consta
Risperidone 131–132

Risperidone (Teva)	131
Risperon	. 131
Ritalin	. 138
Ritalin LA	
Ritonavir	
Rituximab (Mabthera)	
	204
Rituximab (Riximyo)	
Rivaroxaban	
Rivastigmine	
Rivastigmine Patch BNM 10	
Rivastigmine Patch BNM 5	
Rivotril	125
Riximyo	206
RIXUBIS	40
Rizamelt	128
Rizatriptan	128
RoActemra S29	
Ronapreve	191
Ropin	117
Ropinirole hydrochloride	117
Rosuvastatin	56
Rosuvastatin Viatris	56
Rotarix	
Rotavirus oral vaccine	200
Roxane	
Roxane-Propranolol	
Roxithromycin	00 00
Rubifen	120
Rubifen SR	138
Rugby Capsaicin Topical Cream	
Rugby Capsaicin Topical Cream Musculoskeletal	112
Rugby Capsaicin Topical Cream Musculoskeletal Nervous	112 119
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina	112 119 nt
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII]	112 119 nt 41
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib	112 119 nt 41 162
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan	112 119 nt 41 162 49
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rythmodan	112 119 nt 41 162 49
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rythmoorm	112 119 nt 41 162 49 50
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rythmonorm	112 119 nt 41 162 49 50 127
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rythmonorm	112 119 nt 41 42 49 50 127 48
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rytmonorm	112 119 nt 41 42 49 50 127 48 157
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rythmonorm	112 119 nt 41 42 49 50 127 48 157
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rytmonorm - S - Sabril Sacubitril with valsartan Sagent SalAir	112 119 nt 41 162 49 50 127 48 157 238
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rytmonorm - S - Sabril Sacubitril with valsartan Sagent SalAir Salazopyrin	112 119 nt 41 162 49 50 127 48 157 238 8
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rythmodan Rythmonorm - S - Sabril Sacubitril with valsartan Sagent SalAir Salazopyrin Salazopyrin	112 119 nt 41 49 50 127 48 157 238 8
Rugby Capsaicin Topical Cream Musculoskeletal Nervous	112 119 nt 41 49 50 127 48 157 238 8
Rugby Capsaicin Topical Cream Musculoskeletal Nervous Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rythmodan Rythmodan Sabril Sabril Sabril Sacubitril with valsartan Sagent Salazopyrin Salazopyrin EN Salazopyrin EN Salbutamol Salbutamol with ipratropium	112 119 nt 41 40 50 127 48 48 8
Rugby Capsaicin Topical Cream Musculoskeletal Nervous	112 119 nt 41 42 49 50 127 48 157 238
Rugby Capsaicin Topical Cream Musculoskeletal Nervous	112 119 nt 41 41 49 50 48 8 8 8 8 238 239 70
Rugby Capsaicin Topical Cream Musculoskeletal Nervous. Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rytmonorm - S - Sabril Sacubitril with valsartan. Sagent SalAir. Salazopyrin Salazopyrin EN Salbutamol Salbutamol with ipratropium bromide. Salicylic acid.	112 119 nt 41 162 50 127 48 127 238 238 239 70 237
Rugby Capsaicin Topical Cream Musculoskeletal Nervous. Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rytmonorm - S - Sabril Sacubitril with valsartan. Sagent Salazopyrin Salazopyrin EN Salazopyrin EN Salbutamol.	112 119 nt 41 162 102 102 102 49 102 127 238 238 239 70 237 128
Rugby Capsaicin Topical Cream Musculoskeletal Nervous. Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rytmonorm - S - Sabril Sacubitril with valsartan. Sagent Salazopyrin Salazopyrin EN Salazopyrin EN Salbutamol Salbutamol. Salbutamol Salicylic acid Salmeterol Sandonigran Sanofi Primaquine	112 119 nt 41 49 50 127 48 238 238 238 238 239 70 237 128 99
Rugby Capsaicin Topical Cream Musculoskeletal Nervous. Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rytmonorm - S - Sabril Sacubitril with valsartan. Sagent Salazopyrin Salazopyrin EN Salazopyrin EN Salbutamol. Salbutamol. Salbutamol. Salbutamol. Salbutamol. Salbutamol. Salbutamol. Salicylic acid Salmeterol. Sanofi Primaquine Sapropterin dihydrochloride	112 119 nt 41 40 50 127 48 238 8 238 238 239 70 237 128 99 31
Rugby Capsaicin Topical Cream Musculoskeletal Nervous. Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rytmonorm - S - Sabril Sacubitril with valsartan. Sagent Salazopyrin Salazopyrin EN Salazopyrin EN Salazopyrin EN Salbutamol. Salbutam	112 119 nt 41 50 127 48 50 238 238 238 239 70 237 237
Rugby Capsaicin Topical Cream Musculoskeletal Nervous. Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitnib Rythmodan Rytmonorm - S - Sabril Sacubitril with valsartan. Sagent Salazopyrin Salazopyrin EN Salazopyrin EN Salbutamol.	112 119 nt 41 50 127 48 8 8 8 8 8
Rugby Capsaicin Topical Cream Musculoskeletal Nervous. Rurioctocog alfa pegol [Recombina factor VIII] Ruxolitinib Rythmodan Rytmonorm - S - Sabril Sacubitril with valsartan. Sagent Salazopyrin Salazopyrin EN Salazopyrin EN Salazopyrin EN Salbutamol. Salbutam	112 119 nt 41 49 50 48 48 48 48 48 238 70 237 71 71 71 71

Sedatives and Hypnotics 135
Seebri Breezhaler239
Selegiline hydrochloride117
Senna
Senokot27
SensoCard15
Serc 128
Serenace
Seretide
Seretide Accuhaler 238
Serevent
Serevent Accuhaler
Sertraline
Setrona124
Setrona AU124
Sevredol
Sex Hormones Non
Contraceptive
Shield XL
Shingles vaccine
Shingrix
SII-Onco-BCG
Sildenafil
Silhouette MMT-373
Siltuximab
Simvastatin
Simvastatin Mylan
Simvastatin Viatris
Sinemet
Sinemet CR 117
Sirolimus
Siterone
Slow-Lopresor
Smith BioMed Rapid Pregnancy
Test
Sodibic
Sodium acid phosphate
Sodium alginate
Sodium benzoate
Sodium bicarbonate
Blood
Extemporaneous
Sodium calcium edetate
Sodium chloride
Blood
Respiratory
Sodium citrate with sodium lauryl
sulphoacetate
Sodium citro-tartrate
Sodium cromoglicate
Alimentary
Sensory
Sodium Fusidate [fusidic acid]
Dermatological
Infection
Sensory244
,

Sodium hyaluronate [Hyaluronic	
acid]24	8
Sodium phenylbutyrate	1
Sodium picosulfate2	7
Sodium polystyrene sulphonate4	6
Sodium tetradecyl sulphate4	
Sodium valproate12	7
Sofradex	, л
Soframycin24	
Solgar	5
Solifenacin Mylan7	2
Solifenacin succinate7	0
Solifenacia Vietrie	0
Solifenacin Viatris	
Solu-Cortef8	
Solu-Medrol	0
Solu-Medrol-Act-O-Vial8	0
Somatropin (Omnitrope)8	
Sotalol5	
Spacer device24	
Span-K4	6
Spinal Muscular Atrophy13	6
Spinraza13	6
Spiolto Respimat24	0
Spiractin5	4
Spiriva23	9
Spiriva Respimat23	9
Spironolactone5	
Sporanox9	8
Sprycel15	7
Stelara22	
Stemetil12	9
Steril-Gene8	
SteroClear	
Stesolid	
Stimulants/ADHD Treatments	6
Stiripentol	
Stocrin	6
Stomahesive	0
Strattera	
Strides Shasun	1
Stromectol6 Sucralfate1	
Sulfadiazine Silver	
Sulfadiazine sodium9	6
Sulfasalazine	
Sulphur	0
Sulprix12	
Sumagran 12	
Sumatriptan 12	8
Sunitinib16	2
Sunitinib Pfizer16	
Sunscreens7	
Sunscreens, proprietary7	1
Sure-T MMT-8632	1
Sure-T MMT-873 2	1
Survimed OPD26	1

Sustagen Hospital Formula	264
Sustagen Hospital Formula	
Sustagen Hospital Formula Active	264
Sustanon Ampoules	01
Sustantin Ampoules	010
Sylvant	210
Symbicort Turbuhaler 100/6	238
Symbicort Turbuhaler 200/6	238
Symbicort Turbuhaler 400/12	238
Symmetrel	117
Sympathomimetics	57
Synacthen	
Synacthen Depot	
Synacthene Retard	
	01
Synagis	203
Synflorix	
Synthroid	
Syntometrine	
Syrup (pharmaceutical grade)	253
Systane Unit Dose	248
-T-	
Tacrolimus	
Dermatological	70
Opening	70
Oncology	233
Tacrolimus Sandoz	233
Taliglucerase alfa	32
Tambocor	49
Tamoxifen citrate	. 166
Tamoxifen Sandoz	. 166
Tamsulosin hydrochloride	
Tamsulosin-Rex	
Tandem Cartridge	
Tandem t:slim X2 with Basal-IQ	16
Tap water	253
Taro	
Taro-Testosterone	
Tasigna	. 160
Tasmar	117
Taurine	
Tecfidera	
Tegretol	125
Tegretol CR	105
Telfast	
Teligent	95
Temaccord	
Temazepam	
Temozolomide	. 154
Tenofovir disoproxil	102
Tenofovir Disoproxil Emtricitabine	
Mylan	104
Tenofovir Disoproxil Emtricitabine	. 104
	40.4
Viatr	. 104
Tenofovir Disoproxil Mylan	
Tenoxicam	
Tensipine MR10	52
Tepadina	
Terbinafine	

T 1 1 1 1 1	~~~
Terbutaline sulphate	
Teriflunomide	135
Teriparatide	114
Testosterone	. 81
Testosterone cipionate	. 81
Testosterone esters	. 81
Testosterone undecanoate	. 81
Tetrabenazine	118
Tetrabromophenol	78
Tetracosactrin	
Tetracycline	
Teva Lisinopril	47
Thalidomide	
Thalomid	155
Theophylline	241
Thiamine hydrochloride	
Thiamine multichem	34
THIO-TEPA	146
Thioguanine	149
Thiotepa	146
Thyroid and Antithyroid Agents	.83
Ticagrelor	41
Ticagrelor Sandoz	41
Tilcotil	
Timolol	
	240
Timoptol XE	
Tiotropium bromide	239
Tiotropium bromide with	
olodaterol	240
Tivicay	107
Tixagevimab with cilgavimab	219
TMP	. 97
Tobramycin	
Infection	. 96
Sensory	244
Tobramycin BNM	96
Tobramycin Mylan	
Tobrex	244
Tocilizumab	
Tofranil	123
Tolcapone	117
Tolvaptan	54
Topamax	127
Topical Products for Joint and	
Muscular Pain	
Topiramate	127
Topiramate Actavis	127
Total parenteral nutrition (TPN)	. 45
TPN	45
Tramadol hydrochloride	122
Tramal SR 100	
Tramal SR 150	
Tramal SR 200	
Trandate	
Tranexamic acid	
Tranylcypromine sulphate	124

Trastuzumab2	20
Trastuzumab emtansine	
Travatan2	
Travoprost2	47
Treatments for Dementia1	40
Treatments for Substance	
Dependence 1	41
Trental 400	58
Tretinoin	
Dermatological	62
Oncology1	
Trexate	
Triamcinolone acetonide	-0
Alimentary	იი
Allifiendary	00
Dermatological	
Hormone	81
Triamcinolone acetonide with	
gramicidin, neomycin and nystatin	
Dermatological	65
Sensory2	44
Triazolam1	
Trimethoprim	97
Trimethoprim with	
sulphamethoxazole	
[Co-trimoxazole]	97
Trisequens	87 87
Trisul	
Trophic Hormones	
Tropicamide2	47
Trulicity	12
Trusopt2	
TruSteel	
Tryzan	48
Tuberculin PPD [Mantoux] test2	84
Tubersol2	84
Two Cal HN2	66
Tykerb1	59
Tysabri1	
- U -	
UK Synacthen	81
Ultibro Breezhaler2	40
Ultraproct	.8
Umeclidinium2	39
Umeclidinium with vilanterol2	40
Univent	
Upadacitinib2	22
Ural2	
Urea	
Urex Forte	
Urinary Agents	11
Urinary Tract Infections1	
Urinorm1	
Uromitexan1	
Ursodeoxycholic acid	
Ursosan	
Ustekinumab2	25

Jtrogestan	83
/accinations	075
aclovir	
/alaciclovir	
/alganciclovir	
/alganciclovir Mylan	07
/ancomycin	
/annair /arenicline Pfizer	230
/arenicline tartrate /aricella vaccine [Chickenpox	142
vancella vaccine [Chickenpox vaccine]	004
vaccinej	284
Varicella zoster vaccine [Shingles	~~ 4
vaccine]	284
Varicella zoster virus (Oka strain) liv	e
attenuated vaccine [shingles	
vaccine]	
/arious	
/arivax	
/asodilators	58
asopressin Agonists	
/asorex	
/ebulis	
/edafil	
/edolizumab	
/eletri	
/enclexta	
/enetoclax	156
/enlafaxine	
/enomil235-	-236
/ENOX	
/entolin	
/epesid	151
/erapamil hydrochloride	
/ermox	
/ersacloz	
/esanoid	
/exazone	
/fend	99
/iaderm KC	65
Viatris	
Cardiovascular	
Infection96,	
/ictoza	
/igabatrin	
/igisom	
/ildagliptin	12
/ildagliptin with metformin	
hydrochloride	
/impat	
/inblastine sulphate	
/incristine sulphate	
/inorelbine	
/inorelbine Ebewe	
Viramune Suspension	106

ViruPOS244
Vit.D3
VitA-POS
Vitabdeck35
Vital
Vitamin B complex
Vitamin B6 25
Vitamins
Vivonex TEN
Voltaren 111
Voltaren D111
Voltaren Ophtha245
Voltaren SR 111
Volumatic243
Voriconazole
Votrient
Vttack
- W -
Warfarin sodium
Wart Preparations71
Wasp venom allergy treatment
Water
Blood
Extemporaneous253
White Soft Liquid Paraffin AFT
Wool fat with mineral oil
- X -
- X - Xarelto
Xifaxan
XMET Maxamum
Xolair
Xylocaine
Xylocaine 2% Jelly118
Xyntha 40
- Z - Zapril
Zarontin
Zaroxolyn
Zavedos
Zeffix
Zematop
Zetlam
Ziagen
Zidovudine [AZT] 107
Zidovudine [AZT] with
lamivudine 107
Ziextenzo45
Zimybe57
Zinc and castor oil66
Zinc sulphate
Zincaps
Zinnat91
Ziprasidone131
Zista236
Zithromax91

Zoledronic acid	
Hormone	80
Musculoskeletal	115
Zoledronic acid Mylan	80
Zoledronic acid Viatris	
Hormone	<mark>80</mark>
Musculoskeletal	115
Zoledronic-US	115
Zopiclone	136
Zopiclone Actavis	136
Zostavax	
Zostrix	112
Zostrix HP	
Zuclopenthixol decanoate	133
Zuclopenthixol hydrochloride.	131
Zusdone	131
Zyban	141
Zypine	131
Zypine ODT	
Zyprexa Relprevv	
Zytiga	164



