April 2023 Volume 30

Editors:

Kaye Wilson, Doris Chong & Ayeshah Khan email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street

PO Box 10 254 Wellington

Freephone Information Line

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz
@Pharmaceutical Management Agency
ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	,	
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	37
	Cardiovascular System	47
	Dermatologicals	62
	Genito Urinary System	72
	Hormone Preparations – Systemic	79
	Infections – Agents For Systemic Use	90
	Musculoskeletal System	111
	Nervous System	117
	Oncology Agents & Immunosuppressants	144
	Respiratory System & Allergies	237
	Sensory Organs	247
	Various	252
Section C	Extemporaneous Compounds (ECPs)	254
Section D	Special Foods	257
Section I	National Immunisation Schedule	278

Introducing Pharmac

Index

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
Antacids and Antiflatulents			
Antacids and Reflux Barrier Agents			
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg posachet		30	✓ Gaviscon Infant
SODIUM ALGINATE Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (13.61)	60	Gaviscon Extra Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 ml	nl Acidex
Phosphate Binding Agents			
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) –	12.56	100	✓ Alu-Tab
Subsidy by endorsement		500 ml 173 ml	
Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed according		s or w	
Antidiarrhoeals			
Agents Which Reduce Motility			
* Tab 2 mg* Cap 2 mg	10.75	400 400	✓ Nodia✓ Diamide Relief
Rectal and Colonic Anti-inflammatories			
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy	166.50	90	✓ Entocort CIR
■ SA1886 Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant pract the following criteria:	titioner. Approvals va	alid for	or 6 months for applications meeting

1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and

2 Any of the following:

2.1 Diabetes; or

continued...

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	•	Manufacturer

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes: or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture: or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

THE RECORD TO LIKE		
Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP	✓ Colifoam✓ Cortifoam \$29
	21.1 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
OLSALAZINE	<u> </u>			Thailada a
Tab 500 mg	56.02	60	1	Atnahs
				Olsalazine S29
	93.37	100	1	Dipentum
Cap 250 mg	53.00	100	✓	Dipentum
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	/	Essential
				Prednisolone S29
SODIUM CROMOGLICATE				
Cap 100 mg	113.35	100	✓	Ralicrom
SULFASALAZINE				
* Tab 500 mg	16.52	100	1	Salazopyrin
* Tab EC 500 mg	17.86	100	/	Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE				
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and				
cinchocaine hydrochloride 5 mg per g11.06	30 g OP	Ultraproct		
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and				
cinchocaine hydrochloride 1 mg7.30	12	 Ultraproct 		
HYDROCORTISONE WITH CINCHOCAINE		-		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl		
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl		

Management of Anal Fissures

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available on	а		
PSO	19.00	5	✓ Robinul
	65.45	10	✓ Max Health
(Max Health Inj 200 mcg per ml, 1 ml ampoule to be delisted 1 Se	eptember 2023)		
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO	6.35	5	✓ Buscopan
			✓ Buscopan S29 S29
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement.......14.58 14 ✓ Klacid

- a) Maximum of 28 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

FΑ	MOTIDINE - Only on a prescription			
*	Tab 20 mg	4.91	100	✓ Famotidine
	·			Hovid S29
*	Tab 40 mg	8.48	100	✓ Famotidine
	•			Hovid \$29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement	57.02	10	✓ Mylan S29
	Subsidy by andorsement - Subsidised for nations receiving treatment as part of nalliative care			

Proton Pump Inhibitors

LA	NSOPRAZOLE		
*	Cap 15 mg4.2	0 100	✓ Lanzol Relief
*	Cap 30 mg5.2	6 100	✓ Lanzol Relief
ON	MEPRAZOLE		
	For omeprazole suspension refer Standard Formulae, page 254		
*	Cap 10 mg	4 90	✓ Omeprazole actavis 10
*	Cap 20 mg	6 90	✓ Omeprazole actavis 20
*	Cap 40 mg	1 90	✓ Omeprazole actavis 40
*	Powder — Only in combination	0 5 g	✓ Midwest
*	Inj 40 mg ampoule with diluent37.3	8 5	✓ <u>Dr Reddy's</u> Omeprazole
			✓ Ocicure S29
PΑ	NTOPRAZOLE		
*	Tab EC 20 mg	9 90	✓ Panzop Relief
*	Tab EC 40 mg2.7	4 90	✓ Panzop Relief
~	Panzop Relief to be Principal Supply on 1 July 2023	7 30	- I diizop Hellel

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mgSUCRALFATE	14.51	50	✓	Gastrodenol S29
Tab 1 g	35.50 (48.28)	120		Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail pha Tab 550 mg	•	56	/	<u>Xifaxan</u>
▶ SA1461 Special Authority for Subsidy Initial application only from a gastroenterologist, hepatologist on the patologist. Approvals valid for 6 months where the patient hat tolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Practition the patologist. Approvals valid without further renewal unless not be nefiting from treatment.	as hepatic encephalop oner on the recomme	athy d	espite an n of a gast	adequate trial of maximum roenterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE – Special Authority see SA1320 below – Retail pha Cap 25 mg Cap 100 mg Oral liq 50 mg per ml	110.00 280.00	100 100 0 ml 0	✓)P ✓	Proglicem \$29 Proglicem \$29 Proglycem \$29 e5 Pharma \$29
■ SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid withou appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit — Up to 5 kit available on a PSO	t further renewal unle		fied where	
Insulin - Short-acting Preparations				
INSULIN NEUTRAL ▲ Inj human 100 u per ml		0 ml C	√ ✓	Actrapid Humulin R Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
INSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	/	NovoMix 30 FlexPen

	Subsidy (Manufacturer's P	rico) Subci	Fully Brand or dised Generic
	(Wanuacturers F	Per	✓ Manufacturer
INSULIN ISOPHANE			
▲ Inj human 100 u per ml	17.68	10 ml OP	✓ Humulin NPH✓ Protaphane
▲ Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH✓ Protaphane Penfill
INSULIN ISOPHANE WITH INSULIN NEUTRAL			· i i otapilano i onimi
▲ Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
,			✓ Mixtard 30
▲ Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
			✓ PenMix 30✓ PenMix 50
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,			
3 ml		5	✓ Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,			
3 ml	42.66	5	Humalog Mix 50
Insulin - Long-acting Preparations			
INSULIN GLARGINE ▲ Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
▲ Inj 100 u per ml, 3 ml		5	✓ Lantus
▲ Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
ilisuilii - napiu Actilig Freparations			
INSULIN ASPART			_
▲ Inj 100 u per ml, 10 ml		1	✓ NovoRapid
▲ Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
▲ Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
NSULIN GLULISINE ▲ Inj 100 u per ml, 10 ml	27.02	4	√ Anidro
▲ Inj 100 u per ml, 3 ml		1 5	✓ Apidra✓ Apidra
▲ Inj 100 u per ml, 3 ml disposable pen		5	✓ Apidra SoloStar
NSULIN LISPRO			F · · · · · · · · ·
▲ Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ Humalog
▲ Inj 100 u per ml, 3 ml		5	✓ Humalog
Alpha Glucosidase Inhibitors			
ACARBOSE			
# Tab 50 mg	8 95	90	✓ Accarb
* Tab 100 mg		90	✓ Accarb
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE	7.50	400	/ Decarl
* Tab 5 mg	7.50	100	✓ <u>Daonil</u>
GLICLAZIDE	45.40	500	/ Oll-1-1-
* Tab 80 mg	15.18	500	✓ Glizide
GLIPIZIDE	4.50	100	✓ Minidiah
* Tab 5 mg	4.58	100	✓ <u>Minidiab</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg	14.74	1,000		Metformin Mylan Metformin Viatris
* Tab immediate-release 850 mg(Metformin Mylan Tab immediate-release 500 mg to be delisted		500	•	Metformin Mylan
PIOGLITAZONE				
* Tab 15 mg	6.80	90	✓	Vexazone
* Tab 30 mg		90	✓	Vexazone
* Tab 45 mg	12.25	90	1	Vexazone
VILDAGLIPTIN				
Tab 50 mg	35.00	60	✓	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	1	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride		60	1	Galvumet

GLP-1 Agonists

DULAGLUTIDE - Special Authority see SA2065 below - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

* Inj 1.5mg per 0.5 ml prefilled pen115.23 ✓ Trulicity

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*: or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

LIRAGLUTIDE - Special Authority see SA2187 on the next page - Retail pharmacy

a) Maximum of 9 inj per prescription

✓ fully subsidised

- b)
- a) Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.
- ✓ Victoza

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

⇒SA2187 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial Special Authority approval for either an SGLT-2 inhibitor or GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*: or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*: or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*: or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

 a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack,

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 10 mg58.56	30	Jardiance
*	Tab 25 mg	30	 Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	8.56	60	✓ Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips.......20.00 1 OP ✓ CareSens Dual

			_
Subsidy	Fully	Brand or	Ī
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips26	6.20	50 test OP	SensoCard
-----------------------------	------	------------	-----------

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

IIVC	ocina i cia necescipa	OH		
*	29 g × 12.7 mm	10.95	100	✓ B-D Micro-Fine
*	31 g × 5 mm		100	✓ B-D Micro-Fine
*	31 g × 6 mm		100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED	LE - Maximum of 2	:00 dev per p	prescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g x 8 mm needle	13.56	100	B-D Ultra Fine II
		1.36	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓ B-D Ultra Fine II
		1.36	10	
		(1.99)		B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year pe	riod.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			Y2 with Rasal-IO

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	✓	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist: or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Fither
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	I Generic
\$	Per 🗸	Manufacturer

continued...

- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

1 OP

1 OP

✓ Sure-T MMT-863

✓ Sure-T MMT-873

ALIMENTARY TRACT AND METABOLISM					
	Subsidy (Manufacturer's Price \$	e) S Per	Fully Subsidised	Brand or Generic Manufacturer	
continued					
pump therapy; and 4 The patient is continuing to derive benefit from pump the 5 The patient had achieved and is maintaining a HbA1c of 6 The patient has had no increase in severe unexplained 7 The patient's HbA1c has not deteriorated more than 5 m 8 Either: 8.1 Applicant is a relevant specialist; or	f equal to or less than hypoglycaemic episoo nmol/mol from baselin	des from			
8.2 Applicant is a nurse practitioner working within the Renewal — (Previous use before 1 September 2012) only from the control of the contro		lict or nu	rco proctiti	oner Approvale valid for G	
years for applications meeting the following criteria:	oni a relevant specia	iist oi Tiu	ise praciii	orier. Approvais valid for 2	
All of the following:					
1 The patient is continuing to derive benefit according to the	he treatment plan and	l has ma	intained a	HbA1c of equal to or less	
than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 m	amol/mol from initial a	nnlicatio	n: and		
3 The patient has not had an increase in severe unexplair				ne; and	
4 Either:	7, 3,			.,	
4.1 Applicant is a relevant specialist; or					
4.2 Applicant is a nurse practitioner working within the	neir vocational scope.				
INSULIN PUMP CARTRIDGE - Special Authority see SA1985	on page 19 - Retail	pharmac	:y		
a) Maximum of 3 sets per prescription					
b) Only on a prescriptionc) Maximum of 13 packs of cartridge sets will be funded p	or voar				
Cartridge 300 U, t:lock × 10	50.00	1 OP	√ T	andem Cartridge	
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Specia		85 on pa	ge 19 – Re	etail pharmacy	
a) Maximum of 3 sets per prescription	,	•		, ,	
b) Only on a prescription					
c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10	120.00	1 OP	./ N	IiniMed Sure-T	
10 min steer needle, 60 cm tubing x 10	130.00	TOP	• IV	MMT-884A	
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T	
				MMT-886A	
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T	
				MMT-864A	
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T MMT-866A	
8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ N	IiniMed Sure-T	
o min otoo hoodio, oo on tabing x 10		1 01	- 14	MMT-874A	
8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T	
				MMT-876A	

6 mm steel needle; 29 G; manual insertion; 60 cm tubing ×

8 mm steel needle; 29 G; manual insertion; 60 cm tubing ×

Subsidy	H	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
 \$	Per	✓	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles		1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line × 10 with	130 00	1 OP	✓ TruSteel

1 OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

13 mm teflon needle, 45 cm tubing × 10130.00	1 OP
13 mm teflon needle, 60 cm tubing × 10130.00	1 OP

6 mm teflon needle, 80 cm blue tubing.......130.00

6 mm teflon needle, 80 cm clear tubing × 10130.00

- ✓ MiniMed Silhouette MMT-382A
- ✓ MiniMed Silhouette MMT-368A
 ✓ MiniMed Silhouette
- MMT-381A

 MiniMed Silhouette

✓ MiniMed Silhouette

- MMT-383A ✓ MiniMed Silhouette MMT-377A
- MMT-378A ✓ MiniMed Silhouette MMT-384A
- 6 mm teflon needle, 110 cm tubing × 10
 10 P

 6 mm teflon needle, 45 cm blue tubing × 10
 130.00
 1 OP
- ✓ MiniMed Quick-Set MMT-398A
 ✓ MiniMed Mio MMT-941A
- ✓ MiniMed Mio MMT-921A
 - ✓ MiniMed Mio MMT-943A
 - ✓ MiniMed Mio MMT-923A
 - ✓ MiniMed Quick-Set MMT-399A
 - ✓ MiniMed Mio MMT-945A
 - ✓ MiniMed Mio MMT-965A
 - ✓ MiniMed Mio MMT-925A
 - ✓ MiniMed Quick-Set MMT-387A
 - ✓ MiniMed Quick-Set MMT-396A
 - ✓ MiniMed Quick-Set MMT-397A
 - ✓ MiniMed Mio MMT-975A
 - ✓ MiniMed Quick-Set MMT-386A

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price	e) Sub	Fully	Brand or Generic
	\$	Per	1	Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	NSERTION WITH II	NSERTION	I DEVICE)	- Special Authority see
13 mm teflon cannula; angle insertion; insertion device; 110 line × 10 with 10 needles	cm 140.00	1 OP	√ Διι	toSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cl	n	1 OP		toSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 17 mm teflon cannula; angle insertion; 60 cm line × 10 with	NSERTION) - Spe			
10 needles; luer lock	IT INSERTION WIT	1 OP H INSERT		houette MMT-373 ICE) – Special Authority
110 cm line × 10 with 10 needles	m	1 OP		toSoft 90
line x 10 with 10 needles		1 OP		toSoft 90 toSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 c	m	1 OP		toSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH Retail pharmacy a) Maximum of 3 sets per prescription		-		
b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock	130.00	1 OP	√ Qu	ick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock		1 OP	√ Qu	ick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 c a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded per	n page 19 – Retail _I	oharmacy	-	
10 × luer lock conversion cartridges 1.8 ml for Paradigm pun Cartridge for 5 and 7 series pump; 1.8 ml × 10	ps50.00	1 OP 1 OP	✔ Mir 1	R Cartridge 1.8 niMed .8 Reservoir //MT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP		niMed 8.0 Reservoir //MT-332A

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer	
Digestives Including Enzymes					
PANCREATIC ENZYME					
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	/	Creon 10000	
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase 1,250 U protease))		100	/	Panzytrat	
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	/	Creon 25000	
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph					
Eur U)		20 g O	-	Creon Micro	
(Panzytrat Cap pancreatin (175 mg (25,000 U lipase, 22,500 U al	mylase, 1,250 U pro	otease),) to be del	isted 1 June 2023)	
URSODEOXYCHOLIC ACID - Special Authority see SA1739 be	low - Retail pharma	асу			
Cap 250 mg	32.95	100	✓	Ursosan	

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults: and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (**Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

(Ma	Subsidy anufacturer's Price \$	e) Subs	Fully idised	Brand or Generic Manufacturer
continued Renewal — (Chronic severe drug induced cholestatic liver injury months where the patient continues to benefit from treatment. Renewal — (Pregnancy/Primary biliary cholangitis) from any rele treatment remains appropriate and the patient is benefiting from treat renewal — (Total parenteral nutrition induced cholestasis) from where the paediatric patient continues to require TPN and who is benefiting in bilirubin levels.	r) from any relevant practitionement.	evant practitier. Approval	s valid t	Approvals valid for 6 for 2 years where the als valid for 6 months
Laxatives				
Bulk-forming Agents				
SPAGHULA (PSYLLIUM) HUSK - Only on a prescription		05		
* Powder for oral soln	6.00	250 g OP	✓ IV	lacro Organic Psyllium Husk
	12.20	500 g OP	✓ <u>K</u>	onsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS		05		
* Dry	6.02 (17.32)	500 g OP	N	ormacol Plus
(Normacol Plus Dry to be delisted 1 October 2023)	(17.02)		11	omacor ras
Faecal Softeners				
DOCUSATE SODIUM – Only on a prescription				
* Tab 50 mg		100		oloxyl
* Tab 120 mg DOCUSATE SODIUM WITH SENNOSIDES	3.13	100	√ <u>C</u>	<u>oloxyl</u>

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below - Retail ph	armacy		
Inj 12 mg per 0.6 ml vial	36.00	1	1	Relistor
,	246.00	7	1	Relistor

⇒SA1691 Special Authority for Subsidy

POLOXAMER – Only on a prescription Not funded for use in the ear.

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Osmotic Laxatives

GL	YCEROL		
*	Suppos 2.8/4.0 g - Only on a prescription10.39	20	✓ Lax-suppositories
			Glycerol

30 ml OP

✓ Coloxyl

	Subsidy (Manufacturer's Price) Sub	Fully	Brand or Generic
	\$	Per	√	Manufacturer
LACTULOSE – Only on a prescription				
* Oral liq 10 g per 15 ml	3.61	500 ml	✓ <u>L</u> a	aevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIO	CARBONATE AND	SODIUM C	HLORIE	DE
Powder for oral soln 13.125 g with potassium chloride 46.6 m	0,			
sodium bicarbonate 178.5 mg and sodium chloride 350.7	7 mg6.70	30	✓ M	<u>lolaxole</u>
SODIUM ACID PHOSPHATE - Only on a prescription				
Enema 16% with sodium phosphate 8%	2.50	1	✓ F	leet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE	- Only on a presci	ription		
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,				
5 ml	35.89	50	✓ M	licolette
			✓ M	licolette-S29 S29
Micolette to be Principal Supply on 1 June 2023				
Stimulant Laxatives				

*	Tab 5 mg	5.80	200	 Bisacodyl Viatris
*	Suppos 10 mg	3.69	10	✓ Lax-Suppositories
SEI	NNA - Only on a prescription			
*	Tab, standardised	2.17	100	
		(8.21)		Senokot
		0.43	20	
		(2.06)		Senokot
SO	DIUM PICOSULFATE - Special Authority see SA2053 below	v – Retail pharma	су	
	Oral soln 7.5 mg per ml	7.40	30 ml OP	Dulcolax SP Drop

⇒SA2053 Special Authority for Subsidy

BISACODYL - Only on a prescription

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA − Special Authority see SA1986 below − Retail pharmacy
Inj 50 mg vial1,142.60 1 ✓ Myozyme

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
- 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
- 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 below - Retail pharmacy

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg	CBS	50	Solgar
Powder	CBS	400 g	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE - Special Authority see SA1987 below - Retail pharmacy

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
COENZYME Q10 - Special Authority see SA2039 below - Retail	I pharmacy			
Cap 120 mg	CBS	30	✓	Solgar
Cap 160 mg	CBS	60		Go Healthy
01000				•

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
` \$ ´	Per 🗸	Manufacturer

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts: or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 b	oelow - Retail pharmacy		
Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral liq 1 g per 10 ml	CBS	118 ml	✓ Carnitor S29
Oral liq 500 mg per 10 ml		300 ml	✓ Balance

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN – Special Authority see SA2041 below – R	etail pharmacy		
Tab 100 mg	CBS	100	Country Life
Cap 100 mg	CBS	100	✓ Solgar

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see SA1989 below – Retail pharmacy
Tab soluble 100 mg.......1,452.70 30 OP ✓ Kuvan

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

continued...

- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE – Special Authority see SA1599 below	 Retail pharmacy 		
Soln 100 mg per ml	CBS	100 ml	✓ Amzoate S29

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE − Special Authority see SA1990 below − Retail pharmacy
Grans 483 mg per g......2,016.00 174 g OP

✓ Pheburane

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Gaucher's Disease

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

Difflam

	Subsidy (Manufacturer's Pric \$	ce) Sub	Fully Brand or osidised Generic Manufacturer
CARMELLOSE SODIUM WITH GELATIN AND PECTIN	*		
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	•	Orabase
	1.52	5 g OP	
	(3.60)		Orabase
Powder		28 g OP	
	(10.95)		Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
	(6.00)		Bonjela
TRIAMCINOLONE ACETONIDE			
Paste 0.1%	5.33	5 g OP	Kenalog in Orabase
On the control Authority			
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE			
Oral gel 20 mg per g	4.74	40 g OP	✓ <u>Decozol</u>
NYSTATIN			
Oral liq 100,000 u per ml	1.76	24 ml OP	✓ Nilstat
Vitamin B			
HYDROXOCOBALAMIN			
* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a	PSO2.46	3	✓ Cobal-B12 S29 ✓ <u>Hydroxocobalamin</u> <u>Panpharma</u> ✓ Vita-B12 S29
PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription			
* Tab 25 mg - No patient co-payment payable	2 70	90	✓ Vitamin B6 25
* Tab 50 mg		500	✓ Pyridoxine
Tab oo ng		000	multichem
THIAMINE HYDDOCHI ODIDE Only on a proparintion			
THIAMINE HYDROCHLORIDE - Only on a prescription * Tab 50 mg	4.65	100	✓ Thiamine multichem
· ·	4.00	100	Tillallille illulucilelli
VITAMIN B COMPLEX	7.45	500	€ Bulana
* Tab, strong, BPC	/.15	500	✓ Bplex
Vitamin C			
ASCORBIC ACID			
a) No more than 100 mg per dose			
b) Only on a prescription			
* Tab 100 mg	12.50	500	✓ Cvite
-			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pr \$		Fully dised	Brand or Generic Manufacturer
Vitamin D				
ALFACALCIDOL * Cap 0.25 mcg * Cap 1 mcg * Oral drops 2 mcg per ml CALCITRIOL * Cap 0.25 mcg * Cap 0.5 mcg	87.98 60.68 7.89	100 100 20 ml OP 100 100	✓ C	One-Alpha One-Alpha One-Alpha S29 S29 One-Alpha Calcitriol-AFT Calcitriol-AFT
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescripti * Oral liq 188 mcg per ml (7,500 iu per ml)		12 4.8 ml OP		<u>rit.D3</u> Puria
Multivitamin Preparations				
MULTIVITAMIN RENAL — Special Authority see SA1546 below — Retail pharmacy * Cap				
MULTIVITAMINS – Special Authority see SA1036 below – Retail * Powder	72.00 I without further r		notifie	•

VITAMINS

VI	AWIINS		
*	Tab (BPC cap strength)18.50	1,000	✓ Mvite
*	Cap (fat soluble vitamins A, D, E, K) - Special Authority see		
	SA1720 below – Retail pharmacy23.40	60	✓ Vitabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

Eully.

Drand or

Cubaidy

	Subsidy (Manufacturer's Price)	Subsid		Brand or Generic
	\$	Per		Manufacturer
Minerals				
Calcium				
CALCIUM CARBONATE * Tab 1.25 g (500 mg elemental) * Tab eff 1.25 g (500 mg elemental) – Subsidy by endorseme		250 100	✓ Ca	alci-Tab 500 alcium 500 mg Hexal 829
Only when prescribed for patients unable to swallow cal inappropriate and the prescription is endorsed according		ts or where	calciun	n carbonate tablets are
CALCIUM GLUCONATE * Inj 10%, 10 ml ampoule	32.00	10		ax Health - Hameln S29
	64.00	20		ax Health \$29
lodine				
POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓ <u>Ne</u>	euroTabs
Iron				
FERROUS FUMARATE * Tab 200 mg (65 mg elemental)	3.04	100	✓ <u>Fe</u>	erro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	5.98	100	✓ <u>Fe</u>	erro-F-Tabs
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental) * Oral lig 30 mg (6 mg elemental) per 1 ml		30 500 ml		errograd erodan
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority s Inj 50 mg per ml, 10 ml vial	ee SA1840 below – F	Retail pharm 1	acy	rinject
■ SA1840 Special Authority for Subsidy Initial application — (serum ferritin less than or equal to 20 months for applications meeting the following criteria:	mcg/L) from any rele	vant practiti	oner.	Approvals valid for 3

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or

 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Roth:

1 Patient continues to have iron-deficiency anaemia; and

2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE

Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%33.60	355 ml	✓ Phillips Milk of Magnesia S29
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule25.53	10	✓ <u>Martindale</u>
Zinc		
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)11.00	100	✓ Zincaps

✓ Ferrosia

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — **(chronic renal failure)** from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA1775 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe	100.00	6	Binocrit
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	Binocrit
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	Binocrit
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	Binocrit
Inj 10,000 iu in 1 ml, syringe	197.50	6	Binocrit
Inj 40,000 iu in 1 ml, syringe	250.00	1	Binocrit

_		(Manufacturer's Price) Per	Subsidised	
N	Megaloblastic				
	DLIC ACID Tab 0.8 mg	26.60	1,000	✓	Folic Acid multichem
*	Tab 5 mg	5.82	100		Folic Acid Mylan
	Oral liq 50 mcg per ml	28.82	25 ml C	_	Folic Acid Viatris Biomed

Subsidy

Fully

Brand or

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Treaters Group in conjunction with the reational reacting	orinia mariagoriiorii gro	up.	
Inj 250 iu vial	612.50	· 1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	Alprolix
ELTROMBOPAG - Special Authority see SA1743 below - Wastage claimable	Retail pharmacy		
Tab 25 mg	1,550.00	28	Revolade
Tab 50 mg	3,100.00	28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

Inj 30 mg in 1 ml vial3,570).00 1	Hemlibra
Inj 60 mg in 0.4 ml vial	3.00 1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial12,492	2.00 1	✓ Hemlibra
Inj 150 mg in 1 ml vial17,846	5.00 1	✓ Hemlibra

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Fither:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Fither:

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
- 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1.178.30	1	✓ NovoSeven RT
Inj 2 mg syringe		1	✓ NovoSeven RT
Inj 5 mg syringe	·	1	✓ NovoSeven RT
Ini 8 mg syringe	·	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FĚIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

,			
Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe	575.00	1	Xyntha
Inj 1,000 iu prefilled syringe		1	Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	Xyntha

NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	1	✓ RIXUBIS
Inj 1,000 iu vial	1	✓ RIXUBIS
Inj 2,000 iu vial	1	✓ RIXUBIS
Inj 3,000 iu vial	1	✓ RIXUBIS

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [Xpharm]

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

managed by the macmophilia meaters droup in conjuncti	on with the National i	aemopriiid	i manayemen C
Inj 250 iu vial	210.00	1	Advate
Inj 500 iu vial	420.00	1	Advate
Inj 1,000 iu vial	840.00	1	Advate
Inj 1,500 iu vial	1,260.00	1	Advate
Inj 2,000 iu vial	1,680.00	1	Advate
Inj 3,000 iu vial	2,520.00	1	Advate

	Subsidy	Fu	lly Brand or
(I	Manufacturer's Price)	Subsidise	
	\$	Per	/ Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE F	S) – [Xpharm]		
For patients with haemophilia. Rare Clinical Circumstances Br		recombinant	factor VIII. Access to funded
treatment is managed by the Haemophilia Treaters Group in co			
subject to criteria.	nijanotion with the i	adional mach	noprima Management Group,
Inj 250 iu vial	237 50	1 •	✓ Kogenate FS
Inj 500 iu vial			Kogenate FS
•		-	Kogenate FS
Inj 1,000 iu vial			•
Inj 2,000 iu vial	*		Kogenate FS
Inj 3,000 iu vial	,	1 •	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] -	[Xpharm]		
For patients with haemophilia A receiving prophylaxis treatmen		d treatment is	managed by the Haemophilia
Treaters Group in conjunction with the National Haemophilia M			3 , 1
Inj 250 iu vial	0 0 1	1 •	/ Adynovate
Inj 500 iu vial		-	/ Adynovate
Inj 1,000 iu vial		•	/ Adynovate
Inj 2,000 iu vial			Adynovate Adynovate
	2,400.00	'	Auyilovate
SODIUM TETRADECYL SULPHATE			
* Inj 3% 2 ml		5	
	(73.00)		Fibro-vein
TRANEXAMIC ACID			
Tab 500 mg	10.45	60	Mercury Pharma
Mercury Pharma to be Principal Supply on 1 June 2023		-	moroury i marma
Microary Friamia to be Frinoipal Supply of Frounce 2020			
Vitamin K			
VICAIIIII IX			
PHYTOMENADIONE			
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO	8.00	5	Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO		-	Konakion MM
		•	Ronakion illiii
Antithrombotic Agents			
And the on both Agents			
Antiplotolet Agente			
Antiplatelet Agents			
ASPIRIN			
* Tab 100 mg	1/105	990	Ethics Aspirin EC
	14.90	990	Ethics Aspirin EC
CLOPIDOGREL			
* Tab 75 mg	4.60	84	✓ Clopidogrel
			Multichem
	5.07	•	Arrow - Clopid
Arrow - Clopid to be Principal Supply on 1 May 2023			
(Clopidogrel Multichem Tab 75 mg to be delisted 1 May 2023)			
DIPYRIDAMOLE	40.00	00	.
* Tab long-acting 150 mg	13.93	60	Pytazen SR
TICAGRELOR - Special Authority see SA1955 on the next page -	Retail pharmacy		
Brand switch fee payable (Pharmacode 2653206) - see page 2	52 for details		
* Tab 90 mg		56	✓ Ticagrelor Sandoz
Ť			

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment: and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Subsidised

10

Fully

Brand or

Generic

Clexane Forte

	φ	rei	Manuacturer
Heparin and Antagonist Preparations			
ENOXAPARIN SODIUM - Special Authority see SA2152 below	- Retail pharmacy	/	
Inj 20 mg in 0.2 ml syringe	31.28	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe		10	✓ Clexane
Inj 60 mg in 0.6 ml syringe		10	✓ Clexane
Inj 80 mg in 0.8 ml syringe		10	✓ Clexane
Inj 100 mg in 1 ml syringe		10	✓ Clexane
Inj 120 mg in 0.8 ml syringe		10	Clexane Forte

Subsidy

(Manufacturer's Price)

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Low molecular weight heparin treatment is required during a patients pregnancy; or

- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner.
Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery: or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

		Subsidy		Fully	
		(Manufacturer's Price) \$	S Per	Subsidised •	Generic Manufacturer
HEPARIN SODIUM		<u> </u>	1 01		Wallatatata
Inj 1,000 iu per ml, 5 ml ampou	مار	86 11	50	1	Pfizer
Inj 5,000 iu per ml, 5 ml vial			10		Heparin Sodium
inj 3,000 ia per ini, 3 ini viai		00.00	10	•	Panpharma
Heparin Sodium Panpharn	na to be Principal Supply on	1 July 2023			
Inj 5,000 iu per ml, 1 ml			5	1	DBL Heparin
					Sodium S29
		70.33		1	Hospira
Inj 5,000 iu per ml, 5 ml ampou	ıle		50		Pfizer
Inj 25,000 iu per ml, 0.2 ml			5		Hospira
, _5,000 to pot till, 012 till till.		42.40	•		Heparin DBL S29
		482.20	50		Heparin DBL S29
(Pfizer Inj 5,000 iu per ml, 5 ml amp	noule to be delicted 1 July 2		50	•	Hepailli DDL 329
	poule to be delisted 1 July 2	020)			
HEPARINISED SALINE				_	
Inj 10 iu per ml, 5 ml		65.48	50	•	Pfizer
Oral Anticocculanta					
Oral Anticoagulants					
DABIGATRAN					
Cap 75 mg - No more than 2 of	cap per day	76.36	60	1	Pradaxa
Cap 110 mg		76.36	60	✓	Pradaxa
Cap 150 mg		76.36	60	1	Pradaxa
RIVAROXABAN					
Tab 10 mg - No more than 1 t	tah nor day	83 10	30	1	Xarelto
Tab 15 mg – Up to 14 tab avai			28		Xarelto
Tab 20 mg			28		Xarelto
· ·			_0	•	Au oito
WARFARIN SODIUM	ana mat lutanahanan - lala				
Note: Marevan and Coumadin		0.40	F0		0
* Tab 1 mg			50		Coumadin
Mr. Tale O		6.46	100		Marevan
* Tab 2 mg			50		Coumadin
* Tab 3 mg			100		Marevan
* Tab 5 mg			50		Coumadin
		11.48	100	•	Marevan
Blood Colony-stimulating	Factors				
Blood Colony-Stimulating	Taciois				
FILGRASTIM - Special Authority s					
Inj 300 mcg per 0.5 ml prefilled		96.22	10		Nivestim
In: 400 mag nor 0 E ml profilled	l aurinaa	440.50	40	./	Niira adina

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or

Inj 480 mcg per 0.5 ml prefilled syringe......148.58

continued...

10

✓ Nivestim

	BLOOD AND	BLOOD	FOR	MING ORGANS
	Subsidy (Manufacturer's Price) \$	Sub:	Fully sidised	Brand or Generic Manufacturer
continued 3 Peripheral blood stem cell mobilisation or bone marrow of 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10 to 1.5 Treatment of drug-induced prolonged neutropenia (ANC Note: *Febrile neutropenia risk greater than or equal to 20% aft European Organisation for Research and Treatment of Cancer (PEGFILGRASTIM – Special Authority see SA1912 below – Rei	0 ⁹ /L); or < 0.5 ×10 ⁹ /L). er taking into account (EORTC) guidelines.		·	·
Inj 6 mg per 0.6 ml syringe		1	_	iextenzo leulastim
Initial application only from a relevant specialist, vocationally recommendation of a relevant specialist. Approvals valid without neutropenia in patients undergoing high risk chemotherapy for the Note: *Febrile neutropenia risk greater than or equal to 5% after Organisation for Research and Treatment of Cancer (EORTC) of Fluids and Electrolytes Intravenous Administration	ut further renewal unles cancer (febrile neutrope or taking into account o	ss notified enia risk g	where reater the	used for prevention of nan or equal to 5%*).
GLUCOSE [DEXTROSE] # Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO # Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO POTASSIUM CHLORIDE		5 1		iomed iomed
* Inj 75 mg per ml, 10 ml	65.00	50	√ J	uno
SODIUM BICARBONATE Inj 8.4%, 50 ml a) Up to 5 inj available on a PSO b) Not in combination	22.40	1	√ B	liomed
Inj 8.4%, 100 ml	22.95	1	√ B	liomed

SODIUM CHLORIDE

Not funded for use as a nasal drop. Not funded for nebuliser use except when used in conjunction with an antibiotic intended for nebuliser use.

✓ Baxter	500 ml	Inj 0.9%, bag – Up to 2000 ml available on a PSO1.33
✓ Baxter	1.000 ml	1.36

Only if prescribed on a prescription for renal dialysis, maternity or post-natal care in the home of the patient, or on a PSO for emergency use. (500 ml and 1,000 ml packs)

Inj 23.4% (4 mmol/ml), 20 ml ampoule	35.50	5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standar	d Formulae, page	254	
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO	4.00	20	✓ Fresenius Kabi
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PSO	5.25	50	✓ Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	✓ Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)			
Infusion	CBS	1 OP	✓ TPN

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

WATER

- 1) On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or
- 2) On a bulk supply order; or
- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 10 ml ampoule – Up to 5 inj available on a PSO	7.19	50	✓ Pfizer
	7.60		Multichem
Inj 20 ml ampoule - Up to 5 inj available on a PSO	5.00	20	✓ Fresenius Kabi
(Pfizer Ini 10 ml ampoule to be delisted 1 September 2023)			

Oral Administration		
CALCIUM POLYSTYRENE SULPHONATE Powder	300 g OP	✓ Calcium Resonium
Powder for oral soln - Up to 5 sach available on a PSO9.53	50	✓ Electral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]		
Soln with electrolytes (2 × 500 ml)8.55	1,000 ml OP	✓ Pedialyte - Bubblegum
PHOSPHORUS		
Tab eff 500 mg (16 mmol)82.50	100	Phosphate Phebra
POTASSIUM CHLORIDE		
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)5.26	60	
(17.10)		Chlorvescent
* Tab long-acting 600 mg (8 mmol)	200	✓ Span-K
SODIUM BICARBONATE		
Cap 840 mg8.52	100	✓ Sodibic
		✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE		
Powder84.65	454 g OP	✓ Resonium-A

✓ Zapril

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
` \$	Por 🗸	Manufacture

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DO	XAZOSIN		
*	Tab 2 mg17.35	500	 Doxazosin Clinect
*	Tab 4 mg20.94	500	 Doxazosin Clinect
РΗ	ENOXYBENZAMINE HYDROCHLORIDE		
*	Cap 10 mg65.00	30	✓ BNM S29
	216.67	100	✓ Dibenzyline S29
PR	AZOSIN		
*	Tab 1 mg5.53	100	✓ Arrotex-Prazosin
			S29 S29
*	Tab 2 mg7.00	100	✓ Arrotex-Prazosin
	•		S29 S29
*	Tab 5 mg11.70	100	✓ Arrotex-Prazosin
	•		S29 S29

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL

* Tab 0.5 mg

CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

2.69

* Tab 0.5 mg2.05	30	▼ Zapili
* Tab 2.5 mg5.79	90	✓ Zapril
Tab 5 mg10.05	90	✓ Zapril
ENALAPRIL MALEATE		
* Tab 5 mg1.75	90	✓ Acetec
* Tab 10 mg1.97	90	✓ Acetec
* Tab 20 mg	90	✓ Acetec
LISINOPRIL		
* Tab 5 mg11.07	90	✓ Ethics Lisinopril
•		✓ Teva Lisinopril
* Tab 10 mg11.67	90	✓ Ethics Lisinopril
		✓ Teva Lisinopril
* Tab 20 mg14.69	90	✓ Ethics Lisinopril
		✓ Teva Lisinopril
PERINDOPRIL		
* Tab 2 mg	30	✓ Coversyl
* Tab 4 mg2.95	30	✓ Coversyl
* Tab 8 mg5.02	30	✓ Coversyl

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

			Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
# Tab 10 mg	QU	IINAPRIL			
# Tab 20 mg	*	Tab 5 mg	5.97	90	✓ Arrow-Quinapril 5
### AMIPRIL ### Cap 1.25 mg	*	Tab 10 mg	5.18	90	✓ Arrow-Quinapril 10
# Cap 1.25 mg	*	Tab 20 mg	7.95	90	✓ Arrow-Quinapril 20
Tryzan to be Principal Supply on 1 May 2023 Cap 2.5 mg	RA	MIPRIL			
Tryzan to be Principal Supply on 1 May 2023 Cap 2.5 mg	*	Cap 1.25 mg	6.90	90	✓ Tryzan
★ Cap 2.5 mg 6.60 90 ✓ Tryzan Tryzan to be Principal Supply on 1 May 2023 6.75 90 ✓ Tryzan ★ Cap 5 mg 7.05 90 ✓ Tryzan ★ Cap 10 mg 7.05 90 ✓ Tryzan Tryzan to be Principal Supply on 1 May 2023 ACE Inhibitors with Diuretics CUINAPRIL WITH HYDROCHLOROTHIAZIDE – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who were taking quinapril with hydrochlorothiazide prior to 1 May 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of quinapril with hydrochlorothiazide. Tab 10 mg with hydrochlorothiazide 12.5 mg 4.10 30 ✓ Accuretic 10 Tab 20 mg with hydrochlorothiazide 12.5 mg 4.10 30 ✓ Accuretic 20 Angiotensin II Antagonists CANDESARTAN CILEXETIL * Tab 4 mg 2.00 90 ✓ Candestar * Tab 16 mg 3.31 90 ✓ Candestar * Tab 32 mg 5.26 90 ✓ Candestar * Tab 12.5 mg 1.56 84 ✓ Losartan Actavis * Tab 50 mg 1.8					•
** Cap 5 mg	*		6.60	90	✓ Tryzan
Tryzan to be Principal Supply on 1 May 2023 Cap 10 mg		Tryzan to be Principal Supply on 1 May 2023			•
# Cap 10 mg	*	Cap 5 mg	6.75	90	✓ Tryzan
ACE Inhibitors with Diuretics QUINAPRIL WITH HYDROCHLOROTHIAZIDE — Subsidy by endorsement Subsidy by endorsement — Subsidised for patients who were taking quinapril with hydrochlorothiazide prior to 1 May 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of quinapril with hydrochlorothiazide. Tab 10 mg with hydrochlorothiazide 12.5 mg		Tryzan to be Principal Supply on 1 May 2023			
ACE Inhibitors with Diuretics QUINAPRIL WITH HYDROCHLOROTHIAZIDE — Subsidy by endorsement Subsidy by endorsement — Subsidised for patients who were taking quinapril with hydrochlorothiazide prior to 1 May 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of quinapril with hydrochlorothiazide. Tab 10 mg with hydrochlorothiazide 12.5 mg	*		7.05	90	Tryzan
Augiotensin II Antagonists CANDESARTAN CILEXETIL * Tab 16 mg * Tab 13 mg * Tab 12 mg * Tab 12 mg * Tab 12 mg * Tab 12 mg * Tab 15 mg * Tab 15 mg * Tab 10 mg * Tab 50 mg * Tab		Tryzan to be Principal Supply on 1 May 2023			
Subsidy by endorsement – Subsidised for patients who were taking quinapril with hydrochlorothiazide prior to 1 May 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of quinapril with hydrochlorothiazide. Tab 10 mg with hydrochlorothiazide 12.5 mg			y hy andorsament		
Angiotensin II Antagonists CANDESARTAN CILEXETIL K Tab 4 mg		2022 and the prescription is endorsed accordingly. P exists a record of prior dispensing of quinapril with hydrogeneous properties.	harmacists may annotate the drochlorothiazide.	pres	scription as endorsed where ther
ANDESARTAN CILEXETIL Tab 4 mg		Tab 20 mg with hydrochlorothiazide 12.5 mg	5.25	30	✓ Accuretic 20
K Tab 4 mg 2.00 90 ✓ Candestar K Tab 8 mg 2.28 90 ✓ Candestar K Tab 16 mg 3.31 90 ✓ Candestar K Tab 32 mg 5.26 90 ✓ Candestar OSARTAN POTASSIUM I.56 84 ✓ Losartan Actavis K Tab 12.5 mg 1.84 84 ✓ Losartan Actavis K Tab 50 mg 2.25 84 ✓ Losartan Actavis K Tab 100 mg 3.50 84 ✓ Losartan Actavis Angiotensin II Antagonists with Diuretics OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE 4.00 30 ✓ Arrow-Losartan &	Α	ngiotensin II Antagonists			
★ Tab 8 mg 2.28 90 ✓ Candestar ★ Tab 16 mg 3.31 90 ✓ Candestar ★ Tab 32 mg 5.26 90 ✓ Candestar .OSARTAN POTASSIUM ★ Tab 12.5 mg 1.56 84 ✓ Losartan Actavis ★ Tab 25 mg 1.84 84 ✓ Losartan Actavis ★ Tab 50 mg 2.25 84 ✓ Losartan Actavis ★ Tab 100 mg 3.50 84 ✓ Losartan Actavis Angiotensin II Antagonists with Diuretics **OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE **Tab 50 mg with hydrochlorothiazide 12.5 mg 4.00 30 ✓ Arrow-Losartan &	CA	NDESARTAN CILEXETIL			
* Tab 16 mg 3.31 90 ✓ Candestar * Tab 32 mg 5.26 90 ✓ Candestar LOSARTAN POTASSIUM 1.56 84 ✓ Losartan Actavis * Tab 12.5 mg 1.84 84 ✓ Losartan Actavis * Tab 50 mg 2.25 84 ✓ Losartan Actavis * Tab 100 mg 3.50 84 ✓ Losartan Actavis Angiotensin II Antagonists with Diuretics LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE * Tab 50 mg with hydrochlorothiazide 12.5 mg 4.00 30 ✓ Arrow-Losartan &	*	Tab 4 mg	2.00	90	✓ Candestar
★ Tab 16 mg 3.31 90 ✓ Candestar ★ Tab 32 mg 5.26 90 ✓ Candestar .OSARTAN POTASSIUM ★ Tab 12.5 mg 1.56 84 ✓ Losartan Actavis ★ Tab 25 mg 1.84 84 ✓ Losartan Actavis ★ Tab 50 mg 2.25 84 ✓ Losartan Actavis ★ Tab 100 mg 3.50 84 ✓ Losartan Actavis Angiotensin II Antagonists with Diuretics OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE ★ Tab 50 mg with hydrochlorothiazide 12.5 mg 4.00 30 ✓ Arrow-Losartan &	*	S .		90	
K Tab 32 mg 5.26 90 ✓ Candestar OSARTAN POTASSIUM I.56 84 ✓ Losartan Actavis K Tab 12.5 mg 1.84 84 ✓ Losartan Actavis K Tab 50 mg 2.25 84 ✓ Losartan Actavis K Tab 100 mg 3.50 84 ✓ Losartan Actavis Angiotensin II Antagonists with Diuretics OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE 4.00 30 ✓ Arrow-Losartan &	K			90	✓ Candestar
OSARTAN POTASSIUM k Tab 12.5 mg	k				
k Tab 12.5 mg 1.56 84 ✓ Losartan Actavis k Tab 25 mg 1.84 84 ✓ Losartan Actavis k Tab 50 mg 2.25 84 ✓ Losartan Actavis k Tab 100 mg 3.50 84 ✓ Losartan Actavis Angiotensin II Antagonists with Diuretics OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE k Tab 50 mg with hydrochlorothiazide 12.5 mg 4.00 30 Arrow-Losartan &	\sim	•			
★ Tab 25 mg 1.84 84 ✓ Losartan Actavis ★ Tab 50 mg 2.25 84 ✓ Losartan Actavis ★ Tab 100 mg 3.50 84 ✓ Losartan Actavis Angiotensin II Antagonists with Diuretics COSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE ★ Tab 50 mg with hydrochlorothiazide 12.5 mg 4.00 30 ✓ Arrow-Losartan &	-		1.56	0.1	✓ Locartan Actavis
★ Tab 50 mg 2.25 84 ✓ Losartan Actavis ★ Tab 100 mg 3.50 84 ✓ Losartan Actavis Angiotensin II Antagonists with Diuretics OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE ★ Tab 50 mg with hydrochlorothiazide 12.5 mg 4.00 30 ✓ Arrow-Losartan &		•			
★ Tab 100 mg Angiotensin II Antagonists with Diuretics OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE ★ Tab 50 mg with hydrochlorothiazide 12.5 mg 4.00 30 Arrow-Losartan &					
Angiotensin II Antagonists with Diuretics OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE * Tab 50 mg with hydrochlorothiazide 12.5 mg		•			
OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE * Tab 50 mg with hydrochlorothiazide 12.5 mg4.00 30 Arrow-Losartan &	r	Tab Too Hig	3.30	04	LUSARIAN ACIAVIS
★ Tab 50 mg with hydrochlorothiazide 12.5 mg	A	ngiotensin II Antagonists with Diuretics			
				30	

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN – Special Authority see S	A1905 on the next page	 Retail p 	oharmacy
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure: and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III: or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthet	ics, Local, pag	e 117	
AMIODARONE HYDROCHLORIDE			
▲ Tab 100 mg	3.49	30	✓ Aratac
▲ Tab 200 mg		30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a PSC)9.12	6	✓ Cordarone-X
	15.22	10	✓ Max Health
ATROPINE SULPHATE			
* Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a			
PSO	15.09	10	✓ Martindale
DIGOXIN			
* Tab 62.5 mcg - Up to 30 tab available on a PSO	7.80	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO		240	✓ Lanoxin
* Oral liq 50 mcg per ml		60 ml	✓ Lanoxin
			✓ Lanoxin Paediatric
			Elixir S29
			✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE			
▲ Cap 100 mg	23.87	100	✓ Rythmodan
FLECAINIDE ACETATE	20.07		,
▲ Tab 50 mg	10.05	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg		90	✓ Flecainide
Δ Oap long doing 100 mg	05.70	30	Controlled
			Release Teva
▲ Cap long-acting 200 mg	54 28	90	✓ Flecainide
_ cap long doing too mg	0 1.20	00	Controlled
			Release Teva
Inj 10 mg per ml, 15 ml ampoule	.104.00	5	✓ Tambocor
MEXILETINE HYDROCHLORIDE			
	160.00	100	✓ Teva S29
▲ Cap 150 mg			
▲ Cap 250 mg	.202.00	100	✓ Teva S29

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PROPAFENONE HYDROCHLORIDE Tab 150 mg	40.90	50	1	Rytmonorm
Antihypotensives				
MIDODRINE - Special Authority see SA1474 below - Retail pha	ırmacy			
Tab 2.5 mg	38.23	100	•	Midodrine Medsurge
	53.00		1	Gutron
Tab 5 mg	59.98	100	•	Midodrine Medsurge
	79.00		✓	Gutron
(Gutron Tab 2.5 mg to be delisted 1 August 2023)				

(Gutron Tab 2.5 mg to be delisted 1 August 2023) (Gutron Tab 5 mg to be delisted 1 August 2023)

►SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ΑT	ENOLOL			
*	Tab 50 mg	9.33	500	Mylan Atenolol
	-			✓ Viatris
*	Tab 100 mg	14.20	500	Mylan Atenolol
*	Oral liq 25 mg per 5 ml	21.25	300 ml OP	✓ Atenolol AFT
				S29 S29
		38.20		✓ Essential
				Generics S29
		49.85		✓ Atenolol AFT
	Restricted to children under 12 years of ag	e.		
BIS	SOPROLOL FUMARATE			
*	Tab 2.5 mg	1.84	90	✓ Bisoprolol Mylan
	· ·			✓ Bisoprolol Viatris
*	Tab 5 mg	2.55	90	✓ Bisoprolol Mylan
	•			✓ Bisoprolol Viatris
*	Tab 10 mg	3.62	90	✓ Bisoprolol Mylan
				 Bisoprolol Viatris
CA	RVEDILOL			
*	Tab 6.25 mg	2.24	60	✓ Carvedilol Sandoz
*	Tab 12.5 mg		60	✓ Carvedilol Sandoz
*	Tab 25 mg		60	✓ Carvedilol Sandoz

	Subsidy		Fully	
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
LABETALOL				
* Tab 100 mg	14.50	100	✓	<u>Trandate</u>
* Tab 200 mg	27.00	100	✓	<u>Trandate</u>
* Inj 5 mg per ml, 20 ml ampoule	59.06	5		
	(88.60)			Trandate
* inj 5 mg per ml, 20 ml vial	42.29	1		
	(48.20)			Alvogen S29
METOPROLOL SUCCINATE				
* Tab long-acting 23.75 mg	1.45	30	✓	Betaloc CR
* Tab long-acting 47.5 mg	1.43	30	✓	Betaloc CR
* Tab long-acting 95 mg	2.15	30	✓	Betaloc CR
* Tab long-acting 190 mg		30	✓	Betaloc CR
METOPROLOL TARTRATE				
* Tab 50 mg	5.66	100	✓	IPCA-Metoprolol
* Tab 100 mg		60		IPCA-Metoprolol
* Tab long-acting 200 mg		28		Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial		5		Metoprolol IV Mylan
				Metoprolol IV Viatris
NADOLOL				
* Tab 40 mg	19.19	100	✓	Nadolol BNM
* Tab 80 mg	30.39	100	✓	Nadolol BNM
PROPRANOLOL				
* Tab 10 mg	7.04	100	✓	Drofate
* Tab 40 mg		100		IPCA-Propranolol
* Cap long-acting 160 mg		100		Cardinol LA
* Oral liq 4 mg per ml – Special Authority see SA1327 bek				
Retail pharmacy		500 m	ı 🗸	Roxane-
•				Propranolol S29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

S	O	T	P	۱I	()	ı
_	_	٠	•	**	٠,	_	_

*	Tab 80 mg37.50	500	✓ Mylan
*	Tab 160 mg14.00	100	Mylan

		(Manufacturer's Price)		Subsidised	
		\$	Per		Manufacturer
C	alcium Channel Blockers				
	aloidii Olidiiici Blookers				
D	ihydropyridine Calcium Channel Blockers				
AM	LODIPINE				
*	Tab 2.5 mg		90		Vasorex
*	Tab 5 mg		90	/	Vasorex
	Tab 10 mg	1.19	90	•	Vasorex
	ODIPINE				
	Tab long-acting 2.5 mg		30		Plendil ER
	Tab long-acting 5 mg		90	_	Felo 5 ER
	Tab long-acting 10 mg	4.32	90	•	Felo 10 ER
	EDIPINE			_	
*	Tab long-acting 10 mg	18.80	56	•	Tensipine MR10 S29
*	Tab long-acting 20 mg	0 12	50	1	Mylan (12 hr
-1.	rab long doding 20 mg		00	•	release) S29
		17.72	100	1	Nyefax Retard
*	Tab long-acting 30 mg		14		Mylan Italy (24 hr
•	- 1.2. 1.0.1.g 4.0.1.1.g		• •		release) \$29
		34.10	100	/	Mylan (24 hr
		••			release) S29
*	Tab long-acting 60 mg	52 81	100	/	Mylan (24 hr
•••	rab long adding of mg				release) S29
					release) —
0	ther Calcium Channel Blockers				
DII	TIAZEM HYDROCHLORIDE				
	Cap extended-release 120 mg	44 40	100	1	Accord S29
*	Cap long-acting 120 mg		500		Apo-Diltiazem CD
•	oup iong downg i_o ing iiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiii	65.35			Diltiazem CD Clinect
	Diltiazem CD Clinect to be Principal Supply on 1 June 20)23			
*	Cap long-acting 180 mg	7.00	30		Cardizem CD
*	Cap long-acting 240 mg	9.30	30	•	Cardizem CD
•	cord \$29 Cap extended-release 120 mg to be delisted 1 Jun	,			
(Ap	o-Diltiazem CD Cap long-acting 120 mg to be delisted 1 June	2023)			
	RHEXILINE MALEATE				
*	Tab 100 mg	62.90	100	1	Pexsig
۷E	RAPAMIL HYDROCHLORIDE				
*	Tab 40 mg		100		Isoptin
*	Tab 80 mg	11.74	100	/	Isoptin

Subsidy

Fully

Brand or

Tab long-acting 120 mg......36.02

Inj 2.5 mg per ml, 2 ml ampoule - Up to 5 inj available on a

100

30

5

✓ Isoptin Retard S29✓ Isoptin SR

✓ Isoptin SR

✓ Isoptin

				_
	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	I Generic Manufacturer
	Ψ	1 01		Wandactarci
Centrally-Acting Agents				
CLONIDINE				
* Patch 2.5 mg, 100 mcg per day – Only on a prescription	10.24	4	./	Mylan
* Patch 5 mg, 200 mcg per day — Only on a prescription		4		Mylan
* Patch 7.5 mg, 300 mcg per day — Only on a prescription		4		Mylan
	10.33	4	•	wyian
CLONIDINE HYDROCHLORIDE				A =
* Tab 25 mcg		112		Clonidine Teva
* Tab 150 mcg		100		Catapres
* Inj 150 mcg per ml, 1 ml ampoule	29.68	10	•	<u>Medsurge</u>
METHYLDOPA				
* Tab 250 mg	15.10	100	✓	Methyldopa Mylan
·	52.85	500		Methyldopa Mylan
				S29 S29
Diuretics				
Loop Diuretics				
•				
BUMETANIDE			_	
* Tab 1 mg		30		Burinex S29 S29
	16.36	100		Burinex
* Inj 500 mcg per ml, 4 ml vial	7.95	5	/	Burinex
FUROSEMIDE [FRUSEMIDE]				
Tab 40 mg - Up to 30 tab available on a PSO	8.00	1,000	/	IPCA-Frusemide
* Tab 500 mg		50	1	Urex Forte
•	89.48		1	Furosemid-
				Ratiopharm \$29
	169.96	100	1	Furosemid-
				Ratiopharm S29
* Oral liq 10 mg per ml) ml C		Lasix
* Inj 10 mg per ml, 25 ml ampoule		6		Lasix
* Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a f	PSO2.40	5	/	Furosemide-Baxter
Balancian Oundon Blooding				
Potassium Sparing Diuretics				
AMILORIDE HYDROCHLORIDE				
Oral liq 1 mg per ml	32.10 2	5 ml C)P 🗸	Biomed
EPLERENONE - Special Authority see SA1728 below - Retail p				
Tab 25 mg		30	1	Inspra
Tab 50 mg		30		Inspra
	25.00	30	•	Πορια
⇒SA1728 Special Authority for Subsidy			,	
Initial application from any relevant practitioner. Approvals vali	d without further rene	wal u	niess notif	ried for applications meeting
the following criteria:				
Both:				

- 1 Patient has heart failure with ejection fraction less than 40%; and
- 2 Either:
 - 2.1 Patient is intolerant to optimal dosing of spironolactone; or
 - 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

	Subsidy	Ful	ly Brand or
	(Manufacturer's Price	e) Subsidise Per •	d Generic Manufacturer
SPIRONOLACTONE			
₭ Tab 25 mg			Spiractin
★ Tab 100 mg			Spiractin
Oral liq 5 mg per ml	33.00	25 ml OP	Biomed
Potassium Sparing Combination Diuretics			
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE			
* Tab 5 mg with furosemide 40 mg		28	Frumil
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZ	ZIDE		
* Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	Moduretic
Thiazide and Related Diuretics			
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]			
★ Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	Arrow-
			<u>Bendrofluazide</u>
May be supplied on a PSO for reasons other than eme			-
₹ Tab 5 mg	34.55	500	Arrow-
			<u>Bendrofluazide</u>
CHLOROTHIAZIDE			
Oral liq 50 mg per ml	27.82	25 ml OP	Biomed
CHLORTALIDONE [CHLORTHALIDONE]			
Tab 25 mg	6.95	50	/ Hygroton
NDAPAMIDE			
★ Tab 2.5 mg	10.45	90	Dapa-Tabs
ŭ	11.61		/ Mylan
			Indapamide \$29
Mylan Indapamide 👀 Tab 2.5 mg to be delisted 1 August 20	023)		
METOLAZONE			
Tab 5 mg	CBS	1	Metolazone S29
		50	Zaroxolyn S29
Vasopressin receptor antagonists			
OLVAPTAN - Special Authority see SA2166 below - Retail pl	harmacy		
Tab 15 mg			/ Jinarc
Tab 30 mg			/ Jinarc
Tab 45 mg + 15 mg	*		Jinarc
Tab 60 mg + 30 mg	*		Jinarc
Tab 90 mg + 30 mg	1,747.00	56 OP ✓	Jinarc
⇒SA2166 Special Authority for Subsidy			
nitial application — (autosomal dominant polycystic kidner			
n the recommendation of a renal physician. Approvals valid fo	r 12 months for appl	ications meeting	the following criteria:
l of the following:			

1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and

continued...

(Ma	Subsidy anufacturer's Price)	Sub	Fully	Brand or Generic
	\$	Per	•	Manufacturer

continued...

- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Either:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents			
Fibrates			
BEZAFIBRATE * Tab 200 mg * Tab long-acting 400 mg		90 30	✓ <u>Bezalip</u>✓ <u>Bezalip</u> Retard
Other Lipid-Modifying Agents			
ACIPIMOX * Cap 250 mg	21.56 25.44	30	✓ Olbetam S29 S29 ✓ Olbetam
Resins			
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	32.89	30	✓ Colestid
HMG CoA Reductase Inhibitors (Statins)			
ATORVASTATIN			
* Tab 10 mg	6.16	500	✓ Lorstat
* Tab 20 mg		500	Lorstat
* Tab 40 mg		500	Lorstat
* Tab 80 mg	26.54	500	✓ <u>Lorstat</u>
PRAVASTATIN * Tab 20 mg	2.11	28	✓ <u>Pravastatin Mylan</u> ✓ Pravastatin Viatris
* Tab 40 mg	3.61	28	✓ Pravastatin Mylan
ROSUVASTATIN - Special Authority see SA2093 on the next page	– Retail pharma	ICV	
* Tab 5 mg		30	✓ Rosuvastatin Viatris
* Tab 10 mg	2.42	30	✓ Rosuvastatin Viatris
* Tab 20 mg		30	✓ Rosuvastatin Viatris
* Tab 40 mg	5.28	30	✓ Rosuvastatin Viatris

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Pe	r 🗸	Manufacturer

⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient is considered to be at risk of cardiovascular disease; and
 - 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:
 - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
 - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
 - 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:
Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIN	IVASTATIN			
*	Tab 10 mg	1.23	90	Simvastatin Mylan
	Tab 20 mg		90	✓ Simvastatin Mylan
	Tab 40 mg		90	✓ Simvastatin Mylan
	· ·			✓ Simvastatin Viatris
*	Tab 80 mg	7.12	90	✓ Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Special Authority see SA1045 below – Retail pharmacy		
* Tab 10 mg	30	✓ Ezetimibe Sandoz

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 x normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 be	olow – Retail	pharmacy	
Tab 10 mg with simvastatin 10 mg	5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg		30	✓ Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

CI	VCERVI	TRINITRATE	

*	Oral pump spray, 400 mcg per dose – Up to 250 dose			
	available on a PSO	7.48	250 dose OP	✓ Nitrolingual Pump Sprav
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
	Patch 50 mg, 10 mg per day		30	✓ Nitroderm TTS
ISO	OSORBIDE MONONITRATE			
*	Tab 20 mg	19.55	100	✓ Ismo 20
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
	Tab long-acting 60 mg		90	✓ Duride

Sympathomimetics

ADRENALINE

III 1 III 1,000, 1 IIII ampoule – Op to 5 III available on a F504.96	5	Aspen Aurenanne
12.65		DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	✓ Aspen Adrenaline

Ini 1 in 1 000 1 ml ampaula. Un to E ini quailable on a DCO

CARDIOVASCIII AR SYSTEM

CARDIOVASCULAR SYSTEM				
	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
Vasodilators				
HYDRALAZINE HYDROCHLORIDE				
* Tab 25 mg - Special Authority see SA1321 below - Retail				
pharmacy	CBS	1	•	Hydralazine
		56	•	Onelink \$29
		84	1	AMDIPHARM \$29
		100	1	Onelink S29
* Inj 20 mg ampoule	25.90	5	•	Apresoline
Initial application from any relevant practitioner. Approvals valid the following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitr inhibitors and/or angiotensin receptor blockers.				
MINOXIDIL				
▲ Tab 10 mg	78.40	100	•	Loniten
NICORANDIL				
▲ Tab 10 mg	25.57	60	1	Ikorel
▲ Tab 20 mg	32.28	60	•	Ikorel
PAPAVERINE HYDROCHLORIDE				
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	•	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]				
Tab 400 mg	42.26	50	•	Trental 400
Endothelin Receptor Antagonists				
AMBRISENTAN - Special Authority see SA1702 below - Retail	pharmacy			
Tab 5 mg		30		Ambrisentan Mylan
Tab 10 mg	1,550.00	30		Ambrisentan Viatris
			•	Mylan
SA1702 Special Authority for Subsidy	on Daniel			
Special Authority approved by the Pulmonary Arterial Hypertensic Notes: Application details may be obtained from Pharmac's webs		c dov	t nz/SAEc	arme or:
The Coordinator, PAH Panel	site <u>scriedule.priarilla</u>	c.gov	LIIZ/SAFC	orris or.
Pharmac, PO Box 10-254, WELLINGTON				
Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac	.govt.nz			
BOSENTAN - Special Authority see SA1991 below - Retail phar				
Tab 62.5 mg	•	60	✓	Bosentan Dr Reddy's

⇒SA1991 Special Authority for Subsidy

Tab 125 mg119.85

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

60

✓ Bosentan Dr

Reddy's

Reddy's

Subsidy			
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	 Manufacturer 	

continued

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Fither:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

			_
Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
` ¢ ′	Por 🗸	Manufacturor	

Phosphodiesterase Type 5 Inhibitors

		SILDENAFIL - Special Authority see SA1992 below - Retail pharmacy
4 ✓ Vedafil	.0.85	Tab 25 mg
4 ✓ Vedafil	.1.70	Tab 50 mg
12 ✓ Vedafil	10.20	Tab 100 mg

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III; or
 - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 below	 Retail pharmacy 		
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz



Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

ADAPAI FNF

IS

- a) Maximum of 30 g per prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
SOTRETINOIN - Special Authority see SA2023 bel	low – Retail pharmacy		
Cap 5 mg	11.26	60	Oratane
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
- 2 Patient is not of child bearing potential.

TRFTINOIN

✓ ReTrieve Crm 0.5 mg per g - Maximum of 50 g per prescription15.57 50 q OP

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

# Crm 1%	8.56	10 g OP	✓ Crystaderm
MUPIROCIN			
Oint 2%	6.60	15 g OP	
	(11.50)	_	Bactroban

- a) Only on a prescription
- b) Not in combination

	Subsidy (Manufacturer's Pric	ce) Sul	Fully bsidised	Brand or Generic Manufacturer	
SODIUM FUSIDATE [FUSIDIC ACID]	· · · · · · · · · · · · · · · · · · ·				
Crm 2%	1.59	5 g OP	✓	Foban	
a) Maximum of 5 g per prescription		J	-		
b) Only on a prescription					
c) Not in combination		- 0-			
Oint 2%	1.59	5 g OP	✓ [<u>Foban</u>	
a) Maximum of 5 g per prescription b) Only on a prescription					
b) Only on a prescriptionc) Not in combination					
SULFADIAZINE SILVER					
Crm 1%	10.80	50 g OP	✓	Flamazine	
a) Up to 250 g available on a PSO		3 -			
b) Not in combination					
Antifungals Topical					
Antinungais ropical					
For systemic antifungals, refer to INFECTIONS, Antifungals, pa	ge 97				
AMOROLFINE					
a) Only on a prescription					
b) Not in combination Nail soln 5%	14.02	5 ml OP	./ 1	MyssMsil	
	14.93	5 IIII OP	V <u>i</u>	<u>MycoNail</u>	
CLOTRIMAZOLE * Crm 1%	1 10	20 g OP	1	Clomazol	
a) Only on a prescription		20 g Oi	• •	Siomazoi	
b) Not in combination					
* Soln 1%	4.36	20 ml OP			
	(7.55)		(Canesten	
a) Only on a prescription					
b) Not in combination					
ECONAZOLE NITRATE	1.00	00 = OD			
Crm 1%	(7.78)	20 g OP	ı	Pevaryl	
a) Only on a prescription	(1.10)			Ovaryi	
b) Not in combination					
Foaming soln 1%, 10 ml sachets	9.89	3			
	(17.92)		F	Pevaryl	
a) Only on a prescription					
b) Not in combination					
MICONAZOLE NITRATE * Crm 2%	0.01	15 a OD	./ 1	Multichem	
a) Only on a prescription	0.01	15 g OP	• <u>!</u>	<u>Multichem</u>	
b) Not in combination					
* Lotn 2%	4.36	30 ml OP			
	(10.03)		[Daktarin	
a) Only on a prescription					
b) Not in combination	4.00	20 1 00			
* Tinct 2%	4.36 (12.10)	30 ml OP	ı	Daktarin	
a) Only on a prescription	(12.10)		ı	Jantaiiii	
b) Not in combination					
•					

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per ✔ Manufacturer

Antipruritic Preparations

CALAMINE

a) Only on a prescription

b) Not in combination

CROTAMITON

a) Only on a prescription

b) Not in combination

MENTHOL - Only in combination

1) Only in combination with a dermatological base or proprietary Topical Corticosteriod - Plain

2) With or without other dermatological galenicals.

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 80

Corticosteroids - Plain

BETAMETHASONE DIPROPIONATE			
Crm 0.05%	2.96	15 g OP	Diprosone
	36.00	50 g OP	✓ Diprosone
Oint 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	✓ Diprosone
Oint 0.05% in propylene glycol base	4.33	30 g OP	✓ Diprosone OV
BETAMETHASONE VALERATE			
* Crm 0.1%	4.53	50 g OP	✓ Beta Cream
* Oint 0.1%	5.84	50 g OP	✓ Beta Ointment
* Lotn 0.1%	25.00	50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE			
* Crm 0.05%	2.40	30 g OP	✓ Dermol
* Oint 0.05%	2.33	30 g OP	✓ Dermol
CLOBETASONE BUTYRATE			
Crm 0.05%	5.38	30 g OP	
	(10.00)	· ·	Eumovate
HYDROCORTISONE			
* Crm 1% - Only on a prescription	1.78	30 g OP	✓ Ethics
, , ,	17.15	500 g	✓ Hydrocortisone (PSM)
	20.40		✓ Noumed
* Powder – Only in combination	49.95	25 g	✓ ABM
Up to 5% in a dermatological base (not proprietary	Topical Corticosterio	d – Plain) with	or without other dermatologic

Up to 5% in a dermatological base (not proprietary Topical Corticosteriod – Plain) with or without other dermatological galenicals

(Hydrocortisone (PSM) Crm 1% to be delisted 1 August 2023)

	Subsidy	lwine) Cubei	Fully	Brand or Generic
	(Manufacturer's F	Per	uiseu •	Manufacturer
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only o	n			
a prescription		250 ml	1	DP Lotn HC
HYDROCORTISONE BUTYRATE				
Lipocream 0.1%	4.85	100 g OP	1	Locoid Lipocream
Oint 0.1%		100 g OP		Locoid
Milky emul 0.1%	12.33	100 ml OP	1	Locoid Crelo
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.46	15 g OP	1	Advantan
Oint 0.1%	4.46	15 g OP	1	Advantan
MOMETASONE FUROATE				
Crm 0.1%	1.95	15 g OP	1	Elocon Alcohol Free
	3.10	50 g OP		Elocon Alcohol Free
Oint 0.1%		15 g OP		Elocon
	2.90	50 g OP		Elocon
Lotn 0.1%	4.50	30 ml OP	•	<u>Elocon</u>
TRIAMCINOLONE ACETONIDE			_	
Crm 0.02%		100 g OP		Aristocort
Oint 0.02%	6.35	100 g OP	•	Aristocort
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUS	SIDIC ACID]			
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		
	(10.45)			Fucicort
a) Maximum of 15 g per prescription				
b) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE - Only on a prescript		05		
* Crm 1% with miconazole nitrate 2%		15 g OP	•	Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - OI	, , ,			-
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP		Pimafucort
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCII		-IN		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg				
and gramicidin 250 mcg per g - Only on a prescription		15 g OP		Via dama KO
	(9.28)			Viaderm KC
Barrier Creams and Emollients				
Barrier Creams				
DIMETHICONE				
* Crm 5% pump bottle	4.30	500 ml OP	•	healthE Dimethicone 5%
* Crm 10% pump bottle	4 50	500 ml OD	./	<u>Dimethicone 5%</u> healthE
* Crm 10% pump bottle	4.52	500 ml OP	•	Dimethicone 10%
ZINIC AND CACTOD OIL				Dillicultotic 10/0
ZINC AND CASTOR OIL * Oint	165	500 a		Boucher
~ UIII	4.00	500 g	•	Doublici

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy	Delas) Cul-	. ,	Brand or
	(Manufacturer's	Price) Subs Per		Generic Manufacturer
	<u> </u>	1 01		via raraotaro
Emollients				
AOUEOUO ODEAM				
AQUEOUS CREAM * Crm	1 70	F00 =	. OF	1 1
* Cm	1.73	500 g		M Aqueous
055011000001			<u>U</u>	ream
CETOMACROGOL	4.00	500		
* Crm BP	1.99	500 g	✓ <u>Cet</u>	omacrogol-AFT
CETOMACROGOL WITH GLYCEROL				
Crm 90% with glycerol 10%		500 ml OP	✓ Eva	
	2.35			rmacy Health
			-	orbolene with
				lycerin
	3.50	1,000 ml OP	Eva	ra
Evara to be Principal Supply on 1 July 2023				
(Pharmacy Health Sorbolene with Glycerin Crm 90% with glyce	erol 10% to be del	isted 1 July 202	3)	
EMULSIFYING OINTMENT				
* Oint BP	3.40	500 g	✓ Em	ulsifying
			0	intment ADE
OIL IN WATER EMULSION				
* Crm	2.04	500 g	✓ Fatt	y Cream AFT
PARAFFIN		3		
Oint liquid paraffin 50% with white soft paraffin 50%	4 94	500 g OP	✓ Wh	te Soft Liquid
One ilquid paramin 50 /5 with write 501 paramin 50 /5		000 g Oi		araffin AFT
White Soft Liquid Paraffin AFT to be Principal Supply	on 1 May 2023		•	
Oint liquid paraffin 50% with white soft paraffin 50%,		500 ml OP	✓ hea	IthE
(healthE Oint liquid paraffin 50% with white soft paraffin 50%, t				
UREA		.,,		
* Crm 10%	1 27	100 g OP	√ hoo	IthE Urea Cream
	1.37	100 g OF	• IIEa	illic Olea Cleaili
WOOL FAT WITH MINERAL OIL - Only on a prescription	- 00	4 000		
* Lotn hydrous 3% with mineral oil		1,000 ml	DD	
	(14.96)			Lotion
	(20.53) 1.40	050 ml OD	Alþi	na-Keri Lotion
		250 ml OP	DD	Lotion
	(5.87) 5.60	1,000 ml	טף	LUUUII
	(23.91)	1,000 1111	ВK	Lotion
	1.40	250 ml OP	אט	LOUGH
	(7.73)	200 1111 01	BK	Lotion
	(7.70)		DI.	-0.011
Other Dermatological Bases				
•				
PARAFFIN			_	
White soft - Only in combination		450 g	✓ hea	
	19.99	2,500 g	✓ hea	
Only in combination with a dermatological galenical or	as a diluent for a	proprietary Topi	cal Cortic	osteroid – Plain.

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully osidised	Brand or Generic Manufacturer
Minor Skin Infections				
POVIDONE IODINE Oint 10% a) Maximum of 130 g per prescription b) Only on a prescription	7.40 6	55 g OP	√ <u>[</u>	<u>Betadine</u>
Antiseptic Solution 10%		100 ml 15 ml	-	<u>Riodine</u> Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	5.40	500 ml 100 ml	✓ F	Riodine
Skin preparation, povidone iodine 10% with 70% alcohol		100 ml		Betadine Skin Prep Pfizer
Parasiticidal Preparations	(7.78)		·	FIIZGI

a valid Special Authority for patient of that institution.

2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.

200 ml OP

✓ healthE

Dimethicone 4% Lotion

 For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:

DIMETHICONE

- 2.1 Both:
 - 2.1.1 The patient is in the community; and

IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy

- 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
- 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables); or

continued...

DERMATOLOGICALS

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
- 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

- Any of the following: 1 Filaricides: or
 - 2 Cutaneous larva migrans (creeping eruption); or
 - 3 Strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5%5.75	30 g OP	✓ Lyderm
Lotn 5%	30 ml OP	✓ A-Scabies

Psoriasis and Eczema Preparations

Retail pharmacy		
17.86	60	Novatretin
41.36	60	✓ Novatretin
	17.86	17.86 60

DERMATOLOGICAL

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment:
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	39.35	60 g OP	 Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g	15.90	30 g OP	✓ Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR			
Soln BP - Only in combination	36.25	200 ml	✓ Midwest
1) Up to 10% only in combination with a dermatological b	ase or propri	ietary Topical C	orticosteriod - Pl

- Plain
- 2) With or without other dermatological galenicals.

COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and

allantoin crm 2.5%	6.59	75 g OP	
	(8.00)	J	Egopsoryl TA
	3.43	30 g OP	01 7
	(4.35)	- · · · · ·	Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ Coco-Scalp
	7.95	40 g OP	✓ Coco-Scalp

PIMECROLIMUS - Special Authority see SA1970 on the next page - Retail pharmacy

- a) Maximum of 15 g per prescription
- b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks

b) reservation of region presemption and resimilar	p. 000p	o po	,
Cream 1%	28.50	15 g OP	Elidel



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1970 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN – Only on a prescription * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium........4.44 500 ml

SALICYLIC ACID

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain or collodion flexible
- 2) With or without other dermatological galenicals.

SULPHUR

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain
- 2) With or without other dermatological galenicals.

TACROLIMUS

- a) Maximum of 30 g per prescription
- b) Note: a maximum of 30 g per prescription and no more than one prescription per 12 weeks.

⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

BETAMETHASONE VALERATE			
* Scalp app 0.1%9	.84 100	ml OP 🗸	Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%	.26 30 r	ml OP 🗸	<u>Dermol</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%6	.57 100	ml OP 🗸	Locoid
KETOCONAZOLE Shampoo 2%	.23 100	ml OP	Sebizole Sebizole

- a) Maximum of 100 ml per prescription
- b) Only on a prescription

Pinetarsol

DERMATOLOGICALS

Subsidy (Manufacturer's Price) \$ Per

Subsidised er

Fully

Brand or Generic Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

200 g OP

Marine Blue Lotion SPF 50+

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

IMIQUIMOD

Crm 5%, 250 mg sachet......21.72 24 **✓ Perrigo**

PODOPHYLLOTOXIN

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

Contraceptives - Non-hormonal

Condoms

CONDOMS				_
	vailable on a PSO		144	✓ Moments
€ 53 mm			10	✓ Moments
		11.64	144	✓ Moments
 a) Maximum of 60 de 				
b) Up to 60 dev availa				_
53 mm, 0.05 mm thickness	S		10	✓ Moments
		11.42	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
53 mm, chocolate, brown.			10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
53 mm, strawberry, red			10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
56 mm			10	✓ Moments
		11.64	144	✓ Moments
 a) Maximum of 60 de 				
b) Up to 60 dev availa	able on a PSO			
56 mm, 0.05 mm thickness	S		12	✓ Gold Knight
		15.57	144	Gold Knight
 a) Up to 60 dev available 				
b) Maximum of 60 de				
	s (bulk pack)	14.61	144	Gold Knight
 a) Maximum of 60 de 				
b) Up to 60 dev available				
56 mm, 0.08 mm thickness	S	0.97	10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
56 mm, 0.08 mm thickness	s, red	0.97	10	✓ Moments
		11.64	144	✓ Moments
a) Up to 60 dev availa				
b) Maximum of 60 de	v per prescription			
56 mm, chocolate		1.30	12	Gold Knight
		15.57	144	Gold Knight
a) Up to 60 dev availa	able on a PSO			
b) Maximum of 60 de	v per prescription			
56 mm, strawberry		1.30	12	Gold Knight
•		15.57	144	✓ Gold Knight
a) Up to 60 dev availa	able on a PSO			
b) Maximum of 60 de				
		1.42	12	Gold Knight XL
		14.87	144	✓ Shield XL
		17.02		Gold Knight XL

GENITO-URINARY SYSTEM

		Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
		\$	Per	✓	Manufacturer
*	60 mm (bulk pack)	14.87	144	✓ (Gold Knight XL

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IUD 29.1 mm length × 23.2 mm width	29.80	1	✓ 7 MED NSHA Silver/
				Copper Short ✓ Choice 380 7med Nsha Silver/
				copper Short
				✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	29.80	1	✓ Choice
				TT380 Standard

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit: and

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- - 1 Patient is on a Social Welfare benefit; or
 - 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO 10.00 84 ✓ Mercilon 28

✓ Choice Load 375

Subsidy (Manufacturer's Price)		Fully	Brand or Generic
\$	Per	oubsidised ✓	Manufacturer
_			
1.50	84	√ L	o-Oralcon 20 ED
2.18		✓ N	licrogynon 20 ED
6.45	112	√ F	emme-Tab ED
6.62	63		
(16.50)		M	licrogynon 30
hority see SA0500 or	the p	revious pag	ge
_			
1.50	84	√ 0	ralcon 30 ED
1.77		√ L	evlen ED
6.45	112	√ F	emme-Tab ED
	(Manufacturer's Price) \$	(Manufacturer's Price) \$ Per	(Manufacturer's Price) \$ Subsidised Per ✓

(Microgynon 20 ED 1 ab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets to be delisted 1 August 2023) (Femme-Tab ED Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets to be delisted 1 August 2023) (Levlen ED Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets to be delisted 1 August 2023) (Femme-Tab ED Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets to be delisted 1 August 2023)

ETHINYLOESTRADIOL WITH NORETHISTERONE

Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to			
84 tab available on a PSO	12.25	84	✓ Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - Up			
to 84 tab available on a PSO	21.99	84	✓ Norimin
	29.32	112	✓ Norimin

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

GENITO-URINARY SYSTEM

	Subsidy		Fully	Brand or
1)	Manufacturer's Price)		Subsidised	I Generic
	\$	Per	•	Manufacturer
LEVONORGESTREL				
* Tab 30 mcg - Up to 84 tab available on a PSO	16.50	84	1	Microlut
	22.00	112	1	Microlut
* Subdermal implant (2 × 75 mg rods) – Up to 3 pack available				
on a PSO	106.92	1	1	Jadelle
MEDROXYPROGESTERONE ACETATE				
Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PS	O9.18	1	1	Depo-Provera
NORETHISTERONE		·		
	10.05	84	/	Noriday 20
Tab 350 mcg - Up to 84 tab available on a PSO	12.25	04	•	Noriday 28
Emergency Contraceptives				
Linergency Contraceptives				
LEVONORGESTREL				
* Tab 1.5 mg	1.75	1	1	Levonorgestrel
				BNM
	4.95		1	Postinor-1
a) Maximum of 2 tob par proparintian				

- a) Maximum of 2 tab per prescription
- b) Up to 5 tab available on a PSO
- c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.
- d) Levonorgestrel BNM to be Principal Supply on 1 June 2023

(Postinor-1 Tab 1.5 mg to be delisted 1 June 2023)

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- · prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP	A child
(24.15)		Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators	20 g OP	✓ Clomazol
MICONAZOLE NITRATE	-	
* Vaginal crm 2% with applicator	40 g OP	✓ Micreme
•	10 g 01	- Interestine
NYSTATIN		
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ <u>Nilstat</u>
vaginal offir 100,000 a per 3 g with applicator(3)	75 g Oi	· inistat

Brand or

Generic

Manufacturer

Fully

Subsidised

Per

100

✓ Tamsulosin-Rex

Myometrial and Vaginal Hormone Preparations			
ERGOMETRINE MALEATE Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		_	4
PSO	160.00	5	✓ DBL Ergometrine
OESTRIOL	0.00	45 - OD	/ Overally
** Crm 1 mg per g with applicator ** Pessaries 500 mcg		15 g OP 15	✓ <u>Ovestin</u> ✓ Ovestin
OXYTOCIN – Up to 5 inj available on a PSO	0.00	13	• Ovestill
Inj 5 iu per ml, 1 ml ampoule	4 98	5	✓ Oxytocin BNM
Oxytocin BNM to be Principal Supply on 1 June 2023	4.30	3	Oxytociii biliii
Inj 10 iu per ml, 1 ml ampoule	5.98	5	✓ Oxytocin BNM
,			✓ Oxytocin GH S29
Oxytocin BNM to be Principal Supply on 1 June 2023			,
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj availa	ble on a PSO		
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoul		5	✓ Syntometrine
Pregnancy Tests - hCG Urine			
PREGNANCY TESTS - HCG URINE			
a) Up to 200 test available on a PSO			
b) Only on a PSO			
Cassette	12.00	40 test OP	✓ Smith BioMed Rapid
			Pregnancy Test
	16.00		✓ David One Step
			Cassette
			Pregnancy Test
Urinary Agents			
For urinary tract Infections refer to INFECTIONS, Antibacterials, pa	ige 108		
5-Alpha Reductase Inhibitors			
FINASTERIDE - Special Authority see SA0928 below - Retail pha	armacy		
* Tab 5 mg	4.81	100	✓ Ricit
⇒SA0928 Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals valid	without further	r renewal unless	s notified for applications meeting
the following criteria:			
Both:			
1 Patient has symptomatic benign prostatic hyperplasia; and			

Subsidy

(Manufacturer's Price)

\$

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 on the next page - Retail pharmacy

2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

* Cap 400 mcg......22.31

Alpha-1A Adrenoreceptor Blockers

2 Either:

GENITO-URINARY SYSTEM

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 ` \$	Per	✓	Manufacturer

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

* Tab 5 mg	2 100	✓ Alchemy Oxybutynin S29
POTASSIUM CITRATE		
Oral liq 3 mmol per ml - Special Authority see SA1083 below -		
Retail pharmacy31.80	200 ml OP	✓ Biomed

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE			
* Grans eff 4 g sachets	2.22	28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg	2.05	30	✓ Solifenacin Mylan
			✓ Solifenacin Viatris
Tab 10 mg	3.72	30	✓ Solifenacin Mylan
			✓ Solifenacin Viatris

Detection of Substances in Urine

ORTHO-TOLIDINE			
* Compound diagnostic sticks	7.50	50 test OP	
	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	.13.92	100 test OP	✓ Albustix

Obstetric Preparations

Antiprogesterones

MI	FF	PR	IST	ONE	=

Mifegyne	1	O60.00	Tab 200 mg - Up to 15 tab available on a PSO
✓ Mifegyne	3	180.00	

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Price) Subsidised		Generic	
\$	Per	✓	Manufacturer	
				_

Calcium Homeostasis

CALCITONIN			
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ Miacalcic
CINACALCET - Special Authority see SA2170 below -	Retail pharmacy		
Tab 30 mg - Wastage claimable	42.06	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable	84.12	28	✓ Cinacalet Devatis

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
	Per	✓	Manufacturer	

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial	18.00	1	✓ Zoledronic acid
			<u>Mylan</u>
			Zoledronic acid
			Viatris

Corticosteroids and Related Agents for Systemic Use

	Subsidy		Fully	
	(Manufacturer's Price \$) Sub Per	sidised •	
ETHYLPREDNISOLONE ACETATE				
Inj 40 mg per ml, 1 ml vial	47.06	5	1	Depo-Medrol
REDNISOLONE				•
Gral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	/	Redipred
REDNISONE				
€ Tab 1 mg	18.58	500	1	Prednisone Clinect
Tab 2.5 mg		500	1	Prednisone Clinect
Tab 5 mg - Up to 30 tab available on a PSO		500	1	Prednisone Clinect
Tab 20 mg - Up to 30 tab available on a PSO		500		Prednisone Clinect
ETRACOSACTRIN				
FINACOSACTAIN Finj 250 mcg per ml, 1 ml ampoule	75.00	1	J	Synacthen
inj 200 mog per mi, i mi ampoule	13.00	'		UK Synacthen
F Inj 1 mg per ml, 1 ml ampoule	690.00	1		Synacthen Depot
inj i mg per mi, i mi ampoule	090.00	'		Synacthene
			•	Retard \$29
				Hetara 529
RIAMCINOLONE ACETONIDE			_	
Inj 10 mg per ml, 1 ml ampoule		5		Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule	51.10	5	1	Kenacort-A 40
Sex Hormones Non Contraceptive				
Androgen Agonists and Antagonists				
SYPROTERONE ACETATE				•••
Tab 50 mg		50		Siterone
Tab 100 mg	28.03	50	•	<u>Siterone</u>
ESTOSTERONE				
Patch 5 mg per day	225.00	30	1	Androderm
ESTOSTERONE CIPIONATE				
Inj 100 mg per ml, 10 ml vial	85.00	1	1	Depo-Testosterone
, 01- , -	393.00			Taro-
				Testosterone S29
ESTOSTERONE ESTERS				
Inj 250 mg per ml, 1 ml	12.98	1	1	Sustanon Ampoules
ESTOSTERONE UNDECANOATE				
Cap 40 mg — Subsidy by endorsement	21.00	60	/	Andriol Testocaps
Sup to my - Subsidy by Gildorsement	35.00	100		•
Cubaidy by andargament autholdicad for notice to the				Steril-Gene S29
Subsidy by endorsement – subsidised for patients who				
1 November 2021 and the prescription is endorsed acc				
where there exists a record of prior dispensing of testo	sterone undecanoate	cap 40 mg	in the	e preceaing 12 months.

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer **Hormone Replacement Therapy - Systemic** Oestrogens **OESTRADIOL** 28 OP Estrofem Tab 2 mg4.12 28 OP Estrofem Patch 50 mcg per 24 hours7.04 ✓ Climara a) No more than 1 patch per week b) Only on a prescription ✓ Estradot 13.50 ✓ Fetraderm MX S29 a) No more than 2 patch per week b) Only on a prescription Patch 50 mcg per day......7.04 8 ✓ Estradot 50 mcg ✓ Estradiol TDP Mylan S29 14.50 ✓ Estraderm MX S29 a) No more than 2 patch per week b) Only on a prescription Patch 75 mcg per day......7.91 8 ✓ Estradot ✓ Estradiol TDP 10.60 Mylan S29 a) No more than 2 patch per week b) Only on a prescription Patch 100 mcg per day......7.91 ✓ Estradot ✓ Estraderm MX S29 15.50 a) No more than 2 patch per week b) Only on a prescription **OESTRADIOL VALERATE** 84 ✓ Progynova ✓ Progynova **OESTROGENS** Conjugated, equine tab 300 mcg......3.01 28 (17.50)Premarin Conjugated, equine tab 625 mcg......4.12 28 (17.50)Premarin

Progestogens

ME	DROXYPROGESTERONE ACETATE		
*	Tab 2.5 mg	30	✓ Provera
	8.75	56	Provera
*	Tab 5 mg	56	Provera
	17.50	100	Provera
*	Tab 10 mg8.94	30	Provera

	Subsidy (Manufacturer's Pric	e) Sub	Fully Brand or sidised Generic ✓ Manufacturer
Progestogen and Oestrogen Combined Prepara	ntions		
OESTRADIOL WITH NORETHISTERONE			
* Tab 1 mg with 0.5 mg norethisterone acetate		28 OP	171
W. Tab O man with 4 man manathistanana anatata	(18.10)	00.00	Kliovance
* Tab 2 mg with 1 mg norethisterone acetate	(18.10)	28 OP	Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	(10.10)		Miogest
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP	
ocolitation tab (12) and 1 mg ocolitation tab (0)	(18.10)	20 01	Trisequens
	(10110)		· · · · · · · · · · · · · · · · · · ·
Other Oestrogen Preparations			
OESTRIOL			
* Tab 2 mg	7.00	30	✓ Ovestin
-			<u> </u>
Other Progestogen Preparations			
LEVONORGESTREL			
* Intra-uterine device 52 mg	269.50	1	✓ Mirena
* Intra-uterine device 13.5 mg	215.60	1	✓ Jaydess
MEDROXYPROGESTERONE ACETATE			
Tab 100 mg	116.15	100	✓ Provera HD
NORETHISTERONE			
* Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Primolut N
PROGESTERONE			
* Cap 100 mg	14.85	30	✓ Utrogestan
Utrogestan to be Principal Supply on 1 May 2023			
Thyroid and Antithyroid Agents			
Thyroid and Antithyroid Agents			
CARBIMAZOLE			
* Tab 5 mg	7.56	100	✓ Neo-Mercazole
LEVOTHYROXINE			
* Tab 25 mcg	5.55	90	✓ Synthroid
* Tab 50 mcg		28	✓ Mercury Pharma
	5.79	90	Synthroid
¥ Tob 100 mag	64.28	1,000	✓ Eltroxin
* Tab 100 mcg	1./8 6.01	28 90	✓ Mercury Pharma✓ Synthroid
	6.01	1,000	✓ Synthroid ✓ Eltroxin
DDODYLTHOUDAOL Coosiel Authority and CA4400 halves		1,000	- LIUVAIII
PROPYLTHIOURACIL – Special Authority see SA1199 below –		400	/ DTIL
Tab 50 mg	35.00	100	✓ PTU \$29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

_			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA2032 below	- Retail pharm	acy	
*	Inj 5 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 15 mg cartridge	139.50	1	✓ Omnitrope
	, ,			✓ Omnitrope S29 S29

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
<u> </u>	Per	/	Manufacturer	

continued...

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
 - 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
 - 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
 - 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

(Manufacturer ['] s Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred;
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and

(Manufacturer ⁱ s Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

(1	Subsidy	Fully	Brand or
	Manufacturer's Price)	Subsidised	Generic
	\$ F	Per 🗸	Manufacturer

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GOSERELIN	
COCLILLIN	

Implant 3.6 mg, syringe		1	✓ Teva
Implant 10.8 mg, syringe	122.37	1	✓ Teva

I FUPRORFI IN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of

(221.60)

Lucrin Depot 1-month

Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy of \$591.68 per 1 inj with Endorsement.......177.50

Inj 4 mcg per ml, 1 ml67.18

Lucrin Depot 3-month (591.68)

✓ Minirin

10

Vasopressin Agonists

DESMOPRESSIN			
Wafer 120 mcg	47.00	30	Minirin Melt
DESMOPRESSIN ACETATE			
Tab 100 mcg	25.00	30	✓ Minirin
Tab 200 mcg	54.45	30	✓ Minirin
▲ Nasal spray 10 mcg per dose	27.95	6 ml OP	✓ Desmopressin-
• •			PH&T

Other Endocrine Agents

CABERGOI INF

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
Dostinex	2	waived by Special Authority see SA2070 below4.43
✓ Dostinex	8	17.94

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLOMIFFNE CITRATE

✓ Mylan 10 Clomiphen S29

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
METYRAPONE Cap 250 mg	558.00	50	✓ <u>M</u>	etopirone

	Subsidy		Fully Brand or
	(Manufacturer's Price		sidised Generic
	\$	Per	✓ Manufacturer
Anthelmintics			
ALBENDAZOLE - Special Authority see SA1318 below - Retail	nharmacy		
Tab 400 mg		60	✓ Eskazole S29
	403.20	00	LSKAZOIC 625
► SA1318 Special Authority for Subsidy Initial application only from an infectious disease specialist or cl	iniaal miarahialaaiat	Annrovol	a valid for C months where the
patient has hydatids.	inical microbiologist	. Approvai	s valid for 6 months where the
Renewal only from an infectious disease specialist or clinical mic	rohiologiet Annroy	als valid for	6 months where the treatment
remains appropriate and the patient is benefitting from the treatm		ais valia ioi	o months where the treatment
MEBENDAZOLE – Only on a prescription	Orit.		
Tab 100 mg	7 97	6	✓ Vermox
Oral lig 100 mg per 5 ml		15 ml	<u>vermox</u>
Oral ng 100 mg por 0 m	(7.83)	10 1111	Vermox
PRAZIQUANTEL	(1122)		
Tab 600 mg	68.00	8	✓ Biltricide
- Tab 000 mg	00.00	U	Biliticide
Antibacterials			
7 IIII Saoto II III S			
a) For topical antibacterials, refer to DERMATOLOGICALS, page			
b) For anti-infective eye preparations, refer to SENSORY ORGA	NS, page 247		
Cephalosporins and Cephamycins			
Cephalosporins and CephalityCins			
CEFACLOR MONOHYDRATE			
Cap 250 mg	25.85	100	✓ Ranbaxy-Cefaclor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.75	100 ml	✓ Ranbaxy-Cefaclor
CEFALEXIN			
Cap 250 mg	3.85	20	✓ Cephalexin ABM
Cap 500 mg	5.85	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	7.88	100 ml	✓ Flynn
Grans for oral liq 50 mg per ml - Wastage claimable	10.38	100 ml	✓ Flynn
CEFAZOLIN – Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance with a	a Health NZ Hospita	l approved	protocol and the prescription is
endorsed accordingly.			
Inj 500 mg vial	3.39	5	✓ <u>AFT</u>
Inj 1 g vial	3.49	5	✓ <u>AFT</u>
CEFTRIAXONE – Subsidy by endorsement			
a) Up to 10 inj available on a PSO			
 b) Subsidised only if prescribed for a dialysis or cystic fibrosit 			
pelvic inflammatory disease, or the treatment of suspecte	d meningococcal dis	sease, and	the prescription or PSO is
endorsed accordingly.			
Inj 500 mg vial		1	✓ Ceftriaxone-AFT
Inj 1 g vial	3.59	5	✓ Ceftriaxone-AFT
CEFUROXIME AXETIL – Subsidy by endorsement			
Only if prescribed for prophylaxis of endocarditis and the pres			
Tab 250 mg	45.93	50	✓ Zinnat

(Zinnat Tab 250 mg to be delisted 1 March 2024)

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	 Manufacturer 	

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable1	6.97	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Atypical mycobacterial infection; or
 - 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	E-Mycin
a) Up to 20 tab available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
 a) Up to 300 ml available on a PSO 			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
 a) Up to 200 ml available on a PSO 			
b) Wastage claimable			
ROXITHROMYCIN			
Tab 150 mg	13.19	50	Arrow- Roxithromycin
Tab 300 mg	25.00	50	✓ Arrow- Roxithromycin

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully idised	
Penicillins				
AMOXICILLIN				
Cap 250 mg	43.45	500	1	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP			_	
Cap 500 mg	66.44	500		Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	/	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable	4.70	4001	,	All
Grans for oral liq 250 mg per 5 ml	1./3	100 ml	•	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable Inj 250 mg vial	15 07	10	1	Ibiamox
Inj 500 mg vial		10		Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO		10		Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID		. •		
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab available on a PSO	0.80	10	1	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25		10	٠	Culaili Duo 300/123
per ml		100 ml	1	Augmentin
a) Up to 200 ml available on a PSO		1001111	•	Augmontin
b) Wastage claimable				
Grans for oral lig amoxicillin 50 mg with clavulanic acid 12.5	ma			
per ml – Up to 200 ml available on a PSO		00 ml OP	1	Curam
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj				
available on a PSO	375 97	10	1	Bicillin LA
			-	
BENZYLPENICILLIN SODIUM [PENICILLIN G] Inj 600 mg (1 million units) vial – Up to 5 inj available on a P	SO 11.00	10	_	Sandoz
	30 11.09	10	•	Januoz
FLUCLOXACILLIN Cap 250 mg – Up to 30 cap available on a PSO	15.70	250	_	Flucloxacillin-AFT
Cap 500 mg - Up to 30 cap available on a PSO		500		Flucioxacillin-AFT
Grans for oral liq 25 mg per ml		100 ml		AFT
a) Up to 200 ml available on a PSO		1001111	-	<u></u>
b) Wastage claimable				
Grans for oral liq 50 mg per ml	3.68	100 ml	1	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Inj 250 mg vial	17.56	10	✓	Flucloxin
Inj 500 mg vial	18.87	10		Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.70	5	1	Flucil

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised		
PHENOXYMETHYLPENICILLIN (PENICILLIN V)					•
Cap 250 mg - Up to 30 cap available on a PSO	3.84	50	✓	Cilicaine VK	
Cap 500 mg	6.86	50	✓	Cilicaine VK	
a) Up to 20 cap available on a PSO					
b) Up to 2 x the maximum PSO quantity for RFPP					
Grans for oral liq 125 mg per 5 ml	3.40	100 m		<u>AFT</u>	
a) Up to 200 ml available on a PSO					
b) Wastage claimable					
Grans for oral liq 250 mg per 5 ml	4.24	100 m		<u>AFT</u>	
a) Up to 300 ml available on a PSO					
b) Up to 2 x the maximum PSO quantity for RFPP					
c) Wastage claimable					

Tetracyclines

DO	XYCYCLINE			
*	Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	✓ Doxine
MIN	NOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	19.32	100	
		(52.04)		Minomycin

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has

 ${\sf TETRACYCLINE\ - Special\ Authority\ see\ SA1332\ below\ - \ Retail\ pharmacy}$

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 62

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
CLINDAMYCIN				
Cap hydrochloride 150 mg	5.30	24	✓	Dalacin C
Inj 150 mg per ml, 4 ml ampoule	35.10	10	✓	Hameln
	39.00		✓	Dalacin C
(Dalacin C Inj 150 mg per ml, 4 ml ampoule to be delisted 1 A	ugust 2023)			
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist	- Subsidy by endorseme	ent		
Only if prescribed for dialysis or cystic fibrosis patient and	the prescription is endo	rsed	accordingly	<i>I</i> .
Inj 150 mg		1		Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorseme	nt95.00	5	1	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patie		trac	t infection a	and the prescription is
endorsed accordingly.				
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	nt91.00	5	✓	Wockhardt S29
	182.00	10	✓	Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patie	nt or complicated urinary	trac	t infection a	and the prescription is
endorsed accordingly.				
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement		10		Pfizer
	87.50	50		Pfizer
Only if prescribed for a dialysis or cystic fibrosis patie endorsed accordingly.	nt or complicated urinary	trac	t infection a	and the prescription is
MOXIFLOXACIN - Special Authority see SA1740 below - Re	etail pharmacy			
No patient co-payment payable	•			

⇒SA1740 Special Authority for Subsidy

Initial application — (**Tuberculosis**) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*: and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications: or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Fither:

INFECTIONS - AGENTS FOR SYSTEMIC USE				
	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
continued				
2.1 Has tried and failed to clear infection using azithrom2.2 Has laboratory confirmed azithromycin resistance; a3 Treatment is only for 7 days.	•			
Initial application — (Penetrating eye injury) only from an ophti requires prophylaxis following a penetrating eye injury and treatment Note: Indications marked with * are unapproved indications.	0 11		d for 1 mo	nth where the patient
PAROMOMYCIN - Special Authority see SA1689 below - Retail	pharmacy			
Cap 250 mg		16	✓ H	umatin S29
⇒SA1689 Special Authority for Subsidy				
Initial application only from an infectious disease specialist, clinic month for applications meeting the following criteria: Either:	al microbiologist or o	gastroer	nterologist	. Approvals valid for 1
Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage.				
Renewal only from an infectious disease specialist, clinical microb applications meeting the following criteria: Either:	iologist or gastroente	erologis	t. Approv	als valid for 1 month for
1 Patient has confirmed cryptosporidium infection; or2 For the eradication of Entamoeba histolyica carriage.				
PYRIMETHAMINE - Special Authority see SA1328 below - Retai	l pharmacy			
Tab 25 mg	48.00	30	✓ D	araprim S29
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of	a period of 3 months		ss notified	d for applications meeting
SODIUM FUSIDATE [FUSIDIC ACID]				
Tab 250 mg	67.85	36	✓ F	ucidin
SULFADIAZINE SODIUM - Special Authority see SA1331 below	 Retail pharmacy 			
Tab 500 mg ≫SA1331 Special Authority for Subsidy	543.20	56	✓ W	ockhardt \$29
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following:	without further renev	wal unle	ss notified	for applications meeting
For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months or		; or		
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	18.50	5	✓ <u>To</u>	obramycin Mylan iatris
Only if prescribed for dialysis or cystic fibrosis patient and	the prescription is e	ndorsec	d accordin	gly.
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by	005.00	n ala		ahuamusia DAIIA
endorsement	395.00 56	6 dose	→ <u>To</u>	obramycin BNM

b) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly.

a) Wastage claimable

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
TRIMETHOPRIM				
* Tab 300 mg - Up to 30 tab available on a PSO	18.55	50	√ <u>1</u>	<u>IMP</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	AZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - l				
to 30 tab available on a PSO	64.80	500	✓ 1	<u> </u>
★ Oral liq 8 mg sulphamethoxazole 40 mg per ml - Up to 200				
available on a PSO	2.97	100 m	nl 🗸 🛚	Deprim
VANCOMYCIN – Subsidy by endorsement	r nranhulavia af anda	aa ralit	io or for troo	tment of Cleatridium
Only if prescribed for a dialysis or cystic fibrosis patient or for difficile following metronidazole failure and the prescription is			is or for trea	ument of Clostridium
Inj 500 mg vial		1	✓ N	<u>Mylan</u>

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 63
- b) For topical antifungals refer to GENITO URINARY, page 76

FLUCONAZOLE

Cap 50 mg	2.75	28	✓ Dizole
Cap 150 mg	0.65	1	✓ Mylan ✓ Mylan
Cap 200 mg		28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority see SA1359 below - Retail pharmacy	109.34	35 ml	✓ Diflucan

(Dizole Cap 50 mg to be delisted 1 August 2023)

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised: and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

	Subsidy (Manufacturer's Price \$	e) S	Fully Subsidised	Brand or Generic Manufacturer	
ITRACONAZOLE Cap 100 mg	4.27	15	√ II	trazole	
Oral liq 10 mg per ml – Special Authority see SA1322 below Retail pharmacy	-	50 ml O	P 🗸 S	Sporanox	

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE			
Tab 200 mg - PCT	CBS	30	✓ Burel S29
•			✓ Link Healthcare S29
			✓ Nizoral \$29
		100	✓ Strides Shasun S29
		100	✓ Taro \$29
(Link Healthcare S29 Tab 200 mg to be delisted 1 July 2023)			
(Nizoral S29 Tab 200 mg to be delisted 1 July 2023)			
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Re	etail pharmacy		
Tab modified-release 100 mg	206.00	24	✓ Posaconazole Juno
Oral liq 40 mg per ml	342.51	105 ml OP	✓ Devatis
	761.13		✓ Noxafil
Devatis to be Principal Supply on 1 May 2023			

Devatis to be Principal Supply on 1 May 2023

(Noxafil Oral liq 40 mg per ml to be delisted 1 May 2023)

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
TERBINAFINE				
* Tab 250 mg	8.15	84	✓ <u>D</u>	<u>eolate</u>
VORICONAZOLE - Special Authority see SA1273 below - Reta	il pharmacy			
Tab 50 mg	91.00	56	✓ V	ttack
Tab 200 mg	350.00	56	✓ V	ttack
Powder for oral suspension 40 mg per ml - Wastage				
claimable	1,523.22	70 ml	✓ V	fend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient is immunocompromised; and
 - 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
 - 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

INFECTIONS - AGENTS FOR SYSTEMIC US	E		
	Subsidy (Manufacturer's Price \$) Sub Per	Fully Brand or sidised Generic Manufacturer
Antitrichomonal Agents			
METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO		21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml		100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			_
Tab 500 mg	36.16	10	✓ <u>Arrow-Ornidazole</u>
Antituberculotics and Antileprotics			
Note: There is no co-payment charge for all pharmaceuticals lis	ted in the Antituberco	ulotics and	Antileprotics group regardless of
immigration status.			1 0 1 0
CLOFAZIMINE - Retail pharmacy-Specialist			
a) No patient co-payment payable			
 b) Prescriptions must be written by, or on the recommendat dermatologist. 		disease ph	nysician, clinical microbiologist or
* Cap 50 mg	442.00	100	✓ Lamprene S29
CYCLOSERINE - Retail pharmacy-Specialist			
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendat	ion of, an infectious	disease ph	nysician, clinical microbiologist or
respiratory physician.	044.00		(0.1.1.0
Cap 250 mg	344.00	60	✓ Cyclorin S29
DAPSONE - Retail pharmacy-Specialist			
a) No patient co-payment payable		ا اا	
 b) Prescriptions must be written by, or on the recommendat dermatologist 	ion of, an infectious	disease pr	nysician, clinical microbiologist or
Tab 25 mg	268.50	100	✓ Dapsone
Tab 100 mg		100	✓ Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Speciali			·
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendat	ion of, an infectious	disease ph	nysician, clinical microbiologist or
respiratory physician			
Tab 100 mg		100	✓ EMB Fatol S29
Tab 400 mg	49.34	56	✓ Myambutol S29
ISONIAZID - Retail pharmacy-Specialist			
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendat	ion of, an internal m	edicine phy	ysician, paediatrician, clinical
microbiologist, dermatologist or public health physician * Tab 100 mg	23.00	100	✓ PSM
ISONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist	20.00	100	<u> </u>
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendat	ion of, an internal me	edicine phy	vsician, paediatrician, clinical
microbiologist, dermatologist or public health physician	,	,,	, , , ,
* Tab 100 mg with rifampicin 150 mg		100	✓ <u>Rifinah</u>
* Tab 150 mg with rifampicin 300 mg	179.13	100	✓ <u>Rifinah</u>

	Subsidy (Manufacturer's Price \$) Su Per	Fully bsidised	Brand or Generic Manufacturer
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Speciali	st			
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommer respiratory physician				_
Grans for oral liq 4 g sachet	280.00	30	✓ P:	aser S29
PROTIONAMIDE – Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommer respiratory physician 				
Tab 250 mg	305.00	100	→ P	eteha S29
PYRAZINAMIDE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommer respiratory physician		disease p	hysician,	clinical microbiologist or
* Tab 500 mg	64.95	100	✓ A	FT-Pyrazinamide
RIFABUTIN - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommer gastroenterologist				
* Cap 150 mg	353.71	30	✓ M	lycobutin
RIFAMPICIN – Subsidy by endorsement				
 a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus infecting antimicrobial based on susceptibilities and the prescriberation of the payable and in paediatrician, or public health physician. 	ption is endorsed accord nternal medicine physici	dingly; ca an, clinica	n be waiv Il microbio	ed by endorsement - ologist, dermatologist,
* Cap 150 mg		100 100	_	<u>ifadin</u> ifadin
* Cap 300 mg * Oral liq 100 mg per 5 ml		60 ml	_	<u>ifadin</u> ifadin
- Ordering for orinin	12.00	00 1111	• 11	<u>naam</u>
Antivirals				
For eye preparations refer to Eye Preparations, Anti-Infective	Preparations, page 247	7		
Hepatitis B Treatment				
ENTECAVIR			_	
* Tab 0.5 mg	52.00	30		ntecavir Mylan
			√ E	ntecavir Sandoz
LAMIVUDINE – Special Authority see SA1685 below – Reta		00	17	-41- ···
Tab 100 mg Oral liq 5 mg per ml		28 40 ml OP		<u>etlam</u> effix
⇒SA1685 Special Authority for Subsidy	270.00	40 1111 01	V 2.	CIIIA
Initial application only from a relevant specialist or medical	oractitioner on the recon	nmendatio	n of a rel	evant specialist
Approvals valid for 1 year where used for the treatment or pre		mondan	,,, o, a ,o,	ovani oposiansi.
Renewal from any relevant practitioner. Approvals valid for 2	2 years where used for t	he treatm	ent or pre	evention of hepatitis B.
TENOFOVIR DISOPROXIL				
Tenofovir disoproxil prescribed under endorsement for the		cluded in t	he count	of up to 4 subsidised
antiretrovirals for the purposes of Special Authority SA21 * Tab 245 mg (300 mg as a maleate)		30	✓ <u>T</u> (enofovir Disoproxil Mylan

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
Herpesvirus Treatments				
ACICLOVIR				
* Tab dispersible 200 mg	1.78	25	✓ L	ovir
* Tab dispersible 400 mg	5.81	56	✓ L	ovir
* Tab dispersible 800 mg	6.46	35	✓ L	<u>ovir</u>
VALACICLOVIR				
Tab 500 mg	6.50	30	✓ V	aclovir
Tab 1,000 mg	13.76	30	✓ V	aclovir
VALGANCICLOVIR - Special Authority see SA1993 below - Re	tail pharmacy			
Tab 450 mg		60	✓ <u>V</u>	alganciclovir Mylan

⇒SA1993 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient is immunocompromised; and

Subsidy (Manufacturer's Pr	ice)	Fully Subsidised	Brand or Generic	
(Wandacard STT)	Per	✓ Cubsidised	Manufacturer	

continued...

- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Patient is immunocompromised; and
 - 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see \$A2138 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 105 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

*	Tab 200 mg with ter	nofovir disoproxil 245 mg) (300 mg as a

30

✓ Tenofovir Disoproxil **Emtricitabine** Mylan

✓ Tenofovir Disoproxil **Emtricitabine Viatr**

⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Roth:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion: and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV. Viral Hepatitis and Sexual Health Medicine clinical auidelines:

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion: and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on Pharmac's website) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

✓ Lagevrio

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — **(Confirmed HIV)** only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

- Both:
 - 1 Treatment course to be initiated within 72 hours post exposure; and
 - 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (**Percutaneous exposure**) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA2139 on the previous	is page – Retail phai	rmacy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA2139 on the previo	us page – Retail pha	armacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA2139 on the previo	us page – Retail pha	armacy	
Tab 200 mg	84.00	60	✓ Nevirapine
			<u>Alphapharm</u>
			Nevirapine Viatris
Oral suspension 10 mg per ml	203.55	240 ml OP	✓ Viramune
			Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA2139 on the Tab 300 mg Oral lig 20 mg per ml	180.00	Retail pharmad 60 240 ml OP	Σy ✓ Ziagen ✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.	see SA2139 or		
Tab 600 mg with lamivudine 300 mg	29.50	30	✓ Abacavir/ Lamivudine Viatris
About in the section of the Winds in the Department Operation of	75.00		✓ Kivexa

Abacavir/Lamivudine Viatris to be Principal Supply on 1 May 2023 (Kivexa Tab 600 mg with lamivudine 300 mg to be delisted 1 May 2023)

	Subsidy (Manufacturer's Pr \$		Fully Brand or dised Generic Manufacturer
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOP	POVII Special	Authority coo S	
pharmacy	TIOXIL - Special	Authority See C	A2100 oil page 100 - Helali
Note: Efavirenz with emtricitabine and tenofovir disoproxil c anti-retroviral Special Authority	ounts as three an	ti-retroviral med	lications for the purposes of the
Tab 600 mg with emtricitabine 200 mg and tenofovir disopro	xil		
245 mg (300 mg as a maleate)		30	✓ Mylan✓ Viatris
EMTRICITABINE - Special Authority see SA2139 on page 105	- Retail pharmacy	/	
Cap 200 mg	307.20	30	✓ Emtriva
LAMIVUDINE - Special Authority see SA2139 on page 105 - R	etail pharmacy		
Tab 150 mg		60	✓ Lamivudine
ŭ			Alphapharm
			✓ Lamivudine Viatris
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3TC
ZIDOVUDINE [AZT] - Special Authority see SA2139 on page 10	05 – Retail pharma	acv	
Cap 100 mg		100	✓ Retrovir
Oral liq 10 mg per ml		200 ml OP	✓ Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE - Special Authority see	e SA2139 on page	e 105 – Retail n	harmacy
Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority.			•
Tab 300 mg with lamivudine 150 mg	33.00	60	✓ Alphapharm
Protease Inhibitors			
ATAZANAVIR SULPHATE - Special Authority see SA2139 on p	page 105 – Retail	pharmacy	
Cap 150 mg	•	60	✓ Atazanavir Mylan
	141.68		✓ Teva
Atazanavir Mylan to be Principal Supply on 1 May 2023			
Cap 200 mg		60	Atazanavir Mylan
	188.91		✓ Teva
Atazanavir Mylan to be Principal Supply on 1 May 2023 (Teva Cap 150 mg to be delisted 1 May 2023) (Teva Cap 200 mg to be delisted 1 May 2023)			
DARUNAVIR - Special Authority see SA2139 on page 105 - Re	etail nharmacy		
Tab 400 mg		60	✓ <u>Darunavir Mylan</u> ✓ Darunavir Viatris
Tab 600 mg	196.65	60	✓ <u>Darunavir Mylan</u>✓ <u>Darunavir Viatris</u>
(Darunavir Mylan Tab 600 mg to be delisted 1 August 2023)			
LOPINAVIR WITH RITONAVIR - Special Authority see SA2139	on page 105 – R	etail nharmacy	
Tab 100 mg with ritonavir 25 mg		60	✓ Lopinavir/Ritonavir
			Mylan
Tab 200 mg with ritonavir 50 mg	295.00	120	✓ <u>Lopinavir/Ritonavir</u> Mylan
Oral liq 80 mg with ritonavir 20 mg per ml	735.00	300 ml OP	✓ Kaletra
RITONAVIR - Special Authority see SA2139 on page 105 - Ref			
Tab 100 mg		30	✓ Norvir

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Strand Transfer Inhibitors

DOLUTEGRAVIR - Special Authority see SA2139 on page	105 - Retail pharmacy		
Tab 50 mg	1,090.00	30	Tivicay
RALTEGRAVIR POTASSIUM - Special Authority see SA2	139 on page 105 – Reta	il pharmacy	
Tab 400 mg	1,090.00	60	✓ Isentress
Tab 600 mg	1.090.00	60	✓ Isentress HD

Immune Modulators

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

Inj 180 mcg prefilled syringe......500.00

✓ Pegasys

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued...

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Fither:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*: or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*: or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Initial application — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	19.95	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg - Up to 15 cap available on a	ı		
PSO	86.40	100	✓ Macrobid
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	245.00	100	✓ Arrow-Norfloxacin
Only if prescribed for a patient with an uncomplicated up		on that is unre	esponsive to a first line agent o

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per		Manufacturer
Anticholinesterases				
Antichonnesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	✓ M	lax Health
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	50.28	100	✓ M	lestinon
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENIAC CODILIM				
DICLOFENAC SODIUM * Tab EC 25 mg	1.00	50	√ n	iclofenac Sandoz
* Tab EC 25 flig * Tab 50 mg dispersible		20		oltaren D
* Tab EC 50 mg		50		iclofenac Sandoz
* Tab long-acting 75 mg		100	_	oltaren SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a F		5		oltaren
* Suppos 12.5 mg		10		oltaren
* Suppos 25 mg		10		oltaren
* Suppos 50 mg – Up to 10 supp available on a PSO		10		oltaren
* Suppos 100 mg		10	-	oltaren
		10	•	Ollaron
IBUPROFEN	01.40	1 000	/ D	allava
* Tab lang action 800 mg		1,000		elieve
* Tab long-acting 800 mg		30 200 m	_	rufen SR
* Oral liq 20 mg per ml	11.29	200 II	_	thics
	11.29			enpaed 100 mg per 5 ml
KETODDOFFN				J IIII
KETOPROFEN	10.07	00		
* Cap long-acting 200 mg	12.07	28	• 0	ruvail SR
MEFENAMIC ACID				
* Cap 250 mg	1.25	50		
	(10.82)		P	onstan
	0.50	20		
	(7.50)		P	onstan
NAPROXEN				
* Tab 250 mg	32.69	500		oflam 250
* Tab 500 mg	28.71	250	_	oflam 500
* Tab long-acting 750 mg	6.47	28	_	aprosyn SR 750
* Tab long-acting 1 g	8.62	28	✓ <u>N</u>	aprosyn SR 1000
TENOXICAM				
* Tab 20 mg	18.50	100	✓ Ti	ilcotil
* Inj 20 mg vial	9.95	1	✓ A	FT
NSAIDs Other				
CELECOXIB				
Cap 100 mg	3.45	60	1 0	elebrex
Oap 100 mg		00		elecoxib Pfizer
Cap 200 mg	3 20	30	_	elebrex
очр 200 mg		00	_	elecoxib Pfizer
			- 0	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	/	Manufacturer

Topical Products for Joint and Muscular Pain

CAPSAICIN

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary)*, and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroguine. Note: Indication marked with a * is an unapproved indication.

* Tab 200 mg			✓ Plaquenil
LEFLUNOMIDE			
Tab 10 mg	6.00	30	✓ Arava
Tab 20 mg	6.00	30	✓ Arava
PENICILLAMINE			
Tab 125 mg	67.23	100	D-Penamine
Tab 250 mg	110.12	100	D-Penamine

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODIUM		
* Tab 70 mg2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL		
* Tab 70 mg with colecalciferol 5,600 iu	4	✓ Fosamax Plus

Other Treatments

DENOSUMAB – Special Authority see SA1777 below – R	letail pharmacy		
Inj 60 mg prefilled syringe	326.00	1	Prolia

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial		1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial		1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA1	779 below – Retail	pharmacy	
* Tab 60 mg	53.76	28	✓ Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

Tab 35 mg	2.50	4	Risedronate Sandoz
Risedronate Sandoz to be Principal Supply on 1 June 2023			
TERIPARATIDE - Special Authority see SA1139 below - Retail pharm	nacy		
Inj 250 mcg per ml, 2.4 ml	490.00	1	✓ Forteo

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

✓ Aclasta

100 ml OP

	M	USCULO	SKEL	LETAL SYSTEM
(1)	Subsidy Manufacturer's Price \$) Sub Per	Fully sidised	Brand or Generic Manufacturer
continued c) A vertebral fracture is defined as a 20% or greater reduction relative to the posterior height of that body, or a 20% or great body above or below the affected vertebral body. d) A maximum of 18 months of treatment (18 cartridges) will be	ter reduction in a			
ZOLEDRONIC ACID Inj 0.05 mg per ml, 100 ml, bag	22.53 1	00 ml OP		Coledronic Acid Viatris Coledronic-US 829

Hyperuricaemia and Antigout

Zoledronic Acid Viatris to be Principal Supply on 1 June 2023 Inj 0.05 mg per ml, 100 ml, vial.......60.00

(Aclasta Inj 0.05 mg per ml, 100 ml, vial to be delisted 1 June 2023)

		•			
ALL	OPURINOL				
*	Tab 100 mg		11.47	500	✓ DP-Allopurinol
*	Tab 300 mg		28.57	500	✓ DP-Allopurinol
BEI	NZBROMARONE -	- Special Authority see SA1963 belo	w – Retail pharmacy		
	Tab 50 mg		22.50	100	✓ Narcaricin mite \$29
	Tab 100 mg		13.50	30	✓ Desuric S29
					✓ Urinorm S29
			45.00	100	✓ Benzbromaron AL
					100 \$29

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg6.0	0 100	✓ Colgout
FEBUXOSTAT - Special Authority see SA2054 below - Retail pharmacy		
Tab 80 mg20.0	0 28	✓ Febuxostat
		multichem
Tab 120 mg20.0	0 28	✓ Febuxostat
		<u>multichem</u>

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required

WIGGOLOSKELLTAL STOTEM				
	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer
continued				
and serum urate remains greater than 0.36 m maximum tolerated dose; or	nmol/I despite use of probe	enecid at do	ses of	f up to 2 g per day or
2.3 The patient has renal impairment such that pr remains greater than 0.36 mmol/l despite opti	imal treatment with allopu	inol (see N	ote); o	r
2.4 The patient has previously had an initial Spec Initial application — (Tumour lysis syndrome) only from applications meeting the following criteria: Both:				•
1 Patient is scheduled to receive cancer therapy carryi2 Patient has a documented history of allopurinol intole		risk of tum	our lys	sis syndrome; and
Renewal — (Gout) from any relevant practitioner. Approva	als valid for 2 years where	the treatme	ent ren	mains appropriate and the
Renewal — (Tumour lysis syndrome) only from a haema treatment remains appropriate and the patient is benefitting PROBENECID		orovals vali	d for 6	weeks where the
* Tab 500 mg	66.95	100	√ F	Probenecid-AFT
Muscle Relaxants				
BACLOFEN				
* Tab 10 mg		100	-	Pacifen
Inj 0.05 mg per ml, 1 ml ampoule – Subsidy by endorse Subsidised only for use in a programmable pump ir caused intolerable side effects and the prescription	n patients where oral antis	1 pastic agen	_	Lioresal Intrathecal e been ineffective or have
Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorseme Subsidised only for use in a programmable pump ir caused intolerable side effects and the prescription	ent306.82 n patients where oral antis	5 pastic agen	_	Medsurge re been ineffective or have

✓ Dantrium

✓ Dantrium

✓ Norflex

✓ Dantrium S29 S29

100

100

100

DANTROLENE

ORPHENADRINE CITRATE

Cap 25 mg.......112.13

Cap 50 mg......77.00

Tab 100 mg20.76

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg	38.24	60	 Symmetrel
	63.73	100	Symmetrel
APOMORPHINE HYDROCHLORIDE			
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	121.84	5	✓ Movapo
ENTACAPONE			
▲ Tab 200 mg	18.04	100	✓ Comtan
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg	15.80	100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
* Tab 100 mg with carbidopa 25 mg	21.11	100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg	43.65	100	✓ Sinemet CR
* Tab 250 mg with carbidopa 25 mg	38.39	100	✓ Sinemet
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	5.51	100	✓ Ramipex
▲ Tab 1 mg	18.66	100	✓ Ramipex
RASAGILINE			
* Tab 1 mg	53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	4.05	84	✓ Ropin
▲ Tab 1 mg	4.95	84	✓ Ropin
▲ Tab 2 mg	6.48	84	✓ Ropin
▲ Tab 5 mg	14.50	84	✓ Ropin

SELEGILINE HYDROCHLORIDE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of selegiline hydrochloride.

* Tab 5 mg	48.00	100	✓ Eldepryl S29
TOLCAPONE			
▲ Tab 100 mg	152.38	100	✓ Tasmar

Anticholinergics

BENZATROPINE MESYLATE			
Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
a) Up to 10 inj available on a PSO			
b) Only on a PSO			

PROCYCLIDINE HYDROCHLORIDE



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE - Special Authority see SA1403 below - Retail pharmacy

Wastage claimable

⇒SA1403 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Anaesthetics

Local

LIDOCAINE [LIGNOCAINE]

- a) Up to 150 ml available on a PSO
- b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

- a) Up to 5 each available on a PSO
- Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.

	Subsidy	•	Fully	Brand or
	(Manufacturer's Price)	Subsidised	I Generic
	\$	Per	•	Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	44.00	200 m	/	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO		25	1	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	9.00 [°]	25	1	Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO	6.20 [°]	5	/	Lidocaine-Claris
	6.85		1	Lidocaine-Baxter
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	7.15	5	1	Lidocaine-Baxter
(Lidocaine-Claris Inj 1%, 20 ml vial to be delisted 1 June 2023)				
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –				
Subsidy by endorsement	103 32	10	1	Pfizer
· ·	100.32	10	•	F IIZCI
a) Up to 5 each available on a PSO				

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

⇒SA0906 Special Authority for Subsidy

Topical Local Anaesthetics

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDUCAINE [LIGNOCAINE] - Special Authority see SA0906 and	ve – Retali pnar	macy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Author	ority see SA0906	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓ EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 111

Non-opioid	Analgesics
------------	------------

•			
ASPIRIN			
* Tab dispersible 300 mg - Up to 30 tab available on a PS	O4.50	100	Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia of accordingly.	or diabetic peripheral	neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.14	57 g OP	✓ Rugby Capsaicin Topical Cream S29
NEFOPAM HYDROCHLORIDE			
Tab 30 mg	23.40	90	✓ Acupan

	Subsidy		Fully	Brand or
	(Manufacturer's Price)) Subs	idised	Generic Manufacturer
ARACETAMOL	Ψ	101		Wallardolardi
Tab 500 mg - blister pack	19.75	1,000	✓ Pa	acimol
a) Maximum of 300 tab per prescription; of the control of the		1,000	· <u></u>	
b) Up to 30 tab available on a PSO c)	an so warrou sy ondolosmone			
Subsidy by endorsement for high	er quantities is available for patien	its with long	term co	onditions who require
regular daily dosing for one montl	n or greater, and the prescription is	s annotated	I accordi	ingly. Pharmacists may
annotate the prescription as endo	rsed where dispensing history sup	ports a lon	g-term c	ondition.
Maximum of 100 tab per dispensi				
	dispense in repeat dispensings no	ot exceeding	g 100 tal	b per dispensing.
Tab 500 mg - bottle pack – Maximum of 300 ta		4 000		
prescription; can be waived by endorseme	nt 17.92	1,000	_	<u>oumed</u> Paracetamol
 Subsidy by endorsement for higher q 	•	•		, ,
daily dosing for one month or greater				macists may annotate tl
prescription as endorsed where dispe	0 ,			
2) Maximum of 100 tab per dispensing t	•			,
non-endorsed patients), then dispens	se in repeat dispensings not excee	aing 100 ta	ab per di	spensing.
Oral lig 120 mg per 5 ml	3.98	200 ml	✓ Pa	aracetamol
2 1 3 Pr				(Ethics)
	5.45	1,000 ml	✓ Pa	aracare
	10.50 20	00 ml OP	✓ A\	/allon
 a) Maximum of 600 ml per prescription; ca 	an be waived by endorsement			
b) Up to 200 ml available on a PSO				
c) Not in combination				
d)				1000 1/1
Maximum of 200 ml per dispensir				
non-endorsed patients), then dispersion of the control of the cont	ense in repeat dispensing not exc			
, , ,	or greater and the prescription is	·		
3 , 3	escription as endorsed where disp			0,
condition.	sociipiion de endereed where diep	ronomy mot	ory oupp	one a long term
e) Paracetamol (Ethics) to be Principal Su	ipply on 1 June 2023			
Oral liq 250 mg per 5 ml	3.35	200 ml	✓ Pa	amol
a) Maximum of 600 ml per prescription; ca	an be waived by endorsement			
b) Up to 200 ml available on a PSO	•			
 c) Not in combination 				
d)				
	ng for non-endorsed patients. If qu			,
, , ,				
non-endorsed patients), then disp		its with long	•	
non-endorsed patients), then disp 2) Subsidy by endorsement for high		endoread	or annot	ated accordingly
non-endorsed patients), then disp 2) Subsidy by endorsement for high- regular daily dosing for one montl	n or greater and the prescription is			0,
non-endorsed patients), then disp 2) Subsidy by endorsement for high- regular daily dosing for one montl				0,
non-endorsed patients), then disp 2) Subsidy by endorsement for high- regular daily dosing for one montl Pharmacists may annotate the pr condition.	n or greater and the prescription is escription as endorsed where disp			oorts a long-term
non-endorsed patients), then disp 2) Subsidy by endorsement for high- regular daily dosing for one montl Pharmacists may annotate the pr	n or greater and the prescription is escription as endorsed where disp	ensing hist	ory supp	ports a long-term
non-endorsed patients), then disp 2) Subsidy by endorsement for high- regular daily dosing for one montl Pharmacists may annotate the pr condition. Suppos 125 mg	n or greater and the prescription is escription as endorsed where disp3.594.1812.40	pensing historement	ory supp ✓ G a	oorts a long-term acet acet

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	Fully Brand or ised Generic Manufacturer
Opioid Analgesics			
CODEINE PHOSPHATE – Safety medicine; prescriber may determine Tab 15 mg		quency 100	✓ Noumed ✓ PSM
Noumed to be Principal Supply on 1 May 2023 Tab 30 mg	6.98	100	✓ Aspen ✓ Noumed
Tab 60 mg(PSM Tab 15 mg to be delisted 1 May 2023)	13.89	100	✓ Noumed
DIHYDROCODEINE TARTRATE Tab long-acting 60 mg FENTANYL	8.60	60	✓ <u>DHC Continus</u>
a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fr Inj 50 mcg per ml, 2 ml ampoule Inj 50 mcg per ml, 10 ml ampoule Patch 12.5 mcg per hour Patch 25 mcg per hour Patch 50 mcg per hour Patch 75 mcg per hour	3.75 9.41 6.99 7.99 9.49	10 10 5 5 5 5	✓ Boucher and Muir ✓ Boucher and Muir ✓ Fentanyl Sandoz ✓ Fentanyl Sandoz ✓ Fentanyl Sandoz ✓ Fentanyl Sandoz
Patch 100 mcg per hour	equency reimbursed at the rate		Fentanyl Sandoz
Tab 5 mg		10 200 ml 200 ml 200 ml 10	✓ Methadone BNM ✓ Biodone ✓ Biodone Forte ✓ Biodone Extra Forte ✓ AFT
b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fr Oral liq 1 mg per ml Oral liq 2 mg per ml Oral liq 5 mg per ml		200 ml 200 ml 200 ml 200 ml	✓ RA-Morph ✓ RA-Morph ✓ Ordine S29 ✓ RA-Morph ✓ Ordine S29 ✓ RA-Morph

	Subsidy		Euller	Brand or
	(Manufacturer's Price)		Fully Subsidised	Generic
	\$	Per		Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	equency			
Tab immediate-release 10 mg		10	1	Sevredol
Tab immediate-release 20 mg		10		Sevredol
Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 30 mg		10		m-Eslon
Cap long-acting 60 mg		10		m-Eslon
Cap long-acting 100 mg		10		m-Eslon
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5		Medsurge
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		5		Medsurge
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		5		Medsurge
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		5		Medsurge
	00	Ŭ		<u>ouourgo</u>
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre		00	,	0
Tab controlled-release 5 mg		20		Oxycodone Sandoz
Tab controlled-release 10 mg		20		Oxycodone Sandoz
Tab controlled-release 20 mg		20		Oxycodone Sandoz
Tab controlled-release 40 mg		20		Oxycodone Sandoz
Tab controlled-release 80 mg		20		Oxycodone Sandoz
Cap immediate-release 5 mg		20		OxyNorm OxyNorm
Cap immediate-release 10 mg		20		OxyNorm OxyNorm
Cap immediate-release 20 mg		20		OxyNorm OxyNorm
Oral liq 5 mg per 5 ml		250 m		OxyNorm Hamala
Inj 10 mg per ml, 1 ml ampoule		5		Hameln
Inj 10 mg per ml, 2 ml ampoule		5		Hameln
Inj 50 mg per ml, 1 ml ampoule		5		<u>Hameln</u>
PARACETAMOL WITH CODEINE - Safety medicine; prescriber				
* Tab paracetamol 500 mg with codeine phosphate 8 mg	27.50	1,000	/	Paracetamol +
				Codeine (Relieve)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	equency			
Tab 50 mg	4.70	10	✓	PSM
· ·	8.68		✓	Noumed Pethidine
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	SO29.88	5	✓	DBL Pethidine
				Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a P	SO30.72	5	1	DBL Pethidine
,				Hydrochloride
(PSM Tab 50 mg to be delisted 1 August 2023)				,
TRAMADOL HYDROCHLORIDE	1 50	00		Tromal CD 100
Tab sustained-release 100 mg		20		Tramal SR 100
Tab sustained release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20 100		Tramal SR 200 Arrow-Tramadol
Cap 50 mg	2.00	100	•	ALLOW-HAIHAUUI

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

Antidepressants

Cy	clic	and	Rela	ted	Ager	nts
----	------	-----	------	-----	------	-----

AMITRIPTYLINE - Safety medicine; prescriber may dete	ermine dispensing frequer	псу	
Tab 10 mg	2.49	100	✓ Arrow-Amitriptyline
Tab 25 mg	1.51	100	✓ Arrow-Amitriptyline
Tab 50 mg	2.51	100	✓ Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine	; prescriber may determin	e dispensin	g frequency
Tab 10 mg	10.17	30	✓ Clomipramine Teva
Tab 25 mg	11.99	30	✓ Clomipramine Teva
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsid	dy by endorsement		
a) Safety medicine; prescriber may determine dispe	nsing frequency		
b) Subsidy by endorsement - Subsidised for patient	s who were taking dosule	pin [dothiepi	n] hydrochloride prior to 1 June
2019 and the prescription is endorsed accordingly	/ Pharmacists may anno	tata tha nra	scription as andorsed where ther

2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride. ✓ Dosulepin Mylan

✓ Dosulepin Viatris

✓ Dosulepin

Mylan S29 ✓ Dosulepin

Viatris S29

(Dosulepin Mylan Tab 75 mg to be delisted 1 May 2023)

IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber may determine dispensing frequency

Tab 10 mg	5.48	50	Tofranil
•	10.96	100	✓ Tofranil
Tab 25 mg	8.80	50	Tofranil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; preso	criber may determin	ne dispensin	g frequency
Tab 10 mg	2.46	100	✓ Norpress
Norpress to be Principal Supply on 1 May 2023			·
Tab 25 mg	6.29	180	✓ Norpress
Norpress to be Principal Supply on 1 May 2023			-

Monoamine-Oxidase Inhibitors (MAOIs) - Non Selective

TRANYLCYPROMINE SULPHATE

Tab 10 mg	12.85	28	✓ Parnate S29 S29
·	22.94	50	✓ Parnate
	45.88	100	✓ Parnate S29 S29
	96.00		✓ Parnate S29 S29

Monoamine-Oxidase Type A Inhibitors

MOCL	.OBEMI	DΕ	
IVIOUL		ᆫ	

*	Tab 150 mg11	.80	60 •	^ <u>Aurorix</u>
*	Tab 300 mg	9.25	60 •	Aurorix

Selective Serotonin Reuptake Inhibitors

CITALOPRAM HYDROBROMIDE - Brand switch fee payable (Pharmacode 2653222) - see page 252 for details ✓ Celapram

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)	D	Fully Subsidised	Brand or Generic
	\$	Per		Manufacturer
ESCITALOPRAM				
★ Tab 10 mg	1.07	28	✓ [Escitalopram
t. =				(Ethics)
★ Tab 20 mg	1.92	28	✓ <u>F</u>	Escitalopram (Ethics)
LUOXETINE HYDROCHLORIDE				<u> </u>
Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement	2.50	28	√ <u>F</u>	<u>Fluox</u>
When prescribed for a patient who cannot swallow accordingly; or	v whole tablets or caps	ules	and the pre	scription is endorsed
2) When prescribed in a daily dose that is not a mult				
endorsed. Note: Tablets should be combined with	th capsules to facilitate	incre	emental 10	mg doses.
Cap 20 mg	2.22	30	✓ E	Brown & Burk S29
	2.91	84		luox
	3.13	90		Arrow-Fluoxetine
Arrow-Fluoxetine to be Principal Supply on 1 June 2023 Fluox Cap 20 mg to be delisted 1 June 2023) PAROXETINE	3			
★ Tab 20 mg	4.11	90	✓ <u>I</u>	_oxamine
SERTRALINE M. Tala 50 mag	0.00	00		
★ Tab 50 mg	0.99	30	_	Setrona Setrona All
لا حمل 100 mg	1 74	30		Setrona AU Setrona
★ Tab 100 mg	1./4	30	_	Setrona Setrona AU
Setrona AU Tab 50 mg to be delisted 1 October 2023)			• •	Deli Olia AO
Setrona AU Tab 100 mg to be delisted 1 October 2023)				
,				
Other Antidepressants				
//IRTAZAPINE Tab 30 mg	2.60	28		Noumed
Tab 45 mg		28	_	Noumed
•		۷۵	▼ <u>I</u>	<u>voullieu</u>
/ENLAFAXINE	6.00	0.4		Enlefey VD
k Cap 37.5 mg		84	_	Enlafax XR
k Cap 75 mg		84		Enlafax XR
≰ Cap 150 mg	11.16	84	✓ [Enlafax XR
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
DIAZEPAM – Safety medicine; prescriber may determine dispe	nsing frequency			
		5	√	Hospira
	· · · · · · · · · · · · · · · · · · ·	-		p
Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorsement				
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement a) Up to 5 inj available on a PSO				
Inj 5 mg per ml, 2 ml ampoule — Subsidy by endorsement a) Up to 5 inj available on a PSO b) Only on a PSO				
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement a) Up to 5 inj available on a PSO	ures".	5	√ 9	Stesolid

	Subsidy		Fully Brand or
	(Manufacturer's Price \$	e) Per	Subsidised Generic Manufacturer
PHENYTOIN SODIUM			
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	104.58	5	✓ Hospira
* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a PSO	154.01	5	✓ Hospira
Control of Epilepsy			
CARBAMAZEPINE			
* Tab 200 mg	14.53	100	✓ Tegretol
* Tab long-acting 200 mg	16.98	100	✓ Tegretol CR
	33.96	200	✓ Tegretol CR
* Tab 400 mg	34.58	100	✓ Tegretol
* Tab long-acting 400 mg		100	✓ Tegretol CR
* Oral liq 20 mg per ml	26.37	250 m	✓ Tegretol
CLOBAZAM - Safety medicine; prescriber may determine dispe	nsing frequency		
Tab 10 mg	9.12	50	✓ Frisium
CLONAZEPAM - Safety medicine; prescriber may determine dis	spensing frequency		
Oral drops 2.5 mg per ml		10 ml C	P ✓ Rivotril
ETHOSUXIMIDE			
Cap 250 mg	78.89	56	✓ Essential
			Ethosuximide \$29
	140.88	100	✓ Zarontin
Oral liq 250 mg per 5 ml		200 m	
GABAPENTIN			
Note: Not subsidised in combination with subsidised pregab	alin		
* Cap 100 mg		100	✓ Nupentin
* Cap 300 mg		100	✓ Nupentin
* Cap 400 mg		100	✓ Nupentin
LACOSAMIDE – Special Authority see SA1125 below – Retail p			i saponini
▲ Tab 50 mg	,	14	✓ Vimpat
▲ Tab 50 mg		14	✓ Vimpat ✓
_ 100 100 mg	200.24	56	✓ Vimpat
▲ Tab 150 mg		14	✓ Vimpat
	300.40	56	✓ Vimpat
▲ Tab 200 mg		56	✓ Vimpat
			F

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see note).

Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.



		Subsidy (Manufacturer's F	Prico\	Fully Subsidised	
		(Manulacidiei S i	Per	Jubsidised	Manufacturer
Α	MOTRIGINE				
\ \	Tab dispersible 2 mg	55.00	30	1	Lamictal
	Tab dispersible 5 mg		30		Lamictal
k	Tab dispersible 25 mg		56	_	Logem
k	Tab dispersible 50 mg		56		Logem
k	Tab dispersible 100 mg		56	_	Logem
E'	VETIRACETAM				
	Tab 250 mg	5.84	60	1	Everet
	Tab 500 mg	10.51	60	1	Everet
	Tab 750 mg	16.71	60	1	Everet
	Tab 1,000 mg	21.82	60	1	Everet
	Oral liq 100 mg per ml	44.78	300 ml (OP 🗸	Levetiracetam-AFT
Н	ENOBARBITONE				
	For phenobarbitone oral liquid refer Standard Formulae,	•	500	,	DOM
K	Tab 15 mg		500		PSM
ĸ	Tab 30 mg	40.00	500	•	PSM
Ή	ENYTOIN SODIUM				
ĸ	Tab 50 mg	75.00	200	✓	Dilantin Infatab
	Cap 30 mg	74.00	200	1	Dilantin
	Cap 100 mg	37.00	200	1	Dilantin
ĸ	Oral liq 30 mg per 5 ml	22.03	500 m	nl 🗸	Dilantin
PR	EGABALIN				
	Note: Not subsidised in combination with subsidised gab	•	56	./	Drogobalia Dfizor
	Cap 25 mg		50		Pregabalin Pfizer
	0 75	7.80		_	Milpharm S29
K	Cap 75 mg	2.65	56		Pregabalin Pfizer
		8.10			Milpharm S29
	Cap 150 mg	4.01	56		Lyrica
				•	Pregabalin Pfizer
		12.44		1	Milpharm S29
	Cap 300 mg	7.38	56	1	Pregabalin Pfizer
P	IMIDONE				,
	Tab 250 mg	37.35	100	1	Primidone Clinect
	-		100	•	aone omiett
U	DIUM VALPROATE	40.05	400	,	Follow Owner by 1.1
	Tab 100 mg		100		Epilim Crushable
	Tab 200 mg EC		100		Epilim
	Tab 500 mg EC		100		Epilim
6	Oral liq 200 mg per 5 ml	20.48	300 m		Epilim S/F Liquid
					Epilim Syrup
÷	Inj 100 mg per ml, 4 ml	41.50	1	/	Epilim IV
Τ	IRIPENTOL - Special Authority see SA1330 below - Reta	il pharmacy			
	Cap 250 mg		60	1	Diacomit \$29
	Powder for oral liq 250 mg sachet		60		Diacomit S29
	- 0 + 000 101 0101 114 200 1119 3001101		00	•	DidCollin Car

■ SA1330 Special Authority for Subsidy
Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

Subsidy Fully	Brand or
(Manufacturer's Price) Subsidised	Generic
\$ Per ✔	Manufacturer

continued...

TOPIRAMATE

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

10	I II IAWA I L		
\blacktriangle	Tab 25 mg11.07	60	Arrow-Topiramate
	•		✓ Topiramate Actavis
	26.04		✓ Topamax
\blacktriangle	Tab 50 mg	60	✓ Arrow-Topiramate
	ů		✓ Topiramate Actavis
	44.26		✓ Topamax
\blacktriangle	Tab 100 mg31.99	60	✓ Arrow-Topiramate
	ů		✓ Topiramate Actavis
	75.25		✓ Topamax
\blacktriangle	Tab 200 mg55.19	60	✓ Arrow-Topiramate
	·		✓ Topiramate Actavis
	129.85		✓ Topamax
\blacktriangle	Sprinkle cap 15 mg20.84	60	✓ Topamax
\blacktriangle	Sprinkle cap 25 mg	60	✓ Topamax

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and

VIGABATRIN − Special Authority see SA2088 below − Retail pharmacy

Tab 500 mg119.30

- 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or

100

✓ Sabril

- 1.3 Patient has tuberous sclerosis complex: and
- 2 Fither:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

	Subsidy	Fully	Brand or
(M	anufacturer's Price)	Subsidised	Generic
	ф . D.		Manufacturer

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 111

Acute Mig	raine i r	eatment
-----------	-----------	---------

RIZATRIPTAN Tab orodispersible 10 mg	3.65	30	✓ Rizamelt
SUMATRIPTAN			
Tab 50 mg	14.41	90	✓ Sumagran
Tab 100 mg		90	✓ Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per			
prescription	34.00	2 OP	Imigran

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50

PIZOTIFEN

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT − Special Authority see SA0987 below − Retail pharmacy
Cap 2 × 80 mg and 1 × 125 mg......30.00 3 OP

✓ Emend Tri-Pack

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

* Tab 16 mg4.62	100	✓ <u>Serc</u>
CYCLIZINE HYDROCHLORIDE		
Tab 50 mg0.49	10	✓ Nausicalm
CYCLIZINE LACTATE		
Inj 50 mg per ml, 1 ml ampoule - Up to 10 inj available on a		
PSO16.36	10	✓ <u>Hameln</u>
DOMPERIDONE		
* Tab 10 mg2.85	100	Pharmacy Health
4.00		Domperidone Viatris
Domperidone Viatris to be Principal Supply on 1 July 2023		
(Pharmacy Health Tab 10 mg to be delisted 1 June 2023)		
HYOSCINE HYDROBROMIDE		
* Inj 400 mcg per ml, 1 ml ampoule93.00	10	✓ Martindale \$29
Patch 1.5 mg - Special Authority see SA1998 on the next		
page – Retail pharmacy17.70	2	 Scopoderm TTS

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE			
* Tab 10 mg - Up to 30 tab available on a F	PSO1.30	100	✓ <u>Metoclopramide</u> <u>Actavis 10</u>
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 in ONDANSETRON	j available on a PSO7.00	10	✓ <u>Baxter</u>
* Tab 4 mg	2.27	50	✓ Periset
·	2.68		✓ Onrex
Tab disp 4 mg - Up to 10 tab available on	a PSO0.76	10	✓ Ondansetron
			ODT-DRLA
* Tab 8 mg	4.10	50	✓ Periset
•	4.57		✓ Onrex
Tab disp 8 mg - Up to 10 tab available on	a PSO1.13	10	✓ Ondansetron ODT-DRLA
(Onrex Tab 4 mg to be delisted 1 August 2023))		
(Onrex Tab 8 mg to be delisted 1 August 2023))		
PROCHLORPERAZINE			
* Tab 3 mg buccal	5.97	50	
	(30.00)		Buccastem
* Tab 5 mg - Up to 30 tab available on a PS	,	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml - Up to 5 inj avail		10	✓ Stemetil

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine	dispensing frequenc	:у	
Tab 100 mg	5.15	30	✓ Sulprix
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg	29.78	60	✓ Sulprix
ARIPIPRAZOLE - Safety medicine; prescriber may determine	e dispensing frequen	су	
Tab 5 mg	10.50	30	Aripiprazole Sandoz
Tab 10 mg	10.50	30	✓ Aripiprazole Sandoz
Tab 15 mg	10.50	30	✓ Aripiprazole Sandoz
Tab 20 mg	10.50	30	✓ Aripiprazole Sandoz
Tab 30 mg	10.50	30	 Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine;	prescriber may dete	rmine disper	nsing frequency
Tab 10 mg - Up to 30 tab available on a PSO	14.83	100	✓ Largactil
Tab 25 mg - Up to 30 tab available on a PSO	15.62	100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO	36.73	100	✓ Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO	30.79	10	✓ Largactil

	Subsidy	,	Fully	
	(Manufacturer's Price \$	e) Per	Subsidised	
	<u></u>	Per		Manufacturer
CLOZAPINE - Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing frequ	ency			
Tab 25 mg	6.69	50	•	Clopine
			/	Clozaril
	13.37	100	✓	Clopine
			✓	Clozaril
Tab 50 mg	8.67	50	✓	Clopine
	17.33	100	✓	Clopine
Tab 100 mg	17.33	50	1	Clopine
•			1	Clozaril
	34.65	100	1	Clopine
				Clozaril
Tab 200 mg	34.65	50	1	Clopine
	69.30	100		Clopine
Suspension 50 mg per ml		100 m		Versacloz
HALOPERIDOL – Safety medicine; prescriber may determine di			./	Caranasa
Tab 500 mcg – Up to 30 tab available on a PSO		100	_	Serenace
Tab 1.5 mg — Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50		Serenace
0 111 0 11 11 11 11 11 11 11 11 11 11 11	29.72	100	_	Serenace
Oral liq 2 mg per ml – Up to 200 ml available on a PSO		100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	SO21.55	10	•	Serenace
.EVOMEPROMAZINE - Safety medicine; prescriber may deter	mine dispensing fre	quency		
Tab 25 mg (33.8 mg as a maleate)		100		Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan `
Tab 100 mg (135 mg as a maleate)		100	1	Nozinan (Swiss)
Tab 100 mg as a maleate		100		Nozinan
.EVOMEPROMAZINE HYDROCHLORIDE - Safety medicine; p				
Inj 25 mg per ml, 1 ml ampoule	10./5	5		Neuraxpharm \$29
			_	Nozinan S29 S29
	24.48	10	/	<u>Wockhardt</u>
ITHIUM CARBONATE - Safety medicine; prescriber may deter	mine dispensing fre	equency		
Tab long-acting 400 mg		100		Priadel
Cap 250 mg		100	1	Douglas
DLANZAPINE – Safety medicine; prescriber may determine disp				•
Tab 2.5 mg		28	1	Zypine
Tab 5 mg		28		Zypine
•		28		Zypine ODT
Tab 10 mg		28	_	
Tab 10 mg			_	Zypine Zypine ODT
Tab orodispersible 10 mg		28	•	Zypine ODT
PERICYAZINE $-$ Safety medicine; prescriber may determine dis	pensing frequency			
Tab 2.5 mg	10.49	84	/	Neulactil
	12.49	100		Neulactil
Tab 10 mg	37.34	84		Neulactil
	44.45	100	1	Neulactil
QUETIAPINE - Safety medicine; prescriber may determine disp	ensina freauencv			
Tab 25 mg	. ,	90	1	Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90		Quetapel
1 ab 000 mg	12.00	50	•	audiupoi

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
RISPERIDONE – Safety medicine; prescriber may determin	ne dispensing frequency			
Tab 0.5 mg	1.86	60	✓ <u>F</u>	Risperidone (Teva)
Tab 1 mg		60	✓ F	Risperidone (Teva)
Tab 2 mg	2.29	60	✓ F	lisperidone (Teva)
Tab 3 mg		60	✓ F	lisperidone (Teva)
Tab 4 mg		60	✓ F	lisperidone (Teva)
Oral liq 1 mg per ml	8.90	30 ml	✓ F	lisperon
ZIPRASIDONE - Safety medicine; prescriber may determing	e dispensing frequency			
Cap 20 mg	17.90	60	✓ Z	usdone
Cap 40 mg		60	✓ Z	usdone
Cap 60 mg		60	✓ Z	usdone
Cap 80 mg		60	✓ Z	usdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine	; prescriber may determin	e disp	ensing freg	uency
Tab 10 mg		100	• •	Clopixol

Depot Injections

FLUPENTHIXOL DECANOATE - Safety medicine; prescriber m Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	13.14	ensing frequ 5 5 5	ency ✓ Fluanxol ✓ Fluanxol ✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber ma	av determine dispe	ensina freau	encv
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO		5 5	 ✓ Haldol ✓ Haldol Concentrate ✓ Haldol Decanoas (\$29)
OLANZAPINE – Special Authority see SA1428 below – Retail pl Safety medicine; prescriber may determine dispensing frequ	ency	4	/ Zummana Dalamann
Inj 210 mg vial		1	✓ Zyprexa Relprevv
Inj 300 mg vial		1	✓ Zyprexa Relprevv
Inj 405 mg vial	504.00	1	Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA1429 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing	frequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	Manufacturer

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

Inj 175 mg syringe	815.85	1	✓ Invega Trinza
Inj 263 mg syringe	1,072.26	1	✓ Invega Trinza
Inj 350 mg syringe		1	✓ Invega Trinza
Inj 525 mg syringe		1	✓ Invega Trinza

⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO......19.80 5 **Clopixol**

Anxiolytics

3USPIR(ONE HY	'DROCHL	.oride
---------	--------	---------	--------

000	SI INGINE ITI BITOGILEGITIBE				
*	Tab 5 mg	18.50	100	1	Buspirone Viatris
*	Tab 10 mg	12.50	100	✓	Buspirone Viatris

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CLONAZEPAM - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 500 mcg	5.64	100	✓	Paxam
Tab 2 mg	10.78	100	✓	Paxam
DIAZEPAM - Safety medicine; prescriber may determine disper	nsing frequency			
Tab 2 mg	61.07	500	✓.	Arrow-Diazepam
Tab 5 mg	73.60	500	✓.	Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency			•
Tab 1 mg	9.72	250	✓.	Ativan
Tab 2.5 mg	12.50	100	✓	Ativan

Multiple Sclerosis Treatments

⇒SA2176 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 2 Patients has an EDSS score between 0 6.0; and
- 3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 4 All of the following:
 - 4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 4.5 Either:
 - 4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 6 Any of the following:
 - 6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.



	Subsidy (Manufacturer's Price) \$	Subs	Fully sidised	Brand or Generic Manufacturer
DIMETHYL FUMARATE – Special Authority see SA2176 on the a) Wastage claimable b) Note: Treatment on two or more funded multiple scleros		•	•	mitted.
Cap 120 mg Cap 240 mg		14 56	-	ecfidera ecfidera
FINGOLIMOD – Special Authority see SA2176 on the previous a) Wastage claimable b) Note: Treatment on two or more funded multiple scleros Cap 0.5 mg	is treatments simultan		•	mitted. i ilenya
GLATIRAMER ACETATE – Special Authority see SA2176 on the Note: Treatment on two or more funded multiple sclerosis to Inj 40 mg prefilled syringe	reatments simultaneou		permitte	ed. copaxone
INTERFERON BETA-1-ALPHA – Special Authority see SA2176 Note: Treatment on two or more funded multiple sclerosis tr Inj 6 million iu prefilled syringe	reatments simultaneou 1,170.00		permitte A	
INTERFERON BETA-1-BETA – Special Authority see SA2176 of Note: Treatment on two or more funded multiple sclerosis truling 8 million iu per 1 ml	reatments simultaneou		permitte	
NATALIZUMAB – Special Authority see SA2176 on the previous Note: Treatment on two or more funded multiple sclerosis to Inj 20 mg per ml, 15 ml vial	reatments simultaneou			ed. 'ysabri
OCRELIZUMAB – Special Authority see SA2176 on the previous Note: Treatment on two or more funded multiple sclerosis to Inj 30 mg per ml, 10 ml vial	reatments simultaneou			ed. Ocrevus
TERIFLUNOMIDE – Special Authority see SA2176 on the previ a) Wastage claimable b) Note: Treatment on two or more funded multiple scleros Tab 14 mg	is treatments simultan	•	•	mitted. . ubagio
Out of the count of the country of				

Sedatives and Hypnotics

MELATONIN − Special Authority see SA1666 below − Retail pharmacy

Tab modified-release 2 mg − No more than 5 tab per day11.50 30

✓ Vigisom

Restricted to patients under 18 years of age.

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the

			IVL	RVOUS SYSTEM
(Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
continued				
following criteria:				
All of the following:				
 Patient is aged 18 years or under*; and Patient has demonstrated clinically meaningful benefit from Patient has had a trial of funded modified-release melatonir recurrence of persistent and distressing insomnia; and Funded modified-release melatonin is to be given at doses 	n discontinuation wit	hin th	ne past 12 r	
Note: Indications marked with * are unapproved indications.				
MIDAZOLAM - Safety medicine; prescriber may determine disper	nsing frequency			
Inj 1 mg per ml, 5 ml ampoule		10	✓	Midazolam Mylan
	6.10		✓	Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available				
on a PSO		10		Pfizer
On a PSO for status epilepticus use only. PSO must be e				
Inj 5 mg per ml, 3 ml ampoule		5	•	Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule – Up to 5 inj available o a PSO		5	./	Pfizer
On a PSO for status epilepticus use only. PSO must be e (Midazolam Mylan Inj 1 mg per ml, 5 ml ampoule to be delisted 1 S	endorsed for status e September 2023)	epilep	oticus use o	nly.
PHENOBARBITONE SODIUM – Special Authority see SA1386 be	•	асу		
Inj 200 mg per ml, 1 ml ampoule	103.30	10	✓	Max Health S29
 SA1386 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Both: For the treatment of terminal agitation that is unresponsive. 			nless notific	ed for applications meeting
2 The applicant is part of a multidisciplinary team working in p				
TEMAZEPAM – Safety medicine; prescriber may determine disper	•	0.5		Manusia an
Tab 10 mg		25	•	<u>Normison</u>
TRIAZOLAM – Safety medicine; prescriber may determine dispen	•	100		
Tab 125 mcg		100		Hypam
Tab 250 mcg		100		пуран
1 db 200 mog	(11.20)	100		Hypam
ZOPICLONE - Safety medicine; prescriber may determine dispen	,		•	71
Tab 7.5 mg		500	✓ ;	Zopiclone Actavis
Spinal Muscular Atrophy				

NUSINERSEN – PCT only – Special Authority see SA2174 below
Inj 12 mg per 5 ml vial120,000.00 1 ✓ Spinraza

⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

Stimulants/ADHD Treatments

TOMOXETINE		
Cap 10 mg18.41	28	 ✓ APO-Atomoxetine ✓ APO-Atomoxetine S29 S29
		 Generic Partners
107.03		✓ Strattera
Cap 18 mg27.06	28	✓ APO-Atomoxetine✓ Generic Partners
107.03		✓ Strattera
Cap 25 mg	28	✓ APO-Atomoxetine✓ Generic Partners
Cap 40 mg	28	✓ APO-Atomoxetine✓ Generic Partners
107.03		✓ Strattera
Cap 60 mg46.51	28	✓ APO-Atomoxetine ✓ APO-Atomoxetine S29 S29
		✓ Generic Partners
Cap 80 mg56.45	28	✓ APO-Atomoxetine ✓ APO-Atomoxetine S29 S29
		 Generic Partners
Cap 100 mg58.48	28	✓ APO-Atomoxetine ✓ APO-Atomoxetine S29 S29
		✓ Generic Partners

(Strattera Cap 10 mg to be delisted 1 November 2023)

(Strattera Cap 18 mg to be delisted 1 November 2023) (Strattera Cap 40 mg to be delisted 1 November 2023)

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
\$	Per	✓	Manufacturer

DEXAMFETAMINE SULFATE - Special Authority see SA1149 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

✓ PSM	100	21.00	 	 	5 mg	Tab
✓ Aspen		28.50			-	

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 on the next page - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency ✓ Rubifen 30 ✓ Ritalin Tab immediate-release 10 mg......3.00 30 ✓ Rubifen Tab extended-release 18 mg......7.75 ✓ Methylphenidate ER 30 - Teva ✓ Rubifen 30 ✓ Rubifen SR 30 ✓ Methylphenidate ER 30 - Teva ✓ Methylphenidate ER - Teva Tab extended-release 54 mg.......22.25 ✓ Methylphenidate ER 30 - Teva



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Roth:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

a) Only on a controlled drug form

 b) Safety medicine; prescriber may determine dispensin 	g trequency		
Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	Concerta
Tab extended-release 36 mg		30	Concerta
Tab extended-release 54 mg	86.24	30	Concerta
Cap modified-release 10 mg	15.60	30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price) \$	S Per	Fully subsidised	Brand or Generic Manufacturer	
---	----------	---------------------	-------------------------------------	--

continued...

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Fither:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see SA1999 below – Retail ph	narmacy		
Tab 100 mg	29.13	60	✓ Modavigil

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia		
DONEPEZIL HYDROCHLORIDE		
* Tab 5 mg4.34	90	✓ Donepezil-Rex
* Tab 10 mg	90	✓ Donepezil-Rex

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
RIVASTIGMINE – Special Authority see SA1488 below – Retail p Patch 4.6 mg per 24 hour	•	30	✓	Rivastigmine Patch BNM 5
Patch 9.5 mg per 24 hour	38.00	30	✓	Rivastigmine Patch BNM 10

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

- Buprenorphine Naloxone BNM
- Tab sublingual 8 mg with naloxone 2 mg34.00 28
- ✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone: and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the

ubsidy	Fully	Brand or
eturer's Price)	Subsidised	Generic
 \$ Per	✓	

continued...

following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHI ORIDE

Tab modified-release 150 mg	11.00	30	✓ Zyban
DISULFIRAM			
Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see SA	1408 below – Reta	il pharmacy	
Tab 50 mg	133.33	30	✓ Naltraccord


⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.



	(Manufacturer's Price)	Sul Per	bsidised	Generic Manufacturer
	Ψ	rei		Manuacturer
NICOTINE				
 a) Nicotine will not be funded in amounts less than 4 weeks 	of treatment.			
b) Note: Direct Provision by a pharmacist permitted under t	the provisions in Part	I of Secti	ion A.	
Patch 7 mg - Up to 28 patch available on a PSO	19.14	28	✓ H	labitrol
Patch 7 mg for direct distribution only - [Xpharm]	4.13	7	✓ H	labitrol
Patch 14 mg - Up to 28 patch available on a PSO	21.05	28	✓ H	labitrol
Patch 14 mg for direct distribution only - [Xpharm]	6.48	7	✓ H	labitrol
Patch 21 mg - Up to 28 patch available on a PSO	24.12	28	✓ H	labitrol
Patch 21 mg for direct distribution only - [Xpharm]	10.93	7	✓ H	labitrol
Lozenge 1 mg - Up to 216 loz available on a PSO		216	✓ H	labitrol
Lozenge 1 mg for direct distribution only - [Xpharm]		36	✓ H	labitrol
Lozenge 2 mg - Up to 216 loz available on a PSO	21.65	216	✓ H	labitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	3.40	36	✓ H	labitrol
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO	38.21	384	✓ H	labitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]	9.04	96	✓ H	labitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO	21.42	204	✓ H	labitrol
	38.21	384	✓ H	labitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]	9.04	96	✓ H	labitrol
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO	24.17	204	✓ H	labitrol
	44.17	384	✓ H	labitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]	10.47	96	✓ H	labitrol
Gum 4 mg (Mint) - Up to 384 piece available on a PSO	44.17	384	✓ H	labitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]	10.47	96	✓ H	labitrol

Subsidy

Fully

Brand or

Gum 4 mg (Mint) for direct distribution only - [Xpharm]......10.47 VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack

- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 4216.	67 53 OF	✓ <u>Varenicline Pfizer</u>
Tab 1 mg17.	62 56	✓ Varenicline Pfizer

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking;
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to guit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement
 - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	•	

continued...

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHI ORIDE - PCT only - Specialist - Special Authority see SA2153 below

Inj 25 mg vial	 	77.00	1	✓ Ribomustin
Inj 100 mg vial	 	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	 	3.23	1 mg	✓ Baxter

⇒SA2153 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6: and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

	Subsidy acturer's Price) Subs	Fully	Brand or Generic	
(wanua	\$ Per	uiseu •	Manufacturer	

continued...

- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has Hodgkin's lymphoma requiring treatment; and
 - 2 Patient has a ECOG performance status of 0-2; and
 - 3 Patient has received one prior line of chemotherapy; and
 - 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
 - 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

DUCULEAN DCT Patail pharmany Chanielist

BUSULFAN - PCT - Retail pharmacy-Specialist	100	✓ Myleran
Tab 2 mg	100	♥ Wylcian
CARBOPLATIN – PCT only – Specialist		C DDL Oanhandatin
Inj 10 mg per ml, 45 ml vial	1	✓ DBL Carboplatin
45.20		✓ Carboplatin Ebewe
48.50	4	✓ Carbaccord
Inj 1 mg for ECP0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist		
Inj 100 mg vial710.00	1	✓ <u>BiCNU</u>
Inj 100 mg for ECP710.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist		
Tab 2 mg29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist		
Inj 1 mg per ml, 50 ml vial15.00	1	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial21.00	1	✓ Cisplatin Ebewe
29.66	'	✓ DBL Cisplatin
Inj 1 mg for ECP	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE	9	
	50	√ Cycloney
Tab 50 mg - PCT - Retail pharmacy-Specialist	1	✓ <u>Cyclonex</u> ✓ Endoxan
127.80	6	✓ Cytoxan
1-11-1	1	✓ Endoxan
Inj 2 g vial – PCT only – Specialist	1 mg	✓ Baxter
, ,	ring	Daxiei
IFOSFAMIDE – PCT only – Specialist		
lnj 1 g96.00	1	✓ Holoxan
lnj 2 g	. 1	✓ Holoxan
Inj 1 mg for ECP0.10	1 mg	✓ Baxter

	Subsidy	0	Fully	
	(Manufacturer's Price) \$	Per	bsidised •	Generic Manufacturer
LOMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	1	CeeNU
Cap 40 mg		20	✓	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	✓	Alkeran
Inj 50 mg - PCT only - Specialist		1	✓	Melpha
	67.80		1	Alkeran
			1	Alkeran S29 S29
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
, 5				100
	110.00		✓	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1		Oxaliplatin Accord
Inj 1 mg for ECP	0.48	1 mg	✓	Baxter
HIOTEPA - PCT only - Specialist		•		
Inj 15 mg vial	CBS	1	1	Bedford S29
.,		•		Max Health S29
				THIO-TEPA \$29
				Tepadina \$29
Ini 100 ma vial	CDC	4		Max Health \$29
Inj 100 mg vial	OBS	1		
			•	Tepadina S29
Antimetabolites				
ZACITIDINE - PCT only - Specialist - Special Authority s	ee SA2141 below			

AZACITIDINE - PCT only - Specialist - Special Author	ority see SA2141 below		
Inj 100 mg vial	75.06	1	✓ Azacitidine Dr
			Reddy's
Inj 1 mg for ECP	0.83	1 mg	✓ Baxter

⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome: or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

				_
	Subsidy		Fully Brand or	
	(Manufacturer's Price	e) Sul	bsidised Generic	
	\$	Per	✓ Manufactu	ırer
CALCIUM FOLINATE				
Tab 15 mg - PCT - Retail pharmacy-Specialist	135.33	10	✓ DBL Leuco	vorin
Tub 10 mg 1 01 Trotail pharmacy operation	100.00	10	Calcium	VOIIII
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17 10	5	✓ Hospira	
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist		1	✓ Calcium Fo	linato
inj 10 mg per mi, 5 mi viai – POT – Hetali phamacy-Speciali	51	1	Sandoz	illiale
				linata
			✓ Calcium Fo	
			Sandoz S	
Inj 50 mg - PCT - Retail pharmacy-Specialist	72.80	10	Leucovorin	
			Pharmaci	a \$29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	Calcium Fo	linate
			Sandoz	
Inj 100 mg - PCT only - Specialist	7.33	1	✓ Calcium Fo	linate
			Ebewe	
	94.90	10	✓ Leucovorin	
			Pharmaci	a S29
Inj 300 mg - PCT only - Specialist	22.51	1	✓ Calcium Fo	
ing 500 mg - PCT only - Specialist	22.31	1	Ebewe	illiale
	05.44			DDI 000
	25.14		Leucovorin	DBL 829
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	✓ Calcium Fo	linato
ing to mg per mi, 33 mi viai – PCT only – Specialist	25.14	1	Sandoz	illiale
				linata
			✓ Calcium Fo	
			Sandoz S	
Inj 1 g - PCT only - Specialist	67.51	1	Calcium Fo	linate
			Ebewe	
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	Calcium Fo	linate
			Sandoz	
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓ Baxter	
CAPECITABINE - Retail pharmacy-Specialist		-		
Tab 150 mg	10.00	60	✓ Capercit	
Tab 500 mg		120	✓ Capecitabir	ne-
1 db 000 mg		120	DRLA S29	
			DILA	
			✓ Capercit	
CLADDIDINE DOT only Chariolist				
CLADRIBINE – PCT only – Specialist	740.00		/ 1 th-1- 000	
Inj 2 mg per ml, 5 ml		1	✓ Litak S29	
Inj 1 mg per ml, 10 ml		1	✓ Leustatin	
Inj 10 mg for ECP	749.96	10 mg OP	✓ Baxter	
CYTARABINE			_	
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Speciali	st472.00	5	✓ Pfizer	
Inj 100 mg per ml, 20 ml vial - PCT - Retail				
pharmacy-Specialist		1	✓ Pfizer	
Inj 1 mg for ECP - PCT only - Specialist		10 mg	Baxter	
Inj 100 mg intrathecal syringe for ECP - PCT only - Speciali	st94.40 1	00 mg OP	✓ Baxter	
FLUDARABINE PHOSPHATE		-		
Tab 10 mg - PCT - Retail pharmacy-Specialist	412 00	20	✓ Fludara Ora	al
Inj 50 mg vial – PCT only – Specialist		5	✓ Fludarabine	
Inj 50 mg for ECP – PCT only – Specialist		50 mg OP	✓ Baxter	LUCITO
ing so mg for Lot 1 or only opposition	120.00	oo mg Oi	- DUALGI	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price	-) (Fully Subsidised	
	(Manufacturer's Frice	Per	Subsidised •	Manufacturer
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	1	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	29.44	1	1	Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist	0.62	100 mg	/	Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	1	DBL Gemcitabine
Inj 1 g	15.89	1	1	Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg	1	Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	52.57	1	1	Accord
, ,	71.44		✓	Irinotecan Actavis 100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	1	Baxter
MERCAPTOPURINE		•		
Tab 50 mg - PCT - Retail pharmacy-Specialist	25.90	25	1	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialist				
Special Authority see SA1725 below		00 ml O)P 🗸	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist	9.98	90	✓ <u>Trexate</u>
*	Tab 10 mg - PCT - Retail pharmacy-Specialist	33.71	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist	56.05	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe	14.61	1	✓ Methotrexate
				Sandoz
*	Inj 10 mg prefilled syringe	14.66	1	✓ Methotrexate
	, 01			Sandoz
*	Inj 15 mg prefilled syringe	14.77	1	✓ Methotrexate
	, - 3, , 3-			Sandoz
*	Inj 20 mg prefilled syringe	14 88	1	✓ Methotrexate
	, = 5g p. 5 g 5 g 5		·	Sandoz
*	Inj 25 mg prefilled syringe	14 99	1	✓ Methotrexate
-,-	This 20 mg promod dyringo	14.00	•	Sandoz
*	Inj 30 mg prefilled syringe	15.00	1	✓ Methotrexate
~	iiij 50 iiig preiilied syriiige	13.03	'	Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialis	30.00	5	✓ Methotrexate DBL
*	inj 25 mg per mi, 2 mi viar – POT – Netali pharmacy-Specialis	130.00	5	Onco-Vial
	Ini OF man and OO melvial BOT Batail abanessas Crassiali	-+ 45.00		*****
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Speciali	St45.00	1	✓ DBL Methotrexate
	Lides Land DOT Division On the	05.00		Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist.	25.00	1	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial - PCT - Retail			
	pharmacy-Specialist		1	✓ <u>Methotrexate Ebewe</u>
*	Inj 1 mg for ECP - PCT only - Specialist		1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist	4.73	5 mg OP	✓ Baxter

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	•	Manufacturer
PEMETREXED - PCT only - Specialist - Special Authority see	SA1679 below			
Inj 100 mg vial	60.89	1	✓	Juno Pemetrexed
Inj 500 mg vial	217.77	1	✓	Juno Pemetrexed
Inj 1 mg for ECP	0.55	1 mg	✓	Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Roth:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

- All of the following:
 - 1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 40 mg	126.31	25	Lanvis
Other Cytotoxic Agents			
AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharm.	acy-Specialist		
Cap 0.5 mg		100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pri	ice) Su	Fully bsidised	Brand or Generic
	\$	Per	1	Manufacturer
BLEOMYCIN SULPHATE - PCT only - Specialist				
Inj 15,000 iu, vial	185.16	1	✓ [OBL Bleomycin
				Sulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	✓ E	Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see	SA1889 below			
Inj 3.5 mg vial	74.93	1	✓ [OBL Bortezomib
	105.00		✓ E	3ortezomib
				Dr-Reddy's
Inj 1 mg for ECP	22.26	1 mg	✓ E	3axter Saxter
(Bortezomib Dr-Reddy's Inj 3.5 mg vial to be delisted 1 May 2023	3)			

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	72.11	1	✓ DBL Dacarbazine
	580.60	10	✓ Dacarbazine APP \$29
Inj 200 mg for ECP	72.11	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist		· ·	
Inj 2 mg per ml, 10 ml	171.93	1	✓ Pfizer
Inj 20 mg vial1		10	✓ Daunorubicin
			Zentiva S29
Inj 20 mg for ECP	171.93	20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist			
Inj 20 mg	48.75	1	Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	✓ Docetaxel
			Accord S29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP		1 mg	✓ Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist		·	
Inj 2 mg per ml, 5 ml vial	10.00	1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Doxorubicin Ebewe
, ,	17.00		✓ Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	 Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Arrow-Doxorubicin
• •	69.99		✓ Accord \$29
	-		✓ Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	✓ Baxter

	Subsidy		Fully	y Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
EPIRUBICIN HYDROCHLORIDE – PCT only – Specialist				
Inj 2 mg per ml, 5 ml vial	25.00	1	✓	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	•	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	•	Epirubicin Ebewe
Inj 1 mg for ECP		1 mg	•	Baxter
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	1	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	1	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali		1	1	Rex Medical
Inj 1 mg for ECP - PCT only - Specialist		1 mg	•	Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist				
Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg	_	Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail phar	macy-Specialist			
Cap 500 mg	, ,	100	1	Devatis
IBRUTINIB – Special Authority see SA2168 below – Retail pharm				
Tab 140 mg		30	/	Imbruvica
Tab 420 mg		30		Imbruvica
	0,002.00	30	•	IIIDIavica

⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
 - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
 - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
 - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

Inj 5 mg vial - PCT only - Specialist	109.74	1	✓ Zavedos
Inj 10 mg vial - PCT only - Specialist	233.64	1	Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer	
LENALIDOMIDE - Retail pharmacy-Specialist - Special Authori	ty see SA2047 below			
Wastage claimable				
Cap 5 mg	5,122.76	28	✓ Revlimid	
Cap 10 mg	4,655.25	21	✓ Revlimid	
	6,207.00	28	✓ Revlimid	
Cap 15 mg	5,429.39	21	✓ Revlimid	
	7,239.18	28	✓ Revlimid	
Cap 25 mg	7,627.00	21	✓ Revlimid	

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 0.4 1 ---
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MFSNA

Tab 400 mg - PCT - Retail pharmacy-Specialist314.00	50	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist448.50	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.45	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist407.40	15	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist2.96	100 mg	✓ Baxter

	Subsidy		Fully	
	(Manufacturer's Price)		ubsidised	
	\$	Per	•	Manufacturer
MITOMYCIN C - PCT only - Specialist				
Inj 5 mg vial	641.70	1	1	Accord S29
Inj 20 mg vial	1,250.00	1	1	Teva
Inj 1 mg for ECP	269.85	1 mg	/	Baxter
MITOZANTRONE - PCT only - Specialist				
Inj 2 mg per ml, 10 ml vial	97.50	1	1	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	1	Baxter
DLAPARIB - Retail pharmacy-Specialist - Special Authority	y see SA2163 below			
Tab 100 mg	3,701.00	56	1	Lynparza
Tab 150 mg		56	1	Lynparza

⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Fither:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Por 🗸	Manufacturor

continued...

period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or

5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL - PCT only - Specialist		
Inj 30 mg47.	30 5	Paclitaxel Ebewe
Inj 100 mg24.		Paclitaxel Ebewe
91.		Paclitaxel Actavis
Inj 150 mg26.	69 1	Paclitaxel Ebewe
137.	50	✓ Anzatax
		Paclitaxel Actavis
Inj 300 mg44.	00 1	Paclitaxel Ebewe
275.	00	✓ Anzatax
		Paclitaxel Actavis
Inj 1 mg for ECP0.	20 1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see SA1979 below		
lnj 750 iu per ml, 5 ml vial	00 1	✓ Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 The patient has relapsed acute lymphoblastic leukaemia; and

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharmacy-Sp	ecialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 on the next page	– Retail pha	rmacy	
Cap 5 mg	9.13	5	✓ Temaccord
Cap 20 mg	16.38	5	✓ Temaccord
•	18.30		✓ Apo-Temozolomide
Cap 100 mg	35.98	5	✓ Temaccord
•	40.20		✓ Apo-Temozolomide
Cap 140 mg	50.12	5	✓ Temaccord
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ Temaccord

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per
Manufacturer

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Authority see SA1124 on the next page

✓ Thalomid	28	378.00	 · 	Cap 50 mg
✓ Thalomid	28	756.00	 	Cap 100 mg

	bsidy	Fully	Brand or
(Manufact	turer's Price) Sub-	sidised	Generic
·	\$ Per	✓	Manufacturer

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist	479.50	100	Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Aut	hority see SA1868 belo	w	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OP	✓ Venclexta
Tab 10 mg		14 OP	✓ Venclexta
Tab 50 mg	239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable	8,209.41	120	✓ Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

n.	Subsidy		Fully	
(IV	lanufacturer's Price) \$	Subsi Per	uiseu •	Generic Manufacturer
VINBLASTINE SULPHATE				
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist.	270.37	5	1	Hospira
Inj 1 mg for ECP - PCT only - Specialist	6.00	1 mg	1	Baxter
VINCRISTINE SULPHATE				
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist	74.52	5	•	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist	102.73	5	•	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist	12.60	1 mg	✓	Baxter
VINORELBINE - PCT only - Specialist				
Inj 10 mg per ml, 1 ml vial	12.00	1	1	Navelbine
, 01	42.00		1	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial	56.00	1	1	Navelbine
	210.00		1	Vinorelbine Ebewe
	328.65		•	Sagent S29
Inj 1 mg for ECP	1.25	1 mg	1	Baxter
Inj 50 mg for ECP	328.65 50) mg OP	1	Baxter (Sagent)
(Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 October 2024,)			
Navelhine Ini 10 mg per ml 5 ml vial to be delisted 1 October 2024)			

(Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 October 2024)

Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below Wastage claimable 224 ✓ Alecensa

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable		
Tab 20 mg3,7	74.06 60	✓ Sprycel
Tab 50 mg6,2	214.20 60	✓ Sprycel
Tab 70 mg7,6	92.58 60	✓ Sprycel

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

|--|

continued...

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
 - 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
 - 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB – Retail pharmacy-Specialist – Special Authority see SA2115 below Brand switch fee payable (Pharmacode 2651564) - see page 252 for details

Tab 100 mg	329.70	30	✓ Alchemy
Tab 150 mg	569.70	30	✓ Alchemy

⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Fither
 - 3.1 Patient is treatment naive: or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer	
GEFITINIB - Retail pharmacy-Specialist - Special Authority see	SA2116 below				
Tab 250 mg	918.00	30	✓ Ir	essa	

⇒SA2116 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	Tab 100 mg - [Xpharm] - Special Authority see SA1460		
	below2,400.00	60	✓ Glivec
*	Cap 100 mg58.23	60	✓ Imatinib-Rex
*	Cap 400 mg84.79	30	✓ Imatinib-Rex

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website <u>schedule.pharmac.govt.nz/SAForms</u>, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990
Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see \$A2035 on the next page - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage cla	aimable
-------------	---------

Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6,532.00	120	Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day: and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

1/	۸	ast	han	ω	\sim	ai	m	a	hl	Δ
v	٧	ası	ıau	10	u	aı		a	v	ı

Tab 75 mg4,000.00	21	✓ Ibrance
Tab 100 mg4,000.00	21	✓ Ibrance
Tab 125 mg4,000.00	21	✓ Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

state; and

4.2.2 Either:

- 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
- 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

	PAZOPANIB - S	Special Authority see	SA1190 below -	Retail pharmacy
--	---------------	-----------------------	----------------	-----------------

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology, and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal: or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
RUXOLITINIB — Special Authority see SA1890 below — Retai Wastage claimable	il pharmacy				
Tab 5 mg	2,500.00	56	√ Ja	akavi	
Tab 10mg	5,000.00	56	√ Ja	akavi	
Tab 15 mg	5,000.00	56	√ Ja	akavi	
Tab 20 mg	5.000.00	56	✓ Ja	akavi	

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2117 below - Retail pharmacy

Cap 12.5 mg208.38	28	Sunitinib Pfizer
Cap 25 mg416.77	28	✓ Sunitinib Pfizer
Cap 50 mg	28	✓ Sunitinib Pfizer

⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- All of the following:
 - 1 The patient has metastatic renal cell carcinoma; and
 - 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
 - 3 The patient has good performance status (WHO/ECOG grade 0-2); and
 - 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
 - 5 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
- 5.2 Haemoglobin level < lower limit of normal; or
- 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
- 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
- 5.5 Karnofsky performance score of less than or equal to 70; or
- 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Day /	Manufacturer

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 84

ABIRATERONE ACETATE – Retail pharmacy-Specialist – Special Authority see SA2118 below Wastage claimable

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Tab 50 mg	4.21	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
Ÿ	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Aut	hority see SA1895 on	the next pag	e
Ini 50 mg per ml. 5 ml prefilled syringe	,	, ,	

			_
Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Por 🗸	Manufacturer	

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

OCTREOTIDE

Inj 50 mcg per ml, 1 ml ampoule27.58	5	✓ Max Health
Inj 100 mcg per ml, 1 ml ampoule32.71	5	✓ Octreotide GH \$29 ✓ Max Health
Inj 500 mcg per ml, 1 ml ampoule	5	✓ Octreotide GH S29 ✓ Max Health
OCTREOTIDE LONG-ACTING – Special Authority see SA2119 below – Retail plnj depot 10 mg prefilled syringe	narmacy 1	✓ Octreotide Depot Teva
Inj depot 20 mg prefilled syringe	1	✓ Octreotide Depot Teva
Inj depot 30 mg prefilled syringe718.55	1	✓ Octreotide Depot Teva

⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

has failed: or

2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly: and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

TAMOXIFEN CITRATE

*	Tab 10 mg15.0	0 60	•	Tamoxifen Sandoz
*	Tab 20 mg6.6	5 60	•	Tamoxifen Sandoz

				-
	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per	✓	Manufacturer
Aromatase Inhibitors				
ANASTROZOLE * Tab 1 mg	4.55	30	/	Anatrole
EXEMESTANE		00	,	Dinay Francostona
* Tab 25 mg LETROZOLE	14.50	30	•	Pfizer Exemestane
* Tab 2.5 mg	5.84	30	✓	<u>Letrole</u>
Immunosuppressants				
Cytotoxic Immunosuppressants				
AZATHIOPRINE				
* Tab 25 mg	7.36	60	✓	<u>Azamun</u>
* Tab 50 mg	8.10	100	/	<u>Azamun</u>
MYCOPHENOLATE MOFETIL				
Tab 500 mg	35.90	50	✓	Cellcept
Cap 250 mg	35.90	100	1	Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement	187.25 165	5 ml C	OP 🗸	Cellcept
Mycophenolate powder for oral liquid is subsidised only		swall	ow tablets	and capsules, and when

Fusion Proteins

ETANERCEPT - Special Authority see SA2103 below -	- Retail pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector		4	✓ Enbrel
Inj 50 mg prefilled syringe		4	✓ Enbrel

⇒SA2103 Special Authority for Subsidy

the prescription is endorsed accordingly.

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
- 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment: and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Fither:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - POT only - 8	pecialist		
Inj 50 mg per ml, 5 ml	2,774.48	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT	Γonly – Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29

Monoclonal Antibodies

ADALIMUMAB (AMGEVITA) – Special Authority see SA2178 bek	ow – Retail pharr	nacy	
Inj 20 mg per 0.4 ml prefilled syringe	190.00	1	Amgevita
Inj 40 mg per 0.8 ml prefilled pen	375.00	2	✓ Amgevita
Inj 40 mg per 0.8 ml prefilled syringe	375.00	2	✓ Amgevita
			_

⇒SA2178 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's guality of life; and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions: and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Patient has pyoderma gangrenosum*; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — **(Crohn's disease - adults)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — **(Crohn's disease - fistulising)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss: and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose: or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions. or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or 2 Both:
- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema): or

3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	
\$	Per 🗸	Manufacturer

continued...

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects: or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Fither:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate: and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4: or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs: and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the

Subsidy	Ful	y Brand or
(Manufacturer's	Price) Subsidise	d Generic
\$	Per •	Manufacturer

continued...

treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 below - Retail pharmacy

, = = (openia riament coo or in the		
Inj 20 mg per 0.2 ml prefilled syringe	1,599.96	2	✓ Humira
Inj 40 mg per 0.4 ml prefilled syringe	•	2	✓ Humira
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	1,599.96	2	Humira
,g po. e.e p.eea eyge		_	

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved guality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsic	lised	Generic	
\$	Per	✓	Manufacturer	

continued...

treatment: or

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value: or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — **(Crohn's disease - adult)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Fither:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 12 Fither
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	idised	Generic
\$	Per	1	Manufacturer

continued...

meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

the first dose to assess response to treatment; and

- 9 Fither
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Fither:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

Inj 120 mg per ml casirivimab, 11.1 ml vial (1) and inj 120 mg

⇒SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 below

0,	openia riamoni, coo orrivor boion		
lnj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
lnj 5 mg per ml, 100 ml vial	1,820.00	1	✓ Erbitux
ni 1 mg for ECP	3.82	1 ma	✓ Baxter

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

GEMTUZUMAB OZOGAMICIN - PCT only - Specialist - Special Authority see SA2158 on the next page

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2158 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2179 below

 Inj 100 mg
 428.00
 1
 ✓ Remicade

 Inj 1 mg for ECP
 4.40
 1 mg
 ✓ Baxter

⇒SA2179 Special Authority for Subsidy

Initial application — (**Crohn's disease (adults))** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — **(fistulising Crohn's disease)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (**neurosarcoidosis**) only from a neurologist or Practitioner on the recommendation of a neurologist.

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances: or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab. etanercept or secukinumab for severe chronic plaque psoriasis: or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- - 1 Either:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:

1.1 Both:

- 1.2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
- 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis: or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults): or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis: or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis; or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis: and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)		idised	Generic
\$	Per	1	Manufacturer

continued...

- 2.1 Patients SCCAI is greater than or equal to 4; or
- 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab: or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

(Man	Subsidy ufacturer's Price)	Fu Subsidis	ılly	Brand or Generic
(wich	. ,	Per	✓	Manufacturer

continued...

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

MEPOLIZUMAB - Special Authority see SA2154 below - F	Retail pharmacy		
Inj 100 mg prefilled pen	1,638.00	1	Nucala
Inj 100 mg vial	1,638.00	1	Nucala

⇒SA2154 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Author	ority see SA2155 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive: and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL: and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
- 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Specialist - Special Authority see SA2143 below

(Synagis Inj 100 mg per ml, 1 ml vial to be delisted 1 January 2024)

⇒SA2143 Special Authority for Subsidy

Initial application — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Infant was born in the last 2 years and has severe lung, airway, neurological or neuromuscular disease that requires ongoing, life-sustaining community ventilation; or
- 2 Both:
 - 2.1 Infant was born in the last 12 months; and
 - 2.2 Any of the following:
 - 2.2.1 Patient was born at less than 28 weeks gestation; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient was born at less than 32 weeks gestation; and
 - 2222 Fither
 - 2.2.2.2.1 Patient has chronic lung disease; or
 - 2.2.2.2. Patient is Māori or any Pacific ethnicity; or
 - 2.2.3 Both:
 - 2.2.3.1 Patient has haemodynamically significant heart disease; and
 - 2.2.3.2 Any of the following:
 - 2.2.3.2.1 Patient has unoperated simple congenital heart disease with significant left to right shunt (see note a); or
 - 2.2.3.2.2 Patient has unoperated or surgically palliated complex congenital heart disease; or
 - 2.2.3.2.3 Patient has severe pulmonary hypertension (see note b); or
 - 2.2.3.2.4 Patient has moderate or severe LV failure (see note c).

Notes:

a) Patient requires/will require heart failure medication, and/or patient has significant pulmonary hypertension, and/or patient

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

will require surgical palliation/definitive repair within the next 3 months.

- b) Mean pulmonary artery pressure more than 25 mmHg.
- c) LV Ejection Fraction less than 40%.

Renewal — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months where patient still meets initial criteria.

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist -	 Special Authority see SA197 	6 below	
Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin: or

Subsidy (Manufacturer's Price)	(Fully Subsidised	Brand or Generic	
	Per	✓	Manufacturer	

continued...

- 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	✓	Manufacturer

continued...

- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2114 below

Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2114 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*: and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive: or
 - 2.2 Fither:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL: or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL;
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1.000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and

0.1.11			_
Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer
--

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia*

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	1	Manufacturer

continued...

associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment: and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1.000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Fither
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1,73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
 \$	Per	✓	

continued...

experienced intolerable side effects.

d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area: or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites: or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

continued...

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or

Subsidy	F	ılly	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 on the next page - Retail pharmacy

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Inj 100 mg per ml, 1.5 ml vial with cilgavimab 100 mg per ml,1.5 ml vial	0.00	1	✓ Evusheld
TOCILIZUMAB – PCT only – Special Authority see SA2159 below Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra ✓ Actemra S29 S29
			✓ RoActemra S29 S29
	880.00	4	✓ RoActemra S29 S29
Inj 20 mg per ml, 10 ml vial	550.00	1	✓ Actemra
			✓ Actemra S29 S29
Inj 20 mg per ml, 20 ml vial	1,100.00	1	✓ RoActemra S29 S29✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
	4,400.00	4	✓ RoActemra S29 S29
Inj 1 mg for ECP	2.85	1 mg	✓ Baxter

⇒SA2159 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsic	lised	Generic
\$	Per	1	Manufacturer

continued...

following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Fither:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated: or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Fither:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Roth:

- 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Spe	ecial Authority see SA1632 on the ne	xt page	
Inj 150 mg vial	1,350.00	1	✓ Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Ini 1 mg for FCP	9.36	1 ma	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Fither:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria: All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology):
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	 Manufac 	turer

continued...

- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib: or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 4 Fither:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
 - 5 Trastuzumab not to be given in combination with lapatinib; and
 - 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2144 below

Inj 100 mg vial	2,320.00	1	Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	24.52	1 mg	✓ Baxter

⇒SA2144 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

- 5.1 Patient does not have symptomatic brain metastases; or
- 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine;
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe......4,162.00 ✓ Stelara

⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy: or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

continued...

insufficient benefit to meet renewal criteria: or

2.2.2 Both:

2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and

2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active ulcerative colitis: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see \$A2183 below

Inj 300 mg vial3,313.00 1 ✓ Entvvio

⇒SA2183 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
- 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
- 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids: or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids: or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has active ulcerative colitis; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Programmed Cell Death-1 (PD-1) Inhibitors

TEZOLIZUMAB - PO	CT only – Specialist – Special Au	ithority see SA2195 below		
Inj 60 mg per ml, 2	20 ml vial	9,503.00	1	✓ Tecentriq
Inj 1 mg for ECP		8.08	1 mg	✓ Baxter

⇒SA2195 Special Authority for Subsidy

Initial application — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with atezolizumab and met all remaining criteria below prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic non-small cell lung cancer; and
 - 2.2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
 - 2.3 There is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
 - 2.4 Patient has an ECOG 0-2; and
 - 2.5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
 - 2.6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 12 weeks; and
 - 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
<u> </u>	\$ Pe	er 🗸	Manufacturer

continued...

- 1.2 Patient's disease has had a partial response to treatment; or
- 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

DURVALUMAB - PCT only - Specialist - Special Authority s	ee SA2164 below		
Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	Imfinzi
Inj 1 mg for ECP	9.59	1 mg	✓ Baxter

⇒SA2164 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

NIVOLUMAB - PCT only - Specialist - Special Authority see SA2120 below		
Inj 10 mg per ml, 4 ml vial	1	Opdivo
Inj 10 mg per ml, 10 ml vial2,629.96	1	Opdivo
Inj 1 mg for ECP27.62	1 mg	✓ Baxter

⇒SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes: and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

MBROLIZUMAB - PCT only - Specialist - Special Autl	hority see SA2197 on the	next page	
Inj 25 mg per ml, 4 ml vial	4,680.00	1	✓ Keytruda
Inj 1 mg for ECP	47.74	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2197 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

lesions is also considered progression).

 Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
 - 2.2 Patient has not had chemotherapy for their disease in the palliative setting; and
 - 2.3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
 - 2.4 There is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
 - 2.5 Pembrolizumab to be used as monotherapy; and
 - 2.6 Either:
 - 2.6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
 - 2.6.2 Both:
 - 2.6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
 - 2.6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment: and
 - 2.7 Patient has an ECOG 0-2; and
 - 2.8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 12 weeks; and
 - 2.9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment: or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (non-small cell lung cancer first-line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2.2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 2.3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 2.4 There is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 2.5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 2.6 Patient has an ECOG 0-2; and
- 2.7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 12 weeks: and
- 2.8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	✓ Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail pl	harmacy		
Wastage claimable			
Tab 10 mg	6,512.29	30	Afinitor
Tab 5 mg	4,555.76	30	Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

	Subsidy (Manufacturer's Prices)	e) Sub	Fully sidised	Brand or Generic Manufacturer
SIROLIMUS - Special Authority see SA2005 below - Retail phar	rmacy			
Tab 1 mg	749.99	100		Rapamune
Tab 2 mg	1,499.99	100	✓ I	Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	√	Rapamune

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- · Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy: or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note): or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009:45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and

Subs (Manufactur	. ,	Fully lised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS – Special Authority see SA1745 below – Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg		100	✓ Tacrolimus Sandoz
Cap 1 mg		100	✓ Tacrolimus Sandoz
Cap 5 mg		50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – Special Authority see SA2079 on the next page – Retail pharmacy		
Tab 15 mg1,271.00	28	✓ RINVOQ

	Subsidy		Fully	Brand or
(Mar	nufacturer's Price)	Subsic	lised	Generic
	\$	Per	•	Manutacturer

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 ini per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

Inj 0.15 mg per 0.3 ml auto-injector90	0.00	1 OP	✓ Epipen Jr
Epipen Jr to be Principal Supply on 1 July 2023			
Inj 0.3 mg per 0.3 ml auto-injector90	0.00	1 OP	Epipen

Epipen to be Principal Supply on 1 July 2023

■ SA2185 | Special Authority for Subsidy | Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
- 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

_	Subsidy		Fully	
(N	fanufacturer's Price)	Sub Per	sidised •	
SEE VENOM ALLEDOV TREATMENT. Or wish Authority on OAA	Ψ 1007 11 1-			
BEE VENOM ALLERGY TREATMENT – Special Authority see SA1				
Initiation kit - 5 vials freeze dried venom with diluent		1 OP		VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP		VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with				
diluent	285.00	1 OP	•	Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent				
9 ml, 3 diluent 1.8 ml		1 OP		Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent .	305.00	1 OP		Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see S	A1367 on the pre	vious page	– Re	etail pharmacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze				
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	/	Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze				
dried venom, with diluent	305.00	1 OP	•	Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze				
dried venom, with diluent	305.00	1 OP	/	Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze				
dried venom, with diluent	305.00	1 OP	/	Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze				
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	/	Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze				
dried venom, with diluent	305.00	1 OP	•	Venomil \$29
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
₭ Tab 10 mg	1 71	100	/	Zista
♦ Oral liq 1 mg per ml		200 ml	1	Histaclear
			•	<u>Histaclear</u>
CHLORPHENIRAMINE MALEATE	2.84	200 ml		Histaclear Histafen
CHLORPHENIRAMINE MALEATE * Oral liq 2 mg per 5 ml	2.84			
SHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE	9.37	200 ml 500 ml		
SHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE	9.37 2.02	200 ml		Histafen
SHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE	2.84 9.37 2.02 (8.40)	200 ml 500 ml 40		
CHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE	9.37 2.02 (8.40) 1.01	200 ml 500 ml		Histafen
CHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE Tab 2 mg	2.84 9.37 2.02 (8.40) 1.01 (5.99)	200 ml 500 ml 40		Histafen Polaramine
CHLORPHENIRAMINE MALEATE K Oral liq 2 mg per 5 ml DEXTROCHLORPHENIRAMINE MALEATE K Tab 2 mg	2.84 9.37 2.02 (8.40) 1.01 (5.99)	200 ml 500 ml 40 20		Histafen Polaramine
CHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE Tab 2 mg Oral liq 2 mg per 5 ml	2.84 9.37 2.02 (8.40) 1.01 (5.99) 1.77	200 ml 500 ml 40 20		Histafen Polaramine Polaramine
CHLORPHENIRAMINE MALEATE Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE Tab 2 mg Oral liq 2 mg per 5 ml EXOFENADINE HYDROCHLORIDE	2.84 9.37 2.02 (8.40) 1.01 (5.99) 1.77 (10.29)	200 ml 500 ml 40 20		Histafen Polaramine Polaramine
CHLORPHENIRAMINE MALEATE CHLORPHENIRAMINE MALEATE COUNTY OF THE COUNTY	2.84 9.37 2.02 (8.40) 1.01 (5.99) 1.77 (10.29)	200 ml 500 ml 40 20 100 ml		Histafen Polaramine Polaramine
CHLORPHENIRAMINE MALEATE K Oral liq 2 mg per 5 ml CHLORPHENIRAMINE MALEATE K Tab 2 mg CHLORPHENIRAMINE MALEATE K Oral liq 2 mg per 5 ml CHLORPHENIRAMINE MALEATE K Tab 60 mg	2.849.372.02 (8.40) 1.01 (5.99)1.77 (10.29)4.34 (8.23)	200 ml 500 ml 40 20 100 ml		Histafen Polaramine Polaramine Polaramine
CHLORPHENIRAMINE MALEATE K Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE K Tab 2 mg C Oral liq 2 mg per 5 ml EXOFENADINE HYDROCHLORIDE K Tab 60 mg	2.849.372.02 (8.40) 1.01 (5.99)1.77 (10.29)4.34 (8.23)	200 ml 500 ml 40 20 100 ml		Histafen Polaramine Polaramine Polaramine
CHLORPHENIRAMINE MALEATE * Oral liq 2 mg per 5 ml DEXTROCHLORPHENIRAMINE MALEATE * Tab 2 mg * Oral liq 2 mg per 5 ml EEXOFENADINE HYDROCHLORIDE * Tab 60 mg	2.849.372.02 (8.40) 1.01 (5.99)1.77 (10.29)4.34 (8.23)4.74	200 ml 500 ml 40 20 100 ml		Histafen Polaramine Polaramine Polaramine Telfast
CHLORPHENIRAMINE MALEATE K Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE K Tab 2 mg FOR Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE K Tab 60 mg	2.849.372.02 (8.40) 1.01 (5.99)1.77 (10.29)4.34 (8.23)4.74 (8.23)	200 ml 500 ml 40 20 100 ml 20		Histafen Polaramine Polaramine Polaramine Telfast
CHLORPHENIRAMINE MALEATE Foral liq 2 mg per 5 ml FEXTROCHLORPHENIRAMINE MALEATE Foral liq 2 mg Foral liq 2 mg per 5 ml FEXOFENADINE HYDROCHLORIDE Foral liq 2 mg Foral	2.849.372.02 (8.40) 1.01 (5.99)1.77 (10.29)4.34 (8.23)4.74 (8.23) 14.22	200 ml 500 ml 40 20 100 ml 20		Histafen Polaramine Polaramine Polaramine Telfast Telfast
CHLORPHENIRAMINE MALEATE K Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE K Tab 2 mg FOR Oral liq 2 mg per 5 ml EXTROCHLORPHENIRAMINE MALEATE K Tab 60 mg	2.849.372.02 (8.40) 1.01 (5.99)1.77 (10.29)4.34 (8.23)4.74 (8.23) 14.22 (26.44)	200 ml 500 ml 40 20 100 ml 20	•	Histafen Polaramine Polaramine Polaramine Telfast Telfast

	Subsidy		Fully	Brand or
	(Manufacturer's	Price)	Subsidised	I Generic
	` \$	Per	✓	Manufacturer
DDOMETI IA ZINIE LIVODOGI II ODIDE				
PROMETHAZINE HYDROCHLORIDE				
* Tab 10 mg	1.39	50	•	<u>Allersoothe</u>
* Tab 25 mg	1.58	50	✓	Allersoothe
* Oral lig 1 mg per 1 ml	3.39	100 m	· •	Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a P		5	•	Hospira
This 20 mg por mi, 2 mi ampould to the first available on a f	0021.00	Ü	•	Поорши
Inhalad Cartingatoraida				
Inhaled Corticosteroids				
DECLOMETIMOONE DIDDODIONATE				
BECLOMETHASONE DIPROPIONATE				_
Aerosol inhaler, 50 mcg per dose		200 dose	OP 🗸	Qvar
Aerosol inhaler, 50 mcg per dose CFC-free	8.54	200 dose	OP 🗸	Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose	OP 🗸	Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose	_	Beclazone 100
		200 dose	-	Beclazone 250
Aerosol inhaler, 250 mcg per dose CFC-free	22.07	∠uu uuse	UP V	Deciazone 200
BUDESONIDE				
Powder for inhalation, 100 mcg per dose	17 00	200 dose	OP 🗸	Pulmicort
Tomas for inflatation, 100 mag per accommission		200 0000	0.	Turbuhaler
B 1 4 11 1 11 200				
Powder for inhalation, 200 mcg per dose	19.00	200 dose	OP 🗸	Pulmicort
				Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose	OP 🗸	Pulmicort
			•	Turbuhaler
				Turbundici
FLUTICASONE				
Aerosol inhaler, 50 mcg per dose	7.19	120 dose	OP 🗸	Flixotide
Powder for inhalation, 50 mcg per dose	8.61	60 dose	OP 🗸	Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose	OP 🗸	Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose		Flixotide
. 01				
Aerosol inhaler, 250 mcg per dose		120 dose	-	<u>Flixotide</u>
Powder for inhalation, 250 mcg per dose	11.93	60 dose	OP 🗸	Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agonist	s			
milates zeng setting zeta astronocopiet rigerilet				
EFORMOTEROL FUMARATE				
Powder for inhalation, 12 mcg per dose, and monodose device	re 20.64	60 dos	Δ	
1 owder for initial ation, 12 mag per dose, and monodose device	(35.80)	00 003	C	Foradil
(F. and II Decoder to sink alating 40 areas and according	()	P. L. J. J. L. L.	0000)	Forauli
(Foradil Powder for inhalation, 12 mcg per dose, and monodose of	device to be del	listea 1 July	2023)	
EFORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated	, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	00.1	0.0	
(equivalent to eformoterol fumarate 6 mcg metered dose		60 dose	OP	
	(16.90)			Oxis Turbuhaler
INDACATEROL				
	61.00	20 doco	∩D ./	Onbrez Breezhaler
Powder for inhalation 150 mcg		30 dose		
Powder for inhalation 300 mcg	61.00	30 dose	UP 🗸	Onbrez Breezhaler
SALMETEROL				
	26.25	120 dose	∩P √	Serevent
Aerosol inhaler CFC-free, 25 mcg per dose			_	
Powder for inhalation, 50 mcg per dose, breath activated	26.25	60 dose	۷۲ 🗸	Serevent Accuhaler

	Subsidy (Manufacturer's	Price) Subsi	Fully Brand or dised Generic
	\$	Per	✓ Manufacturer
Inhaled Corticosteroids with Long-Acting Beta-	-Adrenocept	tor Agonists	
BUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide v		100 00	(D. D. O.)
6 mcg eformoterol fumarate metered dose)		120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumal per dose (equivalent to 400 mcg budesonide with 12 mc			
eformoterol fumarate metered dose) – No more than 2	⁄9		
dose per day	82.50	120 dose OP	✓ DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 r	ncg33.74	120 dose OP	✓ Symbicort
			Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 r	ncg33.74	120 dose OP	✓ Symbicort
Douglas for inholation 100 mag with aformatoral furnavata			Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg – No more than 2 dose per day	33.74	60 dose OP	✓ Symbicort
12 mg - No more than 2 dose per day		00 0036 01	Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓ Breo Ellipta
FLUTICASONE WITH SALMETEROL		00 0000 0.	2.00 Lp.u.
Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.79	120 dose OP	✓ Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - No)		
more than 2 dose per day	33.74	60 dose OP	 Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg - No			_
more than 2 dose per day	44.08	60 dose OP	✓ Seretide Accuhaler
Beta-Adrenoceptor Agonists			
SALBUTAMOL			4 14 . 11
Oral liq 400 mcg per ml		150 ml	✓ <u>Ventolin</u>
Infusion 1 mg per ml, 5 ml		10 5	✓ Ventolin✓ Ventolin
ing 500 mag per mi, 1 mi – op to 5 mg available on a 1 50		J	Ventonii
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000			
dose available on a PSO	3.80	200 dose OP	✓ Respigen
	(0.00)		✓ SalAir
Nahuliaanaala 1 maa manual 0 5 ml ammaula - Illa ta 00 mah	(6.20)		Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO		20	✓ Asthalin
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb		20	- <u>Admann</u>
available on a PSO		20	✓ Asthalin
TERBUTALINE SULPHATE			
Powder for inhalation, 200 mcg per dose (equivalent to			
250 mcg metered dose), breath activated	22.20	120 dose OP	✓ Bricanyl Turbuhaler
,			•

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	1	Manufacturer	

Anticholinergic Agents

IPRATROPIUM BROMIDE

Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose available on a PSO	16.00	200 dose OP	√ Atrovent
	10.20	200 dose OP	Alrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 neb			
available on a PSO	11.73	20	Univent
	28.20		✓ Accord S29

Inhaled Beta-Adrenoceptor Agonists with Anticholinergic Agents

SALBUTAMOL WITH IPRATROPIUM BROMIDE

Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg per			
dose CFC-free	12.19	200 dose OP	Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per			
vial 2.5 ml amnoule — Un to 20 neh available on a PSO	11 04	20	✓ Duolin

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umedictinium.
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

Powder for inhalation, 18 mcg per dose	50.37	30 dose	✓ Spiriva
Soln for inhalation 2.5 mcg per dose	50.37	60 dose OP	Spiriva Respimat

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

|--|

continued...

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

Antifibrotics

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg	2,554.00	60 OP	Ofev
Cap 150 mg	3,870.00	60 OP	Ofev

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 on the next page

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90	Esbriet
Tab 267 mg	1,215.00	90	Esbriet

Subsidy	e)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor	Antagonists
----------------------	-------------

MC	ONTELUKAST			
*	Tab 4 mg	3.10	28	✓ Montelukast Mylan
*	Tab 5 mg	3.10	28	✓ Montelukast Mylan
	•			✓ Montelukast Viatris
*	Tab 10 mg	2.90	28	✓ Montelukast Mylan
	-			✓ Montelukast Viatris

Methylxanthines

AM	IN	C	P	PHYLLINE	
			_	_	

*	Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available on a			
	PSO	180.00	5	✓ DBL Aminophylline
TH	EOPHYLLINE			
*	Tab long-acting 250 mg	23.94	100	✓ Nuelin-SR
*	Oral liq 80 mg per 15 ml	17.62	500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA - Special Authority see SA1978 below - Re	tail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see \$A2196 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg	q		
and ivacaftor 75 mg	.27,647.39	84	Trikafta
Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg			
and ivacaftor 150 mg	.27,647.39	84	Trikafta

⇒SA2196 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Either:
 - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Either:
 - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
 - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Note:

١

 a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212273s004lbl.pdf.

IVACAFTOR - PCT only - Specialist - Special Author	ority see SA2017 below		
Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

1 Patient has been diagnosed with cystic fibrosis; and

	RESPIRAT	UKT 5151E	IN AND ALLERGIES
	Subsidy (Manufacturer's P		Fully Brand or dised Generic Manufacturer
continued			
2 Either:			
Patient must have G551D mutation in the cystic fibrest 1 allele; or Patient must have other gating (class III) mutation (G1244E, G1349		
and S549R) in the CFTR gene on at least 1 allele;			- in at a ab a a a in a a baa Ma a a a ab a t
3 Patients must have a sweat chloride value of at least 60 m sweat collection system; and			
 4 Treatment with ivacaftor must be given concomitantly with 5 Patient must not have an acute upper or lower respiratory (including antibiotics) for pulmonary disease in the last 4 w 6 The dose of ivacaftor will not exceed one tablet or one sac 7 Applicant has experience and expertise in the managemer 	infection, pulmor reeks prior to cor thet twice daily; a	nary exacerbati mmencing treat and	on, or changes in therapy
SODIUM CHLORIDE			
Not funded for use as a nasal drop. Soln 7%	24.50	90 ml OP	✓ Biomed
Nasal Preparations			
Allergy Prophylactics			
BUDESONIDE			
Metered aqueous nasal spray, 50 mcg per dose		200 dose OP	✓ <u>SteroClear</u>
Metered aqueous nasal spray, 100 mcg per dose	2.84	200 dose OP	✓ <u>SteroClear</u>
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE			
Aqueous nasal spray, 0.03%	5.23	15 ml OP	✓ <u>Univent</u>
Respiratory Devices			
MASK FOR SPACER DEVICE			
a) Up to 50 dev available on a PSO			
b) Only on a PSO c) Only for children aged six years and under	0.70	j	Z a alkamban Masta
SmallPEAK FLOW METER	2.70	1	✓ e-chamber Mask
a) Up to 25 dev available on a PSO			
b) Only on a PSO			
Low range	9.54	1	Mini-Wright AFS Low Range
Normal range	9.54	1	Mini-Wright Standard
SPACER DEVICE			
a) Up to 50 dev available on a PSO			
b) Only on a PSO	0.05	4	
220 ml (single patient)		1	✓ e-chamber Turbo

✓ e-chamber La Grande✓ Volumatic

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Respiratory Stimulants

CAFFEINE CITRATE

Oral liq 20 mg per ml (10 mg base per ml)......15.10 25 ml OP **✔ Biomed**

				<u> </u>
	Subsidy		Fully	Brand or
	(Manufacturer's F	Orion) Qui	osidised	Generic
	\$	Per	Jaiuiaeu	Manufacturer
	Ψ	FEI		Manufacturei
Ear Preparations				
FLUMETASONE PIVALATE				
	4.40	7.5	,	
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	•	Locacorten-Viaform
				ED's
			1	Locorten-Vioform
			•	LOGOTICH VIOLOTHI
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI	N AND NYSTAT	ΓIN		
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate				
2.5 mg and gramicidin 250 mcg per g	E 16	7.5 ml OP	./	Kenacomb
2.5 mg and gramicium 250 mcg per g		7.5 IIII OF	•	Reliacollib
Ear/Eye Preparations				
, '				
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN				
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and				
gramicidin 50 mcg per ml	4.50	8 ml OP		
	(9.27)			Sofradex
EDAM/CETIN CHI DI IATE	` ,			
FRAMYCETIN SULPHATE				
Ear/Eye drops 0.5%	4.13	8 ml OP		
	(8.65)			Soframycin
	· ,			,
Eve Branarations				
Eye Preparations				
Eye preparations are only funded for use in the eye, unless explic	citly stated other	wise.		
Anti-Infective Preparations				
·				
ACICLOVIR				
* Eye oint 3%	14 88	4.5 g OP	1	ViruPOS
		9 0.		
CHLORAMPHENICOL				
Eye oint 1%	1.09	5 g OP	1	Devatis
Eye drops 0.5%		10 ml	1	Chlorsig
Lyo dropo 0.0 /0	7.50	10 ml OP		Chlorafast
			•	Cilioralasi
Funded for use in the ear*. Indications marked with * are	e unapproved in	dications.		
(Chlorafast Eye drops 0.5% to be delisted 1 September 2023)				
CIPROFLOXACIN				
	0.70	- 100	,	o
Eye drops 0.3% - Subsidy by endorsement	9./3	5 ml OP		Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis o	r severe bacteria	al conjunctiviti	s resista	ant to chloramphenicol; or
for the second line treatment of chronic suppurative otitis	media (CSOM)	*: and the pre	scription	n is endorsed accordingly.
Note: Indication marked with a * is an unapproved indication		,		3,
	Allori.			
GENTAMICIN SULPHATE				
Eye drops 0.3%	11.40	5 ml OP	1	Genoptic
(Genoptic Eye drops 0.3% to be delisted 1 August 2023)				
PROPAMIDINE ISETHIONATE				
* Eye drops 0.1%	2 97	10 ml OP		
=, - =, - =, - = = = = = = = = = = = = =				Brolene
	(14.55)			DIOIGIIG
SODIUM FUSIDATE [FUSIDIC ACID]				
Eye drops 1%	5.29	5 g OP	1	Fucithalmic
,		0 g 01	-	
TOBRAMYCIN				
Eye oint 0.3%	10.45	3.5 g OP	1	Tobrex
Eye drops 0.3%		5 ml OP		Tobrex
-, 0 5/0 0/0 0/0 // ······················		5 IIII OI	-	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	•	Manufacturer	
				ě

Corticosteroids and Other Anti-Inflammatory Preparations

DE	XAMETHASONE			
*	Eye oint 0.1%	5.86	3.5 g OP	✓ Maxidex
*	Eye drops 0.1%	4.50	5 ml OP	✓ Maxidex
	Ocular implant 700 mcg - Special Authority see SA1680 below			
	- Retail pharmacy	44.50	1	Ozurdex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per q5.39	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml4.50	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM Eye drops 0.1%8.80	5 ml OP	✓ Voltaren Ophtha
FLUOROMETHOLONE * Eye drops 0.1%	5 ml OP	✓ FML ✓ Flucon

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price) Subs	sidised	Generic
	\$	Per	✓	Manufacturer
EVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
, , ,	(10.34)		L	ivostin
DDOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	✓ L	omide .
REDNISOLONE ACETATE				
Eye drops 1%	6.92	10 ml OP	✓ P	rednisolone-AFT
,	7.00	5 ml OP	✓ P	red Forte
REDNISOLONE SODIUM PHOSPHATE - Special Authority	see SA1715 below	v – Retail phari	macv	
Eve drops 0.5%, single dose (preservative free)		20 dose	•-	linims
7				Prednisolone

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE

Glaucoma Preparations - Beta Blockers

ВE	IAXOLOL
----	---------

*	Eye drops 0.25%11.80	5 mi OP	Betoptic S
*	Eye drops 0.5%	5 ml OP	✓ Betoptic

TIMOLOL

 ★ Eye drops 0.25%
 1.81
 5 ml OP
 ✓ Arrow-Timolol

 ★ Eye drops 0.5%
 2.04
 5 ml OP
 ✓ Arrow-Timolol

 ★ Eye drops 0.5%, gel forming – Subsidy by endorsement
 3.78
 2.5 ml OP
 ✓ Timoptol XE

Subsidised for patients who were taking timolol eye drops 0.5%, gel forming prior to 1 April 2023 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of timolol eye drops 0.5%, gel forming.

(Timoptol XE Eye drops 0.5%, gel forming to be delisted 1 March 2024)

Glaucoma Preparations - Carbonic Anhydrase Inhibitors

BRINZOLAMIDE

DORZOLAMIDE HYDROCHLORIDE - Subsidy by endorsement

Subsidised for patients who were taking dorzolamide hydrochloride eye drops 2% prior to 1 April 2023 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of dorzolamide hydrochloride eye drops 2%.

(17.44) Trusopt

(Trusopt Eye drops 2% to be delisted 1 March 2024)

249

	Subsidy (Manufacturer's Pri \$	ce) Subs	Fully sidised	Brand or Generic Manufacturer
DORZOLAMIDE WITH TIMOLOL * Eye drops 2% with timolol 0.5%	2.73	5 ml OP	✓ Do	rtimopt
Glaucoma Preparations - Prostaglandin Analogu	ues			
BIMATOPROST * Eye drops 0.03%	5.95	3 ml OP	_	matoprost Multichem
LATANOPROST * Eye drops 0.005%TRAVOPROST	1.82	2.5 ml OP	✓ <u>Te</u>	<u>va</u>
* Eye drops 0.004%	9.75	2.5 ml OP	✓ <u>Tra</u>	<u>avatan</u>
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE * Eye drops 0.2% BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE	4.29	5 ml OP	✓ <u>Arı</u>	row-Brimonidine
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Co	mbigan
LATANOPROST WITH TIMOLOL Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	✓ <u>Arı</u>	row - Lattim
PILOCARPINE HYDROCHLORIDE # Eye drops 1% # Eye drops 2% Subsidised for oral use pursuant to the Standard Formula	5.35 7.99	15 ml OP 15 ml OP 15 ml OP	✓ Iso	pto Carpine pto Carpine pto Carpine
* Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy	31.95	20 dose	✓ Mir	nims Pilocarpine

⇒SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

ATROPINE SULPHATE		
* Eye drops 1%17.36	15 ml OP	✓ <u>Atropt</u>
CYCLOPENTOLATE HYDROCHLORIDE		
* Eye drops 1%8.76	15 ml OP	Cyclogyl
TROPICAMIDE		
* Eye drops 0.5%	15 ml OP	Mydriacyl
* Eye drops 1%	15 ml OP	Mydriacyl

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	/	Manufacturer	

Preparations for Tear Deficiency

For acetylcysteine eye drops refer Standard Formulae, page 254

HYPROMELLOSE

*	Eye drops 0.5%	19.50	15 ml OP	Methopt
---	----------------	-------	----------	---------

HYPROMELLOSE WITH DEXTRAN

Preservative Free Ocular Lubricants

⇒SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA2134 above – Retail p	oharmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL	- Special Authority se	e SA2134	above – Retail pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	Systane Unit Dose
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml	10.78	30	✓ Systane Unit Dose
(Systane Unit Dose Eye drops 0.4% and propylene glycol 0.3%	6, 0.4 ml to be delisted	1 June 2	023)
SODIUM HYALURONATE [HYALURONIC ACID] - Special A	uthority see SA2134 al	oove – Re	tail pharmacy
Eye drops 1 mg per ml	13.85	0 ml OP	✓ <u>Hylo-Fresh</u>
Hylo-Fresh has a 6 month expiry after opening. The F	Pharmacy Procedures	Manual re	striction allowing one bottle per
month is not relevant and therefore only the prescribe	d dosage to the neares	st OP may	be claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%4.15 (Naphcon Forte Eye drops 0.1% to be delisted 1 September 2023)	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eva dropp 0.19/	5 ml OP	✓ Olopatadine Teva
Eye drops 0.1%	5 IIII OF	▼ <u>Olopalaulile reva</u>
* Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE		
Eve oint 138 mcg per g	5 a OP	✓ VitA-POS



Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	/	Manufacturer	

Various

PHARMACY SERVICES

May only be claimed once per patient.

* Brand switch fee4.50

1 fee SSF Alchemy

✓ BSF Celapram

✓ BSF Ticagrelor Sandoz

✓ BSF Vebulis

- a) The Pharmacode for BSF Alchemy is 2651564 see also page 158
- b) The Pharmacode for BSF Ticagrelor Sandoz is 2653206 see also page 41
- c) The Pharmacode for BSF Vebulis is 2653214 see also page 61
- d) The Pharmacode for BSF Celapram is 2653222 see also page 123

(BSF Alchemy Brand switch fee to be delisted 1 May 2023)

(BSF Celapram Brand switch fee to be delisted 1 June 2023)

(BSF Ticagrelor Sandoz Brand switch fee to be delisted 1 June 2023)

(BSF Vebulis Brand switch fee to be delisted 1 June 2023)

Agents Used in the Treatment of Poisonings

Antidotes

AC.	FT'	VΙ	\cap	/27	ΓFΙ	M	⊏

Inj 200 mg per ml, 10 ml ampoule	52.88	10	✓ Martindale Pharma
----------------------------------	-------	----	---------------------

NALOXONE HYDROCHLORIDE

- a) Up to 10 inj available on a PSO
- b) Only on a PSO

Removal and Elimination

CHARCOAL

*	Oral lig 50 g per 250 ml	43.50	250 ml OP	•	Carbosorb

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	Exjade
Tab 500 mg dispersible	1.105.00	28	Exiade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or

Subsidy		Fully	Brand or
(Manufacturer		Subsidised	Generic
\$	Per	✓	Manufacturer

- 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
- 3.3 Treatment with deferiprone has resulted in arthritis; or
- 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below - F	Retail pharmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

# Inj 500 mg vial	151.31	10	✓ DBL Desferrioxamine Mesylate for Inj BP ✓ Deferoxamine Pfizer S29 \$29
SODIUM CALCIUM EDETATE * Inj 200 mg per ml, 5 ml	53.31	6	
, 3 , .	(156.71)	-	Calcium Disodium Versenate



Standard Formulae

Standard Formulae			
ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol Preservative Water	60 mg 40 ml qs to 100 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml) Phenobarbitone Sodium Glycerol BP Water	400 mg 4 ml to 40 ml
CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative Water	300 mg 40 ml qs to 100 ml	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative Water	qs qs to 500 ml
FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Water (Preservative should be used if quantity supplied is f than 5 days. Maximum 500 ml per prescription.)	1 tab qs to 500 ml or more	(Preservative should be used if quantity supplied is than 5 days.) SALIVA SUBSTITUTE FORMULA Methylcellulose Preservative Water	5 g qs to 500 ml
METHADONE MIXTURE Methadone powder Glycerol Water	qs qs to 100 ml	(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.) SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml Water	qs qs
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqu	10 g to 100 ml id mixture)	(Only funded if prescribed for treatment of hyponatra VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection	•
OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml	Glycerin with sucrose suspension Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	to 100 ml

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per Manufacturer

	\$	Per	1	Manufacture
Extemporaneously Compounded Preparations and	Galenica	ls		

CODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency

(90.09)Douglas Only in extemporaneously compounded codeine linctus. COLLODION FLEXIBLE Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined. ✓ PSM 100 ml COMPOUND HYDROXYBENZOATE - Only in combination Only in extemporaneously compounded oral mixtures. 100 ml ✓ Midwest GLYCERIN WITH SODIUM SACCHARIN - Only in combination Only in combination with Ora-Plus or when used in the vancomycin oral lquuid Standard Formulae. 473 ml ✓ Ora-Sweet SF GLYCERIN WITH SUCROSE - Only in combination Only in combination with Ora-Plus or when used in the vancomycin oral Iguuid Standard Formulae. 473 ml ✓ Ora-Sweet GI YCFROI 500 ml ✓ healthE Glycerol BP Only in extemporaneously compounded oral liquid preparations. METHADONE HYDROCHLORIDE a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing frequency d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets). ✓ AFT 1 a METHYL HYDROXYBENZOATE ✓ Midwest 25 q METHYLCELLULOSE 100 g ✓ MidWest ✓ Ora-Plus 473 ml METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN - Only in combination 473 ml Ora-Blend SF METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only in combination Suspension......30.95 473 ml Ora-Blend PHENOBARRITONE SODIUM

PROPYLENE GLYCOL

Only in extemporaneously compounded methyl hydroxybenzoate 10% solution.

SODIUM BICARBONATE

Only in extemporaneously compounded omeprazole and lansoprazole suspension

✓ MidWest

✓ MidWest

✓ Midwest

10 a

100 g

325.00

Only in children up to 12 years

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price)	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatiol Liq		500 ml	✓ M	idwest
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

1 cancer in children: or

Both:

- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

Subsic		Fully	Brand or
(Manufacture	r's Price) Subs	idised	Generic
\$	Per	1	Manufacturer

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)12.30		
30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)12.30	200 ml OP	✓ Calogen
Oil30.00	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml114.92	4 OP	✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	rmacy [HP3]	PROTEIN SUPPLEMENT — Special Authority see SA1524 above — Hospital pha
✓ Protifar	225 g OP	Powder7.90
✓ Resource	227 g OP	8.95
Beneprotein		

Subsidy (Manufacturer's Price) \$ Fully Subsidised Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority se	e SA1095 above -	 Hospital pharm 	nacy [HP3]
Liquid	3.75	500 ml OP	✓ Glucerna Select
·	7.50	1,000 ml OP	✓ Nutrison Advanced
			Diason
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA	1095 above – Ho	spital pharmacy	[HP3]
Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	2.10		✓ Nutren Diabetes

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED − Special Authority see SA1525 above − Hospital pharmacy [HP3]
Powder60.48 400 g OP ✓ Monogen

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Powder54.00 400 g OP ✓ Kindergen

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

SPECIAL FOODS

Subsid	y Fu	y Brand or
(Manufacturer	,	
\$	Per	Manufacturer

continued...

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

practitioner and date contacted.			
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority see S Liquid		he previous pag 500 ml OP	e – Hospital pharmacy [HP3] ✓ Nutrini Energy RTH ✓ Frebini Energy
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see SA Liquid		previous page 500 ml OP	Hospital pharmacy [HP3]✓ Nutrini RTH✓ Pediasure RTH
	6.50		✓ Frebini Original
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Ampharmacy [HP3]	uthority see	SA1379 on the	previous page – Hospital
Liquid	6.00	500 ml OP	Nutrini Energy Multi Fibre
	7.00		✓ Frebini Energy Fibre
PAEDIATRIC ENTERAL FEED WITH FIBRE 1KCAL/ML - Special Authorization pharmacy [HP3]	•	·	, ,
Liquid	7.00	500 ml OP	Frebini Original Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA13	79 on the p	revious page - I	Hospital pharmacy [HP3]
Liquid (strawberry)	1.60	200 ml OP	✓ Fortini
Liquid (vanilla)	1.60	200 ml OP	✓ Fortini
	6.99	500 ml OP	✓ Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379	on the pre	vious page – Ho	ospital pharmacy [HP3]
Liquid (chocolate)	1.07	200 ml OP	✓ Pediasure
Liquid (strawberry)	1.07	200 ml OP	✓ Pediasure
Liquid (vanilla)	1.07	200 ml OP	✓ Pediasure
	1.34	250 ml OP	✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Author pharmacy [HP3]	rity see SA	1379 on the pre	vious page – Hospital
Liquid (unflavoured)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)	1.60	200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 on the Powder		age – Hospital _I 400 g OP	pharmacy [HP3] Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML - Special Authority so Liquid	· ·		- Hospital pharmacy [HP3] ✓ Nepro HP RTH
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see S Liquid		ous page – Hos 220 ml OP	
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA	1101 on the previou	s page – Hosp	ital pharmacy [HP3]
Liquid, 200 ml bottle	11.52	4 OP	
	(13.24)		NovaSource Renal
Liquid (apricot) 125 ml	11.52	4 OP	✓ Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	✓ Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 vear for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome: or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Liquid (caramel) 125 ml.......11.52

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - Spec			
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see \$,	
Liquid (grapefruit), 250 ml carton		18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra

	Subsidy (Manufacturer's \$		Fully osidised	Brand or Generic Manufacturer
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see S Powder (unflavoured)		orevious page - 80 g OP		al pharmacy [HP3] /ivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Auth [HP3]	ority see SA13	77 on the previ	ous page	e - Hospital pharmacy
Liquid	9.60	500 ml OP	✓ S	Survimed OPD
	12.04	1,000 ml OP	-	lutrison Advanced Peptisorb Peptisorb
(Pentisorh Liquid to be delisted 1 June 2023)				

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Child aged one to eight years: and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Liquid	4.00	500 ml OP	1	Nutrini Low Energy	
				Multi Eibre	

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	✓	Manufacturer

3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g., to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (**Long-term medical condition**) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia: or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	•	Manufacturer	

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 264 - Hospital pharmacy [HP3]

- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

Liquid	250 ml OP 1,000 ml OP * Ensure Plus HN * Nutrison Energy * Fresubin HP Energy
ENTERAL FEED 1KCAL/ML – Special Authority see SA1859 on page 264 – F Liquid	Hospital pharmacy [HP3] 250 ml OP 1,000 ml OP * Isosource Standard Nutrison Standard RTH * Osmolite RTH
6.50 ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority see SA1856 Liquid	✓ Fresubin Original 9 on page 264 – Hospital pharmacy [HP3] 1,000 ml OP ✓ Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML – Special Authority see SA1859 or Liquid	n page 264 – Hospital pharmacy [HP3] 1,000 ml OP Jevity RTH Nutrison Multi Fibre Fresubin Original Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority see SA1859 c Liquid	1,000 ml OP ✓ Jevity Plus
9.80	✓ Fresubin HP Energy Fibre
ENTERAL FEED WITH PROTEIN 1.2KCAL/ML - Special Authority see SA188 Liquid	500 ml OP ✓ Fresubin Intensive
ORAL FEED (POWDER) – Special Authority see SA1859 on page 264 – Hosp Powder (chocolate)	pital pharmacy [HP3] 840 g OP ✓ Sustagen Hospital Formula
Powder (vanilla)	850 g OP
26.00	850 g OP ✓ Ensure

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 264 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
Lituoisement	(1.26) (1.26)	200 IIII OF	Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 ml			
with Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement		237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 264 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisin Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements: or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML	 Special Authority see SA1195 on the previous p 	age – Hospital p	harmacy [HP3]
Liquid	5.50	500 ml OP	✓ Nutrison
			Concentrated
	6.50		✓ Fresubin 2kcal HP
	11.00	1,000 ml OP	Ensure Two Cal HN
			RTH

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

continued...

SPECIAL FOODS

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX - Special Authority see SA17			
Powder	2.81	1,000 g OP	
	(5.15)	-	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA172	9 above – Hospital p	oharmacy [HP3]	
Powder	3.93 ·	1,000 g OP	
	(7.32)		NZB Low Gluten
	, ,		Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 ab	ove – Hospital pharr	nacy [HP3]	
Powder	5.62	2,000 g OP	
	(18.10)	-	Horlevs Flour

	Subsidy		Fully	Brand or
	(Manufacturer's Pri \$	ce) Sub Per	sidised •	Generic Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page – H	lospital pharr	nacy [HF	P3]
Buckwheat Spirals	2.00	250 g OP		
	(3.11)		0)rgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)		0)rgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)		0)rgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)		0)rgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)	_	0)rgran
Rice and Corn Penne	2.00	250 g OP		
	(2.92)	_	0)rgran
Rice and Maize Pasta Spirals	2.00	250 g OP		•
	(2.92)	•	0)rgran
Rice and Millet Spirals	2.00	250 g OP		•
	(3.11)	•	0)rgran
Rice and corn spaghetti noodles	2.00	375 g OP		•
	(2.92)	•	0)rgran
Vegetable and Rice Spirals	2.00	250 g OP		•
-	(2.92)	,	0)rgran
Italian long style spaghetti	2.00	220 g OP		-
- · · ·	(3.11)	-	0)rgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic S Per ✓ Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE – Special Authority see SA1108 on the previous page – Hospital pharmacy [HP3]

armacy [HP3]			
Tabs		75 OP	Phlexy 10
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex
			Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior
, ,			Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex
		•	Powder
Powder (neutral) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (orange) 28 g sachets		30	✓ PKU Lophlex
1 owder (orange) 20 g sacriets	950.00	30	Powder
Powder (orange) 36 g sachet	202.00	30	✓ PKU Anamix Junior
rowder (orange) so g sacher	393.00	30	
Decides (conflict 00 marches	000.00	00	Orange
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior
			Vanilla
Infant formula		400 g OP	✓ PKU Anamix Infant
Powder (orange)		500 g OP	XP Maxamum
Powder (unflavoured)	320.00	500 g OP	XP Maxamum
Liquid (berry)	13.10	125 ml OP	PKU Anamix Junior
			LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior
			LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior
_ 1 (LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex
Oral contraction (beines) roo g		00 01	Sensation 20
Liquid (juicy berries) 62.5 ml	030.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
		60 OP	✓ PKU Lophiex LQ 10
Liquid (juicy orange) 62.5 ml			•
Liquid (juicy berries) 125 ml.		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING MIX – Special Authority see SA1108 on the previous page – Hospital pharmacy [HP3]

Powder8.22 500 g OP ✓ Loprofin Mix

LOW PROTEIN PASTA - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Animal shapes11.91	500 g OP	Loprofin
Lasagne	250 g OP	✓ Loprofin
Low protein rice pasta11.91	500 g OP	✓ Loprofin
Macaroni	250 g OP	✓ Loprofin
Penne11.91	500 g OP	✓ Loprofin
Spaghetti	500 g OP	✓ Loprofin
Spirals11.91	500 g OP	✓ Loprofin

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

MINO ACID FOHMULA — Special Authority see SA2092 below — Hospital phar Powder43.60	400 g OP	✓ Alfamino
Powder (unflavoured)53.00	400 g OP	✓ Alfamino Junio ✓ Elecare ✓ Elecare LCP
		✓ Neocate Gold ✓ Neocate Junior Unflavoured
Powder (vanilla)53.00	400 g OP	 ✓ Neocate SYNEO ✓ Elecare ✓ Neocate Junior Vanilla

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis: or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency: or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA - Special Auth	nority see SA1953 below -	Hospital pharr	nacy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption: or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

SPECIAL FOODS

Su		Fully	Brand or
(Manufact		dised	Generic
	\$ Per	•	Manufacturer

continued...

2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and

- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
- 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms: or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
 - 2 Severe malabsorption: or
 - 3 Short bowel syndrome: or
 - 4 Intractable diarrhoea: or
 - 5 Biliary atresia: or
 - 6 Cholestatic liver diseases causing malsorption; or
 - 7 Cystic fibrosis: or
 - 8 Proven fat malabsorption; or
 - 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

 Powder (unflavoured)
 35.50
 300 g OP
 KetoCal 4:1

 V Ketocal 3:1
 KetoCal 4:1

 Powder (vanilla)
 35.50
 300 g OP
 KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent.................0.00 10 ✓ BCG Vaccine

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
 - 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old; or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 - 9 above.

Ini	211	J diphtheria	toyoid with	20 11 1	tatanus	tovoid	g mca

pertussis toxoid. 8 mcg pertussis filamentous

10 **Boostrix Boostrix**

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Subsid Per	ised Generic ✓ Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE -	· · · · · · · · · · · · · · · · · · ·		
Funded for any of the following:	[Apriaiii]		
A single dose for children up to the age of 7 who have c	ompleted primary imr	nunisation:	or
2) A course of four vaccines is funded for catch up program			
primary immunisation; or			
An additional four doses (as appropriate) are funded for			
pre- or post splenectomy; pre- or post solid organ transpregimens; or	nant, renai dialysis ar	ia otner sev	rerely immunosuppressive
Five doses will be funded for children requiring solid org	an transplantation.		
Note: Please refer to the Immunisation Handbook for approp		ch up progr	ammes.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin and 80 D-antigen units	0.00	40	4 1 4 1 IDV
poliomyelitis virus in 0.5ml syringe		10	✓ <u>Infanrix IPV</u>
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN	ND HAEMOPHILUS I	NFLUENZA	AE TYPE B VACCINE -
[Xpharm] Funded for patients meeting any of the following criteria:			
Up to four doses for children up to and under the age of	10 for primary immur	nisation; or	
2) An additional four doses (as appropriate) are funded for	, ,	,	to and under the age of
10 who are patients post haematopoietic stem cell trans			
post solid organ transplant, renal dialysis and other seve			
 Up to five doses for children up to and under the age of Note: A course of up-to four vaccines is funded for catch up in 			
to complete full primary immunisation. Please refer to the Imi			
programmes.			nopilate conceane to caton ap
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,	0.00	10	✓ Infanrix-hexa
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10	IIIIaiirix-nexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm] One dose for patients meeting any of the following:			
For primary vaccination in children; or			
An additional dose (as appropriate) is funded for (re-)im	munisation for patien	ts post haei	matopoietic stem cell
transplantation, or chemotherapy; functional asplenic; p			
or post cochlear implants, renal dialysis and other sever			
 For use in testing for primary immunodeficiency disease paediatrician. 	s, on the recommend	iation of an	internal medicine physician or
pasalationi.			
Haemophilus Influenzae type B polysaccharide 10 mcg			
conjugated to tetanus toxoid as carrier protein 20-40 mcg	;		
prefilled syringe plus vial 0.5 ml	0.00	1	✓ Hiberix
HEPATITIS A VACCINE - [Xpharm]			
Funded for patients meeting any of the following criteria:			
 Two vaccinations for use in transplant patients; or Two vaccinations for use in children with chronic liver di 	coaco: or		
One dose of vaccine for close contacts of known hepatil	,		
,			
Inj 1440 ELISA units in 1 ml syringe		1	✓ Havrix
Inj 720 ELISA units in 0.5 ml syringe	0.00	1	✓ <u>Havrix Junior</u>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B RECOMBINANT VACCINE – [Xpharm] Inj 10 mcg per 0.5 ml prefilled syringe Funded for patients meeting any of the following crite		1	√ E	ngerix-B
 for household or sexual contacts of known acut for children born to mothers who are hepatitis B for children up to and under the age of 18 years serology and require additional vaccination or reference for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual int for patients following immunosuppression; or for solid organ transplant patients; or for post-haematopoietic stem cell transplant (HS) following needle stick injury. 	s surface antigen (HBsAgs s inclusive who are consic equire a primary course of ercourse; or) pos derec	itive; or I not to have	
Inj 20 mcg per 1 ml prefilled syringe Funded for patients meeting any of the following crite		1	✓ <u>E</u>	ngerix-B
 for household or sexual contacts of known acut for children born to mothers who are hepatitis B for children up to and under the age of 18 years serology and require additional vaccination or reference of the household of	s surface antigen (HBsAgs s inclusive who are consic equire a primary course of tercourse; or) pos derec	itive; or I not to have	
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AN Any of the following:	,	- [Xpl	narm]	
1) Maximum of two doses for children aged 14 years a 2) Maximum of three doses for patients meeting any o 1) People aged 15 to 26 years inclusive; or 2) Either: People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or 2) Transplant (including stem cell) patients: 3) Maximum of four doses for people aged 9 to 26 years.	f the following criteria: or	nerap	у	
Inj 270 mcg in 0.5 ml syringe	0.00	10	√ <u>G</u>	ardasil 9

Fully

Brand or

		(Manufacturer's Price) \$	Per	Subsidised <	Generic Manufacturer
INFLUENZA	VACCINE				
•	g in 0.25 ml syringe (paediatric quadrivalent vaccine pharm]	,	1	•	Afluria Quad Junior (2023 formulation)
A)	INFLUENZA VACCINE – child aged 6 months to is available each year for patients aged 6 months to i) all children aged 6 months to 35 months from	35 months who mee			criteria, as set by Pharmac:
B)	Doctors are the only Contractors entitled to claim pasyringe (paediatric quadrivalent vaccine) to patients and they may only do so in respect of the influenza	eligible under the ab	ove	criteria for	subsidised immunisation
Inj 60 mo	g in 0.5 ml syringe (quadrivalent vaccine)	110.00	10	•	Afluria Quad (2023 formulation)

Subsidy

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d)

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- People 55 to 64 years of age (inclusive) and is M\u00e4ori or of any Pacific ethnicity from 1 April 2023 to 31 December 2023; or
- c) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) Down syndrome, or
 - vii) are pregnant; or
- d) children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- e) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or
- f) children 3 to 12 years of age (inclusive), from 1 April 2023 to 31 December 2023;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	1	Manufacturer

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression: or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Either:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant: or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Ini 10 mcg of each meningococcal polysaccharide conjugated

NATIONAL IMMONIOATION CONEDCE				
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xphar	m]			
Any of the following:				
 A) Three doses for children up to 12 months of age (inclus B) Up to three doses (dependent on age at first dose) for a of age (inclusive) for primary immunisation, from 1 Marc C) Both: 	catch-up programme	e for c	children from	13 months to 59 months
 Person is one year of age or over; and Any of the following: 				
 i) up to two doses and a booster every five yet functional or anatomic asplenia, HIV, completorgan transplant; or 				
 ii) up to two doses for close contacts of mening iii) up to two doses for person who has previous iv) up to two doses for bone marrow transplant v) up to two doses for person pre- and post-im 	sly had meningococca patients; or	al dise		group; or
D) Both:				
 Person is aged between 13 and 25 years (inclusiv Either: 	re); and			
 i) Two doses for individuals who are entering values boarding school hostels, tertiary education heii) ii) Two doses for individuals who are currently residence, military barracks, or prisons, from 	alls of residence, mili living in boarding sch	tary b	arracks, or postels, tertiar	orisons; or
*Immunosuppression due to corticosteroid or other immunos			•	d of greater than 28 days.
Inj 175 mcg per 0.5 ml prefilled syringe	0.00	1	✓ Bo	exsero
MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both:				
 The child is under 12 months of age; and Any of the following: 				
Up to three doses for patients pre- and post splen HIV, complement deficiency (acquired or inherited)	l), or pre or post solid			
2) Two doses for close contacts of meningococcal ca3) Two doses for child who has previously had mening4) A maximum of two doses for bone marrow transpl	ngococcal disease of	any g	roup; or	
5) A maximum of two doses for child pre- and post-in				
Note: children under 12 months of age require two dos recommended booster schedules with meningococcal A	es 8 weeks apart. Re	efer to	the Immuni	sation Handbook for
*Immunosuppression due to steroid or other immunosu		ıst be	for a period	of greater than 28 days.
Inj 10 mcg in 0.5 ml syringe		1	✓ No	eisvac-C
PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm	•			
A primary course of three doses for previously unvaccir			ū	
Note: please refer to the Immunisation Handbook for the app	•	catch	n up program	nmes
Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6E 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml	ο,			
prefilled syringe	0.00	10	✓ Sy	<u>/nflorix</u>

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
- 2) Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection; or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) cochlear implants or intracranial shunts: or
 - g) cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - I) diabetes; or
 - m) Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note:	please refer to the	Immunisation	Handbook f	or the appropriate	schedule for	catch up	programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

0, 0, 1, 02, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,		
syringe	10	Prevenar 13
	1	✓ Prevenar 13

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE — [Either: 1) Up to three doses (as appropriate) for patients with HIN chemotherapy; pre- or post-splenectomy or with function complement deficiency (acquired or inherited), cochleated 2) All of the following: a) Patient is a child under 18 years for (re-)immunistic b) Treatment is for a maximum of two doses; and c) Any of the following: i) on immunosuppressive therapy or radiation immune response; or iii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; or v) who are immune-suppressed following orgator vi) with cochlear implants or intracranial shuntsic vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more that prednisone of 2 mg/kg per day or greater, of 20 mg or greater; or ix) with chronic pulmonary disease (including a x) pre term infants, born before 28 weeks gest xi) with cardiac disease, with cyanosis or failure xii) with diabetes; or	Xpharm] 7, for patients post hae anal asplenia, pre- or primary ation; and therapy, vaccinate when transplantation (inclust; or an two weeks, and where children who weight restricted with high ation; or	ematopoietic stepost-solid organ immunodeficier seen there is expuding haematop or are on an equence than 10 kg	em cell transplant, or transplant, renal dialysis, ncy; or ected to be a sufficient poietic stem cell transplant); ivalent daily dosage of y on a total daily dosage of
xiv) who are pre-or post-splenectomy, or with fu	nctional asplenia.		
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	: viduals; or oriate schedule for cat	ch-up programr	Pneumovax 23 nes. IPOL
ROTAVIRUS ORAL VACCINE – [Xpharm] Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 14 v 2) no vaccination being administered to children aged 24	veeks of age; and		<u></u>
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator			Rotarix Rotarix

	IIIIIIIIII		7111011 001120022
	Subsidy (Manufacturer's Price) \$	Fi Subsidis Per	ully Brand or sed Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm] Either:			
1) Maximum of one dose for primary vaccination for either	:		
a) Any infant born on or after 1 April 2016; orb) For previously unvaccinated children turning 11 y	ears old on or after 1	July 2017, w	ho have not previously had a
varicella infection (chickenpox), or			
2) Maximum of two doses for any of the following:			
a) Any of the following for non-immune patients:	ha candidataa far tra	nanlantation	
 i) with chronic liver disease who may in future ii) with deteriorating renal function before trans iii) prior to solid organ transplant; or 		nspiantation;	, or
iv) prior to any elective immunosuppression*, o	r		
 v) for post exposure prophylaxis who are immu 			
b) For patients at least 2 years after bone marrow tr			
c) For patients at least 6 months after completion of			
 d) For HIV positive non immune to varicella with mile e) For patients with inborn errors of metabolism at rivaricella, or 			
f) For household contacts of paediatric patients who immune compromise where the household contacts.			
g) For household contacts of adult patients who hav immunocompromised, or undergoing a procedure has no clinical history of varicella.	e no clinical history of	varicella and	d who are severely
* immunosuppression due to steroid or other immunosuppres	ssive therany must be	for a treatm	ent period of greater than
28 days	50.10 morapyao. 50		ioni ponou oi gioutoi utun
Inj 1350 PFU prefilled syringe	0.00	1	✓ Varivax
		10	✓ <u>Varivax</u>
VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] - [Xph	arm]		
Funded for patients meeting the following criteria:			
 Two doses for all people aged 65 years 			
Inj 50 mcg per 0.5 ml vial plus vial	0.00	1	✓ Shingrix
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATE	D VACCINE [SHING	LES VACCIN	NE] - [Xpharm]
Funded for patients meeting the following criteria:			
 One dose for all people aged 65 years 			
Inj 19,400 PFU prefilled syringe plus vial	0.00	1	✓ Zostavax
		10	✓ Zostavax
Diagnostic Agents			
TUBERCULIN PPD [MANTOUX] TEST - [Xpharm]			
Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	✓ <u>Tubersol</u>

- Symbols -	Agents for Parkinsonism and Related	Amoxicillin93
3TC107	Disorders 117	Amoxicillin with clavulanic acid93
7 MED NSHA Silver/Copper	Agents Used in the Treatment of	Amphotericin B33
Short74	Poisonings252	Amsacrine149
- A -	Agrylin149	AmsaLyo149
A-Scabies68	Albendazole90	Amsidine149
Abacavir sulphate106	Albey238	Amzoate31
Abacavir sulphate with	Albustix78	Anaesthetics118
lamivudine 106	Alchemy158	Anagrelide hydrochloride149
Abacavir/Lamivudine Viatris106	Alchemy Oxybutynin78	Analgesics 119
Abiraterone acetate164	Aldurazyme29	Anastrozole167
Acarbose11	Alecensa157	Anatrole167
Accarb11	Alectinib157	Andriol Testocaps81
Accuretic 1048	Alendronate sodium112	Androderm81
Accuretic 2048	Alendronate sodium with	Anoro Ellipta242
Acetazolamide249	colecalciferol112	Antabuse141
Acetec47	Alfacalcidol34	Antacids and Antiflatulents6
Acetic acid with hydroxyquinoline and	Alfamino273	Anthelmintics90
ricinoleic acid76		Antiacne Preparations62
Acetylcysteine252		Antiallergy Preparations237
Aci-Jel76	· ·	Antianaemics37
Aciclovir	Alkeran146	Antiandrogen Oral
Infection102		Contraceptives
Sensory247		Antiarrhythmics49
Acidex6	Allerpro Syneo 1276	Antibacterials90
Acipimox55	Allerpro Syneo 2276	Antibacterials Topical62
Acitretin68		Anticholinergic Agents241
Aclasta115	Allmercap148	Anticholinesterases111
Actemra219	•	Antidepressants
Actemra S29219	•	Antidiarrhoeals6
Actinomycin D150	·	Antiepilepsy Drugs
Actrapid10	•	Antifibrinolytics, Haemostatics and
Actrapid Penfill10		Local Sclerosants38
Acupan119		Antifibrotics242
Adalimumab (Amgevita)174		Antifungals97
Adalimumab (Humira - Alternative	Alu-Tab6	Antifungals Topical63
brand)		Antihistamines238
Adapalene	•	Antihypotensives50
ADR Cartridge 1.824	· ·	Antimalarials 99
Adrenaline	Ambrisentan58	Antimigraine Preparations128
Cardiovascular57	Ambrisentan Mylan58	Antinausea and Vertigo Agents 128
Respiratory237		Antipruritic Preparations64
Advantan65		Antipsychotics129
Advate40	Amiloride hydrochloride53	Antiretrovirals105
Adynovate41	Amiloride hydrochloride with	Antirheumatoid Agents
Afinitor233	•	Antispasmodics and Other Agents
Aflibercept	Amiloride hydrochloride with	Altering Gut Motility8
Afluria Quad	hydrochlorothiazide 54	Antithrombotic Agents41
(2023 formulation)	Aminophylline243	Antithymocyte globulin
Afluria Quad Junior	Amiodarone hydrochloride49	(equine) 174
(2023 formulation)	Amisulpride129	Antitrichomonal Agents
AFT-Pyrazinamide101	Amitriptyline123	Antituberculotics and
Agents Affecting the	Amlodipine	Antileprotics
Renin-Angiotensin System 47		Antiulcerants9

Antivirals	101	Atnahs Olsalazine	8	Beta Cream	6
Anxiolytics		Atomoxetine		Beta Ointment	
Anzatax		Atorvastatin		Beta Scalp	
Apidra		Atropine sulphate		Beta-Adrenoceptor Agonists	
Apidra SoloStar		Cardiovascular	49	Beta-Adrenoceptor Blockers	
APO-Atomoxetine		Sensory		Betadine	
APO-Atomoxetine S29		Atropt		Betadine Skin Prep	
Apo-Azithromycin		Atrovent		Betaferon	
Apo-Diltiazem CD		Aubagio		Betahistine dihydrochloride	
Apo-Temozolomide		Augmentin		Betaine	
Apomorphine hydrochloride		Aurorix		Betaloc CR	
Aprepitant		AutoSoft 30		Betamethasone dipropionate	
Apresoline		AutoSoft 90		Betamethasone dipropionate with	
Aqueous cream		Avallon		calcipotriol	69
Aratac		Avelox		Betamethasone sodium phosphate	
Arava		Avonex		with betamethasone acetate	80
Arginine		Avonex Pen		Betamethasone valerate6	
Aripiprazole		Azacitidine		Betamethasone valerate with sodiur	,
Aripiprazole Sandoz		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Aristocort		Azamun		Betaxolol	
Arrotex-Prazosin S29		Azathioprine		Betnovate	
Arrow - Clopid		Azilect		Betoptic	
Arrow - Lattim		Azithromycin		Betoptic S	
Arrow-Amitriptyline		Azopt		Bexsero	
Arrow-Bendrofluazide		AZT		Bezafibrate	
Arrow-Brimonidine		-B-	107	Bezalip	
Arrow-Diazepam		B-D Micro-Fine	16	Bezalip Retard	5i
Arrow-Doxorubicin		B-D Ultra Fine		Bicalutamide	
Arrow-Fluoxetine		B-D Ultra Fine II		Bicillin LA	
Arrow-Losartan &	124	Bacillus Calmette-Guerin (BCG)	10	BiCNU	
	10	vaccine	174		
Hydrochlorothiazide			174	Bile and Liver TherapyBiltricide	
Arrow-Norfloxacin		Bacillus Calmette-Guerin vaccine	070		
Arrow-Ornidazole				Bimatoprost	
Arrow-Quinapril 10		Baclofen		Bimatoprost Multichem	
Arrow-Quinapril 20		Bactroban		Binarex	
Arrow-Quinapril 5		Balance		Binocrit	
Arrow-Roxithromycin		Barrier Creams and Emollients		Biodone	
Arrow-Timolol		BCG Vaccine		Biodone Extra Forte	
Arrow-Topiramate		Beclazone 100		Biodone Forte	
Arrow-Tramadol		Beclazone 250		Bisacodyl	2
Arsenic trioxide		Beclazone 50		Bisacodyl Viatris	2
Asacol		Beclomethasone dipropionate		Bisoprolol fumarate	
Ascorbic acid		Bee venom allergy treatment		Bisoprolol Mylan	
Aspen Adrenaline	57	Bendamustine hydrochloride		Bisoprolol Viatris	
Aspirin	44	Bendrofluazide	54	BK Lotion	
Blood		Bendroflumethiazide		Bleomycin sulphate	. 150
Nervous		[Bendrofluazide]		Blood Colony-stimulating	
Asthalin		Benralizumab		Factors	4
Atazanavir Mylan		Benzathine benzylpenicillin		Blood glucose diagnostic test	
Atazanavir sulphate		Benzatropine mesylate		meter	1
Atenolol		Benzbromaron AL 100		Blood glucose diagnostic test	
Atenolol AFT		Benzbromarone		strip	1
Atenolol AFT S29		Benztrop		Blood glucose test strips (visually	
Atezolizumab		Benzydamine hydrochloride	32	impaired)	1
ATGAM		Benzylpenicillin sodium [Penicillin		Blood Ketone Diagnostic Test	
Ativan	133	G]	93	Strip	14

Bonjela33	Calcium Folinate Sandoz S29 147	7 Cetomacrogol6
Boostrix278	Calcium gluconate35	Cetomacrogol with glycerol6
Bortezomib150	Calcium Homeostasis79	Cetomacrogol-AFT6
Bortezomib Dr-Reddy's150	Calcium polystyrene sulphonate46	6 Cetuximab19
Bosentan58		
Bosentan Dr Reddy's58	Calogen259	Chemotherapeutic Agents14
Bplex33	Candesartan cilexetil48	B Chickenpox vaccine28
Breo Ellipta240	Candestar48	3 Chlorafast24
Brevinor 1/2875	Canesten63	
Bricanyl Turbuhaler240	Capecitabine147	7 Chloramphenicol24
Brimonidine tartrate250	Capecitabine-DRLA147	7 Chlorothiazide5
Brimonidine tartrate with timolol	Capercit147	
maleate250	Capoten47	7 Chlorpromazine hydrochloride12
Brinzolamide249	Capsaicin	Chlorsig24
Brolene247	Musculoskeletal112	Chlortalidone [Chlorthalidone]5
Brown & Burk124		
Brufen SR111	Captopril47	7 Chlorvescent4
BSF Alchemy252	Carafate10	Choice 380 7med Nsha Silver/copper
BSF Celapram252	Carbaccord145	5 Short 7
BSF Ticagrelor Sandoz252	Carbamazepine125	
BSF Vebulis252		Choice TT380 Short7
Buccastem129	Carbomer25	Choice TT380 Standard7
Budesonide	Carboplatin145	Choline salicylate with cetalkonium
Alimentary6		
Respiratory239, 245		2 Ciclosporin23
Budesonide with eformoterol240		Cilazapril4
Bumetanide53	Cardizem CD52	Cilicaine VK9
Buprenorphine Naloxone BNM 140	CareSens Dual14	Cinacalcet7
Buprenorphine with naloxone140	CareSens N15	5 Cipflox9
Bupropion hydrochloride141	CareSens N POP15	
Burel98	CareSens N Premier15	
Burinex53	CareSens PRO15	5 Sensory24
Burinex S2953	Carmellose sodium with gelatin and	Ciprofloxacin Teva24
Buscopan	pectin33	3 Cisplatin14
Buscopan S29	Carmustine145	
Buspirone hydrochloride132	Carnitor30	Citalopram hydrobromide12
Buspirone Viatris132		Cladribine14
Busulfan145		Clarithromycin
- C -	Casirivimab and imdevimab19	Alimentary
Cabergoline88	Catapres5	Infection9
Caffeine citrate246		
Calamine64	Cefaclor monohydrate90	Clexane Forte4
Calamine-AFT64		Climara8
Calci-Tab 50035		Clindamycin9
Calcipotriol69	Ceftriaxone90	
Calcitonin79	Ceftriaxone-AFT90	
Calcitriol34	Cefuroxime axetil90	
Calcitriol-AFT34	Celapram123	Clobetasol propionate64, 7
Calcium 500 mg Hexal35	Celebrex11	
Calcium carbonate6, 35	Celecoxib11	
Calcium carbonate PAI6		
Calcium Channel Blockers52	Celestone Chronodose80	Dermatological 6
Calcium Disodium Versenate253		Genito-Urinary7
Calcium folinate147	Centrally-Acting Agents53	B Clomifene citrate8
Calcium Folinate Ebewe147	Cephalexin ABM90	Clomipramine hydrochloride12
Calcium Folinate Sandoz147		

Clonazepam12	5, 133	Coumadin	44	DBL Cisplatin	145
Clonidine	53	Country Life	30	DBL Dacarbazine	150
Clonidine hydrochloride	53	Coversyl	47	DBL Desferrioxamine Mesylate for	or Inj
Clonidine Teva	53	Creon 10000	25	BP	253
Clopidogrel	41	Creon 25000	25	DBL Docetaxel	150
Clopidogrel Multichem	41	Creon Micro	25	DBL Ergometrine	7
Clopine		Crotamiton	64	DBL Gemcitabine	148
Clopixol13	1-132	Crystaderm		DBL Gentamicin	95
Clotrimazole		Curam		DBL Heparin Sodium	44
Dermatological	63	Curam Duo 500/125	93	DBL Leucovorin Calcium	
Genito-Urinary		Cvite	33	DBL Methotrexate Onco-Vial	148
Clozapine	130	Cyclizine hydrochloride	128	DBL Pethidine Hydrochloride	122
Clozaril		Cyclizine lactate		DBL Vincristine Sulfate	
Co-trimoxazole	97	Cyclogyl		Decozol	
Coal tar	69	Cyclonex		Deferasirox	252
Coal tar with allantoin, menthol,		Cyclopentolate hydrochloride		Deferiprone	253
phenol and sulphur	69	Cyclophosphamide		Deferoxamine Pfizer S29	253
Coal tar with salicylic acid and		Cyclorin		Denosumab	112
sulphur	69	Cycloserine		Deolate	
Cobal-B12		Cyproterone acetate		Deoxycoformycin	154
Coco-Scalp		Cyproterone acetate with		Depo-Medrol	
Codeine phosphate		ethinyloestradiol	76	Depo-Provera	76
Extemporaneous	255	Cystadane		Depo-Testosterone	81
Nervous		Cytarabine		Deprim	97
Coenzyme Q10	29	Cytotec		Dermol	
Colchicine	115	Cytoxan		Desferrioxamine mesilate	253
Colecalciferol		. D -		Desmopressin	8
Colestid		D-Penamine	112	Desmopressin acetate	
Colestipol hydrochloride	55	Dabigatran	44	Desmopressin-PH&T	
Colgout	115	Dacarbazine		Desuric	
Colifoam		Dacarbazine APP	150	Detection of Substances in	
Colistin sulphomethate		Dactinomycin [Actinomycin D]	150	Urine	<mark>7</mark> 8
Colistin-Link		Daivobet		Dexamethasone	
Collodion flexible		Daivonex		Hormone	80
Colloidal bismuth subcitrate	10	Daktarin	63	Sensory	
Colofac	8	Dalacin C	95	Dexamethasone phosphate	
Coloxyl	26	Dantrium	116	Dexamethasone with framycetin	
Combigan	250	Dantrium S29		gramicidin	
Compound electrolytes		Dantrolene	116	Dexamethasone with neomycin	
Compound electrolytes with glucos	e	Daonil	11	sulphate and polymyxin B	
[Dextrose]	46	Dapa-Tabs	54	sulphate	248
Compound hydroxybenzoate		Dapsone	100	Dexamfetamine sulfate	137
Comtan	117	Daraprim	96	Dexmethsone	80
Concerta	138	Darunavir	107	Dextrochlorpheniramine	
Condoms	73	Darunavir Mylan	107	maleate	238
Condyline	71	Darunavir Viatris	107	Dextrose	45-46
Contraceptives - Hormonal	74	Dasatinib	157	DHC Continus	121
Contraceptives - Non-hormonal	72	Daunorubicin	150	Diabetes	10
Copaxone		Daunorubicin Zentiva	150	Diabetes Management	14
Cordarone-X		David One Step Cassette Pregn		Diacomit	
Corticosteroids and Related Agent		Test	•	Diagnostic Agents	
for Systemic Use		DBL Adrenaline	57	Diamide Relief	
Corticosteroids Topical		DBL Aminophylline	243	Diamox	
Cortifoam		DBL Bleomycin Sulfate		Diasip	
Cosentyx		DBL Bortezomib	150	Diazepam1	24, 133
Cosmegen	150	DBL Carboplatin		Diazoxide	10

Dibenzyline47	Doxorubicin Ebewe150	EMLA11
Diclofenac Sandoz111	Doxorubicin hydrochloride150	Empagliflozin1
Diclofenac sodium	Doxycycline94	Empagliflozin with metformin
Musculoskeletal111	DP Lotion66	hydrochloride1
Sensory248	DP Lotn HC65	Emtricitabine10
Differin	DP-Allopurinol115	Emtricitabine with tenofovir
Difflam32	Dr Reddy's Omeprazole9	disoproxil10
Diflucan97	Drofate51	Emtriva10
Digestives Including Enzymes25	Drugs Affecting Bone	Emulsifying ointment
Digoxin	Metabolism 112	Emulsifying Ointment ADE
Dihydrocodeine tartrate121	Dual blood glucose and blood ketone	Enalapril maleate
Dilantin	diagnostic test meter 14	Enbrel
Dilantin Infatab	Dulaglutide	Endocrine Therapy
Diltiazem CD Clinect	Dulcolax SP Drop27	Endoxan14
Diltiazem hydrochloride	Duocal Super Soluble Powder258	Engerix-B28
		Enlafax XR
Dimethicone	Duolin241 Duolin HFA241	Enoxaparin sodium
Dimethyl fumarate		
Dipentum	DuoResp Spiromax240	Enstilar
Diphtheria, tetanus and pertussis	Duride	Ensure
vaccine	Durvalumab229	Ensure Plus
Diphtheria, tetanus, pertussis and	-E-	Ensure Plus HN
polio vaccine	e-chamber La Grande245	Ensure Plus RTH
Diphtheria, tetanus, pertussis, polio,	e-chamber Mask245	Ensure Two Cal HN RTH26
hepatitis B and haemophilus	e-chamber Turbo245	Entacapone11
influenzae type B vaccine 279	E-Mycin92	Entecavir10
Diprosone64	e5 Pharma10	Entecavir Mylan10
Diprosone OV64	Ear Preparations247	Entecavir Sandoz10
Dipyridamole41	Ear/Eye Preparations247	Entocort CIR
Disopyramide phosphate49	Easiphen Liquid272	Entresto 24/26
Disulfiram141	Econazole nitrate63	Entresto 49/51
Diuretics53	Efavirenz106	Entresto 97/103
Dizole97	Efavirenz with emtricitabine and	Entyvio22
Docetaxel150	tenofovir disoproxil107	Epilim12
Docetaxel Accord150	Eformoterol fumarate239	Epilim Crushable12
Docetaxel Sandoz150	Eformoterol fumarate dihydrate239	Epilim IV12
Docusate sodium26	Eftrenonacog alfa [Recombinant	Epilim S/F Liquid12
Docusate sodium with	factor IX]38	Epilim Syrup12
sennosides26	Efudix71	Epipen23
Dolutegravir108	Egopsoryl TA69	Epipen Jr23
Domperidone 128	Elaprase29	Epirubicin Ebewe15
Domperidone Viatris128	Eldepryl117	Epirubicin hydrochloride 15
Donepezil hydrochloride139	Elecare273	Eplerenone
Donepezil-Rex139	Elecare LCP273	Epoetin alfa
Dornase alfa243	Electral46	Epoprostenol
Dortimopt250	Elelyso32	Eptacog alfa [Recombinant factor
Dorzolamide hydrochloride249	Elemental 028 Extra263	VIIa]
Dorzolamide with timolol250	Elexacaftor with tezacaftor, ivacaftor	Erbitux19
Dostinex	and ivacaftor244	Ergometrine maleate
Dosulepin [Dothiepin]	Elidel 69	Erlotinib15
hydrochloride123	Elocon	Erythrocin IV
Dosulepin Mylan123	Elocon Alcohol Free	Erythromycin (as lactobionate)
Dosulepin Viatris123	Eltrombopag38	Erythromycin ethyl succinate
Dothiepin	Eltroxin	Esbriet24
Doxazosin	EMB Fatol	Escitalopram
Doxazosin Clinect	Emend Tri-Pack	Escitalopram (Ethics)
Doxine94	Emicizumab	Eskazole
DUXING94	EIIIICIZUIIIau	∟oraZUIせ

Essential Ethosuximide	125	Femme-Tab ED	75	Flutamin	164
Essential Generics	50	Fenpaed 100 mg per 5 ml	111	Fluticasone	239
Essential Prednisolone	8	Fentanyl	121	Fluticasone furoate with	
Estraderm MX		Fentanyl Sandoz	121	vilanterol	240
Estradiol TDP Mylan	82	Ferinject	35	Fluticasone propionate	245
Estradot		Ferodan	35	Fluticasone with salmeterol	240
Estradot 50 mcg	82	Ferriprox	253	Flynn	90
Estrofem	82	Ferro-F-Tabs	35	FML	248
Etanercept	167	Ferro-tab	35	Foban	63
Ethambutol hydrochloride	100	Ferrograd	35	Folic acid	38
Ethics Aspirin	119	Ferrosig		Folic Acid multichem	38
Ethics Aspirin EC		Ferrous fumarate		Folic Acid Mylan	38
Ethics Lisinopril	47	Ferrous fumarate with folic acid.	35	Folic Acid Viatris	
Ethinyloestradiol with		Ferrous sulfate	35	Food Thickeners	269
desogestrel	74	Fexofenadine hydrochloride		Foods And Supplements For In	
Ethinyloestradiol with		Fibro-vein		Errors Of Metabolism	
levonorgestrel	75	Filgrastim		Foradil	
Ethinyloestradiol with		Finasteride		Forteo	
norethisterone	75	Fingolimod		Fortini	
Ethosuximide		Firazyr		Fortini Multi Fibre	
Etopophos		Flagyl		Fortisip	
Etoposide		Flagyl-S		Fortisip Multi Fibre	
Etoposide phosphate		Flamazine		Fosamax	
Etravirine		Flecainide acetate		Fosamax Plus	
Eumovate		Flecainide BNM		Framycetin sulphate	
Evara		Flecainide Controlled Release		Frebini Energy	262
Everet		Teva	49	Frebini Energy Fibre	262
Everolimus		Fleet Phosphate Enema		Frebini Original	
Evista		Flixonase Hayfever & Allergy		Frebini Original Fibre	262
Evusheld		Flixotide		Fresubin 2kcal HP	
Exemestane		Flixotide Accuhaler		Fresubin HP Energy	
Exjade		Florinef		Fresubin HP Energy Fibre	
Extemporaneously Compour		Fluanxol		Fresubin Intensive	
Preparations and	laca	Flucil		Fresubin Original	
Galenicals	255	Flucloxacillin		Fresubin Original Fibre	
Eye Preparations		Flucloxacillin-AFT		Frisium	
Eylea		Flucloxin		Frumil	
Ezetimibe		Flucon		Frusemide	
Ezetimibe Sandoz		Fluconazole		Fucicort	
Ezetimibe with simvastatin		Fludara Oral		Fucidin	
- F -		Fludarabine Ebewe		Fucithalmic	
Factor eight inhibitor bypassi	ina	Fludarabine phosphate		Fulvestrant	
fraction		Fludrocortisone acetate		Fungilin	
Famotidine		Fluids and Electrolytes		Furosemid-Ratiopharm	
Famotidine Hovid		Flumetasone pivalate		Furosemide [Frusemide]	
		•	241	Furosemide-Baxter	
FasenraFaslodex		Fluocortolone caproate with fluocortolone pivalate and		fusidic acid	
			0		60 61
Fatty Cream AFT		cinchocaine		Dermatological	ნპ, ნ
Febuxostat multishem		Fluorometholone		Infection	
Febuxostat multichem	115	Fluorouracil		Sensory	24
Feed Thickener Karicare	070	Fluorouracil Accord		- G -	401
Aptamil		Fluorouracil sodium		Gabapentin	120
FEIBA NF		Fluox		Gacet	
Felo 10 ER		Fluoxetine hydrochloride		Galsulfase	
Felo 5 ER		Flupenthixol decanoate		Galvumet	
Felodipine	52	Flutamide	164	Galvus	12

Gardasil 9	280	Havrix279 Hydrocortisone with natamycin and	t
Gastrodenol	10	Havrix Junior279 neomycin	6
Gaviscon Extra Strength	6	Haylor syrup238 Hydrogen peroxide	6
Gaviscon Infant	6	healthE Dimethicone 10%65 Hydroxocobalamin	
Gazyva	201	healthE Dimethicone 4% Lotion 67 Hydroxocobalamin Panpharma	3
Gefitinib		healthE Dimethicone 5%65 hydroxycarbamide	
GEM Aqueous Cream	66	healthE Glycerol BP255 Hydroxychloroquine	11
Gemcitabine Ebewe	148	healthE Urea Cream66 Hydroxyurea	
Gemcitabine hydrochloride	148	Healtheries Simple Baking Mix270 [hydroxycarbamide]	15
Gemtuzumab ozogamicin	191	Hemastix78 Hygroton	
Genoptic	247	Hemlibra	
Gentamicin sulphate		Heparin sodium44 Hymenoptera	23
Infection	95	Heparin Sodium Panpharma44 Hyoscine butylbromide	
Sensory	247	Heparinised saline44 Hyoscine hydrobromide	
Gilenya	134	Heparon Junior261 Hypam	13
Ginet		Hepatitis A vaccine279 Hyperuricaemia and Antigout	11
Glatiramer acetate	134	Hepatitis B recombinant Hypromellose	
Glecaprevir with pibrentasvir	103	vaccine	
Glibenclamide		Herceptin	
Gliclazide	11	Hiberix	9
Glipizide		Hiprex	16
Glivec		Histaclear238 Ibrutinib	15
Glizide		Histafen238 Ibuprofen	
Glucagen Hypokit		Holoxan145 Icatibant	
Glucagon hydrochloride		Horleys Bread Mix270 Idarubicin hydrochloride	
Glucerna Select		Horleys Flour270 Idursulfase	
Glucose [Dextrose]		Hormone Replacement Therapy - Ifosfamide	
Gluten Free Foods		Systemic	
Glycerin with sodium saccharin		HPV280 lloprost	
Glycerin with sucrose		Humalog11 Imatinib mesilate	
Glycerol		Humalog Mix 2511 Imatinib-Rex	
Alimentary	26	Humalog Mix 5011 Imbruvica	15
Extemporaneous		Human papillomavirus (6, 11, 16, 18, Imfinzi	
Glyceryl trinitrate		31, 33, 45, 52 and 58) vaccine Imigran	
Alimentary	8	[HPV]	
Cardiovascular		Humatin96 Imiguimod	
Glycopyrronium		Humira183 Immune Modulators	
Glycopyrronium bromide		HumiraPen183 Immunosuppressants	
Glycopyrronium with		Humulin 30/7011 Incruse Ellipta	
indacaterol	242	Humulin NPH11 Indacaterol	
Go Healthy		Humulin R	
Gold Knight		Hyaluronic acid251 Infanrix IPV	
Gold Knight XL		Hydralazine58 Infanrix-hexa	
Goserelin		Hydralazine hydrochloride58 Infant Formulae	
Gutron		Hydrocortisone Infatrini	
Gynaecological Anti-infectives.		Dermatological64 Infliximab	
- H -		Hormone80 Influenza vaccine	28
Habitrol	142	Hydrocortisone (PSM)64 Inhaled Corticosteroids	
Haemophilus influenzae type B		Hydrocortisone acetate	
vaccine		Hydrocortisone acetate with Beta-adrenoceptor Agonists	23
Haldol		pramoxine hydrochloride	
Haldol Concentrate		Hydrocortisone and paraffin liquid Instillagel Lido	
Haldol Decanoas		and lanolin	
Haloperidol		Hydrocortisone butyrate65, 70 Insulin aspart with insulin aspart	
Haloperidol decanoate		Hydrocortisone with cinchocaine8 protamine8	1
Harvoni		Hydrocortisone with miconazole65 Insulin glargine	
1 IUI VOI II	100	riyarooonaoone war iniconazole	1

					=
Insulin glulisine11	Isopto Carpine	250	Lamictal	1	12
Insulin isophane11	Isosorbide mononitrate	57	Lamivudine		
Insulin isophane with insulin	Isosource Standard	267	Lamivudine Alphapharm	1	0
neutral11	Isotretinoin	62	Lamivudine Viatris	1	0
Insulin lispro11	Ispaghula (psyllium) husk	26	Lamotrigine	1	2
Insulin lispro with insulin lispro	Itch-Soothe	64	Lamprene	1	10
protamine11	Itraconazole	98	Lanoxin		4
Insulin neutral10	Itrazole	98	Lanoxin Paediatric Elixir		
Insulin pen needles16	Ivacaftor	244	Lanoxin PG		4
Insulin pump16	Ivermectin	67	Lanoxin S29		
Insulin pump cartridge21	- J -		Lansoprazole		
Insulin pump infusion set (steel	Jadelle	76	Lantus		1
cannula)21	Jakavi		Lantus SoloStar		
Insulin pump infusion set (steel	Jardiamet	14	Lanvis		
cannula, straight insertion) 22	Jardiance	14	Lanzol Relief		
Insulin pump infusion set (teflon	Jaydess		Lapatinib ditosylate		
cannula)23	Jevity HiCal RTH		Largactil		
Insulin pump infusion set (teflon	Jevity Plus		Laronidase		
cannula, angle insertion with	Jevity RTH		Lasix		
insertion device)24	Jinarc		Latanoprost		
Insulin pump infusion set (teflon	Juno Pemetrexed		Latanoprost with timolol		
cannula, angle insertion) 24	- K -	149	Lax-Suppositories		
		224	Lax-suppositories Glycerol		
Insulin pump infusion set (teflon	Kadcyla				
cannula, straight insertion with	Kaletra		Laxatives		
insertion device)	Kalydeco		Laxsol		
Insulin pump infusion set (teflon	Kemadrin		Ledipasvir with sofosbuvir		
cannula, straight insertion) 24	Kenacomb		Leflunomide		
Insulin pump reservoir24	Kenacort-A 10		Lenalidomide		
Insulin syringes, disposable with	Kenacort-A 40		Letrole		
attached needle16	Kenalog in Orabase		Letrozole		
Intelence	Ketocal 3:1		Leukeran FC	1	4
Interferon beta-1-alpha134	KetoCal 4:1	277	Leukotriene Receptor		
Interferon beta-1-beta134	Ketoconazole		Antagonists	2	24:
Intra-uterine device74	Dermatological		Leuprorelin		
Invega Sustenna131	Infection		Leustatin		
Invega Trinza132	Ketogenic Diet		Levetiracetam		
IPCA-Frusemide53	Ketoprofen	111	Levetiracetam-AFT	1	2
IPCA-Metoprolol51	KetoSens	14	Levlen ED		. 7
IPCA-Propranolol51	Keytruda		Levocabastine	2	249
IPOL286	Kindergen	261	Levocarnitine		3
Ipratropium bromide241, 245	Kivexa	106	Levodopa with benserazide	1	[1]
Iressa159	Klacid		Levodopa with carbidopa		
Irinotecan Actavis 100148	Alimentary	9	Levomepromazine	1	130
Irinotecan hydrochloride148	Infection	91	Levomepromazine		
Irinotecan-Rex148	Kliogest	83	hydrochloride	1	13
Iron (as ferric carboxymaltose)35	Kliovance	83	Levonorgestrel		
Iron polymaltose36	Kogenate FS	41	Genito-Urinary		.70
Isentress108	Konakion MM	41	Hormone		8
Isentress HD108	Konsyl-D		Levonorgestrel BNM		
Ismo 2057	Kuvan		Levothyroxine		
Ismo 40 Retard57	- L -		Lidocaine [Lignocaine]		
Isoniazid100	Labetalol	51	Lidocaine [Lignocaine]		
Isoniazid with rifampicin100	Lacosamide		hydrochloride	1	119
Isoptin52	Lactulose		Lidocaine [Lignocaine] with		
Isoptin Retard52	Laevolac		chlorhexidine	1	110
Isoptin SR	Lagevrio		Lidocaine [Lignocaine] with		
			=ounio [=ignoounio] will		

prilocaine	119	Madopar 125	117	Methotrexate Ebewe	148
Lidocaine-Baxter	119	Madopar 250	117	Methotrexate Sandoz	148
Lidocaine-Claris	119	Madopar 62.5	117	Methyl hydroxybenzoate	255
Life Extension	31	Madopar HBS		Methylcellulose	255
Lignocaine	.118-119	Madopar Rapid	117	Methylcellulose with glycerin and	
Lioresal Intrathecal	116	Magnesium hydroxide		sodium saccharin	255
Lipid-Modifying Agents	55	Magnesium sulphate	36	Methylcellulose with glycerin and	
Liquigen	259	Mantoux	287	sucrose	255
Liraglutide	12	Marevan	44	Methyldopa	53
Lisinopril	47	Marine Blue Lotion SPF 50+	71	Methyldopa Mylan	53
Litak	147	Martindale Pharma	252	Methyldopa Mylan S29	53
Lithium carbonate	130	Mask for spacer device	245	Methylnaltrexone bromide	26
Livostin	249	Maviret	103	Methylphenidate ER - Teva	137
LMX4	119	Maxidex	248	Methylphenidate hydrochloride	137
Lo-Oralcon 20 ED	75	Maxitrol	248	Methylphenidate hydrochloride	
Locacorten-Viaform ED's	247	MCT oil (Nutricia)	259	extended-release	138
Local preparations for Anal and	t	Measles, mumps and rubella		Methylprednisolone	80
Rectal Disorders	8	vaccine	283	Methylprednisolone (as sodium	
Locasol	273	Mebendazole	90	succinate)	80
Locoid	65, 70	Mebeverine hydrochloride	8	Methylprednisolone aceponate	65
Locoid Crelo	65	Medrol	80	Methylprednisolone acetate	81
Locoid Lipocream	65	Medroxyprogesterone acetate		Methylxanthines	243
Locorten-Vioform	247	Genito-Urinary	76	Metoclopramide Actavis 10	129
Lodoxamide		Hormone	82-83	Metoclopramide hydrochloride	
Logem	126	Mefenamic acid		Metolazone	54
Lomide	249	Melatonin	134	Metopirone	89
Lomustine	146	Melpha	146	Metoprolol IV Mylan	51
Loniten	58	Melphalan	146	Metoprolol IV Viatris	51
Loperamide hydrochloride	<mark>6</mark>	Menactra	283	Metoprolol succinate	51
Lopinavir with ritonavir	107	Meningococcal (groups A, C, Y	and	Metoprolol tartrate	51
Lopinavir/Ritonavir Mylan	107	W-135) conjugate vaccine	283	Metrogyl	100
Loprofin	272	Meningococcal B multicompone	nt	Metronidazole	100
Loprofin Mix	272	vaccine	284	Metyrapone	89
Lorafix		Meningococcal C conjugate		Mexiletine hydrochloride	49
Loratadine	238	vaccine	284	Miacalcic	79
Lorazepam	133	MenQuadfi	283	Micolette	27
Lorstat	55	Menthol	64	Micolette-S29	27
Losartan Actavis	48	Mepolizumab	200	Miconazole	33
Losartan potassium	48	Mercaptopurine	148	Miconazole nitrate	
Losartan potassium with		Mercilon 28	74	Dermatological	63
hydrochlorothiazide	48	Mesalazine	7	Genito-Urinary	76
Lovir		Mesna	152	Micreme	76
Loxamine	124	Mestinon	111	Micreme H	65
Lucrin Depot 1-month	88	Metabolic Disorder Agents	27	Microgynon 20 ED	75
Lucrin Depot 3-month	88	Metformin hydrochloride	12	Microgynon 30	75
Lyderm	68	Metformin Mylan	12	Microlut	76
Lynparza	153	Metformin Viatris		Midazolam	135
Lyrica	126	Methadone BNM	121	Midazolam Mylan	135
- M -		Methadone hydrochloride		Midazolam-Baxter	135
m-Eslon	122	Extemporaneous	255	Midodrine	
Mabthera	204	Nervous		Mifegyne	78
Macro Organic Psyllium Husk .	26	Methenamine (hexamine)		Mifepristone	
Macrobid		hippurate	110	Milpharm	
Macrogol 3350 with potassium		Methopt		Minerals	
chloride, sodium bicarbonate	e and	Methotrexate		Mini-Wright AFS Low Range	245
sodium chloride	<mark>27</mark>	Methotrexate DBL Onco-Vial	148	Mini-Wright Standard	

Minidiab	11	Moments	73	Natulan	154
MiniMed 1.8 Reservoir		Mometasone furoate	65	Nausafix	129
MMT-326A	24	Monogen	260	Nausicalm	128
MiniMed 3.0 Reservoir		Montelukast	243	Navelbine	157
MMT-332A	24	Montelukast Mylan	243	Nefopam hydrochloride	119
MiniMed 770G	16	Montelukast Viatris	243	Neisvac-C	
MiniMed Mio MMT-921A	23	Moroctocog alfa [Recombinant fac	tor	Neo-Mercazole	83
MiniMed Mio MMT-923A	23	VIII]	40	Neocate Gold	
MiniMed Mio MMT-925A	23	Morphine hydrochloride		Neocate Junior Unflavoured	273
MiniMed Mio MMT-941A	23	Morphine sulphate		Neocate Junior Vanilla	
MiniMed Mio MMT-943A	23	Motetis		Neocate SYNEO	
MiniMed Mio MMT-945A	23	Mouth and Throat	32	Neoral	
MiniMed Mio MMT-965A	23	Movapo	117	Neostigmine metilsulfate	111
MiniMed Mio MMT-975A	23	Moxifloxacin		Nepro HP (strawberry)	
MiniMed Quick-Set MMT-386A	23	MSUD Maxamum	271	Nepro HP (vanilla)	
MiniMed Quick-Set MMT-387A	23	Mucilaginous laxatives with		Nepro HP RTH	
MiniMed Quick-Set MMT-396A		stimulants	26	Neulactil	
MiniMed Quick-Set MMT-397A		Mucolytics		Neulastim	
MiniMed Quick-Set MMT-398A		Mucosoothe		Neuraxpharm	
MiniMed Quick-Set MMT-399A		Multiple Sclerosis Treatments		NeuroTabs	
MiniMed Silhouette MMT-368A		Multivitamin renal		Nevirapine	
MiniMed Silhouette MMT-377A		Multivitamins		Nevirapine Alphapharm	
MiniMed Silhouette MMT-378A		Mupirocin		Nevirapine Viatris	
MiniMed Silhouette MMT-381A		Muscle Relaxants		Nicorandil	
MiniMed Silhouette MMT-382A		Mvite		Nicotine	
MiniMed Silhouette MMT-383A		Myambutol		Nifedipine	
MiniMed Silhouette MMT-384A		Mycobutin		Nifuran	
MiniMed Sure-T MMT-864A		MycoNail		Nilotinib	
MiniMed Sure-T MMT-866A		Mycophenolate mofetil		Nilstat	100
MiniMed Sure-T MMT-874A		Mydriacyl		Alimentary	20
MiniMed Sure-T MMT-876A		Mylan (12 hr release)		Genito-Urinary	
MiniMed Sure-T MMT-884A				Infection	
MiniMed Sure-T MMT-886A		Mylan (24 hr release)		Nintedanib	
		Mylan Atenolol			
Minims Produissions		Mylan Clomiphen		Nipent Nirmatrelvir with ritonavir	
Minims Prednisolone		Mylan Indapamide		Nitrates	
Minirin		Mylan Italy (24 hr release)			
Minirin Melt		Myleran		Nitroderm TTS	
Mino-tabs		Mylotarg	191	Nitrofurantoin	
Minocycline hydrochloride		Myometrial and Vaginal Hormone	77	Nitrolingual Pump Spray	51
Minorycin		Preparations		Nivestim	44
Minor Skin Infections		Myozyme N -	21	Nivolumab	
Minoxidil				Nizoral	
Mirena		Nadolol		Nodia	
Mirtazapine		Nadolol BNM		Noflam 250	
Misoprostol		Naglazyme		Noflam 500	
Mitomycin C		Naloxone hydrochloride		Non-Steroidal Anti-Inflammatory	
Mitozantrone		Naltraccord		Drugs	111
Mitozantrone Ebewe		Naltrexone hydrochloride		Nonacog gamma, [Recombinant	
Mixtard 30		Naphazoline hydrochloride		Factor IX]	40
MMR II		Naphcon Forte		Norethisterone	
Moclobemide		Naprosyn SR 1000		Genito-Urinary	76
Modafinil		Naprosyn SR 750		Hormone	
Modavigil		Naproxen		Norflex	
Moduretic		Narcaricin mite		Norfloxacin	
Molaxole		Nasal Preparations		Noriday 28	
Molnupiravir	104	Natalizumab	134	Norimin	75

				_
Normacol Plus	26	Octocog alfa [Recombinant factor	Other Endocrine Agents	88
Normison	. 135	VIII] (Advate)40	Other Oestrogen Preparations	8
Norpress	.123	Octocog alfa [Recombinant factor	Other Progestogen	
Nortriptyline hydrochloride	.123	VIII] (Kogenate FS)41	Preparations	83
Norvir		Octreotide165	Other Skin Preparations	
Noumed		Octreotide Depot Teva165	Ovestin	
Dermatological	64	Octreotide GH165	Genito-Urinary	7
Nervous121,		Octreotide long-acting	Hormone	
Noumed Paracetamol		Oestradiol82	Oxaliplatin1	
Noumed Pethidine		Oestradiol valerate	Oxaliplatin Accord1	
NovaSource Renal		Oestradiol with norethisterone83	Oxaliplatin Actavis 1001	
Novatretin		Oestriol	Oxaliplatin Ebewe1	140
NovoMix 30 FlexPen			Oxis Turbuhaler2	
		Genito-Urinary77		
NovoRapid		Hormone83	Oxpentifylline	. 30
NovoRapid FlexPen		Oestrogens82	Oxybutynin	. / (
NovoRapid Penfill		Ofev242	Oxycodone hydrochloride1	122
NovoSeven RT		Oil in water emulsion66	Oxycodone Sandoz1	22
Noxafil		Olanzapine130–131	OxyNorm1	
Nozinan	. 130	Olaparib153	Oxytocin	
Nozinan (Swiss)	. 130	Olbetam55	Oxytocin BNM	7
Nozinan S29		Olbetam S2955	Oxytocin GH	7
Nucala	.200	Olopatadine251	Oxytocin with ergometrine	
Nuelin	.243	Olopatadine Teva251	maleate	7
Nuelin-SR	.243	Olsalazine8	Ozurdex2	248
Nupentin	.125	Omalizumab202	-P-	
Nusinersen		Omeprazole9	Pacifen1	116
Nutilis		Omeprazole actavis 109	Pacimol1	
Nutren Diabetes		Omeprazole actavis 209	Paclitaxel1	
Nutrient Modules		Omeprazole actavis 409	Paclitaxel Actavis1	
Nutrini Energy Multi Fibre		Omnitrope84	Paclitaxel Ebewe1	
Nutrini Energy RTH		Omnitrope S2984	Paediatric Seravit	
Nutrini Low Energy Multi Fibre		Onbrez Breezhaler239	Palbociclib1	
		Oncaspar LYO154	Paliperidone1	
Nutrini Peptisorb		OncoTICE174	Polineridana polmitata	10
Nutrini Peptisorb Energy			Paliperidone palmitate1	
Nutrini RTH	.202	Ondansetron 129	Palivizumab	
Nutrison 800 Complete Multi	007	Ondansetron ODT-DRLA129	Pamidronate disodium1	
Fibre		One-Alpha34	Pamisol	
Nutrison Advanced Diason		One-Alpha S2934	Pamol1	
Nutrison Advanced Peptisorb		Onrex129	Pancreatic enzyme	
Nutrison Concentrated		Opdivo229	Pantoprazole	
Nutrison Energy		Ora-Blend255	Panzop Relief	
Nutrison Energy Multi Fibre		Ora-Blend SF255	Panzytrat	
Nutrison Multi Fibre		Ora-Plus255	Papaverine hydrochloride	
Nutrison Standard RTH	.267	Ora-Sweet255	Para-amino salicylic acid1	0.
Nyefax Retard	52	Ora-Sweet SF255	Paracare1	2
Nystatin		Orabase33	Paracetamol1	
Alimentary	33	Oral and Enteral Feeds260	Paracetamol (Ethics)1	2
Genito-Urinary	76	Oralcon 30 ED75	Paracetamol + Codeine	
Infection		Oratane62	(Relieve) 1	22
NZB Low Gluten Bread Mix		Ordine121	Paracetamol with codeine1	2
-0-		Orgran	Paraffin	
Obinutuzumab	.201	Ornidazole	Paraffin liquid with wool fat2	
Obstetric Preparations		Orphenadrine citrate116	Parasiticidal Preparations	6.
Ocicure		Ortho-tolidine	Parnate1	
Ocrelizumab		Oruvail SR	Parnate S291	
Ocrevus		Osmolite RTH	Paromomycin	
Oci c vuo	. 104	20/	1 atottioniyotti	30

Paroxetine	124	Phosphorus	46	Prednisolone	8
Paser		Phytomenadione		Prednisolone acetate	24
Paxam	133	Pilocarpine hydrochloride	250	Prednisolone sodium	
Paxlovid	105	Pimafucort	65	Prednisolone sodium	
Pazopanib		Pimecrolimus	69	phosphate	
Peak flow meter	245	Pine tar with trolamine laurilsulfate		Prednisolone-AFT	
Pedialyte - Bubblegum	46	and fluorescein	70	Prednisone	
Pediasure	262	Pinetarsol	70	Prednisone Clinect	
Pediasure Plus		Pioglitazone		Pregabalin	
Pediasure RTH		Pirfenidone	242	Pregabalin Pfizer	
Pegaspargase		Pizotifen	128	Pregnancy Tests - hCG Urine	
Pegasys	108	PKU Anamix Infant	272	Premarin	
Pegfilgrastim		PKU Anamix Junior	272	Prevenar 13	28
Pegylated interferon alfa-2a	108	PKU Anamix Junior Chocolate	272	Priadel	130
Pembrolizumab	230	PKU Anamix Junior LQ	272	Primaquine	9
Pemetrexed	149	PKU Anamix Junior Orange	272	Primidone	120
Penicillamine	112	PKU Anamix Junior Vanilla	272	Primidone Clinect	
Penicillin G	93	PKU Lophlex LQ 10		Primolut N	8
PenMix 30	11	PKU Lophlex LQ 20	272	Priorix	28
PenMix 50	11	PKU Lophlex Powder	272	Probenecid	110
Pentasa	7	PKU Lophlex Sensation 20	272	Probenecid-AFT	110
Pentostatin [Deoxycoformycin]	154	Plaquenil	112	Procarbazine hydrochloride	15
Pentoxifylline [Oxpentifylline]	58	Plendil ER	52	Prochlorperazine	129
Peptamen Junior	262	Pneumococcal (PCV10) conjugate		Proctofoam	
Pepti-Junior	276	vaccine	284	Proctosedyl	
Peptisorb		Pneumococcal (PCV13) conjugate		Procyclidine hydrochloride	11
Perhexiline maleate	52	vaccine	285	Progesterone	8
Pericyazine	130	Pneumococcal (PPV23)		Proglicem	
Perindopril		polysaccharide vaccine	286	Proglycem	10
Periset	129	Pneumovax 23		Progynova	8
Perjeta	204	Podophyllotoxin	71	Prolia	112
Permethrin		Polaramine	238	Promethazine hydrochloride	23
Perrigo	71	Poliomyelitis vaccine	286	Propafenone hydrochloride	50
Pertuzumab		Poloxamer		Propamidine isethionate	
Peteha	101	Poly-Gel	251	Propranolol	
Pethidine hydrochloride	122	Poly-Tears		Propylene glycol	
Pevaryl		Poly-Visc		Propylthiouracil	
Pexsig		Polycal		Prostacur	
Pfizer Exemestane		Polyethylene glycol 400 and		Protaphane	1
Pharmacy Health Sorbolene with		propylene glycol	251	Protaphane Penfill	
Glycerin		Ponstan	111	Protifar	25
Pharmacy Services		Posaconazole	98	Protionamide	
Pheburane		Posaconazole Juno	98	Provera	8
Phenasen	149	Postinor-1	76	Provera HD	8
Phenobarbitone	126	Potassium chloride4	15-46	Psoriasis and Eczema	
Phenobarbitone sodium		Potassium citrate	78	Preparations	68
Extemporaneous	255	Potassium iodate	35	PTU	
Nervous		Povidone iodine	67	Pulmicort Turbuhaler	23
Phenoxybenzamine		Pradaxa		Pulmozyme	
hydrochloride	47	Pramipexole hydrochloride		Puri-nethol	
Phenoxymethylpenicillin (Penicil		Pravastatin		Puria	
V)		Pravastatin Mylan	55	Pyrazinamide	
Phenytoin sodium		Pravastatin Viatris	55	Pyridostigmine bromide	
Phillips Milk of Magnesia		Praziquantel		Pyridoxine hydrochloride	
Phlexy 10		Prazosin		Pyridoxine multichem	
Phosphate Phebra		Pred Forte		Pyrimethamine	
					-

Pytazen SR	41	Risperidone (Teva)	131	Secukinumab	21
- Q -		Risperon	131	Sedatives and Hypnotics	13
Quetapel	130	Ritalin		Seebri Breezhaler	24
Quetiapine		Ritalin LA	138	Selegiline hydrochloride	
Quick-Set MMT-392		Ritonavir		Senna	2
Quick-Set MMT-393		Rituximab (Mabthera)		Senokot	2
Quinapril		Rituximab (Riximyo)		SensoCard	
Quinapril with		Rivaroxaban		Serc	
hydrochlorothiazide	48	Rivastigmine		Serenace	
Qvar		Rivastigmine Patch BNM 10		Seretide	
-R-		Rivastigmine Patch BNM 5		Seretide Accuhaler	
RA-Morph	121	Rivotril		Serevent	
Ralicrom		Riximyo		Serevent Accuhaler	
Raloxifene hydrochloride		RIXUBIS		Sertraline	
Raltegravir potassium		Rizamelt		Setrona	
Ramipex		Rizatriptan		Setrona AU	
Ramipril		RoActemra S29		Sevredol	
		Robinul		Sex Hormones Non	12
Ranbaxy-Cefaclor					0
Rapamune		Ronapreve		Contraceptive	
Rasagiline		Ropin		Shield XL	
Reandron 1000		Ropinirole hydrochloride		Shingles vaccine	
Recombinant factor IX	,	Rosuvastatin		Shingrix	28
Recombinant factor VIIa		Rosuvastatin Viatris		SII-Onco-BCG	1/
Recombinant factor VIII		Rotarix		Sildenafil	6
Rectogesic		Rotavirus oral vaccine		Silhouette MMT-373	
Redipred		Roxane		Siltuximab	
Relieve		Roxane-Propranolol		Simvastatin	5
Relistor		Roxithromycin		Simvastatin Mylan	
Remicade		Rubifen		Simvastatin Viatris	
Renilon 7.5		Rubifen SR		Sinemet	
Resonium-A		Rugby Capsaicin Topical Crea		Sinemet CR	
Resource Beneprotein	259	Musculoskeletal	112	Sirolimus	23
Respigen		Nervous	119	Siterone	8
Respiratory Devices	245	Rurioctocog alfa pegol [Recor	nbinant	Slow-Lopresor	5
Respiratory Stimulants		factor VIII]	41	Smith BioMed Rapid Pregnancy	
Retinol palmitate	251	Ruxolitinib	162	Test	
ReTrieve	62	Rythmodan	49	Sodibic	4
Retrovir	107	Rytmonorm		Sodium acid phosphate	2
Revlimid	152	· -S-		Sodium alginate	
Revolade	38	Sabril	127	Sodium benzoate	
Riboflavin	30	Sacubitril with valsartan	48	Sodium bicarbonate	
Ribomustin	144	Sagent	157	Blood	. 45–4
Ricit	77	SalAir	240	Extemporaneous	
Rifabutin	101	Salazopyrin	8	Sodium calcium edetate	
Rifadin	101	Salazopyrin EN		Sodium chloride	
Rifampicin		Salbutamol		Blood	4
Rifaximin		Salbutamol with ipratropium		Respiratory	
Rifinah		bromide	241	Sodium citrate with sodium lauryl	
Rilutek		Salicylic acid	70	sulphoacetate	
Riluzole		Salmeterol	239	Sodium citro-tartrate	
RINVOQ		Sandomigran		Sodium cromoglicate	
Riodine		Sanofi Primaquine		Alimentary	
Risedronate Sandoz		Sapropterin dihydrochloride	30	Sensory	24
Risedronate sodium		Scalp Preparations		Sodium Fusidate [fusidic acid]	24
Risperdal Consta		Scopoderm TTS		Dermatological	6
Risperidone		Sebizole		Dominatological	
1 110PUHUUHU	101-102	OODILUIG	/ 🗸		

Infection	96	Sure-T MMT-873	21	Tensipine MR10	52
Sensory	247	Survimed OPD	264	Tepadina	146
Sodium hyaluronate [Hyaluronic		Sustagen Hospital Formula	267	Terbinafine	99
acid]	251	Sustagen Hospital Formula		Terbutaline sulphate	
Sodium phenylbutyrate	31	Active	267	Teriflunomide	
Sodium picosulfate		Sustanon Ampoules		Teriparatide	
Sodium polystyrene sulphonate		Sylvant		Testosterone	
Sodium tetradecyl sulphate		Symbicort Turbuhaler 100/6		Testosterone cipionate	
Sodium valproate	126	Symbicort Turbuhaler 200/6		Testosterone esters	81
Sofradex		Symbicort Turbuhaler 400/12		Testosterone undecanoate	
Soframycin	247	Symmetrel		Tetrabenazine	118
Solgar	28-31	Sympathomimetics		Tetrabromophenol	78
Solifenacin Mylan		Synacthen		Tetracosactrin	
Solifenacin succinate		Synacthen Depot		Tetracycline	94
Solifenacin Viatris	78	Synacthene Retard		Teva Lisinopril	47
Solu-Cortef	80	Synagis		Thalidomide	
Solu-Medrol	80	Synflorix		Thalomid	155
Solu-Medrol-Act-O-Vial	80	Synthroid		Theophylline	243
Somatropin (Omnitrope)		Syntometrine		Thiamine hydrochloride	
Sotalol		Syrup (pharmaceutical grade)		Thiamine multichem	
Spacer device		Systane Unit Dose		THIO-TEPA	
Span-K		· -Т-		Thioguanine	149
Spinal Muscular Atrophy		Tacrolimus		Thiotepa	
Spinraza		Dermatological	70	Thyroid and Antithyroid Agents	
Spiolto Respimat		Oncology		Ticagrelor	
Spiractin		Tacrolimus Sandoz		Ticagrelor Sandoz	41
Spiriva		Taliglucerase alfa		Tilcotil	
Spiriva Respimat		Tambocor		Timolol	249
Spironolactone		Tamoxifen citrate		Timoptol XE	
Sporanox		Tamoxifen Sandoz		Tiotropium bromide	
Sprycel		Tamsulosin hydrochloride		Tiotropium bromide with	
Stelara		Tamsulosin-Rex		olodaterol	242
Stemetil	129	Tandem Cartridge	21	Tivicay	
Steril-Gene	81	Tandem t:slim X2 with Basal-IQ		Tixagevimab with cilgavimab	
SteroClear	245	Tap water	256	TMP	
Stesolid	124	Taro	98	Tobramycin	
Stimulants/ADHD Treatments	136	Taro-Testosterone	81	Infection	96
Stiripentol	126	Tasigna	160	Sensory	24
Stocrin		Tasmar		Tobramycin BNM	
Stomahesive	33	Taurine	31	Tobramycin Mylan	
Strattera	136	Tecentriq	228	Tobrex	
Strides Shasun	98	Tecfidera	134	Tocilizumab	
Stromectol	67	Tegretol	125	Tofranil	123
Sucralfate	10	Tegretol CR		Tolcapone	117
Sulfadiazine Silver	63	Telfast	238	Tolvaptan	54
Sulfadiazine sodium	96	Teligent	95	Topamax	
Sulfasalazine	8	Temaccord	154	Topical Products for Joint and	
Sulphur	70	Temazepam	135	Muscular Pain	112
Sulprix		Temozolomide		Topiramate	127
Sumagran		Tenofovir disoproxil		Topiramate Actavis	
Sumatriptan		Tenofovir Disoproxil Emtricitabin	ne	Total parenteral nutrition (TPN)	
Sunitinib		Mylan		TPN	
Sunitinib Pfizer		Tenofovir Disoproxil Emtricitabin		Tramadol hydrochloride	
Sunscreens	71	Viatr		Tramal SR 100	122
Sunscreens, proprietary		Tenofovir Disoproxil Mylan		Tramal SR 150	122
Sure-T MMT-863		Tenoxicam		Tramal SR 200	122

Trandate	51	Uromitexan	152	Vincristine sulphate	15
Tranexamic acid	41	Ursodeoxycholic acid	25	Vinorelbine	
Tranylcypromine sulphate	123	Ursosan	25	Vinorelbine Ebewe	
Trastuzumab		Ustekinumab	225	Viramune Suspension	10
Trastuzumab emtansine		Utrogestan	83	ViruPOS	24
Travatan	250	- V -		Vit.D3	
Travoprost	250	Vaccinations	278	Vita-B12	
Treatments for Dementia		Vaclovir	102	VitA-POS	25
Treatments for Substance		Valaciclovir	102	Vitabdeck	3
Dependence	140	Valganciclovir	102	Vital	26
Trental 400		Valganciclovir Mylan	102	Vitamin B complex	3
Tretinoin		Vancomycin	97	Vitamin B6 25	3
Dermatological	62	Vannair	240	Vitamins	.33-3
Oncology		Varenicline Pfizer	142	Vivonex TEN	26
Trexate		Varenicline tartrate	142	Voltaren	11
Triamcinolone acetonide		Varicella vaccine [Chickenpox		Voltaren D	11
Alimentary	33	vaccine]	287	Voltaren Ophtha	24
Dermatological	65	Varicella zoster vaccine [Shingle	S	Voltaren SR	11
Hormone		vaccine]		Volumatic	
Triamcinolone acetonide with		Varicella zoster virus (Oka strain) live	Voriconazole	9
gramicidin, neomycin and nys	statin	attenuated vaccine [shingles	,	Votrient	16
Dermatological	65	vaccine]	287	Vttack	9
Sensory		Various	252	- W -	
Triazolam		Varivax	287	Warfarin sodium	4
Trikafta	244	Vasodilators	58	Wart Preparations	7
Trimethoprim	97	Vasopressin Agonists	88	Wasp venom allergy treatment	
Trimethoprim with		Vasorex		Water	
sulphamethoxazole		Vebulis	61	Blood	4
[Co-trimoxazole]	97	Vedafil	60	Extemporaneous	25
Trisequens		Vedolizumab	226	White Soft Liquid Paraffin AFT	
Trisul	97	Veletri	61	Wool fat with mineral oil	6
Trophic Hormones		Venclexta	156	- X -	
Tropicamide		Venetoclax	156	Xarelto	4
Trulicity		Venlafaxine	124	Xifaxan	1
Trusopt	249	Venomil	238	XMET Maxamum	27
TruSteel		VENOX	238	Xolair	20
Tryzan		Ventolin	240	XP Maxamum	
Tuberculin PPD [Mantoux] test		Vepesid	151	Xylocaine	11
Tubersol		Verapamil hydrochloride		Xylocaine 2% Jelly	11
Two Cal HN		Vermox		Xyntha	4
Tykerb	159	Versacloz		· Z-	
Tysabri		Vesanoid	156	Zapril	4
. U -		Vexazone	12	Zarontin	
UK Synacthen	81	Vfend	99	Zaroxolyn	
Ultibro Breezhaler		Viaderm KC	65	Zavedos	
Ultraproct		Viatris		Zeffix	10
Umeclidinium		Cardiovascular	50	Zematop	7
Umeclidinium with vilanterol	242	Infection		Zetlam	10
Univent		Victoza		Ziagen	10
Upadacitinib	,	Vigabatrin		Zidovudine [AZT]	
Ural		Vigisom		Zidovudine [AZT] with	
Urea		Vildagliptin		lamivudine	10
Urex Forte		Vildagliptin with metformin		Ziextenzo	
Urinary Agents		hydrochloride	12	Zimybe	
Urinary Tract Infections		Vimpat	125	Zinc and castor oil	
Urinorm		Vinblastine sulphate		Zinc sulphate	

