June 2023 Volume 30

Editors:

Kaye Wilson, & Ayeshah Khan email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz
@Pharmaceutical Management Agency
ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written from and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	ind oddonig i narmae i	
Section A	General Rules	5
Section B	Alimanton, Tract C Matabaliana	
Section D	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	38
	Cardiovascular System	48
	Dermatologicals	63
	Genito Urinary System	72
	Hormone Preparations – Systemic	79
	Infections – Agents For Systemic Use	90
	Musculoskeletal System	111
	Nervous System	117
	Oncology Agents & Immunosuppressants	144
	Respiratory System & Allergies	238
	Sensory Organs	248
	Various	253
Section C	Extemporaneous Compounds (ECPs)	255
Section D	Special Foods	258
Section I	National Immunisation Schedule	280
	Index	291

Introducing Pharmac

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
Antacids and Antiflatulents			
Antacids and Reflux Barrier Agents			
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg posachet		30	✓ Gaviscon Infant
SODIUM ALGINATE Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (13.61)	60	Gaviscon Extra Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 ml	nl Acidex
Phosphate Binding Agents			
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) –	12.56	100	✓ Alu-Tab
Subsidy by endorsement		500 ml 173 ml	
Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed according		s or w	
Antidiarrhoeals			
Agents Which Reduce Motility			
* Tab 2 mg* Cap 2 mg	10.75	400 400	✓ Nodia✓ Diamide Relief
Rectal and Colonic Anti-inflammatories			
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy	166.50	90	✓ Entocort CIR
■ SA1886 Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant pract the following criteria:	titioner. Approvals va	alid for	or 6 months for applications meeting

1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and

2 Any of the following:

2.1 Diabetes; or

continued...

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	•	Manufacturer

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes: or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture: or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

THE RECORD TO LIKE		
Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP	✓ Colifoam✓ Cortifoam \$29
	21.1 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
OLSALAZINE				
Tab 500 mg	56.02	60	✓	Atnahs
				Olsalazine S29
	93.37	100	✓	Dipentum
Cap 250 mg	53.00	100	1	Dipentum
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential
				Prednisolone S29
SODIUM CROMOGLICATE				
Cap 100 mg	113.35	100	1	Ralicrom
SULFASALAZINE				
* Tab 500 mg	16.52	100	1	Salazopyrin
* Tab EC 500 mg		100		Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g11.06	30 g OP	Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg7.30	12	Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available on a	ì		
PSO	19.00	5	✓ Robinul
	65.45	10	✓ Max Health
Robinul to be Principal Supply on 1 September 2023			
(Max Health Inj 200 mcg per ml, 1 ml ampoule to be delisted 1 Sep	otember 2023)		
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO	6.35	5	✓ Buscopan
			✓ Buscopan S29 S29
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL - Wastage claimable

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement.......14.58 14 ✓ Klacid

- a) Maximum of 28 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

FΑ	MOTIDINE - Only on a prescription			
*	Tab 20 mg	4.91	100	✓ Famotidine
	·			Hovid S29
*	Tab 40 mg	8.48	100	✓ Famotidine
				Hovid \$29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement	57.02	10	✓ Mylan S29
	Subsidy by andorsament - Subsidised for nations race	ivina treatment as	nart of nallis	ative care

Proton Pump Inhibitors

LA	NSOPRAZOLE			
*	Cap 15 mg4	.20	100	✓ Lanzol Relief
*	Cap 30 mg5	.26	100	✓ Lanzol Relief
ON	IEPRAZOLE			
	For omeprazole suspension refer Standard Formulae, page 255			
*	Cap 10 mg1.	.94	90	✓ Omeprazole actavis 10
*	Cap 20 mg	.86	90	✓ Omeprazole actavis 20
*	Cap 40 mg	.11	90	✓ Omeprazole actavis 40
*	Powder – Only in combination42. Only in extemporaneously compounded omeprazole suspension.	.50	5 g	✓ Midwest
*	Inj 40 mg ampoule with diluent	.38	5	✓ <u>Dr Reddy's</u> Omeprazole
				✓ Ocicure S29
PA	NTOPRAZOLE			
*	Tab EC 20 mg	.99	90	✓ Panzop Relief
*	Tab EC 40 mg	.74	90	✓ Panzop Relief
•	Panzop Relief to be Principal Supply on 1 July 2023			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mgSUCRALFATE	14.51	50	✓	Gastrodenol S29
Tab 1 g	35.50 (48.28)	120		Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail pha Tab 550 mg	•	56	/	<u>Xifaxan</u>
▶ SA1461 Special Authority for Subsidy Initial application only from a gastroenterologist, hepatologist on the patologist. Approvals valid for 6 months where the patient hat tolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Practition the patologist. Approvals valid without further renewal unless not be nefiting from treatment.	as hepatic encephalop oner on the recomme	athy d	espite an n of a gast	adequate trial of maximum roenterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE – Special Authority see SA1320 below – Retail pha Cap 25 mg Cap 100 mg Oral liq 50 mg per ml	110.00 280.00	100 100 0 ml 0	✓)P ✓	Proglicem \$29 Proglicem \$29 Proglycem \$29 e5 Pharma \$29
■ SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valinypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid withou appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit − Up to 5 kit available on a PSO	t further renewal unle		fied where	
Insulin - Short-acting Preparations				
INSULIN NEUTRAL ▲ Inj human 100 u per ml		0 ml C	√ ✓	Actrapid Humulin R Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
INSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	/	NovoMix 30 FlexPen

	Subsidy (Manufacturer's P	rico) Subci	Fully Brand or dised Generic
	(Wanuacturers F	Per	✓ Manufacturer
INSULIN ISOPHANE			
▲ Inj human 100 u per ml	17.68	10 ml OP	✓ Humulin NPH✓ Protaphane
▲ Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH✓ Protaphane Penfill
INSULIN ISOPHANE WITH INSULIN NEUTRAL			· i i otapilano i onimi
▲ Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
,			✓ Mixtard 30
▲ Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
			✓ PenMix 30✓ PenMix 50
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,			
3 ml		5	✓ Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,			
3 ml	42.66	5	Humalog Mix 50
Insulin - Long-acting Preparations			
INSULIN GLARGINE ▲ Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
▲ Inj 100 u per ml, 3 ml		5	✓ Lantus
▲ Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
ilisuilii - napiu Actilig Freparations			
INSULIN ASPART			_
▲ Inj 100 u per ml, 10 ml		1	✓ NovoRapid
▲ Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
▲ Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
NSULIN GLULISINE ▲ Inj 100 u per ml, 10 ml	27.02	4	√ Anidro
▲ Inj 100 u per ml, 3 ml		1 5	✓ Apidra✓ Apidra
▲ Inj 100 u per ml, 3 ml disposable pen		5	✓ Apidra SoloStar
NSULIN LISPRO			F · · · · · · · · ·
▲ Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ Humalog
▲ Inj 100 u per ml, 3 ml		5	✓ Humalog
Alpha Glucosidase Inhibitors			
ACARBOSE			
# Tab 50 mg	8 95	90	✓ Accarb
* Tab 100 mg		90	✓ Accarb
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE	7.50	400	/ Decarl
* Tab 5 mg	7.50	100	✓ <u>Daonil</u>
GLICLAZIDE	45.40	500	/ Oll-1-1-
* Tab 80 mg	15.18	500	✓ Glizide
GLIPIZIDE	4.50	100	✓ Minidiah
* Tab 5 mg	4.58	100	✓ <u>Minidiab</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg	14.74	1,000		Metformin Mylan Metformin Viatris
* Tab immediate-release 850 mg	11.28	500	✓ i	Metformin Mylan Metformin Wylan Metformin Viatris
(Metformin Mylan Tab immediate-release 500 mg to be delisted 1	August 2023)		٠,	wedomin viadis
PIOGLITAZONE				
* Tab 15 mg		90	✓ \	<u>Vexazone</u>
* Tab 30 mg	7.30	90	✓ !	<u>Vexazone</u>
* Tab 45 mg	12.25	90	✓ !	<u>Vexazone</u>
VILDAGLIPTIN				
Tab 50 mg	35.00	60	✓ (Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	1	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	•	Galvumet

GLP-1 Agonists

 ${\tt DULAGLUTIDE\ - Special\ Authority\ see\ SA2065\ below\ -\ Retail\ pharmacy}$

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes: and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*: or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*: or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

LIRAGLUTIDE - Special Authority see SA2187 on the next page - Retail pharmacy

- a) Maximum of 9 inj per prescription
- b)
- a) Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

⇒SA2187 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial Special Authority approval for either an SGLT-2 inhibitor or GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*: or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*: or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*: or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

 a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack,

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 10 mg58.56	30	Jardiance
*	Tab 25 mg	30	 Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	8.56	60	✓ Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips.......20.00 1 OP ✓ CareSens Dual

			_
Subsidy	Fully	Brand or	Ī
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips26	6.20	50 test OP	SensoCard
-----------------------------	------	------------	-----------

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

IIVC	ocini i civ necescio i maximam di 200 dev per prescripti	OH		
*	29 g × 12.7 mm	10.95	100	✓ B-D Micro-Fine
*	31 g × 5 mm		100	✓ B-D Micro-Fine
*	31 g × 6 mm		100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED	LE - Maximum of 2	:00 dev per p	prescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g x 8 mm needle	13.56	100	B-D Ultra Fine II
		1.36	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓ B-D Ultra Fine II
		1.36	10	
		(1.99)		B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year pe	riod.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			Y2 with Rasal-IO

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	✓	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist: or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Fither
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	I Generic
\$	Per 🗸	Manufacturer

continued...

- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

	ALIMENTAR	TRAC	ANL	DMETABOLISM
	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
continued				
pump therapy; and				
4 The patient is continuing to derive benefit from pump the				
5 The patient had achieved and is maintaining a HbA1c of				
6 The patient has had no increase in severe unexplained h			seline;	and
7 The patient's HbA1c has not deteriorated more than 5 m 8 Either:	moi/moi trom baseline	e; and		
8.1 Applicant is a relevant specialist; or				
8.2 Applicant is a nurse practitioner working within the	eir vocational scone			
Renewal — (Previous use before 1 September 2012) only fro	•	st or nurse	practiti	ioner. Approvals valid for 2
years for applications meeting the following criteria:	m a roioram oposiam		p.ao	
All of the following:				
1 The patient is continuing to derive benefit according to the	e treatment plan and	has mainta	ained a	HbA1c of equal to or less
than 80 mmol/mol; and				
2 The patient's HbA1c has not deteriorated more than 5 m				
3 The patient has not had an increase in severe unexplain4 Either:	ed hypoglycaemic epi	sodes from	i baseli	ine; and
. —				
4.1 Applicant is a relevant specialist; or4.2 Applicant is a nurse practitioner working within the	air vocational scope			
	•			
INSULIN PUMP CARTRIDGE – Special Authority see SA1985	on page 19 – Retail p	harmacy		
a) Maximum of 3 sets per prescription				
b) Only on a prescriptionc) Maximum of 13 packs of cartridge sets will be funded per	rvoor			
Cartridge 300 U, t:lock × 10		1 OP	√ 1	Fandem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special		-		•
a) Maximum of 3 sets per prescription	Additionly 300 OATSO	o on page	10 11	ciali priarriacy
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ I	MiniMed Sure-T
				MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ I	MiniMed Sure-T
				MMT-886A
6 mm steel needle; 60 cm tubing x 10	130.00	1 OP	✓ I	MiniMed Sure-T
Commented mandles 00 and tables as 40	100.00	1 OD		MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	•	MiniMed Sure-T

8 mm steel needle; 80 cm tubing × 10130.00 1 OP

8 mm steel needle; 60 cm tubing × 10130.00

✓ MiniMed Sure-T MMT-874A ✓ MiniMed Sure-T

MMT-866A

6 mm steel needle; 29 G; manual insertion; 60 cm tubing x

MMT-876A

✓ Sure-T MMT-873

8 mm steel needle; 29 G; manual insertion; 60 cm tubing x

✓ Sure-T MMT-863 1 OP

1 OP

1 OP

(Sure-T MMT-863 6 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock to be delisted 1 December 2023)

(Sure-T MMT-873 8 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock to be delisted 1 December 2023)

Subsidy	H	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
 \$	Per	✓	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles		1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line × 10 with	130 00	1 OP	✓ TruSteel

1 OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

13 mm teflon needle, 45 cm tubing × 10130.00	1 OP
13 mm teflon needle, 60 cm tubing × 10130.00	1 OP

6 mm teflon needle, 80 cm blue tubing.......130.00

6 mm teflon needle, 80 cm clear tubing × 10130.00

- ✓ MiniMed Silhouette MMT-382A
- ✓ MiniMed Silhouette MMT-368A
 ✓ MiniMed Silhouette
- MMT-381A

 MiniMed Silhouette

✓ MiniMed Silhouette

- MMT-383A ✓ MiniMed Silhouette MMT-377A
- MMT-378A ✓ MiniMed Silhouette MMT-384A
- 6 mm teflon needle, 110 cm tubing × 10
 10 P

 6 mm teflon needle, 45 cm blue tubing × 10
 130.00
 1 OP
- ✓ MiniMed Quick-Set MMT-398A
 ✓ MiniMed Mio MMT-941A
- ✓ MiniMed Mio MMT-921A
 - ✓ MiniMed Mio MMT-943A
 - ✓ MiniMed Mio MMT-923A
 - ✓ MiniMed Quick-Set MMT-399A
 - ✓ MiniMed Mio MMT-945A
 - ✓ MiniMed Mio MMT-965A
 - ✓ MiniMed Mio MMT-925A
 - ✓ MiniMed Quick-Set MMT-387A
 - ✓ MiniMed Quick-Set MMT-396A
 - ✓ MiniMed Quick-Set MMT-397A
 - ✓ MiniMed Mio MMT-975A
 - ✓ MiniMed Quick-Set MMT-386A

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION WITH INSERTION DEVICE) - Special Authority see SA1985 on page 19 - Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon cannula; angle insertion; insertion device; 110 cm line × 10 with 10 needles......140.00 1 OP ✓ AutoSoft 30 13 mm teflon cannula; angle insertion; insertion device: 60 cm line × 10 with 10 needles......140.00 1 OP ✓ AutoSoft 30 INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION) - Special Authority see SA1985 on page 19 -Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 17 mm teflon cannula; angle insertion; 60 cm line × 10 with 1 OP ✓ Silhouette MMT-373 (Silhouette MMT-373 17 mm teflon cannula: angle insertion: 60 cm line x 10 with 10 needles: luer lock to be delisted 1 December 2023) INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) - Special Authority see SA1985 on page 19 - Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device; ✓ AutoSoft 90 1 OP 6 mm teflon cannula: straight insertion: insertion device: 60 cm line × 10 with 10 needles......140.00 1 OP ✓ AutoSoft 90 9 mm teflon cannula: straight insertion: insertion device: 1 OP ✓ AutoSoft 90 9 mm teflon cannula; straight insertion; insertion device; 60 cm 1 OP ✓ AutoSoft 90 INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) - Special Authority see SA1985 on page 19 -Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 1 OP ✓ Quick-Set MMT-393 9 mm teflon cannula: straight insertion: 60 cm tubing × 10 with 1 OP ✓ Quick-Set MMT-392 (Quick-Set MMT-393 6 mm teflon cannula; straight insertion; 60 cm tubing x 10 with 10 needles; luer lock to be delisted 1 December 2023) (Quick-Set MMT-392 9 mm teflon cannula; straight insertion; 60 cm tubing x 10 with 10 needles; luer lock to be delisted 1

December 2023)

Fully

Brand or

	(Manufacturer's Price) \$	Sub Per	sidised •	Generic Manufacturer
INSULIN PUMP RESERVOIR - Special Authority see SA1985 on	page 19 – Retail ph	narmacy		
a) Maximum of 3 sets per prescription		-		
b) Only on a prescription				
c) Maximum of 13 packs of reservoir sets will be funded per y	ear.			
10 x luer lock conversion cartridges 1.8 ml for Paradigm pump	s50.00	1 OP	✓	ADR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 10	50.00	1 OP	✓ N	/liniMed 1.8 Reservoir MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP	✓ N	/liniMed 3.0 Reservoir MMT-332A

Subsidy

Digestives Including Enzymes

PANCREATIC	ENZYME
------------	--------

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase	
10,000 Ph Eur U, total protease 600 Ph Eur U)34.93	✓ <u>Creon 10000</u>
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase	
25,000 Ph Eur U, total protease 1,000 Ph Eur U)94.38 100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase	
3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph	
Eur U)34.93 20 g OP	Creon Micro
URSODEOXYCHOLIC ACID - Special Authority see SA1739 below - Retail pharmacy	
Cap 250 mg32.95 100	✓ <u>Ursosan</u>

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to

	Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic	
	\$	Per 🗸	Manufacturer	
continued				

allogenic stem cell or bone marrow transplantation; and

2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Laxatives

Bull	k-form	ing A	gents

ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln	6.00	250 g OP	✓ Macro Organic Psyllium Husk
	12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS * Dry	6.00	500 g OP	
* Uly	(17.32)	500 g OP	Normacol Plus
(Normacol Plus Dry to be delisted 1 October 2023)			

Faeca	C~4	
Faeca	- 201	ieners

DOCUSATE SODIUM – Only on a prescription			
* Tab 50 mg	2.31	100	✓ Coloxyl
* Tab 120 mg	3.13	100	✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES			
Tab 50 mg with sennosides 8 mg	3.50	200	✓ <u>Laxsol</u>
POLOXAMER – Only on a prescription			
Not funded for use in the ear.			
* Oral drops 10%	3.98	30 ml OP	✓ Coloxyl

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below – Retail ph	armacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

unless notified for applications meeting the following criteria:

Roth:

- 1 The patient is receiving palliative care; and
- 2 Fither:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Osmotic Laxatives

GLYCEROL * Suppos 2.8/4.0 g - Only on a prescription	10.39	20	✓ <u>Lax-suppositories</u> <u>Glycerol</u>
LACTULOSE - Only on a prescription			
* Oral liq 10 g per 15 ml	3.61	500 ml	✓ <u>Laevolac</u>
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM E Powder for oral soln 13.125 g with potassium chloride 46.6	mg,		_
sodium bicarbonate 178.5 mg and sodium chloride 350	0.7 mg 6.70	30	✓ <u>Molaxole</u>
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	✓ Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per m	, ,	escription	
5 ml	35.89	50	✓ <u>Micolette</u> ✓ Micolette-S29 S29

Stimulant Laxatives

Stimulant Laxatives			
BISACODYL - Only on a prescription			
* Tab 5 mg	5.80	200	 Bisacodyl Viatris
* Suppos 10 mg	3.69	10	✓ Lax-Suppositories
SENNA - Only on a prescription			
* Tab, standardised	2.17	100	
	(8.21)		Senokot
	0.43	20	
	(2.06)		Senokot
SODIUM PICOSULFATE - Special Authority see SA2053	below - Retail pharma	су	
Oral soln 7.5 mg per ml	7.40	30 ml OP	✓ Dulcolax SP Drop

⇒SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA - Special Authority see SA	1986 on the next page – Retail p	oharmacy	
Inj 50 mg vial	1,142.60	1	✓ Myozyme

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	1	Manufacturer	

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 below - Retail	ıı pnarmacy		
Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg	CBS	50	✓ Solgar
Powder	CBS	400 a	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE – Special Authority see SA1987 on the next page – Retail pharmacy		
Powder for oral soln575.00	180 g OP	Cystadane

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5.10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below -	- Retail pharmacy		
Cap 120 mg	CBS	30	✓ Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

```
GALSULFASE – Special Authority see SA1988 below – Retail pharmacy
Inj 1 mg per ml, 5 ml vial.......2,234.00 1 ✓ Naglazyme
```

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

DURSULFASE - Special Authority see SA1623 on the	e next page – Retail pharmacy		
Inj 2 mg per ml, 3 ml vial	4,608.30	1	Elaprase

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below - Retail pharmacy Tab 500 mgCBS 30 ✓ Solgar 30 ✓ Solgar 60 ✓ Balance ✓ Carnitor S29 Oral lig 1 g per 10 mlCBS 118 ml ✓ Novitium Sugar Free S29 Oral lig 500 mg per 10 mlCBS 300 ml Balance

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
RIBOFLAVIN – Special Authority see SA2041 below – Retail phar Tab 100 mg	•	100		Country Life Puritan's Pride Vitamin B-2 100 mg \$29
Cap 100 mg	CBS	100	1	Solgar

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see SA1989 below – Retail pharmacy Tab soluble 100 mg1,452.70 30 OP
✓ Kuvan

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Fither:

- 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
- 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer	
SODILIM PHENVI PLITYPATE Special Authority con SA1000 l	holow Dotail pharms	201/			

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retai	l pharmacy		
Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	✓ Life Extension

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and

Subsidy	Fu	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	•	Manufacturer

continued...

liver and spleen size; and

- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Mouth and Throat

Agents Used in Mouth Ulceration

DENIZVOAMINE HYDDOCHI ODIDE

BENZYDAMINE HYDROCHLORIDE			
Soln 0.15% - Higher subsidy of \$21.73 per 500 ml with			
Endorsement	9.00	500 ml	
	(21.73)		Difflam
Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly.	oral mucositis a	as a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	J	Orabase
	1.52	5 a OP	
	(3.60)	- 3 -	Orabase
Powder	, ,	28 g OP	
	(10.95)	- 3 -	Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE	, ,		
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
7 Adilosive gol 6.7 /6 With octamornam ornorido 6.61 /6	(6.00)	10 9 01	Bonjela
TRIANGINGI ONE ACETONIDE	(0.00)		Donjoid
TRIAMCINOLONE ACETONIDE Paste 0.1%	F 00	r = 0D	/ Kanalan in Orahaaa
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	E 06	20	✓ Fungilin
		20	• Fullgilli
MICONAZOLE			
Oral gel 20 mg per g	4.74	40 g OP	✓ <u>Decozol</u>
NYSTATIN			
Oral liq 100,000 u per ml	1.76	24 ml OP	✓ <u>Nilstat</u>

	(Manufacturer's Price)		sed Generic Manufacturer
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN * Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a Po	SO2.46	3	✓ Cobal-B12 \$299 ✓ Hydroxocobalamin Panpharma ✓ Vita-B12
PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose	4.10	5	✓ Cobalin-H 529
b) Only on a prescription * Tab 25 mg - No patient co-payment payable * Tab 50 mg		90 500	✓ <u>Vitamin B6 25</u> ✓ Pyridoxine multichem
THIAMINE HYDROCHLORIDE – Only on a prescription * Tab 50 mg VITAMIN B COMPLEX	4.65	100	✓ <u>Thiamine multichem</u>
* Tab, strong, BPC	11.25	500	✓ Bplex
Vitamin C			
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription * Tab 100 mg	12.50	500	✓ <u>Cvite</u>
Vitamin D			
ALFACALCIDOL * Cap 0.25 mcg * Cap 1 mcg		100 100	 ✓ One-Alpha ✓ One-Alpha ✓ One-Alpha S29 S29
* Oral drops 2 mcg per ml CALCITRIOL * Cap 0.25 mcg * Cap 0.5 mcg	7.89	0 ml OP 100 100	✓ One-Alpha ✓ Calcitriol-AFT ✓ Calcitriol-AFT
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescripi * Oral liq 188 mcg per ml (7,500 iu per ml)		12 .8 ml OP	✓ <u>Vit.D3</u> ✓ Puria
Multivitamin Preparations			
MULTIVITAMIN RENAL – Special Authority see SA1546 on the * Cap		harmacy 30	✓ Clinicians Renal Vit

Subsidy

Fully

Brand or

	ALIMENTAF	RY TRACT	AND	METABOLISM
	Subsidy (Manufacturer's Pric \$	e) Subs Per	Fully idised	Brand or Generic Manufacturer
■ SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals val the following criteria: Either: 1 The patient has chronic kidney disease and is receiving e 2 The patient has chronic kidney disease grade 5, defined	either peritoneal dial	ysis or haem	odialys	is; or
15 ml/min/1.73 m² body surface area (BSA). MULTIVITAMINS – Special Authority see SA1036 below – Reta * Powder	il pharmacy	200 a OP		aediatric Seravit
⇒SA1036 Special Authority for Subsidy		Ŭ		
Initial application from any relevant practitioner. Approvals valinborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid without approval for multivitamins.				·
VITAMINS * Tab (BPC cap strength) * Cap (fat soluble vitamins A, D, E, K) – Special Authority see		1,000	✓ <u>M</u>	<u>vite</u>
SA1720 below – Retail pharmacy		60	√ ∨	itabdeck
■ SA1720 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals val the following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; of 2 Patient is an infant or child with liver disease or short gut 3 Patient has severe malabsorption syndrome.	r	newal unless	notified	d for applications meeting
Minerals				
Calcium				
CALCIUM CARBONATE				
* Tab 1.25 g (500 mg elemental) * Tab eff 1.25 g (500 mg elemental) – Subsidy by endorsemental		250 100	_	alci-Tab 500 alcium 500 mg
Only when prescribed for patients unable to swallow call inappropriate and the prescription is endorsed according to the contract of the contr		lets or where	calciur	Hexal \$29 m carbonate tablets are
CALCIUM GLUCONATE * Inj 10%, 10 ml ampoule	32.00	10	✓ M	ax Health - Hameln S29
	64.00	20	✓ M	ax Health S29
lodine				
POTASSIUM IODATE	4.50	00	<i>a</i>	

Iron

FERROUS FUMARATE

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

^{*}Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)) (Fully Subsidised	
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	5.98	100	/	Ferro-F-Tabs
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental) * Oral liq 30 mg (6 mg elemental) per 1 ml		30 500 ml		<u>Ferrograd</u> Ferodan
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority sinj 50 mg per ml, 10 ml vial	ee SA1840 below – I	Retail p 1	,	Ferinject

⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient continues to have iron-deficiency anaemia; and

2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOS	ίĿ
-----------------	----

Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%33.60	355 ml	✓ Phillips Milk of Magnesia ©29
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule25.53	10	✓ Martindale

5

✓ Ferrosiq

ALIMENTARY TRACT AND METABOLISM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA1775 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	Binocrit
Inj 6,000 iu in 0.6 ml, syringe		6	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	Binocrit
Inj 10,000 iu in 1 ml, syringe		6	Binocrit
Inj 40,000 iu in 1 ml, syringe		1	Binocrit

Fully

Brand or

	(Manufacturer's Price) \$) Subs Per	sidised •	Generic Manufacturer
Megaloblastic				
FOLIC ACID * Tab 0.8 mg	26.60	1,000	√ F	olic Acid multichem
* Tab 5 mg	5.82	100		olic Acid Mylan olic Acid Viatris
Oral liq 50 mcg per ml	28.82 2	5 ml OP	_	iomed

Subsidy

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

rreaters Group in conjunction with the National Haer	noprilia ivianagement gro	up.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	✓ Alprolix
ELTROMBOPAG – Special Authority see SA1743 below Wastage claimable	– Retail pharmacy		
Tab 25 mg	1,550.00	28	✓ Revolade
Tab 50 mg	3,100.00	28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10.000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

Inj 30 mg in 1 ml vial	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial12,492.00	1	✓ Hemlibra
Inj 150 mg in 1 ml vial17,846.00	1	✓ Hemlibra

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Either:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more: and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Fither:

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	Subsidy		Fully	Brand or
\$ Per ✓ Manufacturer	(Manufacturer's Price		Subsidised	Generic
	\$	Per	✓	Manufacturer

continued...

- 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
- 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	· ·	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1	,315.00	1	✓ FEIBA NF
Inj 1,000 U	2	,630.00	1	✓ FEIBA NF
Inj 2,500 U	6	.575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	✓ Xyntha
Inj 1,000 iu prefilled syringe		1	✓ Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	Xyntha
Inj 3,000 iu prefilled syringe		1	✓ Xyntha

NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	1	✓ RIXUBIS
Inj 1,000 iu vial	1	✓ RIXUBIS
Inj 2,000 iu vial	1	✓ RIXUBIS
Inj 3,000 iu vial2,610.00	1	✓ RIXUBIS

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [Xpharm]

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

managed by the Haemophina Heaters Group in conjun	CHOIT WITH THE INCHAINT	acmopinii	i management di
Inj 250 iu vial	210.00	1	Advate
Inj 500 iu vial	420.00	1	Advate
Inj 1,000 iu vial	840.00	1	✓ Advate
Inj 1,500 iu vial	1,260.00	1	✓ Advate
Inj 2,000 iu vial	1,680.00	1	✓ Advate
Inj 3,000 iu vial	2,520.00	1	Advate

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per	✓	Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	FS) - [Xpharm]			
For patients with haemophilia. Rare Clinical Circumstances		e reco	mbinant f	actor VIII. Access to funded
treatment is managed by the Haemophilia Treaters Group in				
subject to criteria.				
Inj 250 iu vial	237.50	1	✓	Kogenate FS
Inj 500 iu vial	475.00	1	✓	Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial	1,900.00	1		Kogenate FS
Inj 3,000 iu vial	2,850.00	1	✓	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]	- [Xpharm]			
For patients with haemophilia A receiving prophylaxis treatme	ent. Access to funder	d trea	tment is m	nanaged by the Haemophilia
Treaters Group in conjunction with the National Haemophilia	Management group.			
Inj 250 iu vial	300.00	1	✓	Adynovate
Inj 500 iu vial	600.00	1		Adynovate
Inj 1,000 iu vial	,	1		Adynovate
Inj 2,000 iu vial	2,400.00	1	✓	Adynovate
SODIUM TETRADECYL SULPHATE				
* Inj 3% 2 ml	28.50	5		
	(73.00)			Fibro-vein
TRANEXAMIC ACID				
Tab 500 mg	10.45	60	1	Mercury Pharma
- 1.22 000 mg				
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO	8.00	5	1	Konakion MM
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO	9.21	5	✓	Konakion MM
, , ,				
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	14 95	990	1	Ethics Aspirin EC
CLOPIDOGREL	14.00	000	•	Euros Aspiriii Eo
* Tab 75 mg	E 07	84	./	Arrow - Clopid
· ·	5.07	04	•	Arrow - Ciopia
DIPYRIDAMOLE				.
* Tab long-acting 150 mg		60	•	Pytazen SR
TICAGRELOR - Special Authority see SA1955 below - Retail pl	,			
* Tab 90 mg	23.85	56	•	Ticagrelor Sandoz
⇒SA1955 Special Authority for Subsidy				
Initial application — (acute coronary syndrome) from any rele	evant practitioner. Ap	prova	als valid fo	r 12 months for applications
meeting the following criteria:				
Both:				

1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and

syndrome; and
2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12

continued...

months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Fither:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA2	152 below – Retail pharmacy		
Inj 20 mg in 0.2 ml syringe	31.28	10	Clexane
Inj 40 mg in 0.4 ml syringe	42.49	10	Clexane
Inj 60 mg in 0.6 ml syringe	60.67	10	Clexane
Inj 80 mg in 0.8 ml syringe	80.89	10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe		10	 Clexane Forte
Inj 150 mg in 1 ml syringe	143.86	10	Clexane Forte

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

	Subsidy		Fully	
(Ma	anufacturer's Price) \$	Sub Per	osidised •	Generic Manufacturer
HEPARIN SODIUM				
Inj 1,000 iu per ml, 5 ml ampoule	86.11	50	1	Pfizer
Inj 5,000 iu per ml, 5 ml vial		10		Heparin Sodium
., -,				Panpharma
Heparin Sodium Panpharma to be Principal Supply on 1 July	2023			
Inj 5,000 iu per ml, 1 ml		5	1	DBL Heparin
, 0,000 10 poi, 1	02.00	Ü		Sodium S29
	70.33		1	Hospira
Inj 5,000 iu per ml, 5 ml ampoule		50		Pfizer
Inj 25,000 iu per ml, 0.2 ml		5		Hospira
πη 20,000 ta μσι πι, σ.2 mi		J		•
	42.40			Heparin DBL S29
(Director 5 000 in course 5 and course 1 to 1 to 1 to 1 to 1 to 2000)	482.20	50	•	Heparin DBL S29
Pfizer Inj 5,000 iu per ml, 5 ml ampoule to be delisted 1 July 2023)				
HEPARINISED SALINE				
Inj 10 iu per ml, 5 ml	65.48	50	✓	Pfizer
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	76.36	60	✓	Pradaxa
Cap 110 mg		60	✓	Pradaxa
Cap 150 mg	76.36	60	✓	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83 10	30	1	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28		Xarelto
Tab 20 mg		28		Xarelto
-	77.00	20	•	, tur onto
VARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.	0.40	- 0		0
* Tab 1 mg		50		Coumadin
Ψ Tob 0 mm	6.46	100		Marevan
* Tab 2 mg		50		Coumadin
* Tab 5 mg		100		Marevan
★ Tab 5 mg		50	_	Coumadin
	11.48	100	•	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM - Special Authority see SA1259 below - Retail pharma			_	
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	/	<u>Nivestim</u>

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or

10

✓ Nivestim

2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or

Inj 480 mcg per 0.5 ml prefilled syringe.......148.58

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC $< 0.5 \times 10^9/L$).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

Inj 6 mg per 0.6 ml syringe - Brand switch fee payable

(Pharmacode 2657066) - see page 253 for details65.00

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

GLUCOSE	[DEXTROSE]
GLOCOCL	DEXILOUE

* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO		5	Biomed
* Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO	15.00	1	✓ Biomed
POTASSIUM CHLORIDE			
* Inj 75 mg per ml, 10 ml	65.00	50	Juno
SODIUM BICARBONATE			
Inj 8.4%, 50 ml2	22.40	1	✓ Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			
Inj 8.4%, 100 ml	22.95	1	✓ Biomed

a) Up to 5 inj available on a PSO

b) Not in combination

SODIUM CHLORIDE

Not funded for use as a nasal drop. Not funded for nebuliser use except when used in conjunction with an antibiotic intended for nebuliser use.

Inj 0.9%, bag - Up to 2000 ml available on a PSO	1.33	500 ml	✓ Baxter
	1.36	1 000 ml	✓ Baxter

Only if prescribed on a prescription for renal dialysis, maternity or post-natal care in the home of the patient, or on a PSO for emergency use. (500 ml and 1,000 ml packs)

Inj 23.4% (4 mmol/ml), 20 ml ampoule	35.50	5	Biomed
For Sodium chloride oral liquid formulation refer Standard	l Formulae, page	255	
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO	4.00	20	✓ Fresenius Kabi
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PSO	5.25	50	✓ Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	✓ Fresenius Kabi
AL PARENTERAL NUTRITION (TPN)			

OTAL	. PAREN	IERAL	NUI	KIII	ON (IPN)	
Infi	ucion						

Infusion	CBS	1 OP	✓ TPN
Intusion	BS	I OP	V IPI

Ziextenzo

Subsidy	Fully	Brand or
(Manufacturer's Pr	ice) Subsidised	Generic
\$	Per 🗸	Manufacturer

WATER

- On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or
- 2) On a bulk supply order; or
- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 10 ml ampoule – Up to 5 inj available on a PSO	7.19 7.60	50	✓ Pfizer✓ Multichem
Multichem to be Principal Supply on 1 September 2023 Inj 20 ml ampoule – Up to 5 inj available on a PSO	5.00	20	✓ Fresenius Kabi
(Pfizer Inj 10 ml ampoule to be delisted 1 September 2023)			

Oral Administration		
CALCIUM POLYSTYRENE SULPHONATE Powder169.85	300 g OP	✓ Calcium Resonium
COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO9.53	50	✓ Electral
•	50	Electral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)8.55	1,000 ml OP	✓ Pedialyte - Bubblegum
PHOSPHORUS		
Tab eff 500 mg (16 mmol)82.50	100	Phosphate Phebra
POTASSIUM CHLORIDE		
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)5.26 (17.10)	60	Chlorvescent
* Tab long-acting 600 mg (8 mmol)	200	✓ Span-K
SODIUM BICARBONATE		
Cap 840 mg8.52	100	✓ Sodibic
		✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE		
Powder84.65	454 g OP	✓ Resonium-A

_			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DOXAZOSIN			
* Tab 2 mg	17.35	500	 Doxazosin Clinect
* Tab 2 mg * Tab 4 mg	20.94	500	✓ Doxazosin Clinect
PHENOXYBENZAMINE HYDROCHLORIDE			
* Cap 10 mg	65.00	30	✓ BNM S29
	216.67	100	✓ Dibenzyline S29
PRAZOSIN			
* Tab 1 mg	5.53	100	✓ Arrotex-Prazosin S29 S29
* Tab 2 mg	7.00	100	✓ Arrotex-Prazosin S29 S29
* Tab 5 mg	11.70	100	✓ Arrotex-Prazosin S29 S29

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CA	РΤ	ΓO	ΡI	٦I	ı

*	Oral liq 5 mg per ml	94.99	95 ml OP	✓ Capoten
	Oral liquid restricted to children under 12 years of age.			

CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril. * Tah 0.5 mg

2 69

* 1ab 0.5 mg2.09 90	▼ Zapili
* Tab 2.5 mg5.79 90	✓ Zapril
Tab 5 mg10.05 90	✓ Zapril
ENALAPRIL MALEATE	
* Tab 5 mg1.75 90	✓ Acetec
Acetec to be Principal Supply on 1 September 2023	
* Tab 10 mg1.97 90	✓ Acetec
Acetec to be Principal Supply on 1 September 2023	
* Tab 20 mg2.35 90	✓ Acetec
Acetec to be Principal Supply on 1 September 2023	
LISINOPRIL	
* Tab 5 mg11.07 90	✓ Ethics Lisinopril
·	✓ Teva Lisinopril
* Tab 10 mg11.67 90	✓ Ethics Lisinopril
v	✓ Teva Lisinopril
* Tab 20 mg14.69 90	✓ Ethics Lisinopril
v	✓ Teva Lisinopril
PERINDOPRIL	
* Tab 2 mg	✓ Coversyl
* Tab 4 mg2.95 30	✓ Coversyl

✓ Coversyl

30

✓ 7anril

	CARDIOVASCULAR SYSTEM			
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
QUINAPRIL				
* Tab 5 mg	5.97	90	✓	Arrow-Quinapril 5
* Tab 10 mg	5.18	90	✓	Arrow-Quinapril 10
* Tab 20 mg	7.95	90	✓	Arrow-Quinapril 20
RAMIPRIL				
* Cap 1.25 mg	6.90	90	✓	Tryzan
* Cap 2.5 mg	6.60	90	1	<u>Tryzan</u>
* Cap 5 mg		90		<u>Tryzan</u>
* Cap 10 mg	7.05	90	•	<u>Tryzan</u>
ACE Inhibitors with Diuretics				
exists a record of prior dispensing of quinapril with hydro Tab 10 mg with hydrochlorothiazide 12.5 mgTab 20 mg with hydrochlorothiazide 12.5 mg	4.10	30 30		Accuretic 10 Accuretic 20
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
* Tab 4 mg	2.00	90		<u>Candestar</u>
* Tab 8 mg		90		<u>Candestar</u>
* Tab 16 mg		90		Candestar
* Tab 32 mg	5.26	90	•	<u>Candestar</u>
LOSARTAN POTASSIUM				
* Tab 12.5 mg		84		Losartan Actavis
* Tab 25 mg		84		Losartan Actavis
* Tab 50 mg		84		Losartan Actavis
* Tab 100 mg	3.50	84	•	Losartan Actavis
Angiotensin II Antagonists with Diuretics				
LOCADTANI DOTACCILIM MITH HVDDOCHI ODOTHIAZIDI	-			

LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE

* Tab 50 mg with hydrochlorothiazide 12.5 mg.......4.00

30

Arrow-Losartan & Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see \$	SA1905 below - Retail pharma	acy
Tab 24.3 mg with valsartan 25.7 mg	190.00 50	6 ✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00 50	6 Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00 5	6 Entresto 97/103

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:

	Subsidy	Fu	,	Brand or
(Manuf	facturer's Price)	Subsidis	ed	Generic
	\$	Per	/	Manufacturer

continued...

- 2.1 Patient is in NYHA/WHO functional class II; or
- 2.2 Patient is in NYHA/WHO functional class III; or
- 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 117

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaestnet	ics, Locai, <mark>page</mark>	117	
AMIODARONE HYDROCHLORIDE			
▲ Tab 100 mg	3.49	30	✓ Aratac
▲ Tab 200 mg	4.49	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a PSC)9.12	6	✓ Cordarone-X
	15.22	10	✓ Max Health
ATROPINE SULPHATE			
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a			
PSO	15.09	10	✓ Martindale
	10.00	10	· <u>martinaalo</u>
DIGOXIN	7.00	040	/ Lamauria DO
* Tab 62.5 mcg – Up to 30 tab available on a PSO		240	✓ <u>Lanoxin PG</u>
* Tab 250 mcg – Up to 30 tab available on a PSO		240	✓ <u>Lanoxin</u>
* Oral liq 50 mcg per ml	16.60	60 ml	✓ Lanoxin
			✓ Lanoxin Paediatric
			Elixir S29
			✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE			
▲ Cap 100 mg	20.05	84	✓ Rythmodan -
			Cheplafarm S29
	23.87	100	✓ Rythmodan
ELECAINIDE ACETATE	20.07	100	- Try annouan
FLECAINIDE ACETATE	10.05	00	/ Florestate DAM
▲ Tab 50 mg		60	✓ Flecainide BNM
▲ Cap long-acting 100 mg	35.78	90	✓ Flecainide
			Controlled
			Release Teva
Flecainide Controlled Release Teva to be Principal Supply or			4
▲ Cap long-acting 200 mg	54.28	90	✓ Flecainide
			Controlled
			Release Teva
Flecainide Controlled Release Teva to be Principal Supply or	•		_
Inj 10 mg per ml, 15 ml ampoule	104.00	5	✓ Tambocor
MEXILETINE HYDROCHLORIDE			
▲ Cap 150 mg	162.00	100	✓ Teva S29
▲ Cap 250 mg		100	✓ Teva S29
- Oup 200 mg		100	- 1010

(Ma	Subsidy anufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
ROPAFENONE HYDROCHLORIDE Tab 150 mg	40.90	50	✓ R	Rytmonorm
Antihypotensives				
AIDODRINE – Special Authority see SA1474 below – Retail pharma Tab 2.5 mg	•	100	✓ N	lidodrine Medsurge
Midadrina Madaurga to be Principal Cumply on 1 August 202	53.00		√ G	Gutron
Midodrine Medsurge to be Principal Supply on 1 August 202 Tab 5 mg		100	✓ N	lidodrine Medsurge
Midodrine Medsurge to be Principal Supply on 1 August 202	79.00 3		√ G	Gutron

(Gutron Tab 2.5 mg to be delisted 1 August 2023)

(Gutron Tab 5 mg to be delisted 1 August 2023)

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATENOLOL			
* Tab 50 mg	9.33	500	✓ Mylan Atenolol
			✓ Viatris
* Tab 100 mg	14.20	500	✓ Mylan Atenolol
* Oral liq 25 mg per 5 ml		300 ml OP	✓ Atenolol AFT
			S29 S29
	38.20		✓ Essential
			Generics S29
	49.85		✓ Atenolol AFT
Restricted to children under 12 years of age.			
(Mylan Atenolol Tab 50 mg to be delisted 1 November 2023)		
BISOPROLOL FUMARATE			
* Tab 2.5 mg	1.8/	90	✓ Bisoprolol Mylan
* Tab 2.5 Hig	1.04	90	✓ Bisoprolol Viatris
* Tab 5 mg	2.55	90	✓ Bisoprolol Mylan
* Tab 5 mg	2.00	90	✓ Bisoprolol Viatris
* Tab 10 mg	3 62	90	✓ Bisoprolol Mylan
* Tab To Tily		90	✓ Bisoprolol Viatris
(Bisoprolol Mylan Tab 2.5 mg to be delisted 1 November 20. (Bisoprolol Mylan Tab 5 mg to be delisted 1 November 2023	,		bisoproior viauris
CARVEDILOL			
* Tab 6.25 mg	2.24	60	✓ Carvedilol Sandoz
* Tab 12.5 mg		60	✓ Carvedilol Sandoz
* Tab 25 mg		60	✓ Carvedilol Sandoz
•			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
LARSTALOL	Ψ	1 61		Manuacturei
LABETALOL	44.50			
* Tab 100 mg		100	_	randate
* Tab 200 mg		100	✓ <u>T</u>	<u>randate</u>
* Inj 5 mg per ml, 20 ml ampoule		5	_	
	(88.60)		T	randate
* inj 5 mg per ml, 20 ml vial	42.29	1		
	(48.20)		Α	Alvogen S29
METOPROLOL SUCCINATE				
* Tab long-acting 23.75 mg	1.45	30	✓ B	Betaloc CR
* Tab long-acting 47.5 mg		30	✓ B	Betaloc CR
* Tab long-acting 95 mg		30	✓ B	Betaloc CR
* Tab long-acting 190 mg		30	✓ B	Betaloc CR
METOPROLOL TARTRATE				
* Tab 50 mg	5.66	100	✓	PCA-Metoprolol
* Tab 100 mg		60	_	PCA-Metoprolol
* Tab long-acting 200 mg		28	_	Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial		5		Metoprolol IV Mylan
, , , , , , , , , , , , , , , , , , , ,				Metoprolol IV Viatris
NADOLOL				
* Tab 40 mg	19.19	100	✓ N	ladolol BNM
* Tab 80 mg	30.39	100	✓ N	ladolol BNM
PROPRANOLOL			_	
* Tab 10 mg	7.04	100	✓ D	Profate
* Tab 40 mg		100	_	PCA-Propranolol
* Cap long-acting 160 mg		100	_	Cardinol LA
* Oral liq 4 mg per ml – Special Authority se				
Retail pharmacy		00 n	nl 🗸 R	Roxane-
··· F ·· ··· ·· · · · · · · · · · · · ·	•			Propranolol \$29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only): or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

*	Tab 80 mg37	7.50 5	i00 •	✓ Mylan
*	Tab 160 mg14	4.00 1	00	✓ Mylan

	Subsidy (Manufacturer's Price) \$	Full Subsidise Per •	•
Calcium Channel Blockers			
Dihydropyridine Calcium Channel Blockers			
AMLODIPINE	1.00	90	^r Vasorex
* Tab 2.5 mg * Tab 5 mg			Vasorex Vasorex
* Tab 3 mg			Vasorex
FELODIPINE		00 0	<u>rusorex</u>
* Tab long-acting 2.5 mg	1.45	30	Plendil ER
* Tab long-acting 2.5 mg			Felo 5 ER
* Tab long-acting 10 mg			Felo 10 ER
NIFEDIPINE		00	T CIO TO EIT
	10.00	FC .4	Tamainina MD10 000
* Tab long-acting 10 mg	18.80	56	Tensipine MR10 S29
* Tab long-acting 20 mg	9.12	50	Mylan (12 hr
1 1 22 10 19 20 11 g			release) \$29
	17.72	100	Nyefax Retard
* Tab long-acting 30 mg			Mylan Italy (24 hr
Tab long doing of mg		17	release) \$29
	34.10	100	Mylan (24 hr
	34.10	100	• '
W. Tab land action CO man	50.04	100	release) S29
* Tab long-acting 60 mg	52.81	100	Mylan (24 hr
			release) S29
Other Calcium Channel Blockers			
DILTIAZEM HYDROCHLORIDE			
* Cap long-acting 120 mg	65.35	500	Diltiazem CD Clinect
* Cap long-acting 180 mg			Cardizem CD
* Cap long-acting 240 mg		30	Cardizem CD
PERHEXILINE MALEATE			
* Tab 100 mg	62.90	100	Pexsig
VERAPAMIL HYDROCHLORIDE			- J
* Tab 40 mg	7 01	100	' Isoptin
* Tab 40 mg			' Isoptin
* Tab long-acting 120 mg			Isoptin Retard \$29
			' Isoptin SR
* Tab long-acting 240 mg	15.12		Isoptin SR
* Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a	- · -	-	r · ·
PSO	25.00	5	' Isoptin
			·
Centrally-Acting Agents			
CLONIDINE	10.04	4	/ Mulan
* Patch 2.5 mg, 100 mcg per day — Only on a prescription			Mylan Mylan
 Patch 5 mg, 200 mcg per day – Only on a prescription Patch 7.5 mg, 300 mcg per day – Only on a prescription 			Mylan Mylan
T aton 7.5 mg, 500 mag per day - Only on a prescription	10.33	→ ▼	<u>mytati</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
CLONIDINE HYDROCHLORIDE				
* Tab 25 mcg	29.32	112	✓	Clonidine Teva
* Tab 150 mcg	37.07	100	✓	Catapres
* Inj 150 mcg per ml, 1 ml ampoule		10	✓	Medsurge
METHYLDOPA				
* Tab 250 mg	15.10	100	✓	Methyldopa Mylan
•	52.85	500	•	Methyldopa Mylan
				S29 S29

Loop Diuretics

BUMETANIDE			
* Tab 1 mg	4.91	30	✓ Burinex S29 S29
	16.36	100	✓ Burinex
* Inj 500 mcg per ml, 4 ml vial	7.95	5	✓ Burinex
FUROSEMIDE [FRUSEMIDE]			
Tab 40 mg - Up to 30 tab available on a PSO	8.00	1,000	✓ IPCA-Frusemide
* Tab 500 mg		50	✓ Urex Forte
	89.48		✓ Furosemid-
			Ratiopharm S29
	169.96	100	✓ Furosemid-
			Ratiopharm S29
* Oral liq 10 mg per ml	11.20	30 ml OP	✓ Lasix
* Inj 10 mg per ml, 25 ml ampoule		6	✓ Lasix
* Ini 10 mg per ml 2 ml ampoule - Up to 5 ini available on		5	✓ Furosemide-Baxter

Potassium Sparing Diuretics

AMILORIDE HYDROCHLORIDE		
Oral liq 1 mg per ml32.10	25 ml OP	Biomed
EPLERENONE – Special Authority see SA1728 below – Retail pharmacy		
Tab 25 mg	30	✓ Inspra
Tab 50 mg25.00	30	✓ Inspra

⇒SA1728 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has heart failure with ejection fraction less than 40%; and
- 2 Either:
 - 2.1 Patient is intolerant to optimal dosing of spironolactone; or
 - 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

SPIRONOLACTONE

*	Tab 25 mg	3.68	100	✓ Spiractin
	Tab 100 mg		100	✓ Spiractin
	Oral lig 5 mg per ml	33.00	25 ml OP	✓ Biomed

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	
Potassium Sparing Combination Diuretics				
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE * Tab 5 mg with furosemide 40 mg AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZI		28	•	Frumil
* Tab 5 mg with hydrochlorothiazide 50 mg		50	•	Moduretic
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	1	Arrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerg * Tab 5 mg		500	•	Arrow- Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml	27.82 2	25 ml C	P 🗸	Biomed
CHLORTALIDONE [CHLORTHALIDONE] Tab 25 mg	6.95	50	1	<u>Hygroton</u>
INDAPAMIDE * Tab 2.5 mg	10.45 11.61	90 100		<u>Dapa-Tabs</u> Mylan Indapamide S29
(Mylan Indapamide 329 Tab 2.5 mg to be delisted 1 August 202 METOLAZONE	23)			
Tab 5 mg	CBS	1 50		Metolazone S29 Zaroxolyn S29
Vasopressin receptor antagonists				
TOLVAPTAN – Special Authority see SA2166 below – Retail pha Tab 15 mg Tab 30 mg Tab 45 mg + 15 mg Tab 60 mg + 30 mg Tab 90 mg + 30 mg	873.50 1,747.00 1,747.00	28 OF 28 OF 56 OF 56 OF 56 OF		Jinarc Jinarc Jinarc Jinarc Jinarc

⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and

- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation: and
- 3 Either:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents

Fibrates

DE.	ZAFIDRATE		
*	Tab 200 mg	90	✓ Bezalip
*	Tab long-acting 400 mg21.21	30	✓ Bezalip Retard

Other Lipid-Modifying Agents

V ∪ IDIM ∪ V			
	 חור	18 44	\sim

*	Cap 250 mg21.56	30	✓ Olbetam S29 S29
	25.44		✓ Olbetam

Resins

ΔΤΟΒΥΔΩΤΔΤΙΝΙ

COLESTIPOL HYDROCHLORIDE			
Grans for oral lig 5 g	32 89	30	✓ Colestid

HMG CoA Reductase Inhibitors (Statins)

ATOTIVASTATIN			
* Tab 10 mg	6.16	500	✓ Lorstat
* Tab 20 mg	9.24	500	✓ Lorstat
* Tab 40 mg		500	✓ Lorstat
* Tab 80 mg		500	✓ Lorstat
PRAVASTATIN			
* Tab 20 mg	2.11	28	✓ Pravastatin Mylan
•			✓ Pravastatin Viatris
* Tab 40 mg	3.61	28	✓ Pravastatin Mylan
ROSUVASTATIN - Special Authority see SA2093 below -	- Retail pharmacy		
* Tab 5 mg		30	✓ Rosuvastatin Viatris
N T-1-40		00	/ December 1 de Michele

*	Tab 10 mg
*	Tab 00 mg

* T	Tab 5 mg	1.70	30	✓	Rosuvastatin Viatris
	Tab 10 mg		30	1	Rosuvastatin Viatris
	Tab 20 mg		30	1	Rosuvastatin Viatris
	Tab 40 mg		30	1	Rosuvastatin Viatris

⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	er
--	----

continued...

- 1 Both:
 - 1.1 Patient is considered to be at risk of cardiovascular disease; and
 - 1.2 Patient is Māori or any Pacific ethnicity: or
 - 2 Both:
 - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
 - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atoryastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIMVASTATIN

*	Tab 10 mg	1.23	90	 Simvastatin Mylan
	Tab 20 mg		90	✓ Simvastatin Mylan
	Tab 40 mg		90	✓ Simvastatin Mylan
	•			✓ Simvastatin Viatris
*	Tab 80 mg	7.12	90	✓ Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

EZETIMIBE - Special Authority see SA1045 below - Retail pharmacy			
* Tab 10 mg1	.95	30	✓ Ezetimibe Sandoz

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 x normal) when treated with one statin; or
- 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
- 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA104	46 below – Retail _I	pharmacy	
Tab 10 mg with simvastatin 10 mg	5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg	7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg		30	✓ Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL T	RINITRATE

*	Oral pump spray, 400 mcg per dose - Up to 250 dose			
	available on a PSO	7.48	250 dose OP	✓ Nitrolingual Pump Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
*	Patch 50 mg, 10 mg per day	18.62	30	✓ Nitroderm TTS
ISC	SORBIDE MONONITRATE			
*	Tab 20 mg	19.55	100	✓ Ismo 20
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
	Tab long-acting 60 mg		90	✓ <u>Duride</u>

Sympathomimetics

ADDENALINE		
ADRENALINE		
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	Aspen Adrenaline
12.65		DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	Aspen Adrenaline

	C	ARDIO	OVAS	CULAR SYSTEM
	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Vasodilators				
HYDRALAZINE HYDROCHLORIDE				
* Tab 25 mg - Special Authority see SA1321 below - Retail				
pharmacy	CBS	1		Hydralazine
		56		Onelink S29
		84		AMDIPHARM \$29
		100		Onelink S29
* Inj 20 mg ampoule SA1321 Special Authority for Subsidy ** Inj 20 mg ampoule	25.90	5	•	Apresoline
 Initial application from any relevant practitioner. Approvals valid the following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitra inhibitors and/or angiotensin receptor blockers. 				
-				
MINOXIDIL A Tab 10 mg	78.40	100	1	Loniten
-	70.40	100	•	Lomen
NICORANDIL ▲ Tab 10 mg	25 57	60	1	Ikorel
▲ Tab 20 mg		60		lkorel
PAPAVERINE HYDROCHLORIDE				
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	1	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]				•
Tab 400 mg	42.26	50	1	Trental 400
•				
Endothelin Receptor Antagonists				
AMBRISENTAN – Special Authority see SA1702 below – Retail p	harmacy			
Tab 5 mg		30		Ambrisentan Mylan
Tab 10 mg	1 550 00	30		Ambrisentan Viatris Ambrisentan Viatris
Tab To mg	1,550.00	30		Mylan
SA1702 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertension Notes: Application details may be obtained from Pharmac's websi The Coordinator, PAH Panel		c.govt.n		•
Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: <u>PAH@pharmac.</u> ;	novt nz			
BOSENTAN - Special Authority see SA1991 on the next page - I	netail pharmacy		,	

✓ Bosentan Dr

✓ Bosentan Dr

Reddy's

Reddy's

60

60

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHAWHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised: or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL - Special Authority see SA1992 below - Retail pharmacy			
Tab 25 mg	85	1 🗸	Vedafil
Tab 50 mg	70 4	1 🗸	Vedafil
Tab 100 mg	20 1	2	Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III; or
 - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
 - 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 below - Retail pharmacy

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST – Special Authority see SA1705 below – Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml185.03 30 **✓ Vebulis**

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

ADAPALENE

- a) Maximum of 30 g per prescription

22.89	30 g OP	Differin
etail pharmacy	· ·	
11.26	60	Oratane
18.75	120	✓ Oratane
26.73	120	✓ Oratane
	etail pharmacy 11.26 18.75	etail pharmacy 60 6018.75 120

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
- 2 Patient is not of child bearing potential.

TRETINOIN

50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

# Crm 1%	8.56	10 g OP	✓ Crystaderm
MUPIROCIN			
Oint 2%	6.60	15 g OP	
	(11.50)		Bactroban

- a) Only on a prescription
- b) Not in combination

	Subsidy		,	and or
	(Manufacturer's F \$	Price) Subs Per		eneric anufacturer
SODILIM ELICIDATE (ELICIDIC ACID)	Ψ	1 01	- 1016	arialaciarei
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.59	5 g OP	✓ Foba	n
a) Maximum of 5 g per prescription		0 9 01	1000	<u></u>
b) Only on a prescription				
c) Not in combination				
Oint 2%	1.59	5 g OP	✓ Foba	<u>n</u>
a) Maximum of 5 g per prescription				
b) Only on a prescriptionc) Not in combination				
SULFADIAZINE SILVER				
Crm 1%	10.80	50 g OP	✓ Flam	azine
a) Up to 250 g available on a PSO		00 9 01	· · · · ·	uziii0
b) Not in combination				
Antifungala Tanical				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals,	, page 97			
AMOROLFINE				
a) Only on a prescription				
b) Not in combination Nail soln 5%	14.02	5 ml OP	√ Myos	Mail
	14.93	3 1111 OF	✓ Mycc	<u>nvan</u>
CLOTRIMAZOLE * Crm 1%	1 10	20 g OP	✓ Clom	1270
a) Only on a prescription	1.10	20 g Oi	Cion	iazoi
b) Not in combination				
* Soln 1%	4.36	20 ml OP		
	(7.55)		Cane	sten
a) Only on a prescription				
b) Not in combination				
ECONAZOLE NITRATE	4.00	00 00		
Crm 1%		20 g OP	Povo	nd
a) Only on a prescription	(7.78)		Peva	ıyı
b) Not in combination				
Foaming soln 1%, 10 ml sachets	9.89	3		
	(17.92)		Peva	ryl
a) Only on a prescription				
b) Not in combination				
MICONAZOLE NITRATE	0.04	45 00		
* Crm 2%	0.81	15 g OP	✓ <u>Multi</u>	<u>cnem</u>
a) Only on a prescriptionb) Not in combination				
* Lotn 2%	4.36	30 ml OP		
	(10.03)		Dakta	arin
a) Only on a prescription	` ,			
b) Not in combination				
* Tinct 2%		30 ml OP	5.1.	
a) Only on a green with the	(12.10)		Dakta	arın
a) Only on a prescriptionb) Not in combination				
b) Not in combination				

✓ Calamine-AFT

✓ Itch-Soothe

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per Manufacturer

100 g

20 g OP

Antipruritic Preparations

CALAN	١IN	ΙE
-------	-----	----

- a) Only on a prescription
- b) Not in combination
- **CROTAMITON**
 - a) Only on a prescription
 - b) Not in combination

MENTHOL - Only in combination

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 80

Corticosteroids - Plain

BETAMETHASONE DIPROPIONATE			<i>a</i> = .
Crm 0.05%		15 g OP	✓ Diprosone
	36.00	50 g OP	✓ <u>Diprosone</u>
Oint 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	✓ Diprosone
Oint 0.05% in propylene glycol base	4.33	30 g OP	✓ Diprosone OV
BETAMETHASONE VALERATE			
* Crm 0.1%	4.53	50 g OP	✓ Beta Cream
* Oint 0.1%	5.84	50 g OP	✓ Beta Ointment
★ Lotn 0.1%	.25.00	50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE			
★ Crm 0.05%	2.40	30 g OP	✓ Dermol
* Oint 0.05%		30 g OP	✓ Dermol
CLOBETASONE BUTYRATE		J	
Crm 0.05%	5.38	30 g OP	
	(10.00)	00 g 0.	Eumovate
HYDROCORTISONE	(10.00)		Lamovato
	1 70	00 = OD	/ Fabine
* Crm 1% – Only on a prescription		30 g OP	✓ Ethics
	17.15	500 g	✓ Hydrocortisone (PSM)
	20.40		✓ Noumed
Noumed to be Principal Supply on 1 August 2023			
★ Powder – Only in combination	40 05	25 g	✓ ABM
Up to 5% in a dermatological base (not proprietary Topical Co			
nalenicals	11003161100 -	- i iaiii) witti C	williout office definatologica

(Hydrocortisone (PSM) Crm 1% to be delisted 1 August 2023)

	Subsidy (Manufacturer's I \$	Price) Subs	Fully sidised	
YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN	<u> </u>	Per		Manufacturer
Loth 1% with paraffin liquid 15.9% and lanolin 0.6% — Only	, on			
a prescription		250 ml	1	DP Lotn HC
YDROCORTISONE BUTYRATE	10.07	200 1111	•	<u>DI LOUITIO</u>
Lipocream 0.1%	4.95	100 g OP	1	Locoid Lipocream
Oint 0.1%		100 g OF		Locoid
Milky emul 0.1%		100 g Ci		Locoid Crelo
IETHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.46	15 g OP	1	Advantan
Oint 0.1%		15 g OP		Advantan Advantan
IOMETASONE FUROATE		10 9 01	•	Auvantan
Crm 0.1%	1.05	15 g OP	1	Elocon Alcohol Free
CIIII 0.1 /6	3.10	50 g OP		Elocon Alcohol Free
Oint 0.1%		15 g OP		Elocon
5 V. /V	2.90	50 g OP		Elocon
Lotn 0.1%		30 ml OP		Elocon
RIAMCINOLONE ACETONIDE		-		
Crm 0.02%	6.30	100 g OP	1	Aristocort
Oint 0.02%		100 g OP	_	Aristocort
ETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FI Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49 (10.45) iption 1.89 Only on a prescri	15 g OP	•	Fucicort Micreme H Pimafucort
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 n				
and gramicidin 250 mcg per g - Only on a prescription	(9.28)	15 g OP		Viaderm KC
Barrier Creams and Emollients				
Barrier Creams				
Darrier Orcanio				
				haalib E
IMETHICONE Crm 5% pump bottle	4.30	500 ml OP	•	healthE Dimethicone 5%
IMETHICONE		500 ml OP 500 ml OP		Dimethicone 5% healthE
IMETHICONE € Crm 5% pump bottle				Dimethicone 5%
IMETHICONE € Crm 5% pump bottle € Crm 10% pump bottle	4.52	500 ml OP	✓	Dimethicone 5% healthE Dimethicone 10%
IMETHICONE € Crm 5% pump bottle	4.52		1	Dimethicone 5% healthE

	Subsidy (Manufacturer's \$	Price) Subs	Fully idised	Brand or Generic Manufacturer
Emollients	*			
AQUEOUS CREAM * Crm	1.73	500 g	√ <u>(</u>	GEM Aqueous Cream
CETOMACROGOL * Crm BP CETOMACROGOL WITH GLYCEROL	1.99	500 g	√ <u>(</u>	Cetomacrogol-AFT
Crm 90% with glycerol 10%	2.13 2.35	500 ml OP	_	Evara Pharmacy Health Sorbolene with Glycerin
Every to be Principal Supply on 1, July 2002	3.50	1,000 ml OP	✓ E	Evara
Evara to be Principal Supply on 1 July 2023 (Pharmacy Health Sorbolene with Glycerin Crm 90% with glyce	rol 10% to be del	listed 1 July 202	3)	
* Oint BP	3.40	500 g	✓ <u>E</u>	Emulsifying Ointment ADE
OIL IN WATER EMULSION * Crm	2.04	500 g	✓ <u>F</u>	Fatty Cream AFT
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50%	4.94	500 g OP	✓ <u>\</u>	White Soft Liquid Paraffin AFT
UREA * Crm 10%	1.37	100 g OP	√ ł	nealthE Urea Cream
WOOL FAT WITH MINERAL OIL − Only on a prescription ★ Lotn hydrous 3% with mineral oil	5.60	1,000 ml		
2 Con Hydrods 6 % With Hilliotal Oil	(14.96) (20.53) 1.40	250 ml OP		DP Lotion Alpha-Keri Lotion
	(5.87) 5.60 (23.91)	1,000 ml	-	DP Lotion BK Lotion
	1.40 (7.73)	250 ml OP	-	BK Lotion
Other Dermatological Bases				
PARAFFIN White soft - Only in combination Only in combination with a dermatological galenical or	19.99	450 g 2,500 g proprietary Top	✓ ł	nealthE nealthE rticosteroid – Plain.



Minor Skin Infections			
OVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ <u>Betadine</u>
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	4.15	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer
Devenitional Dynamoustians			
Parasiticidal Preparations			
DIMETHICONE			
₭ Lotn 4%	4.25	200 ml OP	✓ healthE

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic Manufacturer

Dimethicone 4% Lotion

✓ Stromectol

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- 3) For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA2228 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or

IVERMECTIN - Special Authority see SA2228 below - Retail pharmacy

- - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
- 2.2.1 The person is unable to complete topical therapy: or
- 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

2.2 Either:

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
- 2.2 Either:
 - 2.2.1 The person is unable to complete topical therapy: or
 - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5%5.75	30 g OP	Lyderm
Lotn 5%	30 ml OP	 A-Scabies

Psoriasis and Eczema Preparations

ACITRETIN – Special Authority see SA2024 below – Retail pharmacy			
Cap 10 mg17	7.86	60	✓ Novatretin
Cap 25 mg4	1.36	60	✓ Novatretin

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	39.35	60 g OP	✓ Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g		30 g OP	✓ Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR			
Soln BP - Only in combination	36.25	200 ml	✓ Midwest

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

	Subsidy (Manufacturer's P	rice) Su	Fully bsidised	Brand or Generic
	\$	Per	✓	Manufacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULI	PHUR			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% an	d			
allantoin crm 2.5%		75 g OP		
	(8.00)		l	Egopsoryl TA
	3.43	30 g OP		
	(4.35)			Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR	4.07	05 - 00		0 0
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97 7.95	25 g OP 40 g OP		Coco-Scalp Coco-Scalp
DIMEGRALIMINA O CLARIC CONTROL DE L		40 g OF	•	Coco-Scaip
PIMECROLIMUS – Special Authority see SA1970 below – Retain 1970 be	il pharmacy			
a) Maximum of 15 g per prescription	han ana nragarin	ion nor 10 w	ooko	
b) Note: a maximum of 15 g per prescription and no more t Cream 1%		15 g OP		Elidel
⇒SA1970 Special Authority for Subsidy	20.30	13 g O1	• !	Liluci
Initial application only from a dermatologist, paediatrician, ophtl	halmologist or an	rolovant pr	actitiona	r on the recommendation
of a dermatologist, paediatrician or ophthalmologist. Approvals v				
meeting the following criteria:	raliu Williout Iurili	ei ieliewai ui	11699 110	uneu ioi applications
HOID.				
1 Patient has atopic dermatitis on the eyelid; and	to topical corticos	steroids: per	iorificial	dermatitis, rosacea.
 Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications 				
1 Patient has atopic dermatitis on the eyelid; and				
 Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. 	pical corticostero	ds, cataracts	s, glauco	
2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to	pical corticostero	ds, cataracts	s, glauco	
Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.	pical corticostero	ds, cataracts	s, glauco	ma, or raised intraocula
Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID	pical corticostero SCEIN – Only or 14.44	ds, cataracts n a prescripti 500 ml	s, glauco on ✓ إ	ma, or raised intraocula
Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder — Only in combination	pical corticostero SCEIN - Only of	ds, cataracts a prescripti 500 ml 250 g	s, glauco on ✓ إ	ma, or raised intraocula Pinetarsol Midwest
Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID	pical corticostero SCEIN - Only of	ds, cataracts a prescripti 500 ml 250 g	s, glauco on ✓ إ	ma, or raised intraocula Pinetarsol Midwest
1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN - Only of	ds, cataracts a prescripti 500 ml 250 g	s, glauco on ✓ إ	ma, or raised intraocula Pinetarsol Midwest
1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN - Only of	ds, cataracts a prescripti 500 ml 250 g	s, glauco on ✓ إ	ma, or raised intraocula Pinetarsol Midwest
1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN – Only or n4.4418.88 proprietary Topic	ds, cataracts n a prescripti 500 ml 250 g al Corticoste	s, glauco	ma, or raised intraocula Pinetarsol Midwest
Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN - Only or 14.4418.88 proprietary Topic	ds, cataracts n a prescripti 500 ml 250 g al Corticoste	s, glaucconn on y j roid – P	ma, or raised intraocula Pinetarsol Midwest lain or collodion flexible Midwest
1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN - Only or 14.4418.88 proprietary Topic	ds, cataracts n a prescripti 500 ml 250 g al Corticoste	s, glaucconn on y j roid – P	ma, or raised intraocula Pinetarsol Midwest lain or collodion flexible Midwest
Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN - Only or 14.4418.88 proprietary Topic	ds, cataracts n a prescripti 500 ml 250 g al Corticoste	s, glaucconn on y j roid – P	ma, or raised intraocula Pinetarsol Midwest lain or collodion flexible Midwest
Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN - Only or 14.4418.88 proprietary Topic	ds, cataracts n a prescripti 500 ml 250 g al Corticoste	s, glaucconn on y j roid – P	ma, or raised intraocula Pinetarsol Midwest lain or collodion flexible Midwest
1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN - Only or 14.4418.88 proprietary Topic	ds, cataracts n a prescripti 500 ml 250 g al Corticoste	s, glaucconn on y j roid – P	ma, or raised intraocula Pinetarsol Midwest lain or collodion flexible Midwest
1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder — Only in combination	pical corticostero SCEIN - Only or 1	ds, cataracts a a prescripti 500 ml 250 g al Corticoste 100 g al Corticoste	e, glaucco on	ma, or raised intraocula Pinetarsol Midwest Iain or collodion flexible Midwest Iain
1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN - Only or 1	ds, cataracts n a prescripti 500 ml 250 g al Corticoste	e, glaucco on	ma, or raised intraocula Pinetarsol Midwest lain or collodion flexible Midwest
1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN — Only or 1	ds, cataracts a a prescripti 500 ml 250 g al Corticoste 100 g al Corticoste	on y roid - P	Pinetarsol Midwest Iain or collodion flexible Midwest Jain
1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination	pical corticostero SCEIN — Only or 1	ds, cataracts a a prescripti 500 ml 250 g al Corticoste 100 g al Corticoste	on y roid - P	Pinetarsol Midwest Iain or collodion flexible Midwest Jain

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

DERMATOLOGICALS

	Subsidy (Manufacturer's P			Brand or Generic	
	<u> </u>	Per		Manufacturer	
Scalp Preparations					
BETAMETHASONE VALERATE * Scalp app 0.1%	9.84	100 ml OP	✓ Be	eta Scalp	
CLOBETASOL PROPIONATE * Scalp app 0.05%	6.26	30 ml OP	✓ <u>De</u>	ermol	
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	6.57	100 ml OP	✓ <u>Lo</u>	ocoid	
KETOCONAZOLE					

a) Maximum of 100 ml per prescription

b) Only on a prescription

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

SPF 50+

Sebizole Sebizole

100 ml OP

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 69

PODOPHYLLOTOXIN

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

Other Skin Preparations

Antineoplastics

F	LU	IORO	URACIL	SODIUM

IMIQUIMOD

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

Contraceptives - Non-hormonal

Condoms

CONDOMS				_
	vailable on a PSO		144	✓ Moments
€ 53 mm			10	✓ Moments
		11.64	144	✓ Moments
 a) Maximum of 60 de 				
b) Up to 60 dev availa				_
53 mm, 0.05 mm thickness	S		10	✓ Moments
		11.42	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
53 mm, chocolate, brown.			10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
53 mm, strawberry, red			10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
56 mm			10	✓ Moments
		11.64	144	✓ Moments
 a) Maximum of 60 de 				
b) Up to 60 dev availa	able on a PSO			
56 mm, 0.05 mm thickness	S		12	✓ Gold Knight
		15.57	144	Gold Knight
 a) Up to 60 dev available 				
b) Maximum of 60 de				
	s (bulk pack)	14.61	144	Gold Knight
 a) Maximum of 60 de 				
b) Up to 60 dev available				
56 mm, 0.08 mm thickness	S	0.97	10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
56 mm, 0.08 mm thickness	s, red	0.97	10	✓ Moments
		11.64	144	✓ Moments
a) Up to 60 dev availa				
b) Maximum of 60 de	v per prescription			
56 mm, chocolate		1.30	12	Gold Knight
		15.57	144	Gold Knight
a) Up to 60 dev availa	able on a PSO			
b) Maximum of 60 de	v per prescription			
56 mm, strawberry		1.30	12	Gold Knight
•		15.57	144	✓ Gold Knight
a) Up to 60 dev availa	able on a PSO			
b) Maximum of 60 de				
		1.42	12	Gold Knight XL
		14.87	144	✓ Shield XL
		17.02		Gold Knight XL

		Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
		\$	Per	✓	Manufacturer
*	60 mm (bulk pack)	14.87	144	✓ (Gold Knight XL

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IUD 29.1 mm length × 23.2 mm width	29.80	1	✓ 7 MED NSHA Silver/
				Copper Short ✓ Choice 380 7med Nsha Silver/
				copper Short
				✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	29.80	1	✓ Choice
				TT380 Standard

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit: and

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- - 1 Patient is on a Social Welfare benefit; or
 - 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO 10.00 84 ✓ Mercilon 28

✓ Choice Load 375

✓ Brevinor 1/28

✓ Norimin

✓ Brevinor-1 28 Day
✓ Norimin-1 28 Day

84

112

84

112

16.33

29.32

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	_	Subsidised	
	\$	Per		Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	1.50	84	✓	Lo-Oralcon 20 ED
	2.18			Microgynon 20 ED
	6.45	112	✓	Femme-Tab ED
Lo-Oralcon 20 ED to be Principal Supply on 1 August 20				
* Tab 30 mcg with levonorgestrel 150 mcg		63		
	(16.50)			Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Autb) Up to 63 tab available on a PSO	nority see SA0500 or	the	previous pa	age
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	1.50	84	✓	Oralcon 30 ED
	1.77		1	Levlen ED
	6.45	112	✓	Femme-Tab ED
Oralcon 30 ED to be Principal Supply on 1 August 2023				
(Microgynon 20 ED Tab 20 mcg with levonorgestrel 100 mcg and (Femme-Tab ED Tab 20 mcg with levonorgestrel 100 mcg and 7 (Levlen ED Tab 30 mcg with levonorgestrel 150 mcg and 7 inert (Femme-Tab ED Tab 30 mcg with levonorgestrel 150 mcg and 7	inert tablets to be de ablets to be delisted	listed 1 Au	1 August i gust 2023)	2023)
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to	1			

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and

84 tab available on a PSO......12.25

to 84 tab available on a PSO......21.99

Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - Up

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully bsidised	Brand or Generic Manufacturer
continued				
 on a Social Welfare benefit; or have an income no greater than the benefit. 				
The approval numbers of Special Authorities approved before combined oral contraceptives and progestogen-only contracep LEVONORGESTREL				
* Tab 30 mcg – Up to 84 tab available on a PSO	16.50 22.00	84 112		Aicrolut Aicrolut
Subdermal implant (2 x 75 mg rods) – Up to 3 pack availation a PSO		1	√ J	ladelle
MEDROXYPROGESTERONE ACETATE		•	-	<u>uuoo</u>
Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a NORETHISTERONE	a PSO9.18	1	✓ [Depo-Provera
Tab 350 mcg – Up to 84 tab available on a PSO	12.25	84	√ <u> </u> <u> </u>	loriday 28
Emergency Contraceptives				
LEVONORGESTREL * Tab 1.5 mg	1.75	1	√ <u>L</u>	_evonorgestrel BNM
a) Maximum of 2 tab per prescriptionb) Up to 5 tab available on a PSOc) Note: Direct Provision by a pharmacist permitted	under the provisions in F	Part I of (Section <i>F</i>	
Antiandrogen Oral Contraceptives				
Prescribers may code prescriptions "contraceptive" (code "O") and prescription charge will be as per other contraceptives, as • \$5.00 prescription charge (patient co-payment) will apply • prescription may be written for up to six months supply.	follows:	for conti	raceptior	The period of supply
Prescriptions coded in any other way are subject to the non co of supply. ie. Prescriptions may be written for up to three mor		charges	, and the	non-contraceptive period
CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL * Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – to 168 tab available on a PSO		168	. (Ginet

Gynaecological Anti-infectives				
ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLI	EIC ACID			
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sul	phate			
0.025%, glycerol 5% and ricinoleic acid 0.75% with a	applicator8.43	100 g OP		
	(24.87)		Aci-Jel	
CLOTRIMAZOLE				
* Vaginal crm 1% with applicators	3.50	35 g OP	✓ Clomazol	
* Vaginal crm 2% with applicators	3.85	20 g OP	✓ Clomazol	
MICONAZOLE NITRATE				
* Vaginal crm 2% with applicator	6.89	40 g OP	✓ Micreme	
NYSTATIN				
Vaginal crm 100,000 u per 5 g with applicator(s)	4.00	75 g OP	✓ Nilstat	
5 , 1 - 9 (-)		J		

Brand or

Fully

	(Manufacturer's P	rice) Subsi Per	dised Generic Manufacturer
Myometrial and Vaginal Hormone Preparations			
ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO		5	✓ DBL Ergometrine
OESTRIOL			
* Crm 1 mg per g with applicator * Pessaries 500 mcg		15 g OP 15	✓ <u>Ovestin</u> ✓ Ovestin
OXYTOCIN - Up to 5 inj available on a PSO			<u></u>
Inj 5 iu per ml, 1 ml ampoule		5	✓ Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	5.98	5	✓ Oxytocin BNM ✓ Oxytocin GH S29
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj avai	lable on a PSO		,
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo		5	✓ Syntometrine
Pregnancy Tests - hCG Urine			
PREGNANCY TESTS - HCG URINE a) Up to 200 test available on a PSO b) Only on a PSO Cassette	12.00	40 test OP	✓ Smith BioMed Rapid
Casselle	12.00	40 lesi OF	Pregnancy Test
	16.00		✓ David One Step Cassette Pregnancy Test
Urinary Agents			
For urinary tract Infections refer to INFECTIONS, Antibacterials,	page 108		
5-Alpha Reductase Inhibitors			
FINASTERIDE – Special Authority see SA0928 below – Retail p * Tab 5 mg		100	✓ <u>Ricit</u>
⇒SA0928 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria:	d without further i	renewal unless	notified for applications meeting

Subsidy

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE – Special Authority see	SA1032 on the next pa	ge – Retail _I	pharmacy
* Cap 400 mcg	22.31	100	✓ <u>Tamsulosin-Rex</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price) Subsid \$ Per	Fully dised	Brand or Generic Manufacturer	
--	----------------	-------------------------------------	--

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

OYVBI ITVNIN

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

* Tab 5 mg	100	✓ Alchemy Oxybutynin S29
POTASSIUM CITRATE		
Oral liq 3 mmol per ml - Special Authority see SA1083 below -		
Retail pharmacy31.80	200 ml OP	✓ Biomed
CA1000 Chariel Avabagian for Cubaids		

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

bonontang nom the treatment.			
SODIUM CITRO-TARTRATE			
* Grans eff 4 g sachets	2.22	28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg	2.05	30	✓ Solifenacin Mylan
ů			✓ Solifenacin Viatris
Tab 10 mg	3.72	30	✓ Solifenacin Mylan
Ÿ			✓ Solifenacin Viatris
(California Mulan Tab Constable delisted & Dage			

(Solifenacin Mylan Tab 5 mg to be delisted 1 December 2023) (Solifenacin Mylan Tab 10 mg to be delisted 1 December 2023)

Detection	of Substan	nces in Urine
Detection	oi Substai	ices ili Ullile

ORTHO-TOLIDINE				
* Compound diagnostic sticks	7.50	50 test OP		
	(8.25)		Hemastix	
TETRABROMOPHENOL				
* Blue diagnostic strips	13.92	100 test OP	✓ Albustix	

Obstetric Preparations

Antiprogesterones

			IFEPRISTONE
Mifegyne	1	ailable on a PSO60.00	Tab 200 mg - Up to 15 tab available on a PSO
Mifegyne	3	180.00	

MI

Subsidy	Fully Subsidised		Brand or	
(Manufacturer's Price)			Generic	
\$	Per	✓	Manufacturer	
				_

Calcium Homeostasis

CALCITONIN			
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ Miacalcic
CINACALCET - Special Authority see SA2170 below -	Retail pharmacy		
Tab 30 mg - Wastage claimable	42.06	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable	84.12	28	✓ Cinacalet Devatis

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

	Subsidy		Fully	Brand or
(Mai	nufacturer's Price)	Subsidised		Generic
	\$	Per	1	Manufacturer

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

(Zoledronic acid Mylan Inj 4 mg per 5 ml, vial to be delisted 1 November 2023)

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

Corticosteroids and Related Agents for Systemic Use

DETAMETHASONE SOCIONITHOSITIATE WITH BETAMETHASONE ACE	IAIL	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml19.20		
(36.96))	Celestone
		Chronodose
DEXAMETHASONE		
* Tab 0.5 mg - Up to 60 tab available on a PSO1.50	30	<u>Dexmethsone</u>
* Tab 4 mg – Up to 30 tab available on a PSO2.65	30	<u>Dexmethsone</u>
Oral liq 1 mg per ml49.50	25 ml OP	Biomed
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO7.86	10	/ Hameln
* Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO 13.10	10	Hameln
FLUDROCORTISONE ACETATE		
* Tab 100 mcg	100	Florinef
HYDROCORTISONE		
* Tab 5 mg	100	Douglas
* Tab 20 mg		Douglas Douglas
* Inj 100 mg vial		Solu-Cortef
a) Up to 5 inj available on a PSO	·	<u> </u>
b) Only on a PSO		
METHYLPREDNISOLONE		
	100	Medrol
* Tab 4 mg		Medrol
* Tab 100 Hig223.10	20 ♥	Medioi

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully Brand or sidised Generic Manufacturer
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
lnj 40 mg vial	22.30	1	✓ Solu-Medrol-Act- O-Vial
Inj 125 mg vial	34.10	1	✓ Solu-Medrol-Act- O-Vial
Inj 500 mg vial	26.88	1	✓ Solu-Medrol-Act- O-Vial
Inj 1 g vial	32.84	1	✓ Solu-Medrol
METHYLPREDNISOLONE ACETATE		·	
Inj 40 mg per ml, 1 ml vial	47.06	5	✓ Depo-Medrol
	700	3	• Depo-medioi
PREDNISOLONE ★ Oral liq 5 mg per ml — Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	✓ <u>Redipred</u>
PREDNISONE			
₭ Tab 1 mg		500	Prednisone Clinect
★ Tab 2.5 mg		500	✓ Prednisone Clinect
Fab 5 mg - Up to 30 tab available on a PSO		500	✓ Prednisone Clinect
★ Tab 20 mg – Up to 30 tab available on a PSO	50.51	500	Prednisone Clinect
ETRACOSACTRIN			
★ Inj 250 mcg per ml, 1 ml ampoule	75.00	1	✓ Synacthen
			✓ UK Synacthen
★ Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓ Synacthen Depot ✓ Synacthene Retard S29
TRIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule	20.80	5	✓ Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule	51.10	5	✓ Kenacort-A 40
Sex Hormones Non Contraceptive			
Androgen Agonists and Antagonists			
CYPROTERONE ACETATE			
Tab 50 mg		50	✓ <u>Siterone</u>
Tab 100 mg	28.03	50	✓ <u>Siterone</u>
ESTOSTERONE			
Patch 5 mg per day	225.00	30	✓ Androderm
ESTOSTERONE CIPIONATE			
Inj 100 mg per ml, 10 ml vial	85.00 393.00	1	✓ Depo-Testosterone ✓ Taro- Testosterone \$29
			100100101010
ESTOSTERONE ESTERS			
Inj 250 mg per ml, 1 ml	12.98	1	✓ Sustanon Ampoules
		•	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
TESTOSTERONE UNDECANOATE				
Cap 40 mg - Subsidy by endorsement	21.00	60	1	Andriol Testocaps
	35.00	100	1	Steril-Gene S29
Subsidy by endorsement – subsidised for patients who w 1 November 2021 and the prescription is endorsed according where there exists a record of prior dispensing of testost Inj 250 mg per ml, 4 ml vial	rdingly. Pharmacists erone undecanoate c	may a	annotate t mg in the	the prescription as endorsed

Hormone Replacement Therapy - Systemic

Oestrogens

-	STRADIOL			
*	Tab 1 mg		28 OP	
	T. I. O.	(11.10)	00.00	Estrofem
*	Tab 2 mg		28 OP	Estratoria
	Datab 50 man nav 04 basses	(11.10)	4	Estrofem ✓ Climara
	Patch 50 mcg per 24 hours	7.04	4	Climara
	a) No more than 1 patch per week			
	b) Only on a prescription	6.10	8	✓ Estradot
	Patch 25 mcg per day	9.85	0	✓ Estradiol TDP
		9.00		
				Mylan S29
		13.50		✓ Estraderm MX S29
	a) No more than 2 patch per week			
	b) Only on a prescription	7.04	•	45
	Patch 50 mcg per day		8	✓ Estradot 50 mcg
		10.75		✓ Estradiol TDP
				Mylan S29
		14.50		✓ Estraderm MX S29
	a) No more than 2 patch per week			
	b) Only on a prescription			4=
	Patch 75 mcg per day		8	✓ Estradot
		11.88		✓ Estradiol TDP
				Mylan S29
	a) No more than 2 patch per week			
	b) Only on a prescription		_	
	Patch 100 mcg per day		8	✓ Estradot
		15.50		✓ Estraderm MX S29
	a) No more than 2 patch per week			
	b) Only on a prescription			
ΟE	STRADIOL VALERATE			
*	Tab 1 mg	12.36	84	✓ Progynova
*	Tab 2 mg	12.36	84	Progynova
OE	STROGENS			
*	Conjugated, equine tab 300 mcg	3.01	28	
		(17.50)		Premarin
*	Conjugated, equine tab 625 mcg		28	
		(17.50)		Premarin

	Subsidy		Fully Brand or
	(Manufacturer's Price \$) Sub Per	sidised Generic Manufacturer
rogestogens			
EDROXYPROGESTERONE ACETATE			
* Tab 2.5 mg	4.69	30	✓ Provera
· ·	8.75	56	✓ Provera
÷ Tab 5 mg	9.80	56	✓ Provera
	17.50	100	✓ Provera
Tab 10 mg	8.94	30	✓ Provera
Progestogen and Oestrogen Combined Prepara	ntions		
ESTRADIOL WITH NORETHISTERONE			
Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP	
	(18.10)		Kliovance
Tab 2 mg with 1 mg norethisterone acetate	5.40	28 OP	
	(18.10)		Kliogest
Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg			
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP	
-	(18.10)		Trisequens
Other Oestrogen Preparations			
ESTRIOL			
	7.00	30	✓ Ovestin
·			
Other Progestogen Preparations			
FVONODOCCTDCI			
EVONORGESTREL	000.50	4	/ Minama
Intra-uterine device 52 mg		1 1	✓ Mirena
f Intra-uterine device 13.5 mg	213.00	ı	Jaydess
EDROXYPROGESTERONE ACETATE			
Tab 100 mg	116.15	100	✓ Provera HD
ORETHISTERONE			
₹ Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Primolut N
ROGESTERONE			
Cap 100 mg	14.85	30	✓ <u>Utrogestan</u>
Thyroid and Antithyroid Agents			
ARBIMAZOLE	7.50	100	/ Non Marranala
F Tab 5 mg	/.56	100	✓ Neo-Mercazole
EVOTHYROXINE			
: Tab 25 mcg	5.55	90	Synthroid
Tab 50 mcg		28	✓ Mercury Pharma
	5.79	90	✓ Synthroid
	64.28	1,000	✓ Eltroxin
Tab 100 mcg		28	Mercury Pharma
	6.01	90	✓ Synthroid
	66.78	1,000	✓ Eltroxin
ROPYLTHIOURACIL – Special Authority see SA1199 on the n	ext page - Retail ph	armacy	
Tab 50 mg		100	✓ PTU S29

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub: Per	sidised ✓	Generic Manufacturer	

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

	MATRORIN (OMBITRORE) O CLAMIC II OACCOOL I	D . " .		
	MATROPIN (OMNITROPE) – Special Authority see SA2032 below		nacy	
*	Inj 5 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 15 mg cartridge	.139.50	1	✓ Omnitrope
				✓ Omnitrope S29 S29

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
 - 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
 - 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
 - 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
 - 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months

Subsidy	,		Brand or
(Manufacturer's Price)	Su Per	bsidised	Generic Manufacturer
-			

continued...

for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
- 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria;
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Fither:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater

Subsidy (Manufacturer's Price)	Subs Per	Fully sidised	Brand or Generic Manufacturer
 Ψ	1 01		I VIGITATION CONTRACTOR

continued...

than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or

2 All of the following:

- 2.1 The patient has been treated with somatropin for more than 12 months; and
- 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
- 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
- 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or

3 All of the following:

- 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
- 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3.3 The patient has severe growth hormone deficiency (see notes); and
- 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth

	Subsidy	F	ully	Brand or
(1)	Manufacturer's Price)	Subsidi	sed	Generic
	\$	Per	•	Manufacturer

continued...

hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GOSERELIN			
Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe	122.37	1	✓ Teva

LEUPRORELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

0 1 1			
Inj 3.75 mg prefilled dual chamber syringe - Higher su	bsidy of		
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher s	subsidy		

of \$591.68 per 1 inj with Endorsement.......177.50 (591.68)

Lucrin Depot 3-month

Vasopressin Agonists

DESMOPRESSIN Wafer 120 mcg	47.00	30	✓ Minirin Melt
DESMOPRESSIN ACETATE			
Tab 100 mcg	25.00	30	✓ Minirin
Tab 200 mcg	54.45	30	✓ Minirin
▲ Nasal spray 10 mcg per dose	27.95	6 ml OP	✓ <u>Desmopressin-</u> <u>PH&T</u>
Inj 4 mcg per ml, 1 ml	67.18	10	✓ Minirin

Other Endocrine Agents

CABERGOLINE

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
Dostinex	2	waived by Special Authority see SA2070 below4.43
✓ Dostinex	8	17.94

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

(Mani	Subsidy	Fully	Brand or
	ufacturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLO	MIF	FNF	CITI	RATE

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE Cap 250 mg	558.00	50	✓ <u>Metopirone</u>

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
Anthelmintics			
ALBENDAZOLE - Special Authority see SA1318 below - Retail	pharmacy		
Tab 400 mg	469.20	60	✓ Eskazole S29
■ SA1318 Special Authority for Subsidy Initial application only from an infectious disease specialist or cl patient has hydatids. Renewal only from an infectious disease specialist or clinical mic	robiologist. Approval		
remains appropriate and the patient is benefitting from the treatment	ent.		
MEBENDAZOLE – Only on a prescription Tab 100 mg	7.07	6	√ Vormov
Oral lig 100 mg per 5 ml		15 ml	✓ <u>Vermox</u>
014 nq 100 mg por 0 m	(7.83)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	✓ Biltricide
Antibacterials			
a) For topical antibacterials, refer to DERMATOLOGICALS, pag b) For anti-infective eye preparations, refer to SENSORY ORGA			
Cephalosporins and Cephamycins			
CEFACLOR MONOHYDRATE			
Cap 250 mg		100	✓ Ranbaxy-Cefaclor
Grans for oral liq 125 mg per 5 ml – Wastage claimable	3.75 1	00 ml	✓ Ranbaxy-Cefaclor
CEFALEXIN			
Cap 250 mg		20	✓ Cephalexin ABM
Cap 500 mgGrans for oral lig 25 mg per ml – Wastage claimable		20 00 ml	✓ <u>Cephalexin ABM</u> ✓ Flynn
Grans for oral lig 50 mg per ml — Wastage claimable		00 ml	
CEFAZOLIN – Subsidy by endorsement		00 1111	• <u>- 131111</u>
Only if prescribed for dialysis or cellulitis in accordance with a endorsed accordingly.	a Health NZ Hospital	appro	ved protocol and the prescription is
Inj 500 mg vial	3.39	5	✓ AFT
lnj 1 g vial		5	✓ <u>AFT</u>
CEFTRIAXONE – Subsidy by endorsement			
a) Up to 10 inj available on a PSO			
 Subsidised only if prescribed for a dialysis or cystic fibros pelvic inflammatory disease, or the treatment of suspecte 			

endorsed accordingly.

CEFUROXIME AXETIL - Subsidy by endorsement

(Zinnat Tab 250 mg to be delisted 1 March 2024)

Inj 1 g vial3.59

Tab 250 mg45.93

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly.

5

50

✓ Ceftriaxone-AFT

✓ Ceftriaxone-AFT

✓ Zinnat

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per	 Manufacturer 	

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable1	6.97	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

1 Atypical mycobacterial infection; or

EDVILIDOMYCINI (AC LACTODIONIATE)

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
 - 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial10.00 1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE	
Tab 400 mg16.95	E-Mycin
a) Up to 20 tab available on a PSO	
b) Up to 2 x the maximum PSO quantity for RFPP	4
Grans for oral liq 200 mg per 5 ml	E-Mycin
a) Up to 300 ml available on a PSO	
b) Up to 2 x the maximum PSO quantity for RFPPc) Wastage claimable	
Grans for oral liq 400 mg per 5 ml	✓ E-Mycin
a) Up to 200 ml available on a PSO	, .
b) Wastage claimable	
ROXITHROMYCIN	
Tab 150 mg13.19 50	✓ Arrow-
	Roxithromycin
Arrow-Roxithromycin to be Principal Supply on 1 August 2023	
Tab 300 mg25.00 50	✓ Arrow-
555	Roxithromycin

Arrow-Roxithromycin to be Principal Supply on 1 August 2023

	Subsidy		Fully	Brand or
	(Manufacturer's Pi	rice) Subs Per	idised	Generic Manufacturer
	\$	rei	<u> </u>	Manuacturei
Penicillins				
AMOXICILLIN				
Cap 250 mg	43.45	500	1	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	66.44	500	1	Alphamox
 a) Up to 30 cap available on a PSO 				
b) Up to 10 x the maximum PSO quantity for RFPP			_	
Grans for oral liq 125 mg per 5 ml	1.40	100 ml		Alphamox 125
 a) Up to 200 ml available on a PSO 				
b) Wastage claimable			_	
Grans for oral liq 250 mg per 5 ml	1.73	100 ml	/	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable	15.07	10	./	lhiamay
Inj 250 mg vialInj 500 mg vial		10 10		Ibiamox Ibiamox
Inj 1 g vial — Up to 5 inj available on a PSO		10		Ibiamox
	21.04	10	٠	IDIAIIIOX
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab	2.22	40		0 D 500//05
available on a PSO		10	•	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 i	•	100		A
per ml	0.50	100 ml	٧	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r per ml – Up to 200 ml available on a PSO	•	100 ml OP	./	Curam
·	2.20	100 IIII OF	•	Curain
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj	075 07	40	,	District A
available on a PSO	3/5.9/	10	•	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]			_	
Inj 600 mg (1 million units) vial – Up to 5 inj available on a P	SO 11.09	10		Sandoz
FLUCLOXACILLIN				
Cap 250 mg - Up to 30 cap available on a PSO		250		Flucloxacillin-AFT
Cap 500 mg - Up to 30 cap available on a PSO		500		Flucloxacillin-AFT
Grans for oral liq 25 mg per ml	3.29	100 ml	/	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable	0.00	400	,	A F-T
Grans for oral liq 50 mg per ml	3.68	100 ml	•	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable Inj 250 mg vial	17.56	10		Flucioxin
Inj 500 mg vial		10		Flucioxin
Inj 1 g vial — Up to 5 inj available on a PSO		5	_	Flucil
,		•	-	

	Subsidy (Manufacturer's Price)		Fully Subsidised	Generic	
	\$	Per		Manufacturer	
PHENOXYMETHYLPENICILLIN (PENICILLIN V)					
Cap 250 mg - Up to 30 cap available on a PSO	3.84	50	1	Cilicaine VK	
Cap 500 mg	6.86	50	✓	Cilicaine VK	
a) Up to 20 cap available on a PSO					
b) Up to 2 x the maximum PSO quantity for RFPP					
Grans for oral liq 125 mg per 5 ml	3.40	100 m	✓	<u>AFT</u>	
a) Up to 200 ml available on a PSO					
b) Wastage claimable					
Grans for oral liq 250 mg per 5 ml	4.24	100 m	✓	<u>AFT</u>	
a) Up to 300 ml available on a PSO					
b) Up to 2 x the maximum PSO quantity for RFPP					
c) Wastage claimable					

Tetracyclines

DC	XYCYCLINE			
*	Tab 100 mg - Up to 30 tab available on a PSO	.64.43	500	Doxine
MI	NOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	.19.32	100	
		(52.04)		Minomycin

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

 ${\sf TETRACYCLINE\ - Special\ Authority\ see\ SA1332\ below\ - \ Retail\ pharmacy}$

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 63

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	✓	Manufacturer
CLINDAMYCIN				
Cap hydrochloride 150 mg	5.30	24	1	Dalacin C
Inj 150 mg per ml, 4 ml ampoule	35.10	10	✓	Hameln
	39.00		✓	Dalacin C
Hameln to be Principal Supply on 1 August 2023 (Dalacin C Inj 150 mg per ml, 4 ml ampoule to be delisted 1 August 2023)	gust 2023)			
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist -	Subsidy by endorseme	ent		
Only if prescribed for dialysis or cystic fibrosis patient and t	he prescription is endo	rsed	accordingly	y.
Inj 150 mg	65.00	1	1	Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement	t95.00	5	1	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patien endorsed accordingly.		trac	t infection	and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	t91.00	5	1	Wockhardt S29
, ,	182.00	10	1	Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patien endorsed accordingly.	t or complicated urinary	trac	t infection	and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	t18.38	10	1	Pfizer
, ,	87.50	50	1	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patien endorsed accordingly.	t or complicated urinary	trac	t infection	and the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Reta No patient co-payment payable	ail pharmacy			
Tab 400 mg	42.00	5	1	Avelox

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

SODIUM FUSIDATE [FUSIDIC ACID]

SULFADIAZINE SODIUM - Special Authority see SA1331 below - Retail pharmacy

Tab 500 mg543.20 56 **✓ Wockhardt** 329

⇒SA1331 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

	Subsidy (Manufacturer's Price	e) Sub:	,	Brand or Generic
	\$	Per	1	Manufacturer
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	18.50	5	✓ <u>Tol</u> ✓ Via	bramycin Mylan tris
Only if prescribed for dialysis or cystic fibrosis patient ar	nd the prescription is	endorsed a	according	ly.
Solution for inhalation 60 mg per ml, 5 ml - Subsidy by				
endorsement	395.00	56 dose	✓ <u>Tol</u>	bramycin BNM
a) Wastage claimableb) Only if prescribed for a cystic fibrosis patient and the	prescription is endo	orsed accord	dinalv.	
TRIMETHOPRIM	, , , , , , , , , , , , , , , , , , ,		9.).	
* Tab 300 mg - Up to 30 tab available on a PSO	18.55	50	✓ <u>TM</u>	<u>P</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	(AZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg -				
to 30 tab available on a PSO		500	✓ <u>Tri</u>	sul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml - Up to 200				
available on a PSO	2.97	100 ml	✓ De	prim
VANCOMYCIN – Subsidy by endorsement				
Only if prescribed for a dialysis or cystic fibrosis patient or fo			for treatm	ent of Clostridium
difficile following metronidazole failure and the prescription is Inj 500 mg vial		gıy. 1	✓ Mv	lan
inj 500 mg viai	2.00	'	- <u>IVI y</u>	<u>iuii</u>

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 64
- b) For topical antifungals refer to GENITO URINARY, page 76

FLUCONAZOLE

Cap 50 mg	2.75	28	✓ Dizole✓ Mylan
Cap 150 mg	0.65	1	✓ Mylan
Cap 200 mg		28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority			
see SA1359 below – Retail pharmacy	109.34	35 ml	Diflucan
Wastage claimable			

(Dizole Cap 50 mg to be delisted 1 August 2023)

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE Cap 100 mg 4.27 15 ✓ Itrazole Oral liq 10 mg per ml – Special Authority see SA1322 below – 150 ml OP ✓ Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCT	CBS	30	✓ Burel S29
			✓ Link Healthcare S29
			✓ Nizoral S29
		100	✓ Strides Shasun S29
			✓ Taro S29
(Link Healthcare S29 Tab 200 mg to be delisted 1 July 20	123)		
(Nizoral S29 Tab 200 mg to be delisted 1 July 2023)			
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below	- Retail pharmacy		
Tab modified-release 100 mg	206.00	24	✓ Posaconazole Juno
Oral liq 40 mg per ml	342.51	105 ml OP	✓ Devatis

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Subsidy (Manufacturer's P	rice) S	Fully Subsidised	Brand or Generic	
<u> </u>	Per	1	Manufacturer	

continued...

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy: or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

* Tab 250 mg8.15	84	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 below - Retail pharmacy		
Tab 50 mg91.00	56	✓ Vttack
Tab 200 mg350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage		
claimable1,523.22	70 ml	✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Antitrichomonal Agents

Tab 200 mg − Up to 30 tab available on a PSO	METRONIDAZOLE			
Oral liq benzoate 200 mg per 5 ml	Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Antituberculotics and Antileprotics Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status. CLOFAZIMINE − Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. ** Cap 50 mg			100 ml	✓ Flagyl-S
Antituberculotics and Antileprotics Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status. CLOFAZIMINE − Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. * Cap 50 mg	•		10	✓ Flagyl
Antituberculotics and Antileprotics Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status. CLOFAZIMINE − Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. * Cap 50 mg	ORNIDAZOI F			
Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status. CLOFAZIMINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. ★ Cap 50 mg		36.16	10	✓ Arrow-Ornidazole
immigration status. CLOFAZIMINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. ★ Cap 50 mg	Antituberculotics and Antileprotics			
CLOFAZIMINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. ★ Cap 50 mg	, , , , ,	d in the Antitube	erculotics and	Antileprotics group regardless of
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. * Cap 50 mg	· ·			
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. * Cap 50 mg	. , ,			
** Cap 50 mg	b) Prescriptions must be written by, or on the recommendation	n of, an infection	us disease ph	ysician, clinical microbiologist or
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician. Cap 250 mg	•	442.00	100	✓ Lamprene S29
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician. Cap 250 mg	CYCLOSERINE - Betail pharmacy-Specialist			·
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician. Cap 250 mg				
Cap 250 mg	b) Prescriptions must be written by, or on the recommendation	n of, an infectiou	us disease ph	ysician, clinical microbiologist or
DAPSONE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist Tab 25 mg		344.00	60	✓ Cyclorin \$29
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist Tab 25 mg			00	- Cyclemia
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist Tab 25 mg				
dermatologist Tab 25 mg	, , , , , ,			
Tab 100 mg	dermatologist			
ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg	•			•
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg	Tab 100 mg	329.50	100	✓ Dapsone
 b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg	ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialist			
respiratory physician Tab 100 mg85.73 100 ✓ EMB Fatol 529	a) No patient co-payment payable			
Tab 100 mg		n of, an infectiou	us disease ph	ysician, clinical microbiologist or
Tab 400 mg	, ,,,	85.73	100	✓ EMB Fatol S29
	Tab 400 mg	49.34	56	✓ Myambutol S29

	Subsidy		Fully	
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
SONIAZID - Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommend		edicine	physician	, paediatrician, clinical
microbiologist, dermatologist or public health physician		400	,	DOM
* Tab 100 mg	23.00	100	•	<u>PSM</u>
SONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist				
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommend		edicine	physician	, paediatrician, clinical
microbiologist, dermatologist or public health physician			_	
* Tab 100 mg with rifampicin 150 mg		100		Rifinah Rifinah
* Tab 150 mg with rifampicin 300 mg		100	•	<u>niiinan</u>
PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist				
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommend	lation of an infactious	diaaaa	o opogialia	at alinical microbiologist or
respiratory physician	ation of, an infectious	JISCAS	e specialis	st, cililical microbiologist of
Grans for oral liq 4 g sachet	280.00	30	1	Paser S29
PROTIONAMIDE - Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommend	ation of, an infectious	diseas	e specialis	st, clinical microbiologist or
respiratory physician			•	•
Tab 250 mg	305.00	100	✓	Peteha S29
PYRAZINAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommend	lation of, an infectious	diseas	e physicia	n, clinical microbiologist or
respiratory physician	64.05	100	./	AET Duraninamida
* Tab 500 mg	64.95	100	•	AFT-Pyrazinamide
RIFABUTIN – Retail pharmacy-Specialist				
a) No patient co-payment payable		J:		
b) Prescriptions must be written by, or on the recommend gastroenterologist	ation of, an infectious	useas	e pnysicia	n, respiratory pnysician or
# Cap 150 mg	353.71	30	1	Mycobutin
RIFAMPICIN – Subsidy by endorsement		00	-	, ooba
a) No patient co-payment payable				
b) For confirmed recurrent Staphylococcus aureus infection	on in combination with	other e	effective a	nti-staphylococcal
antimicrobial based on susceptibilities and the prescrip				
Retail pharmacy - Specialist. Specialist must be an inte	ernal medicine physicia	an, clir	nical micro	biologist, dermatologist,
paediatrician, or public health physician.				
* Cap 150 mg		100		Rifadin Bifadin
* Cap 300 mg* Oral liq 100 mg per 5 ml		100 60 m		Rifadin Rifadin
Total liq 100 mg per 3 mil	12.00	00 111	•	niiauiii
Antivirals				
For eye preparations refer to Eye Preparations, Anti-Infective F	Preparations, page 248			
Hepatitis B Treatment				
ENTECAVIR				
	52.00	30		Entecavir Mylan

✓ Entecavir Sandoz

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully idised	Brand or Generic Manufacturer	
LAMIVUDINE - Special Authority see SA1685 below - Retail pha	armacy				Т
Tab 100 mg	6.95	28		<u>'etlam</u>	
Oral liq 5 mg per ml	270.00	240 ml OP	✓ Z	effix	

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA2139., page 105

*	Tab 245 mg (300 mg as a maleate)	15.00	30	✓ Tenofovir Disoproxil
				<u>Mylan</u>
				✓ Tenofovir Disoproxil

Herpesvirus Treatments

ACICLOVIR			
* Tab dispersible 200 mg	1.78	25	✓ Lovir
* Tab dispersible 400 mg	5.81	56	Lovir
* Tab dispersible 800 mg		35	Lovir
VALACICLOVIR			
Tab 500 mg	6.50	30	✓ Vaclovir
Tab 1,000 mg		30	✓ Vaclovir
VALGANCICLOVIR - Special Authority see SA1993 below - Re	tail pharmacy		
Tab 450 mg		60	✓ Valganciclovir
·			Mylan
			✓ Valganciclovir
			Viatris

⇒SA1993 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 Roth:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and

continued...

Viatris

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 ✓ Harvoni

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL — Subsidy by endorsement; can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 105 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

	minorman	on to available on the mannae webelle.	
*	Tab 200 mg	with tenofovir disoproxil 245 mg (300 mg as a	
	maleate)1:	5.45

30

✓ Tenofovir Disoproxil Emtricitabine Mylan

✓ <u>Tenofovir Disoproxil</u>
Emtricitabine Viatr

(Tenofovir Disoproxil Emtricitabine Mylan Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a maleate) to be delisted 1 November 2023)

⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical quidelines:

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical quidelines:

https://ashm.org.au/HIV/PrEP/

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — **(Confirmed HIV)** only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

- Both:
 - 1 Treatment course to be initiated within 72 hours post exposure; and
 - 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required: or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA2139 on the previous	s page – Retail phar	rmacy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA2139 on the previo	us page – Retail pha	armacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA2139 on the previo	us page – Retail pha	armacy	
Tab 200 mg	84.00	60	✓ Nevirapine
			<u>Alphapharm</u>
			Nevirapine Viatris
Oral suspension 10 mg per ml	203.55	240 ml OP	✓ Viramune
			Suspension

Nucleosides Reverse Transcriptase Inhibitors

Tab 300 mg	180.00	60	✓ Ziagen
Oral liq 20 mg per ml	256.31	240 ml OP	✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE - Special Author	ty see SA2139 on	the previous p	age - Retail pharmacy
a) Brand switch fee payable (Pharmacode 2655853) - see	page 253 for detail	ils	
b) Note: abacavir with lamivudine (combination tablets) or	ounts as two anti-re	etroviral medica	ations for the purposes of the
anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	29.50	30	✓ Abacavir/
			Lamivudine

ABACAVIR SULPHATE - Special Authority see SA2139 on the previous page - Retail pharmacy

Viatris

	Subsidy		Fully	y Brand or
	(Manufacturer's Price)		ubsidised	
	\$	Per		Manufacturer
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPRO	OXIL - Special Aut	hority s	ee <mark>SA2</mark> 1	139 on page 105 – Retail
pharmacy				
Note: Efavirenz with emtricitabine and tenofovir disoproxil cou	unts as three anti-re	etroviral	medicat	tions for the purposes of the
anti-retroviral Special Authority				
Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxi		00		Mulan
245 mg (300 mg as a maleate)	100.00	30		Mylan Viatris
(Mylan Tab 600 mg with emtricitabine 200 mg and tenofovir disopr	ovil 245 ma (200 m	a ac a		
(Mylan Tab 000 mg with emitticitabilite 200 mg and teriolovii disopi 2023)	0xii 245 mg (500 m	y as a i	naicaic)	i to be delisted i December
EMTRICITABINE – Special Authority see SA2139 on page 105 –	Rotail pharmacy			
Cap 200 mg		30	/	Emtriva
LAMIVUDINE - Special Authority see SA2139 on page 105 - Ret		00		
Tab 150 mg		60	/	Lamivudine
7ab 130 mg		00	•	Alphapharm
			1	Lamivudine Viatris
Oral lig 10 mg per ml	102.50 24	10 ml O		3TC
(Lamivudine Alphapharm Tab 150 mg to be delisted 1 November 2	2023)			
ZIDOVUDINE [AZT] - Special Authority see SA2139 on page 105	– Retail pharmacy			
Cap 100 mg		100	✓	Retrovir
Oral liq 10 mg per ml	30.45 20	00 ml 0	P 🗸	Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE - Special Authority see	SA2139 on page 10)5 – Re	tail phar	macy
Note: zidovudine [AZT] with lamivudine (combination tablets)	counts as two anti-	retrovira	al medic	ations for the purposes of
the anti-retroviral Special Authority.				
Tab 300 mg with lamivudine 150 mg	33.00	60	✓	Alphapharm
B., 1, 1, 9, 9,				
Protease Inhibitors				
ATAZANAVIR SULPHATE - Special Authority see SA2139 on pa	ge 105 – Retail pha	ırmacy		
Cap 150 mg	•	60	✓	Atazanavir Mylan
Cap 200 mg		60	✓	Atazanavir Mylan
DARUNAVIR - Special Authority see SA2139 on page 105 - Reta	ail pharmacy			
Tab 400 mg		60	✓	Darunavir Mylan
			✓	Darunavir Viatris
Tab 600 mg	196.65	60		Darunavir Mylan
(5)			•	Darunavir Viatris
(Darunavir Mylan Tab 600 mg to be delisted 1 August 2023)				
LOPINAVIR WITH RITONAVIR – Special Authority see SA2139 of		•		
Tab 100 mg with ritonavir 25 mg	150.00	60	•	Lopinavir/Ritonavir
Table 000 was with all an arise 50 was	005.00	400	,	Mylan
Tab 200 mg with ritonavir 50 mg	295.00	120	•	Lopinavir/Ritonavir
DITOMAND OF THE PARTY OF THE PA				<u>Mylan</u>
RITONAVIR – Special Authority see SA2139 on page 105 – Retain Tab. 100 mm.		00		Manudu
Tab 100 mg	43.31	30	•	Norvir
Strand Transfer Inhibitors				
DOLUTEGRAVIR - Special Authority see SA2139 on page 105 - Tab 50 mg		30	/	Tivicay
RALTEGRAVIR POTASSIUM - Special Authority see SA2139 on		oharma	CV	-
Tab 400 mg		60		Isentress
Tab 600 mg		60		Isentress HD

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Series Annufacturer

Immune Modulators

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 2 Patient has 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder, and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist.

Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive: or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Fither
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate: or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Initial application — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE

★ Tab 1 g19.95 100 ✓ <u>Hiprex</u>

INFECTIONS - AGENTS FOR SYSTEMIC USE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
NITROFURANTOIN				
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓	Nifuran
* Tab 100 mg	37.50	100	✓	Nifuran
* Cap modified-release 100 mg - Up to 15 cap available on a PSO	86.40	100	/	Macrobid
NORFLOXACIN				
Tab 400 mg - Subsidy by endorsement	245.00	100	✓	Arrow-Norfloxacin
Only if prescribed for a patient with an uncomplicated uri with proven resistance to first line agents and the prescri				ive to a first line agent or

✓ fully subsidised

Principal Supply

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per		Manufacturer
Anticholinesterases				
Antichonnesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	✓ M	lax Health
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	50.28	100	✓ M	lestinon
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENIAC CODILIM				
DICLOFENAC SODIUM * Tab EC 25 mg	1.00	50	√ n	iclofenac Sandoz
* Tab EC 25 flig * Tab 50 mg dispersible		20		oltaren D
* Tab EC 50 mg		50		iclofenac Sandoz
* Tab long-acting 75 mg		100	_	oltaren SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a F		5		oltaren
* Suppos 12.5 mg		10		oltaren
* Suppos 25 mg		10		oltaren
* Suppos 50 mg – Up to 10 supp available on a PSO		10		oltaren
* Suppos 100 mg		10	-	oltaren
		10	•	Oltaron
IBUPROFEN	01.40	1 000	/ D	allava
* Tab lang action 800 mg		1,000		elieve
* Tab long-acting 800 mg		30 200 m	_	rufen SR
* Oral liq 20 mg per ml	11.29	200 II	_	thics
	11.29			enpaed 100 mg per 5 ml
KETODDOFFN				J IIII
KETOPROFEN	10.07	00		
* Cap long-acting 200 mg	12.07	28	• 0	ruvail SR
MEFENAMIC ACID				
* Cap 250 mg	1.25	50		
	(10.82)		P	onstan
	0.50	20		
	(7.50)		P	onstan
NAPROXEN				
* Tab 250 mg	32.69	500		oflam 250
* Tab 500 mg	28.71	250	_	oflam 500
* Tab long-acting 750 mg	6.47	28	_	aprosyn SR 750
* Tab long-acting 1 g	8.62	28	✓ <u>N</u>	aprosyn SR 1000
TENOXICAM				
* Tab 20 mg	18.50	100	✓ Ti	ilcotil
* Inj 20 mg vial	9.95	1	✓ A	FT
NSAIDs Other				
CELECOXIB				
Cap 100 mg	3.45	60	√ ∩	elebrex
Oap 100 mg		00		elecoxib Pfizer
Cap 200 mg	3 20	30	_	elebrex
очр 200 mg		00	_	elecoxib Pfizer
			- 0	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	/	Manufacturer

Topical Products for Joint and Muscular Pain

CAPSAICIN

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary)*, and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroguine. Note: Indication marked with a * is an unapproved indication.

* Tab 200 mg			✓ Plaquenil
LEFLUNOMIDE			
Tab 10 mg	6.00	30	✓ Arava
Tab 20 mg	6.00	30	✓ Arava
PENICILLAMINE			
Tab 125 mg	67.23	100	D-Penamine
Tab 250 mg	110.12	100	D-Penamine

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODIUM		
* Tab 70 mg2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL		
* Tab 70 mg with colecalciferol 5,600 iu	4	✓ Fosamax Plus

Other Treatments

DENOSUMAB – Special Authority see SA1777 below – R	letail pharmacy		
Inj 60 mg prefilled syringe	326.00	1	Prolia

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial		1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial		1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA1	779 below – Retail	pharmacy	
* Tab 60 mg	53.76	28	✓ Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM			
Tab 35 mg	2.50	4	✓ Risedronate Sandoz
TERIPARATIDE – Special Authority see SA1139 on the next page	e – Retail pharma	су	
Inj 250 mcg per ml, 2.4 ml	490.00	1	✓ Forteo

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
\$	Per	√	Manufacturer

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, bag	22.53 100 ml OP	 Zoledronic Acid
		<u>Viatris</u>
		✓ Zoledronic-US S29

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg	28.57	500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA1963	below - Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite S29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	✓ Benzbromaron AL
			100 829

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg	6.00	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA2054 on the next page -	- Retail pharmacy		
Tab 80 mg	20.00	28	✓ <u>Febuxostat</u> multichem
Tab 120 mg	20.00	28	✓ Febuxostat multichem

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

⇒SA2054 Special Authority for Subsidy

Initial application — (**Gout**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or

2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout.. Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:
Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

PROBENECID

BACLOFFN

Muscle Relaxants

* Tab 10 mg	4.20	100	✓ Pacifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorseme	ent11.55	1	✓ Lioresal Intrathecal
Subsidised only for use in a programmable pump in pa caused intolerable side effects and the prescription is e			ents have been ineffective or have
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement.	306.82	5	✓ Medsurge
Subsidised only for use in a programmable pump in pa caused intolerable side effects and the prescription is of			ents have been ineffective or have
DANTROLENE			
Cap 25 mg	112.13	100	✓ Dantrium
			✓ Dantrium S29 S29
Cap 50 mg	77.00	100	✓ Dantrium
ORPHENADRINE CITRATE			
Tab 100 mg	20.76	100	✓ Norflex

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg	38.24	60	Symmetrel
	63.73	100	Symmetrel
APOMORPHINE HYDROCHLORIDE			
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	121.84	5	✓ Movapo
ENTACAPONE			
▲ Tab 200 mg	18.04	100	✓ Comtan
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg	15.80	100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100	Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
* Tab 100 mg with carbidopa 25 mg		100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg		100	✓ Sinemet CR
* Tab 250 mg with carbidopa 25 mg	38.39	100	✓ Sinemet
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg		100	✓ Ramipex
▲ Tab 1 mg	18.66	100	✓ Ramipex
RASAGILINE			
* Tab 1 mg	53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	4.05	84	✓ Ropin
▲ Tab 1 mg	4.95	84	✓ Ropin
▲ Tab 2 mg		84	✓ Ropin
▲ Tab 5 mg	14.50	84	✓ Ropin

SELEGILINE HYDROCHLORIDE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of selegiline hydrochloride.

* Tab 5 mg	.48.00	100	✓ Eldepryl S29
(Eldepryl S29) Tab 5 mg to be delisted 1 September 2023)			
TOLCAPONE			
▲ Tab 100 mg	152.38	100	✓ Tasmar

Anticholinergics

BENZATROPINE MESYLATE			
Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra

- a) Up to 10 inj available on a PSO
- b) Only on a PSO

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	•	Kemadrin
Agents for Essential Tremor, Chorea and Relate	ed Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharr Wastage claimable Tab 50 mg	•	56	/	Rilutek
Initial application only from a neurologist or respiratory specialist following criteria: All of the following: 1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vita 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow. Renewal from any relevant practitioner. Approvals valid for 18 m All of the following: 1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and	et. Approvals valid for duration of 5 years of al capacity within 2 m	r less onths	s; and prior to the	e initial application; and
3 Any of the following: 3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow. TETRABENAZINE Tab 25 mg	106 59	112	•	Motetis
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE]				

Gel 2%, tube - Subsidy by endorsement14.50 30 ml ✓ Xylocaine 2% Jelly

a) Up to 150 ml available on a PSO

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Gel 2%, 11 ml urethral syringe - Subsidy by endorsement......59.50 10 ✓ Instillagel Lido

a) Up to 5 each available on a PSO

b) Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price) Sub	sidised	Generic
	\$	Per		Manufacturer
IDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	44.00	200 ml	✓	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	9.50	25	✓	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	9.00	25	✓	Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.85	5	✓	Lidocaine-Baxter
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	7.15	5	✓	Lidocaine-Baxter
IDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes -				
Subsidy by endorsement		10	1	Pfizer
a) Up to 5 each available on a PSO				
b) Subsidised only if prescribed for urethral or cervical	administration and th	e prescript	ion is e	endorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] — Special Authority see SA0906	6 above – Retail pharm	acy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special A	Authority see SA0906	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

(Pfizer Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes to be delisted 1 November 2023)

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 111

Non-opi	oid Ana	Igesics
---------	---------	----------------

Tron opiola rinalgooloo		
ASPIRIN		
* Tab dispersible 300 mg - Up to 30 tab available on a PSO4.50	100	Ethics Aspirin
CAPSAICIN - Subsidy by endorsement		
Subsidised only if prescribed for post-herpetic neuralgia or diabetic periph accordingly.	neral neuropathy a	nd the prescription is endorsed
Crm 0.075%11.95	45 g OP	✓ Zostrix HP
15.14	57 g OP	Rugby Capsaicin
		Topical
		Cream S29
NEFOPAM HYDROCHLORIDE		
Tab 30 mg23.40	90	✓ Acupan

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or
	(Manufacturer's Pri	ce) Sub: Per	sidised Generic ✓ Manufacturer
	Ψ	rei	Manufacturer
PARACETAMOL	10.75	1 000	/ Pasimal
Tab 500 mg - blister pack		1,000	✓ <u>Pacimol</u>
 a) Maximum of 300 tab per prescription; can be w b) Up to 30 tab available on a PSO c) 	valved by endorsement		
Subsidy by endorsement for higher quan regular daily dosing for one month or gre annotate the prescription as endorsed wh Maximum of 100 tab per dispensing for n (for non-endorsed patients), then dispensed to the patients of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab per tab 500 mg - bottle pack — Maximum of 300 tab 500 mg - bot	ater, and the prescription nere dispensing history s non-endorsed patients. It	n is annotated upports a lor quantities p	d accordingly. Pharmacists may ng-term condition. rescribed for more than 100 tabs
prescription; can be waived by endorsement	17.92	1,000	✓ <u>Noumed</u> Paracetamol
 Subsidy by endorsement for higher quantitie daily dosing for one month or greater, and the prescription as endorsed where dispensing has aximum of 100 tab per dispensing for non- non-endorsed patients), then dispense in reg 	ne prescription is annotat nistory supports a long-te endorsed patients. If qu	ed according erm condition antities preso	rm conditions who require regular ly. Pharmacists may annotate the cribed for more than 100 tabs (for
Oral liq 120 mg per 5 ml	3.98	200 ml	✓ Paracetamol (Ethics)
	10.50	200 ml OP	✓ Avallon
a) Maximum of 600 ml per prescription; can be web) Up to 200 ml available on a PSO c) Not in combination d) 1) Maximum of 200 ml per dispensing for not approximately a per dispensing for not approximately a per dispensing for not approximately approxim	on-endorsed patients. If		
non-endorsed patients), then dispense in 2) Subsidy by endorsement for higher quan regular daily dosing for one month or gre Pharmacists may annotate the prescription condition.	tities is available for patic ater and the prescription on as endorsed where di	ents with long is endorsed spensing his	g term conditions who require or annotated accordingly. tory supports a long-term
Note: 200 ml presentations of paracetan provisions in Part I of Section A.	. , ,		
Oral liq 250 mg per 5 ml		200 ml	✓ <u>Pamol</u>
 Maximum of 200 ml per dispensing for no non-endorsed patients), then dispense in Subsidy by endorsement for higher quan regular daily dosing for one month or gre Pharmacists may annotate the prescriptic condition. Note: 200 ml presentations of paracetan 	repeat dispensing not e tities is available for patic ater and the prescription on as endorsed where di	xceeding 200 ents with long is endorsed spensing his	O ml per dispensing. g term conditions who require or annotated accordingly. tory supports a long-term
provisions in Part I of Section A.			
* Suppos 250 mg		10	✓ Gacet✓ Gacet
* Suppos 500 mg* Suppos 500 mg		10 50	✓ Gacet ✓ Gacet
	12.70	00	4400 .

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Sub Per	osidised •	Generic Manufacturer
	\$	rei	<u> </u>	Manuaciurei
Opioid Analgesics				
CODEINE PHOSPHATE - Safety medicine; prescriber may det	ermine dispensing fre	quency		
Tab 15 mg	5.92	100	_	<u>loumed</u>
Tab 30 mg	6.98	100		Aspen
T 00	40.00	400	_	loumed
Tab 60 mg	13.89	100	• 1	loumed
DIHYDROCODEINE TARTRATE				
Tab long-acting 60 mg	8.60	60	✓ [OHC Continus
FENTANYL				
 a) Only on a controlled drug form 				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr				
Inj 50 mcg per ml, 2 ml ampoule		10	_	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10	_	Boucher and Muir
Patch 25 mag per hour		5 5	_	entanyl Sandoz entanyl Sandoz
Patch 25 mcg per hourPatch 50 mcg per hour		5 5	_	entanyi Sandoz
Patch 75 mcg per hour		5	_	entanyl Sandoz
Patch 100 mcg per hour		5		entanyl Sandoz
METHADONE HYDROCHLORIDE		ŭ	-	omany. oanuo_
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr	equency			
d) Extemporaneously compounded methadone will only be		e of the c	heapest	form available
(methadone powder, not methadone tablets).				
e) For methadone hydrochloride oral liquid refer Standard F	Formulae, page 255			
Tab 5 mg	1.45	10	✓ N	Nethadone BNM
Oral liq 2 mg per ml		200 ml	_	<u> Biodone</u>
Oral liq 5 mg per ml		200 ml	_	Biodone Forte
Oral liq 10 mg per ml		200 ml	_	Biodone Extra Forte
Inj 10 mg per ml, 1 ml	68.90	10	✓ A	AFT
MORPHINE HYDROCHLORIDE				
 a) Only on a controlled drug form 				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr		200 - 1	, -	NA 84 II
Oral liq 1 mg per ml		200 ml	_	RA-Morph
Oral liq 2 mg per ml		200 ml		RA-Morph
Oral liq 5 mg per ml	19.44	200 ml		Ordine S29
Ovel lie 10 men men mi	07.74	200!		RA-Morph
Oral liq 10 mg per ml	27.74	200 ml	V (Ordine \$29

✓ RA-Morph

	Subsidy		Fully	
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
MORPHINE SULPHATE	· · · · · · · · · · · · · · · · · · ·			
a) Only on a controlled drug form b) No potient as powered payable.				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre		40		Carmadal
Tab immediate-release 10 mg		10		Sevredol Sevredol
Tab immediate-release 20 mg		10		Sevredol m-Esion
Cap long-acting 10 mg		10	_	
Cap long-acting 30 mg		10 10	_	m-Eslon m-Eslon
1 0 0 0		10	_	m-Esion
Cap long-acting 100 mgInj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		5	_	
		5 5	_	Medsurge Medsurge
Inj 10 mg per ml, 1 ml ampoule — Up to 5 inj available on a F		5	_	Medsurge Medsurge
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a F				Medsurge Medsurge
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	2506.28	5	•	<u>weasurge</u>
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre				
Tab controlled-release 5 mg	2.69	20		Oxycodone Sandoz
Tab controlled-release 10 mg	2.69	20		Oxycodone Sandoz
Tab controlled-release 20 mg		20		Oxycodone Sandoz
Tab controlled-release 40 mg	5.49	20		Oxycodone Sandoz
Tab controlled-release 80 mg		20		Oxycodone Sandoz
Cap immediate-release 5 mg		20		<u>OxyNorm</u>
Cap immediate-release 10 mg		20		<u>OxyNorm</u>
Cap immediate-release 20 mg	5.23	20		<u>OxyNorm</u>
Oral liq 5 mg per 5 ml		250 m		<u>OxyNorm</u>
Inj 10 mg per ml, 1 ml ampoule		5	✓	<u>Hameln</u>
Inj 10 mg per ml, 2 ml ampoule	11.49	5	✓	<u>Hameln</u>
Inj 50 mg per ml, 1 ml ampoule	22.92	5	✓	<u>Hameln</u>
PARACETAMOL WITH CODEINE - Safety medicine; prescriber	r mav determine disp	ensino	a frequenc	V
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		Paracetamol +
		,		Codeine (Relieve)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form b) No petient as payment payable.				
b) No patient co-payment payable	naunnau			
c) Safety medicine; prescriber may determine dispensing fre		10		PSM
Tab 50 mg		10		Noumed Pethidine
Noumad Pathiding to be Principal Cumply on 1 August 20	8.68		•	Noumea Petniaine
Noumed Pethidine to be Principal Supply on 1 August 2		_	./	DBL Pethidine
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	25029.88	5	•	
Int FO was a small O and a small of the La F to the small of the small of the La F to the small	200 00 70	_	,	Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a F	28030.72	5	•	DBL Pethidine
(DOM T 50				Hydrochloride
(PSM Tab 50 mg to be delisted 1 August 2023)				
TRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg	1.52	20	✓	Tramal SR 100
Tab sustained-release 150 mg	2.10	20	✓	Tramal SR 150
Tab sustained-release 200 mg		20	✓	Tramal SR 200
Cap 50 mg		100		Arrow-Tramadol
•				

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	/	Manufacturer

Antidepressants

Cyclic and Related Agents

AMITRIPTYLINE - Safety medicine; prescriber may dete	rmine dispensing frequent	СУ	
Tab 10 mg	2.49	100	✓ Arrow-Amitriptyline
Tab 25 mg	1.51	100	✓ Arrow-Amitriptyline
Tab 50 mg	2.51	100	✓ Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine	; prescriber may determine	e dispensing	g frequency
Tab 10 mg	10.17	30	✓ Clomipramine Teva
Tab 25 mg	11.99	30	✓ Clomipramine Teva
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsid	ly by endorsement		

- a) Safety medicine; prescriber may determine dispensing frequency
- b) Subsidy by endorsement Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 June 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.

Tab 75 mg	3.85	30
Cap 25 mg	.7.83	50
•		

- ✓ Dosulepin Viatris
- ✓ Dosulepin Mylan S29
- ✓ Dosulepin Viatris S29

IMIPRAMINE HYDROCHLORIDE - Safety medicine: prescriber may determine dispensing frequency

Tab 10 mg	5.48	50	✓ Tofranil
· ·	10.96	100	✓ Tofranil
Tab 25 mg	8.80	50	✓ Tofranil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicin	ne; prescriber may determir	ne dispensir	ng frequency
Tah 10 mg	2.46	100	✓ Nornress

NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; prescri	iber may determin	e dispensin	g frequency
Tab 10 mg	2.46	100	✓ Norpress
Tab 25 mg	6.29	180	✓ Norpress

Monoamine-Oxidase Inhibitors (MAOIs) - Non Selective

		ANTICTPROMINE SULPHATE	ΙĦ
Parnate	50	Tab 10 mg	*
✓ Parnate S29 S29	100	96.00	

(Parnate S29 S29 Tab 10 mg to be delisted 1 August 2023)

Monoamine-Oxidase Type A Inhibitors

TO A NIVE OVER COMMENT OF IT DELIVED

MO	CLOBEMIDE			
*	Tab 150 mg	.11.80	60	✓ Aurorix
*	Tab 300 mg	. 19.25	60	Aurorix

Selective Serotonin Reuptake Inhibitors

* Tab 20 mg	2.86	84	✓ Celapram
ESCITALOPRAM			4
* Tab 10 mg	1.07	28	Escitalopram (Ethics)

*	Tab 20 mg	28	(Ethics) ✓ Escitalopram (Ethics)

m

123

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

		21.11			
		Subsidy (Manufacturer's Price)		Fully Subsidised	
		(Manufacturer's Price)	Per		
FI !	JOXETINE HYDROCHLORIDE	<u> </u>			
	Tab dispersible 20 mg, scored — Subsidy by endorsement Subsidised by endorsement	2.50	28	•	Fluox
	 When prescribed for a patient who cannot swallow accordingly; or 				
	When prescribed in a daily dose that is not a multiple endorsed. Note: Tablets should be combined with	•			•
	Cap 20 mg	2 22	30	/	Brown & Burk S29
	Oup 20 mg	3.13	90		Arrow-Fluoxetine
РΔ	ROXETINE				
	Tab 20 mg	4.11	90	1	Loxamine
	RTRALINE		-	,	
-	Tab 50 mg	0.99	30	J	Setrona
~	140 00 mg	0.03	00		Setrona AU
*	Tab 100 mg	1.74	30		Setrona
					Setrona AU
(Se	etrona AU Tab 50 mg to be delisted 1 October 2023)				
	trona AU Tab 100 mg to be delisted 1 October 2023)				
0	ther Antidepressants				
MIF	RTAZAPINE				
	Tab 30 mg	2.60	28		Noumed
	Tab 45 mg	3.45	28	/	Noumed
۷E	NLAFAXINE				
*	Cap 37.5 mg	6.38	84		Enlafax XR
	Cap 75 mg		84		Enlafax XR
*	Cap 150 mg	11.16	84	/	Enlafax XR
Α	ntiepilepsy Drugs				
Α	gents for Control of Status Epilepticus				
DIA	ZEPAM - Safety medicine; prescriber may determine dispens	sing frequency			
	Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorsement	27.92	5	✓	Hospira
	a) Up to 5 inj available on a PSO				
	b) Only on a PSO	_			
	c) PSO must be endorsed "not for anaesthetic procedure		_		0. ".
	Rectal tubes 5 mg - Up to 5 tube available on a PSO	54.58	5	•	Stesolid
	ENYTOIN SODIUM				
*	Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a		_	_	
	PSO	104.58	5	/	Hospira
*	Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a	454.04	_	,	Haanina
	PSO	154.01	5	•	Hospira

	Subsidy		Fully	Brand or
	(Manufacturer's Price \$	Per Sub	osidised •	Generic Manufacturer
Control of Epilepsy				
CARBAMAZEPINE				
* Tab 200 mg	14.53	100	✓ T	egretol
* Tab long-acting 200 mg	16.98	100	√ T	egretol CR
	33.96	200	√ T	egretol CR
* Tab 400 mg	34.58	100	√ T	egretol
* Tab long-acting 400 mg	39.17	100	√ T	egretol CR
* Oral liq 20 mg per ml	26.37	250 ml	√ T	egretol
CLOBAZAM - Safety medicine; prescriber may determine dispe	ensing frequency			
Tab 10 mg		50	√ F	risium
CLONAZEPAM – Safety medicine; prescriber may determine d			-	
Oral drops 2.5 mg per ml		10 ml OP	./ D	ivotril
1 01	1.30	IU IIII UP	V N	iivotrii
ETHOSUXIMIDE				
Cap 250 mg	78.89	56	√ E	ssential
				Ethosuximide S29
	140.88	100	✓ Z	arontin
Oral lig 250 mg per 5 ml	56.35	200 ml	✓ Z	arontin
GABAPENTIN				
Note: Not subsidised in combination with subsidised prega	halin			
* Cap 100 mg		100	✓ N	lupentin
* Cap 300 mg		100	_	lupentin
* Cap 400 mg		100	_	lupentin
LACOSAMIDE – Special Authority see SA2223 below – Retail	•	4.4	./ V	'immat
▲ Tab 50 mg		14		impat
▲ Tab 100 mg		14		impat
A Tob 150 mg	200.24	56		impat
▲ Tab 150 mg		14		impat
A Tob 000 mg	300.40	56 56		impat
▲ Tab 200 mg	400.55	56	→ V	impat
⇒SA2223 Special Authority for Subsidy				

SA2223 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam, and any two of carbamazepine, lamotrigine, and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

LAMOTRIGINE

\blacktriangle	Tab dispersible 2 mg	55.00	30	✓ Lamictal
	Tab dispersible 5 mg		30	✓ Lamictal
	Tab dispersible 25 mg		56	✓ Logem
	Tab dispersible 50 mg		56	✓ Logem
	Tab dispersible 100 mg		56	✓ Logem

	Subsidy		Fully Brand or
	(Manufacturer's F		sidised Generic
	\$	Per	✓ Manufacturer
.EVETIRACETAM			
Tab 250 mg	5.84	60	✓ Everet
Tab 500 mg	10.51	60	✓ Everet
Tab 750 mg	16.71	60	✓ Everet
Tab 1,000 mg	21.82	60	✓ Everet
Oral liq 100 mg per ml	44.78	300 ml OP	✓ Levetiracetam-AFT
HENOBARBITONE			
For phenobarbitone oral liquid refer Standard F	ormulae, page 255		
← Tab 15 mg	40.00	500	✓ PSM
Fab 30 mg	40.00	500	✓ PSM
HENYTOIN SODIUM			
₹ Tab 50 mg	75.00	200	✓ Dilantin Infatab
Cap 30 mg		200	✓ Dilantin
Cap 100 mg		200	✓ Dilantin
Oral liq 30 mg per 5 ml		500 ml	✓ Dilantin
			✓ Dilantin Paediatric
PREGABALIN			
Note: Not subsidised in combination with subsi	idicad gahanantin		
Cap 25 mg	0 ,	56	✓ Pregabalin Pfizer
σαρ 20 mg	7.80	30	✓ Milpharm S29
€ Cap 75 mg		56	✓ Pregabalin Pfizer
e Cap 75 mg		30	•
0 150	8.10	50	✓ Milpharm \$29
Cap 150 mg	4.01	56	✓ Lyrica
			✓ Pregabalin Pfizer
	12.44		✓ Milpharm S29
Cap 300 mg	7.38	56	Pregabalin Pfizer
RIMIDONE			
← Tab 250 mg	37.35	100	Primidone Clinect
ODIUM VALPROATE			
Tab 100 mg	13.65	100	✓ Epilim Crushable
Tab 200 mg EC		100	✓ Epilim
Tab 500 mg EC		100	✓ Epilim
FOral lig 200 mg per 5 ml		300 ml	✓ Epilim S/F Liquid
1 3 F	• • • • • • • • • • • • • • • • • • • •		✓ Epilim Syrup
Inj 100 mg per ml, 4 ml	41.50	1	✓ Epilim IV
TIRIPENTOL – Special Authority see SA2217 bel			r
•	· · ·	60	/ Dissemit 520
Cap 250 mg		60	✓ Diacomit S29
Powder for oral liq 250 mg sachet	509.29	60	✓ Diacomit S29

⇒SA2217 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

	Cubaidu		F. III.	Drondor
(A	Subsidy Manufacturer's Price)		Fully Subsidised	Brand or Generic
(1)	s Frice)	Per	Jupaidised	Manufacturer
PIRAMATE				
Tab 25 mg	11.07	60	1	Arrow-Topiramate
1 4 5 1 1 9 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		•		Topiramate Actavis
	26.04			Topamax
Tab 50 mg		60		Arrow-Topiramate
.		•	_	Topiramate Actavis
	44.26			Topamax
Tab 100 mg		60		Arrow-Topiramate
		•		Topiramate Actavis
	75.25			Topamax
Tab 200 mg		60		Arrow-Topiramate
- 145 255 mg		00		Topiramate Actavis
	129.85			Topamax
Sprinkle cap 15 mg		60		Topamax
Sprinkle cap 25 mg		60	_	Topamax
		50	•	· · · · · · · · · · · · · · · · · · ·
GABATRIN – Special Authority see SA2088 below – Retail pharr		400	,	0-1
Tab 500 mg	119.30	100	•	Sabril

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Fither:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
 - 1.3 Patient has tuberous sclerosis complex: and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields..

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 111

Acute Migraine Treatment

RIZATRIPTAN

	Subsidy (Manufacturer's Prio \$	ce) Su Per	Fully ubsidised	Brand or Generic Manufacturer
SUMATRIPTAN				
Tab 50 mg	14.41	90	√ <u>S</u>	Sumagran
Tab 100 mg	22.68	90	√ <u>S</u>	Sumagran .
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 in	j per			
prescription	34.00	2 OP	✓ lı	migran
Prophylaxis of Migraine				
For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR	SYSTEM, page 51			
PIZOTIFEN	· -···· F-1-9 - 0 ·			
* Tab 500 mcg	23.21	100	√ S	Sandomigran
-				
Antinausea and Vertigo Agents				
For Antispasmodics refer to ALIMENTARY TRACT, page 8				
APREPITANT – Special Authority see SA0987 below – Retail	l pharmaou			
Cap 2 × 80 mg and 1 × 125 mg		3 OP	√ F	mend Tri-Pack
Oap 2 × 00 mg and 1 × 120 mg				
- CACCOT Consist Authority for Cubaidu			=	
⇒SA0987 Special Authority for Subsidy			_	adoracina hiahlu
nitial application from any relevant practitioner. Approvals v	valid for 12 months wh	nere the pa	tient is ur	ndergoing highly
nitial application from any relevant practitioner. Approvals we metogenic chemotherapy and/or anthracycline-based chemo	valid for 12 months whatherapy for the treatm	nere the pa	tient is ur	0 0 0 7
nitial application from any relevant practitioner. Approvals v emetogenic chemotherapy and/or anthracycline-based chemo Renewal from any relevant practitioner. Approvals valid for 1:	valid for 12 months whotherapy for the treatm 2 months where the p	nere the pa nent of mal patient is ur	tient is ur	0 0 0 7
nitial application from any relevant practitioner. Approvals v emetogenic chemotherapy and/or anthracycline-based chemo Renewal from any relevant practitioner. Approvals valid for 1: chemotherapy and/or anthracycline-based chemotherapy for t	valid for 12 months whotherapy for the treatm 2 months where the p	nere the pa nent of mal patient is ur	tient is ur	0 0 0 7
nitial application from any relevant practitioner. Approvals we emetogenic chemotherapy and/or anthracycline-based chemo Renewal from any relevant practitioner. Approvals valid for 1: shemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE	valid for 12 months whatherapy for the treatm 2 months where the p he treatment of malig	nere the pa nent of mal patient is ur nancy.	tient is ur ignancy. ndergoing	highly emetogenic
nitial application from any relevant practitioner. Approvals we metogenic chemotherapy and/or anthracycline-based chemo Renewal from any relevant practitioner. Approvals valid for 1: shemotherapy and/or anthracycline-based chemotherapy for the BETAHISTINE DIHYDROCHLORIDE Tab 16 mg	valid for 12 months whatherapy for the treatm 2 months where the p he treatment of malig	nere the pa nent of mal patient is ur	tient is ur	highly emetogenic
nitial application from any relevant practitioner. Approvals we metogenic chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline. Approvals valid for 1: shemotherapy and/or anthracycline-based chemotherapy for the SETAHISTINE DIHYDROCHLORIDE Tab 16 mg	valid for 12 months whatherapy for the treatm 2 months where the phe treatment of malig	nere the pa nent of mal patient is ur nancy.	tient is urignancy.	highly emetogenic
Initial application from any relevant practitioner. Approvals were togenic chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy from any relevant practitioner. Approvals valid for 1: shemotherapy and/or anthracycline-based chemotherapy for the set and th	valid for 12 months whatherapy for the treatm 2 months where the phe treatment of malig	nere the pa nent of mal patient is ur nancy.	tient is urignancy.	highly emetogenic
Initial application from any relevant practitioner. Approvals were togenic chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline. Approvals valid for 1: shemotherapy and/or anthracycline-based chemotherapy for the set of the s	valid for 12 months whatherapy for the treatment 2 months where the phe treatment of malig	nere the pa nent of mal patient is ur nancy.	tient is urignancy.	highly emetogenic
Initial application from any relevant practitioner. Approvals were togenic chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy for the themotherapy for the themotherapy and/or anthracycline-based chemotherapy for the themotherapy and/or anthracycline-based chemotherapy for the themotherapy for the themotherapy and/or anthracycline-based chemotherapy for the themotherapy for the themothe	valid for 12 months whatherapy for the treatment 2 months where the phe treatment of malig	nere the pa nent of mal patient is ur nancy. 100	tient is ur ignancy. ndergoing	highly emetogenic Serc lausicalm
Initial application from any relevant practitioner. Approvals were togenic chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline. Approvals valid for 1: shemotherapy and/or anthracycline-based chemotherapy for the set of the s	valid for 12 months whatherapy for the treatment 2 months where the phe treatment of malig	nere the pa nent of mal patient is ur nancy.	tient is ur ignancy. ndergoing	highly emetogenic
Initial application from any relevant practitioner. Approvals were togenic chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy for the themotherapy for the themotherapy and/or anthracycline-based chemotherapy for the themotherapy for the themothe	valid for 12 months whatherapy for the treatment 2 months where the phe treatment of malig4.62	nere the parent of mal patient is ur nancy. 100 10	tient is ur ignancy. idergoing S L	highly emetogenic Serc Jausicalm Jameln
Initial application from any relevant practitioner. Approvals of the application from any relevant practitioner. Approvals of the application from any relevant practitioner. Approvals valid for 12 shemotherapy and/or anthracycline-based chemotherapy for the application of the ap	valid for 12 months whatherapy for the treatment 2 months where the phe treatment of malig4.62	nere the pa nent of mal patient is ur nancy. 100	tient is ur ignancy. idergoing S L	highly emetogenic Serc Jausicalm Jameln Domperidone
nitial application from any relevant practitioner. Approvals we metogenic chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy for the motherapy and/or anthracycline-based chemotherapy for the motherapy	valid for 12 months whatherapy for the treatment 2 months where the phe treatment of malignum	nere the parent of mal patient is ur nancy. 100 10	tient is ur ignancy. idergoing S L	highly emetogenic Serc Jausicalm Jameln
Initial application from any relevant practitioner. Approvals were togenic chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy for the themotherapy for the themothe	valid for 12 months whatherapy for the treatment 2 months where the phe treatment of malignum	nere the parent of mal patient is ur nancy. 100 10	tient is ur ignancy. idergoing S L	highly emetogenic Serc Jausicalm Jameln Domperidone
nitial application from any relevant practitioner. Approvals we metogenic chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy and/or anthracycline-based chemotherapy for the motherapy and/or anthracycline-based chemotherapy for the motherapy	valid for 12 months whatherapy for the treatment 2 months where the phe treatment of malignum	nere the parent of mal patient is ur nancy. 100 10	tient is ur ignancy. Indergoing	highly emetogenic Serc Jausicalm Jameln Domperidone

⇒SA1998 Special Authority for Subsidy

Patch 1.5 mg - Special Authority see SA1998 below - Retail

pharmacy......17.70

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

✓ Scopoderm TTS

✓ Aripiprazole Sandoz

Aripiprazole \$29

✓ Aripiprazole Sandoz

✓ Aripiprazole Sandoz

✓ Aripiprazole Sandoz

✓ Aripiprazole Sandoz

✓ Ascend

✓ Largactil

✓ Largactil

✓ Largactil✓ Largactil

30

30

30

30

30

100

100

100

10

			IVL	NVOUS STSTEW
(M	Subsidy lanufacturer's Price) \$	Per	Fully Subsidised	Generic
METOCLOPRAMIDE HYDROCHLORIDE				
* Tab 10 mg - Up to 30 tab available on a PSO	1.30	100	/	Metoclopramide Actavis 10
★ Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO ONDANSETRON	7.00	10	✓	Baxter
* Tab 4 mg	2.27	50	1	Periset
1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	2.68			Onrex
Periset to be Principal Supply on 1 August 2023				
Tab disp 4 mg - Up to 10 tab available on a PSO	0.76	10	•	Ondansetron ODT-DRLA
* Tab 8 mg	4.10	50	1	Periset
v	4.57		✓	Onrex
Periset to be Principal Supply on 1 August 2023				
Tab disp 8 mg - Up to 10 tab available on a PSO	1.13	10	✓	Ondansetron ODT-DRLA
(Onrex Tab 4 mg to be delisted 1 August 2023) (Onrex Tab 8 mg to be delisted 1 August 2023)				
PROCHLORPERAZINE				
* Tab 3 mg buccal		50		_
# T. F	(30.00)	050	,	Buccastem
* Tab 5 mg – Up to 30 tab available on a PSO		250		Nausafix Ottoratii
* Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO	25.81	10		Stemetil
Antipsychotics				
General				
AMISULPRIDE – Safety medicine; prescriber may determine disper Tab 100 mg Tab 200 mg Tab 400 mg ARIPIPRAZOLE – Safety medicine; prescriber may determine disper	5.15 14.96 29.78	30 60 60	1	Sulprix Sulprix Sulprix
ducty medicine, precented may determine dispe			_	

A = 0 0 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.
—
*Three months or six months, as applicable, dispensed all-at-once
* Three months of six months, as applicable, dispensed all-ar-once

CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; prescriber may determine dispensing frequency

Tab 10 mg - Up to 30 tab available on a PSO......14.83

Tab 100 mg - Up to 30 tab available on a PSO.......36.73

	Subsidy	,	Fully	
	(Manufacturer's Price \$	e) Per	Subsidised	Generic Manufacturer
	Ψ	rei		iviariulacturei
CLOZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing frequency			_	
Tab 25 mg	6.69	50		Clopine
				Clozaril
	13.37	100		Clopine
				Clozaril
Tab 50 mg		50		Clopine
	17.33	100		Clopine
Tab 100 mg	17.33	50		Clopine
				Clozaril
	34.65	100		Clopine
				Clozaril
Tab 200 mg		50		Clopine
	69.30	100		Clopine
Suspension 50 mg per ml	67.62	100 m	· •	Versacloz
IALOPERIDOL - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 500 mcg - Up to 30 tab available on a PSO		100	1	Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO	9.43	100	1	Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50	1	Serenace
•	29.72	100	1	Serenace
Oral liq 2 mg per ml - Up to 200 ml available on a PSO	23.84	100 m	· •	Serenace
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS		10	1	Serenace
EVOMEPROMAZINE - Safety medicine; prescriber may determ		Vaneur		
Tab 25 mg (33.8 mg as a maleate)		100	1	Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan
Tab 100 mg (135 mg as a maleate)		100		Nozinan (Swiss)
Tab 100 mg as a maleate		100		Nozinan
•				
EVOMEPROMAZINE HYDROCHLORIDE - Safety medicine; p				
Inj 25 mg per ml, 1 ml ampoule	16.75	5	/	Neuraxpharm S29
			1	Nozinan S29 S29
	24.48	10	•	Wockhardt
ITHIUM CARBONATE - Safety medicine; prescriber may deter	mine dispensina fre	auency	,	
Tab long-acting 400 mg		100		Priadel
Cap 250 mg		100	_	Douglas
, •				J
DLANZAPINE - Safety medicine; prescriber may determine disp Tab 2.5 mg		20	./	Zunino
		28		Zypine Zypine
Tab 5 mg		28		Zypine ODT
Tab 10 mg		28	_	
Tab aradianaraible 10 mg		28	_	Zypine Zypine ODT
Tab orodispersible 10 mg		28	•	Zypine ODT
ERICYAZINE – Safety medicine; prescriber may determine dis			_	
Tab 2.5 mg		84		Neulactil
	12.49	100		Neulactil
Tab 10 mg		84		Neulactil
	44.45	100	1	Neulactil
QUETIAPINE - Safety medicine; prescriber may determine dispe	ensing frequency			
Tab 25 mg	. ,	90	1	Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90		Quetapel
1 45 500 mg	12.00	00	•	austupoi

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
RISPERIDONE – Safety medicine; prescriber may determin	ne dispensing frequency			
Tab 0.5 mg	1.86	60	✓ <u>F</u>	Risperidone (Teva)
Tab 1 mg		60	✓ F	Risperidone (Teva)
Tab 2 mg	2.29	60	✓ F	lisperidone (Teva)
Tab 3 mg		60	✓ F	lisperidone (Teva)
Tab 4 mg		60	✓ F	lisperidone (Teva)
Oral liq 1 mg per ml	8.90	30 ml	✓ F	lisperon
ZIPRASIDONE - Safety medicine; prescriber may determing	e dispensing frequency			
Cap 20 mg	17.90	60	✓ Z	usdone
Cap 40 mg		60	✓ Z	usdone
Cap 60 mg		60	✓ Z	usdone
Cap 80 mg		60	✓ Z	usdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine	; prescriber may determin	e disp	ensing freg	uency
Tab 10 mg		100	• •	Clopixol

Depot Injections

FLUPENTHIXOL DECANOATE - Safety medicine; prescriber m Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	13.14	ensing frequ 5 5 5	ency ✓ Fluanxol ✓ Fluanxol ✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber ma	av determine dispe	ensina freau	encv
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO		5 5	 ✓ Haldol ✓ Haldol Concentrate ✓ Haldol Decanoas (\$29)
OLANZAPINE – Special Authority see SA1428 below – Retail pl Safety medicine; prescriber may determine dispensing frequ	ency	4	/ Zummana Dalamann
Inj 210 mg vial		1	✓ Zyprexa Relprevv
Inj 300 mg vial		1	✓ Zyprexa Relprevv
Inj 405 mg vial	504.00	1	Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA1429 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing	frequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

Inj 175 mg syringe	815.85	1	✓ Invega Trinza
Inj 263 mg syringe	1,072.26	1	✓ Invega Trinza
Inj 350 mg syringe		1	✓ Invega Trinza
Inj 525 mg syringe		1	✓ Invega Trinza

⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO......19.80 5 **Clopixol**

Anxiolytics

3USPIR(ONE HY	'DROCHL	.oride
---------	--------	---------	--------

DO	SI INGINE ITI BITOGILEGITIBE				
*	Tab 5 mg	18.50	100	1	Buspirone Viatris
*	Tab 10 mg	12.50	100	✓	Buspirone Viatris

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CLONAZEPAM - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 500 mcg	5.64	100	✓	Paxam
Tab 2 mg	10.78	100	✓	Paxam
DIAZEPAM - Safety medicine; prescriber may determine disper	nsing frequency			
Tab 2 mg	61.07	500	✓.	Arrow-Diazepam
Tab 5 mg	73.60	500	✓.	Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency			•
Tab 1 mg	9.72	250	✓.	Ativan
Tab 2.5 mg	12.50	100	✓	Ativan

Multiple Sclerosis Treatments

⇒SA2176 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 2 Patients has an EDSS score between 0 6.0; and
- 3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 4 All of the following:
 - 4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 4.5 Either:
 - 4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 6 Any of the following:
 - 6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.



	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
DIMETHYL FUMARATE – Special Authority see SA2176 on the a) Wastage claimable b) Note: Treatment on two or more funded multiple scleros			rmitted.
Cap 120 mg Cap 240 mg	520.00	14	Tecfidera Tecfidera
FINGOLIMOD – Special Authority see SA2176 on the previous a) Wastage claimable b) Note: Treatment on two or more funded multiple scleros Cap 0.5 mg	sis treatments simultan	eously is not pe	rmitted. Gilenya
GLATIRAMER ACETATE – Special Authority see SA2176 on the Note: Treatment on two or more funded multiple sclerosis to Inj 40 mg prefilled syringe	reatments simultaneou	usly is not permit	ted. Copaxone
INTERFERON BETA-1-ALPHA – Special Authority see SA2176 Note: Treatment on two or more funded multiple sclerosis t Inj 6 million iu prefilled syringe Injection 6 million iu per 0.5 ml pen injector	reatments simultaneou 1,170.00	usly is not permit	
INTERFERON BETA-1-BETA — Special Authority see SA2176 Note: Treatment on two or more funded multiple sclerosis t Inj 8 million iu per 1 ml	reatments simultaneou	usly is not permit	
NATALIZUMAB – Special Authority see SA2176 on the previou Note: Treatment on two or more funded multiple sclerosis t Inj 20 mg per ml, 15 ml vial	reatments simultaneou	ısly is not permit	ted. Tysabri
OCRELIZUMAB – Special Authority see SA2176 on the previous Note: Treatment on two or more funded multiple sclerosis to Inj 30 mg per ml, 10 ml vial	reatments simultaneou	ısly is not permit	ted. Ocrevus
TERIFLUNOMIDE – Special Authority see SA2176 on the previ a) Wastage claimable b) Note: Treatment on two or more funded multiple scleros Tab 14 mg	sis treatments simultan	eously is not pe	rmitted. Aubagio
· · · · · · · · · · · · · · · · · ·			

Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy Tab modified-release 2 mg - No more than 5 tab per day 11.50 30 Vigisom Restricted to patients aged 18 years or under.

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the

				VOUS SYSTEM
	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
continued following criteria: All of the following:				
 Patient is aged 18 years or under*; and Patient has demonstrated clinically meaningful benefit fro Patient has had a trial of funded modified-release melator recurrence of persistent and distressing insomnia; and Funded modified-release melatonin is to be given at dose 	nin discontinuation wit	hin th	ne past 12 m	
Note: Indications marked with * are unapproved indications.				
MIDAZOLAM – Safety medicine; prescriber may determine disp Inj 1 mg per ml, 5 ml ampoule		10		lidazolam Mylan Iidazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available on a PSO		10	√ P	fizer
On a PSO for status epilepticus use only. PSO must be	e endorsed for status e	epilep	ticus use on	ly.
Inj 5 mg per ml, 3 ml ampouleInj 5 mg per ml, 3 ml plastic ampoule — Up to 5 inj available		5	✓ N	lidazolam-Baxter
a PSO		5	✓ P	fizer
On a PSO for status epilepticus use only. PSO must be (Midazolam Mylan Inj 1 mg per ml, 5 ml ampoule to be delisted to		epilep	ticus use on	ly.
PHENOBARBITONE SODIUM - Special Authority see SA1386	below - Retail pharma	acv		
Inj 200 mg per ml, 1 ml ampoule	•	10	✓ N	lax Health S29
■ SA1386 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Both: 1 For the treatment of terminal agitation that is unresponsive. The applicant is part of a multidisciplinary team working in the substitution of the substitution	re to other agents; and		nless notifie	d for applications meeting
TEMAZEPAM – Safety medicine; prescriber may determine disp Tab 10 mg		25	✓ N	lormison
TRIAZOLAM – Subsidy by endorsement			_	
a) Safety medicine; prescriber may determine dispensing fr b) Subsidised for patients who were taking triazolam prior to Pharmacists may annotate the prescription as endorsed preceding 12 months.	o 1 June 2023 and the			
Tab 125 mcg	5.10 (9.85)	100	Н	lypam
Tab 250 mcg	4.10 [°] (11.20)	100	Н	lypam
ZOPICLONE – Safety medicine; prescriber may determine dispersion 7.5 mg		E00	./7	anialana Astavia
Tab 7.5 mg	10.00	500	<u> </u>	opiclone Actavis
Spinal Muscular Atrophy				
NUSINERSEN – PCT only – Special Authority see SA2174 on the lini 12 mg per 5 ml vial		1	4.0	pinraza



Subsidy (Manufacturer's Price) \$ Per

Fully Subsidised Brand or Generic Manufacturer

⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml, 60 mg per bottle......14,100.00 80 ml OP ✓ Evrysdi

⇒SA2203 Special Authority for Subsidy

Initial application — (**spinal muscular atrophy (SMA)**) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Stimulants/ADHD Treatments				
ATOMOXETINE				
Cap 10 mg	18.41	28		APO-Atomoxetine APO-Atomoxetine S29 S29
			✓	Generic Partners
	107.03		✓	Strattera
Cap 18 mg	27.06	28	1	APO-Atomoxetine Generic Partners
	107.03			Strattera
Cap 25 mg	29.22	28		APO-Atomoxetine Generic Partners
Cap 40 mg	29.22	28	_	APO-Atomoxetine Generic Partners
	107.03		1	Strattera
Cap 60 mg	46.51	28	✓	APO-Atomoxetine APO-Atomoxetine S29 S29
				Generic Partners
Cap 80 mg	56.45	28	✓	APO-Atomoxetine APO-Atomoxetine S29 S29 Generic Partners
Cap 100 mg	58.48	28	<i>y</i>	APO-Atomoxetine APO-Atomoxetine S29 S29 Generic Partners
(Strattera Cap 10 mg to be delisted 1 November 2023) (Strattera Cap 18 mg to be delisted 1 November 2023) (Strattera Cap 40 mg to be delisted 1 November 2023)			·	denent rainers
DEXAMFETAMINE SULFATE – Special Authority see SA1149 b a) Only on a controlled drug form	oelow – Retail pharma	су		
b) Safety medicine; prescriber may determine dispensing fre	equency			
Tab 5 mg	' '	100	✓	PSM Aspen

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.



	Subsidy	F	ully	Brand or
(Ma	anufacturer's Price)	Subsidi	sed	Generic
	\$	Per	✓	Manufacturer

continued...

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine: prescriber may determine dispensing frequency

b) Salety medicine, prescriber may determine dispe	ensing frequency		
Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg	3.00	30	✓ Ritalin
•			✓ Rubifen
Tab extended-release 18 mg	7.75	30	Methylphenidate ER
•			- Teva
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg		30	✓ Rubifen SR
Tab extended-release 27 mg		30	Methylphenidate ER
•			- Teva
Tab extended-release 36 mg	15.50	30	✓ Methylphenidate ER
			- Teva
Tab extended-release 54 mg	22 25	30	✓ Methylphenidate ER
Tab extended release of mg		00	- Teva

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Subsidy		Fully	Brand or	
(Manufacturer's Pr	rice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Roth:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg		30	Concerta
Tab extended-release 36 mg		30	Concerta
Tab extended-release 54 mg		30	Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Fither:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or



	Subsidy	F	ully	Brand or
(Ma	anufacturer's Price)	Subsidi	sed	Generic
	\$	Per	✓	Manufacturer

continued...

4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see SA1999 below – Retail pharmacy

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below	- Retail pharmacy		
Patch 4.6 mg per 24 hour	38.00	30	 Rivastigmine Patch
			<u>BNM 5</u>
Patch 9.5 mg per 24 hour	38.00	30	 Rivastigmine Patch
			BNM 10

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✓ Manufacturer

28

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg11.76

Tab sublingual 8 mg with naloxone 2 mg34.00

✓ Buprenorphine
 Naloxone BNM
 ✓ Buprenorphine

Buprenorphine
Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone);
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg......11.00 30 ✓ Zyban

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
DISULFIRAM Tab 200 mg	236.40	100	√	Antabuse
NALTREXONE HYDROCHLORIDE – Special Authority see SA1 Tab 50 mg		harma 30		Naltraccord

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Patch 7 mg - Up to 28 patch available on a PSO	19.14	28	Habitrol
Patch 7 mg for direct distribution only - [Xpharm]	4.13	7	Habitrol
Patch 14 mg - Up to 28 patch available on a PSO	21.05	28	Habitrol
Patch 14 mg for direct distribution only - [Xpharm]	6.48	7	Habitrol
Patch 21 mg - Up to 28 patch available on a PSO	24.12	28	Habitrol
Patch 21 mg for direct distribution only - [Xpharm]	10.93	7	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO	19.76	216	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	3.35	36	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO		216	Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	3.40	36	Habitrol
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO	21.42	204	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]	9.04	96	Habitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO	21.42	204	Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]	9.04	96	Habitrol
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO	24.17	204	Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]	10.47	96	Habitrol
Gum 4 mg (Mint) - Up to 384 piece available on a PSO	24.17	204	Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]	10.47	96	Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg		56	✓ Varenicline Pfizer

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHI ORIDE - PCT only - Specialist - Special Authority see SA2153 below

Inj 25 mg vial	 	77.00	1	✓ Ribomustin
Inj 100 mg vial	 	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	 	3.23	1 mg	✓ Baxter

⇒SA2153 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6: and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

	Subsidy acturer's Price) Subs	Fully	Brand or Generic	
(wanua	\$ Per	uiseu •	Manufacturer	

continued...

- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has Hodgkin's lymphoma requiring treatment; and
 - 2 Patient has a ECOG performance status of 0-2; and
 - 3 Patient has received one prior line of chemotherapy; and
 - 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
 - 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

PLICITIEAN DCT Patail pharmany Chanielist

BUSULFAN - PCT - Retail pharmacy-Specialist	100	✓ Myleran
Tab 2 mg	100	♥ Wylcian
CARBOPLATIN – PCT only – Specialist		C DDL Oanhandatin
Inj 10 mg per ml, 45 ml vial	1	✓ DBL Carboplatin
45.20		✓ Carboplatin Ebewe
48.50	4	✓ Carbaccord
Inj 1 mg for ECP0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist		
Inj 100 mg vial710.00	1	✓ <u>BiCNU</u>
Inj 100 mg for ECP710.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist		
Tab 2 mg29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist		
Inj 1 mg per ml, 50 ml vial15.00	1	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial21.00	1	✓ Cisplatin Ebewe
29.66	'	✓ DBL Cisplatin
Inj 1 mg for ECP	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE	9	
	50	√ Cycloney
Tab 50 mg - PCT - Retail pharmacy-Specialist	1	✓ <u>Cyclonex</u> ✓ Endoxan
127.80	6	✓ Cytoxan
1-11-1	1	✓ Endoxan
Inj 2 g vial – PCT only – Specialist	1 mg	✓ Baxter
, ,	ring	Daxiei
IFOSFAMIDE – PCT only – Specialist		
lnj 1 g96.00	1	✓ Holoxan
lnj 2 g	. 1	✓ Holoxan
Inj 1 mg for ECP0.10	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price)	S	Fully Subsidised	I Generic
	\$	rei		ivianulacturei
LOMUSTINE – PCT – Retail pharmacy-Specialist			_	
Cap 10 mg		20		CeeNU
Cap 40 mg	399.15	20	/	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist	65.00	1	1	Melpha
, , ,	67.80		1	Alkeran
			1	Alkeran S29 S29
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
iij 100 iig viai	25.01	'	•	100
	110.00		./	Oxaliplatin Ebewe
Ini E ma nor ml. 00 ml viol		1		•
Inj 5 mg per ml, 20 ml vial				Alchemy Oxaliplatin
Ini dana far ECD	46.32	4		Oxaliplatin Accord
Inj 1 mg for ECP	0.35	1 mg	•	Baxter
THIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	✓	Bedford S29
			/	Max Health S29
			/	THIO-TEPA \$29
			1	Tepadina S29
Ini 100 ma vial	CBS	1		•
iij 100 iig viai		- 1		
Inj 100 mg vial	CBS	1		Max Health S29 Tepadina S29

Antimetabolites

ee SA2141 below	
75.06 1 🗸 <u>A</u>	zacitidine Dr
	Reddy's
0.83 1 mg 🗸 B	axter

⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully	Brand or
	(Manufacturer's Price	٠,	Subsidised	
	(Wallulacturer 5 Frice	Per	Jubsiuiseu	Manufacturer
	Ψ	rei		Manuacturer
CALCIUM FOLINATE				
Tab 15 mg - PCT - Retail pharmacy-Specialist	135 33	10	/	DBL Leucovorin
rab to mg . To thous pharmady openianos	100.00	10	•	Calcium
			_	
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	•	Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali	st7.28	1	✓	Calcium Folinate
, , , , , , , , , , , , , , , , , , , ,				Sandoz
			•	Calcium Folinate
				Sandoz S29 S29
	36.48	5	1	Eurofolic S29
Ini FO ma DCT Detail pharmagy Charielist		10		Leucovorin
Inj 50 mg - PCT - Retail pharmacy-Specialist	/ 2.80	10	•	
				Pharmacia S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	/	Calcium Folinate
ing roining portini, roinin vicii i roi only opposition ini		•	-	Sandoz
			_	
	47.45	5	•	Eurofolic S29
Inj 100 mg - PCT only - Specialist	7.33	1	✓	Calcium Folinate
, , , ,				Ebewe
	04.00	40	,	
	94.90	10	•	Leucovorin
				Pharmacia S29
Inj 300 mg - PCT only - Specialist	22 51	1	1	Calcium Folinate
injood ing ToT drily openialist			•	Ebewe
			_	
	25.14		•	Leucovorin DBL §29
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	✓	Calcium Folinate
, , , , , , , , , , , , , , , , , , , ,				Sandoz
			,	
			•	Calcium Folinate
				Sandoz S29 S29
Inj 1 g - PCT only - Specialist	67 51	1	1	Calcium Folinate
ing i g i or only opecialist			•	Ebewe
			_	
Inj 10 mg per ml, 100 ml vial – PCT only – Specialist	72.00	1	✓	Calcium Folinate
				Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	1	Baxter
	0.00	ring	•	Daxter
CAPECITABINE – Retail pharmacy-Specialist				
Tab 150 mg	10.00	60	✓	Capercit
Tab 500 mg		120		Capecitabine-
Tab 300 mg		120	•	•
				DRLA S29
			_	
			•	Capercit
CLADRIBINE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml	749.96	1		Litak S29
Inj 1 mg per ml, 10 ml	749.96	1	✓	Leustatin
Inj 10 mg for ECP	749.96	10 mg C)P 🗸	Baxter
,				
CYTARABINE				
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali	st472.00	5	✓	Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail				
pharmacy-Specialist	40.00	1	.1	Pfizer
			_	
Inj 1 mg for ECP - PCT only - Specialist		10 mg		Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Speciali	st94.40 1	00 mg (OP 🗸	Baxter
• • •		,		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	ce) S	Subsidised	Generic
	\$	Per	•	Manufacturer
FLUDARABINE PHOSPHATE				
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	1	Fludara Oral
Inj 50 mg vial - PCT only - Specialist	634.00	5	1	Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	126.80	50 mg O	P 🗸	Baxter
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	1	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	29.44	1	1	Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist	0.62	100 mg	1	Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	1	DBL Gemcitabine
lnj 1 g		1	1	Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg	1	Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	52.57	1	1	Accord
	71.44		1	Irinotecan Actavis
				100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	1	Baxter
MERCAPTOPURINE		Ü		
Tab 50 mg - PCT - Retail pharmacy-Specialist	25.90	25	1	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialist				
Special Authority see SA1725 below		100 ml O	P 🗸	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

		Subsidy		Fully	
	(1	Manufacturer's Price)		Subsidised	
_		\$	Per		Manufacturer
ME	THOTREXATE				
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist	9.98	90	1	Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist		90	1	Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist	56.05	5	/	Methotrexate DBL
*	Inj 7.5 mg prefilled syringe	14.61	1	✓	Methotrexate
					Sandoz
*	Inj 10 mg prefilled syringe	14.66	1	1	Methotrexate
					Sandoz
*	Inj 15 mg prefilled syringe	14.77	1	1	Methotrexate
					Sandoz
*	Inj 20 mg prefilled syringe	14.88	1	1	Methotrexate
	, 0, , 0				Sandoz
*	Inj 25 mg prefilled syringe	14.99	1	1	Methotrexate
	, 0, , 0				Sandoz
*	Inj 30 mg prefilled syringe	15.09	1	1	Methotrexate
	, 0, , 0				Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist	30.00	5	1	Methotrexate DBL
	, , , , , , , , , , , , , , , , , , , ,				Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialis	st45.00	1	1	DBL Methotrexate
•	m, so my por m, so m mar i or moram pharmacy operani		•		Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist	25.00	1	1	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		•		
	pharmacy-Specialist	79.99	1	1	Methotrexate Ebewe
*	Inj 1 mg for ECP – PCT only – Specialist		1 mg		Baxter
*	Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist		mg O		Baxter
	METREXED – PCT only – Specialist – Special Authority see SA		9 0	-	
r =	Inj 100 mg vial Specialist – Special Authority see SA		1	1	Juno Pemetrexed
	Inj 500 mg vial		1		Juno Pemetrexed
	Inj 1 mg for ECP		1 mg		Baxter
	11 1 11 19 101 LOI		ring	•	DUALGI

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Permetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
- 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

THIOGUANINE - PCT - Retail pharmacy-Specialist	
Tab 40 mg	

Other Cytotoxic Agents

AMSACRINE - PCT only - Specialist		
Inj 50 mg per ml, 1.5 ml ampoule1,500.00	6	✓ Amsidine S29
4,736.00		✓ Amsidine S29
Inj 75 mg1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist		
Cap 0.5 mg1,175.87	100	Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist		
Inj 1 mg per ml, 10 ml vial	10	✓ Phenasen
Inj 10 mg for ECP481.70	10 mg OP	✓ Baxter
BLEOMYCIN SULPHATE - PCT only - Specialist		
Inj 15,000 iu, vial	1	✓ DBL Bleomycin Sulfate
Inj 1,000 iu for ECP14.32	1,000 iu	✓ Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see SA1889 below		
Inj 3.5 mg vial74.93	1	✓ DBL Bortezomib
Inj 1 mg for ECP22.26	1 mg	✓ Baxter

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

DACARBAZINE - PCT only - Specialist

Inj 200 mg vial	72.11	1	✓ DBL Dacarbazine
	580.60	10	 Dacarbazine
			APP S29
Ini 200 mg for ECP	79 11	200 mg OP	✓ Rayter

	Subsidy (Manufacturer's P	rice) Subs	Fully	
	(Manufacturer's P \$	rice) Subs Per	iaised	
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist				
Inj 0.5 mg vial	255.00	1	1	Cosmegen
Inj 0.5 mg for ECP	255.00	0.5 mg OP		Baxter
DAUNORUBICIN - PCT only - Specialist				
Inj 2 mg per ml, 10 ml	171.93	1	/	Pfizer
Inj 20 mg vial		10		Daunorubicin
. •				Zentiva S29
Inj 20 mg for ECP	171.93	20 mg OP	1	Baxter
DOCETAXEL - PCT only - Specialist				
Inj 20 mg	48 75	1	/	Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1		DBL Docetaxel
Inj 20 mg per ml, 4 ml vial		1		Docetaxel
·, · · · g · · · ·				Accord \$29
Inj 80 mg	195.00	1	/	Docetaxel Sandoz
Inj 1 mg for ECP		1 mg		Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist		9		Duntoi
Inj 2 mg per ml, 5 ml vial	10.00	1	_	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		Doxorubicin Ebewe
iiij 2 iiig pei iiii, 25 iiii vial	17.00	'		Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial		1		Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1		Arrow-Doxorubicin
.,g po, .oo	69.99	·		Accord \$29
	00.00			Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	_	Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist		3		
Inj 2 mg per ml, 5 ml vial	25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1		Epirubicin Ebewe
Inj 1 mg for ECP		1 mg		Baxter
ETOPOSIDE		Ü		
Cap 50 mg - PCT - Retail pharmacy-Specialist	340 73	20	/	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10		Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia		1		Rex Medical
Inj 1 mg for ECP - PCT only - Specialist		1 mg		Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist		Ü		
Inj 100 mg (of etoposide base)	40.00	1	/	Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg		Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pha		٠ع		-
ПТОНОХТОВЕА [ПТОВОХТСАВВАМІОЕ] — РСТ — Retail pria 		100	/	Devatis
		100	•	Devalla
IBRUTINIB – Special Authority see SA2168 below – Retail phar	•	00	,	landa and a s
Tab 140 mg	,	30		Imbruvica
Tab 420 mg	9,652.00	30	•	Imbruvica

⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
 - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
 - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
 - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

Inj 5 mg vial – PCT only – Specialist	109.74	1	Zavedos
Inj 10 mg vial - PCT only - Specialist	233.64	1	Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77 1	mg	✓ Baxter
ENALIDOMINE Date: I abandone Canadalist Con	asial Authority and CACOAT halass		

LENALIDOMIDE - Retail pharmacy-Specialist - Special Authority see SA2047 below

Wastage claimable	•		
Cap 5 mg	5,122.76	28	✓ Revlimid
Cap 10 mg	4,655.25	21	✓ Revlimid
, ,	6,207.00	28	✓ Revlimid
Cap 15 mg	5,429.39	21	Revlimid
, -	7,239.18	28	Revlimid
Cap 25 mg	7,627.00	21	✓ Revlimid

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the

Cubaide		F. III.	Drondor	
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00 50	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50 50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Spe	ecialist177.45 15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Sp	pecialist407.40 15	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96 100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist		
Inj 5 mg vial	641.70 1	✓ Accord S29
Inj 20 mg vial		✓ Teva
Inj 1 mg for ECP	269.85 1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist		
Inj 2 mg per ml, 10 ml vial	97.50 1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP	5.51 1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Aut	hority see SA2163 below	
Tab 100 mg	3,701.00 56	✓ Lynparza
Tab 150 mg	3,701.00 56	✓ Lynparza

⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and

Subsidy (Manufacturer's Price)	,		Brand or Generic
 \$	Per	/Siuiseu	Manufacturer

continued...

- 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
- 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
- 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
 - 2 Either:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
 - 3 Treatment to be administered as maintenance treatment; and
 - 4 Treatment not to be administered in combination with other chemotherapy; and
 - 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
 - 5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	Paclitaxel Ebewe
Inj 100 mg	24.00	1	Paclitaxel Ebewe
, ,	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
, ,	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
, ,	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see	SA1979 on the next page		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

Subsidy	Fully		Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Roth:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail ph	armacy-Specialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 below	v – Retail pharmacy		
Cap 5 mg	9.13	5	✓ Temaccord
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
Cap 100 mg	35.98	5	✓ Temaccord
	40.20		✓ Apo-Temozolomide
Cap 140 mg	50.12	5	✓ Temaccord
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ Temaccord

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has

	ubsidy cturer's Price) Subs	Fully	Brand or Generic
·	\$ Per	•	Manufacturer

continued...

relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special	Authority see SA1124 below		
Cap 50 mg	378.00	28	✓ Thalomid
Cap 100 mg	756.00	28	✓ Thalomid

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

Cap 10 mg - PC	T – Retail pharmacy-Specialist	479.50	100	Vesanoid
	ail pharmacy-Specialist - Special Authority		he next page	
Tab 14×10 mg, 7	7×50 mg, 21×100 mg	1,771.86	42 OP	✓ Venclexta
			14 OP	✓ Venclexta
Tab 50 mg		239.44	7 OP	✓ Venclexta
Tab 100 mg - Wa	astage claimable	8,209.41	120	✓ Venclexta

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist270.37	5	✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist6.00	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	✓ DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	✓ DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter

	Subsidy (Manufacturer's Pric	o) Sui	Fully	Brand or Generic
	(Manufacturer's Fric	Per Sui	JSIUISEU ✓	Manufacturer
/INORELBINE				
Cap 20 mg	30.00	1	1	/inorelbine Te Arai
Cap 30 mg		1	1	/inorelbine Te Arai
Cap 80 mg	60.00	1	1	/inorelbine Te Arai
Inj 10 mg per ml, 1 ml vial - PCT only - Specialist		1	✓ N	Navelbine
	42.00		1	/inorelbine Ebewe
Inj 10 mg per ml, 5 ml vial - PCT only - Specialist	56.00	1	✓ N	Navelbine
	210.00		✓ \	/inorelbine Ebewe
	328.65		✓ 9	Sagent S29
Inj 1 mg for ECP - PCT only - Specialist	1.25	1 mg	✓ E	Baxter
Inj 50 mg for ECP - PCT only - Specialist		50 mg OP	✓ E	Baxter (Sagent)
Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe. Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe.	r 2024)	J		, ,

Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable

Tab 20 mg	60	✓ Sprycel
Tab 50 mg	60	✓ Sprycel
Tab 70 mg	60	✓ Sprycel

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or

|--|

continued...

- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special Auth	ority see SA2115 below		
Tab 100 mg	329.70	30	Alchemy
Tab 150 mg	569.70	30	✓ Alchemy

⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Fither:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB – Retail pharmacy-Specialist – Special Authority s	see SA2116 below		
Tab 250 mg	918.00	30	✓ Iressa

⇒SA2116 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
<u> </u>	1 01		Wandactarci

continued...

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESII ATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	Tab 100 mg - [Xpharm] - Special Authority see SA1460		
	below2,400.00	60	✓ Glivec
*	Cap 100 mg58.23	60	✓ Imatinib-Rex
*	Cap 400 mg84.79	30	✓ Imatinib-Rex

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms, and prescriptions

should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST - access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 on the next page - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6,532.00	120	Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

W	astage	claima	ble
---	--------	--------	-----

Tab 75 mg4,000.00	21	Ibrance
Tab 100 mg4,000.00	21	✓ Ibrance
Tab 125 mg4,000.00	21	✓ Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

Subs	sidy Fı	ılly E	Brand or
(Manufactu	urer's Price) Subsidis	ed 0	Generic
\$	\$ Per	\(\sigma\)	Manufacturer

continued...

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state: and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal: or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
RUXOLITINIB – Special Authority see SA1890 below – Retail p Wastage claimable	pharmacy			
Tab 5 mg	2,500.00	56	1	Jakavi
Tab 10mg	5,000.00	56	✓	Jakavi
Tab 15 mg	5,000.00	56	✓	Jakavi
Tab 20 mg	5,000.00	56	/	Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2117 below - Retail pharmacy

Cap 12.5 mg	28	 Sunitinib Pfizer
Cap 25 mg416.77	28	✓ Sunitinib Pfizer
Cap 50 mg694.62	28	✓ Sunitinib Pfizer

⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
- 5.2 Haemoglobin level < lower limit of normal; or
- 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
- 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
- 5.5 Karnofsky performance score of less than or equal to 70; or
- 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 84

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

Tab 250 mg4,276.19 120 **✓ Zytiga**

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

DIGALLITANIDE

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Tab 50 mg	4.21	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
· ·	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Authority	see SA1895 on	the next page	9
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
\$	Per	1	Manufacturer

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

OCTRECTIDE

Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓ Max Health
,			✓ Octreotide GH \$29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓ Max Health
			✓ Octreotide GH S29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓ Max Health
			✓ Octreotide GH S29
OCTREOTIDE LONG-ACTING - Special Authority see SA	12119 below – Retail pha	armacy	
Inj depot 10 mg prefilled syringe	439.97	ĺ	✓ Octreotide Depot Teva
Inj depot 20 mg prefilled syringe	647.03	1	✓ Octreotide Depot Teva
Inj depot 30 mg prefilled syringe	718.55	1	✓ Octreotide Depot Teva

⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	

continued...

- 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
- 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has failed: or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

TAMOXIFEN CITRATE

*	Tab 10 mg15.00	60	✓ Tamoxifen Sandoz
*	Tab 20 mg	60	✓ <u>Tamoxifen Sandoz</u>

	(Manufacturer's Price) \$	Su Per	ıbsidised ✓	Generic Manufacturer
Aromatase Inhibitors				
ANASTROZOLE * Tab 1 mg	4.55	30	✓ <u>A</u> ı	natrole
EXEMESTANE * Tab 25 mg	9.86	30	✓ Pf	fizer Exemestane
LETROZOLE * Tab 2.5 mg	5.84	30	✓ <u>Le</u>	<u>etrole</u>

Subsidy

Fully

Brand or

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE			
* Tab 25 mg	7.36	60	Azamun
* Tab 50 mg	8.10	100	✓ Azamun
MYCOPHENOLATE MOFETIL			
Tab 500 mg	35.90	50	✓ Cellcept
Cap 250 mg	35.90	100	✓ Cellcept
Powder for oral lig 1 g per 5 ml - Subsidy by endorsement	187.25	165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT - Special Authority see SA2103 below	– Retail pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Inj 50 mg prefilled syringe	1,050.00	4	✓ Enbrel

⇒SA2103 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Subsidy	Fu	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	:	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or

Subsidy	F	ully	Brand or	
(Manufacturer's Pr	rice) Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

continued...

2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
- 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
- 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment: and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Fither:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	✓	Manufacturer

continued...

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Specia	alist		
Inj 50 mg per ml, 5 ml	2,774.48	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only	– Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29

Monoclonal Antibodies

ADALIMUMAB (AMGEVITA) – Special Authority see SA21/8	8 below – Retail pharma	acy	
Inj 20 mg per 0.4 ml prefilled syringe	190.00	1	✓ Amgevita
Inj 40 mg per 0.8 ml prefilled pen	375.00	2	✓ Amgevita
Inj 40 mg per 0.8 ml prefilled syringe	375.00	2	✓ Amgevita

⇒SA2178 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
 - - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics: and
- 3 Patient has 3 or more active lesions: and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plague psoriasis: or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plague or plagues have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment: and
 - 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 2.2 Fither:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

1 Patient has pvoderma gangrenosum*; and

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	 Manufacturer 	

continued...

2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
(Manuacturer's Frice)				
\$	Per	/	Manufacturer	

continued...

Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	С
\$	Per	✓	Manufacturer	

continued...

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course iuvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects: or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or

2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or

2 All of the following:

- 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated): and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated): and

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic	
	Per	1	Manufacturer	

continued...

- 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 25 Fither
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate: and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4: or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 below - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe	1,599.96	2	✓ Humira
Inj 40 mg per 0.4 ml prefilled syringe	1,599.96	2	Humira
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	1,599.96	2	Humira

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

treatment: or

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — **(Crohn's disease - adult)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
\$	Per	•	Manufacturer

continued...

- 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Fither:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Subsidy	Fu	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

1 ✓ Evlea

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after

	DNCOLOGY AGENTS AND IWIW	UNUSUFFR	LOGANTO		
			Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
contin	ed the first dose to assess response to treatr Either:	ment; and			
Э	9.1 Patient has not previously received 9.2 Both:	d an anti-IL5 bio	logical therapy for thei	ir severe eosino	philic asthma; or
	9.2.1 Patient was refractory or in 9.2.2 Patient was not eligible to o within 12 months of comme	continue treatme	nt with previous anti-I		erapy and discontinued
	al — (Severe eosinophilic asthma) only or applications meeting the following criter		ory physician or clinica	al immunologist	. Approvals valid for 2
	An increase in the Asthma Control Test (A Either:	ACT) score of at	least 5 from baseline;	and	
	2.1 Exacerbations have been reduced2.2 Reduction in continuous oral cortic control.		•		·
CASIF	VIMAB AND IMDEVIMAB - [Xpharm] - S	Special Authority	see SA2096 below		
In	120 mg per ml casirivimab, 11.1 ml vial (1 per ml imdevimab, 11.1 ml vial (1)	, ,	,	1 OP 🗸	Ronapreve
⇒SA	096 Special Authority for Subsidy				
valid f	pplication — (Treatment of profoundly r 2 weeks for applications meeting the foll		romised patients) fro	om any relevant	practitioner. Approvals
	e following:	D 10: and			
	Patient has confirmed (or probable) COVI The patient is in the community with mild		ase severity*· and		
	Patient is profoundly immunocompromise against COVID-19 or is unvaccinated; and	d** and is at risk		ed an adequate	response to vaccination
4	Patient's symptoms started within the last				
	Patient is not receiving high flow oxygen of				
	Casirivimab and imdevimab is to be admi		•		mg.
** Exa	* Mild to moderate disease severity as den riples include B-cell depletive illnesses or	patients receivin	g treatment that is B-0		
	IMAB - PCT only - Specialist - Special				Ful-14
	5 mg per ml, 20 ml vial 5 mg per ml, 100 ml vial				Erbitux Erbitux
	1 mg for ECP			= -	Baxter
	697 Special Authority for Subsidy			Ü	
	pplication only from a medical oncologis	t or medical pra	ctitioner on the recom	mendation of a	medical oncologist.
	als valid for 6 months for applications mee	eting the following	g criteria:		
	e following: Patient has locally advanced, non-metast	atic cauamous	call cancer of the book	d and nack: and	
	Patient has locally advanced, non-metast Patient is contraindicated to, or is intolera			a and neun, and	
	Patient has good performance status; and				
	To be administered in combination with ra				

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA2158 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2179 below

Inj 100 mg	428.00	1	Remicade
Inj 1 mg for ECP	4.40	1 mg	Baxter

⇒SA2179 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

	Subsidy		Fully	Brand or
(1	Manufacturer's Price)	Subs	sidised	Generic
	\$	Per	1	Manufacturer

continued...

- 1 Paediatric patient has active Crohn's disease; and
- 2 Fither
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (**Graft vs host disease**) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less: and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist.

bsidy	Fully	Brand or
turer's Price) Subsid	dised	Generic
 \$ Per	✓	

continued...

Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

- Either:
 - 1 A withdrawal period has been tried and the patient has relapsed; or
 - 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- uı. 1 Fither
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation; or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis: or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis; or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis: and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:
All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's guality of life (see Notes); and
- 2 Fither
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Fither:

Subsidy (Manufacturer's Price)	s	Fully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2.1 Patients SCCAI is greater than or equal to 4; or
- 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs: and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI: and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

MEPOLIZUMAB - Special Authority see SA2154 below	N – Retail pharmacy		
Inj 100 mg prefilled pen	1,638.00	1	Nucala
Inj 100 mg vial	1,638.00	1	Nucala

⇒SA2154 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA2155 below

Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	Baxter

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive: and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and</p>
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Subsidy	Fu	ılly	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	/	Manufacturer	

continued...

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe	450.00	1	Xolair
Ini 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months. unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
- 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Specialist - Special Authority see SA2143 below

✓ Synagis (Synagis Inj 100 mg per ml, 1 ml vial to be delisted 1 January 2024)

⇒SA2143 Special Authority for Subsidy

Initial application — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Infant was born in the last 2 years and has severe lung, airway, neurological or neuromuscular disease that requires ongoing, life-sustaining community ventilation; or
 - 2 Both:
 - 2.1 Infant was born in the last 12 months; and
 - 2.2 Any of the following:
 - 2.2.1 Patient was born at less than 28 weeks gestation; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient was born at less than 32 weeks gestation; and
 - - 2.2.2.2.1 Patient has chronic lung disease; or
 - 2.2.2.2. Patient is Māori or any Pacific ethnicity; or
 - 2.2.3 Both:
 - 2.2.3.1 Patient has haemodynamically significant heart disease; and
 - 2.2.3.2 Any of the following:
 - 2.2.3.2.1 Patient has unoperated simple congenital heart disease with significant left to right shunt (see note a); or
 - 2.2.3.2.2 Patient has unoperated or surgically palliated complex congenital heart disease; or
 - 2.2.3.2.3 Patient has severe pulmonary hypertension (see note b); or
 - 2.2.3.2.4 Patient has moderate or severe LV failure (see note c).

Notes:

a) Patient requires/will require heart failure medication, and/or patient has significant pulmonary hypertension, and/or patient

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

will require surgical palliation/definitive repair within the next 3 months.

- b) Mean pulmonary artery pressure more than 25 mmHg.
- c) LV Ejection Fraction less than 40%.

Renewal — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months where patient still meets initial criteria.

PERTUZUMAB - PCT only - Specialist - Special Author	rity see SA1606 below		
Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	✓ Baxter

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist -	Special Authority see SA197	'6 below	
Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin: or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept: or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or

Subsidy		Fully	Brand or
(Manufacturer's	Price) Subs	sidised	Generic
<u> </u>	Per	1	Manufacturer

continued...

- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist	cial Authority see SA2114 b	elow	
Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

SA2114 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Fither:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD: and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1.000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects: and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- Patient who was previously treated with rituximab for nephrotic syndrome*: and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — **(haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy): and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia*

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and

S	Subsidy	Fully	Brand or
(Manufa	cturer's Price) Sub	sidised	Generic
	\$ Per	✓	Manufacturer

continued...

- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and

Subsidy	Fully		Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Roth:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment: or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	
--	--

continued...

experienced intolerable side effects.

d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	✓	

continued...

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

 SILTUXIMAB Special Authority see SA1596 on the next page Retail pharmacy

	,		, ,			
Note: Siltuximab is to	be administered at de	oses no greater than	11 mg/kg eve	ry 3 weeks.		
Inj 100 mg vial			770.57	1	1	Sylvant
Inj 400 mg vial			3,082.33	1	1	Sylvant

Subsidy	,		Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Pe	er 🗸	Manufacturer	

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Inj 100 mg per ml, 1.5 ml vial with cilgavimab 100 mg per ml,1.5 ml vial	00 1	✓ Evusheld
TOCILIZUMAB – PCT only – Special Authority see SA2159 below Inj 20 mg per ml, 4 ml vial220.	00 1	✓ Actemra
		✓ Actemra S29 S29
		✓ RoActemra S29 S29
880.	00 4	✓ RoActemra S29 S29
Inj 20 mg per ml, 10 ml vial550.	00 1	✓ Actemra
		✓ Actemra S29 S29
		✓ RoActemra S29 S29
Inj 20 mg per ml, 20 ml vial1,100.	00 1	✓ Actemra
		✓ Actemra S29 S29
		✓ RoActemra S29 S29
4,400.	00 4	✓ RoActemra S29 S29
Inj 1 mg for ECP2.	85 1 mg	✓ Baxter

⇒SA2159 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018:15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Roth
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Fither:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and

- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

Baxter

1 ma

	Subsidy	Fully		Brand or
(Ma	anufacturer's Price)	Subsic	lised	Generic
	\$	Per	✓	Manufacturer

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and

	Subsidy	Fully	Brand or
(Manui	facturer's Price) S	Subsidised	Generic
	\$ Per	✓	Manufacturer

continued...

- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib: or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Fither:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2144 below

Inj 100 mg vial	2,320.00	1	Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	24.52	1 mg	✓ Baxter

⇒SA2144 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 5.1 Patient does not have symptomatic brain metastases; or
- 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine;
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

✓ Stelara

⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 Any of the following:

- 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy: or
- 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
- 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment: or
- 2 Both:
 - 2.1 Patient has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

insufficient benefit to meet renewal criteria: or

2.2.2 Both:

2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and

2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active ulcerative colitis: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB – PCT only – Special Authority see SA2183 below Inj 300 mg vial3,313.00

1 ✓ Entvvio

⇒SA2183 Special Authority for Subsidy

Initial application — (**Crohn's disease - adults**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated): or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
- 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
- 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids: or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has active ulcerative colitis: and

continued...

- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Programmed Cell Death-1 (PD-1) Inhibitors

ATEZOLIZUMAB - PCT only - Specialist - Special Auth	nority see SA2195 below		
Inj 60 mg per ml, 20 ml vial	9,503.00	1	✓ Tecentriq
Inj 1 mg for ECP	8.08	1 mg	✓ Baxter

⇒SA2195 Special Authority for Subsidy

Initial application — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with atezolizumab and met all remaining criteria below prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic non-small cell lung cancer; and
 - 2.2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
 - 2.3 There is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
 - 2.4 Patient has an ECOG 0-2; and
 - 2.5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
 - 2.6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 12 weeks; and
 - 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

- 1.2 Patient's disease has had a partial response to treatment; or
- 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

DURVALUMAB - PCT only - Specialist - Special Authority see SA2	164 below		
Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	Imfinzi
Inj 1 mg for ECP	9.59	1 mg	Baxter

⇒SA2164 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

NIVOLUMAB - PCT only - Specialist - Special Authority see SA2	2120 below		
Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	✓ Opdivo
Inj 1 mg for ECP		1 mg	✓ Baxter

⇒SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes: and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA2197 on the next page	e
Inj 25 mg per ml, 4 ml vial4,680.00	✓ Keytruda
Inj 1 mg for ECP47.74 1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2197 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

lesions is also considered progression).

 Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
 - 2.2 Patient has not had chemotherapy for their disease in the palliative setting; and
 - 2.3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
 - 2.4 There is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
 - 2.5 Pembrolizumab to be used as monotherapy; and
 - 2.6 Either:
 - 2.6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
 - 2.6.2 Both:
 - 2.6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
 - 2.6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment: and
 - 2.7 Patient has an ECOG 0-2: and
 - 2.8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 12 weeks; and
 - 2.9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (non-small cell lung cancer first-line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2.2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 2.3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 2.4 There is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 2.5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 2.6 Patient has an ECOG 0-2; and
- 2.7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 12 weeks; and
- 2.8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail pha	ırmacy		
Wastage claimable			
Tab 10 mg	6,512.29	30	Afinitor
Tab 5 mg	4,555.76	30	Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per	✓	Manufacturer
SIROLIMUS – Special Authority see SA2218 below – Retail pha	rmacy			
Tab 1 mg	749.99	100	✓ R	apamune
Tab 2 mg	1,499.99	100	✓ R	apamune
Oral liq 1 mg per ml	449.99	60 ml O	P 🗸 R	apamune

⇒SA2218 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	

continued...

- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg		100	✓ Tacrolimus Sandoz
Cap 1 mg		100	✓ Tacrolimus Sandoz
Cap 5 mg		50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – Special Authority see SA2079 on the next page – Reta	il pharmacy		
Tab 15 mg	1.00	28	✓ RINVOQ

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 inj per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
 - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy
Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy		Fully Brand or
	(Manufacturer's Prices)	e) Subs	sidised Generic Manufacturer
	*		
BEE VENOM ALLERGY TREATMENT – Special Authority see S			
Initiation kit - 5 vials freeze dried venom with diluent		1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluer		1 OP	✓ Hymenoptera S29
VASP VENOM ALLERGY TREATMENT – Special Authority see		_	
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			_
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze			_
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze			•
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Antihistamines			
Anumstanines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1.71	100	✓ Zista
Zista to be Principal Supply on 1 September 2023			
★ Oral liq 1 mg per ml	2.84	200 ml	✓ <u>Histaclear</u>
CHLORPHENIRAMINE MALEATE			
♦ Oral liq 2 mg per 5 ml	9.37	500 ml	✓ Histafen
DEXTROCHLORPHENIRAMINE MALEATE			
F Tab 2 mg	2.02	40	
ŭ	(8.40)		Polaramine
	`1.01 [′]	20	
	(5.99)		Polaramine
♦ Oral liq 2 mg per 5 ml		100 ml	
	(10.29)		Polaramine
EXOFENADINE HYDROCHLORIDE			
← Tab 60 mg	4.34	20	
	(8.23)		Telfast
₭ Tab 120 mg		10	
	(8.23)	0.0	Telfast
	14.22	30	Talfaat
	(26.44)		Telfast
ORATADINE			
★ Tab 10 mg		100	✓ <u>Lorafix</u>
* Oral liq 1 mg per ml	1.43	100 ml	Haylor syrup

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	(Manufacturer's			Generic
	\$	Per	1	Manufacturer
PROMETHAZINE HYDROCHLORIDE				
₭ Tab 10 mg	1.39	50	1	Allersoothe
₭ Tab 25 mg		50		Allersoothe
FOral liq 1 mg per 1 ml		100 ml	1	Allersoothe
★ Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a		5	1	Hospira
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose	14.01	200 dose OP	1	Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP		Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP		Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP		Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free		200 dose OP		Beclazone 250
	22.07	200 dose Oi	٠	Deciazone 250
BUDESONIDE Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	/	Pulmicort
rowder for initial attorit, 100 micg per dose	17.00	200 00se OF	•	Turbuhaler
Douglas for inholation, 000 may not door	10.00	000 doos OD	./	Pulmicort
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	•	Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	J	Pulmicort
1 owder for initial autori, 400 micg per dose	52.00	200 dose Oi	٠	Turbuhaler
LUTICASONE				
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	1	Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP		Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP		Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP		Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose OP		Flixotide
Powder for inhalation, 250 mcg per dose		60 dose OP		Flixotide Accuhaler
Toward for initial allott, 250 mag per 4550		00 0000 01	_	Thixotiae Addunater
Inhaled Long-acting Beta-adrenoceptor Agonis	sts			
FORMOTEROL FUMARATE				
Powder for inhalation, 12 mcg per dose, and monodose dev		60 dose		
	(35.80)		_,	Foradil
Foradil Powder for inhalation, 12 mcg per dose, and monodose	device to be de	listed 1 July 2023	3)	
FORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated				
(equivalent to eformoterol fumarate 6 mcg metered dos	e)10.32	60 dose OP		
·	(16.90)			Oxis Turbuhaler
NDACATEROL	•			
Powder for inhalation 150 mcg	61.00	30 dose OP	1	Onbrez Breezhaler
Powder for inhalation 300 mcg		30 dose OP		Onbrez Breezhaler
G		00 0000 01	•	JDI DI OULIUIUI
SALMETEROL Association below CEO from CE many new dates	00.05	100 dans OD	,	Camanamat
Aerosol inhaler CFC-free, 25 mcg per dose		120 dose OP		Serevent
Powder for inhalation, 50 mcg per dose, breath activated	26.25	60 dose OP	•	Serevent Accuhaler

120 dose OP

120 dose OP

60 dose OP

60 dose OP

✓ Seretide

✓ Seretide

✓ Seretide Accuhaler

✓ Seretide Accuhaler

(M	Subsidy anufacturer's Price		Fully Brand or ised Generic Manufactur	er
Inhaled Corticosteroids with Long-Acting Beta-Ad	renoceptor A	Agonists		
BUDESONIDE WITH EFORMOTEROL Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide with 6 mcg eformoterol fumarate metered dose) Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mcg eformoterol fumarate metered dose) – No more than 2	41.50 120	O dose OP	✓ DuoResp Sp	iromax
dose per day	82.50 120	O dose OP	✓ DuoResp Sp	iromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg	18.23 120	0 dose OP	✓ Vannair	
Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg	33.74 120	0 dose OP	✓ Symbicort Turbuhale	r 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg	21.40 120	O dose OP	✓ Vannair	
Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg	33.74 120	0 dose OP	✓ Symbicort Turbuhale	r 200/6
Powder for inhalation 400 mcg with eformoterol fumarate				
12 mcg - No more than Ž dose per day	33.74 60	dose OP	✓ Symbicort Turbuhale	r 400/12
FLUTICASONE FUROATE WITH VILANTEROL Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08 30	dose OP	✓ Breo Ellipta	
FLUTICASONE WITH SALMETEROL				

Beta-Adrenoceptor Agonists			
SALBUTAMOL	450	/ Vantalin	
Oral liq 400 mcg per ml	150 ml 10	✓ <u>Ventolin</u> ✓ Ventolin	
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO53.00	5	✓ Ventolin	

Inhaled Beta-Adrenoceptor Agonists

Aerosol inhaler 50 mcg with salmeterol 25 mcg25.79

Aerosol inhaler 125 mcg with salmeterol 25 mcg32.60

more than 2 dose per day......33.74

more than 2 dose per day......44.08

Powder for inhalation 100 mcg with salmeterol 50 mcg - No

Powder for inhalation 250 mcg with salmeterol 50 mcg - No

SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO	3.80	200 dose OP	✓ Respigen ✓ SalAir
	(6.20)		Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO	8.96	20	✓ Asthalin
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO	9.43	20	✓ Asthalin
TERBUTALINE SULPHATE			
Powder for inhalation, 200 mcg per dose (equivalent to			
250 mcg metered dose), breath activated	22.20	120 dose OP	 Bricanyl Turbuhaler

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Anticholinergic Agents

IPRATROPIUM BROMIDE

Aerosol inhaler, 20 mcg per dose CFC-free - Up to 400 dose			
available on a PSO16.	.20 200 do	ose OP 🗸 🗸	trovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 neb			
available on a PSO11.	.73 2	20 🗸 U	Jnivent
28	20	✓ A	ccord \$29

Inhaled Beta-Adrenoceptor Agonists with Anticholinergic Agents

SALBUTAMOL WITH IPRATROPIUM BROMIDE

Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg per		
dose CFC-free12.19	200 dose OP	Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per		
vial 2.5 ml ampoule – Un to 20 neb available on a PSO 11.04	20	✓ Duolin

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umedictinium
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	

continued...

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
 - 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA	A1584 on the previou	s page - Retail pharmacy
Powder for Inhalation 50 mcg with indacaterol 110 mcg	31.00 30 dose C	P VIItibro Breezhaler
TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see	SA1584 on the prev	ious page – Retail pharmacy
Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg	81.00 60 dose C	P Spiolto Respimat
UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 of	on the previous page	- Retail pharmacy
Powder for inhalation 62.5 mcg with vilanterol 25 mcg	77.00 30 dose C	P ✓ Anoro Ellipta

Antifibrotics

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg	2,554.00	60 OP	Ofev
Cap 150 mg	3,870.00	60 OP	Ofev

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 on the next page

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90	Esbriet
Tab 267 mg	1,215.00	90	Esbriet

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists

МО	NTELUKAST			
*	Tab 4 mg	3.10	28	✓ Montelukast Mylan
*	Tab 5 mg	3.10	28	✓ Montelukast Mylan
	•			✓ Montelukast Viatris
*	Tab 10 mg	2.90	28	✓ Montelukast Mylan
	·			✓ Montelukast Viatris

Methylxanthines

AMINIOPHVI I INE

AMINOTITIELINE		
* Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available on a		
PSO180.00	5	✓ DBL Aminophylline
THEOPHYLLINE		

111	LOTTTLLINE			
*	Tab long-acting 250 mg	23.94	100	✓ Nuelin-SR
*	Oral lig 80 mg per 15 ml	17 62	500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA - Special Authority see SA1978 below - Retail	l pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	,	Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2196 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor	37.5 mg		
and ivacaftor 75 mg	27,647.39	84	Trikafta
Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor	or 75 mg		
and ivacaftor 150 mg	27,647.39	84	✓ Trikafta

⇒SA2196 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Either:
 - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Either:
 - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
 - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Note:

 a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212273s004lbl.pdf.

IVACAFTOR - PCT only - Specialist - Special Author	rity see SA2017 below		
Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

1 Patient has been diagnosed with cystic fibrosis; and

	Subsidy (Manufacturer's Pr \$	ice) Subsi Per	Fully dised	Brand or Generic Manufacturer
continued	Ψ	1 01		Waltalactarer
2 Either:				
 Patient must have G551D mutation in the cystic fibrolleast 1 allele; or 	osis transmemb	rane conductar	nce reg	ulator (CFTR) gene on a
2.2 Patient must have other gating (class III) mutation (D, G178R, G5	51S, S	1251N, S1255P, S549N
and S549R) in the CFTR gene on at least 1 allele; a			- !4-	mbanaia an bu Maanahu
3 Patients must have a sweat chloride value of at least 60 mr sweat collection system; and	noi/L by quantite	alive pilocarpin	e iorito	prioresis or by Macroduc
4 Treatment with ivacaftor must be given concomitantly with s				
5 Patient must not have an acute upper or lower respiratory in (including antibiotics) for pulmonary disease in the last 4 was				
6 The dose of ivacaftor will not exceed one tablet or one sach	net twice daily; a	nd	inone v	mir ivadanor, and
7 Applicant has experience and expertise in the management	t of cystic fibrosi	S.		
SODIUM CHLORIDE Not funded for use as a nasal drop.				
Soln 7%	24.50	90 ml OP	✓ B	iomed
Novel Dyenovetions				
Nasal Preparations				
Allergy Prophylactics				
BUDESONIDE				
Metered aqueous nasal spray, 50 mcg per dose Metered aqueous nasal spray, 100 mcg per dose		200 dose OP 200 dose OP	_	<u>teroClear</u> teroClear
FLUTICASONE PROPIONATE	2.04	200 0036 01	• <u>5</u>	<u>terocrear</u>
Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ <u>F</u>	lixonase Hayfever
IDD ATDODUM DDOMDE				& Allergy
IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%	5.23	15 ml OP	√ U	nivent
Respiratory Devices				
MASK FOR SPACER DEVICE				
a) Up to 50 dev available on a PSO b) Only on a PSO				
c) Only for children aged six years and under			_	
Small	2.70	1	√ e	-chamber Mask
PEAK FLOW METER a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range	9.54	1	✓ M	lini-Wright AFS
Normal range	9.54	1	✓ M	Low Range lini-Wright
.				Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSO b) Only on a PSO				
220 ml (single patient)		1	-	-chamber Turbo
510 ml (single patient)	5.95	1	√ e	-chamber La Grande

✓ Volumatic

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Respiratory Stimulants

CAFFEINE CITRATE

Oral liq 20 mg per ml (10 mg base per ml)......15.10 25 ml OP ✓ Biomed



	Subsidy		Fully Brand or
	(Manufacturer's P	,	dised Generic
	\$	Per	✓ Manufacturer
Ear Preparations			
FLUMETASONE PIVALATE			
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform
			ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	IN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ Kenacomb
		7.0 0.	
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4.50	8 ml OP	
3	(9.27)		Sofradex
FRAMYCETIN SULPHATE	` ,		
Ear/Eye drops 0.5%	4.13	8 ml OP	
	(8.65)		Soframycin
	,		•
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expli-	citly stated othery	vise.	
	,		
Anti-Infective Preparations			
ACICLOVIR			
* Eye oint 3%	14.88	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL		ŭ	
Eye oint 1%	1.09	5 g OP	✓ Devatis
Eye drops 0.5%		10 ml OP	✓ Chlorsiq
, ,	7.50		✓ Chlorafast
a) Funded for use in the ear*. Indications marked with	* are unapproved	I indications.	
b) Chlorsig to be Principal Supply on 1 September 2023			
(Chlorafast Eye drops 0.5% to be delisted 1 September 2023)			
CIPROFLOXACIN			
Eye drops 0.3% - Subsidy by endorsement	9.73	5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis of	r severe bacteria	I conjunctivitis	resistant to chloramphenicol; or
for the second line treatment of chronic suppurative otities		; and the preso	cription is endorsed accordingly.
Note: Indication marked with a * is an unapproved indic	ation.		
GENTAMICIN SULPHATE			
Eye drops 0.3%	11.40	5 ml OP	✓ Genoptic
(Genoptic Eye drops 0.3% to be delisted 1 August 2023)			
PROPAMIDINE ISETHIONATE			
* Eye drops 0.1%	2.97	10 ml OP	
	(14.55)		Brolene
SODIUM FUSIDATE [FUSIDIC ACID]			
Eye drops 1%	5.29	5 g OP	✓ Fucithalmic
		-	

	Subsidy		Fully	Brand or	
	(Manufacturer s P	rice) S	Subsidised	Generic	
	\$	Per	•	Manufacturer	
TOBRAMYCIN					
Eye oint 0.3%	10.45	3.5 g OF	•	Tobrex	
Eye drops 0.3%		5 ml OF	•	Tobrex	
Corticosteroids and Other Anti-Inflammatory Pre	parations				
DEXAMETHASONE					
* Eye oint 0.1%	5.86	3.5 g OF	· •	Maxidex	
* Eye drops 0.1%		5 ml OF		Maxidex	
Ocular implant 700 mcg - Special Authority see SA1680 belo		· · · · · ·			
Retail pharmacy		1	1	Ozurdex	
	1,777.00	'	-	Ozuruck	

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not vet completed a family: and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
	sulphate 6,000 u per g5.5	.39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml4.	.50	5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM			
	Eye drops 0.1%	.80	5 ml OP	✓ Voltaren Ophtha

	Subsidy		Fully Brand or
	(Manufacturer's F		sidised Generic
	\$	Per	✓ Manufacturer
FLUOROMETHOLONE			
* Eye drops 0.1%	3.09	5 ml OP	✓ FML
	5.20	· · · · · ·	✓ Flucon
	3.20		· Hucon
LEVOCABASTINE			
Eye drops 0.5 mg per ml	8.71	4 ml OP	
	(10.34)		Livostin
LODOXAMIDE	()		
	0.74	40 100	
Eye drops 0.1%	8./1	10 ml OP	✓ Lomide
NEPAFENAC			
Eye drops 0.3%	8.80	3 ml OP	✓ Ilevro
, ,		0 1111 01	- nevro
PREDNISOLONE ACETATE			
Eye drops 1%	6.92	10 ml OP	Prednisolone-AFT
	7.00	5 ml OP	✓ Pred Forte
DDEDNICOLONE CODILIM DUOCDUATE Cassial Authori	tu ann CA171E balau	Datail above	
PREDNISOLONE SODIUM PHOSPHATE - Special Authori			
Eye drops 0.5%, single dose (preservative free)	41.20	20 dose	✓ Minims
			Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE

Glaucoma Preparations - Beta Blockers

BETAXOLOL		
* Eye drops 0.25%11.8	0 5 ml OP	✓ Betoptic S
* Eye drops 0.5%	0 5 ml OP	✓ Betoptic
TIMOLOL		
* Eye drops 0.25%	1 5 ml OP	✓ Arrow-Timolol
* Eye drops 0.5%	4 5 ml OP	✓ Arrow-Timolol
* Eye drops 0.5%, gel forming – Subsidy by endorsement	8 2.5 ml OP	✓ Timoptol XE

Subsidised for patients who were taking timolol eye drops 0.5%, gel forming prior to 1 April 2023 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of timolol eye drops 0.5%, gel forming.

(Timoptol XE Eye drops 0.5%, gel forming to be delisted 1 March 2024)

Glaucoma Preparations - Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE * Tab 250 mg	17.03	100	✓ Diamox
BRINZOLAMIDE * Eye drops 1%	7.30	5 ml OP	✓ <u>Azopt</u>

	Subsidy (Manufacturer's Pr \$	ice) Subs Per	Fully idised	Brand or Generic Manufacturer
DORZOLAMIDE HYDROCHLORIDE – Subsidy by endorsemen Subsidised for patients who were taking dorzolamide hydroc endorsed accordingly. Pharmacists may annotate the presc dispensing of dorzolamide hydrochloride eye drops 2%.	hloride eye drops			
* Eye drops 2%	9.77 (17.44)	5 ml OP	Trı	usopt
(Trusopt Eye drops 2% to be delisted 1 March 2024)	()			
DORZOLAMIDE WITH TIMOLOL				
* Eye drops 2% with timolol 0.5%	2.73	5 ml OP	✓ <u>Do</u>	ortimopt
Glaucoma Preparations - Prostaglandin Analog	jues			
BIMATOPROST			<i>-</i>	
* Eye drops 0.03%	5.95	3 ml OP		<u>matoprost</u> Multichem
LATANOPROST			-	
* Eye drops 0.005%	1.82	2.5 ml OP	✓ <u>Te</u>	<u>va</u>
TRAVOPROST	0.75	0.5 1.00		
* Eye drops 0.004%	9.75	2.5 ml OP	✓ <u>Tra</u>	<u>avatan</u>
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE	4.00	5l OD		Delegant die
* Eye drops 0.2%	4.29	5 ml OP	V Ar	row-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE * Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	√ Cc	ombigan
LATANOPROST WITH TIMOLOL	10.50	3 IIII OF	• 00	mbigan
Eye drops 0.005% with timolol 0.5%	2 49	2.5 ml OP	✓ Ar	row - Lattim
PILOCARPINE HYDROCHLORIDE		2.0 0.		<u> </u>
* Eye drops 1%	4.26	15 ml OP	✓ Iso	opto Carpine
* Eye drops 2%		15 ml OP	✓ Iso	opto Carpine
* Eye drops 4%	7.99	15 ml OP	✓ Iso	opto Carpine
Subsidised for oral use pursuant to the Standard Formu	lae.			
* Eye drops 2% single dose – Special Authority see SA0895				
below – Retail pharmacy	34.19	20 dose	✓ Mi	nims Pilocarpine
⇒SA0895 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals vali	id for 2 years for a	pplications me	eeting the	e following criteria:
Either:	ray to the process	otivo: or		
 Patient has to use an unpreserved solution due to an alle Patient wears soft contact lenses. 	rgy to the preserv	alive, or		
Note: Minims for a general practice are considered to be "tools of the control of	of trade" and are r	nt annroved a	s snecia	I authority items
Renewal from any relevant practitioner. Approvals valid for 2 ye				
benefiting from treatment.			- appiop	a.c and the patient to

Mydriatics and Cycloplegics			
ATROPINE SULPHATE * Eye drops 1%17.36	15 ml OP	✓ Atropt	
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%	15 ml OP	✓ Cyclogyl	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Pr	rice) Subsid	dised	Generic
	\$	Per	✓	Manufacturer
TROPICAMIDE				
* Eye drops 0.5%	7.15	15 ml OP	✓ M	lydriacyl
* Eye drops 1%		15 ml OP		lydriacyl
The Lyc diops 1/0	0.00	13 1111 01	· IVI	iyunacyi
Preparations for Tear Deficiency				
For acetylcysteine eye drops refer Standard Formulae, page 255	5			
HYPROMELLOSE				
	10.50	45 ml OD		1444
* Eye drops 0.5%	19.50	15 ml OP	✓ M	lethopt
HYPROMELLOSE WITH DEXTRAN				
* Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ P	olv-Tears
Lyc drops 0.0 /6 with doxidan 0.1 /6	2.00	10 1111 01	• •	ory rears

Preservative Free Ocular Lubricants

⇒SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

and the second s			
CARBOMER - Special Authority see SA2134 above - Retail pl	harmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL	- Special Authority see	SA2134 a	bove – Retail pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml	10.78	30	 Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Au	thority see SA2134 abo	ove – Retai	l pharmacy
Eye drops 1 mg per ml			
Hylo-Fresh has a 6 month expiry after opening. The P			
month is not relevant and therefore only the prescribed	I dosage to the nearest	OP may b	e claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g3.80	5 g OP	✓ VitA-POS

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	1	Manufacturer	

Various

PHARMACY SERVICES

May only be claimed once per patient.

 ✓ BSF Abacavir/ Lamivudine Viatris
 ✓ BSF Ziextenzo

- a) The Pharmacode for BSF Abacavir/Lamivudine Viatris is 2655853 see also page 106
- b) The Pharmacode for BSF Ziextenzo is 2657066 see also page 46

(BSF Abacavir/Lamivudine Viatris Brand switch fee to be delisted 1 August 2023)

(BSF Ziextenzo Brand switch fee to be delisted 1 September 2023)

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE		
Inj 200 mg per ml, 10 ml ampoule52.8	38 1	0 ✓ Martindale Pharma
NALOXONE HYDROCHLORIDE		

- a) Up to 10 inj available on a PSO
- b) Only on a PSO

Removal and Elimination

CHARCOAL

*	Oral liq 50 g per 250 ml43.50	0 250 ml Of	✓ Carbosorb-X
	a) I In to 050 ml available on a DCO		

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wa	stage	e cla	ima	ble
Tak	105		ام:	

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	Exiade
Tab 500 mg dispersible		28	Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>



	Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic	
	\$	Per	•	Manufacturer	
continued					

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below - I	Retail pharmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

⇒SA1480 Special Authority for Subsidy

DESFERBIOXAMINE MESII ATE

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

* Inj 500 mg vial	151.31	10	✓ DBL Desferrioxamine Mesylate for Inj BP ✓ Deferoxamine Pfizer S29 S29
SODIUM CALCIUM EDETATE * Inj 200 mg per ml, 5 ml	52 21	6	
* III 200 IIIg per IIII, 3 III	(156.71)	Ü	Calcium Disodium

Versenate

Standard Formulae

ACETYLCYSTEINE EYE DROPS PHENOBARBITONE ORAL LIQUID	
Acetylcysteine inj 200 mg per ml, 10 ml qs Phenobarbitone Sodium	1 g
Suitable eye drop base qs Glycerol BP	70 ml
Water	to 100 ml
CODEINE LINCTUS (3 mg per 5 ml)	
Codeine phosphate 60 mg PHENOBARBITONE SODIUM PAEDIATRIC OF	AL LIQUID (10
Glycerol 40 ml mg per ml)	
Preservative qs Phenobarbitone Sodium	400 mg
Water to 100 ml Glycerol BP	4 ml
Water	to 40 ml
CODEINE LINCTUS (15 mg per 5 ml)	
Codeine phosphate 300 mg PILOCARPINE ORAL LIQUID	
Glycerol 40 ml Pilocarpine 4% eye drops	qs
Preservative qs Preservative	qs
Water to 100 ml Water	to 500 ml
(Preservative should be used if quantity supplied	is for more
FOLINIC MOUTHWASH than 5 days.)	
Calcium folinate 15 mg tab 1 tab	
Preservative qs SALIVA SUBSTITUTE FORMULA	
Water to 500 ml Methylcellulose	5 g
(Preservative should be used if quantity supplied is for more Preservative	qs
than 5 days. Maximum 500 ml per prescription.) Water	to 500 ml
METHADONE MIXTURE (Preservative should be used if quantity supplied	
than 5 days. Maximum 500 mi per prescription.)	
Methadone powder qs	
Glycerol qs SODIUM CHLORIDE ORAL LIQUID	
Water to 100 ml Sodium chloride inj 23.4%, 20 ml	qs
Water METHYL HYDROXYBENZOATE 10% SOLUTION Water (Only funded if proceeding for tree-treet of bypon	qs
METHYL HYDROXYBENZOATE 10% SOLUTION (Only funded if prescribed for treatment of hypon Methyl hydroxybenzoate 10 g	atraemia)
Propylene glycol to 100 ml VANCOMYCIN ORAL SOLUTION (25 mg per m	١
(Use 1 ml of the 10% solution per 100 ml of oral liquid mixture) Vancomycin 500 mg injection	5 vials
Glycerin with sucrose suspension	37.5 ml
OMEPRAZOLE SUSPENSION Water	to 100 ml
Omeprazole capsules or powder qs (Only funded if prescribed for treatment of Clostr	
Sodium bicarbonate powder BP 8.4 g following metronidazole failure)	ululli ullilolle
Water to 100 ml	

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Extemporaneously Compounded Preparations and Galenicals** CODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency (90.09)Douglas Only in extemporaneously compounded codeine linctus. COLLODION FLEXIBLE Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined. 100 ml ✓ PSM COMPOUND HYDROXYBENZOATE - Only in combination Only in extemporaneously compounded oral mixtures. 100 ml ✓ Midwest GLYCERIN WITH SODIUM SACCHARIN - Only in combination Only in combination with Ora-Plus or when used in the vancomycin oral Iguuid Standard Formulae. 473 ml ✓ Ora-Sweet SF GLYCERIN WITH SUCROSE - Only in combination Only in combination with Ora-Plus or when used in the vancomycin oral Iguuid Standard Formulae. Suspension......30.95 473 ml ✓ Ora-Sweet GI YCFROL 500 ml ✓ healthE Glycerol BP Only in extemporaneously compounded oral liquid preparations. METHADONE HYDROCHLORIDE a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing frequency d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets). ✓ AFT 1 a METHYL HYDROXYBENZOATE ✓ Midwest 25 q METHYLCELLULOSE ✓ MidWest 100 q ✓ Ora-Plus 473 ml METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN - Only in combination 473 ml Ora-Blend SF METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only in combination 473 ml ✓ Ora-Blend PHENOBARBITONE SODIUM Powder - Only in combination......52.50 ✓ MidWest 10 a 325.00 ✓ MidWest 100 q Only in children up to 12 years PROPYLENE GLYCOL

✓ fully subsidised	
Principal Supply	

SODIUM BICARBONATE

Only in extemporaneously compounded methyl hydroxybenzoate 10% solution.

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

Liq......11.25

Midwest

✓ Midwest

500 ml

500 a

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$		Fully lised	Brand or Generic Manufacturer
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatio		500 ml	✓ M	lidwest
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT – Special Authority see SA1930 above – Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	✓	Manufacturer	

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA2204 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia: or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT	 Special Authority see 	SA2204 on the pre	evious page – Hospita	I pharmacy [HP3]

Emulsion (neutral)		
30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)12.30	200 ml OP	✓ Calogen
Oil30.00	500 ml OP	✓ MCT oil (Nutricia)
MCT Emulsion, 250 ml	4 OP	Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	armacy [HP3]	PROTEIN SUPPLEMENT – Special Authority see SA1524 above – Hospital pha
✓ Protifar	225 g OP	Powder
✓ Resource	227 g OP	8.95
Beneprotein	-	

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA	1095 above -	 Hospital pharm 	acy [HP3]
Liquid	3.75	500 ml OP	✓ Glucerna Select
·	7.50	1,000 ml OP	✓ Nutrison Advanced
			Diason
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095	above – Hos	spital pharmacy	[HP3]
Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	2.10		✓ Nutren Diabetes

Fat Modified Products

⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism.

Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule. for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED − Special Authority see SA2205 above − Hospital pharmacy [HP3]
Powder60.48 400 g OP ✓ Monogen

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Powder54.00 400 g OP

✓ Kindergen

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

applications meeting the following criteria:

R∩th:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority see Liquid		the previous pag 500 ml OP	ge – Hospital pharmacy [HP3] Nutrini Energy RTH Frebini Energy
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see Stiquid	2.68	ne previous page 500 ml OP	✓ Nutrini RTH✓ Pediasure RTH
	6.50		Frebini Original
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special pharmacy [HP3]	I Authority se	ee SA1379 on the	e previous page – Hospital
Liquid	6.00	500 ml OP	Nutrini Energy Multi Fibre
	7.00		✓ Frebini Energy Fibre
PAEDIATRIC ENTERAL FEED WITH FIBRE 1KCAL/ML - Special Apharmacy [HP3]	Authority see	SA1379 on the p	previous page – Hospital
Liquid	7.00	500 ml OP	✓ Frebini Original Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA	1379 on the	previous page -	Hospital pharmacy [HP3]
Liquid (strawberry)	1.60	200 ml OP	✓ Fortini
Liquid (vanilla)	1.60	200 ml OP	✓ Fortini
	6.99	500 ml OP	✓ Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA13	379 on the pr	evious page – H	ospital pharmacy [HP3]
Liquid (chocolate)	1.07	200 ml OP	✓ Pediasure
Liquid (strawberry)	1.07	200 ml OP	✓ Pediasure
Liquid (vanilla)	1.07	200 ml OP	✓ Pediasure
	1.34	250 ml OP	✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Autoharmacy [HP3]	thority see S	A1379 on the pre	evious page - Hospital
Liquid (unflavoured)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)		200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on	the previous	page – Hospital	pharmacy [HP3]
Powder		400 g OP	✓ Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...

Subsidy)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
	Per	✓	Manufacturer

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML - Special Authority see Liquid		revious page – 500 ml OP	
RENAL ORAL FEED 1.8 KCAL/ML - Special Authority see SA1 Liquid		ous page – Hos 220 ml OP	pital pharmacy [HP3] Nepro HP (strawberry) Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA11	01 on the previous	s <mark>page</mark> – Hospi	tal pharmacy [HP3]
Liquid, 200 ml bottle	11.52	4 OP	
	(13.24)		NovaSource Renal
Liquid (apricot) 125 ml	11.52	4 OP	✓ Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	✓ Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - Special Authority see S.	A1377 above -	Hospital pharmacy [HP3]
Liquid18.06 1,	000 ml OP	✓ Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see SA1377 above - He	lospital pharma	cy [HP3]
Liquid (grapefruit), 250 ml carton171.00	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton171.00	18 OP •	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton171.00	18 OP •	✓ Elemental 028 Extra

	Subsidy (Manufacturer's I \$	Price) Subs Per	Fully idised	Brand or Generic Manufacturer
ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see S Powder (unflavoured)		revious page – I 80 g OP		al pharmacy [HP3] Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Auti [HP3]	nority see SA137	7 on the previou	ıs page	e – Hospital pharmacy
Liquid	9.60	500 ml OP	✓ S	Survimed OPD
	12.04	1,000 ml OP	✓ N	lutrison Advanced Peptisorb

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML - Special Authority see SA1196 above - Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Low Energy Liquid.......4.00 Multi Fibre

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive: or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:

Subsidy (Manufacturer's	Ful Price) Subsidise	,	
` \$	Per	Manufacturer	

- 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
- 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
- 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease; or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube - refer to specific medical condition criteria); or

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or

9 Severe chronic neurological conditions.			
ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on p	•		
Liquid		250 ml OP 1.000 ml OP	✓ Ensure Plus HN ✓ Ensure Plus RTH
	7.00	1,000 1111 OF	✓ Nutrison Energy
	9.60		✓ Fresubin HP Energy
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on page	je 265 – Hosp	oital pharmacy [HP3]
Liquid		250 ml OP	✓ Isosource Standard
	5.29	1,000 ml OP	Nutrison Standard RTH
	0.50		✓ Osmolite RTH
	6.50		✓ Fresubin Original
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority se Liquid		i page 265 – Ho 1.000 ml OP	ospital pharmacy [HP3] Nutrison
Liquid	5.23	1,000 1111 01	800 Complete
			Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority see \$			
Liquid	5.29	1,000 ml OP	✓ Jevity RTH ✓ Nutrison Multi Fibre
	7.00		✓ Fresubin Original
			Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority see	SA1859 on p	age 265 – Hos	
Liquid		1,000 ml OP	Jevity Plus
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see			
Liquid	7.00	1,000 ml OP	✓ Jevity HiCal RTH ✓ Nutrison Energy
			Multi Fibre
	9.80		✓ Fresubin HP Energy Fibre
ENTERAL FEED WITH PROTEIN 1.2KCAL/ML - Special Authority	see SA1859 d	on page 265 – H	Hospital pharmacy [HP3]
Liquid		500 ml OP	✓ Fresubin Intensive
ORAL FEED (POWDER) - Special Authority see SA1859 on page 2	<mark>65</mark> – Hospital	pharmacy [HP:	3]
Powder (chocolate)	14.00	840 g OP	✓ Sustagen Hospital
	00.00	050 · OD	Formula
Powder (vanilla)	26.00 14.00	850 g OP 840 g OP	✓ Ensure✓ Sustagen Hospital
i ortaoi (raillia)	17.00	0-10 y 01	Formula Active
	26.00	850 g OP	✓ Ensure

	Subsidy	Fully	Brand or
(1)	Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 265 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml			
with Endorsement		200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement	0.85	237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML — Special Authority see SA1859 on page 265 — Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

continued...

SPECIAL FOODS

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
 \$	Per	✓	

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and

practitioner and date contacted.

- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

(1.90) Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER - Special Authority see SA1106 o	n the previous page – Hosp	pital pharmacy	[HP3]
Powder	6.53	300 g OP	✓ Nutilis
	7.25	380 g OP	✓ Feed Thickener
			Karicare Aptamil

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA	11/29 above – Hospital pharmacy	[HP3]
Powder	2.81 1,000 g	OP
	(5.15)	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA	1729 above – Hospital pharmacy [HP3]
Powder	3.93 1,000 g	OP
	(7.32)	NZB Low Gluten Bread Mix
	3.51	
	(10.87)	Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729	above - Hospital pharmacy [HP3]	
Powder	5.62 2,000 g	OP
	(18.10)	Horleys Flour

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	ce) Su Per	ibsidised ✓	Generic Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	orevious page – Ho	ospital pha	rmacy [H	P3]
Buckwheat Spirals	2.00	250 g OP		
	(3.11)			Orgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)			Orgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)			Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)			Orgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)			Orgran
Rice and Corn Penne	2.00	250 g OP		
	(2.92)			Orgran
Rice and Maize Pasta Spirals	2.00	250 g OP		
	(2.92)			Orgran
Rice and Millet Spirals	2.00	250 g OP		
	(3.11)			Orgran
Rice and corn spaghetti noodles	2.00	375 g OP		
	(2.92)			Orgran
Vegetable and Rice Spirals	2.00	250 g OP		
	(2.92)			Orgran
Italian long style spaghetti	2.00	220 g OP		
	(3.11)			Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	osidised	Generic
\$	Per	/	Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE -	Special Authority see S	A1108 on the	previous page – Hospital
pharmacy [HP3]			
Tabs	99.00	75 OP	✓ Phlexy 10
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex
, ,, ,			Powder

(,/, g			Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (neutral) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior Orange
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	✓ XP Maxamum
Liquid (berry)		125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ

Liquid (unflavoured)13	3.10	125 ml OP
Liquid (forest berries), 250 ml carton		18 OP 30 OP

✓ PKU Anamix Junior LQ✓ Easiphen Liquid

✓ PKU Lophlex LQ 20
 ✓ PKU Lophlex
 Sensation 20
 ✓ PKU Lophlex LQ 10

36 OP

Liquid (juicy berries) 62.5 ml	60 OP
Liquid (juicy citrus) 62.5 ml939.00	60 OP
Liquid (juicy orange) 62.5 ml	60 OP
Liquid (juicy berries) 125 ml936.00	30 OP
Liquid (juicy orange) 125 ml936.00	30 OP

✓ PKU Lophlex LQ 10
✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 20 ✓ PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING	MIX - Special Authority see SA1108 on the previous	page – Hospital į	oharmacy [HP3]
Powder	8.22	500 g OP	✓ Loprofin Mix
LOW DOCTEN DACTA	On a sial Authority and CA4400 and the monitors were	I I a a mile a la a man	· · [LID0]

LOW PROTEIN PASTA - Special Authority see S	SA1108 on the previous page - I	Hospital pharm	acy [HP3]
Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne	5.95	250 g OP	✓ Loprofin
Low protein rice pasta	11.91	500 g OP	✓ Loprofin
Macaroni	5.95	250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals	11.91	500 g OP	✓ Loprofin

Subsidy (Manufacturer's Price) \$ Fully Subsidised Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Gastrointestinal and Other Malabsorptive Problems

IINO ACID FORMULA - Special Authority see SA2092 below -	Hospital phar	macy [HP3]	
Powder	43.60	400 g OP	✓ Alfamino
		•	 Alfamino Junio
Powder (unflavoured)	53.00	400 g OP	✓ Elecare
,		Ü	✓ Elecare LCP
			✓ Neocate Gold
			✓ Neocate Junior Unflavoured
			✓ Neocate SYNEC
Powder (vanilla)	53.00	400 g OP	✓ Elecare
, ,		3	✓ Neocate Junior Vanilla

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or alleray or malabsorotion; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or

2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption: or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and

Subsidy (Manufacturer's Price)	Subs	Fully idised	Brand or Generic
 \$	Per	✓	Manufacturer

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
- 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption: or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea: or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...



	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)	Sı	ubsidised	Generic
	\$	Per	✓	Manufacturer

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketoenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)35.50	300 g OP	✓ KetoCal 4:1
		✓ Ketocal 3:1
Powder (vanilla)35.50	300 g OP	✓ KetoCal 4:1

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

Other Supplements for PKU

(PKU GMPro Ultra Lemonade Powder (Lemonade) 33.4 g sachets to be delisted 1 December 2023)

⇒SA2229 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient was previously receiving, or would receive PKU Lophlex Sensation Berries under (SA1108); and
- 2 PKU Sensation Berries is unable to be sourced at this time; and
- 3 Patient has trialled the currently funded PKU Lophlex products and these were not tolerated .

Note: These criteria are attached to short term funding to cover an out-of-stock situation on PKU Sensation Berries supplied by Nutricia.

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent.................0.00 10 ✓ BCG Vaccine

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy: or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
 - 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old; or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.
- Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid. 8 mcg pertussis filamentous

10 **Boostrix Boostrix**

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Subsid Per	ised Generic ✓ Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE -	· · · · · · · · · · · · · · · · · · ·		
Funded for any of the following:	[Apriaiii]		
1) A single dose for children up to the age of 7 who have o	ompleted primary imr	nunisation:	or
2) A course of four vaccines is funded for catch up prograr			
primary immunisation; or			
3) An additional four doses (as appropriate) are funded for			
pre- or post splenectomy; pre- or post solid organ transpregimens; or	nant, renai diaiysis ar	ia otner sev	rereiy immunosuppressive
Five doses will be funded for children requiring solid org	an transplantation.		
Note: Please refer to the Immunisation Handbook for approp		ch up progr	ammes.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin and 80 D-antigen units	0.00	40	4.1.4. 1. IDV
poliomyelitis virus in 0.5ml syringe		10	✓ <u>Infanrix IPV</u>
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AI	ND HAEMOPHILUS I	NFLUENZA	AE TYPE B VACCINE -
[Xpharm] Funded for patients meeting any of the following criteria:			
Up to four doses for children up to and under the age of	10 for primary immu	nisation: or	
2) An additional four doses (as appropriate) are funded for	, ,	,	to and under the age of
10 who are patients post haematopoietic stem cell trans			
post solid organ transplant, renal dialysis and other seve			
3) Up to five doses for children up to and under the age of			
Note: A course of up-to four vaccines is funded for catch up to complete full primary immunisation. Please refer to the Im-			
programmes.	That house of the hard of the	tioi illo api	or opinate contradiction outers ap
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,	0.00	40	/ Infamily have
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10	✓ <u>Infanrix-hexa</u>
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]			
One dose for patients meeting any of the following: 1) For primary vaccination in children; or			
2) An additional dose (as appropriate) is funded for (re-)im	munisation for patien	ts post hae	matopoietic stem cell
transplantation, or chemotherapy; functional asplenic; p			
or post cochlear implants, renal dialysis and other sever			
For use in testing for primary immunodeficiency disease	s, on the recommend	lation of an	internal medicine physician or
paediatrician.			
Haemophilus Influenzae type B polysaccharide 10 mcg			
conjugated to tetanus toxoid as carrier protein 20-40 mcg	;		
prefilled syringe plus vial 0.5 ml		1	✓ Hiberix
HEPATITIS A VACCINE - [Xpharm]			
Funded for patients meeting any of the following criteria:			
Two vaccinations for use in transplant patients; or			
Two vaccinations for use in children with chronic liver di One does of vaccine for close centrate of known benetit	,		
One dose of vaccine for close contacts of known hepatil	ils A Cases.		
Inj 1440 ELISA units in 1 ml syringe	0.00	1	✓ Havrix
Inj 720 ELISA units in 0.5 ml syringe		1	✓ Havrix Junior

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer	
HEPATITIS B RECOMBINANT VACCINE – [Xpharm]	0.00	1	√ F	ngerix-B	

- Funded for patients meeting any of the following criteria: 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
 - 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
 - 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
 - 4) for HIV positive patients; or
 - 5) for hepatitis C positive patients; or
 - 6) for patients following non-consensual sexual intercourse; or
 - 7) for patients following immunosuppression; or
 - 8) for solid organ transplant patients; or
 - 9) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 10) following needle stick injury.

Engerix-B Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients following immunosuppression; or
- 8) for solid organ transplant patients; or
- 9) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 10) following needle stick injury; or
- 11) for dialysis patients; or
- 12) for liver or kidney transplant patients.

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV]

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d)
- a) A) Any of the following:
 - 1) Maximum of two doses for children aged 14 years and under; or
 - 2) Maximum of three doses for patients meeting any of the following criteria:
 - 1) People aged 15 to 26 years inclusive; or
 - 2) Either:

People aged 9 to 26 years inclusive

- 1) Confirmed HIV infection; or
- 2) Transplant (including stem cell) patients: or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy
- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which

may be a sub-set of the population described in paragraphs A above. ✓ Gardasil 9

		Subsidy		Fully	Brand or
		(Manufacturer's Price)	Su	bsidised	Generic
		\$	Per	•	Manufacturer
INFLUENZA	VACCINE				
Inj 30 mo	cg in 0.25 ml syringe (paediatric quadrivalent vaccine)			
_ [X	pharm]	11.00	1	√ µ	Afluria Quad Junior (2023 formulation)
A)	INFLUENZA VACCINE - child aged 6 months to	35 months			
.,	is available each year for patients aged 6 months to		et the fol	lowing c	riteria, as set by Pharmac:
	i) all children aged 6 months to 35 months from	1 April 2023 to 31 De	ecember	2023.	
B)	Doctors are the only Contractors entitled to claim pa syringe (paediatric quadrivalent vaccine) to patients and they may only do so in respect of the influenza	eligible under the at	ove crit	eria for s	ubsidised immunisation
Inj 60 mo	eg in 0.5 ml syringe (quadrivalent vaccine)	110.00	10	√	Afluria Quad (2023 formulation)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) People 55 to 64 years of age (inclusive) and is Māori or of any Pacific ethnicity from 1 April 2023 to 31 December 2023; or
- c) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes: or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) Down syndrome, or
 - vii) are pregnant; or
- d) children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- e) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or
- f) children 3 to 12 years of age (inclusive), from 1 April 2023 to 31 December 2023;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$ P	Per	Manufacturer

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

inj, measies virus 1,000 doibbo, mumps virus 5,012 doibbo,		
Rubella virus 1,000 CCID50; prefilled syringe/ampoule of		
diluent 0.5 ml	5	✓ MMR II
	10	✓ Priorix

(Manufacturer's Price) Subsidised Generic	Subsidy		Fully	Brand or	
	(Manufacturer's Price))	Subsidised	Generic	
\$ Per ✔ Manufacturer	\$	Per	1	Manufacturer	

MENINGOCOCCAL (GROUPS A. C. Y AND W-135) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- a) A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients
 with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post
 solid organ transplant: or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
 - B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons.
 - C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A, C, Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
 - D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

to a total of approximately 55 mcg of tetanus toxoid carrier per 0.5 ml vial	0.00	1	✓ MenQuadfi
Inj 4 mcg of each meningococcal polysaccharide conjugated to a total of approximately 48 mcg of diphtheria toxoid carrier			
per 0.5 ml vial	0.00	1	✓ Menactra
•		5	✓ Menactra

	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised Per 🗸	Generic Manufacturer
	Φ	rei •	- Ivianulaciulei
MENINGOCOCCAL B MULTICOMPONENT VACCINE			
a) Only on a prescription			
b) No patient co-payment payable			
c) Any of the following:			
 a) Any of the following: A) Three doses for children up to 12 months of a 	iga (inclusiva) for prima	ary immunication	· or
B) Up to three doses (dependent on age at first of			
59 months of age (inclusive) for primary immu			
C) Both:	,	ŭ	•
1) Person is one year of age or over; and			
Any of the following:			
i) up to two doses and a booster eve			
patients with functional or anatomi		lement deficiency	(acquired or inherited), of
pre- or post-solid organ transplant ii) up to two doses for close contacts		oo of any group:	or
iii) up to two doses for close contacts	•	, , , ,	
iv) up to two doses for bone marrow t		ngoooda alood	o or any group, or
v) up to two doses for person pre- ar		ssion*; or	
D) Both:			
1) Person is aged between 13 and 25 year	rs (inclusive); and		
2) Either:			
i) Two doses for individuals who are	•		•
living in boarding school hostels, to Justice residences or prisons; or	ernary education halls	or residerice, mili	lary Darracks, Toulin
ii) Two doses for individuals who are	currently living in boar	dina school host	els, tertiary education hal
of residence, military barracks, or	, ,	•	
E) Contractors will be entitled to claim payment			
multicomponent vaccine to patients eligible un	nder the above criteria	pursuant to their	contract with Te Whatu
Ora Health New Zealand for subsidised immu		•	spect of the
Meningococcal B multicomponent vaccine list			District Control of Control
 F) Contractors may only claim for patient popula may be a sub-set of the population described 			by their contract, which
*Immunosuppression due to corticosteroid or other immu			eriod of areater than
28 days.	moodpproceive therap	y made bo for a pe	shod of groater than
Inj 175 mcg per 0.5 ml prefilled syringe	0.00	1 ✓ B	exsero
MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm]			
Both:			
1) The child is under 12 months of age; and			
2) Any of the following:			
 Up to three doses for patients pre- and post splet 			
HIV, complement deficiency (acquired or inherite	/· ' '	organ transplant	, or
Two doses for close contacts of meningococcal of the contacts of the contact of the conta	, 0 1	001/ 0701101 07	
3) Two doses for child who has previously had men4) A maximum of two doses for bone marrow transp		arry group; or	
5) A maximum of two doses for child pre- and post-			
Note: children under 12 months of age require two do	• • • • • • • • • • • • • • • • • • • •	efer to the Immun	isation Handbook for
recommended booster schedules with meningococcal			.caorr randbook for
*Immunosuppression due to steroid or other immunosu		st be for a period	of greater than 28 days.

✓ Neisvac-C

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm]

1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B.

7F, 9V, 14 and 23F; 3 mcg of pneumococcal

polysaccharide serotypes 4, 18C and 19F in 0.5 ml

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
- Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection: or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) cochlear implants or intracranial shunts; or
 - g) cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - I) diabetes: or
 - m) Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

1 Prevenar 13

	NATIONAL	IMMUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price)	Fully Subsidised Per 🗸	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – Either:	[Xpharm]		
1) Up to three doses (as appropriate) for patients with HI chemotherapy; pre- or post-splenectomy or with functi complement deficiency (acquired or inherited), cochlet 2) All of the following: a) Patient is a child under 18 years for (re-)immunis b) Treatment is for a maximum of two doses; and c) Any of the following: i) on immunosuppressive therapy or radiation immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; o	onal asplenia, pre- or ar implants, or primary sation; and therapy, vaccinate with	post-solid organ t immunodeficiend then there is expe	ransplant, renal dialysis, by; or cted to be a sufficient
v) who are immune-suppressed following orgor or vi) with cochlear implants or intracranial shunt vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more the prednisone of 2 mg/kg per day or greater, or	s; or an two weeks, and wh	o are on an equiv	valent daily dosage of
20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks ges xi) with cardiac disease, with cyanosis or failu xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with full xiii with Down syndrome; or xiv)	station; or re; or	gh-dose corticost	eroid therapy); or
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	0.00	1 ✓ <u>F</u>	neumovax 23
POLIOMYELITIS VACCINE – [Xpharm] Up to three doses for patients meeting either of the following 1) For partially vaccinated or previously unvaccinated inc 2) For revaccination following immunosuppression.			
Note: Please refer to the Immunisation Handbook for appro Inj 80D antigen units in 0.5 ml syringe		tch-up programm 1 ✓ <u>I</u> I	
ROTAVIRUS ORAL VACCINE – [Xpharm] Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 14 2) no vaccination being administered to children aged 24	•		
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube	0.00	10 ✓ F	totarix
Oral susp live attenuated human rotavirus	0.00	10 🔏	lotoriy

10

✓ Rotarix

1,000,000 CCID50 per dose, prefilled oral applicator......0.00

NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm]			
Either: 1) Maximum of one dose for primary vaccination for either a) Any infant born on or after 1 April 2016; or b) For previously unvaccinated children turning 11 yearicella infection (chickenpox), or 2) Maximum of two doses for any of the following: a) Any of the following for non-immune patients: i) with chronic liver disease who may in future ii) with deteriorating renal function before traniii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, v) for post exposure prophylaxis who are imm b) For patients at least 2 years after bone marrow to c) For patients at least 6 months after completion od) For HIV positive non immune to varicella with mile) For patients with inborn errors of metabolism at a	years old on or after 1 of the period of the	nsplantation; or nts.; or ce of their specia vice of their spec osuppression on	llist, or ialist, or advice of HIV specialist, o
varicella, or f) For household contacts of paediatric patients wh immune compromise where the household contacts of adult patients who hat immunocompromised, or undergoing a procedur has no clinical history of varicella. * immunosuppression due to steroid or other immunosuppre	to are immunocompron act has no clinical histo we no clinical history of e leading to immune co	nised, or undergo ry of varicella, or varicella and wh ompromise where	oing a procedure leading to o are severely e the household contact
28 days Inj 1350 PFU prefilled syringe	0.00	_	arivax arivax
VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] a) Only on a prescription b) No patient co-payment payable c) A) Funded for patients meeting the following criteria: 1) Two doses for all people aged 65 years B) Contractors will be entitled to claim payment from twaccine) to patients eligible under the above criteri Zealand for subsidised immunisation, and they mawaccine] listed in the Pharmaceutical Schedule. C) Contractors may only claim for patient populations a sub-set of the population described in paragraph Inj 50 mcg per 0.5 ml vial plus vial	a pursuant to their cont y only do so in respect within the criteria that a A above.	tract with Te What of the Varicella zare covered by the	atu Ora Health New coster vaccine [Shingles neir contract, which may be chingrix
Inj 19,400 PFU prefilled syringe plus vial	0.00	_	ostavax ostavax
Diagnostic Agents			
TUBERCULIN PPD [MANTOUX] TEST - [Xpharm] Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1 ✓ <u>T</u>	ubersol
fully subsidised	S29 Unapproved	medicine supplied	under Section 29

Principal Supply

Agents Used in the Treatment of Short	- Symbols -	Disorders	117	Amoxicillin	93
Short	3TC107	Agents Used in the Treatment of			
Short	7 MED NSHA Silver/Copper	Poisonings	253	Amphotericin B	33
A-	Short74	Agrylin	150		
Abacavir sulphate with	- A -	Albendazole	90		
Abacavir sulphate with	A-Scabies69				
Lamivudine	Abacavir sulphate106	Albustix	78	Amzoate	31
Abacavir/Lamivudine Viatris	Abacavir sulphate with	Alchemy	159	Anaesthetics	118
Abiratoroe acetate 165	lamivudine 106	Alchemy Oxaliplatin	146	Anagrelide hydrochloride	150
Acarbose .11 Alecensa 158 Anatrole .16 Accurelic 10 .49 Alendronate sodium .112 Androld Testocaps .86 Accuretic 20 .49 Alendronate sodium with .112 Androderm .88 Acetazolamide .250 colecalciferol .112 Antarole .112 Antarole .144 Antacidia .144 Antacidia .144 Antacidia .144 Antacidia .144 Antacidia and Antilifatulents .66 Antilifatulents .66 Antilifatulents .60 Antiliacricia .60 Antilifatulents .60	Abacavir/Lamivudine Viatris106	Alchemy Oxybutynin	78	Analgesics	119
Accare (a) .11 Alectinib. 158 Andriol Testocaps. 86 Accuretic 10 .49 Alendronate sodium. 112 Androderm. 86 Accuretic 20 .49 Alendronate sodium with Anoro Ellipta .24 Acetea Calouride .250 Acetea Calouride .112 Antabuse .14 Acetic acid with hydroxyquinoline and ricinoleic acid acid. .76 Alfaranino. .274 Antabuse .14 Acetylcysteine. .253 Alginic acid .6 Antailerry Preparations. .6 Aciclovir .76 Alginic acid .6 Antiallergy Preparations. .6 Aciclovir .76 Alginic acid .6 Antialarry Preparations. .6 Acidovir .76 Alginic acid .6 Antialierry Preparations. .6 Acidovir .76 Alginic acid .6 Antialierry Preparations. .2 Acidovir .76 Alginic acid .6 Antialierry Preparations. .2 Acidex .6	Abiraterone acetate165	Aldurazyme	30	Anastrozole	168
Accuretic 10. 49 Alendronate sodium	Acarbose11	Alecensa	158	Anatrole	168
Accuretic 10. 49 Alendronate sodium	Accarb11	Alectinib	158	Andriol Testocaps	82
Accetec 20. 49 Alendronate sodium with coetacidiren 250 colecalciferol. 112 Antabuse. 144 Antabuse. 144 Acetec. 48 Alfacalcidol 34 Antacids and Antifilatulents. 66 Acetylcysteine. 253 Alginic acid 66 Antiallergy Preparations. 238 Aci-Jel. 76 Alglucosidase alfa. 27 Antiancen Preparations. 238 Aci-Jel. 76 Alglucosidase alfa. 27 Antianaemics. 38 Antiallergy Preparations. 38 Aciclovir Infection. 102 Alkeran. 146 Antiandrogen Oral Alkeran. 146 Antiandrogen Oral Alkeran. 250 Antibacterials Topical. 56 Acigimox. 56 Allerpro Syneo 1 277 Antibacterials Topical. 56 Actemia S29. 220 Allerpro Syneo 2 277 Antibacterials Topical. 66 Actemia S29. 220 Allopurinol. 115 Anticopressants. 112 Actemia S29. 220 Allopurinol. 115 Anticopressants. 112 Actrapid Penfill. 10 Alpha-Adrenoceptor Blockers. 48 Anticolarinosetases. 111 Actrapid Penfill. 10 Alpha-Morx. 93 Antibinotytics, Haemostatics and Adapalene. 63 Alphamox 250. 93 Antifibrionicytics, Haemostatics and Adapalene. 63 Alpmanx 250. 93 Antifibrionicytics, Haemostatics and Adapalene. 63 Aluminium hydroxide. 66 Antifilation. 234 Antifibriorics. 244 Amibrisentan Wilaris. 59 Antifibriorics. 245 Advogen. 52 Antihypotensives. 55 Antihypotensives. 56 Antihiproric Preparations. 66 Antificency. 120 Antihiproric Preparations. 66 Antihiproric Preparations. 67 Antihiproric Preparations. 67 Antihiproric Prepar	Accuretic 1049	Alendronate sodium	112		
Acetacolamide .250 colecalciferol. .112 Antabuse .144 Acetec .48 Aclacidod .34 Antacids and Antiflatulents .69 Acetic acid with hydroxyquinoline and ricinoleic acid .76 Alfamino .274 Anthelminitics .99 Aci-Jel .76 Alfamino Junior .274 Antiacne Preparations .66 Aci-Jel .76 Alginic acid .6 Antianerina .27 Aciclevir .76 Alginic acid .6 Antianerina .23 Aciclovir .76 Algeran S29 .146 Contraceptives .77 Acidex .6 Allergro Syneo 1 .277 Antiandryhminics .56 Acidex .6 Allerpro Syneo 2 .277 Antibacerials Topical .6 Acidex .6 Allerpro Syneo 2 .277 Antibacerials Topical .6 Acternia .220 Allopurinol .115 Anticolinerials Topical .6 Acternia S29 .220 Allopurinol <td>Accuretic 2049</td> <td>Alendronate sodium with</td> <td></td> <td>Anoro Ellipta</td> <td>243</td>	Accuretic 2049	Alendronate sodium with		Anoro Ellipta	243
Acetec. 48 Afacacidolo .34 Antacids and Antiflatulents .6 Acetic acid with hydroxyquinoline and ricinoleic acid 76 Alfamino. .274 Anthelmintics .6 Acetylcysteine. .253 Alginic acid .6 Antialergy Preparations .6 Aci-Jel. .76 Algucosidase alfa. .27 Antialnergy Preparations .23 Aciclev. .6 Algucosidase alfa. .27 Antialnergy Preparations .23 Aciclovin .102 Alkeran .166 Antiallergy Preparations .6 Aciclovin .102 Alkeran .166 Antiallergy Preparations .23 Aciclovin .102 Alkeran .166 Antiallergy Preparations .23 Acicolovin .102 Alkeran .161 Antiallargy Preparations .23 Acitonion .102 Alkeran .29 .24 Antiallarian .24 Acitonion .102 Allerro .29 .277 Antibalergy Preparations .24	Acetazolamide250	colecalciferol	112		
Acetic acid with hydroxyquinoline and ricinoleic acid. Affamino. 274 Anthalminitics. 99 Acelylcysteine. 253 Alginic acid. 6 Antiallergy Preparations. 66 AcicJol. 76 Alginic acid. 6 Antianalergy Preparations. 236 AcicJovir Alkeran. 146 Antianadrogen Oral Antianadrogen Oral Infection. 102 Alkeran. 146 Contraceptives. 77 Acidox. 6 Allerpro Syneo 1 277 Antiandrogen Oral 77 Acidox. 6 Allerpro Syneo 1 277 Antibacterials 99 Actidox. 6 Allerpro Syneo 1 277 Antibacterials Topical 66 Actiretin. 69 Allersoothe. 240 Anticholinergic Agents. 244 Acterna S29. 220 Alloparop. 115 Anticholinesterases. 117 Actrapid Penfill. 10 Alpha-Keri Lotion. 67 Antifibrinolytics, Haemostatics and Acupan. 119 Alphamox.	Acetec48	Alfacalcidol	34		
ricinoleic acid 76 Alfamino Junior 274 Antiacne Preparations 6 Accl-yelseine 253 Alglinic acid 6 Antiallergy Preparations 238 Aci-Jel 76 Alguosoidase alfa 27 Antianaemics 38 Aciclovir Alkeran 146 Antiandrogen Oral Contraceptives 76 Sensory 248 Allerfix 250 Antiarrhythnics 56 Acidex 6 Allerpro Syneo 1 277 Antiarrhythnics 56 Acimon 56 Allerpro Syneo 2 277 Antibacterials 99 Actima 69 Allersoothe 240 Anticholinesterases 111 Actemra 220 Allersoothe 240 Anticholinesterases 111 Actemra S29 220 Allopurinol 115 Antidepressants 122 Actrapid 10 Alpha-Adrenoceptor Blockers 48 Anticilierrhoeals 6 Actrapid Penfill 10 Alphamox 93 A		Alfamino	274	Anthelmintics	90
Acetylcysteine. 253 Alginic acid. .6 Antiallergy Preparations .236 Aci-lel. .76 Alglucosidase alfa. .27 Antianaemics .33 Aciclovir .14e Antiandrogen Oral .26 Infection .102 Alkeran S29 .14e Antiandrogen Oral Acidex .6 Allerfix .250 Antiarrhythmics .56 Acidex .6 Allerpro Syneo 1 .277 Antibacterials .90 Actimic .69 Allersoothe .240 Anticholinergic Agents .242 Acteman .220 Allerpro Syneo 2 .277 Antibacterials Topical .66 Acterania .220 Allerpro Syneo 2 .277 Antibacterials Topical .66 Acterania .220 Allerpro Syneo 2 .277 Antibacterials Topical .66 Acterania .220 Allerpro Syneo 2 .277 Anticholinersterases .11 Acterania .20 Allerpro Syneo 2 .277 Anticholinersterases					
Aci-el. 76 Alglucosidase alfa. 27 Antianaemics. 36 Aciclovir Alkeran 146 Antiandrogen Oral Infection 102 Alkeran S29 146 Contraceptives. 77 Sensory. 248 Allerix 250 Antiarrhythmics. 56 Acidex. 6 Allerpro Syneo 1 277 Antiabacterials Topical 66 Acipimox. 56 Allerpro Syneo 2 277 Antibacterials Topical 66 Acipimox. 56 Allerpro Syneo 2 277 Anticholinergic Agents 244 Actemra. 220 Allopurinol 115 Anticholinergic Agents 244 Actemra S29. 220 Allopurinol 115 Anticholinergic Agents 122 Actinomycin D. 151 Alpha-Adrenoceptor Blockers 48 Anticholinergic Agents 122 Actrapid 10 Alpha-Keri Lotion 67 Antieplepsy Drugs 122 Actrapid Penfill. 10 Alphamox 25 93 Antifibrin		Alginic acid	6	•	
Aciclovir Alkeran 146 Antiandrogen Oral Infection 102 Alkeran S29 146 Contraceptives 76 Sensory 248 Allerfix 250 Antiarrhythmics 56 Acidex 6 Allerpro Syneo 1 277 Antibacterials 99 Acipimox 56 Allerpro Syneo 2 277 Antibacterials Topical 66 Actimox 69 Allersoothe 240 Anticholinerstic Agents 244 Actemra 220 Allerocap 148 Anticholinerstic Agents 244 Actemra S29 220 Alloprinol 115 Antidepressants 122 Actrapid 10 Alpha-Adrenoceptor Blockers 48 Antidiarrhoeals 6 Actrapid Penfill 10 Alphamox 93 Antifiprolytics, Haemostatics and Acupan 119 Alphamox 93 Antifiprolytics, Haemostatics and Acupan 119 Alphamox 250 93 Antifiprolytics, Haemostatics and Adalimumab (ŭ .			
Infection	Aciclovir				
Sensory. 248 Allerfix. 250 Antiarrhythmics. 56 Acidex. .6 Allerpro Syneo 1 277 Antibacterials .99 Acipimox. .56 Allerpro Syneo 2 277 Antibacterials Topical .66 Actiretin. .69 Allersoothe. .240 Anticholinergic Agents .244 Actemra .220 Allmercap. .148 Anticholinergic Agents .244 Actemra S29. .220 Allopurinol .115 Anticholinesterases .117 Actinomycin D. .151 Alpha-Adrenoceptor Blockers .48 Antidiarrhoeals .6 Actrapid .10 Alpha-Adrenoceptor Blockers .48 Antidiarrhoeals .6 Actrapid Penfill. .10 Alpha-Keri Lotion .67 Antifiprinolytics, Haemostatics and Acupan .19 Alphamox 125 .93 Local Sclerosants .33 Adalimumab (Amgevita) .175 Alphamox 250 .93 Antifipriortics .244 Adalaimumab (Humira - Alternative	Infection102				76
Acidex 6 Allerpro Syneo 1 2777 Antibacterials 96 Aciprimox .56 Allerpro Syneo 2 277 Antibacterials Topical .66 Acitretin .69 Allerscothe .240 Anticholinergic Agents .242 Actemra .220 Allmercap .148 Anticholinesterases .111 Actemra S29 .220 Allopurinol .115 Anticholinesterases .112 Actrapid .10 Alpha-Adrenoceptor Blockers .48 Antidiarrhoeals .6 Actrapid Penfill .10 Alpha-Keri Lotion .67 Antipilepsy Drugs .122 Actrapid Penfill .10 Alphamox .23 Antifibriotytics, haemostatics and Acupan .119 Alphamox 125 .93 Antifibriotytics, haemostatics and Local Sclerosants .33 Antifibriotytics, haemostatics and Local Sclerosants .33 Antifibriotytics, haemostatics and Local Sclerosants .33 Antifibriotytics, haemostatics .34 Adalaimumab (Algentia Ali	Sensory248	Allerfix	250		
Acipimox .56 Allerpro Syneo 2 277 Antibacterials Topical 66 Acitretin .69 Allersoothe .240 Anticholinergic Agents .242 Actemra .220 Allopurinol .151 Anticholinesterases .117 Acterna S29 .220 Allopurinol .115 Antidepressants .127 Actrapid .10 Alpha-Adrenoceptor Blockers .48 Antidiperessants .122 Actrapid Penfill .10 Alphamox .28 Antifibrinolytics, Haemostatics and .24 Adalimumab (Amgevita) .175 Alphamox .250 .33 Antifibrinolo		Allerpro Syneo 1	277		
Acitretin. 69 Allersoothe. 240 Anticholinergic Agents. 242 Actemra 220 Allmercap. 148 Anticholinesterases. 111 Actemra S29. 220 Allopurinol. 115 Antidepressants. 123 Actinomycin D. 151 Alpha-Adrenoceptor Blockers. 48 Antidepressants. 16 Actrapid 10 Alpha-Adrenoceptor Blockers. 48 Antidiarrhoeals. 6 Actrapid Penfill. 10 Alpha-Keri Lotion. 67 Antiepilepsy Drugs. 122 Actrapid Penfill. 10 Alphamox. 93 Antifibrinolytics, Haemostatics and Acupan. 119 Alphamox. 93 Antifibriolos. 48 Adalimumab (Humira - Alternative brand) 175 Alpanox. 93 Antifibrotics. 244 Adapalene. 63 Aluminium hydroxide. 6 Antifungals. 99 Adrapalene. 63 Aluminium hydroxide. 6 Antihinatianines. 238 Adrealine. 25	Acipimox56				
Actemra 220 Allmercap 148 Anticholinesterases 117 Actemra S29 220 Allopurinol 115 Antidepressants 122 Actrapid Domina 151 Alpha-Adrenoceptor Blockers 48 Antidepressants 67 Actrapid Penfill 10 Alpha-Adrenoceptor Blockers 48 Antifibrinolidics 67 Actrapid Penfill 10 Alphamox 93 Antifibrinolytics, Haemostatics and Acupan 119 Alphamox 293 Antifibrinolytics, Haemostatics and Acupan 119 Alphamox 125 93 Antifibrinolytics, Haemostatics and Adalimumab (Amgevita) 175 Alphamox 250 93 Antifibrinolytics 24 Adalimumab (Humira - Alternative brand) 184 Alu-Tab 6 Antifiungals 93 Adapalene 63 Aluminium hydroxide 6 Antifiungals 96 Adapalene 63 Aluminium hydroxide 6 Antifiprotalics 23 Adres (arbidge 1.8 25 Alvogen					
Actrapid					
Actrapid	Actemra S29220	•		Antidepressants	123
Actrapid 10 Alpha-Keri Lotion 67 Antiepilepsy Drugs 124 Actrapid Penfill 10 Alphamox 93 Antifibrinolytics, Haemostatics and Acupan 119 Alphamox 125 93 Local Sclerosants 35 Adalimumab (Amgevita) 175 Alphamox 250 93 Antifibrotics 245 Adalimumab (Humira - Alternative brand) Alprolix 39 Antifibrotics 245 Adapalene 63 Aluminium hydroxide 6 Antifingals Topical 66 Adapalene 63 Aluminium hydroxide 6 Antifingals Topical 66 Adrenaline 25 Alvogen 52 Antihypotensives 55 Adrenaline 40 Ambrisentan 59 Antimigarine Preparations 12 Cardiovascular 58 Ambrisentan Mylan 59 Antimigarine Preparations 12 Advatanan 66 Ambrisentan Viatris 59 Antimigarine Preparations 12 Adyanovate 41 Amgevita <t< td=""><td>Actinomycin D151</td><td>•</td><td></td><td></td><td></td></t<>	Actinomycin D151	•			
Actrapid Penfill	•				
Acupan 119 Alphamox 125 93 Local Sclerosants 36 Adalimumab (Amgevita) 175 Alphamox 250 93 Antifibrotics 245 Adalimumab (Humira - Alternative brand) Alprolix 39 Antifiprotics 245 Adapalene 63 Alu-Tab 6 Antifungals 95 Adapalene 63 Aluminium hydroxide 6 Antihistamines 236 Adrenaline 25 Alvogen 52 Antihypotensives 55 Adrenaline 4mantadine hydrochloride 117 Antimalarials 96 Cardiovascular 58 Ambrisentan 59 Antimigraine Preparations 127 Respiratory 238 Ambrisentan Mylan 59 Antipruritic Preparations 126 Advantan 66 Ambrisentan Viatris 59 Antipruritic Preparations 66 Advate 41 Amgevita 175 Antipruritic Preparations 66 Advate 41 Amgevita 175 Antipruritic Prep	Actrapid Penfill10	Alphamox	93		
Adalimumab (Amgevita) 175 Alphamox 250 93 Antifibrotics 243 Adalimumab (Humira - Alternative brand) 184 Alurab 39 Antifungals 97 Adapalene 63 Aluminium hydroxide 6 Antihistamines 233 ADR Cartridge 1.8 25 Alvogen 52 Antihypotensives 55 Adrenaline Amantadine hydrochloride 117 Antimalarials 98 Cardiovascular 58 Ambrisentan 59 Antimigraine Preparations 127 Respiratory 238 Ambrisentan Mylan 59 Antiportic Preparations 126 Advata 41 Amgevita 175 Antiportic Preparations 126 Advate 41 Amgevita 175 Antiportic Preparations 126 Adynovate 42 Amiloride hydrochloride with 175 Antiretrovirals 105 Afliuria Quad 100 Amiloride hydrochloride with 112 Antirematoid Agents 112 Afluria Quad Junior 101					
Adalimumab (Humira - Alternative brand)	Adalimumab (Amgevita)175			Antifibrotics	243
Adapalene .63 Aluminium hydroxide 6 Antihistamines 236 ADR Cartridge 1.8 .25 Alvogen .52 Antihypotensives .57 Adrenaline Amantadine hydrochloride .117 Antimalarials .96 Cardiovascular .58 Ambrisentan .59 Antimigraine Preparations .127 Respiratory .238 Ambrisentan Mylan .59 Antimigraine Preparations .128 Advata .66 Ambrisentan Viatris .59 Antipuritic Preparations .66 Advate .41 Amgevita .175 Antipsychotics .128 Adynovate .42 Amiloride hydrochloride with hydrochloride with Antiretrovirals .105 Afliitor .234 Amiloride hydrochloride with Antirematoid Agents .112 Afluria Quad furosemide .55 Antithrombotic Agents .42 Afluria Quad hydrochlorothiazide .55 Antithrombotic Agents .42 Afluria Quad Junior Amicophylline .244 Antit				Antifungals	97
Adapalene .63 Aluminium hydroxide 6 Antihistamines 236 ADR Cartridge 1.8 .25 Alvogen .52 Antihypotensives .57 Adrenaline Amantadine hydrochloride .117 Antimalarials .96 Cardiovascular .58 Ambrisentan .59 Antimigraine Preparations .127 Respiratory .238 Ambrisentan Mylan .59 Antimigraine Preparations .128 Advata .66 Ambrisentan Viatris .59 Antipuritic Preparations .66 Advate .41 Amgevita .175 Antipsychotics .128 Adynovate .42 Amiloride hydrochloride with hydrochloride with Antiretrovirals .105 Afliitor .234 Amiloride hydrochloride with Antirematoid Agents .112 Afluria Quad furosemide .55 Antithrombotic Agents .42 Afluria Quad hydrochlorothiazide .55 Antithrombotic Agents .42 Afluria Quad Junior Amicophylline .244 Antit	brand) 184	Alu-Tab	6	Antifungals Topical	64
Adrenaline Cardiovascular 58 Ambrisentan 59 Antimalarials 99 Antimigraine Preparations 127 Antimigraine Preparations 127 Antimigraine Preparations 128 Ambrisentan Mylan 59 Antimigraine Preparations 128 Ambrisentan Mylan 59 Antimigraine Preparations 128 Antimigraine Preparations 128 Antimigraine Preparations 129 Antimigraine 129 Antimig	Adapalene	Aluminium hydroxide	6		
Adrenaline Cardiovascular 58 Ambrisentan 59 Antimalarials 99 Antimigraine Preparations 127 Antimigraine Preparations 127 Antimigraine Preparations 128 Ambrisentan Mylan 59 Antimigraine Preparations 128 Ambrisentan Mylan 59 Antimigraine Preparations 128 Antimigraine Preparations 128 Antimigraine Preparations 129 Antimigraine 129 Antimig	ADR Cartridge 1.825	Alvogen	52	Antihypotensives	51
Respiratory. 238 Ambrisentan Mylan. 59 Antinausea and Vertigo Agents. 126 Advantan 66 Ambrisentan Viatris 59 Antipruritic Preparations. 65 Advate 41 Amgevita. 175 Antipsychotics. 125 Adynovate 42 Amiloride hydrochloride 54 Antiretrovirals. 105 Afinitor 234 Amiloride hydrochloride with Antiretrovirals. 112 Aflibercept 190 furosemide 55 Antispasmodics and Other Agents Afluria Quad Amiloride hydrochloride with Altering Gut Motility 8 Afluria Quad Junior Aminophylline 244 Antithrombotic Agents 42 Afluria Quad Junior Aminophylline 244 Antithrymocyte globulin 42 (2023 formulation) 283 Amiodarone hydrochloride 50 (equine) 175 AFT-Pyrazinamide 101 Amisulpride 129 Antitrichomonal Agents 100 Agents Affecting the Amitriptyline 123 Antileprotics	Adrenaline	Amantadine hydrochloride	117		
Advantan 66 Ambrisentan Viatris 59 Antipruritic Preparations 66 Advate 41 Amgevita 175 Antipsychotics 125 Adynovate 42 Amiloride hydrochloride 54 Antiretrovirals 105 Afinitor 234 Amiloride hydrochloride with Antiretrovirals 112 Aflibercept 190 furosemide 55 Antispasmodics and Other Agents Afluria Quad Amiloride hydrochloride with Altering Gut Motility 8 Afluria Quad Junior Aminophylline 244 Antithrombotic Agents 42 Afluria Quad Junior Aminophylline 244 Antithrombotic Agents 42 AFT-Pyrazinamide 101 Amisulpride 50 (equine) 175 Agents Affecting the Amitriptyline 129 Antitrichomonal Agents 100 Agents Affecting the Amitriptyline 123 Antituberculotics and Antituberculotics Renin-Angiotensin System 48 Amlodipine 53 Antileprotics 100	Cardiovascular58	Ambrisentan	59	Antimigraine Preparations	127
Advate 41 Amgevita 175 Antipsychotics 126 Adynovate 42 Amiloride hydrochloride 54 Antiretrovirals 105 Afinitor 234 Amiloride hydrochloride with Antiretrovirals 105 Aflibercept 190 furosemide 55 Antispasmodics and Other Agents Afluria Quad Amiloride hydrochloride with Altering Gut Motility 8 Afluria Quad Junior Aminophylline 244 Antithrombotic Agents 42 Afluria Quad Junior Aminophylline 244 Antithrombotic Agents 42 AFT-Pyrazinamide 101 Amisulpride 50 (equine) 175 Agents Affecting the Amitriptyline 129 Antithromonal Agents 100 Renin-Angiotensin System 48 Amlodipine 53 Antileprotics 100	Respiratory238	Ambrisentan Mylan	59	Antinausea and Vertigo Agents	128
Advate 41 Amgevita 175 Antipsychotics 126 Adynovate 42 Amiloride hydrochloride 54 Antiretrovirals 105 Afinitor 234 Amiloride hydrochloride with Antiretrovirals 105 Aflibercept 190 furosemide 55 Antispasmodics and Other Agents Afluria Quad Amiloride hydrochloride with Altering Gut Motility 8 Afluria Quad Junior Aminophylline 244 Antithrombotic Agents 42 Afluria Quad Junior Aminophylline 244 Antithrombotic Agents 42 AFT-Pyrazinamide 101 Amisulpride 50 (equine) 175 Agents Affecting the Amitriptyline 129 Antithromonal Agents 100 Renin-Angiotensin System 48 Amlodipine 53 Antileprotics 100	Advantan66	Ambrisentan Viatris	59	Antipruritic Preparations	65
Afinitor	Advate41	Amgevita	175		
Aflibercept	Adynovate42	Amiloride hydrochloride	54		
Aflibercept	Afinitor234	Amiloride hydrochloride with		Antirheumatoid Agents	112
(2023 formulation) 283 hydrochlorothiazide 55 Antithrombotic Agents 42 Afluria Quad Junior Aminophylline 244 Antithymocyte globulin 245 (2023 formulation) 283 Amiodarone hydrochloride 50 (equine) 175 AFT-Pyrazinamide 101 Amisulpride 129 Antitrichomonal Agents 100 Agents Affecting the Amitriptyline 123 Antituberculotics and Renin-Angiotensin System 48 Amlodipine 53 Antileprotics 100	Aflibercept190	furosemide	55		
Afluria Quad Junior Aminophylline 244 Antithymocyte globulin (2023 formulation) 283 Amiodarone hydrochloride 50 (equine) 175 AFT-Pyrazinamide 101 Amisulpride 129 Antitrichomonal Agents 100 Agents Affecting the Amitriptyline 123 Antituberculotics and Renin-Angiotensin System 48 Amlodipine 53 Antileprotics 100	Afluria Quad			Altering Gut Motility	8
Afluria Quad Junior Aminophylline 244 Antithymocyte globulin (2023 formulation) 283 Amiodarone hydrochloride 50 (equine) 175 AFT-Pyrazinamide 101 Amisulpride 129 Antitrichomonal Agents 100 Agents Affecting the Amitriptyline 123 Antituberculotics and Renin-Angiotensin System 48 Amlodipine 53 Antileprotics 100	(2023 formulation) 283	hydrochlorothiazide	55	Antithrombotic Agents	42
(2023 formulation) 283 Amiodarone hydrochloride 50 (equine) 175 AFT-Pyrazinamide 101 Amisulpride 129 Antitrichomonal Agents 100 Agents Affecting the Renin-Angiotensin System 48 Amiodipine 123 Antituberculotics and Amidiprotics 53 Antileprotics 100					
AFT-Pyrazinamide 101 Amisulpride 129 Antitrichomonal Agents 100 Agents Affecting the Renin-Angiotensin System 48 Amlodipine 53 Antileprotics 100 Antituberculotics and 100 Ant					175
Agents Affecting the Amitriptyline	AFT-Pyrazinamide101				
Renin-Angiotensin System	•				
	Renin-Angiotensin System 48	Amlodipine	53	Antileprotics	100
	Agents for Parkinsonism and Related	Amorolfine	64	Antiulcerants	9

Antivirals	101	Atnahs Olsalazine	8	Beta Cream	65
Anxiolytics	132	Atomoxetine	137	Beta Ointment	
Anzatax	154	Atorvastatin	56	Beta Scalp	71
Apidra	11	Atropine sulphate		Beta-Adrenoceptor Agonists	.241
Apidra SoloStar		Cardiovascular	50	Beta-Adrenoceptor Blockers	
APO-Atomoxetine	137	Sensory	251	Betadine	
APO-Atomoxetine S29	137	Atropt		Betadine Skin Prep	68
Apo-Azithromycin	91	Atrovent		Betaferon	134
Apo-Temozolomide		Aubagio		Betahistine dihydrochloride	
Apomorphine hydrochloride		Augmentin		Betaine	
Aprepitant		Aurorix		Betaloc CR	
Apresoline		AutoSoft 30		Betamethasone dipropionate	65
Aqueous cream		AutoSoft 90	24	Betamethasone dipropionate with	
Aratac		Avallon		calcipotriol	69
Arava		Avelox		Betamethasone sodium phosphate	
Arginine		Avonex		with betamethasone acetate	80
Aripiprazole		Avonex Pen		Betamethasone valerate6	
Aripiprazole Sandoz		Azacitidine		Betamethasone valerate with sodiur	,
Aristocort		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Arrotex-Prazosin S29		Azamun		Betaxolol	
Arrow - Clopid		Azathioprine		Betnovate	
Arrow - Lattim		Azilect		Betoptic	
		Azithromycin			
Arrow-Amitriptyline		•		Betoptic S	
Arrow-Bendrofluazide		Azopt		Bexsero	
Arrow-Brimonidine		AZT	107	Bezafibrate	
Arrow-Diazepam		-B-	40	Bezalip	
Arrow-Doxorubicin		B-D Micro-Fine		Bezalip Retard	
Arrow-Fluoxetine	124	B-D Ultra Fine		Bicalutamide	
Arrow-Losartan &		B-D Ultra Fine II		Bicillin LA	
Hydrochlorothiazide		Bacillus Calmette-Guerin (BCG	,	BiCNU	
Arrow-Norfloxacin		vaccine	175	Bile and Liver Therapy	
Arrow-Ornidazole		Bacillus Calmette-Guerin		Biltricide	
Arrow-Quinapril 10		vaccine		Bimatoprost	
Arrow-Quinapril 20		Baclofen		Bimatoprost Multichem	
Arrow-Quinapril 5	49	Bactroban	63	Binarex	165
Arrow-Roxithromycin	92	Balance		Binocrit	38
Arrow-Timolol	250	Barrier Creams and Emollients.	66	Biodone	121
Arrow-Topiramate	127	BCG Vaccine	280	Biodone Extra Forte	121
Arrow-Tramadol	122	Beclazone 100	240	Biodone Forte	121
Arsenic trioxide	150	Beclazone 250	240	Bisacodyl	27
Asacol	7	Beclazone 50	240	Bisacodyl Viatris	27
Ascend Aripiprazole	129	Beclomethasone dipropionate	240	Bisoprolol fumarate	51
Ascorbic acid		Bee venom allergy treatment	239	Bisoprolol Mylan	
Aspen Adrenaline	58	Bendamustine hydrochloride		Bisoprolol Viatris	51
Aspirin		Bendrofluazide		BK Lotion	67
Blood	42	Bendroflumethiazide		Bleomycin sulphate	
Nervous		[Bendrofluazide]	55	Blood Colony-stimulating	
Asthalin		Benralizumab		Factors	45
Atazanavir Mylan		Benzathine benzylpenicillin		Blood glucose diagnostic test	
Atazanavir sulphate		Benzatropine mesylate		meter	15
Atenolol		Benzbromaron AL 100		Blood glucose diagnostic test	10
Atenolol AFT		Benzbromarone		strip	15
Atenolol AFT S29		Benztrop		Blood glucose test strips (visually	13
Atezolizumab		Benzydamine hydrochloride		impaired)	15
ATGAM				Blood Ketone Diagnostic Test	10
Ativan		Benzylpenicillin sodium [Penicil		ů .	4.4
Auvall	133	G]	ყა	Strip	14

Bonjela	33	Calcium gluconate	35	Cetomacrogol with glycerol	67
Boostrix	280	Calcium Homeostasis		Cetomacrogol-AFT	67
Bortezomib	150	Calcium polystyrene sulphonat	e47	Cetuximab	192
Bosentan	59	Calcium Resonium		Charcoal	
Bosentan Dr Reddy's	59	Calogen	260	Chemotherapeutic Agents	144
Bplex	34	Candesartan cilexetil	49	Chickenpox vaccine	290
Breo Ellipta	241	Candestar	49	Chlorafast	
Brevinor 1/28	75	Canesten	64	Chlorambucil	145
Brevinor-1 28 Day	75	Capecitabine	147	Chloramphenicol	248
Bricanyl Turbuhaler	241	Capecitabine-DRLA		Chlorothiazide	
Brimonidine tartrate		Capercit		Chlorpheniramine maleate	239
Brimonidine tartrate with timolol		Capoten	48	Chlorpromazine hydrochloride.	
maleate	251	Capsaicin		Chlorsig	248
Brinzolamide	250	Musculoskeletal	112	Chlortalidone [Chlorthalidone].	
Brolene	248	Nervous	119	Chlorthalidone	
Brown & Burk	124	Captopril	48	Chlorvescent	
Brufen SR	111	Carafate		Choice 380 7med Nsha Silver/	copper
BSF Abacavir/Lamivudine		Carbaccord		Short	
Viatris	253	Carbamazepine		Choice Load 375	
BSF Ziextenzo		Carbimazole		Choice TT380 Short	
Buccastem		Carbomer		Choice TT380 Standard	
Budesonide		Carboplatin		Choline salicylate with cetalkor	
Alimentary	6	Carboplatin Ebewe		chloride	
Respiratory2		Carbosorb-X		Ciclosporin	
Budesonide with eformoterol		Cardinol LA		Cilazapril	
Bumetanide		Cardizem CD		Cilicaine VK	
Buprenorphine Naloxone BNM		CareSens Dual		Cinacalcet	
Buprenorphine with naloxone		CareSens N		Cipflox	
Bupropion hydrochloride		CareSens N POP		Ciprofloxacin	
Burel		CareSens N Premier		Infection	94
Burinex		CareSens PRO		Sensory	
Burinex S29		Carmellose sodium with gelatir		Ciprofloxacin Teva	
Buscopan		pectin		Cisplatin	
Buscopan S29		Carmustine		Cisplatin Ebewe	14
Buspirone hydrochloride		Carnitor		Citalopram hydrobromide	
Buspirone Viatris		Carvedilol		Cladribine	
Busulfan		Carvedilol Sandoz		Clarithromycin	177
- C -	140	Casirivimab and imdevimab		Alimentary	(
Cabergoline	88	Catapres		Infection	
Caffeine citrate		CeeNU		Clexane	
Calamine		Cefaclor monohydrate		Clexane Forte	4
Calamine-AFT		Cefalexin		Climara	
Calci-Tab 500		Cefazolin		Clindamycin	
Calcipotriol		Ceftriaxone		Clinicians	
Calcitonin		Ceftriaxone-AFT		Clinicians Renal Vit	
Calcitriol		Cefuroxime axetil		Clobazam	
Calcitriol-AFT		Celapram		Clobetasol propionate	
Calcium 500 mg Hexal		Celebrex		Clobetasone butyrate	
Calcium carbonate		Celecoxib		Clofazimine	100
Calcium Channel Blackers		Celecoxib Pfizer		Clomazol	0.
Calcium Channel Blockers		Celleant Chronodose		Dermatological	
Calcium Disodium Versenate		Cellcept		Genito-Urinary	
Calcium folinate		Centrally-Acting Agents		Clomifene citrate	89
Calcium Folinate Ebewe		Cephalexin ABM		Clomipramine hydrochloride	
Calcium Folinate Sandoz		Cetirizine hydrochloride		Clomipramine Teva	
Calcium Folinate Sandoz S29	147	Cetomacrogol	b/	Clonazepam	. 125, 133

Clonidine	53	Country Life	31	DBL Dacarbazine	150
Clonidine hydrochloride	54	Coversyl		DBL Desferrioxamine Mesylate	for Inj
Clonidine Teva		Creon 10000		BP	
Clopidogrel		Creon 25000	25	DBL Docetaxel	15 ⁻
Clopine		Creon Micro	25	DBL Ergometrine	7
Clopixol		Crotamiton		DBL Gemcitabine	
Clotrimazole		Crystaderm		DBL Gentamicin	
Dermatological	64	Curam		DBL Heparin Sodium	
Genito-Urinary		Curam Duo 500/125		DBL Leucovorin Calcium	14
Clozapine		Cvite		DBL Methotrexate Onco-Vial	
Clozaril		Cyclizine hydrochloride		DBL Pethidine Hydrochloride	
Co-trimoxazole		Cyclizine lactate		DBL Vincristine Sulfate	
Coal tar		Cyclogyl	251	Decozol	
Coal tar with allantoin, menthol		Cyclonex		Deferasirox	
phenol and sulphur		Cyclopentolate hydrochloride		Deferiprone	
Coal tar with salicylic acid and	70	Cyclophosphamide		Deferoxamine Pfizer S29	
	70			Denosumab	
sulphur		Cyclorin		Deolate	
Cobal-B12		Cycloserine			
Cobalin-H		Cyproterone acetateth	01	Deoxycoformycin	
Coco-Scalp	70	Cyproterone acetate with	70	Depo-Medrol	
Codeine phosphate	050	ethinyloestradiol		Depo-Provera	
Extemporaneous		Cystadane		Depo-Testosterone	8
Nervous		Cytarabine		Deprim	
Coenzyme Q10		Cytotec		Dermol	
Colchicine		Cytoxan	145	Desferrioxamine mesilate	
Colecalciferol		-D-		Desmopressin	88
Colestid		D-Penamine		Desmopressin acetate	88
Colestipol hydrochloride		Dabigatran		Desmopressin-PH&T	
Colgout		Dacarbazine		Desuric	11
Colifoam	7	Dacarbazine APP	150	Detection of Substances in	
Colistin sulphomethate	95	Dactinomycin [Actinomycin D]	151	Urine	78
Colistin-Link	95	Daivobet	69	Dexamethasone	
Collodion flexible	256	Daivonex	69	Hormone	80
Colloidal bismuth subcitrate		Daktarin	64	Sensory	249
Colofac	8	Dalacin C	95	Dexamethasone phosphate	80
Coloxyl	26	Dantrium	116	Dexamethasone with framycetin	and
Combigan	251	Dantrium S29	116	gramicidin	248
Compound electrolytes	47	Dantrolene	116	Dexamethasone with neomycin	
Compound electrolytes with glu	ıcose	Daonil		sulphate and polymyxin B	
[Dextrose]	47	Dapa-Tabs	55	sulphate	249
Compound hydroxybenzoate		Dapsone		Dexamfetamine sulfate	
Comtan		Daraprim		Dexmethsone	80
Concerta	139	Darunavir		Dextrochlorpheniramine	
Condoms		Darunavir Mylan		maleate	239
Condyline		Darunavir Viatris		Dextrose	
Contraceptives - Hormonal		Dasatinib		DHC Continus	
Contraceptives - Non-hormona		Daunorubicin		Diabetes	
Copaxone		Daunorubicin Zentiva		Diabetes Management	
Cordarone-X		David One Step Cassette Preg		Diacomit	
Corticosteroids and Related Ac		Test		Diagnostic Agents	
for Systemic Use	,	DBL Adrenaline		Diamide Relief	
Corticosteroids Topical		DBL Aminophylline		Diamox	250
Cortifoam		DBL Bleomycin Sulfate		Diasip	
Cosentyx		DBL Bortezomib		Diazepam	
		DBL Carboplatin		Diazoxide	
Cosmegen Coumadin		DBL Cisplatin		Dibenzyline	
Outriduiti	40	DDL 013414111	140	DINGHZYIII IC	40

			_
Diclofenac Sandoz111	Doxorubicin Ebewe151	EMLA	119
Diclofenac sodium	Doxorubicin hydrochloride151	Empagliflozin	14
Musculoskeletal111	Doxycycline94	Empagliflozin with metformin	
Sensory249	DP Lotion67	hydrochloride	14
Differin63	DP Lotn HC66	Emtricitabine	10
Difflam33	DP-Allopurinol115	Emtricitabine with tenofovir	
Diflucan97	Dr Reddy's Omeprazole9	disoproxil	104
Digestives Including Enzymes25	Drofate52	Emtriva	
Digoxin50	Drugs Affecting Bone	Emulsifying ointment	
Dihydrocodeine tartrate121	Metabolism 112	Emulsifying Ointment ADE	
Dilantin	Dual blood glucose and blood ketone	Enalapril maleate	4
Dilantin Infatab126	diagnostic test meter14	Enbrel	
Dilantin Paediatric	Dulaglutide12	Endocrine Therapy	
Diltiazem CD Clinect53	Dulcolax SP Drop27	Endoxan	14
Diltiazem hydrochloride53	Duocal Super Soluble Powder259	Engerix-B	
Dimethicone	Duolin242	Enlafax XR	
Dimethyl fumarate	Duolin HFA242	Enoxaparin sodium	
		Enstilar	
Dipentum	DuoResp Spiromax241		
Diphtheria, tetanus and pertussis	Duride	Ensure	
vaccine	Durvalumab230	Ensure Plus	
Diphtheria, tetanus, pertussis and	-E-	Ensure Plus HN	
polio vaccine	e-chamber La Grande246	Ensure Plus RTH	
Diphtheria, tetanus, pertussis, polio,	e-chamber Mask246	Ensure Two Cal HN RTH	
hepatitis B and haemophilus	e-chamber Turbo246	Entacapone	
influenzae type B vaccine 281	E-Mycin92	Entecavir	
Diprosone65	e5 Pharma10	Entecavir Mylan	
Diprosone OV65	Ear Preparations248	Entecavir Sandoz	
Dipyridamole42	Ear/Eye Preparations248	Entocort CIR	
Disopyramide phosphate50	Easiphen Liquid273	Entresto 24/26	
Disulfiram142	Econazole nitrate64	Entresto 49/51	4
Diuretics54	Efavirenz106	Entresto 97/103	49
Dizole97	Efavirenz with emtricitabine and	Entyvio	22
Docetaxel	tenofovir disoproxil107	Epilim	120
Docetaxel Accord151	Eformoterol fumarate240	Epilim Crushable	120
Docetaxel Sandoz151	Eformoterol fumarate dihydrate 240	Epilim IV	
Docusate sodium26	Eftrenonacog alfa [Recombinant	Epilim S/F Liquid	120
Docusate sodium with	factor IX]	Epilim Syrup	
sennosides26	Efudix71	Epipen	
Dolutegravir107	Egopsoryl TA70	Epipen Jr	
Domperidone128	Elaprase29	Epirubicin Ebewe	
Domperidone Viatris128	Eldepryl117	Epirubicin hydrochloride	
Donepezil hydrochloride140	Elecare	Eplerenone	
Donepezil-Rex140	Elecare LCP274	Epoetin alfa	
Dornase alfa244	Electral	Epoprostenol	
Dortimopt	Elelyso	Eptacog alfa [Recombinant factor	
Dorzolamide hydrochloride251	Elemental 028 Extra264	VIIa]	4
Dorzolamide with timolol	Elexacaftor with tezacaftor, ivacaftor	Erbitux	
Dostinex	and ivacaftor245	Ergometrine maleate	
Dosulepin [Dothiepin]	Elidel 70	Erlotinib	150
	Elocon	Erythrocin IV	
hydrochloride			
Dosulepin Mylan	Elocon Alcohol Free	Erythromycin (as lactobionate)	97
Dosulepin Viatris	Eltrowin 39	Erythromycin ethyl succinate	. 97
Dothiepin	Eltroxin	Esbriet	
Doxazosin	EMB Fatol	Escitalopram	
Doxazosin Clinect48	Emend Tri-Pack	Escitalopram (Ethics)	
Doxine94	Emicizumab40	Eskazole	90

Essential Ethosuximide	125	Felo 5 ER	53	Flupenthixol decanoate	13
Essential Generics	51	Felodipine	53	Flutamide	16
Essential Prednisolone	8	Femme-Tab ED	75	Flutamin	16
Estraderm MX	82	Fenpaed 100 mg per 5 ml	111	Fluticasone	24
Estradiol TDP Mylan	82	Fentanyl	121	Fluticasone furoate with	
Estradot		Fentanyl Sandoz	121	vilanterol	24
Estradot 50 mcg	82	Ferinject	36	Fluticasone propionate	24
Estrofem	82	Ferodan	36	Fluticasone with salmeterol	
Etanercept	168	Ferriprox	254	Flynn	9
Ethambutol hydrochloride	100	Ferro-F-Tabs	36	FML	25
Ethics Aspirin	119	Ferro-tab	35	Foban	6
Ethics Aspirin EC	42	Ferrograd	36	Folic acid	
Ethics Lisinopril	48	Ferrosig	36	Folic Acid multichem	
Ethinyloestradiol with		Ferrous fumarate		Folic Acid Mylan	3
desogestrel	74	Ferrous fumarate with folic acid	36	Folic Acid Viatris	3
Ethinyloestradiol with		Ferrous sulfate	36	Food Thickeners	27
levonorgestrel	75	Fexofenadine hydrochloride	239	Foods And Supplements For Int	oorn
Ethinyloestradiol with		Fibro-vein	42	Errors Of Metabolism	27
norethisterone	75	Filgrastim	45	Foradil	24
Ethosuximide	125	Finasteride	<mark>77</mark>	Forteo	
Etopophos	151	Fingolimod	134	Fortini	26
Etoposide	151	Firazyr	238	Fortini Multi Fibre	26
Etoposide phosphate		Flagyl	100	Fortisip	26
Etravirine		Flagyl-S		Fortisip Multi Fibre	
Eumovate	65	Flamazine	64	Fosamax	
Eurofolic	147	Flecainide acetate		Fosamax Plus	11
Evara	66-67	Flecainide BNM	50	Framycetin sulphate	24
Everet	126	Flecainide Controlled Release		Frebini Energy	
Everolimus	234	Teva	50	Frebini Energy Fibre	
Evista	113	Fleet Phosphate Enema	27	Frebini Original	
Evrysdi	136	Flixonase Hayfever & Allergy	246	Frebini Original Fibre	
Evusheld	220	Flixotide		Fresubin 2kcal HP	
Exemestane	168	Flixotide Accuhaler	240	Fresubin HP Energy	26
Exjade		Florinef		Fresubin HP Energy Fibre	26
Extemporaneously Compounded		Fluanxol		Fresubin Intensive	
Preparations and		Flucil	93	Fresubin Original	
Galenicals	256	Flucloxacillin	93	Fresubin Original Fibre	
Eye Preparations	248	Flucloxacillin-AFT		Frisium	
Eylea		Flucloxin		Frumil	5
Ezetimibe		Flucon	250	Frusemide	
Ezetimibe Sandoz	57	Fluconazole	97	Fucicort	6
Ezetimibe with simvastatin	58	Fludara Oral	148	Fucidin	
-F-		Fludarabine Ebewe	148	Fucithalmic	
Factor eight inhibitor bypassing		Fludarabine phosphate		Fulvestrant	16
fraction	41	Fludrocortisone acetate	80	Fungilin	3
Famotidine	9	Fluids and Electrolytes		Furosemid-Ratiopharm	
Famotidine Hovid	9	Flumetasone pivalate		Furosemide [Frusemide]	
Fasenra	191	Fluocortolone caproate with		Furosemide-Baxter	
Faslodex		fluocortolone pivalate and		fusidic acid	
Fatty Cream AFT		cinchocaine	8	Dermatological	64, 6
Febuxostat		Fluorometholone		Infection	
Febuxostat multichem		Fluorouracil		Sensory	24
Feed Thickener Karicare	-	Fluorouracil Accord		- G -	
Aptamil	271	Fluorouracil sodium		Gabapentin	12
FEIBA NF		Fluox		Gacet	12
Felo 10 ER		Fluoxetine hydrochloride		Galsulfase	
		-			

Galvumet	12	Haloperidol decanoate1	Hydrocortisone with cinchocaine	8
Galvus	12	Harvoni10	Hydrocortisone with miconazole	66
Gardasil 9	282	Havrix2	Hydrocortisone with natamycin and	
Gastrodenol	10	Havrix Junior2	neomycin	66
Gaviscon Extra Strength	6	Haylor syrup2	Hydrogen peroxide	60
Gaviscon Infant	6	healthE Dimethicone 10%	66 Hydroxocobalamin	34
Gazyva	202	healthE Dimethicone 4% Lotion	68 Hydroxocobalamin Panpharma	34
Gefitinib	159	healthE Dimethicone 5%		
GEM Aqueous Cream	67	healthE Glycerol BP2	56 Hydroxychloroquine	112
Gemcitabine Ebewe		healthE Urea Cream		
Gemcitabine hydrochloride	148	Healtheries Simple Baking Mix2	[hydroxycarbamide]	. 15
Gemtuzumab ozogamicin	192	Hemastix		
Genoptic		Hemlibra		
Gentamicin sulphate		Heparin sodium		
Infection	95	Heparin Sodium Panpharma		
Sensory	248	Heparinised saline		
Gilenya		Heparon Junior20		
Ginet		Hepatitis A vaccine2		
Glatiramer acetate		Hepatitis B recombinant	Hypromellose	
Glecaprevir with pibrentasvir		vaccine2		
Glibenclamide		Herceptin2		
Gliclazide		Hiberix2		9
Glipizide		Hiprex10		
Glivec		Histaclear2		
Glizide		Histafen2		
Glucagen Hypokit		Holoxan14	·	
Glucagon hydrochloride		Horleys Bread Mix2		
Glucerna Select		Horleys Flour2		
Glucose [Dextrose]		Hormone Replacement Therapy -	Ifosfamide	
Gluten Free Foods		Systemic		
Glycerin with sodium saccharin		HPV2		
Glycerin with sucrose		Humalog		
Glycerol		Humalog Mix 25	Inatinib mesilate	
Alimentary	27	Humalog Mix 50		
Extemporaneous		Human papillomavirus (6, 11, 16, 18,	Imbruvica	
Glyceryl trinitrate		31, 33, 45, 52 and 58) vaccine	Imfinzi	
Alimentary	8	[HPV]2		
Cardiovascular		Humatin	<u> </u>	
Glycopyrronium		Humira1		
Glycopyrronium bromide		HumiraPen1	•	
Glycopyrronium with		Humulin 30/70		
indacaterol	243	Humulin NPH		242
Go Healthy		Humulin R		
Gold Knight		Hyaluronic acid2		
Gold Knight XL		Hydralazine		
Goserelin		Hydralazine hydrochloride		
Gutron		Hydrocortisone	Infant Formulae	
Gynaecological Anti-infectives.		Dermatological		
- H -	70	Hormone		
	1/12	Hydrocortisone (PSM)		
Habitrol Haemophilus influenzae type E		Hydrocortisone acetate		
vaccine		Hydrocortisone acetate with	Inhaled Long-acting	241
Haldol		pramoxine hydrochloride		2//
Haldol Concentrate		Hydrocortisone and paraffin liquid	Inspra	
Haldol Decanoas		and lanolin		
Haloperidol		Hydrocortisone butyrate		
ι ιαιορ ε παοι	100	riyarocorusone bulyrale00,	1 1113uiii1 aspart	

Insulin aspart with insulin aspart	Isoptin	53	Lagevrio	10
protamine10	Isoptin Retard	53	Lamictal	12
Insulin glargine11	Isoptin SR	53	Lamivudine	102, 10
Insulin glulisine11	Isopto Carpine	251	Lamivudine Alphapharm	10
Insulin isophane11	Isosorbide mononitrate	58	Lamivudine Viatris	
Insulin isophane with insulin	Isosource Standard		Lamotrigine	12
neutral11	Isotretinoin	63	Lamprene	
Insulin lispro11	Ispaghula (psyllium) husk	26	Lanoxin	
Insulin lispro with insulin lispro	Itch-Soothe		Lanoxin Paediatric Elixir	
protamine11	Itraconazole		Lanoxin PG	
Insulin neutral10	Itrazole		Lanoxin S29	
Insulin pen needles16	Ivacaftor		Lansoprazole	
Insulin pump16	Ivermectin		Lantus	
Insulin pump cartridge21	- J -		Lantus SoloStar	
Insulin pump infusion set (steel	Jadelle	76	Lanvis	
cannula)21	Jakavi		Lanzol Relief	
Insulin pump infusion set (steel	Jardiamet		Lapatinib ditosylate	
cannula, straight insertion) 22	Jardiance		Largactil	12
Insulin pump infusion set (teflon	Jaydess		Laronidase	12
cannula)23	Jevity HiCal RTH		Lasix	
Insulin pump infusion set (teflon	Jevity Plus		Latanoprost	
cannula, angle insertion with	Jevity RTH Jinarc		Latanoprost with timolol	
insertion device)			Lax-Suppositories	
Insulin pump infusion set (teflon	Juno Pemetrexed	149	Lax-suppositories Glycerol	
cannula, angle insertion)		005	Laxatives	
Insulin pump infusion set (teflon	Kadcyla		Laxsol	
cannula, straight insertion with	Kalydeco		Ledipasvir with sofosbuvir	
insertion device)	Kemadrin		Leflunomide	
Insulin pump infusion set (teflon	Kenacomb		Lenalidomide	
cannula, straight insertion) 24	Kenacort A 10		Letrole	
Insulin pump reservoir25	Kenacort-A 40		Letrozole	
Insulin syringes, disposable with	Kenalog in Orabase		Leukeran FC	14
attached needle	Ketocal 3:1		Leukotriene Receptor	
Intelence	KetoCal 4:1	2/8	Antagonists	
Interferon beta-1-alpha134	Ketoconazole		Leuprorelin	
Interferon beta-1-beta134	Dermatological		Leustatin	
Intra-uterine device74	Infection		Levetiracetam	
Invega Sustenna131	Ketogenic Diet		Levetiracetam-AFT	
Invega Trinza132	Ketoprofen		Levlen ED	
IPCA-Frusemide54	KetoSens		Levocabastine	
IPCA-Metoprolol52	Keytruda		Levocarnitine	
IPCA-Propranolol52	Kindergen	262	Levodopa with benserazide	
IPOL289	Klacid		Levodopa with carbidopa	
Ipratropium bromide242, 246	Alimentary		Levomepromazine	13
Iressa159	Infection		Levomepromazine	
Irinotecan Actavis 100148	Kliogest		hydrochloride	13
Irinotecan hydrochloride148	Kliovance		Levonorgestrel	
Irinotecan-Rex148	Kogenate FS	42	Genito-Urinary	
Iron (as ferric carboxymaltose)36	Konakion MM		Hormone	
Iron polymaltose36	Konsyl-D		Levonorgestrel BNM	
lsentress107	Kuvan	31	Levothyroxine	8
Isentress HD107	-L-		Lidocaine [Lignocaine]	118–11
Ismo 2058	Labetalol		Lidocaine [Lignocaine]	
Ismo 40 Retard58	Lacosamide	125	hydrochloride	11
Isoniazid101	Lactulose	27	Lidocaine [Lignocaine] with	
Isoniazid with rifampicin101	Laevolac	27	chlorhexidine	11

Lidocaine [Lignocaine] with	Madopar 125117	Methotrexate Ebewe	149
prilocaine 119	Madopar 250117	Methotrexate Sandoz	149
Lidocaine-Baxter119	Madopar 62.5117	Methyl hydroxybenzoate	25
Life Extension32	Madopar HBS117	Methylcellulose	
Lignocaine118–119	Madopar Rapid117	Methylcellulose with glycerin and	
Lioresal Intrathecal116	Magnesium hydroxide36	sodium saccharin	25
Lipid-Modifying Agents56	Magnesium sulphate36	Methylcellulose with glycerin and	
Liquigen260	Mantoux290	sucrose	25
Liraglutide12	Marevan45	Methyldopa	
Lisinopril48	Marine Blue Lotion SPF 50+71	Methyldopa Mylan	
Litak	Martindale Pharma253	Methyldopa Mylan S29	
Lithium carbonate130	Mask for spacer device246	Methylnaltrexone bromide	
Livostin250	Maviret103	Methylphenidate ER - Teva	13
LMX4119	Maxidex249	Methylphenidate hydrochloride	13
Lo-Oralcon 20 ED75	Maxitrol249	Methylphenidate hydrochloride	
Locacorten-Viaform ED's248	MCT oil (Nutricia)260	extended-release	13
Local preparations for Anal and	Measles, mumps and rubella	Methylprednisolone	8
Rectal Disorders8	vaccine285	Methylprednisolone (as sodium	
Locasol274	Mebendazole90	succinate)	8 [.]
Locoid66, 71	Mebeverine hydrochloride8	Methylprednisolone aceponate	
Locoid Crelo66	Medrol80	Methylprednisolone acetate	
Locoid Lipocream66	Medroxyprogesterone acetate	Methylxanthines	
Locorten-Vioform248	Genito-Urinary76	Metoclopramide Actavis 10	
Lodoxamide250	Hormone83	Metoclopramide hydrochloride	
Logem125	Mefenamic acid111	Metolazone	
Lomide250	Melatonin134	Metopirone	
Lomustine	Melpha146	Metoprolol IV Mylan	
Loniten59	Melphalan146	Metoprolol IV Viatris	
Loperamide hydrochloride6	Menactra286	Metoprolol succinate	
Lopinavir with ritonavir107	Meningococcal (groups A, C, Y and	Metoprolol tartrate	
Lopinavir/Ritonavir Mylan107	W-135) conjugate vaccine 286	Metrogyl	
Loprofin273	Meningococcal B multicomponent	Metronidazole	
Loprofin Mix273	vaccine	Metyrapone	
Lorafix239	Meningococcal C conjugate	Mexiletine hydrochloride	
Loratadine239	vaccine287	Miacalcic	
Lorazepam133	MenQuadfi286	Micolette	
Lorstat	Menthol	Micolette-S29	2
Losartan Actavis	Mepolizumab201	Miconazole	
Losartan potassium49	Mercaptopurine148	Miconazole nitrate	
Losartan potassium with	Mercilon 2874	Dermatological	6
hydrochlorothiazide	Mesalazine7	Genito-Urinary	
Lovir	Mesna	Micreme	
Loxamine	Mestinon111	Micreme H	
Lucrin Depot 1-month88	Metabolic Disorder Agents27	Microgynon 20 ED	
Lucrin Depot 3-month88	Metformin hydrochloride12	Microgynon 30	7
Lyderm	Metformin Mylan12	Microlut	7
Lynparza153	Metformin Viatris12	Midazolam	
Lyrica	Methadone BNM121	Midazolam Mylan	
- M -	Methadone hydrochloride	Midazolam-Baxter	13
m-Eslon	Extemporaneous256	Midodrine	
Mabthera 205	Nervous121	Mifegyne	
Macro Organic Psyllium Husk26	Methenamine (hexamine)	Mifepristone	
Macrobid110	hippurate109	Milpharm	
Macrogol 3350 with potassium	Methopt252	Minerals	
chloride, sodium bicarbonate and	Methotrexate149	Mini-Wright AFS Low Range	
sodium chloride27	Methotrexate DBL Onco-Vial149	Mini-Wright Standard	
	ITU	Triigin Otaniaala	

Minidiab	11	Moments	73	Natulan	155
MiniMed 1.8 Reservoir		Mometasone furoate	66	Nausafix	129
MMT-326A	25	Monogen	261	Nausicalm	128
MiniMed 3.0 Reservoir		Montelukast	244	Navelbine	158
MMT-332A	25	Montelukast Mylan		Nefopam hydrochloride	119
MiniMed 770G		Montelukast Viatris		Neisvac-C	
MiniMed Mio MMT-921A		Moroctocog alfa [Recombinant fact		Neo-Mercazole	
MiniMed Mio MMT-923A		VIII]		Neocate Gold	
MiniMed Mio MMT-925A		Morphine hydrochloride		Neocate Junior Unflavoured	
MiniMed Mio MMT-941A		Morphine sulphate		Neocate Junior Vanilla	
MiniMed Mio MMT-943A		Motetis		Neocate SYNEO	
MiniMed Mio MMT-945A		Mouth and Throat		Neoral	
MiniMed Mio MMT-965A MiniMed Mio MMT-975A		Movapo		Neostigmine metilsulfate	
		Moxifloxacin		Nepafenac	
MiniMed Quick-Set MMT-386A		MSUD Maxamum	2/2	Nepro HP (strawberry)	
MiniMed Quick-Set MMT-387A		Mucilaginous laxatives with		Nepro HP (vanilla)	
MiniMed Quick-Set MMT-396A		stimulants		Nepro HP RTH	
MiniMed Quick-Set MMT-397A		Mucolytics		Neulactil	
MiniMed Quick-Set MMT-398A		Mucosoothe		Neuraxpharm	
MiniMed Quick-Set MMT-399A	23	Multiple Sclerosis Treatments	133	NeuroTabs	35
MiniMed Silhouette MMT-368A	23	Multivitamin renal	34	Nevirapine	106
MiniMed Silhouette MMT-377A	23	Multivitamins	35	Nevirapine Alphapharm	106
MiniMed Silhouette MMT-378A	23	Mupirocin	63	Nevirapine Viatris	106
MiniMed Silhouette MMT-381A	23	Muscle Relaxants	116	Nicorandil	59
MiniMed Silhouette MMT-382A	23	Mvite	35	Nicotine	142
MiniMed Silhouette MMT-383A	23	Myambutol	100	Nifedipine	
MiniMed Silhouette MMT-384A		Mycobutin		Nifuran	
MiniMed Sure-T MMT-864A		MycoNail		Nilotinib	
MiniMed Sure-T MMT-866A		Mycophenolate mofetil		Nilstat	
MiniMed Sure-T MMT-874A		Mydriacyl		Alimentary	33
MiniMed Sure-T MMT-876A		Mylan (12 hr release)		Genito-Urinary	
MiniMed Sure-T MMT-884A		Mylan (24 hr release)		Infection	
MiniMed Sure-T MMT-886A				Nintedanib	
		Mylan Atenolol			
Minims Produicalone		Mylan Clomiphen		Nipent	
Minims Prednisolone		Mylan Indapamide		Nirmatrelvir with ritonavir	
Minirin		Mylan Italy (24 hr release)		Nitrates	
Minirin Melt		Myleran		Nitroderm TTS	
Mino-tabs		Mylotarg	192	Nitrofurantoin	
Minocycline hydrochloride		Myometrial and Vaginal Hormone		Nitrolingual Pump Spray	
Minomycin		Preparations		Nivestim	
Minor Skin Infections		Myozyme	27	Nivolumab	
Minoxidil		- N -		Nizoral	
Mirena	83	Nadolol	52	Nodia	6
Mirtazapine		Nadolol BNM	52	Noflam 250	
Misoprostol	9	Naglazyme	29	Noflam 500	111
Mitomycin C		Naloxone hydrochloride	253	Non-Steroidal Anti-Inflammatory	
Mitozantrone	153	Naltraccord	142	Drugs	111
Mitozantrone Ebewe	153	Naltrexone hydrochloride	142	Nonacog gamma, [Recombinant	
Mixtard 30		Naphazoline hydrochloride		Factor IX]	41
MMR II		Naphcon Forte		Norethisterone	
Moclobemide		Naprosyn SR 1000		Genito-Urinary	76
Modafinil		Naprosyn SR 750		Hormone	
Modavigil		Naproxen		Norflex	
Moduretic		Narcaricin mite		Norfloxacin	
Molaxole		Nasal Preparations		Noriday 28	
Molnupiravir		Natalizumab		Norimin	75
1410111upiiavii	104	vald Zu au	104	[NOIHIHI]	10

Narimin 1 00 Day	75	Oaranina	104	Osmalita DTU	000
Normanal Dive		Octobra olfo [Basembinant fo		Osmolite RTH Other Endocrine Agents	
Normacol Plus		Octocog alfa [Recombinant fa			
Normison		VIII] (Advate)		Other Oestrogen Preparations	0
Norpress		Octocog alfa [Recombinant fa		Other Progestogen	0.
Nortriptyline hydrochloride		VIII] (Kogenate FS)		Preparations	
Norvir	107	Octreotide		Other Skin Preparations	
Noumed		Octreotide Depot Teva		Other Supplements for PKU	27
Dermatological		Octreotide GH		Ovestin	
Nervous		Octreotide long-acting		Genito-Urinary	
Noumed Paracetamol		Oestradiol		Hormone	
Noumed Pethidine		Oestradiol valerate		Oxaliplatin	
NovaSource Renal		Oestradiol with norethisterone	e83	Oxaliplatin Accord	140
Novatretin		Oestriol		Oxaliplatin Actavis 100	140
Novitium Sugar Free	30	Genito-Urinary	77	Oxaliplatin Ebewe	140
NovoMix 30 FlexPen	10	Hormone	83	Oxis Turbuhaler	240
NovoRapid	11	Oestrogens	82	Oxpentifylline	5
NovoRapid FlexPen	11	Ofev	243	Oxybutynin	7
NovoRapid Penfill	11	Oil in water emulsion	<mark>67</mark>	Oxycodone hydrochloride	12
NovoSeven RT		Olanzapine	130–131	Oxycodone Sandoz	12
Nozinan	130	Olaparib		OxyNorm	12
Nozinan (Swiss)	130	Olbetam		Oxytocin	
Nozinan S29		Olbetam S29	<u>56</u>	Oxytocin BNM	
Nucala		Olopatadine		Oxytocin GH	
Nuelin		Olopatadine Teva		Oxytocin with ergometrine	
Nuelin-SR		Olsalazine		maleate	7 ⁻
Nupentin		Omalizumab		Ozurdex	
Nusinersen		Omeprazole		- P -	
Nutilis		Omeprazole actavis 10		Pacifen	110
Nutren Diabetes		Omeprazole actavis 20		Pacimol	
Nutrient Modules		Omeprazole actavis 40		Paclitaxel	
Nutrini Energy Multi Fibre		Omnitrope		Paclitaxel Actavis	
Nutrini Energy RTH		Omnitrope S29		Paclitaxel Ebewe	
Nutrini Low Energy Multi Fibre		Onbrez Breezhaler		Paediatric Seravit	
Nutrini Peptisorb		Oncaspar LYO		Palbociclib	
		OncoTICE	175	Paliperidone	
Nutrini Peptisorb Energy					
Nutrini RTH	203	OndansetronOndansetron ODT-DRLA		Paliperidone palmitate Palivizumab	
Nutrison 800 Complete Multi	260				
Fibre		One-Alpha		Pamidronate disodium	
Nutrison Advanced Diason		One-Alpha S29		Pamisol	
Nutrison Advanced Peptisorb.		Onrex		Pamol	
Nutrison Concentrated		Opdivo		Pancreatic enzyme	
Nutrison Energy		Ora-Blend		Pantoprazole	
Nutrison Energy Multi Fibre		Ora-Blend SF		Panzop Relief	
Nutrison Multi Fibre		Ora-Plus		Papaverine hydrochloride	
Nutrison Standard RTH		Ora-Sweet		Para-amino salicylic acid	
Nyefax Retard	53	Ora-Sweet SF		Paracetamol	
Nystatin		Orabase		Paracetamol (Ethics)	
Alimentary		Oral and Enteral Feeds		Paracetamol + Codeine	
Genito-Urinary	76	Oralcon 30 ED		(Relieve)	
Infection		Oratane		Paracetamol with codeine	
NZB Low Gluten Bread Mix	271	Ordine		Paraffin	
-0-		Orgran		Paraffin liquid with wool fat	
Obinutuzumab		Ornidazole		Parasiticidal Preparations	6
Obstetric Preparations		Orphenadrine citrate		Parnate	12
Ocicure	9	Ortho-tolidine		Parnate S29	
Ocrelizumab	134	Oruvail SR	111	Paromomycin	9

Paroxetine	124	Phytomenadione	42	Prednisolone acetate	25
Paser	101	Pilocarpine hydrochloride	251	Prednisolone sodium	
Paxam	133	Pimafucort	66	Prednisolone sodium	
Paxlovid	105	Pimecrolimus	70	phosphate	25
Pazopanib	162	Pine tar with trolamine laurilsulfate		Prednisolone-AFT	
Peak flow meter	246	and fluorescein	70	Prednisone	8
Pedialyte - Bubblegum	47	Pinetarsol	70	Prednisone Clinect	8
Pediasure		Pioglitazone		Pregabalin	12
Pediasure Plus		Pirfenidone		Pregabalin Pfizer	
Pediasure RTH	263	Pizotifen	128	Pregnancy Tests - hCG Urine	<mark>7</mark>
Pegaspargase		PKU Anamix Infant	273	Premarin	
Pegasys		PKU Anamix Junior	273	Prevenar 13	28
Pegfilgrastim		PKU Anamix Junior Chocolate		Priadel	13
Pegylated interferon alfa-2a		PKU Anamix Junior LQ		Primaguine	9
Pembrolizumab		PKU Anamix Junior Orange		Primidone	
Pemetrexed		PKU Anamix Junior Vanilla		Primidone Clinect	
Penicillamine		PKU GMPro Ultra Lemonade		Primolut N	
Penicillin G		PKU Lophlex LQ 10		Priorix	
PenMix 30		PKU Lophlex LQ 20		Probenecid	
PenMix 50		PKU Lophlex Powder		Probenecid-AFT	
Pentasa		PKU Lophlex Sensation 20		Procarbazine hydrochloride	
Pentostatin [Deoxycoformycin]		•		Prochlorperazine	
Pentoxifylline [Oxpentifylline]		Plaquenil		Proctofoam	
Pontomon Junior	39	Pneumococcal (PCV10) conjugate	33		
Peptamen Junior		, , , ,	200	Proctosedyl	
Pepti-Junior		vaccine	200	Procyclidine hydrochloride	
Perhexiline maleate		Pneumococcal (PCV13) conjugate	000	Progesterone	
Pericyazine		vaccine	288	Proglicem	
Perindopril		Pneumococcal (PPV23)	000	Proglycem	
Periset		polysaccharide vaccine		Progynova	
Perjeta		Pneumovax 23		Prolia	
Permethrin		Podophyllotoxin		Promethazine hydrochloride	
Perrigo	71	Polaramine		Propafenone hydrochloride	
Pertuzumab	205	Poliomyelitis vaccine	289	Propamidine isethionate	
Peteha		Poloxamer	26	Propranolol	
Pethidine hydrochloride		Poly-Gel		Propylene glycol	25
Pevaryl		Poly-Tears	252	Propylthiouracil	
Pexsig	53	Poly-Visc	252	Prostacur	16
Pfizer Exemestane	168	Polycal	258	Protaphane	1
Pharmacy Health Sorbolene with		Polyethylene glycol 400 and		Protaphane Penfill	
Glycerin	67	propylene glycol	252	Protifar	26
Pharmacy Services	253	Ponstan	111	Protionamide	10
Pheburane	32	Posaconazole	98	Provera	8
Phenasen	150	Posaconazole Juno	98	Provera HD	8
Phenobarbitone	126	Potassium chloride	46-47	Psoriasis and Eczema	
Phenobarbitone sodium		Potassium citrate	78	Preparations	6
Extemporaneous	256	Potassium iodate	35	PTU	8
Nervous		Povidone iodine	68	Pulmicort Turbuhaler	
Phenoxybenzamine		Pradaxa	45	Pulmozyme	24
hydrochloride	48	Pramipexole hydrochloride		Puri-nethol	
Phenoxymethylpenicillin (Penicill		Pravastatin	56	Puria	
V)		Pravastatin Mylan		Puritan's Pride Vitamin	
Phenytoin sodium1	24. 126	Pravastatin Viatris		B-2 100 mg	3
Phillips Milk of Magnesia		Praziquantel		Pyrazinamide	10
Phlexy 10		Prazosin		Pyridostigmine bromide	
Phosphate Phebra		Pred Forte		Pyridoxine hydrochloride	
Phosphorus		Prednisolone		Pyridoxine multichem	
1 1103p1101u3	47	1 100111301011 0	0 1	i yiidoxiiie iiididileiii	o

Pyrimethamine	96	Risperdal Consta	132	Scalp Preparations	71
Pytazen SR	42	Risperidone	131–132	Scopoderm TTS	128
- Q -		Risperidone (Teva)		Sebizole	71
Quetapel	130	Risperon		Secukinumab	
Quetiapine		Ritalin	138	Sedatives and Hypnotics	
Quick-Set MMT-392		Ritalin LA		Seebri Breezhaler	
Quick-Set MMT-393		Ritonavir	107	Selegiline hydrochloride	
Quinapril		Rituximab (Mabthera)	205	Senna	
Quinapril with		Rituximab (Riximyo)		Senokot	
hydrochlorothiazide	49	Rivaroxaban		SensoCard	
Qvar		Rivastigmine		Serc	128
- R -		Rivastigmine Patch BNM 10.	140	Serenace	130
RA-Morph	121	Rivastigmine Patch BNM 5		Seretide	
Ralicrom		Rivotril		Seretide Accuhaler	
Raloxifene hydrochloride		Riximyo		Serevent	
Raltegravir potassium		RIXUBIS		Serevent Accuhaler	240
Ramipex		Rizamelt		Sertraline	
Ramipril		Rizatriptan		Setrona	
Ranbaxy-Cefaclor		RoActemra S29		Setrona AU	12/
Rapamune		Robinul		Sevredol	
Rasagiline	117	Ronapreve		Sex Hormones Non	122
Reandron 1000		Ropin		Contraceptive	81
Recombinant factor IX		Ropinirole hydrochloride		Shield XL	
Recombinant factor VIIa		Rosuvastatin		Shingles vaccine	
Recombinant factor VIII		Rosuvastatin Viatris		Shingrix	
Rectogesic		Rotarix		SII-Onco-BCG	17
Redipred	0	Rotavirus oral vaccine		Sildenafil	173
Relieve		Roxane		Silhouette MMT-373	
Relistor		Roxane-Propranolol		Siltuximab	
Remicade				Simvastatin	
Renilon 7.5		Roxithromycin		Simvastatin Mylan	
				Simvastatin Viatris	
Resonium-A		Rubifen SR			
Resource Beneprotein		Rugby Capsaicin Topical Cre		Sinemet	
Respigen		Musculoskeletal		Sinemet CR	
Respiratory Devices		Nervous		Sirolimus	
Respiratory Stimulants	247	Rurioctocog alfa pegol [Reco		Siterone	
Retinol palmitate		factor VIII]		Slow-Lopresor	
ReTrieve		Ruxolitinib		Smith BioMed Rapid Pregnancy	
Retrovir		Rythmodan		Test	
Revlimid		Rythmodan - Cheplafarm		Sodibic	
Revolade		Rytmonorm	51	Sodium acid phosphate	
Riboflavin		-\$-		Sodium alginate	
Ribomustin		Sabril		Sodium benzoate	31
Ricit		Sacubitril with valsartan		Sodium bicarbonate	
Rifabutin		Sagent	158	Blood	
Rifadin	101	SalAir		Extemporaneous	256
Rifampicin		Salazopyrin	8	Sodium calcium edetate	254
Rifaximin	10	Salazopyrin EN	8	Sodium chloride	
Rifinah	101	Salbutamol		Blood	46
Rilutek	118	Salbutamol with ipratropium		Respiratory	246
Riluzole		bromide	242	Sodium citrate with sodium laur	
RINVOQ	236	Salicylic acid	70	sulphoacetate	27
Riodine		Salmeterol	240	Sodium citro-tartrate	
Risdiplam	136	Sandomigran	128	Sodium cromoglicate	
Risedronate Sandoz		Sanofi Primaquine	99	Alimentary	8
Risedronate sodium		Sapropterin dihydrochloride		Sensory	250

Sodium Fusidate [fusidic acid]		Sunscreens, proprietary	71	Tenofovir Disoproxil Mylan	102
Dermatological	64	Sure-T MMT-863	21	Tenofovir Disoproxil Viatris	
Infection		Sure-T MMT-873	21	Tenoxicam	
Sensory		Survimed OPD	265	Tensipine MR10	
Sodium hyaluronate [Hyaluronic		Sustagen Hospital Formula		Tepadina	146
acid]		Sustagen Hospital Formula		Terbinafine	
Sodium phenylbutyrate		Active	268	Terbutaline sulphate	
Sodium picosulfate		Sustanon Ampoules		Teriflunomide	
Sodium polystyrene sulphonate		Sylvant		Teriparatide	
Sodium tetradecyl sulphate		Symbicort Turbuhaler 100/6		Testosterone	
Sodium valproate		Symbicort Turbuhaler 200/6		Testosterone cipionate	
Sofradex		Symbicort Turbuhaler 400/12		Testosterone esters	
Soframycin		Symmetrel		Testosterone undecanoate	
Solgar		Sympathomimetics		Tetrabenazine	
		Synacthen		Tetrabromophenol	
Solifenacin Mylan Solifenacin succinate				Tetracosactrin	
		Synacthen Depot		Tetracycline	
Solifenacin Viatris		Synacthene Retard			
Solu-Cortef		Synagis		Teva Lisinopril	47
Solu-Medrol		Synflorix		Thalidomide	
Solu-Medrol-Act-O-Vial		Synthroid		Thalomid	
Somatropin (Omnitrope)		Syntometrine		Theophylline	
Sotalol		Syrup (pharmaceutical grade)	257	Thiamine hydrochloride	
Spacer device		Systane Unit Dose	252	Thiamine multichem	
Span-K		-T-		THIO-TEPA	
Spinal Muscular Atrophy		Tacrolimus		Thioguanine	
Spinraza		Dermatological		Thiotepa	
Spiolto Respimat		Oncology		Thyroid and Antithyroid Agents	
Spiractin		Tacrolimus Sandoz		Ticagrelor	
Spiriva		Taliglucerase alfa		Ticagrelor Sandoz	
Spiriva Respimat		Tambocor		Tilcotil	
Spironolactone		Tamoxifen citrate	167	Timolol	
Sporanox	98	Tamoxifen Sandoz	167	Timoptol XE	
Sprycel	158	Tamsulosin hydrochloride	77	Tiotropium bromide	242
Stelara	226	Tamsulosin-Rex	77	Tiotropium bromide with	
Stemetil	129	Tandem Cartridge		olodaterol	243
Steril-Gene	82	Tandem t:slim X2 with Basal-IQ	16	Tivicay	107
SteroClear	246	Tap water	257	Tixagevimab with cilgavimab	220
Stesolid	124	Taro	98	TMP	9
Stimulants/ADHD Treatments	137	Taro-Testosterone	81	Tobramycin	
Stiripentol	126	Tasigna	161	Infection	9
Stocrin	106	Tasmar	117	Sensory	249
Stomahesive	33	Taurine	32	Tobramycin BNM	
Strattera	137	Tecentrig	229	Tobramycin Mylan	9
Strides Shasun	98	Tecfidera	134	Tobrex	
Stromectol	68	Tegretol	125	Tocilizumab	220
Sucralfate	10	Tegretol CR		Tofranil	123
Sulfadiazine Silver	64	Telfast		Tolcapone	
Sulfadiazine sodium		Teligent		Tolvaptan	
Sulfasalazine		Temaccord		Topamax	
Sulphur		Temazepam		Topical Products for Joint and	
Sulprix		Temozolomide		Muscular Pain	119
Sumagran		Tenofovir disoproxil		Topiramate	
Sumatriptan		Tenofovir Disoproxil Emtricitabine		Topiramate Actavis	
Sunitinib		Mylan	104	Total parenteral nutrition (TPN)	اكا 14
Sunitinib Pfizer		Tenofovir Disoproxil Emtricitabine	104	TPN	
Sunscreens		Viatr	104	Tramadol hydrochloride	
Out 1001 CC1 10	/ 1	v Iau	104	riamauoi nyurotillollue	124

Tramal SR 100	122	Urinary Agents	77	Vildagliptin with metformin	
Tramal SR 150	122	Urinary Tract Infections	109	hydrochloride	12
Tramal SR 200	122	Urinorm	115	Vimpat	125
Trandate	52	Uromitexan	153	Vinblastine sulphate	157
Tranexamic acid	42	Ursodeoxycholic acid	25	Vincristine sulphate	157
Tranylcypromine sulphate	123	Ursosan	25	Vinorelbine	
Trastuzumab		Ustekinumab		Vinorelbine Ebewe	
Trastuzumab emtansine		Utrogestan	83	Vinorelbine Te Arai	158
Travatan	251	- V -		Viramune Suspension	106
Travoprost	251	Vaccinations	280	ViruPOS	248
Treatments for Dementia		Vaclovir	102	Vit.D3	34
Treatments for Substance		Valaciclovir	102	Vita-B12	34
Dependence	141	Valganciclovir	102	VitA-POS	252
Trental 400		Valganciclovir Mylan	102	Vitabdeck	35
Tretinoin		Valganciclovir Viatris		Vital	2 <mark>6</mark> 4
Dermatological	63	Vancomycin	97	Vitamin B complex	34
Oncology	156	Vannair	241	Vitamin B6 25	
Trexate	149	Varenicline Pfizer	142	Vitamins	34-35
Triamcinolone acetonide		Varenicline tartrate	142	Vivonex TEN	265
Alimentary	33	Varicella vaccine [Chickenpox		Voltaren	111
Dermatological	66	vaccine]	290	Voltaren D	111
Hormone	81	Varicella zoster vaccine [Shingles	3	Voltaren Ophtha	249
Triamcinolone acetonide with		vaccine]		Voltaren SR	
gramicidin, neomycin and ny	ystatin	Varicella zoster virus (Oka strain)	live	Volumatic	246
Dermatological	66	attenuated vaccine [shingles		Voriconazole	99
Sensory	248	vaccine]	290	Votrient	162
Triazolam	135	Various	253	Vttack	99
Trikafta	245	Varivax	290	- W -	
Trimethoprim	97	Vasodilators	59	Warfarin sodium	
Trimethoprim with		Vasopressin Agonists	88	Wart Preparations	71
sulphamethoxazole		Vasorex	53	Wasp venom allergy treatment	239
[Co-trimoxazole]	97	Vebulis	62	Water	
Trisequens	83	Vedafil	61	Blood	47
Trisul	97	Vedolizumab	227	Extemporaneous	257
Trophic Hormones	84	Veletri		White Soft Liquid Paraffin AFT	67
Tropicamide	252	Venclexta	156	Wool fat with mineral oil	67
Trulicity	12	Venetoclax	156	- X -	
Trusopt	251	Venlafaxine	124	Xarelto	45
TruSteel	<mark>22</mark>	Venomil	239	Xifaxan	10
Tryzan	49	VENOX	239	XMET Maxamum	272
Tuberculin PPD [Mantoux] test	290	Ventolin		Xolair	
Tubersol		Vepesid		XP Maxamum	273
Two Cal HN	270	Verapamil hydrochloride	53	Xylocaine	
Tykerb		Vermox	90	Xylocaine 2% Jelly	118
Tysabri	134	Versacloz	130	Xyntha	41
- U -		Vesanoid	156	- Z -	
UK Synacthen	81	Vexazone		Zapril	48
Ultibro Breezhaler	243	Vfend		Zarontin	125
Ultraproct	8	Viaderm KC	66	Zaroxolyn	55
Umeclidinium	242	Viatris		Zavedos	152
Umeclidinium with vilanterol	243	Cardiovascular	51	Zeffix	
Univent	242, 246	Infection	97, 107	Zematop	<mark>70</mark>
Upadacitinib	236	Victoza	12	Zetlam	102
Ural		Vigabatrin		Ziagen	106
Urea	67	Vigisom	134	Zidovudine [AZT]	107
Urex Forte	54	Vildagliptin		Zidovudine [AZT] with	

lamivudine	107
Ziextenzo	46
Zimybe	
Zinc and castor oil	66
Zinc sulphate	
Zincaps	37
Zinnat	
Ziprasidone	131
Zista	239
Zithromax	91
Zoledronic acid	
Hormone	
Musculoskeletal	115
Zoledronic acid Mylan	80
Zoledronic acid Viatris	
Hormone	
Musculoskeletal	
Zoledronic-US	115
Zopiclone	
Zopiclone Actavis	
Zostavax	
Zostrix	
Zostrix HP	
Zuclopenthixol decanoate	
Zuclopenthixol hydrochloride	
Zusdone	
Zyban	
Zypine	
Zypine ODT	
Zyprexa Relprevv	
7vtiga	165