# October 2023 Volume 30

#### **Editors:**

Kaye Wilson & Ayeshah Khan email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

#### Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at <a href="mailto:pharmac.govt.nz/subscribe">pharmac.govt.nz/subscribe</a>.

#### Production

Typeset automatically from XML and T<sub>E</sub>X. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

#### **Programmers**

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency

ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	ind oddonig i narmae i	
Section A	General Rules	5
Section B	Alimanton, Trast 9 Matabalian	
Section D	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	38
	Cardiovascular System	48
	Dermatologicals	71
	Genito Urinary System	80
	Hormone Preparations – Systemic	87
	Infections – Agents For Systemic Use	98
	Musculoskeletal System	119
	Nervous System	125
	Oncology Agents & Immunosuppressants	154
	Respiratory System & Allergies	249
	Sensory Organs	259
	Various	264
Section C	Extemporaneous Compounds (ECPs)	266
Section D	Special Foods	269
Section I	National Immunisation Schedule	291
	Index	302

Introducing Pharmac

# **Introducing Pharmac**

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

#### Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at <a href="https://www.pharmac.govt.nz/about">https://www.pharmac.govt.nz/about</a>.

# **Purpose of the Pharmaceutical Schedule**

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Te Whatu Ora Hospitals, as well as any access conditions that may apply;
- the Pharmaceuticals, including Medical Devices, used in Te Whatu Ora Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Te Whatu Ora Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Te Whatu Ora Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Te Whatu Ora Hospitals. Section H lists the Pharmaceuticals that that can be used in Te Whatu Ora Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

# **Explaining pharmaceutical entries**

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

#### Example



# Glossary

### **Units of Measure**

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

#### SECTION B: ALIMENTARY TRACT AND METABOLISM

SECTION B: ALIMENTARY TRACT AND ME	TABOLISM			
	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID  Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	<b>✓</b> G	aviscon Infant
* Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (13.61)	60	G	aviscon Extra Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 ml	A	cidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE  * Tab 600 mg  CALCIUM CARBONATE	12.56	100	<b>✓</b> A	lu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement		500 ml 473 ml		oxane alcium carbonate PAI 829
Only when prescribed for patients unable to swallow cale inappropriate and the prescription is endorsed according		ts or whe	ere calciur	n carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on * Tab 2 mg*  Cap 2 mg	10.75	400 400	✓ N ✓ <u>D</u>	odia iamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE  Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy	166.50	90	✓ E	ntocort CIR

# ⇒SA1886 Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
  - 2.1 Diabetes; or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation\*.

Note: Indication marked with \* is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis\*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
  - 3.1 Diabetes: or
  - 3.2 Cushingoid habitus; or
  - 3.3 Osteoporosis where there is significant risk of fracture: or
  - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
  - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
  - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
  - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with \* is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

### HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP	✓ Colifoam ✓ Cortifoam \$29
	21.1 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg	20	✓ Asacol
Suppos 1 a 50.96	28	✓ Pentasa

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
OLSALAZINE				
Tab 500 mg	56.02	60	✓	Atnahs
				Olsalazine S29
	93.37	100	1	Dipentum
Cap 250 mg	53.00	100	1	Dipentum
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential
				Prednisolone S29
SODIUM CROMOGLICATE				
Cap 100 mg	113.35	100	1	Ralicrom
SULFASALAZINE				
* Tab 500 mg	16.52	100	1	Salazopyrin
* Tab EC 500 mg	17.86	100		Salazopyrin EN

# Local preparations for Anal and Rectal Disorders

### **Antihaemorrhoidal Preparations**

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g11.06	30 g OP	<ul><li>Ultraproct</li></ul>
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg7.30	12	<ul><li>Ultraproct</li></ul>
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

# **Management of Anal Fissures**

GLYCERYL TRINITRATE – Special Authority see SA1329 below – Retail pharmacy

★ Oint 0.2%.......22.00 30 g OP

✓ Rectogesic

# ⇒SA1329 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

# **Antispasmodics and Other Agents Altering Gut Motility**

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available or	ıa		
PSO	19.00	5	✓ Robinul
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO	1.91	5	✓ Spazmol
	6.35		✓ Buscopan
			✓ Buscopan S29 S29

Spazmol to be Principal Supply on 1 December 2023

(Buscopan Inj 20 mg, 1 ml to be delisted 1 December 2023)

(Buscopan S29 S29 Inj 20 mg, 1 ml to be delisted 1 December 2023)

	ALIMENTARY	TR.	ACT AND	METABOLISM
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
IEBEVERINE HYDROCHLORIDE  Tab 135 mg  Colofac to be Principal Supply on 1 December 2023	8.50	90	✓ C	olofac
Antiulcerants				
Antisecretory and Cytoprotective				
IISOPROSTOL – Wastage claimable  Tab 200 mcg – Up to 120 tab available on a PSO	47.73	120	<b>✓</b> 0;	ytotec
Helicobacter Pylori Eradication				
ELARITHROMYCIN  Tab 500 mg – Subsidy by endorsement	ri eradication and presci			accordingly.
H2 Antagonists				
AMOTIDINE - Only on a prescription				
₹ Tab 20 mg	4.91	100		amotidine Hovid §29
÷ Tab 40 mg	10.32	100	<b>✓</b> Fa	amotidine Hovid §29
Inj 10 mg per ml, 4 ml — Subsidy by endorsement Subsidy by endorsement — Subsidised for patients rec		10 t of pa		ylan S29
Proton Pump Inhibitors				
ANSOPRAZOLE  Cap 15 mg  Cap 30 mg  MEPRAZOLE  For omeprazole suspension refer Standard Formulae, page	5.26	100 100	_	anzol Relief anzol Relief
Cap 10 mg	•	90		meprazole actavis 10
Cap 20 mg	2.02	90		meprazole actavis 20
€ Cap 40 mg	3.18	90		meprazole actavis 40
Powder – Only in combination  Only in extemporaneously compounded omeprazole:		5 g	<b>✓</b> M	idwest
Inj 40 mg ampoule with diluent	37.38	5		<u>r Reddy's</u> Omeprazole

Panzop Relief to be Principal Supply on 1 December 2023

\* Tab EC 40 mg .......2.74

Panzop Relief to be Principal Supply on 1 December 2023

**PANTOPRAZOLE** 

✓ Ocicure S29

✓ Panzop Relief

✓ Panzop Relief

90

90

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

<sup>\*</sup>Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully Brand or
	(Manufacturer's Price	Per	Subsidised Generic Manufacturer
Site Protective Agents			
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mg	14.51	50	✓ Gastrodenol S29
SUCRALFATE Tab 1 g	35.50 (48.28)	120	Carafate
Bile and Liver Therapy			
RIFAXIMIN – Special Authority see SA1461 below – Re Tab 550 mg	'	56	✓ Xifaxan
Initial application only from a gastroenterologist, hepat hepatologist. Approvals valid for 6 months where the patolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or	atient has hepatic encephalop  Practitioner on the recomme	athy de	espite an adequate trial of maximur
nepatologist. Approvals valid without further renewal ur benefiting from treatment.	less notified where the treath	nent rei	mains appropriate and the patient is
Diabetes			
Hyperglycaemic Agents			
DIAZOXIDE - Special Authority see SA1320 below - R			
Cap 25 mg		100	✓ Proglicem \$29
Cap 100 mg Oral liq 50 mg per ml		100 0 ml O	<ul> <li>✓ Proglicem \$29</li> <li>✓ Proglycem \$29</li> <li>✓ e5 Pharma \$29</li> </ul>
➤ SA1320 Special Authority for Subsidy  Initial application from any relevant practitioner. Appropriate Appropriate Appropriate Appropriate Approvals valid appropriate and the patient is benefiting from treatment.			d for the treatment of confirmed
GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSC	)32.00	1	✓ Glucagen Hypokit
Insulin - Short-acting Preparations			
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26 1	0 ml O	
▲ Inj human 100 u per ml, 3 ml	42.66	5	<ul><li>✓ Humulin R</li><li>✓ Actrapid Penfill</li><li>✓ Humulin R</li></ul>
Insulin - Intermediate-acting Preparations	s		
NSULIN ASPART WITH INSULIN ASPART PROTAMIN	NE		
▲ Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ NovoMix 30 FlexPen

	Subsidy		Fully Brand or
	(Manufacturer's Pr \$	ice) Subsi Per	dised Generic  ✓ Manufacturer
INSULIN ISOPHANE	· · · · · · · · · · · · · · · · · · ·		
▲ Inj human 100 u per ml	17.68	10 ml OP	✓ Humulin NPH
			✓ Protaphane
▲ Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH
			✓ Protaphane Penfill
INSULIN ISOPHANE WITH INSULIN NEUTRAL			•
▲ Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
▲ Inj human with neutral insulin 100 u per ml, 3 ml	12.66	5	<ul><li>✓ Mixtard 30</li><li>✓ Humulin 30/70</li></ul>
Inj numan with neutral insulin 100 u per mi, 5 mi	42.00	J	✓ PenMix 30
			✓ PenMix 50
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,			
3 ml	42.66	5	✓ Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,			_
3 ml	42.66	5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
msdim - Long-acting r reparations			
INSULIN GLARGINE			
▲ Inj 100 u per ml, 10 ml		1	✓ Lantus
▲ Inj 100 u per ml, 3 ml		5 5	<ul><li>✓ Lantus</li><li>✓ Lantus SoloStar</li></ul>
a inj roo u per mi, o mi disposable peri		J	- Lantus solostai
Insulin - Rapid Acting Preparations			
INSULIN ASPART			
▲ Inj 100 u per ml, 10 ml		1	✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
▲ Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
INSULIN GLULISINE  Inj 100 u per ml, 10 ml	27.02	1	✓ Apidra
▲ Inj 100 u per ml, 3 ml		5	✓ Apidra ✓ Apidra
▲ Inj 100 u per ml, 3 ml disposable pen		5	✓ Apidra SoloStar
INSULIN LISPRO			•
▲ Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ Humalog
▲ Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Alpha Glucosidase Inhibitors			
ACARBOSE			
* Tab 50 mg	8.95	90	✓ Accarb
* Tab 100 mg	15.29	90	✓ Accarb
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
* Tab 5 mg	7.50	100	✓ <u>Daonil</u>
GLICLAZIDE			
* Tab 80 mg	20.10	500	✓ Glizide
GLIPIZIDE			
* Tab 5 mg	4.58	100	✓ <u>Minidiab</u>

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg	14.74	1,000	<b>/</b>	Metformin Viatris
* Tab immediate-release 850 mg	11.28	500	1	Metformin Mylan
			✓	Metformin Viatris
(Metformin Mylan Tab immediate-release 850 mg to be delisted	l 1 January 2024)			
PIOGLITAZONE				
* Tab 15 mg	6.80	90	✓	Vexazone
* Tab 30 mg		90	✓	Vexazone
* Tab 45 mg		90	1	Vexazone
VILDAGLIPTIN				
Tab 50 mg	35.00	60	1	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	1	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride		60	1	Galvumet

### **GLP-1 Agonists**

DULAGLUTIDE - Special Authority see SA2065 below - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

#### ⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has type 2 diabetes; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is Māori or any Pacific ethnicity\*: or
    - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
    - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
    - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*, or
    - 2.2.5 Patient has diabetic kidney disease (see note b)\*; and
  - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

LIRAGLUTIDE - Special Authority see SA2187 on the next page - Retail pharmacy

- a) Maximum of 9 inj per prescription
- b)
- a) Not to be given in combination with a funded SGLT-2 inhibitor or other GLP-1 agonist.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

### ⇒SA2187 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Patient has previously received an initial Special Authority approval for either an SGLT-2 inhibitor or GLP-1 agonist; or
- 2 All of the following:
  - 2.1 Patient has type 2 diabetes; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is Māori or any Pacific ethnicity\*; or
    - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
    - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
    - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
    - 2.2.5 Patient has diabetic kidney disease (see note b)\*; and
  - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause.

#### SGLT2 Inhibitors

#### ⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Fither:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
  - 2.1 Patient has type 2 diabetes; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is Māori or any Pacific ethnicity\*; or
    - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
    - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*: or
    - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*: or
    - 2.2.5 Patient has diabetic kidney disease (see note b)\*; and
  - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

 a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack,

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 10 mg58.56	30	Jardiance
*	Tab 25 mg	30	<ul> <li>Jardiance</li> </ul>

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	8.56	60	✓ Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>

# **Diabetes Management**

# **Ketone Testing**

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes; or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

# **Dual Blood Glucose and Blood Ketone Testing**

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes; or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips.......20.00 1 OP ✓ CareSens Dual

				_
Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	sed	Generic	
\$	Per	•	Manufacturer	

# **Blood Glucose Testing**

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
  - 1) is receiving insulin or sulphonylurea therapy; or
  - 2) is pregnant with diabetes: or
  - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
  - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

### BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips26	6.20	50 test OP	SensoCard
-----------------------------	------	------------	-----------

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Insulin Syringes and Needles**

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

#### INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

IIV	octivi Livine Edeco — Maximum of 200 dev per prescription	1		
*	29 g × 12.7 mm	10.95	100	✓ B-D Micro-Fine
*	31 g × 5 mm	12.26	100	✓ B-D Micro-Fine
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	E – Maximum of 2	00 dev per p	orescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.56	100	B-D Ultra Fine II
	, ,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g x 8 mm needle	13.56	100	B-D Ultra Fine II
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
	, -	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	B-D Ultra Fine II
	•	1.36	10	
		(1.99)		B-D Ultra Fine II

# **Insulin Pumps**

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year pe	riod.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			Y2 with Rasal-IO

#### **⇒SA1603** Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
<b>\$</b>	Per	✓	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
  - 6.1 Applicant is a relevant specialist; or
  - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Fither:
  - 4.1 Applicant is a relevant specialist: or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

<del></del>			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
  - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 8.2 The pump is due for replacement; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

#### continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
  - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 4.2 The pump is due for replacement; and
- 5 Either:
  - 5.1 Applicant is a relevant specialist; or
  - 5.2 Applicant is a nurse practitioner working within their vocational scope.

### **Insulin Pump Consumables**

#### ⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
  - 6.1 Applicant is a relevant specialist; or
  - 6.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

#### All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Fither
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	I Generic
\$	Per 🗸	Manufacturer

continued...

- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

	ALIMENTAR	TRAC	ANL	DMETABOLISM
	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
continued				
pump therapy; and				
4 The patient is continuing to derive benefit from pump the				
5 The patient had achieved and is maintaining a HbA1c of				
6 The patient has had no increase in severe unexplained h			seline;	and
7 The patient's HbA1c has not deteriorated more than 5 m 8 Either:	moi/moi trom baseline	e; and		
8.1 Applicant is a relevant specialist; or				
8.2 Applicant is a nurse practitioner working within the	eir vocational scone			
Renewal — (Previous use before 1 September 2012) only fro	•	st or nurse	practiti	ioner. Approvals valid for 2
years for applications meeting the following criteria:	m a roioram oposiam		p.ao	
All of the following:				
1 The patient is continuing to derive benefit according to the	e treatment plan and	has mainta	ained a	HbA1c of equal to or less
than 80 mmol/mol; and				
2 The patient's HbA1c has not deteriorated more than 5 m				
<ul><li>3 The patient has not had an increase in severe unexplain</li><li>4 Either:</li></ul>	ed hypoglycaemic epi	sodes from	i baseli	ine; and
. —				
<ul><li>4.1 Applicant is a relevant specialist; or</li><li>4.2 Applicant is a nurse practitioner working within the</li></ul>	air vocational scope			
	•			
INSULIN PUMP CARTRIDGE – Special Authority see SA1985	on page 19 – Retail p	harmacy		
a) Maximum of 3 sets per prescription				
<ul><li>b) Only on a prescription</li><li>c) Maximum of 13 packs of cartridge sets will be funded per</li></ul>	rvoor			
Cartridge 300 U, t:lock × 10		1 OP	<b>√</b> 1	Fandem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special		-		•
a) Maximum of 3 sets per prescription	Additionly 300 OATSO	o on page	10 11	ciali priarriacy
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	<b>✓</b> I	MiniMed Sure-T
				MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	<b>✓</b> I	MiniMed Sure-T
				MMT-886A
6 mm steel needle; 60 cm tubing x 10	130.00	1 OP	<b>✓</b> I	MiniMed Sure-T
Commented mandles 00 and tables as 40	100.00	1 OD		MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	•	MiniMed Sure-T

8 mm steel needle; 80 cm tubing × 10 .......130.00 1 OP

8 mm steel needle; 60 cm tubing × 10 .......130.00

✓ MiniMed Sure-T MMT-874A ✓ MiniMed Sure-T

MMT-866A

6 mm steel needle; 29 G; manual insertion; 60 cm tubing x

MMT-876A

✓ Sure-T MMT-873

8 mm steel needle; 29 G; manual insertion; 60 cm tubing x

✓ Sure-T MMT-863 1 OP

1 OP

1 OP

(Sure-T MMT-863 6 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock to be delisted 1 December 2023)

(Sure-T MMT-873 8 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock to be delisted 1 December 2023)

Subsidy	H	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
 \$	Per	✓	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles		1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line × 10 with	130 00	1 OP	✓ TruSteel

1 OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

13 mm teflon needle, 45 cm tubing × 10130.00	1 OP
13 mm teflon needle, 60 cm tubing × 10130.00	1 OP

6 mm teflon needle, 80 cm blue tubing.......130.00

6 mm teflon needle, 80 cm clear tubing × 10 ......130.00

- ✓ MiniMed Silhouette MMT-382A
- ✓ MiniMed Silhouette MMT-368A
   ✓ MiniMed Silhouette
- MMT-381A

  MiniMed Silhouette

✓ MiniMed Silhouette

- MMT-383A ✓ MiniMed Silhouette MMT-377A
- MMT-378A ✓ MiniMed Silhouette MMT-384A
- 6 mm teflon needle, 110 cm tubing × 10
   10 P

   6 mm teflon needle, 45 cm blue tubing × 10
   130.00
   1 OP
- ✓ MiniMed Quick-Set MMT-398A
   ✓ MiniMed Mio MMT-941A
- ✓ MiniMed Mio MMT-921A
  - ✓ MiniMed Mio MMT-943A
  - ✓ MiniMed Mio MMT-923A
  - ✓ MiniMed Quick-Set MMT-399A
  - ✓ MiniMed Mio MMT-945A
  - ✓ MiniMed Mio MMT-965A
  - ✓ MiniMed Mio MMT-925A
  - ✓ MiniMed Quick-Set MMT-387A
  - ✓ MiniMed Quick-Set MMT-396A
  - ✓ MiniMed Quick-Set MMT-397A
  - ✓ MiniMed Mio MMT-975A
  - ✓ MiniMed Quick-Set MMT-386A

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. \*Three months or six months, as applicable, dispensed all-at-once

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION WITH INSERTION DEVICE) - Special Authority see SA1985 on page 19 - Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon cannula; angle insertion; insertion device; 110 cm line × 10 with 10 needles......140.00 1 OP ✓ AutoSoft 30 13 mm teflon cannula; angle insertion; insertion device: 60 cm line x 10 with 10 needles......140.00 1 OP ✓ AutoSoft 30 INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION) - Special Authority see SA1985 on page 19 -Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 17 mm teflon cannula; angle insertion; 60 cm line × 10 with 1 OP ✓ Silhouette MMT-373 (Silhouette MMT-373 17 mm teflon cannula: angle insertion: 60 cm line x 10 with 10 needles: luer lock to be delisted 1 December 2023) INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) - Special Authority see SA1985 on page 19 - Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device; ✓ AutoSoft 90 1 OP 6 mm teflon cannula: straight insertion: insertion device: 60 cm line × 10 with 10 needles......140.00 1 OP ✓ AutoSoft 90 9 mm teflon cannula: straight insertion: insertion device: 1 OP ✓ AutoSoft 90 9 mm teflon cannula; straight insertion; insertion device; 60 cm 1 OP ✓ AutoSoft 90 INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) - Special Authority see SA1985 on page 19 -Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 1 OP ✓ Quick-Set MMT-393 9 mm teflon cannula: straight insertion: 60 cm tubing × 10 with 1 OP ✓ Quick-Set MMT-392 (Quick-Set MMT-393 6 mm teflon cannula; straight insertion; 60 cm tubing x 10 with 10 needles; luer lock to be delisted 1 December 2023) (Quick-Set MMT-392 9 mm teflon cannula; straight insertion; 60 cm tubing x 10 with 10 needles; luer lock to be delisted 1

December 2023)

MMT-332A

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer
INSULIN PUMP RESERVOIR – Special Authority see SA1985 on a) Maximum of 3 sets per prescription	page 19 – Retail ph	narmacy		
b) Only on a prescription				
c) Maximum of 13 packs of reservoir sets will be funded per y	/ear.			
10 × luer lock conversion cartridges 1.8 ml for Paradigm pump	s50.00	1 OP	✓ A	DR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 10	50.00	1 OP	✓ N	liniMed
				1.8 Reservoir MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP	✓ N	liniMed 3.0 Reservoir

(MiniMed 1.8 Reservoir MMT-326A Cartridge for 5 and 7 series pump; 1.8 ml x 10 to be delisted 1 November 2023)

# **Digestives Including Enzymes**

#### PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase			
10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase			
25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase			
3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)	34.93	20 g OP	<ul><li>Creon Micro</li></ul>
URSODEOXYCHOLIC ACID - Special Authority see SA1739 below	- Retail pha	armacy	
Cap 250 mg	33.95	100	<ul><li>Ursosan</li></ul>

#### **⇒SA1739** Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

**Initial application** — (**Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IqM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

**Initial application** — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

#### Laxatives

# **Bulk-forming Agents**

ISP	AGHULA (PSYLLIUM) HUSK - Only on a prescription			
*	Powder for oral soln	6.00	250 g OP	✓ Macro Organic
			Ü	Psyllium Husk
		20.00	500 g OP	✓ Konsyl-D
/A A .	and Organia Bardirum Hradi Barralan fan anal aala ta bardalia	4-4 1 F-6 000	24)	-

(Macro Organic Psyllium Husk Powder for oral soln to be delisted 1 February 2024)

Faecal Softeners
------------------

DOCUSATE SODILIM - Only on a prescription

* Tab 50 mg  * Tab 120 mg		100 100	✓ Coloxyl ✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES Tab 50 mg with sennosides 8 mg	3.50	200	✓ <u>Laxsol</u>
POLOXAMER – Only on a prescription  Not funded for use in the ear.			
* Oral drops 10%	4.17	30 ml OP	✓ Coloxyl

# **Opioid Receptor Antagonists - Peripheral**

METHYLNALTREXONE BROMIDE - Special Authority s	see SA1691 below – Retail p	harmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	Relistor
,	246.00	7	✓ Relistor

#### **⇒SA1691** Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)	Ş	Fully Subsidised	Brand or Generic	
<b>\$</b>	Per	✓	Manufacturer	

continued...

- 1 The patient is receiving palliative care; and
- 2 Either:
  - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

#### **Osmotic Laxatives**

GLYCEROL  * Suppos 2.8/4.0 g - Only on a prescription	20	✓ <u>Lax-suppositories</u> <u>Glycerol</u>
LACTULOSE - Only on a prescription		
* Oral liq 10 g per 15 ml	500 ml	✓ Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AN Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg 8.50	ID SODIUM (	CHLORIDE  Molaxole
SODIUM ACID PHOSPHATE - Only on a prescription		
Enema 16% with sodium phosphate 8%	1	✓ Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE - Only on a pre	scription	
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,		
5 ml35.89	50	✓ <u>Micolette</u>
		✓ Micolette-S29 S29

### Stimulant Laxatives

BISACODYL – Only on a prescription			
* Tab 5 mg	5.80	200	<ul> <li>Bisacodyl Viatris</li> </ul>
* Suppos 10 mg		10	✓ Lax-Suppositories
SENNA - Only on a prescription			
* Tab, standardised	2.17	100	
	(8.21)		Senokot
	0.43	20	
	(2.06)		Senokot
SODIUM PICOSULFATE - Special Authority see SA2053 be	low – Retail pharma	асу	
Oral soln 7.5 mg per ml	7.40	30 ml OP	<ul> <li>Dulcolax SP Drop</li> </ul>

### ⇒SA2053 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

# **Metabolic Disorder Agents**

ALGLUCOSIDASE ALFA - Special Authority see SA1986 on the next page - Reta	ail pharmacy	
Inj 50 mg vial1,142.60	1	Myozyme

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

### ⇒SA1986 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
  - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
  - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
  - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
  - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE – Special Authorit	y see SA2042 below – Retail p	harmacy
-----------------------------	-------------------------------	---------

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg		50	✓ Solgar
Powder		400 g	✓ Biomed

#### ⇒SA2042 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE – Special Authority see SA1987 on the next page – Retail pl	harmacy		
Powder for oral soln	575.00	180 g OP	✓ Cystadane

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### ⇒SA1987 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
  - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
  - 2.2 A 5.10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
  - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below -	- Retail pharmacy		
Cap 120 mg	CBS	30	✓ Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

### ⇒SA2039 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

```
GALSULFASE – Special Authority see SA1988 below – Retail pharmacy
Inj 1 mg per ml, 5 ml vial.......2,234.00 1 ✓ Naglazyme
```

#### ⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

DURSULFASE - Special Authority see SA1623 on th	e next page - Retail pharmacy		
Inj 2 mg per ml, 3 ml vial	4,608.30	1	Elaprase

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# ⇒SA1623 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

#### ⇒SA1695 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Fither:
  - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below - Retail pharmacy Tab 500 mg .......CBS 30 ✓ Solgar 30 ✓ Solgar 60 ✓ Balance ✓ Carnitor S29 Oral lig 1 g per 10 ml ......CBS 118 ml ✓ Novitium Sugar Free S29 Oral lig 500 mg per 10 ml ......CBS 300 ml ✓ Balance

#### ⇒SA2040 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
RIBOFLAVIN – Special Authority see SA2041 below – Retail phar Tab 100 mg	•	100		Country Life Puritan's Pride Vitamin B-2 100 mg \$29
Cap 100 mg	CBS	100	1	Solgar

#### ⇒SA2041 Special Authority for Subsidy

**Initial application** only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

#### Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see SA1989 below – Retail pharmacy Tab soluble 100 mg .......1,452.70 30 OP 
✓ Kuvan

### ⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

#### 1 Fither:

- 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
- 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
  - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

# ⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

	(Manu	facturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer	
CODILIM BUENIVI BLITVDATE	Chariel Authority and CA1000 below	Dotail pharma	201/			

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 below - Retail pharmacy
Grans 483 mg per q......2,016,00 174 g OP 

✓ Pheburane

### ⇒SA1990 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retai	l pharmacy		
Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	✓ Life Extension

# **⇒SA2043** Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

#### Gaucher's Disease

### ⇒SA2137 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3\* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
  - 3.1 Patient has haematological complications of Gaucher disease; or
  - 3.2 Patient has skeletal complications of Gaucher disease; or
  - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
  - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
  - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with \* is an unapproved indication

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and

Subsidy	Fu	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
\$	Per	✓	Manufacturer

continued...

liver and spleen size; and

- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

# **Mouth and Throat**

# **Agents Used in Mouth Ulceration**

RENZYDAMINE HYDROCHI ORIDE

BENZYDAMINE HYDROCHLORIDE			
Soln 0.15% - Higher subsidy of \$21.73 per 500 ml with			
Endorsement	9.00	500 ml	
	(21.73)		Difflam
Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly.	oral mucositis	as a result of tr	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)		Orabase
	1.52	5 g OP	
	(3.60)	o g o.	Orabase
Powder	, ,	28 g OP	0145450
1 011001	(10.95)	20 g 0.	Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE	(10.00)		C.C.I.I.G.I.O.I.I.C
	2.06	15 a OB	
* Adhesive gel 8.7% with cetalkonium chloride 0.01%		15 g OP	Doniele
	(6.00)		Bonjela
TRIAMCINOLONE ACETONIDE			
Paste 0.1%	5.49	5 g OP	<ul><li>Kenalog in Orabase</li></ul>
Ovenhammacal Anti infactives			
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE		_0	. <del></del> g
	474	40 a OB	√ Decerel
Oral gel 20 mg per g	4./4	40 g OP	✓ <u>Decozol</u>
NYSTATIN			
Oral liq 100,000 u per ml	2.22	24 ml OP	✓ Nilstat

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	) Per	Subsidised	Generic Manufacturer
	Ψ	rei		Manuacturer
Vitamins				
Training				
Vitamin B				
HYDROXOCOBALAMIN				
* Inj 1 mg per ml, 1 ml ampoule - Up to 6 inj available on a PS	O2.46	3	✓	Cobal-B12 S29
			✓	Hydroxocobalamin
				Panpharma
			1	Vita-B12
	4.10	5	✓	Cobalin-H S29
PYRIDOXINE HYDROCHLORIDE				
a) No more than 100 mg per dose				
b) Only on a prescription				
* Tab 25 mg - No patient co-payment payable	3.43	90	1	Vitamin B6 25
* Tab 50 mg		500		Pyridoxine
				multichem
THIAMINE HYDROCHLORIDE - Only on a prescription				
* Tab 50 mg	4.65	100	1	Thiamine multichem
VITAMIN B COMPLEX			-	
* Tab, strong, BPC	11 25	500	1	Bplex
* Tab, Silong, Bro	11.20	300		phiex
Vitamin C				
ASCORBIC ACID				
a) No more than 100 mg per dose				
b) Only on a prescription				
* Tab 100 mg	12.50	500	✓	Cvite
Vitamin D				
ALFACALCIDOL			_	
* Cap 0.25 mcg		100		One-Alpha
* Cap 1 mcg	87.98	100		One-Alpha
				One-Alpha S29 S29
* Oral drops 2 mcg per ml	60.68 2	0 ml 0	OP 🗸	One-Alpha
CALCITRIOL			_	
* Cap 0.25 mcg		100		Calcitriol-AFT
* Cap 0.5 mcg	13.68	100		Calcitriol-AFT
COLECALCIFEROL				
* Cap 1.25 mg (50,000 iu) - Maximum of 12 cap per prescripti		12		Vit.D3
* Oral liq 188 mcg per ml (7,500 iu per ml)		.8 ml (		Puria
(D. via Onellia 400 man manual (7.500 in manual) ta ba deliste d.4 No.	3	5 ml O	P	Clinicians
(Puria Oral liq 188 mcg per ml (7,500 iu per ml) to be delisted 1 M	arcn 2024)			
Multivitamin Preparations				
MULTIVITAMIN RENAL – Special Authority see SA1546 on the r	nevt nage - Datail al	harma	CV	
* Cap		narma 30		Clinicians Renal Vit
~ Оар	0.43	30	•	Cillicialis Neliai VII

	ALIMENTA	RY TRA	CT AND	METABOLISM
(	Subsidy (Manufacturer's Prio \$	ce) S Per	Fully subsidised	Brand or Generic Manufacturer
■ SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Either:				0
<ol> <li>The patient has chronic kidney disease and is receiving eith</li> <li>The patient has chronic kidney disease grade 5, defined as</li> <li>ml/min/1.73 m² body surface area (BSA).</li> </ol>				
MULTIVITAMINS – Special Authority see SA1036 below – Retail p		200 g OF	· <b>✓</b> P	aediatric Seravit
■ SA1036   Special Authority for Subsidy   Initial application from any relevant practitioner. Approvals valid inborn errors of metabolism.   Renewal from any relevant practitioner. Approvals valid without full approval for multivitamins.   VITAMINS				
* Tab (BPC cap strength)	18.50	1,000	✓ <u>N</u>	lvite
* Cap (fat soluble vitamins A, D, E, K) – Special Authority see SA1720 below – Retail pharmacy	23.40	60	<b>✓</b> V	itabdeck
■ SA1720 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following:  1 Patient has cystic fibrosis with pancreatic insufficiency; or 2 Patient is an infant or child with liver disease or short gut sy 3 Patient has severe malabsorption syndrome.		newal unl	ess notifie	d for applications meeting
Minerals				
Calcium				
CALCIUM CARBONATE  * Tab 1.25 g (500 mg elemental)  * Tab eff 1.25 g (500 mg elemental) – Subsidy by endorsement		250 100	-	calci-Tab 500 calcium 500 mg Hexal \$29
Subsidy by endorsement – Only when prescribed for paed considered unsuitable.	liatric patients (<	5 years) v	vhere calc	
CALCIUM GLUCONATE  * Inj 10%, 10 ml ampoule	32.00	10	✓ N	lax Health - Hameln §29
	64.00	20	✓ N	lax Health S29
lodine				

POTASSIUM IODATE

### Iron

#### FERROUS FUMARATE

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price	) Sı	Fully ubsidised	Brand or Generic	
	\$	Per	•	Manufacturer	
FERROUS FUMARATE WITH FOLIC ACID  * Tab 310 mg (100 mg elemental) with folic acid 350 mcg FERROUS SULFATE	5.98	100	<b>√</b> <u>i</u>	Ferro-F-Tabs	
* Tab long-acting 325 mg (105 mg elemental)      * Oral liq 30 mg (6 mg elemental) per 1 ml		30 250 ml 500 ml	✓ Ē	Ferrograd Ferro-Liquid Ferodan	
IRON (AS FERRIC CARBOXYMALTOSE) — Special Authority so Inj 50 mg per ml, 10 ml vial		Retail pha		Ferinject	

#### ⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
  - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE  * Inj 50 mg per ml, 2 ml ampoule34.50	5	✓ Ferrosig
Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%33.60	355 ml	✓ Phillips Milk of Magnesia ©29
MAGNESIUM SULPHATE  * Inj 2 mmol per ml, 5 ml ampoule25.53	10	✓ Martindale

## **ALIMENTARY TRACT AND METABOLISM**

Subsidy	Fu	lly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	/	Manufacturer	

### **Antianaemics**

## Hypoplastic and Haemolytic

### ⇒SA2266 Special Authority for Subsidy

Initial application — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus; and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus; and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)\*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

**Renewal** — **(chronic renal failure)** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with \* is an unapproved indication

#### EPOETIN ALFA - Special Authority see SA2266 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	<ul><li>Binocrit</li></ul>
Inj 2,000 iu in 1 ml, syringe		6	<ul><li>Binocrit</li></ul>
Inj 3,000 iu in 0.3 ml, syringe		6	<ul><li>Binocrit</li></ul>
Inj 4,000 iu in 0.4 ml, syringe		6	<ul><li>Binocrit</li></ul>
Inj 5,000 iu in 0.5 ml, syringe		6	Binocrit
Inj 6,000 iu in 0.6 ml, syringe		6	<ul><li>Binocrit</li></ul>
Inj 8,000 iu in 0.8 ml, syringe		6	<ul><li>Binocrit</li></ul>
Inj 10,000 iu in 1 ml, syringe		6	<ul><li>Binocrit</li></ul>
Inj 40,000 iu in 1 ml, syringe		1	<ul><li>Binocrit</li></ul>

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID  * Tab 0.8 mg	26.60	1,000	<b>√</b> F	olic Acid multichem
* Tab 5 mg	5.82	100		olic Acid Mylan olic Acid Viatris
Oral liq 50 mcg per ml(Folic Acid Mylan Tab 5 mg to be delisted 1 January 2024)	28.82 2	5 ml OP	_	iomed

# Antifibrinolytics, Haemostatics and Local Sclerosants

#### EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

riodicio di odp ili confancion man allo ridalona ridoni	op.i.i.aa.i.agoo.ii. g. o	~p.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial		1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial		1	✓ Alprolix
Inj 4,000 iu vial		1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 below - Wastage claimable	- Retail pharmacy		
Tab 25 mg	1.550.00	28	✓ Revolade
Tab 50 mg	,	28	✓ Revolade

#### ⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
  - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
  - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  - 3.3 Patient has a platelet count of less than or equal to 10.000 platelets per microlitre.

**Initial application** — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
  - 3.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per

	Subsidy	Fully	Brand or
(Manu	acturer's Price)	Subsidised	Generic
	\$ Per		Manufacturer

continued...

microliter: or

3.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

**Initial application — (severe aplastic anaemia)** only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

#### EMICIZUMAB - [Xpharm] - Special Authority see SA2272 below

Inj 30 mg in 1 ml vial	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	· ·	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial	12,492.00	1	✓ Hemlibra
Inj 150 mg in 1 ml vial	17,846.00	1	✓ Hemlibra

### ⇒SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

	Subsidy		Fully	Brand or
(Mai	nufacturer's Price)	Subsi	dised	Generic
	\$	Per	✓	Manufacturer

#### EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	·	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Ini 8 mg syringe	9.426.40	1	✓ NovoSeven RT

### FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FEIBA NF
	2,630.00	1	✓ FEIBA NF
Ini 2.500 U		1	✓ FEIBA NF

#### MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	Xyntha
Inj 1,000 iu prefilled syringe		1	Xyntha
Inj 2,000 iu prefilled syringe		1	Xyntha
Inj 3,000 iu prefilled syringe		1	✓ Xyntha

#### NONACOG GAMMA. [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial		1	✓ RIXUBIS
Inj 2,000 iu vial		1	✓ RIXUBIS
Ini 3.000 iu vial	2.610.00	1	✓ RIXUBIS

#### OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [Xpharm]

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 250 iu vial	.210.00 1	✓ Advate
lnį 500 iu vial		✓ Advate
lnj 1,000 iu vial		✓ Advate
Inj 1,500 iu vial1	,260.00 1	✓ Advate
Inj 2,000 iu vial1	,680.00 1	✓ Advate
Inj 3,000 iu vial2		✓ Advate

### OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

casjoot to cittoria.			
Inj 250 iu vial	237.50	1	✓ Kogenate FS
Inj 500 iu vial		1	✓ Kogenate FS
Inj 1,000 iu vial	950.00	1	✓ Kogenate FS
Inj 2,000 iu vial	1,900.00	1	✓ Kogenate FS
Inj 3,000 iu vial	2,850.00	1	✓ Kogenate FS

41

	Subsidy (Manufacturer's Price) \$	Fu Subsidise Per	•
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]			
For patients with haemophilia A receiving prophylaxis treatm		d treatment is	managed by the Haemophilia
Treaters Group in conjunction with the National Haemophilia		4	/ A duma
Inj 250 iu vial			Adynovate
Inj 500 iu vial			Adynovate
Inj 1,000 iu vial			Adynovate
Inj 2,000 iu vial	2,400.00	1	Adynovate
SODIUM TETRADECYL SULPHATE			
* Inj 3% 2 ml	28.50	5	
	(73.00)		Fibro-vein
TRANEXAMIC ACID			
Tab 500 mg	10.45	60 •	Mercury Pharma
Vitamin K			
PHYTOMENADIONE			
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO	8.00	5	✓ Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO		-	Konakion MM

## **Antithrombotic Agents**

### Antiplatelet Agents

ASPIRIN	990	✓ Ethics Aspirin EC
CLOPIDOGREL	84	✓ Arrow - Clopid
DIPYRIDAMOLE  * Tab long-acting 150 mg13.93	60	✓ Pytazen SR
TICAGRELOR – Special Authority see SA1955 below – Retail pharmacy  * Tab 90 mg23.85	56	✓ <u>Ticagrelor Sandoz</u>

### ⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Either:
  - 1.1 Patient has had a neurological stenting procedure\* in the last 60 days; or
  - 1.2 Patient is about to have a neurological stenting procedure performed\*; and
- 2 Either:
  - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

continued...

- 2.2 Either:
  - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
  - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

**Initial application** — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

**Initial application — (Stent thrombosis)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

**Renewal — (Percutaneous coronary intervention with stent deployment)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

Notes: indications marked with \* are unapproved indications.

Note: \*\* Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

## **Heparin and Antagonist Preparations**

ENOXAPARIN SODIUM - Special Authority see SA2152 below - Retail pharmacy

Inj 20 mg in 0.2 ml syringe	31 28	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe		10	✓ Clexane
Inj 60 mg in 0.6 ml syringe		10	<ul><li>Clexane</li></ul>
Inj 80 mg in 0.8 ml syringe		10	Clexane
Inj 100 mg in 1 ml syringe		10	✓ Clexane
Inj 120 mg in 0.8 ml syringe		10	<ul> <li>Clexane Forte</li> </ul>
Ini 150 mg in 1 ml syringe		10	Clexane Forte

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 <b>`</b> \$	Per	✓	

continued...

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

**Initial application** — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
  - 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website\*; and
  - 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy: or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

#### HEPARIN SODIUM Inj 1,000 iu per ml, 5 ml ampoule ......86.11 50 ✓ Pfizer 10 ✓ Heparin Sodium Panpharma 5 ✓ DBL Heparin Sodium S29 ✓ Hospira 5 ✓ Hospira ✓ Heparin DBL S29 42.40 482.20 ✓ Heparin DBL S29 50 HEPARINISED SALINE Inj 10 iu per ml, 5 ml .......65.48 ✓ Pfizer 50 **Oral Anticoagulants DABIGATRAN** ✓ Pradaxa 60 ✓ Pradaxa 60 60 ✓ Pradaya

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per		Manufacturer
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	15.60	30	<b>✓</b> )	(arelto
Xarelto to be Principal Supply on 1 December 2023				
Tab 15 mg - Up to 14 tab available on a PSO	14.56	28	<b>✓</b> )	(arelto
Xarelto to be Principal Supply on 1 December 2023				
Tab 20 mg	14.56	28	<b>✓</b> )	(arelto
Xarelto to be Principal Supply on 1 December 2023				
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	✓ (	Coumadin
	6.46	100		Marevan
* Tab 2 mg	4.31	50	✓ (	Coumadin
* Tab 3 mg		100	✓ N	Marevan
* Tab 5 mg		50	✓ (	Coumadin
	11.48	100	✓ N	Marevan
			-	
Blood Colony-stimulating Factors				
Blood oblony damataling radiois				

FILGRAS'	TIM - Special Authority see SA1259 below - Retail	pharmacy		
Inj 30	0 mcg per 0.5 ml prefilled syringe	96.22	10	✓ Nivestim
Inj 48	0 mcg per 0.5 ml prefilled syringe	148.58	10	✓ Nivestim

#### ⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%\*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC  $< 0.5 \times 10^9$ /L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10<sup>9</sup>/L).

Note: \*Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

#### ⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%\*). Note: \*Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

## Fluids and Electrolytes

#### Intravenous Administration

#### GLUCOSE [DEXTROSE]

*	Inj 50%, 10 ml ampoule - Up to 5 inj available on a PSO34.75	5	<ul><li>Biomed</li></ul>
*	Inj 50%, 90 ml bottle - Up to 5 inj available on a PSO	1	<ul><li>Biomed</li></ul>

	Subsidy (Manufacturer's Price \$	e) S Per	Fully Subsidised	
POTASSIUM CHLORIDE				
Inj 75 mg per ml, 10 ml	65.00	50	1	Juno
SODIUM BICARBONATE				
Inj 8.4%, 50 ml	22.40	1	1	Biomed
a) Up to 5 inj available on a PSO				
b) Not in combination				
Inj 8.4%, 100 ml	22.95	1	1	Biomed
a) Up to 5 inj available on a PSO     b) Not in combination				
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Not funded for nebulis for nebuliser use.	•	used in c	onjunctio	n with an antibiotic intende
Inj 0.9%, bag - Up to 2000 ml available on a PSO	1.33	500 ml	1	Baxter
	1.36	1,000 m		Baxter
Only if prescribed on a prescription for renal dialysis, n for emergency use. (500 ml and 1,000 ml packs)				
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	/	Biomed
For Sodium chloride oral liquid formulation refer Stand				
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20		Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO.		50		Fresenius Kabi Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	•	rresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)	200			
				TPN
Infusion NATER		1 OP	•	
	when on the same fo	rm as an	ı injection	listed in the Pharmaceution
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of each of the dilution of sodium chloride soln 79 Inj 10 ml ampoule — Up to 5 inj available on a PSO	when on the same for eye drops; or 6 for cystic fibrosis pa 7.60	rm as an	injection	listed in the Pharmaceution
<ol> <li>On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or</li> <li>On a bulk supply order; or</li> <li>When used in the extemporaneous compounding of e</li> <li>When used for the dilution of sodium chloride soln 79</li> </ol>	when on the same for eye drops; or 6 for cystic fibrosis pa 7.60	rm as an	n injection	
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of each of the dilution of sodium chloride soln 79 Inj 10 ml ampoule — Up to 5 inj available on a PSO	when on the same for eye drops; or 6 for cystic fibrosis pa 7.60	rm as an utients or 50	n injection	<u>Multichem</u>
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	when on the same for eye drops; or 6 for cystic fibrosis pa 7.60	rm as an utients or 50	n injection	<u>Multichem</u>
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of 6 4) When used for the dilution of sodium chloride soln 79 Inj 10 ml ampoule — Up to 5 inj available on a PSO	when on the same for eye drops; or 6 for cystic fibrosis pa	rm as an	nly.	<u>Multichem</u>
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 79 Inj 10 ml ampoule — Up to 5 inj available on a PSO	when on the same for eye drops; or 6 for cystic fibrosis pa	rm as an utients or 50	nly.	<u>Multichem</u> Fresenius Kabi
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 79  Inj 10 ml ampoule — Up to 5 inj available on a PSO	eye drops; or 6 for cystic fibrosis pa7.605.00	rm as an striction as an	nly.	Multichem Fresenius Kabi Calcium Resonium
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 79  Inj 10 ml ampoule — Up to 5 inj available on a PSO	eye drops; or 6 for cystic fibrosis pa7.605.00	rm as an	nly.	Multichem Fresenius Kabi
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 79  Inj 10 ml ampoule — Up to 5 inj available on a PSO	eye drops; or 6 for cystic fibrosis pa7.605.00169.85	rm as an striction as an	nily.	Multichem Fresenius Kabi Calcium Resonium
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 79  Inj 10 ml ampoule — Up to 5 inj available on a PSO	eye drops; or 6 for cystic fibrosis pa7.605.00169.85	rm as an striction as an	nily.	Multichem Fresenius Kabi  Calcium Resonium  Electral  Pedialyte -
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 79  Inj 10 ml ampoule — Up to 5 inj available on a PSO	eye drops; or 6 for cystic fibrosis pa	rm as an striction as an	nly.	Multichem Fresenius Kabi  Calcium Resonium  Electral  Pedialyte -
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 79  Inj 10 ml ampoule — Up to 5 inj available on a PSO	eye drops; or 6 for cystic fibrosis pa	rm as an attients or 50 20 300 g OI 50 000 ml C	nly.	Multichem Fresenius Kabi  Calcium Resonium  Electral  Pedialyte - Bubblegum
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 79  Inj 10 ml ampoule — Up to 5 inj available on a PSO	eye drops; or 6 for cystic fibrosis pa	rm as an attients or 50 20 300 g OI 50 000 ml C	nly.	Multichem Fresenius Kabi  Calcium Resonium  Electral  Pedialyte - Bubblegum
NATER  1) On a prescription or Practitioner's Supply Order only Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 79  Inj 10 ml ampoule — Up to 5 inj available on a PSO	eye drops; or 6 for cystic fibrosis pa	rm as an attients or 50 20 300 g Of 50 000 ml C	nly.	Multichem Fresenius Kabi  Calcium Resonium  Electral  Pedialyte - Bubblegum

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
SODIUM BICARBONATE Cap 840 mg	8.52	100		Sodibic Sodibic
SODIUM POLYSTYRENE SULPHONATE Powder	84.65	454 g C	OP 🗸	Resonium-A

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Alpha-Adrenoceptor Blockers**

## **Alpha Adrenoceptor Blockers**

DOXAZOSIN		
* Tab 2 mg17.35	500	<ul> <li>Doxazosin Clinect</li> </ul>
* Tab 4 mg20.94	500	✓ Doxazosin Clinect
PHENOXYBENZAMINE HYDROCHLORIDE		
* Cap 10 mg65.00	30	✓ BNM S29
216.67	100	✓ Dibenzyline \$29
PRAZOSIN		
* Tab 1 mg5.53	100	✓ Arrotex-Prazosin S29 S29
* Tab 2 mg7.00	100	✓ Arrotex-Prazosin S29 S29
* Tab 5 mg11.70	100	✓ Arrotex-Prazosin S29 S29

# **Agents Affecting the Renin-Angiotensin System**

### **ACE Inhibitors**

CAP.	$T \cap D$	חוו	
LAP	אווו	RII	

\* Tah 0.5 mg

*	Oral liq 5 mg per ml	94.99	95 ml OP	✓ Capoten
	Oral liquid restricted to children under 12 years of age.			

#### CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

2 69

不	1ab 0.5 mg2.09	90	♥ Zaprii
*	Tab 2.5 mg5.79	90	✓ Zapril
	Tab 5 mg10.05	90	✓ Zapril
ΕN	ALAPRIL MALEATE		
*	Tab 5 mg1.75	90	✓ Acetec
	Acetec to be Principal Supply on 1 February 2024		
*	Tab 10 mg1.97	90	✓ Acetec
	Acetec to be Principal Supply on 1 February 2024		
*	Tab 20 mg	90	✓ Acetec
	Acetec to be Principal Supply on 1 February 2024		
LIS	SINOPRIL		
*	Tab 5 mg11.07	90	✓ Ethics Lisinopril
			✓ Teva Lisinopril
*	Tab 10 mg11.67	90	✓ Ethics Lisinopril
	·		✓ Teva Lisinopril
*	Tab 20 mg14.69	90	✓ Ethics Lisinopril
	•		✓ Teva Lisinopril
PE	RINDOPRIL		
*	Tab 2 mg	30	✓ Coversyl
*	· ·	30	✓ Coversyl
*	Tab 8 mg	30	✓ Coversyl
-			

✓ 7anril

			AI II		COLAN STSTEW
	(N	Subsidy flanufacturer's Price) \$	Per	Fully Subsidised	
*** RA ***	JINAPRIL Tab 5 mg Tab 10 mg Tab 20 mg MIPRIL Cap 1.25 mg Cap 2.5 mg Cap 5 mg Cap 1 mg Cap 1 mg	5.18 7.95 6.90 6.60 6.75	90 90 90 90 90 90 90	\ \ \ \	Arrow-Quinapril 5 Arrow-Quinapril 10 Arrow-Quinapril 20 Tryzan Tryzan Tryzan Tryzan Tryzan
A	CE Inhibitors with Diuretics				
QL	JINAPRIL WITH HYDROCHLOROTHIAZIDE – Subsidy by endor Subsidy by endorsement – Subsidised for patients who were tal 2022 and the prescription is endorsed accordingly. Pharmacist exists a record of prior dispensing of quinapril with hydrochlorott Tab 10 mg with hydrochlorothiazide 12.5 mg	king quinapril with s may annotate the hiazide. 4.10		scription a	
A	ngiotensin II Antagonists				
**** LO***	NDESARTAN CILEXETIL  Tab 4 mg	2.28 3.31 5.26 2.00 2.29 2.86	90 90 90 90 84 84 84 84	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Candestar Candestar Candestar Candestar Candestar Losartan Actavis Losartan Actavis Losartan Actavis Losartan Actavis
A	ngiotensin II Antagonists with Diuretics				
CA *	NDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDE Tab 16 mg with hydrochlorothiazide 12.5 mg	4.10	30	✓	APO-Candesartan HCTZ 16/12.5
*	Tab 32 mg with hydrochlorothiazide 12.5 mg	5.25	30	•	APO-Candesartan HCTZ 32/12.5
_	SARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE Tab 50 mg with hydrochlorothiazide 12.5 mg	4.00	30	•	Arrow-Losartan & Hydrochlorothiazide

# Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1905 of	n the next page -	- Retail	pharmacy
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	<ul> <li>Manufac</li> </ul>	turer

## **⇒SA1905** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure: and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III; or
  - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
  - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 125

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

## **Antiarrhythmics**

To lighted the hydrochionide refer to NETTV 000 0101 EW, Anacometics, Eccar, pr	age 120	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg	30	✓ Aratac
▲ Tab 200 mg4.49	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a PSO 9.12	6	✓ Cordarone-X
15.22	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a		
PSO15.09	10	✓ Martindale
DIGOXIN		
* Tab 62.5 mcg - Up to 30 tab available on a PSO	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO	240	✓ Lanoxin
* Oral liq 50 mcg per ml16.60	60 ml	✓ Lanoxin
		<ul> <li>Lanoxin Paediatric</li> </ul>
		Elixir \$29
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg20.05	84	✓ Rythmodan -
		Cheplafarm S29
23.87	100	✓ Rythmodan

Medsurae

Medsurae

✓ Midodrine

100

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
FLECAINIDE ACETATE			
▲ Tab 50 mg	19.95	60	✓ Flecainide BNM ✓ Flecainide
Elegainida PNM ta ha Principal Cupply on 1 December C	1000		Sandoz S29
Flecainide BNM to be Principal Supply on 1 December 2 Cap long-acting 100 mg		90	✓ <u>Flecainide</u> <u>Controlled</u> Release Teva
▲ Cap long-acting 200 mg	54.28	90	✓ Flecainide Controlled
Inj 10 mg per ml, 15 ml ampoule	104.00	5	Release Teva ✓ Tambocor
▲ Cap 150 mg	162.00	100	✓ Teva S29
▲ Cap 250 mg		100	✓ Teva \$29
PROPAFENONE HYDROCHLORIDE  Tab 150 mg	40.90	50	✓ Rytmonorm
Antihypotensives			
MIDODRINE – Special Authority see SA1474 below – Retail pha Brand switch fee payable (Pharmacode 2660741) - see page Tab 2.5 mg	e 264 for details	100	✓ <u>Midodrine</u>

# Tab 5 mg ......59.98

■ SA1474 Special Authority for Subsidy
Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

# **Beta-Adrenoceptor Blockers**

# **Beta Adrenoceptor Blockers**

ATENOLOL			
* Tab 50 mg	9.33	500	✓ Mylan Atenolol
* Tab 100 mg	14.20	500	✓ <u>Viatris</u> ✓ Atenolol Viatris
* Oral liq 25 mg per 5 ml	21.25	300 ml OP	<ul><li>✓ Mylan Atenolol</li><li>✓ Atenolol AFT</li></ul>
	38.20		S29 S29 ✓ Essential
	49.85		Generics S29 ✓ Atenolol AFT

Restricted to children under 12 years of age. (Mylan Atenolol Tab 50 mg to be delisted 1 November 2023)

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
SOPROLOL FUMARATE				
Tab 2.5 mg	1.84	90	✓	Bisoprolol Mylan
			✓	Bisoprolol Viatris
★ Tab 5 mg	2.55	90		Bisoprolol Mylan
				Bisoprolol Viatris
F Tab 10 mg	3.62	90		Bisoprolol Mylan
Piganyalal Miylan Tab 0.5 mg to be delicted 1 Navambay 2002)			•	Bisoprolol Viatris
Bisoprolol Mylan Tab 2.5 mg to be delisted 1 November 2023) Bisoprolol Mylan Tab 5 mg to be delisted 1 November 2023)				
,				
ARVEDILOL	0.04	00		0
← Tab 6.25 mg		60		Carvedilol Sandoz
€ Tab 12.5 mg		60		Carvedilol Sandoz
≰ Tab 25 mg	2.95	60	•	Carvedilol Sandoz
ABETALOL				
€ Tab 100 mg		100		<u>Trandate</u>
€ Tab 200 mg		100	•	<u>Trandate</u>
Inj 5 mg per ml, 20 ml ampoule		5		Tuandata
k ini F and a grant OO and viol	(88.60)			Trandate
k inj 5 mg per ml, 20 ml vial		1		
	(48.20)			Alvogen S29
METOPROLOL SUCCINATE				
Fab long-acting 23.75 mg		30	_	Betaloc CR
Tab long-acting 47.5 mg		30		Betaloc CR
Fab long-acting 95 mg		30		Betaloc CR
Fab long-acting 190 mg	4.27	30	•	Betaloc CR
METOPROLOL TARTRATE				
₹ Tab 50 mg	5.66	100		IPCA-Metoprolol
₹ Tab 100 mg	7.55	60		IPCA-Metoprolol
Fab long-acting 200 mg		28		Slow-Lopresor
Inj 1 mg per ml, 5 ml vial	26.50	5	_	Metoprolol IV Mylan
			•	Metoprolol IV Viatris
IADOLOL				
★ Tab 40 mg	19.19	100		Nadolol BNM
€ Tab 80 mg	30.39	100	✓	Nadolol BNM
ROPRANOLOL				
★ Tab 10 mg	7.04	100	✓	<u>Drofate</u>
★ Tab 40 mg	8.75	100	✓ ]	IPCA-Propranolol
Cap long-acting 160 mg	18.17	100	1	Cardinol LA
♦ Oral liq 4 mg per ml - Special Authority see SA1327 below -	-			
Retail pharmacy	CBS 5	500 m	nl 🗸 l	Roxane-
				Propranolol \$29

### **⇒SA1327** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or

(Man	Subsidy	F	ully	Brand or
	ufacturer's Price)	Subsidi	sed	Generic
	\$	Per	<b>✓</b>	Manufacturer

continued...

2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

#### SOTAL OL

*	Tab 80 mg	37.50	500	✓ Mylan
*	Tab 160 mg	14.00	100	✓ Mylan

## **Calcium Channel Blockers**

## **Dihydropyridine Calcium Channel Blockers**

ΑM	LODIPINE			
*	Tab 2.5 mg	1.45	90	✓ Vasorex
*	Tab 5 mg	1.21	90	✓ Vasorex
	Tab 10 mg		90	✓ Vasorex
FE	LODIPINE			
*	Tab long-acting 2.5 mg	1.45	30	✓ Plendil ER
	Tab long-acting 5 mg		90	✓ Felo 5 ER
	Tab long-acting 10 mg		90	✓ Felo 10 ER
NIF	EDIPINE			
*	Tab long-acting 10 mg - Subsidy by endorsement	19.42	56	✓ Tensipine MR10 S29

Subsidised for patients who were taking nifedipine tab long-acting 10 mg prior to 1 July 2023 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of nifedipine tab long-acting 10 mg.

*	Tab long-acting 20 mg9.12	50	✓ Mylan (12 hr release) S29
	17.72	100	✓ Nyefax Retard
*	Tab long-acting 30 mg4.78	14	✓ Mylan Italy (24 hr release) S29
	34.10	100	✓ Mylan (24 hr release) \$29
*	Tab long-acting 60 mg52.81	100	✓ Mylan (24 hr
			release) S29

(Mylan (12 hr release) \$29 Tab long-acting 20 mg to be delisted 1 December 2023) (Mylan (24 hr release) \$29 Tab long-acting 30 mg to be delisted 1 February 2024)

## **Other Calcium Channel Blockers**

DILTIAZEM HYDROCHLORIDE			
* Cap long-acting 120 mg	65.35	500	✓ <u>Diltiazem CD Clinect</u>
* Cap long-acting 180 mg	7.00	30	✓ Cardizem CD
* Cap long-acting 240 mg	9.30	30	✓ Cardizem CD
PERHEXILINE MALEATE			
★ Tab 100 mg	62 90	100	✓ Paysin

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	
n)	fanufacturer's Price) \$	Per	Subsidised •	
ERAPAMIL HYDROCHLORIDE				
★ Tab 40 mg	7.01	100	✓	Isoptin
★ Tab 80 mg	11.74	100	✓	Isoptin
★ Tab long-acting 120 mg	36.02	100	•	Isoptin Retard S29
			_	Isoptin SR
★ Tab long-acting 240 mg	15.12	30	✓	Isoptin SR
★ Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a				
PSO	25.00	5	/	Isoptin
Controller Asting Asserts				
Centrally-Acting Agents				
CLONIDINE				
Patch 2.5 mg, 100 mcg per day - Only on a prescription	11.70	4	/	Mylan
Patch 5 mg, 200 mcg per day - Only on a prescription		4		Mylan
Patch 7.5 mg, 300 mcg per day - Only on a prescription		4	✓	Mylan
CLONIDINE HYDROCHLORIDE				
★ Tab 25 mcg	29.32	112	/	Clonidine Teva
₹ Tab 150 mcg		100		Catapres
Inj 150 mcg per ml, 1 ml ampoule		10		Medsurge
METHYLDOPA				
₹ Tab 250 mg	15.10	100	/	Methyldopa Mylan
	52.85	500		Methyldopa Mylan
				S29 S29
Diuretics				
Loop Diuretics				
BUMETANIDE				
₭ Tab 1 mg	4 91	30	/	Burinex S29 S29
r tab i mg	16.36	100		Burinex
k Inj 500 mcg per ml, 4 ml vial		5		Burinex
		ŭ		
UROSEMIDE [FRUSEMIDE] Tab 40 mg – Up to 30 tab available on a PSO	8.00	1,000	J	IPCA-Frusemide
Tab To fing Op to ob tab available off a Foot		50		Urex Forte
<b>k</b> Tah 500 mα	25 00			Furosemid-
≰ Tab 500 mg	25.00 89.48	50	<b>✓</b>	
★ Tab 500 mg		30	/	
★ Tab 500 mg		30	•	Ratiopharm S29
≰ Tab 500 mg		100		
★ Tab 500 mg	89.48			Ratiopharm \$29
★ Tab 500 mg	89.48		•	Ratiopharm \$29 Furosemid- Ratiopharm \$29
l⊀ Oral liq 10 mg per ml	89.48 169.96 11.20 30	100 0 ml O	<b>,</b> P <b>,</b>	Ratiopharm \$29 Furosemid- Ratiopharm \$29 Lasix
★ Oral liq 10 mg per ml k Inj 10 mg per ml, 25 ml ampoule	89.48 169.96 11.20 30 60.65	100 0 ml O 6	P	Ratiopharm \$29  Furosemid- Ratiopharm \$29  Lasix Lasix
l⊀ Oral liq 10 mg per ml	89.48 169.96 11.20 30 60.65	100 0 ml O	P	Ratiopharm \$29 Furosemid- Ratiopharm \$29 Lasix
★ Oral liq 10 mg per ml k Inj 10 mg per ml, 25 ml ampoule	89.48 169.96 11.20 30 60.65	100 0 ml O 6	P	Ratiopharm \$29  Furosemid- Ratiopharm \$29  Lasix Lasix
Oral liq 10 mg per ml Inj 10 mg per ml, 25 ml ampoule Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a PS0  Potassium Sparing Diuretics	89.48 169.96 11.20 30 60.65	100 0 ml O 6	P	Ratiopharm \$29  Furosemid- Ratiopharm \$29  Lasix Lasix
Oral liq 10 mg per ml Inj 10 mg per ml, 25 ml ampoule Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO  Potassium Sparing Diuretics  MILORIDE HYDROCHLORIDE	89.48 169.96 11.20 30 60.65 D2.40	100 0 ml O 6	P	Ratiopharm \$29  Furosemid- Ratiopharm \$29  Lasix Lasix
Oral liq 10 mg per ml Inj 10 mg per ml, 25 ml ampoule Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO  Potassium Sparing Diuretics  MILORIDE HYDROCHLORIDE  Oral liq 1 mg per ml	89.48 169.96 11.20 30 30 30 30 30 30 30 30 30 3	100 0 ml O 6 5	P	Ratiopharm \$29  Furosemid- Ratiopharm \$29  Lasix Lasix Furosemide-Baxter
FOral liq 10 mg per ml	89.48  169.96 11.20 36 60.65  D2.40  32.10 28  Retail pharmacy	100 O ml O 6 5	P V	Ratiopharm \$29  Furosemid- Ratiopharm \$29  Lasix Lasix Furosemide-Baxter
Oral liq 10 mg per ml Inj 10 mg per ml, 25 ml ampoule Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO  Potassium Sparing Diuretics  MILORIDE HYDROCHLORIDE  Oral liq 1 mg per ml	89.48  169.9611.20 3060.652.40 32.10 26 Retail pharmacy18.50	100 0 ml O 6 5	P V	Ratiopharm \$29  Furosemid- Ratiopharm \$29  Lasix Lasix Furosemide-Baxter

			O A DDIOV		III AD OVOTEM
			CARDIOV	ASC	ULAR SYSTEM
_		Subsidy (Manufacturer's Price \$		Fully dised	Brand or Generic Manufacturer
Ini	SA1728 Special Authority for Subsidy tial application from any relevant practitioner. Approvals valid following criteria: th:  1 Patient has heart failure with ejection fraction less than 40 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolaci 2.2 Patient has experienced a clinically significant adve	%; and tone; or			•
SP *	IRONOLACTONE Tab 25 mg Tab 100 mg Oral liq 5 mg per ml	10.65	100 100 25 ml OP	<b>✓</b> S	spiractin piractin siomed
P	otassium Sparing Combination Diuretics				
* AN	IILORIDE HYDROCHLORIDE WITH FUROSEMIDE  Tab 5 mg with furosemide 40 mg IILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZI	DE	28		rumil
*	Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	✓ N	Ioduretic
Т	hiazide and Related Diuretics				
	NDROFLUMETHIAZIDE [BENDROFLUAZIDE]	51 50	500	<b>.</b> .	Prow-

BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]  * Tab 2.5 mg – Up to 150 tab available on a PSO51.50	500	✓ Arrow- Bendrofluazide
May be supplied on a PSO for reasons other than emergency.  ★ Tab 5 mg61.00	500	✓ Arrow- Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml27.82	25 ml OP	✓ Biomed
CHLORTALIDONE [CHLORTHALIDONE] Tab 25 mg	50	✓ <u>Hygroton</u>
INDAPAMIDE	90	✓ Dapa-Tabs
METOLAZONE Tab 5 mgCBS	1	✓ Metolazone S29
1ab 3 mg0b3	50	✓ Zaroxolyn \$29

# Vasopressin receptor antagonists

TOLVAPTAN - Special Authority see SA2166 on the next page	- Retail pharmac	v	
Tab 15 mg		, 28 OP	✓ Jinarc
Tab 30 mg		28 OP	✓ Jinarc
Tab 45 mg + 15 mg		56 OP	✓ Jinarc
Tab 60 mg + 30 mg		56 OP	<ul><li>Jinarc</li></ul>
Tab 90 mg + 30 mg		56 OP	<ul><li>Jinarc</li></ul>

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	<ul> <li>Manufac</li> </ul>	turer

### **⇒SA2166** Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m<sup>2</sup> at treatment initiation; and
- 3 Either:
  - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m<sup>2</sup> within one-year; or
  - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m<sup>2</sup> per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m<sup>2</sup>; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents							
Fibrates							
BEZAFIBRATE       * Tab 200 mg       19.46         * Tab long-acting 400 mg       21.21	90 30	✓ <u>Bezalip</u> ✓ <u>Bezalip Retard</u>					
Other Lipid-Modifying Agents							
ACIPIMOX  * Cap 250 mg21.56 25.44	30	✓ Olbetam S29 S29 ✓ Olbetam					
Resins							
COLESTIPOL HYDROCHLORIDE  Grans for oral liq 5 g32.89  COLESTYRAMINE	30	✓ Colestid					
Powder for oral suspension 4 g sachet	50	<ul> <li>✓ Colestyramine -         Mylan \$29</li> <li>✓ Quantalan sugar         free \$29</li> </ul>					
HMG CoA Reductase Inhibitors (Statins)							
ATORVASTATIN  * Tab 10 mg 6.16  * Tab 20 mg 9.24  * Tab 40 mg 14.92  * Tab 80 mg 26.54	500 500 500 500	✓ Lorstat ✓ Lorstat ✓ Lorstat ✓ Lorstat					

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	1	Manufacturer
PRAVASTATIN				
* Tab 20 mg	2.11	28	✓	Pravastatin Mylan
•			✓	Pravastatin Viatris
* Tab 40 mg	3.61	28	✓	Pravastatin Mylan
(Pravastatin Mylan Tab 20 mg to be delisted 1 January 2024)				·
ROSUVASTATIN - Special Authority see SA2093 below - Retail	pharmacy			
* Tab 5 mg		30	1	<b>Rosuvastatin Viatris</b>
Rosuvastatin Viatris to be Principal Supply on 1 December				
* Tab 10 mg	1.69	30	✓	<b>Rosuvastatin Viatris</b>
Rosuvastatin Viatris to be Principal Supply on 1 December	er 2023			
* Tab 20 mg	2.71	30	✓	<b>Rosuvastatin Viatris</b>
Rosuvastatin Viatris to be Principal Supply on 1 December	er 2023			
* Tab 40 mg	4.55	30	✓	<b>Rosuvastatin Viatris</b>
Rosuvastatin Viatris to be Principal Supply on 1 December	er 2023			

#### **⇒SA2093** Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Patient is considered to be at risk of cardiovascular disease; and
  - 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:
  - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
  - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:
Both:

- 1 Any of the following:
  - 1.1 Patient has proven coronary artery disease (CAD); or
  - 1.2 Patient has proven peripheral artery disease (PAD); or
  - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

	Subsidy (Manufacturer's Price)	5	Fully Subsidised	
	\$	Per	•	Manufacturer
SIMVASTATIN				
* Tab 10 mg	1.68	90		Simvastatin Mylan
				Simvastatin Viatris
* Tab 20 mg	2.54	90		Simvastatin Mylan
				Simvastatin Viatris
* Tab 40 mg	4.11	90		Simvastatin Mylan
				Simvastatin Viatris
* Tab 80 mg	8.81	90		Simvastatin Mylan
			✓	Simvastatin Viatris

### Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Special Authority see SA1045 below – Retail pharmacy		
* Tab 10 mg	30	✓ Ezetimibe Sandoz
Ezetimibe Sandoz to be Principal Supply on 1 December 2023		

#### **⇒SA1045** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
  - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than  $10 \times \text{normal}$ ) when treated with one statin; or
  - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
  - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy

ELECTION DE TITLE CONTINUE OPERATION CONTINUE CO	TO DOTOTE TECTAL	nannaoy	
Tab 10 mg with simvastatin 10 mg	5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg	7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	✓ Zimybe

#### ⇒SA1046 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

#### **Nitrates**

#### GLYCFRYL TRINITRATE

*	Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO7.48	250 dose OP	✓ Nitrolingual Pump Spray
	Patch 25 mg, 5 mg per day	30 30	<ul><li>✓ Nitroderm TTS</li><li>✓ Nitroderm TTS</li></ul>

Subsidy (Manufacturer's P	Price) Su	Fully Brand or ubsidised Generic	
\$	Per	✓ Manufacturer	
OSORBIDE MONONITRATE			
Tab 20 mg	100	✓ Ismo 20	
Tab long-acting 40 mg	30	✓ Ismo 40 Retard ✓ Duride	1
Tab long-acting 60 mg13.50	90	Duride	
Sympathomimetics			
DRENALINE			
Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO4.98	5	✓ Aspen Adrena	line
12.65		✓ DBL Adrenaling	
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira	
49.00	10	Aspen Adrena	line
Vasodilators			
YDRALAZINE HYDROCHLORIDE			
Tab 25 mg - Special Authority see SA1321 below - Retail			
pharmacyCBS	1	<ul><li>Hydralazine</li></ul>	
	56	✓ Onelink S29	
	84	✓ AMDIPHARM®	629
	100	✓ Camber S29	
Inj 20 mg ampoule25.90	5	Apresoline	
>SA1321 Special Authority for Subsidy itial application from any relevant practitioner. Approvals valid without further e following criteria: ither:	renewal unle	ess notified for application	ons meeti
itial application from any relevant practitioner. Approvals valid without further e following criteria: ither:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients with a nitrate.			
itial application from any relevant practitioner. Approvals valid without further e following criteria: ther:  1 For the treatment of refractory hypertension; or			
itial application from any relevant practitioner. Approvals valid without further e following criteria: ither:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients with a nitrate with			
<ul> <li>itial application from any relevant practitioner. Approvals valid without further e following criteria: ither:</li> <li>1 For the treatment of refractory hypertension; or</li> <li>2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.</li> </ul>			nded to A
itial application from any relevant practitioner. Approvals valid without further e following criteria:  ther:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL	who are intole	erant or have not respor	nded to A
itial application from any relevant practitioner. Approvals valid without further e following criteria:  ther:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	who are intole	erant or have not respor ✓ Minoxidil Rom	nded to A
Itial application from any relevant practitioner. Approvals valid without further e following criteria:  ither:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	who are intole	✓ Minoxidil Rom ✓ Loniten	nded to A
itial application from any relevant practitioner. Approvals valid without further e following criteria:  ther:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	who are intole 60 100	erant or have not respon  ✓ Minoxidil Rom ✓ Loniten	nded to A
itial application from any relevant practitioner. Approvals valid without further e following criteria:  ther:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	who are intole 60 100 60	✓ Minoxidil Rom ✓ Loniten ✓ Ikorel	nded to A
itial application from any relevant practitioner. Approvals valid without further e following criteria:  ther:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	who are intole 60 100 60	✓ Minoxidil Rom ✓ Loniten	nded to A
itial application from any relevant practitioner. Approvals valid without further e following criteria:  ther:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	60 100 60 60 5	✓ Minoxidil Rom ✓ Loniten ✓ Ikorel ✓ Hospira	nded to A
itial application from any relevant practitioner. Approvals valid without further e following criteria:  ther:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	60 100 60 60 60	✓ Minoxidil Rom ✓ Loniten ✓ Ikorel	nded to A
itial application from any relevant practitioner. Approvals valid without further e following criteria:  ther:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	60 100 60 60 5	✓ Minoxidil Rom ✓ Loniten ✓ Ikorel ✓ Hospira	nded to A
Itial application from any relevant practitioner. Approvals valid without further e following criteria:  ither:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	60 100 60 60 5 50	✓ Minoxidil Rom ✓ Loniten ✓ Ikorel ✓ Hospira ✓ Trental 400	a S29
Itial application from any relevant practitioner. Approvals valid without further e following criteria:  ither:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients v inhibitors and/or angiotensin receptor blockers.  INOXIDIL 1 Tab 10 mg	60 100 60 60 5 50	✓ Minoxidil Rom ✓ Loniten ✓ Ikorel ✓ Hospira ✓ Trental 400	a \$29
Itial application from any relevant practitioner. Approvals valid without further e following criteria:  ither:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients vinhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	60 100 60 60 5 50	✓ Minoxidil Rom ✓ Loniten ✓ Ikorel ✓ Hospira ✓ Trental 400	a \$29
Itial application from any relevant practitioner. Approvals valid without further e following criteria:  ither:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients vinhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	60 100 60 5 50	✓ Minoxidil Rom ✓ Loniten ✓ Ikorel ✓ Ikorel ✓ Hospira ✓ Trental 400 ✓ Ambrisentan N	a \$29  /iatris
Itial application from any relevant practitioner. Approvals valid without further e following criteria:  ither:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients vinhibitors and/or angiotensin receptor blockers.  INOXIDIL Tab 10 mg	60 100 60 60 5 50	✓ Minoxidil Rom ✓ Loniten ✓ Ikorel ✓ Hospira ✓ Trental 400	a \$29  /iatris

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

(Mylan Tab 10 mg to be delisted 1 December 2023)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## ⇒SA2253 Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Ambrisentan is to be used as PAH monotherapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
    - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:

(Ma	Subsidy nufacturer's Price)	F Subsid	ully	Brand or Generic
	\$	Per	1	Manufacturer

continued...

- 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
- 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Ambrisentan is to be used as PAH dual therapy; and
  - 5.2 Either:
    - 5.2.1 Patient has tried a PAH monotherapy (sildenafil or bosentan) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; or
    - 5.2.2 Patient has tried PAH dual therapy including bosentan and has experienced intolerable side effects on bosentan; and
  - 5.3 Both:
    - 5.3.1 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy; and
    - 5.3.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease).

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 5 Both:
  - 5.1 Ambrisentan is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Both:
      - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV; and
      - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN - Special Authority see SA2254 below - Retail pharmacy

Tab 62.5 mg	119.85	60	✓ Bosentan Dr
			Reddy's
Tab 125 mg	119.85	60	✓ Bosentan Dr
			Reddy's

#### ⇒SA2254 Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per		Manufacturer

continued...

- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Bosentan is to be used as PAH monotherapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient has experienced intolerable side effects on sildenafil; or
    - 5.2.2 Patient has an absolute contraindication to sildenafil; or
    - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Either:
  - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool\*\*; or
  - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy.

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has pulmonary arterial hypertension (PAH)\*; and

Subsidy		Fully	Brand or	
(Manufacturer's Price	,	Subsidised	Generic	
\$	Per		Manufacturer	

continued...

- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Bosentan is to be used as part of PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

## **Phosphodiesterase Type 5 Inhibitors**

ILDENAFIL – Special Authority see SA2255 on the next pag	e – Retail pharmacy		
Tab 25 mg	0.85	4	✓ Vedafil
Tab 50 mg	1.70	4	✓ Vedafil
Tab 100 mg	10.20	12	✓ Vedafil

SI

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

### **⇒SA2255** Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon\*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon\*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension\*) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH is confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
    - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Note: Indications marked with \* are Unapproved Indications.

† The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH</u>

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## **Prostacyclin Analogues**

⇒SA2256 Special Authority for Subsidy

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
  - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
  - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † : or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Epoprostenol is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list; or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool\*\*.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

\*\* the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST - Special Authority see SA2257 below - Retail pharmacy

#### ⇒SA2257 Special Authority for Subsidy

**Initial application** — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
- 4.1.5 Any of the following:
  - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
  - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
  - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Iloprost is to be used as PAH monotherapy; and
  - 5.2 Either:
    - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
    - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

**Initial application** — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † : or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*: or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
  - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and

|--|

continued...

- 5.2 Either:
  - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
  - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
- 5.3 Either:
  - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool\*\*; or
  - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

**Initial application** — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
  - 4.1 All of the following:
    - 4.1.1 PAH has been confirmed by right heart catheterisation; and
    - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm<sup>-5</sup>); and
    - 4.1.5 Any of the following:
      - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
      - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool\*\*; or
      - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
  - 5.1 Iloprost is to be used as PAH triple therapy; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is on the lung transplant list: or
    - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
    - 5.2.3 Both:
      - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool\*\*; and
      - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

**Renewal** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool\*\*.

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

<sup>\*\*</sup> the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

# **Antiacne Preparations**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 98

#### **ADAPALENE**

- a) Maximum of 30 g per prescription
- b) Only on a prescription

22.89	30 g OP	<ul><li>Differin</li></ul>
etail pharmacy	· ·	
11.26	60	<ul><li>Oratane</li></ul>
18.75	120	✓ Oratane
	120	✓ Oratane
		etail pharmacy 60 6018.75 120

#### ⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

#### **TRETINOIN**

Crm 0.5 mg per g − Maximum of 50 g per prescription .......15.57 50 g OP ✓ ReTrieve

### Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 98

#### HYDROGEN PEROYIDE

# Crm 1%	8.56	10 g OP	✓ Crystaderm
MUPIROCIN			
Oint 2%	6.60	15 g OP	
	(11.50)		Bactroban

- a) Only on a prescription
- b) Not in combination

# **DERMATOLOGICALS**

	Subsidy		Fully Brand or
	(Manufacturer's F \$	rice) Subs Per	sidised Generic  Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID]	<u> </u>		
Crm 2%	1.59	5 g OP	✓ Foban
a) Maximum of 5 g per prescription		·	
b) Only on a prescription			
c) Not in combination			
Oint 2%	1.59	5 g OP	✓ <u>Foban</u>
a) Maximum of 5 g per prescription			
<ul><li>b) Only on a prescription</li><li>c) Not in combination</li></ul>			
SULFADIAZINE SILVER			
Crm 1%	10.80	50 g OP	✓ Flamazine
	10.00	50 g OF	▼ Fidiliazille
<ul><li>a) Up to 250 g available on a PSO</li><li>b) Not in combination</li></ul>			
b) Not in combination			
Antifungals Topical			
For systemic antifungals, refer to INFECTIONS, Antifunga	als, page 105		
AMOROLFINE			
a) Only on a prescription			
b) Not in combination			
Nail soln 5%	21.87	5 ml OP	✓ MycoNail
CLOTRIMAZOLE			•
* Crm 1%	1.10	20 g OP	✓ Clomazol
a) Only on a prescription			
b) Not in combination			
<b>₭</b> Soln 1%		20 ml OP	_
	(7.55)		Canesten
a) Only on a prescription			
b) Not in combination			
ECONAZOLE NITRATE	4.00	00 00	
Crm 1%		20 g OP	Davismil
a) Only on a proparintian	(7.78)		Pevaryl
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>			
Foaming soln 1%, 10 ml sachets	9.89	3	
. oag oo /o, . o oaoo	(17.92)	· ·	Pevaryl
a) Only on a prescription	,		•
b) Not in combination			
MICONAZOLE NITRATE			
<b>₭</b> Crm 2%	0.81	15 g OP	✓ Multichem
a) Only on a prescription		· ·	
b) Not in combination			
* Lotn 2%		30 ml OP	<b>5</b>
	(10.03)		Daktarin
a) Only on a prescription			
b) Not in combination  ★ Tinct 2%	4.26	30 ml OP	
<b>★</b> Tinct 2%	(12.10)	30 IIII OP	Daktarin
a) Only on a prescription	(12.10)		Duntaiii
b) Not in combination			
-/			

	Subsidy		Fully	Brand or
	(Manufacturer's Pric		ubsidised	Generic
	\$	Per		Manufacturer
Antipruritic Preparations				
CALAMINE				
a) Only on a prescription				
b) Not in combination				
Crm, aqueous, BP	1.08	100 g	<b>√</b> <u>C</u>	Calamine-AFT
	3.45		<b>✓</b> h	ealthE Calamine
				Aqueous
CROTAMITON				
a) Only on a prescription				
b) Not in combination			_	
Crm 10%	3.29	20 g OP	✓ <u>li</u>	tch-Soothe
MENTHOL - Only in combination				
1) Only in combination with a dermatological base or prop	orietary Topical Cort	icosterio	d – Plain	
2) With or without other dermatological galenicals.	, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			
,				

25 g

100 g

29.60

✓ MidWest
✓ MidWest

✓ DP Lotn HC

250 ml

# **Corticosteroids Topical**

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 88

Corticosteroids - Plai
------------------------

BETAMETHASONE DIPROPIONATE			
Crm 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	✓ Diprosone
Oint 0.05%	2.96	15 g OP	✓ Diprosone
	36.00	50 g OP	<ul><li>Diprosone</li></ul>
Oint 0.05% in propylene glycol base	4.33	30 g OP	Diprosone OV
BETAMETHASONE VALERATE			
* Crm 0.1%	4.53	50 g OP	✓ Beta Cream
* Oint 0.1%	5.84	50 g OP	✓ Beta Ointment
* Lotn 0.1%	25.00	50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE			
* Crm 0.05%	2.40	30 g OP	✓ Dermol
* Oint 0.05%	2.33	30 g OP	✓ Dermol
CLOBETASONE BUTYRATE		-	
Crm 0.05%	5.38	30 g OP	
	(10.00)	· ·	Eumovate
HYDROCORTISONE			
* Crm 1% - Only on a prescription	1.78	30 g OP	✓ Ethics
	20.40	500 g	✓ Noumed
* Powder – Only in combination	49.95	25 g	✓ ABM
Up to 5% in a dermatological base (not proprietary Togalenicals	oical Corticosterio	d – Plain) with o	or without other dermatological
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only	v on		
The state of the s	, -		

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

# **DERMATOLOGICALS**

	Subsidy (Manufacturer's F \$	Price) Subs	Fully Brand or sidised Generic Manufacture	r
HYDROCORTISONE BUTYRATE	·			
Lipocream 0.1%	4.85	100 g OP	✓ Locoid Lipoc	ream
Oint 0.1%	10.28	100 g OP	✓ Locoid	
Milky emul 0.1%	12.33	100 ml OP	✓ Locoid Crelo	
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%		15 g OP	✓ Advantan	
Oint 0.1%	4.95	15 g OP	Advantan	
MOMETASONE FUROATE			_	
Crm 0.1%		15 g OP	✓ Elocon Alcoh	
01 - 0 - 01	3.10	50 g OP	✓ Elocon Alcoh	ol Free
Oint 0.1%		15 g OP	Elocon	
Lotn 0.1%	2.90	50 g OP 30 ml OP	✓ <u>Elocon</u> ✓ Elocon	
	4.50	SU IIII UP	EIUCOII	
FRIAMCINOLONE ACETONIDE	0.40	100 - 00	Autatoria	
Crm 0.02% Oint 0.02%		100 g OP 100 g OP	<ul><li>✓ Aristocort</li><li>✓ Aristocort</li></ul>	
Ollit 0.02 %	0.54	100 g OF	Anstocon	
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU	JSIDIC ACID]			
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49	15 g OP		
	(10.45)		Fucicort	
<ul><li>a) Maximum of 15 g per prescription</li><li>b) Only on a prescription</li></ul>				
HYDROCORTISONE WITH MICONAZOLE - Only on a prescri	ption			
★ Crm 1% with miconazole nitrate 2%	1.89	15 g OP	✓ Micreme H	
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - (	Only on a prescrip	otion		
Oint 1% with natamycin 1% and neomycin sulphate 0.5%	3.35	15 g OP	Pimafucort	
RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	ΓΙΝ		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 m	ng			
and gramicidin 250 mcg per g - Only on a prescription	3.49	15 g OP		
	(9.28)		Viaderm KC	
Barrier Creams and Emollients				
Barrier Creams				
DIMETHICONE				
★ Crm 5% pump bottle	4.30	500 ml OP	✓ healthE	
· r · · r · · · · · ·			Dimethicon	e 5%
<b>★</b> Crm 10% pump bottle	4.52	500 ml OP	✓ healthE Dimethicon	e 10%
ZINC AND CASTOR OIL			Dilletilloon	J 10/0
k Oint	4.25	500 g	✓ Evara	
	4.65	9	✓ Boucher	
Evara to be Sole Supply on 1 November 2023				
Boucher Oint to be delisted 1 November 2023)				

	Subsidy (Manufacturer's \$	Price) Subsi Per	Fully Brand or dised Generic Manufacturer
Emollients			
QUEOUS CREAM			
Crm	1.30	100 g	✓ healthE Aqueous  Cream SLS Free
	1.73	500 g	✓ Evara ✓ <u>GEM Aqueous</u> <u>Cream</u>
CETOMACROGOL			
k Crm BP	1.99	500 g	✓ Cetomacrogol-AFT
ETOMACROGOL WITH GLYCEROL			
Crm 90% with glycerol 10%	2.13	500 ml OP	✓ Evara
	3.50	1,000 ml OP	✓ Evara
MULSIFYING OINTMENT			
6 Oint BP	3.40	500 g	<ul><li>Emulsifying Ointment ADE</li></ul>
DIL IN WATER EMULSION			
€ Crm	2.04	500 g	✓ Fatty Cream AFT
ARAFFIN			
Oint liquid paraffin 50% with white soft paraffin 50%	4.94	500 g OP	✓ White Soft Liquid  Paraffin AFT
IREA			4
F Crm 10%	1.37	100 g OP	✓ healthE Urea Cream
/OOL FAT WITH MINERAL OIL – Only on a prescription			
Lotn hydrous 3% with mineral oil		1,000 ml	DDLaffer
	(14.96)		DP Lotion Alpha-Keri Lotion
	(20.53) 1.40	250 ml OP	Alpha-Nell Lollon
	(5.87)	200 IIII OF	DP Lotion
	5.60	1,000 ml	DI LOUOII
	(23.91)	1,000 1111	BK Lotion
	1.40	250 ml OP	
	(7.73)		BK Lotion
Other Dermatological Bases			
PARAFFIN			

450 g

2,500 g

19.99

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

✓ healthE
✓ healthE

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



Oint 10%       7.40       65 g OP       ✓ Betadine         a) Maximum of 130 g per prescription         b) Only on a prescription         Antiseptic Solution 10%       4.15       100 ml       ✓ Riodine         Antiseptic soln 10%       3.83       15 ml       ✓ Riodine         5.40       500 ml       ✓ Riodine         Skin preparation, povidone iodine 10% with 30% alcohol       1.63       100 ml	Ainor Skin Infections			
a) Maximum of 130 g per prescription b) Only on a prescription  Antiseptic Solution 10%	OVIDONE IODINE			45
b) Only on a prescription  Antiseptic Solution 10%		7.40	65 g OP	✓ Betadine
Antiseptic Solution 10%	, 31 1 1			
Antiseptic soln 10%	b) Only on a prescription			
Skin preparation, povidone iodine 10% with 30% alcohol	Antiseptic Solution 10%	4.15	100 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	Antiseptic soln 10%	3.83	15 ml	✓ Riodine
(3.48) Betadine Skin Prep Skin preparation, povidone iodine 10% with 70% alcohol1.63 100 ml		5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 70% alcohol1.63 100 ml	Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
		(3.48)		Betadine Skin Prep
	Skin preparation, povidone iodine 10% with 70% alcohol		100 ml	
,	- p - p - m - 7 p - m - 1 m -			Pfizer
		, ,		
Parasiticidal Preparations	IMETHICONE			
<u> </u>	£ Lotn 4%	4.25	200 ml OP	✓ healthE

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Per

Brand or

Generic

Manufacturer

Dimethicone 4% Lotion

✓ Stromectol

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- 3) For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

### ⇒SA2228 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

### Fither:

1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or

IVERMECTIN - Special Authority see SA2228 below - Retail pharmacy 

- - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and

2.2 Either:

- 2.2.1 The person is unable to complete topical therapy: or
- 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:

### **DERMATOLOGICALS**

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

#### continued...

- 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
- 2.2 Fither:
  - 2.2.1 The person is unable to complete topical therapy: or
  - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

#### **PERMETHRIN**

Crm 5%5.75	30 g OP	<ul><li>Lyderm</li></ul>
Lotn 5%4.28	30 ml OP	<ul><li>A-Scabies</li></ul>
(Lyderm Crm 5% to be delisted 1 February 2024)		

# **Psoriasis and Eczema Preparations**

ACITRETIN - Special Authority see SA2024 below - Retail pharmacy		
Cap 10 mg17.86	60	✓ Novatretin
Cap 25 mg 41.36	60	✓ Novatretin

### ⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Fither:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

### BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per g59.9	95 60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g39.3	35 60 g OP	Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g15.9	90 30 g OP	✓ Daivobet
CALCIPOTRIOL		
Oint 50 mcg per g40.0	00 120 g OP	Daivonex

	Subsidy	Dring) O'	Fully	Brand or
	(Manufacturer's I \$	Price) Sub	sidised •	Generic Manufacturer
OAL TAR				
Soln BP – Only in combination		200 ml		idwest
<ol> <li>Up to 10% only in combination with a dermatol</li> <li>With or without other dermatological galenicals</li> </ol>		ietary Topical (	Corticost	eriod – Plain
OAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND S Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5%				
allantoin crm 2.5%	6.59	75 g OP	г.	ranaamil TA
	(8.00) 3.43	30 g OP	Εί	gopsoryl TA
	(4.35)	50 g Oi	Ed	gopsoryl TA
OAL TAR WITH SALICYLIC ACID AND SULPHUR	, ,		`	,
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP		oco-Scalp
	7.95	40 g OP	✓ C	oco-Scalp
MECROLIMUS - Special Authority see SA1970 below - F	Retail pharmacy			
<ul><li>a) Maximum of 15 g per prescription</li><li>b) Note: a maximum of 15 g per prescription and no mo</li></ul>	ore than one prescri	ntion ner 12 we	eks	
Cream 1%		15 g OP	,ono. ✓ El	idel
■ SA1970 Special Authority for Subsidy nitial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approveneeting the following criteria: ioth:	ophthalmologist or ar	, ,		
■ SA1970 Special Authority for Subsidy nitial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approveneeting the following criteria:	ophthalmologist or an als valid without furth ons to topical cortice	ner renewal uni	less notif orificial de	ied for applications
■ SA1970 Special Authority for Subsidy  iitial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve the eting the following criteria:  ioth:  1 Patient has atopic dermatitis on the eyelid; and  2 Patient has at least one of the following contraindicating documented epidermal atrophy, documented allergy to pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUC	ophthalmologist or an als valid without furth ons to topical cortico to topical corticostero ORESCEIN – Only o	ner renewal uni osteroids: perio oids, cataracts, on a prescriptio	less notif orificial de glaucom	ied for applications ermatitis, rosacea, na, or raised intraocu
■ SA1970 Special Authority for Subsidy Initial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve neeting the following criteria: I Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindicating documented epidermal atrophy, documented allergy to pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUCKS Soln 2.3% with trolamine laurilsulfate and fluorescein soci	ophthalmologist or an als valid without furth ons to topical cortico to topical corticostero ORESCEIN – Only o	ner renewal uni osteroids: perio oids, cataracts,	less notif orificial de glaucom	ied for applications
■ SA1970 Special Authority for Subsidy  iitial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve the eting the following criteria:  ioth:  1 Patient has atopic dermatitis on the eyelid; and  2 Patient has at least one of the following contraindicating documented epidermal atrophy, documented allergy the pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUCK Soln 2.3% with trolamine laurilsulfate and fluorescein social calcidus.	ophthalmologist or an als valid without furth ons to topical cortico topical corticostero ORESCEIN – Only odium5.41	per renewal uni posteroids: perio pids, cataracts, on a prescriptio 500 ml	orificial di glaucom	ied for applications ermatitis, rosacea, na, or raised intraocu
■ SA1970 Special Authority for Subsidy Initial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve neeting the following criteria: I Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindicating documented epidermal atrophy, documented allergy to pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUCKS Soln 2.3% with trolamine laurilsulfate and fluorescein soci	ophthalmologist or an als valid without furth ons to topical cortico topical corticostero DRESCEIN — Only odium	osteroids: perio oids, cataracts, on a prescriptio 500 ml	orificial d glaucom	ied for applications ermatitis, rosacea, na, or raised intraocu inetarsol idwest
■ SA1970 Special Authority for Subsidy  itial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve the eting the following criteria:  ioth:  1 Patient has atopic dermatitis on the eyelid; and  2 Patient has at least one of the following contraindicating documented epidermal atrophy, documented allergy the pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUCK Soln 2.3% with trolamine laurilsulfate and fluorescein social CALICYLIC ACID  Powder — Only in combination	ophthalmologist or an als valid without furth ons to topical cortico topical corticostero DRESCEIN — Only odium	osteroids: perio oids, cataracts, on a prescriptio 500 ml	orificial d glaucom	ied for applications ermatitis, rosacea, na, or raised intraocu inetarsol idwest
SA1970 Special Authority for Subsidy initial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve neeting the following criteria: ioth:  1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindicatid documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUCK Soln 2.3% with trolamine laurilsulfate and fluorescein socialicyLic ACID Powder — Only in combination	ophthalmologist or ar als valid without furth ons to topical cortico o topical corticostero DRESCEIN – Only of dium	osteroids: perio oids, cataracts, on a prescriptio 500 ml	less notificial de glaucom  Pi Moid – Pla	ied for applications ermatitis, rosacea, na, or raised intraocu inetarsol idwest
■ SA1970 Special Authority for Subsidy  itial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve neeting the following criteria:  ioth:  1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindicatid documented epidermal atrophy, documented allergy to pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUCK Soln 2.3% with trolamine laurilsulfate and fluorescein socialicyLIC ACID  Powder − Only in combination	ophthalmologist or ar als valid without furth ons to topical cortico o topical corticostero DRESCEIN — Only of dium	osteroids: perio oids, cataracts, on a prescriptio 500 ml 250 g cal Corticostero	less notificial deglaucom  Pi Moid – Pla	ermatitis, rosacea, na, or raised intraocu inetarsol idwest in or collodion flexible
■ SA1970 Special Authority for Subsidy Initial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve neeting the following criteria: Into Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindicating documented epidermal atrophy, documented allergy to pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUCK Soln 2.3% with trolamine laurilsulfate and fluorescein social LICYLIC ACID Powder — Only in combination	ophthalmologist or ar als valid without furth ons to topical cortico o topical corticostero DRESCEIN — Only of dium	osteroids: perio oids, cataracts, on a prescriptio 500 ml 250 g cal Corticostero	less notificial deglaucom  Pi Moid – Pla	ermatitis, rosacea, na, or raised intraocu inetarsol idwest in or collodion flexible
■ SA1970 Special Authority for Subsidy Initial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve neeting the following criteria: Into Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindicating documented epidermal atrophy, documented allergy the pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUCK Soln 2.3% with trolamine laurilsulfate and fluorescein social LICYLIC ACID Powder — Only in combination	ophthalmologist or are als valid without furth ons to topical cortico topical corticostero or topical corticostero ORESCEIN — Only odium	osteroids: perio oids, cataracts, on a prescriptio 500 ml 250 g cal Corticostero 100 g cal Corticostero	less notificial deglaucom  Pi Moid – Pla  Moid – Pla	ied for applications ermatitis, rosacea, na, or raised intraocu inetarsol idwest in or collodion flexible idwest in
■ SA1970 Special Authority for Subsidy Initial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve neeting the following criteria: Into Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindicating documented epidermal atrophy, documented allergy to pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUCK Soln 2.3% with trolamine laurilsulfate and fluorescein social LICYLIC ACID Powder — Only in combination	ophthalmologist or are als valid without furth ons to topical cortico topical corticostero or topical corticostero ORESCEIN — Only odium	osteroids: perio oids, cataracts, on a prescriptio 500 ml 250 g cal Corticostero	less notificial deglaucom  Pi Moid – Pla  Moid – Pla	ermatitis, rosacea, na, or raised intraocu inetarsol idwest in or collodion flexible
■ SA1970 Special Authority for Subsidy Initial application only from a dermatologist, paediatrician, of a dermatologist, paediatrician or ophthalmologist. Approve neeting the following criteria: Into Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindicating documented epidermal atrophy, documented allergy the pressure.  INE TAR WITH TROLAMINE LAURILSULFATE AND FLUCK Soln 2.3% with trolamine laurilsulfate and fluorescein social LICYLIC ACID Powder — Only in combination	ophthalmologist or are als valid without furth ons to topical cortico topical corticostero DRESCEIN – Only odium	osteroids: perio oids, cataracts, on a prescriptio 500 ml 250 g cal Corticostero 100 g cal Corticostero 30 g OP	less notificial deglaucom  Pi Moid – Pla  Moid – Pla	ermatitis, rosacea, na, or raised intraocu inetarsol idwest in or collodion flexible idwest in

### **DERMATOLOGICALS**

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

### ⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

# **Scalp Preparations**

BETAMETHASONE VALERATE  * Scalp app 0.1%	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE  * Scalp app 0.05%	30 ml OP	✓ Dermol
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%3.23	100 ml OP	✓ Sebizole
		Sebizole

- a) Maximum of 100 ml per prescription
- b) Only on a prescription

### Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

SPF 50+

# **Wart Preparations**

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 77

#### **PODOPHYLLOTOXIN**

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

# Other Skin Preparations

### **Antineoplastics**

LIODO	MIIIODS	

IMIQUIMOD

Crm 5%, 250 mg sachet......21.72 24 **✓ Perrigo** 

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

# **Contraceptives - Non-hormonal**

# **Condoms**

^^	NDOMO			
	NDOMS	44.40		
	49 mm – Up to 144 dev available on a PSO		144	✓ Moments
۴	53 mm		10	✓ Moments
		11.64	144	✓ Moments
	<ul> <li>a) Maximum of 60 dev per prescription</li> </ul>			
	b) Up to 60 dev available on a PSO			_
÷	53 mm, 0.05 mm thickness		10	✓ Moments
		11.42	144	✓ Moments
	<ul> <li>a) Up to 60 dev available on a PSO</li> </ul>			
	<ul> <li>b) Maximum of 60 dev per prescription</li> </ul>			
+	53 mm, chocolate, brown	0.95	10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
+	53 mm, strawberry, red	0.95	10	✓ Moments
	•	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
÷	56 mm	0.97	10	✓ Moments
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
6	56 mm, 0.05 mm thickness	1 30	12	✓ Gold Knight
	oo min, o.oo min unoknood	15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO	10.07	177	- dola kingik
	b) Maximum of 60 dev per prescription			
÷	56 mm, 0.05mm thickness (bulk pack)	1/161	144	✓ Gold Knight
•	, , ,	14.01	144	▼ Gold Killgill
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO	0.07	40	/ W
÷	56 mm, 0.08 mm thickness		10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
÷	56 mm, 0.08 mm thickness, red		10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	<ul> <li>b) Maximum of 60 dev per prescription</li> </ul>			
÷	56 mm, chocolate	1.30	12	Gold Knight
		15.57	144	Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, strawberry	1.30	12	✓ Gold Knight
	•	15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			•
	b) Maximum of 60 dev per prescription			
÷	60 mm	1 42	12	✓ Gold Knight XL
•	V 111111111111111111111111111111111111	17.02	144	✓ Gold Knight XL
	a) Maximum of 60 dev per prescription	17.02	177	- GOIG KINGIN AL
	a) iviaximum of ou dev per prescription			

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

- a) Maximum of 60 dev per prescription
- b) Up to 60 dev available on a PSO

### Contraceptive Devices

### INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

* IUD 29.1 mm length × 23.2 mm width	.29.	80	)
--------------------------------------	------	----	---

✓ 7 MED NSHA Silver/ Copper Short

✓ Choice 380 7med Nsha Silver/ copper Short

✓ Choice TT380 Short

TT380 Standard

# ✓ Choice Load 375

✓ Choice

### Contraceptives - Hormonal

### Combined Oral Contraceptives

# **⇒SA0500** Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 Patient is on a Social Welfare benefit: or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- · on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### ETHINYLOFSTRADIOL WITH DESOGESTREL

\* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO.......10.00 84 ✓ Mercilon 28

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	<b>✓</b>	Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 84 tab available on a PSO	1.50	84	✓	Lo-Oralcon 20 ED
* Tab 30 mcg with levonorgestrel 150 mcg		63		
	(16.50)			Microgynon 30
<ul> <li>a) Higher subsidy of \$15.00 per 63 tab with Special Aut</li> <li>b) Up to 63 tab available on a PSO</li> <li>* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets</li> </ul>	-	the p	orevious pa	age
Up to 84 tab available on a PSO	1.50	84	/	Oralcon 30 ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to	1			
84 tab available on a PSO		84	✓	Brevinor 1/28
	16.33	112	✓	Brevinor-1 28 Day
				Norimin-1 28 Day
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U	n			
to 84 tab available on a PSO	•	84	1	Norimin
10 0 1 100 0 1 10 10 10 10 10 10 10 10 1	29.32	112		Norimin
	,			

# **Progestogen-only Contraceptives**

### ⇒SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
  - 1.1 Patient is on a Social Welfare benefit: or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED LEVONORGESTREI

*	Tab 30 mcg - Up to 84 tab available on a PSO		84  12	<ul><li>✓ Microlut</li><li>✓ Microlut</li></ul>
*	Subdermal implant (2 x 75 mg rods) — Up to 3 pack available on a PSO	.92	1	✓ Jadelle
	Jadelle to be Principal Supply on 1 December 2023	-		
ME	DROXYPROGESTERONE ACETATE			
	Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a PSO9.	.18	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84	✓ <u>N</u>	oriday 28	

# **Emergency Contraceptives**

### LEVONORGESTREL

- 1 ✓ Levonorgestrel BNM
  - a) Maximum of 2 tab per prescription
  - b) Up to 5 tab available on a PSO
  - c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

# Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- A maximum \$5.00 prescription charge (patient co-payment) may apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to any non contraceptive prescription charges that apply, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

### CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

\* Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs - Up 

✓ Ginet

Aci-Jel

✓ Clomazol

✓ Clomazol

# **Gynaecological Anti-infectives**

A(	ÆΊ	ГIС	AC	IJD	W	ITH	H H	ΥD	R	O)	XΥ	'C	)U	IN	0	ΙJ	NE	· A	NE	) R	IIC	ΛK	Ю	LEI	C.	ΑC	ΉL	)
----	----	-----	----	-----	---	-----	-----	----	---	----	----	----	----	----	---	----	----	-----	----	-----	-----	----	---	-----	----	----	----	---

Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate	
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP
(24.87)	

35 a OP

20 q OP

168

*	Vaginal crm 2% with applicators	3.85

MICONAZOLE NITRATE \* Vaginal crm 2% with applicator .......6.89 40 g OP ✓ Micreme

**CLOTRIMAZOLE** 

NYSTATIN Vaginal crm 100,000 u per 5 g with applicator(s) ......5.70 75 q OP Nilstat

# **Myometrial and Vaginal Hormone Preparations**

#### **ERGOMETRINE MALEATE**

Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	160.00	5	✓ DBL Ergometrine
OESTRIOL  * Crm 1 mg per g with applicator  Pessaries 500 mcg		15 g OP 15	✓ Ovestin ✓ Ovestin

OXYTOCIN – Up to 5 inj available on a PSO	
to the first and the first and the second of	

ATTOOM OP to o my available on a too		
Inj 5 iu per ml, 1 ml ampoule4.98	5	✓ 0
Inj 10 iu per ml, 1 ml ampoule5.98	5	✓ 0
		_

✓ Oxytocin GH S29

Subsidy (Manufacturer's Price)	Subsid	ully ised	Brand or Generic
\$	Per	1	Manufacturer

OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 ini available on a PSO

Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoule ....32.40 5 ✓ Syntometrine

# Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

40 test OP ✓ Smith BioMed Rapid **Pregnancy Test** 16.00

✓ David One Step Cassette **Pregnancy Test** 

# **Urinary Agents**

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 116

### 5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy ✓ Ricit \* Tab 5 mg .......4.79 100 Ricit to be Principal Supply on 1 December 2023

### ⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Fither:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

# Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

✓ Tamsulosin-Rex

### ⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

# **Other Urinary Agents**

### **OXYBUTYNIN**

100 ✓ Alchemy Oxybutynin

#### POTASSIUM CITRATE

Oral lig 3 mmol per ml - Special Authority see SA1083 on the Biomed 200 ml OP

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

### ⇒SA1083 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATI	Ł
-----------------------	---

* Grans eff 4 g sachets	3.50	28	✓ Ural
SOLIFENACIN SUCCINATE			
Tab 5 mg	2.05	30	<ul><li>Solifenacin Mylan</li></ul>
•			✓ Solifenacin Viatris
Tab 10 mg	3.72	30	✓ Solifenacin Mylan
			✓ Solifenacin Viatris
(Salifonacin Mulan Tab 5 mg to be delicted 1 December 1	2022)		

(Solifenacin Mylan Tab 5 mg to be delisted 1 December 2023) (Solifenacin Mylan Tab 10 mg to be delisted 1 December 2023)

# **Detection of Substances in Urine**

	C	F	П	Ή	C	)-7	ГС	L	IDI	N	Ε	
--	---	---	---	---	---	-----	----	---	-----	---	---	--

* Compound diagnostic sticks	7.50 (8.25)	50 test OP	Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	.13.92	100 test OP	Albustix

# **Obstetric Preparations**

# Antiprogesterones

MI	FF	PR	IST	ONF	

✓ Mifegyne	1	79.90	Tab 200 mg - Up to 15 tab available on a PSO
✓ Mifegyne	3	180.00	

(	Subsidy (Manufacturer's Price)		Fully ubsidised	Brand or Generic
	\$	Per	1	Manufacturer

# **Calcium Homeostasis**

CALCITONIN  * Inj 100 iu per ml, 1 ml ampoule121.00	5	✓ Miacalcic
CINACALCET - Special Authority see SA2170 below - Retail pharmacy		
Tab 30 mg - Wastage claimable42.06	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable84.12	28	✓ Cinacalet Devatis

**⇒SA2170** Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
  - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
  - 1.3 The patient is symptomatic; or
- 2 All of the following:
  - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
  - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
  - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate

**Renewal — (parathyroid carcinoma or calciphylaxis)** only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
  - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
  - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
  - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
  - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

	Subsidy	Fully	Brand or
(Manut	facturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

**Renewal — (secondary or tertiary hyperparathyroidism)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Fither:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

#### **ZOLEDRONIC ACID**

(Zoledronic acid Mylan Inj 4 mg per 5 ml, vial to be delisted 1 November 2023)

RETAMETHASONE SODILIM PHOSPHATE WITH RETAMETHASONE ACETATE

# Corticosteroids and Related Agents for Systemic Use

DLI	AMETHASONE SOCION FROSFIATE WITH BETAMETHASONE ACE	IAIL	
*	Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml19.20	5	
	(36.96)	)	Celestone
			Chronodose
DEX	(AMETHASONE		
*	Tab 0.5 mg - Up to 60 tab available on a PSO1.50	30	✓ Dexmethsone
*	Tab 4 mg - Up to 30 tab available on a PSO2.65	30	✓ Dexmethsone
	Oral liq 1 mg per ml	25 ml OP	Biomed
DEX	(AMETHASONE PHOSPHATE		
	Dexamethasone phosphate injection will not be funded for oral use.		
*	Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO7.86	10	✓ <u>Hameln</u>
*	Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO 13.10	10	✓ <u>Hameln</u>
FLU	DROCORTISONE ACETATE		
*	Tab 100 mcg11.46	100	✓ Florinef
	DROCORTISONE		
	Tab 5 mg	100	✓ Douglas
	Tab 20 mg20.32		✓ Douglas
	Inj 100 mg vial4.38		✓ Solu-Cortef
	a) Up to 5 inj available on a PSO		
	b) Only on a PSO		
MET	THYLPREDNISOLONE		
	Tab 4 mg112.00	100	✓ Medrol
	Tab 100 mg223.10		✓ Medrol

Tab 1 mg		Subsidy (Manufacturer's Price \$	e) Subs	Fully Brand or sidised Generic  Manufacturer
Inj 40 mg vial	METHYL PREDNISOLONE (AS SODIUM SUCCINATE)			
Inj 500 mg vial	,	22.30	1	
Inj 1 g vial	Inj 125 mg vial	34.10	1	
### ACTION OF THE PREDNISOLONE ACETATE   Inj 40 mg per ml, 1 ml vial	Inj 500 mg vial	26.88	1	
### ACTION OF THE PREDNISOLONE ACETATE   Inj 40 mg per ml, 1 ml vial	lni 1 a vial	32.84	1	✓ Solu-Medrol
Inj 40 mg per ml, 1 ml vial		32.04	1	• Solu-ivieuloi
REDNISOLONE		47.06	_	✓ Dono Modrol
Redipred   Restricted to children under 12 years of age.   Redipred   Restricted to children under 12 years of age.   Redipred   Restricted to children under 12 years of age.   Redipred   Redipred   Restricted to children under 12 years of age.   Redipred   Redi		47.06	5	▼ Depo-Medioi
Tab 1 mg	* Oral liq 5 mg per ml - Up to 30 ml available on a PSO	6.00	30 ml OP	✓ <u>Redipred</u>
Tab 2.5 mg	PREDNISONE			
Tab 5 mg				
Tab 20 mg				
ETRACOSACTRIN    Inj 250 mcg per ml, 1 ml ampoule				
Synacthen   Synacthen   William	* Tab 20 mg - Up to 30 tab available on a PSO	50.51	500	✓ Prednisone Clinect
VIK Synacthen   Synacthen   Synacthen   Synacthen   Synacthen   Synacthen   Synacthene   Retard   Synachhene   Synachhene   Retard   Synachhene   Synachhene   Synachhene   S	TETRACOSACTRIN			
Inj 1 mg per ml, 1 ml ampoule   690.00   1   Synacthen Depot   Synacthene   Retard \$29   Synacthene   Synac	★ Inj 250 mcg per ml, 1 ml ampoule	86.25	1	
FRIAMCINOLONE ACETONIDE  Inj 10 mg per ml, 1 ml ampoule				
Inj 10 mg per ml, 1 ml ampoule	k Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓ Synacthene
Inj 40 mg per ml, 1 ml ampoule	TRIAMCINOLONE ACETONIDE			
Sex Hormones Non Contraceptive  Androgen Agonists and Antagonists  EXPROTERONE ACETATE Tab 50 mg	Inj 10 mg per ml, 1 ml ampoule	21.42	5	✓ Kenacort-A 10
Androgen Agonists and Antagonists  CYPROTERONE ACETATE  Tab 50 mg	Inj 40 mg per ml, 1 ml ampoule	52.63	5	✓ Kenacort-A 40
CYPROTERONE ACETATE       14.37       50       ✓ Siterone         Tab 50 mg       28.03       50       ✓ Siterone         TESTOSTERONE       225.00       30       ✓ Androderm         TESTOSTERONE CIPIONATE       85.00       1       ✓ Depo-Testosterone         Inj 100 mg per ml, 10 ml vial.       85.00       1       ✓ Depo-Testosterone         393.00       ✓ Taro-       Testosterone       329	Sex Hormones Non Contraceptive			
Tab 50 mg       14.37       50       ✓ Siterone         Tab 100 mg       28.03       50       ✓ Siterone         ESTOSTERONE       225.00       30       ✓ Androderm         ESTOSTERONE CIPIONATE       85.00       1       ✓ Depo-Testosterone         Inj 100 mg per ml, 10 ml vial       85.00       1       ✓ Taro-Testosterone         TESTOSTERONE ESTERS       1       Testosterone       529	Androgen Agonists and Antagonists			
Tab 100 mg       28.03       50       ✓ Siterone         ESTOSTERONE       225.00       30       ✓ Androderm         ESTOSTERONE CIPIONATE       85.00       1       ✓ Depo-Testosterone         Inj 100 mg per ml, 10 ml vial       393.00       ✓ Taro-Testosterone         TESTOSTERONE ESTERS       1       1	CYPROTERONE ACETATE			
TESTOSTERONE Patch 5 mg per day	Tab 50 mg	14.37	50	✓ <u>Siterone</u>
Patch 5 mg per day	Tab 100 mg	28.03	50	✓ <u>Siterone</u>
Patch 5 mg per day	ESTOSTERONE			
ESTOSTERONE CIPIONATE Inj 100 mg per ml, 10 ml vial		225.00	30	✓ Androderm
Inj 100 mg per ml, 10 ml vial				
TESTOSTERONE ESTERS			1	✓ Taro-
				1 631031610116 025
	TESTOSTERONE ESTERS			
		12.98	1	✓ Sustanon Ampoules

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
TESTOSTERONE UNDECANOATE				
Cap 40 mg - Subsidy by endorsement	21.00	60	1	Andriol Testocaps
	35.00	100	1	Steril-Gene S29
Subsidy by endorsement – subsidised for patients who w 1 November 2021 and the prescription is endorsed according where there exists a record of prior dispensing of testost Inj 250 mg per ml, 4 ml vial	rdingly. Pharmacists erone undecanoate c	may a	annotate t mg in the	the prescription as endorsed

# **Hormone Replacement Therapy - Systemic**

# Oestrogens

-	STRADIOL			
*	Tab 1 mg		28 OP	
		(11.10)		Estrofem
*	Tab 2 mg		28 OP	
		(11.10)		Estrofem
	Patch 50 mcg per 24 hours	7.04	4	✓ Climara
	a) No more than 1 patch per week			
	b) Only on a prescription		_	<b>.</b>
	Patch 25 mcg per day		8	✓ Estradot
		9.85		<ul><li>Estradiol TDP Mylan</li></ul>
		13.50		✓ Estraderm MX S29
	<ul> <li>a) No more than 2 patch per week</li> </ul>			
	b) Only on a prescription			
	Patch 50 mcg per day	7.04	8	✓ Estradot 50 mcg
		10.75		<ul><li>Estradiol TDP Mylan</li></ul>
				<ul><li>Estradiol Viatris</li></ul>
		14.50		✓ Estraderm MX S29
	a) No more than 2 patch per week			
	b) Only on a prescription			
	Patch 75 mcg per day	7.91	8	✓ Estradot
		11.88		<ul><li>Estradiol TDP Mylan</li></ul>
				<ul><li>Estradiol Viatris</li></ul>
	a) No more than 2 patch per week			
	b) Only on a prescription			
	Patch 100 mcg per day	7.91	8	✓ Estradot
		12.95		<ul><li>Estradiol TDP Mylan</li></ul>
				<ul><li>Estradiol Viatris</li></ul>
		15.50		✓ Estraderm MX S29
	a) No more than 2 patch per week			
	b) Only on a prescription			
OF	STRADIOL VALERATE			
	Tab 1 mg	12.36	84	✓ Progynova
	Tab 2 mg		84	✓ Progynova
	·		•	
-	STROGENS	2.01	00	
本	Conjugated, equine tab 300 mcg	(17.50)	28	Premarin
*	Conjugated, equine tab 625 mcg	, ,	28	FIEIIIaIIII
*	Conjugated, equilie tab 020 mog	(17.50)	20	Premarin
		(17.30)		i IGIIIaiiii

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	(Manufacturer's Frice)	Per	
rogestogens			
EDROXYPROGESTERONE ACETATE			
Tab 2.5 mg	4.69	30	✓ Provera
•	8.75	56	✓ Provera
Tab 5 mg	9.80	56	✓ Provera
	17.50	100	✓ Provera
Tab 10 mg	8.94	30	✓ Provera
Progestogen and Oestrogen Combined Prepara	ntions		
ESTRADIOL WITH NORETHISTERONE			
Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP	
	(18.10)		Kliovance
Tab 2 mg with 1 mg norethisterone acetate		28 OP	
	(18.10)		Kliogest
Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg			
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP	
	(18.10)		Trisequens
Other Oestrogen Preparations			
ESTRIOL			_
Tab 2 mg	7.70	30	Ovestin
Other Progestogen Preparations			
VONORGESTREL			_
Intra-uterine device 52 mg		1	✓ Mirena
Intra-uterine device 13.5 mg	215.60	1	Jaydess
EDROXYPROGESTERONE ACETATE			
Tab 100 mg	116.15	100	Provera HD
ORETHISTERONE			
Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Primolut N
ROGESTERONE			
Cap 100 mg	14.85	30	✓ <u>Utrogestan</u>
hyroid and Antithyroid Agents			
ARBIMAZOLE	7.50	100	✓ Nee Mereersts
Tab 5 mg	/.5b	100	✓ Neo-Mercazole
VOTHYROXINE			
Tab 25 mcg		90	✓ Synthroid
Tab 50 mcg		28	✓ Mercury Pharma
	5.79	90	✓ Synthroid
	64.28	1,000	
Tab 100 mcg		28	✓ Mercury Pharma
	6.01	90	✓ Synthroid
	66.78	1,000	✓ Eltroxin
OPYLTHIOURACIL - Special Authority see SA1199 on the n	ext page - Retail ph	armacı	y
Tab 50 mg		100	✓ PTU S29

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or
(Manufacturer's Pric	ce)	Subsidised	Generic
\$	Pe	r 🗸	Manufacturer

### ⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

# **Trophic Hormones**

### **Growth Hormones**

SO	MATROPIN (OMNITROPE) - Special Authority see SA2032 below	<mark>v</mark> – Retail pharm	acy	
*	Inj 5 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
				✓ Omnitrope S29 S29
*	Inj 15 mg cartridge	139.50	1	✓ Omnitrope
				✓ Omnitrope S29 S29

### ⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g., cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
  - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
  - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
  - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
  - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
  - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
	Per	1	Manufacturer	

continued...

for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

**Renewal — (Turner syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred

**Initial application** — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
  - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- (Height(cm)/plasma creatinine (umol/l)  $\times$  40 = corrected GFR (ml/min/1.73m<sup>2</sup> in a child who may or may not be receiving dialysis; or
- 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Fither:
  - 5.1 Both:
    - 5.1.1 The patient is aged two years or older: and
    - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
  - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
 \$	Per	•	Manufacturer	

continued...

than or equal to 0.5 standard deviations in the preceding 12 months.

**Initial application — (adults and adolescents)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and
  - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
  - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or

### 2 All of the following:

- 2.1 The patient has been treated with somatropin for more than 12 months; and
- 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
- 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
- 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or

### 3 All of the following:

- 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
- 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3.3 The patient has severe growth hormone deficiency (see notes); and
- 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth

	Subsidy	F	ully	Brand or
(1)	Manufacturer's Price)	Subsidi	sed	Generic
	\$	Per	•	Manufacturer

continued...

hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

# **GnRH Analogues**

GOSERELIN			
Implant 3.6 mg, syringe	65.68	1	Teva
Implant 10.8 mg, syringe	122.37	1	✓ Teva

#### I FUPRORFI IN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

injoiro ing promod dadi onambor cyringo	ringilior dubbidy or		
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe	- Higher subsidy		

Ini 3.75 mg prefilled dual chamber syringe — Higher subsidy of

(591.68) Lucrin Depot 3-month

# Vasopressin Agonists

DESMOPRESSIN  Wafer 120 mcg47.00	30	✓ Minirin Melt
DESMOPRESSIN ACETATE		
Tab 100 mcg25.00	30	✓ Minirin
Tab 200 mcg54.45	30	✓ Minirin
▲ Nasal spray 10 mcg per dose34.95	6 ml OP	✓ Desmopressin- PH&T
Inj 4 mcg per ml, 1 ml67.18	10	✓ Minirin

# Other Endocrine Agents

#### **CABERGOLINE**

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
Dostinex	2	waived by Special Authority see SA2070 below4.43
✓ Dostinex	8	17.94

### ⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy (Manufacturer's Price	) Per	Fully Subsidised	Brand or Generic Manufacturer	
•	rei		Manuacturer	

continued...

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly\*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with \* is an unapproved indication.

CLO	MIF	FNF	CITI	RATE

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE Cap 250 mg	558.00	50	✓ Metopirone

	Subsidy	\ 0	Fully Brand or
	(Manufacturer's Pri \$	Per	sidised Generic  Manufacturer
	<u> </u>		
Anthelmintics			
ALBENDAZOLE - Special Authority see SA1318 below - Retail	pharmacy		
Tab 400 mg	469.20	60	✓ Eskazole S29
⇒SA1318 Special Authority for Subsidy			
Initial application only from an infectious disease specialist or c	linical microbiologi	st. Approval	s valid for 6 months where the
patient has hydatids.			0 11 1 11 1 1
Renewal only from an infectious disease specialist or clinical mic remains appropriate and the patient is benefitting from the treatm		ovais valid toi	r 6 months where the treatment
MEBENDAZOLE – Only on a prescription	iciit.		
Tab 100 mg	7 97	6	✓ <u>Vermox</u>
Oral lig 100 mg per 5 ml		15 ml	· <u>voimox</u>
	(7.83)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	✓ Biltricide
Antibacterials			
<ul> <li>a) For topical antibacterials, refer to DERMATOLOGICALS, pag</li> <li>b) For anti-infective eye preparations, refer to SENSORY ORGA</li> </ul>			
Cephalosporins and Cephamycins			
CEFACLOR MONOHYDRATE			
Cap 250 mg		100	✓ Ranbaxy-Cefaclor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.75	100 ml	✓ Ranbaxy-Cefactor
CEFALEXIN			
Cap 250 mg		20 20	✓ Cephalexin ABM
Cap 500 mgGrans for oral liq 25 mg per ml – Wastage claimable		20 100 ml	✓ <u>Cephalexin ABM</u> ✓ Flynn
Grans for oral liq 50 mg per ml — Wastage claimable		100 ml	✓ Flynn
Grans for oral liq 50 mg per mi vvastage dialinable	11.75	100 1111	✓ Cefalexin Sandoz
CEFAZOLIN - Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance with	a Te Whatu Ora H	ospital appro	oved protocol and the prescription
is endorsed accordingly.			_
Inj 500 mg vial		5	✓ Cefazolin-AFT
Inj 1 g vial		5	✓ Cefazolin-AFT
Inj 2 g vial	7.09	5	✓ Cefazolin-AFT
CEFTRIAXONE – Subsidy by endorsement			
<ul><li>a) Up to 10 inj available on a PSO</li><li>b) Subsidised only if prescribed for a dialysis or cystic fibros</li></ul>	is nationt or the tr	eatment of a	onorrhoes or the treatment of
pelvic inflammatory disease, or the treatment of suspecte endorsed accordingly.			
Inj 500 mg vial	0.79	1	✓ Ceftriaxone-AFT
Inj 1 g vial		5	✓ Ceftriaxone-AFT
CEFUROXIME AXETIL – Subsidy by endorsement			
Only if prescribed for prophylaxis of endocarditis and the pre	scription is endors	ed according	ıly.
Tab 250 mg		50	✓ Zinnat
(Zinnet Teh 050 mg to be delicted 1 March 0004)			

(Zinnat Tab 250 mg to be delisted 1 March 2024)

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	✓	Manufacturer

### **Macrolides**

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable1	6.97	15 ml	✓ Zithromax

### ⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome\*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome\*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms\*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with \* are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis\*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with \* are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with \* are unapproved indications

CLARITHROMYCIN – Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

**⇒SA1857** Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy	Fi	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

**Initial application — (Helicobacter pylori eradication)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	<ul><li>E-Mycin</li></ul>
a) Up to 20 tab available on a PSO     b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 200 mg per 5 ml	5.00	100 ml	✓ E-Mycin
<ul> <li>a) Up to 300 ml available on a PSO</li> <li>b) Up to 2 x the maximum PSO quantity for RFPP</li> <li>c) Wastage claimable</li> </ul>			
Grans for oral liq 400 mg per 5 mla) Up to 200 ml available on a PSO b) Wastage claimable	6.77	100 ml	✓ E-Mycin
ROXITHROMYCIN			
Tab 150 mg	13.19	50	✓ <u>Arrow-</u> <u>Roxithromycin</u>
Tab 300 mg	25.00	50	✓ <u>Arrow-</u> <u>Roxithromycin</u>

	Subsidy		Fully	
	(Manufacturer's Price \$	e) Sub Per	osidised •	Generic Manufacturer
Dania III	<u> </u>			
Penicillins				
AMOXICILLIN			_	
Cap 250 mg	43.45	500	/	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP			_	
Cap 500 mg	66.44	500	•	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP	0.00	100		Almhamau 105
Grans for oral liq 125 mg per 5 ml	2.22	100 ml	•	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable	0.01	100 ml	./	Alphamay 050
Grans for oral liq 250 mg per 5 ml	2.01	100 1111	•	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable Inj 250 mg vial	15 07	10	1	Ibiamox
Inj 500 mg vial		10		Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO		10		Ibiamox
, , ,		10		ibiamox
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab available on a PSO	1.50	10	./	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25		10	•	Curam Duo 500/125
per ml	•	100 ml	1	Augmentin
•	0.50	100 1111	•	Augmenum
<ul><li>a) Up to 200 ml available on a PSO</li><li>b) Wastage claimable</li></ul>				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5	ma			
per ml — Up to 200 ml available on a PSO	•	100 ml OP	1	Curam
·	2.20	100 1111 01	•	Ourain
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj	075 07	10	./	Bicillin LA
available on a PSO	3/5.9/	10	•	DICIIIII LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial - Up to 5 inj available on a P	SO 16.50	10	•	Sandoz
FLUCLOXACILLIN			_	
Cap 250 mg - Up to 30 cap available on a PSO		250		Flucloxacillin-AFT
Cap 500 mg – Up to 30 cap available on a PSO		500		Flucloxacillin-AFT
Grans for oral liq 25 mg per ml	3.29	100 ml	•	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable	0.60	100 ml	./	A ET
Grans for oral liq 50 mg per ml	3.00	100 ml	٧	<u>AFT</u>
<ul><li>a) Up to 200 ml available on a PSO</li><li>b) Wastage claimable</li></ul>				
Inj 250 mg vial	17 56	10	1	Flucloxin
Inj 500 mg vial		10		Flucioxin
Inj 1 g vial – Up to 5 inj available on a PSO		5		Flucil
, • ,		-		

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)		Fully Subsidised	Generic	
	\$	Per		Manufacturer	
PHENOXYMETHYLPENICILLIN (PENICILLIN V)					
Cap 250 mg - Up to 30 cap available on a PSO	3.84	50	1	Cilicaine VK	
Cap 500 mg	6.86	50	✓	Cilicaine VK	
a) Up to 20 cap available on a PSO					
b) Up to 2 x the maximum PSO quantity for RFPP					
Grans for oral liq 125 mg per 5 ml	3.40	100 m	<b>✓</b>	<u>AFT</u>	
a) Up to 200 ml available on a PSO					
b) Wastage claimable					
Grans for oral liq 250 mg per 5 ml	4.24	100 m	✓	<u>AFT</u>	
a) Up to 300 ml available on a PSO					
b) Up to 2 x the maximum PSO quantity for RFPP					
c) Wastage claimable					

# **Tetracyclines**

DO	XYCYCLINE			
*	Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	Doxine
MIN	NOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	19.32	100	
		(52.04)		Minomycin

### **⇒SA1355** Special Authority for Manufacturers Price

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

 $\label{temperature} \textbf{TETRACYCLINE} \ - \textbf{Special Authority see} \ \frac{\textbf{SA1332 below}}{\textbf{below}} - \textbf{Retail pharmacy}$ 

### ⇒SA1332 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

### Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 71

#### **CIPROFLOXACIN**

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 050 mg. Unita 5 tab available on a DCO	0.40	00	✓ Cipflox
Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	Cipilox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
• ,	4.25	10	✓ Ciprofloxacin -
			Torrent S29
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	✓ Manufacturer
CLINDAMYCIN			
Cap hydrochloride 150 mg	5.30	24	✓ Dalacin C
Inj 150 mg per ml, 4 ml ampoule	35.10	10	✓ <u>Hameln</u>
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S	ubsidy by endorseme	nt	
Only if prescribed for dialysis or cystic fibrosis patient and the	prescription is endor	sed a	accordingly.
Inj 150 mg	65.00	1	✓ Colistin-Link
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement	95.00	5	✓ DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.	r complicated urinary	tract	et infection and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	✓ Wockhardt S29
	182.00	10	✓ Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.	r complicated urinary	tract	et infection and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	18.38	10	✓ Pfizer
	87.50	50	✓ Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.	r complicated urinary	tract	et infection and the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Retail No patient co-payment payable	pharmacy		
Tab 400 mg	42.00	5	✓ Avelox

### ⇒SA1740 Special Authority for Subsidy

**Initial application** — (**Tuberculosis**) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Active tuberculosis\*: and
  - 1.2 Any of the following:
    - 1.2.1 Documented resistance to one or more first-line medications; or
    - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
    - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
    - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
    - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.\*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with \* are unapproved indications.

**Renewal** only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium\* and is symptomatic; and
- 2 Fither:
  - 2.1 Has tried and failed to clear infection using azithromycin; or
  - 2.2 Has laboratory confirmed azithromycin resistance; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer continued... 3 Treatment is only for 7 days. Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eve injury and treatment is for 5 days only. Note: Indications marked with \* are unapproved indications. PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy ✓ Humatin S29 16 ⇒SA1689 Special Authority for Subsidy Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria: Either: 1 Patient has confirmed cryptosporidium infection; or 2 For the eradication of Entamoeba histolyica carriage. Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria: Fither: 1 Patient has confirmed cryptosporidium infection; or 2 For the eradication of Entamoeba histolyica carriage. PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy 30 ✓ Daraprim S29 ⇒SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of age. SODIUM FUSIDATE [FUSIDIC ACID] ✓ Fucidin Tab 250 mg ......135.70 SULFADIAZINE SODIUM - Special Authority see SA1331 below - Retail pharmacy 56 Tab 500 mg .......543.20 ✓ Wockhardt \$29 ⇒SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of age. **TOBRAMYCIN** Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement......18.50 ✓ Tobramycin Mylan ✓ Viatris Only if prescribed for dialysis or cystic fibrosis patient and the prescription is endorsed accordingly. Solution for inhalation 60 mg per ml. 5 ml - Subsidy by

✓ fully subsidised **Principal Supply** 

a) Wastage claimable

endorsement......395.00

c) Tobramycin BNM to be Principal Supply on 1 December 2023 (Tobramycin Mylan Inj 40 mg per ml, 2 ml vial to be delisted 1 January 2024)

b) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly.

✓ Tobramycin BNM

56 dose

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
TRIMETHOPRIM				
* Tab 300 mg - Up to 30 tab available on a PSO	18.55	50	✓ :	<u>TMP</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	AZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - U	Jo			
to 30 tab available on a PSO		500	1	Trisul
* Oral lig 8 mg sulphamethoxazole 40 mg per ml - Up to 200	ml		,	
available on a PSO	2.97	100 n	nl 🗸	Deprim
VANCOMYCIN - Subsidy by endorsement				•
Only if prescribed for a dialysis or cystic fibrosis patient or for	r prophylaxis of endo	cardit	is or for trea	atment of Clostridium
difficile following metronidazole failure and the prescription is				
Inj 500 mg vial		<b>1</b>	✓	Mylan

### **Antifungals**

- a) For topical antifungals refer to DERMATOLOGICALS, page 72
- b) For topical antifungals refer to GENITO URINARY, page 84

#### FLUCONAZOLE

Cap 50 mg	4.10	28	✓ Mylan
Mylan to be Principal Supply on 1 December 2023 Cap 150 mg		1	✓ Mylan
Mylan to be Principal Supply on 1 December 2023		00	,
Cap 200 mgMylan to be Principal Supply on 1 December 2023	8.90	28	✓ Mylan
Powder for oral suspension 10 mg per ml — Special Authority	100.00	05 1	4 D:0
see SA1359 below – Retail pharmacy	129.02	35 ml	✓ Diflucan

### ⇒SA1359 Special Authority for Subsidy

**Initial application** — **(Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Initial application — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised: and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

### Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer	
ITRACONAZOLE Cap 100 mg	6.83	15	<b>√</b> 1	trazole	
Oral lig 10 mg per ml – Special Authority see SA1322 below		13	• •	li azoie	
Retail pharmacy	141.80 1	50 ml (	OP 🗸 S	Sporanox	

### ⇒SA1322 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

# KETOCONAZOLE

Tab 200 mg - PCT	CBS	30	✓ Burel S29
		100	✓ Strides Shasun S29
			✓ Taro S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - F	etail pharmacy		
Tab modified-release 100 mg	206.00	24	✓ Posaconazole Juno
Oral liq 40 mg per ml	342.51	105 ml OP	✓ Devatis

### **⇒SA1285** Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy\*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression\* and requires on going posaconazole treatment.

Note: \* Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

### TERRINAFINE

* Tab 250 mg	8.97	84	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 on the next page	je – Retail phar	macy	
Tab 50 mg	91.00	56	✓ Vttack
Tab 200 mg	350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml – Wastage			
claimable	1,523.22	70 ml	✓ Vfend

Subsidy		Fully	Brand or	
(Manufacturer's Price	e) (	Subsidised	Generic	
\$	Per	1	Manufacturer	

### ⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient has proven or probable invasive aspergillus infection; or
  - 3.2 Patient has possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

### All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
  - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

### **Antimalarials**

PRIMAQUINE – Special Authority see SA1684 belo	ow – Retail pharmacy		
Tab 15 mg	400.00	100	✓ Sanofi
,			Primaquine S29

### ⇒SA1684 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

**Renewal** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

# **Antitrichomonal Agents**

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.16	10	✓ Arrow-Ornidazole

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Antituberculotics and Antileprotics** Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status. BEDAQUILINE - Special Authority see SA2244 below - Retail pharmacy No patient co-payment payable 24 OP ✓ Sirturo ⇒SA2244 Special Authority for Subsidy Initial application — (multi-drug resistant tuberculosis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both: 1 The person has multi-drug resistant tuberculosis (MDR-TB); and 2 Manatū Hauora - Ministry of Health's Tuberculosis Clinical Network has reviewed the individual case and recommends bedaquiline as part of the treatment regimen. CLOFAZIMINE - Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. ✓ Lamprene S29 \* Cap 50 mg.......442.00 100 CYCLOSERINE - Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician. 60 ✓ Cyclorin S29 DAPSONE - Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist 100 ✓ Dapsone 100 ✓ Dapsone ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician Tab 100 mg ......85.73 100 ✓ EMB Fatol \$29 56 Tab 400 mg .......49.34 ✓ Myambutol S29 ISONIAZID - Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician 100 ✓ PSM ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician 100 Rifinah 100 ✓ Rifinah

INFECTIONS - AGENTS FOR SYSTEMIC USE					
	Subsidy (Manufacturer's Price) \$	Subs Per	Fully Brand or sidised Generic ✓ Manufacturer		
LINEZOLID - Special Authority see SA2234 below - Retail phar	macy				
No patient co-payment payable			<i>4</i> <b>–</b>		
Tab 600 mg Oral lig 20 mg per ml		10 150 ml	✓ Zyvox ✓ Zyvox		
	1,079.00	150 1111	♥ Zyvox		
⇒SA2234   Special Authority for Subsidy   Initial application — (multi-drug resistant tuberculosis) from	any relevant practition	oner Anni	rovals valid for 18 months for		
applications meeting the following criteria:	any reservant praesint				
Both:					
<ol> <li>The person has multi-drug resistant tuberculosis (MDR-TI</li> <li>Manatū Hauora - Ministry of Health's Tuberculosis Clinica linezolid as part of the treatment regimen.</li> </ol>		ed the indi	vidual case and recommends		
PARA-AMINO SALICYLIC ACID - Retail pharmacy-Specialist					
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendation respiratory physician</li> </ul>		disease spe	ecialist, clinical microbiologist or		
Grans for oral liq 4 g sachet	280.00	30	✓ Paser S29		
PROTIONAMIDE - Retail pharmacy-Specialist					
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendation respiratory physician</li> </ul>	ion of, an infectious c	disease spe	ecialist, clinical microbiologist or		
Tab 250 mg	305.00	100	✓ Peteha S29		
PYRAZINAMIDE - Retail pharmacy-Specialist					
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommendation respiratory physician</li> </ul>	ion of, an infectious c	disease phy	ysician, clinical microbiologist or		
* Tab 500 mg	64.95	100	✓ AFT-Pyrazinamide		
RIFABUTIN – Retail pharmacy-Specialist			•		
a) No patient co-payment payable					
b) Prescriptions must be written by, or on the recommendation	ion of, an infectious o	disease phy	sician, respiratory physician or		
gastroenterologist	050.71	20	✓ Musehutin		
* Cap 150 mg	353.71	30	✓ Mycobutin		
RIFAMPICIN – Subsidy by endorsement					
<ul> <li>a) No patient co-payment payable</li> <li>b) For confirmed recurrent Staphylococcus aureus infection antimicrobial based on susceptibilities and the prescription Retail pharmacy - Specialist. Specialist must be an interpaediatrician, or public health physician.</li> </ul>	n is endorsed accord	dingly; can	be waived by endorsement -		
* Cap 150 mg	58.54	100	✓ Rifadin		
Rifadin to be Principal Supply on 1 December 2023					
* Cap 300 mg	122.06	100	✓ Rifadin		
Rifadin to be Principal Supply on 1 December 2023  * Oral liq 100 mg per 5 ml	12.60	60 ml	✓ Rifadin		

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

# **Antivirals**

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 259

# **Hepatitis B Treatment**

* Tab 0.5 mg	12.04	30	✓ Entecavir (Rex)
v	52.00		<ul><li>✓ Entecavir Mylan</li><li>✓ Entecavir Sandoz</li></ul>
(Entecavir Mylan Tab 0.5 mg to be delisted 1 March 2024)			

(Entecavir Sandoz Tab 0.5 mg to be delisted 1 March 2024)

LAMIVUDINE – Special Authority see SA1685 below – Retail pharmacy

 Tab 100 mg
 12.06
 28
 ✓ Zetlam

 Oral liq 5 mg per ml
 270.00
 240 ml OP
 ✓ Zeffix

#### ⇒SA1685 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA2139, page 113

Mylan

✓ Tenofovir Disoproxil

 Tenofovir Disoproxi Viatris

(Tenofovir Disoproxil Mylan Tab 245 mg (300 mg as a maleate) to be delisted 1 February 2024)

# **Herpesvirus Treatments**

ACICLOVIR		
* Tab dispersible 200 mg1	.78 25	✓ Lovir
* Tab dispersible 400 mg5	5.81 56	✓ Lovir
* Tab dispersible 800 mg6	35	✓ Lovir
VALACICLOVIR		
Tab 500 mg6	6.50 30	✓ Vaclovir
Tab 1,000 mg13	3.76 30	✓ Vaclovir
/ALGANCICLOVIR - Special Authority see SA1993 below - Retail pharr	nacy	
Tab 450 mg132	2.00 60	✓ Valganciclovir
-		Mylan
		✓ Valganciclovir
		Viatris

(Valganciclovir Mylan Tab 450 mg to be delisted 1 February 2024)

⇒SA1993 Special Authority for Subsidy

**Initial application** — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for

Subsidy		Fully	Brand or
(Manufacturer's Price)		sidised	Generic
\$	Per	1	Manufacturer

continued...

applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
  - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
  - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
  - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
  - 1 Patient has undergone a lung transplant; and
  - 2 Either:
    - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
    - 2.2 The recipient is cytomegalovirus positive; and
  - 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

# **Hepatitis C Treatment**

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg ......24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 **✓ Harvoni** 

### ⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website <a href="http://www.pharmac.govt.nz/maviret">http://www.pharmac.govt.nz/maviret</a> or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990.

Email: hepcpanel@pharmac.govt.nz

# **HIV Prophylaxis and Treatment**

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disporoxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 113 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

*	Tab 200	mg with	tenofovir	disoproxil	245 r	ng (300	) mg as a

maleate) 15.45

Emtricitabine Mylan ✔ <u>Tenofovir Disoproxil</u> Emtricitabine Viatr

(Tenofovir Disoproxil Emtricitabine Mylan Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a maleate) to be delisted 1 November 2023)

### ⇒SA2138 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

continued...

✓ Tenofovir Disoproxil

30

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical quidelines:

https://ashm.org.au/HIV/PrEP/

**Renewal** from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

# **COVID-19 Treatments**

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

# Antiretrovirals

#### ⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

**Renewal — (Confirmed HIV)** only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

**Initial application — (Percutaneous exposure)** only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

# Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA2139 on the previous pa	age – Retail pharm	nacy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA2139 on the previous p	oage – Retail pharr	macy	
Tab 200 mg	770.00	60	✓ Intelence

	Subsidy (Manufacturer's I \$	Price) Subs Per	Fully Brand or idised Generic  Manufacturer
EVIRAPINE – Special Authority see SA2139 on page 113 – F Tab 200 mg		60	✓ <u>Nevirapine</u> <u>Alphapharm</u>
Oral suspension 10 mg per ml	203.55	240 ml OP	<ul><li>✓ Nevirapine Viatris</li><li>✓ Viramune</li><li>Suspension</li></ul>
Nucleosides Reverse Transcriptase Inhibitors			
BACAVIR SULPHATE - Special Authority see SA2139 on pa	ge 113 – Retail p	narmacy	
Tab 300 mg Oral liq 20 mg per ml	180.00	60 240 ml OP	<ul><li>✓ Ziagen</li><li>✓ Ziagen</li></ul>
BACAVIR SULPHATE WITH LAMIVUDINE – Special Authori Note: abacavir with lamivudine (combination tablets) count anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	29.50	30	✓ <u>Abacavir/</u> <u>Lamivudine</u> Viatris
FAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOF	PROXIL - Specia	Authority see	SA2139 on page 113 – Retail
narmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil anti-retroviral Special Authority	counts as three a	nti-retroviral me	dications for the purposes of t
Tab 600 mg with emtricitabine 200 mg and tenofovir disopr 245 mg (300 mg as a maleate)		30	✓ Mylan ✓ Viatris
Mylan Tab 600 mg with emtricitabine 200 mg and tenofovir dis 023)	oproxil 245 mg (3	00 mg as a mal	
MTRICITABINE – Special Authority see SA2139 on page 113 Cap 200 mg		cy 30	✓ Emtriva
AMIVUDINE - Special Authority see SA2139 on page 113 - F			
Tab 150 mg		60	✓ Lamivudine Alphapharm
Oral liq 10 mg per mlamivudine Alphapharm Tab 150 mg to be delisted 1 Novemb		240 ml OP	✓ Lamivudine Viatris ✓ 3TC
DOVUDINE [AZT] – Special Authority see SA2139 on page 1 Cap 100 mg	152.25	100	✓ Retrovir
Oral liq 10 mg per ml	ee SA2139 on pag		
Tab 300 mg with lamivudine 150 mg	92.40	60	<ul><li>✓ Alphapharm</li><li>✓ Lamivudine/</li><li>Zidovudine Viatris</li></ul>
Protease Inhibitors			
TAZANAVIR SULPHATE - Special Authority see SA2139 on	page 113 - Retai	l pharmacy	

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
DARUNAVIR – Special Authority see SA2139 on page 113 – Re Tab 400 mg  Tab 600 mg	132.00 150.00	60 60	✓ [	Darunavir Mylan Darunavir Viatris Darunavir Viatris
(Darunavir Mylan Tab 400 mg to be delisted 1 January 2024)  LOPINAVIR WITH RITONAVIR – Special Authority see SA2139  Tab 100 mg with ritonavir 25 mg		pharma		_opinavir/Ritonavir
Tab 200 mg with ritonavir 50 mg		120	<b>√</b> <u>I</u>	Mylan ∟opinavir/Ritonavir Mylan
RITONAVIR – Special Authority see SA2139 on page 113 – Ret Tab 100 mg	'	30	<b>✓</b> 1	Norvir

# Strand Transfer Inhibitors

DOLUTEGRAVIR - Special Authority see SA2139 on page 11	3 - Retail pharmacy		
Tab 50 mg	1,090.00	30	✓ Tivicay
RALTEGRAVIR POTASSIUM - Special Authority see SA2139	on page 113 – Reta	il pharmacy	
Tab 400 mg	1,090.00	60	✓ Isentress
Tab 600 mg	1,090.00	60	✓ Isentress HD

# **Immune Modulators**

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

#### ⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
  - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
  - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	1	Manufacturer

continued...

following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; or
  - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma\*; or
- 2 All of the following:
  - 2.1 Patient has a myeloproliferative disorder\*; and
  - 2.2 Patient is intolerant of hydroxyurea; and
  - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
  - 3.1 Patient has a myeloproliferative disorder; and
  - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either
  - 3.1 Patient has a cutaneous T cell lymphoma\*: or
  - 3.2 Both:
    - 3.2.1 Patient has a myeloproliferative disorder\*; and
    - 3.2.2 Fither:
      - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
        - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with \* are unapproved indications.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant\* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with \* are unapproved indications.

# **Urinary Tract Infections**

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	19.95	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg - Up to 15 cap available on a			
PSO	81.20	100	✓ Macrobid
Macrobid to be Principal Supply on 1 December 2023			
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	245.00	100	Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		. ,	rand or
	(Manufacturer's Price)			ieneric
	\$	Per	<b>y</b> 10	1anufacturer
Anticholinesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	✓ <u>Max</u>	Health
PYRIDOSTIGMINE BROMIDE			_	
▲ Tab 60 mg	50.28	100	✓ Mes	tinon
Non-Steroidal Anti-Inflammatory Drugs				
Non-Steroidal Anti-Illianimatory Drugs				
DICLOFENAC SODIUM				
* Tab EC 25 mg		50		ofenac Sandoz
* Tab 50 mg dispersible		20	✓ Volt	
* Tab EC 50 mg		50		ofenac Sandoz
* Tab long-acting 75 mg		100		aren SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a		5	✓ Volt	
* Suppos 12.5 mg		10	✓ Volt	
* Suppos 25 mg		10	✓ Volt	
* Suppos 50 mg - Up to 10 supp available on a PSO		10	✓ Volt	
* Suppos 100 mg	7.00	10	✓ Volt	aren
IBUPROFEN				
* Tab 200 mg	21.40	1,000	✓ Relie	eve
* Tab long-acting 800 mg	3.05	30	✓ Bruf	en SR
* Oral liq 20 mg per ml	2.25	200 m		
	11.29		✓ Fen	paed 100 mg per
			5 i	ml .
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	✓ Oruv	vail SR
MEFENAMIC ACID				
* Cap 250 mg	1.25	50		
очр =00 mg	(10.82)		Pons	stan
	0.50	20		
	(7.50)		Pons	stan
NAPROXEN	(1.55)			
* Tab 250 mg	32 69	500	✓ Nofl	am 250
* Tab 500 mg		250	✓ Nofl	
* Tab long-acting 750 mg		28		rosyn SR 750
* Tab long-acting 1 g		28		rosyn SR 1000
		20	· itup	100011 011 1000
TENOXICAM	10.50	100	✓ Tile	-4!1
* Tab 20 mg		100	✓ <u>Tilco</u>	
* Inj 20 mg vial	9.95	1	✓ AFT	
NSAIDs Other				
CELECOXIB				
Cap 100 mg	3 45	60	✓ Cele	hrey
σαρ του mg		00		ecoxib Pfizer
Cap 200 mg	3 20	30	✓ Cele	
		00		coxib Pfizer
			- 5010	

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

# **Topical Products for Joint and Muscular Pain**

### **CAPSAICIN**

# ⇒SA1289 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

# **Antirheumatoid Agents**

#### HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)\*, sarcoidosis (pulmonary and non-pulmonary)\*, and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroquine. Note: Indication marked with a \* is an unapproved indication.

* Tab 200 mg	8.78	100	Plaquenil
LEFLUNOMIDE			
* Tab 10 mg	6.00	30	✓ Arava
Arava to be Principal Supply on 1 December 2023			
* Tab 20 mg	6.00	30	✓ Arava
Arava to be Principal Supply on 1 December 2023			
PENICILLAMINE			
Tab 125 mg	67.23	100	<ul><li>D-Penamine</li></ul>
Tab 250 mg	110.12	100	✓ D-Penamine

# **Drugs Affecting Bone Metabolism**

# Alendronate for Osteoporosis

* Tab 70 mg2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL		
* Tab 70 mg with colecalciferol 5,600 iu1.51	4	✓ Fosamax Plus

# **Other Treatments**

ALENDRONATE SODIUM

DENOSUMAB – Special Authority see SA1777 below – Re	tail pharmacy		
Inj 60 mg prefilled syringe	326.00	1	Prolia

### ⇒SA1777 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

1 The patient has severe, established osteoporosis; and

Subsidy	F	ully	Brand or
(Manufacturer's I	Price) Subsidis	sed	Generic
\$	Per	•	Manufacturer

#### continued...

- 2 Either:
  - 2.1 The patient is female and postmenopausal; or
  - 2.2 The patient is male or non-binary; and
  - 3 Any of the following:
    - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note): or
    - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
    - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
    - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
    - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
    - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
  - 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
  - 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
  - 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

# PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial		1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial		1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA177	9 on the next pa	ge – Retail	pharmacy
* Tab 60 mg	53.76	28	✓ Evista

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

# ⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or egual to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has guantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

#### RISEDRONATE SODIUM

Tab 35 mg ......2.50 Risedronate Sandoz TERIPARATIDE - Special Authority see SA1139 below - Retail pharmacy Forteo

### ⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

#### Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily;

Subsidy (Manufacturer's Price)	S	Fully	Brand or Generic
 \$	Per	1	Manufacturer

#### continued...

zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

#### ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, bag22.5	53 100 ml OP	✓ Zoledronic Acid
		<u>Viatris</u>
		✓ Zoledronic-US S29

(Zoledronic-US S29 Inj 0.05 mg per ml, 100 ml, bag to be delisted 1 January 2024)

# Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg	28.57	500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA19	63 below – Retail pharmacy		
Tab 50 mg	32.00	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	<ul> <li>Benzbromaron AL</li> </ul>
			100 \$29

### ⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

#### COLCHICINE

* Tab 500 mcg6.00	100	✓ Colgout
FEBUXOSTAT – Special Authority see SA2054 below – Retail pharmacy		
Tab 80 mg	28	✓ Febuxostat multichem
Tab 120 mg20.00	28	✓ Febuxostat

### ⇒SA2054 Special Authority for Subsidy

**Initial application — (Gout)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
  - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 <b>`</b> \$	Per	✓	Manufacturer

continued...

- 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
- 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
- 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

**Renewal — (Tumour lysis syndrome)** only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

PR	OE	BEN	JE(	CI	D
PK!	UE	۱ط	۱E(	از	U

# **Muscle Relaxants**

BACLOFEN		
* Tab 10 mg4.20	100	✓ Pacifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement11.55	1	✓ Lioresal Intrathecal
Subsidised only for use in a programmable pump in patients where oral a caused intolerable side effects and the prescription is endorsed according		ents have been ineffective or have
Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement306.82	5	✓ Medsurge
Subsidised only for use in a programmable pump in patients where oral a caused intolerable side effects and the prescription is endorsed according		ents have been ineffective or have
DANTROLENE		
Cap 25 mg112.13	100	✓ Dantrium
, ,		✓ Dantrium S29 S29
Cap 50 mg77.00	100	✓ Dantrium
ORPHENADRINE CITRATE		
Tab 100 mg20.76	100	✓ Norflex

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

# Agents for Parkinsonism and Related Disorders

38.24	60	Symmetrel
63.73	100	Symmetrel
59.50	5	✓ Movapo
121.84	5	✓ Movapo
18.04	100	✓ Comtan
13.25	100	✓ Madopar Rapid
	100	✓ Madopar 62.5
15.80	100	✓ Madopar 125
22.85	100	✓ Madopar HBS
26.25	100	Madopar 250
21.11	100	✓ Sinemet
43.65	100	Sinemet CR
38.39	100	✓ Sinemet
5.51	100	✓ Ramipex
18.66	100	✓ Ramipex
53.50	30	✓ Azilect S29
4.05	84	✓ Ropin
		✓ Ropin
	84	✓ Ropin
	84	✓ Ropin
	-	
152.38	100	✓ Tasmar
		63.73 100

# Anticholinergics

BENZATROPINE MESYLATE			
Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
<ul><li>a) Up to 10 inj available on a PSO</li><li>b) Only on a PSO</li></ul>			
PROCYCLIDINE HYDROCHLORIDE			
Tah 5 mg	7 40	100	✓ Kemadrin

# Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE – Special Authority see SA1403 on the next page – Re	etail pharmacy		
Wastage claimable			
Tab 50 mg	130.00	56	✓ Rilutek



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### ⇒SA1403 Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

# All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
  - 5.1 The patient is ambulatory; or
  - 5.2 The patient is able to use upper limbs; or
  - 5.3 The patient is able to swallow.

**Renewal** from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
  - 3.1 The patient is ambulatory; or
  - 3.2 The patient is able to use upper limbs; or
  - 3.3 The patient is able to swallow.

#### **TETRABENAZINE**

### Anaesthetics

LIDOCAINE [LIGNOCAINE]

#### Local

Gel 2%, tube – Subsidy by endorsement	14.50	30 ml	Xylocaine 2% Jelly
a) Up to 150 ml available on a PSO			
b) Subsidised only if prescribed for urethral or cervical a	administration and	d the prescript	tion is endorsed accordingly.
Gel 2%, 11 ml urethral syringe - Subsidy by endorsement	59.50	10	✓ Instillagel Lido
a) Up to 5 each available on a PSO			
b) Subsidised only if prescribed for urethral, cervical or	rectal administrat	tion and the pi	rescription is endorsed
accordingly.			·
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Oral (gel) soln 2%	44.00	200 ml	✓ Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	9.50	25	✓ Lidocaine-Baxter
	17.50	50	
	(35.00)		Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	9.00	25	✓ Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5	
	(20.00)		Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO	6.85	5	<ul> <li>Lidocaine-Baxter</li> </ul>
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO	7.15	5	<ul> <li>Lidocaine-Baxter</li> </ul>
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE			
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –			
Subsidy by endorsement	103.32	10	✓ Pfizer
a) Up to 5 each available on a PSO			

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

(Pfizer Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes to be delisted 1 November 2023)

# **NERVOUS SYSTEM**

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

# **Topical Local Anaesthetics**

# ⇒SA0906 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority see SA0906 above -	- Retail pharr	nacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authority	see SA0906	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓ EMLA

# **Analgesics**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 119

TO Anti-initialinitatory NoAIDO Telef to MOSCOLOSKELETAL, page 1	19		
Non-opioid Analgesics			
ASPIRIN			
* Tab dispersible 300 mg - Up to 30 tab available on a PSO	4.50	100	✓ Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or diabetic accordingly.	periphera	al neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.14	57 g OP	✓ Rugby Capsaicin Topical Cream S29
NEFOPAM HYDROCHLORIDE			
Tab 30 mg	23.40	90	✓ Acupan

	Subsidy (Manufacturer's Price	a) Subs	Fully Brand or sidised Generic	
	\$	Per	✓ Manufacturer	
PARACETAMOL				
Tab 500 mg - blister pack	19.75	1,000	✓ Pacimol	
a) Maximum of 300 tab per prescription; can be waived     b) Up to 30 tab available on a PSO	d by endorsement			
c) 1) Subsidy by endorsement for higher quantities regular daily dosing for one month or greater, annotate the prescription as endorsed where c 2) Maximum of 100 tab per dispensing for non-en (for non-endorsed patients), then dispense in to 500 mg - bottle pack — Maximum of 300 tab per prescription; can be waived by endorsement	and the prescription dispensing history sindorsed patients. If repeat dispensings i 17.92 vailable for patients escription is annotate	is annotated upports a lor quantities protected in 1,000 with long tered according	d accordingly. Pharmacing-term condition. rescribed for more than 1 ig 100 tab per dispensing  Noumed Paracetamol rm conditions who requirely. Pharmacists may ani	sts may 100 tabs g. e regular
<ol> <li>Maximum of 100 tab per dispensing for non-endo non-endorsed patients), then dispense in repeat of</li> </ol>	rsed patients. If qua	antities preso	cribed for more than 100	tabs (for
Oral liq 120 mg per 5 ml	3.98	200 ml	✓ Paracetamol (Ethics)	
	10.50	200 ml OP	✓ Avallon	
a) Maximum of 600 ml per prescription; can be waived b) Up to 200 ml available on a PSO c) Not in combination d)  1) Maximum of 200 ml per dispensing for non-en non-endorsed patients), then dispense in repe 2) Subsidy by endorsement for higher quantities regular daily dosing for one month or greater a Pharmacists may annotate the prescription as condition. 3) Note: 200 ml presentations of paracetamol or provisions in Part I of Section A.	dorsed patients. If of at dispensing not exist available for paties and the prescription endorsed where dis	cceeding 200 ents with long is endorsed spensing his	Oml per dispensing. g term conditions who red or annotated accordingly tory supports a long-term	quire /. 1
Oral liq 250 mg per 5 ml		200 ml	✓ <u>Pamol</u>	
c) Not in combination d)  1) Maximum of 200 ml per dispensing for non-en non-endorsed patients), then dispense in repe 2) Subsidy by endorsement for higher quantities regular daily dosing for one month or greater a Pharmacists may annotate the prescription as condition. 3) Note: 200 ml presentations of paracetamol or	at dispensing not exist available for paties and the prescription endorsed where dis	cceeding 200 ents with long is endorsed spensing his	O ml per dispensing. g term conditions who red or annotated accordingly tory supports a long-term	quire /. 1
provisions in Part I of Section A.	. , ,	piled off Do		
* Suppos 125 mg  * Suppos 250 mg  * Suppos 500 mg	5.39	10 10 50	<ul><li>✓ Gacet</li><li>✓ Gacet</li><li>✓ Gacet</li></ul>	

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Sub: Per	sidised •	Generic Manufacturer
	*			
Opioid Analgesics				
CODEINE PHOSPHATE - Safety medicine; prescriber may det	ermine dispensing fre	quency		
Tab 15 mg	5.92	100	_	loumed
Tab 30 mg	6.98	100		Aspen
Tab CO ma	10.00	100	_	loumed
Tab 60 mg	13.89	100	<u> </u>	loumed
DIHYDROCODEINE TARTRATE	0.00	00	./ -	NIO Camtimus
Tab long-acting 60 mg	8.60	60	• 0	OHC Continus
FENTANYL				
a) Only on a controlled drug form				
<ul><li>b) No patient co-payment payable</li><li>c) Safety medicine; prescriber may determine dispensing fr</li></ul>	oguonov			
Inj 50 mcg per ml, 2 ml ampoule		10	<b>√</b> F	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10	_	Boucher and Muir
Patch 12.5 mcg per hour		5		entanyl Sandoz
Patch 25 mcg per hour		5	_	entanyl Sandoz
Patch 50 mcg per hour	9.49	5	✓ F	entanyl Sandoz
Patch 75 mcg per hour		5		entanyl Sandoz
Patch 100 mcg per hour	18.59	5	✓ <u>F</u>	entanyl Sandoz
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr				
d) Extemporaneously compounded methadone will only be	reimbursed at the rate	e of the ch	neapest	form available
(methadone powder, not methadone tablets).	armulaa naga 066			
For methadone hydrochloride oral liquid refer Standard F Tab 5 mg		10	✓ M	lethadone BNM
Oral lig 2 mg per ml		200 ml	_	Biodone
Oral liq 5 mg per ml		200 ml	_	Biodone Forte
Oral liq 10 mg per ml		200 ml	_	Biodone Extra Forte
Inj 10 mg per ml, 1 ml	68.90	10	✓ A	FT
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
<ul> <li>c) Safety medicine; prescriber may determine dispensing fr</li> </ul>				
Oral liq 1 mg per ml		200 ml		RA-Morph
Oral liq 2 mg per ml		200 ml		RA-Morph
Oral liq 5 mg per ml	19.44 2	200 ml		Ordine \$29
				RA-Morph
Oral liq 10 mg per ml	27.74	200 ml	<b>✓</b> 0	Ordine \$29

✓ RA-Morph

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	_	Subsidised	
	\$	Per		Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr	eauencv			
Tab immediate-release 10 mg		10	1	Sevredol
Tab immediate-release 20 mg		10	1	Sevredol
Cap long-acting 10 mg		10	1	m-Eslon
Cap long-acting 30 mg		10	✓	m-Eslon
Cap long-acting 60 mg	9.00	10	✓	m-Eslon
Cap long-acting 100 mg	10.50	10	✓	m-Eslon
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a P	SO5.38	5	✓	Medsurge
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a		5	✓	Medsurge
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO5.53	5	✓	Medsurge
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a		5	✓	Medsurge
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
, ,				
b) No patient co-payment payable	roguenov.			
Safety medicine; prescriber may determine dispensing from Tab controlled-release 5 mg		20		Oxycodone Sandoz
rab controlled-release 5 mg				
Table and the Hard and a second One of	4.04	30		OxyContin S29
Tab controlled-release 10 mg		20		Oxycodone Sandoz
Tab controlled-release 20 mg		20		Oxycodone Sandoz
Tab controlled-release 40 mg		20		Oxycodone Sandoz
Tab controlled-release 80 mg		20		Oxycodone Sandoz
Cap immediate-release 5 mg		20		OxyNorm OxyNorm
Cap immediate-release 10 mg		20		OxyNorm OxyNorm
Cap immediate-release 20 mg		20		OxyNorm OxyNorm
Oral liq 5 mg per 5 ml		250 m		OxyNorm Hamala
Inj 10 mg per ml, 1 ml ampoule		5		Hameln Hameln
Inj 10 mg per ml, 2 ml ampoule		5		<u>Hameln</u>
Inj 50 mg per ml, 1 ml ampoule		5		<u>Hameln</u>
PARACETAMOL WITH CODEINE - Safety medicine; prescribe	,	•		•
* Tab paracetamol 500 mg with codeine phosphate 8 mg	27.50	1,000	/	Paracetamol +
				Codeine (Relieve)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr	requency			
Tab 50 mg		10	✓	Noumed Pethidine
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a		5		DBL Pethidine
ing oo ing por ini, i iii ampoalo op to o ing ataliasio on a		·		Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a	PSO 30.72	5	/	DBL Pethidine
ing so mg por mi, z mi ampoule — op to o ing available on a	1 0000.72	Ü	•	Hydrochloride
TRAMAROL LIVEROCUI ORIES				yaroomonae
TRAMADOL HYDROCHLORIDE	4.05	00		Tramel CD 400
Tab sustained-release 100 mg		20		Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg		100	•	Arrow-Tramadol
Arrow-Tramadol to be Principal Supply on 1 January 20	24			

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

# **Antidepressants**

# **Cyclic and Related Agents**

•			
AMITRIPTYLINE - Safety medicine; prescriber may determ	nine dispensing frequen	cy	
Tab 10 mg	2.99	100	Arrow-Amitriptyline
Tab 25 mg	1.99	100	Arrow-Amitriptyline
Tab 50 mg	3.14	100	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; p	rescriber may determine	e dispensin	g frequency
Tab 10 mg		30	Clomipramine Teva
Tab 25 mg	11.99	30	Clomipramine Teva
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy	by endorsement		
a) Safety medicine; prescriber may determine dispensi	ing frequency		
b) Subsidy by endorsement – Subsidised for patients v	vho were taking dosulep	in [dothiepi	n] hydrochloride prior to 1 June
2019 and the prescription is endorsed accordingly.		ate the pres	scription as endorsed where there
exists a record of prior dispensing of dosulepin [doth			45
Tab 75 mg		30	✓ Dosulepin Viatris
Cap 25 mg	7.83	50	✓ Dosulepin
			Mylan S29
			✓ Dosulepin
			Viatris \$29
IMIPRAMINE HYDROCHLORIDE – Safety medicine; preso			
Tab 10 mg		50	✓ Tofranil
T-1- 05	10.96	100	✓ Tofranil
Tab 25 mg		50	✓ Tofranil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine;	,		0 1 ,
Tab 10 mg		100	✓ <u>Norpress</u>
Tab 25 mg	6.29	180	✓ <u>Norpress</u>
Monoamine-Oxidase Inhibitors (MAOIs) - No	on Selective		
` '	on ociconive		
TRANYLCYPROMINE SULPHATE			
* Tab 10 mg	22.94	50	✓ Parnate
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE			
* Tab 150 mg	11.80	60	✓ Aurorix
* Tab 300 mg		60	✓ Aurorix ✓ Aurorix
			- (MI ALIX
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE			
* Tab 20 mg	2.86	84	✓ Celapram

✓ Escitalopram

(Ethics)

✓ Escitalopram (Ethics)

28

28

**ESCITALOPRAM** 

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
LUOXETINE HYDROCHLORIDE				
<ul> <li>Tab dispersible 20 mg, scored – Subsidy by endorsemer Subsidised by endorsement</li> </ul>	nt2.50	28	•	Fluox
<ol> <li>When prescribed for a patient who cannot swal accordingly; or</li> </ol>	low whole tablets or cap	sules a	and the p	rescription is endorsed
<ol><li>When prescribed in a daily dose that is not a m endorsed. Note: Tablets should be combined</li></ol>				
Cap 20 mg	2.22	30	•	Brown & Burk S29
	3.13	90	•	Arrow-Fluoxetine
AROXETINE				
★ Tab 20 mg	4.11	90	•	<u>Loxamine</u>
ERTRALINE		_	_	
★ Tab 50 mg ★ Tab 100 mg		30		Setrona Setrona
≮ Tab 100 mg	1./4	30	•	Setrona
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg		28		Noumed
Tab 45 mg	3.45	28	•	Noumed
ENLAFAXINE	0.00	0.4		Enlefey VD
€ Cap 37.5 mg € Cap 75 mg		84 84		Enlafax XR Enlafax XR
€ Cap 150 mg		84	_	Enlafax XR
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
NAZEPAM – Safety medicine; prescriber may determine dis Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsemen		5	_	Hospira
a) Up to 5 inj available on a PSO	L	J	•	Поэрна
b) Only on a PSO				
c) PSO must be endorsed "not for anaesthetic proce	edures".			
Rectal tubes 5 mg - Up to 5 tube available on a PSO	54.58	5	✓	Stesolid
HENYTOIN SODIUM				
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on		_		
PSO		5	•	Hospira
Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on		5	/	Hospira
		J	,	
Control of Epilepsy				
ARBAMAZEPINE	44.50	400		T
₹ Tab long acting 200 mg		100		Tegretol Tegretol CR
Fab long-acting 200 mg	33.96	100 200		Tegretol CR
₭ Tab 400 mg		100		Tegretol
★ Tab long-acting 400 mg	39.17	100	•	Tegretol CR

	Subsidy		Fully Brand or	
	(Manufacturer's Pri		sidised Generic	
	\$	Per	✓ Manufacturer	
CLOBAZAM - Safety medicine; prescriber may determine dispe	ensing frequency			
Tab 10 mg		50	✓ Frisium	
CLONAZEPAM - Safety medicine; prescriber may determine di	spensing frequenc	у		
Oral drops 2.5 mg per ml	7.38	10 ml OP	✓ Rivotril	
ETHOSUXIMIDE				
Cap 250 mg	78.89	56	<ul><li>Essential</li></ul>	
			Ethosuximide	S29
	140.88	100	✓ Zarontin	
Oral liq 250 mg per 5 ml	56.35	200 ml	✓ Zarontin	
GABAPENTIN				
Note: Not subsidised in combination with subsidised pregat	palin			
* Cap 100 mg	6.45	100	✓ Nupentin	
* Cap 300 mg		100	✓ Nupentin	
* Cap 400 mg		100	✓ Nupentin	
LACOSAMIDE - Special Authority see SA2267 below - Retail p				
▲ Tab 50 mg	25.04	14	✓ Vimpat	
▲ Tab 100 mg		14	✓ Vimpat	
·	200.24	56	✓ Vimpat	
▲ Tab 150 mg	75.10	14	✓ Vimpat	
· ·	300.40	56	✓ Vimpat	
▲ Tab 200 mg	400.55	56	✓ Vimpat	

# ⇒SA2267 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

### LAMOTRIGINE

•	l ab dispersible 2 mg	55.00	30	<ul> <li>Lamictal</li> </ul>
$\blacktriangle$	Tab dispersible 5 mg	50.00	30	✓ Lamictal
	Tab dispersible 25 mg		56	✓ Logem
	Tab dispersible 50 mg		56	✓ Logem
	Tab dispersible 100 mg		56	✓ Logem
LE	VETIRACETAM			
	Tab 250 mg	5.84	60	✓ Everet
	Tab 500 mg	10.51	60	✓ Everet
	Tab 750 mg	16.71	60	✓ Everet
	Tab 1,000 mg	21.82	60	✓ Everet
	Oral lig 100 mg per ml	44.78	300 ml OP	✓ Levetiracetam-AFT



		Subsidy (Manufacturer's Price	١	Fully Subsidised	
		(Wallulacturer's Frice	Per	Jubsiuiseu	Manufacturer
HI'	ENOBARBITONE				
	For phenobarbitone oral liquid refer Standard Formulae, pag	e 266			
*	Tab 15 mg		500	1	PSM
	Tab 30 mg		500		PSM
•	7 ab 55 mg	398.50	000		Noumed
		000.00			Phenobarbitone
	Noumed Phenobarbitone to be Principal Supply on 1 De	ecember 2023			
PS	M Tab 30 mg to be delisted 1 December 2023)				
PHE	ENYTOIN SODIUM				
*	Tab 50 mg	75.00	200	1	Dilantin Infatab
	Cap 30 mg	74.00	200	/	Dilantin
	Cap 100 mg	37.00	200	1	Dilantin
*	Oral lig 30 mg per 5 ml	22.03	500 m	· •	Dilantin
	1 31			1	<b>Dilantin Paediatric</b>
Dil	antin Oral liq 30 mg per 5 ml to be delisted 1 March 2024)				
	EGABALIN				
111	Note: Not subsidised in combination with subsidised gabape	antin			
	Cap 25 mg		56	1	Pregabalin Pfizer
	Oap 23 mg		50		•
	0 75	7.80			Milpharm \$29
*	Cap 75 mg		56		Pregabalin Pfizer
		8.10			Milpharm \$29
	Cap 150 mg	4.01	56		Lyrica
				/	Pregabalin Pfizer
		12.44		✓	Milpharm S29
	Cap 300 mg	7.38	56	✓	Pregabalin Pfizer
PRI	MIDONE				
	Tab 250 mg	37.35	100	1	Primidone Clinect
	5		100	-	
οUI	DIUM VALPROATE	10.05	100	,	Fulling Once he lete
	Tab 100 mg		100		Epilim Crushable
	Tab 200 mg EC		100		Epilim
	Tab 500 mg EC		100		Epilim
*	Oral liq 200 mg per 5 ml	20.48	300 m		Epilim S/F Liquid
					Epilim Syrup
*	Inj 100 mg per ml, 4 ml	41.50	1	/	Epilim IV
3TI	RIPENTOL - Special Authority see SA2268 below - Retail p	harmacy			
	Cap 250 mg		60	1	Diacomit
	Powder for oral lig 250 mg sachet		60	_	Diacomit

### ⇒SA2268 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		ubsidised	Generic
	\$	Per	•	Manufacturer
TOPIRAMATE				
▲ Tab 25 mg	11.07	60	1	Arrow-Topiramate
				Topiramate Actavis
	26.04			Topamax
▲ Tab 50 mg		60		Arrow-Topiramate
		00		Topiramate Actavis
	44.26		_	Topamax
▲ Tab 100 mg	*	60		Arrow-Topiramate
_ Tab 100 mg		00		Topiramate Actavis
	75.05			
A Tab 000	75.25	00		Topamax
▲ Tab 200 mg	55.19	60		Arrow-Topiramate
			_	Topiramate Actavis
	129.85		✓.	Topamax
▲ Sprinkle cap 15 mg	20.84	60	✓.	Topamax
▲ Sprinkle cap 25 mg	26.04	60	✓.	Topamax
VIGABATRIN - Special Authority see SA2088 below - Retail				-
▲ Tab 500 mg		100	1	Sabril
Powder for oral soln 500 mg per sachet		60		Sabril
Towaci for oral soin soo my per sacriet	7 1.50	00	• ,	Jabin

### **⇒SA2088** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has infantile spasms; or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy: and
    - 1.2.2 Either:
      - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
      - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
  - 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Fither
  - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

# **Antimigraine Preparations**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 119

# **Acute Migraine Treatment**

**RIZATRIPTAN** 

Tab orodispersible 10 mg.......4.84 30 ✓ Rizamelt

# **NERVOUS SYSTEM**

	Subsidy (Manufacturer's Price) \$	S Per	Fully Brand or Gubsidised Generic Manufacturer	
SUMATRIPTAN				
Tab 50 mg Tab 100 mg Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj pe	22.68	90 90	<ul><li>✓ <u>Sumagran</u></li><li>✓ <u>Sumagran</u></li></ul>	
prescription		2 OP	✓ Imigran	
Prophylaxis of Migraine				
For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SY	STEM, page 51			
PIZOTIFEN  * Tab 500 mcg	23.21	100	✓ Sandomigran	
Antinausea and Vertigo Agents				
For Antispasmodics refer to ALIMENTARY TRACT, page 8				
APREPITANT – Special Authority see SA0987 below – Retail ph Cap 2 × 80 mg and 1 × 125 mg		3 OP	✓ Emend Tri-Pack	<u>(</u>
■ SA0987   Special Authority for Subsidy   Initial application from any relevant practitioner. Approvals valid emetogenic chemotherapy and/or anthracycline-based chemother Renewal from any relevant practitioner. Approvals valid for 12 m chemotherapy and/or anthracycline-based chemotherapy for the	rapy for the treatment conths where the pat	nt of ma ient is u	alignancy.	
BETAHISTINE DIHYDROCHLORIDE  * Tab 16 mg  Serc to be Principal Supply on 1 December 2023	3.70	100	✓ Serc	
CYCLIZINE HYDROCHLORIDE				
Tab 50 mg	0.49	10	✓ <u>Nausicalm</u>	
CYCLIZINE LACTATE				
Inj 50 mg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	16.36	10	✓ Hameln	
DOMPERIDONE				
* Tab 10 mg	4.00	100	✓ <u>Domperidone</u> <u>Viatris</u>	
HYOSCINE HYDROBROMIDE				
Inj 400 mcg per ml, 1 ml ampoule		10	✓ Martindale S29	
pharmacy		2	✓ Scopoderm TTS	3

# ⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

**Renewal** from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price)		Fully Subsidised	I Generic
	\$	Per		Manufacturer
METOCLOPRAMIDE HYDROCHLORIDE			_	
* Tab 10 mg - Up to 30 tab available on a PSO	1.57	100	•	Metoclopramide Actavis 10
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PS ONDANSETRON	SO7.00	10	•	Baxter
	2.27	50	_	Periset
* Tab 4 mg  Tab disp 4 mg – Up to 10 tab available on a PSO		10		Periset ODT
Tab disp 4 mg - op to To tab available on a 1 30	0.76	10		Ondansetron
	0.70		•	ODT-DRLA
* Tab 8 mg	4.10	50	1	Periset
Tab disp 8 mg – Up to 10 tab available on a PSO		10		Periset ODT
Tab disp o mg - Op to To tab available on a T Go	1.13	10		Ondansetron
	1.10			ODT-DRLA
(Ondansetron ODT-DRLA Tab disp 4 mg to be delisted 1 March a (Ondansetron ODT-DRLA Tab disp 8 mg to be delisted 1 March a				
PROCHLORPERAZINE	,			
* Tab 3 mg buccal	5 97	50		
7 Tab o mg baccar	(30.00)	50		Buccastem
	(30.00)			Max Health \$29
* Tab 5 mg - Up to 30 tab available on a PSO	` '	250	1	Nausafix
* Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO		10		Stemetil
A 12 1 12				
Antipsychotics				
General				
AMISULPRIDE - Safety medicine; prescriber may determine dis	pensing frequency			
Tab 100 mg		30	/	Sulprix
Tab 200 mg		60		Sulprix
Tab 400 mg		60		Sulprix
ARIPIPRAZOLE – Safety medicine; prescriber may determine di				
Tab 5 mg		30	1	Aripiprazole Sandoz
Tab 5 mg	10.00	00		Ascend
			-	Aripiprazole S29
Tab 10 mg	10.50	30	./	Aripiprazole Sandoz
Tab 15 mg		30		Aripiprazole Sandoz
Tab 20 mg		30		Aripiprazole Sandoz
Tab 30 mg		30		Aripiprazole Sandoz
<u> </u>				• •
CHLORPROMAZINE HYDROCHLORIDE – Safety medicine; pre				
Tab 10 mg — Subsidy by endorsement		100		Largactil
Subsidised for patients who were taking chlorpromazine prescription is endorsed accordingly. Pharmacists may	,			•
record of prior dispensing of chlorpromazine 10 mg table				u where there exists a
Tab 25 mg - Up to 30 tab available on a PSO	, ,	2 11101 100		Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100		Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO		100		Largactil
(Largactil Tab 10 mg to be delisted 1 April 2024)		10	•	Laryaviii
Largadin Tub To my to be denoted 1 April 2027)				

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	
	(Manufacturer's Pric	e) S Per	Subsidised •	Generic Manufacturer
	Ψ	rei		Ivianulaciurei
CLOZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing	frequency			
Tab 25 mg	6.69	50		Clopine
				Clozaril
	13.37	100		Clopine
				Clozaril
Tab 50 mg		50		Clopine
	17.33	100		Clopine
Tab 100 mg	17.33	50		Clopine
				Clozaril
	34.65	100	•	Clopine
				Clozaril
Tab 200 mg	34.65	50	/	Clopine
	69.30	100	/	Clopine
Suspension 50 mg per ml	67.62	100 ml	✓	Versacloz
ALOPERIDOL - Safety medicine; prescriber may determi	ine dispensina frequency	,		
Tab 500 mcg - Up to 30 tab available on a PSO		100	/	Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50		Serenace
rab 5 mg Op to 50 tab available on a 1 00	29.72	100		Serenace
Oral liq 2 mg per ml - Up to 200 ml available on a PSC		100 ml		Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available or		100 1111		Serenace
			•	Selellace
EVOMEPROMAZINE – Safety medicine; prescriber may of			_	
Tab 25 mg (33.8 mg as a maleate)		100		Nozinan (Swiss)
Tab 25 mg as a maleate	16.10	100		Nozinan
Tab 100 mg (135 mg as a maleate)	41.75	100		Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	/	Nozinan
3				
_		rmine di		frequency
EVOMEPROMAZINE HYDROCHLORIDE - Safety medic	ine; prescriber may dete		spensing	
_	ine; prescriber may dete	rmine di	spensing	Neuraxpharm \$29
EVOMEPROMAZINE HYDROCHLORIDE - Safety medic	ine; prescriber may dete	5	spensing	Neuraxpharm §29 Nozinan S29 §29
EVOMEPROMAZINE HYDROCHLORIDE - Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete 16.75 24.48	5 10	spensing	Neuraxpharm \$29
EVOMEPROMAZINE HYDROCHLORIDE - Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete 16.75 24.48	5 10	spensing  ✓	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt
EVOMEPROMAZINE HYDROCHLORIDE – Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete16.75  24.48 determine dispensing fre72.00	5 10	spensing  ✓	Neuraxpharm §29 Nozinan S29 §29
EVOMEPROMAZINE HYDROCHLORIDE – Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete16.75  24.48 determine dispensing fre72.00	5 10 equency	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete16.75  24.48  determine dispensing from72.0022.36	5 10 equency 100	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt
EVOMEPROMAZINE HYDROCHLORIDE – Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete16.75  24.48 determine dispensing fra72.0022.36 e dispensing frequency	5 10 equency 100 100	spensing	Neuraxpharm 529 Nozinan S29 529 Wockhardt Priadel Douglas
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100	spensing	Neuraxpharm 529 Nozinan S29 529 Wockhardt  Priadel Douglas  Zypine
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT
EVOMEPROMAZINE HYDROCHLORIDE – Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT Zypine
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	24.48  determine dispensing frequency	5 10 equency 100 100 28 28 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT Zypine ODT
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT Zypine Zypine ODT Neulactil
EVOMEPROMAZINE HYDROCHLORIDE – Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT Zypine Zypine ODT Neulactil Neulactil
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28 28 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil
EVOMEPROMAZINE HYDROCHLORIDE – Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT Zypine Zypine ODT Neulactil Neulactil
EVOMEPROMAZINE HYDROCHLORIDE – Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28 28 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28 28 28 28 28	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil
EVOMEPROMAZINE HYDROCHLORIDE — Safety medic Inj 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28 28 28 28 28 4 100 84 100	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil Neulactil Neulactil Quetapel
EVOMEPROMAZINE HYDROCHLORIDE — Safety medicing 25 mg per ml, 1 ml ampoule	ine; prescriber may dete	5 10 equency 100 100 28 28 28 28 28 28 28 38 4 100 84 100	spensing	Neuraxpharm \$29 Nozinan \$29 \$29 Wockhardt  Priadel Douglas  Zypine Zypine Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil Neulactil

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	)	Subsidised	Generic
	\$	Per	✓	Manufacturer
RISPERIDONE – Safety medicine; prescriber may determine di	spensing frequency			
Tab 0.5 mg	2.17	60	✓	Risperidone (Teva)
Tab 1 mg		60	✓	Risperidone (Teva)
Tab 2 mg	2.72	60	✓	Risperidone (Teva)
Tab 3 mg	4.50	60	✓	Risperidone (Teva)
Tab 4 mg	6.25	60	✓	Risperidone (Teva)
Oral liq 1 mg per ml	10.29	30 ml	✓	Risperon
	17.80	100 m	ıl 🗸	Risperon
ZIPRASIDONE - Safety medicine; prescriber may determine di	spensing frequency			
Cap 20 mg	17.90	60	✓	Zusdone
Cap 40 mg	27.41	60	✓	Zusdone
Cap 60 mg	38.39	60	✓	Zusdone
Cap 80 mg	46.55	60	✓	Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pro	escriber may determin	ne disp	ensing fre	equency
Tab 10 mg	•	100	•	Clopixol

# **Depot Injections**

Depot injections		
FLUPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispe	ensing freq	uency
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO	5	✓ Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO20.90	5	✓ Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO40.87	5	✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber may determine disper	nsing frequ	iency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO55.90	5	✓ Haldol Concentrate
		✓ Haldol
		Decanoas \$29
OLANZAPINE - Special Authority see SA1428 below - Retail pharmacy		
Safety medicine; prescriber may determine dispensing frequency		
Inj 210 mg vial252.00	1	✓ Zyprexa Relprevv
Inj 300 mg vial414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial504.00	1	Zyprexa Relprevv

# ⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia; and
  - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE – Special Authority see SA1429 on the next page – Retail pharmacy Safaty medicine: prescriber may determine dispensing frequency.

Safety medicine, prescriber may determine dispensing	rrequericy		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe		1	✓ Invega Sustenna

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

### ⇒SA1429 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

### PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

Inj 175 mg syringe	815.85	1	✓ Invega Trinza
Inj 263 mg syringe		1	✓ Invega Trinza
Inj 350 mg syringe		1	✓ Invega Trinza
Inj 525 mg syringe		1	✓ Invega Trinza

# ⇒SA2167 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

### RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

✓ Risperdal Consta

✓ Risperdal Consta✓ Risperdal Consta

# ⇒SA1427 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

# **Anxiolytics**

# BUSPIRONE HYDROCHLORIDE

טטע	SI ITONE ITI DITOCITEDITIDE				
*	Tab 5 mg	18.50	100	1	Buspirone Viatris
*	Tab 10 mg	12.50	100	✓	<b>Buspirone Viatris</b>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
CLONAZEPAM - Safety medicine; prescriber may determine dis	pensing frequency			
Tab 500 mcg	5.64	100	✓	Paxam
Tab 2 mg	10.78	100	✓	Paxam
DIAZEPAM - Safety medicine; prescriber may determine dispension	sing frequency			
Tab 2 mg	95.00	500	✓	Arrow-Diazepam
Tab 5 mg	115.00	500	✓	Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine dispe	ensing frequency			
Tab 1 mg	9.72	250	✓	Ativan
Tab 2.5 mg	12.50	100	✓	Ativan

# **Multiple Sclerosis Treatments**

### **⇒SA2274** Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

### 1 All of the following:

- 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 1.2 Patients has an EDSS score between 0 6.0; and
- 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 1.4 All of the following:
  - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
  - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
  - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
  - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
  - 1.4.5 Either:
    - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
    - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 1.6 Any of the following:
  - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
  - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
  - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
  - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
  - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer continued... 2 Patient has an active approval for ocrelizumab and does not have primary progressive MS. Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. Renewal — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months). Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. DIMETHYL FUMARATE - Special Authority see SA2274 on the previous page - Retail pharmacy a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Tecfidera ✓ Tecfidera Cap 240 mg.......2,000.00 FINGOLIMOD - Special Authority see SA2274 on the previous page - Retail pharmacy a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Gilenva GLATIRAMER ACETATE - Special Authority see SA2274 on the previous page - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Copaxone INTERFERON BETA-1-ALPHA - Special Authority see SA2274 on the previous page - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Avonex ✓ Avonex Pen INTERFERON BETA-1-BETA - Special Authority see SA2274 on the previous page - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. 15 ✓ Betaferon NATALIZUMAB - Special Authority see SA2274 on the previous page - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Tysabri TERIFLUNOMIDE - Special Authority see SA2274 on the previous page - Retail pharmacy a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Aubagio Multiple Sclerosis Treatments - Other OCRELIZUMAB - Special Authority see SA2273 below - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Ocrevus ⇒SA2273 Special Authority for Subsidy Initial application — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

1 All of the following:

1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed

continued...

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

by a neurologist; and

- 1.2 Patients has an EDSS score between 0 6.0; and
- 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 1.4 All of the following:
  - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
  - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
  - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
  - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
  - 1.4.5 Either:
    - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
    - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 1.6 Any of the following:
  - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
  - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
  - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
  - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
  - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

**Renewal — (Multiple Sclerosis - ocrelizumab)** from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and
- 3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 2.0 to 6.5 (inclusive) at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

# Sedatives and Hypnotics

MELATONIN − Special Authority see SA1666 on the next page − Retail pharmacy
Tab modified-release 2 mg − No more than 5 tab per day......11.50 30

✓ Vigisom
Restricted to patients aged 18 years or under.



Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

### ⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)\*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under\*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Patient is aged 18 years or under\*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

MIDAZOLAM - Safety medicine: prescriber may determine dispensing frequency

Note: Indications marked with \* are unapproved indications.

= - =			
Inj 1 mg per ml, 5 ml ampoule	6.10	10	✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available			
on a PSO	17.28	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be el	ndorsed for stat	tus epilepticu	us use only.
Inj 5 mg per ml, 3 ml ampoule	5.00	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available of	n		
a PSO	13.09	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be el	ndorsed for stat	tus epilepticu	us use only.
PHENOBARBITONE SODIUM – Special Authority see SA1386 be	low – Retail pha	armacy	
Inj 200 mg per ml, 1 ml ampoule	113.37	10	✓ Max Health S29
⇒SA1386 Special Authority for Subsidy			

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

Р

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM - Safety medicine; prescriber may determine dispensin	g frequency		
Tab 10 mg		25	✓ Normison

### TRIAZOLAM - Subsidy by endorsement

- a) Safety medicine: prescriber may determine dispensing frequency
- b) Subsidised for patients who were taking triazolam prior to 1 June 2023 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of triazolam in the preceding 12 months.

Tab 125 mcg	5.10	100	
<b>v</b>	(9.85)		Hypam
Tab 250 mcg	4.10	100	
-	(11.20)		Hypam

(Hypam Tab 125 mcg to be delisted 1 February 2024) (Hypam Tab 250 mcg to be delisted 1 February 2024)

#### **NERVOUS SYSTEM**

✓ Zopiclone Actavis

500

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
ZOPICLONE – Safety medicine; prescriber may determine dispensing frequency					

# **Spinal Muscular Atrophy**

NUSINERSEN - PCT only - Special Authority see SA2174 below

Inj 12 mg per 5 ml vial .......120,000.00 1 ✓ Spinraza

## ⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
  - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
  - 3.2 Both:
    - 3.2.1 Patient is pre-symptomatic; and
    - 3.2.2 Patient has three or less copies of SMN2.

**Renewal — (spinal muscular atrophy (SMA))** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

## All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

#### RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml. 60 mg per bottle......14,100.00 80 ml OP ✓ Evrysdi

#### ⇒SA2203 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

## All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
  - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
  - 3.2 Both:
    - 3.2.1 Patient is pre-symptomatic; and
    - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.



	(Manufacturer's Price)	0	Subsidised	Generic	
	(Manufacturer's Frice)	Per	oubsidised •	Manufacturer	
	· · · · · · · · · · · · · · · · · · ·				
Stimulants/ADHD Treatments					
ATOMOXETINE					
Cap 10 mg	18.41	28	✓	APO-Atomoxetine	
			1	APO-Atomoxetine	
				<b>S29</b> S29	
			✓	Generic Partners	
	107.03		1	Strattera	
Cap 18 mg	27.06	28		APO-Atomoxetine	
				Generic Partners	
	107.03			Strattera	
Cap 25 mg	29.22	28		APO-Atomoxetine	
2 42	22.22		_	Generic Partners	
Cap 40 mg	29.22	28		APO-Atomoxetine	
	107.00			Generic Partners	
Can 60 mg	107.03	28		Strattera APO-Atomoxetine	
Cap 60 mg	40.31	20		APO-Atomoxetine	
			•	S29 S29	
				Generic Partners	
Can 90 mg	EG AE	28		APO-Atomoxetine	
Cap 80 mg		20		APO-Atomoxetine	
			•	S29 S29	
				Generic Partners	
Cap 100 mg	E0 40	28		APO-Atomoxetine	
Cap 100 mg		20		APO-Atomoxetine	
			•	S29 S29	
				Generic Partners	
(Ctrattara Can 10 mg to be delicted 1 Nevember 2002)			•	Generic Partners	
(Strattera Cap 10 mg to be delisted 1 November 2023) (Strattera Cap 18 mg to be delisted 1 November 2023)					
(Strattera Cap 40 mg to be delisted 1 November 2023)					
,	halam Data'll abanna				
DEXAMFETAMINE SULFATE – Special Authority see SA1149	below – Retail pharma	су			
a) Only on a controlled drug form     Cofety and disconnection of the controlled drug form					
b) Safety medicine; prescriber may determine dispensing to	, ,	100		PSM	
Tab 5 mg	28.50	100		<u>PSW</u> Aspen	
	20.00		•	Asheii	

Subsidy

Fully

Brand or

## ⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application — (Narcolepsy)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Renewal** — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensing frequency		
Tab immediate-release 5 mg	30	✓ Rubifen
Tab immediate-release 10 mg3.00	30	<ul><li>✓ Ritalin</li><li>✓ Rubifen</li></ul>
Tab extended-release 18 mg7.75	30	✓ Methylphenidate ER  - Teva
Tab immediate-release 20 mg7.85	30	✓ Rubifen
Tab sustained-release 20 mg - Brand switch fee payable		
(Pharmacode 2665956) - see page 264 for details	30	✓ Rubifen SR
Note: Brand Switch Fee applies only to patients who have transferred fror out of stock.	n Methylphe	enidate ER – Teva brand due to an
Tab extended-release 27 mg11.45	30	✓ Methylphenidate ER  - Teva
Tab extended-release 36 mg15.50	30	<ul><li>Methylphenidate ER</li><li>Teva</li></ul>
Tab extended-release 54 mg22.25	30	✓ Methylphenidate ER

#### ⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:

continued...

- Teva



Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.1 Applicant is a paediatrician or psychiatrist; or
- 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA2278 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency
- c) Note: Brand Switch Fee applies only to patients who have transferred from Methylphenidate ER Teva brand due to an out of stock.

Tab extended-release 18 mg - Brand switch fee payable

(Pharmacode 2665948) - see page 264 for details	58.96	30	Concerta
Tab extended-release 27 mg - Brand switch fee payable			
(Pharmacode 2665948) - see page 264 for details	65.44	30	Concerta
Tab extended-release 36 mg - Brand switch fee payable			
(Pharmacode 2665948) - see page 264 for details	71.93	30	Concerta
Tab extended-release 54 mg - Brand switch fee payable			
(Pharmacode 2665948) - see page 264 for details	86.24	30	Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg	20.40	30	Ritalin LA
Cap modified-release 30 mg	25.52	30	Ritalin LA
Cap modified-release 40 mg	30.60	30	Ritalin LA

#### ⇒SA2278 Special Authority for Subsidy

Initial application — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Fither:

## **NERVOUS SYSTEM**

Subsidy	Fully		Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\(\text{(manufacture of 1 nooy)}\)	Per	<b>✓</b>	Manufacturer	

continued...

- 1 All of the following:
  - 1.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
  - 1.2 Diagnosed according to DSM-IV or ICD 10 criteria; and
  - 1.3 Either:
    - 1.3.1 Applicant is a paediatrician or psychiatrist; or
    - 1.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
  - 1.4 Either:
    - 1.4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
    - 1.4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride; or
  - 2 All of the following:
    - 2.1 Patient meets the Special Authority criteria for SA1964 methylphenidate hydrochloride; and
    - 2.2 Patient would have been prescribed Methylphenidate ER Teva brand; and
    - 2.3 Patient is unable to access Methylphenidate ER Teva brand due to an out of stock.

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock on tab extended-release Methylphenidate ER – Teva subsidised under SA1964 (https://schedule.pharmac.govt.nz/latest/SA1964.pdf)

Renewal — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see SA1999 below – Retail pharmacy
Tab 100 mg .......29.13 60 ✓ Modavigil

## ⇒SA1999 Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
  - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamfetamine are contraindicated.

**Renewal** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.



Treatments for Dementia			
DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg		90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below -			•
Patch 4.6 mg per 24 hour	38.00	30	Rivastigmine Patch
			BNM 5
	90.00		Exelon Patch 5
Patch 9.5 mg per 24 hour	38.00	30	Rivastigmine Patch
•			BNM 10
	90.00		✓ Exelon Patch 10

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Per

Brand or

Generic

Manufacturer

### ⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

## Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg ......11.76

✓ Buprenorphine Naloxone BNM

Tab sublingual 8 mg with naloxone 2 mg .......34.00

**Buprenorphine** 

28

Naloxone BNM

#### ⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health;
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following

## **NERVOUS SYSTEM**

Subsid	dy Fi	ılly Brand or	
(Manufacture	r's Price) Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

**Renewal — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Tab modified-release 150 mg	11.00	30	✓ Zyban
DISULFIRAM Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see SA	A1408 below – Reta	ail pharmacy	
Tab 50 mg	77.77	28	✓ Naltrexone AOP S29
Naltraccord to be Principal Supply on 1 December 202	83.33	30	✓ Naltraccord

### ⇒SA1408 Special Authority for Subsidy

DUDDODION LIVEDOCLII ODIDE

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Te Whatu Ora or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

**Renewal** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
  - 2.1 Patient is still unstable and requires further treatment; or
  - 2.2 Patient achieved significant improvement but requires further treatment; or
  - 2.3 Patient is well controlled but requires maintenance therapy.



#### NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

b) Note. Direct i rovision by a pharmacist permitted ander the provisions i	11 1 411 1 01 0001	on A.
Patch 7 mg - Up to 28 patch available on a PSO19.14	28	<ul><li>Habitrol</li></ul>
Patch 7 mg for direct distribution only - [Xpharm]4.13	7	<ul><li>Habitrol</li></ul>
Patch 14 mg – Up to 28 patch available on a PSO21.05	28	<ul><li>Habitrol</li></ul>
Patch 14 mg for direct distribution only - [Xpharm]6.48	7	<ul><li>Habitrol</li></ul>
Patch 21 mg - Up to 28 patch available on a PSO24.12	28	<ul><li>Habitrol</li></ul>
Patch 21 mg for direct distribution only - [Xpharm]10.93	7	<ul><li>Habitrol</li></ul>
Lozenge 1 mg - Up to 216 loz available on a PSO19.76	216	<ul><li>Habitrol</li></ul>
Lozenge 1 mg for direct distribution only - [Xpharm]3.35	36	<ul><li>Habitrol</li></ul>
Lozenge 2 mg - Up to 216 loz available on a PSO21.65	216	<ul><li>Habitrol</li></ul>
Lozenge 2 mg for direct distribution only - [Xpharm]3.40	36	<ul><li>Habitrol</li></ul>
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO21.42	204	<ul><li>Habitrol</li></ul>
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]9.04	96	<ul><li>Habitrol</li></ul>
Gum 2 mg (Mint) – Up to 384 piece available on a PSO21.42	204	<ul><li>Habitrol</li></ul>
Gum 2 mg (Mint) for direct distribution only - [Xpharm]9.04	96	<ul><li>Habitrol</li></ul>
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO24.17	204	<ul><li>Habitrol</li></ul>
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]10.47	96	<ul><li>Habitrol</li></ul>
Gum 4 mg (Mint) - Up to 384 piece available on a PSO24.17	204	<ul><li>Habitrol</li></ul>
Gum 4 mg (Mint) for direct distribution only - [Xpharm]10.47	96	<ul><li>Habitrol</li></ul>

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 421	6.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg1		56	✓ Varenicline Pfizer

## **⇒SA1845** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either
  - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

**Renewal** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking;

## **NERVOUS SYSTEM**

Subsidy (Manufacturer's Pric	ce)	Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

and

- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

# **Chemotherapeutic Agents**

## **Alkylating Agents**

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA2153 below

Inj 25 mg vial	,		✓ Ribomustin
Inj 100 mg vial30	08.00	1	✓ Ribomustin
Inj 1 mg for ECP	3.23	l mg	✓ Baxter

#### **⇒SA2153** Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 Both:
    - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
    - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
  - 3.3 All of the following:
    - 3.3.1 The patient has not received prior bendamustine therapy; and
    - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+): and
    - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
  - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
  - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	(Manufacturer's Price) Sub-		Generic	
\$	Per	•	Manufacturer	

continued...

- 2 Both:
  - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
  - 2.2 Either:
    - 2.2.1 Both:
      - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
      - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
    - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2: and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with \* are unapproved indications.

BUSINEAN - PCT - Retail pharmacy-Specialist

BUSULFAN - PCT - Retail pharmacy-Specialist			•
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	DBL Carboplatin
	45.20		<ul> <li>Carboplatin Ebewe</li> </ul>
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	710.00	1	✓ BiCNU
Inj 100 mg for ECP		100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	<ul><li>Cisplatin Ebewe</li></ul>
Inj 1 mg per ml, 100 ml vial		1	✓ Cisplatin Ebewe
	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP	0.31	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	145.00	50	✓ Cyclonex
Inj 1 g vial - PCT - Retail pharmacy-Specialist	35.65	1	✓ Endoxan
	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist	71.25	1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
Inj 1 g	96.00	1	✓ Holoxan
Inj 2 g		1	✓ Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	<b>✓</b>	Manufacturer
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	✓	CeeNU
Cap 40 mg	399.15	20	✓	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	✓	Alkeran
Inj 50 mg - PCT only - Specialist		1	✓	Melpha
	67.80		✓	Alkeran
			1	Alkeran S29 S29
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
				100
	110.00		✓	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	33.35	1	✓	<b>Alchemy Oxaliplatin</b>
	46.32		✓	Oxaliplatin Accord
Inj 1 mg for ECP	0.35	1 mg	✓	Baxter
THIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	1	Bedford S29
, ,			/	Max Health \$29
			1	THIO-TEPA S29
				Tepadina
Inj 100 mg vial	CBS	1		Max Health \$29
,		•		Tepadina

#### **Antimetabolites**

		FIDINE - PCT only - Specialist - Special Authority see SA2141 below	ΑZ
✓ Azacitidine Dr	1	100 mg vial75.06	
Reddy's			
✓ Baxter	1 ma	1 mg for FCP 0.83	

## ⇒SA2141 Special Authority for Subsidy

**Initial application** only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
  - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
  - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

**Renewal** only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Cubaidu		F. II.	r Brand or
(1)	Subsidy Manufacturer's Pri	ce) S	Fully Subsidised	
,	\$	Per	1	Manufacturer
CALCIUM FOLINATE				
Tab 15 mg - PCT - Retail pharmacy-Specialist	135.33	10	✓	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	/	Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialist	7.28	1	✓	Calcium Folinate Sandoz
			✓	Calcium Folinate Sandoz S29 S29
	36.48	5	1	Eurofolic S29
Inj 50 mg - PCT - Retail pharmacy-Specialist	72.80	10	1	Leucovorin
, , , ,				Pharmacia S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	✓	Calcium Folinate Sandoz
	47.45	5	1	Eurofolic S29
Inj 100 mg - PCT only - Specialist	7.33	1	✓	Calcium Folinate Ebewe
	94.90	10	•	Leucovorin Pharmacia \$29
Inj 300 mg - PCT only - Specialist	22.51	1	✓	Calcium Folinate Ebewe
	25.14		1	Leucovorin DBL S29
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1		Calcium Folinate Sandoz
			✓	Calcium Folinate Sandoz S29 S29
Inj 1 g - PCT only - Specialist	67.51	1	✓	Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	✓	Calcium Folinate Sandoz
Inj 1 mg for ECP – PCT only – Specialist	0.06	1 mg	•	Baxter
CAPECITABINE – Retail pharmacy-Specialist	0.00	60	./	Consoitabine Vietnie
Tab 150 mg	10.00	60		Capecitabine Viatris Capercit
Capecitabine Viatris to be Principal Supply on 1 January 20			•	Caperon
Tab 500 mg		120	/	Capecitabine Viatris
<b>,</b>	49.00			Capecitabine-
			1	Capercit
Capecitabine Viatris to be Principal Supply on 1 January 2( Capercit Tab 150 mg to be delisted 1 January 2024)	024			
Caperon rab 130 mg to be delisted 1 January 2024)  (Capecitabine-DRLA 829 Tab 500 mg to be delisted 1 January 20.	24)			
Capercit Tab 500 mg to be delisted 1 January 2024)	<i>24)</i>			
CLADRIBINE - PCT only - Specialist			_	
Inj 2 mg per ml, 5 ml		1	<b>√</b>	Litak S29
Inj 1 mg per ml, 10 ml		1	· •	Leustatin
Inj 10 mg for ECP	/49.96	10 mg O	<b>/</b>	Baxter

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy	D.:\ 0k-	Fully Brand or
(Manufacturer's \$	Price) Subs	sidised Generic  Manufacturer
YTARABINE		
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialist472.00	5	✓ Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail		
pharmacy-Specialist	1	✓ Pfizer
Inj 1 mg for ECP - PCT only - Specialist	10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specialist94.40	100 mg OP	✓ Baxter
LUDARABINE PHOSPHATE	•	
Tab 10 mg - PCT - Retail pharmacy-Specialist412.00	20	✓ Fludara Oral
Inj 50 mg vial – PCT only – Specialist	5	✓ Fludarabine Ebewe
Inj 50 mg for ECP – PCT only – Specialist	50 mg OP	✓ Baxter
LUOROURACIL	3 -	
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist10.51	1	✓ Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist	1	✓ Fluorouracil Accord
Inj 1 mg for ECP – PCT only – Specialist	100 mg	✓ Baxter
EMCITABINE HYDROCHLORIDE - PCT only - Specialist	3	
Inj 1 g, 26.3 ml vial	1	✓ DBL Gemcitabine
Inj 1 g	i	✓ Gemcitabine Ebewe
Inj 1 mg for ECP	1 mg	✓ Baxter
RINOTECAN HYDROCHLORIDE - PCT only - Specialist	9	
Inj 20 mg per ml, 5 ml vial52.57	1	✓ Accord
71.44	•	✓ Irinotecan Actavis
71.77		100
100.00		✓ Irinotecan-Rex
Inj 1 mg for ECP	1 mg	✓ Baxter
ERCAPTOPURINE	9	
Tab 50 mg - PCT - Retail pharmacy-Specialist25.90	25	✓ Puri-nethol
Oral suspension 20 mg per ml — Retail pharmacy-Specialist —	25	Fun-nethor
Special Authority see SA1725 below	100 ml OP	✓ Allmercap
On 1705 On a sight Authority See SA 1725 Delow	100 IIII OF	- Allilleicap

## **⇒SA1725** Special Authority for Subsidy

**Initial application** only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

**Renewal** only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

		Subsidy (Manufacturer's Price	) ;	Fully Subsidised	
		\$	Per	1	Manufacturer
ME	THOTREXATE				
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist	9.98	90	✓	Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist		90	✓	Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist	56.05	5	1	Methotrexate DBL
*	Inj 7.5 mg prefilled syringe	14.61	1	/	Methotrexate Sandoz
*	Inj 10 mg prefilled syringe	14.66	1	/	Methotrexate Sandoz
*	Inj 15 mg prefilled syringe	14.77	1	✓	Methotrexate Sandoz
*	Inj 20 mg prefilled syringe	14.88	1	✓	Methotrexate Sandoz
*	Inj 25 mg prefilled syringe	14.99	1	✓	Methotrexate Sandoz
*	Inj 30 mg prefilled syringe	15.09	1	✓	Methotrexate Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialis	st30.00	5	✓	Methotrexate DBL Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specia	list45.00	1	✓	DBL Methotrexate Onco-Vial
* *	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist Inj 100 mg per ml, 50 ml vial - PCT - Retail	25.00	1	✓	Methotrexate Ebewe
	pharmacy-Specialist	67.99	1	1	Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓	Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist	4.73	5 mg Ol	•	Baxter
PΕ	METREXED - PCT only - Specialist - Special Authority see S	SA1679 below			
	Inj 100 mg vial	60.89	1	1	Juno Pemetrexed
	Inj 500 mg vial		1	✓	Juno Pemetrexed
	Inj 1 mg for ECP	0.55	1 mg	1	Baxter

#### ⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Fither:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 2.1 Both:
  - 2.1.1 Patient has chemotherapy-naïve disease; and
  - 2.1.2 Permetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
- 2.2 All of the following:
  - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
  - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
  - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m<sup>2</sup> every 21 days.

THIOGUANINE - PCT - Retail pharmacy-Specialist	
Tah 40 mg	

Tab 40 mg .......126.31 25 **✓ Lanvis** 

## Other Cytotoxic Agents

AMSACRINE – PCT only – Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharma	cy-Specialist		
Cap 0.5 mg		100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter
BLEOMYCIN SULPHATE - PCT only - Specialist			
Inj 15,000 iu, vial	185.16	1	✓ DBL Bleomycin  Sulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	✓ Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority	see SA1889 below		
Inj 3.5 mg vial	74.93	1	✓ DBL Bortezomib
Inj 1 mg for ECP	22.26	1 mg	✓ Baxter

#### ⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis \*.

Note: Indications marked with \* are unapproved indications.

DACARBAZINE - PCT only - Specialist

Inj 200 mg vial	72.11	1	✓ DBL Dacarbazine
, .	580.60	10	<ul> <li>Dacarbazine</li> </ul>
			APP S29
Ini 200 mg for ECP	72 11	200 ma OP	✓ Rayter

	Subsidy		Fully	
	(Manufacturer's P \$	rice) Subsi Per	idised •	Generic Manufacturer
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist				
Inj 0.5 mg vial	255.00	1	/	Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP		Baxter
	200.00	0.5 mg Oi	٠	Daxiei
DAUNORUBICIN – PCT only – Specialist	474.00			D."
Inj 2 mg per ml, 10 ml		1		Pfizer
Inj 20 mg vial	1,495.00	10	•	Daunorubicin Zentiva \$29
Inj 20 mg for ECP	171.93	20 mg OP	1	Baxter
DOCETAXEL - PCT only - Specialist				
Inj 20 mg	48.75	1	/	Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1		DBL Docetaxel
Inj 20 mg per ml, 4 ml vial		1	_	Docetaxel
11) 20 119 poi 1111, 1 111 Viai		•		Accord \$29
Ini 00 ma	105.00	1	./	Docetaxel Sandoz
Inj 80 mg				Baxter
Inj 1 mg for ECP	0.35	1 mg	•	Daxier
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml vial		1		Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1		Doxorubicin Ebewe
	17.00			Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial		1		Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	65.00	1	/	Arrow-Doxorubicin
	69.99		1	Accord S29
			1	Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	1	Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml vial	25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1		Epirubicin Ebewe
Inj 1 mg for ECP		1 mg		Baxter
ETOPOSIDE		9		
	240.72	20	./	Vanagid
Cap 50 mg - PCT - Retail pharmacy-Specialist		10		Vepesid Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10		Rex Medical
Inj 1 mg for ECP – PCT only – Specialist		· ·		Baxter
, ,	0.09	1 mg	•	Daxiei
ETOPOSIDE PHOSPHATE - PCT only - Specialist			_	
Inj 100 mg (of etoposide base)		1		Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg	/	Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail phari	macy-Specialist			
Cap 500 mg		100	1	Devatis
Devatis to be Principal Supply on 1 December 2023				
BRUTINIB - Special Authority see SA2168 below - Retail pharm	nacy			
Tab 140 mg	•	30	/	Imbruvica
Tab 420 mg	•	30		Imbruvica
1 au 720 mg	3,002.00	30	•	iiiibi uvica

**⇒SA2168** Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

ubsidised •	Generic Manufacturer	
Per	Per 🗸	Per Manufacturer

continued...

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
  - 4.1 Both:
    - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
    - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
    - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
    - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
  - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- Both:
  - 1 No evidence of clinical disease progression; and
  - 2 The treatment remains appropriate and the patient is benefitting from treatment.

Ini 5 mg vial. – PCT only – Specialist 109.74

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

#### IDARUBICIN HYDROCHLORIDE

Inj 10 mg vial – PCT only – Specialist		1	✓ Zavedos
Inj 1 mg for ECP - PCT only - Specialist		1 mg	✓ Baxter
LENALIDOMIDE - Retail pharmacy-Specialist - Special Authority	see SA2047 be	elow	
Wastage claimable			
Cap 5 mg	5,122.76	28	Revlimid

Cap 10 mg	4,655.25	21	Revlimid
	6,207.00	28	✓ Revlimid
Cap 15 mg	5,429.39	21	✓ Revlimid
	7,239.18	28	✓ Revlimid
Cap 25 mg	7,627.00	21	✓ Revlimid

### ⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Fither:
  - 3.1 Lenalidomide to be used as third line\* treatment for multiple myeloma; or
  - 3.2 Both:
    - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
    - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments: and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or

continued...

✓ Zavedos

Subsidy		Fully	Brand or	
(Manufacturer's Price		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with \* is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

#### **MFSNA**

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	<ul><li>Uromitexan</li></ul>
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	<ul><li>Uromitexan</li></ul>
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialis	t177.45	15	<ul><li>Uromitexan</li></ul>
Inj 100 mg per ml, 10 ml ampoule - PCT only - Special	ist407.40	15	<ul><li>Uromitexan</li></ul>
Inj 1 mg for ECP - PCT only - Specialist	2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial	641.70	1	✓ Accord S29
Inj 20 mg vial		1	✓ Teva
Inj 1 mg for ECP	269.85	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority	see SA2163 below		
Tab 100 mg	3,701.00	56	✓ Lynparza
Tab 150 mg	3,701.00	56	✓ Lynparza

#### ⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Either:
  - 3.1 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.1.1 Patient has newly diagnosed, advanced disease; and
- 3.1.2 Patient has received one line\*\* of previous treatment with platinum-based chemotherapy; and
- 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen: or
- 3.2 All of the following:
  - 3.2.1 Patient has received at least two lines\*\* of previous treatment with platinum-based chemotherapy; and
  - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line\*\* of platinum-based chemotherapy; and
  - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
  - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen: and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

## All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
  - 2.1 No evidence of progressive disease; or
  - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
  - 5.1 Both:
    - 5.1.1 Patient has received one line\*\* of previous treatment with platinum-based chemotherapy; and
    - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
  - 5.2 Patient has received at least two lines\*\* of previous treatment with platinum-based chemotherapy.

Notes: \*Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component. \*\*A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg	24.00	1	✓ Paclitaxel Ebewe
	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see	SA1979 on the next page		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

Sub	bsidy Fu	lly Brand or	_
(Manufactu	urer's Price) Subsidise	ed Generic	
	\$ Per	<ul> <li>Manufacturer</li> </ul>	

### ⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

#### PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharm	acy-Specialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA2275 below - F	Retail pharmacy		
Cap 5 mg	9.13	5	✓ Temaccord
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
Cap 100 mg	35.98	5	✓ Temaccord
	40.20		✓ Apo-Temozolomide
Cap 140 mg	50.12	5	✓ Temaccord
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ Temaccord

#### ⇒SA2275 Special Authority for Subsidy

Initial application — (gliomas) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. Renewal — (gliomas) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

**Initial application — (neuroendocrine tumours)** only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour\*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

**Renewal — (neuroendocrine tumours)** only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

**Initial application** — **(ewing's sarcoma)** only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

**Renewal — (ewing's sarcoma)** only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a \* is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Au	thority see SA1124 below		
Cap 50 mg	378.00	28	✓ Thalomid
Cap 100 mg	756.00	28	✓ Thalomid

### ⇒SA1124 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis\*.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with \* is an unapproved indication.

#### **TRETINOIN**

Cap 10 mg - PCT - Retail pharmacy-Specialist	479.50	100	Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority	see SA1868 bel	ow	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OP	✓ Venclexta
Tab 10 mg		2 OP	✓ Venclexta
Tab 50 mg	239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable	8,209.41	120	✓ Venclexta

#### ⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and

/ Hoonira

✓ Alecensa

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
	Per	1	Manufacturer	

continued...

2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has previously untreated chronic lymphocytic leukaemia; and

Ini 1 ma nor ml 10 ml vial DCT Datail pharmacy Specialist

- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)\* and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

070 07

#### VINBLASTINE SULPHATE

inj i mg per mi, 10 mi viai – PC i – Retail pharmacy-Specialist270.37	5	▼ Hospira
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	<ul><li>DBL Vincristine Sulfate</li></ul>
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist102.73	5	<ul><li>DBL Vincristine Sulfate</li></ul>
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE		
Cap 20 mg30.00	1	✓ Vinorelbine Te Arai
Cap 30 mg40.00	1	✓ Vinorelbine Te Arai
Cap 80 mg60.00	1	✓ Vinorelbine Te Arai
Inj 10 mg per ml, 1 ml vial - PCT only - Specialist12.00	1	✓ Navelbine
42.00		✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial - PCT only - Specialist56.00	1	✓ Navelbine
168.00		✓ Navelbine S29 S29
210.00		✓ Vinorelbine Ebewe
328.65		✓ Sagent S29
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
Inj 50 mg for ECP - PCT only - Specialist328.65	50 mg OP	✓ Baxter (Sagent)
(Neverthing In: 10 man year of male sights be delicated 1 October 2004)	Ū	, ,

(Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 October 2024) (Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 October 2024) (Baxter (Sagent) Inj 50 mg for ECP to be delisted 1 December 2023)

# Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist.

Subsidy		Fully	Brand or
(Manufacturer's Price)	_	Subsidised	Generic
<u> </u>	Per		Manutacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- 3 Patient has an ECOG performance score of 0-2.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

#### DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable

Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel

## ⇒SA1805 Special Authority for Subsidy

**Initial application** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
  - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
  - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
  - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
  - 3.1 The patient has a diagnosis of CML in chronic phase; and
  - 3.2 Maximum dose of 100 mg/day; and
  - 3.3 Any of the following:
    - 3.3.1 Patient has documented treatment failure\* with imatinib: or
    - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
    - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
    - 3.3.4 Patients is enrolled in the KISS study\*\* and requires dasatinib treatment according to the study protocol.

**Renewal** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib\*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: \*treatment failure for CML as defined by Leukaemia Net Guidelines. \*\*Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB – Retaii pnarmacy-Specialist – Special Authority s	see SA2115 on the r	iext page	
Tab 100 mg	329.70	30	✓ Alchemy
Tab 150 mg	569.70	30	✓ Alchemy

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

### ⇒SA2115 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either
  - 3.1 Patient is treatment naive; or
  - 3.2 Both:
    - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
    - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

## All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2116 below

## **⇒SA2116** Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 4 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
  - 2.1 Patient is treatment naive; or
  - 2.2 Both:
    - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
    - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

#### IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg - [Xpharm] - Special Authority see SA1460		
below2,400.00	60	✓ Glivec
Cap 100 mg44.93	60	✓ Imatinib-Rex
Imatinib-Rex to be Principal Supply on 1 December 2023		
Cap 400 mg69.76	30	✓ Imatinib-Rex
Imatinib-Rex to be Principal Supply on 1 December 2023		

(Glivec Tab 100 mg to be delisted 1 December 2023)

#### ⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10, 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

#### Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

#### LAPATINIB DITOSYLATE - Special Authority see SA2035 below - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

70 Tvkerb

#### ⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

#### NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable 120 ✓ Tasigna 120 ✓ Tasigna 

#### ⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### continued...

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
  - 2 Fither
  - 2.1 Patient has documented CML treatment failure\* with imatinib: or
    - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: \*treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines: and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

## PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

Wastage claimable

Tab 75 mg	4,000.00	21	Ibrance
Tab 100 mg	4,000.00	21	Ibrance
Tab 125 mg	4,000.00	21	✓ Ibrance

### ⇒SA1894 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

4 Either:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- seco

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Fither:
  - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
  - 4.2.2.2 All of the following:
    - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
    - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
    - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB – Special Authority	$\prime$ see SA1190 on the next $\mid$	page – Retail pharmacy
-------------------------------	--	------------------------

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	1	Manufacturer	

## ⇒SA1190 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 Both:
    - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Was	stage	claimable	
	_		

Tab 5 mg	2,500.00	56	Jakavi
Tab 10mg		56	<ul><li>Jakavi</li></ul>
Tab 15 mg		56	Jakavi
Tab 20 mg	·	56	Jakavi

#### ⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
  - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
  - 2.2 Both:
    - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and

Subsidy (Manufacturer's Price) \$	) Per	Fully Subsidised	Brand or Generic Manufacturer	
<u> </u>				

continued...

- 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy;
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authorit	see SA2117 below -	Retail pharmacy
------------------------------	--------------------	-----------------

Cap 12.5 mg	28	<ul> <li>Sunitinib Pfizer</li> </ul>
Cap 25 mg416.77	28	✓ Sunitinib Pfizer
Cap 50 mg	28	✓ Sunitinib Pfizer

#### ⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
    - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Fither:
  - 2.1 The patient's disease has progressed following treatment with imatinib; or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	
--	--

continued...

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
  - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

**Renewal — (GIST pandemic circumstances)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

## **Endocrine Therapy**

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 92

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

Tab 250 mg .......4,276.19 120 ✓ Zytiga

#### ⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
  - 4.1 All of the following:
    - 4.1.1 Patient is symptomatic; and

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	<ul> <li>Manufacturer</li> </ul>	

continued...

- 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
- 4.1.3 Patient has ECOG performance score of 0-1; and
- 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
- 4.2 All of the following:
  - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
  - 4.2.2 Patient has ECOG performance score of 0-2; and
  - 4.2.3 Patient has not had prior treatment with abiraterone.

**Renewal — (abiraterone acetate)** only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression, and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

**Renewal — (pandemic circumstances)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

#### **BICALUTAMIDE**

Tab 50 mg	4.18	28	Binarex
Binarex to be Principal Supply on 1 December 2023			
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Authority	y see SA1895 bel	ow	
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex

#### ⇒SA1895 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
OCTREOTIDE				
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓	Max Health
			/	Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	1	Max Health
,			1	Octreotide GH S29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	/	Max Health
., , , ,		•	1	Octreotide GH S29
OCTREOTIDE LONG-ACTING - Special Authority see SA2119	below – Retail pharm	acv		
Inj depot 10 mg prefilled syringe		1	/	Octreotide Depot
,				Teva
	1,152.00		1	Sandostatin LAR
Inj depot 20 mg prefilled syringe	647.03	1	✓	Octreotide Depot
				Teva
	1,539.00		1	Sandostatin LAR
Inj depot 30 mg prefilled syringe	718.55	1	✓	Octreotide Depot
				Teva

## ⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea\* and vomiting\* due to malignant bowel obstruction\*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Initial application — (Acromegaly)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
  - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
  - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
  - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
  - 2.1 Gastrinoma: and
  - 2.2 Fither:
    - 2.2.1 Patient has failed surgery: or
    - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
  - 3.1 Insulinomas: and
  - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
  - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
  - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly: and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

IAMOXIF	EN CH	RAIL
IAMONII		NAIL

*	Tab 10 mg15.00	60	✓ Tamoxifen Sandoz
	Tamoxifen Sandoz to be Principal Supply on 1 December 2023		
*	Tab 20 mg5.32	60	✓ Tamoxifen Sandoz
	Tamoxifen Sandoz to be Principal Supply on 1 December 2023		

# **Aromatase Inhibitors**

ANASTROZOLE  * Tab 1 mg4.39  Anatrole to be Principal Supply on 1 December 2023	9	30	✓ Anatrole
** Tab 25 mg9.86     Pfizer Exemestane to be Sole Supply on 1 November 2023	6	30	✓ Pfizer Exemestane
LETROZOLE	4	30	✓ <u>Letrole</u>

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
ė.	Dor ./	Monufacturar

## **Immunosuppressants**

## Cytotoxic Immunosuppressants

ΑZ	ATHI	OP	RIN	ΙE
	T - 1-	0-		

不	1ab 25 mg	00	▼ <u>Azamun</u>
*	Tab 50 mg8.10	100	✓ <u>Azamun</u>

#### MYCOPHENOLATE MOFETII

TCOPHENOLATE MOPETIL			
Tab 500 mg	35.90	50	<ul><li>Cellcept</li></ul>
Cap 250 mg	35.90	100	✓ Cellcept
Powder for oral lig 1 g per 5 ml - Subsidy by endorsement		165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

### **Fusion Proteins**

ETANERCEPT - Special Authority see SA2103 below - Retail pharmacy

Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Ini 50 ma prefilled syringe	1.050.00	4	✓ Enbrel

### ⇒SA2103 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Either:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a Te Whatu Ora Hospital; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Por 🗸	Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA: or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

# Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

## Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA: or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or

Subsi	idy Fully	Brand or
(Manufacture	er's Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

of the following.

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

- All of the following:
  - 1 Patient has shown clinical improvement; and
  - 2 Patient continues to require treatment; and
  - 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
  - 2.5 Either:
    - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
  - 2.6 Fither:
    - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application** — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a dermatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:

_			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	•	Por 🗸	Manufacturer

continued...

- 2.1 Both:
  - 2.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
  - 2.1.2 Either:
    - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
    - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2.2 Both:
  - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 2.2.2 Either:
    - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
  - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application: or
  - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

	Subsidy	F	ully	Brand or
(Mar	ufacturer's Price)	Subsid	sed	Generic
	\$	Per	✓	Manufacturer

continued...

- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

## Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Speci	ialist		
Inj 50 mg per ml, 5 ml	2,774.48	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only	y – Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29

## **Monoclonal Antibodies**

	armacy	ee SA2178 below – Retail p	ADALIMUMAB (AMGEVITA) - Special Author
✓ Amgevita	1	190.00	Inj 20 mg per 0.4 ml prefilled syringe
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled pen
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled syringe

#### ⇒SA2178 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has severe Behcet's disease\* that is significantly impacting the patient's quality of life; and
  - 2 Either:
    - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
    - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with \* are unapproved indications.

**Initial application — (Hidradenitis suppurativa)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

Subsidy	Full	y Brand or
(Manufacturer's Price)	Subsidise	d Generic
\$	Per •	Manufacturer

continued...

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plague psoriasis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
  - 1.2 Either:
    - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
    - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
  - 2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
  - 2.2 Fither:
    - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with \* are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
- 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
- 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
- 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

**Initial application — (Crohn's disease - children)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

**Initial application — (Crohn's disease - fistulising)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (Ocular inflammation - chronic)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

**Initial application — (ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects; or
  - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
  - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
  - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Fither:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Either:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

**Renewal — (Arthritis - oligoarticular course juvenile idiopathic)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Fither:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects: or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
  - 2.4 Fither:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
    - 2.5.2 Patient has an ESR greater than 25 mm per hour; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (Arthritis - psoriatic)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis: or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate at maximum tolerated doses (unless contraindicated); and
  - 2.5 Fither:
    - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
  - 2.6 Either:
    - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

**Renewal — (Arthritis - rheumatoid)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

1 Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
- 1.2 Fither:
  - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
    - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Initial application — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis: and
- 2 Fither:
  - 2.1 Patient's SCCAI score is greater than or equal to 4; or
  - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
  - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
<u> </u>	\$ Pe	er 🗸	Manufacturer

continued...

2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (inflammatory bowel arthritis – peripheral)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 on the next page - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe	1,599.96	2	✓ Humira
Inj 40 mg per 0.4 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.4 ml prefilled syringe	1,599.96	2	<ul><li>Humira</li></ul>
Inj 40 mg per 0.8 ml prefilled pen		2	✓ HumiraPen
Ini 40 mg per 0.8 ml prefilled syringe	1,599.96	2	✓ Humira

(HumiraPen Inj 40 mg per 0.8 ml prefilled pen to be delisted 1 March 2024) (Humira Inj 40 mg per 0.8 ml prefilled syringe to be delisted 1 March 2024)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

# ⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease - severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- Both:
  - 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
  - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline: and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
    - 1.1.2 Either:
      - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
      - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 1.2 Both:
    - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 1.2.2 Either:
      - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (Pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

**Renewal — (Pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in

Subsidy (Manufacturer's Pric	ce)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

treatment regimen; or

- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
  - 1.2 PCDAI score is 15 or less: or
  - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (Ocular inflammation – chronic)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
  - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1 The patient has had a good clinical response following 3 initial doses; or
- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
  - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
  - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
  - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

⇒SA1772 Special Authority for Subsidy

**Initial application — (wet age related macular degeneration)** only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
  - 1.3 There is no structural damage to the central fovea of the treated eye; and
  - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
  - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
  - 2.2 Patient has previously\* (\*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

**Initial application — (diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

**Renewal — (wet age related macular degeneration)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

# ⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10°9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
  - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
  - 9.2 Both:
    - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
    - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

**Renewal — (Severe eosinophilic asthma)** only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

- Both:
  - 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
  - 2 Fither:
    - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
    - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

Inj 120 mg per ml casirivimab, 11.1 ml vial (1) and inj 120 mg

#### ⇒SA2096 Special Authority for Subsidy

**Initial application** — (**Treatment of profoundly immunocompromised patients**) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity\*; and
- 3 Patient is profoundly immunocompromised\*\* and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: \* Mild to moderate disease severity as described on the Ministry of Health Website

\*\* Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

#### CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 below

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	Erbitux
Ini 1 mg for ECP		1 ma	✓ Baxter

#### ⇒SA1697 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

GEMTUZUMAB OZOGAMICIN - PCT only - Specialist - Special Authority see SA2269 below

#### ⇒SA2269 Special Authority for Subsidy

Initial application only from a haematologist, paediatric haematologist or paediatric oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and

Subsid	dy Fi	ılly Brand or	
(Manufacture	r's Price) Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2179 below

Inj 100 mg	428.00	1	Remicade
Inj 1 mg for ECP	4.40	1 mg	✓ Baxter

⇒SA2179 Special Authority for Subsidy

**Initial application** — (**Crohn's disease (adults))** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Subsidy (Manufacturer's Pr	ice)	Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

continued...

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less: or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — (**Graft vs host disease**) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

**Initial application — (Pulmonary sarcoidosis)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less: and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

**Initial application — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Per 🗸	Manufacturer

continued...

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
  - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
  - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
  - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement: and
- 3 Patient has steroid-refractory disease; and

Su	ubsidy	Fully	Brand or
(Manufac	cturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

- 4 Either:
  - 4.1 IV cyclophosphamide has been tried; or
  - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

**Renewal — (neurosarcoidosis)** only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

#### Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
  - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  - 2.2 There has been a marked reduction in prednisone dose; and
  - 2.3 Fither:
    - 2.3.1 There has been an improvement in MRI appearances; or
    - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plague psoriasis; and
  - 1.2 Fither:
    - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
    - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
  - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Fither:
  - 1.1 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
  - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
  - 1.2.2 Fither:
    - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 Rheumatoid arthritis: or
  - 2.2 Ankylosing spondylitis; or
  - 2.3 Psoriatic arthritis; or
  - 2.4 Severe ocular inflammation; or
  - 2.5 Chronic ocular inflammation: or
  - 2.6 Crohn's disease (adults): or
  - 2.7 Crohn's disease (children); or
  - 2.8 Fistulising Crohn's disease: or
  - 2.9 Severe fulminant ulcerative colitis: or
  - 2.10 Severe ulcerative colitis: or
  - 2.11 Plaque psoriasis; or
  - 2.12 Neurosarcoidosis: or
  - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis: and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
  - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subs (Manufactur		
\$	Per	Manufacturer

continued...

- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

**Renewal — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

**Initial application** — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
  - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	<ul> <li>Manufacturer</li> </ul>	

continued...

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or

#### 2 Both:

- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2.2 Any of the following:
  - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
  - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
  - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
  - 2.1 Patients SCCAI is greater than or equal to 4; or
  - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab: or
  - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs: and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum

|--|

continued...

tolerated dose (unless contraindicated); and

- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (inflammatory bowel arthritis – peripheral)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

#### 

(Nucala Inj 100 mg vial to be delisted 1 August 2024)

## ⇒SA2154 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
  - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
  - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
  - 9.2 Both:

 Subsidy Manufacturer's Price)	٤	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
- 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

**Renewal — (Severe eosinophilic asthma)** only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Fither:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB – PCT only – Specialist – Special Authority see SA2155 below		
Inj 25 mg per ml, 40 ml vial5,910.00	1	✓ Gazyva
Inj 1 mg for ECP6.21	1 mg	✓ Baxter

SA2155 Special Authority for Subsidy

**Initial application — (chronic lymphocytic leukaemia)** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts\* unless the cytopenias are a consequence of marrow infiltration by CLL: and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

\* Neutrophil greater than or equal to  $1.5 \times 10^9$ /L and platelets greater than or equal to  $75 \times 10^9$ /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has follicular lymphoma; or
  - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen\*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy\*.

Note: \* includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	osidised	Generic
\$	Per	•	Manufacturer

continued

All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

#### OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe	450.00	1	✓ Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

## ⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
  - 1 Patient must be aged 12 years or older; and
  - 2 Either:
    - 2.1 Both:
      - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
      - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
    - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
  - 3 Any of the following:
    - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
    - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
    - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
  - 4 Either:
    - 4.1 Treatment to be stopped if inadequate response\* following 4 doses; or
    - 4.2 Complete response\* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded\* to 6 doses of omalizumab; or
- 2 Both:
  - 2.1 Patient has previously had a complete response\* to 6 doses of omalizumab; and
  - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: \*Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Specialist - Special Authority see SA2143 below

(Synagis Inj 100 mg per ml, 1 ml vial to be delisted 1 January 2024)

#### ⇒SA2143 Special Authority for Subsidy

Initial application — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Infant was born in the last 2 years and has severe lung, airway, neurological or neuromuscular disease that requires ongoing, life-sustaining community ventilation; or
- 2 Both:
  - 2.1 Infant was born in the last 12 months; and
  - 2.2 Any of the following:
    - 2.2.1 Patient was born at less than 28 weeks gestation; or
    - 222 Roth
      - 2.2.2.1 Patient was born at less than 32 weeks gestation; and
      - 2.2.2.2 Fither:
        - 2.2.2.2.1 Patient has chronic lung disease; or
        - 2.2.2.2. Patient is Maori or any Pacific ethnicity; or
    - 2.2.3 Both:
      - 2.2.3.1 Patient has haemodynamically significant heart disease; and
      - 2.2.3.2 Any of the following:
        - 2.2.3.2.1 Patient has unoperated simple congenital heart disease with significant left to right shunt (see
        - 2.2.3.2.2 Patient has unoperated or surgically palliated complex congenital heart disease; or
        - 2.2.3.2.3 Patient has severe pulmonary hypertension (see note b); or
        - 2.2.3.2.4 Patient has moderate or severe LV failure (see note c).

#### Notes:

- a) Patient requires/will require heart failure medication, and/or patient has significant pulmonary hypertension, and/or patient will require surgical palliation/definitive repair within the next 3 months.
- b) Mean pulmonary artery pressure more than 25 mmHg.
- c) LV Ejection Fraction less than 40%.

Renewal — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months where patient still meets initial criteria.

	Subsidy (Manufacturer's P \$	rice) Su Per	Fully bsidised	Brand or Generic Manufacturer					
PERTUZUMAB – PCT only – Specialist – Special Authority see SA2276 below									
Inj 30 mg per ml, 14 ml vial	3,927.00	1	<b>✓</b> P	Perjeta					
Inj 420 mg for ECP	3,927.00	420 mg OF	<b>✓</b> E	Baxter					

⇒SA2276 Special Authority for Subsidy

**Initial application — (metastatic breast cancer)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 Patient is chemotherapy treatment naïve; or
  - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

# 1 Both:

- 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Speci	al Authority see SA19	76 below	
Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

# ⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	✓	

continued...

the maximum tolerated dose of ciclosporin; or

- 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Fither:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
    - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:

ibsidy	Fully	Brand or
turer's Price) Sub	osidised	Generic
 \$ Per	/	

continued...

- 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

## RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2233 below

Inj 100 mg per 10 ml vial	275.33 2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20 1	✓ Riximyo
Inj 1 mg for ECP	1.38 1 mg	✓ Baxter (Riximyo)

#### ⇒SA2233 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant\*.

Note: Indications marked with \* are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
  - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
  - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
  - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
  - 3.4 Patient is a female of child-bearing potential; or
  - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

continued...

4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without

further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection\*.

Note: Indications marked with \* are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
  - 2.1 The patient is rituximab treatment naive: or
  - 2.2 Either:
    - 2.2.1 The patient is chemotherapy treatment naive; or
    - 2.2.2 Both:
      - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
      - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy: or
  - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax: and
- 3 The patient has good performance status; and
- 4 Fither:
  - 4.1 The patient does not have chromosome 17p deletion CLL: or
  - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax: or
  - 1.2 All of the following:
    - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL:
    - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment;
    - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
    - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

**Initial application — (Post-transplant)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:
  - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
  - 3.2 Both:
    - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
    - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS\* or FRNS\*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects: and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS\* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of 4 weeks.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

Note: Indications marked with \* are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

**Initial application** — **(haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

**Renewal** — **(haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microlitre; or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

- 1.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
  - 2.1 Treatment with steroids and splenectomy have been ineffective; or
  - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy): and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia\* requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia\*

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least

Note: Indications marked with \* are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease\*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
  - 2.1 Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
  - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

Note: Indications marked with \* are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE\*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and

Subsidy (Manufacturer's Price)	Fu Subsidis	ılly	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia\*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

**Initial application — (severe antisynthetase syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
  - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
  - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Fither:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	idised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Roth:
  - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy\*; or
  - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy\*; and
- 2 Either:
  - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment: or
  - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

#### Notes:

- a) Indications marked with \* are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

experienced intolerable side effects.

d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma\*: and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> per dose for a maximum of 18 doses.

Note: Indications marked with \* are unapproved indications.

**Initial application** — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant\*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with \* are unapproved indications.

Initial application — (pemiphigus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 All of the following:

- 1.1 Patient has severe rapidly progressive pemphigus; and
- 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
- 1.3 Any of the following:
  - 1.3.1 Skin involvement is at least 5% body surface area; or
  - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
  - 1.3.3 Involvement of two or more mucosal sites; or

2 Both:

- 2.1 Patient has pemphigus; and
- 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with \* are unapproved indications.

Renewal — (pemiphigus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with \* are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD\*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD\*; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with \* are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD\*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Treatment with rituximab for IgG4-RD\* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
  - 1.2 Patient is receiving maintenance treatment for IgG4-RD\*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with \* are unapproved indications.

SECUKINUMAB - Special Authority see SA2084 below - Retail pharmacy

#### ⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Te Whatu Ora Hospital, for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
  - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less: and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
    - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis: or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
- 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
  - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
  - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

#### 1 Fither:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

#### SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Inj 100 mg vial	770.57	1	Sylvant
Inj 400 mg vial	3,082.33	1	✓ Sylvant

#### ⇒SA1596 Special Authority for Subsidy

**Initial application** only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

**Renewal** only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

#### TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

		nab 100 mg per	Inj 100 mg per ml, 1.5 ml vial with cilgavim
<ul><li>Evusheld</li></ul>	1	0.00	ml,1.5 ml vial
		y see SA2159 on the next page	OCILIZUMAB - PCT only - Special Authority
/ A stanson	4	000.00	Ind OO man man and A maladal

<ul><li>Actemra</li></ul>	1	ıal220.00	Inj 20 mg per ml, 4 ml vial
✓ Actemra	1	vial550.00	Inj 20 mg per ml, 10 ml via
✓ Actemra	1	vial1,100.00	Inj 20 mg per ml, 20 ml via

TO

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### ⇒SA2159 Special Authority for Subsidy

**Initial application — (cytokine release syndrome)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
  - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
  - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
  - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
  - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
  - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 rheumatoid arthritis; or
  - 2.2 systemic juvenile idiopathic arthritis: or
  - 2.3 adult-onset Still's disease; or
  - 2.4 polyarticular juvenile idiopathic arthritis: or
  - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Te Whatu Ora Hospital; and
    - 3.2.2 Either:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Fither:
  - 3.1 Treatment with methotrexate is contraindicated; or
  - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Fither
  - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
  - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
  - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
  - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Fither:
  - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Fither:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a Te Whatu Ora Hospital; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Initial application — (polyarticular juvenile idiopathic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course iuvenile idiopathic arthritis (JIA): and
  - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
  - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.4 Any of the following:
    - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

**Renewal — (adult-onset Still's disease)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. **Renewal — (polyarticular juvenile idiopathic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a

Subsidy		Fully	Brand or
(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
<b>*</b>	1 61		Manuacturer

continued...

rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA2277 below

Inj 150 mg vial	1,350.00	1	<ul><li>Herceptin</li></ul>
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	9.36	1 mg	✓ Baxter

#### ⇒SA2277 Special Authority for Subsidy

**Initial application** — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
  - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned: or
  - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
  - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
  - 1.3 Any of the following:
    - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or 1.3.2 Both:
      - 1.3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerable side effects; and
      - 1.3.2.2 The cancer did not progress whilst on lapatinib; or
    - $1.3.3 \ \ \text{The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and }$
  - 1.4 Either:
    - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
    - 1.4.2 All of the following:
      - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 1.5 Trastuzumab not to be given in combination with lapatinib; and
- 1.6 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: \* For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

**Initial application — (metastatic breast cancer)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 2.2 Both:
    - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerable side effects; and
    - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
  - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 3.2 All of the following:
    - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
    - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
  - 1.3 Trastuzumab not to be given in combination with lapatinib; and
  - 1.4 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

2.3 Disease has not progressed during previous treatment with trastuzumab.

 TRASTUZUMAB EMTANSINE − PCT only − Specialist − Special Authority see \$A2144 below

 Inj 100 mg vial
 2,320.00
 1
 ✓ Kadcyla

 Inj 160 mg vial
 3,712.00
 1
 ✓ Kadcyla

 Inj 1 mg for ECP
 24.52
 1 mg
 ✓ Baxter

#### ⇒SA2144 Special Authority for Subsidy

**Initial application** — **(early breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
  - 3.1 The patient has received prior therapy for metastatic disease\*; or
  - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy\*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
  - 5.1 Patient does not have symptomatic brain metastases; or
  - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Inj 90 mg per ml, 1 ml pre-filled syringe........................4,162.00 1 ✓ Stelara

#### ⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active Crohn's disease: and
  - 2.2 Either:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

**Initial application** — **(Crohn's disease - children\*)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active Crohn's disease: and
  - 2.2 Either:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
  - 2.1 Patient has active ulcerative colitis; and
  - 2.2 Fither:
    - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria: or
    - 2.2.2 Both:
      - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
      - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
  - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy\*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with \* is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

✓ Entvvio

## ⇒SA2183 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
  - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

assessed: and

2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (Crohn's disease - children\*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
  - 1.2 PCDAI score is 15 or less: or
  - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

**Initial application — (ulcerative colitis)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
  - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
  - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
  - 2.3 Patient's PUCAI score is greater than or equal to 20\*; and
- 3 Any of the following:
  - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
  - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
  - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with \* is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
  - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy \*; and

bsidy urer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with \* is an unapproved indication.

## Programmed Cell Death-1 (PD-1) Inhibitors

ATEZOLIZUMAB - PCT only - Specialist - Special Author	ority see SA2264 below		
Inj 60 mg per ml, 20 ml vial	9,503.00	1	<ul><li>Tecentriq</li></ul>
Inj 1 mg for ECP	8.08	1 mg	✓ Baxter

#### **⇒SA2264** Special Authority for Subsidy

**Initial application** — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment: or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

#### 

#### **⇒SA2164** Special Authority for Subsidy

**Initial application — (Non-small cell lung cancer)** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	<b>✓</b>	Manufacturer

continued...

- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment: and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Fither:
  - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
  - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

		NIVOLUMAB – PCT only – Specialist – Special Authority see SA2120 below	
Opdivo	1	Inj 10 mg per ml, 4 ml vial1,051.98	
✓ Opdivo	1	Inj 10 mg per ml, 10 ml vial2,629.96	
✓ Baxter	1 ma	Ini 1 mg for ECP27.62	

#### ⇒SA2120 Special Authority for Subsidy

**Initial application** only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV: and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded pembrolizumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
    - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or

 Subsidy Manufacturer's Price)	٤	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
- 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
  must have reduction in short axis to < 10 mm.</li>
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA2265 below

Inj 25 mg per ml, 4 ml vial	4,680.00	1	Keytruda
Inj 1 mg for ECP	47.74	1 mg	Baxter

#### ⇒SA2265 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
  - 4.1 Patient has not received funded nivolumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
    - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
    - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
  - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
  - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
  - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
  must have reduction in short axis to < 10 mm</li>
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Fither:
  - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

determined by a validated test unless not possible to ascertain; or

- 6.2 Both:
  - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
  - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2; and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (non-small cell lung cancer first-line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2; and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 Patient's disease has had a complete response to treatment; or
  - 1.2 Patient's disease has had a partial response to treatment; or
  - 1.3 Patient has stable disease: and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period: and

#### continued...

- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

## Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	✓ Neoral
Cap 100 mg	177.81	50	✓ Neoral
Oral liq 100 mg per ml		50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below -	Retail pharmacy		
Wastage claimable			
Tab 10 mg	6,512.29	30	<ul><li>Afinitor</li></ul>
Tab 5 mg	4,555.76	30	<ul><li>Afinitor</li></ul>
<u> </u>	*		

#### ⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

**Renewal** only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

#### SIROLIMUS - Special Authority see SA2270 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	Rapamune

#### ⇒SA2270 Special Authority for Subsidy

**Initial application** from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP: or
- · Leukoencepthalopathy: or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has severe non-malignant lymphovascular malformation\*; and
- 2 Any of the following:
  - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
  - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
  - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
  - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with \* are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex\*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

 $\textbf{Renewal} - \textbf{(renal angiomyolipoma(s) associated with tuberous sclerosis complex*)} \ \ \text{from any relevant practitioner}.$ 

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with \* are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
    - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
  - 2.2 Both:
    - 2.2.1 Vigabatrin is contraindicated; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with \* are unapproved indications

TACROLIMUS - Special Authority see SA2271 below - Retail pharmacy

Cap 0.5 mg	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	100	✓ Tacrolimus Sandoz
Cap 1 mg84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg248.20	50	✓ Tacrolimus Sandoz

#### ⇒SA2271 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications\*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Either:
  - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
  - 2.2 Patient is a child with nephrotic syndrome\*.

Note: Indications marked with \* are unapproved indications

#### **JAK** inhibitors

#### **⇒SA2079** Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:

	Subsidy	Ful	y Brand or
(Man	ufacturer's Price)	Subsidise	d Generic
	\$	Per •	Manufacturer

continued...

- 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or 3.2 Both:
  - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Te Whatu Ora Hospital; and 3.2.2 Either:
    - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
    - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

# **Antiallergy Preparations**

### Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 ini per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

Inj 0.15 mg per 0.3 ml auto-injector	90.00	1 OP	Epipen Jr
Inj 0.3 mg per 0.3 ml auto-injector	90.00	1 OP	Epipen

## ⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
  - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
- 2 Patient is not to be prescribed more than two devices in initial prescription.

#### ⇒SA1558 Special Authority for Subsidy

**Initial application** only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

## **Allergy Desensitisation**

## **⇒SA1367** Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

**Renewal** only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy Initiation kit - 5 yials freeze dried venom with diluent 305.00 1 OP ✓ VENOX S29 Maintenance kit - 1 vial freeze dried venom with diluent................305.00 1 OP ✓ VENOX S29 Maintenance kit - 6 vials 120 mcg freeze dried venom, with 1 OP ✓ Venomil S29 Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 1 OP ✓ Albev Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent ..... 305.00 1 OP ✓ Hymenoptera S29

	Subsidy	, 6 .	Fully	,	
	(Manufacturer's Price \$	) Sub Per	sidised •		
VASP VENOM ALLERGY TREATMENT - Special Authority see	SA1367 on the pre	vious pag	e – Reta	ail pharmacy	
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze					
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	<b>✓</b>	Albey	
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze					
dried venom, with diluent	305.00	1 OP	<b>✓</b>	lymenoptera S29	
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze					
dried venom, with diluent	305.00	1 OP	✓ \	enomil S29	
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze					
dried venom, with diluent	305.00	1 OP	<b>✓</b> F	lymenoptera S29	
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze					
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	<b>✓</b>	Albey	
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze		_	_		
dried venom, with diluent	305.00	1 OP	<b>✓</b> \	enomil \$29	
Antihistamines					
Antinistanines					
ETIRIZINE HYDROCHLORIDE					
★ Tab 10 mg		100	_	<u>Zista</u>	
F Oral liq 1 mg per ml	2.84	200 ml	<b>✓</b> F	<u>listaclear</u>	
CHLORPHENIRAMINE MALEATE					
Oral liq 2 mg per 5 ml	9.37	500 ml	<b>✓</b> F	listafen	
EXTROCHLORPHENIRAMINE MALEATE					
€ Tab 2 mg	2.02	40			
	(8.40)		F	Polaramine	
	1.01	20			
	(5.99)		F	Polaramine	
F Oral liq 2 mg per 5 ml		100 ml			
	(10.29)		F	Polaramine	
EXOFENADINE HYDROCHLORIDE					
₹ Tab 60 mg	4.34	20			
	(8.23)		T	Telfast	
F Tab 120 mg		10	_		
	(8.23)		ī	Telfast	
	14.22	30	-	T-164	
	(26.44)		ı	Telfast	
ORATADINE		100			
₭ Tab 10 mg		100		<u>-orafix</u>	
Gral liq 1 mg per ml	1.43	100 ml	<b>✓</b>	laylor syrup	
ROMETHAZINE HYDROCHLORIDE			_		
₹ Tab 10 mg		50	_	Allersoothe	
k Tab 25 mg		50	_	Allersoothe	
Koral liq 1 mg per 1 ml		100 ml	-	Allersoothe	

\* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO ....21.09

5

✓ Hospira

120 dose OP

60 dose OP

✓ Serevent ✓ Serevent Accuhaler

	Subsidy		Fully Brand or
	(Manufacturer's Pi	rice) Subsid	dised Generic
	\$	Per	✓ Manufacturer
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose		200 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free	8.54	200 dose OP	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose	17.52	200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free	12.50	200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	✓ Pulmicort
•			Turbuhaler
FLUTICASONE			
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose	8.61	60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
	-		
Inhaled Long-acting Beta-adrenoceptor Agonis	ts		
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dose	10.32	60 dose OP	
(equivalent to elormoteror familiarate o mog metered dose	(16.90)	00 003 <del>0</del> 01	Oxis Turbuhaler
INDAGATEROL	(10.30)		ONIS TUIDUITAICI
INDACATEROL			
Powder for inhalation 150 mcg		30 dose OP	<ul> <li>Onbrez Breezhaler</li> </ul>
Powder for inhalation 300 mcg	61.00	30 dose OP	<ul> <li>Onbrez Breezhaler</li> </ul>
SALMETEROL			

	Subsidy		Fully Brand or
	(Manufacturer's I	Price) Subsid	dised Generic
	\$	Per	✓ Manufacturer
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocept	or Agonists	
BUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide w	<i>i</i> ith		
6 mcg eformoterol fumarate metered dose)		120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumal		120 003 <del>0</del> 01	• Buonesp Spirollax
per dose (equivalent to 400 mcg budesonide with 12 mc	g		
eformoterol fumarate metered dose) - No more than 2	00.50	400 de OD	/ December of the many
dose per day		120 dose OP	✓ DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 n	ncg33.74	120 dose OP	✓ Symbicort
			Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 n	ncg33.74	120 dose OP	✓ Symbicort
			Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			
12 mcg - No more than 2 dose per day	33.74	60 dose OP	✓ Symbicort
			Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓ Breo Ellipta
	44.00	30 dose OF	• Bieo Ellipta
FLUTICASONE WITH SALMETEROL			
Aerosol inhaler 50 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - No			
more than 2 dose per day	33.74	60 dose OP	<ul> <li>Seretide Accuhaler</li> </ul>
Powder for inhalation 250 mcg with salmeterol 50 mcg - No			
more than 2 dose per day		60 dose OP	<ul> <li>Seretide Accuhaler</li> </ul>
, ,			
Beta-Adrenoceptor Agonists			
SALBUTAMOL			_
Oral liq 400 mcg per ml	40.00	150 ml	✓ <u>Ventolin</u>
Infusion 1 mg per ml, 5 ml	118.38	10	✓ Ventolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO	53.00	5	✓ Ventolin
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO	2.00	000 doos OD	✓ December
dose available on a PSO	3.80	200 dose OP	✓ Respigen
	(0.00)		✓ SalAir
	(6.20)		Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb			_
available on a PSO	8.96	20	✓ <u>Asthalin</u>
			✓ Ventolin
			Nebules S29
	51.11		✓ Accord \$29
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule - Up to 30 neb			
available on a PSO		20	✓ Asthalin
		-	

	RESPIRA	TORY SYSTE	M AND ALLERGIES
	Subsidy (Manufacturer's \$	Price) Subsi Per	Fully Brand or dised Generic  Manufacturer
TERBUTALINE SULPHATE Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated	22.20	120 dose OP	✓ Bricanyl Turbuhaler
Anticholinergic Agents			
IPRATROPIUM BROMIDE  Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose available on a PSO		200 daga OR	✓ Atrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne available on a PSO	b	200 dose OP 20	✓ Univent ✓ Accord S29
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic .	Agents	
SALBUTAMOL WITH IPRATROPIUM BROMIDE  Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p dose CFC-free  Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO	12.19	200 dose OP 20 60	✓ Duolin HFA  ✓ <u>Duolin</u> ✓ Duolin Respules \$29
<b>Long-Acting Muscarinic Antagonists</b>			
GLYCOPYRRONIUM — Subsidy by endorsement     a) Inhaled glycopyrronium treatment will not be subsidised if umeclidinium.     b) Glycopyrronium powder for inhalation 50 mcg per dose is having COPD using spirometry if spirometry is possible, a Powder for inhalation 50 mcg per dose	subsidised on and the prescri	ly for patients who	o have been diagnosed as
TIOTROPIUM BROMIDE — Subsidy by endorsement  a) Tiotropium treatment will not be subsidised if patient is als umeclidinium.  b) Tiotropium bromide is subsidised only for patients who ha spirometry is possible, and the prescription is endorsed a 1 October 2018 with a valid Special Authority are deemed Powder for inhalation, 18 mcg per dose	so receiving treative been diagnoccordingly. Part endorsed.	osed as having C	OPD using spirometry if
UMECLIDINIUM – Subsidy by endorsement     a) Umeclidinium will not be subsidised if patient is also recei tiotropium bromide.      A subsidised if patient is also receil to the control of the cont	ving treatment		nhaled glycopyrronium or

b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having

30 dose OP

✓ Incruse Ellipta

COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Powder for inhalation 62.5 mcg per dose......61.50

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

### ⇒SA1584 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

**Renewal** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication: and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL – Special Authority see SA Powder for Inhalation 50 mcg with indacaterol 110 mcg		armacy  ✓ Ultibro Breezhaler
TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see		
Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg		
UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 a	bove – Retail pharmacy	
Powder for inhalation 62.5 mcg with vilanterol 25 mcg	77.00 30 dose OP	Anoro Ellipta

#### **Antifibrotics**

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

#### ⇒SA2012 Special Authority for Subsidy

**Initial application — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Generic	
PIRFENIDONE – Retail pharmacy-Specialist – Special Authority Note: Pirfenidone is not subsidised in combination with subs					
Tab 801 mg	3,645.00	90	✓	Esbriet	
Tab 267 mg	1,215.00	90	✓	Esbriet	

⇒SA2013 Special Authority for Subsidy

**Initial application — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib; or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

## **Leukotriene Receptor Antagonists**

MONTELUKAST		
* Tab 4 mg3.10	28	<ul><li>Montelukast Mylan</li></ul>
		✓ Montelukast Viatris
* Tab 5 mg	28	✓ Montelukast Mylan
		✓ Montelukast Viatris
* Tab 10 mg	28	✓ Montelukast Mylan
v		✓ Montelukast Viatris
(Montelukast Mylan Tab 4 mg to be delisted 1 February 2024)		

(Montelukast Mylan Tab 4 mg to be delisted 1 February 2024) (Montelukast Mylan Tab 5 mg to be delisted 1 January 2024) (Montelukast Mylan Tab 10 mg to be delisted 1 February 2024)

## Methylxanthines

AMI	NOP	HYL	LINE
-----	-----	-----	------

*	Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a			
	PSO	180.00	5	✓ DBL Aminophylline
TH	IEOPHYLLINE			
*	Tab long-acting 250 mg	23.94	100	✓ Nuelin-SR
	Oral lig 80 mg per 15 ml		500 ml	✓ Nuelin

### **Mucolytics**

DORNASE ALFA - Special Authority see SA1978 on the ne	ext page - Retail pharmac	/	
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	/	Manufacturer

### ⇒SA1978 Special Authority for Subsidy

**Initial application — (cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
  - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
  - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25: or
  - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

**Renewal** — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2196 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg

Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg

(56) and ivacaftor 150 mg (28) .......27,647.39 84 OP **✓ Trikafta** 

### ⇒SA2196 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Either:
  - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele): or
  - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Either:
  - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
  - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

#### Note:

 a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda\_docs/label/2021/212273s004lbl.pdf.

IVACAFTOR - PCT only - Specialist - Special Authori	ty see SA2017 below		
Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

### ⇒SA2017 Special Authority for Subsidy

**Initial application** only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
continued			
All of the following:			
<ul><li>1 Patient has been diagnosed with cystic fibrosis; and</li><li>2 Fither:</li></ul>			
	a fibracia transmambrana		ulator (CETD) gana an a
2.1 Patient must have G551D mutation in the cysti least 1 allele: or	c librosis transmembrane	conductance reg	julator (CFTR) gene on a
2.2 Patient must have other gating (class III) mutat	tion (G1244F, G1349D, G	178B. G551S. S	1251N, S1255P, S549N
and S549R) in the CFTR gene on at least 1 all		7011, 00010, 0	120111, 012001 , 001011
3 Patients must have a sweat chloride value of at least (	•	oilocarpine ionto	phoresis or by Macroduc
sweat collection system; and			,
4 Treatment with ivacaftor must be given concomitantly			
5 Patient must not have an acute upper or lower respira	tory infection, pulmonary e		
a contract the contract of the	and the second second		
(including antibiotics) for pulmonary disease in the las		cing treatment v	vith ivacaftor; and
(including antibiotics) for pulmonary disease in the las 6 The dose of ivacaftor will not exceed one tablet or one	e sachet twice daily; and	cing treatment v	vith ivacaftor; and
(including antibiotics) for pulmonary disease in the las  6 The dose of ivacaftor will not exceed one tablet or one  7 Applicant has experience and expertise in the manage	e sachet twice daily; and	cing treatment v	vith ivacaftor; and
(including antibiotics) for pulmonary disease in the las 6 The dose of ivacaftor will not exceed one tablet or one	e sachet twice daily; and	cing treatment v	vith ivacaftor; and

ivasai i	reparations
Allergy	Prophylactics

BUDESONIDE  Metered aqueous nasal spray, 50 mcg per dose		✓ SteroClear ✓ SteroClear
FLUTICASONE PROPIONATE  Metered aqueous nasal spray, 50 mcg per dose1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%5.23	15 ml OP	✓ Univent

# **Respiratory Devices**

	SPACER	

- a) Up to 50 dev available on a PSO
- b) Only on a PSO
- c) Only for children aged six years and under

### PEAK FLOW METER

- a) Up to 25 dev available on a PSO
- b) Only on a PSO

✓ Mini-Wright AFS Low Range

✓ Mini-Wright Standard

1

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	3.65	1	<b>√</b> e	-chamber Turbo
510 ml (single patient)	5.95	1	<b>√</b> e	-chamber La Grande
800 ml	6.50	1	✓ \	olumatic

CAFFEINE (	CITRATE
------------	---------

Oral liq 20 mg per ml (10 mg base per ml)......15.10 25 ml OP ✓ Biomed

	Subsidy (Manufacturer's Pr \$	rice) Su Per	Fully bsidised	Brand or Generic Manufacturer
Ear Preparations				
FLUMETASONE PIVALATE	4.46	7.5 ml OP	./	Locacorten-Viaform
Ear drops 0.02% with clioquinol 1%	4.40	7.5 IIII OP		ED's
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI	IN AND NYSTAT	IN	•	Locorten-Vioform
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓	Kenacomb
Ear/Eye Preparations				
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN				
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml		8 ml OP		
	(9.27) (9.27)			Otodex <sup>S29</sup> Sofradex
FRAMYCETIN SULPHATE Ear/Eye drops 0.5%	4.13	8 ml OP		
	(8.65)	· · · · · · ·	;	Soframycin
Eye Preparations				
Eye preparations are only funded for use in the eye, unless explicit	citly stated otherw	rise.		
Anti-Infective Preparations				
ACICLOVIR  * Eye oint 3%	14.88	4.5 g OP	<b>✓</b> 1	ViruPOS
CHLORAMPHENICOL			-	
Eye oint 1%	1.45	5 g OP 10 ml OP	-	<u>Devatis</u> Chlorsig
Funded for use in the ear*. Indications marked with * arc CIPROFLOXACIN	e unapproved ind	ications.		
Eye drops 0.3% - Subsidy by endorsement		5 ml OP	-	Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis of for the second line treatment of chronic suppurative otitis.  Note: Indication marked with a * is an unapproved indication with a * is an unapproved indication with a * is an unappro	s media (CSOM)*			
PROPAMIDINE ISETHIONATE  * Eye drops 0.1%	2.07	10 ml OP		
* Lye urops 0.1%	(14.55)	IU IIII OP	I	Brolene
SODIUM FUSIDATE [FUSIDIC ACID] Eye drops 1%	5.29	5 g OP	<b>√</b>	Fucithalmic
TOBRAMYCIN		ŭ		T-1
Eye oint 0.3% Eye drops 0.3%		3.5 g OP 5 ml OP		Tobrex Tobrex



Subsidy	Fι	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
 \$	Per	•	Manufacturer

## **Corticosteroids and Other Anti-Inflammatory Preparations**

DE	XAMETHASONE			
*	Eye oint 0.1%	5.86	3.5 g OP	✓ Maxidex
*	Eye drops 0.1%	4.50	5 ml OP	✓ Maxidex
	Ocular implant 700 mcg - Special Authority see SA1680 below			
	- Retail pharmacy1,4	144.50	1	<ul><li>Ozurdex</li></ul>

### **⇒SA1680** Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
  - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

#### DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin		· ·	
b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM			
Eye drops 0.1%	8.80	5 ml OP	✓ Voltaren Ophtha
FLUOROMETHOLONE			
* Eve drops 0.1%	3.09	5 ml OP	✓ FML
•	5.20		✓ Flucon

	Subsidy		Fully	Brand or
	(Manufacturer's P	rice) Subs	sidised	Generic
	\$	Per	✓	Manufacturer
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
,	(10.34)		L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	<b>√</b> L	omide
NEPAFENAC				
Eye drops 0.3%	8.80	3 ml OP	<b>✓</b>	evro
PREDNISOLONE ACETATE				
Eve drops 1%	6.92	10 ml OP	<b>√</b> P	rednisolone-AFT
_,0 0.000 //2	7.00	5 ml OP	<b>✓</b> P	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority so	ee SA1715 below	– Retail pharr	nacy	
Eye drops 0.5%, single dose (preservative free)		20 dose	•	linims Prednisolone

### ⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

### SODIUM CROMOGLICATE

### **Glaucoma Preparations - Beta Blockers**

BETAXOLOL

\* Eve drops 0.25%

	Eye drops 0.5%		✓ Betoptic
	MOLOL	• • .	
*	Eye drops 0.25%2.42	5 ml OP	✓ Arrow-Timolol
*	Eye drops 0.5%	5 ml OP	✓ Arrow-Timolol
*	Eye drops 0.5%, gel forming – Subsidy by endorsement	2.5 ml OP	✓ Timoptol XE

Subsidised for patients who were taking timolol eye drops 0.5%, gel forming prior to 1 April 2023 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of timolol eye drops 0.5%, gel forming.

11 80

5 ml OP

✓ Retentic S

(Timoptol XE Eye drops 0.5%, gel forming to be delisted 1 March 2024)

## **Glaucoma Preparations - Carbonic Anhydrase Inhibitors**

ACETAZOLAMIDE  * Tab 250 mg	17.03	100	✓ Diamox
BRINZOLAMIDE			2.0
* Eye drops 1%	7.30	5 ml OP	✓ Azopt

#### DORZOLAMIDE HYDROCHLORIDE - Subsidy by endorsement

Subsidised for patients who were taking dorzolamide hydrochloride eye drops 2% prior to 1 April 2023 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of dorzolamide hydrochloride eye drops 2%.

(Trusopt Eye drops 2% to be delisted 1 March 2024)

	Subsidy (Manufacturer's I \$	Price) Subs	Fully Brand or sidised Generic  ✓ Manufacturer
DORZOLAMIDE WITH TIMOLOL  * Eye drops 2% with timolol 0.5%	2.73	5 ml OP	✓ <u>Dortimopt</u>
Glaucoma Preparations - Prostaglandin Analo	gues		
BIMATOPROST  * Eye drops 0.03%	5.95	3 ml OP	✓ <u>Bimatoprost</u> <u>Multichem</u>
LATANOPROST  * Eye drops 0.005% TRAVOPROST	1.82	2.5 ml OP	✓ <u>Teva</u>
* Eye drops 0.004%	9.75	2.5 ml OP	✓ <u>Travatan</u>
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE  * Eye drops 0.2%BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE	4.29	5 ml OP	✓ <u>Arrow-Brimonidine</u>
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combigan
LATANOPROST WITH TIMOLOL  Eye drops 0.005% with timolol 0.5%  PII OCARPINE HYDROCHI ORIDE	4.95	2.5 ml OP	✓ Arrow - Lattim
# Eye drops 1%  # Eye drops 2%  * Eye drops 4%  Subsidised for oral use pursuant to the Standard Form	5.35 7.99	15 ml OP 15 ml OP 15 ml OP	<ul><li>✓ Isopto Carpine</li><li>✓ Isopto Carpine</li><li>✓ Isopto Carpine</li></ul>
PILOCARPINE NITRATE  * Eye drops 2% single dose – Special Authority see SA0898 below – Retail pharmacy		20 dose	✓ Minims Pilocarpine

## **⇒SA0895** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## **Mydriatics and Cycloplegics**

ATROPINE SULPHATE			
* Eye drops 1%	18.27	15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE			
* Eye drops 1%	8.76	15 ml OP	✓ Cyclogyl
* Eye drops 1%, single dose (preservative free) - Only on a			
prescription	84.85	20 dose	✓ Minims
			Cyclopentolate
TROPICAMIDE			
* Eye drops 0.5%	7.15	15 ml OP	✓ Mydriacyl
* Eye drops 1%	8.66	15 ml OP	✓ Mydriacyl

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	/	Manufacturer	

## **Preparations for Tear Deficiency**

For acetylcysteine eye drops refer Standard Formulae, page 266

**HYPROMELLOSE** 

HYPROMELLOSE WITH DEXTRAN

### **Preservative Free Ocular Lubricants**

#### ⇒SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
  - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
  - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

**Renewal** from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA2134 above – Retail	pharmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL	- Special Authority see	SA2134 abo	ve – Retail pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml	10.78	30	✓ Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special A	uthority see SA2134 abo	ove – Retail p	harmacy
Eye drops 1 mg per ml	13.85 10	ml OP •	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The I			
month is not relevant and therefore only the prescribe	d dosage to the nearest	OP may be o	claimed.

## Other Eve Preparations

NAPHAZOLINE HYDROCHLORIDE  * Eye drops 0.1%4.15	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%2.17	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT  * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE  Eye oint 138 mcg per g3.80	5 g OP	✓ VitA-POS

		Subsidy (Manufacturer's Pric	ce) Per	Fully Subsidised	
٧	arious				
PH	ARMACY SERVICES				
*	a) May only be claimed once per patient. b) The Pharmacode for BSF Midodrine Medsurge is 266		1 fee	✓	BSF Concerta BSF Midodrine Medsurge BSF Rubifen SR
	c) The Pharmacode for BSF Rubifen SR is 2665956 - se	ee also page 147			
*	d) The Pharmacode for BSF Concerta is 2665948 - see COVID-19 Services	1 0	1 fee	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	After Hours Med Mgmt 15 min After Hours Med Mgmt 30 min After Hours Med Mgmt 45 min Antivirals Eligibility Review Compliance Packaging Med Mgmt 15 min Med Mgmt 30 min Med Mgmt 45 min
*	Immunisation administration fee	0.00	1 fee		Medicine Delivery Immunisation
*	Immunisation co-administration fee	0.00	1 fee	•	Administration Immunisation Co-administration
(BS	SF Concerta Brand switch fee to be delisted 31 December 2020 SF Midodrine Medsurge Brand switch fee to be delisted 1 Nove SF Rubifen SR Brand switch fee to be delisted 31 December 20	mber 2023)			
A	gents Used in the Treatment of Poisonings				
A	ntidotes				
NA	ETYLCYSTEINE Inj 200 mg per ml, 10 ml ampoule LOXONE HYDROCHLORIDE a) Up to 10 inj available on a PSO b) Only on a PSO		10	•	Martindale Pharma
*	Inj 400 mcg per ml, 1 ml ampoule	35.26	10	✓	<u>HameIn</u>
R	emoval and Elimination				
	ARCOAL Oral liq 50 g per 250 ml a) Up to 250 ml available on a PSO b) Only on a PSO	43.50	250 ml (	OP 🗸	Carbosorb-X

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
DEFERASIROX – Special Authority see SA1492 below – Retail p Wastage claimable	harmacy			
Tab 125 mg dispersible	276.00	28	1	Exjade
Tab 250 mg dispersible	552.00	28	1	Exjade
Tab 500 mg dispersible	1,105.00	28	•	Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
  - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2\*; or
  - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
  - 3.3 Treatment with deferiprone has resulted in arthritis; or
  - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels.

PRONE – Special Authority see SA1480 below – Retail pharmacy	
500 mg533.17 100	✓ Ferriprox
liq 100 mg per 1 ml	OP <b>✓ Ferriprox</b>

#### ⇒SA1480 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

# Inj 500 mg vial	151.31	10	✓ DBL  Desferrioxamine  Mesylate for Inj  BP  ✓ Deferoxamine Pfizer  \$29 \$29
SODIUM CALCIUM EDETATE  * Inj 200 mg per ml, 5 ml	53.31	6	
1 44 314 74	(156.71)		Calcium Disodium Versenate



## **Standard Formulae**

Otaridard i Orinidiae			
ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol Preservative Water	60 mg 40 ml qs to 100 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml) Phenobarbitone Sodium Glycerol BP Water	400 mg 4 ml to 40 ml
CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative Water	300 mg 40 ml qs to 100 ml	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative Water	qs qs to 500 ml
FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Water (Preservative should be used if quantity supplied is f than 5 days. Maximum 500 ml per prescription.)	1 tab qs to 500 ml for more	(Preservative should be used if quantity supplied is than 5 days.)  SALIVA SUBSTITUTE FORMULA Methylcellulose Preservative Water	5 g qs to 500 ml
METHADONE MIXTURE Methadone powder Glycerol Water	qs qs to 100 ml	(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.) SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml Water	qs qs
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqu	10 g to 100 ml id mixture)	(Only funded if prescribed for treatment of hyponatra VANCOMYCIN ORAL SOLUTION (25 mg per ml) Vancomycin 500 mg injection Glycerin with sucrose suspension	
OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml	Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	to 100 ml

### EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Series Per Manufacturer

	`	\$	Per
Extemporaneously Compounded Preparations	and (	Galenica	ls

(90.09)

9

Douglas

Only in extemporaneously compounded codeine linctus.

#### COLLODION FLEXIBLE

Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.

COMPOUND HYDROXYBENZOATE - Only in combination

Only in extemporaneously compounded oral mixtures.

Soln .......30.00

100 ml ✓ Midwest

GLYCERIN WITH SODIUM SACCHARIN - Only in combination

Only in combination with Ora-Plus or when used in the vancomycin oral Iquuid Standard Formulae.

GLYCERIN WITH SUCROSE - Only in combination

Only in combination with Ora-Plus or when used in the vancomycin oral Iguuid Standard Formulae.

**GLYCEROL** 

#### METHADONE HYDROCHLORIDE

- a) Only on a controlled drug form
- b) No patient co-payment payable
- c) Safety medicine; prescriber may determine dispensing frequency
- d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).

Powder	7.84	1 g	✓ AFT
METHYL HYDROXYBENZOATE	0.00	05 ~	✓ Midwood
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE			
Powder		100 g	✓ MidWest
Suspension – Only in combination	30.95	473 ml	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCI	HARIN - Only in c	ombination	
Suspension	30.95	473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - O	nly in combination		
Suspension	30.95	473 ml	Ora-Blend
PHENOBARBITONE SODIUM			
Powder - Only in combination	52.50	10 g	✓ MidWest
•	325.00	100 g	✓ MidWest
Only in children up to 12 years		ŭ	
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxyben	zoate 10% solution	n.	
Liq	11.25	500 ml	✓ Midwest
SODIUM BICARBONATE			
Powder BP - Only in combination	10.05	500 g	✓ Midwest

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

## EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price)	Sul Per	Fully osidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatio		500 ml	<b>✓</b> M	lidwest	
WATER Tap - Only in combination	0.00	1 ml	<b>✓</b> Ta	ap water	

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

## **Nutrient Modules**

### Carbohydrate

### ⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

## Carbohydrate And Fat

### **⇒SA1376** Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...



Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	bsidised	Generic
\$	Per	✓	Manufacturer

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 cancer in children; or
  - 2.2 faltering growth; or
  - 2.3 bronchopulmonary dysplasia; or
  - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Powder (neutral)	60.31	I 400 g OP	Duocal Super	
			Soluble Pow	dor

#### Fat

#### **⇒SA2204** Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA2204 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)	200 ml OP	✓ Calogen
30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)12.30	200 ml OP	✓ Calogen
Oil	500 ml OP	✓ MCT oil (Nutricia)
MCT Emulsion, 250 ml114.92	4 OP	✓ Liquigen

#### **Protein**

### ⇒SA1524 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	armacy [HP3]	PROTEIN SUPPLEMENT – Special Authority see SA1524 above – Hospital pha
✓ Protifar	225 g OP	Powder
✓ Resource	227 g OP	8.95
Beneprotein	•	

Subsidy (Manufacturer's Price) \$ Fully Subsidised Brand or Generic Manufacturer

## **Oral and Enteral Feeds**

#### **Diabetic Products**

### ⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see	e SA1095 above -	<ul> <li>Hospital pharm</li> </ul>	nacy [HP3]
Liquid	3.75	500 ml OP	✓ Glucerna Select
·	7.50	1,000 ml OP	✓ Nutrison Advanced Diason
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA	1095 above – Ho	spital pharmacy	
Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	2.10		✓ Nutren Diabetes

### **Fat Modified Products**

#### ⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA2205 above - Ho	ospital pharmac	y [HP3]	
Powder	60.48	400 g OP	✓ Monogen

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

## **Paediatric Products For Children Awaiting Liver Transplant**

### ⇒SA1098 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

### Paediatric Products For Children With Chronic Renal Failure

### ⇒SA1099 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

### **Paediatric Products**

### ⇒SA1379 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 any condition causing malabsorption; or
  - 2.3 faltering growth in an infant/child: or
  - 2.4 increased nutritional requirements; or
  - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

## **SPECIAL FOODS**

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	•	Manufacturer

continued...

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

F			
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority see S Liquid		the previous pag 500 ml OP	ge – Hospital pharmacy [HP3]  ✓ Nutrini Energy RTH  ✓ Frebini Energy
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA Liquid		e previous page 500 ml OP	<ul><li>− Hospital pharmacy [HP3]</li><li>✓ Nutrini RTH</li><li>✓ Pediasure RTH</li></ul>
	6.50		✓ Frebini Original
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special A pharmacy [HP3]	uthority se	e SA1379 on the	e previous page – Hospital
Liquid	6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
	7.00		✓ Frebini Energy Fibre
PAEDIATRIC ENTERAL FEED WITH FIBRE 1KCAL/ML - Special Aut pharmacy [HP3]	hority see	SA1379 on the p	previous page – Hospital
Liquid	7.00	500 ml OP	✓ Frebini Original Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA13	79 on the p	orevious page -	Hospital pharmacy [HP3]
Liquid (strawberry)	1.60	200 ml OP	✓ Fortini
Liquid (vanilla)	1.60	200 ml OP	✓ Fortini
	6.99	500 ml OP	✓ Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379	on the pre	evious page – H	ospital pharmacy [HP3]
Liquid (chocolate)		200 ml OP	✓ Pediasure
Liquid (strawberry)	1.07	200 ml OP	✓ Pediasure
Liquid (vanilla)	1.07	200 ml OP	✓ Pediasure
. , ,	1.34	250 ml OP	✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Author pharmacy (HP3)	rity see SA	1379 on the pre	evious page – Hospital
Liquid (unflavoured)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)		200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)		200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the	previous i	page – Hospital	pharmacy [HP3]
Powder		400 g OP	✓ Peptamen Junior
		-	

## **Renal Products**

### ⇒SA1101 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority s Liquid			Hospital pharmacy [HP3]  Nepro HP RTH
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see S Liquid			
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA	1101 on the previou	s page – Hosp	ital pharmacy [HP3]
Liquid, 200 ml bottle	11.52	4 OP	
	(13.24)		NovaSource Renal
Liquid (apricot) 125 ml	11.52	4 OP	✓ Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	✓ Renilon 7.5

### Specialised And Elemental Products

### ⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 vear for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome: or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Liquid (caramel) 125 ml.......11.52

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - Special Authority see \$A1377 above - Hospital pharmacy [HP3] 1.000 ml OP ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see SA1377 above - Hospital pharmacy [HP3] Liquid (grapefruit), 250 ml carton.......171.00 18 OP ✓ Elemental 028 Extra ✓ Elemental 028 Extra 18 OP Liquid (pineapple & orange), 250 ml carton......171.00 18 OP ✓ Elemental 028 Extra

	Subsidy (Manufacturer's	Price) Subsi Per	Fully dised	Brand or Generic Manufacturer
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see S Powder (unflavoured)	•	revious page – F 80 g OP		ıl pharmacy [HP3] 'ivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Auth [HP3]	ority see SA137	7 on the previou	ıs page	e – Hospital pharmacy
Liquid	9.60 12.04	500 ml OP 1.000 ml OP	_	Survimed OPD Jutrison Advanced
	12.04	1,000 1111 01	• 14	Peptisorb

### Paediatric Products For Children With Low Energy Requirements

### ⇒SA1196 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

## Standard Supplements

#### ⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
  - 2.1 The patient has a condition causing malabsorption; or
  - 2.2 The patient has failure to thrive; or
  - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

**Renewal — (Adults)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m<sup>2</sup> and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant: and
  - 5.2 Any of the following:

continued...

Subsidy (Manufacturer's Price)	S	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

- 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
- 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
- 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease; or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm<sup>3</sup>); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube - refer to specific medical condition criteria); or

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	1	Manufacturer	

- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

9 Severe chronic neurological conditions.			
ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page	ie 276 – Ho	spital pharmacy	/ [HP3]
Liquid	1.75	250 ml OP	✓ Ensure Plus HN
·	7.00	1,000 ml OP	✓ Ensure Plus RTH
			✓ Nutrison Energy
	9.60		✓ Fresubin HP Energy
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on page	276 - Host	oital pharmacy [	HP31
Liquid		250 ml OP	✓ Isosource Standard
T 1	5.29	1,000 ml OP	✓ Nutrison Standard RTH
			✓ Osmolite RTH
	6.50		✓ Fresubin Original
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authority see			
Liquid	5.29	1,000 ml OP	✓ Nutrison
			800 Complete
			Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority see SA			
Liquid	5.29	1,000 ml OP	✓ Jevity RTH
	7.00		✓ Nutrison Multi Fibre
	7.00		✓ Fresubin Original
			Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority see S			
Liquid		1,000 ml OP	✓ Jevity Plus
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see S			
Liquid	7.00	1,000 ml OP	✓ Jevity HiCal RTH
			✓ Nutrison Energy
			Multi Fibre
	9.80		✓ Fresubin HP Energy Fibre
ENTERAL FEED WITH PROTEIN 1.2KCAL/ML - Special Authority se	e SA1859 d	on nage 276 – F	Hospital pharmacy [HP3]
Liquid		500 ml OP	✓ Fresubin Intensive
ORAL FEED (POWDER) - Special Authority see SA1859 on page 276	– Hospital	pharmacy [HP:	31
Powder (chocolate)		840 g OP	✓ Sustagen Hospital
·		ŭ	Formula
	26.00	850 g OP	✓ Ensure
Powder (vanilla)	14.00	840 g OP	<ul> <li>Sustagen Hospital</li> </ul>
		-	Formula Active
	26.00	850 g OP	✓ Ensure

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

#### ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 276 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	Frauma Divia
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with	(1.20)		1 Ortioip
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 ml			
with Endorsement		200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement		237 ml OP	
	(1.33)	000 100	Ensure Plus
	0.72	200 ml OP	- 5
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 276 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisin Multi Fibre

## **High Calorie Products**

### ⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 any condition causing malabsorption; or
  - 1.2 faltering growth in an infant/child; or
  - 1.3 increased nutritional requirements: or
  - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

**Renewal — (Cystic fibrosis)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
  - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Authority see SA	1195 on the previous pa	ige – Hospital p	harmacy [HP3]
Liquid	5.50	500 ml OP	✓ Nutrison
			Concentrated
	6.50		✓ Fresubin 2kcal HP
	11.00	1,000 ml OP	<ul><li>Ensure Two Cal HN</li></ul>
			RTH

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

### **Food Thickeners**

#### ⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

continued...

## SPECIAL FOODS

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per		Manufacturer

continued...

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

### **Gluten Free Foods**

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

### ⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX - Special Authority see SA1729 above - Hospital pharmacy [HP3]

riospitai priarriacy [rir o]	
2.81 1,000 g OP	
(5.15)	Healtheries Simple Baking Mix
- Hospital pharmacy [HP3]	
3.93 1,000 g OP	
(7.32)	NZB Low Gluten
	Bread Mix
3.51	
(10.87)	Horleys Bread Mix
5.62 2,000 g OP	
(18.10)	Horleys Flour
	(5.15)  - Hospital pharmacy [HP3]3.93 1,000 g OP (7.32)  3.51 (10.87) spital pharmacy [HP3]5.62 2,000 g OP

	Subsidy		Fully	Brand or
	(Manufacturer's Pri		idised	Generic
	\$	Per	•	Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page – H	ospital pharm	acy [HF	P3]
Buckwheat Spirals		250 g OP		
	(3.11)	•	0	)rgran
Corn and Vegetable Shells	2.00 <sup>°</sup>	250 g OP		·
·	(2.92)	· ·	0	)rgran
Corn and Vegetable Spirals	2.00 <sup>°</sup>	250 g OP		·
	(2.92)	· ·	0	)rgran
Rice and Corn Lasagne Sheets	1.60 <sup>°</sup>	200 g OP		·
·	(3.82)	· ·	0	)rgran
Rice and Corn Macaroni		250 g OP		ŭ
	(2.92)	Ü	0	)rgran
Rice and Corn Penne	, ,	250 g OP		ŭ
	(2.92)	· ·	0	)rgran
Rice and Maize Pasta Spirals	2.00 <sup>°</sup>	250 g OP		·
·	(2.92)	· ·	0	)rgran
Rice and Millet Spirals	2.00 <sup>°</sup>	250 g OP		·
•	(3.11)	· ·	0	)rgran
Rice and corn spaghetti noodles	2.00 <sup>°</sup>	375 g OP		·
. •	(2.92)	· ·	0	)rgran
Vegetable and Rice Spirals	2.00 <sup>°</sup>	250 g OP		·
	(2.92)	· ·	0	)rgran
Italian long style spaghetti	2.00	220 g OP		•
	(3.11)	ŭ	0	)rgran
	. ,			-

## Foods And Supplements For Inborn Errors Of Metabolism

### **⇒SA1108** Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

## **Supplements For Homocystinuria**

## **Supplements For MSUD**

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

## **Supplements For PKU**

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

amacy [m-3]	
	hlexy 10
	KU Lophlex Powder
	KU Anamix Junior
	Chocolate
· , ,	KU Lophlex
	Powder
	KU Anamix Junior
	KU Lophlex Powder
( 0 / 0	KU Anamix Junior Orange
( , 0	KU Anamix Junior Vanilla
Infant formula	KU Anamix Infant
Powder (orange)	P Maxamum
` ' ' ' '	P Maxamum
	KU Anamix Junior
	LQ
Liquid (orange)	KU Anamix Junior
	LQ
Liquid (unflavoured)	KU Anamix Junior
	LQ
Liquid (forest berries), 250 ml carton	asiphen Liquid
Liquid (juicy tropical) 125 ml	KU Lophlex LQ 20
Oral semi-solid (berries) 109 g	KU Lophlex
	Sensation 20
Liquid (juicy berries) 62.5 ml	KU Lophlex LQ 10
	KU Lophlex LQ 10
	KU Lophlex LQ 10
	KU Lophlex LQ 20
	KU Lophlex LQ 20

### **Foods**

LOW PROTEIN BAKING MIX — Special Authority see SA1108 on the previous page — Hospital pharmacy [HP3]
Powder .......8.22 500 g OP ✓ Loprofin Mix

LOW PROTEIN PASTA - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne	5.95	250 g OP	✓ Loprofin
Low protein rice pasta	11.91	500 g OP	✓ Loprofin
Macaroni		250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals	11.91	500 a OP	✓ Loprofin

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

## Infant Formulae

## For Williams Syndrome

### ⇒SA1110 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder .......44.40 400 g OP ✓ Locasol

### **Gastrointestinal and Other Malabsorptive Problems**

AMINO ACID FORMULA - Special Authority see SA2092 below - Hospital phar	macy [HP3]	
Powder	400 g OP	<ul><li>✓ Alfamino</li><li>✓ Alfamino Junior</li></ul>
Powder (unflavoured)	400 g OP	✓ Elecare ✓ Elecare LCP ✓ Neocate Gold ✓ Neocate Junior Unflavoured
Powder (vanilla)53.00	400 g OP	<ul><li>✓ Neocate SYNEO</li><li>✓ Elecare</li><li>✓ Neocate Junior</li><li>✓ Vanilla</li></ul>

#### ⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis: or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency: or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
  - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
  - 6.2 Either:
    - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
      - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has IgE mediated allergy; and
  - 1.2 All of the following:
    - 1.2.1 Patient remains allergic to cow's milk; and
    - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
    - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 1.2.4 Amino acid formula is required for a nutritional deficit; and
    - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
  - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
  - 2.2 All of the following:
    - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
    - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 2.2.3 Amino acid formula is required for a nutritional deficit; and
    - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
      - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA - Special Auth	nority see SA1953 below -	Hospital pharr	nacy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

### ⇒SA1953 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
  - 2.1 Severe malabsorption; or
  - 2.2 Short bowel syndrome; or
  - 2.3 Intractable diarrhoea: or
  - 2.4 Biliary atresia; or
  - 2.5 Cholestatic liver diseases causing malabsorption; or
  - 2.6 Cystic fibrosis; or
  - 2.7 Proven fat malabsorption; or
  - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
  - 2.9 Intestinal failure: or
  - 2.10 Both:
    - 2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

## SPECIAL FOODS

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	<b>✓</b>	Manufacturer

continued...

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
  - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
  - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

### ⇒SA1557 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
- 1.2 Either:
  - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms: or
  - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption: or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea: or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
  - 11.1 For step down from Amino Acid Formula; and
  - 11.2 The infant is currently receiving funded amino acid formula; and
  - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
  - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsid		
(Manufacturer	's Price) Subsidise	I Generic
\$	Per 💌	Manufacturer

continued...

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

### Fluid Restricted

## ⇒SA1698 Special Authority for Subsidy

**Initial application** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

## **Ketogenic Diet**

### ⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

**Renewal** only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

 Powder (unflavoured)
 35.50
 300 g OP
 KetoCal 4:1

 V Ketocal 3:1
 KetoCal 4:1

 Powder (vanilla)
 35.50
 300 g OP
 KetoCal 4:1

✓ fully subsidised 289

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

# Other Supplements for PKU

٩N	INO ACID FORMULA WITHOUT PHENYLALANINE	- Special Authority see SA2	229 below	v – Hospital pharmacy [HP3]
	Powder (Banana) 35 g sachets	930.00	30	✓ PKU
				sphere20 Banana
	Powder (Chocolate) 32 g Sachets	898.56	30	✓ PKU Build
				20 Chocolate
	Powder (Chocolate) 35 g sachets	930.00	30	✓ PKU
				sphere20 Chocolate
	Powder (Lemon) 35 g sachets	930.00	30	✓ PKU
	· · · ·			sphere20 Lemon
	Powder (Lemonade) 33.4 g sachets	936.00	30	✓ PKU GMPro Ultra
				Lemonade
	Powder (Raspberry Lemonade) 32 g Sachets	898.56	30	✓ PKU Build
	, , , , ,			20 Raspberry
				Lemonade
	Powder (Smooth) 32 g Sachets	898.56	30	✓ PKU Build
				20 Smooth
	Powder (Vanilla) 32 g Sachets	898.56	30	PKU Build 20 Vanilla
	Powder (Red Berry) 35 g sachets		30	✓ PKU sphere20 Red
				Berry
	Powder (Vanilla) 35 g sachets	930.00	30	✓ PKU
	· -			sphere20 Vanilla

(PKU Build 20 Chocolate Powder (Chocolate) 32 g Sachets to be delisted 1 January 2024)
(PKU sphere20 Chocolate Powder (Chocolate) 35 g sachets to be delisted 1 January 2024)
(PKU sphere20 Lemon Powder (Lemon) 35 g sachets to be delisted 1 January 2024)
(PKU GMPro Ultra Lemonade Powder (Lemonade) 33.4 g sachets to be delisted 1 December 2023)
(PKU Build 20 Raspberry Lemonade Powder (Raspberry Lemonade) 32 g Sachets to be delisted 1 January 2024)
(PKU Build 20 Smooth Powder (Smooth) 32 g Sachets to be delisted 1 January 2024)
(PKU Build 20 Vanilla Powder (Vanilla) 32 g Sachets to be delisted 1 January 2024)
(PKU sphere20 Red Berry Powder (Red Berry) 35 g sachets to be delisted 1 January 2024)

(PKU sphere20 Banana Powder (Banana) 35 g sachets to be delisted 1 January 2024)

(PKU sphere20 Vanilla Powder (Vanilla) 35 g sachets to be delisted 1 January 2024)

#### ⇒SA2229 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient was previously receiving, or would receive PKU Lophlex Sensation Berries under (SA1108); and
- 2 PKU Sensation Berries is unable to be sourced at this time; and
- 3 Patient has trialled the currently funded PKU Lophlex products and these were not tolerated.

Note: These criteria are attached to short term funding to cover an out-of-stock situation on PKU Sensation Berries supplied by Nutricia

### **SECTION I: NATIONAL IMMUNISATION SCHEDULE**

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

# **Vaccinations**

#### BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.
- Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent.......................0.00

10 🗸 I

✓ BCG Vaccine

#### DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
  - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
  - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
  - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
  - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
  - 5) A single dose for vaccination of patients aged from 65 years old; or
  - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
  - 7) For vaccination of previously unimmunised or partially immunised patients; or
  - 8) For revaccination following immunosuppression; or
  - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.
- Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid. 8 mcg pertussis filamentous

10

✓ Boostrix

	Subsidy	Fully	
	(Manufacturer's Price) \$	Subsidised Per 🗸	d Generic Manufacturer
NIDUTHEDIA TETANILIS DEDTLISSIS AND DOLLO VACCINE			manadata o
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – Funded for any of the following:	[xpriarrii]		
A single dose for children up to the age of 7 who have c	ompleted primary imr	munication: or	
A course of four vaccines is funded for catch up program primary immunisation; or			years) to complete full
<ol> <li>An additional four doses (as appropriate) are funded for pre- or post splenectomy; pre- or post solid organ transpregimens; or</li> </ol>			
Five doses will be funded for children requiring solid org	an transplantation.		
Note: Please refer to the Immunisation Handbook for approp	•	ch up program	mes.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg		p p - 9	
pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units			
poliomyelitis virus in 0.5ml syringe	0.00	10	Infanrix IPV
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN			<del></del>
Xpharm]	ND HAEIWOPHILUS I	NFLUENZAE	I THE B VACCINE -
Funded for patients meeting any of the following criteria:			
1) Up to four doses for children up to and under the age of	10 for primary immur	nisation; or	
2) An additional four doses (as appropriate) are funded for			
10 who are patients post haematopoietic stem cell trans			
post solid organ transplant, renal dialysis and other seve			
3) Up to five doses for children up to and under the age of			
Note: A course of up-to four vaccines is funded for catch up p			
to complete full primary immunisation. Please refer to the Improgrammes	nunisation handbook	tior the approp	priate scriedule for catch up
programmes.  Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,			
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10	Infanrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]			
One dose for patients meeting any of the following:			
For primary vaccination in children; or			
2) An additional dose (as appropriate) is funded for (re-)imi	munisation for patient	ts post haemat	topoietic stem cell
transplantation, or chemotherapy; functional asplenic; pr			
or post cochlear implants, renal dialysis and other sever			
For use in testing for primary immunodeficiency disease	s, on the recommend	lation of an into	ernal medicine physician or
paediatrician.			
Haamanhilua Influenzaa tuna P naluasaaharida 10 maa			
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg			
prefilled syringe plus vial 0.5 ml		1 🗸	Hiberix
HEPATITIS A VACCINE – [Xpharm]		•	
Funded for patients meeting any of the following criteria:			
Two vaccinations for use in transplant patients; or			
Two vaccinations for use in children with chronic liver dia	sease: or		
One dose of vaccine for close contacts of known hepatit			
Inj 1440 ELISA units in 1 ml syringe			<u>Havrix</u>
Inj 720 ELISA units in 0.5 ml syringe	0.00	1	Havrix Junior

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer	
HEPATITIS B RECOMBINANT VACCINE - [Xpharm]					

✓ Engerix-B

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAq) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients following immunosuppression; or
- 8) for solid organ transplant patients; or
- 9) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 10) following needle stick injury.

**Engerix-B** 

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients following immunosuppression; or
- 8) for solid organ transplant patients; or
- 9) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 10) following needle stick injury; or
- 11) for dialysis patients: or
- 12) for liver or kidney transplant patients.

#### HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV]

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d)
- a) A) Any of the following:
  - 1) Maximum of two doses for children aged 14 years and under; or
  - 2) Maximum of three doses for patients meeting any of the following criteria:

3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy

- 1) People aged 15 to 26 years inclusive; or
- 2) Either:

People aged 9 to 26 years inclusive

- 1) Confirmed HIV infection; or
- 2) Transplant (including stem cell) patients: or
- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

✓ Gardasil 9

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
INFLUENZA VACCINE				
Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine	e)			
- [Xpharm]	11.00	1	✓ A	fluria Quad Junior

- A) INFLUENZA VACCINE child aged 6 months to 35 months
  - is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

(2023 formulation)

- i) all children aged 6 months to 35 months from 1 April 2023 to 31 December 2023.
- B) Doctors are the only Contractors entitled to claim payment for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)110.00	10	<ul> <li>Afluria Quad</li> </ul>
		(2023 formulation)

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d

#### A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- People 55 to 64 years of age (inclusive) and is Māori or of any Pacific ethnicity from 1 April 2023 to 31 December 2023; or
- c) people under 65 years of age who:
  - i) have any of the following cardiovascular diseases:
    - a) ischaemic heart disease, or
    - b) congestive heart failure, or
    - c) rheumatic heart disease, or
    - d) congenital heart disease, or
    - e) cerebo-vascular disease; or
  - ii) have either of the following chronic respiratory diseases:
    - a) asthma, if on a regular preventative therapy, or
    - b) other chronic respiratory disease with impaired lung function; or
  - iii) have diabetes; or
  - iv) have chronic renal disease; or
  - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
  - vi) have any of the following other conditions:
    - a) autoimmune disease, or
    - b) immune suppression or immune deficiency, or
    - c) HIV, or
    - d) transplant recipients, or
    - e) neuromuscular and CNS diseases/disorders, or
    - f) haemoglobinopathies, or
    - g) are children on long term aspirin, or
    - h) have a cochlear implant, or
    - i) errors of metabolism at risk of major metabolic decompensation, or
    - i) pre and post splenectomy, or
    - k) Down syndrome, or
  - vii) are pregnant; or
- d) children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- e) people under 65 years of age who:
  - i) have any of the following serious mental health conditions:
    - a) schizophrenia, or
    - b) major depressive disorder, or
    - c) bipolar disorder, or
    - d) schizoaffective disorder, or
  - ii) are currently accessing secondary or tertiary mental health and addiction services; or
- f) children 3 to 12 years of age (inclusive), from 1 April 2023 to 31 December 2023;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine [Xpharm]		5		uQuadri (2023 Formulation)

#### A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) all children aged 6 months to 35 months from 1 July 2023 to 31 December 2023.
- B) Doctors are the only Contractors entitled to claim payment for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

(FluQuadri (2023 Formulation) Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to be delisted 1 January 2024)

### MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)

#### A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children: or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora for subsidised immunisation, and they may only do so in respect of the measles. mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50, Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

Subsidy	ce)	Fully	Brand or
(Manufacturer's Pri		Subsidised	Generic
\$	Per	✓	Manufacturer

### MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- a) A) Any of the following:
  - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
  - 2) One dose for close contacts of meningococcal cases of any group; or
  - 3) One dose for person who has previously had meningococcal disease of any group; or
  - 4) A maximum of two doses for bone marrow transplant patients; or
  - 5) A maximum of two doses for person pre- and post-immunosuppression\*; or
  - B) Both:
    - 1) Person is aged between 13 and 25 years, inclusive; and
    - 2) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons.
  - C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A, C, Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
  - D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

, ,	•	, ,	, 0			
to a total	of approximately 55	mcg of tetani	us toxoid carrier			
		-		0.00	1	✓ MenQuadfi

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE a) Only on a prescription b) No patient co-payment payable c) a) Any of the following: A) Three doses for children up to 12 months of age (inclusive) for primary immunisation; or B) Up to three doses (dependent on age at first dose) for a catch-up programme for children from 13 months to 59 months of age (inclusive) for primary immunisation, from 1 March 2023 to 31 August 2025; or C) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients; or v) up to two doses for person pre- and post-immunosuppression\*; or D) Both: 1) Person is aged between 13 and 25 years (inclusive); and 2) Either: i) Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prisons; or ii) Two doses for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 March 2023 to 28 February 2024. E) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule. F) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-D above. \*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 12 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression\*. Note: children under 12 months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine. \*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

✓ Neisvac-C

1

Subsidy (Manufacturer's Price)		Fully	Brand or Generic	
\$	Per	•	Manufacturer	

#### PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm]

1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B,

7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

10 **✓ Synflorix** 

### PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
- 2) Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
  - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
  - b) primary immune deficiencies; or
  - c) HIV infection: or
  - d) renal failure, or nephrotic syndrome; or
  - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
  - f) cochlear implants or intracranial shunts; or
  - g) cerebrospinal fluid leaks; or
  - receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
  - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
  - j) pre term infants, born before 28 weeks gestation; or
  - k) cardiac disease, with cyanosis or failure; or
  - I) diabetes; or
  - m) Down syndrome; or
  - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml

10

✓ Prevenar 13

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE — [Either:  1) Up to three doses (as appropriate) for patients with HIN chemotherapy; pre- or post-splenectomy or with function complement deficiency (acquired or inherited), cochleated 2) All of the following:  a) Patient is a child under 18 years for (re-)immunistic b) Treatment is for a maximum of two doses; and c) Any of the following:  i) on immunosuppressive therapy or radiation immune response; or  iii) with primary immune deficiencies; or  iii) with HIV infection; or  iv) with renal failure, or nephrotic syndrome; or  v) who are immune-suppressed following orgator  vi) with cochlear implants or intracranial shuntsic vii) with cerebrospinal fluid leaks; or  viii) receiving corticosteroid therapy for more that prednisone of 2 mg/kg per day or greater, of 20 mg or greater; or  ix) with chronic pulmonary disease (including a x) pre term infants, born before 28 weeks gest xi) with cardiac disease, with cyanosis or failure xii) with diabetes; or	Xpharm]  7, for patients post hae anal asplenia, pre- or primary ation; and  therapy, vaccinate when transplantation (inclust; or  an two weeks, and where children who weight restricted with high ation; or	ematopoietic stepost-solid organ immunodeficier seen there is expuding haematop or are on an equence than 10 kg	em cell transplant, or transplant, renal dialysis, ncy; or ected to be a sufficient poietic stem cell transplant); ivalent daily dosage of y on a total daily dosage of
xiv) who are pre-or post-splenectomy, or with fu	nctional asplenia.		
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	: viduals; or oriate schedule for cat	ch-up programr	Pneumovax 23 nes. IPOL
ROTAVIRUS ORAL VACCINE – [Xpharm]  Maximum of two doses for patients meeting the following:  1) first dose to be administered in infants aged under 14 v  2) no vaccination being administered to children aged 24	veeks of age; and		<u></u>
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator			Rotarix Rotarix

Subsidised

Per

Subsidy

(Manufacturer's Price)

\$

Fully

Brand or

Generic

Manufacturer

VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm] Either:			
1) Maximum of one dose for primary vaccination for either:			
a) Any infant born on or after 1 April 2016; or			
<li>b) For previously unvaccinated children turning 11 years old on varicella infection (chickenpox), or</li>	or after 1 Ju	ly 2017, w	ho have not previously had a
2) Maximum of two doses for any of the following:			
a) Any of the following for non-immune patients:			
i) with chronic liver disease who may in future be candidate.	ates for trans	nlantation	or
ii) with deteriorating renal function before transplantation;		piaritation	, 01
iii) prior to solid organ transplant; or			
iv) prior to any elective immunosuppression*, or			
v) for post exposure prophylaxis who are immune compe			
b) For patients at least 2 years after bone marrow transplantation			
c) For patients at least 6 months after completion of chemother			
<ul> <li>d) For HIV positive non immune to varicella with mild or modera</li> <li>e) For patients with inborn errors of metabolism at risk of major</li> </ul>			
varicella, or	metabolic de	compens	alion, with no clinical history of
f) For household contacts of paediatric patients who are immu	nocompromis	sed. or und	dergoing a procedure leading to
immune compromise where the household contact has no cl			
g) For household contacts of adult patients who have no clinical			
immunocompromised, or undergoing a procedure leading to	immune com	npromise v	vhere the household contact
has no clinical history of varicella.			and a select of market them
<ul> <li>immunosuppression due to steroid or other immunosuppressive theral</li> <li>28 days</li> </ul>	py must be to	or a treatm	ent period of greater than
Inj 1350 PFU prefilled syringe0.0	00 -	1	✓ Varivax
,		0	✓ Varivax
VARICELLA ZOSTER VACCINE [SHINGLES VACCINE]			
a) Only on a prescription			
b) No patient co-payment payable			
c)			
A) Funded for patients meeting the following criteria:			
Two doses for all people aged 65 years  October the serial be petitled to plain nowment from the Funday for the serial people aged 65 years.  October the serial people aged 65 years.	مرامعين مطلع	of Varion	la zastar vassina (Chinalas
<ul> <li>B) Contractors will be entitled to claim payment from the Funder f vaccine) to patients eligible under the above criteria pursuant t</li> </ul>			
Zealand for subsidised immunisation, and they may only do so			
vaccine] listed in the Pharmaceutical Schedule.			ona zooto: racomo [o.m.g.co
C) Contractors may only claim for patient populations within the c	riteria that are	e covered	by their contract, which may be
a sub-set of the population described in paragraph A above.			
Inj 50 mcg per 0.5 ml vial plus vial	00 .	1	✓ Shingrix
Diagnostic Agents			
Diagnostio Agento			
TUBERCULIN PPD [MANTOUX] TEST - [Xpharm]			
Inj 5 TU per 0.1 ml, 1 ml vial0.0	00 -	1	✓ <u>Tubersol</u>

- Symbols -	Renin-Angiotensin System 48	Amorolfine7
3TC115	Agents for Parkinsonism and Related	Amoxicillin10
7 MED NSHA Silver/Copper	Disorders 125	Amoxicillin with clavulanic acid10
Short82	Agents Used in the Treatment of	Amphotericin B3
- A -	Poisonings264	Amsacrine16
A-Scabies77	Agrylin160	AmsaLyo16
Abacavir sulphate115	Albendazole98	Amsidine16
Abacavir sulphate with	Albey249–250	Amzoate3
lamivudine115	Albustix86	Anaesthetics12
Abacavir/Lamivudine Viatris115	Alchemy Oxaliplatin156	Anagrelide hydrochloride16
Abiraterone acetate174	Alchemy Oxybutynin85	Analgesics12
Acarbose11	Aldurazyme30	Anastrozole17
Accarb11	Alecensa167	Anatrole17
Accuretic 1049	Alectinib	Andriol Testocaps9
Accuretic 2049	Alendronate sodium120	Androderm8
Acetazolamide261	Alendronate sodium with	Anoro Ellipta25
Acetec48	colecalciferol120	Antabuse
Acetic acid with hydroxyquinoline and	Alfacalcidol34	Antacids and Antiflatulents
ricinoleic acid84	Alfamino285	Anthelmintics9
Acetylcysteine264	Alfamino Junior285	Antiacne Preparations7
Aci-Jel84	Alginic acid6	Antiallergy Preparations24
Aciclovir	Alglucosidase alfa27	Antianaemics3
Infection110	Alkeran156	Antiandrogen Oral
Sensory259	Alkeran S29156	Contraceptives8
Acidex6	Allerfix261	Antiarrhythmics5
Acipimox56	Allerpro Syneo 1288	Antibacterials9
Acitretin77	Allerpro Syneo 2288	Antibacterials Topical7
Actemra230	Allersoothe250	Anticholinergic Agents25
Actinomycin D161	Allmercap	Anticholinesterases11
Actrapid10	Allopurinol	Antidepressants13
Actrapid Penfill10	Alpha-Adrenoceptor Blockers48	Antidiarrhoeals
Acupan127	Alpha-Keri Lotion75	Antiepilepsy Drugs13
Adalimumab (Amgevita)185	Alphamox101	Antifibrinolytics, Haemostatics and
Adalimumab (Humira - Alternative	Alphamox 125101	Local Sclerosants3
brand) 193	Alphamox 250101	Antifibrotics25
Adapalene71	Alprolix39	Antifungals10
ADR Cartridge 1.825	Alu-Tab6	Antifungals Topical7
Adrenaline	Aluminium hydroxide6	Antihistamines25
Cardiovascular59	Alvogen52	Antihypotensives5
Respiratory249	Amantadine hydrochloride125	Antimalarials10
Advantan74	Ambrisentan59	Antimigraine Preparations13
Advate41	Ambrisentan Mylan59	Antinausea and Vertigo Agents 13
Adynovate42	Ambrisentan Viatris59	Antipruritic Preparations7
Afinitor245	Amgevita185	Antipsychotics13
Aflibercept200	Amiloride hydrochloride54	Antiretrovirals11
Afluria Quad	Amiloride hydrochloride with	Antirheumatoid Agents12
(2023 formulation)294	furosemide55	Antispasmodics and Other Agents
Afluria Quad Junior	Amiloride hydrochloride with	Altering Gut Motility
(2023 formulation) 294	hydrochlorothiazide55	Antithrombotic Agents4
AFT-Pyrazinamide109	Aminophylline255	Antithymocyte globulin
After Hours Med Mgmt 15 min264	Amiodarone hydrochloride50	(equine) 18
After Hours Med Mgmt 30 min264	Amisulpride137	Antitrichomonal Agents10
After Hours Med Mgmt 45 min264	Amitriptyline131	Antituberculotics and
Agents Affecting the	Amlodipine53	Antileprotics10

Antiulcerants	9	Atenolol	51	Benzathine benzylpenicillin	10
Antivirals		Atenolol AFT		Benzatropine mesylate	
Antivirals Eligibility Review		Atenolol AFT S29		Benzbromaron AL 100	
Anxiolytics		Atenolol Viatris		Benzbromarone	
Anzatax		Atezolizumab		Benztrop	
Apidra		ATGAM	105	Benzydamine hydrochloride	
Apidra SoloStar		Ativan		Benzylpenicillin sodium [Penicillin	
'					10
APO-Atomoxetine		Atnahs Olsalazine		G]	
APO-Atomoxetine S29		AtomoxetineAtorvastatin		Beta Cream Beta Ointment	
Apo-AzithromycinAPO-Candesartan HCTZ	99		30	Beta Scalp	
16/12.5	40	Atropine sulphate	EO		
	49	Cardiovascular		Beta-Adrenoceptor Agonists	20
APO-Candesartan HCTZ 32/12.5	40	Sensory		Beta-Adrenoceptor Blockers	
		Atropt		Betadine	
Apo-Temozolomide		Atrovent		Betadine Skin Prep	
Apomorphine hydrochloride		Aubagio		Betaferon	
Aprepitant		Augmentin		Betahistine dihydrochloride	. 13
Apresoline		Autorix		Betaine	
Aqueous cream		AutoSoft 30		Betaloc CR	
Aratac		AutoSoft 90		Betamethasone dipropionate	7
Arava		Avallon		Betamethasone dipropionate with	
Arginine		Avelox		calcipotriol	7
Aripiprazole		Avonex		Betamethasone sodium phosphate	
Aripiprazole Sandoz		Avonex Pen		with betamethasone acetate	
Aristocort		Azacitidine		Betamethasone valerate7	-,
Arrotex-Prazosin S29		Azacitidine Dr Reddy's		Betamethasone valerate with sodiur	
Arrow - Clopid		Azamun		fusidate [fusidic acid]	
Arrow - Lattim		Azathioprine		Betaxolol	
Arrow-Amitriptyline	131	Azilect	125	Betnovate	
Arrow-Bendrofluazide	55	Azithromycin		Betoptic	. 26
Arrow-Brimonidine	262	Azopt	261	Betoptic S	
Arrow-Diazepam		AZT	115	Bexsero	. 29
Arrow-Doxorubicin	161	- B -		Bezafibrate	
Arrow-Fluoxetine	132	B-D Micro-Fine	16	Bezalip	
Arrow-Losartan &		B-D Ultra Fine	16	Bezalip Retard	5
Hydrochlorothiazide	49	B-D Ultra Fine II	16	Bicalutamide	. 17
Arrow-Norfloxacin	118	Bacillus Calmette-Guerin (BCG)		Bicillin LA	
Arrow-Ornidazole	107	vaccine	185	BiCNU	. 15
Arrow-Quinapril 10	49	Bacillus Calmette-Guerin		Bile and Liver Therapy	10
Arrow-Quinapril 20	49	vaccine	291	Biltricide	
Arrow-Quinapril 5	49	Baclofen	124	Bimatoprost	. 26
Arrow-Roxithromycin	100	Bactroban	71	Bimatoprost Multichem	. 26
Arrow-Timolol	261	Balance	30	Binarex	
Arrow-Topiramate	135	Barrier Creams and Emollients	74	Binocrit	3
Arrow-Tramadol	130	BCG Vaccine	291	Biodone	. 12
Arsenic trioxide	160	Beclazone 100	251	Biodone Extra Forte	. 129
Asacol	7	Beclazone 250	251	Biodone Forte	. 129
Ascend Aripiprazole	137	Beclazone 50	251	Bisacodyl	2
Ascorbic acid		Beclomethasone dipropionate	251	Bisacodyl Viatris	
Aspen Adrenaline		Bedaquiline	108	Bisoprolol fumarate	
Aspirin		Bee venom allergy treatment		Bisoprolol Mylan	
Blood	42	Bendamustine hydrochloride		Bisoprolol Viatris	5
Nervous	127	Bendrofluazide		BK Lotion	
Asthalin	252	Bendroflumethiazide		Bleomycin sulphate	. 160
Atazanavir Mylan	115	[Bendrofluazide]	55	Blood Colony-stimulating	
Atazanavir sulphate	115	Benralizumab		Factors	4

Blood glucose diagnostic test	Calcium carbonate PAI	6	Ceftriaxone	98
meter 15	Calcium Channel Blockers	.53	Ceftriaxone-AFT	
Blood glucose diagnostic test	Calcium Disodium Versenate	265	Cefuroxime axetil	98
strip15	Calcium folinate	157	Celapram	
Blood glucose test strips (visually	Calcium Folinate Ebewe	157	Celebrex	119
impaired)15	Calcium Folinate Sandoz	157	Celecoxib	
Blood Ketone Diagnostic Test	Calcium Folinate Sandoz S29	157	Celecoxib Pfizer	
Strip 14	Calcium gluconate		Celestone Chronodose	
Bonjela33	Calcium Homeostasis		Cellcept	
Boostrix291	Calcium polystyrene sulphonate		Centrally-Acting Agents	
Bortezomib	Calcium Resonium		Cephalexin ABM	
Bosentan	Calogen		Cetirizine hydrochloride	
Bosentan Dr Reddy's62	Camber		Cetomacrogol	
Bplex34	Candesartan cilexetil		Cetomacrogol with glycerol	
Breo Ellipta252	Candesartan cilexetil with		Cetomacrogol-AFT	
Brevinor 1/2883	hydrochlorothiazide	49	Cetuximab	20:
Brevinor-1 28 Day83	Candestar		Charcoal	
Bricanyl Turbuhaler253	Canesten		Chemotherapeutic Agents	
Brimonidine tartrate	Capecitabine		Chickenpox vaccine	
Brimonidine tartrate with timolol			Chlorambucil	30
	Capecitabine Viatris			
maleate	Capecitabine-DRLA		Chlorathicatida	
Brinzolamide	Capercit	107	Chlorothiazide	
Brolene	Capoten	. 48	Chlorpheniramine maleate	
Brown & Burk	Capsaicin	400	Chlorpromazine hydrochloride	
Brufen SR	Musculoskeletal		Chlorsig	
BSF Concerta	Nervous		Chlortalidone [Chlorthalidone]	
BSF Rubifen SR264	Captopril		Chlorthalidone	
Buccastem137	Carafate		Chlorvescent	
Budesonide	Carbaccord		Choice 380 7med Nsha Silver/cop	
Alimentary6	Carbamazepine		Short	
Respiratory251, 257	Carbimazole		Choice Load 375	
Budesonide with eformoterol252	Carbomer		Choice TT380 Short	
Bumetanide54	Carboplatin		Choice TT380 Standard	
Buprenorphine Naloxone BNM150	Carboplatin Ebewe	155	Choline salicylate with cetalkonium	n
Buprenorphine with naloxone150	Carbosorb-X	264	chloride	33
Bupropion hydrochloride151	Cardinol LA	. 52	Ciclosporin	
Burel106	Cardizem CD	. 53	Cilazapril	
Burinex54	CareSens Dual	.14	Cilicaine VK	102
Burinex S2954	CareSens N	. 15	Cinacalcet	8
Buscopan8	CareSens N POP	. 15	Cinacalet Devatis	8
Buscopan S298	CareSens N Premier	. 15	Cipflox	102
Buspirone hydrochloride140	CareSens PRO	. 15	Ciprofloxacin	
Buspirone Viatris140	Carmellose sodium with gelatin and		Infection	102
Busulfan	pectin	33	Sensory	259
- C -	Carmustine		Ciprofloxacin - Torrent	
Cabergoline96	Carnitor		Ciprofloxacin Teva	
Caffeine citrate258	Carvedilol		Cisplatin	15
Calamine	Carvedilol Sandoz		Cisplatin Ebewe	
Calamine-AFT73	Casirivimab and imdevimab		Citalopram hydrobromide	
Calci-Tab 500	Catapres		Cladribine	
Calcipotriol77	CeeNU		Clarithromycin	
Calcitonin87	Cefaclor monohydrate		Alimentary	(
Calcitriol 34	Cefalexin	98	Infection	
Calcitriol-AFT34	Cefalexiii Sandoz		Clexane	
Calcium 500 mg Hexal	Cefazolin		Clexane Forte	
Calcium carbonate	Cefazolin-AFT		Climara	
Calcium Calbunate	Ociazoilli-Al- I	. 50	OIIIIIaia	91

Clindamycin	103	[Dextrose]	46	Dapa-Tabs	5
Clinicians		Compound hydroxybenzoate		Dapsone	
Clinicians Renal Vit		Comtan		Daraprim	
Clobazam		Concerta		Darunavir	
Clobetasol propionate		Condoms		Darunavir Mylan	
Clobetasone butyrate		Condyline		Darunavir Viatris	
Clofazimine		Contraceptives - Hormonal		Dasatinib	
Clomazol		Contraceptives - Non-hormonal		Daunorubicin	
Dermatological	72	Copaxone		Daunorubicin Zentiva	
Genito-Urinary		Cordarone-X		David One Step Cassette Pregnar	
Clomifene citrate		Corticosteroids and Related Age		Test	
Clomipramine hydrochloride		for Systemic Use		DBL Adrenaline	
Clomipramine Teva		Corticosteroids Topical		DBL Aminophylline	
Clonazepam		Cortifoam		DBL Bleomycin Sulfate	
Clonidine		Cosentyx		DBL Bortezomib	
Clonidine hydrochloride		Cosmegen		DBL Carboplatin	
Clonidine Teva		Coumadin		DBL Cisplatin	
Clopidogrel		Country Life		DBL Dacarbazine	
Clopine		Coversyl		DBL Desferrioxamine Mesylate fo	
Clopixol		Creon 10000		BP	
Clotrimazole		Creon 25000		DBL Docetaxel	
Dermatological	72	Creon Micro		DBL Ergometrine	
Genito-Urinary		Crotamiton		DBL Gemcitabine	
Clozapine		Crystaderm		DBL Gentamicin	
Clozaril		Curam		DBL Heparin Sodium	
Co-trimoxazole		Curam Duo 500/125		DBL Leucovorin Calcium	
Coal tar		Cvite		DBL Methotrexate Onco-Vial	159
Coal tar with allantoin, menth		Cyclizine hydrochloride		DBL Pethidine Hydrochloride	
phenol and sulphur		Cyclizine lactate		DBL Vincristine Sulfate	
Coal tar with salicylic acid and		Cyclogyl		Decozol	
sulphur		Cyclonex		Deferasirox	
Cobal-B12		Cyclopentolate hydrochloride		Deferiprone	
Cobalin-H		Cyclophosphamide		Deferoxamine Pfizer S29	
Coco-Scalp		Cyclorin		Denosumab	
Codeine phosphate		Cycloserine		Deolate	
Extemporaneous	267	Cyproterone acetate		Deoxycoformycin	
Nervous		Cyproterone acetate with		Depo-Medrol	
Coenzyme Q10		ethinyloestradiol	84	Depo-Provera	
Colchicine		Cystadane		Depo-Testosterone	8
Colecalciferol		Cytarabine		Deprim	
Colestid		Cytotec		Dermol	
Colestipol hydrochloride		Cytoxan		Desferrioxamine mesilate	
Colestyramine		- D -		Desmopressin	
Colestyramine - Mylan		D-Penamine	120	Desmopressin acetate	9
Colgout		Dabigatran		Desmopressin-PH&T	9
Colifoam		Dacarbazine		Desuric	
Colistin sulphomethate		Dacarbazine APP		Detection of Substances in	
Colistin-Link		Dactinomycin [Actinomycin D]		Urine	8
Collodion flexible		Daivobet		Dexamethasone	
Colloidal bismuth subcitrate		Daivonex		Hormone	A!
Colofac		Daktarin		Sensory	
Coloxyl		Dalacin C		Dexamethasone phosphate	
Combigan		Dantrium		Dexamethasone with framycetin a	
Compliance Packaging		Dantrium S29		gramicidin	
Compound electrolytes		Dantrolene		Dexamethasone with neomycin	20
Compound electrolytes with g		Daonil		sulphate and polymyxin B	
Sompound Glood of your Willing	140000	Duv	1 1	odipilato ana polymynii D	

sulphate260	Domperidone Viatris	136	Elecare	28
Dexamfetamine sulfate146	Donepezil hydrochloride	150	Elecare LCP	28
Dexmethsone88	Donepezil-Rex	150	Electral	4
Dextrochlorpheniramine	Dornase alfa	255	Elelyso	3
maleate250	Dortimopt	262	Elemental 028 Extra	27
Dextrose45–46	Dorzolamide hydrochloride	261	Elexacaftor with tezacaftor, ivaca	ftor
DHC Continus129	Dorzolamide with timolol	262	and ivacaftor	25
Diabetes10	Dostinex	96	Elidel	7
Diabetes Management14	Dosulepin [Dothiepin]		Elocon	7
Diacomit134	hydrochloride	131	Elocon Alcohol Free	7
Diagnostic Agents301	Dosulepin Mylan		Eltrombopag	3
Diamide Relief6	Dosulepin Viatris		Eltroxin	
Diamox261	Dothiepin		EMB Fatol	10
Diasip272	Doxazosin	48	Emend Tri-Pack	13
Diazepam 132, 141	Doxazosin Clinect	48	Emicizumab	4
Diazoxide10	Doxine	102	EMLA	12
Dibenzyline48	Doxorubicin Ebewe	161	Empagliflozin	1
Diclofenac Sandoz119	Doxorubicin hydrochloride	161	Empagliflozin with metformin	
Diclofenac sodium	Doxycycline		hydrochloride	1
Musculoskeletal119	DP Lotion	75	Emtricitabine	
Sensory260	DP Lotn HC		Emtricitabine with tenofovir	
Differin71	DP-Allopurinol	123	disoproxil	11
Difflam33	Dr Reddy's Omeprazole	9	Emtriva	11
Diflucan105	Drofate		Emulsifying ointment	7
Digestives Including Enzymes25	Drugs Affecting Bone		Emulsifying Ointment ADE	7
Digoxin50	Metabolism	120	Enalapril maleate	4
Dihydrocodeine tartrate129	Dual blood glucose and blood ketor	ne	Enbrel	
Dilantin134	diagnostic test meter	14	Endocrine Therapy	17
Dilantin Infatab134	Dulaglutide	12	Endoxan	15
Dilantin Paediatric134	Dulcolax SP Drop	27	Engerix-B	29
Diltiazem CD Clinect53	Duocal Super Soluble Powder		Enlafax XR	13
Diltiazem hydrochloride53	Duolin	253	Enoxaparin sodium	4
Dimethicone74, 76	Duolin HFA	253	Enstilar	
Dimethyl fumarate142	Duolin Respules	253	Ensure	27
Dipentum8	DuoResp Spiromax		Ensure Plus	28
Diphtheria, tetanus and pertussis	Duride	59	Ensure Plus HN	27
vaccine291	Durvalumab	240	Ensure Plus RTH	27
Diphtheria, tetanus, pertussis and	-E-		Ensure Two Cal HN RTH	28
polio vaccine292	e-chamber La Grande	258	Entacapone	12
Diphtheria, tetanus, pertussis, polio,	e-chamber Mask	257	Entecavir	11
hepatitis B and haemophilus	e-chamber Turbo	258	Entecavir (Rex)	11
influenzae type B vaccine 292	E-Mycin	100	Entecavir Mylan	11
Diprosone	e5 Pharma	10	Entecavir Sandoz	
Diprosone OV73	Ear Preparations	259	Entocort CIR	
Dipyridamole42	Ear/Eye Preparations	259	Entresto 24/26	
Disopyramide phosphate50	Easiphen Liquid	284	Entresto 49/51	4
Disulfiram151	Econazole nitrate	72	Entresto 97/103	4
Diuretics54	Efavirenz	114	Entyvio	23
Docetaxel161	Efavirenz with emtricitabine and		Epilim	13
Docetaxel Accord161	tenofovir disoproxil	115	Epilim Crushable	13
Docetaxel Sandoz161	Eformoterol fumarate dihydrate	251	Epilim IV	13
Docusate sodium26	Eftrenonacog alfa [Recombinant		Epilim S/F Liquid	13
Docusate sodium with	factor IX]	39	Epilim Syrup	13
sennosides26	Efudix	79	Epipen	
Dolutegravir116	Egopsoryl TA		Epipen Jr	24
Domperidone 136	Elaprase	29	Epirubicin Ebewe	

Epirubicin hydrochloride	161	Eye Preparations	259	Flucloxacillin	101
Eplerenone	54	Eylea	200	Flucloxacillin-AFT	
Epoetin alfa	38	Ezetimibe	58	Flucloxin	<mark>10</mark> 1
Epoprostenol	66	Ezetimibe Sandoz	58	Flucon	260
Eptacog alfa [Recombinant fact	or	Ezetimibe with simvastatin	58	Fluconazole	105
VIIa]	41	-F-		Fludara Oral	158
Erbitux	202	Factor eight inhibitor bypassing		Fludarabine Ebewe	158
Ergometrine maleate	84	fraction	41	Fludarabine phosphate	158
Erlotinib		Famotidine	9	Fludrocortisone acetate	8
Erythrocin IV	100	Famotidine Hovid		Fluids and Electrolytes	45
Erythromycin (as lactobionate).	100	Fasenra	201	Flumetasone pivalate	259
Erythromycin ethyl succinate		Faslodex	175	Fluocortolone caproate with	
Esbriet		Fatty Cream AFT	75	fluocortolone pivalate and	
Escitalopram	131	Febuxostat	123	cinchocaine	8
Escitalopram (Ethics)		Febuxostat multichem	123	Fluorometholone	260
Eskazole		Feed Thickener Karicare		Fluorouracil	158
Essential Ethosuximide		Aptamil	282	Fluorouracil Accord	158
Essential Generics	51	FEIBA NF		Fluorouracil sodium	
Essential Prednisolone		Felo 10 ER		Fluox	
Estraderm MX		Felo 5 ER		Fluoxetine hydrochloride	
Estradiol TDP Mylan		Felodipine		Flupenthixol decanoate	
Estradiol Viatris		Fenpaed 100 mg per 5 ml		FluQuadri (2023 Formulation)	
Estradot		Fentanyl		Flutamide	
Estradot 50 mcg		Fentanyl Sandoz		Flutamin	
Estrofem		Ferinject		Fluticasone	
Etanercept		Ferodan		Fluticasone furoate with	
Ethambutol hydrochloride		Ferriprox		vilanterol	252
Ethics Aspirin		Ferro-F-Tabs		Fluticasone propionate	
Ethics Aspirin EC		Ferro-Liquid		Fluticasone with salmeterol	
Ethics Lisinopril		Ferro-tab		Flynn	
Ethinyloestradiol with		Ferrograd		FML	
desogestrel	82	Ferrosig		Foban	
Ethinyloestradiol with		Ferrous fumarate		Folic acid	
levonorgestrel	83	Ferrous fumarate with folic acid.		Folic Acid multichem	
Ethinyloestradiol with		Ferrous sulfate		Folic Acid Mylan	
norethisterone	83	Fexofenadine hydrochloride		Folic Acid Viatris	
Ethosuximide		Fibro-vein		Food Thickeners	
Etopophos		Filgrastim		Foods And Supplements For Inbo	
Etoposide		Finasteride		Errors Of Metabolism	
Etoposide phosphate		Fingolimod		Forteo	
Etravirine		Firazyr		Fortini	
Eumovate		Flagyl	107	Fortini Multi Fibre	
Eurofolic		Flagyl-S	107	Fortisip	
Evara		Flamazine		Fortisip Multi Fibre	
Everet		Flecainide acetate		Fosamax	
Everolimus		Flecainide BNM		Fosamax Plus	
Evista		Flecainide Controlled Release		Framycetin sulphate	
Evrysdi		Teva	51	Frebini Energy	27/
Evusheld		Flecainide Sandoz		Frebini Energy Fibre	
Exelon Patch 10		Fleet Phosphate Enema		Frebini Original	
Exelon Patch 5		Flixonase Hayfever & Allergy		Frebini Original Fibre	
Exemestane		Flixotide		Fresubin 2kcal HP	
Exjade		Flixotide Accuhaler		Fresubin HP Energy	
Extemporaneously Compounde		Florinef		Fresubin HP Energy Fibre	
	u				
Preparations and Galenicals	067	Fluanxol		Fresubin Intensive Fresubin Original	
Gaici iicais	20/	Flucil	IVI	riesuvili Oligiliai	4/8

Fresubin Original Fibre	279	Glycopyrronium bromide	8	Humira	193
Frisium	133	Glycopyrronium with		HumiraPen	193
Frumil	55	indacaterol	254	Humulin 30/70	11
Frusemide	54	Gold Knight	81	Humulin NPH	11
Fucicort	74	Gold Knight XL	81	Humulin R	10
Fucidin	104	Goserelin		Hyaluronic acid	263
Fucithalmic		Gynaecological Anti-infectives		Hydralazine	
Fulvestrant		- H -		Hydralazine hydrochloride	
Fungilin		Habitrol	152	Hydrocortisone	
Furosemid-Ratiopharm		Haemophilus influenzae type B		Dermatological	73
Furosemide [Frusemide]		vaccine	292	Hormone	
Furosemide-Baxter		Haldol		Hydrocortisone acetate	
fusidic acid		Haldol Concentrate		Hydrocortisone acetate with	
Dermatological	72 74	Haldol Decanoas		pramoxine hydrochloride	7
Infection		Haloperidol		Hydrocortisone and paraffin liquid	
Sensory		Haloperidol decanoate		and lanolin	
- G -	200	Harvoni		Hydrocortisone butyrate	
Gabapentin	122	Havrix		Hydrocortisone with cinchocaine	-
Gacet		Havrix Junior		Hydrocortisone with miconazole	
Galsulfase					
Galvumet		Haylor syrup Croom SI S	230	Hydrocortisone with natamycin a	
		healthE Aqueous Cream SLS Free	75	neomycin	
Galvus Gardasil 9				Hydrogen peroxide	
		healthE Calamine Aqueous		Hydroxocobalamin	
Gastrodenol		healthE Dimethicone 10%		Hydroxocobalamin Panpharma	
Gaviscon Extra Strength		healthE Dimethicone 4% Lotion		hydroxycarbamide	
Gaviscon Infant		healthE Olymethicone 5%		Hydroxychloroquine	120
Gazyva		healthE Glycerol BP		Hydroxyurea	404
Gefitinib		healthE Urea Cream		[hydroxycarbamide]	
GEM Aqueous Cream		Healtheries Simple Baking Mix		Hygroton	
Gemcitabine Ebewe		Hemastix		Hylo-Fresh	
Gemcitabine hydrochloride		Hemlibra		Hymenoptera2	
Gemtuzumab ozogamicin		Heparin sodium		Hyoscine butylbromide	
Gentamicin sulphate		Heparin Sodium Panpharma		Hyoscine hydrobromide	
Gilenya		Heparinised saline		Hypam	
Ginet		Heparon Junior		Hyperuricaemia and Antigout	
Glatiramer acetate		Hepatitis A vaccine	292	Hypromellose	
Glecaprevir with pibrentasvir	112	Hepatitis B recombinant		Hypromellose with dextran	263
Glibenclamide	11	vaccine	293	-1-	
Gliclazide	11	Herceptin	234	Ibiamox	101
Glipizide		Hiberix	292	Ibrance	171
Glivec	170	Hiprex	118	lbrutinib	
Glizide	11	Histaclear	250	Ibuprofen	119
Glucagen Hypokit	10	Histafen	250	Icatibant	249
Glucagon hydrochloride	10	Holoxan	155	Idarubicin hydrochloride	162
Glucerna Select	272	Horleys Bread Mix	282	Idursulfase	29
Glucose [Dextrose]	45	Horleys Flour	282	Ifosfamide	155
Gluten Free Foods	282	Hormone Replacement Therapy -		Ikorel	59
Glycerin with sodium saccharir	1267	Systemic	90	llevro	261
Glycerin with sucrose	267	HPV		lloprost	
Glycerol		Humalog		Imatinib mesilate	
Alimentary	27	Humalog Mix 25		Imatinib-Rex	
Extemporaneous		Humalog Mix 50		Imbruvica	
Glyceryl trinitrate	•	Human papillomavirus (6, 11, 16,		Imfinzi	
Alimentary	8	31, 33, 45, 52 and 58) vaccine	- 7	Imigran	
Cardiovascular		[HPV]	293	Imipramine hydrochloride	
Glycopyrronium		Humatin		Imiquimod	
- , ,				1	

Immune Modulators116	Intra-uterine device		Infection	
Immunisation Administration264	Invega Sustenna		Ketogenic Diet	
Immunisation	Invega Trinza		Ketoprofen	
Co-administration264	IPCA-Frusemide		KetoSens	14
Immunosuppressants178	IPCA-Metoprolol	52	Keytruda	24
Incruse Ellipta253	IPCA-Propranolol	52	Kindergen	27
Indacaterol251	IPOL	300	Klacid	
Indapamide55	Ipratropium bromide	253, 257	Alimentary	
Infanrix IPV292	Iressa		Infection	9
Infanrix-hexa292	Irinotecan Actavis 100	158	Kliogest	9
Infant Formulae285	Irinotecan hydrochloride	158	Kliovance	
Infatrini289	Irinotecan-Rex	158	Kogenate FS	
Infliximab203	Iron (as ferric carboxymaltos		Konakion MM	
Influenza vaccine294	Iron polymaltose		Konsyl-D	
Inhaled Corticosteroids251	Isentress		Kuvan	
Inhaled Long-acting	Isentress HD		-L-	
Beta-adrenoceptor Agonists 251	Ismo 20		Labetalol	5
Inspra54	Ismo 40 Retard		Lacosamide	
Instillagel Lido126	Isoniazid		Lactulose	
Insulin aspart11	Isoniazid with rifampicin		Laevolac	
Insulin aspart with insulin aspart	Isoptin		Lagevrio	
protamine10	Isoptin Retard		Lamictal	
Insulin glargine11	Isoptin SR		Lamivudine	
Insulin glulisine11	Isopto Carpine		Lamivudine Alphapharm	
	Isosorbide mononitrate		Lamivudine Viatris	
Insulin isophane	Isosource Standard		Lamivudine/Zidovudine Viatris	
Insulin isophane with insulin				
neutral	Isotretinoinhusle		Lamotrigine	
Insulin lispro	Ispaghula (psyllium) husk		Lamprene	
Insulin lispro with insulin lispro	Itch-Soothe		Lanoxin	
protamine11	Itraconazole		Lanoxin Paediatric Elixir	
Insulin neutral10	Itrazole		Lanoxin PG	
Insulin pen needles16	lvacaftor		Lanoxin S29	
Insulin pump16	Ivermectin	76	Lansoprazole	
Insulin pump cartridge21	- <b>J</b> -		Lantus	
Insulin pump infusion set (steel	Jadelle		Lantus SoloStar	
cannula)21	Jakavi		Lanvis	
Insulin pump infusion set (steel	Jardiamet		Lanzol Relief	
cannula, straight insertion) 22	Jardiance		Lapatinib ditosylate	
Insulin pump infusion set (teflon	Jaydess		Largactil	
cannula)23	Jevity HiCal RTH		Laronidase	
Insulin pump infusion set (teflon	Jevity Plus		Lasix	
cannula, angle insertion with	Jevity RTH	279	Latanoprost	26
insertion device)24	Jinarc	55	Latanoprost with timolol	
Insulin pump infusion set (teflon	Juno Pemetrexed	159	Lax-Suppositories	2
cannula, angle insertion) 24	- K -		Lax-suppositories Glycerol	2
Insulin pump infusion set (teflon	Kadcyla	236	Laxatives	20
cannula, straight insertion with	Kalydeco	256	Laxsol	
insertion device)24	Kemadrin	125	Ledipasvir with sofosbuvir	11
Insulin pump infusion set (teflon	Kenacomb	259	Leflunomide	120
cannula, straight insertion) 24	Kenacort-A 10	89	Lenalidomide	16
Insulin pump reservoir25	Kenacort-A 40	89	Letrole	
Insulin syringes, disposable with	Kenalog in Orabase		Letrozole	
attached needle	Ketocal 3:1		Leukeran FC	
Intelence114	KetoCal 4:1		Leukotriene Receptor	
Interferon beta-1-alpha142	Ketoconazole		Antagonists	25
Interferon beta-1-beta142	Dermatological	79	Leuprorelin	
	J			_

Leustatin	157	Lorazepam	141	vaccine	29
Levetiracetam	133	Lorstat		Meningococcal C conjugate	
Levetiracetam-AFT	133	Losartan Actavis	49	vaccine	29
Levocabastine	261	Losartan potassium	49	MenQuadfi	29
Levocarnitine		Losartan potassium with		Menthol	<mark>7</mark>
Levodopa with benserazide		hydrochlorothiazide	49	Mepolizumab	
Levodopa with carbidopa		Lovir		Mercaptopurine	
Levomepromazine		Loxamine		Mercilon 28	
Levomepromazine		Lucrin Depot 1-month		Mesalazine	
hydrochloride	138	Lucrin Depot 3-month	96	Mesna	
Levonorgestrel		Lyderm		Mestinon	
Genito-Urinary	83–84	Lynparza		Metabolic Disorder Agents	
Hormone		Lyrica		Metformin hydrochloride	
Levonorgestrel BNM		- M -		Metformin Mylan	
Levothyroxine		m-Eslon	130	Metformin Viatris	
Lidocaine [Lignocaine]		Mabthera		Methadone BNM	
Lidocaine [Lignocaine]		Macro Organic Psyllium Husk		Methadone hydrochloride	
hydrochloride	126	Macrobid		Extemporaneous	26
Lidocaine [Lignocaine] with	120	Macrogol 3350 with potassium		Nervous	
chlorhexidine	126	chloride, sodium bicarbonate	and	Methenamine (hexamine)	12
Lidocaine [Lignocaine] with	120	sodium chloride		hippurate	11
prilocaine	127	Madopar 125		Methopt	
Lidocaine-Baxter		Madopar 250		Methotrexate	
Life Extension		Madopar 62.5		Methotrexate DBL Onco-Vial	
Lignocaine		Madopar HBS		Methotrexate Ebewe	
Linezolid		Madopar Rapid		Methotrexate Sandoz	
Lioresal Intrathecal		Magnesium hydroxide		Methyl hydroxybenzoate	
Lipid-Modifying Agents		Magnesium sulphate		Methylcellulose	
Liquigen		Mantoux		Methylcellulose with glycerin and	20
Liraglutide		Marevan		sodium saccharin	26
Lisinopril	12	Marine Blue Lotion SPF 50+		Methylcellulose with glycerin and	20
Litak		Martindale Pharma		SUCTOSE	26
Lithium carbonate		Mask for spacer device		Methyldopa	
Livostin		Maviret		Methyldopa Mylan	
LMX4		Maxidex		Methyldopa Mylan S29	
Lo-Oralcon 20 ED		Maxitrol		Methylnaltrexone bromide	
Locacorten-Viaform ED's		MCT oil (Nutricia)	271	Methylphenidate ER - Teva	
Local preparations for Anal ar		Measles, mumps and rubella	206	Methylphenidate hydrochloride	14
Rectal Disorders		vaccine		Methylphenidate hydrochloride	1.4
Locasol		Mebendazole Mebeverine hydrochloride		extended-release	
Locoid Crelo				Methylprednisolone	0
		Med Mgmt 15 min		Methylprednisolone (as sodium	0
Locoid Lipocream		Med Mgmt 30 min		succinate)	
Locorten-Vioform		Med Mgmt 45 min		Methylprednisolone aceponate	
Lodoxamide		Medicine Delivery		Methylprednisolone acetate	
Logem		Medrol	00	Methylxanthines	
Lomide		Medroxyprogesterone acetate	00	Metoclopramide Actavis 10	
Lomustine		Genito-Urinary		Metoclopramide hydrochloride	
Loniten		Hormone		Metolazone	
Loperamide hydrochloride		Mefenamic acid		Metopirone	
Lopinavir with ritonavir		Melatonin		Metoprolol IV Mylan	5
Lopinavir/Ritonavir Mylan		Melpha		Metoprolol IV Viatris	
Loprofin		Melphalan		Metoprolol succinate	
Loprofin Mix		Meningococcal (groups A, C, Y		Metoprolol tartrate	
Lorafix		W-135) conjugate vaccine		Metrogyl	
Loratadine	250	Meningococcal B multicompone	ent	Metronidazole	10

Metyrapone	97	Minims Pilocarpine		Myleran	15
Mexiletine hydrochloride	51	Minims Prednisolone		Mylotarg	202
Miacalcic	87	Minirin	96	Myometrial and Vaginal Hormone	
Micolette	27	Minirin Melt	96	Preparations	84
Micolette-S29		Mino-tabs	102	Myozyme	2
Miconazole	33	Minocycline hydrochloride		- N -	
Miconazole nitrate		Minomycin	102	Nadolol	52
Dermatological	72	Minor Skin Infections		Nadolol BNM	52
Genito-Urinary	84	Minoxidil		Naglazyme	29
Micreme	84	Minoxidil Roma	59	Naloxone hydrochloride	264
Micreme H	74	Mirena	91	Naltraccord	15
Microgynon 30	83	Mirtazapine	132	Naltrexone AOP	15
Microlut		Misoprostol	9	Naltrexone hydrochloride	15
Midazolam	144	Mitomycin C	163	Naphazoline hydrochloride	
Midazolam-Baxter	144	Mitozantrone		Naphcon Forte	260
Midodrine	51	Mitozantrone Ebewe	163	Naprosyn SR 1000	
Mifegyne	86	Mixtard 30	11	Naprosyn SR 750	
Mifepristone		Moclobemide		Naproxen	
Milpharm		Modafinil		Narcaricin mite	
Minerals		Modavigil		Nasal Preparations	
Mini-Wright AFS Low Range		Moduretic	55	Natalizumab	
Mini-Wright Standard		Molaxole		Natulan	16
Minidiab		Molnupiravir		Nausafix	
MiniMed 1.8 Reservoir		Moments		Nausicalm	
MMT-326A	25	Mometasone furoate		Navelbine	
MiniMed 3.0 Reservoir		Monogen		Navelbine S29	
MMT-332A	25	Montelukast		Nefopam hydrochloride	
MiniMed 770G		Montelukast Mylan		Neisvac-C	
MiniMed Mio MMT-921A		Montelukast Viatris		Neo-Mercazole	
MiniMed Mio MMT-923A		Moroctocog alfa [Recombinant		Neocate Gold	
MiniMed Mio MMT-925A		VIII]		Neocate Junior Unflavoured	
MiniMed Mio MMT-941A		Morphine hydrochloride		Neocate Junior Vanilla	
MiniMed Mio MMT-943A		Morphine sulphate		Neocate SYNEO	
MiniMed Mio MMT-945A		Motetis		Neoral	
MiniMed Mio MMT-965A		Mouth and Throat		Neostigmine metilsulfate	
MiniMed Mio MMT-975A		Movapo		Nepafenac	
MiniMed Quick-Set MMT-386A		Moxifloxacin		Nepro HP (strawberry)	
MiniMed Quick-Set MMT-387A		MSUD Maxamum		Nepro HP (vanilla)	
MiniMed Quick-Set MMT-396A		Mucolytics		Nepro HP RTH	27
MiniMed Quick-Set MMT-397A		Mucosoothe		Neulactil	
MiniMed Quick-Set MMT-398A		Multiple Sclerosis Treatments		Neuraxpharm	
MiniMed Quick-Set MMT-399A		Multivitamin renal		NeuroTabs	
MiniMed Silhouette MMT-368A		Multivitamins		Nevirapine	
MiniMed Silhouette MMT-377A		Mupirocin		Nevirapine Alphapharm	114
MiniMed Silhouette MMT-378A		Muscle Relaxants		Nevirapine Viatris	114
MiniMed Silhouette MMT-381A		Mvite		Nicorandil	
MiniMed Silhouette MMT-382A		Myambutol		Nicotine	
MiniMed Silhouette MMT-383A		Mycobutin		Nifedipine	
MiniMed Silhouette MMT-384A MiniMed Sure-T MMT-864A		MycoNail		Nifuran Nilotinib	
		Mycophenolate mofetil			170
MiniMed Sure-T MMT-866A		Mydriacyl	202	Nilstat	00
MiniMed Sure-T MMT-874A		Mylan (12 hr release)		Alimentary	
MiniMed Sure-T MMT-876A		Mylan (24 hr release)	53 E4	Genito-UrinaryInfection	
MiniMed Sure-T MMT-884A MiniMed Sure-T MMT-886A		Mylan Atenolol		Nintedanib	
		Mylan Clomiphen		Ninteganio Nipent	
Minims Cyclopentolate	202	IVIVIALI ILAIV (24 FIT FEIEASE)	ეკ	NIDEUL	ID:

Nirmatrelvir with ritonavir	113	Nutrison Advanced Diason	272	One-Alpha S29	34
Nitrates	58	Nutrison Advanced Peptisorb	276	Opdivo	
Nitroderm TTS	58	Nutrison Concentrated	281	Ora-Blend	
Nitrofurantoin	118	Nutrison Energy	279	Ora-Blend SF	26
Nitrolingual Pump Spray		Nutrison Energy Multi Fibre		Ora-Plus	
Nivestim		Nutrison Multi Fibre		Ora-Sweet	
Nivolumab		Nutrison Standard RTH		Ora-Sweet SF	
Nodia		Nyefax Retard		Orabase	
Noflam 250		Nystatin		Oral and Enteral Feeds	
Noflam 500		Alimentary	33	Oralcon 30 ED	
Non-Steroidal Anti-Inflammatory		Genito-Urinary		Oratane	
Drugs	119	Infection		Ordine	
Nonacog gamma, [Recombinant	110	NZB Low Gluten Bread Mix		Orgran	
Factor IX]	41	- 0 -	202	Ornidazole	
Norethisterone	71	Obinutuzumab	212	Orphenadrine citrate	
Genito-Urinary	0.4	Obstetric Preparations		Ortho-tolidine	
				Oruvail SR	
Hormone		Ocicure			
Norflex		Ocrelizumab		Osmolite RTH	
Norfloxacin		Ocrevus	142	Other Endocrine Agents	
Noriday 28		Octocog alfa [Recombinant factor	44	Other Oestrogen Preparations	9
Norimin		VIII] (Advate)	41	Other Progestogen	
Norimin-1 28 Day		Octocog alfa [Recombinant factor		Preparations	
Normison		VIII] (Kogenate FS)		Other Skin Preparations	
Norpress		Octreotide		Other Supplements for PKU	
Nortriptyline hydrochloride		Octreotide Depot Teva		Otodex	259
Norvir		Octreotide GH		Ovestin	
Noumed Paracetamol		Octreotide long-acting		Genito-Urinary	
Noumed Pethidine		Oestradiol		Hormone	
Noumed Phenobarbitone		Oestradiol valerate	90	Oxaliplatin	
NovaSource Renal	275	Oestradiol with norethisterone	91	Oxaliplatin Accord	156
Novatretin	77	Oestriol		Oxaliplatin Actavis 100	
Novitium Sugar Free	30	Genito-Urinary	84	Oxaliplatin Ebewe	156
NovoMix 30 FlexPen	10	Hormone	91	Oxis Turbuhaler	25
NovoRapid	11	Oestrogens	90	Oxpentifylline	59
NovoRapid FlexPen	11	Ofev	254	Oxybutynin	
NovoRapid Penfill	11	Oil in water emulsion	75	Oxycodone hydrochloride	130
NovoSeven RT	41	Olanzapine13	8-139	Oxycodone Sandoz	130
Nozinan	138	Olaparib	163	OxyContin	130
Nozinan (Swiss)	138	Olbetam		OxyNorm	
Nozinan S29		Olbetam S29	<mark>56</mark>	Oxytocin	
Nucala	211	Olopatadine	263	Oxytocin BNM	84
Nuelin	255	Olopatadine Teva		Oxytocin GH	
Nuelin-SR	255	Olsalazine		Oxytocin Panpharma	
Nupentin		Omalizumab		Oxytocin with ergometrine	
Nusinersen		Omeprazole		maleate	8
Nutilis		Omeprazole actavis 10		Ozurdex	
Nutren Diabetes		Omeprazole actavis 20		- P -	
Nutrient Modules		Omeprazole actavis 40		Pacifen	124
Nutrini Energy Multi Fibre		Omnitrope		Pacimol	
Nutrini Energy RTH		Omnitrope S29		Paclitaxel	
Nutrini Low Energy Multi Fibre		Onbrez Breezhaler		Paclitaxel Actavis	
Nutrini Peptisorb		Oncaspar LYO		Paclitaxel Ebewe	
Nutrini Peptisorb Energy		OncoTICE		Paediatric Seravit	
Nutrini RTH	214	Ondansetron ODT DDI A		Palbociclib	
Nutrison 800 Complete Multi	070	Ondansetron ODT-DRLA		Paliperidone	139
Fibre	2/9	One-Alpha	34	Paliperidone palmitate	140

Palivizumab	214	Pexsig	53	Pneumococcal (PPV23)	
Pamidronate disodium		Pfizer Exemestane		polysaccharide vaccine	300
Pamisol		Pharmacy Services		Pneumovax 23	
Pamol		Pheburane	32	Podophyllotoxin	
Pancreatic enzyme		Phenasen		Polaramine	
Pantoprazole		Phenobarbitone		Poliomyelitis vaccine	
Panzop Relief		Phenobarbitone sodium		Poloxamer	
Papaverine hydrochloride		Extemporaneous	267	Poly-Gel	
Para-amino salicylic acid		Nervous		Poly-Tears	
Paracetamol	128	Phenoxybenzamine		Poly-Visc	
Paracetamol (Ethics)		hydrochloride	48	Polycal	
Paracetamol + Codeine		Phenoxymethylpenicillin (Penicillin		Polyethylene glycol 400 and	
(Relieve)	130	V)	102	propylene glycol	263
Paracetamol with codeine		Phenytoin sodium132		Ponstan	
Paraffin		Phillips Milk of Magnesia		Posaconazole	
Paraffin liquid with wool fat		Phlexy 10		Posaconazole Juno	
Parasiticidal Preparations		Phosphate Phebra		Potassium chloride	
Parnate		Phosphorus		Potassium citrate	
Paromomycin		Phytomenadione		Potassium iodate	
Paroxetine		Pilocarpine hydrochloride		Povidone iodine	
Paser		Pilocarpine nitrate		Pradaxa	
Paxam		Pimafucort		Pramipexole hydrochloride	
Paxlovid		Pimecrolimus		Pravastatin	
Pazopanib		Pine tar with trolamine laurilsulfate		Pravastatin Mylan	
Peak flow meter		and fluorescein	78	Pravastatin Viatris	
Pedialyte - Bubblegum		Pinetarsol		Praziquantel	98
Pediasure		Pioglitazone		Prazosin	
Pediasure Plus		Pirfenidone		Pred Forte	
Pediasure RTH		Pizotifen		Prednisolone	
Pegaspargase		PKU Anamix Infant		Prednisolone acetate	
Pegasys		PKU Anamix Junior		Prednisolone sodium	
Pegfilgrastim		PKU Anamix Junior Chocolate		Prednisolone sodium	
Pegylated interferon alfa-2a		PKU Anamix Junior LQ		phosphate	261
Pembrolizumab		PKU Anamix Junior Orange		Prednisolone-AFT	
Pemetrexed		PKU Anamix Junior Vanilla		Prednisone	
Penicillamine		PKU Build 20 Chocolate		Prednisone Clinect	
Penicillin G		PKU Build 20 Raspberry	200	Pregabalin	134
PenMix 30		Lemonade	290	Pregabalin Pfizer	
PenMix 50		PKU Build 20 Smooth		Pregnancy Tests - hCG Urine	
Pentasa		PKU Build 20 Vanilla		Premarin	
Pentostatin [Deoxycoformycin].		PKU GMPro Ultra Lemonade		Prevenar 13	
Pentoxifylline [Oxpentifylline]		PKU Lophlex LQ 10		Priadel	
Peptamen Junior		PKU Lophlex LQ 20		Primaquine	
Pepti-Junior		PKU Lophlex Powder		Primidone	
Perhexiline maleate		PKU Lophlex Sensation 20		Primidone Clinect	13/
Pericyazine		PKU sphere20 Banana		Primolut N	
Perindopril		PKU sphere20 Chocolate		Priorix	
Periset		PKU sphere20 Lemon		Probenecid	
Periset ODT		PKU sphere20 Red Berry	200	Probenecid-AFT	
		DKI Lephoro20 Vanilla	200	Procarbazine hydrochloride	
Perjeta		PKU sphere20 Vanilla			
Permethrin		Plaquenil Plendil ER		Prochlorperazine Proctofoam	
Perrigo			33	Proctosedyl	
		Pneumococcal (PCV10) conjugate	200	Proposition hydrophorida	100
Peteha Pethidine hydrochloride		vaccine Pneumococcal (PCV13) conjugate	233	Procyclidine hydrochloride Progesterone	120
			200		
Pevaryl	12	vaccine	255	Proglicem	I

Proglycem	10	Redipred	89	Roxane-Propranolol	52
Progynova	90	Relieve		Roxithromycin	100
Prolia		Relistor	26	Rubifen	
Promethazine hydrochloride	250	Remicade	203	Rubifen SR	147
Propafenone hydrochloride		Renilon 7.5	275	Rugby Capsaicin Topical Cream	
Propamidine isethionate		Resonium-A	47	Musculoskeletal	120
Propranolol		Resource Beneprotein	271	Nervous	12
Propylene glycol		Respigen		Rurioctocog alfa pegol [Recombin	
Propylthiouracil		Respiratory Devices		factor VIII]	
Prostacur		Respiratory Stimulants		Ruxolitinib	
Protaphane		Retinol palmitate		Rythmodan	
Protaphane Penfill		ReTrieve		Rythmodan - Cheplafarm	
Protifar		Retrovir		Rytmonorm	
Protionamide		Revlimid		-S-	
Provera		Revolade		Sabril	13
Provera HD		Riboflavin		Sacubitril with valsartan	
Psoriasis and Eczema		Ribomustin		Sagent	
Preparations	77	Ricit		SalAir	
PTU		Rifabutin		Salazopyrin	
Pulmicort Turbuhaler		Rifadin		Salazopyrin EN	
Pulmozyme		Rifampicin		Salbutamol	
Puri-nethol		Rifaximin		Salbutamol with ipratropium	204
Puria		Rifinah		bromide	05
	34			Coliculia soid	200
Puritan's Pride Vitamin	01	Rilutek		Salicylic acid	
B-2 100 mg		Riluzole		Salmeterol	
Pyrazinamide		RINVOQ		Sandomigran	
Pyridostigmine bromide		Riodine		Sandostatin LAR	
Pyridoxine hydrochloride		Risdiplam		Sanofi Primaquine	
Pyridoxine multichem		Risedronate Sandoz		Sapropterin dihydrochloride	
Pyrimethamine		Risedronate sodium		Scalp Preparations	
Pytazen SR	42	Risperdal Consta		Scopoderm TTS	
- Q -		Risperidone		Sebizole	
Quantalan sugar free		Risperidone (Teva)		Secukinumab	
Quetapel		Risperon		Sedatives and Hypnotics	
Quetiapine	138	Ritalin		Seebri Breezhaler	
Quick-Set MMT-392	<mark>24</mark>	Ritalin LA	148	Senna	
Quick-Set MMT-393		Ritonavir	116	Senokot	2
Quinapril	49	Rituximab (Mabthera)	215	SensoCard	1
Quinapril with		Rituximab (Riximyo)	217	Serc	136
hydrochlorothiazide	49	Rivaroxaban	45	Serenace	138
Qvar	251	Rivastigmine	150	Seretide	252
- R -		Rivastigmine Patch BNM 10	150	Seretide Accuhaler	
RA-Morph	129	Rivastigmine Patch BNM 5	150	Serevent	25
Ralicrom	8	Rivotril		Serevent Accuhaler	25
Raloxifene hydrochloride	121	Riximyo	217	Sertraline	132
Raltegravir potassium		RIXUBIS	41	Setrona	132
Ramipex	125	Rizamelt	135	Sevredol	
Ramipril		Rizatriptan		Sex Hormones Non	
Ranbaxy-Cefaclor		Robinul		Contraceptive	89
Rapamune		Ronapreve		Shingles vaccine	30
Rasagiline		Ropin		Shingrix	
Reandron 1000		Ropinirole hydrochloride	125	SII-Onco-BCG	
Recombinant factor IX		Rosuvastatin	57	Sildenafil	
Recombinant factor VIIa		Rosuvastatin Viatris	57	Silhouette MMT-373	2,
Recombinant factor VIII		Rotarix		Siltuximab	51
Rectogesic		Rotavirus oral vaccine		Simvastatin	200
1 10010g0310	U	i iotaviius olai valtiile		JVaJlatii1	

				_
Simvastatin Mylan58		253	Taliglucerase alfa	
Simvastatin Viatris58		253	Tambocor	
Sinemet			Tamoxifen citrate	
Sinemet CR	Sporanox	106	Tamoxifen Sandoz	
Sirolimus245	Sprycel	168	Tamsulosin hydrochloride	8
Sirturo108	Stelara	236	Tamsulosin-Rex	8
Siterone89	Stemetil	137	Tandem Cartridge	2
Slow-Lopresor52	Steril-Gene	90	Tandem t:slim X2 with Basal-IQ	
Smith BioMed Rapid Pregnancy	SteroClear	257	Tap water	26
Test85	Stesolid	132	Taro	
Sodibic47	Stimulants/ADHD Treatments	146	Taro-Testosterone	8
Sodium acid phosphate27	Stiripentol	134	Tasigna	170
Sodium alginate6			Tasmar	12
Sodium benzoate31		33	Taurine	
Sodium bicarbonate	Strattera		Tecentrig	240
Blood46–47			Tecfidera	
Extemporaneous267			Tegretol	
Sodium calcium edetate265			Tegretol CR	
Sodium chloride	Sulfadiazine Silver		Telfast	25
Blood46			Teligent	
Respiratory257			Temaccord	
Sodium citrate with sodium lauryl	Sulphur		Temazepam	
sulphoacetate			Temozolomide	
Sodium citro-tartrate			Tenofovir disoproxil	
Sodium cromoglicate	Sumatriptan		Tenofovir Disoproxil Emtricitabine	
Alimentary8			Mylan	113
Sensory261			Tenofovir Disoproxil Emtricitabine	
Sodium Fusidate [fusidic acid]	Sunscreens		Viatr	113
Dermatological			Tenofovir Disoproxil Mylan	
Infection104			Tenofovir Disoproxil Viatris	
Sensory259			Tenoxicam	
Sodium hyaluronate [Hyaluronic	Survimed OPD		Tensipine MR10	
acid]263			Tepadina	
Sodium phenylbutyrate32			Terbinafine	
Sodium picosulfate		279	Terbutaline sulphate	
Sodium polystyrene sulphonate47			Teriflunomide	
Sodium tetradecyl sulphate			Teriparatide	
Sodium valproate134	Symbicort Turbuhaler 100/6		Testosterone	
Sofradex			Testosterone cipionate	Q(
Soframycin259			Testosterone esters	
Solgar28–32			Testosterone undecanoate	
Solifenacin Mylan86			Tetrabenazine	
Solifenacin succinate	, ,		Tetrabromophenol	
Solifenacin Viatris			Tetracosactrin	
Solu-Cortef88			Tetracycline	
Solu-Medrol89	•		Teva Lisinopril	
Solu-Medrol-Act-O-Vial			Thalidomide	
			Thalomid	
Somatropin (Omnitrope)				
Sotalol	•		Theophylline	
Spacer device	Syrup (pharmaceutical grade)		Thiamine hydrochloride Thiamine multichem	
Span-K46		203	THIO-TEPA	
Spinol Muscular Atrophy 145				
Spinal Muscular Atrophy145		70	Thioguanine	
Spinraza			Thiotepa	15
Spiolto Respirat			Thyroid and Antithyroid Agents	9
Spiractin55	Tacrolimus Sandoz	247	Ticagrelor	47

Ticagrelor Sandoz	42	Trimethoprim	105	Vebulis	67
Tilcotil	119	Trimethoprim with		Vedafil	64
Timolol	261	sulphamethoxazole		Vedolizumab	238
Timoptol XE	261	[Co-trimoxazole]	105	Veletri	66
Tiotropium bromide		Trisequens		Venclexta	166
Tiotropium bromide with		Trisul		Venetoclax	166
olodaterol	254	Trophic Hormones		Venlafaxine	132
Tivicay		Tropicamide		Venomil2	
Tixagevimab with cilgavimab		Trulicity		VENOX	
TMP		Trusopt		Ventolin	252
Tobramycin		TruSteel		Ventolin Nebules	252
Infection	104	Tryzan	49	Vepesid	161
Sensory	259	Tuberculin PPD [Mantoux] test	301	Verapamil hydrochloride	54
Tobramycin BNM	104	Tubersol		Vermox	
Tobramycin Mylan		Two Cal HN	281	Versacloz	138
Tobrex		Tykerb		Vesanoid	
Tocilizumab		Tysabri		Vexazone	
Tofranil		- U -		Vfend	
Tolcapone		UK Synacthen	89	Viaderm KC	
Tolvaptan		Ultibro Breezhaler		Victoza	
Topamax		Ultraproct		Vigabatrin	
Topical Products for Joint and		Umeclidinium		Vigisom	
Muscular Pain	120	Umeclidinium with vilanterol		Vildagliptin	
Topiramate		Univent2		Vildagliptin with metformin	
Topiramate Actavis		Upadacitinib		hydrochloride	12
Total parenteral nutrition (TPN)		Ural		Vimpat	
TPN		Urea		Vinblastine sulphate	
Tramadol hydrochloride		Urex Forte		Vincristine sulphate	
Tramal SR 100		Urinary Agents		Vinorelbine	
Tramal SR 150		Urinary Tract Infections		Vinorelbine Ebewe	
Tramal SR 200		Urinorm		Vinorelbine Te Arai	
Trandate		Uromitexan		Viriamune Suspension	
Tranexamic acid		Ursodeoxycholic acid		ViruPOS	
Tranylcypromine sulphate		Ursosan		Vit.D3	
Trastuzumab		Ustekinumab		Vita-B12	
Trastuzumab emtansine		Utrogestan		VitA-POS	
Travatan		- V -		Vitabdeck	
Travoprost		Vaccinations	201	Vital	
Treatments for Dementia		Vaclovir		Vitamin B complex	
Treatments for Substance		Valaciclovir		Vitamin B6 25	
Dependence	150	Valganciclovir		Vitamins	
Trental 400		Valganciclovir Mylan		Vivonex TEN	
Tretinoin		Valganciclovir Viatris		Voltaren	
Dermatological	71	Vancomycin		Voltaren D	
Oncology		Vannair		Voltaren Ophtha	
Trexate		Varenicline Pfizer		Voltaren SR	
Triamcinolone acetonide	139	Varenicline tartrate		Volumatic	
	20		132		
Alimentary		Varicella vaccine [Chickenpox	201	Voriconazole	
Dermatological		Varicella zostar vaccina (Shingle		Votrient	
	89	Varicella zoster vaccine [Shingle		Vttack	106
Triamcinolone acetonide with	ototin	vaccine]			45
gramicidin, neomycin and ny		Various		Warfarin sodium	
Dermatological		Varivax		Wart Preparations	
Sensory		Vasodilators		Wasp venom allergy treatment	200
Triazolam		Vasopressin Agonists		Water	40
Trikafta	∠50	Vasorex	ექ	Blood	46

Extemporaneous	268
White Soft Liquid Paraffin AFT	75
Wool fat with mineral oil	10 75
- X -	/ 5
	45
Xarelto	
Xifaxan	10
XMET Maxamum	. 283
Xolair	
XP Maxamum	
Xylocaine	. 126
Xylocaine 2% Jelly	. 126
Xyntha	41
- Z -	
Zapril	48
Zarontin	. 133
Zaroxolyn	55
Zavedos	162
Zeffix	110
Zematop	78
Zetlam	110
Ziagen	115
Zidovudine [AZT]	115
Zidovudine [AZT] with	. 110
lamivudine	445
iamivudine	. 115
Ziextenzo	45
Zimybe	58
Zinc and castor oil	74
Zinc sulphate	37
Zincaps	37
Zinnat	98
Ziprasidone	
Zista	. 250
Zithromax	99
Zoledronic acid	
Hormone	88
Musculoskeletal	.123
Zoledronic acid Mylan	
Zoledronic acid Viatris	
Hormone	88
Musculoskeletal	123
Zoledronic-US	123
Zopiclone	1/15
Zopiclone Actavis	
Zostrix	
Zostrix HP	. 127
Zuclopenthixol decanoate	. 140
Zuclopenthixol hydrochloride	. 139
Zusdone	. 139
Zyban	. 151
Zypine	. 138
Zypine ODT	. 138
Zyprexa Relprevv	. 139
Zytiga	. 174
Zyvox	.109