5

| May 2024 Volume 31 |
|--------------------------------|
| Editors: |
| Kaye Wilson & Ayeshah Khan |
| email: enquiry@pharmac.govt.nz |
| Telephone +64 4 460 4990 |
| Level 9, 40 Mercer Street |
| PO Box 10 254 Wellington |
| Freenhone Information Line |

Freephone Information Line 0800 66 00 50 (9am - 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and TFX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth

CC

email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency

BY

ISSN 1179-3686 (i)

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

| | Index | 308 |
|-----------|---------------------------------------|-----|
| Section I | National Immunisation Schedule | 294 |
| Section D | Special Foods | 271 |
| Section C | Extemporaneous Compounds (ECPs) | 268 |
| | Various | 265 |
| | Sensory Organs | 260 |
| | Respiratory System & Allergies | 249 |
| | choology rigonio a minanocappicocanio | |

Section A General Rules

| Section B | Alimentary Tract & Metabolism | 6 |
|-----------|--------------------------------------|-----|
| | Blood & Blood Forming Organs | 36 |
| | Cardiovascular System | 46 |
| | Dermatologicals | 68 |
| | Genito Urinary System | 77 |
| | Hormone Preparations – Systemic | 84 |
| | Infections – Agents For Systemic Use | 95 |
| | Musculoskeletal System | 116 |
| | Nervous System | 123 |
| | Oncology Agents & Immunosuppressants | 153 |
| | Respiratory System & Allergies | 249 |
| | Sensory Organs | 260 |
| | Various | 265 |
| | | |
| Section C | Extemporaneous Compounds (ECPs) | 268 |
| Section D | Special Foods | 271 |
| | | |
| Section I | National Immunisation Schedule | 294 |
| | | |
| | | |

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ Hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

| gram g | |
|-----------------------|--|
| kilogram kg | |
| international unit iu | |

Abbreviations

| Capsule Cream Device Dispersible Effervescent Emulsion | Amp Cap Crm Dev Disp Eff Emul EC |
|---|---|
| Enteric Coated | EC |
| | |

| microgram me | cg |
|--------------|----|
| milligramn | ng |
| millilitreı | ml |

| millimole | mmol |
|-----------|------|
| unit | u |

| Gelatinous | Gel | SolutionSoln |
|-------------|------|-----------------------|
| Granules | Gran | SuppositorySupp |
| Infusion | Inf | TabletTab |
| Injection | Inj | Tincture Tinc |
| Liquid | Liq | Trans Dermal Delivery |
| Long Acting | LA | SystemTDDS |
| Ointment | Oint | - |
| Sachet | Sach | |

Read the General Rules : https://pharmac.govt.nz/section-a.

SECTION B: ALIMENTARY TRACT AND METABOLISM

| | Subsidy (Manufacturer's Price) | | Fully ubsidised | Brand or Generic |
|--|--|------------------|--------------------|--|
| | \$ | Per | | Manufacturer |
| Antacids and Antiflatulents | | | | |
| Antacids and Reflux Barrier Agents | | | | |
| LGINIC ACID | | | | |
| Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet | | 30 | 1 | Gaviscon Infant |
| ODIUM ALGINATE | | | | |
| Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour | 1.80 | 60 | | |
| 0 1 11 | (13.61) | | | Gaviscon Extra Strength |
| Oral liq 500 mg with sodium bicarbonate 267 mg and calcium | n | | | Strength |
| carbonate 160 mg per 10 ml | 1.50 (7.50) | 500 ml | | Acidex |
| Dhaan hata Dinding Aganta | (1.00) | | | |
| Phosphate Binding Agents | | | | |
| LUMINIUM HYDROXIDE € Tab 600 mg | | 100 | 1 | Alu-Tab |
| ALCIUM CARBONATE | | | | |
| Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement | 39.00 | 500 ml | 1 | Roxane |
| | | | | |
| | 47.30 | 473 ml | | Calcium carbonate PAI S29 |
| Only when prescribed for patients unable to swallow calor inappropriate and the prescription is endorsed according Antidiarrhoeals | cium carbonate table | | 1 | PAI S29 |
| inappropriate and the prescription is endorsed according | cium carbonate table | | 1 | PAI S29 |
| inappropriate and the prescription is endorsed according Antidiarrhoeals | cium carbonate table | | ✔ ere calci | PAI 329 um carbonate tablets ar |
| inappropriate and the prescription is endorsed according Anticliarrhoeals Agents Which Reduce Motility OPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a Tab 2 mg | cium carbonate table ly. a PSO | ts or wh | ✓ ere calci | PAI 529 um carbonate tablets ar Nodia |
| inappropriate and the prescription is endorsed according Antidiarrhoeals Agents Which Reduce Motility OPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a Tab 2 mg | cium carbonate table ly. a PSO | ts or wh | ✓ ere calci | PAI 329 um carbonate tablets ar |
| inappropriate and the prescription is endorsed according Antidiarrhoeals Agents Which Reduce Motility OPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a Cap 2 mg Rectal and Colonic Anti-inflammatories | cium carbonate table ly. a PSO | ts or wh | ✓ ere calci | PAI 529 um carbonate tablets ar Nodia |
| inappropriate and the prescription is endorsed according Antidiarrhoeals Agents Which Reduce Motility OPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a Cap 2 mg Cap 2 mg Rectal and Colonic Anti-inflammatories UDESONIDE | a PSO | ts or wh | ✓ ere calci | PAI 529 um carbonate tablets ar Nodia |
| inappropriate and the prescription is endorsed according Antidiarrhoeals Agents Which Reduce Motility OPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a Cap 2 mg Rectal and Colonic Anti-inflammatories | a PSO | ts or wh | ✓ ere calci | PAI 529 um carbonate tablets ar Nodia |
| inappropriate and the prescription is endorsed according Antidiarrhoeals Agents Which Reduce Motility OPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a | cium carbonate table ily. a PSO 10.75 7.25 | 400 400 90 | vere calci | PAI 529 um carbonate tablets ar Nodia <u>Diamide Relief</u> Budesonide Te Arai |
| inappropriate and the prescription is endorsed according Antidiarrhoeals Agents Which Reduce Motility OPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a | cium carbonate table ily. a PSO | 400 400 90 | vere calci | PAI 529 um carbonate tablets ar Nodia <u>Diamide Relief</u> Budesonide Te Arai |

continued...

| Subsidy | Full | y Brand or | |
|------------------------|-----------|--------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:

- 3.1 Diabetes; or
- 3.2 Cushingoid habitus; or
- 3.3 Osteoporosis where there is significant risk of fracture; or
- 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
- 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).
- Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

| HYDROCORTISONE AGETATE | | | |
|---|------------|---------|------------------------------|
| Rectal foam 10%, CFC-Free (14 applications) | | 15 g OP | Colifoam |
| HYDROCORTISONE ACETATE WITH PRAMOXINE HYDR | ROCHLORIDE | | |
| Topical aerosol foam, 1% with pramoxine hydrochloride | e 1%26.55 | 10 g OP | Proctofoam S29 |
| MESALAZINE | | | |
| Tab 400 mg | | 100 | Asacol |
| Tab long-acting 500 mg | | 100 | Pentasa |
| Tab 800 mg | | 90 | Asacol |
| - | | | Asacol S29 S29 |
| Modified release granules, 1 g | 118.10 | 100 OP | Pentasa |
| Enema 1 g per 100 ml | | 7 | Pentasa |
| Suppos 500 mg | | 20 | Asacol |
| Suppos 1 g | | 28 | Pentasa |
| | | | |

| | Subsidy | | Fully | |
|--|------------------------------|--------|-------------|----------------------------|
| | (Manufacturer's Price) \$ | Per | Subsidised | I Generic Manufacturer |
| OLSALAZINE | | | | |
| Tab 500 mg | 56.02 | 60 | 1 | Atnahs Olsalazine S29 |
| | 93.37 | 100 | 1 | Dipentum |
| Cap 250 mg | | 100 | | Dipentum |
| PREDNISOLONE SODIUM | | | | |
| Rectal foam 20 mg per dose (14 applications) | 74.10 | 1 OP | | Essential |
| | | | | Prednisolone S29 |
| SODIUM CROMOGLICATE Cap 100 mg | 112.25 | 100 | | Ralicrom |
| SULFASALAZINE | 113.35 | 100 | v | Railcroin |
| SULFASALAZINE ★ Tab 500 mg | 16.52 | 100 | 1 | Salazopyrin |
| * Tab EC 500 mg | | 100 | | Salazopyrin EN |
| Local preparations for Anal and Rectal Disorde | *0 | | | |
| Local preparations for Anal and Rectal Disorde | 15 | | | |
| Antihaemorrhoidal Preparations | | | | |
| LUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PI | ALATE AND CINCH | OCAI | NE | |
| Oint 950 mcg, with fluocortolone pivalate 920 mcg, and | | | | |
| cinchocaine hydrochloride 5 mg per g | | 30 g C | P 🗸 | Ultraproct |
| Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg | 7 20 | 12 | 1 | Ultraproct |
| HYDROCORTISONE WITH CINCHOCAINE | | 12 | • | onapioer |
| Oint 5 mg with cinchocaine hydrochloride 5 mg per g | | 30 g O | P 🗸 | Proctosedyl |
| Suppos 5 mg with cinchocaine hydrochloride 5 mg per g | | 12 | 1 | Proctosedyl |
| Management of Anal Fissures | | | | |
| GLYCERYL TRINITRATE - Special Authority see SA1329 below | w – Retail pharmacy | | | |
| * Oint 0.2% | | 30 g O | P 🗸 | Rectogesic |
| SA1329 Special Authority for Subsidy | | | | |
| nitial application from any relevant practitioner. Approvals vali | | ewal u | nless notif | fied where the patient has |
| chronic anal fissure that has persisted for longer than three week | (S. | | | |
| Antispasmodics and Other Agents Altering Gui | t Motility | | | |
| GLYCOPYRRONIUM BROMIDE | | | | |
| Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available or | na | | | |
| PSO | | 5 | 1 | Robinul |
| HYOSCINE BUTYLBROMIDE | | | | |
| * Tab 10 mg | | 100 | | Buscopan |
| Inj 20 mg, 1 ml – Up to 5 inj available on a PSO | 1.91 | 5 | ~ | Spazmol |
| MEBEVERINE HYDROCHLORIDE | 8 50 | 90 | 1 | Colofac |
| | | 50 | • | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|---------------|---------------------|---|
| Antiulcerants | | | | |
| Antisecretory and Cytoprotective | | | | |
| MISOPROSTOL – Wastage claimable * Tab 200 mcg – Up to 120 tab available on a PSO | 47.73 | 120 | √ (| Cytotec |
| Helicobacter Pylori Eradication | | | | |
| CLARITHROMYCIN Tab 500 mg – Subsidy by endorsement a) Maximum of 28 tab per prescription b) Subsidised only if prescribed for helicobacter pylori en Note: the prescription is considered endorsed if clarit inhibitor and either amoxicillin or metronidazole. | radication and presc | | is endorse | |
| H2 Antagonists | | | | |
| FAMOTIDINE – Only on a prescription * Tab 20 mg | 4.91 | 100 | √ I | Famotidine Hovid ©29 |
| * Tab 40 mg | 10.32 | 100 | √ i | Famotidine Hovid S29 |
| Inj 10 mg per ml, 4 ml – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients received | | 10 t of pa | | Mylan ^{S29} Ə. |
| Proton Pump Inhibitors | | | | |
| LANSOPRAZOLE * Cap 15 mg * Cap 30 mg OMEPRAZOLE For omeprazole suspension refer Standard Formulae, page 2 | 5.26 | 100 100 | - | <u>Lanzol Relief</u> Lanzol Relief |
| Cap 10 mg | 2.06 | 90 | ✓ <u>(</u> | <u>Omeprazole actavis</u> 10 |
| Cap 20 mg | 2.02 | 90 | ✓ <u>(</u> | Omeprazole actavis 20 |
| Cap 40 mg | 3.18 | 90 | ✓ <u>(</u> | <u>Omeprazole actavis</u> 40 |
| Powder – Only in combination Only in extemporaneously compounded omeprazole suspansion | | 5 g | √ I | Midwest |
| Inj 40 mg ampoule with diluent | | 5 | - | <u>Dr Reddy's</u> <u>Omeprazole</u> Dcicure s29 |
| PANTOPRAZOLE * Tab EC 20 mg * Tab EC 40 mg | | 90 90 | | Panzop Relief Panzop Relief |
| Site Protective Agents | | | | |
| COLLOIDAL BISMUTH SUBCITRATE Tab 120 mg | 14.51 | 50 | v (| Gastrodenol S29 |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|-----------------------|---|-----|---------------------|-------------------------------------|
| SUCRALFATE Tab 1 g | | 120 | | Carafate |

Bile and Liver Therapy

⇒SA1461 Special Authority for Subsidy

Initial application only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

56

✓ Xifaxan

Renewal only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Diabetes

Hyperglycaemic Agents

| DIAZOXIDE - Special Authority see SA1320 below - Retail pl | harmacy | | |
|--|---------|----------|-----------------------------------|
| Cap 25 mg | | 100 | Proglicem S29 |
| Cap 100 mg | | 100 | Proglicem S29 |
| Oral liq 50 mg per ml | 620.00 | 30 ml OP | Proglycem S29 |
| | | | e5 Pharma S29 |

⇒SA1320 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains

appropriate and the patient is benefiting from treatment.

GLUCAGON HYDROCHLORIDE

| Inj 1 mg syringe kit – Up to 5 kit available on a PSO | 1 | Glucagen Hypokit |
|---|----------|---|
| Insulin - Short-acting Preparations | | |
| INSULIN NEUTRAL ▲ Inj human 100 u per ml25.26 | 10 ml OP | ✓ Actrapid ✓ Humulin B |
| ▲ Ini human 100 u per ml. 3 ml | 5 | Actrapid Penfill |

Insulin - Intermediate-acting Preparations

| INSULIN ASPART WITH INSULIN ASPART PROTAMINE | | |
|---|----------|--|
| ▲ Inj 100 iu per ml, 3 ml prefilled pen | 5 | NovoMix 30 FlexPen |
| INSULIN ISOPHANE ▲ Inj human 100 u per ml17.68 | 10 ml OP | ✓ Humulin NPH |
| , | | Protaphane |
| ▲ Inj human 100 u per ml, 3 ml | 5 | Humulin NPH |
| | | Protaphane Penfill |

✓ Humulin R

| | Cubaidu | | Fully Brand ar |
|--|-------------------------|----------------------------------|---|
| | Subsidy | () () () () () () | Fully Brand or |
| | (Manufacturer's F \$ | Price) Subs | sidised Generic Manufacturer |
| | ψ | Fei | |
| NSULIN ISOPHANE WITH INSULIN NEUTRAL | | | |
| Inj human with neutral insulin 100 u per ml | | 10 ml OP | Humulin 30/70 |
|] · · · · · · · · · · · · · · · · · · · | | | Mixtard 30 |
| Inj human with neutral insulin 100 u per ml, 3 ml | 12 66 | 5 | ✓ Humulin 30/70 |
| | 42.00 | 5 | |
| | | | PenMix 30 |
| | | | PenMix 50 |
| NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE | | | |
| Inj lispro 25% with insulin lispro protamine 75% 100 u per n | nl | | |
| 3 ml | | 5 | Humalog Mix 25 |
| | | 5 | |
| ▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per n | | _ | • • • • • • • • • • |
| 3 ml | | 5 | Humalog Mix 50 |
| | | | |
| Insulin - Long-acting Preparations | | | |
| NSULIN GLARGINE | | | |
| Inj 100 u per ml, 10 ml | 63.00 | 1 | Lantus |
| Inj 100 u per ml, 3 ml | | 5 | ✓ Lantus |
| Inj 100 u per ml, 3 ml disposable pen | | 5 | ✓ Lantus SoloStar |
| | | 5 | |
| Insulin - Rapid Acting Preparations | | | |
| | | | |
| NSULIN ASPART | | | A N A N N |
| Inj 100 u per ml, 10 ml | | 1 | NovoRapid |
| Inj 100 u per ml, 3 ml | | 5 | NovoRapid Penfill |
| Inj 100 u per ml, 3 ml syringe | 51.19 | 5 | NovoRapid FlexPen |
| NSULIN GLULISINE | | | |
| Inj 100 u per ml, 10 ml | 27.02 | 1 | 🖌 Anidra |
| | | | ✓ Apidra |
| Inj 100 u per ml, 3 ml | | 5 | ✓ Apidra |
| Inj 100 u per ml, 3 ml disposable pen | | 5 | Apidra SoloStar |
| NSULIN LISPRO | | | |
| Inj 100 u per ml, 10 ml | 34 92 | 10 ml OP | Humalog |
| Inj 100 u per ml, 3 ml | | 5 | ✓ Humalog |
| | | 5 | • Humaiog |
| Alpha Glucosidase Inhibitors | | | |
| CARBOSE | | | |
| | 0.05 | 00 | Accest |
| € Tab 50 mg | | 90 | ✓ <u>Accarb</u> |
| F Tab 100 mg | | 90 | Accarb |
| Oral Hypoglycaemic Agents | | | |
| | | | |
| LIBENCLAMIDE | | | |
| € Tab 5 mg | 7.50 | 100 | Daonil |
| LICLAZIDE | | | |
| ← Tab 80 mg | 20.10 | 500 | Glizide |
| - | 20.10 | | <u></u> |
| LIPIZIDE | | | |
| | | 100 | Minidiab |
| 🗧 Tab 5 mg | 4.58 | | |
| - | 4.58 | | |
| IETFORMIN HYDROCHLORIDE | | | Metformin Viatris |
| Tab 5 mg IETFORMIN HYDROCHLORIDE Tab immediate-release 500 mg Tab immediate-release 850 mg | 14.74 | 1,000 500 | ✓ <u>Metformin Viatris</u> ✓ Metformin Viatris |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | Fully | Brand or |
|--|---------------------------|-------|-------------|----------------------------|
| | (Manufacturer's Price) | _ | Subsidised | |
| | \$ | Per | | Manufacturer |
| PIOGLITAZONE | | | | |
| * Tab 15 mg | 6.80 | 90 | ✓ | Vexazone |
| * Tab 30 mg | 7.30 | 90 | | Vexazone |
| * Tab 45 mg | | 90 | ~ | Vexazone |
| VILDAGLIPTIN | | | | |
| Tab 50 mg | | 60 | ✓ | Galvus |
| VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE | | | | |
| Tab 50 mg with 1,000 mg metformin hydrochloride | | 60 | ✓ | Galvumet |
| Tab 50 mg with 850 mg metformin hydrochloride | | 60 | ✓ | Galvumet |
| | | | | |
| GLP-1 Agonists | | | | |
| DULAGLUTIDE - Special Authority see SA2338 below - Retai | l pharmacy | | | |
| Note: Not to be given in combination with a funded SGLT-2 | 2 inhibitor or other GLP | -1 ag | jonist. | |
| Inj 1.5mg per 0.5 ml prefilled pen | | 4 | ✓ | Trulicity |
| ► SA2338 Special Authority for Subsidy | | | | |
| Note: Subsidy for patients with existing approvals prior to 1 Ma | y 2024. Approvals vali | d wit | hout furthe | r renewal unless notified. |
| No new patients will be granted from 1 May 2024 until further no | | | | |
| LIRAGLUTIDE - Special Authority see SA2339 below - Retail | pharmacy | | | |
| a) Maximum of 9 inj per prescription | | | | |
| b) | | | | |
| a) Not to be given in combination with a funded SGL | T-2 inhibitor or other GI | _P-1 | agonist. | |
| b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefille | ed pens will be funded | per n | nonth. | |
| Inj 6 mg per ml, 3 ml prefilled pen | | 3 | ✓ | Victoza |

⇒SA2339 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN - Special Authority see SA2068 on the previous page - Retail pharmacy

| Note: Not to be given in combination with a funded GLP-1 agonisi | i. |
|--|----|
| T 1 10 | |

| * | Tab 10 mg | 58.56 | 30 | Jardiance |
|---|-----------|-------|----|-------------------------------|
| * | Tab 25 mg | 58.56 | 30 | Jardiance |

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – Special Authority see SA2068 on the previous page – Retail pharmacy

| | Note: Not to be given in combination with a funded GLP-1 ac | jonist. | | |
|---|---|---------|----|-------------------------------|
| * | Tab 5 mg with 1,000 mg metformin hydrochloride | | 60 | Jardiamet |
| * | Tab 5 mg with 500 mg metformin hydrochloride | | 60 | Jardiamet |
| * | Tab 12.5 mg with 1,000 mg metformin hydrochloride | | 60 | Jardiamet |
| * | Tab 12.5 mg with 500 mg metformin hydrochloride | | 60 | Jardiamet |
| | | | | |

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or

5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Test strips......15.50

.15.50 10 strip OP

KetoSens

| | Subsidy (Manufacturer's Price) \$ | | Fully lised | Brand or Generic Manufacturer |
|---|--|---|---------------------------------|--|
| Dual Blood Glucose and Blood Ketone Testing | | | | |
| UAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A dual blood glucose and blood ketone diagnostic test m 1) type 1 diabetes; or 2) permanent neonatal diabetes; or 3) undergone a pancreatectomy; or 4) cystic fibrosis-related diabetes; or 5) metabolic disease or epilepsy under the care of a p The prescription must be endorsed accordingly. Only 1 | eter is subsidised for paediatrician, neurolog meter per patient will | a patient wh gist or metal be subsidise | no has: polic sp ed (no i | ecialist. 'epeat prescriptions). Fo |
| the avoidance of doubt patients who have previously rec funded CareSens meter. | | , other than | Careo | ens, are eligible for a |
| Meter with 50 lancets, a lancing device and 10 blood glucos diagnostic test strips | | 1 OP | / C: | areSens Dual |
| 5 | | 101 | • • | |
| Blood Glucose Testing | | | | |
| LOOD GLUCOSE DIAGNOSTIC TEST METER – Subsidy by a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A diagnostic blood glucose test meter is subsidised for a is receiving insulin or sulphonylurea therapy; or is pregnant with diabetes; or is on home TPN at risk of hypoglycaemia or hyperg has a genetic or an acquired disorder of glucose he syndrome. The prescription must be endorsed accordingly. Only or prescriptions). Patients already using the CareSens N F meter, unless they have: | patient who: Jlycaemia; or omeostasis, excluding ne CareSens meter pe | er patient wi | I be su | bsidised (no repeat |

| | Subsidy | <u> </u> | Fully | Brand or |
|---|---------------------------|-----------------|-------------|-------------------------|
| | (Manufacturer's Pri \$ | ce) Subs Per | idised | Generic Manufacturer |
| | et eveileble en e DO | | - | |
| BLOOD GLUCOSE DIAGNOSTIC TEST STRIP – Up to 50 te The number of test strips available on a prescription is res | |) | | |
| Prescribed for a patient on insulin or a sulphonylurea | | rdinaly Phar | maciete | may annotate the |
| prescription as endorsed where there exists a record | | | | |
| Prescribed on the same prescription as insulin or a s endorsed; or | | | | |
| 3) Prescribed for a pregnant woman with diabetes and | | | | |
| Prescribed for a patient on home TPN at risk of hypo | | | | |
| 5) Prescribed for a patient with a genetic or an acquired | | homeostasis | excludin | ng type 1 or type |
| 2 diabetes and metabolic syndrome and endorsed ad | ccordingly. | | | |
| Test strips | | 50 test OP | | reSens N reSens PRO |
| | | | • 04 | |
| LOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED) The number of test strips available on a prescription is res | tricted to 50 unless: | | | |
| Prescribed for a patient on insulin or a sulphonylurea | | rdingly. Phar | macists | may annotate the |
| prescription as endorsed where there exists a record | | | | |
| Prescribed on the same prescription as insulin or a s endorsed; or | ulphonylurea in whic | h case the pr | escriptio | n is deemed to be |
| 3) Prescribed for a pregnant woman with diabetes and | endorsed accordingl | y; or | | |
| 4) Prescribed for a patient on home TPN at risk of hypo | | | | |
| 5) Prescribed for a patient with a genetic or an acquired | | homeostasis | excludin | ng type 1 or type |
| 2 diabetes and metabolic syndrome and endorsed ad | coraingly. | | | |
| Blood glucose test strips | | 50 test OP | ✔ Se | ensoCard |
| Insulin Syringes and Needles | | | | |
| Subsidy is available for disposable insulin syringes, needles, a | nd pen needles if pre | escribed on th | ne same | form as the one used fo |
| he supply of insulin or liraglutide or when prescribed for a patie | | | | |
| annotate the prescription as endorsed where there exists a rec | ord of prior dispensi | ng of insulin c | or liraglut | ide. |
| NSULIN PEN NEEDLES – Maximum of 200 dev per prescript | ion | | | |
| ₭ 29 g × 12.7 mm | | 100 | | D Micro-Fine |
| ₩ 31 g × 5 mm | | 100 | | D Micro-Fine |
| ₩ 31 g × 6 mm | 9.50 | 100 | 🗸 Be | erpu |

100

100

- ✓ B-D Micro-Fine
 - ✓ B-D Micro-Fine

* 31 g × 8 mm 10.95

| | Subsidy (Manufacturer's Price) | Der | Fully Subsidised | Generic | | | |
|---|-----------------------------------|-----|---------------------|-------------------|--|--|--|
| | \$ | Per | <i>.</i> | Manufacturer | | | |
| INSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE – Maximum of 200 dev per prescription | | | | | | | |
| * Syringe 0.3 ml with 29 g × 12.7 mm needle | | 100 | 1 | B-D Ultra Fine | | | |
| | 1.36 | 10 | | | | | |
| | (1.99) | | | B-D Ultra Fine | | | |
| * Syringe 0.3 ml with 31 g × 8 mm needle | | 100 | 1 | B-D Ultra Fine II | | | |
| | 1.30 | 10 | | | | | |
| | (1.99) | | | B-D Ultra Fine II | | | |
| * Syringe 0.5 ml with 29 g x 12.7 mm needle | | 100 | 1 | B-D Ultra Fine | | | |
| | 1.36 | 10 | | | | | |
| | (1.99) | | | B-D Ultra Fine | | | |
| * Syringe 0.5 ml with 31 g × 8 mm needle | () | 100 | 1 | B-D Ultra Fine II | | | |
| -, , , , , , , , , , , , , , , , , , , | 1.36 | 10 | | | | | |
| | (1.99) | | | B-D Ultra Fine II | | | |
| * Syringe 1 ml with 29 g × 12.7 mm needle | · · / | 100 | 1 | B-D Ultra Fine | | | |
| | 1.36 | 10 | | | | | |
| | (1.99) | | | B-D Ultra Fine | | | |
| * Syringe 1 ml with 31 g × 8 mm needle | () | 100 | 1 | B-D Ultra Fine II | | | |
| | 1.36 | 10 | | | | | |
| | (1.99) | .0 | | B-D Ultra Fine II | | | |
| Insulin Pumps | | | | | | | |

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

| a) | Maximum of 1 dev per prescription |
|----|-----------------------------------|
| b) | Only on a prescription |

| C) | Maximum of T insulin pump per patient each four yea | ar period. | | |
|----|---|------------|---|-----------------------------------|
| Mi | in basal rate 0.025 U/h | | 1 | MiniMed 770G |
| Mi | in basal rate 0.1 U/h | | 1 | Tandem t:slim |
| | | | | X2 with Basal-IQ |

⇒SA1603 Special Authority for Subsidy

data and a factor of the second

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and

.

- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

16

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | Su | Ibsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

4 Either:

4.1 Applicant is a relevant specialist; or

4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and

8 Either:

- 8.1 Applicant is a relevant specialist; or
- 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price | e) | Subsidised | Generic | |
| \$ | Per | · · · | Manufacturer | |

continued...

9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

| Subsidy | Fully | Brand or |
|------------------------|------------|----------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufact |

ic Manufacturer

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes: and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the followina:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application - (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person): and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 vears for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events: and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or

| Subs | sidy | Fully | Brand or |
|------------|---------------------|-------|--------------|
| (Manufactu | rer's Price) Subsid | lised | Generic |
| \$ | B Per | 1 | Manufacturer |

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

20

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

| | Subsidy (Manufacturer's Price) \$ | Sub Per | Fully osidised | Brand or Generic Manufacturer |
|--|---|------------------------|-------------------|-------------------------------------|
| continued | | | | |
| than 80 mmol/mol; and | | | | |
| 2 The patient's HbA1c has not deteriorated more than 5 mm 3 The patient has not had an increase in severe unexplained 4 Either: | | | | ne; and |
| 4.1 Applicant is a relevant specialist; or4.2 Applicant is a nurse practitioner working within their | r vocational scope. | | | |
| INSULIN PUMP CARTRIDGE – Special Authority see SA1985 o a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per Cartridge 300 U, t:lock × 10 | year. | narmacy 1 OP | ✓ т | andem Cartridge |
| INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special | Authority see SA1988 | 5 on page | e 19 – Re | etail pharmacy |
| a) Maximum of 3 set per prescription | , | 1.0 | | , , |
| b) Only on a prescription | | | | |
| c) Maximum of 13 infusion sets will be funded per year. | | | <i>.</i> | |
| 10 mm steel needle; 60 cm tubing × 10 | | 1 OP | ✓ N | IiniMed Sure-T MMT-884A |
| 10 mm steel needle; 80 cm tubing × 10 | 130.00 | 1 OP | ✓ N | /iniMed Sure-T MMT-886A |
| 6 mm steel needle; 60 cm tubing × 10 | 130.00 | 1 OP | ✓ N | /iniMed Sure-T MMT-864A |
| 6 mm steel needle; 80 cm tubing × 10 | 130.00 | 1 OP | ✓ N | /iniMed Sure-T MMT-866A |
| 8 mm steel needle; 60 cm tubing × 10 | 130.00 | 1 OP | ✓ N | /iniMed Sure-T MMT-874A |
| 8 mm steel needle; 80 cm tubing × 10 | 130.00 | 1 OP | ✓ N | /iniMed Sure-T MMT-876A |
| (MiniMed Sure-T MMT-884A 10 mm steel needle; 60 cm tubing × | 10 to be delisted 1 | luly 2024 |) | |
| (MiniMed Sure-T MMT-886A 10 mm steel needle; 80 cm tubing × | 10 to be delisted 1 | luly 2024 _, |) | |
| INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT Retail pharmacy | INSERTION) – Spe | cial Auth | ority see | SA1985 on page 19 - |
| a) Maximum of 3 sets per prescription | | | | |
| b) Only on a prescription | | | | |
| c) Maximum of 13 infusion sets will be funded per year. | | | | |
| 6 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles | 130.00 | 1 OP | ✓ т | ruSteel |
| 8 mm steel cannula: straight insertion: 80 cm line x 10 with | | | | |

| 8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles | 130.00 | 1 OP | ✓ TruSteel |
|---|--------|------|------------|
| 6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles | 130.00 | 1 OP | ✓ TruSteel |
| 8 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles | 130.00 | 1 OP | ✓ TruSteel |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-------|---------------------|-------------------------------------|
| NSULIN PUMP INFUSION SET (TEFLON CANNULA) – Spec a) Maximum of 3 set per prescription | cial Authority see SA19 | 85 on | page 19 - | - Retail pharmacy |
| b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 10 mm toffice needle, 110 cm toffice u 10. | 100.00 | 1 OP | | MiniMed Citherrothe |
| 13 mm teflon needle, 110 cm tubing × 10 | 130.00 | TOP | v | MiniMed Silhouette MMT-382A |
| 13 mm teflon needle, 45 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-368A |
| 13 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | ~ | MiniMed Silhouette MMT-381A |
| 13 mm teflon needle, 80 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-383A |
| 17 mm teflon needle, 110 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-377A |
| 17 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-378A |
| 17 mm teflon needle, 80 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-384A |
| 6 mm teflon needle, 110 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set MMT-398A |
| 6 mm teflon needle, 45 cm blue tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-941A |
| 6 mm teflon needle, 45 cm pink tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-921A |
| 6 mm teflon needle, 60 cm blue tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-943A |
| 6 mm teflon needle, 60 cm pink tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-923A |
| 6 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set MMT-399A |
| 6 mm teflon needle, 80 cm blue tubing | 130.00 | 1 OP | 1 | MiniMed Mio MMT-945A |
| 6 mm teflon needle, 80 cm clear tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio |
| 6 mm teflon needle, 80 cm pink tubing × 10 | 130.00 | 1 OP | 1 | MMT-965A MiniMed Mio |
| 6 mm teflon needle, 80 cm tubing × 10 | 130.00 | 1 OP | 1 | MMT-925A MiniMed Quick-Set |
| 9 mm teflon needle, 110 cm tubing × 10 | 130.00 | 1 OP | 1 | MMT-387A MiniMed Quick-Set |
| 9 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MMT-396A MiniMed Quick-Set |
| 9 mm teflon needle, 80 cm clear tubing × 10 | 130.00 | 1 OP | 1 | MMT-397A MiniMed Mio MMT-975A |

| | Subsidy |) Out | Fully Brand or |
|---|---------------------------|-----------------|--|
| | (Manufacturer's Pri \$ | ce) Subs Per | sidised Generic Manufacturer |
| 9 mm teflon needle, 80 cm tubing × 10 | | 1 OP | MiniMed Quick-Set MMT-386A |
| (MiniMed Silhouette MMT-382A 13 mm teflon needle, 110 cm tubi | na x 10 to be de | listed 1 July 2 | |
| (MiniMed Silhouette MMT-368A 13 mm teflon needle, 45 cm tubin | | | |
| (MiniMed Silhouette MMT-383A 13 mm teflon needle, 80 cm tubin | | | |
| MiniMed Silhouette MMT-384A 17 mm teflon needle, 80 cm tubin | | | |
| (MiniMed Quick-Set MMT-387A 6 mm teflon needle, 80 cm tubing | | | |
| MiniMed Quick-Set MMT-386A 9 mm teflon needle, 80 cm tubing | | | |
| INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN SA1985 on page 19 – Retail pharmacy | SERTION WITH | INSERTION | DEVICE) - Special Authority see |
| a) Maximum of 3 sets per prescription | | | |
| b) Only on a prescription | | | |
| c) Maximum of 13 infusion sets will be funded per year. | | | |
| 13 mm teflon cannula; angle insertion; insertion device; 110 cr | m | | |
| line × 10 with 10 needles | | 1 OP | AutoSoft 30 |
| 13 mm teflon cannula; angle insertion; insertion device; 60 cm | | | |
| line × 10 with 10 needles | | 1 OP | ✓ AutoSoft 30 |
| INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT | | - | |
| see SA1985 on page 19 – Retail pharmacy | | | ON DEVICE) - Special Authority |
| a) Maximum of 3 sets per prescription | | | |
| b) Only on a prescription | | | |
| c) Maximum of 13 infusion sets will be funded per year. | | | |
| 6 mm teflon cannula; straight insertion; insertion device; | | | |
| 110 cm line × 10 with 10 needles | 140.00 | 1 OP | ✓ AutoSoft 90 |
| | | TOP | Autosoft 90 |
| 6 mm teflon cannula; straight insertion; insertion device; 60 cm line x 10 with 10 needles | | 1.00 | ✓ AutoSoft 90 |
| | 140.00 | 1 OP | V AutoSoft 90 |
| 9 mm teflon cannula; straight insertion; insertion device; | 140.00 | 1.00 | Auto Coff 00 |
| 110 cm line × 10 with 10 needles | | 1 OP | AutoSoft 90 |
| 9 mm teflon cannula; straight insertion; insertion device; 60 cm | | 4.00 | (Auto 0 - # 00 |
| line × 10 with 10 needles | | 1 OP | AutoSoft 90 |
| NSULIN PUMP RESERVOIR – Special Authority see SA1985 on | page 19 - Retai | l pharmacy | |
| Maximum of 3 sets per prescription | | | |
| b) Only on a prescription | | | |
| c) Maximum of 13 packs of reservoir sets will be funded per y | /ear. | | |
| 10 × luer lock conversion cartridges 1.8 ml for Paradigm pump | s50.00 | 1 OP | ADR Cartridge 1.8 |
| Cartridge for 7 series pump; 3.0 ml × 10 | 50.00 | 1 OP | MiniMed |
| | | | 3.0 Reservoir |
| | | | MMT-332A |
| Discritional Industrian Documents | | | |
| Digestives Including Enzymes | | | |
| PANCREATIC ENZYME | | | |
| Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase | | | |
| 10,000 Ph Eur U, total protease 600 Ph Eur U) | 34 93 | 100 | Creon 10000 |
| Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase | | 100 | |
| 25,000 Ph Eur U, total protease 1,000 Ph Eur U) | 04 28 | 100 | <u>Creon 25000</u> |
| | | 100 | • <u>01601123000</u> |
| Modified release granules pancreatin 60.12 mg (amylase | | | |
| 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph | 24.02 | 20 a OP | Creon Micro |
| Eur U) | | 20 g OP | |
| URSODEOXYCHOLIC ACID - Special Authority see SA1739 on t | | | |
| Cap 250 mg | | 100 | ✓ Ursosan |
| | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Si | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | | ubsidised | Generic |
| `\$´´ | Per | 1 | Manufacturer |

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Laxatives

Bulk-forming Agents

| ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln20.00 | 500 g OP | ✓ Konsyl-D |
|---|------------|--------------------------------------|
| Faecal Softeners | | |
| DOCUSATE SODIUM - Only on a prescription * Tab 50 mg | 100 100 | ✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u> |

| 24 | 🗸 fully subsidised |
|----|--------------------|
| 24 | Principal Supply |

S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply

| | Subsidy | | Fully Brand or |
|---|---------------------|----------------|---------------------------------------|
| | (Manufacturer's Pr | | idised Generic |
| | \$ | Per | Manufacturer |
| DOCUSATE SODIUM WITH SENNOSIDES | | | |
| Tab 50 mg with sennosides 8 mg | 3.50 | 200 | ✓ Laxsol |
| POLOXAMER – Only on a prescription | | | |
| Not funded for use in the ear. | | | |
| * Oral drops 10% | 4.17 | 30 ml OP | ✓ Coloxyl |
| Opioid Receptor Antagonists - Peripheral | | | |
| | CO1 holew Det | il nhormoou | |
| METHYLNALTREXONE BROMIDE – Special Authority see SA1 Inj 12 mg per 0.6 ml vial | | | ✓ Relistor |
| | 246.00 | 7 | ✓ Relistor |
| | 240.00 | 1 | • Heliston |
| SA1691 Special Authority for Subsidy Initial application — (Opioid induced constipation) from any unless notified for applications meeting the following criteria: Both: | relevant practitior | ner. Approvals | s valid without further renewal |
| 1 The patient is receiving palliative care; and 2 Either: | | | |
| 2.1 Oral and rectal treatments for opioid induced cons | tination are ineffe | ctive: or | |
| 2.1 Oral and rectal treatments for opioid induced cons 2.2 Oral and rectal treatments for opioid induced cons | | | ed. |
| | apadon are anabi | | |
| Osmotic Laxatives | | | |
| GLYCEROL | | | |
| Suppos 2.8/4.0 g – Only on a prescription | 10.39 | 20 | Lax-suppositories |
| | | | Glycerol |
| LACTULOSE – Only on a prescription | | | |
| * Oral liq 10 g per 15 ml | 3.61 | 500 ml | ✓ Laevolac |
| MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BI | CARBONATE AN | ID SODIUM C | HLORIDE |
| Powder for oral soln 13.125 g with potassium chloride 46.6 n | | | - |
| sodium bicarbonate 178.5 mg and sodium chloride 350. | | 30 | Molaxole |
| SODIUM ACID PHOSPHATE – Only on a prescription | | | |
| Enema 16% with sodium phosphate 8% | 2 50 | 1 | Fleet Phosphate |
| | 2.00 | | Enema |
| | | optintion | Enoma |
| SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE | • • | scription | |
| Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml | | 50 | Missiste |
| 5 ml | | 50 | Micolette |
| | | | ✓ Micolette-S29 S29 |
| (Micolette-S29 S29 Enema 90 mg with sodium lauryl sulphoace | tate 9 mg per ml, | 5 ml to be del | isted 1 August 2024) |
| Stimulant Laxatives | | | |
| | | | |
| BISACODYL – Only on a prescription * Tab 5 mg | E 90 | 200 | Bisacodyl Viatris |
| | | 200 10 | ✓ Lax-Suppositories |
| | | 10 | |
| SENNA – Only on a prescription | 0.17 | 100 | |
| * Tab, standardised | | 100 | Constrat |
| | (8.21) | 00 | Senokot |
| | 0.43 | 20 | Senokot |
| | (2.06) | | JEHUKUL |
| | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | , | Brand or |
|---|-----------------------------|----------------|--|----------------------------|
| | (Manufacturer's Price \$ | e) Sub Per | | Generic Manufacturer |
| ODIUM PICOSULFATE - Special Authority see SA2053 be | | | | |
| Oral soln 7.5 mg per ml | 7.40 | 30 ml OP | 🗸 Du | Icolax SP Drop |
| SA2053 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals Both: | valid for 6 months for a | pplications | meeting th | ne following criteria: |
| The patient is a child with problematic constipation de macrogol where practicable; and The patient would otherwise require a high-volume bo | | | | |
| tenewal from any relevant practitioner. Approvals valid for benefiting from treatment. | • • • | | | |
| Metabolic Disorder Agents | | | | |
| LGLUCOSIDASE ALFA – Special Authority see SA1986 be | elow – Retail pharmacy | | | |
| Inj 50 mg vial | 1,142.60 | 1 | 🖌 My | ozyme |
| SA1986 Special Authority for Subsidy | | | | |
| nitial application only from a metabolic physician. Approva II of the following: | ils valid for 12 months fo | or application | ons meetir | ng the following criteria: |
| The patient is aged up to 24 months at the time of initi and | al application and has b | een diagno | osed with i | infantile Pompe disease |
| 2 Any of the following: | | | | |
| Diagnosis confirmed by documented deficiency villus biopsies and/or cultured amniotic cells; or | | ase by prer | natal diagr | nosis using chorionic |
| 2.2 Documented deficiency of acid alpha-glucoside | | ccharide te | sting indic | ating a diagnostic |
| elevation of glucose tetrasaccharides; or 2.3 Documented deficiency of acid alpha-glucosid | | | enetic testi | ng indicating a |
| disease-causing mutation in the acid alpha-glu 2.4 Documented urinary tetrasaccharide testing in | | | lucose tet | rasaccharides, and |
| molecular genetic testing indicating a disease- | causing mutation in the | GAA gene | ; and | |
| Patient has not required long-term invasive ventilation (ERT); and | for respiratory failure p | rior to start | ing enzym | e replacement therapy |
| 4 Patient does not have another life-threatening or seve or might be reasonably expected to compromise a res | ponse to ERT; and | - | - | be influenced by ERT |
| 5 Alglucosidase alfa to be administered at doses no gre tenewal only from a metabolic physician. Approvals valid for | | | | owing criteria: |
| Il of the following: | | | ing the foll | owing chicha. |
| 1 The treatment remains appropriate for the patient and | | | | |
| 2 Alglucosidase alfa to be administered at doses no gre 3 Patient has not had severe infusion-related adverse re | | | | opriate pre-medication |
| and/or adjustment of infusion rates; and | eactions which were not | preventab | ie by appi | ophate pre-medication |
| 4 Patient has not developed another life threatening or s influenced by ERT; and | severe disease where the | ne long tern | n prognos | is is unlikely to be |
| 5 Patient has not developed another medical condition t ERT; and | hat might reasonably be | e expected | to compro | omise a response to |
| 6 There is no evidence of life threatening progression of invasive ventilation; and | f respiratory disease as | evidenced | by the nee | eded for > 14 days of |
| 7 There is no evidence of new or progressive cardiomy | opathy. | | | |
| | – Rotail nharmaov | | | |
| RGININE – Special Authority see SA2042 on the next page | - netali phannacy | | | |
| Tab 1,000 mg | CBS | 90 50 | ✓ Clin✓ Sol | nicians |

| | Subsidy acturer's Price) \$ | F Subsid Per | ised (| Brand or Generic Manufacturer |
|--|-----------------------------------|--------------------|---------------|-------------------------------------|
| ■SA2042 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals valid for 6 metabolism that may respond to arginine supplementation. | 3 months where | e patient ha | as a sus | pected inborn error of |
| Renewal only from a metabolic physician. Approvals valid for 24 months Both: | s for applicatior | ns meeting | the follo | owing criteria: |
| The patient has a confirmed diagnosis of an inborn error of metable The treatment remains appropriate and the patient is benefiting from the patient is benefiti | | onds to ar | ginine s | upplementation; and |
| BETAINE – Special Authority see SA1987 below – Retail pharmacy Powder for oral soln | 75.00 180 |) g OP | 🗸 Cys | stadane |
| SA1987 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals valid for 1 All of the following: 1 The patient has a confirmed diagnosis of homocystinuria; and | 12 months for a | pplications | meetin | g the following criteria: |
| 2 Any of the following: 2.1 A cystathionine beta-synthase (CBS) deficiency; or 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) de 2.3 A disorder of intracellular cobalamin metabolism; and | ficiency; or | | | |
| 3 An appropriate homocysteine level has not been achieved despite Renewal only from a metabolic physician. Approvals valid for 12 months patient is benefiting from treatment. | | | | |
| COENZYME Q10 – Special Authority see SA2039 below – Retail pharm Cap 120 mg Cap 160 mg | CBS | 30 60 | ✔ Sol ✔ Go | gar Healthy |
| SA2039 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals valid for € metabolism that may respond to coenzyme Q10 supplementation. Renewal only from a metabolic physician. Approvals valid for 24 months | | | | |
| Both: 1 The patient has a confirmed diagnosis of an inborn error of metab and 2 The treatment remains appropriate and the patient is benefiting fr | | onds to co | enzyme | e Q10 supplementation; |
| GALSULFASE – Special Authority see SA1988 below – Retail pharmac Inj 1 mg per ml, 5 ml vial2,23 | y | 1 | 🗸 Nag | glazyme |
| SA1988 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals valid for 1 Both: | 12 months for a | pplications | meetin | g the following criteria: |
| The patient has been diagnosed with mucopolysaccharidosis VI; Either: | | | | |
| 2.1 Diagnosis confirmed by demonstration of N-acetyl-galacto enzyme activity assay in leukocytes or skin fibroblasts; or 2.2 Detection of two disease causing mutations and patient ha VI. | | | | , , , , |
| Renewal only from a metabolic physician. Approvals valid for 12 months All of the following: | s for applicatior | ns meeting | the follo | owing criteria: |

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| | Subsidy | | Fully | Brand or |
|--|------------------------|------------|---------------|-----------------------------|
| | (Manufacturer's Price | | Subsidised | Generic |
| | \$ | Per | v | Manufacturer |
| continued and/or adjustment of infusion rates; and | | | | |
| 3 Patient has not developed another life threatening or sev | ara disaasa whara th | a long | term progn | osis is unlikely to be |
| influenced by Enzyme Replacement Therapy (ERT); and | | ic long | term progra | |
| 4 Patient has not developed another medical condition that | | expec | ted to com | promise a response to |
| ERT. | | | ···· · | |
| IDURSULFASE - Special Authority see SA1623 below - Retail | pharmacy | | | |
| Inj 2 mg per ml, 3 ml vial | | 1 | ✓ E | laprase |
| ➡SA1623 Special Authority for Subsidy | | | | |
| Initial application only from a metabolic physician. Approvals v | alid for 24 weeks for | applic | ations meet | ing the following criteria: |
| All of the following: | | | | |
| The patient has been diagnosed with Hunter Syndrome (Either: | mucopolysaccharido | sis II); a | and | |
| 2.1 Diagnosis confirmed by demonstration of iduronal assay in cultured skin fibroblasts; or | e 2-sulfatase deficie | ncy in v | white blood | cells by either enzyme |
| 2.2 Detection of a disease causing mutation in the idu | ironate 2-sulfatase g | ene; ar | nd | |
| 3 Patient is going to proceed with a haematopoietic stem c idursulfase would be bridging treatment to transplant; and | | within | the next 3 r | months and treatment with |
| 4 Patient has not required long-term invasive ventilation for (ERT); and | | rior to s | starting Enzy | vme Replacement Therapy |
| 5 Idursulfase to be administered for a total of 24 weeks (eq | uivalent to 12 weeks | pre- a | nd 12 week | s post-HSCT) at doses no |
| greater than 0.5 mg/kg every week. | | | | |
| LARONIDASE - Special Authority see SA1695 below - Retail p | | | | |
| Inj 100 U per ml, 5 ml vial | 1,335.16 | 1 | ✓ A | Idurazyme |
| SA1695 Special Authority for Subsidy | | | | |
| Initial application only from a metabolic physician. Approvals v All of the following: | valid for 24 weeks for | applic | ations meet | ing the following criteria: |
| The patient has been diagnosed with Hurler Syndrome (r Either: | nucopolysacchardos | is I-H); | and | |
| 2.1 Diagnosis confirmed by demonstration of alpha-L- assay in cultured skin fibroblasts; or | iduronidase deficien | cy in w | hite blood c | ells by either enzyme |
| 2.2 Detection of two disease causing mutations in the to have Hurler syndrome; and | alpha-L-iduronidase | gene a | and patient | has a sibling who is known |
| 3 Patient is going to proceed with a haematopoietic stem c laronidase would be bridging treatment to transplant; and | | within | the next 3 r | months and treatment with |
| 4 Patient has not required long-term invasive ventilation for | | rior to s | starting Enzy | me Replacement Therapy |
| (ERT); and | | | | |
| 5 Laronidase to be administered for a total of 24 weeks (eq than 100 units/kg every week. | uivalent to 12 weeks | pre- a | nd 12 post-l | HSCT) at doses no greater |
| LEVOCARNITINE - Special Authority see SA2040 on the next | page – Retail pharma | acv | | |
| Tab 500 mg | • | 30 | ✓ s | olgar |
| Cap 250 mg | | 30 | | olgar |
| Cap 500 mg | CBS | 60 | 🖌 E | Balance |
| Oral liq 1 g per 10 ml | CBS | 118 m | I ∕ C | Carnitor S29 |
| | | | 🗸 N | lovitium Sugar |
| | | | | Free S29 |
| Oral liq 500 mg per 10 ml | CBS | 300 m | I | Balance |
| | | | | |

| Subsidy | Fu | | |
|------------------------|-----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidise | ed Generic | |
| \$ | Per | Manufacturer | |

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN – Special Authority see SA2041 below – Retail pharmacy

| Tab 100 mg | CBS | 100 | ✓ Country Life ✓ Puritan's Pride Vitamin B-2 100 mg ^{S29} |
|------------|-----|-----|---|
| Cap 100 mg | CBS | 100 | ✓ Solgar |

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

➡SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and

| | | | | 2 |
|--|--|------------------------------|----------|---|
| | Subsidy (Manufacturer's Price | e) Subsi | Fully | Brand or Generic |
| | \$ | Per | 1 | Manufacturer |
| continued | | | | |
| Sapropterin to be administered at doses no greater than a Sapropterin to be used alone or in combination with PKU Total treatment duration with sapropterin will not exceed 2 becoming pregnant) and treatment will be stopped after de | dietary managemer 2 months for each | it; and | | s time for planning and |
| SODIUM BENZOATE – Special Authority see SA1599 below – F Soln 100 mg per ml | | 100 ml | 🗸 A | mzoate S29 |
| SA1599 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals va cycle disorder. Renewal only from a metabolic physician. Approvals valid for 12 | alid for 12 months w | here the pat | tient ha | s a diagnosis of a urea |
| patient is benefiting from treatment. | | | | |
| SODIUM PHENYLBUTYRATE – Special Authority see SA1990 I Grans 483 mg per g | | nacy 174 g OP | ✓ P | heburane |
| ► SA1990 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals va cycle disorder involving a deficiency of carbamylphosphate synth synthetase. Renewal only from a metabolic physician. Approvals valid for 12 patient is benefiting from treatment. | etase, ornithine trar | nscarbamyla | se or a | rgininosuccinate |
| TAURINE – Special Authority see SA2043 below – Retail pharm | асу | | | |
| Cap 500 mg Cap 1,000 mg Powder | CBS | 50 90 300 g | 🗸 L | olgar ife Extension ife Extension |
| SA2043 Special Authority for Subsidy Initial application only from a metabolic physician. Approvals variation metabolic physician is a provided with the second term of ter | months for applica ndrial disorder whic | tions meetin h responds t | g the fo | bllowing criteria: |
| TRIENTINE – Special Authority see SA2324 below – Retail phar | U U | | | |
| Cap 250 mg | 2,022.00 | 100 | 🗸 Т | rientine Waymade |
| SA2324 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals value the following criteria: All of the following: 1 Patient has confirmed Wilson disease; and 2 Treatment with D-penicillamine has been trialled and disco effects or has not received sufficient benefit; and | ontinued because th | ne person ha | is expe | rienced intolerable side |
| 3 Treatment with zinc has been trailled and discontinued be not received sufficient benefit, or zinc is considered clinica and requires copper chelation. | | | | |
| Gaucher's Disease | | | | |
| TALIGLUCERASE ALFA – Special Authority see SA2137 on the Inj 200 unit vial | | pharmacy 1 | ✔ E | lelyso |

| Subsidy | Fully | | Brand or | |
|------------------------|------------|---|--------------|--|
| (Manufacturer's Price) | Subsidised | | Generic | |
| \$ | Per | 1 | Manufacturer | |

➡SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).
- Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

| Soln 0.15% - Higher subsidy of \$22.60 per 500 ml with | | 500 1 | |
|---|-----------------|-------------------|---------------------------------|
| Endorsement | 9.00 (22.60) | 500 ml | Difflam |
| Additional subsidy by endorsement for a patient who has | · · · · | s a result of tre | Billian |
| prescription is endorsed accordingly. | | | |
| CARMELLOSE SODIUM WITH GELATIN AND PECTIN | | | |
| Paste | 17.20 | 56 g OP | Stomahesive |
| | 4.55 | 15 g OP | |
| | (7.90) | | Orabase |
| | 1.52 | 5 g OP | |
| | (3.60) | | Orabase |
| Powder | | 28 g OP | o |
| | (10.95) | | Stomahesive |

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Pr \$ | rice) Subs Per | Fully Brand or idised Generic ✔ Manufacturer |
|---|-------------------------------------|-------------------|--|
| CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE * Adhesive gel 8.7% with cetalkonium chloride 0.01% | 2.06 (6.00) | 15 g OP | Bonjela |
| TRIAMCINOLONE ACETONIDE Paste 0.1% | 5.49 | 5 g OP | ✓ Kenalog in Orabase |
| Oropharyngeal Anti-infectives | | | |
| AMPHOTERICIN B Lozenges 10 mg | 5.86 | 20 | 🗸 Fungilin |
| MICONAZOLE Oral gel 20 mg per g | 4.74 | 40 g OP | ✓ <u>Decozol</u> |
| NYSTATIN Oral liq 100,000 u per ml | 2.22 | 24 ml OP | ✓ <u>Nilstat</u> |
| Vitamins | | | |
| Vitamin B | | | |
| HYDROXOCOBALAMIN Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a F | 2SO2.46 | 3 | ✓ Cobal-B12 5299 ✓ <u>Hydroxocobalamin</u> <u>Panpharma</u> ✓ Vita-B12 |
| | 4.10 | 5 | Vita-B12 Cobalin-H \$29 Neo-Cytamen \$29 \$29 |
| | 8.20 | 10 | Vitarubin Depot Injection \$29 |
| PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription | | | |
| * Tab 25 mg – No patient co-payment payable * Tab 50 mg | | 90 500 | ✓ <u>Vitamin B6 25</u> ✓ Pyridoxine multichem |
| THIAMINE HYDROCHLORIDE – Only on a prescription * Tab 50 mg | 4.65 | 100 | ✓ Thiamine multichem |
| VITAMIN B COMPLEX * Tab, strong, BPC | 11.25 | 500 | ✓ Bplex |
| Vitamin C | | | |
| ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription | | | |
| * Tab 100 mg | | 500 | ✓ <u>Cvite</u> |

| | Subsidy (Manufacturer's P | rice) Subs | Fully | Brand or Generic |
|--|------------------------------|----------------|------------|---------------------------|
| | \$ | Per | 1 | Manufacturer |
| Vitamin D | | | | |
| ALFACALCIDOL | | | | |
| * Cap 0.25 mcg | | 100 | ✓ 0 | ne-Alpha |
| | | | ✓ 0 | ne-Alpha S29 S29 |
| * Cap 1 mcg | 87.98 | 100 | | ne-Alpha |
| | co.co | 00 00 | | one-Alpha S29 S29 |
| * Oral drops 2 mcg per ml | 00.08 | 20 ml OP | v (| ne-Alpha |
| CALCITRIOL * Cap 0.25 mcg | 7 90 | 100 | | alcitriol-AFT |
| * Cap 0.25 mcg | | 100 100 | - | alcitriol-AFT |
| | | 100 | • • | |
| COLECALCIFEROL * Cap 1.25 mg (50,000 iu) - Maximum of 12 cap per preso | ription 3.65 | 12 | 1 V | ït.D3 |
| Vit.D3 to be Principal Supply on 1 June 2024 | | 12 | • • | 11.00 |
| * Oral liq 188 mcg per ml (7,500 iu per ml) | 9.00 | 5 ml OP | ✓ 0 | linicians |
| | | | | |
| Multivitamin Preparations | | | | |
| MULTIVITAMIN RENAL - Special Authority see SA1546 below | w – Retail pharmac | y | | |
| * Сар | 7.28 | 30 | ✓ 0 | linicians Renal Vit |
| ■ SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals the following criteria: Either: | | | | |
| The patient has chronic kidney disease and is receivin The patient has chronic kidney disease grade 5, define 15 ml/min/1.73 m² body surface area (BSA). | | | | |
| MULTIVITAMINS – Special Authority see SA1036 below – R | etail pharmacy | | | |
| * Powder | 74.88 | 200 g OP | 🗸 P | aediatric Seravit |
| SA1036 Special Authority for Subsidy | | | | |
| Initial application from any relevant practitioner. Approvals | valid without further i | renewal unless | s notifie | d where the patient has |
| inborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid with | out further repowel u | place petified | whore r | ationt has had a proviou |
| approval for multivitamins. | | niess nouneu | where p | allent has had a previot |
| VITAMINS | | | | |
| * Tab (BPC cap strength) | 18.50 | 1,000 | 🗸 N | lvite |
| Cap (fat soluble vitamins A, D, E, K) – Special Authority | | 1,000 | • • | |
| SA1720 below – Retail pharmacy | | 60 | 🗸 V | itabdeck |
| SA1720 Special Authority for Subsidy | | | | |
| Initial application from any relevant practitioner. Approvals | valid without further i | renewal unless | s notifie | d for applications meetir |
| the following criteria: | | | | |
| Any of the following: | | | | |
| 1 Patient has cystic fibrosis with pancreatic insufficiency | | | | |
| 2 Patient is an infant or child with liver disease or short g | jut syndrome; or | | | |

2 Patient is an infant or child with liver disease or short gut syndrome; or

3 Patient has severe malabsorption syndrome.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|---|---|----------------------|---------------------|--|
| Minerals | | | | |
| Calcium | | | | |
| CALCIUM CARBONATE * Tab 1.25 g (500 mg elemental) * Tab eff 1.25 g (500 mg elemental) – Subsidy by endorseme | nt260.00 | 250 100 | 1 | Calci-Tab 500 Calcium 500 mg Hexal 529 |
| Subsidy by endorsement – Only when prescribed for pa considered unsuitable. | ediatric patients (< 5 | years |) where ca | lcium carbonate oral liquid is |
| CALCIUM GLUCONATE * Inj 10%, 10 ml ampoule | | 10 | 1 | Max Health - Hameln 829 |
| | 64.00 | 20 | 1 | Max Health \$29 |
| lodine | | | | |
| POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine) | 5.99 | 90 | 1 | <u>NeuroTabs</u> |
| Iron | | | | |
| FERROUS FUMARATE * Tab 200 mg (65 mg elemental) FERROUS FUMARATE WITH FOLIC ACID | 3.04 | 100 | v | Ferro-tab |
| * Tab 310 mg (100 mg elemental) with folic acid 350 mcg FERROUS SULFATE | 5.98 | 100 | 1 | Ferro-F-Tabs |
| * Tab long-acting 325 mg (105 mg elemental) * Oral liq 30 mg (6 mg elemental) per 1 ml | 9.25 | 30 250 n 500 n | nl 🖌 | <u>Ferrograd</u> Ferro-Liquid <u>Ferodan</u> |
| IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority s Inj 50 mg per ml, 10 ml vial | | Retail 1 | | Ferinject |

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and

- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application - (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist,

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

1 Patient has been diagnosed with iron-deficiency anaemia; and

- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE

| * Inj 50 mg per ml, 2 ml ampoule | 5 | ✓ Ferrosig |
|---|--------|---|
| Magnesium | | |
| MAGNESIUM HYDROXIDE Suspension 8%33.60 | 355 ml | Phillips Milk of Magnesia 529 |
| MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule | 10 | Martindale |
| * Inj 2 mmol per ml, 10 ml ampoule75.06 Zinc | 10 | ✓ Inresa S29 |
| ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)11.00 | 100 | ✓ Zincaps |

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | 5 | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

Antianaemics

Hypoplastic and Haemolytic

► SA2266 Special Authority for Subsidy

Initial application — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.
- Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA2266 above - Retail pharmacy

Wastage claimable

| Tuotago olamabio | | |
|---------------------------------|-------|------------------------------|
| Inj 1,000 iu in 0.5 ml, syringe | 6 | Binocrit |
| Inj 2,000 iu in 1 ml, syringe | 6 | Binocrit |
| Inj 3,000 iu in 0.3 ml, syringe | 6 | Binocrit |
| Inj 4,000 iu in 0.4 ml, syringe | 6 | Binocrit |
| Inj 5,000 iu in 0.5 ml, syringe | 6 | Binocrit |
| Inj 6,000 iu in 0.6 ml, syringe | 6 | Binocrit |
| Inj 8,000 iu in 0.8 ml, syringe | 6 | Binocrit |
| Inj 10,000 iu in 1 ml, syringe | 6 | Binocrit |
| Inj 40,000 iu in 1 ml, syringe | 1 | Binocrit |
| | | |

| | Subsidy (Manufacturer's Price \$ | e) Sub Per | Fully osidised | Brand or Generic Manufacturer |
|---|---|---------------|---------------------------|--|
| Megaloblastic | | | | |
| FOLIC ACID | | | | |
| * Tab 0.8 mg | 26.60 | 1,000 | ✓ F | Folic Acid multichem |
| * Tab 5 mg | 5.82 | 100 | ✓ <u>F</u> | Folic Acid Viatris |
| Oral liq 50 mcg per ml | 30.26 | 25 ml OP | ✓ E | Biomed |
| Antifibringlytics, Hasmostatics and Losal Solar | oconto | | | |
| Antifibrinolytics, Haemostatics and Local Sclere | USams | | | |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha | arm] | | | |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha For patients with haemophilia B receiving prophylaxis treatm | arm] ent. Access to fund | | ent is ma | anaged by the Haemophili |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha For patients with haemophilia B receiving prophylaxis treatm Treaters Group in conjunction with the National Haemophilia | arm] ent. Access to fund Management grou | | | |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha For patients with haemophilia B receiving prophylaxis treatm Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial | arm] ent. Access to fund Management grou 612.50 | | √ µ | Alprolix |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha For patients with haemophilia B receiving prophylaxis treatm Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial Inj 500 iu vial | arm] ent. Access to fund Management grou 612.50 1,225.00 | | ✓ ✓ | Alprolix Alprolix |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha For patients with haemophilia B receiving prophylaxis treatm Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial | arm] ent. Access to fund Management grou 612.50 1,225.00 2,450.00 | | ✓ ✓ ✓ | Alprolix Alprolix Alprolix |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha For patients with haemophilia B receiving prophylaxis treatm Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial | arm] ent. Access to fund Management grou 612.50 1,225.00 2,450.00 4,900.00 | | I | Alprolix Alprolix Alprolix Alprolix Alprolix |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha For patients with haemophilia B receiving prophylaxis treatm Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial | arm] Management grou 612.50 2,450.00 4,900.00 7,350.00 | | ✓ ✓ ✓ ✓ | Alprolix Alprolix Alprolix |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha For patients with haemophilia B receiving prophylaxis treatm Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial Inj 4,000 iu vial ELTROMBOPAG – Special Authority see SA1743 below – Retai | arm] ent. Access to fund Management group 612.50 2,450.00 2,450.00 4,900.00 7,350.00 9,800.00 | | ✓ ✓ ✓ ✓ | Alprolix Alprolix Alprolix Alprolix Alprolix Alprolix |
| EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpha For patients with haemophilia B receiving prophylaxis treatm Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial Inj 3,000 iu vial Inj 4,000 iu vial | arm] Management group 612.50 2,450.00 4,900.00 7,350.00 9,800.00 I pharmacy | | | Alprolix Alprolix Alprolix Alprolix Alprolix Alprolix |

➡SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and

- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA2272 below

| Inj 30 mg in 1 ml vial | | 1 | Hemlibra |
|---------------------------|----------|---|------------------------------|
| Inj 60 mg in 0.4 ml vial | 7,138.00 | 1 | Hemlibra |
| Inj 105 mg in 0.7 ml vial | | 1 | Hemlibra |
| Inj 150 mg in 1 ml vial | | 1 | Hemlibra |

⇒SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

38

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

| Inj 1 mg syringe | 1 | NovoSeven RT |
|--------------------------|---|--------------|
| Inj 2 mg syringe2,356.60 | 1 | NovoSeven RT |
| Inj 5 mg syringe | 1 | NovoSeven RT |
| Inj 8 mg syringe | 1 | NovoSeven RT |

| | Subsidy | | Fully | Brand or |
|---|-----------------------------|---------|-----------------------|-----------------------------|
| | (Manufacturer's Price) | ç | Fully Subsidised | Generic |
| | \$ | Per | ✓ | Manufacturer |
| FACTOR EIGHT INHIBITOR BYPASSING FRACTION – [Xph | arml | | | |
| For patients with haemophilia. Preferred Brand of bypassi | | oredict | ted use A | ccess to funded treatment |
| is managed by the Haemophilia Treaters Group in conjunc | | | | |
| Inj 500 U | | 1 | | EIBANF |
| Inj 1,000 U | | 1 | | |
| Inj 2,500 U | | 1 | | EIBANF |
| MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – [Xp | | | | |
| For patients with haemophilia. Rare Clinical Circumstance | | a rocor | mhinant far | stor VIII Access to funder |
| treatment is managed by the Haemophilia Treaters Group | in conjunction with the | Vation | al Haamon | hilia Management Group |
| subject to criteria. | | Valion | arnaemop | mila management oroup, |
| Inj 250 iu prefilled syringe | 287 50 | 1 | ✓ X | (yntha |
| Inj 500 iu prefilled syringe | | 1 | | (yntha |
| Inj 1,000 iu prefilled syringe | | 1 | | (yntha |
| Inj 2,000 iu prefilled syringe | | 1 | | lyntha |
| Inj 3,000 iu prefilled syringe | | 1 | | lyntha |
| , , , , | | | - / | , yn dia |
| NONACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xphar | | la | | |
| For patients with haemophilia. Access to funded treatmen | t is managed by the Ha | emopr | illa i reate | rs Group in conjunction |
| with the National Haemophilia Management Group. | 105.00 | | | |
| Inj 500 iu vial | | 1 | | RIXUBIS |
| Inj 1,000 iu vial | | 1 | - | RIXUBIS |
| Inj 2,000 iu vial | | 1 | | RIXUBIS |
| Inj 3,000 iu vial | , | 1 | • • | RIXUBIS |
| OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) | | | | |
| For patients with haemophilia. Preferred Brand of short ha | | | | |
| managed by the Haemophilia Treaters Group in conjunction | | • | | |
| Inj 250 iu vial | | 1 | | dvate |
| Inj 500 iu vial | | 1 | | dvate |
| Inj 1,000 iu vial | | 1 | - | dvate |
| Inj 1,500 iu vial | | 1 | | dvate |
| Inj 2,000 iu vial | | 1 | | dvate |
| Inj 3,000 iu vial | | 1 | ✓ µ | Advate |
| OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENAT | | | | |
| For patients with haemophilia. Rare Clinical Circumstance | es Brand of short half-life | e recor | mbinant fac | ctor VIII. Access to funded |
| treatment is managed by the Haemophilia Treaters Group | in conjunction with the | Vation | al Haemop | hilia Management Group, |
| subject to criteria. | | | | |
| Inj 250 iu vial | | 1 | 🗸 k | Cogenate FS |
| Inj 500 iu vial | | 1 | 🗸 k | Cogenate FS |
| Inj 1,000 iu vial | | 1 | 🗸 k | Cogenate FS |
| Inj 2,000 iu vial | 1,900.00 | 1 | | Cogenate FS |
| Inj 3,000 iu vial | 2,850.00 | 1 | 🗸 k | Cogenate FS |
| RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VI | II] – [Xpharm] | | | |
| For patients with haemophilia A receiving prophylaxis treat | | d treat | ment is ma | anaged by the Haemophilia |
| Treaters Group in conjunction with the National Haemophi | | | | |
| Inj 250 iu vial | 0 0 1 | 1 | 🗸 A | Adynovate |
| Inj 500 iu vial | | 1 | | Adynovate |
| Inj 1,000 iu vial | | 1 | | Adynovate |
| Inj 2,000 iu vial | | 1 | | Adynovate |
| SODIUM TETRADECYL SULPHATE | , | | - | |
| * Inj 3% 2 ml | 28 50 | 5 | | |
| 小 IIJ ∪ /0 ∠ III | (73.00) | 5 | - | ibro-vein |
| | (73.00) | | Г | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| (| Subsidy Manufacturer's Price) \$ | Per | Fully Subsidised | Generic |
|---|--|-----------|---------------------|--------------------------------------|
| TRANEXAMIC ACID | | | | |
| Tab 500 mg | 10.45 45.68 | 60 100 | | <u>Mercury Pharma</u> Cyklokapron |
| Vitamin K | | | | |
| PHYTOMENADIONE | | | | |
| Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO | | 5 5 | | Konakion MM Konakion MM |
| Antithrombotic Agents | | | | |
| Antiplatelet Agents | | | | |
| ASPIRIN * Tab 100 mg Ethics Aspirin EC to be Principal Supply on 1 June 2024 | 12.65 | 990 | 1 | Ethics Aspirin EC |
| CLOPIDOGREL | | | | |
| * Tab 75 mg | 5.07 | 84 | ~ | Arrow - Clopid |
| DIPYRIDAMOLE * Tab long-acting 150 mg | 13.93 | 60 | 1 | Pytazen SR |
| TICAGRELOR – Special Authority see SA1955 below – Retail pha * Tab 90 mg | • | 56 | | <u>Ticagrelor Sandoz</u> Brilinta |
| (Brilinta Tab 90 mg to be delisted 1 July 2024) | | | | |

(Brilinta Tab 90 mg to be delisted 1 July 2024)

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

40

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and

2 Either:

- 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
- 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

| Subsidy | | Fully | Brand or | |
|------------------------|------|---------|--------------|--|
| (Manufacturer's Price) | Subs | sidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.
- Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations

| ENOXAPARIN SODIUM - Special Authority see SA2152 b | elow – Retail pharmacy | | |
|--|------------------------|----|-----------------------------------|
| Inj 20 mg in 0.2 ml syringe | | 10 | Clexane |
| Inj 40 mg in 0.4 ml syringe | | 10 | Clexane |
| Inj 60 mg in 0.6 ml syringe | | 10 | Clexane |
| Inj 80 mg in 0.8 ml syringe | | 10 | Clexane |
| Inj 100 mg in 1 ml syringe | | 10 | Clexane |
| Inj 120 mg in 0.8 ml syringe | | 10 | Clexane Forte |
| Inj 150 mg in 1 ml syringe | 143.86 | 10 | Clexane Forte |
| | | | |

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | ully | Brand or |
|----|-----------------------|--------|------|--------------|
| () | Manufacturer's Price) | Subsid | sed | Generic |
| | \$ | Per | ✓ | Manufacturer |

continued...

treatment; or

- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

HEPARIN SODILIM

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

| Inj 1,000 iu per ml, 5 ml ampoule86.11 | 50 | Pfizer |
|--|----|--------------------------------------|
| Inj 5,000 iu per ml, 5 ml vial83.00 | 10 | ✓ <u>Heparin Sodium</u> Panpharma |
| Inj 5,000 iu per ml, 1 ml32.66 | 5 | ✓ DBL Heparin Sodium S29 |
| 70.33 | | ✓ Hospira |
| Inj 25,000 iu per ml, 0.2 ml22.42 | 5 | Hospira |
| 42.40 | | Heparin DBL S29 |
| 482.20 | 50 | ✓ Heparin DBL S29 |
| HEPARINISED SALINE Inj 10 iu per ml, 5 ml | 50 | ✓ Pfizer |
| Oral Anticoagulants | | |
| DABIGATRAN | | |
| Cap 75 mg – No more than 2 cap per day27.99 Pradaxa to be Principal Supply on 1 July 2024 | 60 | Pradaxa |
| Cap 110 mg27.99 Pradaxa to be Principal Supply on 1 July 2024 | 60 | Pradaxa |
| Cap 150 mg27.99 Pradaxa to be Principal Supply on 1 July 2024 | 60 | Pradaxa |
| RIVAROXABAN | | |
| Tab 10 mg – No more than 1 tab per day | 30 | ✓ Xarelto |
| Tab 15 mg – Up to 14 tab available on a PSO | 28 | ✓ Xarelto |
| Tab 20 mg | 28 | ✓ Xarelto |
| 1 ab 20 mg 14.00 | 20 | |

42

| | Subsidy | | Fully | Brand or |
|--|------------------------|-----|------------|--------------|
| | (Manufacturer's Price) | | Subsidised | Generic |
| | \$ | Per | 1 | Manufacturer |
| WARFARIN SODIUM | | | | |
| Note: Marevan and Coumadin are not interchangeable. | | | | |
| * Tab 1 mg | 3.46 | 50 | ✓ | Coumadin |
| v | 7.50 | 100 | ✓ | Marevan |
| * Tab 2 mg | 4.31 | 50 | ✓ | Coumadin |
| * Tab 3 mg | | 100 | ✓ | Marevan |
| * Tab 5 mg | | 50 | ✓ | Coumadin |
| v | 13.50 | 100 | ✓ | Marevan |
| | | | | |
| Blood Colony-stimulating Factors | | | | |
| FILGRASTIM – Special Authority see SA1259 below – Retail p | harmacv | | | |
| Ini 300 mcg per 0.5 ml prefilled svringe | 06.00 | 10 | 1 | Nivestim |

| Inj 300 mcg per 0.5 ml prefilled syringe | 10 | Nivestim |
|--|--------|------------------------------|
| Inj 480 mcg per 0.5 ml prefilled syringe | 10 | Nivestim |

➡SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5×10^9 /L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5×10^{9} /L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM – Special Authority see SA1912 below – Retail pharmacy

| Inj 6 mg per 0.6 ml syringe | 65.00 | 1 | Ziextenzo |
|-----------------------------|-------|---|-------------------------------|
|-----------------------------|-------|---|-------------------------------|

► SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

| GLUCOSE [DEXTROSE] | | |
|---|----|----------------------------|
| * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO | 5 | Biomed |
| ✤ Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO17.50 | 1 | Biomed |
| POTASSIUM CHLORIDE | | |
| * Inj 75 mg per ml, 10 ml65.00 | 50 | 🗸 Juno |

| | Subsidy | | Fully | Brand or |
|--|-----------------------|----------------|------------|-----------------------------------|
| | (Manufacturer's Pric | | sidised | Generic |
| | \$ | Per | / | Manufacturer |
| | 00.50 | | | |
| Inj 8.4%, 50 ml | 23.52 | 1 | ✓ B | iomed |
| a) Up to 5 inj available on a PSO | | | | |
| b) Not in combination | 04.40 | | | · · · · · · · · |
| Inj 8.4%, 100 ml | | 1 | • 8 | iomed |
| a) Up to 5 inj available on a PSOb) Not in combination | | | | |
| ODIUM CHLORIDE | | | | |
| Not funded for use as a nasal drop. Not funded for nebulise for nebuliser use. | er use except when | used in conji | unction | with an antibiotic intende |
| Inj 0.9%, bag – Up to 2000 ml available on a PSO | 1.33 | 500 ml | ✓ B | axter |
| | 1.36 | 1,000 ml | 🗸 В | axter |
| Only if prescribed on a prescription for renal dialysis, ma | aternity or post-nata | al care in the | home of | of the patient, or on a PS |
| for emergency use. (500 ml and 1,000 ml packs) | | | | |
| Inj 23.4% (4 mmol/ml), 20 ml ampoule | | 5 | ✓ B | iomed |
| For Sodium chloride oral liquid formulation refer Standa | | | | |
| Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO | | 20 | | resenius Kabi |
| Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO | | 50 | _ | resenius Kabi |
| Inj 0.9%, 20 ml ampoule | 5.00 | 20 | ✓ F | resenius Kabi |
| OTAL PARENTERAL NUTRITION (TPN) | | | | |
| Infusion | CBS | 1 OP | 🗸 I | PN |
| VATER | | | | |
| 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of e 4) When used for the dilution of sodium chloride soln 7% | | atients only. | | |
| Inj 10 ml ampoule – Up to 5 inj available on a PSO Inj 20 ml ampoule – Up to 5 inj available on a PSO | | 50 20 | _ | <u>lultichem</u> resenius Kabi |
| Oral Administration | | | | |
| ALCIUM POLYSTYRENE SULPHONATE | | | | |
| Powder | | 300 g OP | ✓ 0 | alcium Resonium |
| COMPOUND ELECTROLYTES | | • | | |
| Powder for oral soln – Up to 5 sach available on a PSO | | 50 | ✓ E | lectral |
| COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE | | | _ | <u></u> |
| Soln with electrolytes | - | ,000 ml OP | ✓ н | ydralyte - |
| | 0.00 | ,000 mi Oi | • 1 | Lemonade |
| PHOSPHORUS | | | | |
| Tab eff 500 mg (16 mmol) | 82 50 | 100 | / P | hosphate Phebra |
| | 02.00 | 100 | • F | noophate i nebia |
| POTASSIUM CHLORIDE | | 60 | | |
| | E 00 | | | |
| Fab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq) | | 60 | ~ | blancasant |
| | (17.10) | | - | hlorvescent |
| K Tab long-acting 600 mg (8 mmol) | (17.10) | 200 | - | hlorvescent pan-K |
| ₭ Tab long-acting 600 mg (8 mmol) SODIUM BICARBONATE | (17.10) 15.35 | 200 | ✓ S | pan-K |
| K Tab long-acting 600 mg (8 mmol) | (17.10) 15.35 | | ✓ s | |

| | Subsidy (Manufacturer's Price) | | | | | | Brand or Generic | |
|-------------------------------|-----------------------------------|----------|-------|--------------|--|--|---------------------|--|
| | \$ | Per | 1 | Manufacturer | | | | |
| SODIUM POLYSTYRENE SULPHONATE | | | | | | | | |
| Powder | | 454 g OP | 🗸 🗸 R | Resonium-A | | | | |

| | Subsidy (Manufacturer's Price \$ | e) Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|--|-----------|---------------------|--|
| Alpha-Adrenoceptor Blockers | | | | |
| Alpha Adrenoceptor Blockers | | | | |
| DOXAZOSIN | | | | |
| * Tab 2 mg | | 500 | | Doxazosin Clinect |
| * Tab 4 mg | 20.94 | 500 | ~ | Doxazosin Clinect |
| PHENOXYBENZAMINE HYDROCHLORIDE | | | | |
| * Cap 10 mg | | 30 | | BNM S29 |
| | 216.67 | 100 | - | Dibenzyline S29 |
| PRAZOSIN | | | - | |
| * Tab 1 mg | 5.53 | 100 | ~ | Arrotex-Prazosin |
| | | | | S29 S29 |
| W. Tab O are | 9.98 | 400 | | Minipress S29 |
| * Tab 2 mg | | 100 | ~ | Arrotex-Prazosin |
| | 10.00 | | | S29 S29 |
| * Tab 5 mg | 13.29 | 100 | | Minipress ^{©29} Arrotex-Prazosin |
| * Tab 5 mg | | 100 | • | S29 S29 |
| | 22.00 | | 1 | Minipress S29 |
| * Cap 1 mg | | 100 | | Prazosin Mylan S29 |
| * Cap 2 mg | | 100 | | Prazosin Mylan S29 |
| * Cap 5 mg | | 100 | | Prazosin Mylan S29 |
| | | | | , |
| Agents Affecting the Renin-Angiotensin Syste | m | | | |
| ACE Inhibitors | | | | |
| CAPTOPRIL | | | | |
| Oral liq 5 mg per ml Oral liquid restricted to children under 12 years of age. | | 00 ml (| OP 🗸 | DP-Captopril |
| CILAZAPRIL – Subsidy by endorsement | | | | |
| Subsidy by endorsement – Subsidised for patients who we endorsed accordingly. Pharmacists may annotate the pres | | | | |
| dispensing of cilazapril. | | | | |
| * Tab 0.5 mg | | 90 | | Zapril |
| * Tab 2.5 mg | | 90 | - | Zapril Zopril |
| | | 90 | • | Zapril |
| | 1 75 | 90 | | Acotoo |
| * Tab 5 mg * Tab 10 mg | | 90 90 | - | <u>Acetec</u> Acetec |
| * Tab 10 mg | | 90 90 | - | Acetec |
| LISINOPRIL | | | | |
| * Tab 5 mg | | 90 | 1 | Ethics Lisinopril |
| | | | | Teva Lisinopril |
| * Tab 10 mg | 11.67 | 90 | | Ethics Lisinopril |
| | | | | Teva Lisinopril |
| * Tab 20 mg | 14.69 | 90 | | Ethics Lisinopril Teva Lisinopril |

| | Subsidy | | Fully Brand or |
|--|-------------------------|------------|--|
| | (Manufacturer's Price) | Subsi | |
| | \$ | Per | Manufacturer |
| PERINDOPRIL | | | |
| | | | |
| * Tab 2 mg | 1.58 | 30 | <u>Coversyl</u> |
| * Tab 4 mg | 2.95 | 30 | Coversyl |
| * Tab 8 mg | | 30 | ✓ Coversyl |
| · · · · · · · · · · · · · · · · · · · | | 00 | e eereisyi |
| QUINAPRIL | | | |
| * Tab 5 mg | 5.97 | 90 | Arrow-Quinapril 5 |
| * Tab 10 mg | | 90 | ✓ Arrow-Quinapril 10 |
| | | | |
| * Tab 20 mg | | 90 | Arrow-Quinapril 20 |
| RAMIPRIL | | | |
| * Cap 1.25 mg | 6 90 | 90 | 🗸 Tryzan |
| | | | |
| * Cap 2.5 mg | | 90 | ✓ <u>Tryzan</u> |
| * Cap 5 mg | | 90 | <u>Tryzan</u> |
| * Cap 10 mg | 7.05 | 90 | Tryzan |
| 1 5 | | | |
| ACE Inhibitors with Diuretics | | | |
| ACE Inhibitors with Diuretics | | | |
| | • • | | |
| QUINAPRIL WITH HYDROCHLOROTHIAZIDE – Subsidy by er | | | |
| Subsidy by endorsement – Subsidised for patients who were | e taking guinapril with | hydrochlor | othiazide prior to 1 May |
| 2022 and the prescription is endorsed accordingly. Pharma | | | |
| exists a record of prior dispensing of quinapril with hydroch | | procompa | |
| | | | |
| Tab 10 mg with hydrochlorothiazide 12.5 mg | 4.10 | 30 | Accuretic 10 |
| Tab 20 mg with hydrochlorothiazide 12.5 mg | 5.25 | 30 | Accuretic 20 |
| | | | |
| Angiotensin II Antagonists | | | |
| Angiotensin il Antagonists | | | |
| | | | |
| CANDESARTAN CILEXETIL | | | |
| * Tab 4 mg | 2.00 | 90 | <u>Candestar</u> |
| * Tab 8 mg | 2.28 | 90 | Candestar |
| * Tab 16 mg | 3 31 | 90 | ✓ Candestar |
| * Tab 32 mg | | 90 | ✓ Candestar |
| * Tab 52 Tily | | 90 | |
| LOSARTAN POTASSIUM | | | |
| * Tab 12.5 mg | 2 00 | 84 | Losartan Actavis |
| 5 | | • · | |
| * Tab 25 mg | | 84 | Losartan Actavis |
| * Tab 50 mg | 2.86 | 84 | Losartan Actavis |
| * Tab 100 mg | 4.57 | 84 | Losartan Actavis |
| J. | | | |
| Angiotensin II Antagonists with Diuretics | | | |
| Anyiotensin ii Antayonists with Diuretics | | | |
| | | | |
| CANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDE | | | _ |
| * Tab 16 mg with hydrochlorothiazide 12.5 mg | 4.10 | 30 | APO-Candesartan |
| - · · · | | | HCTZ 16/12.5 |
| * Tab 22 ma with hydrochlorothia-ida 12 5 ma | 5.05 | 20 | ✓ APO-Candesartan |
| Tab 32 mg with hydrochlorothiazide 12.5 mg | | 30 | |
| | | | HCTZ 32/12.5 |
| LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE | | | |
| | 4.00 | 00 | A Museu Lesenten O |
| * Tab 50 mg with hydrochlorothiazide 12.5 mg | 4.00 | 30 | Arrow-Losartan & |
| | | | Hydrochlorothiazide |
| | | | |

| | Subsidy (Manufacturer's Price) | | Fully sidised | Brand or Generic |
|---|-----------------------------------|--------------|------------------|------------------------------|
| | \$ | Per | 1 | Manufacturer |
| Angiotensin II Antagonists with Neprilysin Inhib | oitors | | | |
| SACUBITRIL WITH VALSARTAN - Special Authority see SA230 | 02 below – Retail pha | armacy | | |
| Tab 24.3 mg with valsartan 25.7 mg | | 56 | | ntresto 24/26 |
| Tab 48.6 mg with valsartan 51.4 mg | | 56 | | ntresto 49/51 |
| Tab 97.2 mg with valsartan 102.8 mg | | 56 | ✓ E | ntresto 97/103 |
| ▶ SA2302 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals value the following criteria: | d without further rene | ewal unles | ss notifie | d for applications meeting |
| All of the following: | | | | |
| Patient has heart failure; and Any of the following: | | | | |
| 2.1 Patient is in NYHA/WHO functional class II; or | | | | |
| 2.2 Patient is in NYHA/WHO functional class III; or | | | | |
| 2.3 Patient is in NYHA/WHO functional class IV; and | | | | |
| 3 Either: | | | | |
| 3.1 Patient has a documented left ventricular ejection 13.2 An ECHO is not reasonably practical, and in the operation of the | | | | |
| treatment; and | | | | |
| 4 Patient is receiving concomitant optimal standard chronic | heart failure treatme | nts. | | |
| Antiarrhythmics | | | | |
| For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaes | sthetics Local page | 123 | | |
| AMIODARONE HYDROCHLORIDE | Salouoo, Looda, pago | 120 | | |
| Tab 100 mg | 2 40 | 30 | 1 A | ratac |
| ▲ Tab 100 mg | | 30 | _ | ratac |
| Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a | | 6 | _ | ordarone-X |
| | 15.22 | 10 | | lax Health |
| ATROPINE SULPHATE | 10122 | | | |
| * Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a | | | | |
| * Inj 600 micg per mi, 1 mi ampoule – Op to 5 mj available on a PSO | | 10 | ~ N | lartindale |
| | | 10 | • <u>II</u> | |
| DIGOXIN | 7 90 | 240 | | anavin BC |
| Tab 62.5 mcg – Up to 30 tab available on a PSO Tab 250 mcg – Up to 30 tab available on a PSO | | 240 240 | _ | anoxin PG anoxin |
| * Tab 250 mcg – Op to 30 tab available on a PSO * Oral lig 50 mcg per ml | | 240 60 ml | | anoxin |
| | | 00 111 | | anoxin Paediatric Elixir |
| | | | 🗸 L | anoxin S29 S29 |
| DISOPYRAMIDE PHOSPHATE | | | _ | |
| ▲ Cap 100 mg | 20.05 | 84 | ✔ R | ythmodan - Cheplafarm S29 |
| | 23.87 | 100 | ✓ R | ythmodan |

48

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|--|
| FLECAINIDE ACETATE | | | | |
| ▲ Tab 50 mg | | 60 | | Flecainide BNM Flecatab S29 |
| Cap long-acting 100 mg | | 90 | | Flecainide Controlled |
| ▲ Cap long-acting 200 mg | 54.28 | 90 | 1 | Release Teva Flecainide Controlled Release Teva |
| Inj 10 mg per ml, 15 ml ampoule MEXILETINE HYDROCHLORIDE | 108.16 | 5 | 1 | Tambocor |
| ▲ Cap 150 mg | 162.00 | 100 | 1 | Teva S29 |
| Cap 250 mg | | 100 | 1 | Teva S29 |
| Tab 150 mg | 40.90 | 50 | ✓ | Rytmonorm |
| Antihypotensives | | | | |
| MIDODRINE - Special Authority see SA1474 below - Retail phare | macy | | | |
| Tab 2.5 mg | | 100 | | MAR-Midodrine S29 Midodrine Medsurge |
| Tab 5 mg | 59.98 | 100 | 1 | Midodrine Medsurge |

SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

| ATENOLOL | | |
|---|-----------|--|
| * Tab 50 mg9.33 | 500 | Viatris |
| * Tab 100 mg14.20 | 500 | ✓ Atenolol Viatris ✓ Mylan Atenolol |
| * Oral liq 25 mg per 5 ml21.25 | 300 ml OP | ✓ Atenolol AFT S29 S29 |
| 38.20 | | Essential Generics (\$29) |
| 49.85 | | Atenolol AFT |
| Restricted to children under 12 years of age. | | |
| (Mylan Atenolol Tab 100 mg to be delisted 1 July 2024) | | |
| (Atenolol AFT S29 S29 Oral liq 25 mg per 5 ml to be delisted 1 August 2024) | | |
| BISOPROLOL FUMARATE | | |
| * Tab 2.5 mg | 90 | Ipca-Bisoprolol |
| * Tab 5 mg | 90 | Ipca-Bisoprolol |
| * Tab 10 mg2.71 | 90 | Ipca-Bisoprolol |

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| _ | | Subsidy | | Fully | |
|-----|---|------------------------------|-------|------------|-----------------------|
| | | (Manufacturer's Price) \$ | Per | Subsidised | |
| CAF | RVEDILOL | | | | |
| * | Tab 6.25 mg | 2.24 | 60 | 1 | Carvedilol Sandoz |
| * | Tab 12.5 mg | 2.30 | 60 | 1 | Carvedilol Sandoz |
| * | Tab 25 mg | 2.95 | 60 | ✓ | Carvedilol Sandoz |
| _AE | BETALOL | | | | |
| * | Tab 100 mg | | 100 | 1 | Trandate |
| * | Tab 200 mg | | 100 | 1 | Trandate |
| ₭ | Inj 5 mg per ml, 20 ml ampoule | | 5 | | |
| | | (88.60) | | | Trandate |
| ŧ | inj 5 mg per ml, 20 ml vial | | 1 | | |
| | | (48.20) | | | Alvogen S29 |
| ١E. | TOPROLOL SUCCINATE | | | | - |
| * | Tab long-acting 23.75 mg | 4.20 | 90 | 1 | Myloc CR |
| ŧ | Tab long-acting 47.5 mg | 3.65 | 90 | 1 | Myloc CR |
| ŧ | Tab long-acting 95 mg | 5.24 | 90 | 1 | Myloc CR |
| * | Tab long-acting 190 mg | 9.76 | 90 | 1 | Myloc CR |
| ٨E. | TOPROLOL TARTRATE | | | | |
| * | Tab 50 mg | 5.66 | 100 | ✓ | IPCA-Metoprolol |
| * | Tab 100 mg | 7.55 | 60 | ✓ | IPCA-Metoprolol |
| * | Tab long-acting 200 mg | 23.40 | 28 | ✓ | Slow-Lopresor |
| ¥ | Inj 1 mg per ml, 5 ml vial | | 5 | ✓ | Metoprolol IV Mylan |
| | | | | ✓ | Metoprolol IV Viatris |
| JAI | DOLOL | | | | |
| * | Tab 40 mg | 19.19 | 100 | | Nadolol BNM |
| * | Tab 80 mg | | 100 | 1 | Nadolol BNM |
| R | DPRANOLOL | | | | |
| * | Tab 10 mg | 7.04 | 100 | 1 | Drofate |
| ¥ | Tab 40 mg | 8.75 | 100 | 1 | IPCA-Propranolol |
| * | Cap long-acting 160 mg | | 100 | 1 | Cardinol LA |
| ¥ | Oral liq 4 mg per ml - Special Authority see SA1327 below - | - | | | |
| | Retail pharmacy | CBS ! | 500 m | nl 🗸 | Roxane- |
| | | | | | Propranolol S29 |

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

| * | Tab 80 mg | 500 | 🗸 <u>Mylan</u> |
|---|-----------------|-----|----------------|
| | Tab 160 mg14.00 | 100 | 🗸 <u>Mylan</u> |

| | Subsidy | | Fully | Brand or |
|--|-------------------------|--------------|------------|-------------------------|
| | (Manufacturer's Price) | | Subsidised | Generic |
| | \$ | Per | 1 | Manufacturer |
| | | | | |
| Calcium Channel Blockers | | | | |
| Dihydropyridine Calcium Channel Blockers | | | | |
| AMLODIPINE | | | | |
| * Tab 2.5 mg | 1 45 | 90 | 1 | Vasorex |
| | | 90 90 | | Vasorex |
| | | | - | |
| * Tab 10 mg | | 90 | • | Vasorex |
| FELODIPINE | | | _ | |
| * Tab long-acting 2.5 mg | 1.45 | 30 | ✓ | Plendil ER |
| * Tab long-acting 5 mg | 4.07 | 90 | ✓ | Felo 5 ER |
| * Tab long-acting 10 mg | | 90 | ✓ | Felo 10 ER |
| NIFEDIPINE | | | | |
| Tab long-acting 10 mg – Subsidy by endorsement | 10.40 | 56 | 1 | Tensipine MR10 S29 |
| * Tab long-acting to mg - Subsidy by endorsement | | 50 | • | |
| Subsidised for patients who were taking nifedipine tab | long-acting 10 mg prior | r to 1 | July 2023 | and the prescription is |
| endorsed accordingly. Pharmacists may annotate the | | | | |
| dispensing of nifedipine tab long-acting 10 mg. | | JU 11 | | |
| Tab long-acting 20 mg | 17 72 | 100 | 1 | Nyefax Retard |
| | | 14 | | Mylan Italy (24 hr |
| * Tab long-acting 30 mg | | 14 | • | |
| | | | _ | release) S29 |
| | 10.24 | 30 | ~ | Nifedipine |
| | | | | Viatris S29 |
| | 34.10 | 100 | ✓ | Mylan (24 hr |
| | | | | release) S29 |
| * Tab long-acting 60 mg | EO 01 | 100 | | • |
| * Tab long-acting 60 mg | | 100 | • | Mylan (24 hr |
| | | | | release) S29 |
| Other Calcium Channel Blockers | | | | |
| other Galcium Ghanner Diockers | | | | |
| DILTIAZEM HYDROCHLORIDE | | | | |
| * Cap long-acting 120 mg | 65.35 | 500 | ✓ | Diltiazem CD Clinect |
| * Cap long-acting 180 mg | | 30 | ✓ | Cardizem CD |
| * Cap long-acting 240 mg | | 30 | - | Cardizem CD |
| PERHEXILINE MALEATE | | | | |
| | 62.00 | 100 | | Dovoia |
| * Tab 100 mg | 02.90 | 100 | • | Pexsig |
| VERAPAMIL HYDROCHLORIDE | | | _ | |
| * Tab 40 mg | 7.01 | 100 | ✓ | Isoptin |
| * Tab 80 mg | 11.74 | 100 | ✓ | Isoptin |
| * Tab long-acting 120 mg | | 100 | ✓ | Isoptin Retard S29 |
| | | | | Isoptin SR |
| * Tab long-acting 240 mg | | 30 | - | Isoptin SR |
| Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on | | | | |
| PSO | | 5 | 1 | Isoptin |
| | | 5 | • | loopull |
| Centrally-Acting Agents | | | | |
| oonaany-Adang Agento | | | | |
| CLONIDINE | | | | |
| * Patch 2.5 mg, 100 mcg per day - Only on a prescription | 11.70 | 4 | ✓ | <u>Mylan</u> |
| Patch 5 mg, 200 mcg per day – Only on a prescription | | 4 | | Mylan |
| Patch 7.5 mg, 300 mcg per day – Only on a prescription | | 4 | | Mylan |
| and the may bee may be day to any on a problem. | | • | • | <u>,</u> |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | Fully | |
|--|------------------------------|-----|------------|-------------------------|
| | (Manufacturer's Price) \$ | Per | Subsidised | Generic Manufacturer |
| CLONIDINE HYDROCHLORIDE | | | | |
| * Tab 25 mcg | | 112 | ✓ | Clonidine Teva |
| * Tab 150 mcg | | 100 | 1 | Catapres |
| * Inj 150 mcg per ml, 1 ml ampoule | | 10 | ✓ | Medsurge |
| METHYLDOPA | | | | |
| * Tab 250 mg | | 100 | ✓ | Methyldopa Mylan |
| 0 | | | | Methyldopa Viatris |
| | 52.85 | 500 | | Methyldopa Mylan |
| | | | | S29 S29 |
| (Methyldopa Mylan Tab 250 mg to be delisted 1 September 2024 | 1) | | | |
| (Methyldopa Mylan S29 S29) Tab 250 mg to be delisted 1 Septe | , | | | |

Diuretics

Loop Diuretics

| BUMETANIDE | | | |
|---|---------|----------|---------------------------------------|
| * Tab 1 mg | 4.91 | 30 | Burinex S29 S29 |
| | 16.36 | 100 | Burinex |
| * Inj 500 mcg per ml, 4 ml vial | 7.95 | 5 | Burinex |
| (Burinex S29 S29) Tab 1 mg to be delisted 1 August 2024) | | | |
| FUROSEMIDE [FRUSEMIDE] | | | |
| Tab 40 mg – Up to 30 tab available on a PSO | 8.00 | 1,000 | IPCA-Frusemide |
| * Tab 500 mg | 25.00 | 50 | Urex Forte |
| | 89.48 | | Furosemid- |
| | | | Ratiopharm S29 |
| | 169.96 | 100 | ✓ Furosemid- |
| | | | Ratiopharm S29 |
| * Oral liq 10 mg per ml | | 30 ml OP | ✓ Lasix |
| * Inj 10 mg per ml, 25 ml ampoule | 60.65 | 6 | 🗸 Lasix |
| * Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a | PSO2.40 | 5 | Furosemide-Baxter |

Potassium Sparing Diuretics

| AMILORIDE HYDROCHLORIDE | | | |
|---|-------------------------------------|----------|----------------------------|
| Oral liq 1 mg per ml | | 25 ml OP | Biomed |
| EPLERENONE - Special Authority see SA1728 below | Retail pharmacy | | |
| Tab 25 mg | | 30 | Inspra |
| Tab 50 mg | 25.00 | 30 | Inspra |
| | | | |

► SA1728 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Patient has heart failure with ejection fraction less than 40%; and

- 2 Either:
 - 2.1 Patient is intolerant to optimal dosing of spironolactone; or

2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

| CARDIOVASCUL | AR SYSTEM |
|--------------|-----------|
|--------------|-----------|

| Subsidy (Manufacturer's Pri \$ | ce) S Per | Fully Subsidised | Brand or Generic Manufacturer |
|--------------------------------------|-------------------------|--|--|
| | 100 100 | ✓ | <u>Spiractin</u> <u>Spiractin</u> Biomed |
| | | | |
| | 28 | • | Frumil |
| | 50 | ✓ | Moduretic |
| | | | |
| 51.50 | 500 | • | <u>Arrow-</u> Bendrofluazide |
| • • | 500 | • | <u>Arrow-</u> Bendrofluazide |
| 29.21 | 25 ml O | P 🖌 | Biomed |
| 6.95 | 50 | ✓] | Hygroton |
| 16.00 | 90 | • | Dapa-Tabs |
| CBS | 1 50 | | Metolazone ^{S29} Zaroxolyn ^{S29} |
| | | | |
| | 28 OP 56 OP 56 OP | | Jinarc Jinarc Jinarc Jinarc Jinarc |
| | (Manufacturer's Pri | (Manufacturer's Price) S Per | (Manufacturer's Price) Subsidised 3.68 100 • |

► SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Either:

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price |) 8 | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

- 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
- 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents

| Fibrates | | |
|--|--|--|
| BEZAFIBRATE * Tab 200 mg | | ✓ <u>Bezalip</u> ✓ <u>Bezalip</u> Retard |
| Other Lipid-Modifying Agents | | |
| ACIPIMOX * Cap 250 mg | .56 30 5.44 | ✓ Olbetam S29 \$29 ✓ Olbetam |
| Resins | | |
| COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g | 2.89 30 | ✓ Colestid |
| Powder for oral suspension 4 g sachet | .50 50 | Colestyramine - Mylan \$299 Quantalan sugar free \$299 |
| HMG CoA Reductase Inhibitors (Statins) | | |
| ATORVASTATIN * Tab 10 mg | 0.24 500 0.92 500 0.54 500 7.16 100 | Lorstat Lorstat Lorstat Lorstat Clinect Clinect |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Brand or Subsidised Generic ✓ Manufacturer |
|--|---|-----|--|
| ROSUVASTATIN – Special Authority see SA2093 below – Reta | il pharmacy | | |
| * Tab 5 mg | 1.29 | 30 | Rosuvastatin Viatris |
| Rosuvastatin Viatris to be Principal Supply on 1 Octobe | r 2024 | | |
| * Tab 10 mg | 1.69 | 30 | Rosuvastatin Viatris |
| Rosuvastatin Viatris to be Principal Supply on 1 Octobe | r 2024 | | |
| * Tab 20 mg | 2.71 | 30 | Rosuvastatin Viatris |
| * Tab 40 mg | | 30 | Rosuvastatin Viatris |

⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient is considered to be at risk of cardiovascular disease; and
- 1.2 Patient is Māori or any Pacific ethnicity; or

2 Both:

- 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
- 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

1 Any of the following:

- 1.1 Patient has proven coronary artery disease (CAD); or
- 1.2 Patient has proven peripheral artery disease (PAD); or
- 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke,
- coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

| (Ma | Subsidy anufacturer's Price | e) | Fully Subsidised | |
|--|--------------------------------|---------|---------------------|------------------------|
| | \$ | Per | 1 | Manufacturer |
| SIMVASTATIN | | | | |
| * Tab 10 mg | 1.68 | 90 | 1 | Simvastatin Mylan |
| | | | | Simvastatin Viatris |
| ₭ Tab 20 mg | 2.54 | 90 | | Simvastatin Viatris |
| 🖌 Tab 40 mg | 4.11 | 90 | | Simvastatin Mylan |
| | | | 1 | Simvastatin Viatris |
| Simvastatin Viatris to be Principal Supply on 1 June 2024 | | | | • · · · • |
| ₭ Tab 80 mg | 8.81 | 90 | | Simvastatin Mylan |
| Oliverse table Mistic to be Driveled Orentz and these 0004 | | | ~ | Simvastatin Viatris |
| Simvastatin Viatris to be Principal Supply on 1 June 2024 | | | | |
| Simvastatin Mylan Tab 40 mg to be delisted 1 December 2024) | | | | |
| Simvastatin Mylan Tab 80 mg to be delisted 1 September 2024) | | | | |
| Selective Cholesterol Absorption Inhibitors | | | | |
| ZETIMIBE | | | | |
| ₭ Tab 10 mg | 1 76 | 30 | 1 | Ezemibe Viatris |
| | | 00 | | Ezetimibe Sandoz |
| | | | • | |
| | | | | - |
| Tab 10 mg with simvastatin 10 mg | | 30 | | Zimybe |
| Tab 10 mg with simvastatin 20 mg | | 30 | | Zimybe |
| Tab 10 mg with simulatatin 40 mg | | 30 | | Zimybe |
| Tab 10 mg with simvastatin 80 mg | 8.15 | 30 | • | Zimybe |
| Nitrates | | | | |
| | | | | |
| GLYCERYL TRINITRATE | | | | |
| Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO | 7 40 05 | i0 dose | | Nitrolingual Pump |
| | | o uose | Ur • | • |
| C Datab 05 mg 5 mg par day | 15 70 | 30 | | Spray Nitroderm TTS |
| Patch 25 mg, 5 mg per day Datch 50 mg, 10 mg per day | | | | |
| Patch 50 mg, 10 mg per day | 18.62 | 30 | • | Nitroderm TTS |
| SOSORBIDE MONONITRATE | | | | |
| ₭ Tab 20 mg | | 100 | | Ismo 20 |
| K Tab long-acting 40 mg | | 30 | | Ismo 40 Retard |
| Fab long-acting 60 mg | 13.50 | 90 | ~ | Duride |
| Sympathomimetics | | | | |
| | | | | |
| ADRENALINE | 4.00 | - | | Aanan Advanatin- |
| Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO | | 5 | | Aspen Adrenaline |
| | 12.65 | - | | DBL Adrenaline |
| Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a PSO . | 27.00 49.00 | 5 10 | - | Hospira |
| | | | ~ | Aspen Adrenaline |

56

| Tab 25 mg - Special Authority see SA1321 below - Retail pharmacy | | Subsidy | | Fully | Brand or |
|---|--|---------------------------------------|--------|-----------------|----------------------------|
| Vasodilators HYDRALAZINE HYDROCHLORIDE * Tab 25 mg - Special Authority see SA1321 below - Retail pharmacy | | · · · · · · · · · · · · · · · · · · · | _ | | |
| HYDRALAZINE HYDROCHLORIDE * Tab 25 mg - Special Authority see SA1321 below - Retail pharmacy | | \$ | Per | | Manufacturer |
| HYDRALAZINE HYDROCHLORIDE * Tab 25 mg - Special Authority see SA1321 below - Retail pharmacy | Vasodilators | | | | |
| Tab 25 mg - Special Authority see SA1321 below - Retail pharmacy | Vasoullators | | | | |
| pharmacy | HYDRALAZINE HYDROCHLORIDE | | | | |
| 56 ✓ Onelink S™ 84 ✓ AMDIPHARM S™ 100 ✓ Camber S™ 100 ✓ Camber S™ >SA1321 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meetir he following criteria: Either: 1 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to AG inhibitors and/or angiotensin receptor blockers. WINOXIDIL Tab 10 mg Tab 10 mg 47.04 60 ✓ Minoxidil Roma S™ 78.40 100 ✓ Loniten VICORANDIL – Brand switch fee payable (Pharmacode 2677903) - see page 265 for details Tab 20 mg 21.73 60 ✓ Max Health PAPAVERINE HYDROCHLORIDE 21.73 60 ✓ Max Health PAPAVERINE HYDROCHLORIDE 25.712 5 ✓ Hospira PENTOXIFYLLINE [OXPENTIFYLLINE] 42.26 50 ✓ Trental 400 Endothelin Receptor Antagonists 200.00 30 ✓ Ambrisentan Viatris | * Tab 25 mg – Special Authority see SA1321 below – Retail | | | | |
| 84 ✓ AMDIPHARM 529 100 ✓ Camber 529 >SA1321 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meetir he following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of neart failure in combination with a nitrate, in patients who are intolerant or have not responded to AG inhibitors and/or angiotensin receptor blockers. WINOXIDIL 47.04 60 ✓ Minoxidil Roma 529 1 Tab 10 mg 47.04 60 ✓ Loniten VICORANDIL – Brand switch fee payable (Pharmacode 2677903) - see page 265 for details 4 100 ✓ Loniten VICORANDIL – Brand switch fee payable (Pharmacode 2677903) - see page 265 for details 4 100 ✓ Loniten VICORANDIL – Brand switch fee payable (Pharmacode 2677903) - see page 265 for details 4 100 ✓ Loniten ATab 20 mg 21.73 60 ✓ Max Health 4 PAPAVERINE HYDROCHLORIDE 4 4 4 4 4 * Inj 12 mg per ml, 10 ml ampoule 257.12 5 ✓ Hospira *ENTOXIFYLLINE [OXPENTIFYLLINE] 42.26 50 ✓ Trental 400 | pharmacy | CBS | 1 | ✓ Н | ydralazine |
| 100 Camber 329 ** Inj 20 mg ampoule 25.90 ** SA1321 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meetir the following criteria: Either: 1 2 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to Ad inhibitors and/or angiotensin receptor blockers. VIIOXIDIL Tab 10 mg * Tab 10 mg 47.04 60 * Tab 10 mg 21.73 60 * Tab 20 mg 27.44 60 * Tab 20 mg 257.12 5 * Hospira 257.12 5 * Hospira 257.12 5 * Tab 400 mg 42.26 50 * Trental 400 Candber Mill 92 92 * Hospira * Particular Experimentary 42.26 50 * Trental 400 * Tab 400 mg 200 | | | 56 | √ 0 | nelink S29 |
| Inj 20 mg ampoule | | | 84 | 🗸 A | MDIPHARM S29 |
| SA1321 Special Authority for Subsidy mitial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to Ad inhibitors and/or angiotensin receptor blockers. VINOXIDIL Tab 10 mg 47.04 60 / Minoxidil Roma ^{\$29} 78.40 100 / Loniten VICORANDIL – Brand switch fee payable (Pharmacode 2677903) - see page 265 for details Tab 10 mg 21.73 60 / Max Health Tab 20 mg 27.44 60 / Max Health PAPAVERINE HYDROCHLORIDE * Inj 12 mg per ml, 10 ml ampoule 257.12 5 / Hospira PENTOXIFYLLINE [OXPENTIFYLLINE] Tab 400 mg 42.26 50 / Trental 400 Endothelin Receptor Antagonists | | | 100 | ✓ C | amber S29 |
| nitial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to AC inhibitors and/or angiotensin receptor blockers. WINOXIDIL ▲ Tab 10 mg | * Inj 20 mg ampoule | | 5 | 🗸 A | presoline |
| nitial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to AC inhibitors and/or angiotensin receptor blockers. WINOXIDIL ▲ Tab 10 mg | SA1321 Special Authority for Subsidy | | | | |
| he following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to AC inhibitors and/or angiotensin receptor blockers. WINOXIDIL ▲ Tab 10 mg | | d without further rene | wal u | Inless notified | d for applications meeting |
| 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to AG inhibitors and/or angiotensin receptor blockers. WINOXIDIL Tab 10 mg 47.04 60 ✓ Minoxidil Roma 🐲 Tab 10 mg 78.40 100 ✓ Loniten VICORANDIL – Brand switch fee payable (Pharmacode 2677903) - see page 265 for details ✓ Loniten Tab 10 mg 21.73 60 ✓ Max Health Tab 20 mg 27.44 60 ✓ Max Health PAPAVERINE HYDROCHLORIDE 27.44 60 ✓ Max Health * Inj 12 mg per ml, 10 ml ampoule 257.12 5 ✓ Hospira PENTOXIFYLLINE [OXPENTIFYLLINE] 42.26 50 ✓ Trental 400 Endothelin Receptor Antagonists 200.00 30 ✓ Ambrisentan Viatris AMBRISENTAN – Special Authority see SA2253 below – Retail pharmacy 200.00 30 ✓ Ambrisentan Viatris Tab 10 mg 200.00 30 ✓ Ambrisentan Viatris | the following criteria: | | | | |
| 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to Ad inhibitors and/or angiotensin receptor blockers. VIINOXIDIL Tab 10 mg 47.04 60 ✓ Minoxidil Roma 🖘 A Tab 10 mg 78.40 100 ✓ Loniten VICORANDIL – Brand switch fee payable (Pharmacode 2677903) - see page 265 for details ✓ Max Health A Tab 10 mg 21.73 60 ✓ Max Health A Tab 20 mg 27.44 60 ✓ Max Health PAPAVERINE HYDROCHLORIDE 27.44 60 ✓ Max Health * Inj 12 mg per ml, 10 ml ampoule 257.12 5 ✓ Hospira PENTOXIFYLLINE [OXPENTIFYLLINE] 42.26 50 ✓ Trental 400 Endothelin Receptor Antagonists AMBRISENTAN – Special Authority see SA2253 below – Retail pharmacy 200.00 30 ✓ Ambrisentan Viatris Tab 10 mg 200.00 30 ✓ Ambrisentan Viatris | Either: | | | | |
| inhibitors and/or angiotensin receptor blockers. WINOXIDIL ▲ Tab 10 mg | 1 For the treatment of refractory hypertension; or | | | | |
| inhibitors and/or angiotensin receptor blockers. WINOXIDIL ▲ Tab 10 mg | 2 For the treatment of heart failure in combination with a nit | rate, in patients who a | are in | tolerant or ha | ave not responded to ACE |
| Tab 10 mg | | | | | |
| 78.40 100 ✓ Loniten NICORANDIL – Brand switch fee payable (Pharmacode 2677903) - see page 265 for details ✓ Tab 10 mg 21.73 60 ✓ Tab 20 mg 27.44 60 ✓ PAPAVERINE HYDROCHLORIDE ✓ Max Health * Inj 12 mg per ml, 10 ml ampoule .257.12 5 ✓ PENTOXIFYLLINE [OXPENTIFYLLINE] .42.26 50 ✓ Trental 400 Endothelin Receptor Antagonists AMBRISENTAN – Special Authority see SA2253 below – Retail pharmacy Tab 5 mg .200.00 30 ✓ Ambrisentan Viatris Tab 10 mg .200.00 30 ✓ Ambrisentan Viatris | MINOXIDIL | | | | |
| 78.40 100 ✓ Loniten NICORANDIL – Brand switch fee payable (Pharmacode 2677903) - see page 265 for details ✓ Tab 10 mg 21.73 60 ✓ Tab 20 mg 27.44 60 ✓ PAPAVERINE HYDROCHLORIDE ✓ Max Health * Inj 12 mg per ml, 10 ml ampoule .257.12 5 ✓ PENTOXIFYLLINE [OXPENTIFYLLINE] .42.26 50 ✓ Trental 400 Endothelin Receptor Antagonists AMBRISENTAN – Special Authority see SA2253 below – Retail pharmacy Tab 5 mg .200.00 30 ✓ Ambrisentan Viatris Tab 10 mg .200.00 30 ✓ Ambrisentan Viatris | ▲ Tab 10 mg | 47 04 | 60 | 🖌 M | inoxidil Roma S29 |
| ▲ Tab 10 mg | | | | | |
| ▲ Tab 10 mg | NICORANDII - Brand switch fee payable (Pharmacode 267790 | 3) - see nage 265 for | dota | ile – | |
| ▲ Tab 20 mg | | | | | ax Health |
| PAPAVERINE HyDROCHLORIDE ** Inj 12 mg per ml, 10 ml ampoule | 5 | | | | |
| * Inj 12 mg per ml, 10 ml ampoule | - | | | | |
| PENTOXIFYLLINE [OXPENTIFYLLINE] Tab 400 mg Endothelin Receptor Antagonists AMBRISENTAN – Special Authority see SA2253 below – Retail pharmacy Tab 5 mg Tab 10 mg 200.00 30 ✓ Ambrisentan Viatris Ambrisentan Viatris | | 257 10 | 5 | / u | ocnira |
| Tab 400 mg 42.26 50 Trental 400 Endothelin Receptor Antagonists AMBRISENTAN – Special Authority see SA2253 below – Retail pharmacy 30 <u>Ambrisentan Viatris</u> Tab 5 mg 200.00 30 <u>Ambrisentan Viatris</u> Tab 10 mg 200.00 30 <u>Ambrisentan Viatris</u> | | | 5 | • 1 | ospila |
| Endothelin Receptor Antagonists AMBRISENTAN – Special Authority see SA2253 below – Retail pharmacy Tab 5 mg | | 40.00 | | <i>.</i> | |
| AMBRISENTAN – Special Authority see SA2253 below – Retail pharmacy Tab 5 mg | l ab 400 mg | | 50 | ✓ 1 | rental 400 |
| AMBRISENTAN – Special Authority see SA2253 below – Retail pharmacy Tab 5 mg | Endethelin Decenter Antegoniste | | | | |
| Tab 5 mg30✓Ambrisentan ViatrisTab 10 mg30✓Ambrisentan Viatris | Endothelin Receptor Antagonists | | | | |
| Tab 5 mg30✓Ambrisentan ViatrisTab 10 mg30✓Ambrisentan Viatris | AMBRISENTAN - Special Authority see SA2253 below - Retail | pharmacy | | | |
| 5 | | | 30 | 🗸 A | mbrisentan Viatris |
| xxCA2252 Special Authority for Subcidy | Tab 10 mg | | 30 | 🗸 🖌 | mbrisentan Viatris |
| TOMACON DURING MULTURING DUDSION | ➡SA2253 Special Authority for Subsidy | | | | |

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and

4 Any of the following:

4.1 All of the following:

- 4.1.1 PAH has been confirmed by right heart catheterisation; and
- 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | F | ully Brand or | |
|-----------------|------------------|---------------------------------|----|
| (Manufacturer's | S Price) Subsidi | sed Generic | |
| \$ | Per | Manufacture | er |

cm⁻⁵); and

- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Ambrisentan is to be used as PAH dual therapy; and
 - 5.2 Either:

continued...

58

| Subsidy | | Fully | Brand or | |
|------------------------|------|---------|--------------|--|
| (Manufacturer's Price) | Subs | sidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 5.2.1 Patient has tried a PAH monotherapy (sildenafil or bosentan) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; or
- 5.2.2 Patient has tried PAH dual therapy including bosentan and has experienced intolerable side effects on bosentan; and
- 5.3 Both:
 - 5.3.1 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy; and
 - 5.3.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease).

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Both:
 - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

continued...

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and</u> treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

| Tab 62.5 mg | 60 | ✓ Bosentan Dr Reddy's |
|-------------|--------|--|
| Tab 125 mg | 60 | ✓ <u>Bosentan Dr</u> <u>Reddy's</u> |

SA2254 Special Authority for Subsidy

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:

- 5.1 Bosentan is to be used as PAH monotherapy; and
- 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil; or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | Su | ubsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Either:
 - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool**; or
 - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | 5 | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

- 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:

- 5.1 Bosentan is to be used as part of PAH triple therapy; and
- 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and</u> treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL - Special Authority see SA2255 below - Retail pharmacy

| Tab 25 mg | 4 | 🗸 Vedafil |
|---------------|----|-----------|
| Tab 50 mg1.70 | 4 | 🗸 Vedafil |
| • | 12 | ✓ Vedafil |

➡SA2255 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

1 Patient has pulmonary arterial hypertension (PAH)*; and

continued...

62

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Su | lbsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH is confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Note: Indications marked with * are Unapproved Indications.

† The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and treatment of</u> pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Prostacyclin Analogues

| EPOPROSTENOL - Special Authority see SA2256 below - Re | etail pharmacy | | |
|--|----------------|---|-----------------------------|
| Inj 500 mcg vial | | 1 | Veletri |
| Inj 1.5 mg vial | 73.21 | 1 | 🗸 Veletri |

⇒SA2256 Special Authority for Subsidy

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- $2\;$ PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:

| | | ully Br | and or |
|----------|-------------------------|------------------------|-------------|
| (Manufac | cturer's Price) Subsidi | sed Ge | eneric |
| | \$ Per | Ma | anufacturer |

- 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵): and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these auidelines) + : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and
- 5 All of the following:
 - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initial application - (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV: and
- 4 Any of the following:
 - 4.1 All of the following:

Principal Supply

- 4.1.1 PAH has been confirmed by right heart catheterisation; and
- 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † ; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major

| Subsidy | | Fully | Brand or | |
|------------------------|------|---------|--------------|--|
| (Manufacturer's Price) | Subs | sidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and

- 5 Both:
 - 5.1 Epoprostenol is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and</u> treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST – Special Authority see SA2257 below – Retail pharmacy

🗸 Vebulis

30

⇒SA2257 Special Authority for Subsidy

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

| Subsidy | y Fu | ly Brand or |
|----------------|--------------------|--------------|
| (Manufacturer' | s Price) Subsidise | d Generic |
| \$ | Per | Manufacturer |

5 Both:

- 5.1 Iloprost is to be used as PAH monotherapy; and
- 5.2 Either:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Either:
 - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
 - 5.3 Either:
 - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; or
 - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

66

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Iloprost is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: <u>2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH</u>

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

| | Subsidy (Manufacturer's Price \$ |) Sub Per | Fully osidised | Brand or Generic Manufacturer |
|---|--|--------------|--------------------------------|-------------------------------------|
| Antiacne Preparations | | | | |
| For systemic antibacterials, refer to INFECTIONS, Antibacterial | s, page 95 | | | |
| ADAPALENE | | | | |
| a) Maximum of 30 g per prescription | | | | |
| b) Only on a prescription | | | | |
| Gel 0.1% | | 30 g OP | ✓ [| Differin |
| ISOTRETINOIN - Special Authority see SA2023 below - Retai | l pharmacy | | | |
| Cap 5 mg | | 60 | ✓ (| Dratane |
| Cap 10 mg | | 120 | ✓ | Dratane |
| Cap 20 mg | | 120 | | Dratane |

➡SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

TRETINOIN

| Crm 0.5 mg per g $-$ Maximum of 50 g per prescription . | 15.57 | 50 g OP | ✓ <u>ReTrieve</u> | |
|--|-----------------|---------|-------------------|--|
| Antibacterials Topical | | | | |
| For systemic antibacterials, refer to INFECTIONS, Antibacter | erials, page 95 | | | |
| HYDROGEN PEROXIDE * Crm 1% | 8 56 | 10 g OP | ✓ Crystaderm | |
| MUPIROCIN | | 10 9 01 | • Orystaderin | |
| Oint 2% | | 15 g OP | | |
| | (13.00) | | Bactroban | |
| a) Only on a prescription | | | | |

b) Not in combination

68

DERMATOLOGICALS

| | Qubaidu | | | — |
|---|------------------------------|-------------|-----------------------------------|---|
| | Subsidy (Manufacturer's I | Price) Subs | Fully Brand or sidised Generic | |
| | \$ | Per | Manufacturer | |
| SODIUM FUSIDATE [FUSIDIC ACID] | | | | _ |
| Crm 2% | 1.59 | 5 g OP | Foban | |
| a) Maximum of 5 g per prescription | | - 5 - | | |
| b) Only on a prescription | | | | |
| c) Not in combination | | | | |
| Oint 2% | 1.59 | 5 g OP | Foban | |
| a) Maximum of 5 g per prescription | | | | |
| b) Only on a prescription | | | | |
| c) Not in combination | | | | |
| SULFADIAZINE SILVER | 40.00 | | | |
| Crm 1% | | 50 g OP | Flamazine | |
| a) Up to 250 g available on a PSO | | | | |
| b) Not in combination | | | | |
| Antifungals Topical | | | | |
| | | | | |
| For systemic antifungals, refer to INFECTIONS, Antifungals, | page 102 | | | |
| AMOROLFINE | | | | |
| a) Only on a prescription | | | | |
| b) Not in combination | | | | |
| Nail soln 5% | 21.87 | 5 ml OP | ✓ MycoNail | |
| CLOTRIMAZOLE | | | | |
| * Crm 1% | 1.10 | 20 g OP | Clomazol | |
| a) Only on a prescription | | | | |
| b) Not in combination | 4.00 | | | |
| * Soln 1% | | 20 ml OP | Conastan | |
| a) Only on a proparintian | (7.55) | | Canesten | |
| a) Only on a prescriptionb) Not in combination | | | | |
| , | | | | |
| ECONAZOLE NITRATE Crm 1% | 1.00 | 20 g OP | | |
| 0111176 | (8.09) | 20 9 01 | Pevaryl | |
| a) Only on a prescription | (0.00) | | i ovalji | |
| b) Not in combination | | | | |
| Foaming soln 1%, 10 ml sachets | 9.89 | 3 | | |
| | (18.64) | | Pevaryl | |
| a) Only on a prescription | | | | |
| b) Not in combination | | | | |
| MICONAZOLE NITRATE | | | _ | |
| * Crm 2% | 0.90 | 15 g OP | Multichem | |
| a) Only on a prescription | | | | |
| b) Not in combination | 4.00 | | | |
| * Lotn 2% | 4.36 (10.03) | 30 ml OP | Daktarin | |
| a) Only on a prescription | (10.03) | | Dariann | |
| b) Not in combination | | | | |
| * Tinct 2% | 4.36 | 30 ml OP | | |
| | (12.10) | - | Daktarin | |
| a) Only on a prescription | | | | |
| b) Not in combination | | | | |
| | | | | |

| | Subsidy (Manufacturer's P \$ | rice) Sub: Per | Fully Brand or sidised Generic ✓ Manufacturer |
|---|------------------------------------|--------------------|---|
| Antipruritic Preparations | | | |
| CALAMINE | | | |
| a) Only on a prescription | | | |
| b) Not in combination | | | |
| Crm, aqueous, BP | 3.45 | 100 g | healthE Calamine Aqueous |
| ROTAMITON | | | |
| a) Only on a prescription | | | |
| b) Not in combination | | | 4 |
| Crm 10% | 3.29 | 20 g OP | ✓ <u>Itch-Soothe</u> |
| ENTHOL – Only in combination | | | |
| Only in combination with a dermatological base or p With or without other dermatological galenicals. | roprietary Topical C | orticosteriod – | Plain |
| Crystals | 6.92 | 25 g | ✓ MidWest |
| | 29.60 | 100 g | MidWest |
| Corticosteroids Topical | | | |
| or systemic corticosteroids, refer to CORTICOSTEROIDS A | ND RELATED AGE | NTS, page 85 | |
| Corticosteroids - Plain | | | |
| ETAMETHASONE DIPROPIONATE | | | |
| Crm 0.05% | | 15 g OP | Diprosone |
| | 36.00 | 50 g OP | Diprosone |
| Diprosone to be Principal Supply on 1 July 2024 | 0.00 | | |
| Oint 0.05% | 2.96 36.00 | 15 g OP 50 g OP | Diprosone Diprosone |
| Diprosone to be Principal Supply on 1 July 2024 | 30.00 | 50 y OF | |
| Oint 0.05% in propylene glycol base | | 30 g OP | Diprosone OV |
| ETAMETHASONE VALERATE | | J - | |
| Crm 0.1% | 4.53 | 50 g OP | Beta Cream |
| Cont 0.1% | | 50 g OP | ✓ Beta Ointment |
| € Lotn 0.1% | | 50 ml OP | ✓ Betnovate |
| LOBETASOL PROPIONATE | | | |
| € Cm 0.05% | | 30 g OP | Dermol |
| € Oint 0.05% | | 30 g OP | ✓ Dermol |
| LOBETASONE BUTYRATE | | | |
| Crm 0.05% | 5 38 | 30 g OP | |
| | (10.00) | 50 g O. | Eumovate |
| YDROCORTISONE | () | | · · ···• |
| Crm 1% – Only on a prescription | 1 78 | 30 g OP | Ethics |
| | 20.40 | 500 g | ✓ <u>Lunes</u> ✓ Noumed |
| Powder – Only in combination | | 25 g | ✓ ABM |
| Up to 5% in a dermatological base (not proprietary To | | 0 | |
| galenicals | - | , | 0 |

70

DERMATOLOGICALS

| | Subsidy | | Fully | Brand or |
|--|-------------------------|---------------------|-----------------------|-------------------------|
| | (Manufacturer's I \$ | Price) Subs Per | sidised ✓ | Generic Manufacturer |
| HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN | | | | |
| Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only | on | | | |
| a prescription | | 250 ml | I | DP Lotn HC |
| DP Lotn HC to be Principal Supply on 1 June 2024 | | | | |
| HYDROCORTISONE BUTYRATE | | | | |
| Lipocream 0.1% | | 100 g OP | | Locoid Lipocream |
| Oint 0.1% | | 100 g OP | | Locoid |
| Milky emul 0.1% | 12.33 | 100 ml OP | ✓ | Locoid Crelo |
| METHYLPREDNISOLONE ACEPONATE | | | - | |
| Crm 0.1% | | 15 g OP | - | Advantan |
| Oint 0.1% | 4.95 | 15 g OP | | Advantan |
| MOMETASONE FUROATE | | | - | |
| Crm 0.1% | | 15 g OP | | Elocon Alcohol Free |
| Oint 0.19/ | 3.10 | 50 g OP | | Elocon Alcohol Free |
| Oint 0.1% | 1.95 2.90 | 15 g OP 50 g OP | | <u>Elocon</u> Elocon |
| Lotn 0.1% | | 30 g OP 30 ml OP | | Elocon |
| | 4.50 | 30 111 01 | • ! | |
| TRIAMCINOLONE ACETONIDE Crm 0.02% | 6 40 | 100 g OP | 1 | Aristocort |
| Oint 0.02% | | 100 g OP | | Aristocort |
| | 0.01 | 100 g 01 | | |
| Corticosteroids - Combination | | | | |
| BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FL | JSIDIC ACIDI | | | |
| Crm 0.1% with sodium fusidate (fusidic acid) 2% | | 15 g OP | | |
| | (10.45) | - | I | Fucicort |
| a) Maximum of 15 g per prescription | | | | |
| b) Only on a prescription | | | | |
| HYDROCORTISONE WITH MICONAZOLE - Only on a prescri | otion | | | |
| * Crm 1% with miconazole nitrate 2% | 1.89 | 15 g OP | ✓] | Micreme H |
| HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - C | Only on a prescri | ption | | |
| Oint 1% with natamycin 1% and neomycin sulphate 0.5% | | 15 g OP | I | Pimafucort |
| TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC | IN AND NYSTA | TIN | | |
| Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 m | g | | | |
| and gramicidin 250 mcg per g – Only on a prescription | | 15 g OP | | |
| | (9.28) | | ` | /iaderm KC |
| Parrier Creame and Emellionte | | | | |
| Barrier Creams and Emollients | | | | |
| Barrier Creams | | | | |
| DIMETHICONE | | | | |
| * Crm 5% pump bottle | 4.30 | 500 ml OP | ~ 1 | nealthE |
| | | | • ! | Dimethicone 5% |
| * Crm 10% pump bottle | | 500 ml OP | ✓ 1 | nealthE |
| | | | | Dimethicone 10% |
| ZINC AND CASTOR OIL | | | | |
| * Oint | 4.25 | 500 g | I | Evara |
| | | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

DERMATOLOGICALS

| | Subsidy (Manufacturer's Pr | rice) Subsi | Fully Brand or dised Generic |
|---|-------------------------------|-------------|--|
| | | Per | Manufacturer |
| Emollients | | | |
| AQUEOUS CREAM | | | |
| Crm | 1.30 | 100 g | healthE Aqueous Cream SLS Free |
| | 1.73 | 500 g | ✓ Evara ✓ <u>GEM Aqueous</u> <u>Cream</u> |
| CETOMACROGOL | | | |
| * Crm BP | 1.99 | 500 g | Cetomacrogol-AFT |
| CETOMACROGOL WITH GLYCEROL | | | |
| Crm 90% with glycerol 10% | 2.13 | 500 ml OP | ✓ Evara |
| | 3.50 | 1,000 ml OP | ✓ Evara |
| EMULSIFYING OINTMENT | | | |
| * Oint BP | 3.13 | 500 g | ✓ Emulsifying Ointment ADE |
| DIL IN WATER EMULSION | | | _ |
| * Crm | 2.04 | 500 g | Fatty Cream AFT |
| PARAFFIN | | | |
| Oint liquid paraffin 50% with white soft paraffin 50% | 4.94 | 500 g OP | ✓ White Soft Liquid Paraffin AFT |
| JREA | | | |
| * Crm 10% | 1.37 | 100 g OP | healthE Urea Cream |
| NOOL FAT WITH MINERAL OIL – Only on a prescription | | | |
| * Lotn hydrous 3% with mineral oil | 5.60 | 1,000 ml | |
| | (14.96) | | DP Lotion |
| | (20.53) | | Alpha-Keri Lotion |
| | 1.40 | 250 ml OP | |
| | (5.87) | 4 000 | DP Lotion |
| | 5.60 | 1,000 ml | DIC Lation |
| | (23.91) | | BK Lotion |
| | 1.40 | 250 ml OP | BK Lotion |
| | (7.73) | | |
| Other Dermatological Bases | | | |
| PARAFFIN | | | |
| White soft – Only in combination | 4.74 | 450 g | EVARA White Soft Paraffin |
| | 4.99 | | healthE |
| | 19.00 | 2,500 g | EVARA White Soft Paraffin |
| | 19.99 | | healthE |

(healthE White soft to be delisted 1 June 2024) (healthE White soft to be delisted 1 June 2024)

DERMATOLOGICALS

| | Subsidy | | Fully | Brand or |
|--|----------------------|-----------------|------------|--|
| | (Manufacturer's Pric | e) Subs Per | sidised | Generic Manufacturer |
| | Ψ | | - | Manulacturer |
| Minor Skin Infections | | | | |
| POVIDONE IODINE | | | | |
| Oint 10% | 7.40 | 65 g OP | 🗸 E | Betadine |
| a) Maximum of 130 g per prescriptionb) Only on a prescription | | | | |
| Antiseptic Solution 10% | 4.15 | 100 ml | ✓ <u>F</u> | Riodine |
| Antiseptic soln 10% | 3.83 | 15 ml | - | Riodine |
| | 6.99 | 500 ml | ✓ F | Riodine |
| Skin preparation, povidone iodine 10% with 30% alcohol | | 100 ml | _ | |
| | (3.48) | | E | Betadine Skin Prep |
| Skin preparation, povidone iodine 10% with 70% alcohol | 1.63 (7.78) | 100 ml | F | Pfizer |
| Parasiticidal Preparations | | | | |
| DIMETHICONE | | | | |
| * Lotn 4% | 4.25 | 200 ml OP | ✓ <u>ł</u> | <u>nealthE</u> Dimethicone 4% Lotion |
| IVERMECTIN - Special Authority see SA2294 below - Retail p | harmacy | | | |
| Tab 3 mg – Up to 100 tab available on a PSO | | 4 | √ 9 | Stromectol |
| 1) PSO for institutional use only. Must be endorsed | with the name of the | e institution f | or whic | h the PSO is required an |

- PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

➡SA2294 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Either:

1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or

2 Both:

2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and

2.2 Either:

- 2.2.1 The person is unable to complete topical therapy; or
- 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Initial application — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis; or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Either:

1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or

2 Both:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) | Sub | Fully sidised | Brand or Generic |
|--|-----------------------------------|-------------|------------------|---------------------|
| | (Manulacturer 3 1 1106) \$ | Per | | Manufacturer |
| continued | | | | |
| 2.1 The person has a confirmed diagnosis of scable | s or is a close contact | of a scabie | es case; | and |
| 2.2 Either: | th a wanter want | | | |
| 2.2.1 The person is unable to complete topical 2.2.2 Previous treatment with topical therapy h | | leared the | infesta | tion |
| Renewal — (Other parasitic infections) from any relevant problem ollowing criteria: Any of the following: 1 1 filariasis; or 2 cutaneous larva migrans (creeping eruption); or 3 strongyloidiasis. 1 | | | | |
| PERMETHRIN | | | | |
| Lotn 5% | 4.28 3 | 0 ml OP | ✓ <u>F</u> | A-Scabies |
| Psoriasis and Eczema Preparations | | | | |
| ACITRETIN – Special Authority see SA2024 below – Retail ph | armacy | | | |
| Cap 10 mg | • | 60 | I N I | lovatretin |
| Novatretin to be Principal Supply on 1 July 2024 | | | | |
| Cap 25 mg | 57.37 | 60 | 🗸 N | lovatretin |

Novatretin to be Principal Supply on 1 July 2024

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

| Foam spray 500 mcg with calcipotriol 50 mcg per g Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g | 39.35 | 60 g OP 60 g OP 30 g OP | Enstilar <u>Daivobet</u> <u>Daivobet</u> |
|--|-------|-------------------------------|--|
| CALCIPOTRIOL Oint 50 mcg per g | 40.00 | 120 g OP | Daivonex |

| | Subsidy | | Fully Brand or |
|---|---|---|---|
| | (Manufacturer's Pi \$ | rice) Sub Per | sidised Generic Manufacturer |
| COAL TAR | Ŷ | | manaratian |
| Soln BP – Only in combination | 36.25 | 200 ml | ✓ Midwest |
| 1) Up to 10% only in combination with a dermatologic | | | |
| 2) With or without other dermatological galenicals. | a base of proprie | elary ropicary | |
| COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SUL | PHUR | | |
| Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% an | d | | |
| allantoin crm 2.5% | 6.59 | 75 g OP | |
| | (8.00) | | Egopsoryl TA |
| | 3.43 | 30 g OP | |
| | (4.35) | | Egopsoryl TA |
| COAL TAR WITH SALICYLIC ACID AND SULPHUR | | | |
| Soln 12% with salicylic acid 2% and sulphur 4% oint | 4.97 | 25 g OP | Coco-Scalp |
| | 7.95 | 40 g OP | Coco-Scalp |
| PIMECROLIMUS – Special Authority see SA1970 below – Reta | il pharmacy | | |
| Maximum of 15 g per prescription | | | |
| b) Note: a maximum of 15 g per prescription and no more t | | tion per 12 we | |
| Cream 1% | | 15 g OP | <u>Elidel</u> |
| SA1970 Special Authority for Subsidy | | | |
| nitial application only from a dermatologist, paediatrician, opht of a dermatologist, paediatrician or ophthalmologist. Approvals w | | | |
| neeting the following criteria: | | | |
| Both: | | | |
| Both: Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to | | | prificial dermatitis, rosacea, |
| Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. | pical corticosteroi | ds, cataracts, | prificial dermatitis, rosacea, glaucoma, or raised intraocular |
| Both: Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE | pical corticosteroi SCEIN – Only or | ids, cataracts, n a prescriptio | prificial dermatitis, rosacea, glaucoma, or raised intraoculai n |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ₭ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium | pical corticosteroi SCEIN – Only or | ds, cataracts, | prificial dermatitis, rosacea, glaucoma, or raised intraocula |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ₭ Soln 2.3% with trolamine laurilsulfate and fluorescein sodiun SALICYLIC ACID | pical corticosteroi SCEIN – Only or 15.41 | ids, cataracts, n a prescriptio 500 ml | prificial dermatitis, rosacea, glaucoma, or raised intraocular n Ý <u>Pinetarsol</u> |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodiun SALICYLIC ACID Powder – Only in combination | pical corticosteroi SCEIN – Only or 15.41 | ids, cataracts, n a prescriptio 500 ml 250 g | prificial dermatitis, rosacea, glaucoma, or raised intraocular n <u>✓ Pinetarsol</u> ✓ Midwest |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ₭ Soln 2.3% with trolamine laurilsulfate and fluorescein sodiun SALICYLIC ACID | pical corticosteroi SCEIN – Only or 15.41 | ids, cataracts, n a prescriptio 500 ml 250 g | prificial dermatitis, rosacea, glaucoma, or raised intraocula n ✓ <u>Pinetarsol</u> ✓ Midwest |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder - Only in combination | pical corticosteroi SCEIN – Only or 15.41 | ids, cataracts, n a prescriptio 500 ml 250 g | prificial dermatitis, rosacea, glaucoma, or raised intraocula n ✓ <u>Pinetarsol</u> ✓ Midwest |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder - Only in combination | pical corticosteroi SCEIN – Only or 1 | ids, cataracts, n a prescriptio 500 ml 250 g | prificial dermatitis, rosacea, glaucoma, or raised intraocula n ✓ <u>Pinetarsol</u> ✓ Midwest |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder – Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. | pical corticosteroi SCEIN – Only or 1 | ids, cataracts, n a prescriptio 500 ml 250 g al Corticoster 100 g | prificial dermatitis, rosacea, glaucoma, or raised intraocula n Pinetarsol Midwest oid – Plain or collodion flexible Midwest |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder - Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. SULPHUR Precipitated - Only in combination with a dermatological base or 1) Only in combination 1) Only in combination | pical corticosteroi SCEIN – Only or 1 | ids, cataracts, n a prescriptio 500 ml 250 g al Corticoster 100 g | prificial dermatitis, rosacea, glaucoma, or raised intraocula n Pinetarsol Midwest oid – Plain or collodion flexible Midwest |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder - Only in combination | pical corticosteroi SCEIN – Only or 1 | ids, cataracts, n a prescriptio 500 ml 250 g al Corticoster 100 g | prificial dermatitis, rosacea, glaucoma, or raised intraocula n Pinetarsol Midwest oid – Plain or collodion flexible Midwest |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder - Only in combination | pical corticosteroi SCEIN – Only or 1 | ids, cataracts, n a prescriptio 500 ml 250 g al Corticoster 100 g al Corticoster | prificial dermatitis, rosacea, glaucoma, or raised intraocula n <u>Y Pinetarsol</u> <u>Y Midwest</u> oid – Plain or collodion flexible Midwest oid – Plain |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder - Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. SULPHUR Precipitated - Only in combination with a dermatological base or 1) Only in combination adermatological base or 2) With or without other dermatological galenicals. | pical corticosteroi SCEIN – Only or 1 | ids, cataracts, n a prescriptio 500 ml 250 g al Corticoster 100 g | prificial dermatitis, rosacea, glaucoma, or raised intraocula n Pinetarsol Midwest oid – Plain or collodion flexible Midwest |
| Both: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE ★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium SALICYLIC ACID Powder - Only in combination | pical corticosteroi SCEIN – Only or 1 | ids, cataracts, n a prescriptio 500 ml 250 g al Corticoster 100 g al Corticoster 30 g OP | orificial dermatitis, rosacea, glaucoma, or raised intraocula n <u>Pinetarsol</u> <u>Midwest</u> oid – Plain or collodion flexible <u>Midwest</u> oid – Plain |

DERMATOLOGICALS

| (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer |
|--|
|--|

SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

| Scalp Preparations | | | |
|--|------------------|-------------------|---|
| BETAMETHASONE VALERATE | | | |
| * Scalp app 0.1% | 9.84 | 100 ml OP | <u>Beta Scalp</u> |
| CLOBETASOL PROPIONATE * Scalp app 0.05% | 6.26 | 30 ml OP | ✓ Dermol |
| HYDROCORTISONE BUTYRATE | | | |
| Scalp lotn 0.1% | 6.57 | 100 ml OP | ✓ Locoid |
| KETOCONAZOLE | 0.00 | | |
| Shampoo 2% | 3.23 4.09 | 100 ml OP | ✓ <u>Sebizole</u> ✓ Sebizole |
| a) Maximum of 100 ml per prescription | | | |
| b) Only on a prescription | | | |
| Sunscreens | | | |
| SUNSCREENS, PROPRIETARY – Subsidy by endorsement Only if prescribed for a patient with severe photosensitivity s endorsed accordingly. | econdary to a de | fined clinical co | ndition and the prescription is |
| Lotn, | 6.50 | 200 g OP | ✓ Marine Blue Lotion SPF 50+ |
| Wart Preparations | | | |
| For salicylic acid preparations refer to PSORIASIS AND ECZEM | A PREPARATIO | NS, page 74 | |
| PODOPHYLLOTOXIN | | | _ |
| Soln 0.5%a) Maximum of 3.5 ml per prescription | | 3.5 ml OP | Condyline |
| b) Only on a prescription | | | |
| Other Skin Preparations | | | |
| Antineoplastics | | | |
| FLUOROURACIL SODIUM | 0.05 | 00 × 00 | |
| | | 20 g OP | Efudix |
| IMIQUIMOD Crm 5%, 250 mg sachet | 21.72 | 24 | 🗸 Perrigo |

| Per ✓ Manufacturer |
|--------------------|
|--------------------|

| | | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|-------------|---|---|-------|---------------------|---------------------|
| Con | traceptives - Non-hormonal | | | | |
| Con | doms | | | | |
| COND | OMS | | | | |
| * 49 | mm - Up to 144 dev available on a PSO | 11.42 | 144 | ✓ | Moments |
| ₭ 53 | mm | 0.95 | 10 | ✓ | Moments |
| | | 11.64 | 144 | ✓ | Moments |
| | Maximum of 60 dev per prescription | | | | |
| | b) Up to 60 dev available on a PSO | | | | |
| ₭ 53 | mm, 0.05 mm thickness | 0.95 | 10 | | Moments |
| | | 11.42 | 144 | ✓ | Moments |
| | a) Up to 60 dev available on a PSO | | | | |
| | b) Maximum of 60 dev per prescription | | | | |
| € 53 | mm, chocolate, brown | | 10 | | Moments |
| | | 11.64 | 144 | 1 | Moments |
| | a) Up to 60 dev available on a PSO | | | | |
| | b) Maximum of 60 dev per prescription | | | | |
| € 53 | mm, strawberry, red | | 10 | | Moments |
| | | 11.64 | 144 | 1 | Moments |
| | a) Up to 60 dev available on a PSO | | | | |
| | b) Maximum of 60 dev per prescription | | | | |
| € 56 | mm | | 10 | | Moments |
| | | 11.64 | 144 | 1 | Moments |
| | a) Maximum of 60 dev per prescription | | | | |
| | b) Up to 60 dev available on a PSO | | | | |
| € 56 | mm, 0.05 mm thickness | | 12 | | Gold Knight |
| | | 24.10 | 144 | 1 | Gold Knight |
| | a) Up to 60 dev available on a PSO | | | | |
| | Maximum of 60 dev per prescription | | | | |
| € 56 | mm, 0.05mm thickness (bulk pack) | 20.17 | 144 | 1 | Gold Knight |
| | Maximum of 60 dev per prescription | | | | |
| | b) Up to 60 dev available on a PSO | | | | |
| € 56 | mm, 0.08 mm thickness | 0.97 | 10 | | Moments |
| | | 11.64 | 144 | ✓ | Moments |
| | a) Up to 60 dev available on a PSO | | | | |
| | b) Maximum of 60 dev per prescription | | | | |
| € 56 | mm, 0.08 mm thickness, red | | 10 | | Moments |
| | | 11.64 | 144 | ~ | Moments |
| | a) Up to 60 dev available on a PSO | | | | |
| | b) Maximum of 60 dev per prescription | | | - | |
| ₭ 56 | mm, chocolate | | 12 | - | Gold Knight |
| | | 21.45 | 144 | 1 | Gold Knight |
| | a) Up to 60 dev available on a PSO | | | | |
| | b) Maximum of 60 dev per prescription | | | - | |
| € 56 | mm, strawberry | | 12 | | Gold Knight |
| | | 21.45 | 144 | ~ | Gold Knight |
| | a) Up to 60 dev available on a PSO | | | | |
| | Maximum of 60 dev per prescription | | | - | . |
| ₭ 60 | mm | | 12 | | Gold Knight XL |
| | | 21.89 | 144 | ~ | Gold Knight XL |
| | a) Maximum of 60 dev per prescription | | | | |
| 70 | b) Uprim 69 destance lable on a PSO | S29 Unapproved | d med | icine supplie | ed under Section 29 |
| e 60 | mm_khillsparst.ppiy | Sole Sebsidised | slabb | ly 🗸 | Gold Knight XL |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|---|
| a) Maximum of 60 dev per prescriptionb) Up to 60 dev available on a PSO | | | | |
| Contraceptive Devices | | | | |
| INTRA-UTERINE DEVICE a) Up to 40 dev available on a PSO b) Only on a PSO ★ IUD 29.1 mm length × 23.2 mm width | 29.80 | 1 | ✓ (| MED NSHA Silver/ Copper Short Choice 380 7med Nsha Silver/ copper Short Choice TT380 Short |
| # IUD 33.6 mm length × 29.9 mm width | 29.80 | 1 | | Choice TT380 Standard |
| ₭ IUD 35.5 mm length × 19.6 mm width | | 1 | √ <u>(</u> | Choice Load 375 |

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

| * | Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up | to | | |
|---|--|-------|----|---------------------------------|
| | 84 tab available on a PSO | 10.00 | 84 | Mercilon 28 |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----------|---------------------|--|
| ETHINYLOESTRADIOL WITH LEVONORGESTREL | | | | |
| * Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets · | - | | | |
| Up to 84 tab available on a PSO | 1.50 | 84 | ✓ | Lo-Oralcon 20 ED |
| * Tab 30 mcg with levonorgestrel 150 mcg | 6.62 | 63 | | |
| | (16.50) | | | Microgynon 30 |
| a) Higher subsidy of \$15.00 per 63 tab with Special Auth b) Up to 63 tab available on a PSO * Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets Up to 84 tab available on a PSO ETHINYLOESTRADIOL WITH NORETHISTERONE | - | the 84 | | ige Oralcon 30 ED |
| Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to | | | | |
| 84 tab available on a PSO | | 84 112 | ✓ | Brevinor 1/28 Brevinor-1 28 Day Norimin-1 28 Day |
| Tab 35 mcg with norethisterone 500 mcg and 7 inert tab – U | р | | | • |
| to 84 tab available on a PSO | | 84 | 1 | Norimin |
| (Brevinor-1 28 Day Tab 35 mcg with norethisterone 1 mg and 7 ir (Norimin 1 28 Day Tab 35 mcg with parathistorone 1 mg and 7 in | | | | , |

(Norimin-1 28 Day Tab 35 mcg with norethisterone 1 mg and 7 inert tab to be delisted 1 December 2024)

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

- 1.1 Patient is on a Social Welfare benefit; or
- 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 Patient is on a Social Welfare benefit; or

2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

80

| * Tab 30 mcg – Up to 84 tab available on a PSO | 16.50 | 84 | Microlut |
|--|--------|-----|----------------------------------|
| | 22.00 | 112 | Microlut |
| ✤ Subdermal implant (2 × 75 mg rods) – Up to 3 pack available | | | |
| on a PSO | 106.92 | 1 | ✓ Jadelle |
| MEDROXYPROGESTERONE ACETATE | | | |
| Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO | 9.18 | 1 | Depo-Provera |
| | | | • |

| | Subsidy (Manufacturer's Price) \$ | Subsi Per | Fully dised | Brand or Generic Manufacturer |
|--|---|-------------------|----------------|-------------------------------------|
| NORETHISTERONE Tab 350 mcg – Up to 84 tab available on a PSO | | 84 | ✓ <u>N</u> | oriday 28 |
| Emergency Contraceptives | | | | |
| LEVONORGESTREL * Tab 1.5 mg a) Maximum of 2 tab per prescription b) Up to 5 tab available on a PSO c) Note: Direct Provision by a pharmacist permitted under | | 1 Part I of Se | _ | <u>evonorgestrel</u> <u>BNM</u> |
| Antiandrogen Oral Contraceptives | | | | |
| Prescribers may code prescriptions "contraceptive" (code "O") whe and prescription charge will be as per other contraceptives, as follo • A maximum \$5.00 prescription charge (patient co-payment) r • prescription may be written for up to six months supply | ows: | for contrac | ception | . The period of supply |

• prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to any non contraceptive prescription charges that apply, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

* Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs - Up

| to 168 tab available on a PSO | 5.08 168 | ✓ Ginet | |
|-------------------------------|----------|---------|--|
|-------------------------------|----------|---------|--|

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID

| Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.87) | 100 g OP | Aci-Jel |
|---|----------|------------------------------|
| CLOTRIMAZOLE | | |
| * Vaginal crm 1% with applicators | 35 g OP | Clomazol |
| * Vaginal crm 2% with applicators | 20 g OP | Clomazol |
| MICONAZOLE NITRATE | | |
| * Vaginal crm 2% with applicator | 40 g OP | Micreme |
| NYSTATIN | 0 | |
| Vaginal crm 100,000 u per 5 g with applicator(s) | 75 g OP | ✓ <u>Nilstat</u> |

Myometrial and Vaginal Hormone Preparations

ERGOMETRINE MALEATE

| Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO | 160.00 | 5 | ✓ DBL Ergometrine |
|---|--------|---------|-----------------------------|
| OESTRIOL | | | |
| * Crm 1 mg per g with applicator | 6.95 | 15 g OP | Ovestin |
| * Pessaries 500 mcg | 7.55 | 15 | Ovestin |

| | Subsidy | | Fully Brand or |
|--|---------------------------|--------------|--|
| | (Manufacturer's Pric | e) Sub | sidised Generic |
| | \$ | Per | Manufacturer |
| | Ŧ | | |
| OXYTOCIN – Up to 5 inj available on a PSO | | | |
| Inj 5 iu per ml, 1 ml ampoule | | 5 | Oxytocin BNM |
| Inj 10 iu per ml, 1 ml ampoule | 5.98 | 5 | Oxytocin BNM |
| | | | ✓ Oxytocin GH \$29 |
| | 11.00 | 10 | ✓ Oxytocin |
| | 11.96 | 10 | • |
| | | | Panpharma |
| (Oxytocin GH S29) Inj 10 iu per ml, 1 ml ampoule to be delis | ted 1 August 2024) | | |
| OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj | available on a PSO | | |
| Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml a | | 5 | Syntometrine |
| ing 5 id with ergometrine maleate 500 meg per mi, 1 mi a | inpoule | 5 | • Syntometrine |
| December 7. de 100 libre | | | |
| Pregnancy Tests - hCG Urine | | | |
| | | | |
| PREGNANCY TESTS - HCG URINE | | | |
| a) Up to 200 test available on a PSO | | | |
| b) Only on a PSO | | | |
| Cassette | | 40 test OP | Smith BioMed Rapid |
| | | | Pregnancy Test |
| | 10.00 | | • • |
| | 16.00 | | David One Step |
| | | | Cassette |
| | | | Pregnancy Test |
| | | | |
| Urinary Agents | | | |
| | | | |
| For urinary tract Infections refer to INFECTIONS, Antibacteria | als, page 113 | | |
| E Almha Daduatasa Inhihitara | | | |
| 5-Alpha Reductase Inhibitors | | | |
| FINACTERIDE Creation Authority and CA0000 holowy Bot | ail pharmany | | |
| FINASTERIDE – Special Authority see SA0928 below – Ret | | 400 | |
| * Tab 5 mg | 4.79 | 100 | ✓ <u>Ricit</u> |
| SA0928 Special Authority for Subsidy | | | |
| Initial application from any relevant practitioner. Approvals | valid without further re | newal unles | s notified for applications meeting |
| the following criteria: | | | |
| Both: | | | |
| | | | |
| Patient has symptomatic benign prostatic hyperplasia | ; and | | |
| 2 Either: | | | |
| 2.1 The patient is intolerant of non-selective alpha | blockers or these are o | ontraindicat | ted; or |
| 2.2 Symptoms are not adequately controlled with r | | | |
| | | | |
| Alpha-1A Adrenoreceptor Blockers | | | |
| Alpha-TA Autenoreceptor Diockers | | | |
| TAMSULOSIN HYDROCHLORIDE - Special Authority see | SA1022 bolow Dotail | nharmaou | |
| | | | Tomoulogin Day |
| * Cap 400 mcg | | 100 | Tamsulosin-Rex |
| SA1032 Special Authority for Subsidy | | | |
| Initial application from any relevant practitioner. Approvals | valid without further re- | newal unles | s notified for applications meeting |
| the following criteria: | | | and the second second second second |
| Both: | | | |
| | | | |
| Patient has symptomatic benign prostatic hyperplasia | | | |
| 2. The patient is intelerant of nen calenting alpha blocks | ra ar thaga are contrain | diagtad | |

2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

82

| | Subsidy (Manufacturer's Pr \$ | rice) Subsi Per | Fully dised | Brand or Generic Manufacturer |
|---|-------------------------------------|--------------------|----------------|--|
| Other Urinary Agents | | | | |
| OXYBUTYNIN * Tab 5 mg | 5.42 | 100 | | lchemy Oxybutynin |
| POTASSIUM CITRATE Oral lig 3 mmol per ml – Special Authority see SA1083 belov | v — | | | |
| Retail pharmacy | | 200 ml OP | ✔ В | iomed |
| SA1083 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid Both: | I for 12 months fo | or applications | meetin | g the following criteria: |
| The patient has recurrent calcium oxalate urolithiasis; and The patient has had more than two renal calculi in the two | vears prior to the | annlication | | |
| Renewal from any relevant practitioner. Approvals valid for 2 yea benefitting from the treatment. | | | appro | priate and the patient is |
| SODIUM CITRO-TARTRATE * Grans eff 4 g sachets | 3 50 | 28 | ✓ U | ral |
| SOLIFENACIN SUCCINATE | | | _ | <u> </u> |
| Tab 5 mg Tab 10 mg | | 30 30 | | olifenacin Viatris olifenacin Viatris |
| Detection of Substances in Urine | | | | |
| ORTHO-TOLIDINE | | | | |
| * Compound diagnostic sticks | 7.50 (8.25) | 50 test OP | H | emastix |
| TETRABROMOPHENOL * Blue diagnostic strips | | 100 test OP | 🗸 A | Ibustix |
| Obstetric Preparations | | | | |
| Antiprogesterones | | | | |
| MIFEPRISTONE Tab 200 mg – Up to 15 tab available on a PSO | 79.90 180.00 | 1 3 | | ifegyne ifegyne |

| | Subsidy Manufacturer's Price) | ę | Fully Subsidised | Brand or Generic |
|--|----------------------------------|---------|---------------------|------------------------------|
| | \$ | Per | 1 | Manufacturer |
| Calcium Homeostasis | | | | |
| CALCITONIN | | | | |
| * Inj 100 iu per ml, 1 ml ampoule | 121.00 | 5 | 🗸 M | liacalcic |
| CINACALCET - Special Authority see SA2170 below - Retail pha | irmacy | | | |
| Tab 30 mg – Wastage claimable | | 28 | √ <u>C</u> | inacalet Devatis |
| Tab 60 mg – Wastage claimable | | 28 | ✓ C | inacalet Devatis |
| ► SA2170 Special Authority for Subsidy | | | | |
| Initial application — (parathyroid carcinoma or calciphylaxis) | only from a nephrol | ogist c | or endocrino | ologist. Approvals valid for |
| 6 months for applications meeting the following criteria: | - | | | |

Either:

1 All of the following:

- 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
- 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
- 1.3 The patient is symptomatic; or

2 All of the following:

- 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
- 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
- 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.
- Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (primary hyperparathyroidism) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
- 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:

fully subsidised

Principal Supply

3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

| continued 3.2 Parathyroid tissue is surgically inaccessible; or 3.3 Parathyroid surgery is not feasible. Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months fo applications meeting the following citeria: Either: 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically accept parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate. ZOLEDRONIC ACID In 4 mg per 5 ml, vial | | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|---|----------|---------------------|-------------------------------------|
| 3.3 Parathyroid surgery is not feasible. Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months fo applications meeting the tollowing orteria: The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically accept parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate. ZOLEDRONIC ACID 1 ✓ Zoledronic acid Viatris In J am gp er 5 ml, vial 18.00 1 ✓ Zoledronic acid Viatris Corticosteroids and Related Agents for Systemic Use 5 Celestone BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE * 19.20 5 * Tab 0.5 mg – Up to 60 tab available on a PSO 2.65 30 ✓ Dexmethaone Oral liq 1 mg per ml. 19.20 5 Celestone Chronodose DEXAMETHASONE PHOSPHATE Examethasone phosphate injection will not be funded for oral use. * Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO | continued | | | | |
| applications meeting the following criteria: Ether: 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically accept parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate. ZOLEDRONIC ACID Inj 4 mg per 5 ml, vial | | | | | |
| parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate. ZOLEDRONIC ACID Inj 4 mg per 5 ml, vial EXAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE ** Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml | applications meeting the following criteria: | any relevant practiti | ioner. | Approvals | valid for 12 months for |
| Inj 4 mg per 5 ml, vial 1 ✓ Zoledronic acid Viatris Corticosteroids and Related Agents for Systemic Use DETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE * Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml 19.20 5 (36.96) Celestone Chronodose DEXAMETHASONE * 1.50 30 ✓ Dexmethsone Chronodose DEXAMETHASONE * 1.50 30 ✓ Dexmethsone Chronodose DEXAMETHASONE * 1.50 30 ✓ Dexmethsone Chronodose DEXAMETHASONE PHOSPHATE Dexamethasone phosphate injection will not be funded for oral use. * Biomed VEXAMETHASONE ACETATE * 10 ✓ HameIn PLOPCORTISONE ACETATE * 11.46 100 ✓ Horinef * Inj 4 mg per ml, 1 mampoule – Up to 5 inj available on a PSO | parathyroid hormone (PTH) level to support ongoing cessal | tion of treatment ha | s not b | een reache | d; or |
| Viatris Viatris Viatris Orticosteroids and Related Agents for Systemic Use BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE * Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml 19.20 5 (36.96) Celestone Chronodose DEXAMETHASONE * Tab 4 mg - Up to 60 tab available on a PSO 2.65 30 ✓ Dexmethsone Oral liq 1 mg per ml mg per ml 52.80 25 ml OP ✓ Biomed DEXAMETHASONE PHOSPHATE Dexamethasone phosphate injection will not be funded for oral use. * Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO 7.86 10 ✓ HameIn FLUDRCCORTISONE ACETATE * Tab 100 mg 20 mg 20.32 100 ✓ Douglas * Tab 5 mg 8.10 100 ✓ Elorinef HYDROCORTISONE * Tab 5 mg 8.10 100 ✓ Douglas * Tab 2 mg 20.32 100 ✓ <td>ZOLEDRONIC ACID</td> <td></td> <td></td> <td></td> <td></td> | ZOLEDRONIC ACID | | | | |
| BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE * Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml | Inj 4 mg per 5 ml, vial | | 1 | ✓ <u>Z</u> | |
| BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE ** Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml | Corticosteroids and Related Agents for Systemic | c Use | | | |
| * Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml 19.20 (36.96) 5 Celestone Chronodose Celestone Chronodose DEXAMETHASONE * Tab 4 mg - Up to 80 tab available on a PSO 1.50 2.65 30 30 ✓ Dexmethsone Dexmethsone * Tab 4 mg - Up to 30 tab available on a PSO 2.65 30 30 ✓ Dexmethsone Dexmethasone phosphate injection will not be funded for oral use. * Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO 7.86 10 10 ✓ Hamein Hamein * Tab 100 mcg. 11 46 100 100 ✓ Elorinef *YDROCORTISONE * Tab 20 mg 8.10 100 100 ✓ Douglas * Tab 20 mg 8.10 100 100 ✓ Douglas * Tab 20 mg 20.32 100 100 ✓ Douglas * Tab 20 mg 23.10 20 ✓ Medrol * Tab 20 mg 23.10 20 ✓ Medrol * Tab 20 mg 223.10 100 ✓ Douglas * Tab 100 mg vial 4.38 1 1 ✓ Solu-Cortef a) Not on a BSO b) Up to 5 inj available on a PSO 112.00 100 ✓ Medrol * Tab 100 mg CAS SODIUM SUCCINATE) Inj 40 mg vial 1 ✓ Solu-Medrol-Act- O-Vial In | | | | | |
| (36.96) Celestone Chronodose DEXAMETHASONE 30 ✓ Dexmethsone * Tab 0.5 mg - Up to 30 tab available on a PSO | | | 5 | | |
| DEXAMETHASONE 30 ✓ Dexmethsone ** Tab 0.5 mg - Up to 60 tab available on a PSO | | | U | C | elestone |
| ** Tab 0.5 mg - Up to 60 tab available on a PSO 1.50 30 ✓ Dexmethsone ** Tab 4 mg - Up to 30 tab available on a PSO 2.65 30 ✓ Dexmethsone Oral liq 1 mg per ml 52.80 25 ml OP ✓ Biomed DEXAMETHASONE PHOSPHATE 25 ml OP ✓ Biomed Dexamethasone phosphate injection will not be funded for oral use. * 10 ✓ HameIn * Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO 7.86 10 ✓ HameIn * Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO 13.10 10 ✓ HameIn FLUDROCORTISONE ACETATE 11.46 100 ✓ Elorinef HYDROCORTISONE 8.10 100 ✓ Douglas * Tab 2 0 mg 20.32 100 ✓ Douglas * Tab 2 0 mg 4.38 1 ✓ Solu-Cortef a) Not on a BSO 4.38 1 ✓ Solu-Cortef b) Up to 5 inj available on a PSO 223.10 100 ✓ Medrol WETHYLPREDNISOLONE 112.00 100 ✓ Medrol WETHYLPREDNISOLONE (AS SODIUM SUCCINATE) 11 1 ✓ Solu-Medrol-Act-O-Vial Inj 125 mg vial | | () | | | Chronodose |
| * Tab 4 mg - Up to 30 tab available on a PSO 2.65 30 ✓ Dexmethsone Oral liq 1 mg per ml | | 4 50 | | 1 | |
| Oral liq 1 mg per ml .52.80 25 ml OP ✓ Biomed DEXAMETHASONE PHOSPHATE Dexamethasone phosphate injection will not be funded for oral use. 10 ✓ HameIn Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO .7.86 10 ✓ HameIn * Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO .13.10 10 ✓ HameIn FLUDROCORTISONE ACETATE * 10 ✓ Florinef HYDROCORTISONE .11.46 100 ✓ Florinef HYDROCORTISONE .20.32 100 ✓ Douglas * Tab 5 mg .20.32 100 ✓ Douglas * Tab 20 mg .20.32 100 ✓ Douglas * Tab 20 mg .20.32 100 ✓ Douglas * Tab 4 ng .20.31 .20.22 100 ✓ Solu-Cortef * Tab 4 mg .112.00 100 ✓ Medrol ✓ Solu-Medrol-Act-O-Vial Inj 125 mg vial .22.30 1 ✓ Solu-Medrol-Act-O-Vial O-Vial Inj 500 mg vial .26.88 1 ✓ Solu-Medrol-Act-O-Vial | | | | | |
| DEXAMETHASONE PHOSPHATE Dexamethasone phosphate injection will not be funded for oral use. Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO7.86 10 ✓ HameIn Hinj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO13.10 10 ✓ HameIn StuDROCORTISONE ACETATE 10 ✓ Florinef YDROCORTISONE 11.46 100 ✓ Elorinef YDROCORTISONE 8.10 100 ✓ Douglas K Tab 5 mg 8.10 100 ✓ Douglas K Tab 20 mg 20.32 100 ✓ Douglas K Tab 20 mg vial 4.38 1 ✓ Solu-Cortef a) Not on a BSO b) Up to 5 inj available on a PSO ✓ Medrol ✓ Medrol K Tab 4 mg 112.00 100 ✓ Medrol ✓ Medrol K Tab 4 mg 223.10 20 ✓ Medrol ✓ Solu-Medrol-Act-O-Vial Inj 125 mg vial 24 34.10 1 ✓ Solu-Medrol-Act-O-Vial Inj 500 mg vial 26.88 1 ✓ Solu-Medrol-Act-O-Vial | | | | | |
| ** Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO7.86 10 - Hameln ** Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO13.10 10 - Hameln ** Tab 100 mcg | DEXAMETHASONE PHOSPHATE | | 5 111 01 | | loned |
| * Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO13.10 10 Hameln FLUDROCORTISONE ACETATE * Tab 100 mcg | | | 10 | 🗸 н | ameln |
| FLUDROCORTISONE ACETATE * Tab 100 mcg | | | | _ | |
| * Tab 100 mcg | | | | _ | |
| * Tab 5 mg 8.10 100 ✓ Douglas * Tab 20 mg 20.32 100 ✓ Douglas * Inj 100 mg vial 4.38 1 ✓ Solu-Cortef a) Not on a BSO b) Up to 5 inj available on a PSO 112.00 100 ✓ Medrol * Tab 4 mg 112.00 100 ✓ Medrol ✓ Medrol * Tab 100 mg 223.10 20 ✓ Medrol METHYLPREDNISOLONE 223.10 20 ✓ Medrol METHYLPREDNISOLONE (AS SODIUM SUCCINATE) 1 ✓ Solu-Medrol-Act-O-Vial Inj 40 mg vial 34.10 1 ✓ Solu-Medrol-Act-O-Vial Inj 500 mg vial 26.88 1 ✓ Solu-Medrol-Act-O-Vial | | 11.46 | 100 | ✓ <u>F</u> | lorinef |
| * Tab 20 mg | HYDROCORTISONE | | | | |
| k Inj 100 mg vial | * Tab 5 mg | 8.10 | 100 | 🗸 D | ouglas |
| a) Not on a BSO b) Up to 5 inj available on a PSO METHYLPREDNISOLONE * Tab 4 mg | | | | | |
| b) Up to 5 inj available on a PSO METHYLPREDNISOLONE * Tab 4 mg | | 4.38 | 1 | ✓ <u>s</u> | olu-Cortef |
| METHYLPREDNISOLONE * Tab 4 mg * Tab 4 mg * Tab 100 mg * Tab 100 mg 223.10 20 * Medrol METHYLPREDNISOLONE (AS SODIUM SUCCINATE) Inj 40 mg vial Inj 125 mg vial Inj 500 mg vial 20 21 223.10 20 * Solu-Medrol-Act-O-Vial 1 1 * Solu-Medrol-Act-O-Vial 1 1 * Solu-Medrol-Act-O-Vial 1 1 * Solu-Medrol-Act-O-Vial | | | | | |
| * Tab 4 mg 112.00 100 ✓ Medrol * Tab 100 mg 223.10 20 ✓ Medrol METHYLPREDNISOLONE (AS SODIUM SUCCINATE) 1 ✓ Solu-Medrol-Act-O-Vial Inj 40 mg vial | | | | | |
| * Tab 100 mg 20 ✓ Medrol METHYLPREDNISOLONE (AS SODIUM SUCCINATE) 1 ✓ Solu-Medrol-Act- O-Vial Inj 40 mg vial 34.10 1 ✓ Solu-Medrol-Act- O-Vial Inj 125 mg vial 34.10 1 ✓ Solu-Medrol-Act- O-Vial Inj 500 mg vial 26.88 1 ✓ Solu-Medrol-Act- O-Vial | | 110.00 | 400 | | • • |
| METHYLPREDNISOLONE (AS SODIUM SUCCINATE) 1 ✓ Solu-Medrol-Act-O-Vial Inj 40 mg vial 34.10 1 ✓ Solu-Medrol-Act-O-Vial Inj 500 mg vial 26.88 1 ✓ Solu-Medrol-Act-O-Vial | | | | | |
| Inj 40 mg vial 22.30 1 ✓ Solu-Medrol-Act-O-Vial Inj 125 mg vial 34.10 1 ✓ Solu-Medrol-Act-O-Vial Inj 500 mg vial 26.88 1 ✓ Solu-Medrol-Act-O-Vial | - | 223.10 | 20 | ♥ IV | leuror |
| O-Vial Inj 125 mg vial 1 ✓ Solu-Medrol-Act- O-Vial Inj 500 mg vial 26.88 1 ✓ Solu-Medrol-Act- O-Vial | | 22.20 | 1 | 10 | olu-Medrol-Act- |
| O-Vial Inj 500 mg vial | וון די ווע יומו | | 1 | * 3 | |
| O-Vial | Inj 125 mg vial | 34.10 | 1 | √ S | |
| | Inj 500 mg vial | | 1 | √ S | |
| inj 1 g viai | Inj 1 g vial | | 1 | √ S | olu-Medrol |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price \$ | e) Per | Full Subsidise | / |
|---|--|-----------|-------------------|--------------------|
| METHYLPREDNISOLONE ACETATE | | | | |
| Inj 40 mg per ml, 1 ml vial | 47.06 | 5 | ~ | Depo-Medrol |
| PREDNISOLONE | | | | |
| Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age. | 6.00 | 30 ml O | P 🗸 | Redipred |
| PREDNISONE | | | | |
| * Tab 1 mg | | 500 | ✓ | Prednisone Clinect |
| * Tab 2.5 mg | | 500 | ✓ | Prednisone Clinect |
| * Tab 5 mg – Up to 30 tab available on a PSO | | 500 | | Prednisone Clinect |
| * Tab 20 mg – Up to 30 tab available on a PSO | 50.51 | 500 | ~ | Prednisone Clinect |
| TETRACOSACTRIN | | | | |
| * Inj 250 mcg per ml, 1 ml ampoule | | 1 | 1 | Synacthen |
| Jee She year | | | | UK Synacthen |
| * Inj 1 mg per ml, 1 ml ampoule | | 1 | | Synacthen Depot |
| | | | - | Synacthene |
| | | | | Retard S29 |
| FRIAMCINOLONE ACETONIDE | | | | |
| Ini 10 mg per ml, 1 ml ampoule | 21 42 | 5 | | Kenacort-A 10 |
| Inj 40 mg per ml, 1 ml ampoule | | 5 | | Kenacort-A 40 |
| Sex Hormones Non Contraceptive Androgen Agonists and Antagonists | | | | |
| | | | | |
| CYPROTERONE ACETATE | | | | |
| Tab 50 mg | | 50 | | Siterone |
| Tab 100 mg | | 50 | ~ | Siterone |
| ESTOSTERONE | | | | |
| Gel (transdermal) 16.2 mg per g | 52.00 | 88 g Ol | > √ | Testogel |
| Testogel to be Principal Supply on 1 July 2024 | | | | |
| Patch 5 mg per day | | 30 | ~ | Androderm |
| ESTOSTERONE CIPIONATE | | | | |
| Inj 100 mg per ml, 10 ml vial | | 1 | ~ | Depo-Testosterone |
| | 393.00 | | ✓ | Taro- |
| | | | | Testosterone S29 |
| ESTOSTERONE ESTERS | | | | |
| Inj 250 mg per ml, 1 ml | 10 08 | 1 | | Sustanon Ampoules |
| | 12.30 | I | • | Sustanon Ampoules |
| ESTOSTERONE UNDECANOATE | | | | |
| Cap 40 mg – Subsidy by endorsement | | 100 | | Steril-Gene S29 |
| Subsidy by endorsement – subsidised for patients who | | | | |
| 1 November 2021 and the prescription is endorsed ac | | | | |
| where there exists a record of prior dispensing of testo | | • | | |
| Inj 250 mg per ml, 4 ml vial | 86.00 | 1 | v | Reandron 1000 |

86

| | Subsidy | Fu | ully Brand or |
|----------------------------------|------------------------|----------|---|
| | (Manufacturer's Price) | Subsidis | ed Generic |
| | \$ | Per | Manufacturer |
| Hormone Replacement Therapy - | Systemic | | |
| Oestrogens | · | | |
| | | | |
| OESTRADIOL | | | |
| * Tab 1 mg | | 28 OP | - |
| W. Tab O and | (11.10) | ~~~~ | Estrofem |
| * Tab 2 mg | | 28 OP | F atura fa un |
| Datah 50 mag par 04 haura | (11.10) | 4 | Estrofem ✓ Climara |
| Patch 50 mcg per 24 hours | | 4 | |
| a) No more than 1 patch per week | | | |
| b) Only on a prescription | 0.05 | 0 | |
| Patch 25 mcg per day | | | Estradiol TDP Mylan |
| | 13.50 | | Estraderm MX S29 |
| | 14.50 | | Estradot |
| a) No more than 2 patch per week | | | |
| b) Only on a prescription | | | |
| Patch 50 mcg per day | | | Estradiol TDP Mylan |
| | | | Estradiol Viatris |
| | 14.50 | | Estraderm MX S29 |
| | | | Estradiol |
| | | | Sandoz S29 |
| | | | Estradot |
| a) No more than 2 patch per week | | | |
| b) Only on a prescription | | | |
| Patch 75 mcg per day | | 8 | Estradiol TDP Mylan |
| | | , | Estradiol Viatris |
| | 14.50 | | Estradiol |
| | | | Sandoz S29 |
| | | | Estradot |
| a) No more than 2 patch per week | | | |
| b) Only on a prescription | | | |
| Patch 100 mcg per day | | 8 | Estradiol TDP Mylan |
| · | | | Estradiol Viatris |
| | 14.50 | | Estradiol |
| | | | Sandoz S29 |
| | | | ✓ Estradot |
| | 15.50 | | Estraderm MX \$29 |
| a) No more than 2 patch per week | 10.00 | | Estruction mix de |
| b) Only on a prescription | | | |
| · · · · | | | |
| OESTRADIOL VALERATE | | | <i>.</i> - |
| Tab 1 mg | | | Progynova |
| * Tab 2 mg | | 84 | Progynova |
| OESTROGENS | | | |
| * Conjugated, equine tab 300 mcg | | 28 | |
| | (17.50) | | Premarin |
| * Conjugated, equine tab 625 mcg | 4.12 | 28 | |
| - | (17.50) | | Premarin |
| | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Pubaidu | | Eully | Brand or |
|---|----------------------------------|-----------|-------------------|-----------------------------|
| | Subsidy (Manufacturer's Price |) Sut | Fully osidised | Brand or Generic |
| | \$ | Per | ~ | Manufacturer |
| Progestogens | | | | |
| riogestogens | | | | |
| MEDROXYPROGESTERONE ACETATE | | | | |
| * Tab 2.5 mg | | 30 | | rovera |
| W. Tab Casa | 8.75 | 56 | - | rovera |
| * Tab 5 mg | 9.80 17.50 | 56 100 | | Provera Provera |
| * Tab 10 mg | | 100 30 | - | Provera |
| Progestogen and Oestrogen Combined Prep | | | | |
| r rogestogen and bestrogen bombined r rep | | | | |
| OESTRADIOL WITH NORETHISTERONE | | | | |
| * Tab 1 mg with 0.5 mg norethisterone acetate | · · - · - · | 28 OP | | |
| Mr. Tab O man the dama and the transmission of the | (18.10) | 00.05 | K | lliovance |
| * Tab 2 mg with 1 mg norethisterone acetate | | 28 OP | | 11: |
| | (18.10) | | K | liogest |
| * Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg with 1 mg norethisterone acetate (10). | | 00 O D | | |
| oestradiol tab (12) and 1 mg oestradiol tab (6) | · · - · - · | 28 OP | т | - |
| | (18.10) | | 1 | risequens |
| Other Oestrogen Preparations | | | | |
| Chief Coolicgen Proparations | | | | |
| OESTRIOL | | | | |
| * Tab 2 mg | 7.70 | 30 | ✓ <u>c</u> | Vestin |
| Other Dregesteren Drenerstiene | | | | |
| Other Progestogen Preparations | | | | |
| LEVONORGESTREL | | | | |
| * Intra-uterine device 52 mg | | 1 | 🗸 N | lirena |
| * Intra-uterine device 13.5 mg | | 1 | 🗸 🗸 | aydess |
| MEDROXYPROGESTERONE ACETATE | | | | |
| Tab 100 mg | 116.15 | 100 | 🗸 P | Provera HD |
| NORETHISTERONE | | | | |
| * Tab 5 mg – Up to 30 tab available on a PSO | 5.49 | 30 | 🗸 P | Primolut N |
| PROGESTERONE | | | | |
| * Cap 100 mg | | 30 | 🗸 U | Itrogestan |
| | | | _ | 3 |
| Thyroid and Antithyroid Agents | | | | |
| CARBIMAZOLE | | | | |
| * Tab 5 mg | 7 56 | 100 | 🖌 N | leo-Mercazole |
| - | | 100 | • 1 | |
| LEVOTHYROXINE | E EE | 00 | | unthroid |
| * Tab 25 mcg * Tab 50 mcg | | 90 28 | | Synthroid Jercury Pharma |
| * 1 ab 50 mcg | 5.79 | 28 90 | | lercury Pharma |
| | 12.86 | 200 | | ltroxin |
| | 64.28 | 1.000 | | Itroxin |
| * Tab 100 mcg | | 28 | | lercury Pharma |
| Ŭ | 6.01 | 90 | | synthroid |
| | 13.36 | 200 | | ltroxin |
| | 66.78 | 1,000 | ✓ E | Itroxin |
| | | | | |

| | Subsidy (Manufacturer's Price) \$ | S Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|----------|---------------------|-------------------------------------|
| PROPYLTHIOURACIL – Special Authority see SA1199 below – I | Retail pharmacy | | | |
| Tab 50 mg | | 100 | 🗸 P | TU \$29 |

➡SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

| SOMATROPIN (OMNITROPE) - Special Authority see | SA2032 below – Retail pharmacy | |
|--|--------------------------------|---------------------------------------|
| * Inj 5 mg cartridge | | Omnitrope |
| | | Omnitrope S29 S29 |
| * Inj 10 mg cartridge | | Omnitrope |
| | | Omnitrope S29 S29 |
| * Inj 15 mg cartridge | | Omnitrope |
| | | ✓ Omnitrope S29 S29 |

► SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under ; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

90

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|----------|--------------|--|
| (Manufacturer's Price) | Su | bsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m²) in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| Subsidy | | Fully | Brand or | |
|------------------------|------------|-------|--------------|--|
| (Manufacturer's Price) | Subsidised | | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and

| | Subsidy | F | ully | Brand or |
|----|-----------------------|--------|------|--------------|
| (1 | Manufacturer's Price) | Subsid | sed | Generic |
| | \$ | Per | 1 | Manufacturer |

continued...

3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GOSERELIN

| GOSERELIN | | | |
|--|---------------|---------------|-----------------------------------|
| Implant 3.6 mg, syringe | | 1 | ✓ <u>Zoladex</u> |
| Implant 10.8 mg, syringe | 138.23 | 1 | Zoladex |
| LEUPRORELIN | | | |
| Additional subsidy by endorsement where the patient is a child | or adolescent | and is unable | to tolerate administration of |
| goserelin and the prescription is endorsed accordingly. | | | |
| Inj 3.75 mg prefilled dual chamber syringe – Higher subsidy of | | | |
| \$221.60 per 1 inj with Endorsement | | 1 | |
| • •• •• • • • • • • • | (221.60) | | Lucrin Depot 1-month |
| Inj 11.25 mg prefilled dual chamber syringe – Higher subsidy | (, | | |
| of \$591.68 per 1 inj with Endorsement | 177.50 | 1 | |
| · · · · · · · · · · · · · · · · · · · | (591.68) | | Lucrin Depot 3-month |
| | (, | | , |
| Vasopressin Agonists | | | |
| | | | |
| DESMOPRESSIN | | | _ |
| Wafer 120 mcg | 47.00 | 30 | Minirin Melt |
| DESMOPRESSIN ACETATE | | | |
| Tab 100 mcg | 25.00 | 30 | Minirin |
| Tab 200 mcg | 54.45 | 30 | 🗸 Minirin |
| ▲ Nasal spray 10 mcg per dose | 34.95 | 6 ml OP | Desmopressin- |
| | | | PH&T |
| | | | 4 • • • • • |
| Inj 4 mcg per ml, 1 ml | 67.18 | 10 | Minirin |
| Other Friday in America | | | |
| Other Endocrine Agents | | | |
| CABERGOLINE | | | |
| Tab 0.5 mg – Maximum of 2 tab per prescription; can be | | | |
| waived by Special Authority see SA2070 on the next page | 4 43 | 2 | ✓ Dostinex |
| warrow by openial nationaly see OA2070 on the next page | | 8 | ✓ Dostinex |
| | 17.54 | 0 | Dodinex |

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Sul Per | Fully osidised | Brand or Generic Manufacturer |
|---|---|-----------------------------|---------------------|-------------------------------------|
| SA2070 Special Authority for Waiver of Rule Initial application from any relevant practitioner. Approvals value the following: Hyperprolactinemia; or Acromegaly*; or Inhibition of lactation. Renewal — (for patients who have previously been funded up practitioner. Approvals valid without further renewal unless notifie which has expired and the treatment remains appropriate and the Note: Indication marked with * is an unapproved indication. | nder Special Author | ity form has prev | SA1031 iously he |) from any relevant |
| CLOMIFENE CITRATE Tab 50 mg | 29.84 | 10 | ✓ N | lylan Clomiphen S29 |

| Cap 250 mg | 558.00 | 50 | Metopirone |
|------------|--------|----|--------------------------------|
|------------|--------|----|--------------------------------|

94

METYRAPONE

| | Subsidy | | Fully | Brand or |
|--|-------------------------------|-----------------|-----------|-------------------------|
| | (Manufacturer's Price) | | | Generic |
| | (Manalactarer 3 1 1100) \$ | Per | 1000 1 | Manufacturer |
| | φ | Fei | | Wanulaciulei |
| | | | | |
| Anthelmintics | | | | |
| ALBENDAZOLE - Special Authority see SA1318 below - Retain | il pharmacy | | | |
| Tab 400 mg | | 60 | ✓ E | skazole S29 |
| ► SA1318 Special Authority for Subsidy | | | | |
| Initial application only from an infectious disease specialist or the second | olinical microbiologist | Approvale | valid f | for 6 months whore the |
| | cilinical microbiologist. | Approvais | vallu i | |
| patient has hydatids. | | | | |
| Renewal only from an infectious disease specialist or clinical mi | icrobiologist. Approva | als valid for 6 | 6 mont | ths where the treatment |
| remains appropriate and the patient is benefitting from the treat | ment. | | | |
| | | | | |
| MEBENDAZOLE – Only on a prescription | | | | |
| Tab 100 mg | 7.97 | 6 | ✓ V | /ermox |
| Oral lig 100 mg per 5 ml | 2.18 | 15 ml | | |
| | (7.83) | | V | /ermox |
| | (1.00) | | • | onnox |
| PRAZIQUANTEL | | | | |
| Tab 600 mg | | 8 | ✓ E | Biltricide |
| | | - | | |
| Antibacterials | | | | |
| Antibacteriais | | | | |
| a) For topical antibacterials, refer to DERMATOLOGICALS, pa | ao 68 | | | |
| | | | | |
| b) For anti-infective eye preparations, refer to SENSORY ORG | ANS, page 260 | | | |
| Conheleenevine and Conhemusine | | | | |
| Cephalosporins and Cephamycins | | | | |
| CEFACLOR MONOHYDRATE | | | | |
| | | | | |
| Cap 250 mg | | 100 | | anbaxy-Cefaclor |
| Grans for oral lig 125 mg per 5 ml – Wastage claimable | 3.75 | 100 ml | 🗸 F | anbaxy-Cefaclor |
| CEFALEXIN | | | | - |
| | 0.05 | | | |
| Cap 250 mg | | 20 | | ephalexin ABM |
| Cap 500 mg | 5.85 | 20 | ✓ 0 | ephalexin ABM |
| Grans for oral lig 25 mg per ml - Wastage claimable | | 100 ml | ✓ F | lynn |
| Grans for oral lig 50 mg per ml – Wastage claimable | | 100 ml | | lynn |
| | | 100 111 | | Cefalexin Sandoz |
| | 11.75 | | • (| eralexin Sandoz |
| CEFAZOLIN – Subsidy by endorsement | | | | |
| Only if prescribed for dialysis or cellulitis in accordance with | a Health NZ Hospital | annroved n | rotocc | and the prescription is |
| | | approved p | 101000 | |
| endorsed accordingly. | | _ | | |
| Inj 500 mg vial | | 5 | | Cefazolin-AFT |
| Inj 1 g vial | 3.59 | 5 | ✓ 0 | efazolin-AFT |
| Inj 2 g vial | 7.09 | 5 | √ 0 | efazolin-AFT |
| | | | - | |
| CEFTRIAXONE – Subsidy by endorsement | | | | |
| a) Up to 10 inj available on a PSO | | | | |
| b) Subsidised only if prescribed for a dialysis or cystic fibro | sis patient, or the trea | tment of gor | orrho | ea. or the treatment of |
| pelvic inflammatory disease, or the treatment of suspect | | | | |
| | eu meningococcar uis | | | |
| endorsed accordingly. | | | | |
| Inj 500 mg vial | 0.79 | 1 | _ | Ceftriaxone-AFT |
| Inj 1 g vial | 3.59 | 5 | √ 0 | Ceftriaxone-AFT |
| , . | | | _ | |
| CEFUROXIME AXETIL – Subsidy by endorsement | | | | |
| Only if prescribed for prophylaxis of endocarditis and the pr | | | | |
| Tab 250 mg | CBS | 20 | 🗸 A | scend- |
| | | | | Cefuroxime S29 |
| | | | | |
| | | | | |

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

 $\ensuremath{\textbf{\#}}$ Three months or six months, as applicable, dispensed all-at-once

| Subsidy (Manufacturer's Price) \$ | Subs Per | Fully sidised | Brand or Generic Manufacturer |
|---|--|---|---|
| | | | |
| ystic fibrosis bronchi | • | ll be sul | |
| | 2 | | ithromax |
| | 15 ml | ✓ Z | ithromax |
| | (Manufacturer's Price) \$ n; can be waived by ystic fibrosis bronchi 8.19 | (Manufacturer's Price) Subs \$ Per h; can be waived by Special Au ystic fibrosis bronchiectasis wi | (Manufacturer's Price) \$ Per ✓ h; can be waived by Special Authority ystic fibrosis bronchiectasis will be sul 8.19 30 ✓ A |

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.
- Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).
- The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

| Tab 250 mg | 8.53 | 14 | Klacid |
|--|------|-------|--------|
| Grans for oral lig 250 mg per 5 ml - Wastage claimable | | 50 ml | Klacid |

► SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Either:

| Subsidy | | Fully | Brand or | |
|-----------------------|------|------------|--------------|--|
| (Manufacturer's Price | e) - | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

1 Atypical mycobacterial infection; or

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents. Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and

2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial | 10.00 | 1 | Erythrocin IV |
|---|-------|--------|-----------------------------------|
| ERYTHROMYCIN ETHYL SUCCINATE | | | |
| Tab 400 mg | | 100 | 🗸 E-Mycin |
| a) Up to 20 tab available on a PSO | | | |
| b) Up to 2 x the maximum PSO quantity for RFPP | | | |
| Grans for oral liq 200 mg per 5 ml | 5.00 | 100 ml | E-Mycin |
| a) Up to 300 ml available on a PSO | | | |
| b) Up to 2 x the maximum PSO quantity for RFPP | | | |
| c) Wastage claimable | | | |
| Grans for oral liq 400 mg per 5 ml | 6.77 | 100 ml | E-Mycin |
| a) Up to 200 ml available on a PSO | | | |
| b) Wastage claimable | | | |
| ROXITHROMYCIN | | | |
| Tab 150 mg | | 50 | Arrow- |
| ő | | | Roxithromycin |
| | | | |
| Tab 300 mg | 25.00 | 50 | ✓ Arrow- |
| | | | Roxithromycin |

| | Subsidy (Manufacturer's F \$ | Price) Subs Per | Fully Brand or idised Generic Manufacturer |
|---|------------------------------------|--------------------|--|
| Penicillins | Ŷ | | |
| AMOXICILLIN | | | |
| Cap 250 mg | 27.50 | 500 | Miro-Amoxicillin |
| | 43.45 | | Alphamox |
| a) Up to 30 cap available on a PSO | | | |
| b) Up to 10 x the maximum PSO quantity for RFPP | | | |
| c) Miro-Amoxicillin to be Principal Supply on 1 September | | | · · · · · · · · · · · · · · · · · · · |
| Cap 500 mg | | 500 | Miro-Amoxicillin |
| | 66.44 | | Alphamox |
| a) Up to 30 cap available on a PSO | | | |
| b) Up to 10 x the maximum PSO quantity for RFPP a) Mine Amazini III in to be Dringing Lower to Amazini III in the Amazini III | 0.4 | | |
| c) Miro-Amoxicillin to be Principal Supply on 1 August 20 | | 100 ml | Alphamox 125 |
| Grans for oral liq 125 mg per 5 ml. | | 100 111 | Alphamox 125 |
| a) Up to 200 ml available on a PSOb) Wastage claimable | | | |
| Grans for oral lig 250 mg per 5 ml | 2.81 | 100 ml | Alphamox 250 |
| a) Up to 300 ml available on a PSO | 2.01 | 100 111 | Alphaniox 250 |
| b) Up to 10 x the maximum PSO guantity for RFPP | | | |
| c) Wastage claimable | | | |
| Inj 250 mg vial | 15.97 | 10 | Ibiamox |
| Inj 500 mg vial | | 10 | ✓ Ibiamox |
| Inj 1 g vial – Up to 5 inj available on a PSO | | 10 | ✓ Ibiamox |
| Alphamox Cap 250 mg to be delisted 1 September 2024) | | | |
| Alphamox Cap 500 mg to be delisted 1 August 2024) | | | |
| AMOXICILLIN WITH CLAVULANIC ACID | | | |
| Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab | | | |
| available on a PSO | 1.59 | 10 | Curam Duo 500/125 |
| Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 n | | | <u></u> |
| per ml | | 100 ml | Augmentin |
| a) Up to 200 ml available on a PSO | | | Ū |
| b) Wastage claimable | | | |
| Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 n | ng | | |
| per ml – Up to 200 ml available on a PSO | | 100 ml OP | Curam |
| | | | |
| Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj | | | |
| available on a PSO | | 10 | Bicillin LA |
| BENZYLPENICILLIN SODIUM [PENICILLIN G] | | | |
| Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS | SO 16.50 | 10 | ✓ Sandoz |
| | | 10 | Candoz |

98

| | Subsidy (Manufacturer's Price | | Fully sidised | |
|--|----------------------------------|-------------|------------------|----------------------------|
| | \$ | Per | - | Manufacturer |
| FLUCLOXACILLIN | | | | |
| Cap 250 mg – Up to 30 cap available on a PSO | | 250 | | Flucloxacillin-AFT |
| Cap 500 mg – Up to 30 cap available on a PSO | | 500 | | Flucloxacillin-AFT |
| Grans for oral liq 25 mg per ml | 3.29 | 100 ml | 1 | AFT |
| a) Up to 200 ml available on a PSOb) Wastage claimable | | | | |
| Grans for oral liq 50 mg per ml | 3.68 | 100 ml | 1 | AFT |
| a) Up to 200 ml available on a PSO b) Wastage claimable | | | | |
| Inj 250 mg vial | | 10 | 1 | Flucloxin |
| Flucloxin to be Principal Supply on 1 July 2024 | | | | |
| Inj 500 mg vial | | 10 | 1 | Flucloxin |
| Flucloxin to be Principal Supply on 1 July 2024 | | | | |
| Inj 1 g vial – Up to 5 inj available on a PSO | 6.00 | 5 | - | Flucil |
| PHENOXYMETHYLPENICILLIN (PENICILLIN V) | | | | |
| Cap 250 mg – Up to 30 cap available on a PSO | 3 84 | 50 | 1 | Cilicaine VK |
| Cap 500 mg | | 50 | | Cilicaine VK |
| a) Up to 20 cap available on a PSO | | 00 | • | <u>omounic vit</u> |
| b) Up to 2 x the maximum PSO quantity for RFPP | | | | |
| Grans for oral liq 125 mg per 5 ml | 3.40 | 100 ml | 1 | AFT |
| a) Up to 200 ml available on a PSO | 0.+0 | 100 111 | • | |
| b) Wastage claimable | | | | |
| Grans for oral liq 250 mg per 5 ml | 4 24 | 100 ml | 1 | AFT |
| a) Up to 300 ml available on a PSO | | 100 111 | • | |
| b) Up to 2 x the maximum PSO quantity for RFPP | | | | |
| c) Wastage claimable | | | | |
| Tetracyclines | | | | |
| • | | | | |
| DOXYCYCLINE X Tab 100 mg Line to 20 tab available on a DSO | 64.40 | 500 | | Doxine |
| * Tab 100 mg – Up to 30 tab available on a PSO | | 500 | • | Doxine |
| MINOCYCLINE HYDROCHLORIDE | | | | |
| * Tab 50 mg – Additional subsidy by Special Authority see | | | | |
| SA1355 below – Retail pharmacy | 5.79 | 60 | | |
| | (12.05) | | | Mino-tabs |
| * Cap 100 mg | 19.32 | 100 | | |
| | (52.04) | | | Minomycin |
| ➡SA1355 Special Authority for Manufacturers Price | | | | |
| Initial application from any relevant practitioner. Approvals va | lid without further rer | newal unles | s notif | ied where the patient has |
| rosacea. | | | | |
| TETRACYCLINE - Special Authority see SA1332 below - Reta | ail pharmacy | | | |
| Tab 250 mg | | 28 | 1 | Accord S29 |
| SA1332 Special Authority for Subsidy | | - | | |
| Initial application from any relevant practitioner. Approvals va | lid for 2 months for a | nnligations | mooti | an the following criterie: |
| Both: | ing for 3 months for a | pplications | meetii | ng the following chiefla: |
| 1 For the eradication of helicobacter pylori following unsuc | cossful treatment wit | h annronria | to fire | t line thereby: and |

1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and

2 For use only in combination with bismuth as part of a quadruple therapy regimen.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|--------------|---------------------|---|
| Other Antibiotics | | | | |
| For topical antibiotics, refer to DERMATOLOGICALS, page 68 | | | | |
| CIPROFLOXACIN Recommended for patients with any of the following: i) microbiologically confirmed and clinically significant pse ii) prostatitis; or iii) pyelonephritis; or iv) gonorrhoea. | eudomonas infection; | or | | |
| Tab 250 mg – Up to 5 tab available on a PSO | 2.42 | 28 | 1 | Cipflox |
| | 3.85 | 10 | 1 | Ciprofloxacin - Torrent ©29 |
| Tab 500 mg – Up to 5 tab available on a PSO | 4.25 | 10 | 1 | Ciprofloxacin - Torrent S29 |
| Tab 750 mg | 5.95 | 28 | ~ | Cipflox |
| CLINDAMYCIN | | | | |
| Cap hydrochloride 150 mg Inj 150 mg per ml, 4 ml ampoule | | 24 10 | | Dalacin C Hameln |
| COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – S Only if prescribed for dialysis or cystic fibrosis patient and the Inj 150 mg | e prescription is endo | | | ∕. Colistin-Link |
| GENTAMICIN SULPHATE Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly. | | 5 / trac | | DBL Gentamicin and the prescription is |
| Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement | 91.00 | 5 | ~ | Wockhardt S29 |
| | 182.00 | 10 | | Teligent \$29 |
| Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly. | or complicated urinary | / trac | t infection a | and the prescription is |
| Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement | | 10 | | Pfizer |
| Only if prescribed for a dialysis or cystic fibrosis patient | 87.50 or complicated urinan | 50 / trac | | Pfizer and the prescription is |
| endorsed accordingly. | or complicated unitary | | | |
| MOXIFLOXACIN – Special Authority see SA1740 below – Retai No patient co-payment payable | l pharmacy | | | |
| Tab 400 mg | | 5 | ~ | Avelox |
| SA1740 Special Authority for Subsidy Initial application — (Tuberculosis) only from a respiratory sp for applications meeting the following criteria: Any of the following: Both: | ecialist or infectious d | iseas | e specialis | t. Approvals valid for 1 year |
| Active tuberculosis*; and Any of the following: | | | | |
| 1.2.1 Documented resistance to one or more firs 1.2.2 Suspected resistance to one or more first-li | , | rculos | sis assume | d to be contracted in an |

| Subsidy | | Fully | Brand or |
|-----------------------|------|--------|--------------|
| Manufacturer's Price) | Subs | idised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

area with known resistance), as part of regimen containing other second-line agents; or

- 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
- 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
- 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and

2 Either:

- 2.1 Has tried and failed to clear infection using azithromycin; or
- 2.2 Has laboratory confirmed azithromycin resistance; and

3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only. Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

➡SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria: Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

30

✓ Daraprim S29

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

| SODIUM FUSIDATE [FUSI | 36 | Fucidin |
|-----------------------|--------|-----------------------------|
| Ũ | | |
| Tab 500 mg | 56 | ✓ Wockhardt S29 |

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price \$ |) Sub: Per | Fully sidised | Brand or Generic Manufacturer |
|---|--|-----------------|-------------------|-------------------------------------|
| ■ SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valic the following criteria: Any of the following: | l without further ren | ewal unles | s notified | d for applications meeting |
| For the treatment of toxoplasmosis in patients with HIV for For pregnant patients for the term of the pregnancy; or For infants with congenital toxoplasmosis until 12 months | | ns; or | | |
| TOBRAMYCIN Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement Only if prescribed for dialysis or cystic fibrosis patient and | | 5 endorsed a | | obramycin (Viatris) gly. |
| Solution for inhalation 60 mg per ml, 5 ml – Subsidy by endorsementa) Wastage claimable | | 56 dose | ✓ <u>T</u> | obramycin BNM |
| b) Only if prescribed for a cystic fibrosis patient and the TRIMETHOPRIM | prescription is endo | rsed accor | dingly. | |
| * Tab 300 mg – Up to 30 tab available on a PSO | | 50 | ✓ <u>⊺</u> | MP |
| TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX/ * Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – L to 30 tab available on a PSO | lp 64.80 nl | 500 100 ml | ✓ <u>⊺</u> | risul eprim |
| VANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or for difficile following metronidazole failure and the prescription is Inj 500 mg vial | endorsed according | | for treat ✓ M | |
| Antifungals | | | | |
| a) For topical antifungals refer to DERMATOLOGICALS, page 65 b) For topical antifungals refer to GENITO URINARY, page 81 FLUCONAZOLE |) | | | |
| Cap 50 mg | | 28 | ✓ <u>M</u> | |
| Cap 150 mg Cap 200 mg | | 1 28 | ✓ <u>M</u> ✓ M | |
| Powder for oral suspension 10 mg per ml – Special Authority see SA1359 below – Retail pharmacy Wastage claimable | | 35 ml | | iflucan |
| SA1359 Special Authority for Subsidy Initial application — (Systemic candidiasis) from any relevant meeting the following criteria: Both: | practitioner. Appro | ovals valid f | or 6 wee | eks for applications |
| Patient requires prophylaxis for, or treatment of systemic of Patient is unable to swallow capsules. | andidiasis; and | | | |
| Initial application | t practitioner. Appr | ovals valid | for 6 mc | onths for applications |

meeting the following criteria: All of the following:

| | Subsidy | | Fully | Brand or |
|---|----------------------------------|----------------|------------|--|
| | (Manufacturer's Price \$ | e) Sub Per | osidised | Generic Manufacturer |
| continued | | | | |
| Patient is immunocompromised; and Patient is at moderate to high risk of invasive fungal infecti Patient is unable to swallow capsules. | on; and | | | |
| Renewal — (Systemic candidiasis) from any relevant practition following criteria: Both: | er. Approvals valio | d for 6 wee | ks for a | pplications meeting the |
| Patient requires prophylaxis for, or treatment of systemic c Patient is unable to swallow capsules. | | | | |
| Renewal — (Immunocompromised) from any relevant practition following criteria: All of the following: | ner. Approvals vali | id for 6 moi | nths for | applications meeting the |
| Patient remains immunocompromised; and Patient remains at moderate to high risk of invasive fungal Patient is unable to swallow capsules. | infection; and | | | |
| TRACONAZOLE | | | | |
| Cap 100 mg Oral liq 10 mg per ml – Special Authority see SA1322 below | | 15 | • | Itrazole |
| Retail pharmacy | | 50 ml OP | 1 | Sporanox |
| SA1322] Special Authority for Subsidy Initial application only from an infectious disease specialist, clinic practitioner on the recommendation of a infectious disease physic valid for 6 months where the patient has a congenital immune def Renewal from any relevant practitioner. Approvals valid for 6 mo benefitting from the treatment. | ian, clinical microb iciency. | iologist or o | clinical i | mmunologist. Approvals |
| KETOCONAZOLE | | | | |
| Tab 200 mg – PCT | CBS | 30 100 | 1 : 1 : | Burel S29 Strides Shasun S29 Taro S29 Teva- Ketoconazole S29 |
| NYSTATIN | | | | |
| Tab 500,000 u | (17.09) | 50 | I | Nilstat |
| Cap 500,000 u | 12.81 (15.47) | 50 | I | Nilstat |
| POSACONAZOLE - Special Authority see SA1285 below - Reta | | | | |
| Tab modified-release 100 mg Oral liq 40 mg per ml | | 24 05 ml OP | | Posaconazole Juno Devatis |
| SA1285 Special Authority for Subsidy Initial application only from a haematologist or infectious disease | e specialist. Appro | vals valid f | or 6 we | eks for applications |

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria: Either:

1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Si | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

| * Tab 250 mg | 4.48 | 42 | ✓ Apo-Terbinafine S29 |
|---|-------------------------------------|-------|----------------------------|
| | 8.97 | 84 | ✓ Deolate |
| VORICONAZOLE - Special Authority see SA1273 below | Retail pharmacy | | |
| Tab 50 mg | | 56 | ✓ Vttack |
| Tab 200 mg | | 56 | Vttack |
| Powder for oral suspension 40 mg per ml - Wastage | | | |
| claimable | | 70 ml | Vfend |
| | | | |

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

1 Detient is immerse

- Patient is immunocompromised; and
 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

104

| PRIMAQUINE - Special Authority see SA1684 on | the next page – Retail pharmacy |
|--|---------------------------------|
| Tab 15 mg | |

Sanofi
 Primaguine S29

S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply

100

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

■ SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria: and
- 2 Primaguine is to be given for a maximum of 21 days.

| Antitrichomonal Agents | | | |
|---|-----------------------|----------------|--------------------------------------|
| METRONIDAZOLE | | | |
| Tab 200 mg – Up to 30 tab available on a PSO | | 250 | Metrogyl |
| Tab 400 mg – Up to 15 tab available on a PSO | 5.23 | 21 | Metrogyl |
| Oral liq benzoate 200 mg per 5 ml | 25.00 | 100 ml | Flagyl-S |
| Suppos 500 mg | 24.48 | 10 | 🗸 Flagyl |
| ORNIDAZOLE | | | |
| Tab 500 mg | | 10 | Arrow-Ornidazole |
| - | | | |
| Antituberculotics and Antileprotics | | | |
| Note: There is no co-payment charge for all pharmaceuticals I immigration status. | isted in the Antitube | erculotics and | Antileprotics group regardless of |
| BEDAQUILINE – Special Authority see SA2244 below – Retai No patient co-payment payable | | | |
| Tab 100mg | 3,084.51 | 24 OP | Sirturo |
| ► SA2244 Special Authority for Subsidy Initial application — (multi-drug resistant tuberculosis) fro applications meeting the following criteria: Both: | om any relevant prac | ctitioner. App | rovals valid for 6 months for |
| The person has multi-drug resistant tuberculosis (MDR- 2 Ministry of Health's Tuberculosis Clinical Network has n of the treatment regimen. | <i>,,</i> | ual case and r | ecommends bedaquiline as part |

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist. 100 Lamprene S29

CYCLOSERINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.

| Cap 250 mg | | 60 | Cyclorin S29 |
|------------|--|----|--------------|
|------------|--|----|--------------|

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| | Subsidy (Manufacturer's Pri \$ | ice) Sub Per | Fully Brand or sidised Generic ✓ Manufacturer | |
|--|--------------------------------------|-----------------|---|-----------|
| DAPSONE – Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable | | | | |
| b) Prescriptions must be written by, or on the recommodermatologist | | is disease ph | iysician, clinical microbi | ologist o |
| Tab 25 mg | | 100 | Dapsone | |
| Tab 100 mg | | 100 | Dapsone | |
| THAMBUTOL HYDROCHLORIDE – Retail pharmacy-S | pecialist | | | |
| a) No patient co-payment payableb) Prescriptions must be written by, or on the recommendation | nendation of, an infectiou | ıs disease ph | iysician, clinical microbi | ologist c |
| respiratory physician | | | | Ū |
| Tab 100 mg | 85.73 | 100 | EMB Fatol \$29 | |
| Tab 400 mg | 49.34 | 56 | Myambutol \$29 | |
| ONIAZID – Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payableb) Prescriptions must be written by, or on the recommendation | nendation of, an internal | medicine phy | /sician, paediatrician, cl | inical |
| microbiologist, dermatologist or public health phys | | | | |
| F Tab 100 mg | 23.00 | 100 | ✓ <u>PSM</u> | |
| ONIAZID WITH RIFAMPICIN - Retail pharmacy-Specia | alist | | | |
| a) No patient co-payment payable | | | | |
| b) Prescriptions must be written by, or on the recommicrobiologist, dermatologist or public health phys | nendation of, an internal ician | medicine phy | /sician, paediatrician, cl | inical |
| Tab 100 mg with rifampicin 150 mg | | 100 | <u>Rifinah</u> | |
| Tab 150 mg with rifampicin 300 mg | 179.13 | 100 | <u>Rifinah</u> | |
| NEZOLID – Special Authority see SA2234 below – Reta | ail pharmacy | | | |
| No patient co-payment payable | | | | |
| Tab 600 mg | 276.89 | 10 | 🗸 Zyvox | |
| Oral liq 20 mg per ml | 1,879.00 | 150 ml | Zyvox | |
| Special Authority for Subsidy itial application — (multi-drug resistant tuberculosis pplications meeting the following criteria: oth: 1 The person has multi-drug resistant tuberculosis (N 2 Ministry of Health's Tuberculosis Clinical Network I the treatment regimen. | MDR-TB); and | | | |
| Ũ | : aliat | | | |
| ARA-AMINO SALICYLIC ACID – Retail pharmacy-Spec | alist | | | |
| a) No patient co-payment payableb) Prescriptions must be written by, or on the recommondation | nendation of an infectiou | is disease sn | ecialist clinical microbi | ntonist o |
| respiratory physician | | io diocuoc op | colaist, chindar microsi | ologiot |
| Grans for oral liq 4 g sachet | | 30 | Paser S29 | |
| ROTIONAMIDE – Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable | | | | |
| b) Prescriptions must be written by, or on the recommendation of the second seco | nendation of, an infectiou | ıs disease sp | ecialist, clinical microbi | ologist |
| Tab 250 mg | | 100 | Peteha S29 | |
| RAZINAMIDE – Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable | | | | |
| b) Prescriptions must be written by, or on the recommon respiratory physician | nendation of, an infectiou | is disease ph | iysician, clinical microbi | ologist |
| Tab 500 mg | 64.95 | 100 | AFT-Pyrazinam | ide |
| 06 V fully subsidised Principal Supply | S29 Unappro | | supplied under Section 29 | |

| | Subsidy | | Fully Brand or |
|---|--|--|--|
| | (Manufacturer's F \$ | Price) Sub Per | sidised Generic Manufacturer |
| IFABUTIN – Retail pharmacy-Specialist | . | | |
| a) No patient co-payment payable | | | |
| b) Prescriptions must be written by, or on the recomm gastroenterologist | endation of, an infectio | ous disease ph | ysician, respiratory physician o |
| 🖌 Cap 150 mg | | 30 | Mycobutin |
| IFAMPICIN – Subsidy by endorsement | | | |
| a) No patient co-payment payable | | | |
| b) For confirmed recurrent Staphylococcus aureus infe antimicrobial based on susceptibilities and the pres | | | 1 2 |
| Retail pharmacy - Specialist. Specialist must be an | • | 0.7 | |
| paediatrician, or public health physician. | | , | |
| ₭ Cap 150 mg | | 100 | ✓ <u>Rifadin</u> |
| € Cap 300 mg | | 100 | ✓ <u>Rifadin</u> |
| • Oral lig 100 mg per 5 ml | 12.60 | 60 ml | ✓ Rifadin Sanofi ✓ Rifadin |
| | 12.00 | 00 111 | |
| Antivirals | | | |
| ar ava proparations rafer to Eva Proparations. Anti Infacti | vo Proparations, page | 260 | |
| or eye preparations refer to Eye Preparations, Anti-Infecti | ve Freparations, page | 200 | |
| Hepatitis B Treatment | | | |
| NTECAVIR | | | |
| Tab 0.5 mg | | 30 | Entecavir (Rex) |
| AMIVUDINE – Special Authority see SA1685 below – Re | | ~~ | |
| Tab 100 mg Oral lig 5 mg per ml | | 28 240 ml OP | ✓ <u>Zetlam</u> ✓ Zeffix |
| ■SA1685 Special Authority for Subsidy | | 240 111 01 | ♥ Zellix |
| nitial application only from a relevant specialist or medica | al practitioner on the re | commendatio | n of a relevant specialist. |
| , . | | | · · · · · · · · · · · · · · · · · · · |
| pprovais valid for a year where used for the treatment of t | Jievention of hepatitis | υ. | |
| | | | nt or prevention of hepatitis B. |
| enewal from any relevant practitioner. Approvals valid fo ENOFOVIR DISOPROXIL | r 2 years where used f | or the treatme | |
| enewal from any relevant practitioner. Approvals valid fo ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for | r 2 years where used f | or the treatme | |
| enewal from any relevant practitioner. Approvals valid fo ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA: | r 2 years where used f the treatment of HIV is 2139., page 110 | for the treatme s included in th | e count of up to 4 subsidised |
| enewal from any relevant practitioner. Approvals valid fo ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA | r 2 years where used f the treatment of HIV is 2139., page 110 | or the treatme | |
| enewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA Tab 245 mg (300 mg as a maleate) | r 2 years where used f the treatment of HIV is 2139., page 110 | for the treatme s included in th | e count of up to 4 subsidised Tenofovir Disoproxil |
| enewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA Tab 245 mg (300 mg as a maleate) | r 2 years where used f the treatment of HIV is 2139., page 110 | for the treatme s included in th | e count of up to 4 subsidised Tenofovir Disoproxil |
| antiretrovirals for the purposes of Special Authority SA Tab 245 mg (300 mg as a maleate) Herpesvirus Treatments CICLOVIR | r 2 years where used f the treatment of HIV is 2139., page 110 | ior the treatme s included in th 30 | e count of up to 4 subsidised <u>Tenofovir Disoproxil</u> <u>Viatris</u> |
| Itenewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA: Tab 245 mg (300 mg as a maleate) Herpesvirus Treatments CICLOVIR Tab dispersible 200 mg | r 2 years where used f the treatment of HIV is 2139., page 110 | ior the treatme s included in th 30 25 | e count of up to 4 subsidised |
| tenewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA: Tab 245 mg (300 mg as a maleate) | r 2 years where used f the treatment of HIV is 2139., page 110 | ior the treatme s included in th 30 25 56 | e count of up to 4 subsidised |
| enewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA: Tab 245 mg (300 mg as a maleate) Herpesvirus Treatments CICLOVIR Tab dispersible 200 mg Tab dispersible 400 mg Tab dispersible 800 mg | r 2 years where used f the treatment of HIV is 2139., page 110 | ior the treatme s included in th 30 25 | e count of up to 4 subsidised |
| tenewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA: Tab 245 mg (300 mg as a maleate) Herpesvirus Treatments CICLOVIR E Tab dispersible 200 mg Tab dispersible 400 mg Tab dispersible 800 mg ALACICLOVIR | r 2 years where used f the treatment of HIV is 2139., page 110 | ior the treatme s included in th 30 25 56 35 | e count of up to 4 subsidised |
| enewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA: Tab 245 mg (300 mg as a maleate) Herpesvirus Treatments CICLOVIR Tab dispersible 200 mg Tab dispersible 400 mg ALACICLOVIR Tab 500 mg | r 2 years where used f the treatment of HIV is 2139., page 110 | ior the treatme s included in th 30 25 56 | Count of up to 4 subsidised <u>Tenofovir Disoproxil</u> <u>Viatris</u> <u>Lovir</u> <u>Lovir</u> <u>Lovir</u> <u>Lovir</u> <u>Vaclovir</u> |
| enewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA: Tab 245 mg (300 mg as a maleate) Herpesvirus Treatments CICLOVIR Tab dispersible 200 mg | r 2 years where used f the treatment of HIV is 2139., page 110 | ior the treatme is included in th 30 25 56 35 30 30 30 | e count of up to 4 subsidised |
| enewal from any relevant practitioner. Approvals valid for ENOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for antiretrovirals for the purposes of Special Authority SA: Tab 245 mg (300 mg as a maleate) Herpesvirus Treatments CICLOVIR Tab dispersible 200 mg Tab dispersible 400 mg ALACICLOVIR Tab 500 mg | r 2 years where used f the treatment of HIV is 2139., page 110 | ior the treatme is included in th 30 25 56 35 30 30 30 | Count of up to 4 subsidised <u>Tenofovir Disoproxil</u> <u>Viatris</u> <u>Lovir</u> <u>Lovir</u> <u>Lovir</u> <u>Lovir</u> <u>Vaclovir</u> |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or |
|----------------------|-----|------------|--------------|
| (Manufacturer's Pric | e) | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

⇒SA1993 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or

2 Both:

- 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
- 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|--|----------------------|---------------------|---|
| Hepatitis C Treatment | | | | |
| GLECAPREVIR WITH PIBRENTASVIR – [Xpharm] Note the supply of treatment is via Pharmac's approved dire website <u>https://pharmac.govt.nz/maviret</u> Tab 100 mg with pibrentasvir 40 mg | | Furth | | an be found on Pharmac's /laviret |
| LEDIPASVIR WITH SOFOSBUVIR – [Xpharm] – Special Author No patient co-payment payable Tab 90 mg with sofosbuvir 400 mg | rity see SA1605 belo 24,363.46 HepCTP) CTP). ct to confirmation of e | w 28 ligibilit | √ ⊦ y. | larvoni |

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – Subsidy by endorsement; can be waived by Special Authority see SA2138 below

- a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138.
- b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement, for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 110 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

➡SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

| | Subsidy | Fully | Brand or |
|---|---|--|--------------------------|
| | (Manufacturer's Price) | Subsidised | Generic |
| | \$ | Per 🗸 | Manufacturer |
| continued | | | |
| Patient has tested HIV negative, does not have signs of seroconversion; and | or symptoms of acute HIV | infection and ha | as been assessed for HIV |
| 2 The Practitioner considers the patient is at elevated ris | sk of HIV exposure and us | se of PrEP is clin | ically appropriate. |
| Notes: Refer to local health pathways or the Australasian So guidelines: | ciety for HIV, Viral Hepati | tis and Sexual H | ealth Medicine clinical |
| https://ashm.org.au/HIV/PrEP/ | | | |
| COVID-19 Treatments | | | |
| MOLNUPIRAVIR – [Xpharm] – Subsidy by endorsement a) No patient co-payment payable b) Treatment is funded only if patient meets access crite and has been endorsed accordingly by the prescriber process. Refer to the Pharmac website for more infor Cap 200 mg | . The supply of treatment rmation about this and sto | t is via Pharmac' ock availability. | · |
| NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by | y endorsement | | |
| a) No patient co-payment payable | | | |
| b) Treatment is funded only if patient meets access crite and has been endorsed accordingly by the prescriber process. Refer to the Pharmac website for more infor | . The supply of treatment | t is via Pharmac' | |
| Tab 150 mg with ritonavir 100 mg | | 30 V | Davlavid |

Tab 150 mg with ritonavir 100 mg0.00 30 V Paxlovid

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid

| | Subsidy | Fully | Brand or |
|-------|--------------------|------------|--------------|
| (Manu | ufacturer's Price) | Subsidised | Generic |
| | \$ Per | <i>✓</i> | Manufacturer |

continued...

for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

| EFAVIRENZ – Special Authority see SA2139 on the previous p Tab 200 mg Tab 600 mg | acy 90 30 | ✓ Stocrin ✓ Stocrin ✓ Efavirenz Milpharm [©]29 |
|--|---------------------|---|
| ETRAVIRINE – Special Authority see SA2139 on the previous Tab 200 mg | nacy 60 | ✓ Intelence |

| | Subsidy (Manufacturer's F \$ | Price) Subs Per | Fully Brand or idised Generic ✓ Manufacturer |
|---|-------------------------------------|--|---|
| VEVIRAPINE – Special Authority see SA2139 on page 110 – F Tab 200 mg | • • | 60 | ✓ <u>Nevirapine</u> <u>Alphapharm</u> |
| Oral suspension 10 mg per ml | 203.55 | 240 ml OP | Nevirapine Viatris Viramune Suspension |
| Nevirapine Alphapharm Tab 200 mg to be delisted 1 July 2024 |) | | · |
| Nucleosides Reverse Transcriptase Inhibitors | | | |
| BACAVIR SULPHATE - Special Authority see SA2139 on page | ge 110 – Retail pl | harmacy | |
| Tab 300 mg | | 60 | Ziagen |
| Oral liq 20 mg per ml | 256.31 | 240 ml OP | Ziagen |
| Ziagen Oral liq 20 mg per ml to be delisted 1 July 2024) | | | |
| BACAVIR SULPHATE WITH LAMIVUDINE – Special Authorit Note: abacavir with lamivudine (combination tablets) count acti retraviral Special Authority. | | | |
| anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg | 29.50 | 30 | ✓ <u>Abacavir/</u> Lamivudine Viatris |
| FAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOF | PROXII – Snecia | l Authority see | |
| harmacy | | i / tutilonty bee | one too on page the thetail |
| Note: Efavirenz with emtricitabine and tenofovir disoproxil of anti-retroviral Special Authority | counts as three a | nti-retroviral me | dications for the purposes of th |
| Tab 600 mg with emtricitabine 200 mg and tenofovir disopro 245 mg (300 mg as a maleate) | | 30 | ✓ Viatris |
| MTRICITABINE – Special Authority see SA2139 on page 110 Cap 200 mg | | су 30 | Emtriva |
| AMIVUDINE – Special Authority see SA2139 on page 110 – F | Retail pharmacy | | |
| Tab 150 mg | | 60 | Lamivudine Viatris |
| Oral liq 10 mg per ml | | 240 ml OP | ✓ 3TC |
| IDOVUDINE [AZT] - Special Authority see SA2139 on page 1 | 10 – Retail pharn | nacy | |
| | | 100 | Retrovir |
| Cap 100 mg Oral liq 10 mg per ml | | | ✓ Retrovir✓ Retrovir |
| Cap 100 mg Oral liq 10 mg per ml | 152.25 30.45 ee SA2139 on pag | 100 200 ml OP ge 110 – Retail | Retrovir pharmacy |
| Cap 100 mg Oral liq 10 mg per ml IDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority se Note: zidovudine [AZT] with lamivudine (combination tablet | | 100 200 ml OP ge 110 – Retail | Retrovir pharmacy |
| Cap 100 mg Oral liq 10 mg per ml IDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority se Note: zidovudine [AZT] with lamivudine (combination table the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg | | 100 200 ml OP ge 110 – Retail anti-retroviral m | Retrovir pharmacy nedications for the purposes of Alphapharm Lamivudine/ |
| Cap 100 mg Oral liq 10 mg per ml IDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority se Note: zidovudine [AZT] with lamivudine (combination tablet the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg | | 100 200 ml OP ge 110 – Retail anti-retroviral m | Retrovir pharmacy nedications for the purposes of Alphapharm Lamivudine/ |
| Cap 100 mg Oral liq 10 mg per ml CIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority se Note: zidovudine [AZT] with lamivudine (combination tablet the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg Alphapharm Tab 300 mg with lamivudine 150 mg to be delisted Protease Inhibitors | | 100 200 ml OP ge 110 – Retail anti-retroviral m 60 | Retrovir pharmacy nedications for the purposes of Alphapharm Lamivudine/ |
| Cap 100 mg Oral liq 10 mg per ml CIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority se Note: zidovudine [AZT] with lamivudine (combination tablet the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg | | 100 200 ml OP ge 110 – Retail anti-retroviral m 60 | Retrovir pharmacy nedications for the purposes of Alphapharm Lamivudine/ Zidovudine Viatris <u>Atazanavir Mylan</u> <u>Atazanavir Mylan</u> |
| Cap 100 mg Oral liq 10 mg per ml IDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority se Note: zidovudine [AZT] with lamivudine (combination tablet the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg Alphapharm Tab 300 mg with lamivudine 150 mg to be delisted Protease Inhibitors TAZANAVIR SULPHATE – Special Authority see SA2139 on Cap 150 mg | | 100 200 ml OP ge 110 – Retail anti-retroviral m 60 I pharmacy 60 | Retrovir pharmacy nedications for the purposes of Alphapharm Lamivudine/ Zidovudine Viatris |

| | Subsidy | | Fully | Brand or |
|--|---|--|--|---|
| | (Manufacturer's Price) \$ | Subs | idised | Generic Manufacturer |
| | Ŧ | 1.61 | | |
| DARUNAVIR – Special Authority see SA2139 on page 110 – | | | | - |
| Tab 400 mg | | 60 60 | | Darunavir Viatris |
| Tab 600 mg | | 60 | | Darunavir Viatris |
| LOPINAVIR WITH RITONAVIR - Special Authority see SA21 | | | | |
| Tab 100 mg with ritonavir 25 mg | 150.00 | 60 | ~ | Lopinavir/Ritonavir |
| | 005.00 | 100 | | <u>Mylan</u> Lemineuin/Ditemeuin |
| Tab 200 mg with ritonavir 50 mg | 295.00 | 120 | v | Lopinavir/Ritonavir Mylan |
| | | | | wyian |
| RITONAVIR – Special Authority see SA2139 on page 110 – F | | 00 | | Norvir |
| Tab 100 mg | | 30 | v | NOTVIE |
| Strand Transfer Inhibitors | | | | |
| DOLUTEGRAVIR - Special Authority see SA2139 on page 11 | 0 – Retail pharmacy | | | |
| Tab 50 mg | | 30 | 1 | Tivicay |
| DOLUTEGRAVIR WITH LAMIVUDINE - Special Authority see | SA2139 on page 110 | - Retail pł | arma | CV |
| Tab 50 mg with lamivudine 300 mg | | 30 | | Dovato |
| RALTEGRAVIR POTASSIUM – Special Authority see SA2139 | on page 110 – Retail r | harmacy | | |
| Tab 400 mg | | 60 | 1 | Isentress |
| Tab 600 mg | | 60 | 1 | Isentress HD |
| Immune Modulators | | | | |
| | | | | |
| | SA2034 below - Retai | Inharman | v | |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see | | | , | eed for ribavirin and meet |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small | group of patients who l | nave a clir | ical n | |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see | l group of patients who l Coordinator at Pharmac | nave a clir | , ical n)23-58 | |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C 0 | l group of patients who l Coordinator at Pharmac | nave a clir on 0800-0 | , ical n)23-58 | 8 option 4. |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C C Inj 180 mcg prefilled syringe | l group of patients who l Coordinator at Pharmac 748.50 | nave a clir on 0800-(4 | iical n 123-58 ✓ | 8 option 4. Pegasys |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C C Inj 180 mcg prefilled syringe | l group of patients who l Coordinator at Pharmac | nave a clir on 0800-0 4 fection w | iical no 23-58 V th HIV | 8 option 4. Pegasys / or genotype 2 or 3 post |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C C Inj 180 mcg prefilled syringe | l group of patients who l Coordinator at Pharmac | nave a clir on 0800-0 4 fection w | iical no 23-58 V th HIV | 8 option 4. Pegasys / or genotype 2 or 3 post |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C (Inj 180 mcg prefilled syringe | I group of patients who l Coordinator at Pharmac 748.50 or 6 infection or co-in nonths for applications r | nave a clir on 0800-0 4 fection w | iical no 23-58 V th HIV | 8 option 4. Pegasys / or genotype 2 or 3 post |
| PEGYLATED INTERFERON ALFA-2A - Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C (Inj 180 mcg prefilled syringe | I group of patients who l Coordinator at Pharmac | nave a clir on 0800-0 4 fection w | iical no 23-58 V th HIV | 8 option 4. Pegasys / or genotype 2 or 3 post |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C (Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac 748.50 or 6 infection or co-in nonths for applications r or 6 infection; or d with HIV; or | nave a clir on 0800-0 4 fection w neeting th | th HIV | 8 option 4. Pegasys / or genotype 2 or 3 post |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C (Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac 748.50 or 6 infection or co-in nonths for applications r or 6 infection; or d with HIV; or | nave a clir on 0800-0 4 fection w neeting th | th HIV | 8 option 4. Pegasys / or genotype 2 or 3 post |
| PEGYLATED INTERFERON ALFA-2A - Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C (Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | nave a clir on 0800-(4 fection w neeting th transplan | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C C Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C C Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C C Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C 0 Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A - Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C C Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C 0 Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C 0 Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C 0 Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C 0 Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | th HIV th HIV th and | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: |
| PEGYLATED INTERFERON ALFA-2A – Special Authority see Note: Pharmac will consider funding ribavirin for the small Special Authority criteria. Please contact the Hepatitis C 0 Inj 180 mcg prefilled syringe | I group of patients who I Coordinator at Pharmac | ave a clir on 0800-(4 fection w neeting th transplan ogist, infec | tical nu 123-58 th HIN e follo t; and tious o | 8 option 4. Pegasys / or genotype 2 or 3 post wing criteria: disease specialist or genera |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

| Subsidy (Manufacturer's Price) | Fully Subsidised | |
|-----------------------------------|---------------------|--------------|
| \$ | Per 🗸 | Manufacturer |

continued...

- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

| METHENAMINE (HEXAMINE) HIPPURATE | | | |
|---|--------|-----|-----------------------------|
| * Tab 1 g | 19.95 | 100 | ✓ <u>Hiprex</u> |
| NITROFURANTOIN | | | |
| * Tab 50 mg – Up to 30 tab available on a PSO | | 100 | Nifuran |
| * Tab 100 mg | 37.50 | 100 | ✓ Nifuran |
| * Cap modified-release 100 mg – Up to 15 cap available on a | | | |
| PSO | | 100 | ✓ Macrobid |
| NORFLOXACIN | | | |
| Tab 400 mg – Subsidy by endorsement | 245.00 | 100 | Arrow-Norfloxacin |

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

| | Subsidy | | Fully Brand or |
|---|------------------------------|-------|---|
| | (Manufacturer's Price) \$ | Per | Subsidised Generic Manufacturer |
| Anticholinesterases | | | |
| EOSTIGMINE METILSULFATE | | | |
| Inj 2.5 mg per ml, 1 ml ampoule | | 10 | Max Health |
| YRIDOSTIGMINE BROMIDE | | | |
| Tab 60 mg | 50.28 | 100 | Mestinon |
| Non-Steroidal Anti-Inflammatory Drugs | | | |
| ICLOFENAC SODIUM | | | |
| F Tab EC 25 mg | 1.99 | 50 | Diclofenac Sandoz |
| Tab 50 mg dispersible | 1.50 | 20 | Voltaren D |
| • Tab EC 50 mg | 1.99 | 50 | Diclofenac Sandoz |
| Tab long-acting 75 mg | | 100 | Voltaren SR |
| Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a | PSO 13.20 | 5 | Voltaren |
| Suppos 12.5 mg | | 10 | ✓ Voltaren |
| Suppos 25 mg | | 10 | Voltaren |
| Suppos 50 mg – Up to 10 supp available on a PSO | | 10 | ✓ Voltaren |
| · Suppos 100 mg | 7.00 | 10 | Voltaren |
| UPROFEN | | | |
| Tab 200 mg | | 1,000 | |
| Tab long-acting 800 mg | | 30 | Brufen SR |
| Oral liq 20 mg per ml | | 200 m | |
| | 11.29 | | Fenpaed 100 mg per 5 ml |
| ETOPROFEN | 40.07 | ~~ | |
| Cap long-acting 200 mg | 12.07 | 28 | Oruvail SR |
| EFENAMIC ACID | | | |
| Cap 250 mg | 1.25 | 50 | |
| | (10.82) | | Ponstan |
| | 0.50 | 20 | |
| | (7.50) | | Ponstan |
| APROXEN | | | |
| F Tab 250 mg | | 500 | Noflam 250 |
| • Tab 500 mg | | 250 | Noflam 500 |
| Tab long-acting 750 mg | 6.47 | 28 | Naprosyn SR 750 |
| Tab long-acting 1 g | 8.62 | 28 | Naprosyn SR 1000 |
| ENOXICAM | | | |
| F Tab 20 mg | | 100 | ✓ <u>Tilcotil</u> |
| Inj 20 mg vial | 9.95 | 1 | ✓ AFT |
| NSAIDs Other | | | |
| ELECOXIB | | | |
| Cap 100 mg | 3.45 | 60 | Celebrex |
| | | | Celecoxib Pfizer |
| 0 000 | 2.00 | 30 | ✓ Celebrex |
| Cap 200 mg | 3.20 | 30 | • Celebrex |

| | Subsidy (Manufacturer's Pric \$ | e) Sul Per | Fully Brand or bsidised Generic ✓ Manufacturer |
|--|---------------------------------------|---------------|--|
| Topical Products for Joint and Muscular Pain | ψ | Fei | |
| | | | |
| CAPSAICIN Crm 0.025% – Special Authority see SA1289 below – Retail | | | |
| pharmacy | | 45 g OP | Zostrix |
| | 13.00 | 60 g OP | Rugby Capsaicin |
| | | | Topical Cream ^{S29} |
| SA1289 Special Authority for Subsidy | | | Urcall a |
| itial application from any relevant practitioner. Approvals valid | d without further re | newal unles | ss notified where the patient has |
| steoarthritis that is not responsive to paracetamol and oral non-s | | | |
| Antirheumatoid Agents | | | |
| | | | |
| IYDROXYCHLOROQUINE – Subsidy by endorsement | | | |
| Subsidised only if prescribed for rheumatoid arthritis, systemi | | | - |
| suppression, relevant dermatological conditions (cutaneous f mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmo | | | |
| Pharmacists may annotate the prescription as endorsed whe | | | |
| hydroxychloroquine. Note: Indication marked with a * is an | | | |
| F Tab 200 mg | | 100 | Plaquenil |
| EFLUNOMIDE | | | |
| F Tab 10 mg | 6.00 | 30 | ✓ <u>Arava</u> |
| F Tab 20 mg | 6.00 | 30 | ✓ Arava |
| ENICILLAMINE | | | |
| Tab 125 mg | | 100 | D-Penamine D-Penamine |
| Tab 250 mg | | 100 | D-Penamine |
| Drugs Affecting Bone Metabolism | | | |
| Alendronate for Osteoporosis | | | |
| LENDRONATE SODIUM | | | |
| ← Tab 70 mg | 3.10 | 4 | Fosamax |
| Fosamax to be Principal Supply on 1 July 2024 | | | |
| LENDRONATE SODIUM WITH COLECALCIFEROL | | | |
| Tab 70 mg with colecalciferol 5,600 iu | 1.99 | 4 | Fosamax Plus |
| Fosamax Plus to be Principal Supply on 1 July 2024 | | | |
| Other Treatments | | | |
| ENOSUMAB - Special Authority see SA1777 below - Retail ph | | | |
| Inj 60 mg prefilled syringe | | 1 | Prolia |
| SA1777 Special Authority for Subsidy | | | |
| nitial application from any relevant practitioner. Approvals valid | d without further re | newal unles | ss notified for applications meeting |
| ne following criteria: | | | |

All of the following:

1 The patient has severe, established osteoporosis; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| (| Subsidy (Manufacturer's Price) | | | Brand or Generic |
|---|-----------------------------------|-----|---|---------------------|
| | \$ | Per | 1 | Manufacturer |

continued...

- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

| Inj 3 mg per ml, 10 ml vial | 32.49 | 1 | Pamisol |
|---|------------------|-------------|-----------------------------|
| Inj 6 mg per ml, 10 ml vial | 88.11 | 1 | Pamisol |
| Inj 9 mg per ml, 10 ml vial | | 1 | Pamisol |
| RALOXIFENE HYDROCHLORIDE - Special Authority see SA1779 | on the next page | – Retail pł | narmacy |
| * Tab 60 mg | 53.76 | 28 | Evista |

| Subsidy (Manufacturer's Price) | Sub | Fully sidised | Brand or Generic |
|---------------------------------------|-----|------------------|---------------------|
| \$ | Per | ~ | Manufacturer |

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

| Tab 35 mg | 2.50 | 4 | <u>Risedronate Sandoz</u> |
|---|--------------|---|---|
| TERIPARATIDE - Special Authority see SA1139 below - Ret | ail pharmacy | | |
| Inj 250 mcg per ml, 2.4 ml | | 1 | 🗸 Teriparatide - Teva |
| | 490.00 | | Forteo |
| Tavinavatida – Tava ta ka Dvinainal Oversky an 1 Ivez (| 2004 | | |

Teriparatide - Teva to be Principal Supply on 1 June 2024 (Forteo Inj 250 mcg per ml, 2.4 ml to be delisted 1 June 2024)

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy (Manufacturer's Price) | Subs | Fully sidised | Brand or Generic |
|-----------------------------------|------|------------------|---------------------|
| (Manulactarer 3 Theo) | Per | √ | Manufacturer |

continued...

absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable

- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

| ZOLEDRONIC ACID Inj 0.05 mg per ml, 100 ml, bag | 22.53 | 100 ml OP | ✓ <u>Zoledronic Acid</u> <u>Viatris</u> |
|--|--------------|-----------|--|
| Hyperuricaemia and Antigout | | | |
| ALLOPURINOL | | | |
| * Tab 100 mg | 11.47 | 500 | DP-Allopurinol |
| | 17.99 | 1,000 | Ipca-Allopurinol |
| Ipca-Allopurinol to be Principal Supply on 1 June 2024 | | | |
| * Tab 300 mg | 22.50 | 500 | Ipca-Allopurinol |
| | 28.57 | | DP-Allopurinol |
| Ipca-Allopurinol to be Principal Supply on 1 June 2024 | | | |
| (DP-Allopurinol Tab 100 mg to be delisted 1 June 2024) | | | |
| (DP-Allopurinol Tab 300 mg to be delisted 1 June 2024) | | | |
| BENZBROMARONE - Special Authority see SA1963 below - Ret | ail pharmacy | | |
| Tab 50 mg | | 100 | Narcaricin mite S29 |
| Tab 100 mg | | 30 | Desuric S29 |
| | | | Urinorm S29 |
| | 45.00 | 100 | Benzbromaron AL |
| | | | 100 S29 |
| | | | |

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 The treatment remains appropriate and the patient is benefitting from the treatment; and

2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg......6.00

100

Colgout

| | Subsidy (Manufacturer's Price) \$ | Su Per | Fully Ibsidised | Brand or Generic Manufacturer |
|--|---|-----------|--------------------|-------------------------------------|
| FEBUXOSTAT – Special Authority see SA2054 below – Retail p | pharmacy | | | |
| Tab 80 mg | | 28 | 🖌 F | ebuxostat (Teva) |
| | 20.00 | | ✓ F | ebuxostat multichem |
| Febuxostat (Teva) to be Principal Supply on 1 June 202 | 4 | | | |
| Tab 120 mg | 11.78 | 28 | 🖌 F | ebuxostat (Teva) |
| - | 20.00 | | ✓ F | ebuxostat multichem |

Febuxostat (Teva) to be Principal Supply on 1 June 2024 (Febuxostat multichem Tab 80 mg to be delisted 1 June 2024) (Febuxostat multichem Tab 120 mg to be delisted 1 June 2024)

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout..

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

| | DBENECID Tab 500 mg | 66.95 | 100 | ✓ Probenecid-AFT |
|----|---|--------|-----|--|
| N | uscle Relaxants | | | |
| ΒA | CLOFEN | | | |
| * | Tab 10 mg | 4.20 | 100 | Pacifen |
| | Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement | 11.55 | 1 | Lioresal Intrathecal |
| | Subsidised only for use in a programmable pump in patient caused intolerable side effects and the prescription is endo | | 1 0 | ents have been ineffective or have |
| | Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement | 306.82 | 5 | Medsurge |
| | Subsidised only for use in a programmable pump in patient caused intolerable side effects and the prescription is endo | | | ents have been ineffective or have |
| DA | NTROLENE | | | |
| | Cap 25 mg | 112.13 | 100 | Dantrium |
| | | | | Dantrium S29 S29 |
| | Cap 50 mg | 77.00 | 100 | Dantrium |
| | | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer | |
|------------------------------------|---|-----|---------------------|-------------------------------------|--|
| ORPHENADRINE CITRATE Tab 100 mg | 20.76 | 100 | ✓ N | lorflex | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|--|-------------------------------------|
| Agents for Parkinsonism and Related Disord | ers | | | |
| Dopamine Agonists and Related Agents | | | | |
| AMANTADINE HYDROCHLORIDE | | | | |
| Cap 100 mg | | 60 | | Symmetrel |
| | 63.73 | 100 | | Symmetrel |
| POMORPHINE HYDROCHLORIDE | | | | |
| Inj 10 mg per ml, 2 ml ampoule | | 5 | - | Movapo |
| Inj 10 mg per ml, 5 ml ampoule | 121.84 | 5 | ✓ | Movapo |
| NTACAPONE | | | | _ |
| Tab 200 mg | | 100 | | Comtan |
| EVODOPA WITH BENSERAZIDE | | | | |
| Tab dispersible 50 mg with benserazide 12.5 mg | | 100 | - | Madopar Rapid |
| Cap 50 mg with benserazide 12.5 mg | | 100 | - | Madopar 62.5 |
| Cap 100 mg with benserazide 25 mg | | 100 | - | Madopar 125 |
| Cap long-acting 100 mg with benserazide 25 mg | | 100 | | Madopar HBS |
| Cap 200 mg with benserazide 50 mg | | 100 | ✓ | Madopar 250 |
| EVODOPA WITH CARBIDOPA | | | | |
| Tab 100 mg with carbidopa 25 mg | | 100 | | Sinemet |
| Tab long-acting 200 mg with carbidopa 50 mg | | 100 | | Sinemet CR |
| Tab 250 mg with carbidopa 25 mg | | 100 | v : | Sinemet |
| RAMIPEXOLE HYDROCHLORIDE | | | | |
| Tab 0.25 mg | | 100 | | Ramipex |
| Tab 1 mg | | 100 | v | Ramipex |
| ASAGILINE | | | | |
| 🗧 Tab 1 mg | 53.50 | 30 | ✓ . | Azilect S29 |
| OPINIROLE HYDROCHLORIDE | | | | |
| Tab 0.25 mg | 4.05 | 84 | ✓ | Ropin |
| Tab 1 mg | 4.95 | 84 | ✓ | Ropin |
| Tab 2 mg | 6.48 | 84 | ✓] | Ropin |
| Tab 5 mg | 14.50 | 84 | ✓ [| <u>Ropin</u> |
| OLCAPONE | | | | |
| Tab 100 mg | 152.38 | 100 | Image: A second s | Tasmar |
| Anticholinergics | | | | |
| ENZATROPINE MESYLATE | | | | |
| Tab 2 mg | 9.59 | 60 | ✓ | Benztrop |
| Inj 1 mg per ml, 2 ml | | 5 | - | Phebra |
| a) Up to 10 inj available on a PSO | | - | | |
| b) Only on a PSO | | | | |
| ROCYCLIDINE HYDROCHLORIDE | | | | |
| Tab 5 mg | 7.40 | 100 | ✓ | Kemadrin |
| Agents for Essential Tremor, Chorea and Rel | | | | |
| | | | | |
| ILUZOLE – Special Authority see SA1403 on the next page Wastage claimable | | | | |
| Tab 50 mg | 130.00 | 56 | 1 | Rilutek |
| | | 00 | • | inutor |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

NERVOUS SYSTEM

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ Pe | er ✓ | Manufacturer |

► SA1403 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

| Tab 25 mg106.59 | 112 | ✓ Motetis | |
|-----------------|-----|-----------|--|
| | | | |

Anaesthetics

Local

| IDOCAINE [LIGNOCAINE] Gel 2%, tube – Subsidy by endorsement | 14.50 | 30 ml | Xylocaine 2% Jelly |
|--|-------------------|-----------------|--|
| a) Up to 150 ml available on a PSO | | 00111 | |
| b) Subsidised only if prescribed for urethral or cervical a | dministration an | d the prescript | tion is endorsed accordingly |
| Gel 2%, 11 ml urethral syringe – Subsidy by endorsement | | | Instillagel Lido |
| a) Up to 5 each available on a PSO | | | _ |
| b) Subsidised only if prescribed for urethral, cervical or i | rectal administra | tion and the p | rescription is endorsed |
| accordingly. | | | |
| IDOCAINE [LIGNOCAINE] HYDROCHLORIDE | | | |
| Oral (gel) soln 2% | | 200 ml | Mucosoothe |
| Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO | | 25 | Lidocaine-Baxter |
| | 17.50 | 50 | |
| | (35.00) | | Xylocaine |
| Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO | 9.00 | 25 | Lidocaine-Baxter |
| Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO | | 5 | |
| | (20.00) | | Xylocaine |
| Inj 1%, 20 ml vial – Up to 5 inj available on a PSO | 6.85 | 5 | Lidocaine-Baxter |
| Inj 2%, 20 ml vial – Up to 5 inj available on a PSO | 7.15 | 5 | Lidocaine-Baxter |
| Inj 10%, 5 ml ampoule - Subsidy by endorsement | CBS | 10 | Xylocard 500 S29 |
| Subsidised only for people receiving palliative care service | ces where other | analgesic age | nts haven't been effective. |

| | | | | V003 3131EM |
|--|---------------------------------------|--------------------------|---------------------|---|
| | Subsidy (Manufacturer's Prio \$ | ce) Per | Fully Subsidised | Brand or Generic Manufacturer |
| Topical Local Anaesthetics | | | | |
| SA0906 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid condition requiring frequent injections or venepuncture. Renewal from any relevant practitioner. Approvals valid for 2 yea benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 abor Crm 4%. | rs where the trea | tment rei | mains appro | |
| | 27.00 | 30 g O | | MX4 |
| LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Autho Crm 2.5% with prilocaine 2.5% Crm 2.5% with prilocaine 2.5% (5 g tubes) | | above – 1 30 g O 5 | Р 🧹 Е | nacy MLA MLA |
| Analgesics | | | | |
| or Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, pa | ge 116 | | | |
| Non-opioid Analgesics | | | | |
| ASPIRIN 券 Tab dispersible 300 mg − Up to 30 tab available on a PSO CAPSAICIN − Subsidy by endorsement | 5.65 | 100 | ✓ <u>E</u> | thics Aspirin |
| Subsidised only if prescribed for post-herpetic neuralgia or dia accordingly. | abetic peripheral r | neuropat | hy and the p | prescription is endorsed |
| accolungy. Crm 0.075% | 11.95 15.14 | 45 g O 57 g O | | ostrix HP ugby Capsaicin Topical Cream S29 |
| NEFOPAM HYDROCHLORIDE Tab 30 mg | 23 40 | 90 | ۷ ۵ | cupan |
| 1 ub 00 mg | | 50 | | oupuil |

| | Subsidy | | Fully Brand or | |
|--|-------------------------------|------------------|----------------------------------|-------------|
| | (Manufacturer's Pr | | sidised Generic | |
| | \$ | Per | Manufacturer | |
| PARACETAMOL | | | | |
| Tab 500 mg - blister pack | | 1,000 | Pacimol | |
| a) Maximum of 300 tab per prescription; can be | waived by endorsement | | | |
| b) Up to 30 tab available on a PSO | | | | |
| c) | | | | |
| Subsidy by endorsement for higher qua | intities is available for pat | tients with long | g term conditions who r | equire |
| regular daily dosing for one month or gr | | | | cists may |
| annotate the prescription as endorsed w | | | | |
| Maximum of 100 tab per dispensing for | | | | |
| (for non-endorsed patients), then disper | | s not exceedin | g 100 tab per dispensi | ng. |
| Tab 500 mg - bottle pack – Maximum of 300 tab per | | | * | |
| prescription; can be waived by endorsement | | 1,000 | Noumed | |
| | | | Paracetamol | |
| Subsidy by endorsement for higher quantit | ies is available for patient | ts with long ter | rm conditions who requ | ire regular |
| daily dosing for one month or greater, and | | | | innotate th |
| prescription as endorsed where dispensing | | | | |
| Maximum of 100 tab per dispensing for nor | | | | 0 tabs (for |
| non-endorsed patients), then dispense in re | epeat dispensings not ex | ceeding 100 ta | ab per dispensing. | |
| | | | _ | |
| Oral liq 120 mg per 5 ml | | 200 ml | Paracetamol | |
| | | | (Ethics) | |
| | 10.50 | 200 ml OP | Avallon | |
| a) Maximum of 600 ml per prescription; can be | waived by endorsement | | | |
| b) Up to 200 ml available on a PSO | | | | |
| c) Not in combination | | | | |
| d) | | | | |
| 1) Maximum of 200 ml per dispensing for | | | | nl (for |
| non-endorsed patients), then dispense | | | | |
| Subsidy by endorsement for higher qua transfer to the second sec | | | | |
| regular daily dosing for one month or gr | | | | |
| Pharmacists may annotate the prescrip | tion as endorsed where c | dispensing his | tory supports a long-ter | rm |
| condition. | | | O to a Manalantan (alla | |
| Note: 200 ml presentations of paraceta | | ipplied on BS | J to a vaccinator (othe | er than a |
| Pharmacist) under the provisions in Par | | | | |
| Note: Direct Provision by a pharmacist approximation with immunication of a shift | | | | |
| conjunction with immunisation of a chilo Oral liq 250 mg per 5 ml | | 200 ml | ✓ Pamol | vaccine. |
| | | 200 111 | | |
| a) Maximum of 600 ml per prescription; can be b) Up to 200 ml available on a PSO | walved by endorsement | | | |
| c) Not in combination | | | | |
| | | | | |
| d)1) Maximum of 200 ml per dispensing for the second se | non-endorsed nationte | f auantities or | escribed exceed 200 m | nl (for |
| non-endorsed patients), then dispense | | | | |
| Subsidy by endorsement for higher qua | | | | equire |
| regular daily dosing for one month or gr | | | | |
| Pharmacists may annotate the prescrip | | | | |
| condition. | | | ,, | |
| Note: 200 ml presentations of paraceta | amol oral liquid mav be su | upplied on BS | O to a Vaccinator (othe | r than a |
| Pharmacist) under the provisions in Par | | | | |
| 4) Note: Direct Provision by a pharmacist | | d under the pro | ovisions in Part I of Sec | ction A in |
| conjunction with immunisation of a child | | | | |
| ··· ;· ··· · ························· | , | | | |
| | | | | |

| | | Subsidy | | Fully | |
|-----|---|------------------------------|---------|-------------|----------------------------|
| | | (Manufacturer's Price) \$ | Per | Subsidised | Generic Manufacturer |
| | Suppos 125 mg | 4.29 | 10 | 1 | Gacet |
| | Suppos 250 mg | | 10 | | Gacet |
| | Suppos 500 mg | | 50 | | Gacet |
|) | pioid Analgesics | | | | |
| C | DEINE PHOSPHATE – Safety medicine; prescriber may d | etermine dispensina fre | auen | cv | |
| | Tab 15 mg | | 100 | | Noumed |
| | Tab 30 mg | | 100 | | Aspen |
| | 5 | | | ✓ | Noumed |
| | Tab 60 mg | | 100 | 1 | Noumed |
| Н | IYDROCODEINE TARTRATE | | | | |
| | Tab long-acting 60 mg | 8.60 | 60 | 1 | DHC Continus |
| - 1 | NTANYL | | | | |
| •• | a) Only on a controlled drug form | | | | |
| | b) No patient co-payment payable | | | | |
| | c) Safety medicine; prescriber may determine dispensing | frequency | | | |
| | Inj 50 mcg per ml, 2 ml ampoule | | 10 | 1 | Boucher and Muir |
| | Inj 50 mcg per ml, 10 ml ampoule | | 10 | | Boucher and Muir |
| | Patch 12.5 mcg per hour | | 5 | ✓ | Fentanyl Sandoz |
| | Patch 25 mcg per hour | 7.99 | 5 | ✓ | Fentanyl Sandoz |
| | Patch 50 mcg per hour | 9.49 | 5 | ✓ | Fentanyl Sandoz |
| | Patch 75 mcg per hour | | 5 | ✓ | Fentanyl Sandoz |
| | Patch 100 mcg per hour | | 5 | 1 | Fentanyl Sandoz |
| Ε | THADONE HYDROCHLORIDE | | | | |
| | a) Only on a controlled drug form | | | | |
| | b) No patient co-payment payable | | | | |
| | c) Safety medicine; prescriber may determine dispensing | frequency | | | |
| | d) Extemporaneously compounded methadone will only b | | e of th | ne cheapes | st form available |
| | (methadone powder, not methadone tablets). | | | | |
| | e) For methadone hydrochloride oral liquid refer Standard | Formulae, page 268 | | | |
| | Tab 5 mg | | 10 | 1 | Methadone BNM |
| | Oral liq 2 mg per ml | | 200 m | nl 🗸 | Biodone |
| | Oral liq 5 mg per ml | 6.40 | 200 m | nl 🗸 | Biodone Forte |
| | Oral liq 10 mg per ml | | 200 m | nl 🗸 | Biodone Extra Forte |
| | Inj 10 mg per ml, 1 ml | | 10 | 1 | AFT |
| C | RPHINE HYDROCHLORIDE | | | | |
| 1 | a) Only on a controlled drug form | | | | |
| | b) No patient co-payment payable | | | | |
| | c) Safety medicine; prescriber may determine dispensing | frequency | | | |
| | Oral lig 1 mg per ml | | 200 m | nl 🗸 | RA-Morph |
| | Oral lig 2 mg per ml | | 200 m | | RA-Morph |
| | Oral liq 5 mg per ml | | 200 m | | Ordine S29 |
| | | | | | RA-Morph |
| | | | | | • |
| | Oral liq 10 mg per ml | 27.74 | 200 m | 11 v | Ordine S29 |

(Ordine S29 Oral liq 10 mg per ml to be delisted 1 July 2024)

NERVOUS SYSTEM

| (N) | Subsidy Ianufacturer's Price |) | Fully Subsidised | |
|--|---------------------------------|---------|---------------------|-------------------|
| (14 | \$ | Per | | |
| ORPHINE SULPHATE | | | | |
| a) Only on a controlled drug form | | | | |
| b) No patient co-payment payable | | | | |
| c) Safety medicine; prescriber may determine dispensing frequ | ency | | | |
| Tab immediate-release 10 mg | | 10 | 1 | Sevredol |
| Tab immediate-release 20 mg | 5.52 | 10 | 1 | Sevredol |
| Cap long-acting 10 mg | 3.00 | 10 | ✓ | m-Eslon |
| Cap long-acting 30 mg | 4.30 | 10 | ✓ | m-Eslon |
| Cap long-acting 60 mg | 9.00 | 10 | 1 | m-Eslon |
| Cap long-acting 100 mg | 10.50 | 10 | ✓ | m-Eslon |
| Oral liq 2 mg per ml – Brand switch fee payable (Pharmacode | | | | |
| 2669986) - see page 265 for details | 16.31 | 100 m | l 🗸 | Wockhardt S29 |
| | 29.80 | | ✓ | Oramorph |
| Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO | 5.38 | 5 | 1 | Medsurge |
| Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSC | | 5 | | Medsurge |
| Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSC | | 5 | | Medsurge |
| Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSC | D6.28 | 5 | 1 | Medsurge |
| YCODONE HYDROCHLORIDE | | | | |
| a) Only on a controlled drug form | | | | |
| b) No patient co-payment payable | | | | |
| c) Safety medicine; prescriber may determine dispensing frequ | encv | | | |
| Tab controlled-release 5 mg | | 20 | 1 | Oxycodone Sandoz |
| 5 | 3.77 | 28 | | Oxycodone Sandoz |
| | | | | S29 S29 |
| | 4.04 | 30 | 1 | OxyContin S29 |
| Tab controlled-release 10 mg | | 20 | | Oxycodone Sandoz |
| | 3.77 | 28 | | Oxycodone Sandoz |
| | •••• | | | S29 S29 |
| Tab controlled-release 20 mg | 3 49 | 20 | 1 | Oxycodone Sandoz |
| Tab controlled-release 40 mg. | | 20 | | Oxycodone Sandoz |
| Tab controlled-release 80 mg | | 20 | | Oxycodone Sandoz |
| Cap immediate-release 5 mg | | 20 | | OxyNorm |
| Cap immediate-release 10 mg | | 20 | | OxyNorm |
| Cap immediate-release 20 mg | | 20 | | OxyNorm |
| Oral lig 5 mg per 5 ml | | 250 m | | OxyNorm |
| Inj 10 mg per ml, 1 ml ampoule | | 5 | | Hameln |
| Inj 10 mg per ml, 2 ml ampoule | | 5 | 1 | HameIn |
| Inj 50 mg per ml, 1 ml ampoule | | 5 | 1 | HameIn |
| RACETAMOL WITH CODEINE – Safety medicine; prescriber ma | av determine disr | ensin | n frequenc | V |
| Tab paracetamol 500 mg with codeine phosphate 8 mg | | 1.000 | | Paracetamol + |
| | | .,000 | 2 | Codeine (Relieve) |
| THIDINE HYDROCHLORIDE | | | | |
| | | | | |
| a) Only on a controlled drug form | | | | |
| b) No patient co-payment payablec) Safety medicine; prescriber may determine dispensing frequ | opov | | | |
| c) Safety medicine; prescriber may determine dispensing frequ Tab 50 mg | | 10 | | Noumed Pethidine |
| Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSC | | 10 5 | | DBL Pethidine |
| | 23.00 | 5 | • | Hydrochloride |
| Ini 50 ma normi. O mi omnovilo — Lin to 5 ini ovalishis sa a DOC | 00 70 | F | | • |
| Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSC | 130.72 | 5 | • | DBL Pethidine |
| | | | | Hydrochloride |

| | Subsidy | | Fully | |
|---|--|---|-------------|--|
| | (Manufacturer's Price) | Per | Subsidised | Generic Manufacturer |
| | \$ | rei | | INIGITUIACIUTEI |
| TRAMADOL HYDROCHLORIDE | 4.05 | ~~ | | Turnel OD 400 |
| Tab sustained-release 100 mg | | 20 | | Tramal SR 100 |
| Tab sustained-release 150 mg | | 20 | | Tramal SR 150 |
| Tab sustained-release 200 mg | | 20 100 | | Tramal SR 200 Arrow-Tramadol |
| Cap 50 mg | | 100 | v | Arrow-Tramadol |
| Antidepressants | | | | |
| Cyclic and Related Agents | | | | |
| AMITRIPTYLINE - Safety medicine; prescriber may determine | e dispensina frequency | | | |
| Tab 10 mg | | 100 | 1 | Arrow-Amitriptyline |
| Tab 25 mg | | 100 | | Arrow-Amitriptyline |
| Tab 50 mg | | 100 | | Arrow-Amitriptyline |
| CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; pres | criber may determine d | isner | nsina freau | ency |
| Tab 10 mg | | 30 | | Clomipramine Teva |
| Tab 25 mg | | 30 | | Clomipramine Teva |
| 0 | 39.97 | 100 | 1 | Anafranil S29 |
| Cap 10 mg | | 28 | 1 | Clomipramine - |
| | | | | Teva S29 |
| Cap 25 mg | 11.19 | 28 | 1 | Clomipramine |
| | | | | Teva S29 |
| | | | | 1010 |
| DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by | | | | |
| a) Safety medicine; prescriber may determine dispensing b) Subsidy by endorsement – Subsidised for patients who | | [dath | ionin1 hudr | achlarida prior to 1 lupa |
| , | were taking uusulepin | luouii | | |
| 2019 and the prescription is endorsed accordingly Ph | armacists may annotate | • | nrecorintia | |
| 2019 and the prescription is endorsed accordingly. Ph exists a record of prior dispensing of dosulenin (dothier | | • | prescriptio | |
| exists a record of prior dispensing of dosulepin [dothier | oin] hydrochloride. | the | | n as endorsed where there |
| exists a record of prior dispensing of dosulepin [dothien Tab 75 mg | bin] hydrochloride. 3.85 | • | · · · · | n as endorsed where there Dosulepin Viatris |
| exists a record of prior dispensing of dosulepin [dothier | bin] hydrochloride. 3.85 | the 30 | · · · · | n as endorsed where there Dosulepin Viatris Dosulepin |
| exists a record of prior dispensing of dosulepin [dothien Tab 75 mg | bin] hydrochloride. 3.85 | the 30 | 1 | n as endorsed where there Dosulepin Viatris Dosulepin Mylan ⁶²⁹ |
| exists a record of prior dispensing of dosulepin [dothien Tab 75 mg | bin] hydrochloride. 3.85 | the 30 | 1 | n as endorsed where there Dosulepin Viatris Dosulepin Mylan ⁵²⁹ Dosulepin |
| exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg | 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 | the 30 | 1 | n as endorsed where there Dosulepin Viatris Dosulepin Mylan ⁶²⁹ |
| exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg (Dosulepin Mylan 329) Cap 25 mg to be delisted 1 October 20 | oin] hydrochloride. 3.85 7.83 | 30 50 | J J | n as endorsed where there Dosulepin Viatris Dosulepin Mylan ⁶²⁹ Dosulepin Viatris ⁶²⁹ |
| exists a record of prior dispensing of dosulepin [dothiep Tab 75 mg Cap 25 mg (Dosulepin Mylan 529) Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib | bin] hydrochloride. | the 30 50 | g frequency | n as endorsed where there Dosulepin Viatris Dosulepin Mylan 529 Dosulepin Viatris 529 |
| exists a record of prior dispensing of dosulepin [dothie Tab 75 mg Cap 25 mg (Dosulepin Mylan 329) Cap 25 mg to be delisted 1 October 20 | 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 | the 30 50 nsing 50 | g frequency | n as endorsed where there Dosulepin Mylan ⁶²⁹ Dosulepin Viatris ⁶²⁹ |
| exists a record of prior dispensing of dosulepin [dothiep Tab 75 mg Cap 25 mg (Dosulepin Mylan ^{sze} Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg | 224) er may determine dispe 10.000 10.000 10.000 10.000 10.000 10.000 | the 30 50 nsing 50 100 | g frequency | n as endorsed where there Dosulepin Mylan ⁶²⁹ Dosulepin Viatris ⁶²⁹ Tofranil Tofranil |
| exists a record of prior dispensing of dosulepin [dothiep Tab 75 mg Cap 25 mg (Dosulepin Mylan 329) Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib | 224) er may determine dispe 10.000 10.000 10.000 10.000 10.000 10.000 | the 30 50 nsing 50 | g frequency | n as endorsed where there Dosulepin Viatris Dosulepin Mylan ⁶²⁹ Dosulepin Viatris ⁶²⁹ Tofranil Tofranil Imipramine |
| exists a record of prior dispensing of dosulepin [dothiep Tab 75 mg Cap 25 mg (Dosulepin Mylan ^{sze} Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg | 224) er may determine dispe | nsing 50 100 28 | g frequency | n as endorsed where there Dosulepin Viatris Dosulepin Mylan ⁶²⁹ Dosulepin Viatris ⁶²⁹ Tofranil Tofranil Imipramine Crescent ⁶²⁹ |
| exists a record of prior dispensing of dosulepin [dothien Tab 75 mg Cap 25 mg (Dosulepin Mylan 529 Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg Tab 25 mg | 224) er may determine dispe | the 30 50 50 100 28 50 | g frequenc: | n as endorsed where there Dosulepin Viatris Dosulepin Mylan ⁸²⁹ Dosulepin Viatris ⁸²⁹ Tofranil Tofranil Imipramine Crescent ⁸²⁹ Tofranil |
| exists a record of prior dispensing of dosulepin [dothien Tab 75 mg Cap 25 mg (Dosulepin Mylan 200 Cap 25 mg to be delisted 1 October 200 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg Tab 25 mg NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; pre | 224) er may determine dispe | the 30 50 50 100 28 50 100 28 | g frequence | n as endorsed where there Dosulepin Viatris Dosulepin Mylan \$29 Dosulepin Viatris \$29 Tofranil Tofranil Imipramine Crescent \$29 Tofranil Imipramil Infranil |
| exists a record of prior dispensing of dosulepin [dothien Tab 75 mg Cap 25 mg (Dosulepin Mylan 329) Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; pre Tab 10 mg | 224) er may determine dispe | the 30 50 50 100 28 50 100 100 100 | g frequence | n as endorsed where there Dosulepin Viatris Dosulepin Mylan \$29 Dosulepin Viatris \$29 Tofranil Tofranil Imipramine Crescent \$29 Tofranil Imipramine Crescent \$29 Tofranil |
| exists a record of prior dispensing of dosulepin [dothien Tab 75 mg Cap 25 mg (Dosulepin Mylan 329) Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg Tab 25 mg NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; pre | 224) er may determine dispe | the 30 50 50 100 28 50 100 28 | g frequence | n as endorsed where there Dosulepin Viatris Dosulepin Mylan \$29 Dosulepin Viatris \$29 Tofranil Tofranil Imipramine Crescent \$29 Tofranil Imipramil Infranil |
| exists a record of prior dispensing of dosulepin [dothien Tab 75 mg Cap 25 mg (Dosulepin Mylan 329) Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; pre Tab 10 mg | 224) er may determine dispe | the 30 50 50 100 28 50 100 100 100 | g frequence | n as endorsed where there Dosulepin Viatris Dosulepin Mylan \$29 Dosulepin Viatris \$29 Tofranil Tofranil Imipramine Crescent \$29 Tofranil Imipramine Crescent \$29 Tofranil |
| exists a record of prior dispensing of dosulepin [dothier Tab 75 mg Cap 25 mg (Dosulepin Mylan 200 Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg Tab 25 mg NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; pre Tab 10 mg Tab 25 mg Monoamine-Oxidase Inhibitors (MAOIs) - Non | 224) er may determine dispe | the 30 50 50 100 28 50 100 100 100 | g frequence | n as endorsed where there Dosulepin Viatris Dosulepin Mylan \$29 Dosulepin Viatris \$29 Tofranil Tofranil Imipramine Crescent \$29 Tofranil Imipramine Crescent \$29 Tofranil |
| exists a record of prior dispensing of dosulepin [dothier Tab 75 mg Cap 25 mg (Dosulepin Mylan ⁶²⁹ Cap 25 mg to be delisted 1 October 20 IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescrib Tab 10 mg Tab 25 mg NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; pre Tab 10 mg Tab 25 mg | bin] hydrochloride. | the 30 50 50 100 28 50 100 100 100 | g frequency | n as endorsed where there Dosulepin Viatris Dosulepin Mylan \$29 Dosulepin Viatris \$29 Tofranil Tofranil Imipramine Crescent \$29 Tofranil Imipramine Crescent \$29 Tofranil |

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|--|---|--|--|---|
| Monoamine-Oxidase Type A Inhibitors | | | | |
| MOCLOBEMIDE | | | | |
| * Tab 150 mg | | 60 | | Aurorix |
| * Tab 300 mg | | 60 | 1 | Aurorix |
| Selective Serotonin Reuptake Inhibitors | | | | |
| CITALOPRAM HYDROBROMIDE | | | | |
| * Tab 20 mg | 2.86 | 84 | 1 | Celapram |
| ESCITALOPRAM | | | | |
| * Tab 10 mg | | 28 | | Ipca-Escitalopram |
| | 1.07 | | ~ | Escitalopram (Ethics) |
| * Tab 20 mg | 1.49 | 28 | 1 | Ipca-Escitalopram |
| FLUOXETINE HYDROCHLORIDE | | | | · · · · |
| Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement | 2.50 | 28 | 1 | <u>Fluox</u> |
| When prescribed for a patient who cannot swallow accordingly; or | · | | | |
| 1) When prescribed for a patient who cannot swallow | ple of 20 mg in which | case | the presci | iption is deemed to be |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multij endorsed. Note: Tablets should be combined with | ple of 20 mg in which h capsules to facilitate | case | the presci emental 10 | iption is deemed to be |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multij | ple of 20 mg in which h capsules to facilitate | case incr | the presci emental 10 | iption is deemed to be) mg doses. |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multi endorsed. Note: Tablets should be combined with Cap 20 mg | ple of 20 mg in which h capsules to facilitate | case incr 30 | the presci emental 10 | iption is deemed to be) mg doses. Brown & Burk 529 |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multij endorsed. Note: Tablets should be combined with | ple of 20 mg in which h capsules to facilitate 2.22 3.13 | case incr 30 | the prescr emental 10 | iption is deemed to be) mg doses. Brown & Burk 529 |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multi endorsed. Note: Tablets should be combined with Cap 20 mg PAROXETINE * Tab 20 mg | ple of 20 mg in which h capsules to facilitate 2.22 3.13 | case incr 30 90 | the prescr emental 10 | iption is deemed to be o mg doses. Brown & Burk 529 <u>Arrow-Fluoxetine</u> |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with Cap 20 mg PAROXETINE * Tab 20 mg SERTRALINE | ple of 20 mg in which h capsules to facilitate 2.22 3.13 4.11 | case incr 30 90 | the prescr emental 10 | iption is deemed to be o mg doses. Brown & Burk 529 <u>Arrow-Fluoxetine</u> |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multij endorsed. Note: Tablets should be combined with Cap 20 mg PAROXETINE * Tab 20 mg SERTRALINE * Tab 50 mg | ple of 20 mg in which h capsules to facilitate | case incr 30 90 90 | the prescr emental 10 | iption is deemed to be o mg doses. Brown & Burk 529 <u>Arrow-Fluoxetine</u> Loxamine |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multij endorsed. Note: Tablets should be combined with Cap 20 mg PAROXETINE * Tab 20 mg SERTRALINE * Tab 50 mg | ple of 20 mg in which h capsules to facilitate | case incr 30 90 90 30 | the prescr emental 10 | iption is deemed to be mg doses. Brown & Burk 529 <u>Arrow-Fluoxetine</u> Loxamine Setrona |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multi endorsed. Note: Tablets should be combined with Cap 20 mg PAROXETINE * Tab 20 mg SERTRALINE * Tab 50 mg * Tab 100 mg Other Antidepressants | ple of 20 mg in which h capsules to facilitate | case incr 30 90 90 30 | the prescr emental 10 | iption is deemed to be mg doses. Brown & Burk 529 <u>Arrow-Fluoxetine</u> Loxamine Setrona |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multi endorsed. Note: Tablets should be combined with Cap 20 mg PAROXETINE * Tab 20 mg SERTRALINE * Tab 50 mg * Tab 100 mg Other Antidepressants | ple of 20 mg in which h capsules to facilitate 2.22 3.13 4.11 0.99 1.74 | case incr 30 90 90 30 | the prescience of the prescien | iption is deemed to be o mg doses. Brown & Burk 529 Arrow-Fluoxetine Loxamine Setrona Setrona Noumed |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with Cap 20 mg Cap 20 mg PAROXETINE * Tab 20 mg SERTRALINE * Tab 50 mg * Tab 100 mg MIRTAZAPINE Tab 30 mg | ble of 20 mg in which h capsules to facilitate | 28 30 | the prescuence of the prescuen | iption is deemed to be o mg doses. Brown & Burk 529 Arrow-Fluoxetine Loxamine Setrona Setrona Noumed Noumed |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multi endorsed. Note: Tablets should be combined with Cap 20 mg PAROXETINE * Tab 20 mg SERTRALINE * Tab 50 mg * Tab 100 mg Other Antidepressants WIRTAZAPINE | ble of 20 mg in which h capsules to facilitate | 288 30 28 30 28 | the prescuence of the prescuen | iption is deemed to be o mg doses. Brown & Burk 529 Arrow-Fluoxetine Loxamine Setrona Setrona Noumed Noumed Noumed |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multi endorsed. Note: Tablets should be combined with Cap 20 mg Cap 20 mg PAROXETINE * Tab 20 mg SERTRALINE * Tab 50 mg * Tab 100 mg Other Antidepressants WIRTAZAPINE Tab 30 mg Tab 45 mg Tab 45 mg | ble of 20 mg in which h capsules to facilitate | 28 30 | the prescuence of the prescuen | iption is deemed to be o mg doses. Brown & Burk 529 Arrow-Fluoxetine Loxamine Setrona Setrona Noumed Noumed |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with Cap 20 mg. Cap 20 mg. PAROXETINE Tab 20 mg. SERTRALINE Tab 50 mg. Tab 100 mg. Other Antidepressants WIRTAZAPINE Tab 30 mg. Tab 45 mg. VENLAFAXINE | ble of 20 mg in which h capsules to facilitate 2.22 3.13 4.11 0.99 1.74 2.60 2.60 | case incr 30 90 90 30 30 30 28 30 28 30 | the prescuence of the prescuen | iption is deemed to be o mg doses. Brown & Burk 529 Arrow-Fluoxetine Loxamine Setrona Setrona Noumed Noumed Noumed Noumed |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with Cap 20 mg. PAROXETINE Tab 20 mg. SERTRALINE Tab 50 mg. Tab 100 mg. Other Antidepressants MIRTAZAPINE Tab 45 mg. VENLAFAXINE Cap 37.5 mg. | ple of 20 mg in which h capsules to facilitate 2.22 3.13 4.11 0.99 1.74 2.60 | case incr 30 90 90 30 30 28 30 28 30 28 30 28 30 84 | the prescuence of the prescuen | iption is deemed to be o mg doses. Brown & Burk 529 Arrow-Fluoxetine Loxamine Setrona Setrona Noumed Noumed Noumed Noumed Noumed Noumed Noumed |
| When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with Cap 20 mg PAROXETINE * Tab 20 mg SERTRALINE * Tab 50 mg * Tab 100 mg MIRTAZAPINE Tab 30 mg | ble of 20 mg in which h capsules to facilitate 2.22 3.13 4.11 0.99 1.74 2.60 | case incr 30 90 90 30 30 30 28 30 28 30 | the prescuence of the prescuen | iption is deemed to be o mg doses. Brown & Burk 529 Arrow-Fluoxetine Loxamine Setrona Setrona Noumed Noumed Noumed Noumed |

| | Subsidy (Manufacturer's Pric | | Fully Brand or sidised Generic |
|--|---------------------------------|----------|-----------------------------------|
| | \$ | Per | Manufacturer |
| Antiepilepsy Drugs | | | |
| Agents for Control of Status Epilepticus | | | |
| DIAZEPAM - Safety medicine; prescriber may determine di | | | |
| Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorseme | nt27.92 | 5 | Hospira |
| a) Up to 5 inj available on a PSO | | | |
| b) Only on a PSO | | | |
| c) PSO must be endorsed "not for anaesthetic prod | | 5 | Ctopolid |
| Rectal tubes 5 mg – Up to 5 tube available on a PSO | | Э | ✓ <u>Stesolid</u> |
| PHENYTOIN SODIUM | | | |
| Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available o | | _ | |
| PSO | | 5 | Hospira |
| Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available o | | - | . Ile entire |
| PSO | 154.01 | 5 | Hospira |
| Control of Epilepsy | | | |
| CARBAMAZEPINE | | | |
| * Tab 200 mg | 14.53 | 100 | Tegretol |
| * Tab long-acting 200 mg | | 100 | Tegretol CR |
| | 33.96 | 200 | Tegretol CR |
| * Tab 400 mg | 34.58 | 100 | Tegretol |
| * Tab long-acting 400 mg | | 100 | Tegretol CR |
| Oral liq 20 mg per ml | | 250 ml | Tegretol |
| CLOBAZAM – Safety medicine; prescriber may determine c | | | |
| Tab 10 mg | 9.12 | 50 | Frisium |
| CLONAZEPAM – Safety medicine; prescriber may determin | e dispensing frequency | | |
| Oral drops 2.5 mg per ml | 7.38 | 10 ml OP | Rivotril |
| ETHOSUXIMIDE | | | |
| Cap 250 mg | | 56 | Essential |
| | | | Ethosuximide S29 |
| | 140.88 | 100 | Zarontin |
| Oral liq 250 mg per 5 ml | | 200 ml | Zarontin |
| GABAPENTIN | | | |
| Note: Not subsidised in combination with subsidised pro- | egabalin | | |
| ₭ Cap 100 mg | • | 100 | ✓ <u>Nupentin</u> |
| ₭ Cap 300 mg | 8.45 | 100 | Nupentin |
| ₭ Cap 400 mg | 10.26 | 100 | Nupentin |
| ACOSAMIDE - Special Authority see SA2267 on the next | | / | |
| Tab 50 mg | | 14 | Vimpat |
| Tab 100 mg | 50.06 | 14 | Vimpat |
| | 200.24 | 56 | Vimpat |
| ▲ Tab 150 mg | | 14 | Vimpat |
| * T 000 | 300.40 | 56 | Vimpat |
| ▲ Tab 200 mg | | 56 | Vimpat |

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | 9 | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

➡SA2267 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

| LAN | NOTRIGINE | | | |
|-----|---|---------|---------------|---|
| | Tab dispersible 2 mg | | 30 | Lamictal |
| | Tab dispersible 5 mg | 50.00 | 30 | Lamictal |
| * | Tab dispersible 25 mg | 4.20 | 56 | Logem |
| * | Tab dispersible 50 mg | 5.11 | 56 | Logem |
| * | Tab dispersible 100 mg | 6.75 | 56 | Logem |
| LE\ | /ETIRACETAM | | | |
| | Tab 250 mg | 5.84 | 60 | Everet |
| | Tab 500 mg | | 60 | Everet |
| | Tab 750 mg | | 60 | Everet |
| | Tab 1,000 mg | | 60 | Everet |
| | Oral liq 100 mg per ml | | 300 ml OP | Levetiracetam-AFT |
| PHI | ENOBARBITONE | | | |
| | For phenobarbitone oral liquid refer Standard Formulae, p | age 268 | | |
| | Tab 15 mg | • | 500 | ✓ PSM |
| | · ~ · · · · · · · · · · · · · · · · · · | 248.50 | | ✓ Noumed |
| | | | | Phenobarbitone |
| | Tab 30 mg | 398 50 | 500 | ✓ Noumed |
| | | | 000 | Phenobarbitone |
| (PS | M Tab 15 mg to be delisted 1 August 2024) | | | |
| • | č č <i>j</i> | | | |
| | ENYTOIN SODIUM | 75.00 | 200 | Dilantin Infatab |
| ጥ | Tab 50 mg | | 200 | Dilantin Dilantin |
| | Cap 30 mg | | 200 | ✓ Dilantin |
| × | Cap 100 mg Oral lig 30 mg per 5 ml | | 200 500 ml | Dilantin Dilantin Paediatric |
| | 1 01 | 22.03 | 500 mi | |
| PRI | EGABALIN | | | |
| | Note: Not subsidised in combination with subsidised gaba | • | 50 | |
| | Cap 25 mg | | 56 | Pregabalin Pfizer |
| | • | 7.80 | | Milpharm S29 |
| * | Cap 75 mg | 2.65 | 56 | Pregabalin Pfizer |
| | | 8.10 | | Milpharm S29 |
| | Cap 150 mg | 4.01 | 56 | Lyrica |
| | | | | Pregabalin Pfizer |
| | | 12.44 | | Milpharm S29 |
| | Cap 300 mg | 7.38 | 56 | Pregabalin Pfizer |
| PRI | MIDONE | | | |
| * | Tab 250 mg | | 100 | Primidone Clinect |
| | č | | | |

| | Subsidy | | Fully | |
|--|-----------------------|-------|------------|-------------------|
| | (Manufacturer's Price | / | Subsidised | |
| | \$ | Per | | Manufacturer |
| SODIUM VALPROATE | | | | |
| Tab 100 mg | | 100 | ✓ | Epilim Crushable |
| Tab 200 mg EC | 27.44 | 100 | ✓ | Epilim |
| Tab 500 mg EC | | 100 | ✓ | Epilim |
| * Oral liq 200 mg per 5 ml | | 300 m | nl 🗸 | Epilim S/F Liquid |
| | | | ✓ | Epilim Syrup |
| * Inj 100 mg per ml, 4 ml | 41.50 | 1 | ✓ | Epilim IV |
| STIRIPENTOL - Special Authority see SA2268 below - Retail ph | armacy | | | |
| Cap 250 mg | 509.29 | 60 | 1 | Diacomit |
| Powder for oral lig 250 mg sachet | | 60 | 1 | Diacomit |
| | | | | |

⇒SA2268 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

| ▲ Tab 25 mg | | 60 | Arrow-Topiramate |
|--|--------|-----|--|
| J. J | | | Topiramate Actavis |
| | 26.04 | | Topamax |
| ▲ Tab 50 mg | | 60 | Arrow-Topiramate |
| Ŭ | | | Topiramate Actavis |
| | 44.26 | | Topamax |
| ▲ Tab 100 mg | | 60 | Arrow-Topiramate |
| • | | | Topiramate Actavis |
| | 75.25 | | Topamax |
| ▲ Tab 200 mg | 55.19 | 60 | Arrow-Topiramate |
| • | | | Topiramate Actavis |
| | 129.85 | | Topamax |
| Sprinkle cap 15 mg | 20.84 | 60 | Topamax |
| Sprinkle cap 25 mg | | 60 | Topamax |
| VIGABATRIN – Special Authority see SA2088 below – Retail pha | | | |
| ▲ Tab 500 mg | | 100 | Sabril |
| Powder for oral soln 500 mg per sachet | | 60 | ✓ Sabril |

► SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|---------|--------------|--|
| (Manufacturer's Price) | Sub | sidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and

2 Either:

- 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
- 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 116

Acute Migraine Treatment

Principal Supply

| RIZATRIPTAN Tab orodispersible 10 mg | 4.84 | 30 | ✓ Rizamelt |
|---|------------------------|---------------|---------------------------------|
| SUMATRIPTAN | | | |
| Tab 50 mg | | 90 90 | ✓ <u>Sumagran</u> |
| Tab 100 mg Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum | | 90 | ✓ <u>Sumagran</u> |
| prescription | | 2 OP | ✓ <u>Clustran</u> |
| Prophylaxis of Migraine | | | |
| For Beta Adrenoceptor Blockers refer to CARDIOVAS | CULAR SYSTEM, page 49 | | |
| PIZOTIFEN ₭ Tab 500 mcg | 02.01 | 100 | - Condomigron |
| K Tab 500 mcg | 25.21 | 100 | Sandomigran |
| Antinausea and Vertigo Agents | | | |
| or Antispasmodics refer to ALIMENTARY TRACT, pa | age 8 | | |
| PREPITANT - Special Authority see SA0987 below | , , | | |
| Cap 2 × 80 mg and 1 × 125 mg | | 3 OP | Emend Tri-Pack |
| SA0987 Special Authority for Subsidy | | | |
| nitial application from any relevant practitioner. App metogenic chemotherapy and/or anthracycline-basec | | | |
| Renewal from any relevant practitioner. Approvals va | | | |
| hemotherapy and/or anthracycline-based chemothera | | | acigoing inging chickegoine |
| BETAHISTINE DIHYDROCHLORIDE | ., | , | |
| * Tab 16 mg | 3.70 | 100 | ✓ <u>Serc</u> |
| | | | |
| fully subsidised | S29 Unappro | oved medicine | e supplied under Section 29 |
| 134 Principal Supply | Solo Subsidiend Supply | | |

Sole Subsidised Supply

| | Subsidy (Manufacturer's Price) \$ | Su Per | Fully Ibsidised | Brand or Generic Manufacturer |
|--|---|-----------|--------------------|---|
| CYCLIZINE HYDROCHLORIDE Tab 50 mg CYCLIZINE LACTATE | 0.49 | 10 | ✓ <u>N</u> | ausicalm |
| Inj 50 mg per ml, 1 ml ampoule – Up to 10 inj available on a PSO | | 10 | √ <u>H</u> | ameln |
| DOMPERIDONE * Tab 10 mg | 4.00 | 100 | ✓ <u>D</u> | omperidone <u>Viatris</u> |
| HYOSCINE HYDROBROMIDE | | | | |
| Inj 400 mcg per ml, 1 ml ampoule Patch 1 mg per 72 hours – Special Authority see SA1998 | 93.00 | 10 | 🗸 N | artindale S29 |
| below – Retail pharmacy | 17.70 | 2 | 🗸 S | copoderm TTS |
| | 88.50 | 10 | | copolamine - Mylan copolamine - Mylan S29 ^{©29} |

➡SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

| METOCLOPRAMIDE HYDROCHLORIDE | | |
|--|-----|---|
| * Tab 10 mg – Up to 30 tab available on a PSO1.57 | 100 | Metoclopramide Actavis 10 |
| * Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO7.00 | 10 | ✓ Baxter |
| ONDANSETRON | | |
| * Tab 4 mg2.27 | 50 | Periset |
| Tab disp 4 mg – Up to 10 tab available on a PSO0.56 | 10 | Periset ODT |
| * Tab 8 mg4.10 | 50 | Periset |
| Tab disp 8 mg – Up to 10 tab available on a PSO | 10 | Periset ODT |
| PROCHLORPERAZINE | | |
| * Tab 3 mg buccal | 50 | |
| (30.00) | | Buccastem |
| (30.00) | | Max Health \$29 |
| (30.00) | | Prochlorperazine |
| | | Brown & Burk S29 |
| * Tab 5 mg – Up to 30 tab available on a PSO | 250 | Nausafix |
| | | ✓ Nausafix - S29 S29 |
| ✤ Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO25.81 | 10 | Stemetil |
| | | |

| | Subsidy | | Fully | Brand or |
|--|--|------------|-------------------------|---------------------|
| | (Manufacturer's Price) | Sub | osidised | Generic |
| | \$ | Per | ~ | Manufacturer |
| Authorita | | | | |
| Antipsychotics | | | | |
| General | | | | |
| General | | | | |
| MISULPRIDE - Safety medicine; prescriber may determine c | lispensing frequency | | | |
| Tab 100 mg | 7.21 | 30 | ✓ 9 | Sulprix |
| Tab 200 mg | | 60 | ✓ 9 | Sulprix |
| Tab 400 mg | | 60 | ✓ 9 | Sulprix |
| RIPIPRAZOLE – Safety medicine; prescriber may determine | dispensing frequency | | | - |
| Tab 5 mg | | 30 | 1 | Aripiprazole Sandoz |
| | | 00 | | Ascend |
| | | | • , | |
| Tob 10 mg | 10 50 | 20 | | Aripiprazole S29 |
| Tab 10 mg | | 30 | | Aripiprazole Sandoz |
| Tab 15 mg | | 30 | | Aripiprazole Sandoz |
| Tab 20 mg | | 30 | | Aripiprazole Sandoz |
| Tab 30 mg | | 30 | | Aripiprazole Sandoz |
| HLORPROMAZINE HYDROCHLORIDE – Safety medicine; p | prescriber may determ | ine disper | nsing fre | equency |
| Tab 25 mg – Up to 30 tab available on a PSO | 15.62 | 100 | ✓ I | Largactil |
| Tab 100 mg – Up to 30 tab available on a PSO | | 100 | ✓ I | Largactil |
| Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO | | 10 | ✓ I | Largactil |
| LOZAPINE – Hospital pharmacy [HP4] | | | | |
| Safety medicine; prescriber may determine dispensing freq | liency | | | |
| Tab 25 mg | | 50 | 1 | Clopine |
| | 0.00 | 00 | | Clozaril |
| | 13.37 | 100 | | Clopine |
| | 10.07 | 100 | | Clozaril |
| Tab 50 mg | 8 67 | 50 | | Clopine |
| Tab 50 mg | 17.33 | 100 | | Clopine |
| Tab 100 mg | | 50 | | Clopine |
| Tab 100 mg | | 50 | | Clozaril |
| | 34.65 | 100 | | Clopine |
| | 34.05 | 100 | | Clozaril |
| Tab 200 mg | 24 65 | 50 | | Clopine |
| Tab 200 mg | | 50 100 | | Clopine |
| Succession E0 mg nor ml | | 100 ml | | Versacloz |
| Suspension 50 mg per ml | | 100 mi | • | versacioz |
| ALOPERIDOL – Safety medicine; prescriber may determine | | | | _ |
| Tab 500 mcg – Up to 30 tab available on a PSO | | 100 | | Serenace |
| Tab 1.5 mg – Up to 30 tab available on a PSO | | 100 | | Serenace |
| Tab 5 mg – Up to 30 tab available on a PSO | | 50 | | Serenace |
| | 29.72 | 100 | | Serenace |
| Oral liq 2 mg per ml – Up to 200 ml available on a PSO | | 100 ml | | Serenace |
| Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a | PSO21.55 | 10 | ✓ 9 | Serenace |
| EVOMEPROMAZINE – Safety medicine; prescriber may dete | rmine dispensina frea | uencv | | |
| Tab 25 mg (33.8 mg as a maleate) | | 100 | I | Nozinan (Swiss) |
| Tab 25 mg as a maleate | | 100 | | Nozinan |
| Tab 100 mg (135 mg as a maleate) | | 100 | | Nozinan (Swiss) |
| Tab 100 mg as a maleate | | 100 | | Nozinan |
| | ······································ | | | |

| | Subsidy | | Fully | Brand or |
|---|-----------------------|--------|------------|--------------------|
| | (Manufacturer's Price | | Subsidised | |
| | \$ | Per | | Manufacturer |
| EVOMEPROMAZINE HYDROCHLORIDE - Safety medicine; p | orescriber may deter | mine d | spensing | frequency |
| Inj 25 mg per ml, 1 ml ampoule | | 5 | 1 | Neuraxpharm S29 |
| | | | 1 | Nozinan S29 S29 |
| | 24.48 | 10 | 1 | Wockhardt |
| Neuraxpharm 🐲 Inj 25 mg per ml, 1 ml ampoule to be delisted | d 1 August 2024) | | | |
| Nozinan S29 529 Inj 25 mg per ml, 1 ml ampoule to be delisted | - / | | | |
| ITHIUM CARBONATE – Safety medicine; prescriber may deter | v , | auonov | , | |
| Tab long-acting 400 mg | | 100 | | Priadel |
| Cap 250 mg | | 100 | | Douglas |
| | | 100 | • | Dougias |
| DLANZAPINE – Safety medicine; prescriber may determine disp | • • • | 20 | | Zunina |
| Tab 2.5 mg | 1.40 | 30 | • | Zypine |
| Zypine to be Principal Supply on 1 August 2024 Tab 5 mg | 1.02 | 30 | | Zypine |
| Zypine to be Principal Supply on 1 August 2024 | 1.90 | 30 | • | zypine |
| Tab orodispersible 5 mg | 2 4 2 | 28 | 1 | Zypine ODT |
| Tab 10 mg | | 30 | | Zypine |
| Zypine to be Principal Supply on 1 August 2024 | | 00 | • | Lypine |
| Tab orodispersible 10 mg | 2 89 | 28 | 1 | Zypine ODT |
| | | 20 | - | |
| PERICYAZINE – Safety medicine; prescriber may determine disp | | 100 | | Neulactil |
| Tab 2.5 mg | | 100 | | Neulactil |
| Tab 10 mg | | 100 | • | Neulacui |
| QUETIAPINE – Safety medicine; prescriber may determine dispe | 0 1 7 | | | . |
| Tab 25 mg | | 90 | | Quetapel |
| Tab 100 mg | | 90 | | Quetapel |
| Tab 200 mg | | 90 | | Quetapel |
| Tab 300 mg | | 90 | • | Quetapel |
| RISPERIDONE – Safety medicine; prescriber may determine dis | | | | |
| Tab 0.5 mg | | 60 | | Risperidone (Teva) |
| Tab 1 mg | | 60 | | Risperidone (Teva) |
| Tab 2 mg | | 60 | | Risperidone (Teva) |
| Tab 3 mg | | 60 | | Risperidone (Teva) |
| Tab 4 mg | | 60 | | Risperidone (Teva) |
| Oral liq 1 mg per ml | | 30 ml | | Risperon |
| | 17.80 | 100 m | • | Risperon |
| ZIPRASIDONE – Safety medicine; prescriber may determine dis | | | - | |
| Cap 20 mg | | 60 | | Zusdone |
| Cap 40 mg | | 60 | | Zusdone |
| Cap 60 mg | | 60 | | Zusdone |
| Cap 80 mg | | 60 | / | Zusdone |
| ZUCLOPENTHIXOL HYDROCHLORIDE – Safety medicine; pres | , | | 0 | |
| Tab 10 mg | 31.45 | 100 | ~ | Clopixol |
| | | | | |
| Depot Injections | | | | |
| Deportingeotions | | | | |

| Safety medicine; prescriber may determine dispensing frequence | | | |
|--|--------|---|--|
| Inj 300 mg vial | 273.56 | 1 | Ability Maintena S29 |
| Inj 400 mg vial | 341.96 | 1 | Abilify Maintena S29 |

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

⇒SA2312 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 Patient has a current Special Authority approval for olanzapine depot injection, risperidone depot injection or paliperidone depot injection; and
- 1.2 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with olanzapine depot injection, risperidone depot injection or paliperidone depot injection; or
- 2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been initiated on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection (see Note below for the olanzapine Special Authority criteria for new olanzapine depot injection patients prior to 1 April 2024).

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
 - The patient has schizophrenia; and
 - The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of aripiprazole depot injection has been associated with fewer days of intensive intervention than prior to the initiation of an atypical antipsychotic depot injection.

FLUPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

PALIPERIDONE - Special Authority see SA1429 on the next page - Retail pharmacy

| TEOLENTITIZOE DECANOATE - Salety medicine, prescriber may determine disp | споту печ | ucity |
|--|-------------|--|
| Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO | 5 | Fluanxol |
| Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO | 5 | Fluanxol |
| Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO | 5 | Fluanxol |
| HALOPERIDOL DECANOATE - Safety medicine; prescriber may determine dispe | nsing frequ | iency |
| Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO | 5 | Haldol |
| Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO | 5 | Haldol Concentrate |
| | | Haldol |
| | | Decanoas S29 |
| OLANZAPINE – Special Authority see SA2313 below – Retail pharmacy | | |
| a) Safety medicine; prescriber may determine dispensing frequency | | |
| b) Note – no new patients to be initiated on olanzapine. | | |
| Inj 210 mg vial | 1 | Zyprexa Relprevv |
| Inj 300 mg vial414.00 | 1 | Zyprexa Relprevv |
| lnj 405 mg vial504.00 | 1 | Zyprexa Relprevv |
| | | |

➡SA2313 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

| Safety medicine; prescriber may determine dispensing fr | requency | | |
|---|----------|---|-------------------------------------|
| Inj 25 mg syringe | | 1 | Invega Sustenna |
| Inj 50 mg syringe | | 1 | Invega Sustenna |
| Inj 75 mg syringe | | 1 | Invega Sustenna |
| Inj 100 mg syringe | | 1 | Invega Sustenna |
| Inj 150 mg syringe | | 1 | 🗸 Invega Sustenna |

| Subsidy (Manufacturer's Pri | , | Fully Subsidised | Brand or Generic | |
|--------------------------------|----|---------------------|---------------------|--|
| \$ | Pe | r 🖌 | Manufacturer | |

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

| Inj 175 mg syringe | | 1 | 🗸 Invega Trinza |
|--------------------|----------|---|-----------------|
| Inj 263 mg syringe | | 1 | 🗸 Invega Trinza |
| Inj 350 mg syringe | | 1 | 🗸 Invega Trinza |
| Inj 525 mg syringe | 1,305.36 | 1 | 🗸 Invega Trinza |

⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The patient has schizophrenia; and

2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

| Inj 25 mg vial | | 1 | Risperdal Consta |
|------------------|--------|---|--------------------------------------|
| Inj 37.5 mg vial | 178.71 | 1 | Risperdal Consta |
| Inj 50 mg vial | 217.56 | 1 | Risperdal Consta |

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

| Inj 200 mg per ml, 1 ml – Up to 5 inj available on a PSO 1 | 19.80 5 | Clopixol |
|--|---------|------------------------------|
|--|---------|------------------------------|

| | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully idised | Brand or Generic Manufacturer |
|---|---|-------------|-----------------|-------------------------------------|
| Anxiolytics | | | | |
| BUSPIRONE HYDROCHLORIDE | | | | |
| * Tab 5 mg | | 100 | ✓ В | suspirone Viatris |
| * Tab 10 mg | | 100 | ✓ <u>B</u> | uspirone Viatris |
| CLONAZEPAM - Safety medicine; prescriber may determine dis | spensing frequency | | | |
| Tab 500 mcg | | 100 | ✓ P | axam |
| Tab 2 mg | 10.78 | 100 | 🗸 P | axam |
| DIAZEPAM - Safety medicine; prescriber may determine disper | sing frequency | | | |
| Tab 2 mg | 0 1 2 | 500 | 🗸 A | rrow-Diazepam |
| Tab 5 mg | 115.00 | 500 | ✓ A | rrow-Diazepam |
| LORAZEPAM - Safety medicine; prescriber may determine disp | ensing frequency | | | |
| Tab 1 mg | | 250 | 🗸 🗸 | tivan |
| Tab 2.5 mg | | 100 | ✓ <u>A</u> | tivan |

Multiple Sclerosis Treatments

⇒SA2274 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
 - 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5° C); and
 - 1.4.5 Either:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
 - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
 - 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or

| | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully idised | Brand or Generic Manufacturer |
|--|--|--|------------------------------|-------------------------------------|
| continued | | | | |
| 1.6.2 A sign of that new inflammatory activity is a 1.6.3 A sign of that new inflammatory is a T2 lesion 1.6.4 A sign of that new inflammatory activity is a features of a recent attack that occurred with 1.6.5 A sign of that new inflammatory activity is new inflammatory activity inflammatory activity is new inflammatory activi | n with associated loc prominent T2 lesion t nin the last 2 years; o w T2 lesions compa | cal swelling that clearly or red with a | g; or / is res previo | ponsible for the clinical |
| Note: Treatment on two or more funded multiple sclerosis treatmer Renewal — (Multiple Sclerosis - dimethyl fumarate, fingolimo beta-1-beta, natalizumab and teriflunomide) from any relevant had an EDSS score of 0 to 6.0 (inclusive) with or without the use of the patient has walked 100 metres or more with or without aids in Note: Treatment on two or more funded multiple sclerosis treatment | d, glatiramer acetat practitioner. Approv of unilateral or bilater the last six months). | e, interfer als valid fo al aids at a | on bei or 12 m any tim | nonths where patient has |
| DIMETHYL FUMARATE – Special Authority see SA2274 on the p | previous page – Reta | ul pharmad | су | |
| a) Wastage claimable b) Note: Treatment on two or more funded multiple coloresis | traatmanta aimultan | | ot nor | mittad |
| b) Note: Treatment on two or more funded multiple sclerosis Cap 120 mg | | 14 | | rinned. Tecfidera |
| Cap 240 mg | | 56 | - | ecfidera |
| FINGOLIMOD – Special Authority see SA2274 on the previous p. a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis Cap 0.5 mg | treatments simultan | | not per | mitted. Silenya |
| | | | | allellya |
| GLATIRAMER ACETATE – Special Authority see SA2274 on the Note: Treatment on two or more funded multiple sclerosis tre Inj 40 mg prefilled syringe | atments simultaneou | | permitt | ed. Copaxone |
| INTERFERON BETA-1-ALPHA – Special Authority see SA2274 of Note: Treatment on two or more funded multiple sclerosis tree Inj 6 million iu prefilled syringe Injection 6 million iu per 0.5 ml pen injector | atments simultaneou | | oermitt | |
| INTERFERON BETA-1-BETA – Special Authority see SA2274 or Note: Treatment on two or more funded multiple sclerosis tre Inj 8 million iu per 1 ml | n the previous page – atments simultaneou | | permitt | |
| NATALIZUMAB – Special Authority see SA2274 on the previous Note: Treatment on two or more funded multiple sclerosis tre Inj 20 mg per ml, 15 ml vial | page – Retail pharma atments simultaneou | | | ed. 'ysabri |
| TERIFLUNOMIDE – Special Authority see SA2274 on the previou a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis Tab 14 mg | treatments simultan | | | mitted. Aubagio |
| Multiple Sclerosis Treatments - Other | | | | |

| OCRELIZUMAB - Special Authority see SA2273 on the | next page – Retail pharmacy | |
|--|----------------------------------|-----------------------------|
| Note: Treatment on two or more funded multiple scl | erosis treatments simultaneously | is not permitted. |
| Inj 30 mg per ml, 10 ml vial | | Ocrevus |

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | <u> </u> | Manufacturer |

⇒SA2273 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
 - 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 1.4.5 Either:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
 - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
 - 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and

Viaisom

| Subsidy | | Fully | Brand or | _ |
|------------------------|-----|------------|--------------|---|
| (Manufacturer's Price) | : | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

30

Sedatives and Hypnotics

MELATONIN – Special Authority see SA1666 below – Retail pharmacy

Restricted to patients aged 18 years or under.

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

| MIDAZOLAM - Safety medicine; prescriber may d | letermine dispensing frequency | | |
|--|--|------------|--------------------------------------|
| Inj 1 mg per ml, 5 ml ampoule | 6.10 | 10 | Midazolam-Baxter |
| Inj 1 mg per ml, 5 ml plastic ampoule – Up to | 10 inj available | | |
| on a PSO | | 10 | Pfizer |
| On a PSO for status epilepticus use only. | | epilepticu | us use only. |
| Inj 5 mg per ml, 3 ml ampoule | 5.00 | 5 | Midazolam-Baxter |
| Inj 5 mg per ml, 3 ml plastic ampoule – Up to | 5 inj available on | | |
| a PSO | | 5 | Pfizer |
| On a PSO for status epilepticus use only. | PSO must be endorsed for status | epilepticu | is use only. |
| PHENOBARBITONE SODIUM - Special Authority | y see <mark>SA1386 below</mark> – Retail pharm | nacy | |
| Inj 200 mg per ml, 1 ml ampoule | | 10 | Max Health S29 |
| ➡SA1386 Special Authority for Subsidy | | | |
| Initial application from any relevant practitioner. | Approvals valid without further rene | ewal unle | ss notified for applications meeting |
| the following criteria: | | | |
| Both: | | | |
| For the treatment of terminal agitation that i | is unresponsive to other agents; an | d | |
| 2 The applicant is part of a multidisciplinary te | eam working in palliative care. | | |
| TEMAZEPAM - Safety medicine; prescriber may | determine dispensing frequency | | |
| | | | |
| Tab 10 mg | , | 25 | ✓ Normison |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Fully Subsidised Per ✓ | Brand or Generic Manufacturer |
|---|---|------------------------------|-------------------------------------|
| ZOPICLONE – Safety medicine; prescriber may determine dispe Tab 7.5 mg | | 500 🗸 | Zopiclone Actavis |
| Spinal Muscular Atrophy | | | |
| NUSINERSEN – PCT only – Special Authority see SA2174 below Inj 12 mg per 5 ml vial | | 1 🗸 | Spinraza |
| Initial application — (spinal muscular atrophy (SMA)) from ar applications meeting the following criteria: All of the following: | ny relevant practitione | r. Approvals va | alid for 12 months for |
| Patient has genetic documentation of homozygous SMN1 heterozygous mutation; and Patient is 18 years of age or under; and Either: | gene deletion, homo: | zygous SMN1 p | oint mutation, or compound |
| 3.1 Patient has experienced the defined signs and sym 3.2 Both: 3.2.1 Patient is pre-symptomatic; and 3.2.2 Patient has three or less copies of SMN2. | nptoms of SMA type I | II or IIIa prior to | o three years of age; or |
| Renewal — (spinal muscular atrophy (SMA)) from any relevar meeting the following criteria: All of the following: | nt practitioner. Approv | vals valid for 12 | months for applications |
| There has been demonstrated maintenance of motor miles Patient does not require invasive permanent ventilation (a reversible cause while being treated with nusinersen; and Nusinersen not to be administered in combination other SI | t least 16 hours per d | ay) in the absen | ice of a potentially |
| RISDIPLAM – [Xpharm] – Special Authority see SA2203 below Note: the supply of risdiplam is via Pharmac's approved dire Pharmac's website https://pharmac.govt.nz/risdiplam Powder for oral soln 750 mcg per ml, 60 mg per bottle | | | can be found on Evrysdi |
| ► SA2203 Special Authority for Subsidy Initial application — (spinal muscular atrophy (SMA)) from an applications meeting the following criteria: All of the following: | ny relevant practitione | r. Approvals va | alid for 12 months for |
| Patient has genetic documentation of homozygous SMN1 heterozygous mutation; and Patient is 18 years of age or under; and Either: | gene deletion, homo: | zygous SMN1 p | oint mutation, or compound |
| 3.1 Patient has experienced the defined signs and sym3.2 Both: | ptoms of SMA type I | II or IIIa prior to | o three years of age; or |
| 3.2.1 Patient is pre-symptomatic; and3.2.2 Patient has three or less copies of SMN2. | | | |
| Renewal — (spinal muscular atrophy (SMA)) from any relevant meeting the following criteria: All of the following: | | | |
| There has been demonstrated maintenance of motor miles | stone function since to | reatment initiation | on; and |

- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

NERVOUS SYSTEM

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Brand or Subsidised Generic ✓ Manufacturer |
|--|---|-----|--|
| Stimulants/ADHD Treatments | | | |
| ATOMOXETINE | | | |
| Cap 10 mg | | 28 | APO-Atomoxetine S29 S29 |
| | | | ✓ Generic Partners |
| | 43.02 | | ✓ APO-Atomoxetine |
| APO-Atomoxetine to be Principal Supply on 1 August 202 | | | |
| Cap 18 mg | | 28 | Generic Partners |
| | 45.57 | | APO-Atomoxetine |
| APO-Atomoxetine to be Principal Supply on 1 August 202 | | | |
| Cap 25 mg | | 28 | Generic Partners |
| | 44.30 | | APO-Atomoxetine |
| APO-Atomoxetine to be Principal Supply on 1 August 202 | | | |
| Cap 40 mg | | 28 | Generic Partners |
| | 46.21 | | APO-Atomoxetine |
| APO-Atomoxetine to be Principal Supply on 1 August 202 | | ~~ | |
| Cap 60 mg | | 28 | APO-Atomoxetine S29 S29 |
| | | | Generic Partners |
| | 51.31 | | APO-Atomoxetine |
| APO-Atomoxetine to be Principal Supply on 1 August 202 | | | |
| Cap 80 mg | 56.45 | 28 | APO-Atomoxetine S29 S29 |
| | | | Generic Partners |
| | 65.20 | | APO-Atomoxetine |
| APO-Atomoxetine to be Principal Supply on 1 August 202 | | | |
| Cap 100 mg | 58.48 | 28 | APO-Atomoxetine S29 S29 |
| | | | Generic Partners |
| | 65.71 | | APO-Atomoxetine |
| APO-Atomoxetine to be Principal Supply on 1 August 202 | <u>2</u> 4 | | |
| (APO-Atomoxetine S29 S29 Cap 10 mg to be delisted 1 August 2024) (Generic Partners Cap 10 mg to be delisted 1 August 2024) (Generic Partners Cap 18 mg to be delisted 1 August 2024) (Generic Partners Cap 25 mg to be delisted 1 August 2024) (Generic Partners Cap 40 mg to be delisted 1 August 2024) (APO-Atomoxetine S29 S29 Cap 60 mg to be delisted 1 August 2024) | | | |
| (Generic Partners Cap 60 mg to be delisted 1 August 2024) (APO-Atomoxetine S29 S29 Cap 80 mg to be delisted 1 August 2 (Generic Partners Cap 80 mg to be delisted 1 August 2024) | | | |
| (APO-Atomoxetine S29 S29 Cap 100 mg to be delisted 1 August 2024) (APO-Atomoxetine S29 S29 Cap 100 mg to be delisted 1 August 2024) | t 2024) | | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|-------------------------------------|
| DEXAMFETAMINE SULFATE - Special Authority see SA11 | 49 below – Retail pharma | су | | |
| a) Only on a controlled drug form | | | | |
| b) Safety medicine; prescriber may determine dispensin | g frequency | | | |
| Tab 5 mg - Brand switch fee payable (Pharmacode 267 | 3886) - | | | |
| see page 265 for details | | 100 | 🗸 P | SM |
| | 28.50 | | 🗸 A | spen |
| | 29.80 | | 🗸 N | oumed |
| | | | | Dexamfetamine |
| Noumed Dexamfetamine to be Principal Supply on 1 | June 2024 | | | |

Noumed Dexamfetamine to be Principal Supply on 1 June 2024

(PSM Tab 5 mg to be delisted 1 June 2024) (Aspen Tab 5 mg to be delisted 1 June 2024)

► SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

1 The treatment remains appropriate and the patient is benefiting from treatment; and

2 Either:

- 2.1 Applicant is a paediatrician or psychiatrist; or
- 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

NERVOUS SYSTEM

| | Subsidy (Manufacturer's Price) | | Fully Subsidised | |
|--|-----------------------------------|---------|---------------------|------------------------------|
| | \$ | Per | | Manufacturer |
| METHYLPHENIDATE HYDROCHLORIDE – Special Authority se | e SA1964 below – R | etail p | harmacy | |
| a) Only on a controlled drug form | | | | |
| b) Safety medicine; prescriber may determine dispensing free | quency | | | |
| c) Note: Brand Switch Fee applies only to patients who have | e transferred from Co | ncerta | a brand d | ue to an out of stock. |
| Tab immediate-release 5 mg | 3.20 | 30 | ✓ | Rubifen |
| Tab immediate-release 10 mg | | 30 | ✓ | Ritalin |
| | | | ✓ | Rubifen |
| Tab extended-release 18 mg – Brand switch fee payable | | | | |
| (Pharmacode 2677822) - see page 265 for details | 7.75 | 30 | 1 | Methylphenidate ER - Teva |
| Tab immediate-release 20 mg | | 30 | 1 | Rubifen |
| Tab sustained-release 20 mg. | | 30 | ✓ | Rubifen SR |
| Tab extended-release 27 mg – Brand switch fee payable | | | | |
| (Pharmacode 2677822) - see page 265 for details | 11.45 | 30 | 1 | Methylphenidate ER - Teva |
| Tab extended-release 36 mg – Brand switch fee payable | | | | |
| (Pharmacode 2677822) - see page 265 for details | 15.50 | 30 | 1 | Methylphenidate ER - Teva |
| Tab extended-release 54 mg – Brand switch fee payable | | | | |
| (Pharmacode 2677822) - see page 265 for details | | 30 | ~ | Methylphenidate ER - Teva |

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER - Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been

| e) | Fully Subsidised | Brand or Generic | |
|-----------|---------------------|---------------------|--|
| \$ Per | 1 | Manufacturer | |

continued...

consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal -- (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER - Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA2305 below - Retail pharmacy

a) Only on a controlled drug form

| b) Safety medicine; prescriber may determine dispension | sing frequency | | |
|---|----------------|----|--------------------------------|
| Tab extended-release 18 mg | | 30 | Concerta |
| Tab extended-release 27 mg | 65.44 | 30 | Concerta |
| Tab extended-release 36 mg | 71.93 | 30 | Concerta |
| Tab extended-release 54 mg | | 30 | Concerta |
| Cap modified-release 10 mg | | 30 | Ritalin LA |
| Cap modified-release 20 mg | | 30 | Ritalin LA |
| Cap modified-release 30 mg | | 30 | Ritalin LA |
| Cap modified-release 40 mg | | 30 | Ritalin LA |
| | | | |

SA2305 Special Authority for Subsidy

Initial application - (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or difficulties with adherence; ٥r
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal - (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

| MODAFINIL - Special Authority see SA1999 on the next page - | - Retail pharmacy | | |
|---|-------------------|----|---|
| Tab 100 mg | | 60 | 1 |

Modavigil

| Subsidy | | Fully | Brand or |
|------------------------|-----|---------|--------------|
| (Manufacturer's Price) | Sub | sidised | Generic |
| | Per | siuiseu | Manufacturer |
| φ | rei | • | Wanulaclurer |

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and

3 Either:

- 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
- 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE

| * Tab 5 mg | 3.70 | 84 | Ipca-Donepezil |
|--|----------|----|-----------------------------------|
| 0 | 4.34 | 90 | Donepezil-Rex |
| Ipca-Donepezil to be Principal Supply on 1 June 2024 | | | |
| * Tab 10 mg | 5.50 | 84 | Ipca-Donepezil |
| 5 | 6.64 | 90 | Donepezil-Rex |
| Ipca-Donepezil to be Principal Supply on 1 June 2024 | | | • |
| (Donepezil-Rex Tab 5 mg to be delisted 1 June 2024) | | | |
| (Donepezil-Rex Tab 10 mg to be delisted 1 June 2024) | | | |
| RIVASTIGMINE - Special Authority see SA1488 below - Retail | nharmaov | | |
| 1 , | | | • - • • • • • • • • |
| Patch 4.6 mg per 24 hour | | 30 | Rivastigmine Patch |
| | | | BNM 5 |
| | 90.00 | | Exelon Patch 5 |
| Patch 9.5 mg per 24 hour | | 30 | Rivastigmine Patch |
| 3 | | | BNM 10 |
| | 90.00 | | Exelon Patch 10 |
| | 00.00 | | |

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 The patient has been diagnosed with dementia; and

2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

| | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully sidised | Brand or Generic Manufacturer |
|---|---|-------------|------------------|-------------------------------------|
| Treatments for Substance Dependence | | | | |
| BUPRENORPHINE WITH NALOXONE – Special Authority see S a) No patient co-payment payable b) Safety medicine; prescriber may determine dispensing free | | il pharma | су | |
| Tab sublingual 2 mg with naloxone 0.5 mg | 11.76 | 28 | ✓ <u>B</u> | uprenorphine Naloxone BNM |
| Tab sublingual 8 mg with naloxone 2 mg | | 28 | ✓ <u>B</u> | uprenorphine Naloxone BNM |

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from

any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

| Tab modified-release 150 mg | 15.00 | 30 | Zyban |
|-----------------------------|-------|----|---------------------------|
|-----------------------------|-------|----|---------------------------|

NERVOUS SYSTEM

| | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully Brand or idised Generic Manufacturer |
|--|---|-------------|--|
| DISULFIRAM Tab 200 mg | | 100 | ✓ Antabuse |
| NALTREXONE HYDROCHLORIDE - Special Authority see SA1 | 1408 below – Retail p | harmacy | |
| Tab 50 mg | 77.77 | 28 | ✓ Naltrexone AOP S29 |
| | 83.33 | 30 | ✓ <u>Naltraccord</u> |

► SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

| 3 11 1 41 1 01 0 | COUDIT A. |
|------------------|--|
| 1 28 | Habitrol |
| 3 7 | Habitrol |
| 5 28 | Habitrol |
| 9 7 | Habitrol |
| 2 28 | Habitrol |
| 9 7 | Habitrol |
| 5 216 | Habitrol |
| 9 36 | Habitrol |
| 5 216 | Habitrol |
| 5 36 | Habitrol |
| 2 204 | Habitrol |
| 7 96 | Habitrol |
| 2 204 | Habitrol |
| 7 96 | Habitrol |
| 7 204 | Habitrol |
| 7 96 | Habitrol |
| 7 204 | Habitrol |
| 7 96 | Habitrol |
| | |
| | 3 7 5 28 9 7 2 28 9 7 6 216 9 36 5 216 5 216 5 204 7 96 7 204 7 96 7 204 7 204 |

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.

c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

| Tab 0.5 mg × 11 and 1 mg × 42 16.67 | 53 OP | Varenicline Pfizer |
|-------------------------------------|-------|--|
| Tab 1 mg17.62 | 56 | ✓ Varenicline Pfizer |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or |
|-----------------------|-----|------------|--------------|
| (Manufacturer's Price | e) | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

➡SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

| | Subsidy | C | Fully sidised | Brand or Generic |
|--|---------------------------------|--------------|---------------|---------------------------------------|
| | (Manufacturer's Price) \$ | Per | | Manufacturer |
| Chemotherapeutic Agents | | | | |
| | | | | |
| Alkylating Agents | | | | |
| BENDAMUSTINE HYDROCHLORIDE - PCT only - Speciali | | | | |
| Inj 25 mg vial Inj 100 mg vial | | 1 1 | | ibomustin ibomustin |
| Inj 1 mg for ECP | | 1 mg | - | axter |
| ➡SA2153 Special Authority for Subsidy | | • | | |
| Initial application — (treatment naive CLL) only from a rele | | | | he recommendation of a |
| relevant specialist. Approvals valid for 12 months for applicat All of the following: | ions meeting the followir | ig criteria: | | |
| 1 The patient has Binet stage B or C, or progressive stag | ae A chronic lymphocytic | leukaem | ia requir | ing treatment: and |
| 2 The patient is chemotherapy treatment naive; and | 5 | | | |
| 3 The patient is unable to tolerate toxicity of full-dose FC | R; and | | | |
| 4 Patient has ECOG performance status 0-2; and 5 Patient has a Cumulative Illness Rating Scale (CIRS) : | score of < 6 ; and | | | |
| 6 Bendamustine is to be administered at a maximum do | | 1 and 2 | everv 4 v | weeks for a maximum of |
| 6 cycles. | . | | , | |
| Note: 'Chronic lymphocytic leukaemia (CLL)' includes small ly | | | mothera | py treatment is considered |
| to comprise a known standard therapeutic chemotherapy regi Initial application — (Indolent, Low-grade lymphomas) or | | | dical pr | actitionar on the |
| recommendation of a relevant specialist. Approvals valid for | | | | |
| All of the following: | · · · · · · · · · · · · · · · · | 5 | | 3 |
| 1 The patient has indolent low grade NHL requiring treat | ment; and | | | |
| 2 Patient has a WHO performance status of 0-2; and | | | | |
| 3 Any of the following: 3.1 Both: | | | | |
| 3.1.1 Patient is treatment naive; and | | | | |
| 3.1.2 Bendamustine is to be administered for | a maximum of 6 cycles (| in combir | ation wi | th rituximab when |
| CD20+); or | | | | |
| 3.2 Both: | | | | |
| 3.2.1 Patient is refractory to or has relapsed v | vithin 12 months of a ritu | ximab cor | ntaining | combined |
| chemo-immunotherapy regimen; and 3.2.2 Bendamustine is to be administered in c | combination with obinutur | zumab foi | ' a maxir | num of 6 cvcles: or |
| 3.3 All of the following: | | | | · · · · · · · · · · · · · · · · · · · |
| 3.3.1 The patient has not received prior bend | amustine therapy; and | | | |
| 3.3.2 Bendamustine is to be administered for | a maximum of 6 cycles i | n relapse | d patient | is (in combination with |
| rituximab when CD20+); and 3.3.3 Patient has had a rituximab treatment-fr | an interval of 12 months | or more. | or | |

3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:

2.2.1 Both:

- 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
- 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application --- (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2: and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and

~

5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

| BUSULFAN – PCT – Retail pharmacy-Specialist | | | |
|---|--------|-----------|--|
| Tab 2 mg | | 100 | Myleran |
| CARBOPLATIN – PCT only – Specialist | | | |
| Inj 10 mg per ml, 45 ml vial | | 1 | DBL Carboplatin |
| | 45.20 | | ✓ Carboplatin Ebewe |
| | 48.50 | | Carbaccord |
| Inj 1 mg for ECP | 0.10 | 1 mg | Baxter |
| CARMUSTINE – PCT only – Specialist | | | |
| Inj 100 mg vial | 710.00 | 1 | BiCNU |
| Inj 100 mg for ECP | | 100 mg OP | ✓ Baxter |
| CHLORAMBUCIL – PCT – Retail pharmacy-Specialist | | - | |
| Tab 2 mg | | 25 | Leukeran FC |
| CISPLATIN – PCT only – Specialist | | | |
| Inj 1 mg per ml, 50 ml vial | 15.00 | 1 | Cisplatin Ebewe |
| Inj 1 mg per ml, 100 ml vial | | 1 | Cisplatin Ebewe Cisplatin Ebewe |
| | 29.66 | 1 | ✓ DBL Cisplatin |
| Inj 1 mg for ECP | | 1 mg | ✓ Baxter |
| CYCLOPHOSPHAMIDE | | | Buxton |
| Tab 50 mg – PCT – Retail pharmacy-Specialist | 145.00 | 50 | Cyclonex |
| Inj 1 g vial – PCT – Retail pharmacy-Specialist | | 1 | ✓ Endoxan |
| | | 6 | ✓ Cytoxan |
| Inj 2 g vial – PCT only – Specialist | | 1 | Endoxan |
| Inj 1 mg for ECP – PCT only – Specialist | | 1 mg | ✓ Baxter |
| , , , , | | 1 119 | Buildi |
| IFOSFAMIDE – PCT only – Specialist | 06.00 | 4 | Holoxan |
| lnj 1 g | | 1 | ✓ Holoxan ✓ Holoxan |
| Inj 2 g | | | ✓ Holoxan ✓ Baxter |
| Inj 1 mg for ECP | 0.10 | 1 mg | - Daxler |

| | Subsidy Manufacturer's Price) | Subs | Fully sidised | Brand or Generic |
|--|----------------------------------|------|-----------------------|----------------------------|
| | \$ | Per | ✓ | Manufacturer |
| LOMUSTINE – PCT – Retail pharmacy-Specialist | | | | |
| Cap 10 mg | 132.59 | 20 | 1 | CeeNU |
| Cap 40 mg | 399.15 | 20 | 1 | CeeNU |
| MELPHALAN | | | | |
| Tab 2 mg – PCT – Retail pharmacy-Specialist | 40.70 | 25 | 1 | Alkeran |
| Inj 50 mg - PCT only - Specialist | | 1 | 1 | Megval S29 |
| | | | 1 | Melpha |
| | 67.80 | | 1 | Alkeran |
| | | | ✓ | Alkeran S29 S29 |
| DXALIPLATIN – PCT only – Specialist | | | | |
| Inj 100 mg vial | 25.01 | 1 | • | Oxaliplatin Actavis 100 |
| | 110.00 | | 1 | Oxaliplatin Ebewe |
| Inj 5 mg per ml, 20 ml vial | | 1 | 1 | Alchemy Oxaliplatin |
| | 46.32 | | | Oxaliplatin Accord |
| Inj 1 mg for ECP | 0.35 | 1 mg | 1 | Baxter |
| THIOTEPA – PCT only – Specialist | | | | |
| Inj 15 mg vial | CBS | 1 | 1 | Bedford S29 |
| | | | 1 | Max Health S29 |
| | | | 1 | THIO-TEPA S29 |
| | 398.00 | | ✓ | Tepadina |
| Inj 100 mg vial | CBS | 1 | 1 | Max Health S29 |
| | 1,800.00 | | 1 | Tepadina |
| Antimetabolites | | | | |
| AZACITIDINE – PCT only – Specialist – Special Authority see SA | 2141 below | | | |
| Inj 100 mg vial | | 1 | 1 | Azacitidine Dr Reddy's |
| Inj 1 mg for ECP | 0.83 | 1 mg | 1 | Baxter |
| SA2141 Special Authority for Subsidy | | Ŭ | | |

► SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

155

| | Subsidy (Manufacturer's Pri | ce) Sub | Fully sidised | Brand or Generic |
|--|--------------------------------|-----------|------------------|------------------------------------|
| | \$ | Per | ✓ | Manufacturer |
| LCIUM FOLINATE | | | | |
| Tab 15 mg – PCT – Retail pharmacy-Specialist | | 10 | 1 | DBL Leucovorin Calcium |
| Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist . | 17.10 | 5 | | Hospira |
| Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Spec | ialist7.28 | 1 | | Calcium Folinate Sandoz |
| | | | 1 | Calcium Folinate Sandoz S29 S29 |
| | 36.48 | 5 | ✓ | Eurofolic S29 |
| Inj 50 mg – PCT – Retail pharmacy-Specialist | 72.80 | 10 | • | Leucovorin Pharmacia S29 |
| Inj 10 mg per ml, 10 ml vial - PCT only - Specialist | 9.49 | 1 | 1 | Calcium Folinate Sandoz |
| | 47.45 | 5 | 1 | Eurofolic S29 |
| Inj 100 mg - PCT only - Specialist | 7.33 | 1 | 1 | Calcium Folinate Ebewe |
| | 94.90 | 10 | ✓ | Leucovorin Pharmacia S29 |
| Inj 300 mg - PCT only - Specialist | 22.51 | 1 | 1 | Calcium Folinate Ebewe |
| | 25.14 | | ✓ | Leucovorin DBL S29 |
| Inj 10 mg per ml, 35 ml vial - PCT only - Specialist | 25.14 | 1 | - | Calcium Folinate Sandoz |
| | | | 1 | Calcium Folinate Sandoz S29 S29 |
| Inj 1 g – PCT only – Specialist | 67.51 | 1 | 1 | Calcium Folinate Ebewe |
| Inj 10 mg per ml, 100 ml vial - PCT only - Specialist | 72.00 | 1 | 1 | Calcium Folinate Sandoz |
| Inj 1 mg for ECP – PCT only – Specialist | 0.06 | 1 mg | ✓ | Baxter |
| PECITABINE – Retail pharmacy-Specialist | | | | |
| Tab 150 mg | 9.80 | 60 | 1 | Capecitabine Viatris |
| Tab 500 mg | | 120 | 1 | Capecitabine Viatris |
| ADRIBINE – PCT only – Specialist | | | | |
| Inj 2 mg per ml, 5 ml | | 1 | | Litak S29 |
| Inj 1 mg per ml, 10 ml | | 1 | | Leustatin |
| Inj 10 mg for ECP TARABINE | 749.96 | 10 mg OP | | Baxter |
| Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Spec Inj 100 mg per ml, 20 ml vial – PCT – Retail | ialist472.00 | 5 | ✓ | Pfizer |
| pharmacy-Specialist | | 1 | | Pfizer Pfizer S29 S29 |
| Inj 1 mg for ECP – PCT only – Specialist | 0.29 | 10 mg | | Baxter |
| Inj 100 mg intrathecal syringe for ECP – PCT only – Spec UDARABINE PHOSPHATE | | 100 mg OP | ✓ | Baxter |
| Tab 10 mg - PCT - Retail pharmacy-Specialist | 412 00 | 20 | 1 | Fludara Oral |
| | | 20 | - | |
| Inj 50 mg vial – PCT only – Specialist | 634.00 | 5 | | Fludarabine Ebewe |

| | Subsidy (Manufacturer's Price |) | Fully Subsidised | |
|--|----------------------------------|---------|-----------------------|---------------------------|
| | \$ | Per | ✓ | Manufacturer |
| LUOROURACIL | | | | |
| Inj 50 mg per ml, 20 ml vial – PCT only – Specialist | | 1 | 1 | Fluorouracil Accord |
| Inj 50 mg per ml, 50 ml vial - PCT only - Specialist | 14.72 | 1 | ✓ | Fluorouracil Accord |
| Inj 50 mg per ml, 100 ml vial - PCT only - Specialist | | 1 | ✓ | Fluorouracil Accord |
| Inj 1 mg for ECP – PCT only – Specialist | 0.62 | 100 mg | g 🗸 | Baxter |
| GEMCITABINE HYDROCHLORIDE – PCT only – Specialist Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine) | , | | | |
| 26.3 ml vial | | 1 | 1 | DBL Gemcitabine |
| lnj 1 g | | 1 | 1 | Gemcitabine Ebewe |
| Inj 1 mg for ECP | | 1 mg | 1 | Baxter |
| RINOTECAN HYDROCHLORIDE – PCT only – Specialist | | | | |
| Inj 20 mg per ml, 5 ml vial | | 1 | 1 | Accord |
| | 71.44 | | ~ | Irinotecan Actavis 100 |
| | 100.00 | | 1 | Irinotecan-Rex |
| Inj 1 mg for ECP | 0.54 | 1 mg | 1 | Baxter |
| /ERCAPTOPURINE | | • | | |
| Tab 50 mg - PCT - Retail pharmacy-Specialist | | 25 | 1 | Puri-nethol |
| Oral suspension 20 mg per ml – Retail pharmacy-Specialist | | | | |
| Special Authority see SA1725 below | | 00 ml C | DP 🗸 | Allmercap |

► SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

| | Subsidy (Manufacturer's Price | | Fully Subsidised | Generic |
|---|----------------------------------|---------|---------------------|-----------------------------|
| | \$ | Per | | Manufacturer |
| | 0.00 | 00 | | T |
| Tab 2.5 mg – PCT – Retail pharmacy-Specialist | | 90 | | Trexate |
| Tab 10 mg – PCT – Retail pharmacy-Specialist Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Special | | 90 5 | | Trexate Methotrexate DBL |
| e inj 2.5 mg per mi, 2 mi – POT – Retail pharmacy-Special | 151 | 5 | | Methotrexate DBL |
| | | | • | S29 S29 |
| k Inj 7.5 mg prefilled syringe | | 1 | 1 | Methotrexate |
| | | | | Sandoz |
| Inj 10 mg prefilled syringe | | 1 | 1 | Methotrexate |
| , , , , , , | | | | Sandoz |
| Inj 15 mg prefilled syringe | 14.77 | 1 | 1 | Methotrexate |
| | | | | Sandoz |
| k Inj 20 mg prefilled syringe | 14.88 | 1 | 1 | Methotrexate |
| , ., , . | | | | Sandoz |
| k Inj 25 mg prefilled syringe | 14.99 | 1 | 1 | Methotrexate |
| | | | | Sandoz |
| Inj 30 mg prefilled syringe | 15.09 | 1 | 1 | Methotrexate |
| | | | | Sandoz |
| Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Spe | cialist30.00 | 5 | 1 | Methotrexate DBL |
| | | | | Onco-Vial |
| Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Sp | ecialist45.00 | 1 | 1 | DBL Methotrexate |
| | | | | Onco-Vial |
| Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Speci | alist25.00 | 1 | 1 | Methotrexate Ebewe |
| Inj 100 mg per ml, 50 ml vial – PCT – Retail | | | | |
| pharmacy-Specialist | 67.99 | 1 | 1 | Methotrexate Ebewe |
| Inj 1 mg for ECP – PCT only – Specialist | | 1 mg | | Baxter |
| Inj 5 mg intrathecal syringe for ECP – PCT only – Special | list4.73 | 5 mg O | Р 🗸 | Baxter |
| PEMETREXED – PCT only – Specialist – Special Authority s | ee SA1679 below | | | |
| Inj 100 mg vial | 60.89 | 1 | 1 | Juno Pemetrexed |
| Inj 500 mg vial | | 1 | | Juno Pemetrexed |
| Inj 1 mg for ECP | 0.55 | 1 mg | ✓ | Baxter |

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

| (Man | Subsidy | Fully | | Brand or |
|---------------------------------------|--------------------|------------|---|--------------|
| | ufacturer's Price) | Subsidised | | Generic |
| · · · · · · · · · · · · · · · · · · · | \$ | Per | 1 | Manufacturer |

continued...

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles: or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: All of the followina:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

т

| THIOGUANINE – PCT – Retail pharmacy-Specialist | | | |
|--|--------|----------------------------|--|
| Tab 40 mg | 25 | Lanvis | |
| | | | |

Other Cytotoxic Agents

| AMSACRINE - PCT only - Specialist | | |
|---|----------|---|
| Inj 50 mg per ml, 1.5 ml ampoule1,500.00 | 6 | Amsidine S29 |
| 4,736.00 | | Amsidine S29 |
| Inj 75 mg1,250.00 | 5 | 🗸 AmsaLyo 🖘 |
| ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist | | |
| Cap 0.5 mg1,175.87 | 100 | 🗸 Agrylin |
| ARSENIC TRIOXIDE – PCT only – Specialist | | |
| Inj 1 mg per ml, 10 ml vial4,817.00 | 10 | Phenasen |
| Inj 10 mg for ECP481.70 | 10 mg OP | Baxter |
| BLEOMYCIN SULPHATE – PCT only – Specialist | | |
| Inj 15,000 iu, vial | 1 | DBL Bleomycin Sulfate |
| Inj 1,000 iu for ECP14.32 | 1,000 iu | Baxter |
| BORTEZOMIB - PCT only - Specialist - Special Authority see SA1889 below | | |
| Inj 3.5 mg vial | 1 | DBL Bortezomib |
| Inj 1 mg for ECP22.26 | 1 mg | Baxter |

■ SA1889 Special Authority for Subsidy

Initial application - (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

| | Subsidy | | Fully | Brand or |
|--|------------------|-----------|--------|------------------------|
| (| Manufacturer's F | | idised | |
| | \$ | Per | 1 | Manufacturer |
| ACARBAZINE – PCT only – Specialist | | | | |
| Inj 200 mg vial | 72.11 | 1 | 1 | DBL Dacarbazine |
| , , | 580.60 | 10 | 1 | Dacarbazine |
| | | | | APP S29 |
| Inj 200 mg for ECP | 72 11 | 200 mg OP | 1 | Baxter |
| , . | | 200 | | |
| ACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist | 055.00 | 4 | | Coomoron |
| Inj 0.5 mg vial | | 1 | | Cosmegen |
| Inj 0.5 mg for ECP | 255.00 | 0.5 mg OP | • | Baxter |
| UNORUBICIN – PCT only – Specialist | | | | |
| Inj 2 mg per ml, 10 ml | | 1 | | Pfizer |
| Inj 20 mg vial | 1,495.00 | 10 | 1 | Daunorubicin |
| | | | | Zentiva S29 |
| Inj 20 mg for ECP | 171.93 | 20 mg OP | ✓ | Baxter |
| DCETAXEL - PCT only - Specialist | | - | | |
| Inj 20 mg | 48 75 | 1 | 1 | Docetaxel Sandoz |
| Inj 10 mg per ml, 8 ml vial | | 1 | | DBL Docetaxel |
| Inj 20 mg per ml, 4 ml vial | | 1 | | Docetaxel |
| | 20.00 | I | • | Accord S29 |
| lai 00 ma | 105.00 | | | |
| Inj 80 mg | | 1 | | Docetaxel Sandoz |
| Inj 1 mg for ECP | 0.35 | 1 mg | • | Baxter |
| XORUBICIN HYDROCHLORIDE – PCT only – Specialist | | | | |
| Inj 2 mg per ml, 5 ml vial | 10.00 | 1 | 1 | Doxorubicin Ebewe |
| Inj 2 mg per ml, 25 ml vial | 11.50 | 1 | 1 | Doxorubicin Ebewe |
| | 17.00 | | 1 | Arrow-Doxorubicin |
| Inj 2 mg per ml, 50 ml vial | 23.00 | 1 | | Doxorubicin Ebewe |
| Inj 2 mg per ml, 100 ml vial | 65.00 | 1 | 1 | Arrow-Doxorubicin |
| | 69.99 | | 1 | Accord S29 |
| | | | 1 | Doxorubicin Ebewe |
| Inj 1 mg for ECP | 0.35 | 1 mg | 1 | Baxter |
| IRUBICIN HYDROCHLORIDE - PCT only - Specialist | | - | | |
| Inj 2 mg per ml, 5 ml vial | 25.00 | 1 | 1 | Epirubicin Ebewe |
| Inj 2 mg per ml, 25 ml vial | | 1 | | Epirubicin Ebewe |
| Inj 2 mg per ml, 100 ml vial | | 1 | | Epirubicin Ebewe |
| Inj 1 mg for ECP | | 1 mg | | Baxter |
| | | i iiig | - | Builton |
| OPOSIDE | 0.40 70 | 00 | | Mana al d |
| Cap 50 mg – PCT – Retail pharmacy-Specialist | | 20 | | Vepesid |
| Cap 100 mg – PCT – Retail pharmacy-Specialist | | 10 | | Vepesid Dev Medical |
| Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist | | 1 | | Rex Medical |
| Inj 1 mg for ECP – PCT only – Specialist | 0.09 | 1 mg | • | Baxter |
| OPOSIDE PHOSPHATE – PCT only – Specialist | | | | |
| Inj 100 mg (of etoposide base) | | 1 | | Etopophos |
| Inj 1 mg (of etoposide base) for ECP | 0.47 | 1 mg | 1 | Baxter |
| DROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail pharm | acv-Specialist | | | |
| Cap 500 mg | | 100 | 1 | Devatis |
| | | | | |
| RUTINIB – Special Authority see SA2168 on the next page – Re | | 20 | | Imbruvica |
| Tab 140 mg | | 30 | | |
| Tab 420 mg | 9,052.00 | 30 | • | Imbruvica |

| Subsidy (Manufacturer's Price) | Subsi | Fully dised | Brand or Generic |
|-----------------------------------|-------|----------------|---------------------|
| \$ | Per | 1 | Manufacturer |

⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
 - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
 - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or
 - 4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

| Inj 5 mg vial – PCT only – Specialist | 109.74 | 1 | Zavedos |
|---|-----------------------|------|------------------------------|
| Inj 10 mg vial – PCT only – Specialist | | 1 | Zavedos |
| Inj 1 mg for ECP – PCT only – Specialist | | 1 mg | Baxter |
| LENALIDOMIDE – Retail pharmacy-Specialist – Special Au Wastage claimable | thority see SA2047 be | low | |
| Cap 5 mg | 5,122.76 | 28 | Revlimid |
| Cap 10 mg | 4,655.25 | 21 | Revlimid |
| | 6,207.00 | 28 | Revlimid |
| Cap 15 mg | 5,429.39 | 21 | Revlimid |
| | 7,239.18 | 28 | Revlimid |
| Cap 25 mg | 7,627.00 | 21 | Revlimid |

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

bortezomib or thalidomide that precludes further treatment with either of these treatments; and

4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

| Tab 400 mg – PCT – Retail pharmacy-Specialist | .00 50 | Uromitexan |
|--|------------|--|
| Tab 600 mg - PCT - Retail pharmacy-Specialist | .50 50 | Uromitexan |
| Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist | .45 15 | Uromitexan |
| Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist | .40 15 | Uromitexan |
| Inj 1 mg for ECP – PCT only – Specialist | .96 100 mg | Baxter |
| MITOMYCIN C – PCT only – Specialist | | |
| Inj 5 mg vial641 | .70 1 | Accord S29 |
| Inj 20 mg vial | .00 1 | Omegapharm S29 |
| , , | | 🗸 Teva |
| Inj 1 mg for ECP269 | .85 1 mg | Baxter |
| MITOZANTRONE – PCT only – Specialist | | |
| Inj 2 mg per ml, 10 ml vial97 | .50 1 | Mitozantrone Ebewe |
| Inj 1 mg for ECP5 | .51 1 mg | Baxter |
| NIRAPARIB – Special Authority see SA2325 below – Retail pharmacy | | |
| Wastage claimable | | |
| Cap 100 mg | .84 56 | 🗸 Zejula |
| 13,393 | .50 84 | 🗸 Zejula |

⇒SA2325 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | I Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 1 Patient has advanced high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line** of treatment with platinum-based chemotherapy; and
- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and
- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Either:
 - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen; or
 - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 No evidence of progressive disease; and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and

4 Either:

- 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
- 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: * "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component. **A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB - Retail pharmacy-Specialist - Special Authority see SA2163 below

| | | , | | , | |
|------------------------------|----|---|------|---|----------------|
| 🗸 Lynparza | 56 | | | | Tab 100 mg |
| Lynparza | 56 | | | | Tab 150 mg |

➡SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and

3 Either:

- 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
- 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 5 Treatment to be administered as maintenance treatment: and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Treatment remains clinically appropriate and patient is benefitting from treatment: and

2 Either:

- 2.1 No evidence of progressive disease; or
- 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion: and
- 3 Treatment to be administered as maintenance treatment: and
- 4 Treatment not to be administered in combination with other chemotherapy: and
- 5 Either:

5.1 Both:

- 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
- 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
- 5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component. **A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

| PACLITAXEL – PCT only – Specialist | | | |
|--|------------|------|--------------------------------------|
| Inj 30 mg | | 5 | Paclitaxel Ebewe |
| Inj 6 mg per ml, 16.7 ml vial | | 1 | Anzatax |
| | 24.00 | | Paclitaxel Ebewe |
| | 91.67 | | Paclitaxel Actavis |
| Inj 150 mg | | 1 | Paclitaxel Ebewe |
| | 137.50 | | Anzatax |
| | | | Paclitaxel Actavis |
| Inj 6 mg per ml, 50 ml vial | | 1 | Anzatax |
| | 44.00 | | Paclitaxel Ebewe |
| | 275.00 | | Paclitaxel Actavis |
| Inj 1 mg for ECP | 0.17 | 1 mg | Baxter |
| PEGASPARGASE - PCT only - Special Authority see SA | 1979 below | | |
| Inj 750 iu per ml, 5 ml vial | 3,455.00 | 1 | Oncaspar LYO \$29 |
| SA1070 Special Authority for Subsidy | | | |

SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal - (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of

| | Subsidy (Manufacturer's Price) | Subsi | Fully dised | Brand or Generic |
|--|-----------------------------------|--------------|----------------|---------------------|
| | \$ | Per | 1 | Manufacturer |
| continued | | | | |
| a relevant specialist. Approvals valid for 12 months for application Both: | is meeting the follow | ing criteria | : | |
| The patient has relapsed acute lymphoblastic leukaemia; a Pegaspargase to be used with a contemporary intensive m | | apy treatm | ient pro | otocol. |
| PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist | | | | |
| Inj 10 mg | CBS | 1 | 🗸 N | ipent S29 |
| PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharmacy- | Specialist | | | |
| Cap 50 mg | 980.00 | 50 | 🗸 N | atulan S29 |
| TEMOZOLOMIDE - Special Authority see SA2275 below - Retail | pharmacy | | | |
| Cap 5 mg | 9.13 | 5 | 🗸 T | emaccord |
| Cap 20 mg | 16.38 | 5 | - | emaccord |
| | 18.30 | | | po-Temozolomide |
| Cap 100 mg | | 5 | - | emaccord |
| | 40.20 | _ | | po-Temozolomide |
| Cap 140 mg | | 5 | - | emaccord |
| Cap 180 mg | | 14 | | ccord S29 |
| Cap 250 mg | | 5 | ✓ T | emaccord |

⇒SA2275 Special Authority for Subsidy

Initial application — (gliomas) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. **Renewal** — (gliomas) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

| THALIDOMIDE - Retail pharmacy-Specialist - Special A | Authority see SA1124 on the | next page |) |
|--|-----------------------------|-----------|------------------------------|
| Cap 50 mg | | 28 | Thalomid |
| Cap 100 mg | 756.00 | 28 | Thalomid |

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | Fu | lly | Brand or |
|------|--------------------|----------|-----|--------------|
| (Man | ufacturer's Price) | Subsidis | ed | Generic |
| | \$ | Per | ✓ | Manufacturer |

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Fither

Eitner:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

| Cap 10 mg – PCT – Retail pharmacy-Specialist | 479.50 | 100 | Vesanoid |
|---|-----------------|-------|-------------------------------|
| VENETOCLAX - Retail pharmacy-Specialist - Special Authority s | see SA1868 belo | W | |
| Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg | 1,771.86 | 42 OP | Venclexta |
| Tab 10 mg | | 2 OP | Venclexta |
| Tab 50 mg | 239.44 | 7 OP | Venclexta |
| Tab 100 mg - Wastage claimable | 8,209.41 | 120 | Venclexta |

► SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

| | Subsidy | | Fully Brand or |
|---|---|--|---|
| | (Manufacturer's Pr | | bsidised Generic |
| | \$ | Per | Manufacturer |
| INBLASTINE SULPHATE | | | |
| Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Spe | ecialist270.37 | 5 | Hospira |
| Inj 1 mg for ECP – PCT only – Specialist | 6.00 | 1 mg | Baxter |
| INCRISTINE SULPHATE | | | |
| Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Spec | cialist51.37 | 5 | DBL Vincristine |
| | | | Sulfate |
| Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Spec | cialist 102.73 | 5 | DBL Vincristine |
| | | | Sulfate |
| Inj 1 mg for ECP – PCT only – Specialist | 12.60 | 1 mg | Baxter |
| INORELBINE | | | |
| Cap 20 mg | | 1 | Vinorelbine Te Arai |
| Cap 30 mg | 40.00 | 1 | Vinorelbine Te Arai |
| Cap 80 mg | | 1 | Vinorelbine Te Arai |
| Inj 10 mg per ml, 1 ml vial – PCT only – Specialist | 12.00 | 1 | Navelbine |
| | 42.00 | | Vinorelbine Ebewe |
| Inj 10 mg per ml, 5 ml vial – PCT only – Specialist | 56.00 | 1 | Navelbine |
| | 168.00 | | Navelbine S29 S29 |
| | 210.00 | | Vinorelbine Ebewe |
| | 328.65 | | Sagent S29 |
| | | 1 mg | ✓ Baxter |
| Inj 1 mg for ECP – PCT only – Specialist | | i ing | Bunton |
| Inj 1 mg for ECP – PCT only – Specialist Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe | er 2024) | , ng | |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe | er 2024) | g | |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors | er 2024) er 2024) | | |
| Vavelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Vavelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable | er 2024) er 2024) e see SA1870 below | | |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority | er 2024) er 2024) e see SA1870 below | 224 | ✓ Alecensa |
| Vavelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Vavelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) e see SA1870 below | | |
| Vavelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Vavelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below | 224 | ✓ Alecensa |
| Vavelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Vavelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: | 224 commendatio | ✓ Alecensa on of a relevant specialist. |
| Avelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Avelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu | 224 commendation | Alecensa on of a relevant specialist. |
| Aveibine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Aveibine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu | 224 commendation | Alecensa on of a relevant specialist. |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu | 224 commendation | Alecensa on of a relevant specialist. |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu s an ALK tyrosine kin | 224 commendation ng cancer; an ase gene rea | ✓ Alecensa on of a relevant specialist. nd arrangement using an appropria |
| Vavelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Vavelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu s an ALK tyrosine kin | 224 commendation ng cancer; an ase gene rea | ✓ Alecensa on of a relevant specialist. nd arrangement using an appropria |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu s an ALK tyrosine kin | 224 commendation ng cancer; an ase gene rea | ✓ Alecensa on of a relevant specialist. nd arrangement using an appropria |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu s an ALK tyrosine kin er on the recommend | 224 commendation ng cancer; an ase gene rea | ✓ Alecensa on of a relevant specialist. nd arrangement using an appropria |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu s an ALK tyrosine kin eer on the recommend CIST criteria; and | 224 commendation ng cancer; an ase gene rea | ✓ Alecensa on of a relevant specialist. nd arrangement using an appropria |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below | 224 commendation ng cancer; an ase gene rea | ✓ Alecensa on of a relevant specialist. nd arrangement using an appropria |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu s an ALK tyrosine kin eer on the recommend CIST criteria; and t. ge – Retail pharmacy | 224 commendation ng cancer; an ase gene rea dation of a re | ✓ Alecensa on of a relevant specialist. nd arrangement using an appropria levant specialist. Approvals va |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu s an ALK tyrosine kin eer on the recommend CIST criteria; and t. ge – Retail pharmacy | 224 commendation ng cancer; an ase gene rea | ✓ Alecensa on of a relevant specialist. nd arrangement using an appropria levant specialist. Approvals va |
| Navelbine Inj 10 mg per ml, 1 ml vial to be delisted 1 Octobe Navelbine Inj 10 mg per ml, 5 ml vial to be delisted 1 Octobe Protein-tyrosine Kinase Inhibitors LECTINIB – Retail pharmacy-Specialist – Special Authority Wastage claimable Cap 150 mg | er 2024) er 2024) r see SA1870 below 7,935.00 practitioner on the re owing criteria: able, non-small cell lu s an ALK tyrosine kin er on the recommend CIST criteria; and t. ge – Retail pharmacy | 224 commendation ng cancer; au ase gene rea dation of a re 60 | ✓ Alecensa on of a relevant specialist. nd arrangement using an appropria levant specialist. Approvals va |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy (Manufacturer's Price) \$ | Fu) Subsidis Per | | |
|---|-------------------------|--|--|
|---|-------------------------|--|--|

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or

3 All of the following:

- 3.1 The patient has a diagnosis of CML in chronic phase; and
- 3.2 Maximum dose of 100 mg/day; and
- 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

| ERLOTINIB - Retai | pharmacy-Specia | list - Special Authority | y see SA2115 below |
|-------------------|-----------------|--------------------------|--------------------|
|-------------------|-----------------|--------------------------|--------------------|

| Tab 100 mg | 30 | Alchemy |
|------------|--------|-----------------------------|
| Tab 150 mg | 30 | Alchemy |

➡SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

| | Subsidy (Manufacturer's Price) \$ | Sub Per | Fully osidised | Brand or Generic Manufacturer |
|--|---|-------------|-------------------|-------------------------------------|
| continued | | | | |
| All of the following: 1 The patient is clinically benefiting from treatment and con 2 Erlotinib to be discontinued at progression; and 3 The regular Special Authority renewal requirements cann | | | | |
| GEFITINIB – Retail pharmacy-Specialist – Special Authority see Tab 250 mg | | 30 | ✓ II | ressa |
| ► SA2116 Special Authority for Subsidy Initial application only from a relevant specialist or medical prace Approvals valid for 4 months for applications meeting the following All of the following: | ng criteria: | | | · |
| Patient has locally advanced, or metastatic, unresectable Either: Patient is treatment naive; or Both: | | Small Ce | aii Lung (| Lancer (NSULU); and |
| 2.2.1 The patient has discontinued erlotinib due 2.2.2 The cancer did not progress whilst on erlot | inib; and | | | |
| 3 There is documentation confirming that disease expresse4 Gefitinib is to be given for a maximum of 3 months. | es activating mutations | s of EGFI | R tyrosin | e kinase; and |
| Renewal only from a relevant specialist or medical practitioner of for 6 months where radiological assessment (preferably including Renewal — (pandemic circumstances) from any relevant prac- the following criteria: All of the following: | g CT scan) indicates N | SCLC h | as not p | rogressed. |
| The patient is clinically benefiting from treatment and con Gefitinib to be discontinued at progression; and The regular Special Authority renewal requirements cann | | | • | |
| IMATINIB MESILATE | | | | |
| * Cap 100 mg * Cap 400 mg | | 60 30 | | <u>matinib-Rex</u> matinib-Rex |
| NILOTINIB – Special Authority see SA2301 below – Retail phar Wastage claimable | macy | | | |
| Cap 150 mg Cap 200 mg | | 120 120 | | asigna asigna |
| ► SA2301 Special Authority for Subsidy Initial application only from a haematologist. Approvals valid for All of the following: | or 6 months for applica | ations me | eting the | e following criteria: |
| Patient has a diagnosis of chronic myeloid leukaemia (CM and Either: | /IL) in blast crisis, high | n risk chro | onic pha | se, or in chronic phase; |
| 2.1 Patient has documented CML treatment failure* w2.2 Patient has experienced treatment limiting toxicity and | | | | ecluding further treatment; |
| 3 Maximum nilotinib dose of 800 mg/day; and4 Subsidised for use as monotherapy only. | | | | |
| Note: *treatment failure as defined by Leukaemia Net Guideline | S. | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|-------------------|---------|-----------|--------------|--|
| (Manufacturer's P | rice) S | ubsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

Wastage claimable

| Tab 75 mg4,000.00 | 21 | Ibrance |
|--------------------|----|-----------------------------|
| Tab 100 mg4,000.00 | 21 | Ibrance |
| Tab 125 mg4,000.00 | 21 | Ibrance |

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

| Tab 200 mg | 1,334.70 | 30 | Votrient |
|------------|----------|----|------------------------------|
| Tab 400 mg | 2,669.40 | 30 | Votrient |

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage claimable

| Tab 5 mg2,500.00 | 56 | 🖌 Jakavi |
|------------------|----|----------|
| Tab 10mg5,000.00 | 56 | 🖌 Jakavi |
| Tab 15 mg | 56 | 🖌 Jakavi |
| Tab 20 mg | 56 | 🗸 Jakavi |

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price | e) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 The treatment remains appropriate and the patient is benefiting from treatment; and

2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2117 below - Retail pharmacy

| Cap 12.5 mg | 28 | Sunitinib Pfizer |
|-------------|--------|--------------------------------------|
| Cap 25 mg | 28 | Sunitinib Pfizer |
| Cap 50 mg | 28 | Sunitinib Pfizer |

⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- Both:
 - 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and the patient is benefiting from treatment.

| | Subsidy | Fu | lly | Brand or |
|-----|---------------------|----------|-----|--------------|
| (Ma | nufacturer's Price) | Subsidis | ed | Generic |
| | \$ | Per | / | Manufacturer |

continued...

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 89

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

➡SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|--------------------|------|------------|--------------|--|
| (Manufacturer's Pr | ice) | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE

| Tab 50 mg | 4.18 | 28 | ✓ Apo- Bicalutamide S23 |
|--|-----------------|-----|------------------------------|
| | | | ✓ Binarex |
| FLUTAMIDE | | | |
| Tab 250 mg | 107.55 | 90 | Prostacur S29 |
| ů – | 119.50 | 100 | Flutamin |
| FULVESTRANT - Retail pharmacy-Specialist - Special Authority | see SA1895 belo | w | |
| Inj 50 mg per ml, 5 ml prefilled syringe | | 2 | Faslodex |

➡SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

| () | Subsidy /anufacturer's Price) \$ | Sul Per | Fully osidised | Brand or Generic Manufacturer |
|--|--|------------|-------------------|-------------------------------------|
| OCTREOTIDE | | | | |
| Inj 50 mcg per ml, 1 ml ampoule | 27.58 | 5 | ✓ <u>N</u> | lax Health |
| | | | ✓ 0 | Octreotide GH S29 |
| Inj 100 mcg per ml, 1 ml ampoule | 32.71 | 5 | 🗸 <u>N</u> | lax Health |
| | | | ✓ 0 | Ctreotide GH S29 |
| | | | ✓ s | Sun Pharma S29 |
| Inj 500 mcg per ml, 1 ml ampoule | 113.10 | 5 | 🗸 N | lax Health |
| | | | ✓ 0 | Ctreotide GH S29 |
| | | | ✓ s | un Pharma S29 |
| OCTREOTIDE LONG-ACTING - Special Authority see SA2119 be | low – Retail pharma | acv | | |
| Inj depot 10 mg prefilled syringe | | 1 | ✓ <u>0</u> | <u>)ctreotide Depot</u> Teva |
| | 1,152.00 | | ✓ s | andostatin LAR |
| Inj depot 20 mg prefilled syringe | 647.03 | 1 | ✓ 0 | Ctreotide Depot |
| | | | _ | Teva |
| | 1,539.00 | | 🗸 S | andostatin LAR |
| Inj depot 30 mg prefilled syringe | 718.55 | 1 | ✓ 0 | Octreotide Depot |
| | | | | Teva |

➡SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price | e) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or

3 Both:

- 3.1 Insulinomas; and
- 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or

5 Both:

- 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
- 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

TAMOXIFEN CITRATE

| * Tab 10 mg | 60 60 | ✓ <u>Tamoxifen Sandoz</u> ✓ <u>Tamoxifen Sandoz</u> |
|-------------------------------|----------|--|
| Aromatase Inhibitors | | |
| ANASTROZOLE * Tab 1 mg4.39 | 30 | ✓ <u>Anatrole</u> |
| EXEMESTANE * Tab 25 mg | 30 | ✓ Pfizer Exemestane |
| LETROZOLE * Tab 2.5 mg5.84 | 30 | ✓ Letrole |

| (| Subsidy Manufacturer's Price \$ |) Subs Per | Fully sidised | Brand or Generic Manufacturer |
|--|---------------------------------------|---------------|------------------|-------------------------------------|
| Immunosuppressants | | | | |
| Cytotoxic Immunosuppressants | | | | |
| AZATHIOPRINE | | | | |
| * Tab 25 mg | 7.36 | 60 | 1 | Azamun |
| * Tab 50 mg | 8.10 | 100 | 1 | Azamun |
| MYCOPHENOLATE MOFETIL | | | | |
| Tab 500 mg | 35.90 | 50 | ✓ (| Cellcept |
| Cap 250 mg | | 100 | ✓ (| Cellcept |
| Powder for oral lig 1 g per 5 ml - Subsidy by endorsement | 187.25 1 | 65 ml OP | ✓ (| Cellcept |
| Mycophenolate powder for oral liquid is subsidised only for the prescription is endorsed accordingly. | r patients unable to | o swallow ta | ablets | and capsules, and when |

Fusion Proteins

| ETANERCEPT - Special Authority see SA2103 below - R | etail pharmacy | | |
|---|----------------|---|----------------------------|
| Inj 25 mg | | 4 | Enbrel |
| Inj 25 mg autoinjector | | 4 | Enbrel |
| Inj 50 mg autoinjector | 1,050.00 | 4 | Enbrel |
| Inj 50 mg prefilled syringe | 1,050.00 | 4 | Enbrel |

⇒SA2103 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either: 1 Both:

- 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Ful | y Brand or | |
|------------------------|-----------|--------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per • | Manufacturer | |

continued...

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

- 18-24 years Male: 7.0 cm; Female: 5.5 cm
- 25-34 years Male: 7.5 cm; Female: 5.5 cm
- 35-44 years Male: 6.5 cm; Female: 4.5 cm
- 45-54 years Male: 6.0 cm; Female: 5.0 cm
- 55-64 years Male: 5.5 cm; Female: 4.0 cm
- 65-74 years Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application - (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist.

| S | Subsidy | Fully | Brand or |
|---------|-------------------------|-------|--------------|
| (Manufa | acturer's Price) Subsid | lised | Generic |
| | \$ Per | 1 | Manufacturer |

continued...

Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both:

- The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

2 Either:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and

1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects; or
- 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Dott

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Renewal — (severe chronic plaque psoriasis)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.
- Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
 - 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
 - 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.
- Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

| (M | Subsidy anufacturer's Price) | Subs | Fully idised | Brand or Generic |
|----|---------------------------------|------|-----------------|---------------------|
| · | \$ | Per | ~ | Manufacturer |

continued...

- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

| ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only – Specialist | | |
|--|---|---------------------------------------|
| Inj 50 mg per ml, 5 ml2,774.48 | 5 | 🗸 ATGAM |
| BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only - Specialist | | |
| Subsidised only for bladder cancer. | | |
| Inj 2-8 × 100 million CFU149.37 | 1 | OncoTICE |
| Inj 40 mg per ml, vial176.90 | 3 | SII-Onco-BCG \$29 |

Monoclonal Antibodies

| ADALIMUMAB (AMGEVITA) - Special Authority see SA2178 b | elow – Retail pharn | nacy | |
|--|---------------------|------|------------------------------|
| Inj 20 mg per 0.4 ml prefilled syringe | | 1 | Amgevita |
| Inj 40 mg per 0.8 ml prefilled pen | | 2 | Amgevita |
| Inj 40 mg per 0.8 ml prefilled syringe | | 2 | Amgevita |

► SA2178 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:

2.1 Either:

- 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
 - 1 Patient has pyoderma gangrenosum*; and
 - 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| Subsidy | | Fully | Brand or | |
|---------------------|-----|------------|--------------|--|
| (Manufacturer's Pri | ce) | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
- 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
- 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
- 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and

3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

| | Subsidy | F | ully | Brand or |
|-----|----------------------|--------|------|--------------|
| (Ma | anufacturer's Price) | Subsid | ised | Generic |
| | \$ | Per | ✓ | Manufacturer |

continued...

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

1 Both:

1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | S | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|----------|--------------|--|
| (Manufacturer's Price) | Sub | bsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects; or

1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or

- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 Both:

| Subsidy | Fully | Brand or | |
|-----------------|-------------------|--------------|--|
| (Manufacturer's | Price) Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 below - Retail pharmacy

| Inj 20 mg per 0.2 ml prefilled syringe1,599.96 | 2 | 🗸 Humira |
|--|---|-------------------------------|
| Inj 40 mg per 0.4 ml prefilled pen1,599.96 | 2 | HumiraPen |
| Inj 40 mg per 0.4 ml prefilled syringe1,599.96 | 2 | Humira |

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

All of the followina:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amaevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease - severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- - 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline: and
 - 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and

3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered. Initial application - (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the followina:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amoevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita: and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither

| Subsidy | Ful | y Brand or | |
|------------------------|-----------|--------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

- 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
 - 2 Patient has received a maximum of 6 months treatment with Amgevita; and
 - 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
 - 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Any of the following:

- 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
- 1.2 CDAI score is 150 or less; or
- 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Fu | lly Brand or | |
|------------------------|----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidis | ed Generic | |
| \$ | Per | Manufacturer | |

continued...

- 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.
- Renewal (Ocular inflammation chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

| Subsidy | | Fully | Brand or | |
|------------------------|-----|----------|--------------|--|
| (Manufacturer's Price) | Su | bsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
 - 2 Patient has received a maximum of 6 months treatment with Amgevita; and
 - 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

| Subsidy | | Fully | Brand or | |
|---------------------|---------|-----------|--------------|--|
| (Manufacturer's Pri | ice) Si | ubsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT – Special Authority see SA1772 below – Retail pharmacy

Inj 40 mg per ml, 0.1 ml vial......1,250.00 1 🖌 Eylea

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Su | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 on the next page - Retail pharmacy

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price |) | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10°9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

- Both:
 - 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
 - 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

BRENTUXIMAB VEDOTIN – PCT only – Special Authority see SA2289 below

| Inj 50 mg vial | 5,275.18 | 1 | Adcetris |
|----------------|----------|---|----------|
|----------------|----------|---|----------|

⇒SA2289 Special Authority for Subsidy

Initial application — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and

| Sub | bsidy F | ully | Brand or |
|------------|----------------------|------|--------------|
| (Manufacti | urer's Price) Subsid | sed | Generic |
| | \$ Per | 1 | Manufacturer |

continued...

- 1.1.2 Patient is ineligible for autologous stem cell transplant; or
- 1.2 Both:
 - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
 - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

Initial application — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

- Inj 120 mg per ml casirivimab, 11.1 ml vial (1) and inj 120 mg
 - per ml imdevimab, 11.1 ml vial (1)......0.00 1 OP 🖌 Ronapreve

⇒SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

| lnj 5 mg per ml, | 20 ml vial | | 364.00 | 1 | Erbitux |
|------------------|-------------|------|----------|------|-----------------------------|
| lni 5 ma per ml. | 100 ml vial | | 1.820.00 | 1 | Erbitux |
| , , , | | | , | 1 mg | Baxter |
| , , | | | | • | |

| | Subsidy (Manufacturer's Price) \$ | | Illy Brand or ed Generic Manufacturer |
|--|--|--|--|
| SA1697 Special Authority for Subsidy Initial application only from a medical oncologist or medical p Approvals valid for 6 months for applications meeting the follow All of the following: Patient has locally advanced, non-metastatic, squamou Patient is contraindicated to, or is intolerant of, cisplatin Patient has good performance status; and To be administered in combination with radiation therap | ving criteria: s cell cancer of the hea ; and | | |
| GEMTUZUMAB OZOGAMICIN – PCT only – Specialist – Spe Inj 5 mg vial | cial Authority see SA22 | | ✓ Mylotarg |
| SA2269 Special Authority for Subsidy Initial application only from a haematologist, paediatric haema applications meeting the following criteria: All of the following: Patient has not received prior chemotherapy for this cor Patient has de novo CD33-positive acute myeloid leuka Patient does not have acute promyelocytic leukaemia; a Gemtuzumab ozogamicin will be used in combination w Patient's disease risk has been assessed by cytogeneti Patient must be considered eligible for standard intensity and cytarabine (AraC); and Gemtuzumab ozogamicin to be funded for one course of 5 mg as separate doses). Note: Acute myeloid leukaemia excludes acute promyelocytic another haematological disorder (eg myelodysplasia or myelop | ndition; and emia; and and ith standard anthracycl c testing to be good or ve remission induction of only (one dose at 3 mg leukaemia and acute m | ine and cytara intermediate; chemotherapy per m ² body s | abine (AraC); and and with standard anthracycline urface area or up to 2 vials of |
| INFLIXIMAB – PCT only – Special Authority see SA2179 below Inj 100 mg | w 428.00 | | Remicade |
| Inj 1 mg for ECP | levant practitioner. App al to 300 or HBI score of ecting more than 50 cm | provals valid for of greater than of the small i | or equal to 10; or intestine; or |

- 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 Any of the following:

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | 9 | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
- 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
- 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

- Both:
 - 1 Patient has acute, fulminant ulcerative colitis; and
 - 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

| Subsidy | F | ully | Brand or |
|-----------------------|----------|------|--------------|
| (Manufacturer's Price |) Subsid | ised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patent has complex peri-anal fistula.

Renewal - (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | F | ully | Brand or | Ī |
|------------------------|---------|------|--------------|---|
| (Manufacturer's Price) | Subsidi | sed | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

- Both:
 - 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
 - 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria: Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Either:

itner:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and

| (Mai | Subsidy | F | ully | Brand or |
|------|---------------------|----------|------|--------------|
| | nufacturer's Price) | Subsidis | sed | Generic |
| | \$ | Per | ✓ | Manufacturer |

continued...

- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Renewal — (plaque psoriasis)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or

1.2 Both:

- 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation; or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis; or

continued...

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | F | ully | Brand or |
|------------------------|--------|------|--------------|
| (Manufacturer's Price) | Subsid | ised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 2.12 Neurosarcoidosis; or
- 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
- 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and

1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or

2 Both:

2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and

2.2 Any of the following:

- 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
- 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
- 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

1 The patient has had a good clinical response following 3 initial doses; or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Ful | ly Brand or | |
|-------------------|------------------|----------------------------------|--|
| (Manufacturer's F | Price) Subsidise | d Generic | |
| \$ | Per | Manufacturer | |

continued...

- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patients SCCAI is greater than or equal to 4; or
 - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and

| Subsidy | Ful | y Brand or |
|----------------------|--------------|--------------|
| (Manufacturer's Pric | e) Subsidise | d Generic |
| \$ | Per 🔹 | Manufacturer |

continued...

- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment .

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

MEPOLIZUMAB - Special Authority see SA2331 below - Retail pharmacy

| Inj 100 mg prefilled pen | 1,638.00 | 1 | Nucala |
|--------------------------|----------|---|----------------------------|
| Inj 100 mg vial | 1,638.00 | 1 | Nucala |
| | | | |

(Nucala Inj 100 mg vial to be delisted 1 August 2024)

⇒SA2331 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than $0.5 \times 10^{\circ}9$ cells/L in the last 12 months; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | F | ully | Brand or |
|-------|-------------------|---------|------|--------------|
| (Manu | facturer's Price) | Subsidi | sed | Generic |
| | \$ | Per | ✓ | Manufacturer |

- continued...
 - 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
 - 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
 - 7 Treatment is not to be used in combination with subsidised benralizumab; and
 - 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
 - 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initial application — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has eosinophilic granulomatosis with polyangiitis; and
- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and
- 3 Either:
 - 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
 - 3.2 Corticosteroids are contraindicated.

Renewal — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where patient has no evidence of clinical disease progression.

| OBINUTUZUMAB - PCT only - Specialist - Special Authority | see SA2155 on the | next page | |
|--|-------------------|-----------|----------------------------|
| Inj 25 mg per ml, 40 ml vial | 5,910.00 | 1 | 🖌 Gazyva |
| Inj 1 mg for ECP | 6.21 | 1 mg | Baxter |

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | I Generic | |
| \$ | Per 🗸 | Manufacturer | |

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^{9} /L and platelets greater than or equal to 75×10^{9} /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

| Inj 150 mg prefilled syringe | 450.00 | 1 | 🗸 Xolair |
|------------------------------|--------|---|----------------------------|
| Inj 150 mg vial | 450.00 | 1 | Xolair |

► SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and

continued...

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and

2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient has previously adequately responded* to 6 doses of omalizumab; or

2 Both:

- 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
- 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

| | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully idised | Brand or Generic Manufacturer | |
|---|---|-------------|-----------------|-------------------------------------|--|
| PERTUZUMAB – PCT only – Specialist – Special Authority see SA2276 below | | | | | |
| Inj 30 mg per ml, 14 ml vial | 3,927.00 | 1 | ✓ P | erjeta | |
| Ini 420 mg for ECP | 3.927.00 42 | 0 ma OP | 🗸 В | axter | |

► SA2276 Special Authority for Subsidy

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 below

| Inj 100 mg per 10 ml vial | | 2 | Mabthera |
|---------------------------|----------|------|------------------------------|
| Inj 500 mg per 50 ml vial | 2,688.30 | 1 | Mabthera |
| Inj 1 mg for ECP | 5.64 | 1 mg | 🗸 Baxter (Mabthera) |

► SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | 9 | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- the maximum tolerated dose of ciclosporin; or
- 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

6 Either:

- 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
- 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

7 Either:

- 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and

8 Either:

- 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and 9 Maximum of two 1.000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

3 Maximum of two 1,000 mg infusions of ntuximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and

3 Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2233 below

| Inj 100 mg per 10 ml vial | | 2 | Riximyo |
|---------------------------|--------|------|--------------------------------------|
| Inj 500 mg per 50 ml vial | 688.20 | 1 | Riximyo |
| Inj 1 mg for ECP | 1.38 | 1 mg | Baxter (Riximyo) |

► SA2233 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of

| Subsid | ly Fu | ully Brand or |
|---------------|----------------------|----------------------------------|
| (Manufacturer | r's Price) Subsidise | ed Generic |
| \$ | Per | Manufacturer |

continued...

4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*. Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or

2.2.2 Both:

- 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
- 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
- 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and

4 Either:

- 4.1 The patient does not have chromosome 17p deletion CLL; or
- 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2. **Renewal — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
- 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and

| Subsidy | | Fully | Brand or |
|------------------------|------|-----------|--------------|
| (Manufacturer's Price) |) Si | ubsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.
- Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

- Both:
 - 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and

2 Either:

2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or

2.2 Both:

2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

- All of the following:
 - 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
 - 2 An initial response lasting at least 12 months was demonstrated; and
 - 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS))

only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only

from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or

2 Both:

- 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
- 2.2 To be used for a maximum of 6 treatment cycles.
- Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.
- Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

1 Either:

1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) |) 5 | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.
- Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia*

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) |) 5 | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.
- Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.
- Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.
- Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

| Subsidy | | Fully | Brand or | _ |
|------------------------|-----|-----------|--------------|---|
| (Manufacturer's Price) | Su | Ibsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
- 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has

| Subsidy | F | ully | Brand or |
|------------------------|---------|------|--------------|
| (Manufacturer's Price) | Subsidi | sed | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

experienced intolerable side effects.

d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD*; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or

| Subsidy | | Fully | Brand or | |
|--------------------|-------|------------|--------------|--|
| (Manufacturer's Pr | rice) | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and

3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart. Note: Indications marked with * are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
 - 1.2 Patient is receiving maintenance treatment for IgG4-RD*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and

3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

| | Special Authority | 1 coo \$12094 b | elow – Retail pharmacy | |
|-------------|---------------------|-----------------|------------------------|--|
| SECUNINUMAD | - Special Authonity | / see SA2084 D | elow – Retail pharmacy | |

| | | ······································ | |
|------------------------------|---|--|---|
| Cosentyx | 1 | prefilled syringe799.50 | Inj 150 mg per ml, 1 ml prefilled syringe |
| Cosentyx | 2 | 1,599.00 | |

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole

| Subsidy | Fu | Illy Brand or | |
|-----------------|-----------------|----------------------------------|--|
| (Manufacturer's | Price) Subsidis | ed Generic | |
| \$ | Per | Manufacturer | |

continued...

body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
- 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and

- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:

| Subsidy | | Fully | Brand or |
|-----------------|--------|------------|--------------|
| (Manufacturer's | Price) | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
- 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB – Special Authority see SA1596 below – Retail pharmacy

| Note: Siltuximab is to be administered at doses no gre | ater than 11 mg/kg every 3 | 3 weeks. | |
|--|----------------------------|----------|-----------------------------|
| Inj 100 mg vial | 770.57 | 1 | Sylvant |
| Inj 400 mg vial | | 1 | Sylvant |

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.
- Inj 100 mg per ml, 1.5 ml vial with cilgavimab 100 mg per

| ml,1.5 ml vial | 0.00 | 1 | Evusheld |
|---|------------------------|------|-----------------------------|
| TOCILIZUMAB – PCT only – Special Authority see SA | A2332 on the next page | | |
| Inj 20 mg per ml, 4 ml vial | | 1 | Actemra |
| Inj 20 mg per ml, 10 ml vial | | 1 | Actemra |
| Inj 20 mg per ml, 20 ml vial | 1,100.00 | 1 | Actemra |
| Inj 1 mg for ECP | | 1 mg | Baxter |
| , , | | Ũ | |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

⇒SA2332 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
 - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

| Subsidy | y Fi | ully Brand or | |
|---------------|-------------------|----------------------------------|--|
| (Manufacturer | s Price) Subsidis | ised Generic | |
| \$ | Per | Manufacturer | |

continued...

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

| Subsidy | Full | / Brand or | _ |
|------------------------|-----------|--------------|---|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a

| | Subsidy | Fully | Brand or |
|-------|-------------------|------------|--------------|
| (Manu | facturer's Price) | Subsidised | Generic |
| | \$ Pe | er 🖌 | Manufacturer |

continued...

rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB (HERCEPTIN) - PCT only - Specialist - Special Authority see SA2287 below

| Inj 150 mg vial | , , | | 1 | Herceptin |
|------------------|--------|----------|------|-------------------------------|
| Inj 440 mg vial | | | 1 | ✓ Herceptin |
| Inj 1 mg for ECP | | 9.36 | 1 mg | Baxter |
| | | | | |

(Herceptin Inj 150 mg vial to be delisted 1 June 2024) (Herceptin Inj 440 mg vial to be delisted 1 June 2024) (Baxter Inj 1 mg for ECP to be delisted 1 June 2024)

⇒SA2287 Special Authority for Subsidy

Renewal — (early breast cancer*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerable side effects; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and

4 Either:

- 4.1 Trastuzumab will not be given in combination with pertuzumab; or
- 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting

| Subsi | dy Fully E | Brand or |
|--------------|------------|--------------|
| (Manufacture | , | Generic |
| \$ | Per 🖌 I | Manufacturer |
| | | |

continued...

the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB (HERZUMA) – PCT only – Special Authority see SA2293 below

| Inj 150 mg vial | 100.00 | 1 | 🗸 Herzuma |
|---|--------|------|----------------------------|
| Herzuma to be Principal Supply on 1 June 2024 | | | |
| Inj 440 mg vial | 293.35 | 1 | 🗸 Herzuma |
| Herzuma to be Principal Supply on 1 June 2024 | | | |
| Inj 1 mg for ECP | 0.70 | 1 mg | Baxter |
| | | • | |

⇒SA2293 Special Authority for Subsidy

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Renewal — (early breast cancer*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
 - 1.3 Any of the following:
 - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
 - 1.3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.4 Either:
 - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 1.4.2 All of the following:
 - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
 - 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

| Subsidy | F | Fully | Brand or | |
|------------------------|--------|-------|--------------|--|
| (Manufacturer's Price) | Subsid | lised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initial application - (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an ECOG score of 0-2.

Renewal — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2144 on the next page

| Inj 100 mg vial | ····· | 2,320.00 | 1 | 🖌 Kadcyla |
|------------------|-------|--------------|------|----------------------------|
| Inj 160 mg vial | | | 1 | 🗸 Kadcyla |
| Inj 1 mg for ECP | | 24.52 | 1 mg | Baxter |

| Subsidy | | Fully | Brand or |
|------------------------|-----|---------|--------------|
| (Manufacturer's Price) | | sidised | Generic |
| \$ | Per | 1 | Manufacturer |

⇒SA2144 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
 - 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
 - 4 Patient has a good performance status (ECOG 0-1); and
 - 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
 - 6 Patient has not received prior funded trastuzumab emtansine treatment; and
 - 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB – Special Authority see SA2182 below – Retail pharmacy

⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|--------------------|-----|------------|--------------|--|
| (Manufacturer's Pr | ce) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

intolerable side effects or insufficient benefit to meet renewal criteria; or

2.2.2 Both:

2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and

2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or

2 Both:

- 2.1 Patient has active Crohn's disease; and
- 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and

2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or

2 Both:

- 2.1 Patient has active ulcerative colitis; and
- 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

2.2.2 Both:

2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or

1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and

2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

► SA2183 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and

3 Any of the following:

- 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
- 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
- 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

1 Any of the following:

- 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
- 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
- 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
- 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
- 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or

1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Programmed Cell Death-1 (PD-1) Inhibitors

| ATEZOLIZUMAB - PCT only - Specialist - Special Authority see SA2264 on the next page | | | | |
|--|------|------|-------------------------------|--|
| Inj 60 mg per ml, 20 ml vial | | 1 | Tecentriq | |
| Inj 1 mg for ECP | 8.08 | 1 mg | Baxter | |

| | Subsidy | Fully | Brand or |
|------|---------------------|------------|--------------|
| (Mar | nufacturer's Price) | Subsidised | Generic |
| | \$ P | Per 🗸 | Manufacturer |

► SA2264 Special Authority for Subsidy

Initial application — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

DURVALUMAB - PCT only - Specialist - Special Authority see SA2164 below

| Inj 50 mg per ml, 10 ml vial | 4,700.00 | 1 | 🖌 Imfinzi |
|-------------------------------|----------|------|----------------------------|
| Inj 50 mg per ml, 2.4 ml vial | 1,128.00 | 1 | 🗸 Imfinzi |
| Inj 1 mg for ECP | 9.59 | 1 mg | Baxter |

➡SA2164 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and

7 Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
- 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

NIVOLUMAB - PCT only - Specialist - Special Authority see SA2306 below

| Inj 10 mg per ml, 4 ml vial1,0 | 51.98 1 | Opdivo |
|---------------------------------|------------|----------------------------|
| Inj 10 mg per ml, 10 ml vial2,6 | 29.96 1 | Opdivo |
| Inj 1 mg for ECP | 27.62 1 mg | ✓ Baxter |

⇒SA2306 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal — (less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment; or
 - 1.1.2 Patient's disease has had a partial response to treatment; or
 - 1.1.3 Patient has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
 - 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

Renewal — (more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Any of the following:
 - 2.1.1.1 Patient's disease has had a complete response to treatment; or
 - 2.1.1.2 Patient's disease has had a partial response to treatment; or
 - 2.1.1.3 Patient has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
 - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2.2 All of the following:
 - 2.2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 Patient has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA2307 below

| Inj 25 mg per ml, 4 ml vial | 4,680.00 | 1 | 🗸 Keytruda |
|-----------------------------|----------|------|----------------------------|
| Inj 1 mg for ECP | 47.74 | 1 mg | Baxter |

► SA2307 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment; or
 - 1.1.2 Patient's disease has had a partial response to treatment; or
 - 1.1.3 Patient has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the

| Subsidy | | Fully | Brand or | |
|---------------------|-----|------------|--------------|--|
| (Manufacturer's Pri | ce) | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

most recent treatment period; and

- 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Any of the following:
 - 2.1.1.1 Patient's disease has had a complete response to treatment; or
 - 2.1.1.2 Patient's disease has had a partial response to treatment; or
 - 2.1.1.3 Patient has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
 - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2.2 All of the following:
 - 2.2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 Patient has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Either:
 - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
 - 6.2 Both:
 - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
 - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2; and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

| Subsidy | | Fully | Brand or |
|------------------------|------|-----------|--------------|
| (Manufacturer's Price) |) Si | ubsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (non-small cell lung cancer first-line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2; and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Other Immunosuppressants

| CICL | OSPORIN | |
|------|----------------|--|
| | | |

| Cap 25 mg | 50 | Neoral |
|------------------------------|----------|----------------------------|
| Cap 50 mg | 50 | Neoral |
| Cap 100 mg | 50 | Neoral |
| Oral liq 100 mg per ml198.13 | 50 ml OP | Neoral |

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|----------|---------------------|-------------------------------------|
| EVEROLIMUS – Special Authority see SA2008 below – Retail p Wastage claimable | harmacy | | | |
| Tab 10 mg Tab 5 mg | | 30 30 | | Afinitor Afinitor |

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

SIROLIMUS - Special Authority see SA2270 below - Retail pharmacy

| Tab 1 mg | | 100 | Rapamune |
|----------------------|----------|----------|------------------------------|
| Tab 2 mg | 1,499.99 | 100 | Rapamune |
| Oral liq 1 mg per ml | | 60 ml OP | Rapamune |

⇒SA2270 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- · Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- · HUS or TTP; or
- · Leukoencepthalopathy; or
- Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly

| | Subsidy | Fully | Brand or |
|-------|--------------------|------------|--------------|
| (Manu | ufacturer's Price) | Subsidised | Generic |
| | \$ Per | ✓ | Manufacturer |

continued...

documents in patient notes; and

- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

| TACROLIMUS - Special Authority see SA2271 on the next | t page – Retail pharmad | су | |
|---|-------------------------|-----|---------------------------------------|
| Cap 0.5 mg | | 100 | Tacrolimus Sandoz |
| Cap 0.75 mg | | 100 | Tacrolimus Sandoz |
| Cap 1 mg | | 100 | Tacrolimus Sandoz |
| Cap 5 mg | | 50 | Tacrolimus Sandoz |

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Fu Subsidis Per | ully ed ✔ | Brand or Generic Manufacturer |
|--|---|-----------------------|-----------------|-------------------------------------|
| ■SA2271 Special Authority for Subsidy Initial application — (organ transplant) only from a relevant si where the patient is an organ transplant recipient. Note: Subsidy applies for either primary or rescue therapy. Initial application — (non-transplant indications*) only from a unless notified for applications meeting the following criteria: | | | | |
| Both: 1 Patient requires long-term systemic immunosuppression; 2 Either: | and | | | |
| 2.1 Ciclosporin has been trialled and discontinued trea clinical response; or 2.2 Patient is a child with nephrotic syndrome*. | atment because of una | acceptable si | de eff | iects or inadequate |
| Note: Indications marked with * are unapproved indications | | | | |

JAK inhibitors

UPADACITINIB – Special Authority see SA2079 below – Retail pharmacy Tab 15 mg1,271.00

RINVOQ

28

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer \$ Antiallergy Preparations Allergic Emergencies ADRENALINE - Special Authority see SA2185 below - Retail pharmacy a) Maximum of 2 ini per prescription b) Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis. 1 OP Epipen Jr Inj 0.3 mg per 0.3 ml auto-injector......90.00 1 OP Epipen ⇒SA2185 Special Authority for Subsidy Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both: 1 Either: 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department: or 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and 2 Patient is not to be prescribed more than two devices in initial prescription. ICATIBANT - Special Authority see SA1558 below - Retail pharmacy Ini 10 ma per ml. 3 ml prefilled svringe......2.668.00 1 Firazvr SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of larvngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

■ SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy

| 05.00 | 1 OP | VENOX S29 |
|-------|-------------------------|--|
| 05.00 | 1 OP | VENOX S29 |
| | | |
| 85.00 | 1 OP | Venomil S29 |
| | | |
| 05.00 | 1 OP | Albey |
| 05.00 | 1 OP | ✓ Hymenoptera S29 |
| | 05.00 85.00 05.00 | 05.00 1 OP 85.00 1 OP 05.00 1 OP |

| | Subsidy | | Fully | |
|--|-----------------------|---------|-----------------------|---------------------|
| | (Manufacturer's Price | | Subsidised | |
| | \$ | Per | | Manufacturer |
| WASP VENOM ALLERGY TREATMENT - Special Authority see | e SA1367 on the pre | vious p | <mark>age</mark> – Re | tail pharmacy |
| Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze | | | | |
| dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml | | 1 OP | 1 | Albey |
| Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze | | | | • |
| dried venom, with diluent | | 1 OP | 1 | Hymenoptera S29 |
| Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze | | - | | , |
| dried venom, with diluent | | 1 OP | 1 | Venomil S29 |
| Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze | | | | |
| dried venom, with diluent | 305.00 | 1 OP | 1 | Hymenoptera S29 |
| Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze | | 1 01 | | nymeneptera 😅 |
| dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml | 305.00 | 1 OP | 1 | Albey |
| Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze | | 101 | • | Albey |
| | | 1 OP | | Venomil S29 |
| dried venom, with diluent | | 100 | - | VEHOIIIII |
| Antihistamines | | | | |
| Antimistanimes | | | | |
| CETIRIZINE HYDROCHLORIDE | | | | |
| * Tab 10 mg | 1.71 | 100 | 1 | Zista |
| * Oral lig 1 mg per ml | 2.84 | 200 ml | 1 | Histaclear |
| | | | | |
| * Oral liq 2 mg per 5 ml | 9.37 | 500 ml | 1 | Histafen |
| (Histafen Oral liq 2 mg per 5 ml to be delisted 1 August 2024) | | 000 111 | | Inotation |
| | | | | |
| DEXTROCHLORPHENIRAMINE MALEATE * Tab 2 mg | 0.00 | 40 | | |
| * Tab 2 mg | | 40 | | Polaramine |
| | (8.40) 1.01 | 20 | | Foldramme |
| | (5.99) | 20 | | Polaramine |
| * Oral liq 2 mg per 5 ml | | 100 ml | | FUIdIdITIITE |
| | (10.29) | 100 111 | | Polaramine |
| | (10.23) | | | 1 Ularamine |
| FEXOFENADINE HYDROCHLORIDE | | | | |
| * Tab 60 mg | | 20 | | - <i>v</i> . |
| | (8.23) | | | Telfast |
| * Tab 120 mg | | 10 | | T - 16 1 |
| | (8.23) | 00 | | Telfast |
| | 14.22 | 30 | | Tolfoot |
| | (26.44) | | | Telfast |
| LORATADINE | | | - | |
| * Tab 10 mg | | 100 | | Lorafix |
| * Oral liq 1 mg per ml | 1.43 | 100 ml | 1 | Haylor syrup |
| PROMETHAZINE HYDROCHLORIDE | | | | |
| * Tab 10 mg | 1.39 | 50 | ✓ | Allersoothe |
| * Tab 25 mg | 1.58 | 50 | 1 | Allersoothe |
| * Oral liq 1 mg per 1 ml | 3.39 | 100 ml | 1 | Allersoothe |
| Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a F | 00 01 00 | 5 | 1 | Hospira |

| | Subsidy | | Fully | |
|--|----------------|----------------------|------------|---------------------|
| () | Manufacturer's | Price) Per | Subsidised | |
| | \$ | rei | | Manulaciulei |
| Inhaled Corticosteroids | | | | |
| BECLOMETHASONE DIPROPIONATE | | | | |
| Aerosol inhaler, 50 mcg per dose | 14.01 | 200 dose | | Qvar |
| Aerosol inhaler, 50 mcg per dose CFC-free | | 200 dose 200 dose | | Beclazone 50 |
| Aerosol inhaler, 100 mcg per dose ch c-nee | | 200 dose 200 dose | • | Qvar |
| Aerosol inhaler, 100 mcg per dose CFC-free | | 200 dose | • | Beclazone 100 |
| | | 200 dose 200 dose | • | Beclazone 250 |
| Aerosol inhaler, 250 mcg per dose CFC-free | 22.07 | 200 0056 | UP V | Deciazone 200 |
| BUDESONIDE | | | | |
| Powder for inhalation, 100 mcg per dose | 17.00 | 200 dose | OP 🗸 | Pulmicort |
| | | | | Turbuhaler |
| Powder for inhalation, 200 mcg per dose | 19.00 | 200 dose | OP 🗸 | Pulmicort |
| | | | | Turbuhaler |
| Powder for inhalation, 400 mcg per dose | 32.00 | 200 dose | OP 🗸 | Pulmicort |
| | | | | Turbuhaler |
| FLUTICASONE | | | | |
| Aerosol inhaler, 50 mcg per dose | 7.19 | 120 dose | OP 🗸 | Flixotide |
| Powder for inhalation, 50 mcg per dose | 8.61 | 60 dose | OP 🗸 | Flixotide Accuhaler |
| Powder for inhalation, 100 mcg per dose | | 60 dose | OP 🗸 | Flixotide Accuhaler |
| Aerosol inhaler, 125 mcg per dose | | 120 dose | OP 🗸 | Flixotide |
| Aerosol inhaler, 250 mcg per dose | | 120 dose | OP 🗸 | Flixotide |
| Powder for inhalation, 250 mcg per dose | | 60 dose | OP 🗸 | Flixotide Accuhaler |
| Inhaled Long-acting Beta-adrenoceptor Agonists | | | | |
| | | | | |
| EFORMOTEROL FUMARATE DIHYDRATE | | | | |
| Powder for inhalation 4.5 mcg per dose, breath activated | | | | |
| (equivalent to eformoterol fumarate 6 mcg metered dose) | | 60 dose | OP | |
| | (16.90) | | | Oxis Turbuhaler |
| NDACATEROL | | | | |
| Powder for inhalation 150 mcg | 61.00 | 30 dose | OP 🗸 | Onbrez Breezhaler |
| Powder for inhalation 300 mcg | 61.00 | 30 dose | OP 🗸 | Onbrez Breezhaler |
| SALMETEROL | | | | |
| Aerosol inhaler CFC-free, 25 mcg per dose | 26.25 | 120 dose | | Serevent |
| Powder for inhalation, 50 mcg per dose, breath activated | | 60 dose | ••• | Serevent Accuhaler |
| i owder for inflatation, so meg per dose, breath activated | 20.20 | ou uuse | Ui⁼ ▼ | Serevenit Accunater |

| | Subsidy (Manufacturer's \$ | | Fully Brand or dised Generic ✓ Manufacturer | | |
|--|----------------------------------|----------------------------|---|--|--|
| Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists | | | | | |
| BUDESONIDE WITH EFORMOTEROL Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide v 6 mcg eformoterol fumarate metered dose) Powder for inhalation 320 mcg with 9 mcg eformoterol fuma per dose (equivalent to 400 mcg budesonide with 12 mc | 41.50 Irate | 120 dose OP | ✓ DuoResp Spiromax | | |
| eformoterol fumarate metered dose) - No more than 2 | | 120 dose OP | | | |
| dose per day Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg | | 120 dose OP 120 dose OP | ✓ DuoResp Spiromax ✓ Vannair | | |
| Powder for inhalation 100 mcg with eformoterol fumarate 6 | | 120 dose OP | Symbicort Turbuhaler 100/6 | | |
| Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg | 21.40 | 120 dose OP | 🗸 Vannair | | |
| Powder for inhalation 200 mcg with eformoterol fumarate 6 | mcg 33.74 | 120 dose OP | Symbicort Turbuhaler 200/6 | | |
| Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg – No more than 2 dose per day | | 60 dose OP | Symbicort Turbuhaler 400/12 | | |
| LUTICASONE FUROATE WITH VILANTEROL | | | | | |
| Powder for inhalation 100 mcg with vilanterol 25 mcg | | 30 dose OP | Breo Ellipta | | |
| LUTICASONE WITH SALMETEROL | | | | | |
| Aerosol inhaler 50 mcg with salmeterol 25 mcg | 25.79 | 120 dose OP | Seretide | | |
| Aerosol inhaler 125 mcg with salmeterol 25 mcg | | 120 dose OP | Seretide | | |
| Powder for inhalation 100 mcg with salmeterol 50 mcg – No more than 2 dose per day | | 60 dose OP | ✓ Seretide Accuhaler | | |
| Powder for inhalation 250 mcg with salmeterol 50 mcg – No more than 2 dose per day | | 60 dose OP | Seretide Accuhaler | | |
| Beta-Adrenoceptor Agonists | | | | | |
| ALBUTAMOL | | | | | |
| Oral liq 400 mcg per ml | | 150 ml | ✓ Ventolin | | |
| Infusion 1 mg per ml, 5 ml | | 10 | Ventolin | | |

| Oral liq 400 mcg per ml | 40.00 |
|---|--------|
| Infusion 1 mg per ml, 5 ml | 130.00 |
| Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO | 130.00 |

5

✓ Ventolin

| | Subsidy | Drian) Suba | Fully Brand or idised Generic |
|--|-----------------------|-------------|---|
| | (Manufacturer's \$ | Per Per | Manufacturer |
| Inhaled Beta-Adrenoceptor Agonists | | | |
| SALBUTAMOL | | | |
| Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO | | 200 dose OP | Respigen |
| | (6.20) | | ✓ SalAir Ventolin |
| Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb | · · · | | |
| available on a PSO | 8.96 | 20 | ✓ <u>Asthalin</u> ✓ PMS- |
| | | | Salbutamol S29 |
| | | | Teva-Salbutamol Sterinebs P.F. 629 |
| | | | ✓ Ventolin |
| | 51.11 | | Vebules s29 ✓ Accord s29 |
| Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb | | 00 | |
| available on a PSO | 9.43 | 20 | ✓ <u>Asthalin</u> ✓ PMS- |
| | | | Salbutamol S29 |
| | 14.15 | 30 | ✓ Salbutamol |
| (Respigen Aerosol inhaler, 100 mcg per dose CFC free to be del | isted 1 Septem | ber 2024) | Cipla S29 |
| TERBUTALINE SULPHATE | | | |
| Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated | 22.20 | 120 dose OP | Bricanyl Turbuhaler |
| Anticholinergic Agents | | | |
| PRATROPIUM BROMIDE | | | |
| Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dos available on a PSO | | 200 dose OP | ✓ Atrovent |
| Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne available on a PSO | | 20 | ✓ Univent |
| | 11.73 28.20 | 20 | Accord \$29 |
| Inhaled Beta-Adrenoceptor Agonists with Antic | holinergic / | Agents | |
| SALBUTAMOL WITH IPRATROPIUM BROMIDE | | | |
| Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p dose CFC-free | | 200 dose OP | ✓ Duolin HFA |
| Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per | | | |
| vial, 2.5 ml ampoule – Up to 20 neb available on a PSO | 11.04 33.12 | 20 60 | ✓ <u>Duolin</u> ✓ Duolin |
| | | | Respules S29 |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully sidised | Brand or Generic Manufacturer |
|--|---|--|--|--|
| Long-Acting Muscarinic Antagonists | | | | |
| GLYCOPYRRONIUM – Subsidy by endorsement a) Inhaled glycopyrronium treatment will not be subsidised i umeclidinium. b) Glycopyrronium powder for inhalation 50 mcg per dose is having COPD using spirometry if spirometry is possible, Powder for inhalation 50 mcg per dose | s subsidised only for and the prescription i | patients wh s endorsed dose OP nt with sub- as having 0 who had t 30 dose dose OP | io have l accord sidised COPD t totropiu sotropiu sotropiu | been diagnosed as dingly. Seebri Breezhaler inhaled glycopyrronium or using spirometry if m dispensed before Spiriva spiriva Respimat |
| a) Umeclidinium will not be subsidised if patient is also rece tiotropium bromide. b) Umeclidinium powder for inhalation 62.5 mcg per dose is COPD using spirometry if spirometry is possible, and the Powder for inhalation 62.5 mcg per dose | subsidised only for prescription is endo | patients wh | o have lingly. | |
| Long-Acting Muscarinic Antagonists with Long | -Acting Beta-Ac | drenoce | otor A | gonists |

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

➡SA1584 Special Authority for Subsidy

| Initial application from any relevant practitioner. | Approvals valid for 2 years for applications meeting the following criteria: |
|---|--|
| Both: | |

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).
- GLYCOPYRRONIUM WITH INDACATEROL Special Authority see SA1584 above Retail pharmacy Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

UMECLIDINIUM WITH VILANTEROL – Special Authority see SA1584 above – Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg77.00 30 dose OP 🖌 Anoro Ellipta

Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – Special Authority see SA2326 on the next page – Retail pharmacy

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

⇒SA2326 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^{9} cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

Antifibrotics

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

| Cap 100 mg | 2,554.00 | 60 OP | Ofev |
|------------|----------|-------|--------------------------|
| Cap 150 mg | 3,870.00 | 60 OP | Ofev |

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) | S | Fully ubsidised | Brand or Generic |
|---|-----------------------------------|-------|--------------------|---------------------|
| | \$ | Per | 1 | Manufacturer |
| PIRFENIDONE - Retail pharmacy-Specialist - Special Authori | ty see SA2013 below | | | |
| Note: Pirfenidone is not subsidised in combination with sul | bsidised nintedanib. | | | |
| Tab 801 mg | | 90 OP | 🖌 E | sbriet |
| Tab 267 mg | 1,215.00 | 90 | 🗸 E | sbriet |
| ►SA2013 Special Authority for Subsidy | | | | |

013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib: or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib: and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists

MONTELUKAST

| * | Tab 4 mg | 28 | Montelukast Viatris |
|---|---------------|----|---|
| | Tab 5 mg | 28 | Montelukast Viatris |
| * | Tab 10 mg2.90 | 28 | Montelukast Viatris |

Methylxanthines

AMINOPHYLLINE

| * Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available o | na | | |
|---|-------|--------|---------------------------------------|
| PSO | | 5 | DBL Aminophylline |
| THEOPHYLLINE | | | |
| * Tab long-acting 250 mg | 24.90 | 100 | Nuelin-SR |
| * Oral liq 80 mg per 15 ml | 17.95 | 500 ml | Nuelin |

Mucolytics

| DORNASE ALFA - Special Authority see SA1978 below - Reta | il pharmacy |
|--|-------------|
| Nebuliser soln. 2.5 mg per 2.5 ml ampoule | |

Pulmozvme

SA1978 Special Authority for Subsidy

Initial application - (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

continued...

6

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2196 below

| Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 3 | 37.5 mg | | |
|--|-----------|-------|------------|
| (56) and ivacaftor 75 mg (28) | 27,647.39 | 84 OP | 🗸 Trikafta |
| Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor | 75 mg | | |
| (56) and ivacaftor 150 mg (28) | 27,647.39 | 84 OP | 🗸 Trikafta |

⇒SA2196 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Either:
 - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and

4 Either:

- 4.1 Patient has a heterozygous or homozygous F508del mutation; or
- 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and

6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Notes:

 a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information <u>https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212273s004lbl.pdf</u>

IVACAFTOR - PCT only - Specialist - Special Authority see SA2017 below

| Tab 150 mg | | 56 | Kalydeco |
|-----------------------------|---|----|------------------------------|
| Oral granules 50 mg, sachet | | 56 | Kalydeco |
| Oral granules 75 mg, sachet | - | 56 | Kalydeco |

➡SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

| | Subsidy | Fu | Ily Brand o | r |
|----|-----------------------|----------|----------------------------|--------|
| (N | lanufacturer's Price) | Subsidis | ed Generic | : |
| | \$ | Per | Manufa | cturer |

continued...

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

| Not funded for use as a nasal drop. Soln 7% | 90 ml OP | ✓ Biomed |
|--|----------------------------|--|
| Nasal Preparations | | |
| Allergy Prophylactics | | |
| BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose | 200 dose OP 200 dose OP | ✓ SteroClear✓ SteroClear |
| FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose1.98 | 120 dose OP | Flixonase Hayfever <u>& Allergy</u> |
| IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%5.23 | 15 ml OP | ✓ Univent |
| Respiratory Devices | | |
| MASK FOR SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO c) Only for children aged six years and under Small | 1 | ✓ e-chamber Mask |
| PEAK FLOW METER a) Up to 25 dev available on a PSO b) Only on a PSO | | |
| Low range | 1 | Mini-Wright AFS Low Range |
| Normal range9.54 | 1 | ✓ Mini-Wright Standard |

| | Subsidy (Manufacturer's Price) \$ |) S Per | Fully ubsidised | Brand or Generic Manufacturer |
|---|---|------------|--------------------|-------------------------------------|
| SPACER DEVICE | | | | |
| a) Up to 50 dev available on a PSO | | | | |
| b) Only on a PSO | | | | |
| 220 ml (single patient) | 3.65 | 1 | 🗸 e | e-chamber Turbo |
| 510 ml (single patient) | 5.95 | 1 | √ € | e-chamber La Grande |
| 800 ml | 6.50 | 1 | ۷ ۱ | /olumatic |
| Respiratory Stimulants | | | | |
| CAFFEINE CITRATE Oral liq 20 mg per ml (10 mg base per ml) | | 5 ml OP | ✓ E | Biomed |

| | Subsidy | | Fully Brand or |
|---|---------------------|---------------------|---|
| | (Manufacturer's Pri | ice) Subs | sidised Generic |
| | \$ | Per | Manufacturer |
| Ear Preparations | | | |
| | | | |
| LUMETASONE PIVALATE Ear drops 0.02% with clioquinol 1% | 1 16 | | I accortan Visform |
| Ear drops 0.02% with choquinor 1% | 4.40 | 7.5 ml OP | Locacorten-Viaform ED's |
| | | | ✓ Locorten-Vioform |
| RIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC | IN AND NYSTATI | N | |
| Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate | - | | |
| 2.5 mg and gramicidin 250 mcg per g | 5.16 | 7.5 ml OP | Kenacomb |
| Ear/Evo Droporationa | | | |
| Ear/Eye Preparations | | | |
| DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN | | | |
| Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and | | | |
| gramicidin 50 mcg per ml | | 8 ml OP | Otoday 520 |
| | (9.27) (9.27) | | Otodex S29 Sofradex |
| FRAMYCETIN SULPHATE | (0.27) | | Conduck |
| Ear/Eye drops 0.5% | 4.13 | 8 ml OP | |
| | (8.65) | | Soframycin |
| Eve Proportions | | | |
| Eye Preparations | | | |
| Eye preparations are only funded for use in the eye, unless expli | citly stated otherw | ise. | |
| Anti-Infective Preparations | | | |
| ACICLOVIR | | | |
| ₭ Eye oint 3% | 14.88 | 4.5 g OP | ✓ ViruPOS |
| CHLORAMPHENICOL | | | |
| Eye oint 1% | 1.09 | 5 g OP | ✓ <u>Devatis</u> |
| Eye drops 0.5% | | 10 ml OP | Chlorsig |
| Funded for use in the ear*. Indications marked with * a | re unapproved indi | cations. | |
| CIPROFLOXACIN Eye drops 0.3% – Subsidy by endorsement | 9.73 | 5 ml OP | Ciprofloxacin Teva |
| When prescribed for the treatment of bacterial keratitis | | | |
| for the second line treatment of chronic suppurative otiti | | | |
| Note: Indication marked with a * is an unapproved indic | cation. | | |
| SODIUM FUSIDATE [FUSIDIC ACID] | | | 6 |
| Eye drops 1% | 5.29 | 5 g OP | Fucithalmic |
| FOBRAMYCIN | 10.45 | 2 5 a OP | |
| Eye oint 0.3% Eye drops 0.3% | | 3.5 g OP 5 ml OP | ✓ Tobrex ✓ Tobrex |
| Corticosteroids and Other Anti-Inflammatory P | | | |
| | | | |
| DEXAMETHASONE ₩ Eye oint 0.1% | 5.86 | 3.5 g OP | Maxidex |
| ★ Eye on 0.1% | | 3.5 g OP 5 ml OP | ✓ Maxidex ✓ Maxidex |
| Ocular implant 700 mcg – Special Authority see SA1680 on | | 0 | |
| the next page - Retail pharmacy | | 1 | ✓ Ozurdex |
| | | | |
| fully subsidised | S29 Linannr | oved medicine s | supplied under Section 29 |

| Subsidy | | Fully | Brand or | |
|------------------------|------|-------|--------------|--|
| (Manufacturer's Price) | Subs | dised | Generic | |
| \$ | Per | 1 | Manufacturer | |

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Either:
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

| * Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b | | |
|---|----------|-------------------------------------|
| sulphate 6,000 u per g5.39 | 3.5 g OP | Maxitrol |
| * Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin | | |
| b sulphate 6,000 u per ml 4.50 | 5 ml OP | Maxitrol |
| DICLOFENAC SODIUM | | |
| Eye drops 0.1% | 5 ml OP | Voltaren Ophtha |
| (Voltaren Ophtha Eye drops 0.1% to be delisted 1 December 2024) | | i |
| FLUOROMETHOLONE | | |
| * Eye drops 0.1% | 5 ml OP | 🖌 FML |
| 5.20 | | Flucon |
| LEVOCABASTINE | | |
| Eye drops 0.5 mg per ml8.71 | 4 ml OP | |
| _); = ================================== |) | Livostin |
| LODOXAMIDE | | |
| Eye drops 0.1% | 10 ml OP | Lomide |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

SENSORY ORGANS

| | Subsidy | | Fully Brand or |
|---|-------------------------|---------------------|--|
| | (Manufacturer's F \$ | Price) Subs Per | sidised Generic Manufacturer |
| IEPAFENAC | | 101 | |
| Eye drops 0.3% | 8.80 | 3 ml OP | ✓ Ilevro |
| REDNISOLONE ACETATE | | | |
| Eye drops 1% | 6.92 7.00 | 10 ml OP 5 ml OP | Prednisolone-AFT Pred Forte |
| REDNISOLONE SODIUM PHOSPHATE - Special Authority | | | |
| Eye drops 0.5%, single dose (preservative free) | 41.20 | 20 dose | Minims Prednisolone |
| SA1715 Special Authority for Subsidy nitial application only from an ophthalmologist or optometrist ollowing criteria: Both: | . Approvals valid f | or 6 months for | r applications meeting the |
| Patient has severe inflammation; and Patient has a confirmed allergic reaction to preservative Renewal from any relevant practitioner. Approvals valid for 6 in | , , | treatment rema | ins appropriate and the patient |
| penefiting from treatment. | | | |
| SODIUM CROMOGLICATE Eye drops 2% | 2.62 | 10 ml OP | ✓ <u>Allerfix</u> |
| Glaucoma Preparations - Beta Blockers | | | |
| ETAXOLOL | | | |
| ₭ Eye drops 0.25% | | 5 ml OP | Betoptic S Betoptic |
| ₭ Eye drops 0.5% | 7.50 | 5 ml OP | Betoptic |
| IMOLOL | | | 4 - - - - - - - - - - |
| € Eye drops 0.25% € Eye drops 0.5% | | 5 ml OP 5 ml OP | <u>Arrow-Timolol</u> <u>Arrow-Timolol</u> |
| Glaucoma Preparations - Carbonic Anhydrase | Inhibitors | | |
| NCETAZOLAMIDE ₭ Tab 250 mg | 17.03 | 100 | Diamox |
| RINZOLAMIDE ∉ Eye drops 1% | 7.30 | 5 ml OP | ✓ <u>Azopt</u> |
| OORZOLAMIDE WITH TIMOLOL ₭ Eye drops 2% with timolol 0.5% | 2.73 | 5 ml OP | ✓ Dortimopt |
| Glaucoma Preparations - Prostaglandin Analo | gues | | |
| BIMATOPROST ≰ Eye drops 0.03% | 5.95 | 3 ml OP | ✓ <u>Bimatoprost</u> Multichem |
| ATANOPROST € Eye drops 0.005% | | 2.5 ml OP | ✓ <u>Teva</u> |
| | | | |

SENSORY ORGANS

| | Subsidy | | Fully | Brand or |
|---|-------------------|------------|------------|--------------------|
| | (Manufacturer's P | rice) Subs | idised | Generic |
| | \$ | Per | 1 | Manufacturer |
| Glaucoma Preparations - Other | | | | |
| BRIMONIDINE TARTRATE | | | | |
| * Eye drops 0.2% | 4.29 | 5 ml OP | ✓ <u>P</u> | Arrow-Brimonidine |
| BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE | | | | |
| * Eye drops 0.2% with timolol maleate 0.5% | | 5 ml OP | ✓ (| Combigan |
| LATANOPROST WITH TIMOLOL | | | | |
| * Eye drops 0.005% with timolol 0.5% | 4.95 | 2.5 ml OP | ✓ <u>A</u> | Arrow - Lattim |
| PILOCARPINE HYDROCHLORIDE | | | | |
| * Eye drops 1% | 4.26 | 15 ml OP | 🗸 s | sopto Carpine |
| * Eye drops 2% | 5.35 | 15 ml OP | ✓ s | sopto Carpine |
| * Eye drops 4% | | 15 ml OP | ✓ Is | sopto Carpine |
| Subsidised for oral use pursuant to the Standard Formu | | | | |
| PILOCARPINE NITRATE | | | | |
| * Eye drops 2% single dose – Special Authority see SA0895 | | | | |
| below – Retail pharmacy | | 20 dose | 🗸 N | linims Pilocarpine |
| ➡ SA0895 Special Authority for Subsidy | | | | |

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

| ATROPINE SULPHATE * Eye drops 1% | 15 ml OP | ✓ Atropt |
|--|----------------------|--|
| CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1% | 15 ml OP | ✓ Cyclogyl |
| Eye drops 1%, single dose (preservative free) – Only on a prescription | 20 dose | Minims Cyclopentolate |
| TROPICAMIDE * Eye drops 0.5% * Eye drops 1% 8.66 | 15 ml OP 15 ml OP | Mydriacyl Mydriacyl |
| Preparations for Tear Deficiency | | |
| For acetylcysteine eye drops refer Standard Formulae, page 268 HYPROMELLOSE * Eye drops 0.5% | 15 ml OP | ✓ Methopt |

| | Subsidy (Manufacturer's Price \$ | Full) Subsidise Per ✔ | d Generic |
|--|--|------------------------------|---|
| Preservative Free Ocular Lubricants | | | |
| SA2134 Special Authority for Subsidy | | | |
| nitial application from any relevant practitioner. Approvals va Both: | lid for 12 months for a | applications mee | ing the following criteria: |
| Confirmed diagnosis by slit lamp or Schirmer test of seve Either: | ere secretory dry eye; | and | |
| 2.1 Patient is using eye drops more than four times d 2.2 Patient has had a confirmed allergic reaction to p | | | |
| Renewal from any relevant practitioner. Approvals valid for 24 drops and has benefited from treatment. | | • | o require lubricating eye |
| ARBOMER – Special Authority see SA2134 above – Retail pl Ophthalmic gel 0.3%, 0.5 g | | 30 🖌 | ' Poly-Gel |
| OLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL - Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml | | | e – Retail pharmacy ´ Systane Unit Dose |
| SODIUM HYALURONATE [HYALURONIC ACID] – Special Au Eye drops 1 mg per ml Hylo-Fresh has a 6 month expiry after opening. The Pl month is not relevant and therefore only the prescribed | 13.85 1 harmacy Procedures | 0 ml OP | Hylo-Fresh on allowing one bottle per |
| Other Eye Preparations | | | |
| NAPHAZOLINE HYDROCHLORIDE ★ Eye drops 0.1% | 4.15 1 | 5 ml OP 🗸 | Naphcon Forte |
| DLOPATADINE Eye drops 0.1% | 2.17 | 5 ml OP 🖌 | Olopatadine Teva |
| PARAFFIN LIQUID WITH WOOL FAT ₭ Eye oint 3% with wool fat 3% | | 3.5 g OP 🖌 | Poly-Visc |
| | | | |

RETINOL PALMITATE

5 g OP 🖌 VitA-POS

VARIOUS

| | Outside | | Fully Drand an | |
|--|-----------------------------------|-------------|---|------------|
| | Subsidy (Manufacturer's Price) | | Fully Brand or dised Generic | |
| | \$ | Per | Manufact | urer |
| | | | | |
| Various | | | | |
| PHARMACY SERVICES | | | | |
| Brand switch fee | 4 50 | 1 fee | 🗸 BSF Max H | ealth |
| | | 1100 | ✓ BSF | cann |
| | | | Methylph | enidate |
| | | | ER - Teva | |
| | | | BSF Noum | ed |
| | | | Dexamfe | |
| | | | BSF Wock | nardt |
| a) May only be claimed once per patient. | | | | |
| b) The Pharmacode for BSF Wockhardt is 2669986 - se | e also <mark>page 128</mark> | | | |
| c) The Pharmacode for BSF Noumed Dexamfetamine is | 2673886 - see also | page 146 | | |
| d) The Pharmacode for BSF Methylphenidate ER - Teva | | lso page 14 | 7 | |
| e) The Pharmacode for BSF Max Health is 2677903 - se | | | | |
| * COVID-19 Services | 0.00 | 1 fee | After Hours | |
| | | | Mgmt 15 | |
| | | | ✓ After Hours | |
| | | | Mgmt 30 | |
| | | | ✓ After Hours | |
| | | | Mgmt 45 | |
| | | | Antivirals I Review | Ingromity |
| | | | | |
| | | | Complianc Packagir | |
| | | | ✓ Med Mgmt | • |
| | | | Med Mgmt Med Mgmt | |
| | | | Med Mgmt Med Mgmt | |
| | | | ✓ Medicine D | |
| * Immunisation administration fee | 0.00 | 1 fee | Immunisati | |
| | | | Administ | ration |
| * Immunisation co-administration fee | 0.00 | 1 fee | Immunisati | ion |
| | | | Co-admir | nistration |
| (BSF Max Health Brand switch fee to be delisted 1 August 2024) | | | | |
| (BSF Methylphenidate ER - Teva Brand switch fee to be delisted | | | | |
| (BSF Noumed Dexamfetamine Brand switch fee to be delisted 1 | June 2024) | | | |
| (BSF Wockhardt Brand switch fee to be delisted 1 June 2024) | | | | |
| Agente Llood in the Treatment of Reisenings | | | | |
| Agents Used in the Treatment of Poisonings | | | | |
| Antidotes | | | | |
| Annuoles | | | | |
| ACETYLCYSTEINE | | | | |
| Inj 200 mg per ml, 10 ml ampoule | | 10 | ✓ Martindale | Pharma |
| NALOXONE HYDROCHLORIDE | | | | |
| a) Up to 10 inj available on a PSO | | | | |
| b) Only on a PSO | | | | |
| * Inj 400 mcg per ml, 1 ml ampoule | | 10 | ✓ Hameln | |
| | | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Prio \$ | ce) Subs Per | Fully sidised | Brand or Generic Manufacturer |
|--|--|---------------------------------|-------------------|-------------------------------------|
| Removal and Elimination | | | | |
| IARCOAL | | | | |
| Oral liq 50 g per 250 ml a) Up to 250 ml available on a PSO b) Only on a PSO | 43.50 | 250 ml OP | ✓ (| Carbosorb-X |
| FERASIROX – Special Authority see SA1492 below – Ret Wastage claimable | ail pharmacy | | | |
| Tab 125 mg dispersible | | 28 | ✓ E | xjade |
| Tab 250 mg dispersible | | 28 | ✓ E | xjade |
| Tab 500 mg dispersible | 1,105.00 | 28 | ✓ E | xjade |
| combination therapy have proven ineffective as 3.2 Treatment with deferiprone has resulted in seve 3.3 Treatment with deferiprone has resulted in arthu 3.4 Treatment with deferiprone is contraindicated d count (ANC) of < 0.5 cells per μL) or recurrent e 0.5 - 1.0 cells per μL). | ere persistent vomiting itis; or ue to a history of agra | g or diarrhoe | a; or (defined | as an absolute neutrop |
| newal only from a haematologist. Approvals valid for 2 yea her: | ars for applications m | eeting the fol | lowing | criteria: |
| For the first renewal following 2 years of therapy, the tr improvement in all three parameters namely serum fer For subsequent renewals, the treatment has been tole in all three parameters namely serum ferritin, cardiac N | ritin, cardiac MRI T2* rated and has resulted | and liver MF d in clinical s | l T2* le | vels; or |
| | ail pharmacy | | | |

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

| * | Inj 500 mg vial | | |
|---|-----------------|--|--|
|---|-----------------|--|--|

 ✓ DBL Desferrioxamine Mesylate for Inj BP
 ✓ Deferoxamine Pfizer

10

| | Subsidy | Fully | Brand or |
|---------------------------|------------------------|------------|-------------------------------|
| | (Manufacturer's Price) | Subsidised | Generic |
| | \$ | Per 🗸 | Manufacturer |
| SODIUM CALCIUM EDETATE | | | |
| * Inj 200 mg per ml, 5 ml | | 6 | |
| | (156.71) | | Calcium Disodium Versenate |

VARIOUS

Standard Formulae

| ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base | qs qs | PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water | 1 g 70 ml to 100 ml |
|--|---|---|---|
| CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol Preservative Water | 60 mg 40 ml qs to 100 ml | PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml) Phenobarbitone Sodium Glycerol BP Water | LIQUID (10 400 mg 4 ml to 40 ml |
| CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative Water FOLINIC MOUTHWASH Calcium folinate 15 mg tab | 300 mg 40 ml qs to 100 ml 1 tab | PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative Water (Preservative should be used if quantity supplied is than 5 days.) | qs qs to 500 ml for more |
| Preservative Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.) METHADONE MIXTURE Methadone powder | qs to 500 ml for more qs | SALIVA SUBSTITUTE FORMULA Methylcellulose Preservative Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.) | 5 g qs to 500 ml for more |
| Glycerol Water METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol | qs to 100 ml 10 g to 100 ml | SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml Water (Only funded if prescribed for treatment of hyponatr VANCOMYCIN ORAL SOLUTION (25 mg per ml) | |
| (Use 1 ml of the 10% solution per 100 ml of oral liqu OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water | uid mixture) qs 8.4 g to 100 ml | Vancomycin 500 mg injection Glycerin with sucrose suspension Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure) | 5 vials 37.5 ml to 100 ml um difficile |

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

| | Subsidy | | Fully Brand or |
|---|----------------------|----------------|----------------------------------|
| | (Manufacturer's Pric | e) Sub: Per | sidised Generic Manufacturer |
| | φ | Per | Manufacturer |
| Extemporaneously Compounded Preparations a | nd Galenicals | 6 | |
| CODEINE PHOSPHATE - Safety medicine; prescriber may deter | rmine dispensing f | requency | |
| Powder – Only in combination | 63.09 | 25 g | |
| | (90.09) | | Douglas |
| Only in extemporaneously compounded codeine linctus. | | | |
| COLLODION FLEXIBLE | | | |
| Note: This product is no longer being manufactured by the su | pplier and will be | delisted fror | n the Schedule at a date to be |
| determined. Collodion flexible | 10.00 | 1001 | |
| | | 100 ml | ✓ PSM |
| COMPOUND HYDROXYBENZOATE – Only in combination | | | |
| Only in extemporaneously compounded oral mixtures. Soln | 20.00 | 100 ml | ✓ Midwest |
| | | 100 111 | • midwest |
| GLYCERIN WITH SODIUM SACCHARIN – Only in combination | music and louvid (| Standard Fa | mulaa |
| Only in combination with Ora-Plus or when used in the vancor Suspension | | 473 ml | ✓ Ora-Sweet SF |
| | | 475111 | • Ola-Sweet Si |
| GLYCERIN WITH SUCROSE – Only in combination Only in combination with Ora-Plus or when used in the vancor | music and launid (| Standard Ea | rmulaa |
| Suspension | | 473 ml | ✓ Ora-Sweet |
| GLYCEROL | | 470111 | |
| KOLICEROL KOLICEROL | 3 23 | 500 ml | healthE Glycerol BP |
| Only in extemporaneously compounded oral liquid prepar | | 500 m | |
| METHADONE HYDROCHLORIDE | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing free | quency | | |
| d) Extemporaneously compounded methadone will only be re | | ate of the ch | neapest form available |
| (methadone powder, not methadone tablets). | | | |
| Powder | 7.84 | 1 g | ✓ AFT |
| (AFT Powder to be delisted 1 July 2024) | | | |
| METHYL HYDROXYBENZOATE | | | |
| Powder | 8.98 | 25 g | Midwest |
| METHYLCELLULOSE | | | |
| Powder | | 100 g | MidWest |
| Suspension – Only in combination | | 473 ml | Ora-Plus |
| METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHA | | | 6 a b b b a |
| Suspension | | 473 ml | Ora-Blend SF |
| METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only | | | |
| Suspension | | 473 ml | Ora-Blend |
| PHENOBARBITONE SODIUM | | | 4 • • • • • • • |
| Powder – Only in combination | | 10 g | MidWest |
| Only in children up to 10 years | 325.00 | 100 g | MidWest |
| Only in children up to 12 years | | | |
| PROPYLENE GLYCOL | ata 100/ | | |
| Only in extemporaneously compounded methyl hydroxybenzo Lig | | 500 ml | ✓ Midwest |
| Lid | 11.20 | 500 mi | • Miuwest |
| | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

| | Subsidy (Manufacturer's Pric \$ | e) Si Per | Fully ubsidised | Brand or Generic Manufacturer |
|--|---------------------------------------|-------------------|--------------------|-------------------------------------|
| SODIUM BICARBONATE | Ŷ | 1.01 | | Manufacturer |
| Powder BP – Only in combination Only in extemporaneously compounded omeprazole and | | 500 g pension. | 🗸 N | lidwest |
| SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatio | ns. | | | |
| Liq | 14.95 | 500 ml | 🗸 N | lidwest |
| WATER | | | | |
| Tap – Only in combination | 0.00 | 1 ml | 🗸 Т | ap water |

SECTION D: SPECIAL FOODS

Fully

Subsidy (Manufacturer's Price)

\$

Subsidised Per ✓ Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Either:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children; or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Initial application — (Inborn errors of metabolism)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. **Renewal — (Cystic fibrosis or renal failure)** only from a dietitian, relevant specialist, vocationally registered general

practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

. Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

| Powder6.72 | 400 g OP | Polycal |
|------------|----------|-----------------------------|
|------------|----------|-----------------------------|

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

| Subsidy | | Fully | Brand or | |
|-------------------|----------|-----------|--------------|--|
| (Manufacturer's F | Price) S | ubsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| CARBOHYDRATE AND FAT S | SUPPLEMENT – Special Author | ity see SA1376 on | the previous pag | je - | Hospital pharmacy [HP3] |
|------------------------|-----------------------------|-------------------|------------------|------|-------------------------|
| Powder (neutral) | | | 400 g OP | 1 | Duocal Super |
| | | | - | | Soluble Powder |

Fat

⇒SA2204 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 10 ascites; or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Indications other than inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| FAT SUPPLEMENT - Special Authority see SA2204 on the | ne previous page – Hos | spital pharmacy | [HP3] |
|--|------------------------|-----------------|--|
| Emulsion (neutral) | | 200 ml OP | Calogen |
| | 38.44 | 500 ml OP | Calogen |
| Emulsion (strawberry) | | 200 ml OP | Calogen |
| Oil | | 500 ml OP | MCT oil (Nutricia) |
| MCT Emulsion, 250 ml | | 4 OP | Liquigen |

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| PROTEIN SUPPLEMENT - Specia | al Authority see SA1524 above – Hospital ph | armacy [HP3] | |
|-----------------------------|---|--------------|------------------------------|
| Powder | | 227 g OP | Resource |
| | | | Beneprotein |
| | 13.82 | 225 g OP | Protifar |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| DIABETIC ENTERAL FEED 1KCAL/ML – Special Authorit Liquid | • | 500 ml OP | Glucerna Select Nutrison Advanced |
|---|---------------------|-----------------|--|
| (Nutrison Advanced Diason Liquid to be delisted 1 July 202 | 24) | | Diason |
| | , | | |
| DIABETIC ORAL FEED 1KCAL/ML – Special Authority se | e SA1095 above – Ho | spital pharmacy | [HP3] |
| Liquid (strawberry) | | 200 ml OP | Diasip |
| Liquid (vanilla) | 1.50 | 200 ml OP | ✓ Diasip |
| | 2.10 | | Nutren Diabetes |

Fat Modified Products

⇒SA2205 Special Authority for Subsidy

Initial application — (**Inborn errors of metabolism**) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. **Initial application** — (**Indications other than errors of inborn metabolism**) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Monogen

| Subsidy | | | |
|------------------------|------------|---|--------------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | ✓ | Manufacturer |

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see | SA1099 above – Hos | oital pharmacy | [HP3] |
|--|--------------------|----------------|-------------------------------|
| Powder | 64.26 | 400 g OP | Kindergen |

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

| | Subsidy | | Fully | Brand or |
|---|-----------------------------|--|--------------|---|
| | (Manufacturer's Price \$ | e) Sub Per | sidised ✓ | Generic Manufacturer |
| ontinued | | | | |
| pplications meeting the following criteria: Both: | | | | |
| The treatment remains appropriate and the patient is benuezed. General Practitioners must include the name of the dietitian practitioner and date contacted. | | | onally re | egistered general |
| AEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority Liquid | | e <mark>previous </mark> 500 ml OP | ✓ F | Hospital pharmacy [HP3 Frebini Energy Iutrini Energy RTH |
| AEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority s Liquid | | previous pa 500 ml OP | ✓ F ✓ N | spital pharmacy [HP3] Pediasure RTH Nutrini RTH Frebini Original |
| AEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Sp harmacy [HP3] | - | SA1379 on | the prev | vious page – Hospital |
| Liquid | 7.00 § 7.14 | 500 ml OP | | rebini Energy Fibre lutrini Energy Multi Fibre |
| PAEDIATRIC ENTERAL FEED WITH FIBRE 1KCAL/ML - Spectharmacy [HP3] | cial Authority see SA | A1379 on th | e previc | ous page – Hospital |
| Liquid | 7.00 § | 500 ml OP | ✓ F | rebini Original Fibre |
| AEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see | | | | |
| Liquid (strawberry) | | 200 ml OP | | ortini |
| Liquid (vanilla) | | 200 ml OP | - | ortini |
| | | 500 ml OP | | Pediasure Plus |
| PAEDIATRIC ORAL FEED 1KCAL/ML – Special Authority see S | | | | |
| Liquid (chocolate) | | 200 ml OP | - | Pediasure |
| Liquid (strawberry) | | 200 ml OP | | Pediasure |
| Liquid (vanilla) | | 200 ml OP | - | Pediasure |
| | | 250 ml OP | - | Pediasure |
| PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML – Special harmacy [HP3] | Authority see SA1 | 379 on the | previous | s page – Hospital |
| Liquid (unflavoured) | | 200 ml OP | 🖌 F | Fortini Multi Fibre |
| Liquid (chocolate) | | 200 ml OP | | ortini Multi Fibre |
| Liguid (strawberry) | | 200 ml OP | - | ortini Multi Fibre |
| | 100 0 | 200 ml OP | 🗸 F | Fortini Multi Fibre |
| Liquid (vanilla) | | | | |
| Liquid (vanilla) PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 | | ge – Hospi | tal pharr | nacy [HP3] |

Renal Products

■ SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

| | Subsidy (Manufacturer's Price) \$ | Fully Subsidised Per ✔ | |
|--|---|---------------------------------------|--|
| continued recommendation of a dietitian, relevant specialist or vocationally applications meeting the following criteria: Both: | registered general pra | actitioner. Appr | ovals valid for 3 years for |
| The treatment remains appropriate and the patient is ben General Practitioners must include the name of the dietitic practitioner and date contacted. | 0 | | registered general |
| RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see Liquid | | | bital pharmacy [HP3] Nepro HP RTH |
| RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid | | 0 ml OP | pharmacy [HP3] Nepro HP (strawberry) Nepro HP (vanilla) |
| RENAL ORAL FEED 2 KCAL/ML – Special Authority see SA110 Liquid, 200 ml bottle | | <mark>ge</mark> – Hospital pł 4 OP | narmacy [HP3] |
| Liquid (apricot) 125 ml Liquid (caramel) 125 ml | (13.24) 13.72 | | NovaSource Renal Renilon 7.5 Renilon 7.5 |

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Any of the following:
 - 1 malabsorption; or
 - 2 short bowel syndrome; or
 - 3 enterocutaneous fistulas; or
 - 4 eosinophilic oesophagitis; or
 - 5 inflammatory bowel disease; or
 - 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – Sp Liquid | | e SA1377 abov 1,000 ml OP | |
|---|--------|------------------------------|--|
| ORAL ELEMENTAL FEED 0.8KCAL/ML – Special Authority se Liquid (grapefruit), 250 ml carton | | – Hospital phar 18 OP | macy [HP3] ✓ Elemental 028 Extra |
| Liquid (pineapple & orange), 250 ml carton Liquid (summer fruits), 250 ml carton | 179.46 | 18 OP 18 OP | ✓ Elemental 028 Extra ✓ Elemental 028 Extra |

| | Subsidy (Manufacturer's Pri \$ | ice) Subs Per | Fully sidised | Brand or Generic Manufacturer |
|--|--------------------------------------|-------------------------|------------------|---------------------------------------|
| ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see S Powder (unflavoured) | | vious page – 80 g OP | | |
| SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Auth [HP3] | ority see SA1377 | on the previo | us page | Hospital pharmacy |
| Liquid | 7.47 | 500 ml OP | | utrison Advanced Peptisorb |
| | 9.60 | | | urvimed OPD |

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML | - Special Authority | see SA1196 | above · | - Hospital pharmacy [HP3] |
|---|---------------------|------------|---------|---------------------------|
| Liquid | 6.27 | 500 ml OP | 1 | Nutrini Low Energy |
| | | | | Multi Fibre |

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on

| Subsidy | Fully | Brand or | _ |
|------------------------|------------|--------------|---|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal - (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the

recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and

3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:
 - Patient is Malnourished
 - 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
 - 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:

| Subsidy | Full | y Brand or | _ |
|------------------------|-----------|--------------|---|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

- 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
- 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
- 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications

meeting the following criteria:

Any of the following:

1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube - refer to specific medical condition criteria); or

SPECIAL FOODS

| | Subsidy (Manufacturer's F | | Fully Subsidised | Brand or Generic |
|--|------------------------------|--------------|---------------------|-----------------------|
| | \$ | Per | | Manufacturer |
| ontinued | | | | |
| 2 Cystic Fibrosis; or | | | | |
| 3 Liver disease; or | | | | |
| 4 Chronic Renal failure; or | | | | |
| 5 Inflammatory bowel disease; or | | | | |
| 6 Chronic obstructive pulmonary disease with hypercapnia; of | or | | | |
| 7 Short bowel syndrome; or | | | | |
| 8 Bowel fistula; or | | | | |
| 9 Severe chronic neurological conditions. | | | | |
| NTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 o | 1 0 | | , . | |
| Liquid | | 250 ml (| | Ensure Plus HN |
| | 8.68 | 1,000 ml | OP 🗸 | Ensure Plus HN RTH |
| | 9.00 | | | Nutrison Energy |
| | 9.60 | | 1 | Fresubin HP Energy |
| NTERAL FEED 1KCAL/ML – Special Authority see SA1859 on | page 278 – Hos | spital pharr | nacy [HP3 |] |
| Liquid | 1.24 | 250 ml (| OP 🗸 | Isosource Standard |
| | 6.50 | 1,000 ml | OP 🗸 | Fresubin Original |
| | 6.56 | | | Osmolite RTH |
| | 6.90 | | ✓ | Nutrison RTH |
| NTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authority | / see <mark>SA1859</mark> c | n page 27 | 8 – Hospit | al pharmacy [HP3] |
| Liquid | 9.05 | 1,000 ml | OP 🗸 | Nutrison |
| | | | | 800 Complete |
| | | | | Multi Fibre |
| NTERAL FEED WITH FIBRE 1 KCAL/ML – Special Authority se | e SA1859 on p | age 278 - | Hospital p | harmacy [HP3] |
| Liquid | 6.56 | 1,000 ml | OP 🗸 | Jevity RTH |
| | 7.00 | | ✓ | Fresubin Original |
| | | | | Fibre |
| | 7.21 | | ~ | Nutrison Multi Fibre |
| NTERAL FEED WITH FIBRE 1.2KCAL/ML – Special Authority s | | | | |
| Liquid | 7.87 | 1,000 ml | OP 🗸 | Jevity Plus RTH |
| NTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority s | | | | |
| Liquid | 8.68 | 1,000 ml | - | Jevity HiCal RTH |
| | | | 1 | Nutrison Energy |
| | | | | Multi Fibre |
| | 9.80 | | 1 | Fresubin HP Energy |
| | | | | Fibre |
| NTERAL FEED WITH PROTEIN 1.2KCAL/ML – Special Author | ity see SA1859 | on page 2 | | |
| Liquid | 9.60 | 500 ml (| OP 🗸 | Fresubin Intensive |
| RAL FEED (POWDER) – Special Authority see SA1859 on pag | e 278 – Hospita | al pharmac | y [HP3] | |
| Powder (chocolate) | 14.00 | ່ 840 g C | | Sustagen Hospital |
| . , | | 5 | | Formula |
| | 26.00 | 850 g C | DP 🗸 | Ensure |
| Powder (vanilla) | | 840 g C | | Sustagen Hospital |
| · · · / | | 5 - | | Formula Active |
| | 26.00 | 850 g C | P 🗸 | Ensure |
| | _0.00 | | | |

| | Subsidy | | ully Brand or |
|--|--------------------------|----------------------|-----------------------------|
| | (Manufacturer's Pr \$ | ice) Subsidis Per | sed Generic Manufacturer |
| | Ŧ | | |
| ORAL FEED 1.5KCAL/ML – Special Authority see SA1859 on pa | | | |
| Additional subsidy by endorsement is available for patients b | | | |
| epidermolysis bullosa, or as exclusive enteral nutrition in child disease, or for patients with COPD and hypercapnia, defined | | | |
| endorsed accordingly. | as CO2 value ex | | g. The prescription must be |
| | | | |
| Liquid (banana) – Higher subsidy of up to \$1.56 per 200 ml with Endorsement | 0.70 | 200 ml OP | |
| | 0.72 (1.26) | 200 MI OP | Fortisip |
| | (1.26) | | Ensure Plus |
| Liquid (abaaalata) Higher subsidy of up to \$1.56 per 200 m | · / | | LIISUIE FIUS |
| Liquid (chocolate) – Higher subsidy of up to \$1.56 per 200 m with Endorsement | | 200 ml OP | |
| | 0.72 (1.26) | 200 111 0F | Fortisip |
| | (1.26) | | Ensure Plus |
| Liquid (fruit of the forest) – Higher subsidy of \$1.56 per 200 i | · / | | |
| with Endorsement | | 200 ml OP | |
| | (1.56) | 200 III OF | Ensure Plus |
| Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with | () | | Ensure Flus |
| Endorsement | | 200 ml OP | |
| | (1.26) | 200 111 01 | Fortisip |
| Liquid (vanilla) – Higher subsidy of up to \$1.65 per 237 ml w | · · · | | i onop |
| Endorsement | | 200 ml OP | |
| | (1.26) | 200 111 01 | Fortisip |
| | 0.85 | 237 ml OP | i onioip |
| | (1.65) | | Ensure Plus |
| | 0.72 | 200 ml OP | |
| | (1.56) | | Ensure Plus |
| ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see | , | 278 – Hospital r | |
| Additional subsidy by endorsement is available for patients b | | | |
| epidermolysis bullosa. The prescription must be endorsed a | | ough a recurry t | abo, or who have severe |
| Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with | | | |
| Endorsement | | 200 ml OP | |
| | (1.26) | 200 111 01 | Fortisip Multi Fibre |
| Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with | () | | |
| Endorsement | | 200 ml OP | |
| | (1.26) | | Fortisip Multi Fibre |
| Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with | (| | |
| Endorsement | 0.72 | 200 ml OP | |
| | (1.26) | | Fortisip Multi Fibre |
| | · -/ | | - p |
| High Calorie Products | | | |
| • | | | |

► SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 on | the previous pa | <mark>age</mark> – Hospital p | harmacy [HP3] |
|--|-------------------|-------------------------------|--|
| Liquid | 6.50 | 500 ml OP | Fresubin 2kcal HP |
| | 6.82 | | Nutrison Concentrated |
| | 13.64 | 1,000 ml OP | Ensure Two Cal HN RTH |
| ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients be epidermolysis bullosa. The prescription must be endorsed are Liquid (vanilla) – Higher subsidy of \$2.34 per 200 ml with | eing bolus fed tl | | |
| Endorsement | 0.96 | 200 ml OP | |
| | (2.34) | | Two Cal HN |

Food Thickeners

SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| FOOD THICKENER | - Special Authority see SA1106 on the previous page - H | ospital pharmacy | / [HP3] |
|----------------|---|------------------|----------------------------------|
| Powder | | 300 g OP | Nutilis |
| | 24.00 | 380 g OP | Aptamil Feed |
| | | | Thickener |

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

| GLUTEN FREE BAKING MIX – Special Authority see SA1729 above – Hosp Powder | | |
|--|--------------------|----------------------------------|
| (5.15 |) | Healtheries Simple Baking Mix |
| GLUTEN FREE BREAD MIX - Special Authority see SA1729 above - Hospit | tal pharmacy [HP3] | |
| Powder | 1,000 g OP | |
| (7.32 |) | NZB Low Gluten Bread Mix |
| 3.51 | | |
| (10.87 |) | Horleys Bread Mix |
| GLUTEN FREE FLOUR - Special Authority see SA1729 above - Hospital ph | narmacy [HP3] | |
| Powder | 2,000 g OP | |
| (18.10 |) | Horleys Flour |

| | Subsidy | | Fully | Brand or |
|---|---------------------------|-----------------|---------------|-------------------------|
| | (Manufacturer's Pri \$ | ice) Sul Per | osidised ✓ | Generic Manufacturer |
| GLUTEN FREE PASTA - Special Authority see SA1729 on the | previous page - H | lospital phar | macy [H | P3] |
| Buckwheat Spirals | | 250 g OP | | |
| | (3.11) | | C | Drgran |
| Corn and Vegetable Shells | 2.00 | 250 g OP | | |
| | (2.92) | | C | Drgran |
| Corn and Vegetable Spirals | 2.00 | 250 g OP | | |
| | (2.92) | | C | Drgran |
| Rice and Corn Lasagne Sheets | 1.60 | 200 g OP | | |
| | (3.82) | | C | Drgran |
| Rice and Corn Macaroni | | 250 g OP | | • |
| | (2.92) | • | C | Drgran |
| Rice and Corn Penne | | 250 g OP | | • |
| | (2.92) | - | C | Drgran |
| Rice and Maize Pasta Spirals | 2.00 | 250 g OP | | - |
| | (2.92) | - | C | Drgran |
| Rice and Millet Spirals | 2.00 | 250 g OP | | - |
| | (3.11) | • | C | Drgran |
| Rice and corn spaghetti noodles | 2.00 | 375 g OP | | - |
| | (2.92) | - | C | Drgran |
| Vegetable and Rice Spirals | | 250 g OP | | • |
| | (2.92) | • | C | Drgran |
| Italian long style spaghetti | | 220 g OP | | - |
| | (3.11) | • | C | Drgran |

Foods And Supplements For Inherited Metabolic Disease

⇒SA2300 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Dietary management of inherited metabolic disease; or
- 2 For use as a supplement to a Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

| AMINOACID FORMULA WITHOUT METHIONINE - | Special Authority see SA2300 | above - He | ospital pharmacy [HP3] |
|--|------------------------------|------------|----------------------------------|
| Powder, 12.5 g sachets | | 30 | HCU Explore 5 |
| Powder, 25 g sachets | 1,048.95 | 30 | HCU Express 15 |
| Powder | | 500 g OP | XMET Maxamum |

Supplements For MSUD and short chain enoyl coA hydratase deficiency

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE – Special Authority see SA2300 above – Hospital pharmacy [HP3]

| Powder, 12.5 g sachets | | 30 | MSUD Explore 5 |
|------------------------|----------|----------|-------------------------------------|
| Powder, 25 g sachets | 1,048.95 | 30 | MSUD Express 15 |
| Powder | | 500 g OP | MSUD Maxamum |

| | Subsidy (Manufacturer's Pric \$ | e) Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---------------------------------------|-----------|---------------------|-------------------------------------|
| Supplements For PKU | | | | |
| MINOACID FORMULA WITHOUT PHENYLALANINE – Spectarracy [HP3] | cial Authority see SA2 | 2300 on | the previo | <mark>us page</mark> – Hospital |
| Tabs | 99 00 | 75 OF | . . | Phlexy 10 |
| Powder (Lemon), 34 g sachets | | 30 | | PKU Express 20 |
| Powder (Neutral), 12.5 g sachets | | 30 | | PKU Explore 5 |
| Powder (Neutral), 34 g sachets | | 30 | | PKU Express 20 |
| Powder (Orange), 25 g sachets | | 30 | | PKU Explore 10 |
| Powder (Orange), 34 g sachets | | 30 | | PKU Express 20 |
| Powder (Raspberry), 25 g sachets | | 30 | | PKU Explore 10 |
| Powder (Tropical), 34 g sachets | | 30 | | PKU Express 20 |
| Powder (berry) 28 g sachets | | 30 | | PKU Lophlex |
| | | 00 | - | Powder |
| Powder (chocolate) 36 g sachet | 303.00 | 30 | 1 | PKU Anamix Junior |
| | | 50 | • | Chocolate |
| Dourder (noutral) 00 a cochata | 026.00 | 30 | | PKU Lophlex |
| Powder (neutral) 28 g sachets | | 30 | v | Pro Lopniex Powder |
| | | ~~ | | |
| Powder (neutral) 36 g sachets | | 30 | | PKU Anamix Junior |
| Powder (orange) 28 g sachets | | 30 | • | PKU Lophlex |
| | | | | Powder |
| Powder (orange) 36 g sachet | | 30 | ~ | PKU Anamix Junior |
| | | | | Orange |
| Powder (vanilla) 36 g sachet | | 30 | ✓ | PKU Anamix Junior |
| | | | | Vanilla |
| Infant formula | 174.72 | 400 g C | DP 🗸 | PKU Anamix Infant |
| Powder (neutral) | | 400 g C | | PKU Start |
| Powder (orange) | | 500 g C | DP 🗸 | XP Maxamum |
| Powder (unflavoured) | | 500 g C | DP 🗸 | XP Maxamum |
| Liquid (berry) | | 125 ml (| OP 🗸 | PKU Anamix Junior |
| | | | | LQ |
| Liquid (orange) | | 125 ml (| OP 🗸 | PKU Anamix Junior |
| | | | | LQ |
| Liquid (unflavoured) | | 125 ml (| OP 🗸 | PKU Anamix Junior |
| (| | | | LQ |
| Liquid (forest berries), 250 ml carton | 540.00 | 18 OF | | Easiphen Liquid |
| Liquid (juicy tropical) 125 ml | | 30 OF | | PKU Lophlex LQ 20 |
| Oral semi-solid (berries) 109 g | | 36 OF | | PKU Lophlex |
| | | 00 01 | - | Sensation 20 |
| Liquid (juicy berries) 62.5 ml | 030 00 | 60 OF | . . | PKU Lophlex LQ 10 |
| Liquid (juicy bernes) 62.5 ml | | 60 OF | | PKU Lophlex LQ 10 |
| Liquid (juicy crange) 62.5 ml | | 60 OF | | PKU Lophlex LQ 10 |
| Liquid (juicy berries) 125 ml | | 30 OF | | PKU Lophlex LQ 20 |
| | | 30 OF | | |
| Liquid (juicy orange) 125 ml | | 30 OF | • | PKU Lophlex LQ 20 |

SPECIAL FOODS

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|--|---|-------|---------------------|------------------------------|
| LYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOM | ME PHENYLALANINE | – Spe | cial Autho | ority see SA2300 on |
| age 285 – Hospital pharmacy [HP3] | | | | |
| Powder (Banana) 35 g sachets | 930.00 | 30 | 1 | PKU |
| | | | | sphere20 Banana |
| Powder (Berry), 20 g sachets | | 60 | ✓ | PKU Restore |
| | | | | Powder |
| Powder (Chocolate) 32 g Sachets | | 30 | ✓ | PKU Build |
| | | | | 20 Chocolate |
| Powder (Chocolate) 35 g sachets | | 30 | 1 | PKU |
| | | | | sphere20 Chocolate |
| Powder (Lemon) 35 g sachets | 930.00 | 30 | 1 | PKU |
| | | 00 | | sphere20 Lemon |
| Powder (Lemonade) 33.4 g sachets | 936.00 | 30 | 1 | PKU GMPro Ultra |
| | | 00 | • | Lemonade |
| Powder (Neutral), 16 g sachets | 449.28 | 30 | 1 | PKU Build 10 |
| Powder (Orange), 20 g sachets | | 60 | | PKU Restore |
| | | 00 | • | Powder |
| Powder (Raspberry Lemonade) 32 g Sachets | 898 56 | 30 | 1 | PKU Build |
| | | 00 | | 20 Raspberry |
| | | | | Lemonade |
| Powder (Smooth) 32 g Sachets | 898 56 | 30 | 1 | PKU Build |
| | | 00 | | 20 Smooth |
| Powder (Vanilla) 32 g Sachets | 898.56 | 30 | 1 | PKU Build 20 Vanilla |
| Powder (neutral), 40 g sachets | | 30 | | Camino Pro |
| · • · · · · · · · · · · · · · · · · · · | | | | Bettermilk |
| Powder (Red Berry) 35 g sachets | 930.00 | 30 | 1 | PKU sphere20 Red |
| | | 00 | - | Berry |
| Powder (Vanilla) 35 g sachets | 930.00 | 30 | 1 | PKU |
| | | 00 | • | sphere20 Vanilla |
| Liquid (Coffee Mocha), 250 ml carton | 684.45 | 30 OF | · · | PKU Glytactin RTD |
| | | 00 01 | • | 15 Lite |
| Liquid (chocolate), 250 ml carton | 684 45 | 30 OF | | PKU Glytactin RTD |
| בוקטוט (טוטטטומוש), בטט וווו טמונטוו | | 50 Or | • | 15 |
| Liquid (neutral), 250 ml carton | 684 45 | 30 OF | | PKU Glytactin RTD |
| בוקטוט (ווכטוומו), בטט וווו טמונטוו | | 50 Or | • | 15 |
| Liquid (vanilla), 250 ml carton | 694 45 | 30 OF | | |
| Liquiu (vanilia), 200 mii Ganon | 004.40 | 30 OF | v | PKU Glytactin RTD 15 Lite |

Foods

| LOW PROTEIN BAKING MIX – Special Authority see SA2300 or Powder | | | y [HP3] ✓ Loprofin Mix |
|--|------------------|---------------|------------------------------|
| LOW PROTEIN PASTA - Special Authority see SA2300 on page | e 285 – Hospital | pharmacy [HP: | 3] |
| Animal shapes | | 500 g OP | Loprofin |
| Lasagne | 6.19 | 250 g OP | Loprofin |
| Low protein rice pasta | | 500 g OP | Loprofin |
| Macaroni | 6.19 | 250 g OP | Loprofin |
| Penne | | 500 g OP | Loprofin |
| Spaghetti | | 500 g OP | Loprofin |
| Spirals | | 500 g OP | Loprofin |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-------------|------------------------|---|
| Supplements for Tyrosinaemia | | | | |
| AMINOACID FORMULA WITHOUT PHENYLALANINE AND TYR pharmacy [HP3] | OSINE – Special Au | thority | / see <mark>SA2</mark> | 300 on page 285 – Hospita |
| Powder (Neutral), 12.5 g sachets | | 30 | | TYR Explore 5 |
| GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME SA2300 on page 285 – Hospital pharmacy [HP3] | TYROSINE AND PH | ENYL | ALANINE | Special Authority see |
| Powder (Red Berry), 35 g sachets | 1,398.60 | 30 | 1 | TYR Sphere 20 |
| Powder (Vanilla), 35 g sachets | | 30 | | TYR Sphere 20 |
| Supplements for Organic Acidaemias | | | | |
| AMINOACID FORMULA WITHOUT METHIONINE, THREONINE Hospital pharmacy [HP3] | | cial Aı | uthority se | e SA2300 on page 285 – |
| Powder, 12.5 g sachets | | 30 | - | MMA/PA Explore 5 |
| Powder, 25 g sachets | 1,048.95 | 30 | • | MMA/PA Express 15 |
| Supplements for Glutaric Aciduria type 1 | | | | |
| AMINOACID FORMULA WITHOUT LYSINE - Special Authority s | | | | |
| Powder, 12.5 g sachets | 349.65 | 30 | v | GA Explore 5 |
| Supplements for Glycogen Storage Disease | | | | |
| HIGH AMYLOPECTIN CORN-STARCH – Special Authority see S Powder. 60 g sachets | | – Hos 30 | | macy [HP3] Glycosade |
| Fowder, 60 g sachets | 241.02 | 30 | • | diycosade |
| Single dose amino acids | | | | |
| ARGININE – Special Authority see SA2300 on page 285 – Hospit Powder, 4 g sachets | 211.45 | 30 | • | Arginine2000 |
| CITRULLINE – Special Authority see SA2300 on page 285 – Hos Powder, 4 g sachets | |] 30 | ~ (| Citrulline1000 |
| ISOLEUCINE – Special Authority see SA2300 on page 285 – Hos Powder, 4 g sachets | | 3] 30 | ✓ | Isoleucine50 |
| LEUCINE – Special Authority see SA2300 on page 285 – Hospita Powder, 4 g sachets | | 30 | ✓ | Leucine100 |
| PHENYLALANINE – Special Authority see SA2300 on page 285 - Powder, 4 g sachets | | [HP3] 30 | | Phenylalanine50 |
| TYROSINE – Special Authority see SA2300 on page 285 – Hospi Powder, 4 g sachets | | 30 | 1 | Tyrosine1000 |
| VALINE – Special Authority see SA2300 on page 285 – Hospital Powder, 4 g sachets | | 30 | ~ | Valine50 |

| S | Subsidy | | Fully | Brand or |
|---------|-----------------|-----|---------|----------|
| (Manufa | cturer's Price) | Sub | sidised | Generic |
| | \$ | Per | 1 | Manufact |

Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 vear where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| LOW CALCIUM INFANT FORMULA – Special Authority see | SA1110 above – Ho | spital pharmad | y [HP3] |
|--|-------------------|----------------|-----------------------------|
| Powder | | 400 g OP | Locasol |

Gastrointestinal and Other Malabsorptive Problems

| AMINO ACID FORMULA - Special Authority see SA2092 below - Hospital phan | macy [HP3] | |
|---|------------|---|
| Powder | 400 g OP | ✓ Alfamino ✓ Alfamino Junior |
| Powder (unflavoured)55.61 | 400 g OP | Neocate Gold Neocate Junior Unflavoured |
| 65.72 | | ✓ Neocate SYNEO ✓ Elecare ✓ Elecare LCP |
| Powder (vanilla)55.61 | 400 g OP | Neocate Junior Vanilla |
| 65.72 | | Elecare |

■ SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis: or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency: or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and

6.2 Fither:

- 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
- 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

continued...

| Subsidy | F | ully | Brand or |
|------------------------|--------|------|----------|
| (Manufacturer's Price) | Subsid | ised | Generic |
| \$ | Per | ✓ | |

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

| Subsidy | Fully Subsidised | | Brand or | |
|------------------------|---------------------|---|--------------|--|
| (Manufacturer's Price) | | | Generic | |
| \$ | Per | ✓ | Manufacturer | |

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

| ENTERAL LIQUID PEPTIDE FORMULA | Special Authority see SA1953 below | Hospital pharm | nacy [HP3] |
|--------------------------------|--|------------------------------------|--------------------|
| Linuial di Longi /mai | 10.14 | | All student Dariet |

| Liquid 1 kcal/ml | 12.44 | 500 ml OP | Nutrini Peptisorb |
|--------------------|-------|-----------|---------------------------------------|
| Liquid 1.5 kcal/ml | 18.66 | 500 ml OP | Nutrini Peptisorb |
| | | | Energy |

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure; or

2.10 Both:

2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

| Subsidy | Fully | | Brand or |
|------------------------|-------------------|---|----------|
| (Manufacturer's Price) | Price) Subsidised | | Generic |
| \$ | Per | 1 | |

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

| EXTENSIVELY HYDROLYSED FORMULA | - Special Authority see SA1557 be | low – Hospital pl | narmacy [HP3] |
|--------------------------------|-----------------------------------|-------------------|--------------------------------------|
| Powder | | 450 g OP | Pepti-Junior |
| | 36.20 | 900 g OP | Allerpro Syneo 1 |
| | | - | Allerpro Syneo 2 |

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

| PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML | - Special Authority see SA1698 | below | – Hospita | al pharmacy [HP3] |
|--|--------------------------------|-------|-----------|-------------------------------|
| Liquid | | | | Infatrini |

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

➡SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

| HIGH FAT LOW CARBOHYDRATE FORMULA - S | pecial Authority see SA1197 | above - Retail | pharmacy |
|---------------------------------------|-----------------------------|----------------|---------------------------------|
| Powder (unflavoured) | | 300 g OP | KetoCal 4:1 |
| | | | Ketocal 3:1 |
| Powder (vanilla) | | 300 g OP | KetoCal 4:1 |

| | Subsidy (Manufacturer's Price) \$ | Fully Subsidised Per ✓ | Brand or Generic Manufacturer |
|---|--|---|--|
| Vaccinations | | | |
| BACILLUS CALMETTE-GUERIN VACCINE – [Xpharm] For infants at increased risk of tuberculosis. Increased risk i 1) living in a house or family with a person with current or 2) having one or more household members or carers who equal to 40 per 100,000 for 6 months or longer; or 3) during their first 5 years will be living 3 months or longer Note a list of countries with high rates of TB are available at www.bcgatlas.org/index.php. Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial with diluent | past history of TB; or within the last 5 year er in a country with a r www.health.govt.nz/tu | rs lived in a count ate of TB > or eq uberculosis (sear | ual to 40 per 100,000 |
| DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE a) Only on a prescription b) No patient co-payment payable c) | | | |
| A) Funded for any of the following criteria: A single dose for pregnant women in the sect A single dose for parents or primary caregiver Specialist Care Baby Unit for more than 3 day 14 days prior to birth; or A course of up to four doses is funded for chil full primary immunisation; or An additional four doses (as appropriate) are stem cell transplantation or chemotherapy; pr dialysis and other severely immunosuppressi A single dose for vaccination of patients aged A single dose for vaccination of patients aged For vaccination of previously unimmunised or For revaccination following immunosuppressi For the patients with tetanus-prone w Notes: Please refer to the Immunisation Handbool Contractors will be entitled to claim payment from the vaccine to patients eligible under the above criteria for subsidised immunisation, and they may only do listed in the Pharmaceutical Schedule. Contractors may only claim for patient populations a sub-set of the population described in paragraphs Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous | rs of infants admitted t vs, who had not been dren from age 7 up to funded for (re-)immun e or post splenectomy ve regimens; or from 65 years old; or from 45 years old wh partially immunised p on; or bounds. a for appropriate scheo he Funder for the sup pursuant to their cont so in respect of the di within the criteria that | to a Neonatal Inte exposed to mate the age of 18 ye isation for patien r; pre- or post sol to have not had 4 vatients; or dule for catch up ply of diphtheria, tract with Health I iphtheria, tetanus | nsive Care Unit or rnal vaccination at least ars inclusive to complete ts post haematopoietic id organ transplant, renal previous tetanus doses; or programmes. tetanus and pertussis New Zealand (Health NZ) and pertussis vaccine |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) Funded for any of the following:
 - 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
 - A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
 - 3) An additional four doses (as appropriate) are funded for (re-)immunisation for people post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 4) Five doses will be funded for children requiring solid organ transplantation.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis and polio vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis and polio vaccine listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

poliomyelitis virus in 0.5ml syringe0.00 10

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE

a) Only on a prescription

I

- b) No patient co-payment payable
- c) A) Funded for children meeting any of the following criteria
 - 1) Up to four doses for children up to and under the age of 10 for primary immunisation; or
 - 2) An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of 10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 3) Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

| nj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg | | |
|--|----|---------------|
| pertussis toxoid, 25 mcg pertussis filamentous | | |
| haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus, | | |
| 10 mcg hepatitis B surface antigen in 0.5 ml syringe0.00 | 10 | Infanrix-hexa |
| | | |

| | Subsidy | Fully | Brand or |
|--|------------------------------|---------------------|----------------------------|
| | (Manufacturer's Price) \$ | Subsidised Per ✓ | Generic Manufacturer |
| HAEMOPHILUS INFLUENZAE TYPE B VACCINE | | | |
| a) Only on a prescription | | | |
| b) No patient co-payment payable | | | |
| c) A) One dose for people meeting any of the followin | g: | | |
| 1) For primary vaccination in children; or | • | | |
| 2) An additional dose (as appropriate) is fund | led for (re-)immunisatio | n for people pos | t haematopoietic stem cell |
| transplantation, or chemotherapy; function | al asplenic; pre or post | splenectomy; p | re- or post solid organ |
| transplant, pre or post cochlear implants, r | enal dialysis and other | severely immun | osuppressive regimens; or |
| For use in testing for primary immunodefic | iency diseases, on the | recommendation | n of an internal medicine |
| physician or paediatrician. | | | |
| B) Contractors will be entitled to claim payment from | | | |
| vaccine to people eligible under the above criter | | | |
| for subsidised immunisation, and they may only listed in the Pharmaceutical Schedule. | do so in respect of the | Haemophilus in | nuenzae type o vaccine |
| C) Contractors may only claim for populations withi | n the criteria that are co | overed by their c | contract which may be a |
| sub-set of the population described in paragraph | | | onitiaet, which may be a |
| | Trabove. | | |
| Haemophilus Influenzae type B polysaccharide 10 mcg | | | |
| conjugated to tetanus toxoid as carrier protein 20-40 m | cg; | | |
| prefilled syringe plus vial 0.5 ml | 0.00 | 1 🖌 | Hiberix |
| HEPATITIS A VACCINE – [Xpharm] | | | |
| Funded for patients meeting any of the following criteria: | | | |
| 1) Two vaccinations for use in transplant patients; or | | | |
| 2) Two vaccinations for use in children with chronic liver | disease; or | | |
| 3) One dose of vaccine for close contacts of known hepa | atitis A cases. | | |
| Inj 1440 ELISA units in 1 ml syringe | 0.00 | 1 🖌 | Havrix |

| Inj 1440 ELISA units in 1 mi syringe | 0.00 | 1 | • Havrix |
|---------------------------------------|------|---|-----------------------------------|
| Inj 720 ELISA units in 0.5 ml syringe | 0.00 | 1 | Havrix Junior |

| | Subsidy | 0 | Fully | Brand or |
|---|-----------------------------------|--------------|------------|------------------------------|
| | (Manufacturer's Price) \$ | Per | sidised | Generic Manufacturer |
| | Ψ | 1.01 | - | Manalastaron |
| HEPATITIS B RECOMBINANT VACCINE – [Xpharm] | 0.00 | | | n martin D |
| Inj 10 mcg per 0.5 ml prefilled syringe | | 1 | ♥ E | ngerix-B |
| Funded for patients meeting any of the following criteria: | | | | |
| for household or sexual contacts of known acute h | | | | s; or |
| for children born to mothers who are hepatitis B su | | | | |
| for children up to and under the age of 18 years in | | | | achieved a positive |
| serology and require additional vaccination or requ | ire a primary course c | of vaccina | tion; or | |
| for HIV positive patients; or | | | | |
| 5) for hepatitis C positive patients; or | | | | |
| for patients following non-consensual sexual intercontent | ourse; or | | | |
| for patients following immunosuppression; or | | | | |
| for solid organ transplant patients; or | | | | |
| for post-haematopoietic stem cell transplant (HSC⁻ | Γ) patients; or | | | |
| following needle stick injury. | | | | |
| | | | | |
| Inj 20 mcg per 1 ml prefilled syringe | | 1 | ✓ <u>E</u> | ngerix-B |
| Funded for patients meeting any of the following criteria: | | | | |
| for household or sexual contacts of known acute h | | | | s; or |
| for children born to mothers who are hepatitis B su | | | | |
| for children up to and under the age of 18 years in | | | | achieved a positive |
| serology and require additional vaccination or requ | ire a primary course c | of vaccina | tion; or | |
| for HIV positive patients; or | | | | |
| for hepatitis C positive patients; or | | | | |
| for patients following non-consensual sexual intercontent | ourse; or | | | |
| for patients following immunosuppression; or | | | | |
| 8) for solid organ transplant patients; or | | | | |
| 9) for post-haematopoietic stem cell transplant (HSC | F) patients; or | | | |
| 10) following needle stick injury; or | | | | |
| 11) for dialysis patients; or | | | | |
| 12) for liver or kidney transplant patients. | | | | |
| | | | | |
| HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 5 | 8) VACCINE [HPV] | | | |
| Maximum of 1 inj per prescription | | | | |
| b) Only on a prescription | | | | |
| No patient co-payment payable | | | | |
| d) | | | | |
| A) Any of the following: | | | | |
| Maximum of two doses for children aged | | | | |
| Maximum of three doses for patients me | | ring criteri | a: | |
| People aged 15 to 26 years inclusion | ve; or | | | |
| 2) Either: | | | | |
| People aged 9 to 26 years inclusiv | e | | | |
| Confirmed HIV infection; or | | | | |
| 2) Transplant (including stem ce | | | | |
| 3) Maximum of four doses for people aged | | | | |
| B) Contractors will be entitled to claim payment f | | | | |
| to patients eligible under the above criteria pu | | | | () |
| subsidised immunisation, and they may only o | to so in respect of the | Human p | apilloma | avirus vaccine listed in the |
| Pharmaceutical Schedule. | | | | |
| C) Contractors may only claim for patient populat | | | covered | by their contract, which |
| may be a sub-set of the population described | | | | |
| Inj 270 mcg in 0.5 ml syringe | 0.00 | 10 | ✓ <u>G</u> | ardasil 9 |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|-------------------------------------|
| INFLUENZA VACCINE Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine) | | 10 | ✔ Ir | nfluvac Tetra (2024 formulation) |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- d)

A) INFLUENZA VACCINE

is available each year for patients who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) Down syndrome, or
 - vii) are pregnant; or
- c) children 4 years of age and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- d) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- C)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.
- Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

10 Y Priorix

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- C)
- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or
 - 2) One dose for individuals who turn 13 years of age while living in boarding school hostels.
- C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A, C, Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
- D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. Ini 10 mcg of each meningococcal polysaccharide conjugated

300

| Sub | sidy | Fully I | Brand or |
|------------|------------------|-----------------------|--------------|
| (Manufactu | irer's Price) Su | ubsidised (| Generic |
| | B Per | I | Manufacturer |

MENINGOCOCCAL B MULTICOMPONENT VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) Any of the following:
 - A) Three doses for children up to 12 months of age (inclusive) for primary immunisation; or
 - B) Up to three doses (dependent on age at first dose) for a catch-up programme for children from 13 months to
 - 59 months of age (inclusive) for primary immunisation, from 1 March 2023 to 31 August 2025; or
 - C) Both:
 - 1) Person is one year of age or over; and
 - 2) Any of the following:
 - i) up to two doces and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
 - ii) up to two doses for close contacts of meningococcal cases of any group; or
 - iii) up to two doses for person who has previously had meningococcal disease of any group; or
 - iv) up to two doses for bone marrow transplant patients; or
 - v) up to two doses for person pre- and post-immunosuppression*; or
 - D) Both:
 - 1) Person is aged between 13 and 25 years (inclusive); and
 - 2) Either:
 - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prisons; or
 - ii) Two doses for individuals who turn 13 years of age while living in boarding school hostels.
 - E) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule.
 - F) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-D above.

*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

| Inj 175 mcg per 0.5 ml prefilled syringe | 0.00 | 1 | Bexsero |
|--|------|----|-----------------------------|
| | | 10 | Bexsero |

MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm]

Both:

- 1) The child is under 12 months of age; and
- 2) Any of the following:
 - 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) Two doses for close contacts of meningococcal cases of any group; or
 - 3) Two doses for child who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for child pre- and post-immunosuppression*.

Note: children under 12 months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 10 mcg in 0.5 ml syringe......0.00 1 🗸 Neisvac-C

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully idised | Brand or Generic Manufacturer |
|---|---|-------------|-----------------|-------------------------------------|
| PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpha | rm] | | | |
| 1) A primary course of three doses for previously unvace | cinated individuals up to | the age o | of 59 m | onths inclusive |
| Note: please refer to the Immunisation Handbook for the a | ppropriate schedule for | catch up | prograr | nmes |
| Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, | 6B, | | | |
| 7F, 9V, 14 and 23F; 3 mcg of pneumococcal | | | | |
| polysaccharide serotypes 4, 18C and 19F in 0.5 ml | | | | |
| prefilled syringe | 0.00 | 10 | 🗸 🗸 S | vnflorix |

| Subsidy Fu Manufacturer's Price) Subsidis | | Brand or Generic |
|--|-------|---------------------|
| \$ | Per 🗸 | Manufacturer |

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) Any of the following:
 - 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
 - Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
 - 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection; or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) cochlear implants or intracranial shunts; or
 - g) cerebrospinal fluid leaks; or
 - receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - I) diabetes; or
 - m) Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
 - 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
 - 5) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Pneumococcal (PCV13) conjugate vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Pneumococcal (PCV13) conjugate vaccine listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

| 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml | | |
|---|----|-------------|
| syringe0.00 | 10 | Prevenar 13 |
| | 1 | Prevenar 13 |

| | Subsidy (Manufacturer's Price) \$ | Fully Subsidised Per 🖌 | Brand or Generic Manufacturer |
|---|---|------------------------------|-------------------------------------|
| PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – Either: | [Xpharm] | | |
| Up to three doses (as appropriate) for patients with HI chemotherapy; pre- or post-splenectomy or with functi complement deficiency (acquired or inherited), cochle All of the following: | onal asplenia, pre- or p | oost-solid organ t | ransplant, renal dialysis, |
| a) Patient is a child under 18 years for (re-)immunisb) Treatment is for a maximum of two doses; andc) Any of the following: | sation; and | | |
| i) on immunosuppressive therapy or radiation immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; ov who are immune-suppressed following org | r | | |
| or vi) with cochlear implants or intracranial shunt vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more th prednisone of 2 mg/kg per day or greater, v 20 mg or greater; or | an two weeks, and wh | | |
| ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks ges xi) with cardiac disease, with cyanosis or failu xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with finance | tation; or re; or | gh-dose corticost | eroid therapy); or |
| Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype) POLIOMYELITIS VACCINE – [Xpharm] Up to three doses for patients meeting either of the following 1) For partially vaccinated or previously unvaccinated ind 2) For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for appro Inj 80D antigen units in 0.5 ml syringe | g: lividuals; or ppriate schedule for cat | _ | |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | 5 | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

ROTAVIRUS ORAL VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) Maximum of two doses for people meeting the following:
 - 1) first dose to be administered in infants aged under 14 weeks of age; and
 - 2) no vaccination being administered to children aged 24 weeks or over.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Rotavirus oral vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Rotavirus oral vaccine listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

| Oral susp live attenuated human rotavirus | | | |
|--|------|----|-----------------------------|
| 1,000,000 CCID50 per dose, squeezable tube | 0.00 | 10 | Rotarix |
| Oral susp live attenuated human rotavirus | | | |
| 1,000,000 CCID50 per dose, prefilled oral applicator | 0.00 | 10 | Rotarix |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

VARICELLA VACCINE [CHICKENPOX VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable
- c) A) Either:
 - 1) Maximum of one dose for primary vaccination for either:
 - a) Any infant born on or after 1 April 2016; or
 - b) For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
 - 2) Maximum of two doses for any of the following:
 - a) Any of the following for non-immune individuals:
 - i) with chronic liver disease who may in future be candidates for transplantation; or
 - ii) with deteriorating renal function before transplantation; or
 - iii) prior to solid organ transplant; or
 - iv) prior to any elective immunosuppression*; or
 - v) for post exposure prophylaxis who are immune competent inpatients; or
 - b) For individuals at least 2 years after bone marrow transplantation, on advice of their specialist; or
 - c) For individuals at least 6 months after completion of chemotherapy, on advice of their specialist; or
 - For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
 - e) For individuals with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
 - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
 - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
 - B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella vaccine [Chickenpox vaccine] vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella vaccine [Chickenpox vaccine] listed in the Pharmaceutical Schedule.
 - C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

| Inj 1350 PFU prefilled syringe | 0.0 |
|--------------------------------|-----|
|--------------------------------|-----|

| ✓ | Varivax |
|---|---------|
| ✓ | Varivax |

1

10

VARICELLA ZOSTER VACCINE [SHINGLES VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable
- C)
- A) Funded for patients meeting the following criteria:
 - 1) Two doses for all people aged 65 years
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella zoster vaccine (Shingles vaccine) to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella zoster vaccine [Shingles vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

| Inj 50 mcg per 0.5 ml vial plus vial | 0.00 | 1 | Shingrix |
|--------------------------------------|------|----|------------------------------|
| | | 10 | Shingrix |

| | Subsidy (Manufacturer's Price) \$ | Sub Per | Fully sidised | Brand or Generic Manufacturer |
|--|---|------------|------------------|-------------------------------------|
| Diagnostic Agents | | | | |
| TUBERCULIN PPD [MANTOUX] TEST – [Xpharm] Inj 5 TU per 0.1 ml, 1 ml vial | 0.00 | 1 | ✓ <u>⊺</u> | ubersol |

| - Symbols - |
|-------------|
|-------------|

| 3TC112 |
|---------------------------------------|
| 3TC112 7 MED NSHA Silver/Copper |
| Short |
| - A - |
| A-Scabies |
| Abacavir sulphate |
| Abacavir sulphate with |
| lamivudine 112 |
| Abacavir/Lamivudine Viatris |
| |
| Abilify Maintena |
| Abiraterone acetate 173 |
| Acarbose11 |
| Accarb |
| Accuretic 1047 |
| Accuretic 2047 |
| Acetazolamide262 |
| Acetec46 |
| Acetic acid with hydroxyquinoline and |
| ricinoleic acid |
| Acetylcysteine |
| Aci-Jel81 |
| Aciclovir |
| Infection107 |
| Sensory260 |
| Acidex |
| Acipimox54 |
| Acitretin74 |
| Actemra |
| Actinomycin D160 |
| Actrapid10 |
| Actrapid Penfill10 |
| Acupan |
| Adalimumab (Amgevita) |
| Adalimumab (Humira - Alternative |
| brand) 192 |
| Adapalene |
| Adapaiene |
| |
| ADR Cartridge 1.823 |
| Adrenaline Cardiovascular |
| |
| Respiratory |
| Advantan |
| Advate |
| Adynovate |
| Afinitor |
| Aflibercept199 |
| AFT-Pyrazinamide106 |
| After Hours Med Mgmt 15 min 265 |
| After Hours Med Mgmt 30 min 265 |
| After Hours Med Mgmt 45 min 265 |
| Agents Affecting the |
| Renin-Angiotensin System 46 |
| Agents for Parkinsonism and Related |

| Disorders Agents Used in the Treatment of | . 123 |
|--|-----------------|
| Agents Used in the Treatment of | |
| Poisonings | . 265 |
| Agrylin | 159 |
| Albendazole | 95 |
| Albey249 | -250 |
| Albustix | 83 |
| Alchemy Oxaliplatin | 155 |
| Alchemy Oxybutynin | 83 |
| Aldurazyme | 28 |
| Alecensa | 167 |
| Alectinib | 167 |
| Alendronate sodium | 117 |
| Alendronate sodium with | |
| colecalciferol | . 117 |
| Alfacalcidol | |
| Alfamino | 289 |
| Alfamino Junior | |
| Alginic acid | 6 |
| Alglucosidase alfa | 26 |
| Alkeran | 155 |
| Alkeran S29 | |
| Allerfix | 262 |
| Allerpro Syneo 1 | 292 |
| Allerpro Syneo 2 | |
| Allersoothe | 250 |
| Allmercap | |
| Allopurinol | |
| Alpha-Adrenoceptor Blockers | 46 |
| Alpha-Keri Lotion | 72 |
| Alphamox | 98 |
| Alphamox 125 | 98 |
| Alphamox 250 | |
| Alprolix | |
| Alu-Tab | |
| Aluminium hydroxide | 6 |
| Alvogen | 50 |
| Amantadine hydrochloride | 123 |
| Ambrisentan | 57 |
| Ambrisentan Viatris | 57 |
| Amgevita | 184 |
| Amiloride hydrochloride | 52 |
| Amiloride hydrochloride with | |
| furosemide | 53 |
| Amilarida hydrochlarida with | |
| hydrochlorothiazide | 53 |
| Aminophylline Amiodarone hydrochloride | 256 |
| Amiodarone hydrochloride | 48 |
| Amisulpride | |
| Amitriptyline | 129 |
| Amlodipine | |
| Amorolfine | |
| Amoxicillin | |
| Amoxicillin with clavulanic acid | <mark>98</mark> |
| | |

| Amphotericin B | 32 |
|---|------------|
| Amsacrine | . 159 |
| AmsaLyo | . 159 |
| Amsidine | . 159 |
| Amzoate | 30 |
| Anaesthetics | . 124 |
| Anafranil | . 129 |
| Anafranil Anagrelide hydrochloride | . 159 |
| Analgesics | . 125 |
| Anastrozole | . 176 |
| Anatrole | . 176 |
| Androderm | 86 |
| Anoro Ellipta | .254 |
| Antabuse | . 151 |
| Antacids and Antiflatulents | 6 |
| Anthelmintics | 95 |
| Antiacne Preparations | 68 |
| Antiallergy Preparations | 249 |
| Antianaemics | |
| Antiandrogen Oral | |
| Contraceptives | 81 |
| Antiarrhythmics | |
| Antibacterials | 95 |
| Antibacterials Topical | |
| Anticholinergic Agents | |
| Anticholinesterases | 116 |
| Antidepressants | 129 |
| Antidiarrhoeals | |
| Antiepilepsy Drugs | 121 |
| Antifibrinolytics, Haemostatics and | |
| Local Sclerosants | 37 |
| Antifibrotics | 255 |
| Antifungals | 102 |
| Antifungals Topical | 60 |
| Antihistamines | 250 |
| Antihypotensives | <u>200</u> |
| Antimalarials | 104 |
| Antimigraine Preparations | 134 |
| Antinausea and Vertigo Agents | 134 |
| Antipruritic Preparations | 70 |
| Antipsychotics | 136 |
| Antiretrovirals | 110 |
| Antirheumatoid Agents | 117 |
| Antispasmodics and Other Agents | |
| Altering Gut Motility | 8 |
| Antithrombotic Agents | 40 |
| Апшинопроце Адения | 40 |
| Antithymocyte alohulin | 18/ |
| Antithymocyte globulin | 104 |
| (equine) | |
| (equine) Antitrichomonal Agents | |
| (equine) Antitrichomonal Agents Antituberculotics and | 105 |
| (equine) Antitrichomonal Agents Antituberculotics and Antileprotics | |
| (equine) Antitrichomonal Agents Antituberculotics and Antileprotics Antiulcerants | 9 |
| (equine) Antitrichomonal Agents Antituberculotics and Antileprotics | 9 107 |

| Anxiolytics140 |
|---|
| Anzatax |
| Apidra |
| Apidra SoloStar11 |
| |
| APO-Atomoxetine 145 |
| APO-Atomoxetine S29 145 |
| Apo-Azithromycin |
| Apo-Bicalutamide 174 |
| APO-Candesartan HCTZ |
| 16/12.5 |
| APO-Candesartan HCTZ |
| 32/12.5 |
| Apo-Temozolomide 165 |
| Apo-Terbinafine 104 |
| Apomorphine hydrochloride123 |
| Aprepitant134 |
| Apresoline57 |
| Aptamil Feed Thickener284 |
| Aqueous cream72 |
| Aratac |
| Arava117 |
| Arginine |
| Arginine2000 |
| Aripiprazole136–137 |
| Aripiprazole Sandoz |
| Aristocort |
| Arrotex-Prazosin S29 |
| Arrow - Clopid |
| Arrow - Lattim |
| Arrow-Amitriptyline |
| Arrow-Bendrofluazide |
| Arrow-Brimonidine |
| Arrow-Diazepam |
| Arrow-Doxorubicin |
| Arrow-Fluoxetine |
| |
| Arrow-Losartan & Hydrochlorothiazide |
| Hydrochlorothlazide |
| Arrow-Norfloxacin |
| Arrow-Ornidazole |
| Arrow-Quinapril 1047 |
| Arrow-Quinapril 2047 |
| Arrow-Quinapril 547 |
| Arrow-Roxithromycin |
| Arrow-Timolol |
| Arrow-Topiramate133 |
| Arrow-Tramadol129 |
| Arsenic trioxide 159 |
| Asacol7 |
| Asacol S297 |
| Ascend Aripiprazole136 |
| Ascend-Cefuroxime |
| Ascorbic acid32 |
| Aspen Adrenaline56 |
| Aspirin |
| Blood40 |
| Nervous125 |

| Asthalin253 |
|---|
| Atazanavir Mylan 112 |
| Atazanavir sulphate 112 |
| Atazanavir Viatris112 |
| Atenolol |
| Atenolol AFT |
| Atenolol AFT S29 |
| Atenolol Viatris |
| Atezolizumab |
| ATGAM |
| Ativan |
| Atnahs Olsalazine |
| |
| Atomoxetine |
| Atorvastatin |
| Atropine sulphate |
| Cardiovascular |
| Sensory263 |
| Atropt |
| Atrovent |
| Aubagio141 |
| Augmentin |
| Aurorix |
| AutoSoft 3023 |
| AutoSoft 9023 |
| Avallon126 |
| Avelox100 |
| Avonex141 |
| Avonex Pen 141 |
| Azacitidine |
| Azacitidine Dr Reddy's 155 |
| Azamun |
| Azathioprine |
| Azilect |
| Azithromycin |
| Azopt |
| AZOpt |
| -B- |
| - р - B-D Micro-Fine15 |
| B-D Ultra Fine |
| |
| B-D Ultra Fine II16 Bacillus Calmette-Guerin (BCG) |
| Bacilius Calmette-Guerin (BCG) |
| vaccine |
| Bacillus Calmette-Guerin |
| vaccine |
| Baclofen121 |
| Bactroban68 |
| Balance |
| Barrier Creams and Emollients71 |
| BCG Vaccine 294 |
| Beclazone 100251 |
| Beclazone 250251 |
| Beclazone 50251 |
| Beclomethasone dipropionate |
| Bedaquiline105 |
| Bee venom allergy treatment249 |
| Bendamustine hydrochloride 153 |
| • |

| Bendrofluazide | 53 |
|--|---|
| Bendroflumethiazide | |
| [Bendrofluazide] | |
| Benralizumab | |
| Benzathine benzylpenicillin | |
| Benzatropine mesylate | 123 |
| Benzbromaron AL 100 | 120 |
| Benzbromarone | |
| Benztrop | |
| Benzydamine hydrochloride | 31 |
| Benzylpenicillin sodium [Penicillin | |
| G] | |
| Beta Cream | |
| Beta Ointment | |
| Beta Scalp | 76 |
| Beta-Adrenoceptor Agonists | 252 |
| Beta-Adrenoceptor Blockers | 49 |
| Betadine | 73 |
| Betadine Skin Prep | 73 |
| Betaferon | 141 |
| Betahistine dihydrochloride | 134 |
| Betaine | |
| Betamethasone dipropionate | 70 |
| Betamethasone dipropionate with | |
| calcipotriol | 74 |
| Betamethasone sodium phosphat | е |
| with betamethasone acetate | 85 |
| Betamethasone valerate | 70, 76 |
| | |
| Betamethasone valerate with sodi | um |
| fusidate [fusidic acid] | 71 |
| fusidate [fusidic acid] Betaxolol | 71 262 |
| fusidate [fusidic acid] Betaxolol Betnovate | 71 262 70 |
| fusidate [fusidic acid] Betaxolol Betnovate Betoptic | 71 262 70 262 |
| fusidate [fusidic acid] Betaxolol Betnovate Betoptic Betoptic S | 71 262 70 262 262 |
| fusidate [fusidic acid] Betaxolol Betnovate Betoptic Betoptic S Bexsero | 71 262 70 262 262 301 |
| fusidate [fusidic acid] Betaxolol Betnovate Betoptic Betoptic S Bexsero Bezafibrate | 71 262 70 262 262 301 54 |
| fusidate [fusidic acid] Betaxolol Betnovate Betoptic Betoptic S Bexsero Bezafibrate Bezalip | 71 262 70 262 262 301 54 54 |
| fusidate [fusidic acid] Betaxolol Betoptic. Betoptic S Bexsero Bezafibrate Bezalip. Bezalip Retard. | 71 262 70 262 262 301 54 54 54 |
| fusidate [fusidic acid] Betaxolol Betoptic. Betoptic S Bexsero Bezafibrate Bezalip Retard Bicalutamide | 71 262 70 262 262 301 54 54 54 54 |
| fusidate [fusidic acid] Betaxolol Betoptic. Betoptic S Bexsero Bezafibrate Bezalip Retard. Bicalutamide Bicillin LA | 71 262 70 262 301 54 54 54 174 98 |
| fusidate [fusidic acid] Betaxolol Betoptic Betoptic S Bexsero Bezafibrate Bezalip Retard Bicalutamide Bicillin LA BiCNU | 71 262 70 262 301 54 54 54 174 98 154 |
| fusidate [fusidic acid] Betaxolol Betnovate Betoptic. Betoptic S Bezafibrate Bezalip Retard Bicalutamide Bicalutamide BicNU Bile and Liver Therapy | 71 70 262 262 301 54 54 174 98 154 10 |
| fusidate [fusidic acid] Betaxolol Betoptic Betoptic S Bexsero Bezafibrate Bezalip Retard Bicalutamide Bicillin LA BiCNU Bile and Liver Therapy Biltricide | 71 262 262 262 301 54 54 54 174 98 154 95 |
| fusidate [fusidic acid] Betaxolol Betoptic Betoptic S Bexsero Bezafibrate Bezalip Retard Bicalutamide Bicillin LA Bile and Liver Therapy Biltricide Bimatoprost | 71 262 262 262 301 54 54 54 174 98 154 10 95 95 952 |
| fusidate [fusidic acid] Betaxolol Betoptic Betoptic S Bexsero Bezafibrate Bezalip Retard Bicalutamide Bicillin LA Bile and Liver Therapy Biltricide Bimatoprost Multichem | 71 262 70 262 301 54 54 54 174 98 154 98 154 95 262 262 |
| fusidate [fusidic acid] Betaxolol Betoptic Betoptic S Bezafibrate Bezalip Bezalip Bezalip Bezalip Bezalip Bezalip Bezalip Bezalip Bezalip Bezalip Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bitricide Bimatoprost Bimatoprost Multichem Binarex | 71 262 70 262 301 54 54 54 54 174 98 154 95 262 262 262 174 |
| fusidate [fusidic acid] Betaxolol Betoptic Betoptic S Bezafibrate Bezalip. Bezalip Retard Bicalutamide Bicillin LA Bile and Liver Therapy Biltricide Bimatoprost Multichem Binarex Binocrit | 71 262 70 262 301 54 54 54 54 174 98 154 95 262 262 262 262 262 301 |
| fusidate [fusidic acid]BetaxololBetopticBetoptic SBetoptic SBexseroBezafibrateBezalip.RetardBicalutamideBicalutamideBicNUBile and Liver TherapyBiltricideBimatoprost MultichemBinarexBinocritBiodone | |
| fusidate [fusidic acid] Betaxolol Betoptic Betoptic S Bezafibrate Bezalip Bezalip Bezalip Retard Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Binarex Binarex Binocrit Biodone Biodone Extra Forte | |
| fusidate [fusidic acid] Betaxolol Betoptic. Betoptic S Bexsero Bezafibrate Bezalip Retard. Bicalutamide Bicillin LA BiCNU. Bile and Liver Therapy. Biltricide Bimatoprost Bimatoprost Multichem Binarex Biodone Extra Forte. Biodone Forte | |
| fusidate [fusidic acid]Betaxolol Betaxolol Betoptic Betoptic S Bezafibrate Bezafibrate Bezalip Retard Bicalutamide Bicalutamide Bicalutamide BicNU Bile and Liver Therapy Biltricide Bimatoprost Multichem Binatoprost Multichem Binarex Biodone Biodone Extra Forte Biodone Forte Bisacodyl | |
| fusidate [fusidic acid]Betaxolol Betaxolol Betoptic Betoptic S Bexsero Bezafibrate Bezalip Retard Bicalutamide Bicalutamide Bicalutamide BicAlutamide Binacon Biodone Extra Forte Bisacodyl Bisacodyl Viatris | |
| fusidate [fusidic acid] Betaxolol Betoptic Betoptic S Bexsero Bezafibrate Bezalip Retard Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bisacodyl Bisacodyl Viatris Bisoprolol fumarate | |
| fusidate [fusidic acid]Betaxolol Betaxolol Betoptic Betoptic S Bezafibrate Bezafibrate Bezalip Retard Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bile and Liver Therapy Bile and Liver Therapy Binatoprost Binatoprost Biodone Extra Forte Bisacodyl Viatris Bisoprolol fumarate BK Lotion | |
| fusidate [fusidic acid] Betaxolol Betoptic Betoptic S Bexsero Bezafibrate Bezalip Retard Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bicalutamide Bisacodyl Bisacodyl Viatris Bisoprolol fumarate | |

| Factors43 |
|-------------------------------------|
| Blood glucose diagnostic test |
| meter 14 |
| Blood glucose diagnostic test |
| strip |
| Blood glucose test strips (visually |
| impaired)15 |
| Blood Ketone Diagnostic Test |
| Strip 13 |
| Sup |
| Bonjela |
| |
| Bortezomib |
| Bosentan60 |
| Bosentan Dr Reddy's60 |
| Bplex |
| Brentuximab Vedotin |
| Breo Ellipta252 |
| Brevinor 1/28 80 |
| Brevinor-1 28 Day80 |
| Bricanyl Turbuhaler253 |
| Brilinta40 |
| Brimonidine tartrate263 |
| Brimonidine tartrate with timolol |
| maleate 263 |
| Brinzolamide262 |
| Brown & Burk 130 |
| Brufen SR 116 |
| BSF Max Health265 |
| BSF Methylphenidate ER - |
| Teva |
| BSF Noumed Dexamfetamine |
| Buccastem135 |
| Budesonide |
| Alimentary |
| Respiratory251, 258 |
| Budesonide Te Arai |
| Budesonide with eformoterol |
| Bumetanide |
| Buprenorphine Naloxone BNM 150 |
| Buprenorphine with naloxone |
| Bupropion hydrochloride |
| Burel |
| Burinex |
| Burinex S29 |
| Buscopan |
| Buspirone hydrochloride140 |
| Buspirone Viatris140 |
| |
| Busulfan154 |
| - C - Cabergoline93 |
| Caffeine citrate |
| |
| Calamine |
| Calci-Tab 500 |
| Calcipotriol |
| Calcitonin |
| Calcitriol |

| Calcitriol-AFT | 33 |
|------------------------------------|----------------------|
| Calcium 500 mg Hexal | 34 |
| Calcium carbonate6, | 34 |
| Calcium carbonate PAI | . 6 |
| Calcium Channel Blockers | 51 |
| Calcium Disodium Versenate | 967 |
| Calcium folinate 1 | |
| Calcium Folinate Ebewe1 | 56 |
| Calcium Folinate Sandoz1 | 56 |
| Calcium Folinate Sandoz S29 | |
| Calcium gluconate | |
| Calcium Homeostasis | 84 |
| Calcium polystyrene sulphonate | |
| Calcium Resonium | 44 |
| Calogen | |
| Camber | |
| Camino Pro Bettermilk2 | 987 |
| Candesartan cilexetil | 47 |
| Candesartan cilexetil with | |
| hydrochlorothiazide | 47 |
| Candestar | 47 |
| Canesten | |
| Capecitabine1 | |
| Capecitabine Viatris1 | |
| Capsaicin | 00 |
| Musculoskeletal1 | 17 |
| Nervous1 | |
| Captopril | |
| Carafate | 10 |
| Carbaccord 1 | |
| Carbamazepine1 | |
| Carbimazole | |
| Carbomer | |
| Carboplatin 1 | 54 |
| Carboplatin Ebewe 1 | 54 |
| Carbosorb-X | 880 |
| Cardinol LA | |
| Cardizem CD | |
| CareSens Dual | 14 |
| CareSens N 14- | |
| CareSens N POP | 14 |
| CareSens N Premier | 14 |
| CareSens PRO | 15 |
| Carmellose sodium with gelatin and | 10 |
| pectin | 31 |
| Carmustine1 | |
| Carnitor | |
| Carvedilol | |
| Carvedilol Sandoz | |
| Casirivimab and imdevimab | |
| | |
| Catapres1 | 52 |
| | |
| Cofaclor monohydrato | 55 |
| Cefaclor monohydrate | 55 95 |
| Cefaclor monohydrate Cefalexin | 55 95 95 |
| Cefaclor monohydrate | 55 95 95 95 |

| | _ |
|---|--|
| Cefazolin-AFT | 95 |
| Ceftriaxone | 95 |
| Ceftriaxone-AFT | |
| Cefuroxime axetil | 95 |
| Celapram 13 | 30 |
| Celebrex11 | 16 |
| Celecoxib11 | |
| Celecoxib Pfizer11 | 16 |
| Celestone Chronodose | 35 |
| Cellcept17 | 77 |
| Centrally-Acting Agents | 51 |
| Cephalexin ABM | 95 |
| Cetirizine hydrochloride 25 | 50 |
| Cetomacrogol | 72 |
| Cetomacrogol with glycerol | 72 |
| Cetomacrogol-AFT | 72 |
| Cetuximab20 | |
| Charcoal | |
| Chemotherapeutic Agents 15 | |
| Chickenpox vaccine |)6 |
| Chlorambucil15 | 54 |
| Chloramphenicol26 | |
| Chlorothiazide | |
| Chlorpheniramine maleate25 | |
| Chlorpromazine hydrochloride13 | 36 |
| Chlorsig | 50 |
| Chlortalidone [Chlorthalidone] | |
| Chlorthalidone | 53 |
| | |
| Chlorvescent | |
| Chlorvescent Choice 380 7med Nsha Silver/copper | 14 |
| Chlorvescent Choice 380 7med Nsha Silver/copper Short | 14 79 |
| Chlorvescent Choice 380 7med Nsha Silver/copper Short | 14 79 79 |
| Chlorvescent Choice 380 7med Nsha Silver/copper Short7 Choice Load 3757 Choice TT380 Short7 | 14 79 79 79 |
| Chlorvescent Choice 380 7med Nsha Silver/copper Short | 14 79 79 79 |
| Chlorvescent Choice 380 7med Nsha Silver/copper Short | 14 79 79 79 79 |
| Chlorvescent Choice 380 7med Nsha Silver/copper Short | 14 79 79 79 79 |
| Chlorvescent | 14 79 79 79 79 79 32 |
| Chlorvescent | 44 79 79 79 79 32 45 |
| Chlorvescent | 44 79 79 79 79 79 79 79 79 79 79 79 79 79 |
| Chlorvescent | 44 79 79 79 79 79 32 45 46 99 34 |
| Chlorvescent | 44 79 79 79 79 32 45 46 99 34 |
| Chlorvescent | 44 79 79 79 79 32 45 46 99 34 |
| Chlorvescent | 14 79 79 79 79 32 15 16 99 34 34 00 |
| Chlorvescent | 44 79 < |
| Chlorvescent | 14 79 79 79 32 15 16 99 34 34 00 00 60 |
| Chlorvescent | 14 79 79 79 79 32 15 16 99 34 34 00 50 50 |
| Chlorvescent | 14 79 79 79 79 32 15 16 99 34 34 00 50 00 50 |
| Chlorvescent | 14 79 70 70 |
| Chlorvescent | 14 79 79 79 79 325 469 340 500 500 500 545 |
| Chlorvescent | 14 79 79 79 79 79 325 469 345 39 345 345 345 345 345 345 345 345 345 345 345 345 345 345 345 345 345 346 346 347 348 349 340 350 350 354 354 357 358 359 350 350 350 350 350 350 350 350 350 350 350 350 350 350 |
| Chlorvescent | 144 79 79 79 79 32 15 6 99 34 15 6 99 34 46 99 34 46 99 93 4 45 60 00 00 00 00 00 00 00 00 00 00 00 00 |
| Chlorvescent | 144 79 79 79 79 32 15 6 99 34 15 6 99 34 46 99 34 46 99 93 4 45 60 00 00 00 00 00 00 00 00 00 00 00 00 |
| Chlorvescent | 14 79 79 79 79 32 15 60 99 34 34 00 00 00 00 00 00 00 00 00 00 00 00 00 |
| Chlorvescent | 14 79 79 79 79 32 15 6 99 34 34 00 00 00 00 00 00 00 00 00 00 00 00 00 |
| Chlorvescent | 14 79 79 79 32 15 16 99 34 34 30 00 00 00 00 00 00 00 00 00 00 00 00 |

| Clexane Forte41 |
|-----------------------------------|
| Climara87 |
| Clindamycin100 |
| Clinicians |
| Clinicians Renal Vit |
| Clobazam |
| Clobetasol propionate |
| Clobetasone butyrate |
| Clofazimine |
| |
| Clomazol |
| Dermatological |
| Genito-Urinary81 |
| Clomifene citrate94 |
| Clomipramine - Teva 129 |
| Clomipramine hydrochloride129 |
| Clomipramine Teva 129 |
| Clonazepam 131, 140 |
| Clonidine |
| Clonidine hydrochloride |
| Clonidine Teva |
| Clopidogrel |
| Clopine |
| Clopixol |
| Clotrimazole |
| Dermatological |
| Genito-Urinary81 |
| |
| Clozapine |
| Clozaril |
| Clustran |
| Co-trimoxazole |
| Coal tar |
| Coal tar with allantoin, menthol, |
| phenol and sulphur75 |
| Coal tar with salicylic acid and |
| sulphur75 |
| Cobal-B12 |
| Cobalin-H32 |
| Coco-Scalp75 |
| Codeine phosphate |
| Extemporaneous269 |
| Nervous |
| Coenzyme Q1027 |
| Colchicine |
| Colecalciferol |
| Colestid |
| Colestipol hydrochloride |
| Colestyramine |
| Colectivering Mulen E4 |
| Colestyramine - Mylan |
| Colgout |
| Colifoam |
| Colistin sulphomethate |
| Colistin-Link100 |
| Collodion flexible |
| Colloidal bismuth subcitrate9 |
| Colofac8 |
| Coloxyl |

| Combigan | 263 |
|------------------------------------|-----------|
| Compliance Packaging | 265 |
| Compound electrolytes | . 44 |
| Compound electrolytes with glucose | |
| [Dextrose] | . 44 |
| Compound hydroxybenzoate | 269 |
| Comtan | |
| Concerta | |
| Condoms | 78 |
| Condyline | |
| Contraceptives - Hormonal | 79 |
| Contraceptives - Non-hormonal | 77 |
| Copaxone | |
| Cordarone-X | . 48 |
| Corticosteroids and Related Agents | |
| for Systemic Use | . 85 |
| Corticosteroids Topical | 70 |
| Cosentyx | |
| Cosmegen | |
| Coumadin | 43 |
| Country Life | |
| Coversyl | |
| Creon 10000 | |
| Creon 25000 | |
| Creon Micro | |
| Crotamiton | |
| Crystaderm | 68 |
| Curam | |
| Curam Duo 500/125 | |
| Cvite | |
| Cyclizine hydrochloride | |
| Cyclizine lactate | |
| Cyclogyl | |
| Cyclonex | |
| Cyclopentolate hydrochloride | 263 |
| Cyclophosphamide | |
| Cyclorin | |
| Cycloserine | |
| Cyklokapron | |
| Cyproterone acetate | |
| Cyproterone acetate with | 00 |
| ethinyloestradiol | Q1 |
| Cystadane | 01 |
| Cytarabine | 156 |
| Cytotec | |
| Cytoxan | |
| - D - | 104 |
| D-Penamine | 117 |
| | 117 |
| Dabigatran | |
| Dacarbazine Dacarbazine APP | 100 |
| Dactinomycin [Actinomycin D] | 100 |
| | |
| Daivobet | |
| Daivonex | |
| Daktarin | |
| Dalacin C | 100 |

| Dantrium 1 | 21 |
|--------------------------------------|----|
| Dantrium S291 | |
| Dantrolene 1 | 21 |
| Daonil | 11 |
| Dapa-Tabs | 53 |
| Dapsone | 06 |
| Daraprim1 | 01 |
| Darunavir1 | 13 |
| Darunavir Viatris1 | 13 |
| Dasatinib1 | |
| Daunorubicin1 | 60 |
| Daunorubicin Zentiva1 | |
| David One Step Cassette Pregnancy | 00 |
| Test | 80 |
| DBL Adrenaline | 56 |
| DBL Aminophylline2 | |
| DBL Bleomycin Sulfate1 | |
| | |
| DBL Bortezomib1 | |
| DBL Carboplatin1 | 04 |
| DBL Cisplatin1 | |
| DBL Dacarbazine1 | 60 |
| DBL Desferrioxamine Mesylate for Inj | ~~ |
| BP2 | |
| DBL Docetaxel1 | 60 |
| DBL Ergometrine | 81 |
| DBL Gemcitabine1 | 5/ |
| DBL Gentamicin1 | 00 |
| DBL Heparin Sodium | 42 |
| DBL Leucovorin Calcium 1 | 56 |
| DBL Methotrexate Onco-Vial1 | |
| DBL Pethidine Hydrochloride1 | 28 |
| DBL Vincristine Sulfate1 | |
| Decozol | 32 |
| Deferasirox2 | 66 |
| Deferiprone | 66 |
| Deferoxamine Pfizer S292 | 66 |
| Denosumab1 | 17 |
| Deolate 1 | 04 |
| Deoxycoformycin 1 | |
| Depo-Medrol | 86 |
| Depo-Provera | |
| Depo-Testosterone | 86 |
| Deprim 1 | 02 |
| Dermol70, | 76 |
| Desferrioxamine mesilate2 | |
| Desmopressin | 93 |
| Desmopressin acetate | |
| Desmopressin-PH&T | |
| Desuric1 | 20 |
| Detection of Substances in | |
| Urine | 83 |
| Dexamethasone | |
| Hormone | 85 |
| Sensory2 | |
| Dexamethasone phosphate | 85 |
| Dexamethasone with framycetin and | |

| gramicidin 260 |
|--|
| Dexamethasone with neomycin |
| sulphate and polymyxin B |
| sulphate 261 |
| Dexamfetamine sulfate |
| Dexmethsone |
| Dextrochlorpheniramine |
| maleate |
| |
| Dextrose |
| DHC Continus127 |
| Diabetes 10 |
| Diabetes Management13 |
| Diacomit133 |
| Diagnostic Agents |
| Diamide Relief6 |
| Diamox |
| Diasip274 |
| Diazepam 131, 140 |
| Diazoxide10 |
| Dibenzyline |
| Diclofenac Sandoz116 |
| Diclofenac sodium |
| Musculoskeletal |
| Sensory |
| Differin |
| Difflam |
| |
| Diflucan |
| Digestives Including Enzymes |
| Digoxin |
| Dihydrocodeine tartrate 127 |
| Dilantin132 |
| Dilantin Infatab132 |
| Dilantin Paediatric132 |
| Diltiazem CD Clinect51 |
| Diltiazem hydrochloride51 |
| Dimethicone71, 73 |
| Dimethyl fumarate 141 |
| Dipentum8 |
| Diphtheria, tetanus and pertussis |
| vaccine |
| Diphtheria, tetanus, pertussis and |
| polio vaccine |
| Diphtheria, tetanus, pertussis, polio, |
| hepatitis B and haemophilus |
| influenzae type B vaccine |
| |
| Diprosone |
| Diprosone OV |
| Dipyridamole |
| Disopyramide phosphate |
| Disulfiram151 |
| Diuretics |
| Docetaxel160 |
| Docetaxel Accord 160 |
| Docetaxel Sandoz 160 |
| Docusate sodium24 |
| Docusate sodium with |

| sennosides 25 |
|--|
| Dolutegravir 113 |
| Dolutegravir with lamivudine113 |
| Domperidone 135 |
| Domperidone Viatris 135 |
| Donepezil hydrochloride 149 |
| Donepezil-Rex149 |
| Dornase alfa256 |
| Dortimopt |
| Dorzolamide with timolol |
| Dostinex |
| Dosulepin [Dothiepin] |
| hydrochloride |
| Dosulepin Mylan |
| Dosulepin Viatris |
| Dothiepin |
| Dovato |
| Doxazosin |
| Doxazosin Clinect |
| Doxazosin Cilinect |
| Doxorubicin Ebewe |
| Doxorubicin Ebewe |
| Doxorubiciii nyurochionde |
| Doxycycline |
| DP Lotin HC |
| DP-Allopurinol |
| DP-Captopril |
| Dr Reddy's Omeprazole |
| Di neuuy s Omepiazoie |
| Drofoto |
| Drofate |
| Drugs Affecting Bone |
| Drugs Affecting Bone Metabolism 117 |
| Drugs Affecting Bone Metabolism |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 diagnostic test meter 14 Dulaglutide 12 Dulcolax SP Drop 26 Duocal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 253 DuoResp Spiromax 252 |
| Drugs Affecting Bone Metabolism |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 Dulaglutide 12 Dulcolax SP Drop 26 Duocal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 253 Duordes Spiromax 252 Durde 56 Durvalumab 241 |
| Drugs Affecting Bone Metabolism |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 diagnostic test meter 14 Dulaglutide 12 Dulcolax SP Drop 26 Duocal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 252 Duride 56 Durvalumab 241 - E - e-chamber La Grande 259 e-chamber Turbo 259 E-Mycin 97 e5 Pharma 10 |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 diagnostic test meter 14 Dulaglutide 12 Dulcolax SP Drop 26 Duocal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 253 DuoResp Spiromax 252 Duride 56 Durvalumab 241 - E - e-chamber La Grande 259 e-chamber Mask 258 e-chamber Turbo 259 E-Mycin 97 e5 Pharma 10 Ear Preparations 260 |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 Dulaglutide 12 Dulcolax SP Drop 26 Duocal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 252 Duride 56 Durvalumab 241 - E - - e-chamber La Grande 259 e-chamber Mask 258 e-chamber Turbo 259 E-Mycin 97 e5 Pharma 10 Ear Preparations 260 Ear/Eye Preparations 260 |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 Dulaglutide 12 Dulcolax SP Drop 26 Duocal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 253 Durde 56 Durvalumab 241 - E - - e-chamber La Grande 259 e-chamber Turbo 259 E-Mycin 97 e5 Pharma 10 Ear Preparations 260 Ear/Eye Preparations 260 Easiphen Liquid 286 |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 Dulaglutide 12 Dulcolax SP Drop 26 Duocal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 253 Durkesp Spiromax 252 Duride 56 Durvalumab 241 - E - - e-chamber La Grande 259 E-Mycin 97 e5 Pharma 10 Ear /Peparations 260 Easiphen Liquid 286 Econazole nitrate 69 |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 Dulaglutide 12 Dulcolax SP Drop 26 Duocal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 253 Durde 56 Durvalumab 241 - E - - e-chamber La Grande 259 e-chamber Turbo 259 E-Mycin 97 e5 Pharma 10 Ear Preparations 260 Ear/Eye Preparations 260 Ear/Eye Preparations 260 Eaconazole nitrate 69 Efavirenz 111 |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 Dulaglutide 12 Dulcoal SP Drop 26 Duccal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 252 Durde 56 Durvalumab 241 - E - - e-chamber La Grande 259 e-chamber Turbo 259 E-Mycin 97 e5 Pharma 10 Ear Preparations 260 Ear/Eye Preparations 260 Easiphen Liquid 286 Econazole nitrate 69 Efavirenz 111 |
| Drugs Affecting Bone 117 Metabolism 117 Dual blood glucose and blood ketone 14 Dulaglutide 12 Dulcolax SP Drop 26 Duocal Super Soluble Powder 272 Duolin 253 Duolin HFA 253 Duolin Respules 253 Durde 56 Durvalumab 241 - E - - e-chamber La Grande 259 e-chamber Turbo 259 E-Mycin 97 e5 Pharma 10 Ear Preparations 260 Ear/Eye Preparations 260 Ear/Eye Preparations 260 Eaconazole nitrate 69 Efavirenz 111 |

| Eformoterol fumarate dihydrate | 251 |
|---|-----|
| Eftrenonacog alfa [Recombinant factor IX] | 27 |
| | |
| Efudix | |
| Egopsoryl TA | |
| Elaprase | |
| Elecare | 289 |
| Elecare LCP | |
| Electral | |
| Elelyso | 30 |
| Elemental 028 Extra | 277 |
| Elexacaftor with tezacaftor, ivacaftor | or |
| and ivacaftor | 257 |
| Elidel | 75 |
| Elocon | |
| Elocon Alcohol Free | 71 |
| Eltrombopag | |
| Eltroxin | |
| EMB Fatol | |
| Emend Tri-Pack | |
| Emicizumab | |
| EMLA | |
| Empagliflozin | 13 |
| Empagliflozin with metformin | 10 |
| hydrochloride | 13 |
| Emtricitabine | 110 |
| Emtricitabine with tenofovir | 112 |
| disoproxil | 100 |
| | |
| Emtriva | |
| Emulsifying ointment | |
| Emulsifying Ointment ADE | 72 |
| Enalapril maleate | |
| Enbrel | 177 |
| Endocrine Therapy | 173 |
| Endoxan | 154 |
| Engerix-B | |
| Enlafax XR | |
| Enoxaparin sodium | |
| Enstilar | |
| Ensure | |
| Ensure Plus | 282 |
| Ensure Plus HN | 281 |
| Ensure Plus HN RTH | 281 |
| Ensure Two Cal HN RTH | 283 |
| Entacapone | |
| Entecavir | |
| Entecavir (Rex) | 107 |
| Entresto 24/26 | 48 |
| Entresto 49/51 | |
| Entresto 97/103 | |
| Entyvio | |
| Epilim | |
| Epilim Crushable | |
| Epilim IV | |
| Epilim S/F Liquid | 133 |
| Epilim Syrup | |
| | |

| Epipen |
|----------------------------------|
| |
| Epipen Jr |
| Epirubicin Ebewe160 |
| Epirubicin hydrochloride 160 |
| Eplerenone52 |
| Epoetin alfa |
| Epoprostenol |
| Eptacog alfa [Recombinant factor |
| VIIa] |
| |
| Erbitux |
| Ergometrine maleate |
| Erlotinib168 |
| Erythrocin IV97 |
| Erythromycin (as lactobionate)97 |
| Erythromycin ethyl succinate97 |
| Esbriet |
| Escitalopram130 |
| Escitalopram (Ethics) |
| |
| Eskazole |
| Essential Ethosuximide |
| Essential Generics |
| Essential Prednisolone8 |
| Estraderm MX87 |
| Estradiol Sandoz87 |
| Estradiol TDP Mylan87 |
| Estradiol Viatris |
| Estradot |
| Estrofem |
| Etanercept |
| Ethambutol hydrochloride |
| |
| Ethics Aspirin |
| Ethics Aspirin EC40 |
| Ethics Lisinopril46 |
| Ethinyloestradiol with |
| desogestrel79 |
| Ethinyloestradiol with |
| levonorgestrel |
| Ethinyloestradiol with |
| norethisterone |
| Ethosuximide |
| |
| Etopophos |
| Etoposide |
| Etoposide phosphate160 |
| Etravirine 111 |
| Eumovate70 |
| Eurofolic156 |
| Evara |
| EVARA White Soft Paraffin72 |
| Everet 132 |
| Everolimus |
| Evista |
| |
| Evrysdi |
| Evusheld |
| Exelon Patch 10 |
| Exelon Patch 5 149 |
| Exemestane176 |

| Exjade266 |
|------------------------------------|
| Extemporaneously Compounded |
| Preparations and |
| Galenicals |
| Eve Bronarationa 200 |
| Eye Preparations |
| Eylea |
| Ezemibe Viatris56 |
| Ezetimibe |
| Ezetimibe Sandoz56 |
| Ezetimibe with simvastatin56 |
| - F - |
| Factor eight inhibitor bypassing |
| fraction |
| Famotidine9 |
| Famotidine Hovid9 |
| Fasenra200 |
| Faslodex174 |
| Fatty Cream AFT72 |
| Febuxostat121 |
| Febuxostat (Teva) |
| Febuxostat multichem |
| FEIBA NF |
| Felo 10 ER51 |
| Felo 5 ER51 |
| Felo 5 ER |
| Felodipine51 |
| Fenpaed 100 mg per 5 ml 116 |
| Fentanyl |
| Fentanyl Sandoz 127 |
| Ferinject34 |
| Ferodan |
| Ferriprox |
| Ferro-F-Tabs34 |
| Ferro-Liquid34 |
| Ferro-tab34 |
| Ferrograd34 |
| Ferrosig35 |
| Ferrous fumarate |
| Ferrous fumarate with folic acid34 |
| Ferrous sulfate34 |
| Fexofenadine hydrochloride250 |
| Fibro-vein |
| Filgrastim43 |
| Finasteride |
| Fingolimod141 |
| Firazyr |
| Flagyl |
| FlagyI-S 105 |
| ••• |
| Flamazine |
| Flecainide acetate |
| Flecainide BNM |
| Flecainide Controlled Release |
| Teva |
| Flecatab |
| Fleet Phosphate Enema25 |
| Flixonase Hayfever & Allergy |
| Flixotide251 |

| Flixotide Accuhaler | |
|--|---------------------------------|
| Florinef | |
| Fluanxol | |
| Flucil | 99 |
| Flucloxacillin | |
| Flucloxacillin-AFT | |
| Flucloxin | |
| Flucon | |
| Fluconazole | 102 |
| Fludara Oral | |
| Fludarabine Ebewe | 156 |
| Fludarabine phosphate | 156 |
| Fludrocortisone acetate | 85 |
| Fluids and Electrolytes | |
| Flumetasone pivalate | 260 |
| Fluocortolone caproate with | |
| fluocortolone pivalate and | |
| cinchocaine | 8 |
| Fluorometholone | 261 |
| Fluorouracil | 157 |
| Fluorouracil Accord | |
| Fluorouracil sodium | 76 |
| Fluox | |
| Fluoxetine hydrochloride | 130 |
| Flupenthixol decanoate | 138 |
| Flutamide | 174 |
| Flutamin | 174 |
| Fluticasone | 251 |
| Fluticasone furoate with | |
| umeclidinium and vilanterol | 254 |
| Fluticasone furoate with | |
| vilanterol | 252 |
| Fluticasone propionate | 258 |
| Fluticasone with salmeterol | |
| Flynn | 95 |
| FML | |
| Foban | <mark>69</mark> |
| Folic acid | 37 |
| Folic Acid multichem | 37 |
| Folic Acid Viatris | 37 |
| Food Thickeners | 283 |
| Foods And Supplements For | |
| Inherited Metabolic Disease | 285 |
| Forteo | |
| Fortini | |
| Fortini Multi Fibre | 276 |
| Fortisip | 282 |
| Fortisip Multi Fibre | 282 |
| Fosamax | 117 |
| Fosamax Plus | |
| Framycetin sulphate | 117 |
| i i danij oo din o dipilato initilitii ili | 260 |
| Frebini Energy | 260 276 |
| Frebini Energy Frebini Energy Fibre | 260 276 276 |
| Frebini Energy Frebini Energy Fibre Frebini Original | 260 276 276 276 |
| Frebini Energy Frebini Energy Fibre | 260 276 276 276 276 |

INDEX: Generic Chemicals and Brands

| Fresubin HP Energy281 |
|------------------------------------|
| Fresubin HP Energy Fibre |
| Fresubin Intensive |
| Fresubin Original |
| Fresubin Original Fibre |
| Frisium131 |
| Frumil53 |
| Frusemide52 |
| Fucicort71 |
| Fucidin 101 |
| Fucithalmic260 |
| Fulvestrant174 |
| Fungilin32 |
| Furosemid-Ratiopharm52 |
| Furosemide [Frusemide]52 |
| Furosemide-Baxter |
| fusidic acid |
| Dermatological 69, 71 |
| Infection |
| Sensory260 |
| - G - |
| GA Explore 5 |
| Gabapentin131 |
| Gacet |
| Galsulfase27 |
| Galvumet12 |
| Galvus12 |
| Gardasil 9 |
| Gastrodenol9 |
| Gaviscon Extra Strength6 |
| Gaviscon Infant6 |
| Gazyva212 |
| Gefitinib169 |
| GEM Aqueous Cream72 |
| Gemcitabine Ebewe 157 |
| Gemcitabine hydrochloride157 |
| Gemtuzumab ozogamicin203 |
| Gentamicin sulphate 100 |
| Gilenya141 |
| Ginet81 |
| Glatiramer acetate 141 |
| Glecaprevir with pibrentasvir 109 |
| Glibenclamide11 |
| Gliclazide11 |
| Glipizide11 |
| Glizide11 |
| Glucagen Hypokit 10 |
| Glucagon hydrochloride10 |
| Glucerna Select 274 |
| Glucose [Dextrose]43 |
| Gluten Free Foods284 |
| Glycerin with sodium saccharin 269 |
| Glycerin with sucrose269 |
| Glycerol |
| Alimentary25 |
| Extemporaneous269 |

| Glyceryl trinitrate | |
|--------------------------------|-------|
| Alimentary | 8 |
| Cardiovascular | 56 |
| Glycopyrronium | |
| Glycopyrronium bromide | |
| Glycopyrronium with | |
| indacaterol | 254 |
| Glycosade | |
| Gold Knight | 78 |
| Gold Knight XL | |
| Goserelin | |
| Gynaecological Anti-infectives | |
| -H- | |
| Habitrol | 151 |
| Haemophilus influenzae type B | |
| vaccine | 296 |
| Haldol | |
| Haldol Concentrate | 138 |
| Haldol Decanoas | |
| Haloperidol | |
| Haloperidol decanoate | 138 |
| Harvoni | |
| Havrix | |
| Havrix Junior | |
| Haylor syrup | |
| HCU Explore 5 | 285 |
| HCU Express 15 | 285 |
| healthE Aqueous Cream SLS | |
| Free | 72 |
| healthE Calamine Aqueous | |
| healthE Dimethicone 10% | 71 |
| healthE Dimethicone 4% Lotion | 73 |
| healthE Dimethicone 5% | 71 |
| healthE Glycerol BP | |
| healthE Urea Cream | |
| Healtheries Simple Baking Mix | .284 |
| Hemastix | |
| Hemlibra | |
| Heparin sodium | |
| Heparin Sodium Panpharma | |
| Heparinised saline | |
| Heparon Junior | |
| Hepatitis A vaccine | .296 |
| Hepatitis B recombinant | |
| vaccine | . 297 |
| Herceptin | |
| Herzuma | |
| Hiberix | |
| Hiprex | . 115 |
| Histaclear | |
| Histafen | |
| Holoxan | |
| Horleys Bread Mix | |
| Horleys Flour | |
| Hormone Replacement Therapy - | |
| Systemic | |
| 0,0001110 | |

| HPV | 297 |
|---|--|
| Humalog | . 11 |
| Humalog Mix 25 | . 11 |
| Humalog Mix 50 | |
| Human papillomavirus (6, 11, 16, 18, | |
| 31, 33, 45, 52 and 58) vaccine | |
| [HPV] | 297 |
| Humatin | |
| Humira | |
| HumiraPen | |
| Humulin 30/70 | |
| Humulin NPH | |
| Humulin R | |
| Hyaluronic acid | 264 |
| Hydralazine | |
| Hydralazine hydrochloride | |
| Hydralyte - Lemonade | |
| Hydrocortisone | |
| Dermatological | 70 |
| Hormone | |
| Hydrocortisone acetate | |
| Hydrocortisone acetate with | ••••• |
| pramoxine hydrochloride | 7 |
| Hydrocortisone and paraffin liquid | |
| and lanolin | 71 |
| Hydrocortisone butyrate71 | 76 |
| Hydrocortisone with cinchocaine | |
| Hydrocortisone with miconazole | |
| Hydrocortisone with natamycin and | |
| neomycin | |
| | . 71 |
| Hvdrogen peroxide | . 71 . 68 |
| Hydrogen peroxide | . 68 |
| Hydrogen peroxide Hydroxocobalamin | 68 32 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma | . 68 . 32 . 32 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma hydroxycarbamide | 68 32 32 160 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma | 68 32 32 160 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma hydroxycarbamide Hydroxychloroquine Hydroxyurea | 68 32 32 160 117 |
| Hydrogen peroxide Hydroxocobalamin Panpharma hydroxycarbamide Hydroxychloroquine Hydroxyurea [hydroxycarbamide] | 68 32 32 160 117 160 |
| Hydrogen peroxide Hydroxocobalamin Panpharma hydroxycarbamide Hydroxychloroquine Hydroxyurea [hydroxycarbamide] Hygroton | 68 32 160 117 160 53 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma hydroxycarbamide Hydroxychloroquine Hydroxyurea [hydroxycarbamide] Hygroton Hylo-Fresh | 68 32 160 117 160 53 264 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma hydroxycarbamide Hydroxychloroquine Hydroxyurea [hydroxycarbamide] Hygroton Hylo-Fresh Hymenoptera | 68 32 160 117 160 53 264 -250 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma hydroxycarbamide Hydroxycarbamide Hydroxyurea [hydroxycarbamide] Hygroton Hylo-Fresh Hymenoptera | 68 32 32 160 117 160 53 264 250 8 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma hydroxycarbamide Hydroxychloroquine Hydroxyurea [hydroxycarbamide] Hygroton Hylo-Fresh Hymoptera Hyoscine butylbromide Hyoscine hydrobromide Hyoscine hydrobromide | 68 32 160 117 160 53 264 250 8 135 120 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma hydroxycarbamide Hydroxychloroquine Hydroxyurea [hydroxycarbamide] Hygroton Hylo-Fresh Hymoptera Hyoscine butylbromide Hyoscine hydrobromide Hyoscine hydrobromide | 68 32 160 117 160 53 264 250 8 135 120 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Panpharma hydroxycarbamide Hydroxycarbamide Hydroxyurea [hydroxyurea [hydroxycarbamide] Hygroton Hylo-Fresh Hymenoptera | 68 32 160 1177 160 53 264 250 8 135 120 263 |
| Hydrogen peroxide | 68 32 32 160 117 160 53 264 250 8 135 120 263 263 |
| Hydrogen peroxide Hydroxocobalamin Hydroxocobalamin Hydroxycarbamide Hydroxycarbamide Hydroxyurea [hydroxyurea [hydroxycarbamide] Hygroton Hylo-Fresh Hymoptera 249- Hyoscine butylbromide Hyoscine hydrobromide Hyperuricaemia and Antigout Hypromellose Hypromellose with dextran | 68 32 32 160 117 160 53 264 250 8 135 120 263 263 |
| Hydrogen peroxide | 68 32 32 160 117 160 53 264 250 8 135 120 263 263 98 |
| Hydrogen peroxide | 68 32 160 117 160 53 264 250 8 135 120 263 263 263 263 170 160 |
| Hydrogen peroxide | 68 32 160 117 160 53 264 250 8 135 120 263 263 263 263 170 160 116 |
| Hydrogen peroxide | 68 32 32 160 117 160 53 264 250 8 135 120 263 263 120 263 170 160 116 249 |
| Hydrogen peroxide | 68 32 32 160 117 160 53 264 250 8 135 120 263 263 120 263 170 160 116 249 |
| Hydrogen peroxide | 68 32 160 117 160 53 264 250 8 135 120 263 263 170 160 116 249 161 28 |
| Hydrogen peroxide | 68 32 160 117 160 53 264 250 8 135 120 263 263 263 98 170 160 116 249 161 28 154 |
| Hydrogen peroxide | 68 32 32 160 117 160 53 264 250 8 135 120 263 263 263 170 160 116 249 161 28 154 262 |

| Imatinib mesilate169 |
|------------------------------------|
| Imatinib-Rex169 |
| Imbruvica160 |
| Imfinzi |
| Imipramine Crescent 129 |
| Imipramine hydrochloride 129 |
| Imiguimod |
| Immune Modulators113 |
| Immunisation Administration |
| Immunisation |
| Co-administration |
| Immunosuppressants |
| Incruse Ellipta |
| Indacaterol |
| Indapamide |
| Infanrix IPV |
| |
| Infanrix-hexa |
| Infant Formulae |
| Infatrini |
| Infliximab |
| Influenza vaccine |
| Influvac Tetra |
| (2024 formulation) 298 |
| Inhaled Corticosteroids251 |
| Inhaled Long-acting |
| Beta-adrenoceptor Agonists 251 |
| Inresa35 |
| Inspra52 |
| Instillagel Lido 124 |
| Insulin aspart11 |
| Insulin aspart with insulin aspart |
| protamine10 |
| Insulin glargine11 |
| Insulin glulisine11 |
| Insulin isophane10 |
| Insulin isophane with insulin |
| neutral |
| Insulin lispro11 |
| Insulin lispro with insulin lispro |
| protamine |
| Insulin neutral |
| Insulin pen needles |
| Insulin pump |
| Insulin pump cartridge |
| |
| Insulin pump infusion set (steel |
| cannula)21 |
| Insulin pump infusion set (steel |
| cannula, straight insertion) 21 |
| Insulin pump infusion set (teflon |
| cannula)22 |
| Insulin pump infusion set (teflon |
| cannula, angle insertion with |
| insertion device) 23 |
| Insulin pump infusion set (teflon |
| cannula, straight insertion with |
| insertion device) 23 |
| |

| nsulin pump reservoir | 23 |
|--|---------------|
| nsulin syringes, disposable with | |
| attached needle | 16 |
| ntelence | |
| nterferon beta-1-alpha | |
| nterferon beta-1-beta | 141 |
| ntra-uterine device | 79 |
| nvega Sustenna | 138 |
| nvega Trinza | 139 |
| pca-Allopurinol | 120 |
| pca-Bisoprolol | 49 |
| pca-Donepezil | 1/0 |
| pca-Escitalopram | |
| PCA-Frusemide | 100 |
| PCA-Metoprolol | |
| PCA-Propranolol | |
| POL | |
| pratropium bromide253 | |
| |), ∠00 100 |
| ressa | |
| rinotecan Actavis 100 rinotecan hydrochloride | 15/ |
| rinotecan nyarochioride | 15/ |
| rinotecan-Rex | 15/ |
| ron (as ferric carboxymaltose) | |
| ron polymaltose | 35 |
| sentress | 113 |
| sentress HD | 113 |
| smo 20 | 56 |
| smo 40 Retard | 56 |
| soleucine50 | 288 |
| soniazid | 106 |
| soniazid with rifampicin | |
| soptin | 51 |
| soptin Retard | 51 |
| soptin SR | 51 |
| sopto Carpine | |
| sosorbide mononitrate | 56 |
| sosource Standard | 281 |
| sotretinoin | 68 |
| spaghula (psyllium) husk | 24 |
| tch-Soothe | 70 |
| traconazole | 103 |
| trazole | |
| vacaftor | 257 |
| vermectin | 73 |
| - J - | |
| Jadelle | 80 |
| Jakavi | |
| Jardiamet | |
| Jardiance | |
| Jaydess | |
| Jevity HiCal RTH | 281 |
| Jevity Plus RTH | |
| Jevity RTH | 281 |
| Jinarc | |
| Juno Pemetrexed | 158 |
| - K - | |
| | |

| Kadcyla236 |
|----------------------------------|
| Kalydeco257 |
| Kemadrin 123 |
| Kenacomb260 |
| Kenacort-A 1086 |
| Kenacort-A 4086 |
| Kenalog in Orabase |
| Ketocal 3:1 |
| KetoCal 4:1 |
| Ketoconazole |
| Dermatological |
| Infection |
| Ketogenic Diet |
| Ketoprofen 116 |
| KetoSens13 |
| Keytruda |
| Kindergen |
| • |
| Klacid |
| Alimentary9 |
| Infection |
| Kliogest |
| Kliovance |
| Kogenate FS |
| Konakion MM40 |
| Konsyl-D24 |
| Kuvan 29 |
| -L- |
| Labetalol50 |
| Lacosamide 131 |
| Lactulose25 |
| Laevolac25 |
| Lagevrio110 |
| Lamictal |
| Lamivudine 107, 112 |
| Lamivudine Viatris 112 |
| Lamivudine/Zidovudine Viatris112 |
| Lamotrigine |
| Lamprene 105 |
| Lanoxin |
| Lanoxin Paediatric Elixir |
| Lanoxin PG48 |
| Lanoxin S2948 |
| Lansoprazole9 |
| Lantus |
| Lantus SoloStar11 |
| Lanvis 159 |
| Lanzol Relief9 |
| Largactil |
| Laronidase |
| Lasix |
| Latanoprost |
| Latanoprost with timolol |
| Lax-Suppositories |
| Lax-suppositories Glycerol |
| Lax-suppositones Gryceror |
| Laxauves |
| Langel |

| Ledipasvir with sofosbuvir 109 |
|---|
| Leflunomide 117 |
| Lenalidomide 161 |
| Letrole176 |
| Letrozole176 |
| Leucine100288 |
| Leukeran FC 154 |
| Leukotriene Receptor |
| Antagonists |
| Leuprorelin93 |
| Leustatin 156 |
| Levetiracetam 132 |
| Levetiracetam-AFT132 |
| Levocabastine |
| Levocarnitine |
| Levodopa with benserazide123 |
| Levodopa with carbidopa123 |
| Levomepromazine 136 |
| Levomepromazine |
| hydrochloride 137 |
| Levonorgestrel |
| Genito-Urinary80-81 |
| Hormone |
| Levonorgestrel BNM |
| Levothyroxine |
| Lidocaine [Lignocaine] |
| Lidocaine [Lignocaine] hydrochloride |
| |
| hydrochloride |
| Lidocaine [Lignocaine] with |
| Lidocaine [Lignocaine] with prilocaine |
| Lidocaine [Lignocaine] with prilocaine 125 Lidocaine-Baxter 124 124 Life Extension 30 30 Lignocaine 124-125 124 Linezolid 106 106 Lioresal Intrathecal 121 121 Lipid-Modifying Agents 54 121 |
| Lidocaine [Lignocaine] with prilocaine 125 Lidocaine-Baxter 124 125 Life Extension 30 30 Lignocaine 124–125 124–125 Linezolid 106 Lioresal Intrathecal 121 Lipid-Modifying Agents 54 Liquigen 273 Liraglutide 12 12 12 |
| Lidocaine [Lignocaine] with prilocaine 125 Lidocaine-Baxter 124 125 Lidocaine-Baxter 124 125 Lignocaine 124-125 126 Linezolid 106 106 Lioresal Intrathecal 121 121 Liquigen 273 54 Liquigen 121 12 Lisnopril 46 46 |
| Lidocaine [Lignocaine] with 125 prilocaine 124 Lidocaine-Baxter 124 Life Extension 30 Lignocaine 124–125 Linezolid 106 Lioresal Intrathecal 121 Lipid-Modifying Agents 54 Liquigen 273 Liraglutide 12 Lisinopril 46 Litak 156 |
| Lidocaine [Lignocaine] with 125 prilocaine 124 Lide Extension 30 Lignocaine 124-125 Linezolid 106 Lioresal Intrathecal 121 Lipid-Modifying Agents 54 Liquigen 273 Liraglutide 12 Lisinopril 46 Litak 156 Lithium carbonate 137 |
| Lidocaine [Lignocaine] with 125 prilocaine 124 Life Extension 30 Lignocaine 124–125 Linezolid 106 Lioresal Intrathecal 121 Lipid-Modifying Agents 54 Liquigen 273 Liraglutide 12 Lisinopril 46 Litak 156 Lithium carbonate 137 Livostin 261 |
| Lidocaine [Lignocaine] with 125 prilocaine 124 Life Extension 30 Lignocaine 124-125 Linezolid 106 Lioresal Intrathecal 121 Lipid-Modifying Agents 54 Liquigen 273 Liraglutide 12 Litak 156 Lithium carbonate 137 Livostin 261 LMX4 125 |
| Lidocaine [Lignocaine] with 125 prilocaine 124 Life Extension 30 Lignocaine 124-125 Linezolid 106 Lioresal Intrathecal 121 Lingen 273 Liraglutide 12 Lisinopril 46 Litak 156 Lithium carbonate 137 Livostin 261 LMX4 125 Lo-Oralcon 20 ED 80 |
| Lidocaine [Lignocaine] with 125 prilocaine 124 Life Extension 30 Lignocaine 124-125 Linezolid 106 Lioresal Intrathecal 121 Lipid-Modifying Agents 54 Liquigen 273 Liraglutide 12 Litak 156 Lithium carbonate 137 Livostin 261 LMX4 125 Lo-Oralcon 20 ED 80 Locacorten-Viaform ED's 260 |
| Lidocaine [Lignocaine] with 125 prilocaine 124 Life Extension 30 Lignocaine 124-125 Linezolid 106 Lioresal Intrathecal 121 Lingen 273 Liraglutide 12 Lisinopril 46 Litak 156 Lithium carbonate 137 Livostin 261 LMX4 125 Lo-Oralcon 20 ED 80 |
| Lidocaine [Lignocaine] with prilocaine 125 Lidocaine-Baxter 124 125 Life Extension 30 30 Lignocaine 124–125 121 Linezolid 106 106 Lioresal Intrathecal 121 121 Lipid-Modifying Agents 54 124 Lizinopril 124 125 Litak 156 137 Litvostin 261 144 LMX4 125 260 Locacorten-Viaform ED's 260 260 Local preparations for Anal and Rectal Disorders 88 289 |
| Lidocaine [Lignocaine] with prilocaine 125 Lidocaine-Baxter 124 125 Life Extension 30 30 Lignocaine 124–125 121 Linezolid 106 106 Lioresal Intrathecal 121 121 Lipid-Modifying Agents 54 124 Ligaigen 273 127 Liraglutide 12 125 Litak 156 137 Livostin 261 137 Livostin 261 80 Locacorten-Viaform ED's 260 Local preparations for Anal and 8 Locasol 289 Loccoid Locasol 289 Locoid |
| Lidocaine [Lignocaine] with prilocaine 125 Lidocaine-Baxter 124 125 Life Extension 30 30 Lignocaine 124–125 121 Linezolid 106 106 Lioresal Intrathecal 121 121 Lipid-Modifying Agents 54 124 Ligaigen 273 127 Liraglutide 12 125 Lisinopril 46 46 Litak 156 137 Livostin 261 137 Livostin 261 80 Locacorten-Viaform ED's 260 260 Local preparations for Anal and Rectal Disorders 88 Locasol 289 Locoid 71, 76 Locoid Crelo 71 71 71 |
| Lidocaine [Lignocaine] with prilocaine 125 Lidocaine-Baxter 124 125 Life Extension 30 30 Lignocaine 124–125 124–125 Linezolid 106 106 Lioresal Intrathecal 121 121 Lipid-Modifying Agents 54 124 Liraglutide 121 121 Liraglutide 121 123 Liraglutide 122 124 Litak 156 115 Lithium carbonate 137 125 Lo-Oralcon 20 ED 80 100 Locacorten-Viaform ED's 260 260 Local preparations for Anal and Rectal Disorders 8 Locasol 289 1000 289 Locoid Crelo 71 76 100 71 |
| Lidocaine [Lignocaine] with 125 prilocaine 124 Life Extension 30 Lignocaine 124-125 Linezolid 106 Lioresal Intrathecal 121 Lignotifying Agents 54 Liquigen 273 Liraglutide 12 Lisinopril 46 Litak 156 Lithium carbonate 137 Livostin 261 LMX4 125 Lo-Oralcon 20 ED 80 Locacorter-Viaform ED's 260 Locaid 71, 76 Locoid Crelo 71 Locoid Crelo 71 Locorten-Vioform 260 |
| Lidocaine [Lignocaine] with prilocaine 125 Lidocaine-Baxter 124 125 Life Extension 30 30 Lignocaine 124–125 124–125 Linezolid 106 106 Lioresal Intrathecal 121 121 Lipid-Modifying Agents 54 124 Liquigen 273 124 Lisinopril 46 126 Litak 156 137 Livostin 261 137 Livostin 261 100 Locacorten-Viaform ED's 260 Locacorten-Viaform Sfor Anal and 8 Locasol 289 10 Locoid Crelo 71 76 Locoid Crelo 71 71 Locorten-Vioform 260 10 Locoxid Lipocream 71 71 Locoxid Crelo 71 71 Locoxid Crelo 71 71 |
| Lidocaine [Lignocaine] with 125 prilocaine 124 Life Extension 30 Lignocaine 124 Life Extension 30 Lignocaine 124 Life Extension 30 Lignocaine 124 Linezolid 106 Lioresal Intrathecal 121 Lipid-Modifying Agents 54 Liquigen 273 Liraglutide 12 Lisinopril 46 Litak 156 Lithium carbonate 137 Livostin 261 LMX4 125 Lo-Oralcon 20 ED 80 Locacorten-Viaform ED's 260 Locasol 289 Locoid 71 Locoid Crelo 71 Locoid Crelo 71 Locoid Crelo 71 Locoid Lipocream 71 Lodoxamide 261 Logem 132 |
| Lidocaine [Lignocaine] with prilocaine 125 Lidocaine-Baxter 124 125 Life Extension 30 30 Lignocaine 124–125 124–125 Linezolid 106 106 Lioresal Intrathecal 121 121 Lipid-Modifying Agents 54 124 Liquigen 273 124 Lisinopril 46 126 Litak 156 137 Livostin 261 137 Livostin 261 100 Locacorten-Viaform ED's 260 Locacorten-Viaform Sfor Anal and 8 Locasol 289 10 Locoid Crelo 71 76 Locoid Crelo 71 71 Locorten-Vioform 260 10 Locoxid Lipocream 71 71 Locoxid Crelo 71 71 Locoxid Crelo 71 71 |

| Loniten57 |
|----------------------------------|
| Loperamide hydrochloride6 |
| Lopinavir with ritonavir113 |
| Lopinavir/Ritonavir Mylan 113 |
| Loprofin |
| Loprofin Mix |
| Lorafix |
| Loratadine |
| Lorazepam140 |
| Lorstat |
| Losartan Actavis |
| Losartan potassium |
| Losartan potassium with |
| hydrochlorothiazide |
| Lovir |
| Lucrin Depot 1-month |
| |
| Lucrin Depot 3-month |
| Lyrica |
| - M - |
| m-Eslon |
| Mabthera |
| Macrobid115 |
| Macrogol 3350 with potassium |
| chloride, sodium bicarbonate and |
| sodium chloride |
| Madopar 125 123 |
| Madopar 250 123 |
| Madopar 62.5 123 |
| Madopar HBS123 |
| Madopar Rapid |
| Magnesium hydroxide |
| Magnesium sulphate |
| Mantoux |
| MAR-Midodrine |
| Marevan43 |
| Marine Blue Lotion SPF 50+76 |
| Martindale Pharma265 |
| Mask for spacer device258 |
| Maviret |
| Maxidex |
| Maxitrol261 |
| MCT oil (Nutricia)273 |
| Measles, mumps and rubella |
| vaccine |
| Mebendazole95 |
| Mebeverine hydrochloride8 |
| Med Mgmt 15 min265 |
| Med Mgmt 30 min265 |
| Med Mgmt 45 min265 |
| Medicine Delivery |
| Medrol85 |
| Medroxyprogesterone acetate |
| Genito-Urinary80 |
| Hormone |
| |

| Mefenamic acid | |
|-----------------------------------|-------|
| Megval | |
| Melatonin | . 143 |
| Melpha | . 155 |
| Melphalan | . 155 |
| Meningococcal (groups A, C, Y and | |
| W-135) conjugate vaccine | |
| Meningococcal B multicomponent | |
| vaccine | 301 |
| Meningococcal C conjugate | |
| wassing | 201 |
| vaccine | |
| MenQuadfi | |
| Menthol | |
| Mepolizumab | |
| Mercaptopurine | |
| Mercilon 28 | 79 |
| Mesalazine | 7 |
| Mesna | . 162 |
| Mestinon | . 116 |
| Metabolic Disorder Agents | |
| Metformin hydrochloride | |
| Metformin Viatris | |
| Methadone BNM | |
| Methadone hydrochloride | |
| Extemporaneous | 260 |
| Nervous | |
| Methenamine (hexamine) | . 121 |
| hippurate | 115 |
| | |
| Methopt | |
| Methotrexate | . 158 |
| Methotrexate DBL Onco-Vial | . 158 |
| Methotrexate DBL S29 | |
| Methotrexate Ebewe | |
| Methotrexate Sandoz | . 158 |
| Methyl hydroxybenzoate | . 269 |
| Methylcellulose | . 269 |
| Methylcellulose with glycerin and | |
| sodium saccharin | . 269 |
| Methylcellulose with glycerin and | |
| sucrose | . 269 |
| Methyldopa | |
| Methyldopa Mylan | |
| Methyldopa Mylan S29 | |
| Methyldopa Viatris | |
| Methylnaltrexone bromide | |
| Methylphenidate ER - Teva | |
| Methylphenidate hydrochloride | |
| Methylphenidate hydrochloride | |
| extended-release | |
| Methylprednisolone | |
| Methylprednisolone (as sodium | 00 |
| succinate) | 05 |
| Methylprednisolone aceponate | |
| | |
| Methylprednisolone acetate | |
| Methylxanthines | .256 |
| Metoclopramide Actavis 10 | . 135 |

| Metoclopramide hydrochloride 135 |
|--|
| Metolazone53 |
| Metopirone94 |
| Metoprolol IV Mylan50 |
| Metoprolol IV Viatris |
| Metoprolol succinate |
| Metoprolol tartrate |
| Metrogyl105 |
| Metronidazole 105 |
| Metyrapone94 |
| Mexiletine hydrochloride |
| Miacalcic |
| Micolette |
| Micolette-S2925 |
| Miconazole |
| Miconazole nitrate |
| Dermatological |
| Genito-Urinary |
| Micreme |
| Micreme H |
| Microgynon 30 |
| Microlut |
| Midazolam |
| Midazolam-Baxter |
| Midodrine |
| Mifegyne |
| Mifepristone |
| |
| Milpharm |
| Minerals |
| Mini-Wright AFS Low Range |
| |
| Minidiab |
| MiniMed 3.0 Reservoir |
| MMT-332A |
| MiniMed 770G |
| MiniMed Mio MMT-921A 22 |
| MiniMed Mio MMT-923A |
| MiniMed Mio MMT-925A |
| MiniMed Mio MMT-941A 22 |
| MiniMed Mio MMT-943A |
| MiniMed Mio MMT-945A22 |
| MiniMed Mio MMT-965A22 |
| MiniMed Mio MMT-975A 22 |
| MiniMed Quick-Set MMT-386A23 |
| MiniMed Quick-Set MMT-387A22 |
| MiniMed Quick-Set MMT-396A22 |
| MiniMed Quick-Set MMT-397A22 |
| MiniMed Quick-Set MMT-398A 22 |
| MiniMed Quick-Set MMT-399A22 |
| MiniMed Silhouette MMT-368A22 |
| MiniMed Silhouette MMT-377A 22 |
| MiniMed Silhouette MMT-378A22 |
| MiniMed Silhouette MMT-381A22 |
| Willinivied Silliouelle Wilvi I-38TA |
| MiniMed Silhouette MMT-382A 22 |
| MiniMed Silhouette MMT-381A22 MiniMed Silhouette MMT-382A22 MiniMed Silhouette MMT-383A22 MiniMed Silhouette MMT-384A22 |

| MiniMed Sure-T MMT-864A | 21 |
|-------------------------------------|-------|
| MiniMed Sure-T MMT-866A | 21 |
| MiniMed Sure-T MMT-874A | 21 |
| MiniMed Sure-T MMT-876A | 21 |
| MiniMed Sure-T MMT-884A | 21 |
| MiniMed Sure-T MMT-886A | 21 |
| Minims Cyclopentolate | 263 |
| Minims Pilocarpine | 263 |
| Minims Prednisolone | 262 |
| Minipress | |
| Minipless | |
| | |
| Minirin Melt | |
| Mino-tabs | 99 |
| Minocycline hydrochloride | 99 |
| Minomycin | 99 |
| Minor Skin Infections | |
| Minoxidil | |
| Minoxidil Roma | |
| Mirena | 88 |
| Miro-Amoxicillin | 98 |
| Mirtazapine | .130 |
| Misoprostol | 9 |
| Mitomycin C | . 162 |
| Mitozantrone | |
| Mitozantrone Ebewe | |
| Mixtard 30 | |
| MMA/PA Explore 5 | |
| MMA/PA Express 15 | 288 |
| Moclobemide | 130 |
| Modafinil | 148 |
| Modavigil | |
| Moduretic | 52 |
| Molaxole | |
| Molnupiravir | |
| | |
| Moments | /8 |
| Mometasone furoate | |
| Monogen | 274 |
| Montelukast | |
| Montelukast Viatris | .256 |
| Moroctocog alfa [Recombinant factor | or |
| VIII] | 39 |
| Morphine hydrochloride | |
| Morphine sulphate | . 128 |
| Motetis | . 124 |
| Mouth and Throat | 31 |
| Мочаро | . 123 |
| Moxifloxacin | . 100 |
| MSUD Explore 5 | .285 |
| MSUD Express 15 | .285 |
| MSUD Maxamum | |
| Mucolytics | |
| Mucosoothe | |
| Multiple Sclerosis Treatments | |
| Multivitamin renal | |
| Multivitamins | |
| Mupirocin | |
| | |

| Muscle Relaxants | |
|--------------------------------|-----------------|
| Mvite | 33 |
| Myambutol | |
| Mycobutin | 107 |
| MycoNail | <mark>69</mark> |
| Mycophenolate mofetil | 177 |
| Mydriacyl | 263 |
| Mylan (24 hr release) | 51 |
| Mylan Atenolol | 49 |
| Mylan Clomiphen | 94 |
| Mylan Italy (24 hr release) | 51 |
| Myleran | 154 |
| Myloc CR | 50 |
| Mylotarg | 203 |
| Myometrial and Vaginal Hormone | |
| Preparations | 81 |
| Myozyme | 26 |
| - N - | |
| Nadolol | 50 |
| Nadolol BNM | 50 |
| Naglazyme | 27 |
| Naloxone hydrochloride | 265 |
| Naltraccord | 151 |
| Naltrexone AOP | 151 |
| Naltrexone hydrochloride | 151 |
| Naphazoline hydrochloride | |
| Naphcon Forte | 264 |
| Naprosyn SR 1000 | 116 |
| Naprosyn SR 750 | 116 |
| Naproxen | |
| Narcaricin mite | |
| Nasal Preparations | |
| Natalizumab | |
| Natulan | |
| Nausafix | 135 |
| Nausafix - S29 | 135 |
| Nausicalm | |
| Navelbine | |
| Navelbine S29 | 167 |
| Nefopam hydrochloride | |
| Neisvac-C | 301 |
| Neo-Cytamen S29 | |
| Neo-Mercazole | |
| Neocate Gold | |
| Neocate Junior Unflavoured | 289 |
| Neocate Junior Vanilla | 289 |
| Neocate SYNEO | |
| Neoral | 245 |
| Neostigmine metilsulfate | . 116 |
| Nepafenac | 262 |
| Nepro HP (strawberry) | 277 |
| Nepro HP (vanilla) | 277 |
| Nepro HP RTH | 277 |
| Neulactil | |
| Neuraxpharm | |
| NeuroTabs | 34 |

| Maximum data data |
|--|
| Nevirapine112 |
| Nevirapine Alphapharm112 |
| Nevirapine Viatris 112 |
| Nicorandil |
| Nicotine |
| Nicourie |
| Nifedipine51 |
| Nifedipine Viatris51 |
| Nifuran 115 |
| Nilotinib169 |
| Nilstat |
| Alimentary32 |
| Alimentary |
| Genito-Urinary81 |
| Infection 103 |
| Nintedanib255 |
| Nipent 165 |
| Niraparib162 |
| Nirmatrelvir with ritonavir |
| |
| Nitrates56 |
| Nitroderm TTS56 |
| Nitrofurantoin115 |
| Nitrolingual Pump Spray56 |
| Nivestim |
| |
| Nivolumab242 |
| Nodia6 |
| Noflam 250 116 |
| Noflam 500 116 |
| Non-Steroidal Anti-Inflammatory |
| Drugs |
| |
| Nanagag gamma [Decembinant |
| Nonacog gamma, [Recombinant |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary. 81 Hormone 88 Norflex. 122 Norfloxacin. 115 Noriday 28 81 Norimin. 80 Norimin-1 28 Day. 80 Norpress. 129 Nortriptyline hydrochloride. 129 Norvir. 113 Noured Dexamfetamine. 146 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin 80 Norimin-1 28 Day 80 Norpress 129 Nortriptyline hydrochloride 129 Norvir 113 Nourmed Dexamfetamine 146 Nourmed Paracetamol 126 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin 80 Norimin-1 28 Day 80 Norpress 129 Nortriptyline hydrochloride 129 Norvir 113 Nourmed Dexamfetamine 146 Nourmed Paracetamol 126 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin 80 Norimin-1 28 Day 80 Norrison 143 Norpress 129 Nortriptyline hydrochloride 129 Norvir 113 Noured Dexamfetamine 146 Noumed Paracetamol 126 Noumed Pethidine 128 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary. 81 Hormone 88 Norflex. 122 Norfloxacin. 115 Noriday 28 81 Norimin. 80 Norimin.1 28 Day. 80 Normison 143 Norpress 129 Nortriptyline hydrochloride. 129 Norvir. 113 Noured Dexamfetamine. 146 Noumed Paracetamol. 126 Noumed Pethidine. 128 Noumed Pethidine. 128 Noumed Pethidine. 128 Noumed Pethidine. 128 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary. 81 Hormone 88 Norflex. 122 Norfloxacin. 115 Noriday 28 81 Norimin. 80 Norimin.1 28 Day. 80 Norrison 143 Norpress 129 Nortvir. 113 Noured Dexamfetamine. 146 Noumed Paracetamol. 126 Noumed Pethidine. 128 Noumed Pethidine. 132 NovaSource Renal 277 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin 80 Norimin-1 28 Day 80 Norrison 143 Norpress 129 Nortriptyline hydrochloride 129 Noured Dexamfetamine 146 Noumed Dexamfetamine 126 Noumed Pethidine 128 Noumed Pethidine 128 Noured Pethidine 132 NovaSource Renal 277 Novatretin 74 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin-1 28 Day Norrinion 80 Norrinion 143 Norpress 129 Nortriptyline hydrochloride 129 Norivir 113 Noured Dexamfetamine 146 Noumed Paracetamol 126 Noumed Pethidine 128 Noumed Pethidine 127 NovaSource Renal 277 Novatretin 74 Novitium Sugar Free 28 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin 80 Norimin-1 28 Day 80 Norrison 143 Norpress 129 Nortriptyline hydrochloride 129 Norivir 113 Noured Dexamfetamine 146 Noumed Paracetamol 126 Noumed Pethidine 128 Noured Petholarbitone 128 NovaSource Renal 277 Novatretin 74 Novitium Sugar Free 28 NovoMix 30 FlexPen 10 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin-1 28 Day Norrinion 80 Norrinion 143 Norpress 129 Nortriptyline hydrochloride 129 Norivir 113 Noured Dexamfetamine 146 Noumed Paracetamol 126 Noumed Pethidine 128 Noumed Pethidine 127 NovaSource Renal 277 Novatretin 74 Novitium Sugar Free 28 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin 80 Norimin-1 28 Day Norrison 143 Norpress 129 Nortriptyline hydrochloride 129 Norivir 113 Noumed Dexamfetamine 146 Noumed Paracetamol 126 Noumed Phenobarbitone 122 Novatretin 74 Novittium Sugar Free 28 NovoMix 30 FlexPen 10 NovoRapid 11 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin 80 Norimin-1 28 Day 80 Normison 143 Norpress 129 Nortriptyline hydrochloride 129 Norivir 113 Noumed Dexamfetamine 146 Noumed Paracetamol 126 Noumed Pethidine 128 NovaGource Renal 277 NovaSource Renal 277 Novatretin 74 Novitium Sugar Free 28 NovoRapid 11 NovoRapid 11 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary. 81 Hormone 88 Norflex 122 Norfloxacin. 115 Noriday 28 81 Norimin 80 Norimin. 80 Norrison 143 Norpress 129 Nortriptyline hydrochloride 129 Norvir 113 Noumed Dexamfetamine 146 Noumed Paracetamol 126 Noumed Pethidine 128 Novasource Renal 277 Novatretin 74 Novitium Sugar Free 28 NovoRapid 11 NovoRapid FlexPen 11 NovoRapid FlexPen 11 NovoRapid Penfill 11 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary 81 Hormone 88 Norflex 122 Norfloxacin 115 Noriday 28 81 Norimin 80 Norimin-1 28 Day 80 Norrison 143 Norpress 129 Norvir 113 Noumed Dexamfetamine 146 Noumed Paracetamol 126 Noumed Pethidine 128 Novasource Renal 277 Novastretin 74 Novitium Sugar Free 28 NovoRapid 11 NovoRapid FlexPen 11 NovoRapid Penfill 11 NovoRapid Penfill 11 |
| Nonacog gamma, [Recombinant Factor IX] 39 Norethisterone 81 Genito-Urinary. 81 Hormone 88 Norflex 122 Norfloxacin. 115 Noriday 28 81 Norimin 80 Norimin. 80 Norrison 143 Norpress 129 Nortriptyline hydrochloride 129 Norvir 113 Noumed Dexamfetamine 146 Noumed Paracetamol 126 Noumed Pethidine 128 Novasource Renal 277 Novatretin 74 Novitium Sugar Free 28 NovoRapid 11 NovoRapid FlexPen 11 NovoRapid FlexPen 11 NovoRapid Penfill 11 |

| Nozinan S29 | 137 |
|--|---|
| Nucala | 211 |
| Nuelin | 256 |
| Nuelin-SR | 256 |
| Nupentin | 131 |
| Nusinersen | 144 |
| Nutilis | 284 |
| Nutren Diabetes | |
| Nutrient Modules | 071 |
| | |
| Nutrini Energy Multi Fibre | 270 |
| Nutrini Energy RTH | 276 |
| Nutrini Low Energy Multi Fibre | 278 |
| Nutrini Peptisorb | 291 |
| Nutrini Peptisorb Energy | 291 |
| Nutrini RTH | 276 |
| Nutrison 800 Complete Multi | |
| Fibre | |
| Nutrison Advanced Diason | 274 |
| Nutrison Advanced Peptisorb | 278 |
| Nutrison Concentrated | 283 |
| Nutrison Energy | 281 |
| Nutrison Energy Multi Fibre | 281 |
| Nutrison Multi Fibre | 281 |
| Nutrison RTH | 281 |
| Nyefax Retard | 51 |
| Nystatin | |
| Alimentary | 20 |
| | |
| | 01 |
| Genito-Urinary | 81 |
| Infection | 103 |
| Infection NZB Low Gluten Bread Mix | 103 |
| Infection NZB Low Gluten Bread Mix - O - | 103 284 |
| Infection NZB Low Gluten Bread Mix - O - Obinutuzumab | 103 284 212 |
| Infection NZB Low Gluten Bread Mix • O - Obinutuzumab Obstetric Preparations | 103 284 212 83 |
| Infection NZB Low Gluten Bread Mix - O - Dbinutuzumab Dbstetric Preparations Dcicure | .103 .284 .212 83 9 |
| Infection VZB Low Gluten Bread Mix - O - Dbinutuzumab Dstetric Preparations Dcicure Dcrelizumab | .103 .284 .212 83 9 .141 |
| Infection NZB Low Gluten Bread Mix - O - Dbinutuzumab Dbstetric Preparations Dcicure | .103 .284 .212 83 9 .141 |
| Infection NZB Low Gluten Bread Mix - O - Dbinutuzumab Dbstetric Preparations Dcicure Dcrelizumab Dcrevus | .103 .284 .212 83 9 .141 |
| Infection NZB Low Gluten Bread Mix - O - Dbinutuzumab Dostetric Preparations Diccure Dcrelizumab Dcrevus Dcrevos Dctocog alfa [Recombinant factor | 103 284 212 83 9 141 141 |
| Infection NZB Low Gluten Bread Mix O - Dbinutuzumab Dostetric Preparations Distetric Preparations Distetric Preparations Disterve . | 103 284 212 83 9 141 141 |
| Infection | 103 284 .212 83 9 .141 .141 39 |
| Infection | 103 284 212 83 9 141 141 39 39 |
| Infection | 103 284 212 83 9 141 141 39 39 175 |
| Infection | 103 284 212 9 141 141 39 39 175 175 |
| Infection | 103 284 212 83 9 141 .141 39 39 .175 .175 .175 |
| Infection | 103 284 212 83 9 .141 141 39 39 .175 .175 .175 .175 |
| Infection | 103 284 212 83 9 141 141 39 39 175 175 175 87 |
| Infection | 103 284 212 9 141 141 39 175 175 175 87 87 |
| Infection | 103 284 212 9 141 141 39 175 175 175 87 87 |
| Infection | 103 284 212 83 9 141 .141 39 175 175 175 175 87 87 87 |
| Infection | 103 284 212 83 9 141 141 39 39 175 175 175 175 87 87 88 81 |
| Infection | 103 284 212 83 9 141 141 39 39 175 175 175 87 87 87 88 81 88 |
| Infection | 103 284 212 83 9 141 141 39 39 175 175 175 175 87 87 87 88 81 88 81 |
| Infection | 103 284 212 83 9 141 141 39 39 175 175 175 87 87 88 81 88 81 88 87 255 |
| Infection | 103 284 212 83 9 141 141 39 39 175 175 175 175 87 87 88 81 88 81 88 87 255 72 |
| Infection | 103 284 212 83 9 141 141 39 39 175 175 175 175 87 87 88 81 88 81 88 87 255 72 |
| Infection | 103 284 212 83 9 141 141 39 175 175 175 175 87 87 88 81 88 81 88 72 138 |
| Infection | 103 284 212 83 9 141 141 39 175 175 175 175 87 88 88 81 88 81 72 -138 163 |

| Olbetam S29 | 54 |
|------------------------------|-----|
| Olopatadine | 264 |
| Olopatadine Teva | |
| Olsalazine | 8 |
| Omalizumab | 213 |
| Omeprazole | 9 |
| Omeprazole actavis 10 | 9 |
| Omeprazole actavis 20 | 9 |
| Omeprazole actavis 40 | 9 |
| Omnitrope | 89 |
| Omnitrope S29 | 89 |
| Onbrez Breezhaler | 251 |
| Oncaspar LYO | 164 |
| OncoTICE | 184 |
| Ondansetron | 135 |
| One-Alpha | 33 |
| One-Alpha S29 | 33 |
| Opdivo | |
| Ora-Blend | |
| Ora-Blend SF | |
| Ora-Plus | 269 |
| Ora-Sweet | |
| Ora-Sweet SF | 269 |
| Orabase | |
| Oral and Enteral Feeds | 274 |
| Oralcon 30 ED | |
| Oramorph | 128 |
| Oratane | |
| Ordine | |
| Orgran | |
| Ornidazole | 105 |
| Orphenadrine citrate | 122 |
| Ortho-tolidine | |
| Oruvail SR | |
| Osmolite RTH | |
| Other Endocrine Agents | |
| Other Oestrogen Preparations | 88 |
| Other Breggetegen | |
| Preparations | 88 |
| Other Skin Preparations | 76 |
| Otodex | |
| Ovestin | |
| Genito-Urinary | 81 |
| Hormone | |
| Oxaliplatin | 155 |
| Oxaliplatin Accord | 155 |
| Oxaliplatin Actavis 100 | 155 |
| Oxaliplatin Ebewe | |
| Oxis Turbuhaler | |
| Oxpentifylline | .57 |
| Oxybutynin | .83 |
| Oxycodone hydrochloride | 128 |
| Oxycodone Sandoz | 128 |
| Oxycodone Sandoz S29 | |
| OxyContin | |
| OxyNorm | |
| , · | |

| Oxytocin | 82 |
|---------------------------------|-----|
| Oxytocin BNM | 82 |
| Oxytocin GH | 82 |
| Oxytocin Panpharma | |
| Oxytocin with ergometrine | 02 |
| maleate | 00 |
| | |
| Ozurdex | 260 |
| | ~ |
| Pacifen1 | |
| Pacimol1 | |
| Paclitaxel1 | |
| Paclitaxel Actavis1 | |
| Paclitaxel Ebewe 1 | |
| Paediatric Seravit | |
| Palbociclib1 | |
| Paliperidone1 | |
| Paliperidone palmitate 1 | 39 |
| Pamidronate disodium1 | |
| Pamisol1 | |
| Pamol 1 | |
| Pancreatic enzyme | |
| Pantoprazole | . 9 |
| Panzop Relief | 9 |
| Papaverine hydrochloride | |
| Para-amino salicylic acid1 | |
| Paracetamol1 | |
| Paracetamol (Ethics) 1 | 26 |
| Paracetamol + Codeine | |
| (Relieve) 1 | 28 |
| Paracetamol with codeine1 | |
| Paraffin | |
| Paraffin liquid with wool fat | 264 |
| Parasiticidal Preparations | |
| Parnate 1 | |
| Paromomycin1 | |
| Paroxetine 1 | |
| Paser 1 | |
| Paxam 1 | |
| Paxlovid 1 | |
| Pazopanib1 | |
| Peak flow meter | |
| Pediasure | 276 |
| Pediasure Plus2 | |
| Pediasure RTH2 | |
| Pegaspargase1 | 64 |
| Pegasys1 | |
| Pegfilgrastim | |
| Pegylated interferon alfa-2a1 | 13 |
| Pembrolizumab2 | |
| Pemetrexed1 | |
| Penicillamine1 | |
| Penicillin G | 98 |
| PenMix 30 | 11 |
| PenMix 50 | |
| Pentasa | |
| Pentostatin [Deoxycoformycin] 1 | 65 |

| Pentoxifylline [Oxpentifylline] | 57 |
|---------------------------------------|-------|
| Peptamen Junior | 276 |
| Pepti-Junior | 292 |
| Perhexiline maleate | |
| Pericyazine | |
| Perindopril | 47 |
| Periset | 135 |
| Periset ODT | 135 |
| Perjeta | |
| Permethrin | 213 |
| Perrigo | 79 |
| Pertuzumab | 215 |
| Peteha | |
| Pethidine hydrochloride | 100 |
| Pevaryl | . 120 |
| | |
| Pexsig | |
| Pfizer Exemestane | |
| Pfizer S29 | 156 |
| Pharmacy Services | |
| Pheburane | |
| Phenasen | |
| Phenobarbitone | 132 |
| Phenobarbitone sodium | |
| Extemporaneous | 269 |
| Nervous | 143 |
| Phenoxybenzamine | |
| hydrochloride | 46 |
| Phenoxymethylpenicillin (Penicillin | |
| V) | |
| Phenylalanine50 | 288 |
| Phenytoin sodium131 | -132 |
| Phillips Milk of Magnesia | 35 |
| Phlexy 10 | 286 |
| Phosphate Phebra | 44 |
| Phosphorus | |
| Phytomenadione | |
| Pilocarpine hydrochloride | |
| Pilocarpine nitrate | 263 |
| Pimafucort | |
| Pimecrolimus | |
| Pine tar with trolamine laurilsulfate | |
| and fluorescein | 75 |
| Pinetarsol | |
| Pioglitazone | |
| Pirfenidone | 256 |
| Pizotifen | 12/ |
| PKU Anamix Infant | |
| | |
| PKU Anamix Junior | |
| PKU Anamix Junior Chocolate | |
| PKU Anamix Junior LQ | |
| PKU Anamix Junior Orange | |
| PKU Anamix Junior Vanilla | |
| PKU Build 10 | |
| PKU Build 20 Chocolate | 287 |
| PKU Build 20 Raspberry | |
| Lemonade | . 287 |
| | |

| PKU Build 20 Smooth | .287 |
|--------------------------------|-----------|
| PKU Build 20 Vanilla | |
| PKU Explore 10 | |
| PKU Explore 5 | 286 |
| PKU Express 20 | 286 |
| PKU Glytactin RTD 15 | 287 |
| PKU Glytactin RTD 15 Lite | |
| PKU GMPro Ultra Lemonade | |
| PKU Lophlex LQ 10 | |
| PKU Lophlex LQ 20 | |
| PKU Lophlex Powder | |
| | |
| PKU Lophlex Sensation 20 | |
| PKU Restore Powder | |
| PKU sphere20 Banana | |
| PKU sphere20 Chocolate | |
| PKU sphere20 Lemon | .287 |
| PKU sphere20 Red Berry | .287 |
| PKU sphere20 Vanilla | .287 |
| PKU Start | |
| Plaquenil | |
| Plendil ER | |
| PMS-Salbutamol | .253 |
| Pneumococcal (PCV10) conjugate | |
| vaccine | . 302 |
| Pneumococcal (PCV13) conjugate | |
| vaccine | . 303 |
| Pneumococcal (PPV23) | |
| polysaccharide vaccine | . 304 |
| Pneumovax 23 | .304 |
| Podophyllotoxin | 76 |
| Polaramine | .250 |
| Poliomyelitis vaccine | .304 |
| Poloxamer | 25 |
| Poly-Gel | .264 |
| Poly-Tears | |
| Poly-Visc | .264 |
| Polycal | .271 |
| Polyethylene glycol 400 and | |
| propylene glycol | 264 |
| Ponstan | .116 |
| Posaconazole | |
| Posaconazole Juno | |
| Potassium chloride4 | 3-44 |
| Potassium citrate | |
| Potassium iodate | |
| Povidone iodine | |
| Pradaxa | |
| Pramipexole hydrochloride | 123 |
| Pravastatin | |
| Praziguantel | |
| Prazosin | |
| Prazosin Mylan | |
| Pred Forte | 40 060 |
| Prednisolone | |
| Prednisolone acetate | |
| Prednisolone acetate | |
| | X |

Prednisolone sodium

| phosphate262 |
|-------------------------------|
| Prednisolone-AFT262 |
| Prednisone86 |
| Prednisone Clinect |
| Pregabalin132 |
| Pregabalin Pfizer |
| Pregnancy Tests - hCG Urine |
| Premarin |
| Prevenar 13 |
| Priadel |
| Primaquine104 |
| Primidone132 |
| Primidone Clinect |
| Primolut N88 |
| Priorix |
| Probenecid121 |
| Probenecid-AFT 121 |
| Procarbazine hydrochloride |
| Prochlorperazine |
| Prochlorperazine Brown & |
| Burk 135 |
| Proctofoam |
| Proctosedyl |
| Procyclidine hydrochloride123 |
| Progesterone |
| Proglicem |
| Proglycem10 |
| Progynova |
| Prolia |
| Promethazine hydrochloride |
| Propafenone hydrochloride |
| Propranolol |
| Propylene glycol |
| Propylthiouracil |
| Prostacur |
| Protaphane 10 |
| Protaphane Penfill |
| Protifar |
| Protionamide |
| Provera |
| Provera HD |
| Deariasis and Ference |
| Preparations |
| PTU |
| Pulmicort Turbuhaler |
| Pulmozyme |
| Puri-nethol 157 |
| Puritan's Pride Vitamin |
| B-2 100 mg 29 |
| Pyrazinamide |
| Pyridostigmine bromide |
| Pyridoxine hydrochloride |
| Pyridoxine multichem |
| Pyrimethamine101 |
| Pytazen SR40 |
| 1 ytuzon OI t |

- 0 -Quantalan sugar free......54 Quetapel......137 Quinapril47 Quinapril with hydrochlorothiazide 47 - R -RA-Morph127 Ralicrom8 Raloxifene hydrochloride118 Raltegravir potassium......113 Ramipex123 Ramipril47 Ranbaxy-Cefaclor.....95 Rapamune.....246 Rasagiline.....123 Reandron 1000......86 Recombinant factor IX 37, 39 Rectogesic.....8 Redipred......86 Relieve.....116 Resonium-A......45 Resource Beneprotein......273 Respigen253 Respiratory Stimulants259 Retinol palmitate......264 Revlimid......161 Riboflavin......29 Ribomustin......153 Rifabutin107 Rifadin107 Rifadin Sanofi 107 Rifampicin.....107 Rifaximin......10 Rifinah106 Riluzole.....123 Risdiplam.....144 Risedronate Sandoz 119 Risedronate sodium......119 Risperdal Consta 139 Risperidone 137, 139

| Risperidone (Teva) | 137 |
|-----------------------------------|-------|
| Risperon | 137 |
| Ritalin | . 147 |
| Ritalin LA | |
| Ritonavir | |
| Rituximab (Mabthera) | |
| Rituximab (Riximyo) | |
| Rivaroxaban | |
| Rivastigmine | |
| Rivastigmine Patch BNM 10 | 149 |
| Rivastigmine Patch BNM 5 | |
| Rivotril | |
| Riximvo | |
| RIXUBIS | |
| Rizamelt | 134 |
| Rizatriptan | 134 |
| Robinul | |
| Ronapreve | |
| Ropin | |
| Ropinirole hydrochloride | 123 |
| Rosuvastatin | |
| Rosuvastatin Viatris | |
| Rotarix | |
| Rotavirus oral vaccine | |
| Roxane-Propranolol | |
| Roxithromycin | |
| Rubifen | |
| Rubifen SR | |
| Rugby Capsaicin Topical Cream | |
| Musculoskeletal | . 117 |
| Nervous | |
| Rurioctocog alfa pegol [Recombina | |
| factor VIII] | |
| Ruxolitinib | |
| Rythmodan | |
| Rythmodan - Cheplafarm | |
| Rytmonorm | |
| - S - | |
| Sabril | 133 |
| Sacubitril with valsartan | |
| Sagent | |
| SalAir | |
| Salazopyrin | |
| Salazopyrin EN | |
| Salbutamol252 | -253 |
| Salbutamol Cipla | 253 |
| Salbutamol with ipratropium | |
| bromide | 253 |
| Salicylic acid | |
| Salmeterol | |
| Sandomigran | |
| Sandostatin LAR | |
| Sanofi Primaquine | |
| Sapropterin dihydrochloride | |
| Scalp Preparations | |
| ooup i iopaialloilð | 10 |
| Scopoderm TTS | |

| Scopolamine - Mylan 13 | 5 |
|-----------------------------------|----------|
| Scopolamine - Mylan S29 13 | 5 |
| Sebizole | 6 |
| Secukinumab22 | |
| Sedatives and Hypnotics14 | 3 |
| Seebri Breezhaler | |
| Senna | |
| Senokot | |
| SensoCard1 | |
| Serc | |
| Serenace | |
| Seretide | |
| Seretide Accuhaler | |
| Serevent | |
| Serevent Accuhaler | |
| | |
| Sertraline | |
| Setrona13 | |
| Sevredol | 8 |
| Sex Hormones Non | |
| Contraceptive | |
| Shingles vaccine | |
| Shingrix | 6 |
| SII-Onco-BCG 18 | |
| Sildenafil6 | |
| Siltuximab23 | |
| Simvastatin5 | 6 |
| Simvastatin Mylan | 6 |
| Simvastatin Viatris | 6 |
| Sinemet 12 | |
| Sinemet CR 12 | 3 |
| Sirolimus24 | 6 |
| Sirturo 10 | 5 |
| Siterone8 | 6 |
| Slow-Lopresor | 0 |
| Smith BioMed Rapid Pregnancy | |
| Test | 2 |
| Sodibic4 | |
| Sodium acid phosphate2 | 5 |
| Sodium alginate | 6 |
| Sodium benzoate | 0 |
| Sodium bicarbonate | |
| Blood4 | 4 |
| Extemporaneous27 | |
| Sodium calcium edetate | |
| Sodium chloride | |
| Blood4 | 4 |
| Respiratory25 | 8 |
| Sodium citrate with sodium lauryl | Ĩ |
| sulphoacetate | 5 |
| Sodium citro-tartrate | |
| Sodium cromoglicate | ` |
| Alimentary | 8 |
| Sensory | |
| Sodium Fusidate [fusidic acid] | - |
| Dermatological | ۵ |
| | J |

| Infection | |
|--------------------------------|-------|
| Sensory | 260 |
| Sodium hyaluronate [Hyaluronic | |
| acid] | 264 |
| Sodium phenylbutyrate | 30 |
| | |
| Sodium picosulfate | 20 |
| Sodium polystyrene sulphonate | 45 |
| Sodium tetradecyl sulphate | 39 |
| Sodium valproate | 133 |
| Sodium valproate Sofradex | 260 |
| Soframycin | 260 |
| Solgar | 26 20 |
| Solifenacin succinate | 00-00 |
| | |
| Solifenacin Viatris | 83 |
| Solu-Cortef | 85 |
| Solu-Medrol | 85 |
| Solu-Medrol-Act-O-Vial | 85 |
| Somatropin (Omnitrope) | 89 |
| Sotalol | 50 |
| Spacer device | 250 |
| | 209 |
| Span-K | 44 |
| Spazmol | 8 |
| Spinal Muscular Atrophy | 144 |
| Spinraza | 144 |
| Spiolto Respimat | 254 |
| Spiractin | 53 |
| Spiriva | 254 |
| Spiriva Respimat | 204 |
| | 204 |
| Spironolactone | 53 |
| Sporanox | 103 |
| Sprycel | 167 |
| Stelara | 237 |
| Stemetil | 135 |
| Steril-Gene | |
| SteroClear | 258 |
| Stesolid | 101 |
| | 101 |
| Stimulants/ADHD Treatments | 145 |
| Stiripentol | |
| Stocrin | 111 |
| Stomahesive | 31 |
| Strides Shasun | 103 |
| Stromectol | 73 |
| Sucralfate | 10 |
| Sulfadiazine Silver | 60 |
| | |
| Sulfadiazine sodium | 101 |
| Sulfasalazine | 8 |
| Sulphur | 75 |
| Sulprix | 136 |
| Sumagran | 134 |
| Sumatriptan | 134 |
| Sunitinib | 170 |
| Sunitinib Pfizer | 170 |
| | 172 |
| Sunscreens | /6 |
| Sunscreens, proprietary | 76 |
| Survimed OPD | 278 |
| | |

| Sustagen Hospital Formula | 281 |
|------------------------------------|-------------------|
| Sustagen Hospital Formula | |
| Active | 281 |
| Sustanon Ampoules | 86 |
| Sylvant | 230 |
| Symbicort Turbuhaler 100/6 | 252 |
| Symbicort Turbuhaler 200/6 | 252 |
| Symbicort Turbuhaler 400/12 | 252 |
| Symmetrel | 123 |
| Sympathomimetics | <mark>56</mark> |
| Synacthen | <mark>86</mark> |
| Synacthen Depot | <mark>86</mark> |
| Synacthene Retard | <mark>86</mark> |
| Synflorix | 302 |
| Synthroid | 88 |
| Syntometrine | |
| Syrup (pharmaceutical grade) | |
| Systane Unit Dose | |
| -T- | |
| Tacrolimus | |
| Dermatological | 75 |
| Oncology | 247 |
| Tacrolimus Sandoz | <u>2</u> 77 |
| Taliglucerase alfa | <u>2</u> 47 20 |
| Tambocor | |
| Tamoxifen citrate | |
| Tamoxifen Sandoz | 170 |
| Tamoxileri Sanuoz | 170 |
| Tamsulosin hydrochloride | |
| Tamsulosin-Rex | 82 |
| Tandem Cartridge | 21 |
| Tandem t:slim X2 with Basal-IQ | 16 |
| Tap water | |
| Taro | |
| Taro-Testosterone | |
| Tasigna | 169 |
| Tasmar | |
| Taurine | |
| Tecentriq | |
| Tecfidera | |
| Tegretol | 131 |
| Tegretol CR | |
| Telfast | 250 |
| Teligent | 100 |
| Temaccord | 165 |
| Temazepam | |
| Temozolomide | 165 |
| Tenofovir disoproxil | |
| Tenofovir Disoproxil Emtricitabine | |
| Viatr | . 109 |
| Tenofovir Disoproxil Viatris | |
| Tenoxicam | |
| Tensipine MR10 | |
| Tepadina | |
| Terbinafine | 104 |
| Terbutaline sulphate | |
| Teriflunomide | |
| | |

| Teriparatide119 |
|--|
| Teriparatide - Teva 119 |
| Testogel |
| Testosterone86 |
| Testosterone cipionate |
| Testosterone esters |
| Testosterone undecanoate |
| Tetrabenazine |
| Tetrabromophenol |
| Tetracosactrin |
| Tetracycline |
| Teva Lisinopril |
| Teva-Ketoconazole |
| Teva-Salbutamol Sterinebs |
| P.F253 |
| Thalidomide165 |
| Thalomid165 |
| Theophylline |
| Theophylline |
| Thiamine hydrochloride |
| Thiamine multichem |
| THIO-TEPA |
| Thioguanine |
| Thiotepa155 Thyroid and Antithyroid Agents |
| Ticagrelor |
| Ticagrelor Sandoz40 |
| Tilcotil |
| Timolol |
| |
| Tiotronium bromide 254 |
| Tiotropium bromide254 |
| Tiotropium bromido with |
| Tiotropium bromide with olodaterol |
| Tiotropium bromide with olodaterol254 Tivicay |
| Tiotropium bromide with olodaterol |
| Tiotropium bromide with olodaterol 254 Tivicay 113 Tixagevimab with cilgavimab 230 TMP 102 Tobramycin 102 Infection 102 Sensory 260 Tobramycin (Viatris) 102 |
| Tiotropium bromide with olodaterol |
| Tiotropium bromide with olodaterol |
| Tiotropium bromide with olodaterol |
| Tiotropium bromide with olodaterol 254 Tivicay 113 Tixagevimab with cilgavimab 230 TMP 102 Tobramycin 102 Infection 102 Sensory 260 Tobramycin (Viatris) 102 Tobramycin BNM 102 Tobrac 260 Tobramycin BIM 102 Tobrac 260 Tocilizumab 230 Tofranil 129 |
| Tiotropium bromide with olodaterol |

| Tranexamic acid40 |
|--|
| |
| Franylcypromine sulphate129 |
| Frastuzumab (Herceptin)234 |
| Frastuzumab (Herzuma)235 |
| Frastuzumab emtansine |
| Fravatan262 |
| Fravoprost262 |
| Freatments for Dementia149 |
| Freatments for Substance |
| Dependence 150 |
| Frelegy Ellipta254 |
| Frental 400 57 |
| Fretinoin |
| Dermatological 68 |
| Oncology166 |
| Trexate |
| Triamcinolone acetonide |
| Alimentary32 |
| Dermatological71 |
| Hormone86 |
| Friamcinolone acetonide with |
| gramicidin, neomycin and nystatin |
| Dermatological71 |
| Sensory260 |
| Trientine30 |
| Trientine Waymade30 |
| Trikafta257 |
| Trimethoprim102 |
| Trimethoprim with |
| sulphamethoxazole |
| [Co-trimoxazole] 102 |
| Trisequens88 |
| |
| Trisul |
| Frophic Hormones89 |
| Frophic Hormones |
| Frophic Hormones |
| Frophic Hormones |
| Frophic Hormones 88 Fropicamide 263 Frulicity 12 FruSteel 21 Fryzan 47 |
| Frophic Hormones 88 Fropicamide 263 Frulicity 12 FruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 |
| Frophic Hormones 88 Fropicamide 263 Frulicity 12 FruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 FruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 Fwo Cal HN 283 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 FruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 Fwo Cal HN 283 FYR Explore 5 286 |
| Frophic Hormones 86 Fropicamide 260 Frulicity 12 FruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 Two Cal HN 283 FYR Explore 5 286 FYR Sphere 20 286 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 FruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 Fwo Cal HN 283 FYR Explore 5 288 FYR Sphere 20 286 Fyrosine1000 286 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 TruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 Fyr Saphere 5 286 FYR Sphere 20 286 Fyrosine 1000 286 Fyrosine 1000 286 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 TruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 FVR Cal HN 286 FYR Explore 5 286 FVR Sphere 20 286 Fysabri 141 - U - - |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 TruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 FWR Cal HN 286 TYR Explore 5 286 Tyrosine 1000 286 Tysabri 141 - U - 365 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 TruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fuberculin PPD [Mantoux] test 307 Five Cal HN 283 FYR Explore 5 286 Fyrsine 1000 286 Fysabri 141 JK Synacthen 86 Jltibro Breezhaler 254 |
| Frophic Hormones 89 Fropicamide 263 Frulicity 12 FruSteel 21 Tryzan 47 Fuberculin PPD [Mantoux] test 307 Fuberculin PPD [Mantoux] test 307 Frwo Cal HN 283 FYR Explore 5 286 Fyrosine 1000 286 Fysabri 141 JK Synacthen 86 Jltibro Breezhaler 254 |
| Frophic Hormones 89 Fropicamide 263 Frulicity 12 FruSteel 21 Tryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 FVR Explore 5 286 Fyrsphere 20 286 Fysabri 141 Jtibro Breezhaler 254 Jltiaproct 254 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 FruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 FWR Sphere 20 286 Fyrosine 1000 286 Fysabri 141 Jtibro Breezhaler 254 Jtraproct 254 Jmeclidinium 254 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 Frulicity 12 Frusteel 21 Frusteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fubersol 307 Fwo Cal HN 263 FYR Sphere 20 286 Fyrosine1000 286 Fysabri 141 - U - JK JK Synacthen 86 Jithoro Breezhaler 254 Jitraproct 26 Jmeclidinium 254 Jnivent 253, 256 |
| Frophic Hormones 86 Fropicamide 260 Frulicity 12 FruSteel 21 FruSteel 21 Fruberculin PPD [Mantoux] test 307 Fubersol 307 Fwo Cal HN 283 FYR Explore 5 286 FYR Sphere 20 286 Fysosine 1000 286 Fysabri 141 - U - 307 JK Synacthen 306 Jitraproct 254 Jmeclidinium 254 Jnivent 253, 255 Jpadacitinib 246 |
| Frophic Hormones 86 Fropicamide 260 Frulicity 12 TruSteel 21 Frusterel 21 Frusterel 21 Frusterel 21 Frusterel 21 Frusterel 307 Fubersol 307 Fwo Cal HN 283 FYR Explore 5 286 FYR Sphere 20 286 Fysabri 141 - U - 307 JK Synacthen 86 Jitraproct 254 Jmeclidinium with vilanterol 254 Jnivent 253, 256 Jpadacitinib 244 Jral 86 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 TruSteel 21 Frusteel 21 Frusteel 21 Frusterel 21 Frusterel 307 Fubersol 307 Fubersol 307 Fwo Cal HN 283 FYR Sphere 20 286 Fyrosine 1000 286 Fysabri 141 - U - 307 JK Synacthen 36 Just Synacthen 36 Just Synacthen 254 Jitraproct 254 Jnivent 253, 256 Jpadacitinib 244 Jral 36 Jrea 72 |
| Frophic Hormones 86 Fropicamide 263 Frulicity 12 TruSteel 21 Fryzan 47 Fuberculin PPD [Mantoux] test 307 Fuberculin PPD [Mantoux] test 307 Five Cal HN 283 FYR Explore 5 286 Fyrsine 1000 286 Fysabri 141 JK Synacthen 86 Jltibro Breezhaler 254 |

| Urinary Agents | 82 |
|------------------------------------|------------|
| Urinary Tract Infections | 115 |
| Urinorm | |
| Uromitexan | . 162 |
| Ursodeoxycholic acid | |
| Ursosan | |
| Ustekinumab | |
| litrogestan | 207 |
| Utrogestan | |
| Vaccinations | |
| Vaclovir | |
| Valaciclovir | |
| Valganciclovir | |
| Valganciclovir Viatris | 107 |
| Valine50 | 107 000 |
| Vancomycin | 100 |
| Vancomycin | 102 |
| Vannair | 202 |
| Varenicline Pfizer | 151 |
| Varenicline tartrate | 151 |
| Varicella vaccine [Chickenpox | |
| vaccine] | . 306 |
| Varicella zoster vaccine [Shingles | |
| vaccine] | . 306 |
| Various | |
| Varivax | |
| Vasodilators | |
| Vasopressin Agonists | 93 |
| Vasorex | 51 |
| Vebulis | 65 |
| Vedafil | |
| Vedolizumab | 239 |
| Veletri | 63 |
| Venclexta | |
| Venetoclax | |
| Venlafaxine | |
| Venomil249 | -250 |
| VENOX | 249 |
| Ventolin252 | -253 |
| Ventolin Nebules | 253 |
| Vepesid | 160 |
| Verapamil hydrochloride | 100 |
| Vermox | |
| Versacloz | |
| Vesanoid | |
| Vexazone | |
| Vfend | ۲۵ ۱۵۸ |
| Viederm KC | . 104 |
| Viaderm KC | |
| Victoza | |
| Vigabatrin | |
| Vigisom | |
| Vildagliptin | 12 |
| Vildagliptin with metformin | |
| hydrochloride | |
| Vimpat | |
| Vinblastine sulphate | |
| Vincristine sulphate | 167 |

| Vinorelbine167 |
|---|
| Vinorelbine Ebewe167 |
| Vinorelbine Te Arai167 |
| Viramune Suspension112 |
| ViruPOS260 |
| Vit.D333 |
| Vita-B1232 |
| VitA-POS |
| Vitabdeck33 |
| Vital |
| Vitamin B complex32 |
| Vitamin B6 25 32 |
| Vitamins |
| Vitarubin Depot Injection32 |
| Vivonex TEN |
| Voltaren 116 |
| Voltaren D116 |
| Voltaren Ophtha261 |
| Voltaren SR 116 |
| Volumatic259 |
| Voriconazole104 |
| Votrient170 |
| Vttack104 |
| - W - |
| Warfarin sodium |
| Wart Preparations76 |
| Wasp venom allergy treatment250 |
| Water |
| |
| Blood44 |
| Extemporaneous 270 |
| Extemporaneous |
| Blood |
| Extemporaneous270 White Soft Liquid Paraffin AFT72 Wool fat with mineral oil72 - X - |
| Extemporaneous |
| Extemporaneous270 White Soft Liquid Paraffin AFT72 Wool fat with mineral oil72 - X - |
| Extemporaneous |
| Extemporaneous |
| Extemporaneous |
| Extemporaneous 270 White Soft Liquid Paraffin AFT 72 Wool fat with mineral oil 72 - X - Xarelto Xarelto 42 Xifaxan 10 XMET Maxamum 285 Xolair 213 XP Maxamum 286 Xylocaine 124 |
| Extemporaneous 270 White Soft Liquid Paraffin AFT 72 Wool fat with mineral oil 72 - X - Xarelto Xarelto 42 Xifaxan 10 XMET Maxamum 285 Xolair 213 XP Maxamum 124 Xylocaine 124 |
| Extemporaneous |
| Extemporaneous 270 White Soft Liquid Paraffin AFT 72 Wool fat with mineral oil 72 - X - Xarelto Xarelto 42 Xifaxan 10 XMET Maxamum 285 Xolair 213 XP Maxamum 124 Xylocaine 124 |
| Extemporaneous |
| Extemporaneous 270 White Soft Liquid Paraffin AFT 72 Wool fat with mineral oil 72 - X - Xarelto Xarelto 42 Xifaxan 10 XMET Maxamum 285 Xolair 213 XP Maxamum 286 Xylocaine 124 Xylocaine 2% Jelly 124 Xylocaine 500 124 Xyntha 39 |
| Extemporaneous |

| Zimybe | 56 |
|-------------------------------|----|
| Zinc and castor oil | 71 |
| Zinc sulphate | 35 |
| Zincaps | 35 |
| Ziprasidone1 | 37 |
| Zista2 | 50 |
| Zithromax | 96 |
| Zoladex | 93 |
| Zoledronic acid | |
| Hormone | |
| Musculoskeletal1 | 20 |
| Zoledronic acid Viatris | |
| Hormone | 85 |
| Musculoskeletal1 | 20 |
| Zopiclone 1 | |
| Zopiclone Actavis1 | 44 |
| Zostrix1 | |
| Zostrix HP1 | 25 |
| Zuclopenthixol decanoate1 | 39 |
| Zuclopenthixol hydrochloride1 | 37 |
| Zusdone1 | 37 |
| Zyban1 | 50 |
| Zypine1 | |
| Zypine ODT 1 | 37 |
| Zyprexa Relprevv1 | 38 |
| Zytiga1 | 73 |
| Zyvox1 | 06 |













