March 2025 Volume 32

Editors:

Kaye Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth

email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written from and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	gg.	_
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	35
	Cardiovascular System	44
	Dermatologicals	66
	Genito Urinary System	75
	Hormone Preparations – Systemic	82
	Infections – Agents For Systemic Use	93
	Musculoskeletal System	115
	Nervous System	121
	Oncology Agents & Immunosuppressants	151
	Respiratory System & Allergies	255
	Sensory Organs	266
	Various	271
Section C	Extemporaneous Compounds (ECPs)	273
Section D	Special Foods	275
Section I	National Immunisation Schedule	299
	Index	315

Introducing Pharmac

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ Hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ Hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the General Rules: https://pharmac.govt.nz/section-a.

SECTION B: ALIMENTARY TRACT AND METABOLISM

SECTION B: ALIMENTARY TRACT AND MET	TABOLISM			
	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg pe sachet		30	√ G	aviscon Infant
* Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (14.39)	60	G	aviscon Extra Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 ml	A	cidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) –		100		lu-Tab
Subsidy by endorsement	47.30	500 ml 473 ml	√ C	oxane alcium carbonate PAI ⁸²⁹
Only when prescribed for patients unable to swallow calc inappropriate and the prescription is endorsed accordingl		ts or whe	ere calciur	n carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a * Tab 2 mg * Cap 2 mg	10.75	400 400	✓ N ✓ <u>D</u>	odia iamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap modified-release 3 mg - Special Authority see SA1886 below - Retail pharmacy	87.60	90	✓ <u>B</u>	udesonide Te Arai

⇒SA1886 Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or

Subsidy (Manufacturer's P	Price) S	Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes: or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

HYDROCORTISONE ACETATE Rectal foam 10%, CFC-Free (14 applications)57.09	15 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
		✓ Asacol S29 S29
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa
(Asacol S29 S29 Tab 800 mg to be delisted 1 July 2025)		

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
OLSALAZINE				
Tab 500 mg	56.02	60	/	Atnahs Olsalazine S29
	93.37	100	1	Dipentum
Cap 250 mg	53.00	100	•	Dipentum
SODIUM CROMOGLICATE Cap 100 mgSULFASALAZINE	113.35	100	•	Ralicrom
* Tab 500 mg	19.49	100	✓	Salazopyrin
* Tab EC 500 mg	20.54	100	1	Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE				
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and				
cinchocaine hydrochloride 5 mg per g13.05	30 g OP	Ultraproct		
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and				
cinchocaine hydrochloride 1 mg8.61	12	Ultraproct		
HYDROCORTISONE WITH CINCHOCAINE				
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl		
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl		

Management of Anal Fissures

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

CL VCODVDDONII IM DDOMIDE

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	19.00	5	✓ Robinul
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	2.25	20	✓ Hyoscine Butylbromide (Adiramedica)
	6.35	100	✓ Buscopan
Hyoscine Butylbromide (Adiramedica) to be Principal Supply on	1 April 2025		
* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO(Buscopan Tab 10 mg to be delisted 1 April 2025)	1.91	5	✓ <u>Spazmol</u>
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	8.50	90	✓ Colofac

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL - Wastage claimable

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement.......14.58 14 ✓ Klacid

- a) Maximum of 28 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

FΑ	MOTIDINE – Only on a prescription			
*	Tab 20 mg	4.91	100	✓ Famotidine
	· ·			Hovid S29
*	Tab 40 mg	10.32	100	✓ Famotidine
	-			Hovid S29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement	CBS	10	✓ Mylan S29
	Subsidy by endorsement – Subsidised for nationts rece	viving treatment as	nart of nallia	ative care

Proton Pump Inhibitors

LANSOPRAZOLE					
* Cap 15 mg4.04	100	✓ Lanzol Relief			
* Cap 30 mg5.43	100	✓ Lanzol Relief			
OMEPRAZOLE					
For omeprazole suspension refer Standard Formulae, page 273					
* Cap 10 mg2.06	90	Omeprazole Teva			
· ·		✓ Omeprazole actavis 10			
* Cap 20 mg2.02	90	Omeprazole Teva			
. •		✓ Omeprazole actavis			
		<u>20</u>			
* Cap 40 mg3.18	90	Omeprazole Teva			
		✓ Omeprazole actavis 40			
* Powder – Only in combination42.50	5 g	✓ Midwest			
Only in extemporaneously compounded omeprazole suspension.	·				
* Inj 40 mg ampoule with diluent	5	✓ <u>Dr Reddy's</u> Omeprazole			
		✓ Ocicure S29			
PANTOPRAZOLE					
	00	✓ Danzon Poliof			
* Tab EC 20 mg	90	✓ Panzop Relief			
* Tab EC 40 mg2.74	90	✓ Panzop Relief			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mgSUCRALFATE	14.51	50	✓ G	astrodenol \$29
Tab 1 g	35.50 (48.28)	120	С	arafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail pl Tab 550 mg SA1461 Special Authority for Subsidy	,	56	√ <u>X</u>	<u>ifaxan</u>
Initial application only from a gastroenterologist, hepatologist nepatologist. Approvals valid for 6 months where the patient tolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Pract nepatologist. Approvals valid without further renewal unless repending from treatment.	has hepatic encephalo	oathy d	lespite an ad n of a gastro	lequate trial of maximum enterologist or
Diabetes				
Hyperglycaemic Agents	la a mara a su			
DIAZOXIDE - Special Authority see SA1320 below - Retail p Cap 25 mg Cap 100 mg Oral liq 50 mg per ml → SA1320 Special Authority for Subsidy	110.00	100 100 30 ml C	✓ P	roglicem \$29 roglicem \$29 5 Pharma \$29
nitial application from any relevant practitioner. Approvals hypoglycaemia caused by hyperinsulinism.	valid for 12 months whe	ere use	d for the trea	atment of confirmed
Renewal from any relevant practitioner. Approvals valid with appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE	out further renewal unle	ess noti	ified where th	he treatment remains
Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	✓ G	lucagen Hypokit
Insulin - Short-acting Preparations				
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml C		ctrapid lumulin R
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓ A	ctrapid Penfill umulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ N	ovoMix 30 FlexPen

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully Brand or idised Generic Manufacturer
NSULIN ISOPHANE			
Inj human 100 u per ml	17.68	10 ml OP	✓ Humulin NPH✓ Protaphane
Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH✓ Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70 ✓ PenMix 30 ✓ PenMix 50
Mixtard 30 Inj human with neutral insulin 100 u per ml to be deli PenMix 50 Inj human with neutral insulin 100 u per ml, 3 ml to b NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE	e delisted 1 June		Penwix 50
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml	42.66	5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml 3 ml		5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
NSULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
Inj 100 u per ml, 3 ml	94.50	5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen	94.50	5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
Inj 100 u per ml, 10 ml		1_	✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
▲ Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
Inj 100 u per ml, 10 ml	27.03	1	✓ Apidra
Inj 100 u per ml, 3 ml		5	✓ Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	✓ Apidra SoloStar
		10 ml OP 5	✓ Humalog✓ Humalog
Alpha Glucosidase Inhibitors			
•			
CARBOSE ★ Tab 50 mg ★ Tab 100 mg		90 90	✓ <u>Accarb</u> ✓ <u>Accarb</u>
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
k Tab 5 mg	7 50	100	✓ Daonil
v	7.50	100	- Duyiiii
GLICLAZIDE 烙 Tab 80 mg	20.10	500	✓ Glizido
r 1 au 00 mg	∠0.10	500	✓ <u>Glizide</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
GLIPIZIDE				
* Tab 5 mg	6.86	100	✓	<u>Minidiab</u>
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg	14.74	1,000	/ [Metformin Viatris
* Tab immediate-release 850 mg		500	✓]	Metformin Viatris
PIOGLITAZONE				
* Tab 15 mg	6.15	90	✓ 1	Vexazone
* Tab 30 mg	7.25	90	✓ :	Vexazone
* Tab 45 mg	12.00	90	✓ :	Vexazone
VILDAGLIPTIN				
Tab 50 mg	35.00	60	✓ (Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	✓ (Galvumet
Tab 50 mg with 850 mg metformin hydrochloride		60	1	Galvumet

GLP-1 Agonists

DULAGLUTIDE - Special Authority see SA2338 below - Retail pharmacy

Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.

Inj 1.5mg per 0.5 ml prefilled pen115.23 ✓ Trulicity

⇒SA2338 Special Authority for Subsidy

Note: Subsidy for patients with existing approvals prior to 1 May 2024. Approvals valid without further renewal unless notified. No new patients will be granted from 1 May 2024 until further notice.

LIRAGLUTIDE - Special Authority see SA2440 below - Retail pharmacy

- a) Maximum of 9 inj per prescription
- b)
- a) Note: Not to be given in combination with another funded GLP-1 agonist or empaqliflozin / empaqliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.
- b) Maximum of 1 pack of 3 (6 mg per ml, 3 ml) prefilled pens will be funded per month.

Inj 6 mg per ml, 3 ml prefilled pen383.72 ✓ Victoza

⇒SA2440 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empadiflozin, metformin, and vildagliptin: and
- 3 Any of the following:
 - 3.1 Patient is Māori or any Pacific ethnicity*: or
 - 3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or

continued...

3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with (empagliflozin /empagliflozin with metformin hydrochloride) unless receiving (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

SGLT2 Inhibitors

⇒SA2408 Special Authority for Subsidy

Initial application — (heart failure reduced ejection fraction) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure; and
- 2 Patient is in NYHA functional class II or III or IV; and
- 3 Either
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or
 - 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment: and
- 4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initial application — (Type 2 Diabetes) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Maori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride] for the treatment of heart failure.

EMPAGLIFLOZIN - S	pecial Authority see SA2408 on the previo	us page - Retail p	harmacy		
* Tab 10 mg		58.56	30	✓	Jardiance
* Tab 25 mg		58.56	30	✓	Jardiance
	TH METFORMIN HYDROCHLORIDE - Sp	ecial Authority see	SA2408 or	n the p	revious page – Retail
pharmacy					
* Tab 5 mg with 1,0	00 mg metformin hydrochloride	58.56	60	✓	Jardiamet
* Tab 5 mg with 500) mg metformin hydrochloride	58.56	60	✓	Jardiamet
* Tab 12.5 mg with	1,000 mg metformin hydrochloride	58.56	60	✓	Jardiamet
* Tab 12.5 mg with	500 ma metformin hydrochloride	58.56	60	1	Jardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

				_
Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes: or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips	33.69	50 test OP	✓ SensoCard
---------------------------	-------	------------	-------------

Subsidy	Fu	ly B	rand or
(Manufacturer's Price)	Subsidise	ed G	Generic
\$	Per	/ N	Manufacturer

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or liraglutide or when prescribed for a patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or liraglutide.

IIV	ocini i chi necoleo – maximum di 200 dev per prescripti	1011		
*	29 g × 12.7 mm	10.95	100	✓ B-D Micro-Fine
*	31 g × 5 mm		100	✓ B-D Micro-Fine
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED	LE - Maximum of 2	200 dev per p	orescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g x 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓ B-D Ultra Fine II
		1.36	10	
		(1.99)		B-D Ultra Fine II
		` '		

Insulin Pumps

INSULIN PUMP WITH ALGORITHM - Special Authority see SA2367 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four y	ear period.
Min basal rate 0.02 U/h	8,970.00
Min boool rate 0.1 LI/b	7 652 00

✓ mylife YpsoPump with CamAPS FX

✓ Tandem t:slim X2 with Basal-IQ

✓ Tandem t:slim
X2 with Control-IQ

⇒SA2367 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

continued...

1

(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	Subsidy	Full	y Brand or
\$ Per ✓ Manufacturer	(Manufacturer's	s Price) Subsidise	d Generic
	\$	Per ✓	Manufacturer

continued...

- 1.1 The patient has type 1 diabetes; or
- 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
- 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 6 months where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

Insulin Pump Consumables

⇒SA2380 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has type 1 diabetes; or
 - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist as likely to benefit; or
 - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
 - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 Patient has been evaluated by a diabetes multidisciplinary team for their suitability for insulin pump therapy; and
- 3 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

INSULIN PUMP CARTRIDGE - Special Authority see SA2380 above - Retail pharmacy

- a) Maximum of 5 sets per prescription
- b) Only on a prescription
- c) Maximum of 19 packs of cartridge sets will be funded per year.
- **★** Cartridge 300 U, t:lock × 10......86.00 1 OP **✓ Tandem Cartridge**

		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer	
INS	SULIN PUMP INFUSION SET (STEEL CANNULA) - Special	•		ne previous		
	a) Maximum of 5 set per prescription	·		·		
	b) Only on a prescription					
*	c) Maximum of 19 infusion sets will be funded per year. 6 mm steel needle; 60 cm tubing × 10	120.00	1 OP	1	MiniMed Sure-T	
~	o min steer needie, oo cin tubing x 10	150.00	1 01	•	MMT-864A	
*	6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	1	MiniMed Sure-T	
	•				MMT-866A	
*	8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	/	MiniMed Sure-T MMT-874A	
*	8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	•	MiniMed Sure-T MMT-876A	
(Mi	(MiniMed Sure-T MMT-866A 6 mm steel needle; 80 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-874A 8 mm steel needle; 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Sure-T MMT-876A 8 mm steel needle; 80 cm tubing × 10 to be delisted 1 October 2026)					
	SULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT le – Retail pharmacy	INSERTION) – Spe	ecial A	uthority se	e SA2380 on the previous	
μαί	a) Maximum of 5 sets per prescription					
	b) Only on a prescription					
	c) Maximum of 19 infusion sets will be funded per year.					
	5.5 mm steel cannula; straight insertion; 45 cm line × 10 with		4 00			
	10 needles	136.00	1 OP	•	mylife Orbit micro	
	5.5 mm steel needle; straight insertion; 60 cm line × 10 with 10 needles	136 00	1 OP	1	mylife Orbit micro	
*	5.5 mm steel needle; straight insertion; 80 cm line × 10 with		. 0.	-	mymo orbit mioro	
	10 needles	136.00	1 OP	1	mylife Orbit micro	
*	8.5 mm steel needle; straight insertion; 60 cm line \times 10 with			_		
	10 needles	136.00	1 OP	•	mylife Orbit micro	
*	8.5 mm steel needle; straight insertion; 80 cm line × 10 with 10 needles	136.00	1 OP	1	mylife Orbit micro	

* 6 mm steel cannula; straight insertion; 80 cm line x 10 with

* 8 mm steel cannula; straight insertion; 80 cm line × 10 with

* 6 mm steel cannula; straight insertion; 60 cm line x 10 with

* 8 mm steel cannula; straight insertion; 60 cm line x 10 with

10 needles......182.00

1 OP

1 OP

1 OP

1 OP

✓ TruSteel

✓ TruSteel

✓ TruSteel

✓ TruSteel

		Subsidy		Fully	Brand or			
		(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer			
_		T						
INS	INSULIN PUMP INFUSION SET (TEFLON CANNULA) – Special Authority see SA2380 on page 17 – Retail pharmacy							
	a) Maximum of 5 set per prescription							
	b) Only on a prescription							
*	c) Maximum of 19 infusion sets will be funded per year. 13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Silhouette			
~	To minitenon needle, oo em tabing x to	100.00	1 01	•	MMT-381A			
*	17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	1	MiniMed Silhouette			
•	, , , , , , , , , , , , , , , , , , ,				MMT-377A			
*	17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Silhouette			
	-				MMT-378A			
*	6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓	MiniMed Quick-Set			
					MMT-398A			
*	6 mm teflon needle, 45 cm blue tubing \times 10	130.00	1 OP	•	MiniMed Mio			
					MMT-941A			
*	6 mm teflon needle, 45 cm pink tubing × 10	130.00	1 OP	•	MiniMed Mio MMT-921A			
*	6 mm toflan needle 60 em blue tubing 10	100.00	1 OP	./	MiniMed Mio			
不	6 mm teflon needle, 60 cm blue tubing x 10	130.00	I OF	•	MMT-943A			
*	6 mm teflon needle, 60 cm pink tubing × 10	130.00	1 OP	1	MiniMed Mio			
	o him tollon noodio, oo on piin tabiilg x ro		. 0.		MMT-923A			
*	6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Quick-Set			
	·				MMT-399A			
*	6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓	MiniMed Mio			
					MMT-945A			
*	6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓	MiniMed Mio			
					MMT-965A			
*	6 mm teflon needle, 80 cm pink tubing x 10	130.00	1 OP	•	MiniMed Mio			
*	0 mm toflan needle 110 em tubing 10	100.00	1 OP	./	MMT-925A MiniMed Quick-Set			
不	9 mm teflon needle, 110 cm tubing × 10	130.00	I OF	•	MMT-396A			
*	9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	1	MiniMed Quick-Set			
-,-	o min tollon hoodio, oo om tabilig x 10		. 0.	•	MMT-397A			
*	9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	1	MiniMed Mio			
	,				MMT-975A			

(MiniMed Silhouette MMT-381A 13 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-377A 17 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Silhouette MMT-378A 17 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-398A 6 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-941A 6 mm teflon needle, 45 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-921A 6 mm teflon needle, 45 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-943A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-933A 6 mm teflon needle, 60 cm pink tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 60 cm blue tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-945A 6 mm teflon needle, 80 cm blue tubing to be delisted 1 October 2026) (MiniMed Mio MMT-965A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-925A 6 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-936A 9 mm teflon needle, 110 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-396A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Quick-Set MMT-397A 9 mm teflon needle, 60 cm tubing × 10 to be delisted 1 October 2026) (MiniMed Mio MMT-975A 9 mm teflon needle, 80 cm clear tubing × 10 to be delisted 1 October 2026)

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
INS	SULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION WITH INS	SERT	ON DEVICE	Special Authority see
	2380 on page 17 – Retail pharmacy				, ,
	a) Maximum of 5 sets per prescription				
	b) Only on a prescription				
	c) Maximum of 19 infusion sets will be funded per year.				
*	13 mm teflon cannula; angle insertion; insertion device; 110 c	m			
	line × 10 with 10 needles	182.00	1 OP	✓ A	utoSoft 30
*	13 mm teflon cannula; angle insertion; insertion device; 60 cn line × 10 with 10 needles	n	1 OP	✓ A	utoSoft 30
INIS	SULIN PUMP INFUSION SET (TEFLON CANNULA, FLEXIBLE	INSERTION WITH	INSFI	RTION DEV	ICE) - Special Authority
	SA2380 on page 17 – Retail pharmacy	INOLITION WITH	IIVOLI	THONDEV	opecial Authority
	a) Maximum of 5 set per prescription				
	b) Only on a prescription				
	c) Maximum of 19 infusion sets will be funded per year.				
*	6 mm teflon cannula; flexible insertion; insertion device; 46 cr	n			
•	line × 10 with 10 needles		1 OP	✓ m	ylife Inset soft
*	6 mm teflon cannula; flexible insertion; insertion device; 60 cr	n			•
•	line with integrated inserter × 10 with 10 needles		1 OP	✓ m	ylife Inset soft
*	6 mm teflon cannula; flexible insertion; insertion device; 80 cr				,
•	line x 10 with 10 needles		1 OP	✓ m	ylife Inset soft
*	9 mm teflon cannula; flexible insertion; insertion device; 60 cr				,
•	line × 10 with 10 needles		1 OP	✓ m	ylife Inset soft
*	9 mm teflon cannula; flexible insertion; insertion device; 80 cr				,
•	line × 10 with 10 needles		1 OP	✓ m	ylife Inset soft
INI	SULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH		INICE		•
	e SA2380 on page 17 – Retail pharmacy	I INSLITTION WITH	IIVOL	ITTION DE	riot) - Special Authority
500	a) Maximum of 5 sets per prescription				
	b) Only on a prescription				
	c) Maximum of 19 infusion sets will be funded per year.				
*	6 mm teflon cannula; straight insertion; insertion device;				
•••	110 cm line × 10 with 10 needles	182.00	1 OP	✓ A	utoSoft 90
*	6 mm teflon cannula; straight insertion; insertion device; 60 ci				
-,-	line × 10 with 10 needles		1 OP	✓ Δ	utoSoft 90
*	9 mm teflon cannula; straight insertion; insertion device;			- 70	4.000.1.00
-,-	110 cm line × 10 with 10 needles	182 00	1 OP	✓ A	utoSoft 90
*	9 mm teflon cannula; straight insertion; insertion device; 60 cr			- 70	4.0001.00
-,-	line × 10 with 10 needles		1 OP	✓ Δ	utoSoft 90
INIC			. •.		
	SULIN PUMP INFUSION SET (TEFLON CANNULA, VARIABLI tail pharmacy	E INSERTION) - SP	eciai	Authority See	e SA2360 on page 17 -
ne	a) Maximum of 5 set per prescription				
	b) Only on a prescription				
	c) Maximum of 19 infusion sets will be funded per year.				
*	13 mm teflon cannula; variable insertion; 60 cm line × 10 with				
*	10 needles		1 OP	✓ V	ariSoft
	TO TICCUICS	102.00	1 01	- V	anoon

MMT-332A

	Subsidy (Manufacturer's Price) \$	Su Per	Fully Brand or ibsidised Generic Manufacture	r
INSULIN PUMP RESERVOIR - Special Authority see SA2380	on page 17 – Retail ph	armacy		
a) Maximum of 9 sets per prescription				
b) Only on a prescription				
 c) Maximum of 36 packs of resevoir sets will be funded per 	,			
* 10 × 1.6 ml glass reservoir for YpsoPump	50.00	1 OP	✓ mylife YpsoP Reservoir	ump
* 10 × luer lock conversion cartridges 1.8 ml for Paradigm p	umps50.00	1 OP	✓ ADR Cartridg	e 1.8
* Cartridge for 7 series pump; 3.0 ml x 10	50.00	1 OP	✓ MiniMed	
			3.0 Reservo	oir

(ADR Cartridge 1.8 10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps to be delisted 1 October 2026) (MiniMed 3.0 Reservoir MMT-332A Cartridge for 7 series pump; 3.0 ml × 10 to be delisted 1 October 2026)

Continuous Glucose Monitor

CONTINUOUS GLUCOSE MONITOR (INTEROPERABLE) – Special Authority see SA2371 below – Retail pharmacy Only on a prescription

Maximum of 28 dev will be funded per year.

⇒SA2371 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has type 1 diabetes; or
 - 1.2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
 - 1.3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
 - 1.4 The patient has atypical inherited forms of diabetes; and
- 2 In the opinion of the treating relevant practitioner the patient would benefit from an Automated Insulin Delivery (AID) system.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

CONTINUOUS GLUCOSE MONITOR (STANDALONE) - Special Authority see SA2370 on the next page - Retail pharmacy Only on a prescription

★ Sensor (Dexcom ONE+) - Maximum of 9 dev per prescription81.00
 Maximum of 40 dev will be funded per year.

Subsid	idy Fully	Brand or
(Manufacture		
\$	Per 🗸	Manufacturer

⇒SA2370 Special Authority for Subsidy

Initial application — (type 1 diabetes) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 The patient has type 1 diabetes; or
- 2 The patient has permanent neonatal diabetes or specific monogenic diabetes subtypes with insulin deficiency, considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit; or
- 3 The patient has Type 3c diabetes considered by the treating endocrinologist or relevant secondary health care professional as practicable, as likely to benefit (Type 3c diabetes includes insulin deficiency due to pancreatectomy, insulin deficiency secondary to cystic fibrosis or pancreatitis); or
- 4 The patient has atypical inherited forms of diabetes.

Renewal — (type 1 diabetes) from any relevant practitioner. Approvals valid for 2 years where the patient is continuing to derive benefit according to the treatment plan agreed at induction.

Digestives Including Enzymes

PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase		
10,000 Ph Eur U, total protease 600 Ph Eur U)34.93	100	✓ Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase		
25,000 Ph Eur U, total protease 1,000 Ph Eur U)94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase		
3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph		
Eur U)	20 g OP	Creon Micro
URSODEOXYCHOLIC ACID - Special Authority see SA1739 below - Retail pharm	nacy	
Cap 250 mg33.95	100	✓ <u>Ursosan</u>

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has been diagnosed with Alagille syndrome: or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications

ColoxvI

Subsidy	e)	Fully	Brand or
(Manufacturer's Pric		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels

Laxatives

Bulk-forming Agents

ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription			
W Douglas for oral calm	20.00	E00 ~ OD	./ Vana

* Powder for oral soln	20.00	500 g OP	✓ Konsyl-D
Faecal Softeners			

DO	CUSATE SODIUM - Only on a prescription		
*	Tab 50 mg	100	✓ (

Tab 50 mg with sennosides 8 mg.......3.50 200 ✓ Laxsol POLOXAMER – Only on a prescription

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE - Special Authority see SA1691 below - Retail pharmacy

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Fither:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)) S Per	ubsidised •	Generic Manufacturer
Osmotic Laxatives				
GLYCEROL * Suppos 2.8/4.0 g - Only on a prescription	10.39	20	1	Lax-suppositories Glycerol
LACTULOSE – Only on a prescription * Oral liq 10 g per 15 ml	3.61	500 ml	•	<u>Laevolac</u>
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIO		SODIUN	I CHLOF	RIDE
Powder for oral soln 13.125 g with potassium chloride 46.6 m sodium bicarbonate 178.5 mg and sodium chloride 350.7	•	30	✓	<u>Molaxole</u>
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	•	Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE		iption		
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml		50	✓	Micolette
Stimulant Laxatives				
BISACODYL – Only on a prescription * Tab 5 mg * Suppos 10 mg SENNA – Only on a prescription * Tab, standardised		200 10 100 20	•	Bisacodyl Viatris Lax-Suppositories Senokot
SODIUM PICOSULFATE – Special Authority see SA2053 below Oral soln 7.5 mg per ml		0 ml OF		Senokot Dulcolax SP Drop
		· ···· • ·		

⇒SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and

|--|

continued...

- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
 - 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
 - 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
 - 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority see SA2042 below - Retail pharmacy

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg	CBS	50	✓ Solgar
Powder		400 g	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Por 🗸	Manufacturer

continued...

- 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
- 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below - Retail pharmacy

Cap 120 mg	CBS	30	Solgar
Cap 160 mg	CBS	60	Go Healthy

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts: or

	Subsidy (Manufacturer's Price) \$	F Subsidis Per	ully sed	Brand or Generic Manufacturer
continued				

- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
 - 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
 - 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
 - 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE - Special Authority see SA1695 below	- Retail pharmacy		
Inj 100 U per ml, 5 ml vial	1,335.16	1	Aldurazyme

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Fither:

L

- 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
- 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA204	10 below - Retail pharmacy		
Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg		60	✓ Balance
		300	✓ Metabolics
Oral lig 1 g per 10 ml	CBS	118 ml	✓ Carnitor S29
1 31			✓ Novitium Sugar
			Free S29
Oral lig 500 mg per 10 ml	CBS	300 ml	✓ Balance

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN – Special Authority see SA2041 on the next Tab 100 mg	, ,	100	✓ Country Life ✓ Puritan's Pride
			Vitamin B-2 100 mg S29
Cap 100 mg	CBS	100	✓ Solgar

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE - Special Authority see SA1989 below - Retail pharmacy

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy
Soln 100 mg per mlCBS

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 on the next page - Retail pharmacy

Grans 483 mg per g......2,016.00 174 g OP ✓ Pheburane

100 ml

✓ Amzoate S29

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	✓ Life Extension

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE − Special Authority see SA2324 below − Retail pharmacy
Cap 250 mg......2,022.00 100 ✓ Trientine Waymade

⇒SA2324 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Wilson disease: and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit: and
- 3 Treatment with zinc has been trailled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

Gaucher's Disease

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher

	Subsidy (Manufacturer's Pri	ice) Subs	Fully sidised	Brand or Generic Manufacturer
continued	Ψ	1 61		iviai idiactal Ci
disease; or				
3.5 Patient is a child and has experienced growth failure6-12 month period; and	re with significant	decrease in p	ercenti	le linear growth over a
4 Taliglucerase alfa is to be administered at a dose no great whole vial (200 units).	ter than 30 unit/kg	every other	week ro	unded to the nearest
Note: Indication marked with * is an unapproved indication				
Renewal only from a metabolic physician or any relevant practitic Approvals valid for 3 years for applications meeting the following All of the following:		imendation of	a meta	bolic physician.
 Patient has demonstrated a symptomatic improvement an symptoms for which therapy was started; and 		•		
2 Patient has demonstrated a clinically objective improveme	ent or no deteriora	tion in haemo	globin l	evels, platelet counts and
liver and spleen size; and 3 Radiological (MRI) signs of bone activity performed at two demonstrate no deterioration shown by the MRI, compare or adjusted dose; and				
Patient has not developed another medical condition that l ERT; and	might reasonably	be expected	to comp	promise a response to
5 Patient is adherent with regular treatment and taligluceras every other week rounded to the nearest whole vial (200 u		ministered at a	a dose i	no greater than 30 unit/kg
Mouth and Throat				
Agents Used in Mouth Ulceration				
BENZYDAMINE HYDROCHLORIDE				
Soln 0.15% - Higher subsidy of \$22.60 per 500 ml with				
Endorsement		500 ml	-	s:##1 =
Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly.	(22.60) s oral mucositis as	a result of tr		oifflam It for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN				
Paste	17.20	56 g OP	√ S	tomahesive
	4.55	15 g OP	_	A b
	(7.90) 1.52	5 g OP	C)rabase
	(3.60)	Jyor	C)rabase
Powder	` ,	28 g OP		
	(10.95)	-	S	tomahesive
TRIAMCINOLONE ACETONIDE Paste 0.1%	5.49	5 g OP	✓ <u>K</u>	enalog in Orabase
Oropharyngeal Anti-infectives				
AMPHOTERICIN B				
Lozenges 10 mg	5.86	20	√ F	ungilin
MICONAZOLE				

Oral gel 20 mg per g......5.19

Oral liq 100,000 u per ml2.22

✓ Decozol

✓ Nilstat

40 g OP

24 ml OP

MICONAZOLE

NYSTATIN

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
Vitamins				
Vitamin B				
HYDROXOCOBALAMIN				
* Inj 1 mg per ml, 1 ml ampoule - Up to 6 inj available on a	PSO2.46	3		Cobal-B12 S29 Vita-B12
	3.95			Hydroxocobalamin Panpharma
	4.10	5		Cobalin-H S29 Neo-Cytamen S29 S29
	8.20	10	•	Vitarubin Depot Injection \$29
(Cobal-B12 S29 Inj 1 mg per ml, 1 ml ampoule to be delisted (Vita-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 July 2 (Cobalin-H S29 Inj 1 mg per ml, 1 ml ampoule to be delisted (Neo-Cytamen S29 S29 Inj 1 mg per ml, 1 ml ampoule to be (Vitarubin Depot Injection S29 Inj 1 mg per ml, 1 ml ampoule PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose	2025) 1 July 2025) delisted 1 July 2025)	025)		
 b) Only on a prescription Tab 25 mg - No patient co-payment payable 	3 43	90	/	Vitamin B6 25
* Tab 50 mg		500		Pyridoxine multichem
THIAMINE HYDROCHLORIDE – Only on a prescription * Tab 50 mg	4.65	100	,	Thiamine multichem
VITAMIN B COMPLEX	4.00	100	•	Thamme manuchem
* Tab, strong, BPC	11.25	500	✓	Bplex
Vitamin C				
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription				
* Tab 100 mg	12.50	500	•	<u>Cvite</u>
Vitamin D				
ALFACALCIDOL * Cap 0.25 mcg	26.32	100		One-Alpha One-Alpha S29 S29
* Cap 1 mcg * Oral drops 2 mcg per ml	60.68	100 20 ml C	1	One-Alpha One-Alpha

Subs (Manufactur			Fully Subsidised	Generic
\$		Per		Manufacturer
CALCITRIOL				
* Cap 0.25 mcg7.8	9	100		Calcitriol-AFT Calcitriol-AFT
* Cap 0.5 mcg13.6	8	100		S29 S29 Calcitriol-AFT Calcitriol-AFT
00/ 50 // 0/55 00/				S29 S29
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescription3.6 * Oral liq 188 mcg per ml (7,500 iu per ml)9.0		12 ml OF		<u>Vit.D3</u> Clinicians
Multivitamin Preparations				
MULTIVITAMIN RENAL - Special Authority see SA1546 below - Retail pha	ırmacy			

✓ Clinicians Renal Vit.

⇒SA1546 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy

200 a OP ✓ Paediatric Seravit

⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINS

*	Tab (BPC cap strength)	1,000	✓ Mvite
*	Cap (fat soluble vitamins A, D, E, K) - Special Authority see		
	SA1720 below – Retail pharmacy23.40	60	✓ Vitabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency: or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

	ALIMENTAN	IINACI	ANI	DIVIETABOLISIVI
	Subsidy (Manufacturer's Price) \$) Subsi	Fully dised	Brand or Generic Manufacturer
Minerals				
Calcium				
CALCIUM CARBONATE * Tab 1.25 g (500 mg elemental) * Tab eff 1.25 g (500 mg elemental) – Subsidy by endorsem	ent260.00	250 100	1	Calci-Tab 500 Calcium 500 mg Hexal §29
Subsidy by endorsement – Only when prescribed for p considered unsuitable.	aediatric patients (< 5	years) wher	e cal	cium carbonate oral liquid is
CALCIUM GLUCONATE * Inj 10%, 10 ml ampoule	32.00	10	√	Max Health - Hameln 629
lodine				
POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine)	5.99	90	√ <u>!</u>	NeuroTabs
Iron				
FERROUS FUMARATE * Tab 200 mg (65 mg elemental)	3.49	100	✓ <u>I</u>	Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg FERROUS SULFATE	5.98	100	√ <u>[</u>	Ferro-F-Tabs
** Tab long-acting 325 mg (105 mg elemental) ** Oral liq 30 mg (6 mg elemental) per 1 ml	9.25	30 250 ml 500 ml	✓ i	Ferrogra <u>d</u> Ferro-Liquid Ferodan
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority Inj 50 mg per ml, 10 ml vial		Retail pharm 1		Ferinject
SA2394 Special Authority for Subsidy Initial application — (Anaemia) from any relevant practitioner following criteria: All of the following: 1 Patient has been diagnosed with anaemia; and 2 Any of the following: 2.1 Serum ferritin level is 20 mcg/L or less; or 2.2 Both:	r. Approvals valid for a	3 months for	appli	ications meeting the
2.2.1 Sarum farritin is hatween 20 and 50 mgg/	l · and			

- 2.2.1 Serum ferritin is between 20 and 50 mcg/L; and
- 2.2.2 C-Reactive Protein (CRP) is at least 5 mg/L; or
- 2.3 Patient has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 3 Any of the following:
 - 3.1 Oral iron treatment has proven ineffective; or
 - 3.2 Oral iron treatment has resulted in dose-limiting intolerance; or
 - 3.3 Rapid correction of anaemia is required.

Renewal — (Anaemia) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Both:

- 1 Patient continues to have anaemia with a serum ferritin level of 20 mcg/L, or less or between 20 and 50 mcg/L with CRP of at least 5 mg/L, or has chronic inflammatory disease with symptoms of anaemia despite normal iron levels; and
- 2 A trial (or re-trial) with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
- 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%33.60	355 ml	✓ Phillips Milk of Magnesia ©29
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule	10 10	✓ <u>Martindale</u> ✓ Inresa \$29
Zinc		
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)11.00	100	✓ Zincaps

✓ Ferrosia

5

BLOOD AND BLOOD FORMING ORGANS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA2266 Special Authority for Subsidy

Initial application — (chronic renal failure) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — **(chronic renal failure)** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA2266 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	Binocrit
Inj 10,000 iu in 1 ml, syringe		6	Binocrit
Inj 40,000 iu in 1 ml, syringe		1	Binocrit

BLOOD AND BLOOD FORMING ORGANS

	Subsidy (Manufacturer's Price) \$		Fully ised ✓	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID * Tab 0.8 mg	26.60	1,000		olic Acid multichem
* Tab 5 mg Oral liq 50 mcg per ml		100 5 ml OP	_	olic Acid Viatris omed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

reaters Group in conjunction with the National	Haemophilia Management gro	up.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
lnj 4,000 iu vial	9,800.00	1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 b Wastage claimable	pelow – Retail pharmacy		
Tab 25 mg	1,550.00	28	✓ Revolade
Tab 50 mg	3,100.00	28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

Ε

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopad treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre

Sub	sidy Fu	ully Brand or
(Manufactu		sed Generic
	S Per	✓ Manufacturer

continued...

and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA2272 below

Inj 30 mg in 1 ml vial	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	•	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial	12,492.00	1	✓ Hemlibra
Inj 150 mg in 1 ml vial	17,846.00	1	✓ Hemlibra

⇒SA2272 Special Authority for Subsidy

Initial application — (Severe Haemophilia A with or without FVIII inhibitors) only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	S Per	ubsidised •	Generic Manufacturer
FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xph	narm]			
For patients with haemophilia. Preferred Brand of bypass				
is managed by the Haemophilia Treaters Group in conjunc	ction with the National H	aemop	hilia Man	agement Group.
Inj 500 U	1,315.00	1	1	FEIBA NF
Inj 1,000 U	2,630.00	1	✓	FEIBA NF
Inj 2,500 U	6,575.00	1	✓	FEIBA NF
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xp	harm]			
For patients with haemophilia. Rare Clinical Circumstance		e recon	nbinant fa	ctor VIII. Access to funded
treatment is managed by the Haemophilia Treaters Group	in conjunction with the	Nationa	ıl Haemo	ohilia Management Group
subject to criteria.	•		·	
Inj 250 iu prefilled syringe	287.50	1	1	Xyntha
Inj 500 iu prefilled syringe	575.00	1	1	Xyntha
Inj 1,000 iu prefilled syringe	1,150.00	1	1	Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	✓ :	Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	1	Xyntha
NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xphai	rm]			
For patients with haemophilia. Access to funded treatmer		emophi	ilia Treate	ers Group in conjunction
with the National Haemophilia Management Group.				, , ,
Inj 1,000 iu vial	870.00	1	1	RIXUBIS
Ini 2.000 iu vial		1	_	RIXUBIS
Inj 3,000 iu vial	2,610.00	1		RIXUBIS
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE)	_ [Ynharm]			
For patients with haemophilia. Preferred Brand of short h		or VIII	Access to	funded treatment is
managed by the Haemophilia Treaters Group in conjunction				
Inj 500 iu vial		1		Advate
Inj 1,000 iu vial		1		Advate
Inj 2,000 iu vial		1	_	Advate
Inj 3,000 iu vial		1		Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENA	•			
For patients with haemophilia. Rare Clinical Circumstance		a racon	nhinant fa	ctor VIII Access to funder
treatment is managed by the Haemophilia Treaters Group				
subject to criteria.	in conjunction with the	valione	ii i iaciiio _i	orilla mariagement Group
Inj 250 iu vial	237 50	1	1	Kogenate FS
Inj 500 iu vial		i		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial		i		Kogenate FS
Inj 3,000 iu vial		1		Kogenate FS
• •	•	•	- '	Acgoniato i o
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR V For patients with haemophilia A receiving prophylaxis trea		d +=aa+=	nantia m	anagad by the Heamanhill
Treaters Group in conjunction with the National Haemoph		u ireair	nent is m	anaged by the Haemophii
		4	./	A duma vata
Inj 1,000 iu vial		1		Adynovate
Inj 2,000 iu vial	∠,400.00	1	•	Adynovate
SODIUM TETRADECYL SULPHATE		_		
* Inj 3% 2 ml		5		
	(73.00)			Fibro-vein
TRANEXAMIC ACID				
TRANEXAMIC ACID Tab 500 mg	10.45	60	•	Mercury Pharma

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer	
Vitamin K					
PHYTOMENADIONE Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO		5 5	• •	onakion MM onakion MM	

Antithrombotic Agents

Antiplatelet Agents

•			
ASPIRIN	12.65	990	✓ Ethics Aspirin EC
CLOPIDOGREL	12.00	990	Ettiles Aspiriii Ee
* Tab 75 mg	5.07	84	✓ Arrow - Clopid
DIPYRIDAMOLE	10.00	00	4 P. danser O.D.
* Tab long-acting 150 mg		60	✓ Pytazen SR
TICAGRELOR – Special Authority see SA1955 below	, ,		
* Tab 90 mg	20.35	56	 Ticagrelor Sandoz

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

oun.

- 1 Fither:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (**Percutaneous coronary intervention with stent deployment**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA215	2 below – Retail pharmacy		
Inj 20 mg in 0.2 ml syringe	21.90	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe	29.74	10	✓ Clexane
Inj 60 mg in 0.6 ml syringe		10	✓ Clexane
Inj 80 mg in 0.8 ml syringe		10	✓ Clexane
Inj 100 mg in 1 ml syringe	70.91	10	✓ Clexane
Inj 120 mg in 0.8 ml syringe	88.11	10	✓ Clexane Forte
Ini 150 mg in 1 ml syringe		10	✓ Clexane Forte

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant

(Mani	Subsidy	Fully	Brand or
	ufacturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM

Inj 1,000 iu per ml, 10 ml vial	127.44	25	✓ Pfizer S29
Inj 1,000 iu per ml, 5 ml ampoule	25.49	10	✓ Wockhardt S29
	103.70		✓ Wockhardt PSF S29
	127.44	50	✓ Pfizer
Inj 5,000 iu per ml, 5 ml vial	83.00	10	✓ <u>Heparin Sodium</u> <u>Panpharma</u>
Inj 5,000 iu per ml, 1 ml	70.33	5	✓ Hospira
Inj 25,000 iu per ml, 0.2 ml		5	✓ Hospira
	42.40		✓ Heparin DBL S29
	482.20	50	✓ Heparin DBL S29
HEPARINISED SALINE			•
Inj 10 iu per ml, 5 ml	96.91	50	✓ Pfizer
Oral Anticoagulants			
DABIGATRAN			
Cap 75 mg - No more than 2 cap per day	27.99	60	✓ Pradaxa
Cap 110 mg	27.99	60	✓ Pradaxa
Cap 150 mg	27.99	60	✓ Pradaxa
RIVAROXABAN			
Tab 10 mg - No more than 1 tab per day	15.60	30	✓ Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28	✓ Xarelto
Tab 20 mg	14.56	28	✓ Xarelto
WARFARIN SODIUM			
Note: Marevan and Coumadin are not interchangeable.			
* Tab 1 mg	3.46	50	✓ Coumadin
	7.50	100	✓ Marevan
* Tab 2 mg	4.31	50	✓ Coumadin
* Tab 3 mg	12.00	100	✓ Marevan
* Tab 5 mg	5.93	50	Coumadin

100

13.50

✓ Marevan

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

Blood Colony-stimulating Factors

FILGRASTIM - Special Authority see SA1259 below - Retail pharm	acy		
Inj 300 mcg per 0.5 ml prefilled syringe	86.60	10	✓ Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	133.72	10	✓ Nivestim

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×109/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

GLUCOSE [DEXTROSE] * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO POTASSIUM CHLORIDE		5 1	✓ Biomed ✓ Biomed
* Inj 75 mg per ml, 10 ml	65.00	50	✓ Juno✓ LumaCina✓ Pfizer S29
SODIUM BICARBONATE			
Inj 8.4%, 50 ml	24.70	1	Biomed
a) Up to 5 inj available on a PSO b) Not in combination Inj 8.4%, 100 ml	25.31	1	✓ Biomed
a) Up to 5 inj available on a PSO b) Not in combination			

	Subsidy (Manufacturer's Pri	ce) Subs	Fully	Brand or Generic
	\$	Per	√	Manufacturer
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Not funded for nebulise	r use except when	used in conju	unction	with an antibiotic intende
for nebuliser use.	4.50	500 ··· l		N4
Inj 0.9%, bag – Up to 2000 ml available on a PSO		500 ml	_	Baxter
Out of an analysis of an analysis of an analysis of	1.58	1,000 ml		Baxter
Only if prescribed on a prescription for renal dialysis, ma for emergency use. (500 ml and 1,000 ml packs)	iternity or post-nat	ai care in the	nome (or the patient, or on a PS
Inj 23.4% (4 mmol/ml), 20 ml ampoule	40.15	5	√ E	Biomed
For Sodium chloride oral liquid formulation refer Standar		273		
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO	4.00	20	√ F	resenius Kabi
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PSO	5.25	50	√ F	resenius Kabi
Inj 0.9%, 20 ml ampoule		20	√ F	resenius Kabi
OTAL PARENTERAL NUTRITION (TPN)				
Infusion	CBS	1 OP	√ T	'PN
VATER			•	• • • • • • • • • • • • • • • • • • • •
On a prescription or Practitioner's Supply Order only w				
3) When used in the extemporaneous compounding of ey4) When used for the dilution of sodium chloride soln 7%		atients only.		
	for cystic fibrosis p	patients only. 50 20		Multichem Fresenius Kabi
When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule – Up to 5 inj available on a PSO	for cystic fibrosis p	50		
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule – Up to 5 inj available on a PSOInj 20 ml ampoule – Up to 5 inj available on a PSO	for cystic fibrosis p	50		
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule – Up to 5 inj available on a PSOInj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE	for cystic fibrosis p7.60 5.00	50 20	√ <u>F</u>	resenius Kabi
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule – Up to 5 inj available on a PSO Inj 20 ml ampoule – Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder	for cystic fibrosis p7.60 5.00	50	√ <u>F</u>	
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule – Up to 5 inj available on a PSO	for cystic fibrosis p	50 20 300 g OP	✓ <u>F</u>	resenius Kabi Calcium Resonium
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO Inj 20 ml ampoule — Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder	for cystic fibrosis p7.605.00169.859.53	50 20	✓ <u>F</u>	resenius Kabi
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.605.00169.859.53	50 20 300 g OP	✓ <u>F</u>	calcium Resonium
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO Inj 20 ml ampoule — Up to 5 inj available on a PSO Oral Administration CALCIUM POLYSTYRENE SULPHONATE Powder	for cystic fibrosis p7.605.00169.859.53	50 20 300 g OP	✓ <u>F</u>	calcium Resonium
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.605.00169.859.53	50 20 300 g OP	✓ <u>F</u>	calcium Resonium
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.605.00169.859.53	50 20 300 g OP	✓ <u>F</u>	Calcium Resonium Clectral Lydralyte - Lemonade
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.605.00169.859.53	50 20 300 g OP	✓ <u>F</u>	calcium Resonium
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.605.00169.859.53	50 20 300 g OP 50 1,000 ml OP	✓ <u>F</u>	Calcium Resonium Clectral Lydralyte - Lemonade
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.60169.859.536.53	50 20 300 g OP 50 1,000 ml OP	✓ <u>F</u>	Calcium Resonium Clectral Lydralyte - Lemonade
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.605.00169.859.536.5382.5082.50	50 20 300 g OP 50 1,000 ml OP	✓ F	Calcium Resonium Clectral Lydralyte - Lemonade
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.605.00169.859.536.5382.5082.50	50 20 300 g OP 50 1,000 ml OP	✓ E ✓ E ✓ E	Calcium Resonium Clectral Hydralyte - Lemonade Phosphate Phebra
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.605.00169.859.536.5382.5082.50	50 20 300 g OP 50 1,000 ml OP	✓ E ✓ E ✓ E	Calcium Resonium Clectral Hydralyte - Lemonade Phosphate Phebra
4) When used for the dilution of sodium chloride soln 7% Inj 10 ml ampoule — Up to 5 inj available on a PSO	for cystic fibrosis p7.60169.859.536.5382.5082.5015.35	50 20 300 g OP 50 1,000 ml OP	✓ Ē ✓ C ✓ E ✓ F	Calcium Resonium Clectral Hydralyte - Lemonade Phosphate Phebra

✓ Resonium-A

454 g OP

Powder84.65

SODIUM POLYSTYRENE SULPHONATE

Alpha-Adrenoceptor Blockers		(Manufacturer's Price)		ıbsidised	Generic
Alpha Adrenoceptor Blockers DOXAZOSIN * Tab 2 mg		\$	Per		Manufacturer
DOXAZOSIN # Tab 2 mg	Alpha-Adrenoceptor Blockers				
# Tab 2 mg	Alpha Adrenoceptor Blockers				
# Tab 4 mg					
## Cap 10 mg	•				
# Cap 10 mg	· ·	20.94	500	•	Doxazosin Clinect
Clibenzyline Cap 10 mg to be delisted 1 July 2025					
Cibenzyline	* Cap 10 mg				
PRAZOSIN * Tab 1 mg 5.53 100	(Discounting of the control of the delicated to the 0005)	216.67	100	•	Dibenzyline \$29
* Tab 1 mg					
# Tab 2 mg		F F0	100		Awatau Duanasin
* Tab 2 mg	* Tab T mg	5.53	100	•	
* Tab 2 mg		0.00		./	
# Tab 5 mg	* Tab 2 mg		100		•
# Tab 5 mg	- 145 E 119		100	-	
* Tab 5 mg		13.29		1	
S29 S29 Minipress Mini	* Tab 5 mg		100		•
* Cap 1 mg	· ·				S29 S29
* Cap 2 mg		22.00		1	Minipress S29
* Cap 5 mg	* Cap 1 mg	15.40	100	1	Prazosin Mylan S29
ACE Inhibitors CAPTOPRIL * Oral liq 5 mg per ml	* Cap 2 mg	15.58	100	1	Prazosin Mylan S29
ACE Inhibitors CAPTOPRIL * Oral liq 5 mg per ml	* Cap 5 mg	23.32	100	✓	Prazosin Mylan S29
ACE Inhibitors CAPTOPRIL * Oral liq 5 mg per ml	Agents Affecting the Benin-Angiotensin System	m			
CAPTOPRIL * Oral liq 5 mg per ml	Agents Affecting the Refill-Affglotensin System	11			
 ★ Oral liq 5 mg per ml	ACE Inhibitors				
Oral liquid restricted to children under 12 years of age. CILAZAPRIL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril. ** Tab 0.5 mg					
CILAZAPRIL — Subsidy by endorsement Subsidy by endorsement — Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril. * Tab 0.5 mg		86.00 10	0 ml OP		DP-Captopril
Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril. * Tab 0.5 mg					
endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril. # Tab 0.5 mg		- 4-1-1	1 11-	0001	
dispensing of cilazapril. # Tab 0.5 mg					
* Tab 0.5 mg		Silption as endorsed w	niere ure	TIC CAISI	s a record or prior
Tab 5 mg		2.69	90	✓	Zapril
(Zapril Tab 0.5 mg to be delisted 1 April 2025) (Zapril Tab 2.5 mg to be delisted 1 April 2025) (Zapril Tab 5 mg to be delisted 1 April 2025) ENALAPRIL MALEATE ★ Tab 5 mg	* Tab 2.5 mg	5.79	90	✓	Zapril
(Zapril Tab 2.5 mg to be delisted 1 April 2025) (Zapril Tab 5 mg to be delisted 1 April 2025) ENALAPRIL MALEATE ★ Tab 5 mg 1.75 90 ✓ Acetec ★ Tab 10 mg 1.97 90 ✓ Acetec	G	10.05	90	•	Zapril
(Zapril Tab 5 mg to be delisted 1 April 2025) ENALAPRIL MALEATE ★ Tab 5 mg 1.75 90 ✓ Acetec ★ Tab 10 mg 1.97 90 ✓ Acetec	, ,				
ENALAPRIL MALEATE * Tab 5 mg					
* Tab 5 mg 1.75 90 ✓ Acetec * Tab 10 mg 1.97 90 ✓ Acetec					
★ Tab 10 mg		1 75	90	1	Acatac
•	•				
	ŭ				

Subsidy

Fully

Brand or

		Subsidy		Fully Brand or
		(Manufacturer's Price)	Per	Subsidised Generic
		\$	Per	✓ Manufacturer
	INOPRIL	44.07	••	
K	Tab 5 mg	11.07	90	Ethics Lisinopril
	T 1 40	44.07	•	Teva Lisinopril
*	Tab 10 mg	11.6/	90	Ethics Lisinopril
	Tab 00	14.00	00	✓ <u>Teva Lisinopril</u>
*	Tab 20 mg	14.69	90	✓ Ethics Lisinopril
				✓ <u>Teva Lisinopril</u>
	RINDOPRIL			
	Tab 2 mg		30	✓ <u>Coversyl</u>
	Tab 4 mg		30	Coversyl
K	Tab 8 mg	3.94	30	✓ Coversyl
Ų	INAPRIL			
ĸ	Tab 5 mg	10.24	90	✓ Arrow-Quinapril 5
K	Tab 10 mg	12.51	90	✓ Arrow-Quinapril 10
K	Tab 20 mg	14.83	90	✓ Arrow-Quinapril 20
RΑ	MIPRIL			
	Cap 1.25 mg	17.25	90	✓ Tryzan
	Cap 2.5 mg		90	✓ Tryzan
	Cap 5 mg		90	✓ Tryzan
	Cap 10 mg		90	✓ Tryzan
A	ngiotensin II Antagonists			
CA	NDESARTAN CILEXETIL			
	Tab 4 mg	2.68	90	✓ Candestar
	Tab 8 mg		90	✓ Candestar
k	Tab 16 mg		90	✓ Candestar
k	Tab 32 mg		90	✓ Candestar
	SARTAN POTASSIUM			
-	Tab 12.5 mg	2 00	84	✓ Losartan Actavis
	Tab 25 mg		84	✓ Losartan Actavis
	Tab 50 mg		84	✓ Losartan Actavis
	Tab 100 mg		84	✓ Losartan Actavis
	145 166 Hg		٠.	= <u>=oourtain /iotavio</u>
A	ngiotensin II Antagonists with Diuretics			
À	NDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDI			
K	Tab 16 mg with hydrochlorothiazide 12.5 mg	4.10	30	✓ APO-Candesartan HCTZ 16/12.5
K	Tab 32 mg with hydrochlorothiazide 12.5 mg	5.25	30	✓ APO-Candesartan
				HCTZ 32/12.5
0	SARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE			
	Tab 50 mg with hydrochlorothiazide 12.5 mg	4.00	30	✓ <u>Arrow-Losartan &</u> <u>Hydrochlorothiazide</u>
A	ngiotensin II Antagonists with Neprilysin Inh	ibitors		
Δ	CUBITRIL WITH VALSARTAN - Special Authority see SA2	302 on the next nage —	Ret	ail nharmacy
<i>γ</i> ¬'	Tab 24.3 mg with valsartan 25.7 mg		56	✓ Entresto 24/26
	Tab 40.6 mg with valoartan E1.4 mg	400.00	50	Fintreste 40/E1

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Tab 48.6 mg with valsartan 51.4 mg190.00

Tab 97.2 mg with valsartan 102.8 mg190.00

✓ Entresto 49/51

✓ Entresto 97/103

56

56

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

⇒SA2302 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 121

Antiarrhythmics

To inglifedante hydrochiende foldr to treat to occupant, fundoculous, ecodi, pe	ago III	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg3.49	30	✓ Aratac
▲ Tab 200 mg4.49	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a PSO9.12	6	✓ Cordarone-X
15.22	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a		
PSO16.10	10	✓ Hikma S29
		✓ Juno S29
		✓ <u>Martindale</u>
DIGOXIN		
* Tab 62.5 mcg - Up to 30 tab available on a PSO	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO16.90	240	✓ Lanoxin
* Oral lig 50 mcg per ml	60 ml	✓ Lanoxin
		Lanoxin Paediatric Elixir
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan
55.90	84	✓ Rythmodan -
33.00	3.	Cheplafarm S29

(1	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	
FLECAINIDE ACETATE				
▲ Tab 50 mg		60		Flecainide BNM
▲ Cap long-acting 100 mg	35.78	90	•	Flecainide Controlled Release Teva
▲ Cap long-acting 200 mg	54.28	90	✓	Flecainide Controlled Release Teva
Inj 10 mg per ml, 15 ml ampoule	102.79	5	1	Almarytm \$29
	108.16			Tambocor Tambocor German S29
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg	162.00	100	1	Teva S29
A Cap 250 mgPROPAFENONE HYDROCHLORIDE	202.00	100	✓	Teva S29
▲ Tab 150 mg	40.90	50	✓	Rytmonorm
Antihypotensives MIDODRINE – Special Authority see SA1474 below – Retail pharm	nacy			
Tab 2.5 mg	•	100		MAR-Midodrine S29 <u>Midodrine</u> <u>Medsurge</u>

■SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

100

✓ MAR-Midodrine S29
✓ Midodrine
Medsurge

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATENOLOL * Tab 50 mg 11.00 * Tab 100 mg 18.50 * Oral liq 25 mg per 5 ml 49.85 Restricted to children under 12 years of age.	500 500 300 ml OP	✓ <u>Viatris</u> ✓ <u>Atenolol Viatris</u> ✓ Atenolol AFT
BISOPROLOL FUMARATE * Tab 2.5 mg	90 90 90	✓ Ipca-Bisoprolol ✓ Ipca-Bisoprolol ✓ Ipca-Bisoprolol
CARVEDILOL * Tab 6.25 mg	60 60 60	✓ Carvedilol Sandoz ✓ Carvedilol Sandoz ✓ Carvedilol Sandoz

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
ABETALOL				
* Tab 100 mg	14.50	100	✓	Trandate
* Tab 200 mg	27.00	100	✓	Trandate
* Inj 5 mg per ml, 20 ml ampoule		5		
	(88.60)			Trandate
METOPROLOL SUCCINATE				
* Tab long-acting 23.75 mg	4.20	90	1	Myloc CR
* Tab long-acting 47.5 mg		90		Myloc CR
* Tab long-acting 95 mg		90	1	Myloc CR
* Tab long-acting 190 mg		90	1	Myloc CR
METOPROLOL TARTRATE				
* Tab 50 mg	5.66	100	1	IPCA-Metoprolol
* Tab 100 mg		60		IPCA-Metoprolol
* Tab long-acting 200 mg		28		Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial		5		Metoprolol IV Mylan
,, ., ., ., ., ., ., ., ., ., .,		-		Metoprolol IV Viatris
NADOLOL				
* Tab 40 mg	19.19	100	1	Nadolol BNM
* Tab 80 mg		100	✓	Nadolol BNM
PROPRANOLOL				
* Tab 10 mg	7.04	100	/	Drofate
* Tab 40 mg		100		IPCA-Propranolol
* Cap long-acting 160 mg		100		Cardinol LA
* Oral liq 4 mg per ml - Special Authority see SA1327 be				-
Retail pharmacy		500 m	n 🗸	Roxane-
• •				Propranolol \$29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

AMI ODIDINI

*	Tab 80 mg37.50	500	✓ Mylan
	Tab 160 mg14.00		✓ Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AIV	LODIPINE			
*	Tab 2.5 mg	1.45	90	✓ Vasorex
	Tab 5 mg		90	✓ Vasorex
*	Tob 10 mg	1 21	00	/ Vacaray

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
FLODIDINE	Ψ	1 01		Warranacturer
ELODIPINE Tob long acting 2.5 mg	2.10	30	1	Plendil ER
Tab long-acting 2.5 mg Tab long-acting 5 mg		90		Felo 5 ER
Tab long-acting 3 mg		90		Felo 10 ER
	0.90	90	•	reio io En
IFEDIPINE				
← Tab long-acting 10 mg — Subsidy by endorsement	19.42	56	•	Tensipine MR10 S29
Subsidised for patients who were taking nifedipine endorsed accordingly. Pharmacists may annotate dispensing of nifedipine tab long-acting 10 mg.	0 0 01		•	
Fab long-acting 20 mg	17.72	100		Nyefax Retard
Fab long-acting 30 mg	4.78	14	✓	Mylan Italy (24 hr release) §29
	34.10	100	✓	Mylan (24 hr release) S29
★ Tab long-acting 60 mg	52.81	100	✓	Mylan (24 hr
Table long doming of mg				release) \$29
				1010000)
Other Calcium Channel Blockers				
ILTIAZEM HYDROCHLORIDE				
★ Cap long-acting 120 mg	65.35	500		Diltiazem CD Clinect
Cap long-acting 180 mg	7.00	30	1	Cardizem CD
Cap long-acting 240 mg	9.30	30	✓ :	Cardizem CD
ERHEXILINE MALEATE				
← Tab 100 mg	62.90	100	✓	Pexsig
ERAPAMIL HYDROCHLORIDE				· ·
F Tab 40 mg	7.01	100	✓	Isoptin
€ Tab 80 mg		100		Isoptin
₹ Tab long-acting 120 mg		100		Isoptin Retard S29
Tab long-acting 120 mg	00.02	100		Isoptin SR
Fab long-acting 240 mg	15 12	30		Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available		00		isopaii ori
PSO		5	1	Isoptin
1 00	25.00	J		130рин
Centrally-Acting Agents				
CLONIDINE				
Fatch 2.5 mg, 100 mcg per day — Only on a prescription		4		<u>Mylan</u>
★ Patch 5 mg, 200 mcg per day − Only on a prescription		4		Mylan
 Patch 7.5 mg, 300 mcg per day – Only on a prescription 	on17.90	4	✓	<u>Mylan</u>
CLONIDINE HYDROCHLORIDE				
← Tab 25 mcg	29.32	112	1	Clonidine Teva
← Tab 150 mcg	40.41	100	1	Catapres
foliation in the first f	14.10	5	✓	Catapres
IETHYLDOPA			•	
F Tab 250 mg	15 10	100	√	Methyldopa Viatris
r 100 200 mg	15.10	100	•	meniyidopa viadis

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

		Subsidy (Manufacturer's P	rice) Subs	sidised (Brand or Generic Manufacturer
		\$	Per	V 1	wanulacturer
ט	iuretics				
L	pop Diuretics				
	METANIDE	40.00	100	4 D	
*	Tab 1 mg		100 5	✓ Bur ✓ Bur	
FU	ROSEMIDE [FRUSEMIDE]			_	
*	Tab 40 mg — Up to 30 tab available on a PSO Tab 500 mg		1,000 50		<u>A-Frusemide</u> x Forte
*	Oral liq 10 mg per ml	11.20	30 ml OP	✓ Las	ix
	Inj 10 mg per ml, 25 ml ampoule		6 5	✓ Las	ix osemide-Baxter
		002.40	<u> </u>	- <u>- u</u>	OSCIIIUC BUXICI
P	otassium Sparing Diuretics				
AM	ILORIDE HYDROCHLORIDE	04.07	100	45	
	Tab 5 mg	81.07	100 28		agis S29 ckhardt S29
	Oral liq 1 mg per ml		25 ml OP	✓ Bio	
EP	LERENONE - Special Authority see SA1728 below - Retail p		00	. In a	
	Tab 25 mg		30 30	✓ <u>Ins</u> p	
	SA1728 Special Authority for Subsidy				
	ial application from any relevant practitioner. Approvals valid following criteria:	d without further i	renewal unless	s notified f	or applications meeting
Bot	•				
	1 Patient has heart failure with ejection fraction less than 402 Either:	%; and			
	2.1 Patient is intolerant to optimal dosing of spironolact				
	2.2 Patient has experienced a clinically significant adve	erse effect while	on optimal dos	sing of spir	onolactone.
	RONOLACTONE Tab 25 mg	3.68	100	✓ Spir	ractin
	Tab 100 mg	10.65	100	✓ Spir	ractin
	Oral liq 5 mg per ml	35.70	25 ml OP	✓ Bio	med
P	otassium Sparing Combination Diuretics				
	ILORIDE HYDROCHLORIDE WITH FUROSEMIDE	0.00			
	Tab 5 mg with furosemide 40 mgILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZI		28	✓ Fru	mii
	Tab 5 mg with hydrochlorothiazide 50 mg		50	✓ Mod	duretic

Subsidy

Fully

Brand or

				_
	Subsidy (Manufacturer's Price) \$) Subs Per	Fully idised	Brand or Generic Manufacturer
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg - Up to 150 tab available on a PSO	51.50	500	✓ <u>Arr</u>	<u>row-</u> Bendrofluazide
May be supplied on a PSO for reasons other than emerg * Tab 5 mg	•	500	✓ <u>Ari</u>	<u>row-</u> Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml	30.67 2	5 ml OP	✓ Bio	omed
CHLORTALIDONE [CHLORTHALIDONE] * Tab 25 mg	6.95	50	✓ <u>H</u> y	<u>groton</u>
INDAPAMIDE * Tab 2.5 mg METOLAZONE	16.00	90	✓ <u>Da</u>	pa-Tabs
Tab 5 mg	CBS	1 50		tolazone S29 roxolyn S29
(Metolazone S29 Tab 5 mg to be delisted 1 July 2025)				
Vasopressin receptor antagonists				
TOLVAPTAN – Special Authority see SA2166 below – Retail pha		28 OP	√ .lir	arc

TOLVAF TAIN — Special Authority see SA2 100 below — netall prial	macy		
Tab 15 mg	873.50	28 OP	Jinarc
Tab 30 mg	873.50	28 OP	Jinarc
Tab 45 mg + 15 mg	1,747.00	56 OP	Jinarc
Tab 60 mg + 30 mg	1,747.00	56 OP	Jinarc
Tab 90 mg + 30 mg		56 OP	Jinarc

⇒SA2166 Special Authority for Subsidy

Initial application — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m² at treatment initiation; and
- 3 Either:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Renewal — (autosomal dominant polycystic kidney disease) only from a renal physician or any relevant practitioner on the recommendation of a renal physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
L	ipid-Modifying Agents				
F	ibrates				
*	ZAFIBRATE Tab 200 mg Tab long-acting 400 mg		90 30		Bezalip Bezalip Retard
C	ther Lipid-Modifying Agents				
	IPIMOX Cap 250 mg	38.19	30	•	Olbetam
R	esins				
CC	DLESTYRAMINE Powder for oral suspension 4 g sachet	61.50	50		Colestyramine - Mylan \$29 Quantalan sugar free \$29
Н	MG CoA Reductase Inhibitors (Statins)				
АТ	ORVASTATIN THE 10 AMERICAN	0.04	00	,	Louis
	Tab 10 mg	0.31 5.16	30 500		<u>Lorstat</u> Lorstat
	Tab 20 mg	0.45	28		Lipitor
	Tab 40 mm	8.12	500		Lorstat
	Tab 40 mg Tab 80 mg		500 30		<u>Lorstat</u> <u>Lorstat</u>
	7ab 00 mg	25.39	500	_	Lorstat
PR	AVASTATIN				
*	Tab 20 mg	7.16	100	✓	Clinect
*	Tab 40 mg	12.25	100	/	Clinect
	SUVASTATIN - Special Authority see SA2093 below - Retain			_	
*	Tab 5 mg		30		Rosuvastatin Viatris
	Tab 10 mg Tab 20 mg		30 30		Rosuvastatin Viatris Rosuvastatin Viatris
	Tab 40 mg		30		Rosuvastatin Viatris
≫ Ini t	SA2093 Special Authority for Subsidy tial application — (cardiovascular disease risk) from any rees notified for applications meeting the following criteria: her: 1 Both:		Appr	ovals valid	without further renewal
	1.1 Patient is considered to be at risk of cardiovascula 1.2 Patient is Māori or any Pacific ethnicity; or	r disease; and			

2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and

continued...

2 Both:

Subsidy		Fully	Brand or	
(Manufacturer's P	rice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atoryastatin and/or simyastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Any of the following:

SIMVASTATIN

- 1.1 Patient has proven coronary artery disease (CAD); or
- 1.2 Patient has proven peripheral artery disease (PAD); or
- 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

* Tab 10 mg1.68	90	✓ <u>Simvastatin Mylan</u>✓ Simvastatin Viatris
* Tab 20 mg	90	✓ Simvastatin Viatris
* Tab 40 mg4.11	90	✓ Simvastatin Viatris
* Tab 80 mg8.81	90	✓ <u>Simvastatin Viatris</u>
Selective Cholesterol Absorption Inhibitors		
EZETIMIBE		
* Tab 10 mg1.76	30	✓ Ezemibe Viatris✓ Ezetimibe Sandoz
(Ezemibe Viatris Tab 10 mg to be delisted 1 July 2025)		
EZETIMIBE WITH SIMVASTATIN		
Tab 10 mg with simvastatin 10 mg5.15	30	✓ Zimybe
Tab 10 mg with simvastatin 20 mg6.15	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg8.15	30	✓ Zimybe

	Subsidy		Fully Brand or
	(Manufacturer's		idised Generic
	\$	Per	✓ Manufacturer
Nitrates			
OLVOEDVI TRINITRATE			
GLYCERYL TRINITRATE			
* Oral pump spray, 400 mcg per dose – Up to 250 dose			
available on a PSO	7.48	250 dose OP	Nitrolingual Pump
			Spray
* Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
* Patch 50 mg, 10 mg per day	18.62	30	✓ Nitroderm TTS
ISOSORBIDE MONONITRATE			
* Tab 20 mg	22.40	100	✓ Ismo 20
* Tab long-acting 40 mg		30	✓ Ismo 40 Retard
		90	_
* Tab long-acting 60 mg	13.30	90	✓ <u>Duride</u>
Sympathomimetics			
Sympathonimetics			
ADRENALINE			
Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PS	SO4.98	5	✓ Aspen Adrenaline
.,,	13.27	-	✓ DBL Adrenaline
	25.30	10	✓ Hameln S29
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a		5	✓ Hospira
ing this to,000, to the ampoule - op to 5 ing available on a	49.00	10	✓ Aspen Adrenaline
	49.00	10	Aspen Autenanne
Vasodilators			
Vasouliators			
HYDRALAZINE HYDROCHLORIDE			
	1		
* Tab 25 mg - Special Authority see SA1321 below - Retai		1	✓ Hvdralazine
		1 56	✓ Hydralazine
* Tab 25 mg - Special Authority see SA1321 below - Retai		56	✓ Onelink S29
* Tab 25 mg - Special Authority see SA1321 below - Retai		56 84	✓ Onelink \$29 ✓ AMDIPHARM \$29
* Tab 25 mg - Special Authority see SA1321 below - Retai pharmacy	CBS	56 84 100	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29
** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule	CBS	56 84	✓ Onelink \$29 ✓ AMDIPHARM \$29
* Tab 25 mg - Special Authority see SA1321 below - Retai pharmacy	CBS	56 84 100	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29
** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule ** SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals va	CBS	56 84 100 5	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline
** Tab 25 mg - Special Authority see SA1321 below - Retai pharmacy ** Inj 20 mg ampoule SA1321 Special Authority for Subsidy	CBS	56 84 100 5	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline
** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule ** SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals va	CBS	56 84 100 5	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline
Tab 25 mg - Special Authority see SA1321 below - Retain pharmacy Inj 20 mg ampoule	CBS	56 84 100 5	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline
** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule ** SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals value following criteria: Either: 1 For the treatment of refractory hypertension; or		56 84 100 5 r renewal unless	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting
** Tab 25 mg – Special Authority see SA1321 below – Retai pharmacy ** Inj 20 mg ampoule ** SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vathe following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a result of the second sec		56 84 100 5 r renewal unless	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting
* Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy		56 84 100 5 r renewal unless	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting
* Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy	25.90 alid without furthe	56 84 100 5 r renewal unless who are intolera	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting and or have not responded to AC
* Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy	25.90 alid without furthe hitrate, in patients	56 84 100 5 r renewal unless who are intolera	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29
* Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy	25.90 alid without furthe	56 84 100 5 r renewal unless who are intolera	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting and or have not responded to AC
* Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy	25.90 alid without furthe hitrate, in patients	56 84 100 5 r renewal unless who are intolera	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten
# Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy # Inj 20 mg ampoule		56 84 100 5 r renewal unless who are intolera	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health
Tab 25 mg – Special Authority see SA1321 below – Retain pharmacy		56 84 100 5 r renewal unless who are intolera 60 100	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline s notified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten
* Inj 20 mg ampoule * Inj 20 mg ampoule * Inj 20 mg ampoule * SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vathe following criteria: Either: 1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a rinhibitors and/or angiotensin receptor blockers. MINOXIDIL A Tab 10 mg		56 84 100 5 r renewal unless who are intolera 60 100	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health
* Inj 20 mg ampoule		56 84 100 5 r renewal unless who are intolera 60 100 60	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health ✓ Max Health
* Tab 25 mg - Special Authority see SA1321 below - Retain pharmacy		56 84 100 5 r renewal unless who are intolera 60 100	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health
* Inj 20 mg ampoule		56 84 100 5 r renewal unless who are intolera 60 100 60 60	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health ✓ Max Health ✓ Hospira
* Tab 25 mg - Special Authority see SA1321 below - Retain pharmacy		56 84 100 5 r renewal unless who are intolera 60 100 60	✓ Onelink \$29 ✓ AMDIPHARM \$29 ✓ Camber \$29 ✓ Apresoline snotified for applications meeting ant or have not responded to AC ✓ Minoxidil Roma \$29 ✓ Loniten ✓ Max Health ✓ Max Health

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Endothelin Receptor Antagonists

AMBRISENTAN - Special Authority see SA2253 below - Retail	pharmacy		
Tab 5 mg	200.00	30	✓ Ambrisentan Viatris
Tab 10 mg	200.00	30	✓ Ambrisentan Viatris

⇒SA2253 Special Authority for Subsidy

Initial application — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease: or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and

				_
Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Ambrisentan is to be used as PAH dual therapy; and
 - 5.2 Either:
 - 5.2.1 Patient has tried a PAH monotherapy (sildenafil or bosentan) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; or
 - 5.2.2 Patient has tried PAH dual therapy including bosentan and has experienced intolerable side effects on bosentan; and
 - 5.3 Both:
 - 5.3.1 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy; and
 - 5.3.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease).

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH 2022 (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Both:
 - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV: and
 - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where the patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN - Special Authority see SA2254 below - Retail pharmacy

Tab 62.5 mg	100.00	60	✓ Bosentan Dr
			Reddy's
Tab 125 mg	100.00	60	✓ Bosentan Dr
			Reddy's

⇒SA2254 Special Authority for Subsidy

Initial application — **(PAH monotherapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease: or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil: or
 - 5.2.2 Patient has an absolute contraindication to sildenafil: or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) †: or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Bosentan is to be used as part of PAH dual therapy; and
- 6 Either:
 - 6.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool**; or
 - 6.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely

Subsidy		Fully	Brand or	
(Manufacturer's Pric	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

benefit from initial dual therapy.

Initial application — (PAH triple therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as part of PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA2255 on the next page – Retail p	oharmacy		
Tab 25 mg	0.72	4	✓ Vedafil
Tab 50 mg		4	✓ Vedafil
Tab 100 mg1		12	✓ Vedafil

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Por 🗸	Manufacturer

⇒SA2255 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II. III or IV: and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH is confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Notes: Note: Indications marked with * are Unapproved Indications.

† The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	✓	Manufacturer	

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA2256 below - Retail pharm	macy		
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri

⇒SA2256 Special Authority for Subsidy

Initial application — **(PAH dual therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist;
 - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

Subsidy	F	ully	Brand or
Manufacturer's Price)	Subsid	sed	Generic
 \$	Per	✓	Manufacturer

continued...

- 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Epoprostenol is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST – Special Authority see SA2257 below – Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml185.03 Vebulis

⇒SA2257 Special Authority for Subsidy

Initial application — (PAH monotherapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these quidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Iloprost is to be used as PAH monotherapy; and
 - 5.2 Either:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

Initial application — (PAH dual therapy) only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † : or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**: or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 5.2 Either:
 - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
- 5.3 Either:
 - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; or
 - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initial application — **(PAH triple therapy)** only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease: or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Iloprost is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Renewal only from a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist. Approvals valid for 2 years where patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool**.

	Subsidy	F	ully	Brand or
(N	lanufacturer's Price)	Subsidi	sed	Generic
	\$	Per	1	Manufacturer

continued...

Notes: † The European Respiratory Journal Guidelines can be found here: 2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH

^{**} the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 93

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a procomption			
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA2023 below - Retai	I pharmacy		
Cap 5 mg	11.26	60	Oratane
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

TRETINOIN

Crm 0.5 mg per g − Maximum of 50 g per prescription16.82 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 93

HYDROGEN PEROXIDE

HYDROGEN PEROXIDE			
* Crm 1%	8.56	10 g OP	Crystaderm
MUPIROCIN			
Oint 2%	6.60	15 g OP	
	(13.00)	•	Bactroban

- a) Only on a prescription
- b) Not in combination

		L	JEKIVI	ATOLOGICALS
	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.69	5 g OP	✓ <u>F</u>	oban
b) Only on a prescriptionc) Not in combination	4.00	5 ~ OD		ah an
Oint 2%	1.69	5 g OP	V <u>F</u>	<u>oban</u>
SULFADIAZINE SILVER	40.00	50 OD		
Crm 1%	10.80	50 g OP	-	lamazine scend 829
a) Up to 250 g available on a PSOb) Not in combination	13.44		• ^	Scena •
(Ascend S29 Crm 1% to be delisted 1 July 2025)				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals,	page 100			
AMOROLFINE				
a) Only on a prescription b) Not in combination				
Nail soln 5%	21.87	5 ml OP	✓ M	lycoNail
CLOTRIMAZOLE				
* Crm 1%	1.10	20 g OP	✓ C	lomazol
a) Only on a prescription b) Not in combination				
* Soln 1%	4.36 (7.55)	20 ml OP	C	anesten
a) Only on a prescriptionb) Not in combination	(1.00)			u
ECONAZOLE NITRATE				
Crm 1%	8.04	20 g OP	✓ P	evaryl
a) Only on a prescription b) Not in combination				

3

Pevaryl

(18.64)

c) Pevaryl to be Principal Supply on 1 June 2025

a) Only on a prescriptionb) Not in combination

Foaming soln 1%, 10 ml sachets.....9.89

	Subsidy	Ol-	Fully	Brand or
	(Manufacturer's F \$	Per Subs	sidised •	Generic Manufacturer
ICONAZOLE NITRATE				
F Crm 2%	0.90	15 g OP	✓ M	lultichem
a) Only on a prescription				
b) Not in combination				
£ Lotn 2%		30 ml OP		
	(10.03)		D	aktarin
a) Only on a prescription				
b) Not in combination Tinct 2%	4.26	30 ml OP		
1 IIICt 2 /6	(12.10)	30 IIII OF	D	aktarin
a) Only on a prescription	(12.10)		D	anam
b) Not in combination				
,				
Antipruritic Preparations				
ALAMINE				
a) Only on a prescription				
b) Not in combination				
Crm, aqueous, BP	3.45	100 g	✓ h	ealthE Calamine
has the Colombia Assessments had be been also construction				Aqueous
healthE Calamine Aqueous to be Principal Supply of	on 1 April 2025			
ROTAMITON				
a) Only on a prescription b) Not in combination				
D) NOUR COMBINATION				
	3 /10	20 a OP	J It	ch-Soothe
Crm 10%	3.49	20 g OP	✓ <u>lt</u>	ch-Soothe
Crm 10% ENTHOL – Only in combination		J	_	ch-Soothe
Cmr 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or		J	_	<u>ch-Soothe</u>
Crm 10% ENTHOL – Only in combination		J	_	ch-Soothe
Cmr 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or	proprietary Topical C	J	Plain	ch-Soothe lidWest
Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals.	proprietary Topical C	Corticosteriod –	Plain	
Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. Crystals	proprietary Topical C	Corticosteriod – 25 g	Plain	lidWest
Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. Crystals Corticosteroids Topical	proprietary Topical C	Corticosteriod – 25 g 100 g	Plain	lidWest
Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. Crystals	proprietary Topical C	Corticosteriod – 25 g 100 g	Plain	lidWest
Crm 10% ENTHOL – Only in combination 1) Only in combination with a dermatological base or 2) With or without other dermatological galenicals. Crystals Corticosteroids Topical	proprietary Topical C	Corticosteriod – 25 g 100 g	Plain	lidWest
Crm 10%	proprietary Topical C	Corticosteriod – 25 g 100 g	Plain	lidWest
Crm 10%	proprietary Topical C6.92 29.60 AND RELATED AGE	Corticosteriod – 25 g 100 g	Plain M M	lidWest
Crm 10%	proprietary Topical C	Corticosteriod – 25 g 100 g NTS, page 83	Plain M M	lidWest lidWest
Crm 10%	proprietary Topical C	Corticosteriod – 25 g 100 g NTS, page 83 15 g OP 50 g OP 15 g OP	Plain M M D D D D D	ijprosone iprosone iprosone
Crm 10%	proprietary Topical C	Corticosteriod – 25 g 100 g NTS, page 83 15 g OP 50 g OP 15 g OP 50 g OP	Plain M M D D C D D	iprosone iprosone iprosone iprosone iprosone
Crm 10%	proprietary Topical C	Corticosteriod – 25 g 100 g NTS, page 83 15 g OP 50 g OP 15 g OP	Plain M M D D C D D	ijprosone iprosone iprosone
Crm 10%	2.96 36.00 2.96 36.00 4.33	25 g 100 g NTS, page 83 15 g OP 50 g OP 15 g OP 50 g OP 30 g OP	Plain M M D D D D D D D	iprosone iprosone iprosone iprosone iprosone iprosone iprosone
Crm 10%	2.96 36.00 2.96 36.00 4.33	25 g 100 g NTS, page 83 15 g OP 50 g OP 15 g OP 50 g OP 30 g OP	Plain M M D D D D D D D D D D D	iprosone iprosone iprosone iprosone iprosone iprosone iprosone
Crm 10%	2.96 36.00 2.96 36.00 4.33	25 g 100 g NTS, page 83 15 g OP 50 g OP 15 g OP 50 g OP 30 g OP 50 g OP 50 g OP	Plain M M D D D D D D D D D D D	iprosone iprosone iprosone iprosone iprosone iprosone iprosone outlier iprosone iprosone
Crm 10%	2.96 36.00 2.96 36.00 4.33	25 g 100 g NTS, page 83 15 g OP 50 g OP 15 g OP 50 g OP 30 g OP	Plain M M D D D D D D D D D D D	iprosone iprosone iprosone iprosone iprosone iprosone iprosone
Crm 10%	2.96 36.00 2.96 36.00 4.33	25 g 100 g NTS, page 83 15 g OP 50 g OP 15 g OP 50 g OP 30 g OP 50 g OP 50 g OP	Plain M M D D D D D D D D D D D	iprosone iprosone iprosone iprosone iprosone iprosone iprosone outlier iprosone
Crm 10%	2.96 AND RELATED AGE 2.96 36.00 2.96 36.00 4.33 5.85 7.90 30.00	25 g 100 g NTS, page 83 15 g OP 50 g OP 15 g OP 50 g OP 30 g OP 50 g OP 50 g OP	Plain M D D D D D D D D D D D D	iprosone iprosone iprosone iprosone iprosone iprosone iprosone outlier iprosone

	Subsidy		. ,	Brand or
	(Manufacturer's Pi	rice) Sub: Per		Generic Manufacturer
OLODETA GOLIE DUTI (DATE	Ψ	1 61		IVIATIUIACIUTEI
CLOBETASONE BUTYRATE	T 00	00 = OD		
Crm 0.05%		30 g OP	г	mayata
	(10.00)		Eui	movate
HYDROCORTISONE				
* Crm 1% – Only on a prescription		30 g OP	✓ Eth	
W Daviday Only in combination	20.40	500 g	✓ <u>No</u>	
* Powder – Only in combination	49.95	25 g	✓ AB	
galenicals	ai Corticosteriou	- Flaili) Willi	Ji Williout	otilei deiliatologicai
•				
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only o		0E0 ml	./ DD	Loto UC
a prescription	12.03	250 ml	♥ <u>DP</u>	Lotn HC
HYDROCORTISONE BUTYRATE		05	٠.	
Lipocream 0.1%		100 g OP		coid Lipocream
Oint 0.1%		100 g OP 100 ml OP	✓ Lo	cola coid Crelo
Milky emul 0.1%	12.33	100 mi OP	♥ LO	cold Creio
METHYLPREDNISOLONE ACEPONATE	4.05	45 × OD		
Crm 0.1%		15 g OP		vantan
Oint 0.1%	4.95	15 g OP	✓ Ad	<u>vantan</u>
MOMETASONE FUROATE			<i>a</i>	
Crm 0.1%		15 g OP		con Alcohol Free
Oint 0.1%	3.50	50 g OP	✓ Eld	con Alcohol Free
OIIII 0.1%	3.50	15 g OP 50 g OP	✓ Eld	
Lotn 0.1%		30 ml OP	✓ Eld	
		00 1111 01	- 10	<u></u>
TRIAMCINOLONE ACETONIDE Crm 0.02%	6.40	100 g OP	√ Ari	stocort
Oint 0.02%		100 g OP		stocort
OIII 0.02/0		100 g O1	* <u>All</u>	3100011
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU				
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP	F	.:
A Mariana of 45 and a second of the	(10.45)		Fuc	cicort
a) Maximum of 15 g per prescription				
b) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE – Only on a prescrip		15 = OD	./ M:	
* Crm 1% with miconazole nitrate 2%		15 g OP	V IVIIC	creme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - O	, , ,		<i>4</i> =.	
Oint 1% with natamycin 1% and neomycin sulphate 0.5%	3.35	15 g OP	✓ Pin	nafucort
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI	N AND NYSTAT	ΊN		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg				
and gramicidin 250 mcg per g – Only on a prescription .		15 g OP	,	
	(9.28)		Via	derm KC

Subsidy (Manufacturer's Price) Per

Fully Subsidised

500 g

✓ Evara

Paraffin AFT

Brand or Generic Manufacturer

D		^-		
Barri	ıer	Cr	eai	ms

DIN	METHICONE		
*	Crm 5% pump bottle4.30	500 ml OP	✓ healthE
			Dimethicone 5%
*	Crm 10% pump bottle4.52	500 ml OP	✓ healthE
			Dimethicone 10%

ZINC AND CASTOR OIL

	our g	
Emollients		
AQUEOUS CREAM		
* Crm	500 g	✓ Evara
CETOMACROGOL		
* Crm BP2.29	500 g	✓ Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL		
Crm 90% with glycerol 10%2.13	500 ml OP	✓ Evara
3.50	1,000 ml OP	✓ Evara
EMULSIFYING OINTMENT		
* Oint BP	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION		
* Crm	500 g	✓ Fatty Cream AFT
2.10		✓ Fatty Emulsion Cream (Evara)
Fatty Emulsion Cream (Evara) to be Principal Supply on 1 April 2025 (Fatty Cream AFT Crm to be delisted 1 April 2025)		
PARAFFIN		
Oint liquid paraffin 50% with white soft paraffin 50%4.94	500 g OP	✓ White Soft Liquid

UREA	

*	Crm 10%	1.37	100 g OP	✓ healthE Urea Cream
WO	OL FAT WITH MINERAL OIL - Only on a prescription			
	Lotn hydrous 3% with mineral oil	5.60	1,000 ml	
		(14.96)		DP Lotion
		(20.53)		Alpha-Keri Lotion
		1.40	250 ml OP	
		(5.87)		DP Lotion
		5.60	1,000 ml	
		(23.91)		BK Lotion
		1.40	250 ml OP	
		(7.73)		BK Lotion

Sub (Manufactu		Fully	Brand or Generic	
`	\$ Per	✓	Manufacturer	

Other Dermatological Bases

P	Δ	R	Δ	F	F	INI
Γ	н	п	н	Г	П	HΝ

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid - Plain.

Minor Skin Infections

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	4.99	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
	6.99	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep

Parasiticidal Preparations

DIMETHICONE

IVERMECTIN - Special Authority see SA2294 below - Retail pharmacy

Tab 3 mg − Up to 100 tab available on a PSO......17.20 4 Stromectol

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA2294 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
 - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
 - 2.2 Either:
 - 2.2.1 The person is unable to complete topical therapy: or
 - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Initial application — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

DERMATOLOGICALS

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 filariasis: or
 - 2 cutaneous larva migrans (creeping eruption); or
 - 3 strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Fither:

- 1 The person has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2 Both:
 - 2.1 The person has a confirmed diagnosis of scabies or is a close contact of a scabies case; and
 - 2.2 Fither:
 - 2.2.1 The person is unable to complete topical therapy: or
 - 2.2.2 Previous treatment with topical therapy has been tried and not cleared the infestation.

Renewal — (Other parasitic infections) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 filariasis: or
- 2 cutaneous larva migrans (creeping eruption); or
- 3 strongyloidiasis.

PERMETHRIN

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA2024 below - Retail	pharmacy		
Cap 10 mg	26.20	60	✓ Novatretin
Cap 25 mg	57.37	60	✓ Novatretin

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per g59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g40.92	60 g OP	✓ Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g14.31	30 g OP	Daivobet

	Subsidy (Manufacturer's Pr	rico) Subc	Fully Brand or sidised Generic
	(Manulacturers Fi	Per	✓ Manufacturer
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	✓ Daivonex
COAL TAR		•	
Soln BP - Only in combination	36.25	200 ml	✓ Midwest
 Up to 10% only in combination with a dermatologic With or without other dermatological galenicals. 	cal base or proprie	etary Topical C	Corticosteriod – Plain
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULI	PHUR		
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% an	d		
allantoin crm 2.5%		75 g OP	
	(8.00) 3.43	20 ~ OD	Egopsoryl TA
	(4.35)	30 g OP	Egopsoryl TA
COAL TAR WITH CALLOVI IO AGIR AND GUI BUILD	(4.33)		Lgopsoryi TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint	4.07	25 a OB	✓ Coco-Scalp
3011 12% with saircylic actu 2% and sulphur 4% offit	7.95	25 g OP 40 g OP	✓ Coco-Scalp
PIMECROLIMUS - Special Authority see SA1970 below - Retai		10 9 01	- Coo Coulp
a) Maximum of 15 g per prescription	ii priarriacy		
b) Note: a maximum of 15 g per prescription and no more t	than one prescript	ion per 12 wee	eks.
Cream 1%		15 g OP	✓ Elidel
Initial application only from a dermatologist, paediatrician, ophtl of a dermatologist, paediatrician or ophthalmologist. Approvals weeting the following criteria: Both:			
 Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. 			
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE	SCEIN - Only or	a prescription	1
* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium	n5.41	500 ml	✓ Pinetarsol
SALICYLIC ACID			
Powder - Only in combination	18.88	250 g	✓ Midwest
 Only in combination with a dermatological base or With or without other dermatological galenicals. 	proprietary Topic	al Corticostero	id – Plain or collodion flexible
SULPHUR			
Precipitated - Only in combination	6.35	100 g	✓ Midwest
 Only in combination with a dermatological base or With or without other dermatological galenicals. 	proprietary Topic	al Corticostero	oid – Plain
TACROLIMUS			
Oint 0.1% – Special Authority see SA2074 on the next page) –		
Retail pharmacy		30 g OP	✓ Zematop
a) Maximum of 30 g per prescription		-	-
b) Note: a maximum of 30 g per prescription and no m	ore than one pres	cription per 12	2 weeks.

DERMATOLOGICALS

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

BETAMETHASONE VALERATE * Scalp app 0.1%	12.95	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%		30 ml OP	✓ Dermol
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%		100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%	3.23	100 ml OP	✓ Sebizole
a) Maximum of 100 ml per prescription	4.09		✓ Sebizole

a) Maximum of 100 ml per prescription

b) Only on a prescription

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

SPF 50+

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 72

PODOPHYLLOTOXIN

- a) Maximum of 3.5 ml per prescription
- b) Only on a prescription

Other Skin Preparations

Antineoplastics

Crm 5%	5.56	20 g OP	✓ <u>Efudix</u>
IMIQUIMOD			
Crm 5% 250 mg sachet	21 72	24	✓ Parrigo

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or	_
(Manufacturer's Price) \$	Per	Subsidised ✓	Generic Manufacturer	
· · · · · · · · · · · · · · · · · · ·				_

b) Prin Substance lable on a PSO

76 60 mm Hallingar Supply....

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Per	Subsidised Generic Manufacturer
Contraceptives - Non-hormonal			
Condoms			
ONDOMS			
∮ 49 mm – Up to 144 dev available on a PSO	14.25	144	✓ Moments
€ 53 mm	1.15	10	✓ Moments
	14.25	144	✓ Moments
 a) Maximum of 60 dev per prescription 			
b) Up to 60 dev available on a PSO			
€ 53 mm, 0.05 mm thickness	1.15	10	✓ Moments
	14.25	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
€ 53 mm, chocolate, brown	1.15	10	✓ Moments
	14.25	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
← 53 mm, strawberry, red	1.15	10	✓ Moments
	14.25	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
₹ 56 mm	1.15	10	✓ Moments
	14.50	144	✓ Moments
 a) Maximum of 60 dev per prescription 			
b) Up to 60 dev available on a PSO			
← 56 mm, 0.05 mm thickness	2.00	12	Gold Knight
	24.10	144	Gold Knight
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
€ 56 mm, 0.05mm thickness (bulk pack)	20.17	144	Gold Knight
a) Maximum of 60 dev per prescription			
b) Up to 60 dev available on a PSO			
€ 56 mm, 0.08 mm thickness	1.15	10	✓ Moments
	14.25	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
56 mm, 0.08 mm thickness, red	1.15	10	✓ Moments
	14.25	144	✓ Moments
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
	1.79	12	Gold Knight
	21.45	144	✓ Gold Knight
a) Up to 60 dev available on a PSO			
b) Maximum of 60 dev per prescription			
56 mm, strawberry	1.79	12	Gold Knight
-	21.45	144	✓ Gold Knight
a) Up to 60 dev available on a PSO			-
b) Maximum of 60 dev per prescription			
€ 60 mm	1.82	12	Gold Knight XL
	21.89	144	✓ Gold Knight XL
a) Maximum of 60 dev per prescription			
h) Idrate 60 dov.evailable on a DSO			

S29 Unapproved medicine supplied under Section 29
Sole ₹8bsidised Striply ✓ Gold Knight XL

GENITO-URINARY SYSTEM

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

- a) Maximum of 60 dev per prescription
- b) Up to 60 dev available on a PSO

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IUD 29.1 mm length x 23.2 mm width	29.80	1	✓ Choice 380 7med Nsha Silver/ copper Short
*	IUD 33.6 mm length × 29.9 mm width	26.80	1	✓ TCu 380 Plus
*	ILID 35.5 mm length × 19.6 mm width	33.00	1	Normal ✓ Cu 375 Standard

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

★ Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO.......10.00 84 Mercilon 28

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 84 tab available on a PSO	1.50	84	√ <u>L</u>	o-Oralcon 20 ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
	(16.50)		N	/licrogynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Autb) Up to 63 tab available on a PSO	hority see SA0500 on	the	previous pag	ge
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets Up to 84 tab available on a PSO		84	√ <u>c</u>	Oralcon 30 ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to 84 tab available on a PSO		84		Alyacen Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - L	Jp			
to 84 tab available on a PSO	21.99	84	√ N	lorimin

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONONGESTREL			
* Tab 30 mcg - Up to 112 tab available on a PSO	22.00	112	✓ Microlut
* Subdermal implant (2 x 75 mg rods) - Up to 3 pack ava	ilable		
on a PSO	106.92	1	✓ <u>Jadelle</u>
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available or	n a PSO 10.56	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price)	S Per	Fully ubsidised	Brand or Generic Manufacturer
NORETHISTERONE Tab 350 mcg – Up to 84 tab available on a PSO	12.25	84	✓ N	Vorethinderone - CDC Voriday Voriday 28
Emergency Contraceptives				

LEVONORGESTREL

- ✓ Levonorgestrel **BNM**
 - a) Maximum of 2 tab per prescription
 - b) Up to 5 tab available on a PSO
 - c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- A maximum \$5.00 prescription charge (patient co-payment) may apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to any non contraceptive prescription charges that apply, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

* Tab 2 mg with ethinvloestradiol 35 mcg and 7 inert tabs - Up

to 168 tab available on a PSO.......5.08

168

100 g OP

Ginet

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID	
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate	
0.025% alvegral 5% and riginalais asid 0.75% with applicator	

0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator.... 8.43 CLOTRIMAZOLE

35 q OP

20 a OP

MICONAZOLE NITRATE

40 g OP

Clomazol Micreme

Aci-Jel

Clomazol

NYSTATIN

OESTRIOL

Vaginal crm 100,000 u per 5 g with applicator(s)5.70 75 g OP Nilstat

Myometrial and Vaginal Hormone Preparations

FRGOMETRINE MAI FATE

Inj 500 mcg per ml	1 ml ampoule – Up to 5 inj available on a
PSO	160.00

5 15 g OP 15

Ovestin

✓ Ovestin

✓ DBL Ergometrine

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price)	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
OXYTOCIN – Up to 5 inj available on a PSO				
Inj 5 iu per ml, 1 ml ampoule	4.98	5	✓ (Dxytocin BNM
Inj 10 iu per ml, 1 ml ampoule		5	1	Dxytocin BNM
	11.96	10	1	Dxytocin
				Panpharma
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj avai	lable on a PSO			
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampo		5	√ <u>9</u>	Syntometrine .

Pregnancy Tests - hCG Urine

ALGINANCI ILSIS-IICG UNINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 112

5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy

★ Tab 5 mg4.79 100 ✓ <u>Ricit</u>

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoreceptor Blockers

SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Pregnancy Test

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's F	Price) Subs	Fully Brand or sidised Generic
	\$	Per	✓ Manufacturer
Other Urinary Agents			
OXYBUTYNIN			
* Tab 5 mg	5.42	100	Alchemy Oxybutynin
POTASSIUM CITRATE			ON July IIII
Oral liq 3 mmol per ml – Special Authority see SA1083 belov Retail pharmacy		200 ml OP	✓ Biomed
⇒SA1083 Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals valid	I for 12 months t	for applications	meeting the following criteria:
Both:			
 The patient has recurrent calcium oxalate urolithiasis; and The patient has had more than two renal calculi in the two 	veare prior to th	a annlication	
Renewal from any relevant practitioner. Approvals valid for 2 year			s appropriate and the patient is
benefitting from the treatment.			o appropriate and the patient is
SODIUM CITRO-TARTRATE			
* Grans eff 4 g sachets	3.50	28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg	1.95	30	✓ Solifenacin succinate Max Health
	2.05		✓ Solifenacin Viatris
Solifenacin succinate Max Health to be Principal Supply			
Tab 10 mg	3.53	30	✓ Solifenacin succinate Max Health
	3.72		✓ Solifenacin Viatris
Solifenacin succinate Max Health to be Principal Supply	on 1 June 2025		
(Solifenacin Viatris Tab 5 mg to be delisted 1 June 2025)			
(Solifenacin Viatris Tab 10 mg to be delisted 1 June 2025)			
Detection of Substances in Urine			
ORTHO-TOLIDINE			
* Compound diagnostic sticks	7.50	50 test OP	
	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	13.92	100 test OP	✓ Albustix
Obstetric Preparations			
Antiprogesterones			
. •			

Tab 200 mg - Up to 15 tab available on a PSO......83.90

180.00

✓ Mifegyne✓ Mifegyne

MIFEPRISTONE

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Calcium Homeostasis

CALCITONIN		_	
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	Miacalcic
CINACALCET - Special Authority see SA2170 below - Reta	il pharmacy		
Tab 30 mg - Wastage claimable	25.24	28	 Cinacalet Devatis
Tab 60 mg - Wastage claimable	50.47	28	✓ Cinacalet Devatis

⇒SA2170 Special Authority for Subsidy

Initial application — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

- Either:
 - 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
 - 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate

Renewal — (parathyroid carcinoma or calciphylaxis) only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
 - 1 The patient's serum calcium level has fallen to < 3mmol/L; and
 - 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initial application — (**primary hyperparathyroidism**) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initial application — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or

Subsidy (Manufacturer's Price)	F Subsid	ully	Brand or Generic
 \$	Per	√	Manufacturer

continued...

- 3.2 Parathyroid tissue is surgically inaccessible; or
- 3.3 Parathyroid surgery is not feasible.

Renewal — (secondary or tertiary hyperparathyroidism) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial	15.65	1	✓ Zoledronic acid
			Viatris

Corticosteroids and Related Agents for Systemic Use

DETAMETHAÇONE CODILIM DHOCDHATE WITH DETAMETHAÇONE ACETATE

BETAMETHASON	E SODIUM PHOSPHATE WITH BETAMETH	ASONE ACETAT	E		
* Inj 3.9 mg with	betamethasone acetate 3 mg per ml, 1 ml	19.20	5		
, 0	31 ,	(36.96)			Celestone
		(55.55)			Chronodose
	_				Officioaccc
DEXAMETHASON	—			_	
* Tab 0.5 mg -	Up to 60 tab available on a PSO	1.80	30		Dexmethsone
* Tab 4 mg - U	p to 30 tab available on a PSO	3.18	30	✓	<u>Dexmethsone</u>
Oral lig 1 mg p	er ml	53.86	25 ml OP	1	Biomed
DEXAMETHASON					
		ral usa			
	ne phosphate injection will not be funded for o		40	,	Hamada
	I, 1 ml ampoule - Up to 5 inj available on a P		10		Hameln
* Inj 4 mg per m	I, 2 ml ampoule $-$ Up to 5 inj available on a P	SO 13.10	10	•	<u>Hameln</u>
FLUDROCORTISC	ONE ACETATE				
* Tab 100 mcg		11.46	100	1	Florinef
=			.00		<u>- 10111101</u>
HYDROCORTISOI	· · -				
			100		Douglas
* Tab 20 mg		20.32	100		Douglas
* Inj 100 mg vial		3.96	1	•	Solu-Cortef
a) Not on	a BSO				
b) Up to 5	5 inj available on a PSO				
, .	•				
METHYLPREDNIS		440.00	400	,	
•			100		Medrol
* Tab 100 mg		223.10	20	/	Medrol
METHYLPREDNIS	SOLONE (AS SODIUM SUCCINATE)				
	,	22.30	1	1	Solu-Medrol-Act-
,			•		O-Vial
					O viui
lni 125 mg vial		34.10	1	1	Solu-Medrol-Act-
ing 125 mg viai			'	•	O-Vial
					O-viai
lni 500 ma vial		43.01	1	1	Solu-Medrol-Act-
ing 500 mg viai		45.01	,	•	O-Vial
					O-VIAI
lni 1 a vial		52.54	1	1	Solu-Medrol
iiij i y vial		32.34	ı	•	Joiu-Meuroi

	Subsidy	, ,	Fully	Brand or
	(Manufacturer's Price \$	e) Sub: Per	sidised •	Generic Manufacturer
METHYLPREDNISOLONE ACETATE				
Inj 40 mg per ml, 1 ml vial	47.06	5	✓ I	Depo-Medrol
PREDNISOLONE				
Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	√ <u>j</u>	Redipred
REDNISONE				
₹ Tab 1 mg	18.58	500	✓	Prednisone Clinect
★ Tab 2.5 mg	21.04	500	✓	Prednisone Clinect
Tab 5 mg - Up to 30 tab available on a PSO	19.30	500	✓	Prednisone Clinect
Tab 20 mg - Up to 30 tab available on a PSO	50.51	500	✓	Prednisone Clinect
ETRACOSACTRIN				
Inj 250 mcg per ml, 1 ml ampoule	86.25	1	1	Synacthen
, ,				JK Synacthen
Finj 1 mg per ml, 1 ml ampoule	690.00	1		Synacthen Depot
, , , , ,				Synacthene '
				Retard S29
RIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	21.42	5	√ 1	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5		Kenacort-A 40
, , ,			-	
Sex Hormones Non Contraceptive				
Androgen Agonists and Antagonists				
YPROTERONE ACETATE				
Tab 50 mg		50		Siterone
Tab 100 mg	31.00	50	√ 9	Siterone
ESTOSTERONE				
Gel (transdermal) 16.2 mg per g	52.00	88 g OP	✓.	Testogel .
ESTOSTERONE CIPIONATE				
Inj 100 mg per ml, 10 ml vial	85.00	1	✓ I	Depo-Testosterone
ESTOSTERONE ESTERS		•		- оро постолого
	10.00	4	./ (Sustanan Amnaulas
Inj 250 mg per ml, 1 ml	1∠.∀ō	1	• ;	Sustanon Ampoules
ESTOSTERONE UNDECANOATE				
Cap 40 mg - Subsidy by endorsement		100		Steril-Gene S29
Subsidy by endorsement – subsidised for patients who				
1 November 2021 and the prescription is endorsed acc				
where there exists a record of prior dispensing of testo	sterone undecanoate	cap 40 mg	in the	preceding 12 months.

Inj 250 mg per ml, 4 ml vial.......86.00

✓ Reandron 1000

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Hormone Replacement Therapy - Systemic

·	Collogono			
OE	STRADIOL			
-	Tab 1 mg	4.12	28 OP	
		(11.10)		Estrofem
*	Tab 2 mg	\ -/	28 OP	
		(11.10)		Estrofem
*	Gel (transdermal) 0.06% (750 mcg/actuation)		80 g OP	✓ Estrogel
	Patch 25 mcg per day		8	✓ Estradiol TDP Mylan
		13.50		✓ Estraderm MX S29
		14.50		✓ Estradot
		21.35		✓ Lyllana
	a) No more than 2 patch per week	21.00		- Lynana
	b) Only on a prescription			
	Patch 50 mcg per day	10.75	8	✓ Estradiol TDP Mylan
	r atch 50 mag per day	10.75	Ü	✓ Estradiol Viatris
		14.50		
		14.50		✓ Estraderm MX S29
				✓ Estradiol Sandoz✓ Estradot
		21.55		
	\	21.33		✓ Lyllana
	a) No more than 2 patch per week			
	b) Only on a prescription	44.00	•	4 T
	Patch 75 mcg per day	11.88	8	✓ Estradiol TDP Mylan
		44.50		✓ Estradiol Viatris
		14.50		✓ Estradiol Sandoz
		00.07		✓ Estradot
		22.37		✓ Lyllana
	a) No more than 2 patch per week			
	b) Only on a prescription			
	Patch 100 mcg per day	12.95	8	✓ Estradiol TDP Mylan
				✓ Estradiol Viatris
		14.50		✓ Estradiol Sandoz
				✓ Estradot
		15.50		✓ Estraderm MX S29
		22.77		✓ Lyllana
	a) No more than 2 patch per week			
	b) Only on a prescription			
OE	STRADIOL VALERATE			
*	Tab 1 mg	12.36	84	✓ Progynova
*	Tab 2 mg		84	✓ Progynova
	STROGENS			- 37
	Conjugated, equine tab 300 mcg	2.01	28	
不	Conjugated, equille lab 300 micg		20	Premarin
*	Conjugated, equine tab 625 mcg	(19.25)	28	FIEIIIaIIII
不	Ourijugateu, equille tab 625 mcg		20	Premarin
		(19.25)		FIEIIIaiiii

	Subsidy (Manufacturer's Price) \$	Per	Subsidised G	rand or eneric anufacturer
Progestogens				
EDROXYPROGESTERONE ACETATE				
Tab 2.5 mg	6.56	30	✓ Prov	era
	8.75	56	✓ Prov	
Tab 5 mg		56	✓ Prov	
	20.13	100	✓ Prov	
Tab 10 mg		30	✓ Prov	
Progestogen and Oestrogen Combined Prepara	itions			
ESTRADIOL WITH NORETHISTERONE				
Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OF	1	
3 0 111 1 111111111	(18.10)		Kliov	ance
Tab 2 mg with 1 mg norethisterone acetate	` '	28 OF		-
J	(18.10)		Kliog	est
Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	(/		09	
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OF	1	
ossitudioi tab (12) and 1 mg ossitudioi tab (0)	(18.10)	_0 01		quens
	(10.10)		11130	quens
Other Oestrogen Preparations				
ESTRIOL				
Tab 2 mg	7.70	30	✓ Oves	tin
•	-			
Other Progestogen Preparations				
EVONORGESTREL				
Intra-uterine device 52 mg	269.50	1	✓ Mire	na
Intra-uterine device 13.5 mg	215.60	1	✓ Jayd	ess
EDROXYPROGESTERONE ACETATE			-	
Tab 100 mg	133 57	100	✓ Prov	era HD
	100.07	100	5 110V	oiu IID
ORETHISTERONE	F 15			
Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Prim	olut N
ROGESTERONE	44.05	00	4	
Cap 100 mg	14.85	30	✓ <u>Utro</u>	gestan
Thyroid and Antithyroid Agents				
ARBIMAZOLE				
Tab 5 mg	7.56	100	✓ Neo-	Mercazole
· ·				
EVOTHYROXINE Tab 05 mag	E	00	./ 0	امامسا
Tab 25 mcg		90	✓ Synt	
Tab 50 mcg		28		ury Pharma
	5.79	90	✓ Synt	
T 11 150	64.28	1,000	✓ Eltro	
Tablet 50 mcg		200	✓ Eltro	
Tab 100 mcg		28		ury Pharma
	6.01	90	✓ Synt	
	66.78	1,000		
Tablet 100 mcg	13.36	200	✓ Eltro	vin

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PROPYLTHIOURACIL - Special Authority see SA1199 below -	Retail pharmacy			
Tab 50 mg	35.00	100	✓ P	TU \$29
⇒SA1199 Special Authority for Subsidy				

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA2032 b	elow - Retail pha	ırmacy	
*	Inj 5 mg cartridge	80.21	i	✓ Omnitrope
*	Inj 10 mg cartridge	80.21	1	✓ Omnitrope
	Inj 15 mg cartridge		1	✓ Omnitrope

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (**Turner syndrome**) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

continued...

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m²) in a child who may or may not be receiving dialysis; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	

continued...

6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred;
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria:
 - 2 The patient is aged six months or older; and
 - 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
 - 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9

Subsidy	Subsidy Fully		Brand or
(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
Ψ	1 01		Wandacturer

continued...

months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GOSERELIN			
Implant 3.6 mg, syringe	66.48	1	✓ Zoladex
Implant 10.8 mg, syringe	138.23	1	✓ Zoladex

LEUPRORELIN

OOOEDELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

good of the proscription is characted accordingly.			
Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of			
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsidy			
of \$591.68 per 1 inj with Endorsement	177.50	1	
•	(591.68)		Lucrin Depot 3-month

Vasopressin Agonists

DESMOPRESSIN Wafer 120 mcg	47.00	30	✓ Minirin Melt
DESMOPRESSIN ACETATE			
Tab 100 mcg	25.00	30	✓ Minirin
Tab 200 mcg	54.45	30	✓ Minirin
▲ Nasal spray 10 mcg per dose	34.95	6 ml OP	✓ <u>Desmopressin-</u> <u>PH&T</u>
Inj 4 mcg per ml, 1 ml	67.18	10	✓ Minirin

Other Endocrine Agents

CABERGOLINE

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
Dostinex	2	waived by Special Authority see SA2070 below4.43
Dostinex	8	17.94

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 Hyperprolactinemia; or

Subsidy	Subsidy Fully		Brand or
(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
Ψ	rei		Manuacturer

continued...

- 2 Acromegaly*; or
 - 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLOM		11 □

Tab 50 mg	29.84	10	✓ Mylan Clomiphen \$29
METYRAPONE Cap 250 mg	558.00	50	✓ Metopirone

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Anthelmintics** ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy 60 ✓ Eskazole S29 **⇒SA1318** Special Authority for Subsidy Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids. Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment. MEBENDAZOLE - Only on a prescription 6 Vermox 15 ml (7.83)Vermox **PRAZIQUANTEL** Biltricide

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 66
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 266

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE			_
Cap 250 mg		100	Ranbaxy-Cefactor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.75	100 ml	 Ranbaxy-Cefactor
CEFALEXIN			
Cap 250 mg	3.85	20	✓ Cephalexin ABM
Cap 500 mg	5.85	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable		100 ml	✓ Flynn
Grans for oral liq 50 mg per ml - Wastage claimable		100 ml	✓ Flynn
1 01	11.75		✓ Cefalexin Sandoz
CEFAZOLIN - Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance with endorsed accordingly.	a Health NZ Hos	pital approved	I protocol and the prescription is
Inj 500 mg vial	3.39	5	✓ Cefazolin-AFT
Inj 1 g vial	3.59	5	✓ Cefazolin-AFT
Inj 2 g vial	7.09	5	✓ Cefazolin-AFT
CEFTRIAXONE - Subsidy by endorsement			
a) Up to 10 inj available on a PSO			
 Subsidised only if prescribed for a dialysis or cystic fibros pelvic inflammatory disease, or the treatment of suspecte endorsed accordingly. 	ed meningococca		the prescription or PSO is
Inj 500 mg vial	0.79	1	✓ Ceftriaxone-AFT
lnj 1 g vial	3.59	5	✓ Ceftriaxone-AFT
CEFUROXIME AXETIL - Subsidy by endorsement			
Only if prescribed for prophylaxis of endocarditis and the pre	scription is endo	rsed according	gly.
Tab 250 mg	CBS	20	✓ Ascend-

Cefuroxime \$29

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	16.97	15 ml	Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

	0 1			,	, ,		,
Tab 250 mg					7.31	12	✓ Klaricid S29
-					8.53	14	✓ Klacid
Grans for oral liq 250 mg per 5 ml -	Wasta	age cla	aimal	ole	192.00	50 ml	✓ Klacid

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Fither:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated.

Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician.

Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	35.82	100	E-Mycin
a) Up to 20 tab available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP Cross for each lin 200 ma not 5 ml.	6.50	100 ml	√ E Musin
Grans for oral liq 200 mg per 5 ml	0.53	100 ml	✓ E-Mycin
Grans for oral liq 400 mg per 5 ml	9.41	100 ml	✓ E-Mycin
ROXITHROMYCIN			
Tab 150 mg	13.19	50	Arrow- Roxithromycin
Tab 300 mg	25.00	50	✓ <u>Arrow-</u> <u>Roxithromycin</u>

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully Brand or sidised Generic Manufacturer
Penicillins			
AMOXICILLIN			
Cap 250 mg	27.50	500	✓ Miro-Amoxicillin
a) Up to 30 cap available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPP			
Cap 500 mg	41.00	500	✓ <u>Miro-Amoxicillin</u>
a) Up to 30 cap available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPP			
Grans for oral liq 125 mg per 5 ml	2.22	100 ml	✓ Alphamox 125
a) Up to 200 ml available on a PSO			
b) Wastage claimable	0.04	400	. Alabaman 050
Grans for oral liq 250 mg per 5 ml	2.81	100 ml	✓ Alphamox 250
a) Up to 300 ml available on a PSO			
b) Up to 10 x the maximum PSO quantity for RFPPc) Wastage claimable			
Inj 250 mg vial	15 97	10	✓ Ibiamox
Inj 500 mg vial		10	✓ Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO		10	✓ Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab			
available on a PSO	1.59	10	✓ Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 r		. •	<u> </u>
per ml	-	100 ml	✓ Augmentin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
c) Augmentin to be Principal Supply on 1 May 2025			
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r	mg		
per ml - Up to 200 ml available on a PSO	4.65 1	100 ml OP	✓ Curam
	5.61		Amoxiclav Devatis
			Forte
Amoxiclav Devatis Forte to be Principal Supply on 1 June			
(Curam Grans for oral liq amoxicillin 50 mg with clavulanic acid 12	2.5 mg per ml to be	delisted 1	June 2025)
BENZATHINE BENZYLPENICILLIN			
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj			<i>4</i>
available on a PSO	432.37	10	✓ Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]			
Inj 600 mg (1 million units) vial - Up to 5 inj available on a PS	SO 16.50	10	✓ Sandoz

	Subsidy (Manufacturer's Price) 8	Fully Brand or Subsidised Generic	
	\$	Per	✓ Manufacti	urer
FLUCLOXACILLIN				
Cap 250 mg - Up to 30 cap available on a PSO	15.79	250	✓ Flucloxacil	lin-AFT
	22.58		Staphlex	
Cap 500 mg - Up to 30 cap available on a PSO	52.99	500	✓ Flucloxacil	lin-AFT
	72.71		Staphlex	
Grans for oral liq 25 mg per ml	4.89	100 ml	✓ <u>AFT</u>	
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 50 mg per ml	5.89	100 ml	✓ AFT	
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Inj 250 mg vial	42.60	10	✓ Flucloxin	
Inj 500 mg vial	45.63	10	✓ Flucloxin	
Inj 1 g vial - Up to 5 inj available on a PSO	6.00	5	✓ Flucil	
(Flucloxacillin-AFT Cap 250 mg to be delisted 1 August 2025)				
(Flucloxacillin-AFT Cap 500 mg to be delisted 1 August 2025)				
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg – Up to 30 cap available on a PSO	7 68	50	✓ Cilicaine V	K
Cap 500 mg		50	✓ Cilicaine V	_
, ,	10.72	50	• Cilicanie vi	<u>r</u>
a) Up to 20 cap available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml	2.40	100 ml	√ AET	
	3.40	100 1111	✓ <u>AFT</u>	
a) Up to 200 ml available on a PSO				
b) Wastage claimable	4.04	4001	(AFT	
Grans for oral liq 250 mg per 5 ml	4.24	100 ml	✓ <u>AFT</u>	
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Tetracyclines				
DOXYCYCLINE				
* Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	Doxine	
MINOCYCLINE HYDROCHLORIDE				
* Tab 50 mg - Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy	5.79	60		
	(12.05)	•••	Mino-tabs	
* Cap 100 mg		100		
	(52.04)		Minomycin	

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

	Subsidy (Manufacturer's Price) \$	Per	Full Subsidised	
Other Antibiotics				
For topical antibiotics, refer to DERMATOLOGICALS, page 66 CIPROFLOXACIN Recommended for patients with any of the following:				
 i) microbiologically confirmed and clinically significant pse ii) prostatitis; or iii) pyelonephritis; or iv) gonorrhoea. 	eudomonas iniection,	OI		
Tab 250 mg - Up to 5 tab available on a PSO	1.95	28	/	Ipca-Ciprofloxacin
Tab 500 mg - Up to 5 tab available on a PSO	3.10	28		Ipca-Ciprofloxacin
Tab 750 mg	4.80	28	•	Ipca-Ciprofloxacin
CLINDAMYCIN				
Cap hydrochloride 150 mg		24		Dalacin C
Inj 150 mg per ml, 4 ml ampoule	35.10	10	•	<u>Hameln</u>
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S				
Only if prescribed for dialysis or cystic fibrosis patient and the		rsed		
Inj 150 mg		1	•	Colistin-Link
Inj 2 million iu, 10 ml vial	216.67	10	•	Colomycin S29
GENTAMICIN SULPHATE				
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement	36.70	5	•	Cidomycin P/Free S29
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	t infection	and the prescription is
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement		5		DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	t infection	and the prescription is
Inj 10 mg per ml, 2 ml ampoule – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.		5 trac		Wockhardt \$29 and the prescription is
Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement	18.38	10		Gentamicin Amdipharm S29
				Pfizer
	91.90	50	•	Gentamicin Noridem S29
Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.	or complicated urinary	trac	t infection	and the prescription is
MOXIFLOXACIN - Special Authority see SA1740 below - Retail	pharmacy			

No patient co-payment payable

Tab 400 mg42.00 5 ✓ Avelox

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Both:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.1 Active tuberculosis*; and
- 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Fither:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

⇒SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

	Subsidy (Manufacturer's Price) \$	Per		Brand or Generic Manufacturer
SODIUM FUSIDATE [FUSIDIC ACID] Tab 250 mg		36	√ Fu	cidin
Tab 500 mg	, ,	100	✓ Su	lfadiazin-Heyl S29
	543.20	56	✓ Wo	ockhardt \$29

SA1331 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy: or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

TOBRAMYCIN

Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement15.50	5	✓ Tobramycin (Viatris)
Only if prescribed for dialysis or cystic fibrosis patient and the prescripti	on is endorsed	accordingly.
Solution for inhalation 60 mg per ml, 5 ml - Subsidy by		
endorsement395.00	56 dose	✓ Tobramycin BNM

a) Wastage claimable

b) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly.

TRIMETHOPRIM

*	Tab 300 mg - Up to 30 tab available on a PSO27.83	50	✓ <u>TMP</u>
---	---	----	--------------

TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]

*	Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - Up	•		
	to 30 tab available on a PSO	115.74	500	✓ Trisul

∀ Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 ml
 available on a PSO.......4.95
 100 ml
 ✓ Deprim

VANCOMYCIN - Subsidy by endorsement

Only if prescribed for a dialysis or cystic fibrosis patient or for prophylaxis of endocarditis or for treatment of Clostridium difficile following metronidazole failure and the prescription is endorsed accordingly.

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 67
- b) For topical antifungals refer to GENITO URINARY, page 79

FLUCONAZOLE

Cap 50 ı	mg	4.10	28	•	Mylan
Cap 150) mg	0.45	1	✓	Mylan
) mg		28	✓	Mylan
Powder	for oral suspension 10 mg per ml - Special Authority				
	SA1359 below – Retail pharmacy	129.02	35 ml	✓	Diflucan

Wastage claimable

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOI F

Cap 100 mg	3 15	✓ Itrazole
27.3	2 60	✓ Itracap S29
Oral liq 10 mg per ml — Special Authority see SA1322 below — Retail pharmacy141.8	0 150 ml OP	✓ Itraconazole Kent \$29

(Sporanox Oral liq 10 mg per ml to be delisted 1 April 2025)

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCT	CBS	30	✓ Burel S29
•		100	✓ Strides Shasun S29
			✓ Taro S29
			✓ Teva-
			Ketoconazole S29
NYSTATIN			

N

Tab 500,000 u	14.16	50	
•	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat

	Subsidy (Manufacturer's Price	e) Subsi	Fully idised	Brand or Generic
	\$	Per	1	Manufacturer
POSACONAZOLE - Special Authority see SA2383 below - Ret	ail pharmacy			
Tab modified-release 100 mg	206.00	24	✓ <u>P</u>	osaconazole Juno
Oral liq 40 mg per ml	342.51	105 ml OP	✓ <u>D</u>	<u>levatis</u>

⇒SA2383 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

TERBINAFINE			
* Tab 250 mg	8.97	84	✓ <u>Deolate</u>
VORICONAZOLE - Special Authority see SA2384 on the next page	je – Retail phar	macy	
Tab 50 mg	71.00	56	✓ Vttack
Tab 200 mg	263.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage			
claimable	1,523.22	70 ml	✓ Vfend

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2384 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Initial application — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Fither:
 - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Renewal — (Invasive fungal infection prophylaxis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Antimalarials

	Subsidy	Fully	Brand or
(Manufa	, ,	sidised	Generic
	\$ Per	· ·	Manutacturer

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	25.86	250	✓ <u>Metronidamed</u>
Tab 400 mg - Up to 15 tab available on a PSO	4.29	21	✓ <u>Metronidamed</u>
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg		10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.52	10	✓ <u>Arrow-Ornidazole</u>

Antituberculotics and Antileprotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

BEDAQUILINE - Special Authority see SA2244 below - Retail pharmacy

No patient co-payment payable

⇒SA2244 Special Authority for Subsidy

Initial application — (multi-drug resistant tuberculosis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The person has multi-drug resistant tuberculosis (MDR-TB); and
- 2 Ministry of Health's Tuberculosis Clinical Network has reviewed the individual case and recommends bedaquiline as part of the treatment regimen.

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

CYCLOSERINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.

	Subsidy (Manufacturer's Price	۱ ۹	Fully Subsidised	
	(Manufacturer's Frice	Per	oubsidised •	Manufacturer
DAPSONE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommend	dation of, an infectious	disease	physicia	n, clinical microbiologist o
dermatologist				_
Tab 25 mg		100		Dapsone
Tab 100 mg		100	•	Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Speci	alist			
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend	dation of, an infectious	disease	physicia	n, clinical microbiologist o
respiratory physician Tab 100 mg	85.73	100	_	EMB Fatol S29
Tab 400 mg		56	_	Myambutol S29
· ·	43.34	30	•	Wyambutor
ISONIAZID – Retail pharmacy-Specialist				
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommend	Nation of an internal me	odicino r	hycioian	naodiatrioian olinical
microbiologist, dermatologist or public health physiciar		edicine p	niysician	, paeulatriciari, ciiriicai
* Tab 100 mg		100	1	PSM
· · · · · · · · · · · · · · · · · · ·	94.50			Isoniazid Teva S29
	327.41			Noumed Isoniazid
Noumed Isoniazid to be Principal Supply on 1 May 20)25			
(PSM Tab 100 mg to be delisted 1 May 2025)				
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist				
a) No patient co-payment payable				
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommend		edicine p	ohysician	, paediatrician, clinical
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physician	١			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physician * Tab 100 mg with rifampicin 150 mg 	n 89.82	100	/	Rifinah
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physician * Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg 	n 89.82 179.13		/	
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physician * Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg LINEZOLID – Special Authority see SA2234 below – Retail pl 	n 89.82 179.13	100	/	Rifinah
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physician * Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg LINEZOLID – Special Authority see SA2234 below – Retail pl No patient co-payment payable 	n 89.82 179.13 narmacy	100 100	1	Rifinah Rifinah
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg LINEZOLID – Special Authority see SA2234 below – Retail pl No patient co-payment payable Tab 600 mg 	n89.82 179.13 narmacy194.60	100 100	<i>*</i>	Rifinah Rifinah
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physician * Tab 100 mg with rifampicin 150 mg	n89.82 179.13 narmacy194.60	100 100	<i>*</i>	Rifinah Rifinah
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg LINEZOLID – Special Authority see SA2234 below – Retail pl No patient co-payment payable Tab 600 mg Oral liq 20 mg per ml * SA2234 Special Authority for Subsidy 	89.82 179.13 narmacy 194.60 1,879.00	100 100 10 10 150 ml	<i>y y y y</i>	Rifinah Rifinah Zyvox Zyvox
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg LINEZOLID – Special Authority see SA2234 below – Retail pl No patient co-payment payable Tab 600 mg Oral liq 20 mg per ml *SA2234 Special Authority for Subsidy Initial application — (multi-drug resistant tuberculosis) for 	89.82 179.13 narmacy 194.60 1,879.00	100 100 10 10 150 ml	<i>y y y y</i>	Rifinah Rifinah Zyvox Zyvox
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg LINEZOLID – Special Authority see SA2234 below – Retail pl No patient co-payment payable Tab 600 mg Oral liq 20 mg per ml * SA2234 Special Authority for Subsidy 	89.82 179.13 narmacy 194.60 1,879.00	100 100 10 10 150 ml	<i>y y y y</i>	Rifinah Rifinah Zyvox Zyvox
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg LINEZOLID – Special Authority see SA2234 below – Retail pl No patient co-payment payable Tab 600 mg Oral liq 20 mg per ml * SA2234 Special Authority for Subsidy Initial application — (multi-drug resistant tuberculosis) fro applications meeting the following criteria: Both:		100 100 10 10 150 ml	<i>y y y y</i>	Rifinah Rifinah Zyvox Zyvox
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg		100 100 10 10 150 ml	pprovals	Rifinah Rifinah Zyvox Zyvox valid for 18 months for
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg		100 100 10 10 150 ml	pprovals	Rifinah Rifinah Zyvox Zyvox valid for 18 months for
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg		100 100 10 10 150 ml	pprovals	Rifinah Rifinah Zyvox Zyvox valid for 18 months for
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg	n	100 100 10 150 ml oner. A	pprovals	Rifinah Rifinah Zyvox Zyvox valid for 18 months for mends linezolid as part of
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg	n	100 100 10 150 ml oner. A	pprovals	Rifinah Rifinah Zyvox Zyvox valid for 18 months for mends linezolid as part of
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg	n	100 100 100 150 ml oner. A	pprovals d recom	Rifinah Rifinah Zyvox Zyvox valid for 18 months for mends linezolid as part of
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg	n	100 100 10 150 ml oner. A	pprovals d recom	Rifinah Rifinah Zyvox Zyvox valid for 18 months for mends linezolid as part of
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg	n	100 100 100 150 ml oner. A	pprovals d recom	Rifinah Rifinah Zyvox Zyvox valid for 18 months for mends linezolid as part of
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg	1	100 100 100 150 ml oner. A case an	pprovals d recom	Rifinah Rifinah Zyvox Zyvox valid for 18 months for mends linezolid as part of
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg	1	100 100 100 150 ml oner. A case an	pprovals d recom	Rifinah Rifinah Zyvox Zyvox valid for 18 months for mends linezolid as part of
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physiciar * Tab 100 mg with rifampicin 150 mg		100 100 100 150 ml oner. A case an	pprovals d recom	Rifinah Rifinah Zyvox Zyvox valid for 18 months for mends linezolid as part of

		Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	I Generic				
PY	RAZINAMIDE - Retail pharmacy-Specialist								
	 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendar respiratory physician 	tion of, an infectious	diseas	e physicia	n, clinical microbiologist or				
*	Tab 500 mg	64.95	100	•	AFT-Pyrazinamide				
RIF	ABUTIN - Retail pharmacy-Specialist								
	a) No patient co-payment payable								
	b) Prescriptions must be written by, or on the recommenda gastroenterologist			, ,					
*	Cap 150 mg	353.71	30	•	Mycobutin				
RIF	AMPICIN - Subsidy by endorsement								
	a) No patient co-payment payable								
	b) For confirmed recurrent Staphylococcus aureus infection								
	antimicrobial based on susceptibilities and the prescriptic Retail pharmacy - Specialist. Specialist must be an inter								
	paediatrician, or public health physician.	mai medicine priyalei	uii, oiii	iloai ililoi o	biologist, dermatologist,				
*	Cap 150 mg	58.54	100	1	Rifadin				
*	Cap 300 mg		100		<u>Rifadin</u>				
*	Oral liq 100 mg per 5 ml	12.60	60 m		Rifadin Sanofi Rifadin				
A	ntivirals								
For	eye preparations refer to Eye Preparations, Anti-Infective Pre	eparations, page 266)						
Н	epatitis B Treatment								
	TECAVIR Tab 0.5 mg	12.04	30	/	Entecavir (Rex)				
	MIVUDINE - Special Authority see SA1685 below - Retail pl								
_,	Tab 100 mg		28	1	Zetlam				
	Oral liq 5 mg per ml	270.00 2	40 ml (OP 🗸	Zeffix				
>	SA1685 Special Authority for Subsidy								
Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.									
Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.									
	Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B.								
Rei	IOCOVID DICODDOVII								
Rei	NOFOVIR DISOPROXIL Tenofovir disporovil prescribed under endorsement for the tr	reatment of HIV is in	hahuk	in the cou	nt of up to 4 subsidised				
Rei	NOFOVIR DISOPROXIL Tenofovir disoproxil prescribed under endorsement for the to antiretrovirals for the purposes of Special Authority SA2139.		cluded	in the cou	nt of up to 4 subsidised				

Herpesvirus	Treatment	ts
-------------	-----------	----

ACICLOVIR			
* Tab dispersible 200 mg	1.78	25	✓ Lovir
* Tab dispersible 400 mg	5.81	56	✓ Lovir
* Tab dispersible 800 mg		35	✓ Lovir
VALACICLOVIR			
Tab 500 mg	9.64	30	✓ Vaclovir
Tab 1,000 mg	17.78	30	✓ Vaclovir
-			

	Subsidy (Manufacturer's Price)	Fully Subsidised		Brand or Generic
	\$	Per	1	Manufacturer
VALGANCICLOVIR - Special Authority see SA1993 below - F	Retail pharmacy			
Tab 450 mg	140.89	60	✓ <u>v</u>	alganciclovir Viatris

⇒SA1993 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Roth:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Fither:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for a months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or

INFECTIONS - AGENTS FOR SYSTEMIC USE							
		Subsidy (Manufacturer's Price \$) Sub	Fully sidised	Brand or Generic Manufacturer		
	ned 2.2 Patient has rapidly rising plasma CMV DNA in a 2.3 Patient has cytomegalovirus retinitis. for the purpose of this Special Authority "immunocomprosuppressive diseases (e.g. HIV) or those receiving im	omised" includes trans					
Нер	atitis C Treatment						
No we Tai LEDIP. No Tai Special Notes: Applica	APREVIR WITH PIBRENTASVIR – [Xpharm] the the supply of treatment is via Pharmac's approved disestite https://pharmac.govt.nz/maviret the 100 mg with pibrentasvir 40 mg	24,750.00 nority see SA1605 belo24,363.46 I (HepCTP) DCTP). ect to confirmation of 6	84 OP	✓ N	an be found on Pharmac's laviret larvoni		
The Co	pordinator, Hepatitis C Treatment Panel ac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990, hepcpanel@pharmac.govt.nz		T. T. Z. T. T. C. T.	<u>or</u> on			
HIV	Prophylaxis and Treatment						
EMTRICITABINE WITH TENOFOVIR DISOPROXIL — Subsidy by endorsement; can be waived by Special Authority see SA2138 below a) Funding for emtricitabine with tenofovir disoproxil for use as PrEP, should be applied using Special Authority SA2138. b) Endorsement for treatment of conditions approved via Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure): Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.							
	Note: Emtricitabine with tenofovir disoproxil prescribed	d under endorsement,	for treatme	ent of co	nditions approved via		

Special Authority SA2139 (antiretrovirals for confirmed HIV, prevention of maternal transmission, post-exposure prophylaxis following exposure to HIV and percutaneous exposure), is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 109 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website

*	Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a maleate)	15.45	30	•	Tenofovir Disoproxil Emtricitabine Viatr
*	Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a				
	succinate)	15.45	30	1	Teva
(Te	eva Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a	succinate) to b	e delisted 1 /	August 2	2025)
-	SA2138 Special Authority for Subsidy				

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

COVID-19 Treatments

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml: or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (**Percutaneous exposure**) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ – Special Authority see SA2139 on the previous pa Tab 600 mg	•	rmacy 30	✓ Efavirenz Milpharm 529
ETRAVIRINE – Special Authority see SA2139 on the previous partial 200 mg	•	armacy 60	✓ Intelence
NEVIRAPINE - Special Authority see SA2139 on the previous pa	age – Retail pha	armacy	
Tab 200 mg	198.25	60	 Nevirapine Viatris
Oral suspension 10 mg per ml	203.55	240 ml OP	✓ Viramune Suspension

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully Brand or dised Generic Manufacturer
Nucleosides Reverse Transcriptase Inhibitors			
ABACAVIR SULPHATE - Special Authority see SA2139 on page Tab 300 mg		acy 60	✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE — Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.			, ,
Tab 600 mg with lamivudine 300 mg	29.50	30	✓ <u>Abacavir/</u> <u>Lamivudine</u> Viatris
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPR pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil co anti-retroviral Special Authority		-	SA2139 on page 109 – Retail
Tab 600 mg with emtricitabine 200 mg and tenofovir disoprox 245 mg (300 mg as a fumarate)	106.88 :il	30	✓ Triovir \$29
245 mg (300 mg as a maleate)		30	✓ Viatris
EMTRICITABINE - Special Authority see SA2139 on page 109 - Cap 200 mg		30	✓ Emtriva
LAMIVUDINE – Special Authority see SA2139 on page 109 – Re Tab 150 mg Oral liq 10 mg per ml	98.00	60 10 ml OP	✓ <u>Lamivudine Viatris</u> ✓ 3TC
ZIDOVUDINE [AZT] - Special Authority see SA2139 on page 108 Cap 100 mg Oral liq 10 mg per ml	152.25	100 00 ml OP	✓ Retrovir ✓ Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE - Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets) the anti-retroviral Special Authority.			
Tab 300 mg with lamivudine 150 mg	92.40	60	Lamivudine/ Zidovudine Viatris
Protease Inhibitors			
ATAZANAVIR SULPHATE - Special Authority see SA2139 on pa	age 109 – Retail pha	armacy	
Cap 150 mg	85.00	60	 ✓ Atazanavir Mylan ✓ Atazanavir Viatris
Cap 200 mg	110.00	60	✓ <u>Atazanavir Viatris</u>
DARUNAVIR - Special Authority see SA2139 on page 109 - Ret			
Tab 400 mg		60	✓ <u>Darunavir Viatris</u>
Tab 600 mg		60	✓ Darunavir Viatris
LOPINAVIR WITH RITONAVIR – Special Authority see SA2139 of Tab 100 mg with ritonavir 25 mg		il pharmacy 60	✓ Lopinavir/Ritonavir Mylan
Tab 200 mg with ritonavir 50 mg	875.00	120	✓ <u>Lopinavir/Ritonavir</u> Mylan
RITONAVIR – Special Authority see SA2139 on page 109 – Reta	ail pharmacy	00	- Namela

30

✓ Norvir

Tab 100 mg43.31

•				
	Subsidy	Fully	Brand or	
	(Manufacturer's Price)	Subsidised	l Generic	
	\$	Per 🗸	Manufacturer	

Strand Transfer Inhibitors

DOLUTEGRAVIR – Special Authority see SA2139 on page	109 – Retail pharmacy		
Tab 50 mg	1,090.00	30	Tivicay
DOLUTEGRAVIR WITH LAMIVUDINE - Special Authority	see SA2139 on page 10	9 – Retail p	harmacy
Tab 50 mg with lamivudine 300 mg	1,090.00	30	✓ Dovato
RALTEGRAVIR POTASSIUM - Special Authority see SA21	139 on page 109 – Reta	il pharmacy	
Tab 400 mg	1,090.00	60	Isentress
Tah 600 mg	1 090 00	60	✓ Isontross HD

Immune Modulators

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

Inj 180 mcg prefilled syringe......748.50

✓ Pegasys

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	osidised	Generic
\$	Per	1	Manufacturer

continued...

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV. HIV or HDV: and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*: or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

⇒SA2406 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli: and
- 2 Either:
 - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
 - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

Renewal from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria: Both:

- 1 Patient has an acute, symptomatic, bacteriologically-proven uncomplicated urinary tract infection (UTI)/cystitis with Escherichia Coli; and
- 2 Either:
 - 2.1 Microbiological testing confirms the pathogen is resistant to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin; or
 - 2.2 The patient has a contraindication or intolerance to all of: trimethoprim, nitrofurantoin, amoxicillin, cefaclor, cefalexin, amoxicillin with clavulanic acid, and norfloxacin that the pathogen is susceptible to.

METHENAMINE (HEXAMINE) HIPPURATE

* Tab 1 g	19.95	100	✓ <u>Hiprex</u>
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg - Up to 15 cap available on a			
PSO	81.20	100	✓ Macrobid
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	245.00	100	✓ Arrow-Norfloxacin
On health and a suite and the second and suite and an arrangement and a suite			

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy (Manufacturer's Pri	ica) Sub	Fully Brand or sidised Generic
	(Manufacturer's Fri	Per Sul.	✓ Manufacturer
Anticholinesterases			
OSTIGMINE METILSULFATE	40.05	10	√ May Haalth
Inj 2.5 mg per ml, 1 ml ampoule	46.25	10	✓ <u>Max Health</u>
/RIDOSTIGMINE BROMIDE	E0 00	100	./ Maatinan
Tab 60 mg	50.28	100	✓ Mestinon
Ion-Steroidal Anti-Inflammatory Drugs			
CLOFENAC SODIUM			
Tab EC 25 mg	2.19	50	✓ Diclofenac Sandoz
Tab 50 mg dispersible	1.50	20	✓ Voltaren D
Tab EC 50 mg	2.19	50	✓ Diclofenac Sandoz
Tab long-acting 75 mg		100	✓ Voltaren SR
Inj 25 mg per ml, 3 ml ampoule - Up to 5 inj available on a		5	✓ Voltaren
Suppos 12.5 mg	2.04	10	✓ Voltaren
Suppos 25 mg		10	✓ Voltaren
Suppos 50 mg - Up to 10 supp available on a PSO		10	✓ Voltaren
Suppos 100 mg	7.00	10	✓ Voltaren
UPROFEN			
Tab 200 mg	21.40	1,000	✓ Relieve
Tab long-acting 800 mg	3.05	30	✓ Brufen SR
	3.65		Ibuprofen SR BNM
Ibuprofen SR BNM to be Principal Supply on 1 April 202			
Oral liq 20 mg per ml	2.85	200 ml	Ethics
Ethics to be Principal Supply on 1 April 2025			
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN			
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) TOPROFEN	12.07	28	✓ Oruvail SR
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg	12.07	28	✓ Oruvail SR
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID		28 50	✓ Oruvail SR
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) TOPROFEN Cap long-acting 200 mg EFENAMIC ACID			✓ Oruvail SR Ponstan
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) TOPROFEN Cap long-acting 200 mg EFENAMIC ACID	1.25		
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) TOPROFEN Cap long-acting 200 mg EFENAMIC ACID	1.25 (10.82)	50	
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg	1.25 (10.82) 0.50	50	Ponstan
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg	1.25 (10.82) 0.50 (7.50)	50	Ponstan
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025)	1.25 (10.82) 0.50 (7.50)	50 20	Ponstan Ponstan
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg	1.25 (10.82) 0.50 (7.50) 39.23 34.45	50 20 500	Ponstan Ponstan ✓ Noflam 250
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg Tab long-acting 750 mg	1.25 (10.82) 0.50 (7.50) 39.23 34.45 10.40	50 20 500 250	Ponstan Ponstan Noflam 250 Noflam 500
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg Tab long-acting 750 mg Tab long-acting 1 g	1.25 (10.82) 0.50 (7.50) 39.23 34.45 10.40	50 20 500 250 28	Ponstan Ponstan Noflam 250 Noflam 500 Naprosyn SR 750
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg	1.25 (10.82) 0.50 (7.50) 39.23 34.45 10.40 11.50	50 20 500 250 28 28	Ponstan Ponstan Noflam 250 Noflam 500 Naprosyn SR 750 Naprosyn SR 1000
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg Tab long-acting 750 mg Tab long-acting 1 g		50 20 500 250 28	Ponstan Ponstan Noflam 250 Noflam 500 Naprosyn SR 750
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg Tab long-acting 750 mg Tab long-acting 1 g ENOXICAM Tab 20 mg Inj 20 mg vial		50 20 500 250 28 28 100	Ponstan Ponstan Noflam 250 Noflam 500 Naprosyn SR 750 Naprosyn SR 1000 Tilcotil
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg Tab long-acting 750 mg Tab long-acting 1 g ENOXICAM Tab 20 mg Inj 20 mg vial		50 20 500 250 28 28 100	Ponstan Ponstan Noflam 250 Noflam 500 Naprosyn SR 750 Naprosyn SR 1000 Tilcotil
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg Tab long-acting 750 mg Tab long-acting 1 g ENOXICAM Tab 20 mg Inj 20 mg vial INSAIDS Other ELECOXIB		50 20 500 250 28 28 100 1	Ponstan Ponstan Noflam 250 Noflam 500 Naprosyn SR 750 Naprosyn SR 1000 Tilcotil AFT
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg Tab long-acting 750 mg Tab long-acting 1 g ENOXICAM Tab 20 mg Inj 20 mg vial		50 20 500 250 28 28 100	Ponstan Ponstan Noflam 250 Noflam 500 Naprosyn SR 750 Naprosyn SR 1000 Tilcotil AFT
rufen SR Tab long-acting 800 mg to be delisted 1 April 2025) ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg Tab long-acting 750 mg Tab long-acting 1 g ENOXICAM Tab 20 mg Inj 20 mg vial INSAIDS Other ELECOXIB		50 20 500 250 28 28 100 1	Ponstan Ponstan Noflam 250 Noflam 500 Naprosyn SR 750 Naprosyn SR 1000 Tilcotil AFT

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or	
(Manufacturer's Pr	rice) Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

Topical Products for Joint and Muscular Pain

CAPSAICIN

(Rugby Capsaicin Topical Cream S29 Crm 0.025% to be delisted 1 July 2025)

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE SULPHATE ★ Tab 200 mg7.80	100	✓ lpca- Hydroxychloroquine
8.78		✓ Plaquenil
Ipca-Hydroxychloroquine to be Principal Supply on 1 May 2025 (Plaquenil Tab 200 mg to be delisted 1 May 2025)		
LEFLUNOMIDE		
* Tab 10 mg6.00	30	✓ Arava
* Tab 20 mg6.00	30	✓ Arava
PENICILLAMINE		
Tab 125 mg	100	✓ D-Penamine
Tab 250 mg110.12	100	✓ D-Penamine

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODIUM * Tab 70 mg	3.10	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL			
* Tab 70 mg with colecalciferol 5,600 iu	1.99	4	✓ Fosamax Plus

Other Treatments

DENOSUMAB - Special Authority see SA2441 on the next page - Retail pharmacy

Note: Denosumab inj 60 mg per 1 ml pre-filled syringe is Medsafe approved for use in osteoporosis. Denosumab inj 120 mg per 1.7 ml vial is Medsafe approved for use in hypercalcaemia of malignancy.

h			
Inj 120 mg per 1.7 ml vial	500.00	1	Xgeva
Inj 60 mg per 1 ml prefilled syringe		1	✓ Prolia

Cream S29

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

⇒SA2441 Special Authority for Subsidy

Initial application — (Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has established osteoporosis; and
- 2 Any of the following:
 - 2.1 History of one significant osteoporotic fracture demonstrated radiologically, with a documented T-Score less than or equal to -2.5, that incorporates BMD measured using dual-energy x-ray absorptiometry (DEXA); or
 - 2.2 History of one significant osteoporotic fracture, demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of logistical, technical or pathophysiological reasons; or
 - 2.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 2.4 Documented T-Score less than or equal to -3.0; or
 - 2.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm that incorporates BMD measured using DEXA; and
- 3 Any of the following:
 - 3.1 Bisphosphonates are contraindicated because the patient's creatinine clearance or eGFR is less than 35 mL/min; or
 - 3.2 The patient has experienced at least two symptomatic new fractures or a BMD loss greater than 2% per year, after at least 12 months' continuous therapy with a funded antiresorptive agent; or
 - 3.3 Bisphosphonates result in intolerable side effects; or
 - 3.4 Intravenous bisphosphonates cannot be administered due to logistical or technical reasons.

Initial application — (Hypercalcaemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has hypercalcaemia of malignancy; and
- 2 Patient has severe renal impairment.

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	32.49	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial		1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	94.34	1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE – Special Authority see SA177		pharmacy	
* Tab 60 mg	53.76	28	Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or

	Subsidy (Manufacturer's Price)	Fully Subsidised	
	\$	Per 🗸	Manufacturer
continued			

continued...

6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause - Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

Tab 35 mg	2.50	4	✓ Risedronate Sandoz
TERIPARATIDE - Special Authority see SA1139 below -	Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml	195.00	1	✓ Teriparatide - Teva

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

$7 \cap 1$		\neg	MIIC	AC	ın
∠∪∟	.ロレ	טחי	NIC	AU	טו

Inj 0.05 mg per ml, 100 ml, bag22.53	100 ml OP	Zoledronic Acid
		Viatris

Hyperuricaemia and Antigout

ALI	LOPURINOL			
*	Tab 100 mg	17.99	1,000	✓ <u>Ipca-Allopurinol</u>
*	Tab 300 mg	22.50	500	✓ <u>Ipca-Allopurinol</u>

Subsidy (Manufacturer's Price) \$ Pe BENZBROMARONE — Special Authority see SA1963 below — Retail pharmacy Tab 50 mg	ng the follow; and least every for applications at the following to least every for applications at treatment at treatment of the following to least every for a f	Generic Manufacturer Narcaricin mite \$29 wing criteria: three months) liver function Colgout Febuxostat (Teva) Febuxostat (Teva) ions meeting the following purinol at doses of at least ated dose; or t discontinuation is required
Tab 50 mg	g the follow ; and least every for applicati	three months) liver function Colgout Febuxostat (Teva) Febuxostat (Teva) ions meeting the following purinol at doses of at least ated dose; or t discontinuation is required
Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting Both: 1 The treatment remains appropriate and the patient is benefitting from the treatment 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at tests. COLCHICINE * Tab 500 mcg	; and least every for application to least every for application to least every for at treatment to least every for at treatment every for a least every for	three months) liver function Colgout Febuxostat (Teva) Febuxostat (Teva) ions meeting the following purinol at doses of at least ated dose; or t discontinuation is required
COLCHICINE * Tab 500 mcg	for application for application for application for application for a formal fo	Febuxostat (Teva) Febuxostat (Teva) ions meeting the following purinol at doses of at least ated dose; or t discontinuation is required
FEBUXOSTAT - Special Authority see SA2054 below - Retail pharmacy Tab 80 mg	for application for application for application for application for a formal formal for a formal	Febuxostat (Teva) Febuxostat (Teva) ions meeting the following purinol at doses of at least ated dose; or t discontinuation is required
Tab 120 mg	for application for applicatio	rebuxostat (Teva) ions meeting the following purinol at doses of at least ated dose; or t discontinuation is required
Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months criteria: Both: 1 Patient has been diagnosed with gout; and 2 Any of the following: 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatm 600 mg/day and addition of probenecid at doses of up to 2 g per day or max 2.2 The patient has experienced intolerable side effects from allopurinol such th and serum urate remains greater than 0.36 mmol/l despite use of probeneci maximum tolerated dose; or 2.3 The patient has renal impairment such that probenecid is contraindicated or remains greater than 0.36 mmol/l despite optimal treatment with allopurinol 2.4 The patient has previously had an initial Special Authority approval for benzi Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. applications meeting the following criteria: Both: 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk	ent with allo imum tolera at treatmen	opurinol at doses of at least ated dose; or t discontinuation is required
Both: 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk	likely to be (see Note); bromarone t	or for treatment of gout
	of tumour ly	ysis syndrome; and
2 Patient has a documented history of allopurinol intolerance. Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the patient is benefitting from treatment.	treatment re	emains appropriate and the
Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approva treatment remains appropriate and the patient is benefitting from treatment. PROBENECID	als valid for	6 weeks where the
* Tab 500 mg66.95 100	/	Probenecid-AFT
Muscle Relaxants		
BACLOFEN * Tab 10 mg		Pacifen Lioresal Intrathecal we been ineffective or have Sintetica Baclofen

caused intolerable side effects and the prescription is endorsed accordingly.

Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have

Intrathecal

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	•	Manufacturer
DANTROLENE				
Cap 25 mg	112.13	100	1	Dantrium
	145.77		1	Dantrium S29 S29
Cap 50 mg	77.00	100	1	Dantrium
(Dantrium Cap 25 mg to be delisted 1 April 2025)				
ORPHENADRINE CITRATE				
Tab 100 mg	23.25	100	1	Norflex

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

AMANTADINE HYDROCHLORIDE		
▲ Cap 100 mg	60	✓ Symmetrel
63.73	100	Symmetrel
APOMORPHINE HYDROCHLORIDE		
▲ Inj 10 mg per ml, 2 ml ampoule59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule121.84	5	✓ Movapo
ENTACAPONE		
▲ Tab 200 mg	100	✓ Entacapone Viatris
18.04		✓ Comtan
(Comtan Tab 200 mg to be delisted 1 July 2025)		
LEVODOPA WITH BENSERAZIDE		
* Tab dispersible 50 mg with benserazide 12.5 mg	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg13.75	100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg	100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg22.85	100	Madopar HBS
* Cap 200 mg with benserazide 50 mg26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA		
* Tab 100 mg with carbidopa 25 mg26.49	100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg44.99	100	✓ Sinemet CR
* Tab 250 mg with carbidopa 25 mg	100	✓ Sinemet
LEVODOPA WITH CARBIDOPA AND ENTACAPONE		
* Tab 50 mg with carbidopa 12.5 mg and entacapone 200 mg27.01	100	✓ Stalevo
* Tab 100 mg with carbidopa 25 mg and entacapone 200 mg34.18	100	✓ Stalevo
* Tab 150 mg with carbidopa 37.5 mg and entacapone 200 mg44.96	100	✓ Stalevo
* Tab 200 mg with carbidopa 50 mg and entacapone 200 mg51.23	100	✓ Stalevo
PRAMIPEXOLE HYDROCHLORIDE		
▲ Tab 0.25 mg	100	✓ Ramipex
▲ Tab 1 mg	100	✓ Ramipex
RASAGILINE		
* Tab 1 mg53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE		
▲ Tab 0.25 mg	84	✓ Ropin
▲ Tab 1 mg	84	✓ Ropin
▲ Tab 2 mg	84	✓ Ropin
▲ Tab 5 mg	84	✓ Ropin
TOLCAPONE		
▲ Tab 100 mg	100	✓ Tasmar

Anticholinergics

BENZATROPINE MESYLATE

Tab 2 mg9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml95.00	5	✓ Phebra

- a) Up to 10 inj available on a PSO
- b) Only on a PSO

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	✓ I	Kemadrin
Agents for Essential Tremor, Chorea and Relate	ed Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharr Wastage claimable Tab 50 mg	•	56	√ 1	Rilutek
■ SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialis following criteria: All of the following:			_	
1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vita 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow.				initial application; and
Renewal from any relevant practitioner. Approvals valid for 18 m All of the following: 1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and 3 Any of the following: 3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow.	nonths for applications	s meef	ting the foll	owing criteria:
TETRABENAZINE Tab 25 mg	106.59	112	✓ <u>I</u>	Motetis
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE] Gel 2% tripe — Subsidy by endorsement	14.50	30 ml	√)	(vlocaine 2% Jelly

Gel 2%, tube – Subsidy by endorsement14.50 30 ml ✓ Xylocaine 2% Jelly

a) Up to 150 ml available on a PSO

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Gel 2%, 11 ml urethral syringe - Subsidy by endorsement......59.50 10 ✓ Instillagel Lido

a) Up to 5 each available on a PSO

b) Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.

00	Per 200 ml 25 50		Generic Manufacturer lucosoothe idocaine-Baxter
00	25		
00	25		
		√ L	idocaine-Baxter
. ^	50		
50	00		
00)		Х	ylocaine
50 [°]	25	✓ L	idocaine-Baxter
00	5		
00)		Х	ylocaine
50 [°]	5	✓ L	idocaine-Baxter
00	5	✓ L	idocaine-Baxter
	10	✓ X	ylocard 500 S29
	50 00 S	00 5 S 10	00 5 ~ L

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority s	see SA0906 above – Retail pl	harmacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE	- Special Authority see SA0	906 above – Ret	ail pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 115

Non-opioid Analgesics

ASPIRIN * Tab dispersible 300 mg - Up to 30 tab available on a F	PSO5.65	100	✓ Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia accordingly.	a or diabetic peripnera	i neuropatny a	ina the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zo-Rub HP S29 ✓ Zostrix HP
	15.14	57 g OP	✓ Rugby Capsaicin Topical Cream \$29
(Rugby Capsaicin Topical Cream S29 Crm 0.075% to be of	lelisted 1 July 2025)		
NEFOPAM HYDROCHLORIDE			
Tah 30 mg	23.40	an	✓ Acupan

		Subsidy (Manufacturer's Price		Fully Brandised Gene	
		\$	Per		ufacturer
PARACETAMOL					
	pack		1,000	✓ Pacimo	<u>) </u>
b) Up to 30 tab	f 300 tab per prescription; can be wai available on a PSO	ived by endorsement			
regula annota 2) Maxim (for no Tab 500 mg - bottle prescription; ca 1) Subsidy b	dy by endorsement for higher quantiti r daily dosing for one month or greate te the prescription as endorsed when lum of 100 tab per dispensing for nor n-endorsed patients), then dispense pack — Maximum of 300 tab per n be waived by endorsement	er, and the prescription re dispensing history sun-endorsed patients. If or in repeat dispensings numbers, 17.92	is annotated a pports a long- quantities presot exceeding 1,000 with long term	accordingly. Iterm condition Scribed for note that the condition Noume Paract Conditions	Pharmacists may ion. nore than 100 tabs dispensing. detamol who require regular
2) Maximum	on as endorsed where dispensing his of 100 tab per dispensing for non-en rsed patients), then dispense in repea	dorsed patients. If qua	ntities prescri		
Oral liq 120 mg per	5 ml	3.98	200 ml	✓ Parace	
	f 600 ml per prescription; can be waiv Il available on a PSO ination	ved by endorsement		<u>,=</u>	<u>,</u>
1) Maxim non-er 2) Subsic regula Pharm conditi 3) Note:	200 ml presentations of paracetamo	epeat dispensing not ex es is available for patier er and the prescription is as endorsed where dis I oral liquid may be supp	ceeding 200 r nts with long to s endorsed or pensing histor	nl per disper erm condition annotated a ry supports a	nsing. ons who require accordingly. a long-term
4) Note:	nacist) under the provisions in Part I on Direct Provision by a pharmacist of under action with immunisation of a child under	up to 200 ml permitted u			
Oral liq 250 mg per a) Maximum of	5 ml f 600 ml per prescription; can be waiv Il available on a PSO	3.35	200 ml	✓ <u>Pamol</u>	
Maxim non-er Subsic regula Pharm conditi		epeat dispensing not exest is available for patient and the prescription is as endorsed where dispensions.	ceeding 200 r nts with long to s endorsed or pensing histor	nl per disper erm conditio annotated a y supports a	nsing. ons who require accordingly. a long-term
Pharm	200 ml presentations of paracetamolacist) under the provisions in Part I of	of Section A			`
	Direct Provision by a pharmacist of unction with immunisation of a child unction				

10

✓ Gacet

				IVL	11V0033131EW
		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
*	Suppos 250 mg		10 50		Gacet Gacet
C	pioid Analgesics				
CC	DEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensina fre	auen	CV	
	Tab 15 mg		100		Noumed
	Tab 30 mg	6.98	100	✓	Noumed
	Tab 60 mg	13.89	100	1	Noumed
DIF	HYDROCODEINE TARTRATE				
	Tab long-acting 60 mg	8.60	60	1	DHC Continus
FF	NTANYL				
. –	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre	equency			
	Inj 50 mcg per ml, 2 ml ampoule		10	✓	Boucher and Muir
	Boucher and Muir to be Principal Supply on 1 May 2025				
	Inj 50 mcg per ml, 10 ml ampoule	9.41	10	•	Boucher and Muir
	Boucher and Muir to be Principal Supply on 1 May 2025			_	
	Patch 12.5 mcg per hour		5		Fentanyl Sandoz
	Patch 25 mcg per hour		5		Fentanyl Sandoz
	Patch 50 mcg per hour		5		Fentanyl Sandoz
	Patch 75 mcg per hour Patch 100 mcg per hour		5 5		Fentanyl Sandoz Fentanyl Sandoz
	• 1	10.37	5	•	remanyi Sandoz
ME	THADONE HYDROCHLORIDE				
	a) Only on a controlled drug form				
	b) No patient co-payment payable	au anau			
	c) Safety medicine; prescriber may determine dispensing fre Tab 5 mg		10	_	Methadone BNM
	Oral lig 2 mg per ml		200 m		Biodone
	Oral lig 5 mg per ml		200 n		Biodone Forte
	Oral lig 10 mg per ml		200 n		Biodone Extra Forte
	Inj 10 mg per ml, 1 ml		10		AFT
МС	ORPHINE HYDROCHLORIDE				
IVIC	a) Only on a controlled drug form				
	b) No patient co-payment payable				
	c) Safety medicine; prescriber may determine dispensing fre	equency			
	Oral liq 1 mg per ml		200 n	nl 🗸	RA-Morph
	Oral liq 2 mg per ml	23.55	200 n	nl 🗸	RA-Morph
	Ovel lie 5 was now and	00.00	000		DA Marriele

✓ RA-Morph

✓ RA-Morph

200 ml

200 ml

Oral liq 10 mg per ml40.25

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
MORPHINE SULPHATE	Ψ	rei		Manuacturer
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	equency			
Tab immediate-release 10 mg	2.80	10	✓	Sevredol
Tab immediate-release 20 mg	5.52	10		Sevredol
Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 30 mg		10		m-Eslon
Cap long-acting 60 mg		10		m-Eslon
Cap long-acting 100 mg		10		m-Eslon
Oral liq 2 mg per ml		100 m		Wockhardt S29
	29.80			Oramorph
			•	Oramorph CDC
		_	_	S29 S29
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5		<u>Medsurge</u>
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a F		5		Medsurge Medsurge
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a F Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a F		5 5		Medsurge Medsurge
	′3Ub.28	э	•	<u>Medsurge</u>
XYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre			,	
Tab controlled-release 5 mg		20		Oxycodone Sandoz
	3.77	28	•	Oxycodone Sandoz
			_	S29 \$29
	4.04	30		OxyContin S29
Tab immediate-release 5 mg		100		Oxycodone Amneal
Tab controlled-release 10 mg		20		Oxycodone Sandoz
	3.77	28	•	Oxycodone Sandoz S29 S29
Tab immediate-release 10 mg	18.77	100	1	Oxycodone Amneal
Tab controlled-release 20 mg	3.41	20	✓	Oxycodone Sandoz
Tab immediate-release 20 mg	26.77	100	✓	Oxycodone Amneal
Tab controlled-release 40 mg	6.67	20	✓	Oxycodone Sandoz
Tab controlled-release 80 mg	12.99	20		Oxycodone Sandoz
Oral liq 1 mg per ml		250 m		Oxycodone Lucis
Inj 10 mg per ml, 1 ml ampoule		5		<u>Hameln</u>
Inj 10 mg per ml, 2 ml ampoule		5		Hameln
Inj 50 mg per ml, 1 ml ampoule		5	•	<u>Hameln</u>
Oxycodone Sandoz S29 S29 Tab controlled-release 5 mg to be	,	5)		
OxyContin S29 Tab controlled-release 5 mg to be delisted 1 Ju	ly 2025)			
Oxycodone Sandoz S29 S29 Tab controlled-release 10 mg to b	oe delisted 1 July 202	?5)		
PARACETAMOL WITH CODEINE - Safety medicine; prescriber	may determine disp	ensing	g frequence	y
* Tab paracetamol 500 mg with codeine phosphate 8 mg	27.50	1,000		Paracetamol +
• • •				Codeine (Relieve)

			NE	RVOUS SYSTEM
	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	' '	10		Naumad Dathidina
Tab 50 mgInj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a F		10 5		Noumed Pethidine DBL Pethidine
inj 50 mg per mi, i mi ampodie – op to 5 mj avaliable on a r	73029.00	5	•	Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a F	'SO30.72	5	✓	DBL Pethidine
TRAMAROL LIVEROCUII ORIDE				Hydrochloride
TRAMADOL HYDROCHLORIDE Tab sustained-release 100 mg	1.05	20		Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg		100		Arrow-Tramadol
σαρ σσ mg		100		7110W Hamado
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine d	lispensina frequency			
Tab 10 mg		100	1	Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg	3.14	100	✓	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; prescr	iber mav determine d	ispen		
Tab 10 mg	•	30		Clomipramine Teva
Tab 25 mg		30		Clomipramine Teva
· ·	16.99	50	1	APO Clomipramine
	39.97	100	1	Anafranil S29
Cap 10 mg	35.50	28	1	Clomipramine Teva
Cap 25 mg	35.50	28	✓	Clomipramine Teva
(Clomipramine Teva Tab 10 mg to be delisted 1 July 2025)				
(Clomipramine Teva Tab 25 mg to be delisted 1 July 2025)				
(Anafranil S29 Tab 25 mg to be delisted 1 July 2025)				
(Clomipramine Teva Cap 10 mg to be delisted 1 July 2025)				
(Clomipramine Teva Cap 25 mg to be delisted 1 July 2025)				
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by en	dorsement			
a) Safety medicine; prescriber may determine dispensing fre	equency			
b) Subsidy by endorsement – Subsidised for patients who w				
2019 and the prescription is endorsed accordingly. Phart		the	prescriptio	n as endorsed where there
exists a record of prior dispensing of dosulepin [dothiepin				5
Tab 75 mg		30		Dosulepin Viatris
Cap 25 mg		50	•	Dosulepin Viatris \$29
IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber	may determine diene	neina	frequenc	
Tab 10 mg		50		y Tofranil
10 mg	10.96	100		Tofranil
Tab 25 mg		28		Imipramine
•				Crescent S29

8.80

50

✓ Tofranil

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; presci	riber may determine o	disper	nsing frequ	iency
Tab 10 mg		100	_	Norpress
Tab 25 mg	6.29	180	•	Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	elective			
TRANYLCYPROMINE SULPHATE				
* Tab 10 mg	22.94	50	1	Parnate
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg	23.60	60	•	<u>Aurorix</u>
* Tab 300 mg	38.50	60	✓	<u>Aurorix</u>
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE				
* Tab 20 mg	2.86	84	•	<u>Celapram</u>
ESCITALOPRAM				
* Tab 10 mg	0.79	28	✓	Ipca-Escitalopram
-	1.07		1	Escitalopram
				(Ethics)
* Tab 20 mg	1.49	28	•	Ipca-Escitalopram
FLUOXETINE HYDROCHLORIDE				
* Tab dispersible 20 mg, scored - Subsidy by endorsement	2.50	28	✓	Fluox
Subsidised by endorsement				
When prescribed for a patient who cannot swallow	whole tablets or caps	sules	and the pr	escription is endorsed
accordingly; or	alo of 20 ma in which	0000	tha proces	intion is doomed to be
When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with				
* Cap 20 mg	3 13	90	1	Arrow-Fluoxetine
PAROXETINE		00	_	7 THOM THUO KOUND
* Tab 20 mg	4 11	90	1	Loxamine
SERTRALINE		00	-	<u> LOXAIIIIIO</u>
* Tab 50 mg	0.00	30	1	Setrona
* Tab 100 mg		30	_	Setrona
* Tab 100 mg	1.74	30	•	<u>Jetrona</u>
Other Antidepressants				
MIRTAZAPINE			_	
Tab 30 mg	2.60	30		Noumed
Tab 45 mg	3.45	30	•	Noumed
VENLAFAXINE				
Cap 37.5 mg		84	_	Enlafax XR
Cap 75 mg		28	_	Enlafax XR
O-n 450	10.32	84		Enlafax XR
Cap 150 mg		28		Enlafax XR
	13.95	84	•	Enlafax XR

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

Antiepilepsy Drugs

Agents for 0	Control of	Status	Epilepticus
--------------	------------	--------	-------------

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency		
Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorsement27.92	5	Hospira
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
c) PSO must be endorsed "not for anaesthetic procedures".		
Rectal tubes 5 mg - Up to 5 tube available on a PSO54.58	5	✓ Stesolid
PHENYTOIN SODIUM		
* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a		
PSO	5	✓ Hospira
* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a		
PSO	5	✓ Hospira
. 33	•	. roop.i.u

Control of Epilepsy

CARRAMAZEDINE

	Tab 200 mg	
*	Tab long-acting 200 mg	

Tab long-acting 200 mg	16.98	100
	33.96	200
Tab 400 mg	34.58	100

Tab long-acting 400 mg.......39.17 * Oral liq 20 mg per ml26.37

CLOBAZAM - Safety medicine; prescriber may determine dispensing frequency Tab 10 mg9.12 CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency

FTHOSUXIMIDE

140.88 **GABAPENTIN**

Note: Not subsidised in combination with subsidised pregabalin * Cap 300 mg......8.45

* Cap 400 mg......10.26 LACOSAMIDE - Special Authority see SA2267 on the next page - Retail pharmacy Tab 100 mg50.06

200.24 300.40

✓ Tegretol

100

100

250 ml

50

10 ml OP

56

100

200 ml

100

100

100

14

14

56

14

56

56

✓ Tearetol AU ✓ Tegretol CR

✓ Tegretol CR ✓ Tegretol

✓ Tegretol CR ✓ Tegretol

✓ Frisium

✓ Rivotril

✓ Essential

Ethosuximide \$29

✓ Zarontin ✓ Zarontin

✓ Nupentin

Nupentin Nupentin

✓ Vimpat ✓ Vimpat

✓ Vimpat

✓ Vimpat ✓ Vimpat

✓ Vimpat

129

Subsidy		Fully	Brand or
(Manufacturer's Price)	;	Subsidised	Generic
\$	Per	/	Manufacturer

⇒SA2267 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has focal epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

LA	MOTRIGINE				
\blacktriangle	Tab dispersible 2 mg	55.00	30	1	Lamictal
\blacktriangle	Tab dispersible 5 mg	50.00	30	1	Lamictal
*	Tab dispersible 25 mg	4.20	56	1	Logem
*	Tab dispersible 50 mg	5.11	56	/	Logem
*	Tab dispersible 100 mg	6.75	56	1	Logem
LE	VETIRACETAM				
	Tab 250 mg	5.84	60	/	Everet
	Tab 500 mg		60	1	Everet
	Tab 750 mg	16.71	60	/	Everet
	Tab 1,000 mg	21.82	60	1	Everet
	Oral liq 100 mg per ml	44.78	300 ml OP	1	Levetiracetam-AFT
	Inj 100 mg per ml, 5 ml vial	38.95	10	/	Levetiracetam-AFT
PH	IENOBARBITONE				
•	For phenobarbitone oral liquid refer Standard Formulae, page	ie 273			
	Tab 15 mg	•	500	1	Noumed
					Phenobarbitone
	Tab 30 mg	398.50	500	/	Noumed
	3				Phenobarbitone
PΗ	IENYTOIN SODIUM				
	Tab 50 mg	75.00	200	/	Dilantin Infatab
-,-	Cap 30 mg		200		Dilantin
	Cap 100 mg		200		Dilantin
*	Oral lig 30 mg per 5 ml		500 ml	/	Dilantin Paediatric
	REGABALIN				
ГГ	Note: Not subsidised in combination with subsidised gabap	ontin			
*	Cap 25 mg		56	1	Pregabalin Pfizer
~	σαρ 25 mg	7.80	30		Milpharm S29
*	Cap 75 mg		56		Pregabalin Pfizer
~	Οάρ 73 mg	8.10	30		Milpharm S29
*	Con 150 mg		56		Lyrica
不	Cap 150 mg	4.01	30		Pregabalin Pfizer
*	Cap 300 mg	7 39	56		Pregabalin Pfizer
W.	, ,	7.30	30	٠	ricyaballi riizei
	IIMIDONE	07.05	400	,	Dulandalana Ollaraat
*	Tab 250 mg	37.35	100	•	Primidone Clinect

,	Subsidy	:aa\ C	Fully	Brand or
(Manufacturer's Pr \$	Per	Subsidised	Generic Manufacturer
ODIUM VALPROATE				
Tab 100 mg	13.65	100	√ E	pilim Crushable
Tab 200 mg EC	27.44	100	✓ E	pilim
Tab 500 mg EC		100	√ E	pilim
★ Oral lig 200 mg per 5 ml		300 ml	√ E	pilim S/F Liquid
				pilim Syrup
₭ Inj 100 mg per ml, 4 ml	41.50	1		pilim IV
STIRIPENTOL - Special Authority see SA2268 below - Retail pha				•
Cap 250 mg	,	60	√ Γ	Diacomit
Powder for oral liq 250 mg sachet		60		Diacomit
Fowder for oral liq 250 flig Sacriet	509.29	00	• 6	naconni

⇒SA2268 Special Authority for Subsidy

TODIRAMATE

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

OPIRAMATE			_
▲ Tab 25 mg	11.07	60	Arrow-Topiramate
			✓ Topiramate Actavis
	26.04		✓ Topamax
▲ Tab 50 mg	18.81	60	Arrow-Topiramate
•			✓ Topiramate Actavis
	44.26		✓ Topamax
▲ Tab 100 mg	31.99	60	Arrow-Topiramate
•			✓ Topiramate Actavis
	75.25		✓ Topamax
▲ Tab 200 mg	55.19	60	Arrow-Topiramate
•			✓ Topiramate Actavis
	129.85		✓ Topamax
Sprinkle cap 15 mg	20.84	60	✓ Topamax
Sprinkle cap 25 mg	26.04	60	✓ Topamax
/IGABATRIN - Special Authority see SA2088 below -	Retail pharmacy		
▲ Tab 500 mg		100	✓ Sabril
Powder for oral soln 500 mg per sachet		60	✓ Sabril

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms: or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy: and
 - 1.2.2 Either:



continued...

- 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 115

Acute Migraine Treatment

Tab orodispersible 10 mg4	1.84	30	Rizamelt
SUMATRIPTAN			
Tab 50 mg14	.41	90	Sumagran
Tab 100 mg22	2.68	90	Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per			-
prescription29	9.80 2	OP ✓	Clustran

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 47

PIZOTIFFN

★ Tab 500 mcg......23.21 100 **✓ Sandomigran**

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT - Special Authority see SA0987 below - Retail pharmacy

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CYCLIZINE HYDROCHLORIDE				
Tab 50 mg	0.66	10	✓ <u>N</u>	<u>Nausicalm</u>
CYCLIZINE LACTATE				
Inj 50 mg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	16.36	10	✓ <u>F</u>	lameln
DOMPERIDONE				
* Tab 10 mg	4.00	100	✓ [<u>Domperidone</u> <u>Viatris</u>
HYOSCINE HYDROBROMIDE				
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	✓ N	Martindale S29
Patch 1 mg per 72 hours – Special Authority see SA1998				
below – Retail pharmacy	88.50	10	✓ 9	Scopolamine - Mylan

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

1.57	100	✓ <u>Metoclopramide</u> <u>Actavis 10</u>
7.00	10	✓ Baxter
2.27	50	✓ Periset
0.56	10	✓ Periset ODT
4.10	50	✓ Periset
0.90	10	✓ Periset ODT
5.97	50	
(30.00)		Buccastem
(30.00)		Max Health S29
(30.00)		Prochlorperazine
		Brown & Burk S29
(30.00)		Prochlorperazine Max Health
25.00	250	✓ Nausafix
	10	✓ Stemetil
	(30.00)	7.00 102.27 500.56 104.10 500.90 105.97 50 (30.00) (30.00) (30.00) (30.00)

(Prochlorperazine Brown & Burk 329 Tab 3 mg buccal to be delisted 1 July 2025)

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine dis			
Tab 100 mg	5.84	30	✓ Sulprix
Tab 200 mg	14.47	60	✓ Sulprix
Tab 400 mg	35.06	60	✓ Sulprix
ARIPIPRAZOLE - Safety medicine; prescriber may determine di	snensina frequer	ncv	
Tab 5 mg		30	✓ Aripiprazole Sandoz
1 ab 5 mg	10.50	00	✓ Ampiprazole dandoz
			Aripiprazole S29
Tab 10 mg	10.50	30	✓ Aripiprazole Sandoz
ŭ			
Tab 15 mg		30	✓ Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz
Tab 30 mg	10.50	30	Aripiprazole Sandoz
(Ascend Aripiprazole S29 Tab 5 mg to be delisted 1 July 2025)			
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; pre	scriber may dete	ermine dispen	sing frequency
Tab 25 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO		10	✓ Largactil
		10	Largaotti
CLOZAPINE – Hospital pharmacy [HP4]			
Safety medicine; prescriber may determine dispensing frequency	•		
Tab 25 mg	6.69	50	✓ Clopine
			✓ Clozaril
	13.37	100	Clopine
			Clozaril
Tab 50 mg	8.67	50	Clopine
	17.33	100	✓ Clopine
Tab 100 mg	17.33	50	✓ Clopine
-			✓ Clozaril
	34.65	100	✓ Clopine
			✓ Clozaril
Tab 200 mg	34.65	50	✓ Clopine
ŭ	69.30	100	✓ Clopine
Suspension 50 mg per ml	147.30	100 ml	✓ Versacloz
HALOPERIDOL – Safety medicine; prescriber may determine dis		1011	
Tab 500 mcg — Up to 30 tab available on a PSO		100	✓ Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO		100	✓ Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50	✓ Serenace
0 11 0 11 1 200 1 11 11 200	29.72	100	✓ Serenace
Oral liq 2 mg per ml — Up to 200 ml available on a PSO		100 ml	✓ Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	O21.55	10	✓ Serenace
LEVOMEPROMAZINE - Safety medicine; prescriber may detern	nine dispensing f	requency	
Tab 25 mg (33.8 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 25 mg as a maleate		100	✓ Nozinan `
Tab 100 mg (135 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 100 mg as a maleate		100	✓ Nozinan
		100	· · · · · · · · · · · · · · · · · · · ·

	Subsidy			and or
	(Manufacturer's Price) \$	Sub Per		eneric anufacturer
	nrescriber may deterr	nine disne		
Inj 25 mg per ml, 1 ml ampoule		10	• .	nan S29 S29
ing 20 mg per mi, 1 mi ampodie	27.70	10	✓ Wool	
(Nozinan S29 S29 Inj 25 mg per ml, 1 ml ampoule to be deliste	ed 1 July 2025)			
LITHIUM CARBONATE - Safety medicine; prescriber may dete	ermine dispensina frec	uencv		
Tab long-acting 400 mg		100	✓ Priac	lel
Cap 250 mg		100	✓ Doug	
OLANZAPINE - Safety medicine; prescriber may determine dis	spensina frequency			
Tab 2.5 mg		30	✓ Zypir	ne
Tab 5 mg		30	✓ Zypiı	
Tab orodispersible 5 mg		28	✓ Zypiı	
Tab 10 mg		30	✓ Zypiı	
Tab orodispersible 10 mg		28	✓ Zypiı	
PERICYAZINE - Safety medicine; prescriber may determine d				
Tab 2.5 mg		100	✓ Neula	actil
Tab 10 mg		100	✓ Neula	
QUETIAPINE – Safety medicine; prescriber may determine dis				
Tab 25 mg		90	✓ Quet	anel
Tab 100 mg		90	✓ Quet	
Tab 200 mg		90	✓ Quet	
Tab 300 mg		90	✓ Quet	
· ·		00	<u>uuot</u>	<u>upo:</u>
RISPERIDONE – Safety medicine; prescriber may determine of		20	√ Dion	ordol
Tab 0.5 mg	2.17	20 60	✓ Risp	eridone (Teva)
	4.01	60	✓ Risp	
	4.01			ndoz S29
Tab 4 man	0.44	00		
Tab 1 mg	2.44	60	✓ Risp	
	0.00			eridone (Teva)
	3.68		✓ Risp	
				ndoz S29
Tab 2 mg	2.72	60	✓ Risp	
	5.00		. —	eridone (Teva)
	5.38		✓ Risp	
				ndoz S29
Tab 3 mg	4.50	60	✓ Risp	
				eridone (Teva)
	8.57		✓ Risp	
				ndoz S29
Tab 4 mg	6.25	60	✓ Risp	
	40.00	00 1		eridone (Teva)
Oral liq 1 mg per ml		30 ml	✓ Risp	<u>eron</u>
ZIPRASIDONE – Safety medicine; prescriber may determine d			-	
Cap 20 mg		60	✓ Zusd	
Cap 40 mg		60	✓ Zusd	
Cap 60 mg		60	✓ Zusd	
Cap 80 mg	46.55	60	✓ Zusd	one
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pr	escriber may determin	ne dispens		
Tab 10 mg	31.45	100	✓ Clop	ixol



Subsidy		Fully	Brand or	_
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

Depot Injections

ARIPIPRAZOLE – Special Authority see SA2395 below – Safety medicine; prescriber may determine dispensing			
Inj 300 mg vial	273.56	1	Abilify Maintena
, ,			✓ Abilify Maintena S29 S29
Inj 400 mg vial	341.96	1	Abilify Maintena
, ,			Abilify Maintena
			S29 S29

⇒SA2395 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Either:
 - 1.1 The patient has had an initial Special Authority approval for risperidone depot injection, paliperidone depot injection or olanzapine depot injection; or
 - 1.2 All of the following:
 - 1.2.1 The patient has schizophrenia or other psychotic disorder; and
 - 1.2.2 The patient has received treatment with oral atypical antipsychotic agents but has been unable to adhere; and
 - 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or
- 2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been started on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection.

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
 - The patient has schizophrenia; and
 - The patient has not been able to adhere with treatment using oral atypical antipsychotic agents; and
 - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

spensing freq	uency
5	✓ Fluanxol
5	✓ Fluanxol
5	✓ Fluanxol
pensing frequ	ency
5	✓ Haldol
5	 Haldol Concentrate
	✓ Haldol
	Decanoas S29
асу	
1	✓ Zyprexa Relprevv
1	✓ Zyprexa Relprevv
1	✓ Zyprexa Relprevv
	5 5 pensing frequ 5

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2313 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA2396 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg syringe	25 1	✓ Invega Sustenna
Inj 50 mg syringe271.	95 1	✓ Invega Sustenna
Inj 75 mg syringe	42 1	✓ Invega Sustenna
Inj 100 mg syringe		✓ Invega Sustenna
Inj 150 mg syringe435.	12 1	✓ Invega Sustenna

⇒SA2396 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has been unable to adhere to treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE - Special Authority see SA2167 below - Retail pharmacy

Inj 175 mg syringe	815.85	1	Invega Trinza
Inj 263 mg syringe	1,072.26	1	✓ Invega Trinza
Inj 350 mg syringe	1,305.36	1	✓ Invega Trinza
Inj 525 mg syringe	1,305.36	1	✓ Invega Trinza

⇒SA2167 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA2397 below - Retail pharmacy

Safety medicine: prescriber may determine dispensing frequency

Inj 25 mg vial	98 1	1	Risperdal Consta
Inj 37.5 mg vial178.7	71 1	✓	Risperdal Consta
Inj 50 mg vial217.5	56 1	✓	Risperdal Consta

⇒SA2397 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or



Subsidy (Manufacturer's Price)	9	Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has not been able to adhere with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

✓ Clopixol

Anxiolytics

BUSPIRONE HYDROCHI ORIDE

* Tab 5 mg	13.95	100	 Buspirone Viatris
* Tab 10 mg	12.50	100	✓ Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may determin	e dispensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine di	spensing frequency		
Tab 2 mg	95.00	500	Arrow-Diazepam

LORAZEPAM - Safety medicine: prescriber may determine dispensing frequency

Ativan 250 100

Ativan

500

Arrow-Diazepam

Multiple Sclerosis Treatments

⇒SA2274 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months: and
 - 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a


		NEC	RVOUS SYSTEM
		INEF	TVOUS STSTEM
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per 🗸	Brand or Generic Manufacturer
continued			
previous attack (where relevant); and			
1.4.4 Each significant attack can be distinguished	ed from the effects of ge	eneral fatigue; a	nd is not associated with a
fever (T> 37.5°C); and 1.4.5 Either:			
1.4.5.1 Each significant attack is severe en	ough to change either t	he FDSS or at l	east one of the Kurtze
Functional System scores by at least		inc EDOO of at h	cast one of the Number
1.4.5.2 Each significant attack is a recurren		of multiple sclei	rosis (tonic
seizures/spasms, trigeminal neuralg	jia, Lhermitte's symptor	m); and [.]	,
1.5 Evidence of new inflammatory activity on an MRI	scan within the past 24	months; and	
1.6 Any of the following:			
1.6.1 A sign of that new inflammatory activity or	MRI scanning (in crite	rion 5 immediate	ely above) is a gadolinium
enhancing lesion; or 1.6.2 A sign of that new inflammatory activity is	a lecion chowing diffue	ion restriction: o	r
1.6.3 A sign of that new inflammatory is a T2 les	•		I
1.6.4 A sign of that new inflammatory activity is			sponsible for the clinical
features of a recent attack that occurred w	, ,		
1.6.5 A sign of that new inflammatory activity is			ous MRI scan; or
2 Patient has an active approval for ocrelizumab and does			
Note: Treatment on two or more funded multiple sclerosis treat Renewal — (Multiple Sclerosis - dimethyl fumarate, fingolin			ta-1-alpha interferen
beta-1-beta, natalizumab and teriflunomide) from any releva			
had an EDSS score of 0 to 6.0 (inclusive) with or without the use			
the patient has walked 100 metres or more with or without aids			
Note: Treatment on two or more funded multiple sclerosis treat	•	•	
DIMETHYL FUMARATE – Special Authority see SA2274 on the	e previous page – Reta	il pharmacy	
a) Wastage claimable	de transferante aboutton		
b) Note: Treatment on two or more funded multiple scleros			mittea. Fecfidera
Cap 240 mg			Tecfidera Tecfidera
FINGOLIMOD – Special Authority see SA2274 on the previous	•	·V	
a) Wastage claimable	pago motan phamia	,	
b) Note: Treatment on two or more funded multiple scleros	sis treatments simultane	eously is not per	mitted.
Cap 0.5 mg	2,200.00	28	Gilenya
GLATIRAMER ACETATE - Special Authority see SA2274 on the	ne previous page – Ret	ail pharmacy	
Note: Treatment on two or more funded multiple sclerosis t			
Inj 40 mg prefilled syringe		_	Copaxone
INTERFERON BETA-1-ALPHA – Special Authority see SA227			
Note: Treatment on two or more funded multiple sclerosis to Inj 6 million iu prefilled syringe			ted. Avonex
Injection 6 million iu per 0.5 ml pen injector			Avonex Pen
(Avonex Pen Injection 6 million iu per 0.5 ml pen injector to be o			

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

INTERFERON BETA-1-BETA – Special Authority see SA2274 on the previous page – Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

✓ Betaferon

✓ Tysabri



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

TERIFLUNOMIDE - Special Authority see SA2274 on page 138 - Retail pharmacy

a) Wastage claimable

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

✓ Teriflunomide Sandoz

659.90

✓ Aubagio

Teriflunomide Sandoz to be Principal Supply on 1 April 2025 (Aubagio Tab 14 mg to be delisted 1 April 2025)

Multiple Sclerosis Treatments - Other

OCRELIZUMAB - Special Authority see SA2273 below - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

✓ Ocrevus

⇒SA2273 Special Authority for Subsidy

Initial application — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months: and
 - 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic): and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 1.4.5 Fither:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
 - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
 - 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon

NERVOUS SYSTEM

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple Sclerosis - ocrelizumab) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initial application — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and
- 3 Patient has no history of relapsing remitting multiple sclerosis.

Renewal — (Primary Progressive Multiple Sclerosis) from any relevant practitioner. Approvals valid for 12 months where patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
MIDAZOLAM – Safety medicine; prescriber may determine dispe Inj 1 mg per ml, 5 ml ampoule		10		Midazolam-Baxter Midazolam Viatris
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available on a PSO		10		Pfizer
On a PSO for status epilepticus use only. PSO must be only in 5 mg per ml, 1 ml plastic ampoule — Up to 10 inj available on a PSO		10		Midazolam-Pfizer
On a PSO for status epilepticus use only. PSO must be only in 5 mg per ml, 3 ml ampoule		epilep 5	✓	only. Midazolam-Baxter Midazolam Viatris
Inj 5 mg per ml, 3 ml plastic ampoule — Up to 5 inj available of a PSO	22.50	5		Pfizer
On a PSO for status epilepticus use only. PSO must be a (Midazolam Viatris Inj 1 mg per ml, 5 ml ampoule to be delisted 1 (Midazolam Viatris Inj 5 mg per ml, 3 ml ampoule to be delisted 1	May 2025)	epiiep	oticus use	only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 b Inj 200 mg per ml, 1 ml ampoule		acy 10	•	Max Health \$29
▶ SA1386 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Both:	without further rene	wal u	nless notif	ied for applications meeting
1 For the treatment of terminal agitation that is unresponsive2 The applicant is part of a multidisciplinary team working in	•	i		
TEMAZEPAM – Safety medicine; prescriber may determine dispertable 10 mg		25	•	Normison
ZOPICLONE – Safety medicine; prescriber may determine disper Tab 7.5 mg		500	•	Zopiclone Actavis
Spinal Muscular Atrophy				

NUSINERSEN − PCT only − Special Authority see SA2174 below
Inj 12 mg per 5 ml vial120,000.00 1 ✓ Spinraza

⇒SA2174 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

RISDIPLAM - [Xpharm] - Special Authority see SA2203 below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

Powder for oral soln 750 mcg per ml, 60 mg per bottle......14,100.00 80 ml OP ✓ Evrysdi

⇒SA2203 Special Authority for Subsidy

Initial application — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Renewal — (spinal muscular atrophy (SMA)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day) in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

Stimulants/ADHD Treatments

ATOMOXETINE			
Cap 10 mg	43.02	28	✓ APO-Atomoxetine
Cap 18 mg	45.57	28	✓ APO-Atomoxetine
Cap 25 mg	44.30	28	✓ APO-Atomoxetine
Cap 40 mg	46.21	28	✓ APO-Atomoxetine
Cap 60 mg	51.31	28	✓ APO-Atomoxetine
Cap 80 mg	65.20	28	✓ APO-Atomoxetine
Cap 100 mg	65.71	28	✓ APO-Atomoxetine
DEXAMFETAMINE SULFATE – Special Authority see SA2410 (a) Only on a controlled drug form b) Safety medicine; prescriber may determine dispensing fr		- Retail phar	macy
Tab 5 mg	29.80	100	✓ <u>Noumed</u> Dexamfetamine



Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
\$	Per	✓	Manufacturer

⇒SA2410 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

LISDEXAMFETAMINE DIMESILATE - Special Authority see SA2415 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Cap 30 mg - No more than 1 cap per day	30	✓ Vyvanse
Cap 50 mg60.00	30	✓ Vyvanse
Cap 70 mg60.00	30	✓ Vyvanse

⇒SA2415 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
 - 2.2 Diagnosed according to DSM-V or ICD 11 criteria; and
 - 2.3 Either:
 - 2.3.1 Applicant is a paediatrician or psychiatrist; or
 - 2.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
 - 2.4 Any of the following:
 - 2.4.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) and has not received sufficient benefit or has experienced intolerable side effects; or
 - 2.4.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
 - 2.4.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate; or
 - 2.4.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment

NERVOUS SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

adherence difficulties: or

- 2.4.5 There is significant concern regarding the risk of diversion or abuse of immediate release methylphenidate hydrochloride; or
- 2.4.6 Both:
 - 2.4.6.1 Patient would have been prescribed a subsidised formulation of methylphenidate hydrochloride (extended-release) but has been unable to access due to supply issues with methylphenidate hydrochloride (extended-release); and
 - 2.4.6.2 Other alternative stimulant presentations (methylphenidate or dexamfetamine) are not appropriate; and
- 2.5 Lisdexamfetamine dimesilate is not to be used in combination with another funded methylphenidate presentation.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA2411 below - Retail pharmacy

- a) Only on a controlled drug form
- h) Safety medicine: prescriber may determine dispensing frequency

b) Calcty medicine, prescriber may actermine dispens	oning inequently		
Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg		30	✓ Rubifen
•	4.00		✓ Ritalin
Tab extended-release 18 mg	7.75	30	✓ Methylphenidate ER - Teva
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg		30	✓ Rubifen SR
Tab extended-release 27 mg		30	Methylphenidate ERTeva
Tab extended-release 36 mg	15.50	30	Methylphenidate ERTeva
Tab extended-release 54 mg	22.25	30	✓ Methylphenidate ER - Teva

⇒SA2411 Special Authority for Subsidy

Initial application — (ADHD in patients aged 5 years or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria: All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients aged under 5 years) only from a paediatrician or psychiatrist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
 - 1 ADHD (Attention Deficit and Hyperactivity Disorder) in patients under 5 years of age; and
 - 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.



Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
 \$	Per	✓	Manufacturer	

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA2446 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

b) Safety medicine, prescriber may determine dispensing	rrequericy		
Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	Concerta
Tab extended-release 36 mg	71.93	30	Concerta
Tab extended-release 54 mg	86.24	30	Concerta
Cap modified-release 10 mg	19.41	30	Ritalin LA
Cap modified-release 20 mg	27.72	30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA2446 Special Authority for Subsidy

Initial application — (ADHD) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 ADHD (Attention Deficit and Hyperactivity Disorder); and
 - 1.2 Diagnosed according to DSM-IV or ICD 10 criteria; and
 - 1.3 Either:
 - 1.3.1 Applicant is a paediatrician or psychiatrist; or
 - 1.3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
 - 1.4 Either:
 - 1.4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or difficulties with adherence; or
 - 1.4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride; or
- 2 Both:
 - 2.1 Patient meets the Special Authority criteria for SA2411 methylphenidate hydrochloride; and
 - 2.2 Patient is unable to access other methylphenidate hydrochloride presentations under Special Authority criteria SA2411 due to an out of stock (see note).

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock on tab extended-release Methylphenidate ER – Teva and tab sustained-release 20 mg Rubifen SR subsidised under

SA2411 (https://schedule.pharmac.govt.nz/2025/02/01/SA2411.pdf).

MODAFINIL – Special Authority see SA2413 below – Retail pharmacy
Tab 100 mg

Modafinil Max Health to be Principal Supply on 1 May 2025

(Modavigil Tab 100 mg to be delisted 1 May 2025)

⇒SA2413 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	•	Manufacturer

continued...

almost daily for three months or more; and

- 2 Fither
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eve movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	3.70	84	✓ Ipca-Donepezil
* Tab 10 mg	5.50	84	✓ Ipca-Donepezil
RIVASTIGMINE - Special Authority see SA1488 below - Re	etail pharmacy		
Patch 4.6 mg per 24 hour	49.40	30	✓ Rivastigmine Patch
			BNM 5
	90.00		✓ Exelon Patch 5
Patch 9.5 mg per 24 hour	49.40	30	✓ Rivastigmine Patch
			BNM 10
	90.00		✓ Exelon Patch 10

(Exelon Patch 5 Patch 4.6 mg per 24 hour to be delisted 1 June 2025) (Exelon Patch 10 Patch 9.5 mg per 24 hour to be delisted 1 June 2025)

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate: and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg11.76 28


Tab sublingual 8 mg with naloxone 2 mg34.00 28 ✓

✓ Buprenorphine Naloxone BNM

✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:



	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health..

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (**Detoxification**) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE Tab modified-release 150 mg	15.00	30	✓ <u>Zyban</u>
DISULFIRAM			
Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see	SA1408 on the next p	age – Retai	pharmacy
Tab 50 mg	77.77	28	✓ Naltrexone AOP S29
	83.33	30	✓ Naltraccord
	102.60		✓ Naltrexone Max Health (\$29)
	138.88	50	✓ Revia S29
(Revia S29 Tab 50 mg to be delisted 1 July 2025)			

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

 b) Note: Direct Provision by a pharmacist permitted under the prov 	risions in Part I	of Section A	
Patch 7 mg - Up to 28 patch available on a PSO	19.62	28	✓ Habitrol
Patch 14 mg - Up to 28 patch available on a PSO		28	✓ Habitrol
Patch 14 mg for direct distribution only - [Xpharm]	12.49	7	✓ Habitrol
Patch 21 mg - Up to 28 patch available on a PSO	24.72	28	✓ Habitrol
Patch 21 mg for direct distribution only - [Xpharm]	13.19	7	✓ Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO	22.53	216	✓ Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	12.89	36	✓ Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO	24.68	216	✓ Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	13.25	36	✓ Habitrol
Gum 2 mg (Fruit) - Up to 204 piece available on a PSO	23.02	204	✓ Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]	17.57	96	✓ Habitrol
Gum 2 mg (Mint) - Up to 204 piece available on a PSO	23.02	204	✓ Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]	17.57	96	✓ Habitrol
Gum 4 mg (Fruit) - Up to 204 piece available on a PSO	25.98	204	✓ Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]	23.87	96	✓ Habitrol
Gum 4 mg (Mint) - Up to 204 piece available on a PSO	25.98	204	✓ Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]	23.87	96	✓ Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg	17.62	56	✓ Varenicline Pfizer

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme,

NERVOUS SYSTEM

Subsidy (Manufacturer's Price)	Fu Subsidis	ılly	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

which includes prescriber or nurse monitoring; and

- 3 Fither
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

	Subsidy		Fully	Brand or
(M	anufacturer's Price)	5	Subsidised	Generic
	\$	Per	/	Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Spe	cialist - Special Authority	see SA2398	3 below
Inj 25 mg vial	50.05	1	Bendamustine Sandoz
	77.00		✓ Ribomustin
Inj 100 mg vial	200.20	1	✓ Bendamustine Sandoz
	308.00		✓ Ribomustin
Inj 1 mg for ECP	3.23	1 mg	✓ Baxter

⇒SA2398 Special Authority for Subsidy

Initial application — (CLL*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 The patient has chronic lymphocytic leukaemia requiring treatment; and
 - 2 Patient has ECOG performance status of 0-2; and
 - 3 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a * includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 The patient has ECOG performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m² twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSINE FAN - PCT - Botail pharmacy-Specialist

Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN – PCT only – Specialist			,
Inj 10 mg per ml, 45 ml vial	25.73	1	✓ Carboplatin Accord ✓ DBL Carboplatin S29 S29
	32.59		✓ DBL Carboplatin
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.06	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist		-	
Inj 100 mg vial	710.00	1	✓ BiCNU
, ,			✓ BiCNU S29 S29
			✓ Novadoz S29
Inj 100 mg for ECP	710.00	100 mg OP	✓ Baxter
(BiCNU S29 S29 Inj 100 mg vial to be delisted 1 July 2025)		3 ·	
(Novadoz S29 Inj 100 mg vial to be delisted 1 July 2025)			
, , ,			
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist Tab 2 mg	20.06	25	✓ Leukeran FC
-	29.00	23	Leukeraniio
CISPLATIN – PCT only – Specialist	0.45		40: 1:: 4
Inj 1 mg per ml, 50 ml vial		1	✓ Cisplatin Accord
	15.00		Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	 Cisplatin Accord
	21.00		Cisplatin Ebewe
	29.66		DBL Cisplatin
Inj 1 mg for ECP	0.19	1 mg	✓ Baxter

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
DVOLODUO ODLIA AUDE	Ψ	1 61		Manuacturer
CYCLOPHOSPHAMIDE Tab 50 mg - PCT - Retail pharmacy-Specialist	145.00	50	./	Cyclonex
Inj 1 g vial – PCT – Retail pharmacy-Specialist		1		<u>Cyclonex</u> Endoxan
iiij i g viai – PO i – netaii pilaimacy-specialist	127.80	6		<u>Cytoxan</u>
Inj 2 g vial - PCT only - Specialist		1		Endoxan
Inj 1 mg for ECP - PCT only - Specialist		1 mg		Baxter
FOSFAMIDE - PCT only - Specialist		9		
	96.00	1	1	Holoxan
Inj 2 g		1		Holoxan
Inj 1 mg for ECP		1 mg		Baxter
OMUSTINE - PCT - Retail pharmacy-Specialist		9		
Cap 40 mg	880.00	20	/	Medac S29
ΛΕΓΡΗΥΓΕΙ ΤΕΙΡΕΙΡΙΑΙ ΤΟ ΤΕ		_0		
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist		1		Megval S29
IIIJ 30 IIIg = 1 01 OIIIy = Specialist	40.23	'		Melpha
	67.80			Alkeran
Megval S29 Inj 50 mg to be delisted 1 July 2025)	0.100			
, ,				
DXALIPLATIN – PCT only – Specialist	05.01	1	./	Ovaliniatin Astavia
Inj 100 mg vial	25.01	'	•	Oxaliplatin Actavis 100
	110.00			Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	33.35	1		Alchemy Oxaliplatin
	46.32			Oxaliplatin Accord
Inj 1 mg for ECP	0.35	1 mg	/	Baxter
THIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	✓	Bedford S29
			✓	Max Health S29
			1	THIO-TEPA \$29
	398.00			Tepadina
Inj 100 mg vial		1		Max Health S29
-, · · · · · · · · · · · · · · · · · · ·	1,800.00	-		Tepadina
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see	SA2141 below			
Inj 100 mg vial		1	✓	Azacitidine Dr Reddy's
				ricauy 3

⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder);

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

continued...

Ω

- 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

- · · · · · · · · · · · · · · · · · · ·		
CALCIUM FOLINATE Tab 15 mg - PCT - Retail pharmacy-Specialist135.33	10	✓ DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist17.10	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist7.28	1	✓ Calcium Folinate Sandoz
		✓ Calcium Folinate Sandoz S29 S29
36.48	5	✓ Eurofolic S29
Inj 50 mg - PCT - Retail pharmacy-Specialist72.80	10	✓ Leucovorin Pharmacia S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist9.49	1	✓ Calcium Folinate Sandoz
47.45	5	✓ Eurofolic S29
Inj 100 mg – PCT only – Specialist7.33	1	✓ Calcium Folinate Ebewe
94.90	10	✓ Leucovorin Pharmacia §29
Inj 300 mg - PCT only - Specialist21.55	1	✓ Leucovorin DBL S29
22.51		✓ Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist25.14	1	Calcium Folinate Sandoz
		✓ Calcium Folinate Sandoz S29 S29
Inj 1 g - PCT only - Specialist67.51	1	Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist72.00	1	Calcium Folinate Sandoz
		✓ Eurofolic S29
Inj 1 mg for ECP - PCT only - Specialist0.06	1 mg	✓ Baxter
CAPECITABINE - Retail pharmacy-Specialist		
Tab 150 mg	60	✓ Capecitabine Viatris
Tab 500 mg46.50	120	✓ Capecitabine Viatris
CLADRIBINE - PCT only - Specialist		
Inj 1 mg per ml, 10 ml749.96	1	✓ Leustatin
Inj 10 mg for ECP749.96	10 mg OP	✓ Baxter
,		

	Subsidy		Fully	Brand or
	(Manufacturer's F		idised	Generic
	\$	Per		Manufacturer
CYTARABINE				
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specia	list472.00	5	✓ F	Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail				
pharmacy-Specialist	48.80	1	1	Cytarabine DBL
F				Pfizer
			√ F	Pfizer S29 S29
Inj 1 mg for ECP - PCT only - Specialist	0.29	10 mg	_	Baxter
Inj 100 mg intrathecal syringe for ECP – PCT only – Specia		100 mg OP		Baxter
FLUDARABINE PHOSPHATE			-	
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	./ [Fludara Oral
Inj 50 mg vial – PCT only – Specialist		20 1	_	Fludara Orai Fludarabine
III 50 IIIg viai – POT OIIIy – Specialist	120.00	1	• r	Sagent S29
	634.00	5	. / [Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist		50 mg OP	_	Baxter
	120.00	30 mg Oi	٠.	Jaktei
FLUOROURACIL	10.51			
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist		1	_	luorouracil Accord
Inj 50 mg per ml, 50 ml vial – PCT only – Specialist		1	_	luorouracil Accord
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1		Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist	0.41	100 mg	• [Baxter
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist				
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine).				
26.3 ml vial		1	_	OBL Gemcitabine
Inj 1 g		1	-	Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg	✓ E	Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	52.57	1	1	Accord
	71.44		✓ I	rinotecan Actavis
				100
	100.00		✓ I	rinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	✓ E	Baxter
MERCAPTOPURINE				
Tab 50 mg - PCT - Retail pharmacy-Specialist	25.90	25	✓ F	Puri-nethol
Oral suspension 20 mg per ml - Retail pharmacy-Specialist			_	
Special Authority see SA1725 below		100 ml OP	1	Allmercap
, , , , , , , , , , , , , , , , , , , ,				(aluprine S29)
			• •	

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

	Subsidy	,	Fully	
(Manufacturer's Price \$	e) Per	Subsidised •	
VETUOTDE VATE	Ψ	1 61		Manuacturer
IETHOTREXATE	7.00	00	./	Travete
Tab 2.5 mg — PCT — Retail pharmacy-Specialist		90		Trexate Trexate
Tab 10 mg - PCT - Retail pharmacy-Specialist		90 5		Trexate Methotrexate DBL
 Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist Inj 7.5 mg prefilled syringe 		1	_	Methotrexate
inj 7.5 mg premied synnge	29.17	'	•	Sandoz
: Inj 10 mg prefilled syringe	19.09	1	•	Methotrexate Sandoz
f Inj 15 mg prefilled syringe	24.53	1	•	Methotrexate Sandoz
f Inj 20 mg prefilled syringe	16.64	1	•	Methotrexate Sandoz
f Inj 25 mg prefilled syringe	20.72	1	✓	Methotrexate
€ Inj 30 mg prefilled syringe	55.00	1	1	Sandoz Methotrexate
				<u>Sandoz</u>
f Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist	t30.00	5	/	Methotrexate DBL Onco-Vial
Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Speciali	st45.00	1	•	DBL Methotrexate Onco-Vial
Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist.	25.00	1	1	Methotrexate Ebewe
Inj 100 mg per ml, 50 ml vial – PCT – Retail				
pharmacy-Specialist	67.99	1	1	Methotrexate Ebewe
Inj 1 mg for ECP - PCT only - Specialist		1 mg	1	Baxter
Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist		5 mg Öl	P 🗸	Baxter
EMETREXED - PCT only - Specialist		Ü		
Inj 100 mg vial	8.99	1	1	Pemetrexed-AFT
,	60.89	•		Juno Pemetrexed
Inj 500 mg vial		1		Pemetrexed-AFT
"I 000 TIG TO THE TOTAL TH	217.77			Juno Pemetrexed
Inj 1 mg for ECP		1 mg		Baxter
HIOGUANINE – PCT – Retail pharmacy-Specialist		g		Duxtor
Tab 40 mg	126.31	25	/	Lanvis
•	120.01	20	•	Lanvis
Other Cytotoxic Agents				
MSACRINE - PCT only - Specialist				
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	•	Amsidine S29
	4,736.00		✓	Amsidine S29
Inj 75 mg	1,250.00	5	1	AmsaLyo S29
· •	6,218.00		1	AmsaLyo S29
NAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Spec	•			•
Cap 0.5 mg		100	1	Agrylin
•	1,170.07	100	•	9.7
RSENIC TRIOXIDE – PCT only – Specialist	4 017 00	10		Dhanasan
Inj 1 mg per ml, 10 ml vial		10		Phenasen
Inj 10 mg for ECP	481.70	I0 mg C		Baxter
LEOMYCIN SULPHATE - PCT only - Specialist Inj 15,000 iu, vial	185.16	1	•	DBL Bleomycin
				Sulfate
Inj 1,000 iu for ECP	14.32	1,000 it	· •	Baxter

	Subsidy (Manufacturer's P	Prico) Sub	Fully	
	(Manulacturer's F	Per	JSIUISEU	Manufacturer
BORTEZOMIB - PCT only - Specialist - Special Authority see	SA2355 below			
Inj 3.5 mg vial		1	1	DBL Bortezomib
Inj 1 mg for ECP	22.26	1 mg	1	Baxter
⇒SA2355 Special Authority for Subsidy		•		
Initial application — (plasma cell dyscrasia) from any relevan	nt practitioner. Ap	pprovals valid	withou	it further renewal unless
notified where the patient has plasma cell dyscrasia, not includin	g Waldenström n	nacroglobulina	aemia,	requiring treatment.
DACARBAZINE - PCT only - Specialist				
Inj 200 mg vial	72.11	1	1	DBL Dacarbazine
Inj 200 mg for ECP		200 mg OP	1	Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist		-		
Inj 0.5 mg vial	255.00	1	1	Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP		Baxter
DAUNORUBICIN - PCT only - Specialist		J		
Inj 2 mg per ml, 10 ml	171.93	1	1	Pfizer
Inj 20 mg for ECP		20 mg OP		Baxter
DOCETAXEL - PCT only - Specialist				
Inj 20 mg	48 75	1	1	Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		i		DBL Docetaxel
Inj 20 mg per ml, 4 ml vial		1		Docetaxel
., ,		•		Accord \$29
Inj 80 mg	195.00	1	/	Docetaxel Sandoz
Inj 1 mg for ECP		1 mg		Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist		9		
Inj 2 mg per ml, 5 ml vial	10.00	1	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		Doxorubicin Ebewe
11) 2 11) por 111, 20 111 va	17.00			Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1		Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	1	Arrow-Doxorubicin
, •,	69.99		1	Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	1	Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml vial	25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial	30.00	1		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1		Epirubicin Ebewe
Inj 1 mg for ECP	0.50	1 mg		Baxter
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist		20		Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10		Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia		1		Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	•	Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist			_	
Inj 100 mg (of etoposide base)		. 1		Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg	•	Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pha	rmacy-Specialist			
Cap 500 mg	20.72	100	•	<u>Devatis</u>
IBRUTINIB - Special Authority see SA2168 on the next page -	Retail pharmacy			
Tab 140 mg	,	30		Imbruvica
Tab 420 mg	9,652.00	30	1	Imbruvica

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	0.1.	Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
\$	Per	1	Manufacturer

⇒SA2168 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Patient has not previously received funded ibrutinib; and
- 3 Ibrutinib is to be used as monotherapy; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 There is documentation confirming that patient has 17p deletion or TP53 mutation; and
 - 4.1.2 Patient has experienced intolerable side effects with venetoclax monotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient has received at least one prior immunochemotherapy for CLL; and
 - 4.2.2 Patient's CLL has relapsed within 36 months of previous treatment; and
 - 4.2.3 Patient has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or

109 74

4.3 Patient's CLL is refractory to or has relapsed within 36 months of a venetoclax regimen.

Renewal — (chronic lymphocytic leukaemia (CLL)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of clinical disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IDARUBICIN HYDROCHLORIDE

Ini 5 mg vial - PCT only - Specialist

ing strig viai – FCT only – Specialist	109.74		▼ Zaveuos
Inj 10 mg vial - PCT only - Specialist	233.64	1	✓ Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	✓ Baxter
LENALIDOMIDE (VIATRIS) - Special Authority se	ee SA2353 below - Retail pharm	nacy	
Cap 5 mg	76.92	21	✓ Lenalidomide
			<u>Viatris</u>
Cap 10 mg	50.30	21	✓ Lenalidomide
			<u>Viatris</u>
Cap 15 mg	62.13	21	✓ Lenalidomide
			<u>Viatris</u>
Cap 25 mg	65.09	21	✓ Lenalidomide
			Viatris

⇒SA2353 Special Authority for Subsidy

Initial application — (Plasma cell dyscrasia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient is not refractory to prior lenalidomide use.

Initial application — (Myelodysplastic syndrome) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5)

continued...

✓ Zavedos

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

associated with a deletion 5q cytogenetic abnormality; and

2 Patient has transfusion-dependent anaemia.

Renewal — (Myelodysplastic syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has not needed a transfusion in the last 4 months; and
- 2 No evidence of disease progression.

MFSNA

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist		50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule – PCT only – Specia		15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule – PCT only – Spec		15	✓ Uromitexan
Inj 1 mg for ECP – PCT only – Specialist		100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist		J	
Inj 5 mg vial	517 65	1	✓ Mitomycin
", O " (g Tul			(Fresenius Kabi) \$29
	526.00		✓ Mitomycin
	020.00		(Sagent) S29
	641.70		✓ Accord S29
Inj 20 mg vial	1,250.00	1	✓ Omegapharm S29
,	,		✓ Teva
Inj 1 mg for ECP	269.85	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP	5.51	1 mg	✓ Baxter
NIRAPARIB – Special Authority see SA2325 below – Ret Wastage claimable		· ·	
Tab 100 mg	13.393.50	84	✓ Zejula
Cap 100 mg		56	✓ Zejula
, ,	13,393.50	84	✓ Zejula

⇒SA2325 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has advanced high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line** of treatment with platinum-based chemotherapy; and
- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and
- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Fither
 - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen; or
 - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	,	Fully	Brand or	
(Manufacturer's Price	·) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 No evidence of progressive disease; and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Either:
 - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
 - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: * "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB – Retail pharmacy-Specialist – Special Author	ority see SA2163 below		
Tab 100 mg	3,701.00	56	Lynparza
Tab 150 mg	3,701.00	56	Lynparza

⇒SA2163 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Fither:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and

Subsidy	Full		
 (Manufacturer's Price) \$	Subsidise Per 🗸	d Generic Manufacturer	

continued...

- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
 - 5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 6 mg per ml, 16.7 ml vial		1	✓ Anzatax
, •	24.00		✓ Paclitaxel Ebewe
	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
, -	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 6 mg per ml, 50 ml vial	37.89	1	✓ Anzatax
, , ,	44.00		✓ Paclitaxel Ebewe
	275.00		✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.17	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see SA1979 b	elow		
Inj 750 iu per ml, 5 ml vial		1	Oncaspar LYO

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMII F).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

alist		
CBS	1	✓ Nipent S29
t page – Retail pharr	macy	
47.45	14	✓ Pomolide
71.18	21	✓ Pomolide
94.90	14	✓ Pomolide
142.35	21	✓ Pomolide
142.35	14	✓ Pomolide
213.53	21	✓ Pomolide
189.81	14	✓ Pomolide
284.71	21	✓ Pomolide

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic	
` \$ ´	Per	1	Manufacturer	

⇒SA2354 Special Authority for Subsidy

Initial application — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has relapsed or refractory plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient has not received prior funded pomalidomide.

Renewal — (Relapsed/refractory plasma cell dyscrasia) from any relevant practitioner. Approvals valid for 12 months where there is no evidence of disease progression.

t	
00 50	✓ Natulan S29
СУ	
13 5	✓ Temaccord
	✓ Temozolomide-
	Taro S29
38 5	✓ Temaccord
30	✓ Apo-Temozolomide
98 5	✓ Temaccord
20	✓ Apo-Temozolomide
12 5	✓ Temaccord
34 5	✓ Temaccord
	00 50 by 13 5 38 5 30 98 5 20 12 5

⇒SA2275 Special Authority for Subsidy

Initial application — (**gliomas**) only from a relevant specialist. Approvals valid for 12 months where the patient has a glioma. **Renewal** — (**gliomas**) only from a relevant specialist. Approvals valid for 12 months where treatment remains appropriate and patient is benefitting from treatment.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*: and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
THALIDOMIDE – Retail pharmacy-Specialist – Special Authority Cap 50 mg	378.00	28	-	'halomid
Cap 100 mg	756.00	28	√ 1	'halomid

⇒SA2356 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

TRETINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist	479.50	100	Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority se		ı	
Tab 14 \times 10 mg, 7 \times 50 mg, 21 \times 100 mg	1,771.86	42 OP	✓ Venclexta
Tab 10 mg		2 OP	✓ Venclexta
Tab 50 mg	239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable	8,209.41	120	✓ Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

(1	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	
VINBLASTINE SULPHATE				
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist		5	1	Hospira
Inj 1 mg for ECP - PCT only - Specialist	6.00	1 mg	✓	Baxter
VINCRISTINE SULPHATE				
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist	74.52	5	•	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist	102.73	5	•	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist	12.60	1 mg	✓	Baxter
VINORELBINE				
Cap 20 mg	30.00	1	✓	Vinorelbine Te Arai
Cap 30 mg		1	✓	Vinorelbine Te Arai
Cap 80 mg		1	✓	Vinorelbine Te Arai
Inj 10 mg per ml, 1 ml vial - PCT only - Specialist	42.00	1	✓	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial - PCT only - Specialist	168.00	1	✓	Navelbine S29 S29
	210.00			Vinorelbine Ebewe
Inj 1 mg for ECP - PCT only - Specialist	3.80	1 mg	✓	Baxter

Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA2385 below - Retail pharmacy

a) Brand switch fee payable (Pharmacode 2700441) - see page 271 for details

b) Wastage claimable

b) Wastage claimable			
Tab 20 mg	132.88	60	✓ Dasatinib-Teva
Tab 50 mg	304.13	60	✓ Dasatinib-Teva
Tab 70 mg	415.75	60	✓ Dasatinib-Teva

⇒SA2385 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or
- 3 Both:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Any of the following:
 - 3.2.1 Patient has documented treatment failure* with imatinib; or
 - 3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines.

······································			
ERLOTINIB - Retail pharmacy-Specialist - Special Auth	ority see SA2422 below		
Tab 100 mg	280.84	30	✓ Alchemy
Tab 150 mg	484.24	30	✓ Alchemy
3			

⇒SA2422 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR; and
- 3 Any of the following:
 - 3.1 Patient is treatment naive; or
 - 3.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 3.3 Both:
 - 3.3.1 The patient has discontinued osimertinib or gefitinib due to intolerance; and
 - 3.3.2 The cancer did not progress while on osimertinib or gefitinib.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB – Retail pharmacy-Specialist – Special Authority see SA2423 below
Tab 250 mg918.00 30 ✓ Iressa

⇒SA2423 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Any of the following:
 - 2.1 Patient is treatment naive: or
 - 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued osimertinib or erlotinib due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on osimertinib or erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR.

Renewal from any relevant practitioner. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

	IMAT	INIB N	1ESIL	ATE
--	------	--------	-------	-----

*	Cap 100 mg44.93	60	✓ <u>Imatinib-Rex</u>
*	Cap 400 mg69.76	30	✓ <u>Imatinib-Rex</u>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
LENVATINIB – Special Authority see SA2442 below – Retail pha Wastage claimable	armacy			
Cap 4 mg Cap 10 mg		30 30		envima envima

⇒SA2442 Special Authority for Subsidy

Initial application — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
 - 2.2 Either:
 - 2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
 - 2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures; and
 - 2.3 Any of the following:
 - 2.3.1 A lesion without iodine uptake in a RAI scan; or
 - 2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
 - 2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
 - 2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other; and
 - 2.4 Patient has thyroid stimulating hormone (TSH) adequately supressed; and
 - 2.5 Patient is not a candidate for radiotherapy with curative intent; and
 - 2.6 Surgery is clinically inappropriate; and
 - 2.7 Patient has an ECOG performance status of 0-2.

Renewal — (thyroid cancer) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable hepatocellular carcinoma; and
- 2 Patient has preserved liver function (Childs-Pugh A); and
- 3 Transarterial chemoembolisation (TACE) is unsuitable; and
- 4 Patient has an ECOG performance status of 0-2: and
- 5 Either:
 - 5.1 Patient has not received prior systemic therapy for their disease in the palliative setting; or
 - 5.2 Both:
 - 5.2.1 Patient has experienced treatment-limiting toxicity from treatment with atezolizumab with bevacizumab; and 5.2.2 No disease progression since initiation of atezolizumab with bevacizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma; and
 - 1.2 The disease is of predominant clear-cell histology; and
 - 1.3 The patient has documented disease progression following one previous line of treatment; and

Subsidy	Ful	,	
(Manufacturer's Price)	Subsidise Per •		
	Per •	Manufacturer	

continued...

- 1.4 The patient has an ECOG performance status of 0-2; and
- 1.5 Lenvatinib is to be used in combination with everolimus; or
- 2 All of the following:
 - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
 - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
 - 2.3 Lenyatinib is to be used in combination with everolimus; and
 - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

MIDOSTAURIN − PCT only − Special Authority see SA2342 below

Cap 25 mg......10,981.00 56 ✓ Rydapt

⇒SA2342 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of acute myeloid leukaemia; and
- 2 Condition must be FMS tyrosine kinase 3 (FLT3) mutation positive; and
- 3 Patient must not have received a prior line of intensive chemotherapy for acute myeloid leukaemia; and
- 4 Patient is to receive standard intensive chemotherapy in combination with midostaurin only; and
- 5 Midostaurin to be funded for a maximum of 4 cycles.

NILOTINIB - Special Authority see SA2301 below - Retail pharmacy

⇒SA2301 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
- 2 Fither:
 - 2.1 Patient has documented CML treatment failure* with a tyrosine kinase inhibitor (TKI); or
 - 2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

OSIMERTINIB – Special Authority see SA2418 on the ne	ext page – Retail pharmacy	1	
Tab 40 mg	9,310.00	30	✓ Tagrisso
Tab 80 mg	9,310.00	30	✓ Tagrisso

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA2418 Special Authority for Subsidy

Initial application — (NSCLC – first line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
 - 2.2 Any of the following:
 - 2.2.1 Patient is treatment naïve; or
 - 2.2.2 Patient has received prior chemotherapy in the adjuvant setting and/or while awaiting EGFR results; or
 - 2.2.3 Both:
 - 2.2.3.1 The patient has discontinued gefitinib or erlotinib due to intolerance; and
 - 2.2.3.2 The cancer did not progress while on gefitinib or erlotinib; and
 - 2.3 There is documentation confirming that the cancer expresses activating mutations of EGFR; and
 - 2.4 Patient has an ECOG performance status 0-3; and
 - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – first line) from any relevant practitioner. Approvals valid for 6 months where response to or stable disease with treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Initial application — (NSCLC – second line) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with osimertinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
 - 2.2 Patient has an ECOG performance status 0-3; and
 - 2.3 The patient must have received previous treatment with erlotinib or gefitinib; and
 - 2.4 There is documentation confirming that the cancer expresses T790M mutation of EGFR following progression on or after erlotinib or gefitinib; and
 - 2.5 The treatment must be given as monotherapy; and
 - 2.6 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (NSCLC – second line) from any relevant practitioner. Approvals valid for 6 months where response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

PALBOCICLIB - Special Authority see SA2345 below - Retail pharmacy

Wastage claimable			
Tab 75 mg	4,000.00	21	✓ Ibrance
Tab 100 mg	4,000.00	21	✓ Ibrance
Tab 125 mg		21	✓ Ibrance

⇒SA2345 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
 - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
 - 1.3 Patient has an ECOG performance score of 0-2; and
 - 1.4 Fither:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 1.4.2 Both:
 - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
 - 1.4.2.2 Patient has not received prior systemic treatment for metastatic disease; and
- 1.5 Treatment must be used in combination with an endocrine partner; and
- 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has an active Special Authority approval for ribociclib; and
 - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to ribociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
 - 2.3 Treatment must be used in combination with an endocrine partner; and
 - 2.4 There is no evidence of progressive disease since initiation of ribociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of palbociclib.

PAZOPANIB - Special Authority se	ee SA2429 below – Retail pharmacy
----------------------------------	-----------------------------------

opanib Teva
rient
opanib Teva
rient

Pazopanib Teva to be Principal Supply on 1 May 2025

(Votrient Tab 200 mg to be delisted 1 May 2025)

(Votrient Tab 400 mg to be delisted 1 May 2025)

⇒SA2429 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
 - 1.2 Either:
 - 1.2.1 The patient is treatment naive: or
 - 1.2.2 The patient has only received prior cytokine treatment; and
 - 1.3 The patient has an ECOG performance score of 0-2; and The patient has intermediate or poor prognosis defined as:
 - 1.4 Any of the following:
 - 1.4.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 1.4.2 Haemoglobin level < lower limit of normal; or
 - 1.4.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 1.4.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 1.4.5 Karnofsky performance score of less than or equal to 70; or
 - 1.4.6 2 or more sites of organ metastasis; and
 - 1.5 Pazopanib to be used for a maximum of 3 months; or
- 2 All of the following:

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ P	Per 🗸	Manufacturer

continued...

- 2.1 The patient has metastatic renal cell carcinoma; and
- 2.2 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
- 2.3 The cancer did not progress whilst on sunitinib; and
- 2.4 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months where there is no evidence of disease progression.

RIBOCICLIB - Special Authority see SA2343 below - Retail pharmacy

Wastage	

✓ Kisqali	21	Tab 200 mg1,883.00
✓ Kisqali	42	3,767.00
✓ Kisqali	63	5.650.00

⇒SA2343 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
 - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
 - 1.3 Patient has an ECOG performance score of 0-2; and
 - 1.4 Any of the following:
 - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
 - 1.4.2 Both:
 - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
 - 1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; or
 - 1.4.3 Both:
 - 1.4.3.1 Patient commenced treatment with ribociclib in combination with an endocrine partner prior to 1 July 2024; and
 - 1.4.3.2 There is no evidence of progressive disease; and
 - 1.5 Treatment to be used in combination with an endocrine partner; and
 - 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has an active Special Authority approval for palbociclib; and
 - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
 - 2.3 Treatment must be used in combination with an endocrine partner; and
 - 2.4 There is no evidence of progressive disease since initiation of palbociclib.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of ribociclib.

RUXOLITINIB – Special Authority see SA1890 on the next page – Retail pharmacy Wastage claimable

Tab 5 mg	2,500.00	56	Jakavi
Tab 10mg		56	Jakavi
Tab 15 mg	·	56	Jakavi
Tab 20 mg	·	56	Jakavi

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Fither:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 22 Both
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

Cap 12.5 mg	208.38	28	 Sunitinib Pfizer
Cap 25 mg		28	✓ Sunitinib Pfizer
Cap 50 mg		28	 Sunitinib Pfizer

⇒SA2430 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval: or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
 - 3 The patient has an ECOG performance score of 0-2; and
 - 4 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib: or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal — (RCC) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months where there is no evidence of disease progression.

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease): or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 87

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

Tab 250 mg4,276.19 120 ✓ Zvtiga

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases: and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or

continued...

- 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE

Tab 50 mg	4.18	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Author	rity see SA1895 bel	OW	
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	✓	Manufacturer
OCTREOTIDE				
Inj 50 mcg per ml, 1 ml vial	27.58	5	✓	Omega S29
Inj 100 mcg per ml, 1 ml vial	48.50	5	✓	Omega S29
Inj 500 mcg per ml, 1 ml vial	113.10	5	✓	Omega S29
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓	Max Health
			✓	Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓	Max Health
			✓	Octreotide GH S29
			✓	Sun Pharma S29
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓	Max Health
			✓	Octreotide GH S29
			✓	Sun Pharma S29
TAMOXIFEN CITRATE				
* Tab 10 mg	15.00	60	✓	Tamoxifen Sandoz
* Tab 20 mg	5.32	60	•	Tamoxifen Sandoz

Long-acting Somatostatin Analogues

⇒SA2445 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has not been successful; and
- 3 Treatment to be given for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has acromegaly; and
- 2 Either:
 - 2.1 Treatment with surgery and radiotherapy is not suitable or was unsuccessful; or
 - 2.2 Treatment is for an interim period while awaiting the beneficial effects of radiotherapy; and
- 3 Treatment with a dopamine agonist has been unsuccessful.

Renewal — (Acromegaly) from any relevant practitioner. Approvals valid for 2 years where iGF1 levels have decreased since starting treatment.

Note: In patients with acromegaly, treatment should be discontinued if IGF1 levels have not decreased 3 months after treatment. In patients treated with radiotherapy treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following treatment withdrawal for at least 4 weeks

Initial application — (pre-operative acromegaly) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

Initial application — (Other Indications) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Surgery has been unsuccessful; or
 - 2.2.2 Patient has metastatic disease after treatment with H2 antagonist or proton pump inhibitors has been unsuccessful; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has not been successful; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of a long-acting somatostatin analogue in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded under Special Authority

Renewal — (Other Indications) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LANREOTIDE	- Special Authority	see SA2445 on the previous	page - Retail pharmacy

Inj 60 mg per 0.5 ml, 0.5 ml syringe	382.77 1	Mytolac
Inj 120 mg per 0.5 ml, 0.5 ml syringe	646.70 1	Mytolac

OCTREOTIDE LONG-ACTING - Special Authority see SA2445 on the previous page - Retail pharmacy

Inj depot 10 mg prefilled syringe	438.40	1	✓ Sandostatin LAR
Inj depot 20 mg prefilled syringe		1	 Sandostatin LAR
Ini denot 30 ma prefilled syringe	670.80	1	✓ Sandostatin I AR

Aromatase Inhibitors

ΑN	ASTROZOLE		
*	Tab 1 mg4.39	30	✓ Anatrole

	LIVILGTANE			
*	Tab 25 mg	9.86	30	✓ Pfizer Exemestane

LETROZOLE

AZATI IIODDINIE

CVEMECTANE

*	Tab 2.5 mg	4.36	28	✓ Accord S29
		4.67	30	✓ Letrole

Immunosuppressants

Cytotoxic Immunosuppressants

AZI	ATHIOPHINE			
*	Tab 25 mg	7.36	60	✓ Azamun
*	Tab 50 mg	8.10	100	✓ Azamun

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer
MYCOPHENOLATE MOFETIL				
Tab 500 mg	35.90	50	1	Cellcept
Cap 250 mg	35.90	100	✓ (Cellcept
Powder for oral lig 1 g per 5 ml - Subsidy by endorsement	187.25	165 ml OP	1	Cellcept
Mycophenolate powder for oral liquid is subsidised only	for patients unab	ole to swallow t	ablets a	and capsules, and when
the prescription is endorsed accordingly.	·			•

Fusion Proteins

ETANERCEPT - Special Authority see SA2399 below - R	etail pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	Enbrel
Inj 50 mg autoinjector		4	Enbrel
Inj 50 mg prefilled syringe		4	Enbrel

⇒SA2399 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 12 Fither
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

```
18-24 years - Male: 7.0 cm; Female: 5.5 cm
25-34 years - Male: 7.5 cm; Female: 5.5 cm
35-44 years - Male: 6.5 cm; Female: 4.5 cm
45-54 years - Male: 6.0 cm; Female: 5.0 cm
55-64 years - Male: 5.5 cm; Female: 4.0 cm
65-74 years - Male: 4.0 cm; Female: 4.0 cm
75+ years - Male: 3.0 cm; Female: 2.5 cm
```

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA): and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA: or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)			Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price	·)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	/	Manufacturer

continued...

- 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 1.2.2 Fither:
 - 1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
- 1.3 Both:
 - 1.3.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
 - 1.3.2 Fither:
 - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing etanercept; and
- 2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose): and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose): and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:

Subsidy Fully (Manufacturer's Price) Subsidised		
 \$	Per 🗸	Manufacturer

continued...

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Specia	llist		
Inj 50 mg per ml, 5 ml	4,439.17	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only	Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29

Monoclonal Antibodies

ADALIMUMAB (AMGEVITA) - Special Authority see SA2400 below	- Retail pharmacy		
Inj 20 mg per 0.4 ml prefilled syringe	190.00	1	✓ Amgevita
Inj 40 mg per 0.8 ml prefilled pen	375.00	2	✓ Amgevita
Inj 40 mg per 0.8 ml prefilled syringe	375.00	2	✓ Amgevita

⇒SA2400 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist or any relevant practitioner on the

Subsidy	y Full	Brand or
(Manufacturer	's Price) Subsidise	I Generic
\$	Per 🗸	Manufacturer

continued...

recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 12 Fither
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has experienced reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has experienced reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value; or
- 3 Both:
 - 3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
 - 3.2 Either:
 - 3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — **(Crohn's disease - fistulising)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has confirmed Crohn's disease: and

Subsidy		ully	Brand or
 (Manufacturer's Price)	Subsid	isea	Generic
\$	Per	✓	Manufacturer

continued...

- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Fither
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Fither:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right): or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA): and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects: or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both
 - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Fither
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
- 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis: and
 - 1.2 Fither
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Fither:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiation on biologic therapy.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:

Subsidy	Ful	ly Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
 \$	Per •	 Manufacturer 	

continued...

- 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
- 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application: or
- 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Å BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease – severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved guality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:
All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	idised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued...

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment: or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or

Subsidy (Manufacturer's Price)	Full Subsidise		
\$	Per 🗸	Manufacturer	

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Fither:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

0.00

1

✓ Eylea

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

- Fither:
 - 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy: or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Fither:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
 - 2 Fither:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

BEVACIZUMAB - PCT only - Special Authority see SA2444 below

Inj 25 mg per ml, 4 ml vial	69.00	1	✓ Vegzelma
Inj 25 mg per ml, 16 ml vial	276.00	1	✓ Vegzelma
Inj 1 mg for ECP	0.71	1 mg	✓ Baxter

⇒SA2444 Special Authority for Subsidy

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with bevacizumab, and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
 - 2.2 Patient has preserved liver function (Child-Pugh A); and
 - 2.3 Transarterial chemoembolisation (TACE) is unsuitable: and
 - 2.4 Any of the following:
 - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
 - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
 - 2.4.3 Both:
 - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
 - 2.4.3.2 No disease progression since initiation of lenvatinib; and
 - 2.5 Patient has an ECOG performance status of 0-2; and
 - 2.6 To be given in combination with atezolizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

Initial application — (advanced or metastatic ovarian cancer) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has FIGO Stage IV epithelial ovarian, fallopian tube, or primary peritoneal cancer; or
 - 1.2 Both:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.2.1 The patient has previously untreated advanced (FIGO Stage IIIB or IIIC) epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 1.2.2 Either:
 - 1.2.2.1 Debulking surgery is inappropriate; or
 - 1.2.2.2 The cancer is sub-optimally debulked (maximum diameter of any gross residual disease greater than 1cm); and
- 2 Bevacizumab to be administered at a maximum dose of 7.5 mg/kg every three weeks; and
- 3 18 weeks concurrent treatment with chemotherapy is planned.

Renewal — (advanced or metastatic ovarian cancer) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

Initial application — (Recurrent Respiratory Papillomatosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Maximum of 6 doses; and
- 2 The patient has recurrent respiratory papillomatosis; and
- 3 The treatment is for intra-lesional administration.

Renewal — (Recurrent Respiratory Papillomatosis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Maximum of 6 doses: and
- 2 The treatment is for intra-lesional administration; and
- 3 There has been a reduction in surgical treatments or disease regrowth as a result of treatment.

Initial application — (Ocular Conditions) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Ocular neovascularisation: or
- 2 Exudative ocular angiopathy.

BRENTUXIMAB VEDOTIN - PCT only - Special Authority see \$A2289 below

⇒SA2289 Special Authority for Subsidy

Initial application — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and
 - 1.1.2 Patient is ineligible for autologous stem cell transplant; or
 - 1.2 Both:
 - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
 - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
Ψ	1 01		Wandacturer

continued...

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

Initial application — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Renewal — (anaplastic large cell lymphoma) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

⇒SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA2401 below

		and openial realitionity dec on E-101 below	- 10 Milling 10 10 10 iiiy opeolalio
Erbitux	1	364.00	Inj 5 mg per ml, 20 ml vial
✓ Erbitux	1	1,820.00	Inj 5 mg per ml, 100 ml vial
✓ Baxter	1 ma	3.82	Ini 1 mg for ECP

⇒SA2401 Special Authority for Subsidy

Initial application — (head and neck cancer, locally advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Cisplatin is contraindicated or has resulted in intolerable side effects; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 To be administered in combination with radiation therapy.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and
- 2 There is documentation confirming disease is RAS and BRAF wild-type; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Patient has not received prior funded treatment with cetuximab; and
- 5 Either:
 - 5.1 Cetuximab is to be used in combination with chemotherapy; or
 - 5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

Renewal — (colorectal cancer, metastatic) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where there is no evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

GEMTUZUMAB OZOGAMICIN − PCT only − Specialist − Special Authority see SA2269 below Inj 5 mg vial12,973.00 1 ✓ Mylotarg

⇒SA2269 Special Authority for Subsidy

Initial application only from a haematologist, paediatric haematologist or paediatric oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2402 below

 Inj 100 mg
 428.00
 1
 ✓ Remicade

 Inj 1 mg for ECP
 4.40
 1 mg
 ✓ Baxter

⇒SA2402 Special Authority for Subsidy

Initial application — (**Crohn's disease (adults)**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3 Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (adults)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Crohn's disease (children))** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Renewal — (Crohn's disease (children)) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the qut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patent has complex peri-anal fistula.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement: and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		idised	Generic	
 \$	Per	✓	Manufacturer	

continued...

- 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; or
 - 1.3 Both:
 - 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
 - 1.3.2 Either:
 - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing infliximab; and

Subsidy	Fully	/ Brand or
(Manufacturer's Price)	Subsidised	d Generic
\$	Per 🗸	Manufacturer

continued...

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis: or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults): or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis: or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Subsidy	Ful	ly Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	 Manufacturer 	

continued...

Renewal — **(rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (fulminant ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and

Subsidy (Manufactureria Drice)	Cubai	Fully	Brand or
 (Manufacturer's Price) \$	Per	dised ✓	Generic Manufacturer

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses: or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < 1/2+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patients SCCAI is greater than or equal to 4; or
 - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0 10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

MEPOLIZUMAB - Special Authority see SA2331 below - Retail pharmacy

⇒SA2331 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initial application — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has eosinophilic granulomatosis with polyangiitis; and

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and
- 3 Fither:
 - 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
 - 3.2 Corticosteroids are contraindicated.

Renewal — (eosinophilic granulomatosis with polyangiitis) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where patient has no evidence of clinical disease progression.

OBINUTUZUMAB - PCT only - Specialist - Special Au	uthority see SA2155 on the	next page	
Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

				
	Subsidy	Fully	Brand or	_
	(Manufacturer's Price)	Subsidised	Generic	
	\$	Por 🗸	Manufacturer	

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive: and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*: and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy ✓ Xolair ✓ Xolair AU Inj 150 mg vial450.00 ✓ Xolair 1

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
(Manuacturer's Frice)		oubsidised		
\$	Per	/	Manufacturer	

continued...

- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Special Authority see SA2419 below

Synagis

SA2419 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Infant was born in the last 12 months; and
 - 2.1.2 Infant was born at less than 32 weeks zero days' gestation; or
 - - 2.2.1 Child was born in the last 24 months; and
 - 2.2.2 Any of the following:
 - 2.2.2.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
 - 2.2.2.2 Both:
 - 2.2.2.2.1 Child has haemodynamically significant heart disease; and
 - 2.2.2.2. Any of the following:
 - 2.2.2.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
 - 2.2.2.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
 - 2.2.2.2.3 Child has severe pulmonary hypertension (see Note C); or
 - 2.2.2.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
 - 2.2.2.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
 - 2.2.2.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Child was born in the last 24 months; and
- 3 Any of the following:
 - 3.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
 - 3.2 Both:
 - 3.2.1 Child has haemodynamically significant heart disease; and
 - 3.2.2 Any of the following:
 - 3.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B);
 - 3.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
 - 3.2.2.3 Child has severe pulmonary hypertension (see Note C); or
 - 3.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
 - 3.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant: or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

continued...

3.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Notes:

- a) Ventilatory/respiratory support includes those on home oxygen, CPAP/VPAP and those with tracheostomies in situ managed at home
- b) Child requires/will require heart failure medication, and/or child has significant pulmonary hypertension, and/or infant will require surgical palliation/definitive repair within the next 3 months
- c) Mean pulmonary artery pressure more than 25 mmHg
- d) LV Ejection Fraction less than 40%
- e) Inborn errors of immunity include, but are not limited to, IFNAR deficiencies

PERTUZUMAB - PCT only - Specialist - Special Authority see SA2276 below

Inj 30 mg per ml, 14 ml vial	3,927.00	1	/	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	1	Baxter

⇒SA2276 Special Authority for Subsidy

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Fither:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 on the next page

inj 100 mg per 10	mi viai	1,075.50	2	•	Madthera
Inj 500 mg per 50	ml vial	2,688.30	1	1	Mabthera
Inj 1 mg for ECP		5.64	1 mg	✓	Baxter (Mabthera)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

6 Either:

- 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
- 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

7 Either:

- 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and

8 Fither:

- 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Both:

- 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and

2 Fither:

- 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2233 below

Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2233 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks: and
- 3 Any of the following:

continued...

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
- 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
- 3.3 Cyclophosphamide and methotrexate are contraindicated; or
- 3.4 Patient is a female of child-bearing potential; or
- 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL;
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Fither:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

3 Fither:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity: and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
 - 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
 - 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective;
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy;
 - 2.2 To be used for a maximum of 6 treatment cycles.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*: and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Fither:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange: or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — **(treatment refractory systemic lupus erythematosus (SLE))** only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Subsidy	Fı	ılly	Brand or	_
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	/	Manufacturer	

continued...

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	•	Manufacturer

continued...

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Fither
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

Lilliel.

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

- Both:
 - 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
 - 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment: or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — **(desensisation prior to transplant)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions: or
 - 1.3.3 Involvement of two or more mucosal sites: or
- 2 Both:
 - 2.1 Patient has pemphigus; and

Subsic		Fully	Brand or
(Manufacture	r's Price) Su	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphiqus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

Initial application — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD*; and
- - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or
 - 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance: and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart. Note: Indications marked with * are unapproved indications.

Renewal — (immunoglobulin G4-related disease (IgG4-RD*)) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
 - 1.2 Patient is receiving maintenance treatment for IgG4-RD*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2403 below - Retail pharmacy Inj 150 mg per ml, 1 ml prefilled syringe.......799.50 Cosentyx 1.599.00 Cosentvx

⇒SA2403 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plague psoriasis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand. foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Either:
 - 1.1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic localised genital or flexural plague psoriasis at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.2.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and

Subsidy	y Fully	Brand or
(Manufacturer	's Price) Subsidised	I Generic
\$	Per 🗸	Manufacturer

continued...

2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Inj 100 mg vial	770.57	1	Sylvant
Inj 400 mg vial	3,082.33	1	✓ Sylvant

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA2404 below

Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra
Inj 20 mg per ml, 10 ml vial	550.00	1	✓ Actemra
Inj 20 mg per ml, 20 ml vial		1	✓ Actemra
Inj 1 mg for ECP	,	1 mg	✓ Baxter

⇒SA2404 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
- 1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
 - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 ` \$	Per	✓	Manufacturer

continued...

the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Fither:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Fither:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
<u> </u>	Per		Manufacturer

continued...

- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB (HERZUMA) - PCT only - Special Authority see \$A2293 below

✓ Herzuma	1	100.00	 	 Inj 150 mg vial
✓ Herzuma	1	293.35	 	 Inj 440 mg vial
✓ Baxter	1 ma	0.70	 	 Ini 1 ma for ECP

⇒SA2293 Special Authority for Subsidy

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Renewal — (early breast cancer*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 1.3 Any of the following:
 - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
 - 1.3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and

1.4 Fither:

- 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
- 1.4.2 All of the following:
 - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

Initial application — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and

(Man	Subsidy ufacturer's Price)	Fu Subsidis	ılly	Brand or Generic
(wich	. ,	Per	✓	Manufacturer

continued...

- 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression: and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initial application — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an FCOG score of 0-2.

Renewal — (gastric, gastro-oesophageal junction and oesophageal cancer) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB DERUXTECAN - PCT only - Special Authority see \$A2420 below 1

✓ Enhertu Inj 1 mg for ECP.......27.05 ✓ Baxter 1 ma

⇒SA2420 Special Authority for Subsidy

Initial application only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with trastuzumab deruxtecan and met all remaining criteria prior to commencing treatment: or
- 2 All of the following:
 - 2.1 Patient has metastatic breast cancer expressing HER-2 IHC3+ or ISH+ (including FISH or other current
 - 2.2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
 - 2.3 Either:
 - 2.3.1 The patient has received prior therapy for metastatic disease; or
 - 2.3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy; and
 - 2.4 Patient has a good performance status (ECOG 0-1); and
 - 2.5 Patient has not received prior funded trastuzumab deruxtecan treatment; and
 - 2.6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab deruxtecan; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2424 on the next page

Inj 100 mg vial	<i>,</i>	 2,320.00	1	Kadcyla
Inj 160 mg vial		 3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP		 24.52	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✔ Manufacturer

⇒SA2424 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or axiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadiuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Either:
 - 6.1 Patient has not received prior funded trastuzumab emtansine or trastuzumab deruxtecan treatment; or
 - 6.2 Both:
 - 6.2.1 Patient has discontinued trastuzumab deruxtecan due to intolerance; and
 - 6.2.2 The cancer did not progress while on trastuzumab deruxtecan; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB - Special Authority see SA2182 below - Retail pharmacy

Ini 90 mg per ml, 1 ml pre-filled syringe........................4,162.00 1 ✓ Stelara

⇒SA2182 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 2 Both:
 - 2.1 Patient has active Crohn's disease: and
 - 22 Fither:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed: and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2.1 Patient has active ulcerative colitis; and
- 2.2 Fither:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB - PCT only - Special Authority see SA2183 below

⇒SA2183 Special Authority for Subsidy

Initial application — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initial application — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 6 months for applications

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (Crohn's disease - children*) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initial application — (ulcerative colitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

Programmed Cell Death-1 (PD-1) Inhibitors

ATEZOLIZUMAB - PCT only - Specialist - Special Aut	hority see SA2443 below		
Inj 60 mg per ml, 20 ml vial	9,503.00	1	✓ Tecentriq
Inj 1 mg for ECP	8.08	1 mg	✓ Baxter

⇒SA2443 Special Authority for Subsidy

Initial application — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer second line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient is currently on treatment with atezolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
 - 2.2 Patient has preserved liver function (Child-Pugh A); and
 - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
 - 2.4 Any of the following:
 - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
 - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
 - 2.4.3 Both:
 - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
 - 2.4.3.2 No disease progression since initiation of lenvatinib; and

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic	
` \$ ´	Per	1	Manufacturer	

continued...

- 2.5 Patient has an ECOG performance status of 0-2; and
- 2.6 To be given in combination with bevacizumab.

Renewal — (unresectable hepatocellular carcinoma) from any relevant practitioner. Approvals valid for 6 months where there is no evidence of disease progression.

DURVALUMAB - PCT only - Specialist - Special Authorit	ty see SA2425 below		
Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	Imfinzi
Inj 1 mg for ECP	9.59	1 mg	Baxter

⇒SA2425 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); or
 - 1.2 Patient has histologically or cytologically documented stage IIb (T1N2a only), locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

NIVOLUMAB - PCT only - Specialist - Special Authority see SA240	05 below		
Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	✓ Opdivo
Inj 1 mg for ECP		1 mg	✓ Baxter

⇒SA2405 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal — (less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment; or
 - 1.1.2 Patient's disease has had a partial response to treatment; or
 - 1.1.3 Patient has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
 - 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Renewal — (more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Any of the following:
 - 2.1.1.1 Patient's disease has had a complete response to treatment; or
 - 2.1.1.2 Patient's disease has had a partial response to treatment; or
 - 2.1.1.3 Patient has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
 - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2.2 All of the following:
 - 2.2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 Patient has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

Initial application — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

Subs (Manufactur	.,	ully Brand or sed Generic	
	Per	✓ Manufacture	er

continued...

- 1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has metastatic renal-cell carcinoma; and
 - 2.2 The disease is of predominant clear-cell histology; and
 - 2.3 Patient has an ECOG performance score of 0-2; and
 - 2.4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and
 - 2.5 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

Renewal — (Renal cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA2386 below

Inj 25 mg per ml, 4 ml vial	4,680.00	1	Keytruda
Inj 1 mg for ECP	47.74	1 mg	✓ Baxter

⇒SA2386 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma, less than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment; or
 - 1.1.2 Patient's disease has had a partial response to treatment; or
 - 1.1.3 Patient has stable disease; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	I Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Renewal — (unresectable or metastatic melanoma, more than 24 months on treatment) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Any of the following:
 - 2.1.1.1 Patient's disease has had a complete response to treatment; or
 - 2.1.1.2 Patient's disease has had a partial response to treatment; or
 - 2.1.1.3 Patient has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
 - 2.1.3 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2.2 All of the following:
 - 2.2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 Patient has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initial application — (non-small cell lung cancer first-line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Either:
 - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
 - 6.2 Both:
 - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
 - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2: and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and

continued...

9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line monotherapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease: and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (non-small cell lung cancer first-line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2: and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Renewal — (non-small cell lung cancer first line combination therapy) only from a medical oncologist or any relevant practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (breast cancer, advanced) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
 - 2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); and
 - 2.2 Patient is treated with palliative intent; and
 - 2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
 - 2.4 Patient has received no prior systemic therapy in the palliative setting; and
 - 2.5 Patient has an ECOG score of 0-2: and
 - 2.6 Pembrolizumab is to be used in combination with chemotherapy; and
 - 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (breast cancer, advanced) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period; and
- 4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (head and neck squamous cell carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies; and
 - 2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
 - 2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
 - 2.4 Patient has an ECOG performance score of 0-2; and
 - 2.5 Either:
 - 2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or
 - 2.5.2 Pembrolizumab to be used as monotherapy; and
 - 2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (head and neck squamous cell carcinoma) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Subsidy	Full	y Brand or	
(Manufacturer's Pr	rice) Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 1.1 Patient's disease has had a complete response to treatment; or
- 1.2 Patient's disease has had a partial response to treatment; or
- 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (MSI-H/dMMR advanced colorectal cancer) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal
 - 2.1.2 Patient has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer; and
 - 2.2 Patient is treated with palliative intent; and
 - 2.3 Patient has not previously received funded treatment with pembrolizumab; and
 - 2.4 Patient has an ECOG performance score of 0-2; and
 - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (MSI-H/dMMR advanced colorectal cancer) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
 - 2.2 Patient has an ECOG performance score of 0-2; and
 - 2.3 Patient has documented disease progression following treatment with chemotherapy; and
 - 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Renewal — (Urothelial carcinoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	,	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 No evidence of disease progression; and
- 3 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initial application — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Both:
 - 2.1.1.1 Patient has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and
 - 2.1.1.2 Patient is ineligible for autologous stem cell transplant; or
 - 2.1.2 Patient has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant; and
 - 2.2 Patient has not previously received funded pembrolizumab; and
 - 2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Renewal — (relapsed/refractory Hodgkin lymphoma) only from a relevant specialist or any relevant practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has received a partial or complete response to pembrolizumab; and
- 2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2414 below - Reta	il pharmacy		
Wastage claimable			
Tab 10 mg	6,512.29	30	Afinitor
Tab 5 mg	4,555.76	30	✓ Afinitor

⇒SA2414 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — **(renal cell carcinoma)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma; and
 - 1.2 The disease is of predominant clear-cell histology; and
 - 1.3 The patient has documented disease progression following one previous line of treatment; and
 - 1.4 The patient has an ECOG performance status of 0-2; and
 - 1.5 Everolimus is to be used in combination with lenvatinib; or
- 2 All of the following:
 - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
 - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
 - 2.3 Everolimus is to be used in combination with lenvatinib; and
 - 2.4 There is no evidence of disease progression.

Renewal — (renal cell carcinoma) from any relevant practitioner. Approvals valid for 4 months where there is no evidence of disease progression.

SIROLIMUS - Special Authority see SA2270 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg		100	✓ Rapamune
Oral lig 1 mg per ml	449.99	60 ml OP	✓ Rapamune

⇒SA2270 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis: or
- . HUS or TTP; or
- · Leukoencepthalopathy: or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Fither

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy	Fully	/ Brand or	_
(Manufacturer's Price)	Subsidised	d Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
- 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease: and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing age potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$		Subsidised	Generic
TACROLIMUS – Special Authority see SA2271 below – Retail ph	,	100		Tanadimus Candan
Cap 0.75 mg		100		Tacrolimus Sandoz Tacrolimus Sandoz
Cap 1 mg		100		Tacrolimus Sandoz
Cap 5 mg		50	✓	Tacrolimus Sandoz

⇒SA2271 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Either:
 - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
 - 2.2 Patient is a child with nephrotic syndrome*.

Note: Indications marked with * are unapproved indications

JAK inhibitors

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 1 The patient 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Fither:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

=itner:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Antiallergy Preparations

Allergic Emergencies

ADRENALINE - Special Authority see SA2185 below - Retail pharmacy

- a) Maximum of 2 ini per prescription
- Additional prescriptions limited to replacement of up to two devices prior to expiry, or replacement of used device for treatment of anaphylaxis.

Inj 0.15 mg per 0.3 ml auto-injector	90.00	1 OP	 Epipen Jr
Inj 0.3 mg per 0.3 ml auto-injector	90.00	1 OP	✓ Epipen

⇒SA2185 Special Authority for Subsidy

Initial application — (anaphylaxis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has experienced an anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
 - 1.2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner; and
 - 2 Patient is not to be prescribed more than two devices in initial prescription.

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy
Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy		Fully Brand or
(Manufacturer's Price		sidised Generic
	\$	Per	✓ Manufacturer
EE VENOM ALLERGY TREATMENT - Special Authority see SA			
Initiation kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Initiation kit - 5 vials freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with			
diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	334.39	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
VENOX S29 Initiation kit - 5 vials freeze dried venom with diluent	to be delisted 1 l	May 2025)	
VASP VENOM ALLERGY TREATMENT - Special Authority see	SA1367 on the pre	evious page	- Retail pharmacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		13	1,
dried polistes venom, 1 diluent 9 ml, 3 diluent 1.8 ml	382.23	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			•
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze			
dried vespula venom, 1 diluent 9 ml, 3 diluent 1.8 ml	431.24	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Aut 19 de Level de la			
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
₭ Tab 10 mg	1.71	100	✓ Zista
FOral liq 1 mg per ml		200 ml	✓ Histaclear
EXTROCHLORPHENIRAMINE MALEATE			
€ Tab 2 mg	2.02	40	
•	(8.40)		Polaramine
	`1.01 [′]	20	
	(5.99)		Polaramine
F Oral liq 2 mg per 5 ml		100 ml	
	(10.29)		Polaramine
EXOFENADINE HYDROCHLORIDE			
★ Tab 60 mg	4.34	20	
	(8.23)		Telfast
€ Tab 120 mg		30	✓ Fexaclear
	14.22		-
h. T. L. 400	(26.44)		Telfast
₭ Tab 180 mg	4.10	30	✓ Fexaclear
Telfast Tab 120 mg to be delisted 1 July 2025)			
ORATADINE			
₭ Tab 10 mg		100	✓ <u>Lorafix</u>
* Oral liq 1 mg per ml	1 43	100 ml	Haylor syrup

	Subsidy		Fully Brand or
	(Manufacturer's	Price) Subsi	
	` \$	Per	✓ Manufacturer
PROMETHAZINE HYDROCHLORIDE			
	1.00	50	✓ Allersoothe
1		50 50	✓ Allersoothe
* Tab 25 mg		• • • • • • • • • • • • • • • • • • • •	
* Oral liq 1 mg per 1 ml		100 ml	✓ Allersoothe
	10.47	_	✓ Phenergan Elixir
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a F	2SO21.09	5	✓ Hospira
Inhaled Corticosteroids			
DECLOMETUA CONE DIDDODIONATE			
BECLOMETHASONE DIPROPIONATE	14.04	000 doss OD	✓ Qvar
Aerosol inhaler, 50 mcg per dose		200 dose OP	
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ Pulmicort
· · · · · · · · · · · · · · · · · · ·			Turbuhaler
Powder for inhalation, 200 mcg per dose	10.00	200 dose OP	✓ Pulmicort
1 Owder for illinatation, 200 fflog per dose	19.00	200 dose Oi	Turbuhaler
December for introduction 400 many many days	00.00	000 de e OD	
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	✓ Pulmicort
			Turbuhaler
FLUTICASONE			
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose	8.61	60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose	7.81	60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose	24.62	120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
, 01			
Inhaled Long-acting Beta-adrenoceptor Agonis	s		
milition zong doming zona daronosopio. Agomo			
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dose	10.32	60 dose OP	
(-1	(16.90)		Oxis Turbuhaler
INDACATEDOL	(10.00)		
INDACATEROL Revides for inhelation 150 mans	04.00	00 dese 00	/ Onbres Breeshele
Powder for inhalation 150 mcg		30 dose OP	✓ Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	 Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose	26.25	120 dose OP	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated		60 dose OP	✓ Serevent Accuhaler
<u> </u>			

	Subsidy		Fully	Brand or
	(Manufacturer's I	Price) Subsi	dised	Generic
	` \$	Per	1	Manufacturer
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocept	or Agonists		
BUDESONIDE WITH EFORMOTEROL				
Powder for inhalation 160 mcg with 4.5 mcg eformoterol				
fumarate per dose (equivalent to 200 mcg budesonide w	ith			
6 mcg eformoterol fumarate metered dose)		120 dose OP	1	DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumara				- actioop opiioiiian
per dose (equivalent to 400 mcg budesonide with 12 mcg				
eformoterol fumarate metered dose) – No more than 2	9			
	92.50	120 dose OP	./	DuoResp Spiromax
dose per day				Vannair
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP		
Powder for inhalation 100 mcg with eformoterol fumarate 6 m	icg33.74	120 dose OP	•	Symbicort
A	04.40	100 de - OD		Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP		Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 m	icg33.74	120 dose OP		Symbicort
				Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			_	
12 mcg - No more than 2 dose per day	33.74	60 dose OP		Symbicort
				Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL				
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	1	Breo Ellipta
FLUTICASONE WITH SALMETEROL				·
Aerosol inhaler 50 mcg with salmeterol 25 mcg	25 79	120 dose OP	1	Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP		Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg – No		120 0000 01	- '	Soronas
more than 2 dose per day	33.74	60 dose OP	1	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No		oo dose Oi	•	Seletiue Acculialei
	44.00	60 dose OP	./	Seretide Accuhaler
more than 2 dose per day	44.00	ou dose OF	•	Seretide Accumater
Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Oral liq 400 mcg per ml	50.00	150 ml	1	Ventolin
Ventolin to be Principal Supply on 1 May 2025				
Infusion 1 mg per ml, 5 ml		10	1	Ventolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO	130.00	5	1	Ventolin
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	4.18	200 dose OP	1	SalAir
	(6.80)		,	Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb				
available on a PSO	8.96	20	1	Asthalin
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule - Up to 30 neb				
available on a PSO		20	1	Asthalin
TERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to				
250 mcg metered dose), breath activated	22 20	120 dose OP	✓ 1	Bricanyl Turbuhaler
_55 mag material 4055/, broadin dollydiod		.20 0000 01	- 1	

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

20

✓ Univent

Anticholinergic Agents

IDD V	TROP	11 11 1	MIDE

(Pharmascience 29 Nebuliser soln, 250 mcg per ml, 2 ml ampoule to be delisted 1 May 2025)

Inhaled Beta-Adrenoceptor Agonists with Anticholinergic Agents

SALBUTAMOL WITH IPRATROPIUM BROMIDE

(Duolin Cipla S29) Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule to be delisted 1 April 2025)

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00 60 dose OP Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

BUDESONIDE WITH GLYCOPYRRONIUM AND EFORMOTEROL – Special Authority see SA2421 below – Retail pharmacy Aerosol inhaler budesonide 160 mcg with glycopyrronium

⇒SA2421 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long-acting muscarinic antagonist and long-acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler therapy.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – Special Authority see SA2326 on the next page – Retail pharmacy

Powder for inhalation fluticasone furoate 100 mcg with

umeclidinium 62.5 mcg and vilanterol 25 mcg......104.24 30 dose OP ✓ Trelegy Ellipta

⇒SA2326 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

- 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
- 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
- 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
- 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months: or
- 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

Antifibrotics

NINTEDANIB - Special Authority see SA2012 below - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and

3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90 OP	 Esbriet
Tab 267 mg	1,215.00	90	Esbriet

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib)

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Leukotriene Receptor Antagonists

MO	NTELUKAST			
*	Tab 4 mg	3.10	28	✓ Montelukast Viatris
*	Tab 5 mg	3.10	28	✓ Montelukast Viatris
*	Tab 10 mg	2.90	28	✓ Montelukast Viatris

Methylxanthines

AM	INC	PH	/LLI	NE
----	-----	----	------	----

*	Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO180.00	5	•	DBL Aminophylline
TH	EOPHYLLINE			
*	Tab long-acting 250 mg25.65	100	✓	Nuelin-SR
*	Oral liq 80 mg per 15 ml	500 ml	1	Nuelin

Mucolytics

DORNASE ALFA − Special Authority see SA1978 below − Retail pharmacy
Nebuliser soln, 2.5 mg per 2.5 ml ampoule.......250.00 6 ✓ Pulmozyme

Subsidy (Manufacturer's Pr	ice)	Fully Subsidised	Brand or Generic	
\$	Per	•	Manufacturer	

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR - PCT only - Special Authority see SA2196 below

Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg

Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg

⇒SA2196 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Either:
 - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Either:
 - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
 - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Notes:

a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212273s004lbl.pdf
 CASTOR - POT only - Cookelist - Coo

/ACAFTOR – PCT only – Specialist – Special Auth	iority see SA2017 below		
Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

continued...

	Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic
	\$	Per 🗸	Manufacturer
continued			

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele: or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

Not funded for use as a nasal drop.

90 ml OP ✓ Biomed

Nasal Preparations

Allergy Prophylactics

BUDESONIDE		
Metered aqueous nasal spray, 50 mcg per dose2.59	200 dose OP	✓ SteroClear
Metered aqueous nasal spray, 100 mcg per dose2.89	200 dose OP	✓ SteroClear
FLUTICA CONF. PROPIONATE		

FLUTICASONE PROPIONATE

Metered aqueous nasal spray, 50 mcg per dose1.98 120 dose OP ✓ Flixonase Hayfever & Allergy

IPRATROPIUM BROMIDE

15 ml OP ✓ Univent

Respiratory Devices

MASK FOR SPACER DEVICE

- a) Up to 50 dev available on a PSO
- b) Only on a PSO

b) Only on a PSO

c) Only for children aged six years and under

✓ e-chamber Mask

PEAK FLOW METER

a) Up to 25 dev available on a PSO

Low range	9.54	1	Mini-Wright AFS Low Range
Normal range	9.54	1	✓ Mini-Wright

Standard

25 ml OP

✓ Biomed

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
SPACER DEVICE				
a) Up to 50 dev available on a PSOb) Only on a PSO				
220 ml (single patient)	3.65	1	√ e	-chamber Turbo
510 ml (single patient)	5.95	1	√ e	-chamber La Grande
800 ml	6.50	1	✓ V	olumatic //
Respiratory Stimulants				
CAFFEINE CITRATE				

Oral liq 20 mg per ml (10 mg base per ml)......16.91

	Subsidy	Duite - \ Out -	Fully Brand or
	(Manufacturer's F	Price) Subs Per	idised Generic Manufacturer
			- Indianacial Ci
Ear Preparations			
FLUMETASONE PIVALATE	4.46	7.5 ml OD	✓ Locacorten-Viaform
Ear drops 0.02% with clioquinol 1%	4.40	7.5 ml OP	ED's
			✓ Locorten-Vioform
TRIANCINOLONE ACETONIDE WITH CRANICIDIN NEOMYC	IN AND NIVOTA	TINI	2 Edebiten-Violoniii
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND IN 151A	HIN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5 16	7.5 ml OP	✓ Kenacomb
2.5 mg and gramicidin 250 mcg per g		7.5 1111 01	Renacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4.50	8 ml OP	
•	(9.27)		Otodex S29
	(9.27)		Sofradex
FRAMYCETIN SULPHATE			
Ear/Eye drops 0.5%		8 ml OP	
	(8.65)		Soframycin
Eye Preparations			
Eye Freparations			
Eye preparations are only funded for use in the eye, unless expli	citly stated other	wise.	
	,		
Anti-Infective Preparations			
ACICLOVIR			
* Eye oint 3%	15.89	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL			
Eye oint 1%	1.09	5 g OP	✓ <u>Devatis</u>
Eye drops 0.5%		10 ml OP	✓ Chlorsig
Funded for use in the ear*. Indications marked with * ar	e unapproved in	idications.	
CIPROFLOXACIN			
Eye drops 0.3% – Subsidy by endorsement		5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis of			
for the second line treatment of chronic suppurative otiti Note: Indication marked with a * is an unapproved indic) ; and the presi	cription is endorsed accordingly.
• • • • • • • • • • • • • • • • • • • •	auon.		
SODIUM FUSIDATE [FUSIDIC ACID] Eye drops 1%	5 20	5 g OP	✓ Fucithalmic
2,5 diopo 1/5		0 g Oi	✓ Fucithalmic S29 S29
			- I dominimile 323
TOBRAMYCIN			
Eye oint 0.3%	10.45	3.5 g OP	✓ Tobrex
Eye drops 0.3%	11.48	5 ml OP	✓ Tobrex

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Corticosteroids and Other Anti-Inflammatory Preparations

DE	XAMETHASONE		
*	Eye oint 0.1%	3.5 g OP	Maxidex
*	Eye drops 0.1%	5 ml OP	Maxidex
	Ocular implant 700 mcg - Special Authority see SA1680 below		
	- Retail pharmacy1,444.50	1	Ozurdex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eve sint 0.19/ with reconvoir authors 0.259/ and polymorin b

sulphate 6,000 u per g		3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxi b sulphate 6,000 u per ml		5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM			
Eye drops 0.1%, single dose	1.85	10 dose	✓ Diclofenac Devatis
, , , ,	5.54	30 dose	✓ Diclofenac Devatis
FLUOROMETHOLONE			
* Eye drops 0.1%	3.09	5 ml OP	✓ FML
, ,	5.20		✓ Flucon

	Subsidy		Fully	Brand or
	(Manufacturer's P	rice) Subs	sidised	Generic
	\$	Per	•	Manufacturer
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
_,,	(10.34)		L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	✓ L	omide .
NEPAFENAC				
Eye drops 0.3%	8.80	3 ml OP	✓	evro
(Ilevro Eye drops 0.3% to be delisted 1 July 2025)				
PREDNISOLONE ACETATE				
Eye drops 1%	6.92	10 ml OP	✓ P	rednisolone-AFT
_,o	7.00	5 ml OP	-	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority	see SA1715 below	– Retail nhar	macv	
Eye drops 0.5%, single dose (preservative free)		20 dose		linims
Lye drops 0.0 /o, single dose (preservative free)	43.20	20 d086	3 IV	Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE Eye drops 2%	2.62 1	10 ml OP	✓ <u>Allerfix</u>
Glaucoma Preparations - Beta Blockers			
BETAXOLOL * Eye drops 0.25%		5 ml OP 5 ml OP	✓ Betoptic S ✓ Betoptic
* Eye drops 0.25% * Eye drops 0.5%		5 ml OP 5 ml OP	✓ <u>Arrow-Timolol</u> ✓ <u>Arrow-Timolol</u>
Glaucoma Preparations - Carbonic Anhydrase Inh	ibitors		
ACETAZOLAMIDE * Tab 250 mg	17.03	100	✓ Diamox
***		100 5 ml OP	✓ Diamox ✓ <u>Azopt</u>
* Tab 250 mg BRINZOLAMIDE * Eye drops 1%	5.11		
* Tab 250 mg BRINZOLAMIDE * Eye drops 1% DORZOLAMIDE WITH TIMOLOL	5.11	5 ml OP	✓ <u>Azopt</u>

	Subsidy (Manufacturer's Price	ce) Subsi	Fully dised	Brand or Generic
	\$	Per	✓	Manufacturer
LATANOPROST				
* Eye drops 0.005%	2.08	2.5 ml OP	✓ T	eva
TRAVOPROST				
* Eye drops 0.004%	6.80	2.5 ml OP	√ T	ravatan
4 2/5 dispo 5:55 1/5		2.0 11.11 01		- uvuun
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE				
* Eye drops 0.2%	5.16	5 ml OP	./ A	rrow-Brimonidine
		3 1111 01	• -	illow-billionidille
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE	7.40	5l OD		
* Eye drops 0.2% with timolol maleate 0.5%	7.13	5 ml OP	√ C	<u>ombigan</u>
LATANOPROST WITH TIMOLOL				
* Eye drops 0.005% with timolol 0.5%	4.95	2.5 ml OP	✓ <u>A</u>	rrow - Lattim
PILOCARPINE HYDROCHLORIDE				
* Eye drops 1%	4.26	15 ml OP		sopto Carpine
* Eye drops 2%		15 ml OP		opto Carpine
* Eye drops 4%		15 ml OP	✓ Is	opto Carpine
Subsidised for oral use pursuant to the Standard Formula	ie.			
PILOCARPINE NITRATE				
* Eye drops 2% single dose – Special Authority see SA0895				
below – Retail pharmacy	35.90	20 dose	✓ M	linims Pilocarpine

⇒SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics		
ATROPINE SULPHATE * Eye drops 1%	15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE	10 1111 01	- Mapt
* Eye drops 1%	15 ml OP	✓ Cyclogyl
TROPICAMIDE * Eye drops 0.5%	15 ml OP 15 ml OP	✓ Mydriacyl✓ Mydriacyl
Preparations for Tear Deficiency		
For acetylcysteine eye drops refer Standard Formulae, page 273		
HYPROMELLOSE * Eye drops 0.5%	15 ml OP	✓ Methopt
HYPROMELLOSE WITH DEXTRAN	15 IIII OF	wethopt

15 ml OP

✓ Poly-Tears

Eye drops 0.3% with dextran 0.1%......2.30

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

Preservative Free Ocular Lubricants

⇒SA2431 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

CARBOMER – Special Authority see SA2431 above – Retail ph	narmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
(Poly-Gel Ophthalmic gel 0.3%, 0.5 g to be delisted 1 July 2025,)		•
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL -	- Special Authority se	e SA2431 a	bove – Retail pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml	10.78	30	✓ Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Aut	hority see SA2431 a	bove – Retai	l pharmacy
Eye drops 1 mg per ml	13.58	10 ml OP	✓ <u>Hylo-Fresh</u>
Hylo-Fresh has a 6 month expiry after opening. The Ph	narmacy Procedures	Manual restr	riction allowing one bottle per
month is not relevant and therefore only the prescribed	dosage to the neare	st OP may be	e claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE		
* Eye drops 0.1%	15 ml OP	✓ <u>Albalon</u>
OLOPATADINE		
Eye drops 0.1%2.17	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT		
* Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE		
Eve oint 138 mca per a	5 a OP	✓ VitA-POS

					VARIOUS
					VARIOUS
_		Subsidy (Manufacturer's Pric \$	e) Subs Per	idised	Brand or Generic Manufacturer
٧	arious				
PH	ARMACY SERVICES				
	Brand switch fee	4.50	1 fee	✓ BS	F Dasatinib-Teva
	 a) May only be claimed once per patient. 				
.14	b) The Pharmacode for BSF Dasatinib-Teva is 270044				
*	Immunisation administration fee - flu		1 fee 1 fee		munisation - Flu munisation Other
	Immunisation co-administration fee - flu and shingles		1 fee		nunisation Flu
~	minumodion co duministration rec- na ana simigros		1100		nd Shingles
(BS	SF Dasatinib-Teva Brand switch fee to be delisted 1 June 202	5)			3
A	gents Used in the Treatment of Poisonings				
_	utidataa				
A	ntidotes				
AC	ETYLCYSTEINE				
	Inj 200 mg per ml, 10 ml ampoule	42.99 52.88	10		L Acetylcysteine rtindale Pharma
	DBL Acetylcysteine to be Principal Supply on 1 April 202	25			
(M	artindale Pharma Inj 200 mg per ml, 10 ml ampoule to be delis	sted 1 November 20	025)		
NA	LOXONE HYDROCHLORIDE				
	a) Up to 10 inj available on a PSO				
	b) Only on a PSO		_		
*	Inj 400 mcg per ml, 1 ml ampoule	13.29	5		L Naloxone
		35.26	10	г ✓ Ha	lydrochloride
	DBL Naloxone Hydrochloride to be Principal Supply on		10	• па	mein
(На	ameln Inj 400 mcg per ml, 1 ml ampoule to be delisted 1 April 3				
		,			
R	emoval and Elimination				
	ARCOAL				
*	Oral liq 50 g per 250 ml	43.50	250 ml OP	✓ Ca	rbosorb-X
	a) Up to 250 ml available on a PSO				
	b) Only on a PSO				
DE	FERASIROX – Special Authority see SA1492 below – Retail	pharmacy			
	Wastage claimable	076.00	00	./	lada.
	Tab 125 mg dispersible		28	✓ Exj	aue

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and

- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine

continued...

✓ Exjade

✓ Exjade

28

28



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or

- 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
- 3.3 Treatment with deferiprone has resulted in arthritis; or
- 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below - F	Retail pharmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERBIOXAMINE I	

* Inj 500 mg vial	151.31	10	✓ Deferoxamine Pfizer S29 S29
	332.88		✓ DBL Desferrioxamine Mesylate for Inj BP
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate

Standard Formulae

Water

otaliaala i olillalac			
ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
Suitable eye drop base	qs	Phenobarbitone Sodium	400 mg
		Glycerol BP	4 ml
CODEINE LINCTUS (3 mg per 5 ml)		Water	to 40 ml
Codeine phosphate	60 mg		
Glycerol	40 ml	PILOCARPINE ORAL LIQUID	
Preservative	qs	Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
		Water	to 500 ml
CODEINE LINCTUS (15 mg per 5 ml)		(Preservative should be used if quantity supplied is	for more
Codeine phosphate	300 mg	than 5 days.)	
Glycerol	40 ml	2 22,27	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	
Water	to 100 ml	Methylcellulose	5 g
		Preservative	qs
FOLINIC MOUTHWASH		Water	to 500 ml
Calcium folinate 15 mg tab	1 tab	(Preservative should be used if quantity supplied is	for more
Preservative	qs	than 5 days. Maximum 500 ml per prescription.)	
Water	to 500 ml	, , , , , , , , , , , , , , , , , , , ,	
(Preservative should be used if quantity supplied is	for more	SODIUM CHLORIDE ORAL LIQUID	
than 5 days. Maximum 500 ml per prescription.)		Sodium chloride inj 23.4%, 20 ml	qs
		Water	qs
METHYL HYDROXYBENZOATE 10% SOLUTION		(Only funded if prescribed for treatment of hyponatr	aemia)
Methyl hydroxybenzoate	10 g	NAMES OF A COLUMN	
Propylene glycol	to 100 ml	VANCOMYCIN ORAL SOLUTION (25 mg per ml)	
(Use 1 ml of the 10% solution per 100 ml of oral liqu	iid mixture)	Vancomycin 500 mg injection	5 vials
OMEPRAZOLE SUSPENSION		Glycerin with sucrose suspension	37.5 ml
		Water	to 100 ml
Omeprazole capsules or powder	qs	(Only funded if prescribed for treatment of Clostridiu	ım difficile
Sodium bicarbonate powder BP	8.4 g	following metronidazole failure)	
Water	to 100 ml		
PHENOBARBITONE ORAL LIQUID			
Phenobarbitone Sodium	1 g		
Glycerol BP	70 ml		
Cilycerol Di	701111		

to 100 ml

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✓ Manufacturer

Extemporaneously	Compounded Prep	parations and Galenicals
------------------	------------------------	--------------------------

CODEINE PHOSPHATE – Safety medicine; prescriber may determine	e dispensina f	requency	
Powder - Only in combination		25 g	
	(90.09)		Douglas
Only in extemporaneously compounded codeine linctus.			
COLLODION FLEXIBLE			
Note: This product is no longer being manufactured by the suppl	er and will be	delisted from	the Schedule at a date to be
determined. Collodion flexible	10 30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE – Only in combination	10.00	100 1111	· I OM
Only in extemporaneously compounded oral mixtures.			
Soln	30.00	100 ml	✓ Midwest
GLYCERIN WITH SODIUM SACCHARIN - Only in combination			
Only in combination with Ora-Plus or when used in the vancomyo	in oral Iguuid S	Standard Form	nulae.
Suspension		473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE - Only in combination			
Only in combination with Ora-Plus or when used in the vancomyo	in oral Iquuid S	Standard Forn	nulae.
Suspension	30.95	473 ml	✓ Ora-Sweet
GLYCEROL			
* Liquid – Only in combination		500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid preparation	ns.		
METHYL HYDROXYBENZOATE			
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE	00.05	400	/ MI-MV 4
Powder Suspension – Only in combination		100 g 473 ml	✓ MidWest✓ Ora-Plus
			V Ola-Flus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN Suspension		mbination 473 ml	✓ Ora-Blend SF
·		4/3 1111	V Ola-biellu 3F
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE – Only in Suspension		473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM	00.30	4/3 1111	V Ola-Dieliu
Powder – Only in combination	52 50	10 g	✓ MidWest
1 Gwdoi Gilly in Goribination	325.00	100 g	✓ MidWest
Only in children up to 12 years		3	
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxybenzoate	10% solution.		
Liq	11.25	500 ml	✓ Midwest
SODIUM BICARBONATE			
Powder BP – Only in combination		500 g	✓ Midwest
Only in extemporaneously compounded omeprazole and lans	soprazole susp	ension.	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination			
Only in extemporaneously compounded oral liquid preparations.	14.05	E00 ml	√ Midweet
Liq	14.95	500 ml	✓ Midwest
WATER	0.00	1 ml	√ Ton water
Tap - Only in combination	0.00	1 ml	✓ Tap water

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA2204 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA2204 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)	15.38	200 ml OP	✓ Calogen
	38.44	500 ml OP	✓ Calogen
Emulsion (strawberry)	15.38	200 ml OP	✓ Calogen
Oil	37.50	500 ml OP	✓ MCT oil (Nutricia)
MCT Emulsion, 250 ml	143.65	4 OP	✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT - Special Authorit	ty see SA1524 above - Hospital pha	rmacy [HP3]	
Powder	8.95	227 g OP	✓ Resource
		-	Beneprotein
	13.82	225 a OP	✓ Protifar

Subsidy (Manufacturer's Price) Fully Subsidised Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see	SA1095 above -	- Hospital pharm	acy [HP3]	
Liquid	4.65	500 ml OP	✓ Glucerna Select	
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]				
Liquid (strawberry)	2.25	200 ml OP	✓ Diasip	
Liquid (vanilla)	2.10	200 ml OP	✓ Nutren Diabetes	
	2.25		✓ Diasip	

Fat Modified Products

⇒SA2205 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has an inborn error of metabolism. Initial application — (Indications other than errors of inborn metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient has a chyle leak; or
- 2 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

P			
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see Liquid		the previous pa 500 ml OP	
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see Saliquid		ne previous page 500 ml OP	e – Hospital pharmacy [HP3] ✓ Pediasure RTH ✓ Nutrini RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special pharmacy [HP3]	Authority se	ee SA1379 on th	ne previous page – Hospital
Liquid	7.14	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA1 Liquid (strawberry) Liquid (vanilla)	1.90	previous page – 200 ml OP 200 ml OP 500 ml OP	- Hospital pharmacy [HP3] ✓ Fortini ✓ Fortini ✓ Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA137 Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.33 1.33	revious page – F 200 ml OP 200 ml OP 200 ml OP 250 ml OP	Hospital pharmacy [HP3] Pediasure Pediasure Pediasure Pediasure Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Auth pharmacy [HP3]	ority see S	A1379 on the pr	evious page – Hospital
Liquid (unflavoured) Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.90 1.90	200 ml OP 200 ml OP 200 ml OP 200 ml OP	✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 on the Powder		page – Hospital 400 g OP	pharmacy [HP3] Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ORAL FEED 1.8 KCAL/ML - Special Auth	ority see SA1101 above – Hospital	pharmacy [HP	'3]
Liquid	3.31 22	20 ml OP ✓	Nepro HP
			(strawberry)
		•	Nepro HP (vanilla)

Brand or

Fully

	(Manufacturer's Price)	Subsic	lised	Generic
	\$	Per	1	Manufacturer
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110	1 on the previous page	ge – Hospita	al pha	rmacy [HP3]
Liquid, 200 ml bottle	13.24	4 OP	✓ N	lovaSource Renal
Liquid (apricot) 125 ml	13.72	4 OP	✓ R	tenilon 7.5
Liquid (caramel) 125 ml	13.72	4 OP	✓ R	enilon 7.5

Subsidy

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption: or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - Sp. Liquid		e SA1377 abov 1,000 ml OP	
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority se	ee SA1377 above	- Hospital phari	macy [HP3]
Liquid (grapefruit), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	179.46	18 OP	✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see	SA1377 above -	Hospital pharma	acy [HP3]
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Au	thority see SA137	7 above – Hosp	pital pharmacy [HP3]
Liquid	7.47	500 ml OP	✓ Nutrison Advanced
			Pentisorh

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

continued...

(Mar	Subsidy nufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML - Special Authority see SA1196 on the previous page - Hospital pharmacy [HP3]

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

continued...

	Subsidy		Fully	Brand or	
(Man	ufacturer's Price)	Subs	sidised	Generic	
	\$	Per	✓	Manufacturer	

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or

continued...

rsidy F Irer's Price) Subsidi	. ,	rand or eneric
 \$ Per	✓ Ma	anufacturer

- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see	e SA1859 on page 282 – Hos	pital pharmacy	/ [HP3]
Liquid	2.17	250 ml OP	 Ensure Plus HN
	8.68	,000 ml OP	 Ensure Plus HN
			RTH
	9.00		✓ Nutrison Energy

			_
	Subsidy		Fully Brand or
	(Manufacturer's F	Price) Subsi	idised Generic
	\$	Per	 Manufacturer
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on	nage 282 - Ho	snital nharmacy	[HP3]
Liquid		250 ml OP	✓ Isosource Standard
Liquid	6.56	1,000 ml OP	✓ Osmolite RTH
		1,000 IIII OP	• • • • • • • • • • • • • • • • • • • •
	6.90		✓ Nutrison RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authorit	y see SA1859 o	on page 282 – H	ospital pharmacy [HP3]
Liquid	9.05	1,000 ml OP	✓ Nutrison
			800 Complete
			Multi Fibre
ENTERAL FEED WITH FIRRY 1 I/CAL/MI Chasial Authority a	00 CA10E0 on r	000 Lloom	
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority s		•	, ,, ,
Liquid		1,000 ml OP	Jevity RTH
	7.21		✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority	see SA1859 on	page 282 - Hos	spital pharmacy [HP3]
Liquid		1,000 ml OP	✓ Jevity Plus RTH
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority		nago 202 Hos	•
Liquid	8.08	1,000 ml OP	✓ Jevity HiCal RTH
			✓ Nutrison Energy
			Multi Fibre
ORAL FEED (POWDER) - Special Authority see SA1859 on page	ge 282 – Hospita	al pharmacy [HP	23]
Powder (chocolate)	•	840 g OP	✓ Sustagen Hospital
		- · · · · · · · · · · · · · · · · · · ·	Formula
	26.00	850 g OP	✓ Ensure
Davidar (vanilla)			
Powder (vanilla)	14.00	840 g OP	✓ Sustagen Hospital
			Formula Active
	26.00	850 g OP	✓ Ensure
ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on pa	age 282 – Hospi	ital pharmacy [H	P31
Additional subsidy by endorsement is available for patients b			
epidermolysis bullosa, or as exclusive enteral nutrition in chile			
disease, or for patients with COPD and hypercapnia, defined			
endorsed accordingly.	as OOZ value c	DAGGGGIIIG DOITIII	ing. The prescription must be
· · · · · · · · · · · · · · · · · · ·			
Liquid (banana) – Higher subsidy of up to \$1.76 per 200 ml	0.70	000 0.D	
with Endorsement		200 ml OP	
	(1.56)		Ensure Plus
	(1.76)		Fortisip
Liquid (chocolate) - Higher subsidy of up to \$1.76 per 200 m	ıl		
with Endorsement	0.72	200 ml OP	
	(1.56)		Ensure Plus
	(1.76)		Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.56 per 200 i			- · · · •
with Endorsement		200 ml OP	
With Endorsement		200 1111 01	Ensure Plus
	(1.56)		Erisure Pius
Liquid (strawberry) – Higher subsidy of \$1.76 per 200 ml with			
Endorsement		200 ml OP	
	(1.76)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.76 per 200 ml w	ith		
Endorsement	0.85	237 ml OP	
	(1.65)	-	Ensure Plus
	0.72	200 ml OP	
	(1.56)	200 1111 01	Ensure Plus
	(1.76)		Fortisip

Subsidy	Ful	ly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per •	Manufacturer	

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 282 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.76 per 200 ml with

Liquid (chocolate) — Higher Subsidy of \$1.70 per 200 Hil With			
Endorsement	0.72	200 ml OP	
	(1.76)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.76 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.76)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.76 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.76)		Fortisio Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully sidised	Brand or Generic Manufacturer
ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 o	n the previous pa	age – Hospital p	harma	cy [HP3]
Liquid	6.82	500 ml OP	✓ N	lutrison
	13.64	1,000 ml OP	✓ E	Concentrated Insure Two Cal HN RTH
ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the	e previous page	- Hospital phari	macy [H	HP3]
Additional subsidy by endorsement is available for patients I epidermolysis bullosa. The prescription must be endorsed a Liquid (vanilla) — Higher subsidy of \$2.34 per 200 ml with	U	hrough a feedin	g tube,	or who have severe
Endorsement	0.96	200 ml OP		
	(2.34)		Т	wo Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER - Special Authority see SATTU6 at	ove – Hospitai pnarmacy	[HP3]	
Powder	8.29	300 g OP	✓ Nutilis
	24.00	380 g OP	✓ Aptamil Feed
			Thickener

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA	1729 above – Hospital բ	oharmacy [HP3]	
Powder	2.81	1,000 g OP	
	(5.15)		Healtheries Simple
			Baking Mix

	Subsidy	Ful	lv Brand or
	(Manufacturer's Price		,
	\$		Manufacturer
GLUTEN FREE BREAD MIX - Special Authority see SA1729 or	n the previous page -	- Hospital pharr	macv [HP3]
Powder		.000 g OP	, 1
	(7.32)		NZB Low Gluten
			Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 on the	previous page - Hos	spital pharmacy	[HP3]
Powder	5.62 2,	,000 g OP	
	(18.10)		Horleys Flour
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page - Hos	spital pharmacy	[HP3]
Buckwheat Spirals	2.00 2	250 g OP	
•	(3.11)	•	Orgran
Corn and Vegetable Shells	2.00 2	250 g OP	
	(2.92)		Orgran
Corn and Vegetable Spirals		250 g OP	
	(2.92)		Orgran
Rice and Corn Lasagne Sheets		200 g OP	
Discount Own Massauri	(3.82)	250 - OD	Orgran
Rice and Corn Macaroni		250 g OP	0
Disc and Care Danna	(2.92)	050 ~ OD	Orgran
Rice and Corn Penne	2.00 2 (2.92)	250 g OP	Orgran
Rice and Maize Pasta Spirals	` '	250 g OP	Olylali
Tilde and Maize Lasta Spirais	(2.92)	250 g Oi	Orgran
Rice and Millet Spirals		250 g OP	Orgium
	(3.11)	-00 g 0.	Orgran
Rice and corn spaghetti noodles		375 g OP	- 3 ··
1 0	(2.92)	Ü	Orgran
Vegetable and Rice Spirals	2.00 2	250 g OP	v
	(2.92)	-	Orgran
Italian long style spaghetti		220 g OP	
	(3.11)		Orgran

(Orgran Buckwheat Spirals to be delisted 1 July 2025)

(Orgran Corn and Vegetable Shells to be delisted 1 July 2025)

(Orgran Corn and Vegetable Spirals to be delisted 1 July 2025)

(Orgran Rice and Corn Lasagne Sheets to be delisted 1 July 2025)

(Orgran Rice and Corn Macaroni to be delisted 1 July 2025)

(Orgran Rice and Corn Penne to be delisted 1 July 2025)

(Orgran Rice and Maize Pasta Spirals to be delisted 1 July 2025)

(Orgran Rice and Millet Spirals to be delisted 1 July 2025)

(Orgran Rice and corn spaghetti noodles to be delisted 1 July 2025)

(Orgran Vegetable and Rice Spirals to be delisted 1 July 2025)

(Orgran Italian long style spaghetti to be delisted 1 July 2025)

Foods And Supplements For Inherited Metabolic Disease

⇒SA2357 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where patient requires dietary management of inherited metabolic disorders.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

Supplements For Homocystinuria

AMINOACID FORMULA WITHOUT METHIONINE - Special Authority see SA2357 on the previous page - Hospital pharmacy [HP3]

Powder (neutral), 36 g sachets	750.30	30	HCU Anamix Junior
Powder, 12.5 g sachets	349.65	30	✓ HCU Explore 5
Powder, 25 g sachets		30	✓ HCU Express 15
Powder (neutral), can	480.42	500 g OP	✓ XMET Maxamum
Powder (unflavoured), can		400 g OP	✓ HCU Anamix Infant
Liquid (juicy berries), 125 ml bottle	1,684.80	30	✓ HCU Lophlex LQ
Liquid (orange), 125 ml bottle		36	✓ HCU Anamix Junior
. , • ,			LQ

Supplements For MSUD and short chain enoyl coA hydratase deficiency

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA2357 on the previous page - Hospital pharmacy [HP3]

rioophai priarriaoy [rir o]			
Powder (neutral) 36 g sachets	750.00	30	MSUD Anamix Junior
Powder, 12.5 g sachets	349.65	30	✓ MSUD Explore 5
Powder, 25 g sachets	1,048.95	30	✓ MSUD Express 15
Powder (neutral), can	454.71	500 g OP	✓ MSUD Maxamum
Powder (orange), can	454.71	500 g OP	MSUD Maxamum
Powder (unflavoured), can	260.00	400 g OP	MSUD Anamix Infant
Liquid (orange) 125 ml bottles	941.40	36	MSUD Anamix Junior LQ
Liquid (juicy berries) 125 ml pouches	1,684.80	30	✓ MSUD Lophlex LQ 20

✓ fully subsidised 289

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Supplements For PKU

• •			
AMINOACID FORMULA WITHOUT PHENYLALANINE	- Special Authority see	SA2357 on page	288 – Hospital pharmacy [HP3]
Tabs		75 OP	✓ Phlexy 10
Powder (Lemon), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Neutral), 12.5 g sachets	220.88	30	✓ PKU Explore 5
Powder (Neutral), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Orange), 25 g sachets	441.75	30	✓ PKU Explore 10
Powder (Orange), 34 g sachets	883.50	30	✓ PKU Express 20
Powder (Raspberry), 25 g sachets	441.75	30	✓ PKU Explore 10
Powder (Tropical), 34 g sachets		30	✓ PKU Express 20
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (neutral) 36 g sachets	393.00	30	PKU Anamix Junior
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior Orange
Powder (unflavoured) 12.5 g sachets	234.00	30	✓ PKU First Spoon
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)		500 g OP	✓ XP Maxamum
Liquid (berry)		125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex Sensation 20
Powder (neutral), 400 g can	715.16	4 OP	✓ PKU Start
Liquid (juicy berries) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml		30 OP	✓ PKU Lophlex LQ 20
			•

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully	Brand or Generic Manufacturer
OLYGOMA OPOPERTIDE AND AMINO ACID CONTAING COM	<u> </u>		. A catha a si	
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME page 288 – Hospital pharmacy [HP3]	PHENYLALANINE	– Special	Authori	ty see SA2357 on
Powder (Banana) 35 g sachets	030.00	30	✓ F	NII
Fowder (Danana) 33 y Sacriets	930.00	30	• -	sphere20 Banana
Powder (Berry), 20 g sachets	440.20	60	./ 0	'KU Restore
Fowder (Derry), 20 g Sacriets	449.20	00	• -	Powder
Powder (Chocolate) 32 g sachets	808 56	30	/ 0	KU Build
1 Owder (Oriocolate) 32 g sacriets	090.30	50	• 1	20 Chocolate
Powder (Chocolate) 35 g sachets	930.00	30	√ F	
1 Owder (Orlocoldie) oo g saariote		00	• •	sphere20 Chocolate
				opiioi ozo oiioooiato
Powder (Lemon) 35 g sachets	930.00	30	✓ F	PKU
				sphere20 Lemon
Powder (Lemonade) 33.4 g sachets	936.00	30	✓ F	KU GMPro Ultra
				Lemonade
Powder (Neutral), 15 g sachets	449.28	30	✓ F	KU Build 10
Powder (Orange), 20 g sachets	449.28	60	✓ F	KU Restore
				Powder
Powder (Raspberry Lemonade) 31 g sachets	898.56	30	✓ F	KU Build
				20 Raspberry
			_	Lemonade
Powder (Smooth) 31 g sachets	898.56	30	✓ F	KU Build
			_	20 Smooth
Powder (Vanilla) 33 g sachets		30		KU Build 20 Vanilla
Powder (neutral), 40 g sachets		30		Slytactin Bettermilk
Powder (unflavoured) 12.5 g sachets		30		KU GMPro Mix-In
Powder (vanilla) 33.4 g sachets	936.00	30	✓ F	KU GMPro Ultra
December (Ded December) OF a contrate	000 00	00		Vanilla
Powder (Red Berry) 35 g sachets	930.00	30	• 1	KU sphere20 Red
Davidas (Vanilla) OF a analysta	000 00	00		Berry
Powder (Vanilla) 35 g sachets	930.00	30	✓ F	
Liquid (noutral) QEO ml corton	000.00	18	./ -	sphere20 Vanilla PKU GMPro LQ
Liquid (neutral), 250 ml carton Liquid (original), 250 ml carton		16 30 OP		KU Glytactin RTD
Liquid (original), 250 mi carton	004.45	30 OF	• -	15
Liquid (Coffee Mocha), 250 ml carton	684.45	30 OP	/ 0	KU Glytactin RTD
Liquid (Conee Mocha), 250 mi Carton	004.45	30 OF	• -	15 Lite
Liquid (chocolate), 250 ml carton	684.45	30 OP	/ 0	KU Glytactin RTD
Liquia (viivoolate), 200 IIII valtoi!		00 01	• [15
Liquid (vanilla), 250 ml carton	684 45	30 OP	√ □	KU Glytactin RTD
Elquia (variilla), 200 IIII varioii		00 01	• 1	15 Lite

Foods

LOW PROTEIN BAKING MIX - Special Authority see SA2357 on pag	e 288 –	Hospital pharmacy	/ [HP3]	
Powder	8.55	500 g OP	✓ Loprofin M	İΧ

✓ fully subsidised 291

	Subsidy		Fully Brand or
	(Manufacturer's P		sidised Generic
	\$	Per	✓ Manufacturer
LOW PROTEIN PASTA - Special Authority see SA2357 on page 1	age 288 – Hospital		
Animal shapes	12.39	500 g OP	✓ Loprofin
Lasagne	6.19	250 g OP	✓ Loprofin
Low protein rice pasta	12.39	500 g OP	✓ Loprofin
Macaroni	6.19	250 g OP	✓ Loprofin
Penne	12.39	500 g OP	✓ Loprofin
Spaghetti	12.39	500 g OP	✓ Loprofin
Spirals	12.39	500 g OP	✓ Loprofin
Supplements for Tyrosinaemia			
AMINOACID FORMULA WITHOUT PHENYLALANINE AND T Dharmacy [HP3]	YROSINE - Specia	al Authority see	e SA2357 on page 288 – Hosp
Powder (Neutral), 12.5 g sachets	3/0 65	30	✓ TYR Explore 5
Powder (neutral) 36 g sachets		30	✓ TYR Anamix Junior
, , ,			
Powder, can		400 g OP	✓ TYR Anamix Infant
Liquid (juicy berries) 125 ml pouches		30	✓ TYR Lophlex LQ 20
Liquid (orange) 125 ml bottle	941.40	36	✓ TYR Anamix Junior LQ
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOI	ME TYROSINE ANI	Ο ΡΗΕΝΙΎΙ ΔΙ Δ	
SA2357 on page 288 – Hospital pharmacy [HP3]	WIL THOUSINE AIN) I IILINI LALA	ANINE - Special Authority see
Powder (Red Berry), 35 g sachets	1 200 60	30	✓ TVP Sphare 20
		30	✓ TYR Sphere 20
Powder (Vanilla), 35 g sachets	1,396.00	30	✓ TYR Sphere 20
Supplements for Organic Acidaemias			
AMINOACID FORMULA WITHOUT ISOLEUCINE, METHIONI	INE, THREONINE A	ND VALINE -	- Special Authority see SA2357
on page 288 – Hospital pharmacy [HP3]			
Powder, can	260.00	400 g OP	MMA/PA Anamix Infant
AMINOACID FORMULA WITHOUT METHIONINE, THREONI Hospital pharmacy [HP3]	NE AND VALINE -	Special Autho	rity see SA2357 on page 288 -
Powder (neutral), 18 g sachets	750.30	30	MMA/PA Anamix Junior
Powder, 12.5 g sachets	349.65	30	✓ MMA/PA Explore 5
Powder, 25 g sachets	1,048.95	30	✓ MMA/PA Express 15
Supplements for Glutaric Aciduria type 1			
AMINOACID FORMULA WITHOUT LYSINE - Special Author	rity see SA2357 on t	page 288 – Hos	spital pharmacy [HP3]
Powder (neutral), 18 g sachets		30	✓ GA1 Anamix Junior
Powder, 12.5 g sachets		30	✓ GA Explore 5
Powder, can		400 g OP	✓ GA1 Anamix Infant
1 Owder, editional and a second a second and	200.00	400 g O1	• WAT AHUMIN IIIUM
Supplements for Glycogen Storage Disease			
HIGH AMYLOPECTIN CORN-STARCH – Special Authority se Powder, 60 g sachets		288 – Hospita 30	ll pharmacy [HP3] ✓ Glycosade
Single dose amino acids			
ARGININE - Special Authority see SA2357 on page 288 - Ho	ospital pharmacy [Hi	P3]	
		00	/ A
Powder, 4 g sachets	211.45	30	✓ Arginine2000

				_
Subsidy (Manufacturer's \$	Price) Per	Fully Subsidised	Generic	
CITRULLINE – Special Authority see SA2357 on page 288 – Hospital pharmacy Powder, 4 g sachets211.45	y [HP3] 30	✓	Citrulline1000	
ISOLEUCINE - Special Authority see SA2357 on page 288 - Hospital pharmac Powder, 4 g sachets141.05	y [HP3] 30	/	Isoleucine50	
LEUCINE – Special Authority see SA2357 on page 288 – Hospital pharmacy [H Powder, 4 g sachets141.05	IP3] 30	/	Leucine100	
PHENYLALANINE – Special Authority see SA2357 on page 288 – Hospital pha Powder, 4 g sachets141.05	rmacy [HP3 30	-	Phenylalanine50	
TYROSINE – Special Authority see SA2357 on page 288 – Hospital pharmacy Powder, 4 g sachets211.45	[HP3] 30	/	Tyrosine1000	
VALINE – Special Authority see SA2357 on page 288 – Hospital pharmacy [HP Powder, 4 g sachets141.05	3]	✓	Valine50	
Other Fat Modified Products				
ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCERIDES - Special pharmacy [HP3]	Authority s	ee SA2357	7 on page 288 – Hosp	ital
Powder (neutral), 100 g sachets47.01	10	✓	Emsogen	
Carbohydrate and Fat with added vitamins and minerals				
PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDRATE, FAT WITH Authority see SA2357 on page 288 – Hospital pharmacy (HP3)	ADDED VIT	TAMINS A	ND MINERALS - Spe	ecial
Powder (neutral), can	400 g C	OP 🗸	Energivit	
Essential Amino Acids				
ESSENTIAL AMINOACID FORMULA - Special Authority see SA2357 on page Powder (neutral), can313.73	288 – Hosp 200 g C	•	acy [HP3] Essential Amino Acid Mix	
Infant Formulae				

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]

Powder46.18 400 g OP ✓ Locasol

✓ fully subsidised 293

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA - Special Authority see SA209	2 below – Hospital pharn	nacy [HP3]	
Powder	43.60	400 g OP	✓ Alfamino✓ Alfamino Junior
Powder (unflavoured)	55.61	400 g OP	✓ Neocate Gold ✓ Neocate Junior Unflavoured
	65.72		✓ Neocate SYNEO✓ Elecare✓ Elecare LCP
Powder (vanilla)	55.61	400 g OP	✓ Neocate Junior Vanilla
	65.72		✓ Elecare

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

Α

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency: or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

number; or

2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or

2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

continued...

✓ fully subsidised 295

bsidy	Fully	Brand or
turer's Price) Subsid	dised	Generic
 \$ Per	✓	

continued...

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA - Special Authority see SA1953 below - Hospital pharmacy [HP3]

Liquid 1 kcal/ml12.44	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml18.66	500 ml OP	✓ Nutrini Peptisorb
		Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA -	 Special Authority see SA1557 on 	the next page - I	Hospital pharmacy [HP3]
Powder	18.10	450 g OP	✓ Pepti-Junior
	36.20	900 g OP	✓ Allerpro Syneo 1
			✓ Allerpro Syneo 2

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 12 Fither
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis: or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula: and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3] Liquid.......2.80 125 ml OP ✓ Infatrini

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

continued...

✓ fully subsidised 297



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
<u> </u>	Per		Manufacturer

continued...

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA119	7 above – Hospita	al pharmacy [HP3]
Powder (unflavoured)36.92	300 g OP	✓ KetoCal 4:1
	_	✓ Ketocal 3:1
Powder (vanilla)36.92	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

	Danish	strain '	1331.	live a	ttenua	ted. v	vial with	diluent.	
			,			,			

uent......0.00 10 **✓ <u>BCG Vaccine AJV</u>**

COVID-19 VACCINE - [Xpharm]

10 ✓ Comirnaty Omicron (JN.1)

Up to three doses for previously unvaccinated children aged 6 months - 4 years at high risk of severe illness.

Inj 10 mcg bretovameran per 0.3 ml, 0.48 ml vial; paediatric	
vaccine, light blue cap0	.00

10 ✓ Comirnaty Omicron (JN.1)

Either:

- 1) One dose for previously unvaccinated children aged 5-11 years old; or
- 2) Up to three doses for immunocompromised children aged 5-11 years old.

10 Comirnaty Omicron

(JN.1)

Any of the following:

- 1) One dose for previously unvaccinated people aged 12-15 years old: or
- 2) Up to three doses for immunocompromised people aged 12-15 years old; or
- 3) Up to two doses for previously unvaccinated people 16-29 years old; or
- 4) Up to four doses for people aged 16-29 at high risk of severe illness; or
- 5) One dose for previously unvaccinated people aged 30 and older; or
- 6) One additional dose every 6 months for previously vaccinated people aged 30 years and over additional dose is given at least 6 months after last dose.

Subsid	dy Full	/ Brand or
(Manufacture	r's Price) Subsidise	d Generic
\$	Per 💌	Manufacturer

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy: or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
 - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old: or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.

syringe	0.00	10	✓ Roostriy
haemagglutinin and 2.5 mcg pertactin in 0.5 ml prefilled			
pertussis toxoid, 8 mcg pertussis filamentous			
Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg			

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for any of the following:
 - 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
 - 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation: or
 - 3) An additional four doses (as appropriate) are funded for (re-)immunisation for people post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 4) Five doses will be funded for children requiring solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis and polio vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis and polio vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

10 ✓ Infanrix IPV

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

- A) Funded for children meeting any of the following criteria
 - 1) Up to four doses for children under the age of 10 years for primary immunisation; or
 - 2) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
 - 3) An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens: or
 - 4) Up to five doses for children under the age of 10 years receiving solid organ transplantation.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Diphtheria, tetanus, pertussis, polio. hepatitis B and haemophilus influenzae type b vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type b vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen, 10mcg H. influenzae type b with tetanus toxoid

✓ Infanrix-hexa 10

		Subsidy (Manufacturer's Price)	D	Fu Subsidise	ed Ge	nd or neric
		\$	Per		IVIa	nufacturer
	OPHILUS INFLUENZAE TYPE B VACCINE					
	Only on a prescription					
b) c)	No patient co-payment payable					
,	A) One dose for people meeting any of the following:					
	 For primary vaccination in children; or 					
	 An additional dose (as appropriate) is funded transplantation, or chemotherapy; functional a 	` '				
	transplant, pre or post cochlear implants, rena	I dialysis and other se	vere	ly immur	nosuppre	ssive regimens; or
	 For use in testing for primary immunodeficient physician or paediatrician. 	cy diseases, on the re	comi	mendatio	n of an i	nternal medicine
	B) Contractors will be entitled to claim payment from the	ne Funder for the supp	oly of	Haemor	hilus inf	luenzae type b
	vaccine to people eligible under the above criteria p					, ,
	for subsidised immunisation, and they may only do in the Pharmaceutical Schedule.	so in respect of the H	aemo	ophilus in	ıfluenzae	type b vaccine list
	 C) Contractors may only claim for populations within the sub-set of the population described in paragraph A 		ered	by their	contract,	which may be a
Ini	10 mcg vial with diluent syringe		1		Act-H	IR
			•		AUCH	<u></u>
	ITIS A VACCINE – [Xpharm]					
	nded for patients meeting any of the following criteria:					
) Two vaccinations for use in transplant patients; or	linnana: or				
	 Two vaccinations for use in children with chronic liver of the content of the chronic liver of the content of the chronic liver of the chronic					
J	7 One access a vaccine for close contacts of known nepal	itio / t bases.				

1

✓ <u>Havrix 1440</u> ✓ Havrix Junior

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
LIEDATITIO D DECOMPINIANT VACCINE	IV. I1					

HEPATITIS B RECOMBINANT VACCINE - [Xpharm]

Funded for patients meeting any of the following criteria:

- 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4) for HIV positive patients; or
- 5) for hepatitis C positive patients; or
- 6) for patients following non-consensual sexual intercourse; or
- 7) for patients prior to planned immunosuppression for greater than 28 days; or
- 8) for patients following immunosuppression; or
- 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury.

- Funded for patients meeting any of the following criteria:
 - 1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
 - 2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
 - 3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
 - 4) for HIV positive patients; or
 - 5) for hepatitis C positive patients; or
 - 6) for patients following non-consensual sexual intercourse; or
 - 7) for patients prior to planned immunosuppression for greater than 28 days; or
 - 8) for patients following immunosuppression; or
 - 9) for solid organ transplant patients; or
- 10) for post-haematopoietic stem cell transplant (HSCT) patients; or
- 11) following needle stick injury; or
- 12) for dialysis patients; or
- 13) for liver or kidney transplant patients.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV]

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- a) A) Any of the following:
 - 1) Maximum of two doses for children aged 14 years and under; or
 - 2) Maximum of three doses for people meeting any of the following criteria:
 - 1) People aged 15 to 26 years inclusive; or

People aged 9 to 26 years inclusive who have

- 1) Confirmed HIV infection; or
- 2) Received a transplant (including stem cell): or
- 3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy
- B) Contractors will be entitled to claim payment from the Funder for the supply of Human papillomavirus vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Human papillomavirus vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

10 ✓ Gardasil 9

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
INFLUENZA VACCINE Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	120.00	10		fluvac Tetra (2025 formulation)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

- a) Maximum of 1 inj per prescription
- b) Only on a prescription
- c) No patient co-payment payable
- ď

A) INFLUENZA VACCINE

is available each year for patients who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) Down syndrome, or
 - vii) are pregnant; or
- c) children 4 years of age and under who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- d) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,			
Rubella virus 1,000 CCID50; prefilled syringe/ampoule of			
diluent 0.5 ml	0.00	10	✓ Priorix

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE

Inj 10 mcg of each meningococcal polysaccharide conjugated

a) Only on a prescription

b) No patient co-payment payable

C)

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients
 with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post
 solid organ transplant; or

MenQuadfi

- 2) One dose for close contacts of meningococcal cases of any group; or
- 3) One dose for person who has previously had meningococcal disease of any group; or
- 4) A maximum of two doses for bone marrow transplant patients; or
- 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or
 - 2) One dose for individuals who turn 13 years of age while living in boarding school hostels.
- C) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal A, C, Y and W-135 vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal A, C, Y and W-135 vaccine listed in the Pharmaceutical Schedule.
- D) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-B above.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 5 mcg of each meningococcal polysaccharide conjugated to a total of approximately 44 mcg of tetanus toxoid carrier

- A) Both:
 - 1) The child is under 12 months of age; and
 - 2) Any of the following:
 - A maximum of three doses (dependant on age at first dose) for patients pre- and post- splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post- solid organ transplant; or
 - A maximum of three doses (dependant on age at first dose) for close contacts of meningococcal cases of any group; or
 - A maximum of three doses (dependant on age at first dose) for child who has previously had meningococcal disease of any group; or
 - 4) A maximum of three doses (dependant on age at first dose) for bone marrow transplant patients; or
 - A maximum of three doses (dependant on age at first dose) for child pre- and post-immunosuppression*.

Note: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	(Manufacturer's Price) Subsidised		Generic	
\$	Per	✓	Manufacturer	

MENINGOCOCCAL B MULTICOMPONENT VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c) Any of the following:
 - A) Three doses for children up to 12 months of age (inclusive) for primary immunisation; or
 - B) Up to three doses (dependent on age at first dose) for a catch-up programme for children from 13 months to 59 months of age (inclusive) for primary immunisation, from 1 March 2023 to 31 August 2025; or
 - C) Both:
 - 1) Person is one year of age or over; and
 - 2) Any of the following:
 - i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
 - ii) up to two doses for close contacts of meningococcal cases of any group; or
 - iii) up to two doses for person who has previously had meningococcal disease of any group; or
 - iv) up to two doses for bone marrow transplant patients; or
 - v) up to two doses for person pre- and post-immunosuppression*; or
 - D) Both:
 - 1) Person is aged between 13 and 25 years (inclusive); and
 - 2) Either:
 - Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences or prisons; or
 - ii) Two doses for individuals who turn 13 years of age while living in boarding school hostels.
 - E) Contractors will be entitled to claim payment from the Funder for the supply of Meningococcal B multicomponent vaccine to patients eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Meningococcal B multicomponent vaccine listed in the Pharmaceutical Schedule.
 - F) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A-D above.

*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Any of the following:
 - 1) A course of three doses for previously unvaccinated children up to the age of 59 months inclusive; or
 - Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10: or
 - 3) Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high risk children aged under 5 years with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) primary immune deficiencies; or
 - c) HIV infection: or
 - d) renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) cochlear implants or intracranial shunts; or
 - g) cerebrospinal fluid leaks; or
 - n) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) cardiac disease, with cyanosis or failure; or
 - diabetes; or
 - m) Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
 - 4) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
 - For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Pneumococcal (PCV13) conjugate vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Pneumococcal (PCV13) conjugate vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml

	Subsidy	Fully	Brand or
	Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - [Xp	harm]		
Either:			

- Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 2) All of the following:
 - a) Patient is a child under 18 years for (re-)immunisation; and
 - b) Treatment is for a maximum of two doses; and
 - c) Any of the following:
 - i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - ii) with primary immune deficiencies; or
 - iii) with HIV infection; or
 - iv) with renal failure, or nephrotic syndrome; or
 - v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant);
 or
 - vi) with cochlear implants or intracranial shunts; or
 - vii) with cerebrospinal fluid leaks; or
 - viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - x) pre term infants, born before 28 weeks gestation; or
 - xi) with cardiac disease, with cyanosis or failure; or
 - xii) with diabetes: or
 - xiii) with Down syndrome; or

Ini 575 mcg in 0.5 ml prefilled syringe (25 mcg of each

xiv) who are pre-or post-splenectomy, or with functional asplenia.

23 pneumococcal serotype)	1	✓ Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm]		
Up to three doses for patients meeting either of the following:		
1) For partially vaccinated or previously unvaccinated individuals; or		
For revaccination following immunosuppression.		

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

BOTAVIBUS OBAL VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Maximum of two doses for people meeting the following:
 - 1) first dose to be administered in infants aged under 14 weeks of age; and
 - 2) no vaccination being administered to children aged 24 weeks or over.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Rotavirus oral vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Rotavirus oral vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, squeezable tube	0.00	10	Rotarix
Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	✓ Rotarix

Subsidy (Manufacturer's Price)	Fully e) Subsidised Per		Brand or Generic Manufacturer
φ	rei		Manuacturer

VARICELLA VACCINE [CHICKENPOX VACCINE]

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Either:
 - 1) Maximum of one dose for primary vaccination for either:
 - a) Any infant born on or after 1 April 2016; or
 - For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
 - 2) Maximum of two doses for any of the following:
 - a) Any of the following for non-immune individuals:
 - i) with chronic liver disease who may in future be candidates for transplantation; or
 - ii) with deteriorating renal function before transplantation; or
 - iii) prior to solid organ transplant; or
 - iv) prior to any elective immunosuppression*; or
 - v) for post exposure prophylaxis who are immune competent inpatients; or
 - b) For individuals at least 2 years after bone marrow transplantation, on advice of their specialist; or
 - c) For individuals at least 6 months after completion of chemotherapy, on advice of their specialist; or
 - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
 - e) For individuals with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
 - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
 - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.
- B) Contractors will be entitled to claim payment from the Funder for the supply of Varicella vaccine [Chickenpox vaccine] vaccine to people eligible under the above criteria pursuant to their contract with Health New Zealand (Health NZ) for subsidised immunisation, and they may only do so in respect of the Varicella vaccine [Chickenpox vaccine] listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs A above.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	(I	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
ARICE	ELLA ZOSTER VACCINE [SHINGLES VACCINE]				
a) (Only on a prescription				
b) I	No patient co-payment payable				
c)					
	 A) Funded for patients meeting the following criteria: 				
	1) Either:				
	 Two doses for all people aged 65 years, or 				
	Two doses for people 18 years of age or o	der with any of the	follow	/ing:	
	 a) pre- and post-haematopoietic stem c 	ell transplant or cell	ular th	nerapy; or	
	b) pre- or post-solid organ transplant; or				
	 c) haematological malignancies; or 				
	 d) people living with poorly controlled H 				
	 e) planned or receiving disease modifying 	ng anti-rheumatic dr	rugs (DMARDs -	targeted synthetic,
	biologic, or conventional synthetic) fo	r polymyalgia rheun	natica	ı, systemic lı	upus erythematosus or
	rheumatoid arthritis; or				
	f) end stage kidney disease (CKD 4 or	5); or			
	g) primary immunodeficiency				
	B) Contractors will be entitled to claim payment from the				
	vaccine) to patients eligible under the above criteria p	ursuant to their cont	tract v	vith Health N	New Zealand (Health NZ)
	for subsidised immunisation, and they may only do so	in respect of the Va	aricell	a zoster vac	cine [Shingles vaccine]
	listed in the Pharmaceutical Schedule				

C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be

_		_	
Diag	MACHIC	Aganta	
		: Agents	

a sub-set of the population described in paragraph A above. Inj 50 mcg per 0.5 ml vial plus vial.......0.00

✓ Shingrix

✓ Shingrix

1

- Symbols -	Albendazole		Amzoate	
3TC111	Albey	256	Anaesthetics	122
- A -	Albustix	81	Anafranil	127
A-Scabies72	Alchemy Oxaliplatin	153	Anagrelide hydrochloride	
Abacavir sulphate111	Alchemy Oxybutynin		Analgesics	123
Abacavir sulphate with	Aldurazyme	<mark>27</mark>	Anastrozole	175
lamivudine111	Alecensa	164	Anatrole	175
Abacavir/Lamivudine Viatris111	Alectinib	164	Anoro Ellipta	260
Abilify Maintena136	Alendronate sodium	116	Antabuse	148
Abilify Maintena S29136	Alendronate sodium with		Antacids and Antiflatulents	6
Abiraterone acetate172	colecalciferol	116	Anthelmintics	93
Acarbose11	Alfacalcidol	31	Antiacne Preparations	66
Accarb11	Alfamino	294	Antiallergy Preparations	255
Acetazolamide268	Alfamino Junior	294	Antianaemics	35
Acetec44	Alginic acid	6	Antiandrogen Oral	
Acetic acid with hydroxyquinoline and	Alglucosidase alfa	24	Contraceptives	79
ricinoleic acid79	Alkeran	153	Antiarrhythmics	
Acetylcysteine271	Allerfix	268	Antibacterials	
Aci-Jel79	Allerpro Syneo 1	296	Antibacterials Topical	66
Aciclovir	Allerpro Syneo 2	296	Anticholinergic Agents	
Infection106	Allersoothe	257	Anticholinesterases	
Sensory266	Allmercap	155	Antidepressants	127
Acidex6	Allopurinol		Antidiarrhoeals	
Acipimox52	Almarytm	47	Antiepilepsy Drugs	129
Acitretin72	Alpha-Adrenoceptor Blockers		Antifibrinolytics, Haemostatics and	
Act-HIB302	Alpha-Keri Lotion		Local Sclerosants	
Actemra233	Alphamox 125		Antifibrotics	261
Actinomycin D157	Alphamox 250	96	Antifungals	
Actrapid10	Alprolix		Antifungals Topical	
Actrapid Penfill10	Alu-Tab	6	Antihistamines	
Acupan123	Aluminium hydroxide		Antihypotensives	
Adalimumab (Amgevita)183	Alyacen	78	Antimalarials	103
Adalimumab (Humira - Alternative	Amantadine hydrochloride		Antimigraine Preparations	
brand) 192	Ambrisentan	55	Antinausea and Vertigo Agents	132
Adapalene66	Ambrisentan Viatris	55	Antipruritic Preparations	
Adcetris201	Amgevita	183	Antipsychotics	
ADR Cartridge 1.821	Amiloride hydrochloride		Antiretrovirals	
Adrenaline	Amiloride hydrochloride with		Antirheumatoid Agents	
Cardiovascular54	furosemide	<u>50</u>	Antispasmodics and Other Agents	
Respiratory255	Amiloride hydrochloride with		Altering Gut Motility	
Advantan69	hydrochlorothiazide	50	Antithrombotic Agents	39
Advate38	Aminophylline		Antithymocyte globulin	
Adynovate38	Amiodarone hydrochloride		(equine)	183
Afinitor251	Amisulpride	134	Antitrichomonal Agents	
Aflibercept198	Amitriptyline		Antituberculotics and	
AFT-Pyrazinamide106	Amlodipine		Antileprotics	104
Agents Affecting the	Amorolfine		Antiulcerants	
Renin-Angiotensin System 44	Amoxicillin	96	Antivirals	
Agents for Parkinsonism and Related	Amoxicillin with clavulanic acid		Anxiolytics	
Disorders 121	Amoxiclav Devatis Forte		Anzatax	
Agents Used in the Treatment of	Amphotericin B		Apidra	
Poisonings	Amsacrine		Apidra SoloStar	
Agrylin156	AmsaLyo		APO Clomipramine	
Albalon270	Amsidine		APO-Atomoxetine	

Apo-Azithromycin	94	Atezolizumab	243	Benzydamine hydrochloride	3
APO-Candesartan HCTZ		ATGAM		Benzylpenicillin sodium [Penicillir	
16/12.5	45	Ativan	138	G]	
APO-Candesartan HCTZ		Atnahs Olsalazine	8	Beta Cream	
32/12.5	45	Atomoxetine		Beta Ointment	
Apo-Temozolomide		Atorvastatin		Beta Scalp	
Apomorphine hydrochloride		Atropine sulphate		Beta-Adrenoceptor Agonists	
Aprepitant		Cardiovascular	46	Beta-Adrenoceptor Blockers	
Apresoline		Sensory		Beta-hCG low sensitivity urine tes	
Aptamil Feed Thickener		Atropt		kit	
Aqueous cream		Atrovent		Betadine	
Aratac		Aubagio		Betadine Skin Prep	
Arava	116	Augmentin	96	Betaferon	
Arginine		Aurorix		Betahistine dihydrochloride	
Arginine2000		AutoSoft 30		Betaine	
Aripiprazole		AutoSoft 90		Betamethasone dipropionate	
Aripiprazole Sandoz		Avelox		Betamethasone dipropionate with	
Aristocort		Avonex		calcipotriol	
Arrotex-Prazosin S29		Avonex Pen		Betamethasone sodium phospha	
Arrow - Clopid		Azacitidine		with betamethasone acetate	
Arrow - Lattim		Azacitidine Dr Reddy's		Betamethasone valerate	
Arrow-Amitriptyline		Azamun		Betamethasone valerate with soc	,
Arrow-Bendrofluazide		Azathioprine		fusidate [fusidic acid]	
Arrow-Brimonidine		Azilect		Betaxolol	
Arrow-Diazepam		Azithromycin		Betnovate	
Arrow-Doxorubicin		Azopt		Betoptic	
Arrow-Fluoxetine		AZT		Betoptic S	
Arrow-Losartan &	120	-B-	111	Bevacizumab	
Hydrochlorothiazide	45	B-D Micro-Fine	16	Bexsero	
Arrow-Norfloxacin		B-D Ultra Fine		Bezafibrate	
Arrow-Ornidazole		B-D Ultra Fine II		Bezalip	
Arrow-Quinapril 10		Bacillus Calmette-Guerin (BCG		Bezalip Retard	5
Arrow-Quinapril 20		vaccine		Bicalutamide	
Arrow-Quinapril 5		Bacillus Calmette-Guerin	100	Bicillin LA	
Arrow-Roxithromycin		vaccine	200	BiCNU	
•		Baclofen		BiCNU S29	
Arrow-Timolol		Bactroban			
Arrow-Topiramate				Bile and Liver Therapy	
Arrow-Tramadol		Balance		Biltricide	
Arsenic trioxide		Barrier Creams and Emollients.		Bimatoprost	
Asacol		BCG Vaccine AJV		Binarex	
Asacol S29		Beclazone 100		Binocrit	
Ascend		Beclazone 250		Biodone	
Ascend Aripiprazole		Beclazone 50		Biodone Extra Forte	
Ascend-Cefuroxime		Beclomethasone dipropionate		Biodone Forte	
Ascorbic acid		Bedaquiline		Bisacodyl	
Aspen Adrenaline	54	Bee venom allergy treatment		Bisacodyl Viatris	2
Aspirin		Bendamustine hydrochloride		Bisoprolol fumarate	
Blood		Bendamustine Sandoz		BK Lotion	
Nervous		Bendrofluazide	51	Bleomycin sulphate	15
Asthalin		Bendroflumethiazide		Blood Colony-stimulating	
Atazanavir Mylan		[Bendrofluazide]		Factors	4
Atazanavir sulphate		Benralizumab		Blood glucose diagnostic test	
Atazanavir Viatris		Benzathine benzylpenicillin		meter	1
Atenolol		Benzatropine mesylate	121	Blood glucose diagnostic test	
Atenolol AFT		Benzbromarone		strip	
Atenolol Viatris	47	Benztrop	121	Blood glucose test strips (visually	/

increasing all	Onlaine elmanata	00	Ostivisios budus ablavida	05/
impaired)	Calcium Illumanataria		Cetirizine hydrochloride	
Blood Ketone Diagnostic Test	Calcium Homeostasis		Cetomacrogol	
Strip	Calcium polystyrene sulphonat		Cetomacrogol with glycerol	
Boostrix300	Calcium Resonium		Cetomacrogol-AFT	
Bortezomib157	Calogen		Cetuximab	
Bosentan57	Camber		Charcoal	
Bosentan Dr Reddy's57	Candesartan cilexetil	45	CheckTop	
Bplex31	Candesartan cilexetil with		Chemotherapeutic Agents	151
Brentuximab Vedotin201	hydrochlorothiazide	45	Chickenpox vaccine	
Breo Ellipta258	Candestar	45	Chlorambucil	
Brevinor 1/2878	Canesten	67	Chloramphenicol	
Breztri Aerosphere260	Capecitabine	154	Chlorothiazide	51
Bricanyl Turbuhaler258	Capecitabine Viatris	154	Chlorpromazine hydrochloride	134
Brimonidine tartrate269	Capsaicin		Chlorsig	266
Brimonidine tartrate with timolol	Musculoskeletal	116	Chlortalidone [Chlorthalidone]	51
maleate269	Nervous	123	Chlorthalidone	
Brinzolamide268	Captopril	44	Chlorvescent	
Brufen SR115	Carafate		Choice 380 7med Nsha Silver/cop	
BSF Dasatinib-Teva271	Carbaccord		Short	
Buccastem133	Carbamazepine		Ciclosporin	
Budesonide	Carbimazole		Cidomycin P/Free	
Alimentary6	Carbomer		Cilazapril	1/
Respiratory257, 264	Carbonletin		Cilicaine VK	
	Carboplatin Accord		Cinacalcet	
Budesonide Te Arai	Carbosorb-X			
Budesonide with eformoterol258			Cinacalet Devatis	04
Budesonide with glycopyrronium and	Cardinol LA		Ciprofloxacin	0.0
eformoterol	Cardizem CD		Infection	
Bumetanide50	CareSens Dual		Sensory	
Buprenorphine Naloxone BNM 147	CareSens N		Ciprofloxacin Teva	
Buprenorphine with naloxone147	CareSens N POP		Cisplatin	
Bupropion hydrochloride148	CareSens N Premier		Cisplatin Accord	152
Burel 101	CareSens PRO		Cisplatin Ebewe	
Burinex50	Carmellose sodium with gelating	n and	Citalopram hydrobromide	128
Buscopan8	pectin	30	Citrulline1000	
Buspirone hydrochloride138	Carmustine		Cladribine	154
Buspirone Viatris138	Carnitor	<mark>27</mark>	Clarithromycin	
Busulfan152	Carvedilol	47	Alimentary	9
- C -	Carvedilol Sandoz	47	Infection	94
Cabergoline91	Casirivimab and imdevimab	202	Clexane	40
Caffeine citrate265	Catapres	49	Clexane Forte	
Calamine	Cefaclor monohydrate		Clindamycin	98
Calci-Tab 50033	Cefalexin		Clinicians	. 25. 32
Calcipotriol73	Cefalexin Sandoz		Clinicians Renal Vit	
Calcitonin82	Cefazolin		Clobazam	
Calcitriol32	Cefazolin-AFT		Clobetasol propionate	
Calcitriol-AFT32	Ceftriaxone		Clobetasone butyrate	
Calcitriol-AFT S2932	Ceftriaxone-AFT		Clofazimine	
Calcium 500 mg Hexal33	Cefuroxime axetil		Clomazol	10-
Calcium carbonate	Celapram			67
Calcium carbonate PAI	Celebrex		Dermatological	
	Celebrex		Genito-Urinary	
Calcium Channel Blockers			Clomifene citrate	
Calcium Disodium Versenate272	Celecoxib Pfizer	115	Clomipramine hydrochloride	
Calcium folinate	Celestone Chronodose		Clanarana Teva	
Calcium Folinate Ebewe	Cellcept		Clonazepam1	
Calcium Folinate Sandoz154	Centrally-Acting Agents		Clonidine	
Calcium Folinate Sandoz S29 154	Cephalexin ABM	93	Clonidine hydrochloride	49

Clonidine Teva49	Corticosteroids Topical	68	DBL Adrenaline	54
Clopidogrel39	Cosentyx	230	DBL Aminophylline	
Clopine134	Cosmegen	157	DBL Bleomycin Sulfate	156
Clopixol135, 138	Coumadin	41	DBL Bortezomib	157
Clotrimazole	Country Life	27	DBL Carboplatin	152
Dermatological67	Coversyl	45	DBL Carboplatin S29	
Genito-Urinary79	COVID-19 vaccine	299	DBL Cisplatin	
Clozapine134	Creon 10000		DBL Dacarbazine	
Clozaril134	Creon 25000		DBL Desferrioxamine Mesylate for	
Clustran	Creon Micro		BP	
Co-trimoxazole100	Crotamiton		DBL Docetaxel	
Coal tar	Crystaderm		DBL Ergometrine	
Coal tar with allantoin, menthol,	Cu 375 Standard		DBL Gemcitabine	
phenol and sulphur73	Curam		DBL Gentamicin	
Coal tar with salicylic acid and	Curam Duo 500/125		DBL Leucovorin Calcium	
sulphur	Cvite		DBL Methotrexate Onco-Vial	
Cobal-B1231	Cyclizine hydrochloride		DBL Naloxone Hydrochloride	
Cobalin-H	•		DBL Pethidine Hydrochloride	
Coco-Scalp	Cyclizine lactate Cyclogyl		DBL Vincristine Sulfate	
•			Decozol	
Codeine phosphate	Cyclonex			
Extemporaneous274	Cyclopentolate hydrochloride		Deferasirox	
Nervous	Cyclophosphamide		Deferiprone	
Coenzyme Q1026	Cyclorin		Deferoxamine Pfizer S29	
Colchicine	Cycloserine		Denosumab	
Colecalciferol32	Cyklokapron		Deolate	
Colestyramine52	Cyproterone acetate	84	Deoxycoformycin	
Colestyramine - Mylan52	Cyproterone acetate with		Depo-Medrol	
Colgout119	ethinyloestradiol		Depo-Provera	
Colifoam7	Cystadane		Depo-Testosterone	
Colistin sulphomethate98	Cytarabine		Deprim	
Colistin-Link98	Cytotec		Dermol	
Collodion flexible274	Cytoxan	153	Desferrioxamine mesilate	272
Colloidal bismuth subcitrate10	- D -		Desmopressin	9
Colofac8	D-Penamine	116	Desmopressin acetate	9
Colomycin98	Dabigatran	41	Desmopressin-PH&T	9
Coloxyl23	Dacarbazine	157	Detection of Substances in	
Combigan269	Dactinomycin [Actinomycin D]	157	Urine	8 [.]
Comirnaty Omicron (JN.1)299	Daivobet	72	Dexamethasone	
Compound electrolytes43	Daivonex	73	Hormone	8
Compound electrolytes with glucose	Daktarin	68	Sensory	26
[Dextrose] 43	Dalacin C	98	Dexamethasone phosphate	8
Compound hydroxybenzoate274	Dantrium	120	Dexamethasone with framycetin a	and
Comtan121	Dantrium S29	120	gramicidin	
Concerta146	Dantrolene	120	Dexamethasone with neomycin	
Condoms	Daonil	11	sulphate and polymyxin B	
Condyline74	Dapa-Tabs		sulphate	26
Continuous glucose monitor	Dapsone		Dexamfetamine sulfate	
(interoperable)21	Daraprim		Dexcom G6	_
Continuous glucose monitor	Darunavir		Dexcom G7	
(standalone)21	Darunavir Viatris		Dexcom ONE+	
Contraceptives - Hormonal77	Dasatinib		Dexmethsone	
Contraceptives - Non-hormonal75	Dasatinib-Teva		Dextrochlorpheniramine	
Copaxone	Daunorubicin		maleate	250
Cordarone-X	David One Step Cassette Pregr		Dextrose	
Corticosteroids and Related Agents	Test	R∩	DHC Continus	
for Systemic Use83	DBL Acetylcysteine	271	Diabetes	
101 Oystellile Ose00	DDF UCEINICASIGNIE	211	DIUDEIE9	10

			_
Diabetes Management14	Dosulepin [Dothiepin]	Elidel	
Diacomit131	hydrochloride 127	Elocon	6
Diagnostic Agents314	Dosulepin Viatris127	Elocon Alcohol Free	
Diamide Relief6	Dothiepin127	Eltrombopag	3
Diamox268	Dovato112	Eltroxin	8
Diasip278	Doxazosin44	EMB Fatol	
Diazepam129, 138	Doxazosin Clinect44	Emend Tri-Pack	
Diazoxide10	Doxine97	Emicizumab	3
Dibenzyline44	Doxorubicin Ebewe157	EMLA	. 12
Diclofenac Devatis267	Doxorubicin hydrochloride157	Empagliflozin	14
Diclofenac Sandoz115	Doxycycline97	Empagliflozin with metformin	
Diclofenac sodium	DP Lotion70	hydrochloride	14
Musculoskeletal115	DP Lotn HC69	Emsogen	. 29
Sensory267	DP-Captopril44	Emtricitabine	.11
Differin66	Dr Reddy's Omeprazole9	Emtricitabine with tenofovir	
Difflam30	Drofate48	disoproxil	. 10
Diflucan100	Drugs Affecting Bone	Emtriva	.11
Digestives Including Enzymes22	Metabolism 116	Emulsifying ointment	70
Digoxin46	Dual blood glucose and blood ketone	Emulsifying Ointment ADE	
Dihydrocodeine tartrate 125	diagnostic test meter 14	Enalapril maleate	
Dilantin	Dulaglutide12	Enbrel	
Dilantin Infatab130	Dulcolax SP Drop24	Endocrine Therapy	.17
Dilantin Paediatric130	Duocal Super Soluble Powder276	Endoxan	
Diltiazem CD Clinect49	Duolin259	Energivit	
Diltiazem hydrochloride49	Duolin Cipla259	Engerix-B	
Dimethicone70–71	Duolin HFA259	Enhertu	
Dimethyl fumarate	DuoResp Spiromax258	Enlafax XR	
Dipentum8	Duride54	Enoxaparin sodium	40
Diperituiri			
•	Durvalumab244		
Diphtheria, tetanus and pertussis vaccine		Enstilar	7
Diphtheria, tetanus and pertussis vaccine300	Durvalumab244	Enstilar	72 .28
Diphtheria, tetanus and pertussis	Durvalumab244	Enstilar Ensure	7 .28 .28
Diphtheria, tetanus and pertussis vaccine	Durvalumab	Enstilar Ensure Ensure Plus	72 .28! .28!
Diphtheria, tetanus and pertussis vaccine	Durvalumab	Enstilar Ensure Ensure Plus Ensure Plus HN	72 .285 .286 .286
Diphtheria, tetanus and pertussis vaccine	Durvalumab	Enstilar Ensure Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH	72 .285 .285 .286 .286
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - 265 e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10	Enstilar Ensure Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH	75 .285 .286 .286 .286 .286
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266	Enstilar Ensure Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone	72 .285 .286 .286 .286 .287 .127 .127
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 - E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266	Enstilar Ensure Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entacapone Viatris	72 .285 .286 .286 .287 .127 .120
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266	Enstilar Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entacapone Viatris Entecavir	72 .283 .284 .284 .284 .281 .121 .100 .100
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E • e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67	Enstilar Ensure Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entacapone Viatris	77 .283 .284 .284 .284 .12 .100 .100 44
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290	Enstilar Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entacapone Viatris Entecavir Entecavir (Rex) Entresto 24/26 Entresto 49/51	77 .283 .284 .284 .284 .12 .100 .100 44
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110	Enstilar Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entacapone Viatris Entecavir Entecavir (Rex) Entresto 24/26 Entresto 49/51 Entresto 97/103	77 .283 .284 .284 .284 .12 .100 .100 44 44
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and	Enstilar Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entacapone Viatris Entecavir Entecavir (Rex) Entresto 24/26 Entresto 49/51	77 .283 .284 .284 .285 .122 .100 .100 44 44 44
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and tenofovir disoproxil 111	Enstilar Ensure Plus Ensure Plus HN Ensure Plus HN RTH Ensure Two Cal HN RTH Entacapone Entacapone Viatris Entecavir Entecavir (Rex) Entresto 24/26 Entresto 49/51 Entresto 97/103 Entyvio	77 .283 .284 .284 .284 .12 .100 .100 44 44 44 44 44
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and tenofovir disoproxil 111 Eformoterol fumarate dihydrate 257	Enstilar	77 .283 .284 .284 .287 .127 .100 .100444444414141
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and tenofovir disoproxil 111 Eformoterol fumarate dihydrate 257 Eftrenonacog alfa [Recombinant	Enstilar	7; .283 .284 .284 .285 .122 .100 .100444444434343
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and tenofovir disoproxil 111 Eformoterol fumarate dihydrate 257	Enstilar	7; .285 .286 .286 .287 .121 .100 .1004!4!4!4!4!4!4!4!
Diphtheria, tetanus and pertussis vaccine 300 Diphtheria, tetanus, pertussis and polio vaccine 301 Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine 301 Diprosone 68 Diprosone OV 68 Dipyridamole 39 Disopyramide phosphate 46 Disulfiram 148 Diuretics 50 Docetaxel 157 Docetaxel Accord 157 Docusate Sandoz 157 Docusate sodium 23 Docusate sodium with sennosides 23	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and tenofovir disoproxil 111 Eformoterol fumarate dihydrate 257 Eftrenonacog alfa [Recombinant factor IX] 36 Efudix 74	Enstilar	7; 288; 284; 284; 284; 12; 100; 100;44; 13; 13; 13; 13; 13;
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and tenofovir disoproxil 111 Eformoterol fumarate dihydrate 257 Eftrenonacog alfa [Recombinant factor IX] 36 Efudix 74 Egopsoryl TA 73	Enstilar	7; 288; 284; 284; 287; 100; 100; 100; 133; 133; 133; 255;
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and tenofovir disoproxil 111 Eformoterol fumarate dihydrate 257 Eftrenonacog alfa [Recombinant factor IX] 36 Efudix 74 Egopsoryl TA 73 Elaprase 26	Enstilar	7; 288; 288; 288; 288; 122; 100; 100; 133; 133; 133; 255; 255;
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and tenofovir disoproxil 111 Eformoterol fumarate dihydrate 257 Eftrenonacog alfa [Recombinant factor IX] 36 Efudix 74 Egopsoryl TA 73 Elaprase 26 Elecare 294	Enstilar	72 .283 .284 .284 .122 .100 .100 44
Diphtheria, tetanus and pertussis vaccine	Durvalumab 244 • E - e-chamber La Grande 265 e-chamber Mask 264 e-chamber Turbo 265 E-Mycin 95 e5 Pharma 10 Ear Preparations 266 Ear/Eye Preparations 266 Easiphen Liquid 290 Econazole nitrate 67 Efavirenz 110 Efavirenz Milpharm 110 Efavirenz with emtricitabine and tenofovir disoproxil 111 Eformoterol fumarate dihydrate 257 Eftrenonacog alfa [Recombinant factor IX] 36 Efudix 74 Egopsoryl TA 73 Elaprase 26	Enstilar	72 .283 .284 .284 .121 .100 .100 44 44 44 44 44 44 44 44 45 45 13
Diphtheria, tetanus and pertussis vaccine	Durvalumab	Enstilar	72 .283 .284 .284 .287 .122 .100 .100 44 44 44 44 44 44 41 41 43 44 43 43 43 43 43 43 43 43 43 43
Diphtheria, tetanus and pertussis vaccine	Durvalumab	Enstilar	72 .283 .284 .284 .287 .122 .100 .1004444 .13 .13 .13 .13 .13 .15 .155656
Diphtheria, tetanus and pertussis vaccine	Durvalumab	Enstilar	72 .283 .284 .284 .287 .122 .100 .1004444 .13 .13 .13 .13 .13 .15 .155656
Diphtheria, tetanus and pertussis vaccine	Durvalumab	Enstilar	72 .283 .284 .284 .285 .122 .100 .1004444444343565656

Erbitux	202	- F -		Fludarabine phosphate	15
Ergometrine maleate		Factor eight inhibitor bypassing		Fludarabine Sagent	
Erlotinib		fraction	38	Fludrocortisone acetate	
Erythrocin IV		Famotidine	9	Fluids and Electrolytes	
Erythromycin (as lactobionate)		Famotidine Hovid	9	Flumetasone pivalate	
Erythromycin ethyl succinate		Fasenra		Fluocortolone caproate with	
Esbriet		Faslodex	173	fluocortolone pivalate and	
Escitalopram		Fatty Cream AFT	70	cinchocaine	
Escitalopram (Ethics)		Fatty Emulsion Cream (Evara).		Fluorometholone	
Eskazole		Febuxostat		Fluorouracil	
Essential Amino Acid Mix		Febuxostat (Teva)		Fluorouracil Accord	15
Essential Ethosuximide	129	FEIBA NF		Fluorouracil sodium	
Estraderm MX	85	Felo 10 ER		Fluox	12
Estradiol Sandoz	85	Felo 5 ER	49	Fluoxetine hydrochloride	12
Estradiol TDP Mylan		Felodipine	49	Flupenthixol decanoate	
Estradiol Viatris		Fentanyl		Flutamide	17
Estradot		Fentanyl Sandoz		Flutamin	
Estrofem		Ferinject		Fluticasone	
Estrogel		Ferodan		Fluticasone furoate with	
Etanercept		Ferriprox		umeclidinium and vilanterol	26
Ethambutol hydrochloride		Ferro-F-Tabs		Fluticasone furoate with	
Ethics Aspirin		Ferro-Liquid		vilanterol	25
Ethics Aspirin EC		Ferro-tab		Fluticasone propionate	
Ethics Lisinopril		Ferrograd		Fluticasone with salmeterol	
Ethinyloestradiol with		Ferrosig		Flynn	
desogestrel	77	Ferrous fumarate	33	FML	
Ethinyloestradiol with		Ferrous fumarate with folic acid		Foban	
levonorgestrel	78	Ferrous sulfate		Folic acid	
Ethinyloestradiol with		Fexaclear		Folic Acid multichem	
norethisterone	78	Fexofenadine hydrochloride		Folic Acid Viatris	
Ethosuximide		Fibro-vein		Food Thickeners	
Etopophos		Filgrastim		Foods And Supplements For	
Etoposide		Finasteride		Inherited Metabolic Disease	28
Etoposide phosphate		Fingolimod		Fortini	
Etravirine		Firazyr	255	Fortini Multi Fibre	
Eumovate		Flagyl		Fortisip	
Eurofolic		Flagyl-S		Fortisip Multi Fibre	
Evara		Flamazine		Fosamax	
EVARA White Soft Paraffin		Flecainide acetate		Fosamax Plus	
Everet		Flecainide BNM		Fosfomycin	
Everolimus		Flecainide Controlled Release		Framycetin sulphate	
Evista		Teva	47	Freestyle Libre 2	20
Evrysdi		Fleet Phosphate Enema		Freestyle Libre 3 Plus	2
Exelon Patch 10		Flixonase Hayfever & Allergy		Frisium	
Exelon Patch 5		Flixotide		Frumil	
Exemestane		Flixotide Accuhaler		Frusemide	
Exjade		Florinef		Fucicort	
Extemporaneously Compounded		Fluanxol		Fucidin	
Preparations and	ı	Flucil		Fucithalmic	
Galenicals	27/	Flucloxacillin		Fucithalmic S29	
Eye Preparations	266	Flucloxacillin-AFT		Fulvestrant	
Eylea		Flucioxaciiiii -Ai-i		Fungilin	
Ezemibe Viatris		Flucon		Furosemide [Frusemide]	۰۰۰۰۰۰
Ezetimibe		Fluconazole		Furosemide-Baxter	5
Ezetimibe Sandoz		Fludara Oral		fusidic acid	
Ezetimibe Sandoz		Fludarabine Ebewe		Dermatological	67 6
LECULLING WILL SHIIVASIAIII		I IUUGIADIIIE EDEWE		DEITIGIUIUUIUAI	. u . n

Infection	100	Haemophilus influenzae type B		Hydralazine	54
Sensory		vaccine	302	Hydralazine hydrochloride	
´ - G -		Haldol	136	Hydralyte - Lemonade	
GA Explore 5	292	Haldol Concentrate	136	Hydrocortisone	
GA1 Anamix Infant		Haldol Decanoas	136	Dermatological	6
GA1 Anamix Junior	292	Haloperidol	134	Hormone	8
Gabapentin	129	Haloperidol decanoate	136	Hydrocortisone acetate	
Gacet		Harvoni	108	Hydrocortisone acetate with	
Galsulfase		Havrix 1440	302	pramoxine hydrochloride	
Galvumet	12	Havrix Junior	302	Hydrocortisone and paraffin liquid	
Galvus	12	Haylor syrup	256	and lanolin	
Gardasil 9	304	HCU Anamix Infant	289	Hydrocortisone butyrate	69, 74
Gastrodenol	10	HCU Anamix Junior	289	Hydrocortisone with cinchocaine	
Gaviscon Extra Strength	6	HCU Anamix Junior LQ	289	Hydrocortisone with miconazole	6
Gaviscon Infant	6	HCU Explore 5	289	Hydrocortisone with natamycin and	t
Gazyva	213	HCU Express 15	289	neomycin	6
Gefitinib	165	HCU Lophlex LQ	289	Hydrogen peroxide	
Gemcitabine Ebewe	155	healthE Calamine Aqueous	68	Hydroxocobalamin	3
Gemcitabine hydrochloride		healthE Dimethicone 10%	70	Hydroxocobalamin Panpharma	3
Gemtuzumab ozogamicin	203	healthE Dimethicone 4% Lotion .	71	hydroxycarbamide	15
Gentamicin Amdipharm	98	healthE Dimethicone 5%	70	Hydroxychloroquine sulphate	110
Gentamicin Noridem		healthE Glycerol BP	274	Hydroxyurea	
Gentamicin sulphate		healthE Urea Cream	70	[hydroxycarbamide]	15
Gilenya	139	Healtheries Simple Baking Mix	287	Hygroton	
Ginet	79	Hemastix	81	Hylo-Fresh	
Glatiramer acetate	139	Hemlibra	37	Hymenoptera	25
Glecaprevir with pibrentasvir	108	Heparin sodium		Hyoscine butylbromide	
Glibenclamide		Heparin Sodium Panpharma	41	Hyoscine Butylbromide	
Gliclazide	11	Heparinised saline	41	(Adiramedica)	
Glipizide		Heparon Junior	279	Hyoscine hydrobromide	
Glizide		Hepatitis A vaccine	302	Hyperuricaemia and Antigout	
Glucagen Hypokit		Hepatitis B recombinant		Hypromellose	
Glucagon hydrochloride		vaccine		Hypromellose with dextran	26
Glucerna Select		Herzuma		-1-	
Glucose [Dextrose]		Hikma		Ibiamox	
Gluten Free Foods		Hiprex		Ibrance	
Glycerin with sodium saccharin		Histaclear		Ibrutinib	
Glycerin with sucrose	274	Holoxan		Ibuprofen	
Glycerol		Horleys Bread Mix		Ibuprofen SR BNM	
Alimentary		Horleys Flour		Icatibant	
Extemporaneous	274	Hormone Replacement Therapy		Idarubicin hydrochloride	
Glyceryl trinitrate	_	Systemic		Idursulfase	
Alimentary		HPV		Ifosfamide	
Cardiovascular		Humalog		llevro	
Glycopyrronium		Humalog Mix 25		lloprost	
Glycopyrronium bromide	8	Humalog Mix 50		Imatinib mesilate	
Glycopyrronium with	000	Human papillomavirus (6, 11, 16,		Imatinib-Rex	
indacaterol		31, 33, 45, 52 and 58) vaccine		Imbruvica	
Glycosade		[HPV]		Imfinzi	
Glytactin Bettermilk		Humatin		Imipramine Crescent	
Gold Knight		Humira		Imipramine hydrochloride	
Gold Knight XL		HumiraPen		Imiquimod	
Goserelin		Humulin 30/70		Immune Modulators	
Gynaecological Anti-infectives	/9	Humulin NPH		Immunisation - Flu	
- H -	440	Humulin R		Immunisation Flu and Shingles	
Habitrol	149	Hyaluronic acid	2/0	Immunisation Other	2/

Immunosuppressants	175	Intra-uterine device	77	Kalydeco	26
Incruse Ellipta	259	Invega Sustenna	137	Kemadrin	
Indacaterol	257	Invega Trinza	137	Kenacomb	
Indapamide	51	Ipca-Allopurinol		Kenacort-A 10	8
Infanrix IPV		Ipca-Bisoprolol		Kenacort-A 40	8
Infanrix-hexa		Ipca-Ciprofloxacin		Kenalog in Orabase	3
Infant Formulae		Ipca-Donepezil		Ketocal 3:1	
Infatrini	297	lpca-Escitalopram		KetoCal 4:1	
Infliximab		IPCA-Frusemide		Ketoconazole	
Influenza vaccine		Ipca-Hydroxychloroquine		Dermatological	7
Influvac Tetra		IPCA-Metoprolol		Infection	
(2025 formulation)	305	IPCA-Propranolol		Ketogenic Diet	
Inhaled Corticosteroids		IPOL		Ketoprofen	
Inhaled Long-acting		Ipratropium bromide		KetoSens	
Beta-adrenoceptor Agonists	257	Iressa		Keytruda	
Inresa		Irinotecan Actavis 100		Kindergen	
Inspra		Irinotecan hydrochloride		Kisqali	
Instillagel Lido		Irinotecan-Rex		Klacid	17
Insulin aspart		Iron (as ferric carboxymaltose).		Alimentary	
Insulin aspart with insulin aspart		Iron polymaltose		Infection	
		Isentress		Klaricid	
protamine		Isentress HD		Kliogest	
Insulin glargine					
Insulin glulisine		Ismo 20		Kliovance	
Insulin isophane		Ismo 40 Retard		Kogenate FS	
Insulin isophane with insulin	4.4	Isoleucine50		Konakion MM	
neutral		Isoniazid		Konsyl-D	
Insulin lispro	11	Isoniazid Teva		Kuvan	2
Insulin lispro with insulin lispro		Isoniazid with rifampicin		-L-	
protamine		Isoptin		Labetalol	
Insulin neutral		Isoptin Retard		Lacosamide	
Insulin pen needles		Isoptin SR		Lactulose	
Insulin pump cartridge	17	Isopto Carpine		Laevolac	
Insulin pump infusion set (steel		Isosorbide mononitrate		Lamictal	
cannula)	18	Isosource Standard		Lamivudine10	
Insulin pump infusion set (steel		Isotretinoin	66	Lamivudine Viatris	11
cannula, straight insertion)	18	Ispaghula (psyllium) husk	23	Lamivudine/Zidovudine Viatris	
Insulin pump infusion set (teflon		Itch-Soothe	68	Lamotrigine	13
cannula)	19	Itracap	101	Lamprene	10
Insulin pump infusion set (teflon		Itraconazole	101	Lanoxin	
cannula, angle insertion with		Itraconazole Kent	101	Lanoxin Paediatric Elixir	4
insertion device)	20	Itrazole	101	Lanoxin PG	4
Insulin pump infusion set (teflon		Ivacaftor	263	Lanoxin S29	4
cannula, flexible insertion with	1	Ivermectin	71	Lanreotide	17
insertion device)		- J -		Lansoprazole	
Insulin pump infusion set (teflon		Jadelle	78	Lantus	
cannula, straight insertion with	h	Jakavi	170	Lantus SoloStar	1
insertion device)		Jardiamet		Lanvis	15
Insulin pump infusion set (teflon		Jardiance		Lanzol Relief	
cannula, variable insertion)	20	Jaydess		Largactil	
Insulin pump reservoir		Jevity HiCal RTH		Laronidase	
Insulin pump with algorithm		Jevity Plus RTH		Lasix	
Insulin syringes, disposable with		Jevity RTH		Latanoprost	
attached needle		Jinarc		Latanoprost with timolol	
Intelence		Juno Pemetrexed		Lax-Suppositories	
Interferon beta-1-alpha		- K -	100	Lax-suppositories Glycerol	
Interferon beta-1-beta		Kadcyla	228	Laxatives	
ווונטוופוטוו טפומ־ו־טפומ		radoyia	200	∟ u∧aliv c 3	4

Laxsol	23	Locoid Crelo	69	Mebendazole	9:
Ledipasvir with sofosbuvir		Locoid Lipocream		Mebeverine hydrochloride	
Leflunomide		Locorten-Vioform		Medac	
Lenalidomide (Viatris)		Lodoxamide		Medrol	
Lenalidomide Viatris		Logem		Medroxyprogesterone acetate	
Lenvatinib		Lomide		Genito-Urinary	7
Lenvima		Lomustine		Hormone	
Letrole		Loniten		Mefenamic acid	
Letrozole		Loperamide hydrochloride		Megval	
Leucine100		Lopinavir with ritonavir		Melatonin	
Leukeran FC		Lopinavir/Ritonavir Mylan		Melpha	
Leukotriene Receptor	102	Loprofin		Melphalan	
Antagonists	262	Loprofin Mix		Meningococcal (groups A, C, Y and	
Leuprorelin	01	Lorafix		W-135) conjugate vaccine	
Leustatin		Loratadine		Meningococcal B multicomponent	. 500
Levetiracetam		Lorazepam		vaccine	300
Levetiracetam-AFT		Lorstat		MenQuadfi	
Levocabastine		Losartan Actavis		Menthol	
Levocarnitine				Mepolizumab	
		Losartan potassium	43	Mercaptopurine	
Levodopa with benserazide		Losartan potassium with hydrochlorothiazide	45	Mercilon 28	13
Levodopa with carbidopa	121			Mesalazine	
Levodopa with carbidopa and	101	Lovir Loxamine		Mesna	
entacapone				Mestinon	
Levomepromazine	134	Lucrin Depot 1-month Lucrin Depot 3-month			
Levomepromazine hydrochloride	105	•		Metabolic Disorder Agents	
	133	LumaCina		Metabolics	
Levonorgestrel	70 70	Lumigan		Metformin hydrochloride	
Genito-Urinary		Lyllana		Metformin Viatris	
Hormone		Lynparza		Methadone BNM	
Levenorgestrel BNM		Lyrica	130	Methadone hydrochloride	12
Levothyroxine		***	100	Methenamine (hexamine)	44
Lidocaine [Lignocaine]1	22-123	m-Eslon		hippurate	
Lidocaine [Lignocaine]	100	Mabthera		Methopt	
hydrochloride	123	Macrobid	114	Methotrexate	
Lidocaine [Lignocaine] with	400	Macrogol 3350 with potassium		Methotrexate DBL Onco-Vial	
prilocaine		chloride, sodium bicarbonate a		Methotrexate Ebewe	
Lidocaine-Baxter		sodium chloride		Methotrexate Sandoz	
Life Extension		Madopar 125		Methyl hydroxybenzoate	
Lignocaine1		Madopar 250		Methylcellulose	2/4
Linezolid		Madopar 62.5		Methylcellulose with glycerin and	07
Lioresal Intrathecal		Madopar HBS		sodium saccharin	. 2/4
Lipid-Modifying Agents		Madopar Rapid	121	Methylcellulose with glycerin and	
Lipitor		Magnesium hydroxide		sucrose	
Liquigen		Magnesium sulphate		Methyldopa	
Liraglutide		Mantoux		Methyldopa Viatris	
Lisdexamfetamine dimesilate		MAR-Midodrine		Methylnaltrexone bromide	
Lisinopril		Marevan		Methylphenidate ER - Teva	
Lithium carbonate		Marine Blue Lotion SPF 50+		Methylphenidate hydrochloride	14
Livostin		Martindale Pharma		Methylphenidate hydrochloride	
LMX4			064	extended-release	. 14
Lo-Oralcon 20 ED		Mask for spacer device			
	78	Maviret	108	Methylprednisolone	
Locacorten-Viaform ED's	78	Maviret Maxidex	108 267	Methylprednisolone Methylprednisolone (as sodium	8
Locacorten-Viaform ED'sLocal preparations for Anal and	78 266	Maxidex Maxitrol	108 267 267	Methylprednisolone Methylprednisolone (as sodium succinate)	8: 8:
Locacorten-Viaform ED's Local preparations for Anal and Rectal Disorders	78 266	Maviret Maxidex Maxitrol MCT oil (Nutricia)	108 267 267	Methylprednisolone Methylprednisolone (as sodium succinate) Methylprednisolone aceponate	8: 8: 6!
Locacorten-Viaform ED'sLocal preparations for Anal and	78 266 8 293	Maxidex Maxitrol	108 267 267 277	Methylprednisolone Methylprednisolone (as sodium succinate)	8; 8; 6;

Metoclopramide Actavis 10	133	Minims Pilocarpine	269	Mupirocin	6
Metoclopramide hydrochloride		Minims Prednisolone		Muscle Relaxants	11
Metolazone	51	Minipress	44	Mvite	
Metopirone	92	Minirin	91	Myambutol	10
Metoprolol IV Mylan	48	Minirin Melt	91	Mycobutin	10
Metoprolol IV Viatris		Mino-tabs	97	MycoNail	
Metoprolol succinate		Minocycline hydrochloride	97	Mycophenolate mofetil	
Metoprolol tartrate		Minomycin		Mydriacyl	
Metronidamed		Minor Skin Infections		Mylan (24 hr release)	
Metronidazole		Minoxidil		Mylan Clomiphen	
Metyrapone		Minoxidil Roma		Mylan Italy (24 hr release)	
Mexiletine hydrochloride		Mirena		Myleran	
Miacalcic		Miro-Amoxicillin	96	mylife Inset soft	
Micolette		Mirtazapine		mylife Orbit micro	1
Miconazole		Misoprostol		mylife YpsoPump Reservoir	
Miconazole nitrate		Mitomycin (Fresenius Kabi)		mylife YpsoPump with CamAPS	
Dermatological	68	Mitomycin (Sagent)		FX	1
Genito-Urinary		Mitomycin C		Myloc CR	
Micreme		Mitozantrone		Mylotarg	
Micreme H		Mitozantrone Ebewe		Myometrial and Vaginal Hormone	20
Microgynon 30		Mixtard 30			7
Microlut		MMA/PA Anamix Infant		Preparations Myozyme	
Midazolam		MMA/PA Anamix Junior		Mytolac	
				- N -	17
Midazolam Viatris		MMA/PA Explore 5			4
Midazolam-Baxter		MMA/PA Express 15		Nadolol	
Midazolam-Pfizer		Moclobemide		Nadolol BNM	
Midodrine		Modafinil		Naglazyme	
Midostaurin		Modafinil Max Health		Naloxone hydrochloride	
Mifegyne		Modavigil		Naltraccord	
Mifepristone		Moduretic		Naltrexone AOP	
Milpharm		Molaxole		Naltrexone hydrochloride	
Minerals		Moments		Naltrexone Max Health	
Mini-Wright AFS Low Range		Mometasone furoate		Naphazoline hydrochloride	
Mini-Wright Standard		Monogen		Naprosyn SR 1000	
Minidiab	12	Montelukast	262	Naprosyn SR 750	
MiniMed 3.0 Reservoir		Montelukast Viatris	262	Naproxen	11
MMT-332A	21	Moroctocog alfa [Recombinate	nt factor	Narcaricin mite	11
MiniMed Mio MMT-921A	19	VIII]	38	Nasal Preparations	26
MiniMed Mio MMT-923A	19	Morphine hydrochloride	125	Natalizumab	13
MiniMed Mio MMT-925A	19	Morphine sulphate	126	Natulan	16
MiniMed Mio MMT-941A	19	Motetis	122	Nausafix	13
MiniMed Mio MMT-943A	19	Mouth and Throat	30	Nausicalm	13
MiniMed Mio MMT-945A	19	Movapo	121	Navelbine S29	16
MiniMed Mio MMT-965A		Moxifloxacin		Nefopam hydrochloride	12
MiniMed Mio MMT-975A		MSUD Anamix Infant		Neo-Cytamen S29	
MiniMed Quick-Set MMT-396A		MSUD Anamix Junior		Neo-Mercazole	
MiniMed Quick-Set MMT-397A		MSUD Anamix Junior LQ		Neocate Gold	
MiniMed Quick-Set MMT-398A		MSUD Explore 5		Neocate Junior Unflavoured	
MiniMed Quick-Set MMT-399A		MSUD Express 15	289	Neocate Junior Vanilla	
MiniMed Silhouette MMT-377A		MSUD Lophlex LQ 20	280	Neocate SYNEO	
MiniMed Silhouette MMT-378A		MSUD Maxamum	280	Neoral	
MiniMed Silhouette MMT-381A		Mucolytics		Neostigmine metilsulfate	
MiniMed Sure-T MMT-864A		Mucosoothe		Nepafenac	
MiniMed Sure-T MMT-866A		Multiple Sclerosis Treatments		Nepro HP (strawberry)	20
		Multivitamin renal	الان السند. د مو		
MiniMed Sure-T MMT-874A				Nepro HP (vanilla)	
MiniMed Sure-T MMT-876A	I 🎖	Multivitamins	52	Neulactil	1ರ

NeuroTabs		NovoSeven RT	37	Olbetam	
Nevirapine		Nozinan		Olopatadine	
Nevirapine Viatris		Nozinan (Swiss)		Olopatadine Teva	27
Nicorandil		Nozinan S29		Olsalazine	
Nicotine		Nucala		Omalizumab	
Nifedipine	49	Nuelin		Omeprazole	
Nifuran		Nuelin-SR		Omeprazole actavis 10	
Nilotinib	167	Nupentin		Omeprazole actavis 20	
Nilstat		Nusinersen		Omeprazole actavis 40	
Alimentary	30	Nutilis		Omeprazole Teva	
Genito-Urinary	79	Nutren Diabetes	278	Omnitrope	
Infection		Nutrient Modules		Onbrez Breezhaler	
Nimenrix		Nutrini Energy Multi Fibre		Oncaspar LYO	
Nintedanib		Nutrini Energy RTH	280	OncoTICE	
Nipent	161	Nutrini Low Energy Multi Fibre		Ondansetron	
Niraparib		Nutrini Peptisorb	296	One-Alpha	
Nirmatrelvir with ritonavir	109	Nutrini Peptisorb Energy	296	One-Alpha S29	
Nitrates	54	Nutrini RTH	280	Opdivo	24
Nitroderm TTS	54	Nutrison 800 Complete Multi		Ora-Blend	27
Nitrofurantoin	114	Fibre	285	Ora-Blend SF	27
Nitrolingual Pump Spray	54	Nutrison Advanced Peptisorb	281	Ora-Plus	
Nivestim	42	Nutrison Concentrated	287	Ora-Sweet	27
Nivolumab	244	Nutrison Energy	284	Ora-Sweet SF	27
Nodia	6	Nutrison Energy Multi Fibre	285	Orabase	30
Noflam 250	115	Nutrison Multi Fibre		Oral and Enteral Feeds	27
Noflam 500	115	Nutrison RTH	285	Oralcon 30 ED	
Non-Steroidal Anti-Inflammatory	•	Nyefax Retard	49	Oramorph	120
Drugs	115	Nystatin		Oramorph CDC S29	
Nonacog gamma, [Recombinant		Alimentary	30	Oratane	6
Factor IX]	38	Genito-Urinary	79	Orgran	
Norethinderone - CDC	79	Infection	101	Ornidazole	
Norethisterone		NZB Low Gluten Bread Mix	288	Orphenadrine citrate	120
Genito-Urinary	79	- O -		Ortho-tolidine	8
Hormone		Obinutuzumab	213	Oruvail SR	11
Norflex	120	Obstetric Preparations	81	Osimertinib	16
Norfloxacin	114	Ocicure	9	Osmolite RTH	28
Noriday	79	Ocrelizumab	140	Other Endocrine Agents	9
Noriday 28	79	Ocrevus	140	Other Oestrogen Preparations	8
Norimin	78	Octocog alfa [Recombinant factor	or	Other Progestogen	
Normison	142	VIII] (Advate)	38	Preparations	80
Norpress	128	Octocog alfa [Recombinant factor		Other Skin Preparations	
Nortriptyline hydrochloride	128	VIII] (Kogenate FS)	38	Otodex	
Norvir		Octreotide		Ovestin	
Noumed Dexamfetamine	143	Octreotide GH	174	Genito-Urinary	79
Noumed Isoniazid	105	Octreotide long-acting	175	Hormone	
Noumed Paracetamol	124	Oestradiol	85	Oxaliplatin	
Noumed Pethidine	127	Oestradiol valerate		Oxaliplatin Accord	
Noumed Phenobarbitone	130	Oestradiol with norethisterone	86	Oxaliplatin Actavis 100	15
Novadoz		Oestriol		Oxaliplatin Ebewe	
NovaSource Renal		Genito-Urinary	79	Oxis Turbuhaler	
Novatretin		Hormone		Oxpentifylline	
Novitium Sugar Free		Oestrogens		Oxybutynin	
NovoMix 30 FlexPen		Ofev		Oxycodone Amneal	120
NovoRapid		Oil in water emulsion		Oxycodone hydrochloride	
NovoRapid FlexPen		Olanzapine		Oxycodone Lucis	
NovoRapid Penfill		Olaparib		Oxycodone Sandoz	120

INDEX: Generic Chemicals and Brands

Oxycodone Sandoz S29	126	Penicillin G	96	PKU Anamix Junior LQ	290
OxyContin		PenMix 30	11	PKU Anamix Junior Orange	290
Oxytocin		PenMix 50	11	PKU Anamix Junior Vanilla	290
Oxytocin BNM		Pentasa	7	PKU Build 10	
Oxytocin Panpharma		Pentostatin [Deoxycoformycin]		PKU Build 20 Chocolate	291
Oxytocin with ergometrine		Pentoxifylline [Oxpentifylline]		PKU Build 20 Raspberry	
maleate	80	Peptamen Junior		Lemonade	291
Ozurdex		Pepti-Junior		PKU Build 20 Smooth	
-P-		Perhexiline maleate		PKU Build 20 Vanilla	
Pacifen	119	Pericyazine		PKU Explore 10	
Pacimol		Perindopril		PKU Explore 5	
Paclitaxel		Periset		PKU Express 20	
Paclitaxel Actavis		Periset ODT		PKU First Spoon	
Paclitaxel Ebewe		Perjeta		PKU Glytactin RTD 15	291
Padagis		Permethrin		PKU Glytactin RTD 15 Lite	
Paediatric Seravit		Perrigo		PKU GMPro LQ	
Palbociclib		Pertuzumab		PKU GMPro Mix-In	
Paliperidone		Peteha		PKU GMPro Ultra Lemonade	
				PKU GMPro Ultra Vanilla	
Paliperidone palmitate		Pethidine hydrochloride		PKU Lophlex LQ 10	
Palivizumab		Pevaryl		PKU Lophley LQ 10	200
Pamidronate disodium		Pexsig		PKU Lophlex LQ 20	
Pamisol		Pfizer Exemestane		PKU Lophlex Powder	
Pamol		Pfizer S29		PKU Lophlex Sensation 20	
Pancreatic enzyme		Pharmacy Services		PKU Restore Powder	
Pantoprazole		Pharmascience		PKU sphere20 Banana	
Panzop Relief		Pheburane		PKU sphere20 Chocolate	
Papaverine hydrochloride		Phenasen		PKU sphere20 Lemon	
Para-amino salicylic acid		Phenergan Elixir		PKU sphere20 Red Berry	
Paracetamol		Phenobarbitone	130	PKU sphere20 Vanilla	
Paracetamol (Ethics)	124	Phenobarbitone sodium		PKU Start	
Paracetamol + Codeine		Extemporaneous		Plaquenil	
(Relieve)		Nervous	142	Plendil ER	
Paracetamol with codeine		Phenoxybenzamine		Pneumococcal (PCV13) conjugate	
Paraffin		hydrochloride		vaccine	310
Paraffin liquid with wool fat		Phenoxymethylpenicillin (Penicilli		Pneumococcal (PPV23)	
Parasiticidal Preparations	71	V)		polysaccharide vaccine	311
Parnate		Phenylalanine50	293	Pneumovax 23	
Paromomycin	99	Phenytoin sodium1	29–130	Podophyllotoxin	
Paroxetine	128	Phillips Milk of Magnesia	34	Polaramine	256
Paser	105	Phlexy 10	290	Poliomyelitis vaccine	311
Paxam		Phosphate Phebra		Poloxamer	
Paxlovid		Phosphorus	43	Poly-Gel	
Pazopanib	169	Phytomenadione	39	Poly-Tears	269
Pazopanib Teva	169	Pilocarpine hydrochloride	269	Poly-Visc	270
Peak flow meter		Pilocarpine nitrate	269	Polycal	275
Pediasure	280	Pimafucort	69	Polyethylene glycol 400 and	
Pediasure Plus	280	Pimecrolimus	73	propylene glycol	270
Pediasure RTH	280	Pine tar with trolamine laurilsulfat	ie	Pomalidomide	
Pegaspargase		and fluorescein		Pomolide	
Pegasys		Pinetarsol	73	Ponstan	115
Pegfilgrastim		Pioglitazone	12	Posaconazole	102
Pegylated interferon alfa-2a		Pirfenidone		Posaconazole Juno	
Pembrolizumab		Pizotifen		Potassium chloride	
Pemetrexed		PKU Anamix Infant		Potassium citrate	
Pemetrexed-AFT		PKU Anamix Junior		Potassium iodate	
Penicillamine		PKU Anamix Junior Chocolate		Povidone iodine	

Pradaxa	41	Puri-nethol	155	RINVOQ	25
Pramipexole hydrochloride		Puritan's Pride Vitamin		Riodine	
Pravastatin		B-2 100 mg	27	Risdiplam	
Praziquantel		Pyrazinamide		Risedronate Sandoz	119
Prazosin		Pyridostigmine bromide		Risedronate sodium	
Prazosin Mylan		Pyridoxine hydrochloride		Risperdal	
Pred Forte		Pyridoxine multichem		Risperdal Consta	
Prednisolone		Pyrimethamine		Risperidone1	
Prednisolone acetate	208	Pytazen SR	39	Risperidone (Teva)	ال
Prednisolone sodium	000	- Q -	50	Risperidone Sandoz	13
phosphate		Quantalan sugar free		Risperon	ال
Prednisolone-AFT		Quetapel		Ritalin	
Prednisone		Quetiapine		Ritalin LA	
Prednisone Clinect		Quinapril		Ritonavir	
Pregabalin		Qvar	25/	Rituximab (Mabthera)	
Pregabalin Pfizer		-R-		Rituximab (Riximyo)	
Pregnancy Tests - hCG Urine		RA-Morph		Rivaroxaban	
Premarin		Ralicrom		Rivastigmine	
Prevenar 13	310	Raloxifene hydrochloride	117	Rivastigmine Patch BNM 10	
Priadel		Raltegravir potassium		Rivastigmine Patch BNM 5	
Primaquine	103	Ramipex	121	Rivotril	129
Primidone	130	Ramipril	45	Riximyo	219
Primidone Clinect	130	Ranbaxy-Cefaclor	93	RIXUBIS	
Primolut N	86	Rapamune	252	Rizamelt	13
Priorix	307	Rasagiline	121	Rizatriptan	13
Probenecid	119	Reandron 1000		Robinul	
Probenecid-AFT	119	Recombinant factor IX	36, 38	Ronapreve	20
Procarbazine hydrochloride	162	Recombinant factor VIIa	37	Ropin	
Prochlorperazine		Recombinant factor VIII	38	Ropinirole hydrochloride	12
Prochlorperazine Brown &		Rectogesic	8	Rosuvastatin	
Burk	133	Redipred		Rosuvastatin Viatris	5
Prochlorperazine Max Health.		Relieve		Rotarix	
Proctofoam		Relistor	23	Rotavirus oral vaccine	
Proctosedyl		Remicade		Roxane-Propranolol	
Procyclidine hydrochloride		Renilon 7.5		Roxithromycin	9!
Progesterone		Resonium-A		Rubifen	14
Proglicem		Resource Beneprotein		Rubifen SR	
Progynova		Respiratory Devices		Rugby Capsaicin Topical Cream	
Prolia		Respiratory Stimulants		Musculoskeletal	
Promethazine hydrochloride		Retinol palmitate		Nervous	
Propafenone hydrochloride		ReTrieve		Rurioctocog alfa pegol [Recombi	
Propranolol		Retrovir		factor VIII]	
Propylene glycol		Revia		Ruxolitinib	
Propylthiouracil		Revolade		Rydapt	
Prostacur		Ribociclib		Rythmodan	
Protaphane		Riboflavin		Rythmodan - Cheplafarm	
Protaphane Penfill		Ribomustin		Rytmonorm	
Protifar		Ricit		-S-	
				•	
Protionamide		Rifabutin		Sabril	
Provera UD		Rifadin		Sacubitril with valsartan	
Provera HD	86	Rifadin Sanofi		SalAir	
Psoriasis and Eczema	70	Rifampicin		Salazopyrin	
Preparations		Rifaximin		Salazopyrin EN	
PTU		Rifinah		Salbutamol	25
Pulmicort Turbuhaler		Rilutek		Salbutamol with ipratropium	
Pulmozyme	262	Riluzole	122	bromide	259

INDEX: Generic Chemicals and Brands

Salicylic acid	73	Sodium cromoglicate		Sulprix	13
Salmeterol	257	Alimentary	8	Sumagran	13
Sandomigran	132	Sensory	268	Sumatriptan	13
Sandostatin LAR	175	Sodium Fusidate [fusidic acid]		Sunitinib	
Sanofi Primaquine	103	Dermatological		Sunitinib Pfizer	
Sapropterin dihydrochloride		Infection		Sunscreens	7
Scalp Preparations		Sensory	266	Sunscreens, proprietary	7
Scopolamine - Mylan	133	Sodium hyaluronate [Hyaluron	ic	Sustagen Hospital Formula	28
Sebizole		acid]		Sustagen Hospital Formula	
Secukinumab	230	Sodium phenylbutyrate	28	Active	28
Sedatives and Hypnotics	141	Sodium picosulfate		Sustanon Ampoules	
Seebri Breezhaler		Sodium polystyrene sulphonate	e43	Sylvant	23
Senna		Sodium tetradecyl sulphate		Symbicort Turbuhaler 100/6	
Senokot	24	Sodium valproate		Symbicort Turbuhaler 200/6	
SensoCard	15	Sofradex		Symbicort Turbuhaler 400/12	
Serc	132	Soframycin	266	Symmetrel	
Serenace		Solgar		Sympathomimetics	5
Seretide		Solifenacin succinate		Synacthen	
Seretide Accuhaler		Solifenacin succinate Max		Synacthen Depot	
Serevent		Health	81	Synacthene Retard	
Serevent Accuhaler		Solifenacin Viatris		Synagis	
Sertraline		Solu-Cortef		Synthroid	
Setrona		Solu-Medrol		Syntometrine	
Sevredol		Solu-Medrol-Act-O-Vial		Syrup (pharmaceutical grade)	27
Sex Hormones Non	120	Somatropin (Omnitrope)		Systane Unit Dose	
Contraceptive	84	Sotalol		- T -	21
Shingles vaccine		Spacer device		Tacrolimus	
Shingrix		Span-K		Dermatological	7
SII-Onco-BCG		Spazmol		Oncology	25
Sildenafil		Spinal Muscular Atrophy		Tacrolimus Sandoz	
Siltuximab		Spinraza		Tagrisso	
Simvastatin		Spiolto Respimat		Taliglucerase alfa	0ا
				Tambocor	۲
Simvastatin Mylan		Spiractin		Tambocor German	
Simvastatin Viatris		Spiriva Danimat			
Sinemet		Spiriva Respimat		Tamoxifen citrate	
Sinemet CR		Spironolactone		Tamoxifen Sandoz	
Sintetica Baclofen Intrathecal		Sporanox		Tamsulosin hydrochloride	
Sirolimus		Stalevo		Tamsulosin-Rex	
Sirturo		Staphlex		Tandem Cartridge	
Siterone		Stelara		Tandem t:slim X2 with Basal-IQ	1
Slow-Lopresor		Stemetil		Tandem t:slim X2 with	
Sodibic		Steril-Gene		Control-IQ	
Sodium acid phosphate		SteroClear		Tap water	
Sodium alginate		Stesolid		Taro	
Sodium benzoate	28	Stimulants/ADHD Treatments		Tasigna	
Sodium bicarbonate		Stiripentol		Tasmar	
Blood		Stomahesive		Taurine	
Extemporaneous	274	Strides Shasun		TCu 380 Plus Normal	7
Sodium calcium edetate	272	Stromectol		Tecentriq	
Sodium chloride		Sucralfate		Tecfidera	
Blood		Sulfadiazin-Heyl		Tegretol	
Respiratory		Sulfadiazine Silver		Tegretol AU	12
Sodium citrate with sodium laur		Sulfadiazine sodium	100	Tegretol CR	12
sulphoacetate		Sulfasalazine		Telfast	
Sodium citro-tartrate		Sulphur	73	Temaccord	
		•		Temazepam	
				· ·	

Temozolomide	162	Muscular Pain	116	TYR Lophlex LQ 20	299
Temozolomide-Taro		Topiramate		TYR Sphere 20	
Tenofovir disoproxil		Topiramate Actavis		Tyrosine1000	
Tenofovir Disoproxil Emtricitabil		Total parenteral nutrition (TPN)		Tysabri	
Viatr		TPN		- U -	10
Tenofovir Disoproxil Viatris		Tramadol hydrochloride		UK Synacthen	8,
Tenoxicam		Tramal SR 100		Ultibro Breezhaler	
Tensipine MR10		Tramal SR 150		Ultraproct	
Tepadina		Tramal SR 200		Umeclidinium	
Terbinafine		Trandate		Umeclidinium with vilanterol	
Terbutaline sulphate		Tranexamic acid		Univent25	
Teriflunomide		Tranylcypromine sulphate		Upadacitinib	
Teriflunomide Sandoz		Trastuzumab (Herzuma)		Ural	
Teriparatide		Trastuzumab deruxtecan		Urea	
Teriparatide - Teva		Trastuzumab emtansine		Urex Forte	
Testogel		Travatan		Urinary Agents	
Testosterone		Travoprost		Urinary Tract Infections	
Testosterone cipionate		Treatments for Dementia		UroFos	
Testosterone esters		Treatments for Substance	147	Uromitexan	
Testosterone undecanoate		Dependence	1.47	Ursodeoxycholic acid	
Tetrabenazine		•		Ursosan	
		Trelegy Ellipta		Ustekinumab	
Tetrabromophenol Tetracosactrin		Trental 400	54		
Tetracycline		Tretinoin	66	Utrogestan	
•		Dermatological		Vaccinations	201
Teva Lisinopril		Oncology		Vaclovir	
Teva-Ketoconazole Thalidomide		Trexate Triamcinolone acetonide	130	Valaciclovir	
			20		
Thalomid		Alimentary		Valganciclovir	
Theophylline	202	Dermatological		Valganciclovir Viatris	
Thiamine hydrochloride		Hormone Triamcinolone acetonide with	64	Valine50	
Thiamine multichem			tatia	Vancomycin	
THIO-TEPA		gramicidin, neomycin and nyst		Vannair	
Thioguanine		Dermatological		Varenicline Pfizer	
Thiotepa		Sensory		Variable variate	14
Thyroid and Antithyroid Agents		Trientine		Varicella vaccine [Chickenpox	044
Ticagrelor		Trientine Waymade		vaccine]	
Ticagrelor Sandoz		Trikafta		Varicella zoster vaccine [Shingles	
Tilcotil		Trimethoprim	100	vaccine]	
Timolol		Trimethoprim with		Varilrix	
Tiotropium bromide	259	sulphamethoxazole	400	Various	
Tiotropium bromide with	000	[Co-trimoxazole]		VariSoft	
olodaterol		Triovir		Vasodilators	
Tivicay		Trisequens		Vasopressin Agonists	
TMP	100	Trisul		Vasorex	
Tobramycin	400	Trophic Hormones		Vebulis	
Infection		Tropicamide		Vedafil	
Sensory	266	Trulicity		Vedolizumab	
Tobramycin (Viatris)		TruSteel		Vegzelma	20
Tobramycin BNM		Tryzan		Veletri	
Tobrex		Tuberculin PPD [Mantoux] test		Venclexta	
Tocilizumab		Tubersol		Venetoclax	
Tofranil		Two Cal HN		Venlafaxine	
Tolcapone		TYR Anamix Infant		Venomil	
Tolvaptan		TYR Anamix Junior		VENOX	
Topamax	131	TYR Anamix Junior LQ		Ventolin	
Topical Products for Joint and		TYR Explore 5	292	Vepesid	157

INDEX: Generic Chemicals and Brands

Verapamil hydrochloride49	XP Maxamum290
Vermox93	Xylocaine123
Versacloz134	Xylocaine 2% Jelly122
Vesanoid163	Xylocard 500123
Vexazone12	Xyntha38
Vfend102	- Z -
Viaderm KC69	Zapril44
Victoza12	Zarontin129
Vigabatrin131	Zaroxolyn51
Vigisom141	Zavedos
Vildagliptin12	Zeffix
Vildagliptin with metformin	Zejula159
hydrochloride12	Zematop73
	Zetlam
Vimpat	
Vinblastine sulphate	Ziagen111
Vincristine sulphate164	Zidovudine [AZT]111
Vinorelbine164	Zidovudine [AZT] with
Vinorelbine Ebewe164	lamivudine111
Vinorelbine Te Arai164	Ziextenzo42
Viramune Suspension110	Ziextenzo AU42
ViruPOS266	Zimybe53
Vit.D332	Zinc and castor oil70
Vita-B1231	Zinc sulphate34
VitA-POS270	Zincaps34
Vitabdeck32	Ziprasidone135
Vital281	Zista
Vitamin B complex31	Zithromax94
Vitamin B6 2531	Zo-Rub HP123
Vitamins31–32	Zo-Rub Osteo116
Vitarubin Depot Injection31	Zoladex91
Vivonex TEN281	Zoledronic acid
Voltaren	Hormone83
Voltaren D	Musculoskeletal118
Voltaren SR115	Zoledronic acid Viatris
Volumatic265	Hormone83
Voriconazole102	Musculoskeletal118
Votrient 169	Zopiclone142
Vttack102	Zopiclone Actavis142
Vyvanse144	Zostrix116
. W -	Zostrix HP123
Warfarin sodium41	Zuclopenthixol decanoate138
Wart Preparations74	Zuclopenthixol hydrochloride135
Wasp venom allergy treatment256	Zusdone135
Water	Zyban148
Blood43	•
	Zypine
Extemporaneous274	Zypine ODT135
White Soft Liquid Paraffin AFT70	Zyprexa Relprevv136
Wool fat with mineral oil70	Zytiga172
- X -	Zyvox105
Xaluprine155	
Xarelto41	
Xgeva116	
Xifaxan10	
XMET Maxamum289	
Xolair214	
Xolair AU214	