

August 2026

Volume 14

Editor:

Kaye Wilson, Doris Chong, Teresa Tay
email: enquiry@pharmac.govt.nz
Telephone +64 4 460 4990
Level 9, 40 Mercer Street
PO Box 10 254 Wellington 6143

Freephone Information Line

0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule, Section H for Hospital Pharmaceuticals (link to PDF copy) emailed to your nominated email address each month by subscribing at schedule.pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX.
XML version of the Schedule available from schedule.pharmac.govt.nz/pub/HML

Programmers

Anrik Drenth

email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency



ISSN 1179-3708 pdf

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/.

Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

Introducing Pharmac 2

Part I General Rules 4

Part II Alimentary Tract and Metabolism 5

Blood and Blood Forming Organs 30

Cardiovascular System 44

Dermatologicals 67

Genito-Urinary System 74

Hormone Preparations 78

Infections 89

Musculoskeletal System 115

Nervous System 121

Oncology Agents and Immunosuppressants 150

Respiratory System and Allergies 263

Sensory Organs 274

Various 281

Extemporaneous Compounds (ECPs) 289

Special Foods 292

Vaccines 310

Part III Optional Pharmaceuticals 321

Index 322

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at <https://pharmac.govt.nz/about>.

Glossary

Units of Measure

gram	g	microgram.....	mcg	millimole.....	mmol
kilogram.....	kg	milligram.....	mg	unit.....	u
international unit.....	iu	millilitre.....	ml		

Abbreviations

application	app	enteric coated.....	EC	solution.....	soln
capsule	cap	granules.....	grans	suppository.....	suppos
cream.....	crm	injection	inj	tablet.....	tab
dispersible	disp	liquid	liq	tincture.....	tinc
effervescent.....	eff	lotion	lotn		
emulsion	emul	ointment.....	oint		

HSS Hospital Supply Status

Guide to Section H listings

Example

ANATOMICAL HEADING			
	Price (ex man. Excl. GST) \$	Per	Brand or Generic Manufacturer
THERAPEUTIC HEADING			
Generic name listed by therapeutic group and subgroup	CHEMICAL A - Restricted see terms below ⚡ Presentation A.....10.00	100	Brand A
	➡ Restricted Only for use in children under 12 years of age		Brand or manufacturer's name
Indicates only presentation B1 is Restricted	CHEMICAL B - Some items restricted see terms below ⚡ Presentation B1.....1,589,00 Presentation B2 ➡ Restricted Oncologist or haematologist	1	Brand B1 e.g. Brand B2
From 1 January 2012 to 30 June 2014, at least 99% of the total volume of this item purchased must be Brand C	CHEMICAL C Presentation C - -1% DV Limit Jan-12 to 201415.00	28	Brand C
	CHEMICAL D - Restricted see terms below ⚡ Presentation D - -1% DV Limit Mar-13 to 201438.65	500	Brand D
Standard national price excluding GST	➡ Restricted <i>Limited to five weeks' treatment</i> Either: 1 For the prophylaxis of venous thromboembolism following a total hip replacement; or 2 For the prophylaxis of venous thromboembolism following a total knee replacement.		Quantity the Price applies to
Form and strength	CHEMICAL E Presentation E		e.g. Brand E
			Not a contracted product
⚡ Item restricted (see above); ⚡ Item restricted (see below) Products with Hospital Supply Status (HSS) are in bold			

PART I: GENERAL RULES

General Rules for Section H of the Pharmaceutical Schedule are included in Section A.

Read the [General Rules](https://pharmac.govt.nz/section-a) : <https://pharmac.govt.nz/section-a>.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antacids and Antiflatulents			
Antacids and Reflux Barrier Agents			
ALUMINIUM HYDROXIDE WITH MAGNESIUM HYDROXIDE AND SIMETICONE			
Tab 200 mg with magnesium hydroxide 200 mg and simeticone 20 mg			<i>e.g. Mylanta</i>
Oral liq 400 mg with magnesium hydroxide 400 mg and simeticone 30 mg per 5 ml			<i>e.g. Mylanta Double Strength</i>
SIMETICONE			
Oral drops 100 mg per ml			
Oral drops 20 mg per 0.3 ml			
Oral drops 40 mg per ml			
SODIUM ALGINATE WITH MAGNESIUM ALGINATE			
Powder for oral soln 225 mg with magnesium alginate 87.5 mg, sachet			<i>e.g. Gaviscon Infant</i>
SODIUM ALGINATE WITH SODIUM BICARBONATE AND CALCIUM CARBONATE			
Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg			<i>e.g. Gaviscon Extra Strength</i>
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml.....	7.50	500 ml	Acidex
SODIUM CITRATE			
Oral liq 8.8% (300 mmol/l).....	25.00	90 ml	Biomed
Phosphate Binding Agents			
ALUMINIUM HYDROXIDE			
Tab 600 mg			
CALCIUM CARBONATE – Restricted see terms below			
↓ Oral liq 250 mg per ml (100 mg elemental per ml).....	47.30	473 ml	Calcium carbonate PAI
	39.00	500 ml	Roxane
➔ Restricted (RS1698)			
Initiation			
Only when prescribed for patients unable to swallow calcium carbonate tablets or where calcium carbonate tablets are inappropriate..			
Antidiarrhoeals and Intestinal Anti-Inflammatory Agents			
Antipropulsives			
DIPHENOXYLATE HYDROCHLORIDE WITH ATROPINE SULPHATE			
Tab 2.5 mg with atropine sulphate 25 mcg			
LOPERAMIDE HYDROCHLORIDE			
Tab 2 mg.....	10.75	400	Nodia
Cap 2 mg – 5% DV Feb-26 to 2028	12.00	400	Diamide Relief
Rectal and Colonic Anti-Inflammatories			
BUDESONIDE – Restricted see terms on the next page			
↓ Cap modified-release 3 mg – 5% DV Dec-25 to 2028	33.38	90	Budesonide Te Arai
Products with Hospital Supply Status (HSS) are in bold			
Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS1723)

Initiation – Crohn's disease

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or
 - 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initiation – Collagenous and lymphocytic colitis (microscopic colitis)

Patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initiation – Gut Graft versus Host disease

Patient has gut Graft versus Host disease following allogenic bone marrow transplantation.

Initiation – non-cirrhotic autoimmune hepatitis

Re-assessment required after 6 months

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indications marked with * are unapproved indications.

Continuation – non-cirrhotic autoimmune hepatitis

Re-assessment required after 6 months

Treatment remains appropriate and the patient is benefitting from the treatment.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC free (14 applications)	57.09	15 g	Colifoam
---	-------	------	----------

HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE

Topical Aerosol foam, 1% with pramoxine hydrochloride 1%

MESALAZINE

Tab EC 400 mg	49.50	100	Asacol
	71.00	90	Octasa
Tab long-acting 500 mg	56.10	100	Pentasa
Tab 800 mg	85.50	90	Asacol
Tab 1,600 mg	85.50	60	Asacol
Modified release granules 1 g	118.10	100 g	Pentasa
Suppos 500 mg	22.80	20	Asacol
Suppos 1 g	50.96	28	Pentasa
Enema 1 g per 100 ml	41.30	7	Pentasa

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
OLSALAZINE			
Tab 500 mg	93.37	100	Dipentum
Cap 250 mg	53.00	100	Dipentum
SODIUM CROMOGLICATE			
Cap 100 mg			
SULFASALAZINE			
Tab 500 mg	19.49	100	Salazopyrin
Tab EC 500 mg	20.54	100	Salazopyrin EN

Local Preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

CINCHOCAINE HYDROCHLORIDE WITH HYDROCORTISONE			
Oint 5 mg with hydrocortisone 5 mg per g	17.03	30 g	Proctosedyl
Suppos 5 mg with hydrocortisone 5 mg per g	11.24	12	Proctosedyl
FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE			
Oint 950 mcg with fluocortolone pivalate 920 mcg and cinchocaine hydrochloride 5 mg per g	13.05	30 g	Ultraproct
Suppos 630 mcg with fluocortolone pivalate 610 mcg and cinchocaine hydrochloride 1 mg	8.61	12	Ultraproct
PREDNISOLONE HEXANOATE WITH CINCHOCAINE HYDROCHLORIDE			
Suppos 1.3 mg with cinchocaine hydrochloride 1 mg per g	8.61	12	Scheriproct

Management of Anal Fissures

GLYCERYL TRINITRATE			
Oint 0.2%	22.00	30 g	Rectogesic

Rectal Sclerosants

OILY PHENOL [PHENOL OILY]			
Inj 5%, 5 ml vial			

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – 5% DV Jul-26 to 2028	11.99	10	Glycopyrronium-AFT
HYOSCINE BUTYLBROMIDE			
Tab 10 mg – 5% DV Apr-25 to 2027	2.25	20	Hyoscine Butylbromide (Adiramedica)
Inj 20 mg, 1 ml ampoule	1.91	5	Spazmol
MEBEVERINE HYDROCHLORIDE			
Tab 135 mg – 5% DV Dec-26 to 2029	8.00	90	Colofac

Antulcerants

Antisecretory and Cytoprotective

MISOPROSTOL			
Tab 200 mcg	47.73	120	Cytotec

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
H2 Antagonists			
CIMETIDINE			
Tab 200 mg			
Tab 400 mg			
FAMOTIDINE			
Tab 20 mg			
Tab 40 mg	10.27	100	Famotidine Hovid MY
Inj 10 mg per ml, 2 ml vial			
Inj 10 mg per ml, 4 ml vial			
RANITIDINE – Restricted see terms below			
↓ Tab 150 mg			
↓ Tab 300 mg			
↓ Inj 25 mg per ml, 2 ml ampoule			
➔ Restricted (RS1703)			
Initiation			
Either:			
1 For continuation use; or			
2 Routine prevention of allergic reactions..			
Proton Pump Inhibitors			
LANSOPRAZOLE			
Cap 15 mg – 5% DV Feb-25 to 2027	1.21	30	Lanzol Relief
	4.04	100	Lanzol Relief
Cap 30 mg – 5% DV Feb-25 to 2027	5.43	100	Lanzol Relief
Tab orodispersible 30 mg	5.43	100	Lansoprazole (Viatris)
OMEPRAZOLE			
↓ Tab dispersible 10 mg			
➔ Restricted (RS1027)			
Initiation			
Only for use in tube-fed patients.			
↓ Tab dispersible 20 mg			
➔ Restricted (RS1027)			
Initiation			
Only for use in tube-fed patients.			
Cap 10 mg	2.06	90	Omeprazole Teva
			Omeprazole actavis 10
Cap 20 mg	2.02	90	Omeprazole Teva
			Omeprazole actavis 20
Cap 40 mg	3.18	90	Omeprazole Teva
			Omeprazole actavis 40
Powder for oral liq.....	52.00	5 g	Midwest
Inf 40 mg vial – 5% DV Aug-26 to 2029	14.50	5	Omezol IV
Inj 40 mg ampoule with diluent.....	37.38	5	Dr Reddy's Omeprazole
PANTOPRAZOLE			
Tab EC 20 mg – 5% DV May-26 to 2028	1.81	90	Panzop Relief
Tab EC 40 mg – 5% DV May-26 to 2028	2.70	90	Panzop Relief
Inj 40 mg vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Site Protective Agents

COLLOIDAL BISMUTH SUBCITRATE

Tab 120 mg 14.51 50 Gastrodenol

SUCRALFATE

Tab 1 g

Bile and Liver Therapy

L-ORNITHINE L-ASPARTATE – **Restricted** see terms [below](#)

↓ Grans for oral liquid 3 g

→ **Restricted (RS1261)**

Initiation

For patients with chronic hepatic encephalopathy who have not responded to treatment with, or are intolerant to lactulose, or where lactulose is contraindicated.

RIFAXIMIN – **Restricted** see terms [below](#)

↓ Tab 550 mg – 5% DV **Dec-24 to 2027** 625.00 56 **Xifaxan**

→ **Restricted (RS1416)**

Initiation

For patients with hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Diabetes

Alpha Glucosidase Inhibitors

ACARBOSE

Tab 50 mg – 5% DV **Feb-25 to 2027** 11.20 90 **Accarb**

Tab 100 mg – 5% DV **Feb-25 to 2027** 17.38 90 **Accarb**

Hyperglycaemic Agents

DIAZOXIDE – **Restricted** see terms [below](#)

↓ Cap 25 mg 110.00 100 Proglidem

↓ Cap 100 mg 280.00 100 Proglidem

↓ Oral liq 50 mg per ml 620.00 30 ml Proglycem

→ **Restricted (RS1028)**

Initiation

For patients with confirmed hypoglycaemia caused by hyperinsulinism.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit 32.00 1 Glucagen Hypokit

GLUCOSE [DEXTROSE]

Tab 1.5 g

Tab 3.1 g

Tab 4 g

Oral soln 15 g per 80 ml sachet 75.00 50 HypoPak Glucose

Gel 40%

GLUCOSE WITH SUCROSE AND FRUCTOSE

Gel 19.7% with sucrose 35% and fructose 19.7%, 18 g sachet

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Insulin - Intermediate-Acting Preparations			
INSULIN ASPART WITH INSULIN ASPART PROTAMINE			
Inj insulin aspart 30% with insulin aspart protamine 70%, 100 u per ml, 3 ml prefilled pen	52.15	5	NovoMix 30 FlexPen
INSULIN DEGLUDEC WITH INSULIN ASPART			
Inj degludec 70 u with insulin aspart 30 u, 100 u per ml, 3 ml.....	80.00	5	Ryzodeg 70/30 Penfill
INSULIN ISOPHANE			
Inj insulin human 100 u per ml, 10 ml vial			
Inj insulin human 100 u per ml, 3 ml cartridge			
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj insulin lispro 25% with insulin lispro protamine 75%, 100 u per ml, 3 ml cartridge.....	42.66	5	Humalog Mix 25
Inj insulin lispro 50% with insulin lispro protamine 50%, 100 u per ml, 3 ml cartridge.....	42.66	5	Humalog Mix 50
INSULIN NEUTRAL WITH INSULIN ISOPHANE			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 10 ml vial			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 40% with insulin isophane 60%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 50% with insulin isophane 50%, 100 u per ml, 3 ml cartridge			
Insulin - Long-Acting Preparations			
INSULIN GLARGINE			
Inj 100 u per ml, 3 ml disposable pen.....	94.50	5	Lantus SoloStar
Inj 100 u per ml, 3 ml cartridge.....	94.50	5	Lantus
Inj 100 u per ml, 10 ml vial.....	63.00	1	Lantus
Insulin - Rapid-Acting Preparations			
INSULIN ASPART			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
Inj 100 u per ml, 3 ml syringe	51.19	5	NovoRapid FlexPen
INSULIN GLULISINE			
Inj 100 u per ml, 10 ml vial.....	27.03	1	Apidra
Inj 100 u per ml, 3 ml cartridge	46.07	5	Apidra
Inj 100 u per ml, 3 ml disposable pen.....	46.07	5	Apidra Solostar
INSULIN LISPRO			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
Insulin - Short-Acting Preparations			
INSULIN NEUTRAL			
Inj human 100 u per ml, 10 ml vial			
Inj human 100 u per ml, 3 ml cartridge			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
Tab 5 mg	7.50	100	Daonil
GLICLAZIDE			
Tab 80 mg	20.10	500	Glizide
GLIPIZIDE			
Tab 5 mg – 5% DV Mar-25 to 2027	6.86	100	Minidiab
METFORMIN HYDROCHLORIDE			
Tab immediate-release 500 mg – 1% DV Mar-23 to 2027	14.74	1,000	Metformin Viatris
Tab immediate-release 850 mg – 1% DV Aug-23 to 2027	11.28	500	Metformin Viatris
PIOGLITAZONE			
Tab 15 mg – 5% DV Dec-24 to 2027	6.15	90	Vexazone
Tab 30 mg – 5% DV Dec-24 to 2027	7.25	90	Vexazone
Tab 45 mg – 5% DV Dec-24 to 2027	12.00	90	Vexazone
VILDAGLIPTIN			
Tab 50 mg	35.00	60	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE			
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	Galvumet

GLP-1 Agonists

DULAGLUTIDE – **Restricted** see terms [below](#)

Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.

↓ Inj 1.5 mg per 0.5 ml prefilled pen 115.23 4 Trulicity

→ **Restricted (RS2135)**

Initiation

Either:

- 1 For continuation use; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and
 - 2.3 Any of the following:
 - 2.3.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack,

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin / empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

LIRAGLUTIDE – **Restricted** see terms [below](#)

Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.

⚡ Inj 6 mg per ml, 3 ml prefilled pen383.72 3 Victoza

➡ **Restricted (RS2136)**

Initiation

Either:

- 1 For continuation use; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and
 - 2.3 Any of the following:
 - 2.3.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin / empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

SGLT2 Inhibitors

➡ **Restricted (RS2069)**

Initiation – heart failure reduced ejection fraction

All of the following:

- 1 Patient has heart failure; and
- 2 Patient is in NYHA functional class II or III or IV; and

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

3 Either:

- 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or
- 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initiation – Type 2 Diabetes

Any of the following:

- 1 For continuation use; or
- 2 Patient has previously had an initial approval for a GLP-1 agonist; or
- 3 All of the following:
 - 3.1 Patient has type 2 diabetes; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 3.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 3.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 3.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 3.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 3.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause.
- c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride) for the treatment of heart failure.

EMPAGLIFLOZIN – **Restricted** see terms [on the previous page](#)

† Tab 10 mg	58.56	30	Jardiance
† Tab 25 mg	58.56	30	Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – **Restricted** see terms [on the previous page](#)

† Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
† Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
† Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
† Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Digestives Including Enzymes

PANCREATIC ENZYME

Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph Eur U)	34.93	20 g	Creon Micro
Powder pancreatin 60.12 mg (3,600 Ph. Eur. u/amylase, 5,000 Ph. Eur. u/lipase and 200 Ph. Eur. u/protease)			

URSODEOXYCHOLIC ACID – **Restricted** see terms [below](#)

⚡ Cap 250 mg.....	62.00	100	APO-Ursodeoxycholic Acid Ursosan
	33.95		

➡ **Restricted** (RS2103)

Initiation – **Alagille syndrome or progressive familial intrahepatic cholestasis**

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initiation – **Chronic severe drug induced cholestatic liver injury**

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initiation – **Primary biliary cholangitis**

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initiation – **Pregnancy**

Patient diagnosed with cholestasis of pregnancy.

Initiation – **Haematological transplant**

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogeneic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initiation – **Total parenteral nutrition induced cholestasis**

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by TPN; and
- 2 Liver function has not improved with modifying the TPN composition.

Initiation – **prevention of sinusoidal obstruction syndrome**

The individual has leukaemia/lymphoma and requires prophylaxis for medications/therapies with a high risk of sinusoidal obstruction syndrome.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Laxatives			
Bowel-Cleansing Preparations			
CITRIC ACID WITH MAGNESIUM CARBONATE HYDRATE AND SODIUM PICOSULFATE			
Powder for oral soln 12 g with magnesium carbonate hydrate 7.4 g and sodium picosulfate 10 mg per sachet			<i>e.g. PicoPrep Orange</i>
MACROGOL 3350 WITH ASCORBIC ACID, POTASSIUM CHLORIDE, SODIUM CHLORIDE AND CITRIC ACID WITH MAGNESIUM CARBONATE HYDRATE AND SODIUM PICOSULFATE			
Powder for oral soln 52.9 g with ascorbic acid 6 g, potassium chloride 740 mg, sodium chloride 2.6 g and sodium sulphate 5.6 g per sachet (1) and powder for oral soln citric acid 12 g with magnesium carbonate hydrate 7.4 g and sodium picosulfate 10 mg per sachet (2)			<i>e.g. Prepkit Orange</i>
MACROGOL 3350 WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE			
Powder for oral soln 755.68 mg with potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 70 g sachet – 5% DV Feb-25 to 2027	16.10	3	Glycoprep Orange
	64.32	12	Glycoprep Orange
Powder for oral soln 755.68 mg with potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 210 g sachet			<i>e.g. Glycoprep Orange</i>
MACROGOL 3350 WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE WITH/WITHOUT SODIUM SULFATE, SODIUM ASCORBATE, ASCORBIC ACID			
Powd for oral soln 100g with potassium chloride 1g, sodium chloride 2g and sodium sulfate 9g per sach(1), powd for oral soln 40g with potassium chloride 1.2g and sodium chloride 3.2g per sach(1) and powd for oral soln ascorbic acid 7.54g and sodium ascorbate 48.11g per sach(1) – 5% DV Dec-26 to 2029	18.52	3	Plenvu
Bulk-Forming Agents			
ISPAGHULA (PSYLLIUM) HUSK			
Powder for oral soln	22.10	500 g	Konsyl-D
STERCULIA WITH FRANGULA – Restricted: For continuation only			
➡ Powder for oral soln			
Faecal Softeners			
DOCUSATE SODIUM			
Tab 50 mg	3.20	100	Coloxyl
Tab 120 mg	4.98	100	Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES			
Tab 50 mg with sennosides 8 mg – 5% DV May-26 to 2028	1.50	100	Solax
PARAFFIN			
Oral liquid 1 mg per ml			
Enema 133 ml			
POLOXAMER			
Oral drops 10%	4.17	30 ml	Coloxyl

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Opioid Receptor Antagonists - Peripheral			
METHYLNALTREXONE BROMIDE – Restricted see terms below			
↓ Inj 12 mg per 0.6 ml vial	36.00	1	Relistor
	246.00	7	Relistor

➔ **Restricted (RS2057)**

Initiation – Opioid induced constipation

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Initiation – Opioid induced constipation outside of palliative care

Limited to 14 days treatment

All of the following:

- 1 Individual has opioid induced constipation; and
- 2 Oral and rectal treatments for opioid induced constipation, including bowel-cleansing preparations, are ineffective or inappropriate; and
- 3 Mechanical bowel obstruction has been excluded.

Osmotic Laxatives

GLYCEROL

Suppos 2.8/4.0 g – 5% DV Feb-26 to 2028	12.39	20	Lax-suppositories Glycerol
---	-------	----	---------------------------------------

Note: DV limit applies to glycerol suppository presentations.

LACTULOSE

Oral liq 10 g per 15 ml – 5% DV Feb-26 to 2028	6.16	500 ml	Laevolac
--	------	--------	-----------------

MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE

Powder for oral soln 6.563 g with potassium chloride 23.3 mg, sodium bicarbonate 89.3 mg and sodium chloride 175.4 mg			
Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg	10.15	30	Molaxole Movicol

SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE

Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml – 5% DV Feb-26 to 2028	36.89	50	Micolette
---	-------	----	------------------

SODIUM PHOSPHATE WITH PHOSPHORIC ACID

Oral liq 16.4% with phosphoric acid 25.14%			
Enema 10% with phosphoric acid 6.58%	3.70	1	Fleet Phosphate Enema

Stimulant Laxatives

BISACODYL

Tab 5 mg – 5% DV Jul-26 to 2028	6.28	200	Bisacodyl-AFT
Suppos 10 mg – 5% DV Feb-25 to 2027	4.14	10	Lax-Suppositories

SENNOSIDES

Tab 7.5 mg

SODIUM PICOSULFATE – **Restricted** see terms [on the next page](#)

↓ Oral soln 7.5 mg per ml	7.40	30 ml	Dulcolax SP Drop
---------------------------------	------	-------	-------------------------

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

➔ **Restricted (RS1843)**

Initiation

Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation.

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA – **Restricted** see terms [below](#)

↓ Inj 50 mg vial 1,142.60 1 Myozyme

➔ **Restricted (RS1793)**

Initiation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Continuation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

ARGININE

- Tab 1,000 mg
- Cap 500 mg
- Powder
- Inj 500 mg per ml, 10 ml vial
- Inj 600 mg per ml, 25 ml vial

BETAININE – **Restricted** see terms [below](#)

↓ Powder for oral soln.....575.00 180 g Cystadane

→ **Restricted (RS1794)**

Initiation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Continuation

Metabolic physician

Re-assessment required after 12 months

The treatment remains appropriate and the patient is benefiting from treatment.

BIOTIN – **Restricted** see terms [below](#)

- ↓ Cap 50 mg
- ↓ Cap 100 mg
- ↓ Inj 10 mg per ml, 5 ml vial
- ↓ Inj 5 mg per ml, 1 ml ampoule

→ **Restricted (RS1330)**

Metabolic physician or metabolic disorders dietitian

CARGLUMIC ACID – **Restricted** see terms [below](#)

- ↓ Tab disp 200 mg

→ **Restricted (RS1831)**

Initiation

Metabolic physician

For the acute in-patient treatment of organic acidaemias as an alternative to haemofiltration.

COENZYME Q10 – **Restricted** see terms [below](#)

- ↓ Cap 120 mg
- ↓ Cap 160 mg

→ **Restricted (RS1832)**

Initiation

Metabolic physician

Re-assessment required after 6 months

The patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Continuation

Metabolic physician

Re-assessment required after 24 months

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE – **Restricted** see terms [below](#)

↓ Inj 1 mg per ml, 5 ml vial.....2,234.00 1 Naglazyme

→ **Restricted (RS1795)**

Initiation

Metabolic physician

Re-assessment required after 12 months

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency confirmed by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Continuation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

HAEM ARGINATE

Inj 25 mg per ml, 10 ml ampoule

IDURSULFASE – **Restricted** see terms [below](#)

↓ Inj 2 mg per ml, 3 ml vial.....4,608.30 1 Elaprase

→ **Restricted (RS1546)**

Initiation

Metabolic physician

Limited to 24 weeks treatment

All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE – **Restricted** see terms [on the next page](#)

↓ Inj 100 U per ml, 5 ml vial.....1,335.16 1 Aldurazyme

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS1607)

Initiation

Metabolic physician

Limited to 24 weeks treatment

All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysaccharidosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE – **Restricted** see terms [below](#)

- ⚡ Tab 500 mg
- ⚡ Cap 250 mg
- ⚡ Cap 500 mg
- ⚡ Oral liq 500 mg per 10 ml
- ⚡ Oral soln 1,000 mg per 10 ml
- ⚡ Oral soln 1,100 mg per 15 ml
- ⚡ Inj 200 mg per ml, 5 ml vial

➔ Restricted (RS1035)

Neurologist, metabolic physician or metabolic disorders dietitian

NITISINONE – **Restricted** see terms [below](#)

⚡ Cap 2 mg – 5% DV Jul-26 to 2028	676.00	60	Nitisinone LogixX Pharma
⚡ Cap 5 mg – 5% DV Jul-26 to 2028	1,302.00	60	Nitisinone LogixX Pharma
⚡ Cap 10 mg – 5% DV Jul-26 to 2028	1,704.00	60	Nitisinone LogixX Pharma

➔ Restricted (RS2164)

Initiation

Patient requires nitisinone for the management of inherited metabolic disorders.

PYRIDOXAL-5-PHOSPHATE – **Restricted** see terms [below](#)

- ⚡ Tab 50 mg

➔ Restricted (RS1331)

Neurologist, metabolic physician or metabolic disorders dietitian

RIBOFLAVIN – **Restricted** see terms [below](#)

- ⚡ Tab 100 mg
- ⚡ Cap 100 mg

➔ Restricted (RS1833)

Initiation

Metabolic physician or neurologist

Re-assessment required after 6 months

The patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation

Metabolic physician or neurologist

Re-assessment required after 24 months

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE – **Restricted** see terms [below](#)

↓ Tab soluble 100 mg 1,452.70 30 Kuvan

→ **Restricted (RS1796)**

Initiation

Metabolic physician

Re-assessment required after 1 month

All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Continuation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE

Cap 500 mg

Powder

Soln 100 mg per ml

Inj 20%, 10 ml ampoule

SODIUM PHENYLBUTYRATE – **Some items restricted** see terms [on the next page](#)

Tab 500 mg

↓ Grans 483 mg per g 2,016.00 174 g Pheburane

Oral liq 250 mg per ml

Inj 200 mg per ml, 10 ml ampoule

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS1797)

Initiation

Metabolic physician

Re-assessment required after 12 months

For the chronic management of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Continuation

Metabolic physician

Re-assessment required after 12 months

The treatment remains appropriate and the patient is benefiting from treatment.

TALIGLUCERASE ALFA – **Restricted** see terms [below](#)

‡ Inj 200 unit vial.....	1,072.00	1	ElELYso
--------------------------	----------	---	---------

➔ Restricted (RS1897)

Initiation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Continuation

Metabolic physician or any relevant practitioner on the recommendation of a metabolic physician

Re-assessment required after 3 years

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

TAURINE – **Restricted** see terms [below](#)

- ↓ Cap 500 mg
- ↓ Cap 1,000 mg
- ↓ Powder

→ **Restricted** (RS1834)

Initiation

Metabolic physician

Re-assessment required after 6 months

The patient has a suspected specific mitochondrial disorder that may respond to taurine supplementation.

Continuation

Metabolic physician

Re-assessment required after 24 months

Both:

- 1 The patient has a confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE – **Restricted** see terms [below](#)

↓ Cap 250 mg 2,022.00 100 Trientine Waymade

→ **Restricted** (RS2026)

Initiation

All of the following:

- 1 Patient has confirmed Wilson disease; and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
- 3 Treatment with zinc has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

Minerals

Calcium

CALCIUM CARBONATE

Tab 1.25 g (500 mg elemental) 7.28 250 Calci-Tab 500
 Tab eff 1.25 g (500 mg elemental)
 Tab eff 1.75 g (1 g elemental)

Copper

→ **Restricted** (RS1928)

Initiation – Moderate to severe burns

Limited to 3 months treatment

Both:

- 1 Patient has been hospitalised with moderate to severe burns; and
- 2 Treatment is recommended by a National Burns Unit specialist.

COPPER – **Restricted** see terms [above](#)

↑ Tab 2.5 mg, chelated

COPPER CHLORIDE – **Restricted** see terms [above](#)

↑ Inj 0.4 mg per ml, 10 ml vial

ALIMENTARY TRACT AND METABOLISM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Fluoride			
SODIUM FLUORIDE			
Tab 1.1 mg (0.5 mg elemental)			
Iodine			
POTASSIUM IODATE			
Tab 253 mcg (150 mcg elemental iodine)	5.99	90	NeuroTabs
POTASSIUM IODATE WITH IODINE			
Oral liq 10% with iodine 5%			
Iron			
FERRIC DERISOMALTOSE – Restricted see terms below			
⚡ Inj 500 mg per 5 ml vial	249.99	1	Monofer
➡ Restricted (RS2175)			
Initiation			
Patient had previously developed iron-infusion related hypophosphataemia or other severe adverse reaction.			
FERROUS FUMARATE			
Tab 200 mg (65.7 mg elemental) – 5% DV Feb-25 to 2027	3.49	100	Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID			
Tab 310 mg (100 mg elemental) with folic acid 350 mcg – 5% DV Dec-24 to 2027	5.98	100	Ferro-F-Tabs
FERROUS GLUCONATE WITH ASCORBIC ACID			
Tab 170 mg (20 mg elemental) with ascorbic acid 40 mg			
FERROUS SULFATE			
Oral liq 30 mg (6 mg elemental) per ml – 5% DV Feb-26 to 2028	10.25	250 ml	Ferro-Liquid
FERROUS SULFATE WITH ASCORBIC ACID			
Tab long-acting 325 mg (105 mg elemental) with ascorbic acid 500 mg			
IRON (AS FERRIC CARBOXYMALTOSE) – Restricted see terms below			
⚡ Inj 50 mg per ml, 10 ml vial	150.00	1	Ferinject
➡ Restricted (RS1417)			
Initiation			
Treatment with oral iron has proven ineffective or is clinically inappropriate.			
IRON (AS SUCROSE)			
Inj 20 mg per ml, 5 ml ampoule	100.00	5	Venofer
<i>(Venofer Inj 20 mg per ml, 5 ml ampoule to be delisted 1 September 2026)</i>			
IRON POLYMALTOSE			
Inj 50 mg per ml, 2 ml ampoule	41.75	5	Ferrosig
Magnesium			
MAGNESIUM AMINO ACID CHELATE			
Cap 750 mg (150 mg elemental)			
MAGNESIUM CHLORIDE			
Inj 1 mmol per 1 ml, 100 ml bag			
MAGNESIUM HYDROXIDE			
Tab 311 mg (130 mg elemental)			
Suspension 8%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MAGNESIUM OXIDE			
Cap 663 mg (400 mg elemental)			
Cap 696 mg (420 mg elemental)			
MAGNESIUM OXIDE WITH MAGNESIUM ASPARTATE, MAGNESIUM AMINO ACID CHELATE AND MAGNESIUM CITRATE			
Cap 500 mg with magnesium aspartate 100 mg, magnesium amino acid chelate 100 mg and magnesium citrate 100 mg (345 mg elemental magnesium)			
Cap 500 mg with magnesium aspartate 100 mg, magnesium amino acid chelate 100 mg and magnesium citrate 100 mg (360 mg elemental magnesium)			
<i>(Any Cap 500 mg with magnesium aspartate 100 mg, magnesium amino acid chelate 100 mg and magnesium citrate 100 mg (360 mg elemental magnesium) to be delisted 1 November 2026)</i>			
MAGNESIUM SULPHATE			
Inj 100 mg per ml, 40 ml bag			
Inj 0.4 mmol per ml, 250 ml bag			
Inj 2 mmol per ml, 10 ml ampoule	75.06	10	Inresa
Inj 2 mmol per ml, 5 ml ampoule – 5% DV Feb-27 to 2029	26.27	10	Martindale
Inj 100 mg per ml, 50 ml bag			

Selenium

SELENIUM – **Restricted** see terms [below](#)

↓ Oral liq 150 mcg per 3 drops

e.g. Clinicians selenium oral drops

↓ Inj 300 mcg per ml, 1 ml ampoule

→ **Restricted (RS1929)**

Initiation – Moderate to severe burns

Limited to 3 months treatment

Both:

- 1 Patient has been hospitalised with moderate to severe burns; and
- 2 Treatment is recommended by a National Burns Unit specialist.

Zinc

ZINC

Oral liq 5 mg per 5 drops

ZINC CHLORIDE

Inj 5.3 mg per ml (5.1 mg per ml elemental), 2 ml ampoule

ZINC SULPHATE

Cap 220 mg (50 mg elemental)	29.14	100	Rugby
Cap 137.4 mg (50 mg elemental)	11.00	100	Zincaps

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

Soln 0.15%

Spray 0.15%

Spray 0.3%

ALIMENTARY TRACT AND METABOLISM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BENZDAMINE HYDROCHLORIDE WITH CETYLPYRIDINIUM CHLORIDE Lozenge 3 mg with cetylpyridinium chloride			
CARBOXYMETHYLCELLULOSE Oral spray			
CARMELLOSE SODIUM WITH PECTIN AND GELATINE Paste Powder			
CHLORHEXIDINE GLUCONATE Mouthwash 0.2% – 5% DV Jan-25 to 2027	3.99	200 ml	healthE
DICHLOROBENZYL ALCOHOL WITH AMYLMETACRESOL Lozenge 1.2 mg with amylmetacresol 0.6 mg			
TRIAMCINOLONE ACETONIDE Paste 0.1% – 5% DV Dec-26 to 2029	3.57	5 g	Kenalog in Orabase

Oropharyngeal Anti-Infectives

AMPHOTERICIN B Lozenge 10 mg	5.86	20	Fungilin
MICONAZOLE Oral gel 20 mg per g – 5% DV Feb-25 to 2027	5.19	40 g	Decozol
NYSTATIN Oral liquid 100,000 u per ml	2.22	24 ml	Nilstat

Other Oral Agents

HYALURONIC ACID WITH LIDOCAINE [LIGNOCAINE] Inj 20 mg per ml			
SILVER DIAMINE FLUORIDE Oral application 38%	139.00	5 ml	Topamine
SODIUM HYALURONATE [HYALURONIC ACID] – Restricted see terms below ⚡ Inj 20 mg per ml, 1 ml syringe ➡ Restricted (RS1175) Otolaryngologist			

Vitamins

Multivitamin Preparations

MULTIVITAMIN AND MINERAL SUPPLEMENT – Restricted see terms below ⚡ Cap	23.35	180	Clinicians Multivit & Mineral Boost
--	-------	-----	--

➡ **Restricted (RS1498)**

Initiation

Limited to 3 months treatment

Both:

- 1 Patient was admitted to hospital with burns; and
- 2 Any of the following:
 - 2.1 Burn size is greater than 15% of total body surface area (BSA) for all types of burns; or
 - 2.2 Burn size is greater than 10% of BSA for mid-dermal or deep dermal burns; or
 - 2.3 Nutritional status prior to admission or dietary intake is poor.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MULTIVITAMIN RENAL – Restricted see terms below			
↓ Cap	7.28	30	Clinicians Renal Vit
→ Restricted (RS1499)			
Initiation			
Either:			
1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or			
2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73m ² body surface area (BSA).			
MULTIVITAMINS			
Tab (BPC cap strength) – 5% DV Dec-26 to 2029	21.65	1,000	Mvite
↓ cap vitamin A 2500 u, betacarotene 3 mg, cholecalciferol 11 mcg, alpha tocopherol 150 u, phytomenadione 150 mcg, folic acid 0.2 mg, ascorbic acid 100 mg, thiamine 1.5 mg, pantothenic acid 12 mg, riboflavin 1.7 mg, niacin 20 mg, pyridoxine hydrochloride 1.9 mg, cyanocobalamin 3 mcg, zinc 7.5 mg and biotin 100 mcg			<i>e.g. Vitabdeck</i>
→ Restricted (RS1620)			
Initiation			
Any of the following:			
1 Patient has cystic fibrosis with pancreatic insufficiency; or			
2 Patient is an infant or child with liver disease or short gut syndrome; or			
3 Patient has severe malabsorption syndrome.			
↓ Powder vitamin A 3200 mcg with vitamin D 100 mcg, vitamin E 54.2 mg, vitamin C 400 mg, vitamin K1 108 mcg thiamine 3.2 mg, riboflavin 4.4 mg, niacin 41 mg, vitamin B6 3.6 mg, folic acid 600 mcg, vitamin B12 9 mcg, biotin 120 mcg, pantothenic acid 24 mg, choline 1250 mg and inositol 700 mg	74.88	200 g	Paediatric Seravit
→ Restricted (RS1178)			
Initiation			
Patient has inborn errors of metabolism.			
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg and glucose 1000 mg, 5 ml ampoule (1)			<i>e.g. Pabrinex IV</i>
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg, 2 ml ampoule (1)			
Inj thiamine hydrochloride 500 mg with riboflavin 8 mg and pyridoxine hydrochloride 100 mg, 10 ml ampoule (1) and inj ascorbic acid 1000 mg with nicotinamide 320 mg and glucose 2000 mg, 10 ml ampoule (1)			

Vitamin A

RETINOL

Tab 10,000 iu

Cap 25,000 iu

Oral liq 150,000 iu per ml

Oral liq 666.7 mcg per 2 drops, 10 ml

Oral liq 5,000 iu per drop, 30 ml

ALIMENTARY TRACT AND METABOLISM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Vitamin B			
HYDROXOCOBALAMIN			
Inj 1 mg per ml, 1 ml ampoule – 5% DV Jul-25 to 2027	3.95	3	Hydroxocobalamin Panpharma
PYRIDOXINE HYDROCHLORIDE			
Tab 25 mg	3.43	90	Vitamin B6 25
Tab 50 mg	23.45	500	Pyridoxine multichem
Inj 100 mg per ml, 2 ml vial			
Inj 100 mg per ml, 1 ml ampoule			
Inj 100 mg per ml, 30 ml vial			
THIAMINE HYDROCHLORIDE			
Tab 50 mg – 5% DV Dec-26 to 2028	4.65	100	Thiamine multichem
	3.89		Thiamine-REX
Tab 100 mg			
Inj 100 mg per ml, 1 ml vial			<i>e.g. Benerva</i>
Inj 50 mg per ml, 2 ml ampoule – 5% DV Sep-26 to 2028	49.95	10	Thiamine Sterop
Inj 100 mg per ml, 2 ml vial			
Inj 125 mg per ml, 2 ml vial			
<i>(Thiamine multichem Tab 50 mg to be delisted 1 December 2026)</i>			
VITAMIN B COMPLEX			
Tab strong, BPC	11.25	500	Bplex
Vitamin C			
ASCORBIC ACID			
Tab 100 mg – 5% DV Mar-26 to 2028	16.00	500	Cvite
Tab chewable 250 mg			
Vitamin D			
ALFACALCIDOL			
Cap 0.25 mcg	26.32	100	One-Alpha
Cap 1 mcg	87.98	100	One-Alpha
Oral drops 2 mcg per ml	60.68	20 ml	One-Alpha
CALCITRIOL			
Cap 0.25 mcg	7.89	100	Calcitriol XL
			Calcitriol-AFT
Cap 0.5 mcg	13.68	100	Calcitriol XL
			Calcitriol-AFT
Oral liq 1 mcg per ml			
Inj 1 mcg per ml, 1 ml ampoule			
COLECALCIFEROL			
Cap 1.25 mg (50,000 iu)	3.65	12	Vit.D3
Oral liq 188 mcg per ml (7,500 iu per ml)	9.00	5 ml	Clinicians
Vitamin E			
ALPHA TOCOPHERYL – Restricted see terms on the next page			
⚡ Oral liq 156 u per ml			

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

➔ **Restricted (RS1632)**

Initiation – Cystic fibrosis

Both:

- 1 Cystic fibrosis patient; and
- 2 Either:
 - 2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
 - 2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

Initiation – Osteoradionecrosis

For the treatment of osteoradionecrosis.

Initiation – Other indications

All of the following:

- 1 Infant or child with liver disease or short gut syndrome; and
- 2 Requires vitamin supplementation; and
- 3 Either:
 - 3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
 - 3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

ALPHA TOCOPHERYL ACETATE – **Restricted** see terms [below](#)

↓ Cap 100 u

↓ Cap 500 u

↓ Oral liq 156 u per ml

➔ **Restricted (RS1176)**

Initiation – Cystic fibrosis

Both:

- 1 Cystic fibrosis patient; and
- 2 Either:
 - 2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
 - 2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

Initiation – Osteoradionecrosis

For the treatment of osteoradionecrosis.

Initiation – Other indications

All of the following:

- 1 Infant or child with liver disease or short gut syndrome; and
- 2 Requires vitamin supplementation; and
- 3 Either:
 - 3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
 - 3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Antianaemics

Hypoplastic and Haemolytic

EPOETIN ALFA – **Restricted** see terms [below](#)

⚡ Inj 1,000 iu in 0.5 ml syringe.....	250.00	6	Binocrit
⚡ Inj 2,000 iu in 1 ml syringe.....	100.00	6	Binocrit
⚡ Inj 3,000 iu in 0.3 ml syringe.....	150.00	6	Binocrit
⚡ Inj 4,000 iu in 0.4 ml syringe.....	96.50	6	Binocrit
⚡ Inj 5,000 iu in 0.5 ml syringe.....	125.00	6	Binocrit
⚡ Inj 6,000 iu in 0.6 ml syringe.....	145.00	6	Binocrit
⚡ Inj 8,000 iu in 0.8 ml syringe.....	175.00	6	Binocrit
⚡ Inj 10,000 iu in 1 ml syringe.....	197.50	6	Binocrit
⚡ Inj 40,000 iu in 1 ml syringe.....	250.00	1	Binocrit

➡ **Restricted (RS1660)**

Initiation – chronic renal failure

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initiation – myelodysplasia*

Re-assessment required after 2 months

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Continuation – myelodysplasia*

Re-assessment required after 12 months

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with epoetin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Initiation – all other indications

Haematologist

For use in patients where blood transfusion is not a viable treatment alternative.

Note: Indications marked with * are unapproved indications

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

EPOETIN BETA – Restricted see terms [below](#)

Note: Epoetin beta is considered a Discretionary Variance Pharmaceutical for epoetin alfa.

- ↓ Inj 2,000 iu in 0.3 ml syringe
- ↓ Inj 3,000 iu in 0.3 ml syringe
- ↓ Inj 4,000 iu in 0.3 ml syringe
- ↓ Inj 5,000 iu in 0.3 ml syringe
- ↓ Inj 6,000 iu in 0.3 ml syringe
- ↓ Inj 10,000 iu in 0.6 ml syringe

→ **Restricted (RS1661)**

Initiation – chronic renal failure

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initiation – myelodysplasia*

Re-assessment required after 12 months

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Continuation – myelodysplasia*

Re-assessment required after 2 months

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with epoetin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Initiation – all other indications

All of the following:

- 1 Haematologist; and
- 2 For use in patients where blood transfusion is not a viable treatment alternative; and
- 3 *Note: Indications marked with * are unapproved indications.

Megaloblastic

FOLIC ACID

Tab 0.8 mg	26.60	1,000	Folic Acid multichem
Tab 5 mg – 1% DV Mar-23 to 2027	5.82	100	Folic Acid Viatrix
Oral liq 50 mcg per ml	31.77	25 ml	Biomed
Inj 5 mg per ml, 10 ml vial			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

Antifibrinolytics, Haemostatics and Local Sclerosants

ALUMINIUM CHLORIDE – **Restricted** see terms [below](#)

↓ Topical soln 20% w/v

e.g. Driclor

→ **Restricted** (RS1500)

Initiation

For use as a haemostatis agent.

APROTININ – **Restricted** see terms [below](#)

↓ Inj 10,000 kIU per ml (equivalent to 200 mg per ml), 50 ml vial

→ **Restricted** (RS1332)

Initiation

Cardiac anaesthetist

Either:

- 1 Paediatric patient undergoing cardiopulmonary bypass procedure; or
- 2 Adult patient undergoing cardiac surgical procedure where the significant risk of massive bleeding outweighs the potential adverse effects of the drug.

ELTROMBOPAG – **Restricted** see terms [below](#)

↓ Tab 25 mg 1,550.00

28 Revolade

↓ Tab 50 mg 3,100.00

28 Revolade

→ **Restricted** (RS1648)

Initiation – idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

Re-assessment required after 6 weeks

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initiation – idiopathic thrombocytopenic purpura - preparation for splenectomy

Haematologist

Limited to 6 weeks treatment

The patient requires eltrombopag treatment as preparation for splenectomy.

Continuation – idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

Re-assessment required after 12 months

The patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre

Initiation – idiopathic thrombocytopenic purpura contraindicated to splenectomy

Haematologist

Re-assessment required after 3 months

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Continuation – idiopathic thrombocytopenic purpura contraindicated to splenectomy

Haematologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Initiation – severe aplastic anaemia

Haematologist

Re-assessment required after 3 months

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Continuation – severe aplastic anaemia

Haematologist

Re-assessment required after 12 months

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB – Restricted see terms [below](#)

⚡ Inj 30 mg in 1 ml vial.....	3,570.00	1	Hemlibra
⚡ Inj 60 mg in 0.4 ml vial.....	7,138.00	1	Hemlibra
⚡ Inj 105 mg in 0.7 ml vial.....	12,492.00	1	Hemlibra
⚡ Inj 150 mg in 1 ml vial.....	17,846.00	1	Hemlibra

→ Restricted (RS1998)

Initiation – Severe Haemophilia A with or without FVIII inhibitors

Haematologist

Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

FERRIC SUBSULFATE

Gel 25.9%

Soln 500 ml

POLIDOCANOL

Inj 0.5%, 30 ml vial

BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM TETRADECYL SULPHATE			
Inj 3%, 2 ml ampoule			
THROMBIN			
Powder			
TRANEXAMIC ACID			
Tab 500 mg – 5% DV May-26 to 2028.....	9.93	60	Mercury Pharma
Inj 100 mg per ml, 5 ml ampoule – 5% DV Mar-25 to 2027	5.39	5	Tranexamic-AFT
Inj 100 mg per ml, 10 ml ampoule – 5% DV Mar-25 to 2027	7.99	5	Tranexamic-AFT

Anticoagulant Reversal Agents

IDARUCIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 50 mg per ml, 50 ml vial.....4,250.00 2 Praxbind

➡ **Restricted (RS1535)**

Initiation

For the reversal of the anticoagulant effects of dabigatran when required in situations of life-threatening or uncontrolled bleeding, or for emergency surgery or urgent procedures.

Blood Factors

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – **Restricted** see terms [below](#)

⚡ Inj 250 iu vial.....612.50 1 Alprolix
 ⚡ Inj 500 iu vial.....1,225.00 1 Alprolix
 ⚡ Inj 1,000 iu vial.....2,450.00 1 Alprolix
 ⚡ Inj 2,000 iu vial.....4,900.00 1 Alprolix
 ⚡ Inj 3,000 iu vial.....7,350.00 1 Alprolix
 ⚡ Inj 4,000 iu vial.....9,800.00 1 Alprolix

➡ **Restricted (RS1684)**

Initiation

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] – **Restricted** see terms [below](#)

⚡ Inj 1 mg syringe.....1,178.30 1 NovoSeven RT
 ⚡ Inj 2 mg syringe.....2,356.60 1 NovoSeven RT
 ⚡ Inj 5 mg syringe.....5,891.50 1 NovoSeven RT
 ⚡ Inj 8 mg syringe.....9,426.40 1 NovoSeven RT

➡ **Restricted (RS1704)**

Initiation

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

FACTOR EIGHT INHIBITOR BYPASSING FRACTION – **Restricted** see terms [below](#)

⚡ Inj 500 U1,315.00 1 FEIBA NF
 ⚡ Inj 1,000 U2,630.00 1 FEIBA NF
 ⚡ Inj 2,500 U6,575.00 1 FEIBA NF

➡ **Restricted (RS1705)**

Initiation

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – Restricted see terms below			
⚡ Inj 250 iu prefilled syringe.....	287.50	1	Xyntha
⚡ Inj 500 iu prefilled syringe.....	575.00	1	Xyntha
⚡ Inj 1,000 iu prefilled syringe.....	1,150.00	1	Xyntha
⚡ Inj 2,000 iu prefilled syringe.....	2,300.00	1	Xyntha
⚡ Inj 3,000 iu prefilled syringe.....	3,450.00	1	Xyntha
➔ Restricted (RS1706)			
Initiation			
For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.			
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – Restricted see terms below			
⚡ Inj 1,000 iu vial.....	870.00	1	RIXUBIS
⚡ Inj 2,000 iu vial.....	1,740.00	1	RIXUBIS
⚡ Inj 3,000 iu vial.....	2,610.00	1	RIXUBIS
➔ Restricted (RS1679)			
Initiation			
For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.			
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – Restricted see terms below			
⚡ Inj 500 iu vial.....	420.00	1	Advate
⚡ Inj 1,000 iu vial.....	840.00	1	Advate
⚡ Inj 2,000 iu vial.....	1,680.00	1	Advate
⚡ Inj 3,000 iu vial.....	2,520.00	1	Advate
➔ Restricted (RS1707)			
Initiation			
For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.			
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) – Restricted see terms below			
⚡ Inj 250 iu vial.....	237.50	1	Kogenate FS
⚡ Inj 500 iu vial.....	475.00	1	Kogenate FS
⚡ Inj 1,000 iu vial.....	950.00	1	Kogenate FS
⚡ Inj 2,000 iu vial.....	1,900.00	1	Kogenate FS
⚡ Inj 3,000 iu vial.....	2,850.00	1	Kogenate FS
➔ Restricted (RS1708)			
Initiation			
For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.			
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] – Restricted see terms below			
⚡ Inj 1,000 iu vial.....	1,200.00	1	Adynovate
⚡ Inj 2,000 iu vial.....	2,400.00	1	Adynovate
➔ Restricted (RS1682)			
Initiation			
For patients with haemophilia A receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.			
Vitamin K			
PHYTOMENADIONE			
Inj 2 mg in 0.2 ml ampoule	8.00	5	Konakion MM Paediatric
Inj 10 mg per ml, 1 ml ampoule	9.21	5	Konakion MM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antithrombotics			
Anticoagulants			
BIVALIRUDIN – Restricted see terms below			
⚡ Inj 250 mg vial			
➡ Restricted (RS1181)			
Initiation			
Either:			
1 For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance; or			
2 For use in patients undergoing endovascular procedures.			
CITRATE SODIUM			
Inj 4% (200 mg per 5 ml), 5 ml ampoule			
Inj 46.7% (1.4 g per 3 ml), 3 ml syringe			
Inj 46.7% (2.36 g per 5 ml), 5 ml ampoule			
DABIGATRAN			
Cap 75 mg	27.99	60	Pradaxa
Cap 110 mg	27.99	60	Pradaxa
Cap 150 mg	27.99	60	Pradaxa
DANAPAROID – Restricted see terms below			
⚡ Inj 750 u in 0.6 ml ampoule			
➡ Restricted (RS1182)			
Initiation			
For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance.			
DEFIBROTIDE – Restricted see terms below			
⚡ Inj 80 mg per ml, 2.5 ml ampoule			
➡ Restricted (RS1183)			
Initiation			
Haematologist			
Patient has moderate or severe sinusoidal obstruction syndrome as a result of chemotherapy or regimen-related toxicities.			
DEXTROSE WITH SODIUM CITRATE AND CITRIC ACID [ACID CITRATE DEXTROSE A]			
Inj 24.5 mg with sodium citrate 22 mg and citric acid 7.3 mg per ml, 100 ml bag			
ENOXAPARIN SODIUM			
Inj 20 mg in 0.2 ml syringe – 5% DV Feb-25 to 2027	21.90	10	Clexane
Inj 40 mg in 0.4 ml ampoule			
Inj 40 mg in 0.4 ml syringe – 5% DV Feb-25 to 2027	29.74	10	Clexane
Inj 60 mg in 0.6 ml syringe – 5% DV Feb-25 to 2027	42.47	10	Clexane
Inj 80 mg in 0.8 ml syringe – 5% DV Feb-25 to 2027	56.62	10	Clexane
Inj 100 mg in 1 ml syringe – 5% DV Feb-25 to 2027	70.91	10	Clexane
Inj 120 mg in 0.8 ml syringe – 5% DV Feb-25 to 2027	88.11	10	Clexane Forte
Inj 150 mg in 1 ml syringe – 5% DV Feb-25 to 2027	100.70	10	Clexane Forte
FONDAPARINUX SODIUM – Restricted see terms below			
⚡ Inj 2.5 mg in 0.5 ml syringe			
⚡ Inj 7.5 mg in 0.6 ml syringe			
➡ Restricted (RS1184)			
Initiation			
For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
HEPARIN SODIUM			
Inj 10 iu per ml, 5 ml ampoule (flushing solution)	19.38	10	Wockhardt
Inj 100 iu per ml, 250 ml bag			
Inj 1,000 iu per ml, 1 ml ampoule	362.98	50	Hospira
Inj 1,000 iu per ml, 5 ml ampoule	164.40	50	Pfizer
Inj 5,000 iu in 0.2 ml ampoule			
Inj 5,000 iu per ml, 1 ml ampoule	70.33	5	Hospira
Inj 5,000 iu per ml, 5 ml ampoule – 5% DV May-26 to 2028	406.15	50	Pfizer
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml ampoule	96.91	50	Pfizer
Inj 100 iu per ml, 2 ml ampoule			
Inj 100 iu per ml, 5 ml ampoule			
PHENINDIONE			
Tab 10 mg			
Tab 25 mg			
Tab 50 mg			
PROTAMINE SULPHATE			
Inj 10 mg per ml, 5 ml ampoule			
RIVAROXABAN			
Tab 10 mg	15.60	30	Xarelto
Tab 15 mg	14.56	28	Xarelto
Tab 20 mg	14.56	28	Xarelto
SODIUM CITRATE WITH SODIUM CHLORIDE AND POTASSIUM CHLORIDE			
Inj 4.2 mg with sodium chloride 5.7 mg and potassium chloride 74.6 mcg per ml, 5,000 ml bag			
WARFARIN SODIUM			
Tab 1 mg	7.50	100	Marevan
Tab 2 mg			
Tab 3 mg	12.00	100	Marevan
Tab 5 mg	13.50	100	Marevan
Antiplatelets			
ASPIRIN			
Tab 100 mg	12.65	990	Ethics Aspirin EC
Suppos 300 mg			
CLOPIDOGREL			
Tab 75 mg – 5% DV Dec-25 to 2028	5.07	84	Arrow - Clopid
DIPYRIDAMOLE – Restricted: For continuation only			
➡ Tab 25 mg			
➡ Cap modified-release 200 mg	55.13	60	Dipyridamole - Strides
➡ Inj 5 mg per ml, 2 ml ampoule			
EPTIFIBATIDE – Restricted see terms below			
↓ Inj 2 mg per ml, 10 ml vial.....	180.38	1	Eptifibatide Viatrix
↓ Inj 750 mcg per ml, 100 ml vial.....	526.50	1	Eptifibatide Viatrix
➡ Restricted (RS1759)			

Initiation

Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
- 2 For use in patients with definite or strongly suspected intra-coronary thrombus on coronary angiography; or
- 3 For use in patients undergoing intra-cranial intervention.

LYSINE ACETYSALICYLATE [LYSINE ASPIRIN] – **Restricted** see terms [below](#)

⚡ Inj 500 mg

e.g. Aspegic

➡ **Restricted** (RS1689)

Initiation

Both:

- 1 For use when an immediate antiplatelet effect is required prior to an urgent interventional neuro-radiology or interventional cardiology procedure; and
- 2 Administration of oral aspirin would delay the procedure.

TICAGRELOR – **Restricted** see terms [below](#)

⚡ Tab 90 mg – **5% DV Dec-24 to 2027** 23.85
20.35

56

Ticagrelor Mylan
Ticagrelor Sandoz
Ticagrelor Sandoz S29

➡ **Restricted** (RS2142)

Initiation

Restricted to treatment of acute coronary syndromes specifically for patients who have recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome, and in whom fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initiation – thrombosis prevention neurological stenting

Re-assessment required after 12 months

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent..

Continuation – thrombosis prevention neurological stenting

Re-assessment required after 12 months

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Initiation – Percutaneous coronary intervention with stent deployment

Limited to 12 months treatment

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initiation – Stent thrombosis

Patient has experienced cardiac stent thrombosis whilst on clopidogrel.

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

Initiation – Myocardial infarction

Limited to 1 week treatment

For short term use while in hospital following ST-elevated myocardial infarction.

Initiation – acute minor stroke or high-risk transient ischemic attack (TIA)*

All of the following:

- 1 Patient has been diagnosed with a minor stroke (NIHSS† score 3 or less), high-risk TIA (ABCD2 score 4 or more) or Crescendo TIA; and
- 2 Either:
 - 2.1 Patient is expected to be a poor metaboliser of clopidogrel, with documented clinical rationale; or
 - 2.2 Patient is allergic to clopidogrel**; and
- 3 Ticagrelor to be prescribed for a maximum of 21 days following minor stroke or TIA.

Continuation – subsequent minor stroke or high-risk transient ischemic attack

Re-assessment required after 1 month

Patient has been diagnosed with a minor stroke (NIHSS score 3 or less), high-risk transient ischemic attack (ABCD2 score 4 or more) or Crescendo TIA.

Notes: Indications marked with * are unapproved indications.

Note:** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment

Note:NIHSS† National Institutes of Health Stroke Scale.

TICLOPIDINE

Tab 250 mg

Fibrinolytic Agents

ALTEPLASE

Inj 2 mg vial
Inj 10 mg vial
Inj 50 mg vial

TENECTEPLASE

Inj 50 mg vial with diluent2,390.00 1 Metalyse

UROKINASE

Inj 5,000 iu vial
Inj 10,000 iu vial
Inj 50,000 iu vial
Inj 100,000 iu vial
Inj 250,000 iu vial
Inj 500,000 iu vial

Colony-Stimulating Factors

Drugs Used to Mobilise Stem Cells

PLERIXAFOR – **Restricted** see terms [below](#)

⚡ Inj 20 mg per ml, 1.2 ml vial – 5% DV Oct-26 to 20288,740.00 1 Mozobil
869.00 **Plerixafor-AFT**

(Mozobil Inj 20 mg per ml, 1.2 ml vial to be delisted 1 October 2026)

➡ **Restricted (RS2157)**

Initiation – Stem cell transplant

Haematologist

Limited to 3 days treatment

All of the following:

continued...

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

1 Either:

- 1.1 Patient is to undergo stem cell transplantation; or
- 1.2 Patient is a donor for stem cell transplantation; and

2 Patient has not had more than one previous unsuccessful mobilisation attempt with plerixafor; and

3 Any of the following:

3.1 Both:

- 3.1.1 Patient is undergoing G-CSF mobilisation; and

3.1.2 Either:

- 3.1.2.1 Has a suboptimal peripheral blood CD34 count of less than or equal to $20 \times 10^6/L$ on day 5 after 4 days of G-CSF treatment; or
- 3.1.2.2 Efforts to collect $> 1 \times 10^6$ CD34 cells/kg have failed after one apheresis procedure; or

3.2 Both:

- 3.2.1 Patient is undergoing chemotherapy and G-CSF mobilisation; and

3.2.2 Any of the following:

3.2.2.1 Both:

- 3.2.2.1.1 Has rising white blood cell counts of $> 2 \times 10^9/L$; and
- 3.2.2.1.2 Has a suboptimal peripheral blood CD34 count of less than or equal to $20 \times 10^6/L$; or
- 3.2.2.2 Efforts to collect $> 1 \times 10^6$ CD34 cells/kg have failed after one apheresis procedure; or
- 3.2.2.3 The peripheral blood CD34 cell counts are decreasing before the target has been received; or

3.3 A previous mobilisation attempt with G-CSF or G-CSF plus chemotherapy has failed.

Granulocyte Colony-Stimulating Factors

FILGRASTIM – **Restricted** see terms [below](#)

⚡ Inj 300 mcg in 0.5 ml prefilled syringe – 5% DV Dec-24 to 2027	86.60	10	Nivestim
⚡ Inj 300 mcg in 1 ml vial	520.00	4	Neupogen
⚡ Inj 480 mcg in 0.5 ml prefilled syringe – 5% DV Dec-24 to 2027	133.72	10	Nivestim

➡ **Restricted (RS1188)**

Haematologist or oncologist

PEGFILGRASTIM – **Restricted** see terms [below](#)

⚡ Inj 6 mg per 0.6 ml syringe – 5% DV Feb-26 to 2028	69.50	1	Ziextenzo
--	-------	---	------------------

➡ **Restricted (RS1743)**

Initiation

For prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*).

Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines

Fluids and Electrolytes

Intravenous Administration

CALCIUM CHLORIDE

Inj 100 mg per ml, 10 ml vial

Inj 100 mg per ml, 50 ml syringe

e.g. Baxter

CALCIUM GLUCONATE

Inj 10%, 10 ml ampoule

e.g. Max Health

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
COMPOUND ELECTROLYTES			
Inj sodium 140 mmol/l, potassium 5 mmol/l, magnesium 1.5 mmol/l, chloride 98 mmol/l, acetate 27 mmol/l, gluconate 23 mmol/l, 500 ml bag.....	62.82	18	Plasma-Lyte 148
Inj sodium 140 mmol/l, potassium 5 mmol/l, magnesium 1.5 mmol/l, chloride 98 mmol/l, acetate 27 mmol/l, gluconate 23 mmol/l, 1,000 ml bag.....	30.72	12	Plasma-Lyte 148
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]			
Inj sodium 140 mmol/l, 5 mmol/l potassium, 1.5 mmol/l magnesium, 98 mmol/l chloride, 27 mmol/l acetate and 23 mmol/l gluconate, glucose 23 mmol/l (5%), 1,000 ml bag	239.04	12	Plasma-Lyte 148 & 5% Glucose
COMPOUND SODIUM LACTATE [HARTMANN'S SOLUTION]			
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bicarbonate 29 mmol/l, chloride 111 mmol/l, 500 ml bag	27.90	18	Baxter
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bicarbonate 29 mmol/l, chloride 111 mmol/l, 1,000 ml bag	19.32	12	Baxter
GLUCOSE [DEXTROSE]			
Inj 5%, 1,000 ml bag.....	53.10	10	Fresenius Kabi
Inj 5%, 100 ml bag.....	97.00	50	Fresenius Kabi
Inj 5%, 250 ml bag.....	63.00	30	Fresenius Kabi
Inj 5%, 50 ml bag.....	162.00	60	Baxter Glucose 5%
Inj 5%, 500 ml bag.....	67.40	20	Fresenius Kabi
Inj 10%, 1,000 ml bag.....	162.00	12	Baxter Glucose 10%
Inj 10%, 500 ml bag.....	126.00	18	Baxter Glucose 10%
Inj 50%, 10 ml ampoule	34.75	5	Biomed
Inj 50%, 500 ml bag.....	423.00	18	Baxter Glucose 50%
Inj 50%, 90 ml bottle.....	17.50	1	Biomed
GLUCOSE WITH POTASSIUM CHLORIDE			
Inj 10% glucose with 20 mmol/l potassium chloride, 500 ml bag			
GLUCOSE WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE			
Inj 2.5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 3,000 ml bag			
Inj 10% glucose with potassium chloride 10 mmol/l and sodium chloride 15 mmol/l, 500 ml bag			
Inj 4% glucose with potassium chloride 20 mmol/l and sodium chloride 0.18%, 1,000 ml bag.....	240.36	12	Baxter
Inj 5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 1,000 ml bag.....	189.00	12	Baxter
Inj 5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.9%, 1,000 ml bag.....	334.08	12	Baxter
GLUCOSE WITH SODIUM CHLORIDE			
Inj glucose 2.5% with sodium chloride 0.45%, 500 ml bag.....	318.78	18	Baxter
Inj 4% glucose and sodium chloride 0.18%, 1,000 ml bag.....	192.96	12	Baxter
Inj 5% glucose and sodium chloride 0.45%, 1,000 ml bag.....	192.84	12	Baxter
Inj 5% glucose and sodium chloride 0.9%, 1,000 ml bag.....	204.84	12	Baxter
POTASSIUM CHLORIDE			
Inj 75 mg (1 mmol) per ml, 10 ml ampoule			
Inj 225 mg (3 mmol) per ml, 20 ml ampoule			

BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
POTASSIUM CHLORIDE WITH SODIUM CHLORIDE			
Inj 10 mmol potassium chloride with 0.29% sodium chloride, 100 ml bag	563.52	48	Baxter
Inj 20 mmol potassium chloride with 0.9% sodium chloride, 1,000 ml bag....	192.72	12	Baxter
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 1,000 ml bag....	299.40	12	Baxter
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 100 ml bag	912.96	48	Baxter
POTASSIUM DIHYDROGEN PHOSPHATE			
Inj 1 mmol per ml, 10 ml ampoule	174.57	10	Hospira
RINGER'S SOLUTION			
Inj sodium 147 mmol/l with potassium 4 mmol/l, calcium 2.2 mmol/l, chloride 156 mmol/l, 1,000 ml bag.....	227.52	12	Baxter
SODIUM ACETATE			
Inj 4 mmol per ml, 20 ml ampoule			
SODIUM BICARBONATE			
Inj 8.4%, 10 ml vial			
Inj 8.4%, 50 ml vial	24.70	1	Biomed
Inj 8.4%, 100 ml vial	25.31	1	Biomed
SODIUM CHLORIDE			
Inj 0.9%, 5 ml ampoule – 5% DV Feb-26 to 2028	4.12	20	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – 5% DV Feb-26 to 2028	7.50	50	Fresenius Kabi
⚡ Inj 0.9%, 3 ml, prefilled syringe, non-sterile pack – 5% DV Jul-26 to 2028	12.60	30	Polyflush
➡ Restricted (RS1297)			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⚡ Inj 0.9%, 5 ml, prefilled syringe, non-sterile pack – 5% DV Jul-26 to 2028	12.60	30	Polyflush
➡ Restricted (RS1297)			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⚡ Inj 0.9%, 10 ml, prefilled syringe, non-sterile pack – 5% DV Jul-26 to 2028	12.60	30	Polyflush
➡ Restricted (RS1297)			
Initiation			
For use in flushing of in-situ vascular access devices only.			
Inj 0.9%, 20 ml ampoule – 5% DV Feb-26 to 2028	5.20	20	Fresenius Kabi
Inj 23.4% (4 mmol/ml), 20 ml ampoule	40.15	5	Biomed
Inj 0.45%, 500 ml bag.....	84.42	18	Baxter
Inj 3%, 1,000 ml bag.....	165.84	12	Baxter
Inj 0.9%, 50 ml bag.....	124.20	60	Baxter
	147.75	75	Baxter-Viaflo
Inj 0.9%, 100 ml bag.....	88.80	48	Baxter
	105.60	60	Baxter-Viaflo
Inj 0.9%, 250 ml bag.....	50.40	24	Baxter
Inj 0.9%, 500 ml bag.....	27.54	18	Baxter
Inj 0.9%, 1,000 ml bag.....	18.96	12	Baxter
Inj 1.8%, 500 ml bottle			
SODIUM DIHYDROGEN PHOSPHATE [SODIUM ACID PHOSPHATE]			
Inj 1 mmol per ml, 20 ml ampoule	59.10	5	Biomed

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
WATER			
Inj 10 ml ampoule – 5% DV Nov-26 to 2028	9.50	50	Multichem
Inj 20 ml ampoule	5.00	20	Fresenius Kabi
Inj 250 ml bag			
Inj 500 ml bag			
Inj, 1,000 ml bag	24.12	12	Baxter
Oral Administration			
CALCIUM POLYSTYRENE SULPHONATE			
Powder	169.85	300 g	Calcium Resonium Roma
COMPOUND ELECTROLYTES			
Powder for oral soln – 5% DV Dec-25 to 2028	9.50	50	Electral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]			
Soln with electrolytes (2 x 500 ml) – 5% DV Jul-26 to 2028	8.45	1	Pedialyte
PHOSPHORUS			
Tab eff 500 mg (16 mmol)			
POTASSIUM CHLORIDE			
Tab eff 548 mg (14 mmol) with chloride 285 mg (8 mmol)			
Tab long-acting 600 mg (8 mmol) – 5% DV Feb-26 to 2028	16.15	200	Span-K
Oral liq 2 mmol per ml			
SODIUM BICARBONATE			
Cap 840 mg	8.52	100	Sodibic
SODIUM CHLORIDE			
Tab 600 mg			
Oral liq 2 mmol/ml			
SODIUM POLYSTYRENE SULPHONATE			
Powder	84.65	454 g	Resonium A
Plasma Volume Expanders			
GELATINE, SUCCINYLATED			
Inj 4%, 500 ml bag	139.10	10	Gelofusine

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Agents Affecting the Renin-Angiotensin System			
ACE Inhibitors			
CAPTOPRIL			
↓ Oral liq 5 mg per ml	86.00	100 ml	DP-Captopril
➔ Restricted (RS1263)			
Initiation			
Any of the following:			
1 For use in children under 12 years of age; or			
2 For use in tube-fed patients; or			
3 For management of rebound transient hypertension following cardiac surgery.			
ENALAPRIL MALEATE			
Tab 5 mg – 5% DV Jul-26 to 2028	1.40	90	Ipca-Enalapril
Tab 10 mg – 5% DV Jul-26 to 2028	1.58	90	Ipca-Enalapril
Tab 20 mg – 5% DV Jul-26 to 2028	2.00	90	Ipca-Enalapril
LISINOPRIL			
Tab 5 mg – 5% DV Mar-26 to 2028	12.00	90	Teva Lisinopril
Tab 10 mg – 5% DV Mar-26 to 2028	12.00	90	Teva Lisinopril
Tab 20 mg – 5% DV Mar-26 to 2028	16.00	90	Teva Lisinopril
PERINDOPRIL			
Tab 2 mg – 5% DV Dec-24 to 2027	1.79	30	Coversyl
Tab 4 mg – 5% DV Dec-24 to 2027	2.44	30	Coversyl
Tab 8 mg – 5% DV Dec-24 to 2027	3.94	30	Coversyl
QUINAPRIL			
Tab 5 mg – 5% DV Mar-25 to 2027	10.24	90	Arrow-Quinapril 5
Tab 10 mg – 5% DV Mar-25 to 2027	12.51	90	Arrow-Quinapril 10
Tab 20 mg – 5% DV Mar-25 to 2027	14.38	90	Arrow-Quinapril 20
RAMIPRIL			
Tab 1.25 mg	5.75	30	Ramipril Viatris
Tab 2.5 mg	5.50	30	Ramipril Viatris
Tab 5 mg	5.25	28	Ramipril Mylan Generics
Tab 10 mg	5.48	28	Ramipril Mylan Generics
Cap 1.25 mg – 5% DV Feb-25 to 2027	17.25	90	Tryzan
Cap 2.5 mg – 5% DV Feb-25 to 2027	16.50	90	Tryzan
Cap 5 mg – 5% DV Feb-25 to 2027	16.88	90	Tryzan
Cap 10 mg – 5% DV Feb-25 to 2027	17.63	90	Tryzan

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Angiotensin II Antagonists			
CANDESARTAN CILEXETIL			
Tab 4 mg – 5% DV Feb-25 to 2027	2.92	98	Candesartan Mylan
	3.53	30	Candesartan Sandoz
			S29 AU
	2.68	90	Candesartan Viatrix
			Candesartan Viatrix
			Sante
			Candestar
Tab 8 mg – 5% DV Feb-25 to 2027	1.91	30	Apo-Candesartan
	2.67	90	Candesartan Viatrix
			Candestar
Tab 16 mg – 5% DV Feb-25 to 2027	2.84	30	Apo-Candesartan
	4.22	90	Candesartan Viatrix
			Candestar
Tab 32 mg – 5% DV Feb-25 to 2027	3.95	30	Apo-Candesartan
	5.24	90	Candesartan Viatrix
			Candestar
LOSARTAN POTASSIUM			
Tab 12.5 mg	2.00	84	Losartan Actavis
Tab 25 mg	2.29	84	Losartan Actavis
Tab 50 mg	2.86	84	Losartan Actavis
Tab 100 mg	4.57	84	Losartan Actavis
Angiotensin II Antagonists with Diuretics			
CANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDE			
Tab 16 mg with hydrochlorothiazide 12.5 mg.....	4.10	30	APO-Candesartan HCTZ
			16/12.5
Tab 32 mg with hydrochlorothiazide 12.5 mg.....	5.25	30	APO-Candesartan HCTZ
			32/12.5
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE			
Tab 50 mg with hydrochlorothiazide 12.5 mg – 5% DV Jul-26 to 2028.....	7.25	90	Losartan & Hydrochlorothiazide (Ipc)
Angiotensin II Antagonists with Neprilysin Inhibitors			
SACUBITRIL WITH VALSARTAN – Restricted see terms below			
↓ Tab 24.3 mg with valsartan 25.7 mg	190.00	56	Entresto 24/26
↓ Tab 48.6 mg with valsartan 51.4 mg	190.00	56	Entresto 49/51
↓ Tab 97.2 mg with valsartan 102.8 mg	190.00	56	Entresto 97/103
➔ Restricted (RS2014)			
Initiation			
All of the following:			
1 Patient has heart failure; and			
2 Any of the following:			
2.1 Patient is in NYHA/WHO functional class II; or			
2.2 Patient is in NYHA/WHO functional class III; or			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Alpha-Adrenoceptor Blockers

DOXAZOSIN

Tab 2 mg	17.35	500	Doxazosin Clinect
Tab 4 mg	20.94	500	Doxazosin Clinect

PHENOXYBENZAMINE HYDROCHLORIDE

- Cap 10 mg
- Inj 50 mg per ml, 1 ml ampoule
- Inj 50 mg per ml, 2 ml ampoule

PHENTOLAMINE MESYLATE

- Inj 5 mg per ml, 1 ml ampoule
- Inj 10 mg per ml, 1 ml ampoule

PRAZOSIN

Tab 1 mg	5.53	100	Arrotex-Prazosin S29
Tab 2 mg	7.00	100	Arrotex-Prazosin S29
Tab 5 mg	11.70	100	Arrotex-Prazosin S29
Cap 1 mg	15.40	100	Prazosin Mylan
Cap 2 mg	15.58	100	Prazosin Mylan
Cap 5 mg	23.32	100	Prazosin Mylan

TERAZOSIN – **Restricted:** For continuation only

- ➔ Tab 1 mg

Antiarrhythmics

ADENOSINE

Inj 3 mg per ml, 2 ml vial – 5% DV Dec-24 to 2027	34.50	5	Adsine
⚡ Inj 3 mg per ml, 10 ml vial – 5% DV Dec-24 to 2027	100.00	5	Adenosine Baxter

➔ **Restricted (RS2158)**

Initiation

For use in cardiac catheterisation, myocardial perfusion scans, electrophysiology and MRI.

AJMALINE – **Restricted** see terms [below](#)

- ⚡ Inj 5 mg per ml, 10 ml ampoule

➔ **Restricted (RS1001)**

Cardiologist

AMIODARONE HYDROCHLORIDE

Tab 100 mg – 5% DV Feb-26 to 2028	4.95	30	Aratac
Tab 200 mg – 5% DV Feb-26 to 2028	5.86	30	Aratac
Inj 50 mg per ml, 3 ml ampoule – 5% DV Feb-26 to 2028	17.96	10	Max Health

ATROPINE SULPHATE

Inj 600 mcg per ml, 1 ml ampoule – 5% DV Feb-25 to 2027	16.10	10	Hikma Martindale
---	-------	----	---------------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DIGOXIN			
Tab 62.5 mcg – 5% DV Feb-26 to 2028	8.58	240	Lanoxin PG
Tab 250 mcg – 5% DV Feb-26 to 2028	18.75	240	Lanoxin
Oral liq 50 mcg per ml			
Inj 250 mcg per ml, 2 ml vial			
DISOPYRAMIDE PHOSPHATE			
Cap 100 mg			
FLECAINIDE ACETATE			
Tab 50 mg	19.95	60	Flecainide BNM
Cap long-acting 100 mg	35.78	90	Flecainide Controlled Release Teva
Cap long-acting 200 mg	54.28	90	Flecainide Controlled Release Teva
Inj 10 mg per ml, 15 ml ampoule	102.79	5	Almarytm
	108.16		Tambocor
			Tambocor German

IVABRADINE – **Restricted** see terms [below](#)

↓ Tab 5 mg

→ **Restricted (RS1566)**

Initiation

Both:

- 1 Patient is indicated for computed tomography coronary angiography; and
- 2 Either:
 - 2.1 Patient has a heart rate of greater than 70 beats per minute while taking a maximally tolerated dose of beta blocker; or
 - 2.2 Patient is unable to tolerate beta blockers.

MEXILETINE HYDROCHLORIDE

Cap 150 mg	162.00	100	Teva
Cap 250 mg	202.00	100	Teva

PROPAFENONE HYDROCHLORIDE

Tab 150 mg

Antihypertensives

MIDODRINE – **Restricted** see terms [below](#)

↓ Tab 2.5 mg – 5% DV Feb-25 to 2027	36.68	100	Midodrine Medsurge
↓ Tab 5 mg – 5% DV Feb-25 to 2027	58.88	100	Midodrine Medsurge

→ **Restricted (RS1427)**

Initiation

Patient has disabling orthostatic hypotension not due to drugs.

Beta-Adrenoceptor Blockers

ATENOLOL

Tab 50 mg – 5% DV Feb-25 to 2027	11.00	500	Viatrix
Tab 100 mg – 5% DV Feb-25 to 2027	18.50	500	Atenolol Viatrix
Oral liq 5 mg per ml	49.85	300 ml	Atenolol-AFT

BISOPROLOL FUMARATE

Tab 2.5 mg	1.36	90	Ipca-Bisoprolol
Tab 5 mg	1.91	90	Ipca-Bisoprolol
Tab 10 mg	2.71	90	Ipca-Bisoprolol

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CARVEDILOL			
Tab 6.25 mg – 5% DV Aug-26 to 2029.....	1.97	60	Carvedilol Sandoz
Tab 12.5 mg – 5% DV Aug-26 to 2029.....	2.03	60	Carvedilol Sandoz
Tab 25 mg – 5% DV Aug-26 to 2029.....	2.46	60	Carvedilol Sandoz
CELIPROLOL – Restricted: For continuation only			
➔ Tab 200 mg			
ESMOLOL HYDROCHLORIDE			
Inj 10 mg per ml, 10 ml vial			
LABETALOL			
Tab 50 mg			
Tab 100 mg	49.54	100	Biocon
	14.50		Trandate
Tab 200 mg	42.07	100	Presolol
	27.00		Trandate
Inj 5 mg per ml, 20 ml ampoule			
METOPROLOL SUCCINATE			
Tab long-acting 23.75 mg.....	4.20	90	Myloc CR
Tab long-acting 47.5 mg.....	3.65	90	Myloc CR
Tab long-acting 95 mg.....	5.24	90	Myloc CR
Tab long-acting 190 mg.....	9.76	90	Myloc CR
METOPROLOL TARTRATE			
Tab 50 mg – 1% DV Mar-22 to 2027	5.66	100	IPCA-Metoprolol
Tab 100 mg – 1% DV Mar-22 to 2027	7.55	60	IPCA-Metoprolol
Tab long-acting 200 mg.....	23.40	28	Slow-Lopresor
Inj 1 mg per ml, 5 ml vial.....	26.50	5	Metoprolol IV Mylan
			Metoprolol IV Viatris
NADOLOL			
Tab 40 mg – 1% DV Mar-22 to 2027	19.19	100	Nadolol BNM
Tab 80 mg – 1% DV Mar-22 to 2027	30.39	100	Nadolol BNM
PROPRANOLOL			
Tab 10 mg – 1% DV Mar-22 to 2027	7.04	100	Drofate
Tab 40 mg – 1% DV Mar-22 to 2027	8.75	100	IPCA-Propranolol
Cap long-acting 160 mg	18.17	100	Cardinol LA
	35.15		Propranolol LA Lupin
Oral liq 4 mg per ml			<i>e.g. Hikma-Propranolol</i>
Inj 1 mg per ml, 1 ml ampoule			
SOTALOL			
Tab 80 mg – 5% DV Feb-26 to 2028	40.00	500	Mylan
Tab 160 mg – 5% DV Feb-26 to 2028	20.00	100	Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMLODIPINE			
Tab 2.5 mg	1.45	90	Vasorex
Tab 5 mg	1.21	90	Vasorex
Tab 10 mg	1.31	90	Vasorex

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FELODIPINE			
Tab long-acting 2.5 mg – 5% DV Feb-25 to 2027	2.18	30	Plendil ER
Tab long-acting 5 mg – 5% DV Feb-25 to 2027	6.57	90	Felo 5 ER
Tab long-acting 10 mg – 5% DV Feb-25 to 2027	6.95	90	Felo 10 ER
ISRADIPINE			
Tab 2.5 mg			
Cap 2.5 mg			
NICARDIPINE HYDROCHLORIDE – Restricted see terms below			
↓ Inj 2.5 mg per ml, 10 ml vial			
➔ Restricted (RS1699)			
Initiation			
Anaesthetist, intensivist, cardiologist or paediatric cardiologist			
Any of the following:			
1 Patient has hypertension requiring urgent treatment with an intravenous agent; or			
2 Patient has excessive ventricular afterload; or			
3 Patient is awaiting or undergoing cardiac surgery using cardiopulmonary bypass.			
NIFEDIPINE			
Tab long-acting 10 mg.....	19.42	56	Tensipine MR10
Tab long-acting 20 mg.....	17.72	100	Nyefax Retard
	9.92	56	Valni Retard
Tab long-acting 30 mg.....	34.10	100	Mylan (24 hr release)
	4.78	14	Mylan Italy (24 hr release)
Tab long-acting 60 mg.....	52.81	100	Mylan (24 hr release)
Cap 5 mg			
NIMODIPINE			
Tab 30 mg – 5% DV Feb-26 to 2028	344.83	100	Nimorhage
	350.00		Nimotop
Inj 0.2 mg per ml, 50 ml vial – 5% DV Feb-26 to 2028	337.50	5	Nimotop
Other Calcium Channel Blockers			
DILTIAZEM HYDROCHLORIDE			
Tab 30 mg			
Cap long-acting 120 mg – 5% DV Dec-25 to 2028	65.35	500	Diltiazem CD Clinect
Cap long-acting 180 mg – 1% DV Mar-22 to 2027	7.00	30	Cardizem CD
Cap long-acting 240 mg – 1% DV Mar-22 to 2027	9.30	30	Cardizem CD
Inj 5 mg per ml, 5 ml vial			
PERHEXILINE MALEATE			
Tab 100 mg.....	62.90	100	Pexsig
VERAPAMIL HYDROCHLORIDE			
Tab 40 mg.....	7.01	100	Isoptin
Tab 80 mg.....	11.74	100	Isoptin
Tab long-acting 120 mg.....	36.02	100	Isoptin SR
Tab long-acting 240 mg.....	15.12	30	Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule.....	25.00	5	Isoptin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Centrally-Acting Agents			
CLONIDINE			
Patch 2.5 mg, 100 mcg per day.....	11.70	4	Mylan
Patch 5 mg, 200 mcg per day.....	12.80	4	Mylan
Patch 7.5 mg, 300 mcg per day.....	17.90	4	Mylan
CLONIDINE HYDROCHLORIDE			
Tab 25 mcg – 5% DV Feb-26 to 2028	29.74	112	Clonidine Teva
Tab 150 mcg – 5% DV Feb-25 to 2027	40.41	100	Catapres
Inj 150 mcg per ml, 1 ml ampoule – 5% DV Jan-25 to 2027	14.10	5	Catapres
METHYLDOPA			
Tab 250 mg	15.10	100	Methyldopa Viatris
Diuretics			
Loop Diuretics			
BUMETANIDE			
Tab 1 mg	16.36	100	Burinex
Inj 500 mcg per ml, 4 ml vial			
FUROSEMIDE [FRUSEMIDE]			
Tab 40 mg – 5% DV Feb-25 to 2027	12.80	1,000	IPCA-Frusemide
Tab 500 mg	25.00	50	Urex Forte
Oral liq 10 mg per ml	11.20	30 ml	Lasix
Inj 10 mg per ml, 2 ml ampoule – 5% DV Aug-26 to 2028	3.97	10	Furosemide-AFT
Inj 10 mg per ml, 25 ml ampoule	60.65	6	Lasix
Osmotic Diuretics			
MANNITOL			
Inj 10%, 1,000 ml bag.....	882.84	12	Baxter
Inj 20%, 500 ml bag.....	1,296.00	18	Baxter
Potassium Sparing Combination Diuretics			
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE			
Tab 5 mg with furosemide 40 mg			
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE			
Tab 5 mg with hydrochlorothiazide 50 mg			
Potassium Sparing Diuretics			
AMILORIDE HYDROCHLORIDE			
Tab 5 mg			
Oral liq 1 mg per ml	35.40	25 ml	Biomed
EPLERENONE – Restricted see terms below			
↓ Tab 25 mg – 5% DV Dec-24 to 2027	15.84	30	Inspira
↓ Tab 50 mg – 5% DV Dec-24 to 2027	25.00	30	Inspira
➔ Restricted (RS1640)			
Initiation			
Both:			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has heart failure with ejection fraction less than 40%; and
- 2 Either:
 - 2.1 Patient is intolerant to optimal dosing of spironolactone; or
 - 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

SPIRONOLACTONE

Tab 25 mg – 5% DV Mar-26 to 2028	4.20	100	Spiractin
Tab 100 mg – 5% DV Mar-26 to 2028	11.40	100	Spiractin
Oral liq 5 mg per ml	35.70	25 ml	Biomed

Thiazide and Related Diuretics

BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]

Tab 2.5 mg	51.50	500	Arrow-Bendrofluazide
Tab 5 mg	61.00	500	Arrow-Bendrofluazide

CHLOROTHIAZIDE

Oral liq 50 mg per ml	30.67	25 ml	Biomed
-----------------------------	-------	-------	--------

CHLORTALIDONE [CHLORTHALIDONE]

Tab 25 mg – 5% DV Feb-26 to 2028	6.95	50	Hygroton
---	------	----	-----------------

INDAPAMIDE

Tab 2.5 mg	16.00	90	Dapa-Tabs
------------------	-------	----	-----------

METOLAZONE

Tab 5 mg			
----------	--	--	--

Vasopressin receptor antagonists

TOLVAPTAN – **Restricted** see terms [below](#)

↓ Tab 15 mg	873.50	28	Jinarc
↓ Tab 30 mg	873.50	28	Jinarc
↓ Tab 45 mg + 15 mg	1,747.00	56	Jinarc
↓ Tab 60 mg + 30 mg	1,747.00	56	Jinarc
↓ Tab 90 mg + 30 mg	1,747.00	56	Jinarc

→ **Restricted (RS1930)**

Initiation – autosomal dominant polycystic kidney disease

Renal physician or any relevant practitioner on the recommendation of a renal physician

Re-assessment required after 12 months

All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 mL/min/1.73 m² at treatment initiation; and
- 3 Either:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – autosomal dominant polycystic kidney disease

Renal physician or any relevant practitioner on the recommendation of a renal physician

Re-assessment required after 12 months

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents

Fibrates

BEZAFIBRATE

Tab 200 mg – 5% DV Mar-25 to 2027	22.65	90	Bezalip
Tab long-acting 400 mg – 5% DV Mar-25 to 2027	21.54	30	Bezalip Retard

HMG CoA Reductase Inhibitors (Statins)

ATORVASTATIN

Tab 10 mg – 5% DV Dec-24 to 2027	0.31	30	Lorstat
	5.16	500	Lorstat
Tab 20 mg – 5% DV Dec-24 to 2027	8.12	500	Lorstat
Tab 40 mg – 5% DV Dec-24 to 2027	13.79	500	Lorstat
Tab 80 mg – 5% DV Dec-24 to 2027	1.52	30	Lorstat
	25.39	500	Lorstat

PRAVASTATIN

Tab 10 mg			
Tab 20 mg	7.16	100	Clinect
Tab 40 mg	12.25	100	Clinect

ROSUVASTATIN – Restricted see terms [below](#)

⚡ Tab 5 mg	4.21	30	Rosuvastatin - Sandoz
	1.29		Rosuvastatin Mylan
			Rosuvastatin Viatris
⚡ Tab 10 mg	4.21	30	Rosuvastatin - Sandoz
	1.69		Rosuvastatin Mylan
			Rosuvastatin Viatris
⚡ Tab 20 mg	4.21	30	Rosuvastatin - Sandoz
	2.71		Rosuvastatin Mylan
			Rosuvastatin Viatris
⚡ Tab 40 mg	4.55	30	Rosuvastatin - Sandoz
			Rosuvastatin Mylan
			Rosuvastatin Viatris

➡ **Restricted (RS1868)**

Initiation – cardiovascular disease risk

Either:

- 1 Both:
 - 1.1 Patient is considered to be at risk of cardiovascular disease; and
 - 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
- 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initiation – familial hypercholesterolemia

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initiation – established cardiovascular disease

Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initiation – recurrent major cardiovascular events

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIMVASTATIN

Tab 10 mg – 5% DV Dec-26 to 2029	1.68	90	Simvastatin Mylan
	1.67		Simvastatin Viatris
Tab 20 mg – 5% DV Dec-26 to 2029	2.35	90	Simvastatin Viatris
Tab 40 mg – 5% DV Dec-26 to 2029	3.85	90	Simvastatin Viatris
Tab 80 mg – 5% DV Dec-26 to 2029	8.75	90	Simvastatin Viatris

Resins

COLESTYRAMINE

Powder for oral liq 4 g

COLESTIPOL HYDROCHLORIDE

Grans for oral liq 5 g

COLESTYRAMINE

Powder for oral suspension 4 g sachet	61.50	50	Colestyramine - Mylan Questran Light (Neon)
---	-------	----	--

Selective Cholesterol Absorption Inhibitors

EZETIMIBE

Tab 10 mg – 5% DV Dec-26 to 2029	1.44	30	Ezetimibe Sandoz
---	------	----	-------------------------

EZETIMIBE WITH SIMVASTATIN

Tab 10 mg with simvastatin 10 mg	11.86	30	Zimybe
Tab 10 mg with simvastatin 20 mg	12.55	30	Zimybe
Tab 10 mg with simvastatin 40 mg	11.55	30	Vytorin
			Zimybe
Tab 10 mg with simvastatin 80 mg	14.27	30	Vytorin
			Zimybe

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Other Lipid-Modifying Agents

ACIPIMOX
Cap 250 mg

Nitrates

GLYCERYL TRINITRATE

Inj 1 mg per ml, 5 ml ampoule			
Inj 1 mg per ml, 10 ml ampoule			
Inj 1 mg per ml, 50 ml vial			
Inj 5 mg per ml, 10 ml ampoule	118.00	5	Hospira
Oral pump spray, 400 mcg per dose	7.48	250 dose	Nitrolingual Pump Spray
Patch 25 mg, 5 mg per day	15.73	30	Nitroderm TTS 5
Patch 50 mg, 10 mg per day	18.62	30	Nitroderm TTS 10

ISOSORBIDE MONONITRATE

Tab 20 mg	22.49	100	Ismo 20
Tab long-acting 40 mg.....	9.80	30	Ismo 40 Retard
Tab long-acting 60 mg.....	13.50	90	Duride

Other Cardiac Agents

LEVOSIMENDAN – **Restricted** see terms [below](#)

⚠ Inj 2.5 mg per ml, 5 ml vial – 5% DV Nov-24 to 2027	509.60	1	Simdax
⚠ Inj 2.5 mg per ml, 10 ml vial			

➡ **Restricted (RS1007)**

Initiation – Heart transplant

Either:

- 1 For use as a bridge to heart transplant, in patients who have been accepted for transplant; or
- 2 For the treatment of heart failure following heart transplant.

Initiation – Heart failure

Cardiologist or intensivist

For the treatment of severe acute decompensated heart failure that is non-responsive to dobutamine.

Sympathomimetics

ADRENALINE

Inj 1 in 1,000, 1 ml ampoule	4.98	5	Aspen Adrenaline
	17.78		DBL Adrenaline
Inj 1 in 1,000, 30 ml vial			
Inj 1 in 10,000, 10 ml ampoule	49.00	10	Aspen Adrenaline
	36.18	5	DBL Adrenaline
Inj 1 in 10,000, 10 ml syringe			

DOBUTAMINE

Inj 12.5 mg per ml, 20 ml ampoule – 5% DV Dec-24 to 2027	61.13	5	Dobutamine-hameln
---	-------	---	--------------------------

DOPAMINE HYDROCHLORIDE

Inj 40 mg per ml, 5 ml ampoule – 5% DV Feb-25 to 2027	46.38	10	Dopamine Basi Max Health Ltd
--	-------	----	--

EPHEDRINE

Inj 3 mg per ml, 10 ml syringe – 5% DV Dec-26 to 2029	142.00	10	Ephedrine Juno
Inj 30 mg per ml, 1 ml ampoule – 5% DV Dec-26 to 2029	42.89	10	Max Health

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ISOPRENALINE [ISOPROTERENOL]			
Inj 200 mcg per ml, 1 ml ampoule			
Inj 200 mcg per ml, 5 ml ampoule			
METARAMINOL			
Inj 0.5 mg per ml, 10 ml syringe			
Inj 0.5 mg per ml, 20 ml syringe			
Inj 0.5 mg per ml, 5 ml syringe			
Inj 1 mg per ml, 1 ml ampoule			
Inj 1 mg per ml, 10 ml syringe			
Inj 10 mg per ml, 1 ml ampoule – 5% DV Dec-26 to 2029	50.35	10	Torbay
NORADRENALINE			
Inj 0.06 mg per ml, 100 ml bag			
Inj 0.06 mg per ml, 50 ml syringe			
Inj 0.1 mg per ml, 100 ml bag			
Inj 0.1 mg per ml, 50 ml syringe			
Inj 0.12 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 50 ml syringe			
Inj 0.16 mg per ml, 50 ml syringe			
Inj 1 mg per ml, 100 ml bag			
Inj 1 mg per ml, 4 ml ampoule – 5% DV Apr-26 to 2028	32.78	10	Noradrenaline Medsurge
PHENYLEPHRINE HYDROCHLORIDE			
Inj 10 mg per ml, 1 ml ampoule	310.42	25	Neosynephrine HCL

Vasodilators

ALPROSTADIL – Restricted see terms below			
↓ Inj 10 mcg vial			
↓ Inj 20 mcg vial			
➔ Restricted (RS1992)			
Initiation			
Both:			
1 Patient has erectile dysfunction; and			
2 Patient is to receive a penile Doppler ultrasonography.			
ALPROSTADIL HYDROCHLORIDE			
Inj 500 mcg per ml, 1 ml ampoule	2,030.33	5	Prostin VR
DIAZOXIDE			
Inj 15 mg per ml, 20 ml ampoule			
HYDRALAZINE HYDROCHLORIDE			
↓ Tab 25 mg			
➔ Restricted (RS1008)			
Initiation			
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure, in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.			
Inj 20 mg ampoule	25.90	5	Apresoline Hydrapres
Inj 20 mg per ml, 1 ml vial.....	283.00	25	Eugia

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MILRINONE			
Inj 1 mg per ml, 10 ml ampoule – 5% DV Dec-24 to 2027	68.00	10	Milrinone-Baxter
MINOXIDIL			
Tab 10 mg	78.40	100	Loniten
NICORANDIL			
Tab 10 mg – 5% DV Feb-26 to 2028	27.81	60	Max Health
Tab 20 mg – 5% DV Feb-26 to 2028	35.12	60	Max Health
PAPAVERINE HYDROCHLORIDE			
Inj 30 mg per ml, 1 ml vial			
Inj 12 mg per ml, 10 ml ampoule	257.12	5	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg			
SODIUM NITROPRUSSIDE			
Inj 50 mg vial			

Endothelin Receptor Antagonists

AMBRISENTAN – Restricted see terms [below](#)

⚡ Tab 5 mg	200.00	30	Ambrisentan Viatris
⚡ Tab 10 mg	200.00	30	Ambrisentan Viatris

➡ **Restricted (RS2159)**

Initiation – PAH monotherapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH monotherapy; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

5.2 Any of the following:

- 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
- 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
- 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH dual therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has tried bosentan (either as PAH monotherapy, or PAH dual therapy with sildenafil) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; or
 - 5.2.2 Patient has experienced intolerable side effects on bosentan; or
 - 5.2.3 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.4 Patient is presenting in NYHA/WHO functional class III or IV, and would benefit from initial dual therapy in the opinion of the treating clinician and has an absolute or relative contraindication to bosentan (eg. due to current liver disease or use of a combined oral contraceptive).

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has pulmonary arterial hypertension (PAH); and
 - 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
 - 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
 - 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Both:
 - 5.2.2.1 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

The patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**.

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults.

Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN – Restricted see terms [below](#)

⚡ Tab 62.5 mg – 5% DV Jan-25 to 2027	100.00	60	Bosentan Dr Reddy's
⚡ Tab 125 mg – 5% DV Jan-25 to 2027	100.00	60	Bosentan Dr Reddy's

➡ **Restricted (RS2160)**

Initiation – PAH monotherapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist,

continued...

	Price	Brand or
	(ex man. excl. GST)	Generic
	\$	Manufacturer
	Per	

continued...
cardiologist or rheumatologist
Limited to 6 months treatment
All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
 - 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
 - 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
 - 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
- 5.1 Bosentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil; or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist
Limited to 6 months treatment
All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as part of PAH dual therapy; and
 - 5.2 Either:
 - 5.2.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool**; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy.

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as part of PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

Patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**.

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults.

Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – **Restricted** see terms [below](#)

↓ Tab 25 mg – 5% DV Dec-24 to 2027	0.72	4	Vedafil
↓ Tab 50 mg – 5% DV Dec-24 to 2027	1.45	4	Vedafil
↓ Tab 100 mg – 5% DV Dec-24 to 2027	11.22	12	Vedafil
↓ Inj 0.8 mg per ml, 12.5 ml vial			

→ **Restricted (RS2161)**

Initiation – tablets Raynaud's Phenomenon

All of the following:

- 1 Patient has Raynaud's phenomenon; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (proper body insulation, avoidance of cold exposure, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient has persisting severe symptoms despite treatment with calcium channel blockers and nitrates (unless contraindicated or not tolerated).

Initiation – tablets Pulmonary arterial hypertension

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH is confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or at least 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initiation – tablets other conditions

Any of the following:

- 1 For use in weaning patients from inhaled nitric oxide; or
- 2 For perioperative use in cardiac surgery patients; or
- 3 For use in intensive care as an alternative to nitric oxide; or
- 4 For use in the treatment of erectile dysfunction secondary to spinal cord injury in patients being treated in a spinal unit.

Initiation – injection

Both:

- 1 For use in the treatment of pulmonary hypertension in infants or children being treated in paediatric intensive care units and neonatal intensive care units when the enteral route is not accessible; and
- 2 Any of the following:
 - 2.1 For perioperative use following cardiac surgery; or
 - 2.2 For use in persistent pulmonary hypertension of the newborn (PPHN); or
 - 2.3 For use in congenital diaphragmatic hernia.

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults.

Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Prostacyclin Analogues

EPOPROSTENOL – **Restricted** see terms [below](#)

⚡ Inj 500 mcg vial.....	36.61	1	Veletri
⚡ Inj 1.5 mg vial	73.21	1	Veletri

➡ **Restricted (RS2162)**

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Epoprostenol is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

Patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool.

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults.

Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST

Inj 50 mcg in 0.5 ml ampoule.....	380.00	5	Ilomedin
↓ Nebuliser soln 10 mcg per ml, 2 ml – 5% DV Dec-25 to 2028.....	166.53	30	Vebulis

→ **Restricted (RS2163)**

Initiation – PAH monotherapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
 - 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
 - 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
 - 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and
- 5 Both:
 - 5.1 Iloprost is to be used as PAH monotherapy; and
 - 5.2 Either:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist,

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and
- 5 All of the following:
 - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Either:
 - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
 - 5.3 Either:
 - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; or
 - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and
- 5 Both:
- 5.1 Iloprost is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

Patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool.

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults.

Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anti-Infective Preparations			
Antibacterials			
HYDROGEN PEROXIDE			
Crm 1% – 5% DV Jan-26 to 2028	4.89	15 g	Crystaderm
Soln 3% (10 vol)			
MAFENIDE ACETATE – Restricted see terms below			
↓ Crm 8.5%			
↓ Powder 5%			
➔ Restricted (RS1299)			
Initiation			
For the treatment of burns patients.			
MUPIROCIN			
Oint 2%			
SODIUM FUSIDATE [FUSIDIC ACID]			
Crm 2% – 5% DV Feb-25 to 2027	1.69	5 g	Foban
Oint 2% – 5% DV Feb-25 to 2027	1.69	5 g	Foban
SULFADIAZINE SILVER			
Crm 1%.....	10.80	50 g	Flamazine
Antifungals			
AMOROLFINE			
Nail soln 5%.....	21.87	5 ml	MycoNail
CICLOPIROX OLAMINE			
Nail soln 8%			
➔ Soln 1% – Restricted: For continuation only			
CLOTRIMAZOLE			
Crm 1% – 5% DV Jul-26 to 2028	1.15	20 g	Clomazol
➔ Soln 1% – Restricted: For continuation only			
ECONAZOLE NITRATE			
Crm 1% – 5% DV Jun-25 to 2027	8.04	20 g	Pevaryl
Foaming soln 1%			
KETOCONAZOLE			
Shampoo 2%.....	4.09	100 ml	Sebizole
METRONIDAZOLE			
Gel 0.75%			
MICONAZOLE NITRATE			
Crm 2%.....	0.90	15 g	Multichem
➔ Lotn 2% – Restricted: For continuation only			
Tinc 2%			
NYSTATIN			
Crm 100,000 u per g			
Antiparasitics			
DIMETHICONE			
Lotn 4% – 5% DV Jun-26 to 2028	4.60	200 ml	healthE Dimethicone 4% Lotion

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MALATHION [MALDISON] Lotn 0.5% Shampoo 1%			
PERMETHRIN Lotn 5%	4.28	30 ml	A-Scabies
PHENOTHRIN Shampoo 0.5%			

Antiacne Preparations

ADAPALENE Crm 0.1% Gel 0.1%			
BENZOYL PEROXIDE Soln 5%			
ISOTRETINOIN Cap 5 mg – 5% DV Dec-24 to 2027	11.26	60	Oratane
Cap 10 mg – 5% DV Dec-24 to 2027	18.75	120	Oratane
Cap 20 mg – 5% DV Dec-24 to 2027	26.73	120	Oratane
TRETINOIN Crm 0.05% – 5% DV Feb-25 to 2027	16.82	50 g	ReTrieve

Antipruritic Preparations

CALAMINE Crm, aqueous, BP – 5% DV Apr-25 to 2027	3.45	100 g	healthE Calamine Aqueous
CROTAMITON Crm 10% – 5% DV Feb-25 to 2027	3.49	20 g	Itch-Soothe

Barrier Creams and Emollients

Barrier Creams

DIMETHICONE Crm 10% pump bottle.....	4.52	460 g	healthE Dimethicone 10%
Crm 5% pump bottle – 5% DV Jun-26 to 2028.....	4.14	460 g	Hydralock
Crm 5% tube – 5% DV Jun-26 to 2028	1.52	100 g	healthE Dimethicone 5%
ZINC Crm			e.g. Zinc Cream (Orion-) ;Zinc Cream (PSM)
Oint Paste			e.g. Zinc oxide (PSM)
ZINC AND CASTOR OIL Crm.....	1.63	20 g	Orion
Oint.....	4.25	500 g	Evava
Note: DV limit applies to the pack sizes of greater than 30 g.			
Oint, BP	1.26	20 g	healthE
Note: DV limit applies to the pack sizes of 30 g or less.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ZINC WITH WOOL FAT			
Crm zinc 15.25% with wool fat 4%			<i>e.g. Sudocrem</i>
Emollients			
AQUEOUS CREAM			
Crm 100 g – 5% DV Mar-25 to 2027	1.25	100 g	Evara
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm 500 g – 5% DV Mar-25 to 2027	1.65	500 g	Evara
Note: DV limit applies to the pack sizes of greater than 100 g.			
CETOMACROGOL			
Crm BP, 100 g – 5% DV Jun-25 to 2027	0.99	100 g	Cetomacrogol Cream
Crm BP, 500 g – 5% DV Feb-25 to 2027	2.29	500 g	AFT Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL			
Crm 90% with glycerol 10% – 5% DV Dec-25 to 2028	1.92	460 g	Evara
	3.25	920 g	Evara
Note: DV limit applies to the pack sizes of greater than 100 g.			
Crm 90% with glycerol 10%,	1.65	100 g	healthE
Note: DV limit applies to the pack sizes of 100 g or less.			
EMULSIFYING OINTMENT			
Oint BP	2.30	100 g	Jaychem
Note: DV limit applies to pack sizes of less than 200 g.			
Oint BP, 500 g – 5% DV Dec-26 to 2029	2.95	500 g	Evara Emulsifying Ointment
Note: DV limit applies to pack sizes of greater than 200 g.			
GLYCEROL WITH PARAFFIN			
Crm glycerol 10% with white soft paraffin 5% and liquid paraffin 10%			<i>e.g. QV cream</i>
OIL IN WATER EMULSION			
Crm, 100 g – 5% DV Apr-25 to 2027	1.43	100 g	Fatty Emulsion Cream (Evara)
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm, 500 g – 5% DV Apr-25 to 2027	2.10	500 g	Fatty Emulsion Cream (Evara)
Note: DV limit applies to the pack sizes of greater than 100 g.			
PARAFFIN			
Oint liquid paraffin 50% with white soft paraffin 50% – 5% DV Aug-26 to 2028	1.78	100 g	healthE
Note: DV limit applies to the pack sizes of 100 g or less.			
White soft.....	0.79	10 g	healthE
Note: DV limit applies to pack sizes of 30 g or less, and to both white soft paraffin and yellow soft paraffin.			
White soft,.....	4.74	450 g	EVARA White Soft Paraffin
Note: DV limit applies to the pack sizes of 500 g or less and greater than 30 g.			
Yellow soft			
Lotn liquid paraffin 85%			<i>e.g QV Bath Oil</i>
PARAFFIN WITH WOOL FAT			
Lotn liquid paraffin 15.9% with wool fat 0.6%			<i>e.g. AlphaKeri;BK ;DP; Hydroderm Lotn</i>
Lotn liquid paraffin 91.7% with wool fat 3%			<i>e.g. Alpha Keri Bath Oil</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
UREA			
Crm 10%.....	1.37	100 g	healthE Urea Cream
WOOL FAT			
Crm			
Corticosteroids			
BETAMETHASONE DIPROPIONATE			
Crm 0.05%.....	36.00	50 g	Diprosone
Note: DV limit applies to the pack sizes of greater than 30 g.			
Oint 0.05%.....	36.00	50 g	Diprosone
Note: DV limit applies to the pack sizes of greater than 30 g.			
BETAMETHASONE VALERATE			
Crm 0.1% – 5% DV Feb-25 to 2027	5.85	50 g	Beta Cream
Oint 0.1% – 5% DV Feb-25 to 2027	7.90	50 g	Beta Ointment
Lotn 0.1% – 5% DV May-25 to 2027	30.00	50 ml	Betnovate
CLOBETASOL PROPIONATE			
Crm 0.05% – 5% DV Feb-26 to 2028	3.75	30 g	Dermol
Oint 0.05% – 5% DV Feb-26 to 2028	3.68	30 g	Dermol
CLOBETASONE BUTYRATE			
Crm 0.05%			
DIFLUCORTOLONE VALERATE – Restricted: For continuation only			
➡ Crm 0.1%			
➡ Fatty oint 0.1%			
HYDROCORTISONE			
Crm 1%, 30 g – 5% DV Jul-26 to 2028	1.75	30 g	Ethics
Note: DV limit applies to the pack sizes of less than or equal to 100 g.			
Crm 1%, 500 g – 5% DV Feb-26 to 2028	20.40	500 g	Noumed
Note: DV limit applies to the pack sizes of greater than 100 g.			
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6%.....	12.83	250 ml	DP Lotn HC
HYDROCORTISONE BUTYRATE			
Crm 0.1%.....	4.85	100 g	Locoid Lipocream
Oint 0.1%.....	10.28	100 g	Locoid
Milky emul 0.1%	12.33	100 ml	Locoid Crelo
METHYLPREDNISOLONE ACEPONATE			
Crm 0.1% – 5% DV Dec-26 to 2029	4.95	15 g	Advantan
Oint 0.1% – 5% DV Dec-26 to 2029	4.95	15 g	Advantan
MOMETASONE FUROATE			
Crm 0.1% – 5% DV Feb-25 to 2027	2.25	15 g	Elocon Alcohol Free
	3.50	50 g	Elocon Alcohol Free
Oint 0.1% – 5% DV Feb-25 to 2027	2.25	15 g	Elocon
	3.50	50 g	Elocon
Lotn 0.1% – 5% DV Feb-25 to 2027	4.99	30 ml	Elocon
TRIAMCINOLONE ACETONIDE			
Crm 0.02%.....	6.49	100 g	Aristocort
Oint 0.02%.....	6.54	100 g	Aristocort

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Corticosteroids with Anti-Infective Agents

BETAMETHASONE VALERATE WITH CLIOQUINOL – **Restricted** see terms [below](#)

↓ Crm 0.1% with clioquinol 3%

→ **Restricted (RS1125)**

Initiation

Either:

- 1 For the treatment of intertrigo; or
- 2 For continuation use.

BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID]

Crm 0.1% with sodium fusidate (fusidic acid) 2%

HYDROCORTISONE WITH MICONAZOLE

Crm 1% with miconazole nitrate 2% – **5% DV Feb-25 to 2027** 2.85 15 g **Micreme H**

HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN

Oint 1% with natamycin 1% and neomycin sulphate 0.5%..... 4.34 15 g Pimafucort

TRIAMCINOLONE ACETONIDE WITH NEOMYCIN SULPHATE, GRAMICIDIN AND NYSTATIN

Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and
gramicidin 250 mcg per g

Psoriasis and Eczema Preparations

ACITRETIN

Cap 10 mg 26.20 60 Novatrelin

Cap 25 mg 57.37 60 Novatrelin

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per g 59.95 60 g Enstilar

Gel 500 mcg with calcipotriol 50 mcg per g – **5% DV Dec-24 to 2027** 40.92 60 g **Daivobet**

Oint 500 mcg with calcipotriol 50 mcg per g – **5% DV Dec-24 to 2027** 14.31 30 g **Daivobet**

CALCIPOTRIOL

Oint 50 mcg per g 40.00 120 g Daivonex

COAL TAR WITH SALICYLIC ACID AND SULPHUR

Oint 12% with salicylic acid 2% and sulphur 4%

METHOXSALEN [8-METHOXYPsorALEN]

Tab 10 mg

Lotn 1.2%

PIMECROLIMUS – **Restricted** see terms [below](#)

↓ Crm 1% 33.00 15 g Elidel

→ **Restricted (RS1781)**

Initiation

Dermatologist, paediatrician or ophthalmologist

Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN

Soln 2.3% with trolamine laurilsulfate and fluorescein sodium 5.41 500 ml Pinetarsol

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

POTASSIUM PERMANGANATE

Tab 400 mg
Crystals

TACROLIMUS

¶ Oint 0.1%.....33.00 30 g Zematop

➡ **Restricted (RS1859)**

Initiation

Dermatologist or paediatrician

Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

BETAMETHASONE VALERATE

Scalp app 0.1% – 5% DV Feb-25 to 202712.95 100 ml **Beta Scalp**

CLOBETASOL PROPIONATE

Scalp app 0.05% – 5% DV Feb-26 to 20286.90 30 ml **Dermol**

HYDROCORTISONE BUTYRATE

Scalp lotn 0.1%.....6.57 100 ml Locoid

Wart Preparations

PODOPHYLLOTOXIN

Soln 0.5%33.60 3.5 ml Condylone

SILVER NITRATE

Sticks with applicator

Other Skin Preparations

DIPHEMANIL METILSULFATE

Powder 2%

IMIQUIMOD

Crm 5%, 250 mg sachet34.50 12 Aldara
21.72 24 Padagis
Perrigo

SUNSCREEN, PROPRIETARY

Lotn.....6.50 200 g Marine Blue Lotion SPF 50+

Antineoplastics

FLUOROURACIL SODIUM

Crm 5% – 5% DV Dec-24 to 20275.56 20 g **Efudix**

METHYL AMINOLEVULINATE HYDROCHLORIDE – **Restricted** see terms [below](#)

¶ Crm 16%

➡ **Restricted (RS1127)**

Dermatologist or plastic surgeon

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Wound Management Products

CALCIUM GLUCONATE
Gel 2.5%

e.g. Orion

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Anti-Infective Agents

ACETIC ACID

Soln 3%

Soln 5%

ACETIC ACID WITH HYDROXYQUINOLINE, GLYCEROL AND RICINOLEIC ACID

Jelly 0.94% with hydroxyquinoline sulphate 0.025%, glycerol 5% and
ricinoleic acid 0.75% with applicator

CHLORHEXIDINE GLUCONATE

Crm 1%

Lotn 1%

CLOTRIMAZOLE

Vaginal crm 1% with applicator – 5% DV Jul-26 to 2028 4.20

35 g

Clomazol

Vaginal crm 2% with applicator – 5% DV Jul-26 to 2028 4.60

20 g

Clomazol

MICONAZOLE NITRATE

Vaginal crm 2% with applicator 6.89

40 g

Micreme

NYSTATIN

Vaginal crm 100,000 u per 5 g with applicator(s) 5.70

75 g

Nilstat

Contraceptives

Antiandrogen Oral Contraceptives

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tablets..... 5.08

168

Ginet

Combined Oral Contraceptives

ETHINYLOESTRADIOL WITH DESOGESTREL

Tab 20 mcg with desogestrel 150 mcg

Tab 30 mcg with desogestrel 150 mcg

ETHINYLOESTRADIOL WITH LEVONORGESTREL

Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets – 5% DV

Apr-26 to 2028 2.00

84

Lo-Oralcon 20 ED

Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets – 5% DV

Apr-26 to 2028 2.30

84

Oralcon 30 ED

Tab 20 mcg with levonorgestrel 100 mcg

Tab 30 mcg with levonorgestrel 150 mcg

ETHINYLOESTRADIOL WITH NORETHISTERONE

Tab 35 mcg with norethisterone 1 mg

Tab 35 mcg with norethisterone 1 mg and 7 inert tab 14.09

84

Brevinor 1/28

Tab 35 mcg with norethisterone 500 mcg

NORETHISTERONE WITH MESTRANOL

Tab 1 mg with mestranol 50 mcg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Contraceptive Devices			
INTRA-UTERINE DEVICE			
IUD 29.1 mm length x 23.2 mm width	29.80	1	Choice 380 7med Nsha Silver/copper Short
IUD 33.6 mm length x 29.9 mm width	26.80	1	TCu 380 Plus Normal
IUD 35.5 mm length x 19.6 mm width	33.00	1	Cu 375 Standard
Emergency Contraception			
LEVONORGESTREL			
Tab 1.5 mg – 5% DV Jun-26 to 2028	1.31	1	Levonorgestrel -1 (Lupin)
Progestogen-Only Contraceptives			
DESOGESTREL			
Tab 75 mcg.....	24.50	84	Cerazette
LEVONORGESTREL			
Tab 30 mcg.....	22.00	112	Microlut
Intra-uterine device 52 mg.....	269.50	1	Mirena
Intra-uterine device 13.5 mg.....	215.60	1	Jaydess
Subdermal implant (2 x 75 mg rods)	106.92	2	Jadelle
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe	10.56	1	Depo-Provera
NORETHISTERONE			
Tab 350 mcg.....	12.25	84	Noriday
	13.23		Noriday 28
Obstetric Preparations			
Antiprogestogens			
MIFEPRISTONE			
Tab 200 mg			
Oxytocics			
CARBOPROST TROMETAMOL			
Inj 250 mcg per ml, 1 ml ampoule			
DINOPROSTONE			
Pessaries 10 mg			
Vaginal gel 1 mg in 3 g	65.39	1	Prostin E2
Vaginal gel 2 mg in 3 g	82.33	1	Prostin E2
ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule	160.00	5	DBL Ergometrine
OXYTOCIN			
Inj 5 iu per ml, 1 ml ampoule – 5% DV Mar-26 to 2028	5.98	5	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule – 5% DV Mar-26 to 2028	7.18	5	Oxytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE			
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028.....	41.47	5	Syntometrine

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Tocolytics

PROGESTERONE

Cap 100 mg 14.85 30 Utrogestan

TERBUTALINE – **Restricted** see terms [below](#)

⚡ Inj 500 mcg ampoule

➡ **Restricted** ([RS1130](#))

Obstetrician

Oestrogens

OESTRIOL

Crn 1 mg per g with applicator 6.95 15 g Ovestin

Pessaries 500 mcg 7.55 15 Ovestin

Urologicals

5-Alpha Reductase Inhibitors

FINASTERIDE – **Restricted** see terms [below](#)

⚡ Tab 5 mg 4.79 100 Ricit

➡ **Restricted** ([RS1131](#))

Initiation

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoceptor Blockers

TAMSULOSIN HYDROCHLORIDE – **Restricted** see terms [below](#)

⚡ Cap 400 mcg – **5% DV Feb-26 to 2028** 28.56 100 Tamsulosin-Rex

➡ **Restricted** ([RS1132](#))

Initiation

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Urinary Alkalisers

POTASSIUM CITRATE – **Restricted** see terms [below](#)

⚡ Oral liq 3 mmol per ml 37.49 200 ml Biomed

➡ **Restricted** ([RS1133](#))

Initiation

Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

SODIUM CITRO-TARTRATE

Grans eff 4 g sachets – **5% DV Dec-26 to 2029** 3.33 28 Ural

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Urinary Antispasmodics			
OXYBUTYNIN			
Tab 5 mg – 5% DV Dec-26 to 2029	5.42	100	Alchemy Oxybutynin
Oral liq 5 mg per 5 ml			
SOLIFENACIN SUCCINATE			
Tab 5 mg – 5% DV Jun-25 to 2027	1.95	30	Solifenacin succinate Max Health
Tab 10 mg – 5% DV Jun-25 to 2027	3.53	30	Solifenacin succinate Max Health

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Anabolic Agents

OXANDROLONE

↓ Tab 2.5 mg

→ **Restricted (RS1302)**

Initiation

For the treatment of burns patients.

Androgen Agonists and Antagonists

CYPROTERONE ACETATE

Tab 50 mg – **5% DV Jul-25 to 2027** 17.05 50 **Siterone**

Tab 100 mg – **5% DV Jul-25 to 2027** 31.00 50 **Siterone**

TESTOSTERONE

Gel (transdermal) 16.2 mg per g, 88 g – **5% DV Apr-25 to 2027** 52.00 60 **Testogel**

TESTOSTERONE CIPIONATE

Inj 100 mg per ml, 10 ml vial..... 97.75 1 Depo-Testosterone

TESTOSTERONE ESTERS

Inj testosterone decanoate 100 mg, testosterone isocarproate 60 mg,
testosterone phenylpropionate 60 mg and testosterone propionate
30 mg per ml, 1 ml ampoule

TESTOSTERONE UNDECANOATE

→ Cap 40 mg – **Restricted:** For continuation only

Inj 250 mg per ml, 4 ml vial – **5% DV Dec-26 to 2028** 27.50 1 **Gonadron**
86.00 Reandron 1000

(Reandron 1000 Inj 250 mg per ml, 4 ml vial to be delisted 1 December 2026)

Calcium Homeostasis

CALCITONIN

Inj 100 iu per ml, 1 ml ampoule 121.00 5 Miacalcic

CINACALCET – **Restricted** see terms [below](#)

↓ Tab 30 mg – **5% DV Dec-24 to 2027** 25.24 28 **Cinacalel Devatis**

↓ Tab 60 mg – **5% DV Dec-24 to 2027** 50.47 28 **Cinacalel Devatis**

→ **Restricted (RS1931)**

Initiation – parathyroid carcinoma or calciphylaxis

Nephrologist or endocrinologist

Re-assessment required after 6 months

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 3 mmol/L); and
- 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Continuation – parathyroid carcinoma or calciphylaxis

Nephrologist or endocrinologist

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initiation – primary hyperparathyroidism

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initiation – secondary or tertiary hyperparathyroidism

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or
 - 3.2 Parathyroid tissue is surgically inaccessible; or
 - 3.3 Parathyroid surgery is not feasible.

Continuation – secondary or tertiary hyperparathyroidism

Re-assessment required after 12 months

Either:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial – 5% DV Dec-24 to 2027	15.65	1	Zoledronic Acid Injection Mylan Zoledronic acid Viatrix
---	-------	---	--

(Zoledronic Acid Injection Mylan Inj 4 mg per 5 ml, vial to be delisted 1 November 2026)

Corticosteroids

BETAMETHASONE

- Tab 500 mcg
- Inj 4 mg per ml, 1 ml ampoule

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

- Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml ampoule

HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DEXAMETHASONE			
Tab 0.5 mg – 5% DV Feb-25 to 2027	1.80	30	Dexamethasone
Tab 4 mg – 5% DV Feb-25 to 2027	3.18	30	Dexamethasone
Oral liq 1 mg per ml	53.86	25 ml	Biomed
DEXAMETHASONE PHOSPHATE			
Inj 4 mg per ml, 1 ml ampoule – 5% DV Mar-26 to 2028	6.88	10	Dexamethasone Medsurge
Inj 4 mg per ml, 2 ml ampoule – 5% DV Mar-26 to 2028	10.98	10	Dexamethasone Medsurge
FLUDROCORTISONE ACETATE			
Tab 100 mcg – 5% DV Dec-25 to 2028	8.05	100	Florinef
HYDROCORTISONE			
Tab 5 mg	8.10	100	Douglas
Tab 20 mg	20.32	100	Douglas
Inj 100 mg vial – 5% DV Dec-24 to 2027	3.96	1	Solu-Cortef
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Tab 4 mg	112.00	100	Medrol
Tab 100 mg	223.10	20	Medrol
Inj 40 mg vial	22.30	1	Solu-Medrol Act-O-Vial
Inj 125 mg vial	34.10	1	Solu-Medrol Act-O-Vial
Inj 500 mg vial	43.01	1	Solu-Medrol Act-O-Vial
Inj 1 g vial	52.54	1	Solu-Medrol
METHYLPREDNISOLONE ACETATE			
Inj 40 mg per ml, 1 ml vial	47.06	5	Depo-Medrol
PREDNISOLONE			
Oral liq 5 mg per ml – 5% DV Dec-24 to 2027	6.00	30 ml	Redipred
Enema 200 mcg per ml, 100 ml			
PREDNISONE			
Tab 1 mg	18.58	500	Prednisone Clinect
Tab 2.5 mg	21.04	500	Prednisone Clinect
Tab 5 mg	19.30	500	Prednisone Clinect
Tab 20 mg	50.51	500	Prednisone Clinect
TRIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule – 5% DV Dec-26 to 2028	21.42	5	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule – 5% DV Dec-26 to 2028	52.63	5	Kenacort-A 40
TRIAMCINOLONE HEXACETONIDE			
Inj 20 mg per ml, 1 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Hormone Replacement Therapy			
Oestrogens			
OESTRADIOL			
Tab 1 mg			
Gel (transdermal) 0.06% (750 mcg/actuation) – 5% DV Nov-24 to 31 Oct 2027	14.25	80 g	EstroGel
Patch 25 mcg per day – 5% DV Dec-25 to 2027	8.89	8	Estradiol TDP Mylan
	16.23		Estradot
Patch 50 mcg per day – 5% DV Dec-25 to 2027	9.26	8	Estradiol TDP Mylan
	15.79		Estradot
Patch 75 mcg per day – 5% DV Dec-25 to 2027	10.33	8	Estradiol TDP Mylan
	16.53		Estradot
Patch 100 mcg per day – 5% DV Dec-25 to 2027	10.59	8	Estradiol TDP Mylan
	16.18		Estradot
OESTRADIOL VALERATE			
Tab 1 mg – 5% DV Dec-25 to 2028	12.36	84	Progynova
Tab 2 mg – 5% DV Dec-25 to 2028	8.24	56	Progynova
	12.36	84	Progynova
OESTROGENS (CONJUGATED EQUINE)			
Tab 300 mcg			
Tab 625 mcg			

Progestogen and Oestrogen Combined Preparations

OESTRADIOL WITH NORETHISTERONE ACETATE			
Tab 1 mg with 0.5 mg norethisterone acetate			
Tab 2 mg with 1 mg norethisterone acetate			
Tab 2 mg with 1 mg norethisterone acetate (10), and tab 2 mg oestradiol (12) and tab 1 mg oestradiol (6)			
OESTROGENS WITH MEDROXYPROGESTERONE ACETATE			
Tab 625 mcg conjugated equine with 2.5 mg medroxyprogesterone acetate			
Tab 625 mcg conjugated equine with 5 mg medroxyprogesterone acetate			

Progestogens

MEDROXYPROGESTERONE ACETATE			
Tab 2.5 mg	7.54	30	Provera
Tab 5 mg	23.15	100	Provera
Tab 10 mg	11.10	30	Provera

Other Endocrine Agents

CABERGOLINE – Restricted see terms on the next page			
↓ Tab 0.5 mg	4.43	2	Dostinex
	17.94	8	Dostinex

HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ **Restricted** (RS1855)

Initiation

Any of the following:

- 1 Inhibition of lactation; or
- 2 Patient has hyperprolactinemia; or
- 3 Patient has acromegaly.

Note: Indication marked with * is an unapproved indication.

CLOMIFENE CITRATE

Tab 50 mg	29.84	10	Mylan Clomiphen
-----------------	-------	----	-----------------

GESTRINONE

Cap 2.5 mg

METYRAPONE

Cap 250 mg

PENTAGASTRIN

Inj 250 mcg per ml, 2 ml ampoule

Other Oestrogen Preparations

OESTRADIOL

Implant 50 mg

OESTRIOL

Tab 2 mg	7.70	30	Ovestin
----------------	------	----	---------

Other Progestogen Preparations

MEDROXYPROGESTERONE

Tab 100 mg	153.60	100	Provera HD
------------------	--------	-----	------------

NORETHISTERONE

Tab 5 mg	5.49	30	Primolut N
----------------	------	----	------------

Pituitary and Hypothalamic Hormones and Analogues

CORTICORELIN (OVINE)

Inj 100 mcg vial

THYROTROPIN ALFA

Inj 900 mcg vial

Adrenocorticotrophic Hormones

TETRACOSACTIDE [TETRACOSACTRIN]

Inj 250 mcg per ml, 1 ml ampoule	86.25	1	Synacthen
Inj 1 mg per ml, 1 ml ampoule	690.00	1	UK Synacthen Synacthen Depot

GnRH Agonists and Antagonists

BUSERELIN

Inj 1 mg per ml, 5.5 ml vial

GONADORELIN

Inj 100 mcg vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GOSERELIN			
Implant 3.6 mg, syringe	66.48	1	Zoladex
Implant 10.8 mg, syringe	138.23	1	Zoladex
LEUPRORELIN ACETATE			
Inj 3.75 mg prefilled dual chamber syringe.....	221.60	1	Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe.....	591.68	1	Lucrin Depot 3-month

Gonadotrophins

CHORIOGONADOTROPIN ALFA

Inj 250 mcg in 0.5 ml syringe

Growth Hormone

SOMATROPIN – Restricted see terms [below](#)

↓ Inj 5 mg cartridge – 5% DV Feb-25 to 2027	80.21	1	Omnitrope Omnitrope AU
↓ Inj 10 mg cartridge – 5% DV Feb-25 to 2027	80.21	1	Omnitrope
↓ Inj 15 mg cartridge – 5% DV Feb-25 to 2027	139.50	1	Omnitrope

→ **Restricted (RS1826)**

Initiation – growth hormone deficiency in children

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

Either:

- Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- All of the following:
 - Height velocity < 25th percentile for age; and adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - Appropriate imaging of the pituitary gland has been obtained.

Continuation – growth hormone deficiency in children

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- No malignancy has developed since starting growth hormone.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Continuation – Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initiation – short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Continuation – short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 Current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initiation – short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of a endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73 m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73 m²) in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months.

Continuation – short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of an endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initiation – Prader-Willi syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Continuation – Prader-Willi syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initiation – adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until it is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Continuation – adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in the Quality of Life Assessment defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have increased to within ± 1 SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin does not exceed 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ± 1 SD of the mean of the normal range for age and sex (other than for obvious external factors); and

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or

3 All of the following:

- 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
- 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3.3 The patient has severe growth hormone deficiency (see notes); and
- 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Thyroid and Antithyroid Preparations

CARBIMAZOLE		
Tab 5 mg – 5% DV Dec-25 to 2028	7.56	100 Neo-Mercazole
IODINE		
Soln BP 50 mg per ml		
LEVOTHYROXINE		
Tab 25 mcg		
Tab 50 mcg		
Tab 100 mcg		
LIOTHYRONINE SODIUM		
↓ Tab 20 mcg		
➔ Restricted (RS1301)		
Initiation		
For a maximum of 14 days' treatment in patients with thyroid cancer who are due to receive radioiodine therapy.		
Inj 10 mcg vial		
Inj 20 mcg vial		
Inj 100 mcg vial		
POTASSIUM IODATE		
Tab 170 mg		
POTASSIUM PERCHLORATE		
Cap 200 mg		
PROPYLTHIOURACIL – Restricted see terms on the next page		
↓ Tab 50 mg	35.00	100 PTU

HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS1276)

Initiation

Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

PROTIRELIN

Inj 100 mcg per ml, 2 ml ampoule

Vasopressin Agents

ARGIPRESSIN [VASOPRESSIN]

Inj 20 u per ml, 1 ml ampoule

DESMOPRESSIN

Wafer 120 mcg 47.00 30 Minirin Melt

DESMOPRESSIN ACETATE

Tab 100 mcg 25.00 30 Minirin

Tab 200 mcg 54.45 30 Minirin

Inj 4 mcg per ml, 1 ml ampoule

Inj 15 mcg per ml, 1 ml ampoule

Nasal drops 100 mcg per ml

Nasal spray 10 mcg per dose, 6 ml 34.95 60 Desmopressin-PH&T

TERLIPRESSIN

Inj 0.2 mg per ml, 5 ml vial – 5% DV Feb-25 to 2027 110.00 5 Terlipressin Ever Pharma

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Antibacterials

Aminoglycosides

AMIKACIN – **Restricted** see terms [below](#)

↓ Inj 5 mg per ml, 10 ml syringe

↓ Inj 5 mg per ml, 5 ml syringe

↓ Inj 15 mg per ml, 5 ml syringe

↓ Inj 250 mg per ml, 2 ml vial – **5% DV Dec-24 to 2027** 169.97

5

DBL Amikacin
Medsurge

→ **Restricted (RS1041)**

Clinical microbiologist, infectious disease specialist or respiratory specialist

GENTAMICIN SULPHATE

Inj 10 mg per ml, 1 ml ampoule 102.60

5

DBL Gentamicin

Inj 10 mg per ml, 2 ml ampoule

Inj 40 mg per ml, 2 ml ampoule 18.38

10

Pfizer

PAROMOMYCIN – **Restricted** see terms [below](#)

↓ Cap 250 mg 126.00

16

Humatin

→ **Restricted (RS1603)**

Clinical microbiologist, infectious disease specialist or gastroenterologist

STREPTOMYCIN SULPHATE – **Restricted** see terms [below](#)

↓ Inj 400 mg per ml, 2.5 ml ampoule

→ **Restricted (RS1043)**

Clinical microbiologist, infectious disease specialist or respiratory specialist

TOBRAMYCIN

↓ Powder

→ **Restricted (RS1475)**

Initiation

For addition to orthopaedic bone cement.

↓ Inj 40 mg per ml, 2 ml vial – **5% DV Dec-24 to 2027** 15.50

5

Tobramycin (Viatris)

→ **Restricted (RS1044)**

Clinical microbiologist, infectious disease specialist or respiratory specialist

↓ Inj 100 mg per ml, 5 ml vial

→ **Restricted (RS1044)**

Clinical microbiologist, infectious disease specialist or respiratory specialist

↓ Solution for inhalation 60 mg per ml, 5 ml 395.00

56 dose

Tobramycin BNM

→ **Restricted (RS1435)**

Initiation

Patient has cystic fibrosis.

Carbapenems

ERTAPENEM – **Restricted** see terms [below](#)

↓ Inj 1 g vial 70.00

1

Invanz

→ **Restricted (RS1045)**

Clinical microbiologist or infectious disease specialist

IMIPENEM WITH CILASTATIN – **Restricted** see terms [on the next page](#)

↓ Inj 500 mg with 500 mg cilastatin vial 60.00

1

Imipenem+Cilastatin
RBX

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS1046)

Clinical microbiologist or infectious disease specialist

MEROPENEM – **Restricted** see terms [below](#)

‡ Inj 500 mg vial – 5% DV Dec-26 to 2029	28.87	10	Meropenem-AFT
‡ Inj 1 g vial – 5% DV Dec-26 to 2029	38.70	10	Meropenem-AFT

➔ Restricted (RS1047)

Clinical microbiologist or infectious disease specialist

Cephalosporins and Cephamycins - 1st Generation
CEFALEXIN

Cap 250 mg – 5% DV Jul-26 to 2028	3.90	20	Cefalexin Lupin
Cap 500 mg – 5% DV Jul-26 to 2028	3.33	20	Cefalexin Sandoz
Grans for oral liq 25 mg per ml – 5% DV Dec-26 to 2028	5.41	100 ml	Cefalexin Sandoz Flynn
	7.88		
Grans for oral liq 50 mg per ml – 5% DV Dec-26 to 2028	7.54	100 ml	Cefalexin Sandoz Flynn
	10.38		

(Flynn Grans for oral liq 25 mg per ml to be delisted 1 December 2026)

(Flynn Grans for oral liq 50 mg per ml to be delisted 1 December 2026)

CEFAZOLIN

Inj 500 mg vial	3.39	5	Cefazolin-AFT
Inj 1 g vial	3.59	5	Cefazolin-AFT
Inj 2 g vial	7.09	5	Cefazolin-AFT

Cephalosporins and Cephamycins - 2nd Generation
CEFACLOR

Cap 250 mg – 5% DV Feb-26 to 2028	29.73	100	Ranbaxy-Cefaclor
Grans for oral liq 25 mg per ml – 5% DV Feb-26 to 2028	5.83	100 ml	Ranbaxy-Cefaclor

CEFOXITIN

Inj 1 g vial

CEFUROXIME

Tab 250 mg			
Inj 750 mg vial	8.16	10	Cefuroxime Devatis
Inj 1.5 g vial	13.01	10	Cefuroxime Devatis

Cephalosporins and Cephamycins - 3rd Generation
CEFOTAXIME

Inj 500 mg vial	1.90	1	Cefotaxime Sandoz
Inj 1 g vial	38.98	10	DBL Cefotaxime

CEFTAZIDIME – **Restricted** see terms [below](#)

‡ Inj 1 g vial	25.80	10	Ceftazidime Kabi
----------------------	-------	----	------------------

➔ Restricted (RS1048)

Clinical microbiologist, infectious disease specialist or respiratory specialist

CEFTAZIDIME WITH AVIBACTAM – **Restricted** see terms [below](#)

‡ Inj ceftazidime 2,000 mg with avibactam 500 mg, vial	2,250.00	10	Zavicefta
--	----------	----	-----------

➔ Restricted (RS2104)

Initiation

Both:

continued...

	Price (ex man. excl. GST)		Brand or Generic Manufacturer
	\$	Per	

continued...

- 1 Prescribed by, or recommended by a clinical microbiologist or infectious disease specialist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital; and
- 2 Either:
 - 2.1 Proven infection with a carbapenem-resistant micro-organism, based on microbiology report; or
 - 2.2 Probable infection with a carbapenem-resistant micro-organism, based on assessment by a clinical microbiologist or infectious disease specialist..

CEFTRIAXONE

Inj 500 mg vial – 5% DV Feb-26 to 2028	0.94	1	Ceftriaxone-AFT
Inj 1 g vial – 5% DV Feb-26 to 2028	3.49	5	Ceftriaxone-AFT
Inj 2 g vial – 5% DV Feb-26 to 2028	8.15	5	Ceftriaxone-AFT

Cephalosporins and Cephamycins - 4th Generation

CEFEPIME – **Restricted** see terms [below](#)

↓ Inj 1 g vial – 5% DV Dec-24 to 2027	3.19	1	Cefepime-AFT
↓ Inj 2 g vial – 5% DV Dec-24 to 2027	4.99	1	Cefepime-AFT

→ **Restricted (RS1049)**

Clinical microbiologist or infectious disease specialist

Cephalosporins and Cephamycins - 5th Generation

CEFTAROLINE FOSAMIL – **Restricted** see terms [below](#)

↓ Inj 600 mg vial	1,834.25	10	Zinforo
-------------------------	----------	----	---------

→ **Restricted (RS1446)**

Initiation – multi-resistant organism salvage therapy

Clinical microbiologist or infectious disease specialist

Either:

- 1 for patients where alternative therapies have failed; or
- 2 for patients who have a contraindication or hypersensitivity to standard current therapies.

Macrolides

AZITHROMYCIN – **Restricted** see terms [below](#)

↓ Tab 250 mg			
↓ Tab 500 mg – 5% DV Jan-26 to 2027	2.80	2	Zithromax
↓ Grans for oral liq 200 mg per 5 ml (40 mg per ml).....	16.97	15 ml	Zithromax

→ **Restricted (RS1598)**

Initiation – bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections

Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with *Pseudomonas aeruginosa* or *Pseudomonas* related gram negative organisms*; or
- 4 Patient has an atypical *Mycobacterium* infection.

Note: Indications marked with * are unapproved indications

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – non-cystic fibrosis bronchiectasis*

Respiratory specialist or paediatrician

Re-assessment required after 12 months

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis will be subsidised in the community.

Continuation – non-cystic fibrosis bronchiectasis*

Respiratory specialist or paediatrician

Re-assessment required after 12 months

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

Note: Indications marked with * are unapproved indications. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis will be subsidised in the community.

Initiation – other indications

Re-assessment required after 5 days

For any other condition.

Continuation – other indications

Re-assessment required after 5 days

For any other condition.

CLARITHROMYCIN – **Restricted** see terms [below](#)

⚡ Tab 250 mg – 1% DV Feb-22 to 2027	8.53	14	Klacid
	7.31	12	Klaricid
⚡ Tab 500 mg – 1% DV Feb-22 to 2027	14.58	14	Klacid
⚡ Grans for oral liq 50 mg per ml	192.00	50 ml	Klacid
⚡ Inj 500 mg vial	9.10	1	Klacid IV

➡ **Restricted (RS1709)**

Initiation – Tab 250 mg and oral liquid

Any of the following:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents; or
- 3 Helicobacter pylori eradication; or
- 4 Prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated.

Initiation – Tab 500 mg

Helicobacter pylori eradication.

Initiation – Infusion

Any of the following:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents; or
- 3 Community-acquired pneumonia.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ERYTHROMYCIN (AS ETHYLSUCCINATE)			
Tab 400 mg	35.82	100	E-Mycin
Grans for oral liq 200 mg per 5 ml	6.53	100 ml	E-Mycin
Grans for oral liq 400 mg per 5 ml	9.41	100 ml	E-Mycin
ERYTHROMYCIN (AS LACTOBIONATE)			
Inj 500 mg vial	59.95	1	Eritrotex
Inj 1 g vial – 5% DV Dec-25 to 2028	125.95	1	Amdipharm
	10.00		Erythrocin IV
ERYTHROMYCIN (AS STEARATE) – Restricted: For continuation only			
➔ Tab 250 mg			
➔ Tab 500 mg			
ROXITHROMYCIN – Some items restricted see terms below			
↓ Tab dispersible 50 mg			
Tab 150 mg	13.19	50	Arrow-Roxithromycin
Tab 300 mg	25.00	50	Arrow-Roxithromycin
➔ Restricted (RS1569)			
Initiation			
Only for use in patients under 12 years of age.			
Penicillins			
AMOXICILLIN			
Cap 250 mg – 5% DV Feb-26 to 2028	54.00	500	Miro-Amoxicillin
Cap 500 mg – 5% DV Nov-26 to 2028	1.14	20	Amoxycillin Sandoz
	41.00	500	Miro-Amoxicillin
Grans for oral liq 125 mg per 5 ml	2.22	100 ml	Alphamox 125
Grans for oral liq 250 mg per 5 ml	2.81	100 ml	Alphamox 250
Inj 250 mg vial	15.97	10	Ibiamox
Inj 500 mg vial	17.43	10	Ibiamox
Inj 1 g vial	21.64	10	Ibiamox
<i>(Miro-Amoxicillin Cap 500 mg to be delisted 1 November 2026)</i>			
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg	1.59	10	Curam Duo 500/125
Grans for oral liq 25 mg with clavulanic acid 6.25 mg per ml – 5% DV			
May-25 to 2027	8.50	100 ml	Augmentin
Grans for oral liq 50 mg with clavulanic acid 12.5 mg per ml – 5% DV			
Jun-25 to 2027	5.61	100 ml	Amoxiclav Devatis
Inj 500 mg with clavulanic acid 100 mg vial – 5% DV Sep-25 to 2027	22.48	10	Forte Synermox
Inj 1,000 mg with clavulanic acid 200 mg vial – 5% DV Sep-25 to 2027	26.90	10	Cerobact
	29.61		Synermox
BENZATHINE BENZYL PENICILLIN			
Inj 900 mg (1.2 million units) vial	43.24	1	Benzetacil
Inj 900 mg (1.2 million units) in 2.3 ml syringe	432.37	10	Bicillin LA
BENZYL PENICILLIN SODIUM [PENICILLIN G]			
Inj 600 mg (1 million units) vial	16.50	10	Sandoz

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUCLOXACILLIN			
Cap 250 mg – 5% DV Aug-25 to 2027	22.58	250	Staphlex
Cap 500 mg – 5% DV Aug-25 to 2027	72.71	500	Staphlex
Grans for oral liq 25 mg per ml – 5% DV Feb-25 to 2027	4.89	100 ml	AFT
Grans for oral liq 50 mg per ml – 5% DV Feb-25 to 2027	5.89	100 ml	AFT
Inj 250 mg vial	42.60	10	Flucloxin
Inj 500 mg vial	45.63	10	Flucloxin
Inj 1 g vial	6.00	5	Flucil
PHENOXYMETHYLPENICILLIN [PENICILLIN V]			
Cap 250 mg – 5% DV Feb-25 to 2027	7.68	50	Cilicaine VK
Cap 500 mg – 5% DV Feb-25 to 2027	13.72	50	Cilicaine VK
Grans for oral liq 125 mg per 5 ml – 5% DV Feb-26 to 2028	5.75	100 ml	AFT
Grans for oral liq 250 mg per 5 ml – 5% DV Feb-26 to 2028	5.89	100 ml	AFT
PIPERACILLIN WITH TAZOBACTAM – Restricted see terms below			
⚡ Inj 4 g with tazobactam 0.5 g vial – 5% DV Dec-25 to 2028	3.15	1	PipTaz-AFT
➡ Restricted (RS1053)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PROCAINE PENICILLIN			
Inj 1.5 g in 3.4 ml syringe			
TICARCILLIN WITH CLAVULANIC ACID – Restricted see terms below			
⚡ Inj 3 g with clavulanic acid 0.1 mg vial			
➡ Restricted (RS1054)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
Quinolones			
CIPROFLOXACIN – Restricted see terms below			
⚡ Tab 250 mg – 5% DV Dec-26 to 2029	1.95	28	Ipca-Ciprofloxacin
⚡ Tab 500 mg – 5% DV Dec-26 to 2029	2.70	28	Ipca-Ciprofloxacin
⚡ Tab 750 mg – 5% DV Dec-26 to 2029	4.80	28	Ipca-Ciprofloxacin
⚡ Oral liq 50 mg per ml			
⚡ Oral liq 100 mg per ml			
⚡ Inj 2 mg per ml, 100 ml bag			
⚡ Inj 2 mg per ml, 100 ml bottle	166.50	10	Ciprofloxacin Kabi
➡ Restricted (RS1055)			
Clinical microbiologist or infectious disease specialist			
MOXIFLOXACIN – Restricted see terms below			
⚡ Tab 400 mg – 5% DV Dec-26 to 2029	42.00	5	Avelox
	25.98		Moximed
⚡ Inj 1.6 mg per ml, 250 ml bottle – 5% DV Dec-26 to 2029	372.10	10	Moxifloxacin Kabi
<i>(Avelox Tab 400 mg to be delisted 1 December 2026)</i>			
➡ Restricted (RS2129)			
Initiation – Mycobacterium infection			
Infectious disease specialist, clinical microbiologist or respiratory specialist			
Any of the following:			
1 Both:			
1.1 Active tuberculosis; and			
1.2 Any of the following:			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2.1 Documented resistance to one or more first-line medications; or
- 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
- 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
- 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
- 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Initiation – Pneumonia

Infectious disease specialist or clinical microbiologist

Either:

- 1 Immunocompromised patient with pneumonia that is unresponsive to first-line treatment; or
- 2 Pneumococcal pneumonia or other invasive pneumococcal disease highly resistant to other antibiotics.

Initiation – Penetrating eye injury

Ophthalmologist

Five days treatment for patients requiring prophylaxis following a penetrating eye injury.

Initiation – Mycoplasma genitalium

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initiation – severe delayed beta-lactam allergy

Infectious disease specialist or clinical microbiologist

Individual has a history of severe delayed beta-lactam allergy.

NORFLOXACIN

Tab 400 mg	245.00	100	Arrow-Norfloxacin
------------------	--------	-----	-------------------

Tetracyclines

DEMECLOCYCLINE HYDROCHLORIDE

Tab 150 mg
Cap 150 mg
Cap 300 mg

DOXYCYCLINE

➔ Tab 50 mg – **Restricted:** For continuation only

Tab 100 mg	64.43	500	Doxine
Inj 5 mg per ml, 20 ml vial			

MINOCYCLINE

Tab 50 mg

➔ Cap 100 mg – **Restricted:** For continuation only

TETRACYCLINE

Tab 250 mg	68.44	28	Accord
Cap 250 mg	35.90	100	Tetracycline AA
Cap 500 mg			

TIGECYCLINE – **Restricted** see terms [on the next page](#)

⚡ Inj 50 mg vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted (RS1059)			
Clinical microbiologist or infectious disease specialist			
Other Antibacterials			
AZTREONAM – Restricted see terms below			
⚡ Inj 1 g vial	364.92	10	Azactam Aztreonam (Fresenius Kabi)
➔ Restricted (RS1277)			
Clinical microbiologist or infectious disease specialist			
CHLORAMPHENICOL – Restricted see terms below			
⚡ Cap 250 mg			
⚡ Cap 500 mg			
⚡ Inj 1 g vial			
➔ Restricted (RS1277)			
Clinical microbiologist or infectious disease specialist			
CLINDAMYCIN – Restricted see terms below			
⚡ Cap 150 mg – 5% DV Dec-24 to 2027	4.94	24	Dalacin C
⚡ Oral liq 15 mg per ml			
⚡ Inj 150 mg per ml, 4 ml ampoule – 5% DV Mar-26 to 2028	48.78	10	Dalacin C
➔ Restricted (RS1061)			
Clinical microbiologist or infectious disease specialist			
COLISTIN SULPHOMETHATE [COLESTIMETHATE] – Restricted see terms below			
⚡ Inj 2 million iu, 10 ml vial	216.67	10	Colomycin
➔ Restricted (RS1062)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
DAPTOMYCIN – Restricted see terms below			
⚡ Inj 500 mg vial – 5% DV Oct-26 to 2028	115.36 44.90	1	Daptomycin Dr Reddy's Daptomycin-AFT
<i>(Daptomycin Dr Reddy's Inj 500 mg vial to be delisted 1 October 2026)</i>			
➔ Restricted (RS1063)			
Clinical microbiologist or infectious disease specialist			
FOSFOMYCIN – Restricted see terms below			
⚡ Powder for oral solution, 3 g sachet – 5% DV Apr-25 to 2027	18.70	1	UroFos
➔ Restricted (RS1315)			
Clinical microbiologist or infectious disease specialist			
LINCOMYCIN – Restricted see terms below			
⚡ Inj 300 mg per ml, 2 ml vial			
➔ Restricted (RS1065)			
Clinical microbiologist or infectious disease specialist			
LINEZOLID – Restricted see terms below			
⚡ Tab 600 mg – 5% DV Dec-24 to 2027	194.60	10	Zyvox
⚡ Oral liq 20 mg per ml	1,879.00	150 ml	Zyvox
⚡ Inj 2 mg per ml, 300 ml bottle – 5% DV Dec-24 to 2027	155.00	10	Linezolid Kabi
➔ Restricted (RS1066)			
Clinical microbiologist or infectious disease specialist			
METHENAMINE (HEXAMINE) HIPPURATE			
Tab 1 g	19.95	100	Hiprex

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NITROFURANTOIN			
Tab 50 mg – 5% DV Dec-24 to 2027	22.20	100	Nifuran
Tab 100 mg	37.50	100	Nitrofurantoin Clinect
Cap modified-release 100 mg	81.20	100	Nifuran
<i>(Nifuran Tab 50 mg to be delisted 1 November 2026)</i>			
<i>(Nifuran Tab 100 mg to be delisted 1 November 2026)</i>			
PIVMECILLINAM – Restricted see terms below			
↓ Tab 200 mg			
→ Restricted (RS1322)			
Clinical microbiologist or infectious disease specialist			
SODIUM FUSIDATE [FUSIDIC ACID] – Restricted see terms below			
↓ Tab 250 mg	135.70	36	Fucidin
→ Restricted (RS1064)			
Clinical microbiologist or infectious disease specialist			
SULFADIAZINE SODIUM – Restricted see terms below			
↓ Tab 500 mg			<i>e.g. Sulfadiazin-Heyl; Wockhardt</i>
→ Restricted (RS1067)			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
TEICOPLANIN – Restricted see terms below			
↓ Inj 400 mg vial – 5% DV Apr-25 to 2027	38.85	1	Teicoplanin Medsurge
→ Restricted (RS1068)			
Clinical microbiologist or infectious disease specialist			
TRIMETHOPRIM			
Tab 100 mg			
Tab 300 mg – 5% DV Feb-25 to 2027	27.83	50	TMP
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]			
Tab 80 mg with sulphamethoxazole 400 mg – 5% DV Feb-25 to 2027	115.74	500	Trisul
Oral liq 8 mg with sulphamethoxazole 40 mg per ml – 5% DV Aug-25 to 2028	4.95	100 ml	Deprim
Inj 16 mg with sulphamethoxazole 80 mg per ml, 5 ml ampoule			
VANCOMYCIN – Restricted see terms below			
↓ Inj 500 mg vial – 5% DV Feb-27 to 2029	28.50	10	Vancomycin Viatrix
→ Restricted (RS1069)			
Clinical microbiologist or infectious disease specialist			

Antifungals

Imidazoles

KETOCONAZOLE

↓ Tab 200 mg

→ **Restricted (RS1410)**

Oncologist

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Polyene Antimycotics

AMPHOTERICIN B

⚡ Inj (liposomal) 50 mg vial – 5% DV Apr-26 to 2028	125.00	1	Amphotericin Liposomal SUN
---	--------	---	---------------------------------------

➡ **Restricted (RS1071)**

Initiation

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist
Either:

- 1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
- 2 Both:
 - 2.1 Possible invasive fungal infection; and
 - 2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

⚡ Inj 50 mg vial

➡ **Restricted (RS1316)**

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist

NYSTATIN

Tab 500,000 u	17.09	50	Nilstat
Cap 500,000 u	15.47	50	Nilstat

Triazoles

FLUCONAZOLE – **Restricted** see terms [below](#)

⚡ Cap 50 mg	4.10	28	Mylan
⚡ Cap 150 mg	0.45	1	Flucazole
			Mylan
⚡ Cap 200 mg	8.90	28	Mylan
⚡ Oral liquid 50 mg per 5 ml	129.02	35 ml	Diflucan
⚡ Inj 2 mg per ml, 50 ml vial	17.66	1	Diflucan
	11.20		Fluconazole-Baxter
⚡ Inj 2 mg per ml, 100 ml vial	33.84	1	Diflucan
	5.20		Fluconazole-Baxter

➡ **Restricted (RS1072)**

Consultant

ITRACONAZOLE – **Restricted** see terms [below](#)

⚡ Cap 100 mg	6.83	15	Itraconazole Crescent
			Itrazole
⚡ Oral liquid 10 mg per ml			

➡ **Restricted (RS1073)**

Clinical immunologist, clinical microbiologist, dermatologist or infectious disease specialist

POSACONAZOLE – **Restricted** see terms [below](#)

⚡ Tab modified-release 100 mg – 5% DV Dec-25 to 2028	123.60	24	Posaconazole Juno
⚡ Oral liq 40 mg per ml – 5% DV Dec-25 to 2028	308.26	105 ml	Devatis

➡ **Restricted (RS2052)**

Initiation

Haematologist or infectious disease specialist

Re-assessment required after 6 weeks

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Either:
 - 1.1 Patient has acute myeloid leukaemia; or
 - 1.2 Patient is planned to receive a stem cell transplant and is at high risk for aspergillus infection; and
- 2 Patient is to be treated with high dose remission induction therapy or re-induction therapy.

Continuation

Haematologist or infectious disease specialist

Re-assessment required after 6 weeks

Both:

- 1 Patient has previously received posaconazole prophylaxis during remission induction therapy; and
- 2 Any of the following:
 - 2.1 Patient is to be treated with high dose remission re-induction therapy; or
 - 2.2 Patient is to be treated with high dose consolidation therapy; or
 - 2.3 Patient is receiving a high risk stem cell transplant.

Initiation – Invasive fungal infection prophylaxis

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Continuation – Invasive fungal infection prophylaxis

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

VORICONAZOLE – **Restricted** see terms [below](#)

↓ Tab 50 mg – 5% DV Aug-25 to 2028	71.00	56	Vttack
↓ Tab 200 mg – 5% DV Aug-25 to 2028	263.00	56	Vttack
↓ Powder for oral suspension 40 mg per ml	1,523.22	70 ml	Vfend
↓ Inj 200 mg vial – 5% DV Dec-25 to 2028	16.89	1	AFT

→ **Restricted (RS2053)**

Initiation – Proven or probable aspergillus infection

Clinical microbiologist, haematologist or infectious disease specialist

Both:

- 1 Patient is immunocompromised; and
- 2 Patient has proven or probable invasive aspergillus infection.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – Possible aspergillus infection

Clinical microbiologist, haematologist or infectious disease specialist

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient has possible invasive aspergillus infection; and
- 3 A multidisciplinary team (including an infectious disease physician) considers the treatment to be appropriate.

Initiation – Resistant candidiasis infections and other moulds

Clinical microbiologist, haematologist or infectious disease specialist

All of the following:

- 1 Patient is immunocompromised; and
- 2 Either:
 - 2.1 Patient has fluconazole resistant candidiasis; or
 - 2.2 Patient has mould strain such as Fusarium spp. and Scedosporium spp; and
- 3 A multidisciplinary team (including an infectious disease physician or clinical microbiologist) considers the treatment to be appropriate.

Initiation – Invasive fungal infection prophylaxis

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Continuation – Invasive fungal infection prophylaxis

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Other Antifungals

CASPOFUNGIN – **Restricted** see terms [below](#)

⚡ Inj 50 mg vial – 5% DV Mar-26 to 2028	110.00	1	Alchemy Caspofungin
⚡ Inj 70 mg vial – 5% DV Mar-26 to 2028	135.00	1	Alchemy Caspofungin

➡ **Restricted (RS1076)**

Initiation

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
- 2 Both:
 - 2.1 Possible invasive fungal infection; and
 - 2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

FLUCYTOSINE – **Restricted** see terms [below](#)

↓ Tab 500 mg

↓ Cap 500 mg

→ **Restricted** ([RS1279](#))

Clinical microbiologist or infectious disease specialist

TERBINAFINE

Tab 250 mg	8.97	84	Deolate
------------------	------	----	---------

Antimycobacterials

Antileprotics

CLOFAZIMINE – **Restricted** see terms [below](#)

↓ Cap 50 mg

→ **Restricted** ([RS1077](#))

Clinical microbiologist, dermatologist or infectious disease specialist

DAPSONE – **Restricted** see terms [below](#)

↓ Tab 25 mg	268.50	100	Dapsone
-------------------	--------	-----	---------

↓ Tab 100 mg	329.50	100	Dapsone
--------------------	--------	-----	---------

→ **Restricted** ([RS1078](#))

Clinical microbiologist, dermatologist or infectious disease specialist

Antituberculars

BEDAQUILINE – **Restricted** see terms [below](#)

↓ Tab 100 mg	3,084.51	24	Sirturo
--------------------	----------	----	---------

→ **Restricted** ([RS1977](#))

Initiation – multi-drug resistant tuberculosis

Limited to 6 months treatment

Both:

- 1 The person has multi-drug resistant tuberculosis (MDR-TB); and
- 2 Ministry of Health's Tuberculosis Clinical Network has reviewed the individual case and recommends bedaquiline as part of the treatment regimen.

CYCLOSERINE – **Restricted** see terms [below](#)

↓ Cap 250 mg

→ **Restricted** ([RS1079](#))

Clinical microbiologist, infectious disease specialist or respiratory specialist

ETHAMBUTOL HYDROCHLORIDE – **Restricted** see terms [below](#)

↓ Tab 100 mg

↓ Tab 400 mg	49.34	56	Myambutol
--------------------	-------	----	-----------

→ **Restricted** ([RS1080](#))

Clinical microbiologist, infectious disease specialist or respiratory specialist

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ISONIAZID – Restricted see terms below			
⚡ Tab 100 mg – 5% DV May-25 to 2027	94.50	100	Isoniazid Teva
	327.41		Noumed Isoniazid
➡ Restricted (RS1281)			
Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician			
ISONIAZID WITH RIFAMPICIN – Restricted see terms below			
⚡ Tab 100 mg with rifampicin 150 mg – 5% DV Feb-25 to 2027	89.82	100	Rifinah
⚡ Tab 150 mg with rifampicin 300 mg – 5% DV Feb-25 to 2027	179.13	100	Rifinah
⚡ Cap 100 mg with rifampicin 150 mg	199.00	100	Rifamazid
➡ Restricted (RS1282)			
Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician			
PARA-AMINOSALICYLIC ACID – Restricted see terms below			
⚡ Grans for oral liq 4 g	280.00	30	Paser
➡ Restricted (RS1083)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PROTIONAMIDE – Restricted see terms below			
⚡ Tab 250 mg	305.00	100	Peteha
➡ Restricted (RS1084)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PYRAZINAMIDE – Restricted see terms below			
⚡ Tab 500 mg			
➡ Restricted (RS1085)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
RIFABUTIN – Restricted see terms below			
⚡ Cap 150 mg	353.71	30	Mycobutin
➡ Restricted (RS1086)			
Clinical microbiologist, gastroenterologist, infectious disease specialist or respiratory specialist			
RIFAMPICIN – Restricted see terms below			
⚡ Cap 150 mg	58.54	100	Rifadin
⚡ Cap 300 mg	122.06	100	Rifadin
⚡ Oral liq 100 mg per 5 ml	12.60	60 ml	Rifadin
⚡ Inj 600 mg vial – 5% DV Dec-26 to 2029	134.98	1	Rifadin
➡ Restricted (RS1087)			
Clinical microbiologist, dermatologist, internal medicine physician, paediatrician or public health physician			

Antiparasitics

Anthelmintics

ALBENDAZOLE – Restricted see terms [below](#)

⚡ Tab 200 mg

⚡ Tab 400 mg

➡ **Restricted (RS1088)**

Clinical microbiologist or infectious disease specialist

IVERMECTIN – Restricted see terms [below](#)

⚡ Tab 3 mg 17.20 | 4 | Stromectol |

➡ **Restricted (RS1283)**

Clinical microbiologist, dermatologist or infectious disease specialist

102

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)
e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MEBENDAZOLE			
Tab 100 mg – 5% DV Dec-24 to 2027	5.18	6	Vermox
Oral liq 100 mg per 5 ml			
PRAZIQUANTEL			
Tab 600 mg			
Antiprotazoals			
ARTEMETHER WITH LUMEFANTRINE – Restricted see terms below			
↓ Tab 20 mg with lumefantrine 120 mg			
→ Restricted (RS1090)			
Clinical microbiologist or infectious disease specialist			
ARTESUNATE – Restricted see terms below			
↓ Inj 60 mg vial			
→ Restricted (RS1091)			
Clinical microbiologist or infectious disease specialist			
ATOVAQUONE WITH PROGUANIL HYDROCHLORIDE – Restricted see terms below			
↓ Tab 62.5 mg with proguanil hydrochloride 25 mg.....	29.50	12	Malarone Junior
↓ Tab 250 mg with proguanil hydrochloride 100 mg.....	72.00	12	Malarone
→ Restricted (RS1092)			
Clinical microbiologist or infectious disease specialist			
CHLOROQUINE PHOSPHATE – Restricted see terms below			
↓ Tab 250 mg			
→ Restricted (RS1093)			
Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist			
MEFLOQUINE – Restricted see terms below			
↓ Tab 250 mg			
→ Restricted (RS1094)			
Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist			
METRONIDAZOLE			
Tab 200 mg	25.86	250	Metronidamed
Tab 400 mg	4.29	21	Metronidamed
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	Flagyl-S
Inj 5 mg per ml, 100 ml bag	18.00	10	Baxter
Suppos 500 mg	24.48	10	Flagyl
NITAZOXANIDE – Restricted see terms below			
↓ Tab 500 mg			
↓ Oral liq 100 mg per 5 ml			
→ Restricted (RS1095)			
Clinical microbiologist or infectious disease specialist			
ORNIDAZOLE			
Tab 500 mg – 5% DV Mar-25 to 2027	36.52	10	Arrow-Ornidazole
PENTAMIDINE ISETHIONATE – Restricted see terms below			
↓ Inj 300 mg vial	216.00	5	Pentacarinat
	638.69		Tillomed
→ Restricted (RS1096)			
Clinical microbiologist or infectious disease specialist			
PRIMAQUINE – Restricted see terms on the next page			
↓ Tab 15 mg			
↓ Tab 7.5 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted (RS1097)			
Clinical microbiologist or infectious disease specialist			
PYRIMETHAMINE – Restricted see terms below			
⚡ Tab 25 mg			
➔ Restricted (RS1098)			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
QUININE DIHYDROCHLORIDE – Restricted see terms below			
⚡ Inj 60 mg per ml, 10 ml ampoule			
⚡ Inj 300 mg per ml, 2 ml vial			
➔ Restricted (RS1099)			
Clinical microbiologist or infectious disease specialist			
SODIUM STIBOGLUCONATE – Restricted see terms below			
⚡ Inj 100 mg per ml, 1 ml vial			
➔ Restricted (RS1100)			
Clinical microbiologist or infectious disease specialist			
SPIRAMYCIN – Restricted see terms below			
⚡ Tab 500 mg			
➔ Restricted (RS1101)			
Maternal-foetal medicine specialist			

Antiretrovirals

Non-Nucleoside Reverse Transcriptase Inhibitors

➔ Restricted (RS1898)			
Initiation – Confirmed HIV			
Patient has confirmed HIV infection.			
Initiation – Prevention of maternal transmission			
Either:			
1 Prevention of maternal foetal transmission; or			
2 Treatment of the newborn for up to eight weeks.			
Initiation – Post-exposure prophylaxis following exposure to HIV			
Both:			
1 Treatment course to be initiated within 72 hours post exposure; and			
2 Any of the following:			
2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or			
2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or			
2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or			
2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.			
Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).			
Initiation – Percutaneous exposure			
Patient has percutaneous exposure to blood known to be HIV positive.			
EFAVIRENZ – Restricted see terms above			
➔ Tab 600 mg – Restricted: For continuation only	65.38	30	Efavirenz Milpharm
⚡ Oral liq 30 mg per ml			
<i>(Efavirenz Milpharm Tab 600 mg to be delisted 1 November 2026)</i>			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ETRAVIRINE – Restricted see terms on the previous page			
† Tab 200 mg	770.00	60	Intelence
NEVIRAPINE – Restricted see terms on the previous page			
† Tab 200 mg – 5% DV Feb-25 to 2027	198.25	60	Nevirapine Viatris
† Oral suspension 10 mg per ml	203.55	240 ml	Viramune Suspension

Nucleoside Reverse Transcriptase Inhibitors

→ **Restricted (RS1899)**

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (<https://www.ashm.org.au/hiv/hiv-management/pep/>).

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

ABACAVIR SULPHATE – **Restricted** see terms [above](#)

† Tab 300 mg	180.00	60	Ziagen
† Oral liq 20 mg per ml			

ABACAVIR SULPHATE WITH LAMIVUDINE – **Restricted** see terms [above](#)

† Tab 600 mg with lamivudine 300 mg – 5% DV Feb-26 to 2028	35.00	30	Abacavir/lamivudine Viatris
---	-------	----	------------------------------------

EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPROXIL – **Restricted** see terms [above](#)

† Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate)	106.88	30	Viatris
† Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a fumarate)	106.88	30	TEEVIR TEEVIR (India) Tenofovir Efavirenz Emtricitabine (-Mylan)

EMTRICITABINE – **Restricted** see terms [above](#)

† Cap 200 mg	307.20	30	Emtriva
--------------------	--------	----	---------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LAMIVUDINE – Restricted see terms on the previous page			
† Tab 150 mg.....	98.00	60	Lamivudine Viatris
† Oral liq 10 mg per ml			
STAVUDINE – Restricted see terms on the previous page			
† Cap 30 mg			
† Cap 40 mg			
† Powder for oral soln 1 mg per ml			
ZIDOVUDINE – Restricted see terms on the previous page			
† Cap 100 mg.....	152.25	100	Retrovir
† Oral liq 10 mg per ml.....	30.45	200 ml	Retrovir
† Inj 10 mg per ml, 20 ml vial.....	750.00	5	Retrovir IV
ZIDOVUDINE WITH LAMIVUDINE – Restricted see terms on the previous page			
† Tab 300 mg with lamivudine 150 mg.....	92.40	60	Lamivudine/Zidovudine Viatris

Protease Inhibitors

➡ **Restricted (RS1900)**

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (<https://www.ashm.org.au/hiv/hiv-management/pep/>).

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

ATAZANAVIR SULPHATE – Restricted see terms [above](#)

† Cap 150 mg – 5% DV Feb-26 to 2028.....	102.50	60	Atazanavir Viatris
† Cap 200 mg – 5% DV Feb-26 to 2028.....	152.30	60	Atazanavir Viatris

DARUNAVIR – Restricted see terms [above](#)

† Tab 400 mg – 5% DV Dec-26 to 2029.....	115.00	60	Darunavir Viatris
† Tab 600 mg – 5% DV Dec-26 to 2029.....	130.00	60	Darunavir Viatris

INDINAVIR – Restricted see terms [above](#)

- † Cap 200 mg
- † Cap 400 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LOPINAVIR WITH RITONAVIR – Restricted see terms on the previous page			
† Tab 100 mg with ritonavir 25 mg			
† Tab 200 mg with ritonavir 50 mg – 5% DV Feb-25 to 2027	875.00	120	Lopinavir/Ritonavir Mylan
† Oral liq 80 mg per ml with ritonavir 20 mg per ml			
RITONAVIR – Restricted see terms on the previous page			
† Tab 100 mg	43.31	30	Norvir

Strand Transfer Inhibitors

→ **Restricted (RS1901)**

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (<https://www.ashm.org.au/hiv/hiv-management/pep/>).

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

DOLUTEGRAVIR – **Restricted** see terms [above](#)

† Tab 50 mg	1,090.00	30	Tivicay
-------------------	----------	----	---------

DOLUTEGRAVIR WITH LAMIVUDINE – **Restricted** see terms [above](#)

† Tab 50 mg with lamivudine 300 mg	1,090.00	30	Dovato
--	----------	----	--------

RALTEGRAVIR POTASSIUM – **Restricted** see terms [above](#)

† Tab 400 mg	1,090.00	60	Isentress
† Tab 600 mg	1,090.00	60	Isentress HD

Antivirals

Hepatitis B

ENTECAVIR

Tab 0.5 mg	12.04	30	Entecavir (Rex)
------------------	-------	----	-----------------

LAMIVUDINE

Tab 100 mg	12.06	28	Zetlam
Oral liq 5 mg per ml	270.00	240 ml	Zeffix

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
TENOFOVIR DISOPROXIL			
Tab 245 mg (300 mg as a maleate) – 5% DV Dec-25 to 2028	13.80	30	Tenofovir Disoproxil
Tab 245 mg (300 mg as a fumarate)	13.80	30	Viartis Ricovir

Hepatitis C

GLECAPREVIR WITH PIBRENTASVIR

Note: the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website <https://www.pharmac.govt.nz/maviret>.

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 Maviret

LEDIPASVIR WITH SOFOSBUVIR – **Restricted** see terms [below](#)

⚡ Tab 90 mg with sofosbuvir 400 mg.....24,363.46 28 Harvoni

➡ **Restricted (RS1528)**

Note: Only for use in patients with approval by the Hepatitis C Treatment Panel (HepCTP). Applications will be considered by HepCTP at its regular meetings and approved subject to eligibility according to the Access Criteria (set out in Section B of the Pharmaceutical Schedule).

Herpesviridae

ACICLOVIR

Tab dispersible 200 mg – **5% DV Feb-26 to 2028**2.05 25 **Lovir**
Tab dispersible 400 mg – **5% DV Feb-26 to 2028**7.55 56 **Lovir**
Tab dispersible 800 mg – **5% DV Feb-26 to 2028**7.43 35 **Lovir**
Inj 250 mg vial – **5% DV Feb-25 to 2027**13.75 5 Aciclovir Injection DBL
Aciclovir-Baxter

CIDOFOVIR – **Restricted** see terms [below](#)

⚡ Inj 75 mg per ml, 5 ml vial

➡ **Restricted (RS1108)**

Clinical microbiologist, infectious disease specialist, otolaryngologist or oral surgeon

FOSCARNET SODIUM – **Restricted** see terms [below](#)

⚡ Inj 24 mg per ml, 250 ml bottle

➡ **Restricted (RS1109)**

Clinical microbiologist or infectious disease specialist

GANCICLOVIR – **Restricted** see terms [below](#)

⚡ Inj 500 mg vial380.00 5 Cymevene

➡ **Restricted (RS1110)**

Clinical microbiologist or infectious disease specialist

LETTERMOVIR – **Restricted** see terms [below](#)

⚡ Tab 240 mg6,664.00 28 Prevymis

➡ **Restricted (RS2189)**

Initiation – CMV prophylaxis post HSCT

Limited to 4 months treatment

All of the following:

- 1 Patient has undergone an allogeneic haematopoietic stem cell transplant; and
- 2 The patient has confirmed presence of cytomegalovirus-specific antibodies; and
- 3 Treatment to commence within 28 days of an allogeneic haematopoietic stem cell transplant; and
- 4 Maximum treatment duration of 100 days post-transplant.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – CMV prophylaxis second or subsequent HSCT

Re-assessment required after 4 months

All of the following:

- 1 Patient has undergone an allogeneic haematopoietic stem cell transplant; and
- 2 The patient has confirmed presence of cytomegalovirus-specific antibodies; and
- 3 Treatment to commence within 28 days of an allogeneic haematopoietic stem cell transplant; and
- 4 Maximum treatment duration of 100 days post-transplant.

Initiation – CMV prophylaxis in severe immunosuppression*

Clinical microbiologist or infectious disease specialist

Re-assessment required after 6 months

Both:

- 1 Patient has severe immunosuppression requiring prophylaxis of CMV; and
- 2 Either:
 - 2.1 Patient is contraindicated to all other funded CMV active oral antiviral agents; or
 - 2.2 Patients CMV is resistant to all other funded CMV active oral antiviral agents.

Continuation – CMV prophylaxis in severe immunosuppression*

Clinical microbiologist or infectious disease specialist

Re-assessment required after 6 months

Both:

- 1 Patient has severe immunosuppression requiring prophylaxis of CMV; and
- 2 Either:
 - 2.1 Patient is contraindicated to all other funded CMV active oral antiviral agents; or
 - 2.2 Patients CMV is resistant to all other funded CMV active oral antiviral agents.

Note: Indications marked with * are unapproved indications.

VALACICLOVIR

Tab 500 mg – 5% DV Feb-25 to 2027	9.64	30	Vaclovir
	13.50	42	Valaciclovir Viatris
Tab 1,000 mg – 5% DV Feb-25 to 2027	17.78	30	Vaclovir
	12.45	21	Valaciclovir Mylan
			Valaciclovir Viatris

VALGANCICLOVIR – Restricted see terms [below](#)

↓ Tab 450 mg – 5% DV Feb-25 to 2027	140.89	60	Valganciclovir Viatris
→ Restricted (RS2137)			

Initiation – Transplant cytomegalovirus prophylaxis

Re-assessment required after 3 months

Patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Continuation – Transplant cytomegalovirus prophylaxis

Re-assessment required after 3 months

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – Lung transplant cytomegalovirus prophylaxis

Re-assessment required after 12 months

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Continuation – Lung transplant cytomegalovirus prophylaxis

Re-assessment required after 12 months

All of the following:

- 1 Patient has undergone a lung re-transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initiation – Cytomegalovirus in immunocompromised patients

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – **Restricted** see terms [below](#)

⚡ Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a maleate) – 5% DV Dec-25 to 2028	13.45	30	Tenofovir Disoproxil Emtricitabine Viatr
⚡ Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a fumarate).....	13.45	30	Tenofovir Disoproxil Emtricitabine Mylan

➡ **Restricted (RS1902)**

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

continued...

	Price		Brand or
(ex man. excl. GST)			Generic
\$	Per		Manufacturer

continued...

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

Initiation – Pre-exposure prophylaxis

Re-assessment required after 24 months

Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines (<https://ashm.org.au/HIV/PrEP/>)

Continuation – Pre-exposure prophylaxis

Re-assessment required after 24 months

Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines (<https://ashm.org.au/HIV/PrEP/>)

Influenza

OSELTAMIVIR – **Restricted** see terms [below](#)

Note: The restriction on the use of oseltamivir to hospitalised patients means that supply into the community for a new course is not permitted. Supply of a part original pack on discharge where initiated as a hospital inpatient is permitted.

↓ Tab 75 mg

↓ Powder for oral suspension 6 mg per ml

→ **Restricted** ([RS1307](#))

Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a Health NZ Hospital approved infections control plan.

ZANAMIVIR

Note: The restriction on the use of zanamivir to hospitalised patients means that supply into the community for a new course is not permitted. Supply of a part original pack on discharge where initiated as a hospital inpatient is permitted.

↓ Powder for inhalation 5 mg 37.38 20 dose Relenza Rotadisk

→ **Restricted** ([RS1369](#))

Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a Health NZ Hospital approved infections control plan.

COVID-19 Treatments

NIRMATRELVIR WITH RITONAVIR – **Restricted** see terms [below](#)

↓ Tab 150 mg with ritonavir 100 mg 1,274.00 30 Paxlovid

→ **Restricted** ([RS1894](#))

Initiation

Only if patient meets access criteria (as per <https://pharmac.govt.nz/covid-oral-antivirals>). Note the supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
REMDESIVIR			
Note: For patients meeting access criteria for oral antiviral treatments (as on Pharmac's website).			
Inj 100 mg vial	615.23	1	Veklury

Immune Modulators

INTERFERON ALFA-2B

- Inj 18 m iu, 1.2 ml multidose pen
- Inj 30 m iu, 1.2 ml multidose pen
- Inj 60 m iu, 1.2 ml multidose pen

INTERFERON GAMMA – **Restricted** see terms [below](#)

- ⚡ Inj 100 mcg in 0.5 ml vial

➔ **Restricted (RS1113)**

Initiation

Patient has chronic granulomatous disease and requires interferon gamma.

PEGYLATED INTERFERON ALFA-2A – **Restricted** see terms [below](#)

⚡ Inj 135 mcg prefilled syringe.....	887.35	1	Pegasys (S29)
⚡ Inj 180 mcg prefilled syringe.....	1,074.79	4	Pegasys
	1,355.71		Pegasys

➔ **Restricted (RS1827)**

Initiation – Chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant

Limited to 48 weeks treatment

Any of the following:

- 1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
- 2 Patient has chronic hepatitis C and is co-infected with HIV; or
- 3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant.

Notes: Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.

Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml.

Continuation – Chronic hepatitis C - genotype 1 infection

Gastroenterologist, infectious disease specialist or general physician

Re-assessment required after 48 weeks

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir.

Initiation – Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 3.1 Patient has responder relapsed; or
- 3.2 Patient was a partial responder; or
- 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir.

Initiation – Chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV

Limited to 6 months treatment

Patient has chronic hepatitis C, genotype 2 or 3 infection.

Initiation – Hepatitis B

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naïve; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log₁₀ IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 Serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (greater than or equal to Metavir Stage F2 or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon.

Initiation – myeloproliferative disorder or cutaneous T cell lymphoma

Re-assessment required after 12 months

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Continuation – myeloproliferative disorder or cutaneous T cell lymphoma

Re-assessment required after 12 months

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications

Initiation – ocular surface squamous neoplasia

Ophthalmologist

Re-assessment required after 12 months

Patient has ocular surface squamous neoplasia*.

Continuation – ocular surface squamous neoplasia

Ophthalmologist

Re-assessment required after 12 months

The treatment remains appropriate and patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initiation – post-allogenic bone marrow transplant

Re-assessment required after 3 months

Patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Continuation – post-allogenic bone marrow transplant

Re-assessment required after 3 months

Patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Anticholinesterases

EDROPHONIUM CHLORIDE – **Restricted** see terms [below](#)

↓ Inj 10 mg per ml, 15 ml vial

↓ Inj 10 mg per ml, 1 ml ampoule

→ **Restricted (RS1015)**

Initiation

For the diagnosis of myasthenia gravis.

NEOSTIGMINE METILSULFATE

Inj 2.5 mg per ml, 1 ml ampoule – **5% DV Feb-25 to 2027** 48.25 10 **Max Health**

NEOSTIGMINE METILSULFATE WITH GLYCOPYRRONIUM BROMIDE

Inj 2.5 mg with glycopyrronium bromide 0.5 mg per ml, 1 ml ampoule –
5% DV Feb-26 to 2028 26.13 10 **Max Health**

PYRIDOSTIGMINE BROMIDE

Tab 60 mg 50.28 100 Mestinon

Antirheumatoid Agents

HYDROXYCHLOROQUINE SULPHATE

Tab 200 mg – **5% DV May-25 to 2027** 7.80 100 **Ipca-
Hydroxychloroquine**

LEFLUNOMIDE

Tab 10 mg 6.00 30 Arava

Tab 20 mg 6.00 30 Arava

PENICILLAMINE

Tab 125 mg 67.23 100 D-Penamine

Tab 250 mg 110.12 100 D-Penamine

SODIUM AUROTHIOMALATE

Inj 10 mg in 0.5 ml ampoule

Inj 20 mg in 0.5 ml ampoule

Inj 50 mg in 0.5 ml ampoule

Drugs Affecting Bone Metabolism

Bisphosphonates

ALENDRONATE SODIUM

Tab 70 mg 3.10 4 Fosamax

ALENDRONATE SODIUM WITH COLECALCIFEROL

Tab 70 mg with colecalciferol 5,600 iu 1.99 4 Fosamax Plus

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial 32.49 1 Pamisol

Inj 6 mg per ml, 10 ml vial 88.11 1 Pamisol

Inj 9 mg per ml, 10 ml vial 94.34 1 Pamisol

RISEDRONATE SODIUM

Tab 35 mg – **5% DV Feb-26 to 2028** 3.00 4 **Risedronate Sandoz**

ZOLEDRONIC ACID

Inj 5 mg per 100 ml, bag – **5% DV Feb-26 to 2028** 19.45 1 **Zoledronic Acid Viatrix**

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Other Drugs Affecting Bone Metabolism

DENOSUMAB – **Restricted** see terms [below](#)

Note: Denosumab inj 60 mg per 1 ml pre-filled syringe is Medsafe approved for use in osteoporosis. Denosumab inj 120 mg per 1.7 ml vial is Medsafe approved for use in hypercalcaemia of malignancy.

⚡ Inj 120 mg per 1.7 ml vial	375.00	1	Xgeva
⚡ Inj 60 mg per 1 ml prefilled syringe.....	187.50	1	Prolia

➡ **Restricted (RS2097)**

Initiation – Osteoporosis

All of the following:

- 1 The patient has established osteoporosis; and
- 2 Any of the following:
 - 2.1 History of one significant osteoporotic fracture demonstrated radiologically, with a documented T-Score less than or equal to -2.5, that incorporates BMD measured using dual-energy x-ray absorptiometry (DEXA); or
 - 2.2 History of one significant osteoporotic fracture, demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of logistical, technical or pathophysiological reasons; or
 - 2.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 2.4 Documented T-Score less than or equal to -3.0; or
 - 2.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm that incorporates BMD measured using DEXA; and
- 3 Any of the following:
 - 3.1 Bisphosphonates are contraindicated because the patient's creatinine clearance or eGFR is less than 35 mL/min; or
 - 3.2 The patient has experienced at least two symptomatic new fractures or a BMD loss greater than 2% per year, after at least 12 months' continuous therapy with a funded antiresorptive agent; or
 - 3.3 Bisphosphonates result in intolerable side effects; or
 - 3.4 Intravenous bisphosphonates cannot be administered due to logistical or technical reasons.

Initiation – Hypercalcaemia

Both:

- 1 Patient has hypercalcaemia of malignancy; and
- 2 Patient has severe renal impairment.

TERIPARATIDE – **Restricted** see terms [below](#)

⚡ Inj 250 mcg per ml, 2.4 ml	490.00	1	Forteo
	200.27		Teriparatide - Teva

➡ **Restricted (RS1143)**

Initiation

Limited to 18 months treatment

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this restriction are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid

continued...

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$ Per	

continued...

5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Enzymes

HYALURONIDASE

Inj 1,500 iu ampoule

Hyperuricaemia and Antigout

ALLOPURINOL

Tab 100 mg	17.99	1,000	Ipca-Allopurinol
Tab 300 mg	22.50	500	Ipca-Allopurinol

BENZBROMARONE – **Restricted:** For continuation only

➔ Tab 50 mg			
➔ Tab 100 mg	45.00	100	Benzbromaron AL 100

COLCHICINE

Tab 500 mcg – 5% DV Dec-26 to 2028	3.15	100	Colchicine Indoco
	6.30		Colgout

(Colgout Tab 500 mcg to be delisted 1 December 2026)

FEBUXOSTAT – **Restricted** see terms [below](#)

↓ Tab 80 mg – 5% DV Dec-26 to 2029	4.66	28	Febuxostat (Teva)
↓ Tab 120 mg – 5% DV Dec-26 to 2029	11.67	28	Febuxostat (Teva)

➔ **Restricted (RS1844)**

Initiation – Gout

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout..

Initiation – Tumour lysis syndrome

Haematologist or oncologist

Re-assessment required after 6 weeks

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Continuation – Tumour lysis syndrome

Haematologist or oncologist

Re-assessment required after 6 weeks

The treatment remains appropriate and patient is benefitting from treatment.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PROBENECID Tab 500 mg			
RASBURICASE – Restricted see terms below			
⚡ Inj 1.5 mg vial			
➡ Restricted (RS1016)			
Haematologist			

Muscle Relaxants and Related Agents

ATRACURIUM BESYLATE			
Inj 10 mg per ml, 2.5 ml ampoule	7.69	5	Medsurge
Inj 10 mg per ml, 5 ml ampoule	9.86	5	Medsurge
BACLOFEN			
Tab 10 mg – 5% DV Dec-24 to 2027	3.70	100	Pacifen
Oral liq 1 mg per ml			
Inj 0.05 mg per ml, 1 ml ampoule	11.55	1	Lioresal Intrathecal
Inj 2 mg per ml, 5 ml ampoule – 5% DV Mar-25 to 2027	490.91	10	Sintetica Baclofen Intrathecal
CLOSTRIDIUM BOTULINUM TYPE A TOXIN			
Inj 100 u vial	467.50	1	Botox
Inj 300 u vial	388.50	1	Dysport
Inj 500 u vial	1,295.00	2	Dysport
DANTROLENE			
Cap 25 mg	145.77	100	Dantrium
Cap 50 mg	77.00	100	Dantrium
Inj 20 mg vial	1,143.74	6	Dantrium IV
MIVACURIUM CHLORIDE			
Inj 2 mg per ml, 10 ml ampoule			
ORPHENADRINE CITRATE			
Tab 100 mg – 5% DV Feb-25 to 2027	23.25	100	Norflex
PANCURONIUM BROMIDE			
Inj 2 mg per ml, 2 ml ampoule			
ROCURONIUM BROMIDE			
Inj 10 mg per ml, 5 ml vial – 5% DV May-26 to 2028	28.96	10	Medsurge
SUXAMETHONIUM CHLORIDE			
Inj 50 mg per ml, 2 ml ampoule – 5% DV Dec-26 to 2029	27.61	10	Martindale
VECURONIUM BROMIDE			
Inj 10 mg vial – 5% DV Apr-25 to 2027	380.00	10	Vecure

Reversers of Neuromuscular Blockade

SUGAMMADEX – Restricted see terms below			
⚡ Inj 100 mg per ml, 2 ml vial – 5% DV Dec-24 to 2027	80.64	10	Sugammadex BNM
⚡ Inj 100 mg per ml, 5 ml vial – 5% DV Dec-24 to 2027	201.60	10	Sugammadex BNM
➡ Restricted (RS1370)			

Initiation

Any of the following:

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$ Per	Manufacturer

continued...

- 1 Patient requires reversal of profound neuromuscular blockade following rapid sequence induction that has been undertaken using rocuronium (i.e. suxamethonium is contraindicated or undesirable); or
- 2 Severe neuromuscular degenerative disease where the use of neuromuscular blockade is required; or
- 3 Patient has an unexpectedly difficult airway that cannot be intubated and requires a rapid reversal of anaesthesia and neuromuscular blockade; or
- 4 The duration of the patient's surgery is unexpectedly short; or
- 5 Neostigmine or a neostigmine/anticholinergic combination is contraindicated (for example the patient has ischaemic heart disease, morbid obesity or COPD); or
- 6 Patient has a partial residual block after conventional reversal.

Non-Steroidal Anti-Inflammatory Drugs

CELECOXIB

Cap 100 mg – 5% DV Feb-26 to 2028.....	3.60	60	Celebrex
Cap 200 mg – 5% DV Jul-26 to 2028.....	2.55	30	Celostea

DICLOFENAC SODIUM

Tab EC 25 mg – 5% DV Feb-25 to 2027.....	2.19	50	Diclofenac Sandoz
Tab 50 mg dispersible.....	1.50	20	Voltaren D
Tab EC 50 mg – 5% DV Feb-25 to 2027.....	2.19	50	Diclofenac Sandoz
Tab long-acting 75 mg – 5% DV Aug-25 to 2028.....	10.00	100	Voltaren SR
Inj 25 mg per ml, 3 ml ampoule.....	13.20	5	Voltaren
Suppos 12.5 mg.....	2.04	10	Voltaren
Suppos 25 mg.....	2.44	10	Voltaren
Suppos 50 mg.....	4.22	10	Voltaren
Suppos 100 mg.....	7.00	10	Voltaren

ETORICOXIB – Restricted see terms [below](#)

↓ Tab 30 mg

↓ Tab 60 mg

↓ Tab 90 mg

↓ Tab 120 mg

→ **Restricted (RS1592)**

Initiation

For in-vivo investigation of allergy only.

IBUPROFEN

Tab 200 mg - 1,000 tablet pack.....	21.40	1,000	Relieve
→ Tab 400 mg – Restricted: For continuation only			
→ Tab 600 mg – Restricted: For continuation only			
Tab long-acting 800 mg – 5% DV Apr-25 to 2027.....	3.65	30	Ibuprofen SR BNM
Oral liq 20 mg per ml – 5% DV Apr-25 to 2027.....	2.85	200 ml	Ethics
Inj 5 mg per ml, 2 ml ampoule			
Inj 10 mg per ml, 2 ml vial			

INDOMETACIN [INDOMETHACIN]

Cap 25 mg
Cap 50 mg
Cap long-acting 75 mg
Inj 1 mg vial
Suppos 100 mg

KETOPROFEN

Cap long-acting 200 mg.....	12.07	28	Oruvail SR
<i>(Oruvail SR Cap long-acting 200 mg to be delisted 1 October 2026)</i>			

MUSCULOSKELETAL SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MEFENAMIC ACID – Restricted: For continuation only			
➡ Cap 250 mg			
NAPROXEN			
Tab 250 mg – 5% DV Feb-25 to 2027	3.92	50	Inza
	7.06	90	Noflam 250
	39.23	500	Noflam 250
Tab 500 mg – 5% DV Feb-25 to 2027	6.89	50	Inza
	34.45	250	Noflam 500
Tab long-acting 750 mg – 5% DV Feb-25 to 2027	10.40	28	Naprosyn SR 750
Tab long-acting 1 g – 5% DV Feb-25 to 2027	11.50	28	Naprosyn SR 1000
PARECOXIB			
Inj 40 mg vial – 5% DV Dec-24 to 2027	46.00	10	Dynastat
SULINDAC			
Tab 100 mg			
Tab 200 mg			
TENOXICAM			
Tab 20 mg – 5% DV Feb-26 to 2028	23.50	100	Tilcotil
Inj 20 mg vial	9.95	1	AFT

Topical Products for Joint and Muscular Pain

CAPSAICIN – Restricted see terms below			
⚠ Crm 0.025%	9.75	45 g	Zo-Rub Osteo Zostrix

➡ **Restricted** ([RS1309](#))

Initiation

Patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

Agents for Parkinsonism and Related Disorders

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE – **Restricted** see terms [below](#)

↓ Tab 50 mg – **5% DV Feb-25 to 2027** 117.00 56 **Rilutek**
 → **Restricted (RS1351)**

Initiation

Neurologist or respiratory specialist

Re-assessment required after 6 months

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Continuation

Re-assessment required after 18 months

All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg – **5% DV Feb-26 to 2028** 126.02 112 **Motetis**

Anticholinergics

BENZATROPINE MESYLATE

Tab 2 mg 10.99 60 Benztrop
 Inj 1 mg per ml, 2 ml ampoule 95.00 5 Phebra

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE

Cap 100 mg 38.24 60 Symmetrel

APOMORPHINE HYDROCHLORIDE

Inj 10 mg per ml, 2 ml ampoule 59.50 5 Movapo
 Inj 10 mg per ml, 5 ml ampoule 121.84 5 Movapo

BROMOCRIPTINE

Cap 5 mg

ENTACAPONE

Tab 200 mg – **5% DV Jul-25 to 2027** 13.73 100 **Entacapone Viartis**

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LEVODOPA WITH BENSERAZIDE			
Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	Madopar Rapid
Cap 50 mg with benserazide 12.5 mg	13.75	100	Madopar 62.5
Cap 100 mg with benserazide 25 mg	15.80	100	Madopar 125
Cap long-acting 100 mg with benserazide 25 mg	22.85	100	Madopar HBS
Cap 200 mg with benserazide 50 mg	26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
Tab 100 mg with carbidopa 25 mg – 5% DV Feb-25 to 2027	26.49	100	Sinemet
Tab long-acting 100 mg with carbidopa 25 mg			
Tab long-acting 200 mg with carbidopa 50 mg – 5% DV Feb-25 to 2027	44.99	100	Sinemet CR
Tab 250 mg with carbidopa 25 mg – 5% DV Feb-25 to 2027	39.49	100	Sinemet
LEVODOPA WITH CARBIDOPA AND ENTACAPONE			
Tab 50 mg with carbidopa 12.5 mg and entacapone 200 mg – 5% DV Jul-25 to 2027	27.01	100	Stalevo
Tab 100 mg with carbidopa 25 mg and entacapone 200 mg – 5% DV Jul-25 to 2027	34.18	100	Stalevo
Tab 150 mg with carbidopa 37.5 mg and entacapone 200 mg – 5% DV Jul-25 to 2027	44.96	100	Stalevo
Tab 200 mg with carbidopa 50 mg and entacapone 200 mg – 5% DV Jul-25 to 2027	51.23	100	Stalevo
PRAMIPEXOLE HYDROCHLORIDE			
Tab 0.25 mg – 5% DV Dec-25 to 2028	5.23	100	Ramipex
Tab 1 mg – 5% DV Dec-25 to 2028	17.73	100	Ramipex
RASAGILINE			
Tab 1 mg	53.50	30	Azilect
ROPINIROLE HYDROCHLORIDE			
Tab 0.25 mg	8.83	84	Ropin
Tab 1 mg	10.09	84	Ropin
Tab 2 mg	12.29	84	Ropin
Tab 5 mg	25.94	84	Ropin
SELEGILINE HYDROCHLORIDE – Restricted: For continuation only			
➡ Tab 5 mg			
TOLCAPONE			
Tab 100 mg	152.38	100	Tasmar

Anaesthetics

General Anaesthetics

DESFLURANE			
Soln for inhalation 100%, 240 ml bottle	1,350.00	6	Suprane
DEXMETETOMIDINE			
Inj 100 mcg per ml, 2 ml vial	42.00	5	Dexmedetomidine Viatris
ETOMIDATE			
Inj 2 mg per ml, 10 ml ampoule			
ISOFLURANE			
Soln for inhalation 100%, 250 ml bottle	2,730.00	6	Aerrane

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
KETAMINE			
Inj 1 mg per ml, 100 ml bag	146.00	5	Biomed
Inj 10 mg per ml, 10 ml syringe	76.00	5	Biomed
Inj 100 mg per ml, 2 ml vial – 5% DV Jan-27 to 2029	36.23	5	Ketalar
	91.98		Ketalar
METHOHEXITAL SODIUM			
Inj 10 mg per ml, 50 ml vial			
PROPOFOL			
Inj 10 mg per ml, 20 ml ampoule – 5% DV Feb-26 to 2028	5.75	5	Fresofol 1% MCT/LCT
Inj 10 mg per ml, 50 ml vial – 5% DV Feb-26 to 2028	27.50	10	Fresofol 1% MCT/LCT
Inj 10 mg per ml, 100 ml vial – 5% DV Feb-26 to 2028	39.90	10	Fresofol 1% MCT/LCT
SEVOFLURANE			
Soln for inhalation 100%, 250 ml bottle	930.00	6	Baxter
THIOPENTAL [THIOPENTONE] SODIUM			
Inj 500 mg ampoule			
Local Anaesthetics			
ARTICAINE HYDROCHLORIDE			
Inj 1%			
ARTICAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 4% with adrenaline 1:100,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:100,000, 1.8 ml dental cartridge			
Inj 4% with adrenaline 1:100,000, 2.2 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:200,000 1.8 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 2.2 ml dental cartridge			
BENZOCAINE			
Gel 20%			
BENZOCAINE WITH TETRACAINE HYDROCHLORIDE			
Gel 18% with tetracaine hydrochloride 2%			<i>e.g. ZAP Topical Anaesthetic Gel</i>
BUPIVACAINE HYDROCHLORIDE			
Inj 5 mg per ml, 4 ml ampoule	62.50	5	Marcaïn Isobaric
Inj 2.5 mg per ml, 20 ml ampoule			
Inj 2.5 mg per ml, 20 ml ampoule sterile pack	28.00	5	Marcaïn
Inj 5 mg per ml, 10 ml ampoule sterile pack	16.20	5	Marcaïn
Inj 5 mg per ml, 20 ml ampoule			
Inj 5 mg per ml, 20 ml ampoule sterile pack	16.56	5	Marcaïn
Inj 1.25 mg per ml, 100 ml bag			
Inj 1.25 mg per ml, 200 ml bag			
Inj 2.5 mg per ml, 200 ml bag			
Inj 1.25 mg per ml, 500 ml bag			
BUPIVACAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 2.5 mg per ml with adrenaline 1:200,000, 10 ml ampoule			
Inj 2.5 mg per ml with adrenaline 1:400,000, 20 ml vial – 5% DV Aug-26 to 2029	57.00	5	Marcaïn with Adrenaline
Inj 5 mg per ml with adrenaline 1:200,000, 20 ml vial – 5% DV Aug-26 to 2029	61.68	5	Marcaïn with Adrenaline

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BUPIVACAINE HYDROCHLORIDE WITH FENTANYL			
Inj 0.625 mg with fentanyl 2 mcg per ml, 100 ml bag			
Inj 0.625 mg with fentanyl 2 mcg per ml, 200 ml bag	165.00	5	Biomed
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml bag – 5% DV Feb-26 to 2028	135.00	5	Bupafen
Inj 1.25 mg with fentanyl 2 mcg per ml, 200 ml bag – 5% DV Feb-26 to 2028	138.00	5	Bupafen
Inj 1.25 mg with fentanyl 2 mcg per ml, 50 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 20 ml syringe	57.35	5	Biomed
BUPIVACAINE HYDROCHLORIDE WITH GLUCOSE			
Inj 0.5% with glucose 8%, 4 ml ampoule – 5% DV Dec-25 to 2028	21.40	5	Marcain Heavy
COCAINE HYDROCHLORIDE			
Paste 5%			
Soln 15%, 2 ml syringe			
Soln 4%, 2 ml syringe	30.77	1	Biomed
COCAINE HYDROCHLORIDE WITH ADRENALINE			
Paste 15% with adrenaline 0.06%			
Paste 25% with adrenaline 0.06%			
ETHYL CHLORIDE			
Spray 100%			
LIDOCAINE [LIGNOCAINE]			
Crm 4%	7.60	5 g	LMX4
	30.00	30 g	LMX4
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Gel 2%	4.87	20 g	Orion
Soln 4%			
Spray 10% – 5% DV Feb-26 to 2028	82.90	50 ml	Xylocaine
Oral (gel) soln 2% – 5% DV Apr-26 to 2028	30.80	200 ml	Xylocaine Viscous
Inj 1%, 20 ml ampoule, sterile pack			
Inj 2%, 20 ml ampoule, sterile pack			
Inj 1%, 5 ml ampoule	15.00	25	Lidocaine-Baxter
Inj 1%, 20 ml vial	19.50	5	Lidocaine-Baxter
Inj 2%, 5 ml ampoule	27.50	25	Lidocaine-Baxter
Inj 2%, 20 ml vial	14.00	5	Lidocaine-Baxter
Inj 10%, 5 ml ampoule			
Gel 2%, 11 ml urethral syringe – 5% DV Feb-26 to 2028	65.45	10	Instillagel Lido
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE			
Inj 1% with adrenaline 1:100,000, 20 ml vial			
Inj 1% with adrenaline 1:100,000, 5 ml ampoule – 5% DV Dec-25 to 2028	32.00	10	Xylocaine
Inj 1% with adrenaline 1:200,000, 20 ml vial	50.00	5	Xylocaine
Inj 2% with adrenaline 1:100,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 2.2 ml dental cartridge			
Inj 2% with adrenaline 1:200,000, 20 ml vial	60.00	5	Xylocaine
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE AND TETRACAINE HYDROCHLORIDE			
Soln 4% with adrenaline 0.1% and tetracaine hydrochloride 0.5%, 5 ml syringe	20.50	1	Topicaïne

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH PHENYLEPHRINE HYDROCHLORIDE			
Nasal spray 5% with phenylephrine hydrochloride 0.5%			
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE			
Crm 2.5% with prilocaine 2.5%.....	45.00	30 g	EMLA
Patch 25 mcg with prilocaine 25 mcg.....	115.00	20	EMLA
Crm 2.5% with prilocaine 2.5%, 5 g.....	45.00	5	EMLA
MEPIVACAINE HYDROCHLORIDE			
Inj 3%, 1.8 ml dental cartridge.....	43.60	50	Scandonest 3%
Inj 3%, 2.2 ml dental cartridge.....	43.60	50	Scandonest 3%
MEPIVACAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 2% with adrenaline 1:100,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:100,000, 2.2 ml dental cartridge			
PRILOCAINE HYDROCHLORIDE			
Inj 0.5%, 50 ml vial.....	100.00	5	Citanest
Inj 2%, 5 ml ampoule			
PRILOCAINE HYDROCHLORIDE WITH FELYPRESSIN			
Inj 3% with felypressin 0.03 iu per ml, 1.8 ml dental cartridge			
Inj 3% with felypressin 0.03 iu per ml, 2.2 ml dental cartridge			
ROPIVACAINE HYDROCHLORIDE			
Inj 2 mg per ml, 10 ml ampoule.....	9.80	5	Ropivacaine Kabi
Inj 2 mg per ml, 20 ml ampoule.....	10.25	5	Ropivacaine Kabi
Inj 2 mg per ml, 100 ml bag.....	32.85	5	Ropivacaine Kabi
Inj 2 mg per ml, 200 ml bag.....	43.40	5	Ropivacaine Kabi
Inj 7.5 mg per ml, 10 ml ampoule.....	11.00	5	Ropivacaine Kabi
Inj 7.5 mg per ml, 20 ml ampoule.....	13.50	5	Ropivacaine Kabi
Inj 10 mg per ml, 10 ml ampoule.....	11.75	5	Ropivacaine Kabi
Inj 10 mg per ml, 20 ml ampoule.....	17.60	5	Ropivacaine Kabi
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Gel 4%			

Analgesics

Non-Opioid Analgesics

ASPIRIN			
Tab dispersible 300 mg.....	5.65	100	Ethics Aspirin
CAPSAICIN – Restricted see terms below			
↓ Crm 0.075%.....	11.95	45 g	Zo-Rub HP Zostrix HP
→ Restricted (RS1145)			
Initiation			
For post-herpetic neuralgia or diabetic peripheral neuropathy.			
METHOXYFLURANE – Restricted see terms below			
↓ Soln for inhalation 99.9 mg per g, 3 ml bottle.....	276.00	10	Penthrox
↓ Soln for inhalation 999.9 mg per g, 3 ml bottle with inhaler device and activated carbon chamber.....	54.00	1	Penthrox
→ Restricted (RS1292)			
Initiation			
Both:			

continued...

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient is undergoing a painful procedure with an expected duration of less than one hour; and
- 2 Only to be used under supervision by a medical practitioner or nurse who is trained in the use of methoxyflurane.

NEFOPAM HYDROCHLORIDE

Tab 30 mg

PARACETAMOL – **Some items restricted** see terms [below](#)

Tab soluble 500 mg

Tab 500 mg - blister pack - 1,000 tablet pack – **5% DV Dec-26 to 2029** 12.06 1,000 **Pacimol**

Tab 500 mg - blister pack - 12 tablet pack

Tab 500 mg - blister pack - 20 tablet pack

Tab 500 mg - bottle pack..... 17.92 1,000 Noured Paracetamol

Oral liq 120 mg per 5 ml – **5% DV Nov-26 to 2028**..... 3.18 200 ml **Noured**

Oral liq 250 mg per 5 ml – **5% DV Jul-26 to 2028** 3.98 200 ml Paracetamol (Ethics)

↓ Inj 10 mg per ml, 100 ml vial – **5% DV Feb-26 to 2028** 15.30 10 **Pamol**

Suppos 25 mg

Suppos 50 mg

Suppos 125 mg 4.29 10 Gacet

Suppos 250 mg 5.39 10 Gacet

Suppos 500 mg 16.55 50 Gacet

(Paracetamol (Ethics) Oral liq 120 mg per 5 ml to be delisted 1 November 2026)

➔ **Restricted (RS1146)**

Initiation

Intravenous paracetamol is only to be used where other routes are unavailable or impractical, or where there is reduced absorption. The need for IV paracetamol must be re-assessed every 24 hours.

SUCROSE

Oral liq 25% 14.61 25 ml Biomed

↓ Oral liq 66.7% (preservative free)

➔ **Restricted (RS1763)**

Initiation

For use in neonatal patients only.

Opioid Analgesics

ALFENTANIL

Inj 0.5 mg per ml, 2 ml ampoule – **5% DV Dec-26 to 2029** 6.99 5 **Medsurge**

CODEINE PHOSPHATE

Tab 15 mg – **5% DV Dec-25 to 2028** 5.82 100 **Noured**

Tab 30 mg – **5% DV Dec-25 to 2028** 6.88 100 **Noured**

Tab 60 mg – **5% DV Dec-25 to 2028** 13.89 100 **Noured**

DIHYDROCODEINE TARTRATE

Tab long-acting 60 mg – **5% DV Feb-26 to 2028** 9.20 60 **DHC Continus**

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FENTANYL			
Inj 10 mcg per ml, 10 ml syringe – 5% DV Feb-25 to 2027	44.50	5	Biomed Fentanyl
Inj 50 mcg per ml, 2 ml ampoule – 5% DV May-25 to 2027	4.25	10	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule – 5% DV May-25 to 2027	9.41	10	Boucher and Muir
Inj 10 mcg per ml, 100 ml bag	114.25	5	Biomed
Inj 20 mcg per ml, 50 ml syringe – 5% DV Feb-25 to 2027	136.50	5	Biomed
Inj 20 mcg per ml, 100 ml bag			
Patch 12 mcg per hour – 5% DV May-25 to 2027	6.02	5	Fentanyl Sandoz
Patch 25 mcg per hour – 5% DV Dec-24 to 2027	6.91	5	Fentanyl Sandoz
Patch 50 mcg per hour – 5% DV Dec-24 to 2027	9.28	5	Fentanyl Sandoz
Patch 75 mcg per hour – 5% DV Dec-24 to 2027	15.50	5	Fentanyl Sandoz
Patch 100 mcg per hour – 5% DV Dec-24 to 2027	16.37	5	Fentanyl Sandoz
METHADONE HYDROCHLORIDE			
Tab 5 mg – 5% DV Sep-26 to 2028	1.38	10	Methadone BNM
Oral liq 2 mg per ml – 5% DV Feb-25 to 2027	7.80	200 ml	Biodone
Oral liq 5 mg per ml – 5% DV Feb-25 to 2027	7.80	200 ml	Biodone Forte
Oral liq 10 mg per ml – 5% DV Feb-25 to 2027	9.65	200 ml	Biodone Extra Forte
Inj 10 mg per ml, 1 ml vial	72.99	10	AFT
MORPHINE HYDROCHLORIDE			
Oral liq 1 mg per ml	19.00	200 ml	RA-Morph
Oral liq 2 mg per ml	23.55	200 ml	RA-Morph
Oral liq 5 mg per ml	28.20	200 ml	RA-Morph
Oral liq 10 mg per ml	40.25	200 ml	RA-Morph
MORPHINE SULPHATE			
Tab immediate-release 10 mg	2.80	10	Sevredol
Tab immediate-release 20 mg	5.52	10	Sevredol
Cap long-acting 10 mg – 5% DV Jul-26 to 2028	4.10	10	m-Eslon
Cap long-acting 30 mg – 5% DV Jul-26 to 2028	6.05	10	m-Eslon
Cap long-acting 60 mg – 5% DV Jul-26 to 2028	12.10	10	m-Eslon
Cap long-acting 100 mg – 5% DV Jul-26 to 2028	14.50	10	m-Eslon
Oral liq 2 mg per ml	42.56	300 ml	Oramorph
	29.80	100 ml	Oramorph CDC S29
	16.31		Wockhardt
Inj 1 mg per ml, 100 ml bag	114.25	5	Biomed
Inj 1 mg per ml, 10 ml syringe	27.25	5	Biomed
Inj 1 mg per ml, 50 ml syringe	63.75	5	Biomed
Inj 1 mg per ml, 2 ml syringe			
Inj 5 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	5.96	5	Medsurge
Inj 10 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	4.99	5	Medsurge
Inj 10 mg per ml, 100 mg cassette			
Inj 10 mg per ml, 100 ml bag			
Inj 15 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	6.93	5	Medsurge
Inj 30 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	7.28	5	Medsurge
Inj 200 mcg in 0.4 ml syringe			
Inj 300 mcg in 0.3 ml syringe			
MORPHINE TARTRATE			
Inj 80 mg per ml, 1.5 ml ampoule			

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
OXYCODONE HYDROCHLORIDE			
Tab controlled-release 5 mg – 5% DV Dec-24 to 2027	2.49	20	Oxycodone Sandoz
Tab immediate-release 5 mg.....	13.77	100	Oxycodone Amneal
Tab controlled-release 10 mg – 5% DV Dec-24 to 2027	2.49	20	Oxycodone Sandoz
Tab immediate-release 10 mg.....	18.77	100	Oxycodone Amneal
Tab controlled-release 20 mg – 5% DV Dec-24 to 2027	3.41	20	Oxycodone Sandoz
Tab immediate-release 20 mg.....	26.77	100	Oxycodone Amneal
Tab controlled-release 40 mg – 5% DV Dec-24 to 2027	6.67	20	Oxycodone Sandoz
Tab controlled-release 80 mg – 5% DV Dec-24 to 2027	12.99	20	Oxycodone Sandoz
Oral liq 1 mg per ml	37.08	250 ml	Rosemont
Inj 1 mg per ml, 100 ml bag			
Inj 10 mg per ml, 1 ml ampoule – 5% DV Dec-24 to 2027	4.37	5	Hameln
Inj 10 mg per ml, 2 ml ampoule – 5% DV Dec-24 to 2027	8.62	5	Hameln
Inj 50 mg per ml, 1 ml ampoule – 5% DV Dec-24 to 2027	14.90	5	Hameln
PARACETAMOL WITH CODEINE			
Tab paracetamol 500 mg with codeine phosphate 8 mg – 5% DV Feb-26 to 2028.....	31.95	1,000	Paracetamol + Codeine (Relieve)
PETHIDINE HYDROCHLORIDE			
Tab 50 mg – 5% DV Feb-26 to 2028	8.68	10	Noumed Pethidine
Inj 5 mg per ml, 10 ml syringe			
Inj 5 mg per ml, 100 ml bag			
Inj 10 mg per ml, 100 ml bag			
Inj 10 mg per ml, 50 ml syringe			
Inj 50 mg per ml, 1 ml ampoule	29.88	5	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule	30.72	5	DBL Pethidine Hydrochloride
REMIFENTANIL			
Inj 1 mg vial	14.95	5	Remifentanil-AFT
Inj 2 mg vial	20.95	5	Remifentanil-AFT
TRAMADOL HYDROCHLORIDE			
Tab sustained-release 100 mg.....	1.95	20	Tramal SR 100
Tab sustained-release 150 mg.....	2.95	20	Tramal SR 150
Tab sustained-release 200 mg.....	3.80	20	Tramal SR 200
Cap 50 mg	3.33	100	Arrow-Tramadol
Oral soln 10 mg per ml			
Inj 10 mg per ml, 100 ml bag			
Inj 50 mg per ml, 1 ml ampoule	10.00	5	Tramal 50
Inj 50 mg per ml, 2 ml ampoule	9.00	5	Tramal 100

Antidepressants

Cyclic and Related Agents

AMITRIPTYLINE			
Tab 10 mg	2.99	100	Arrow-Amitriptyline
Tab 25 mg	1.99	100	Arrow-Amitriptyline
Tab 50 mg	3.14	100	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE			
Tab 25 mg – 5% DV Jul-25 to 2027	16.99	50	APO Clomipramine

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Restricted: For continuation only			
➔ Tab 75 mg	3.85	30	Dosulepin Viatris
➔ Cap 25 mg	7.83	50	Dosulepin Viatris
DOXEPIN HYDROCHLORIDE – Restricted: For continuation only			
➔ Cap 10 mg			
➔ Cap 25 mg			
➔ Cap 50 mg			
IMIPRAMINE HYDROCHLORIDE			
Tab 10 mg	5.48	50	Tofranil
	6.58	60	Tofranil
Tab 25 mg	4.93	28	Imipramine Crescent
	8.80	50	Tofranil
MAPROTILINE HYDROCHLORIDE – Restricted: For continuation only			
➔ Tab 25 mg			
➔ Tab 75 mg			
MIANSERIN HYDROCHLORIDE – Restricted: For continuation only			
➔ Tab 30 mg			
NORTRIPTYLINE HYDROCHLORIDE			
Tab 10 mg	2.24	50	Allegron
Tab 25 mg	2.95	50	Allegron
Monoamine-Oxidase Inhibitors - Non-Selective			
PHENELZINE SULPHATE			
Tab 15 mg			
TRANLYCYPROMINE SULPHATE			
Tab 10 mg			
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE			
Tab 150 mg – 5% DV Feb-25 to 2027	23.60	60	Aurorix
Tab 300 mg – 5% DV Feb-25 to 2027	38.50	60	Aurorix
Other Antidepressants			
MIRTAZAPINE			
Tab 30 mg – 5% DV Jan-26 to 2028	2.34	30	Noumed
Tab 45 mg – 5% DV Jan-26 to 2028	3.10	30	Noumed
VENLAFAXINE			
Cap 37.5 mg	8.29	84	Enlafax XR
Cap 75 mg	10.32	84	Enlafax XR
Cap 150 mg	13.95	84	Enlafax XR
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE			
Tab 20 mg – 5% DV May-26 to 2028	3.55	84	Celapram
ESCITALOPRAM			
Tab 10 mg	0.79	28	Ipca-Escitalopram
Tab 20 mg	1.49	28	Ipca-Escitalopram

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUOXETINE HYDROCHLORIDE			
Tab dispersible 20 mg, scored – 5% DV Mar-26 to 2028	2.37	28	Fluox
Cap 20 mg – 5% DV Mar-26 to 2028	3.50	90	Arrow-Fluoxetine
PAROXETINE			
Tab 20 mg – 5% DV Feb-26 to 2028	4.98	90	Loxamine
SERTRALINE			
Tab 50 mg – 5% DV Apr-26 to 2028	1.24	30	Setrona
Tab 100 mg – 5% DV Apr-26 to 2028	2.00	30	Setrona

Antiepilepsy Drugs

Agents for the Control of Status Epilepticus

CLONAZEPAM			
Inj 1 mg per ml, 1 ml ampoule			
DIAZEPAM			
Inj 5 mg per ml, 2 ml ampoule	27.92	5	Hospira
Rectal tubes 5 mg.....	54.58	5	Stesolid
Rectal tubes 10 mg			
LORAZEPAM			
Inj 2 mg vial			
Inj 4 mg per ml, 1 ml vial			
PARALDEHYDE			
Soln 97%			
Inj 5 ml ampoule			
PHENYTOIN SODIUM			
Inj 50 mg per ml, 5 ml ampoule	154.01	5	Hospira

Control of Epilepsy

CARBAMAZEPINE			
Tab 200 mg	14.53	100	Tegretol
Tab long-acting 200 mg.....	16.98	100	Tegretol CR
Tab 400 mg	34.58	100	Tegretol
Tab long-acting 400 mg.....	39.17	100	Tegretol CR
Oral liq 20 mg per ml	26.37	250 ml	Tegretol
CLOBAZAM			
Tab 10 mg			
CLONAZEPAM			
Oral drops 2.5 mg per ml			
ETHOSUXIMIDE			
Cap 250 mg.....	140.88	100	Zarontin
Oral liq 50 mg per ml	56.35	200 ml	Zarontin
GABAPENTIN			
Note: Gabapentin not to be given in combination with pregabalin			
Cap 100 mg – 1% DV Feb-22 to 2027	6.45	100	Nupentin
Cap 300 mg – 1% DV Feb-22 to 2027	8.45	100	Nupentin
Cap 400 mg – 1% DV Feb-22 to 2027	10.26	100	Nupentin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LACOSAMIDE – Restricted see terms below			
↓ Tab 50 mg	25.04	14	Vimpat
↓ Tab 100 mg	50.06	14	Vimpat
	200.24	56	Vimpat
↓ Tab 150 mg	75.10	14	Vimpat
	300.40	56	Vimpat
↓ Tab 200 mg	400.55	56	Vimpat
↓ Inj 10 mg per ml, 20 ml vial			
→ Restricted (RS1988)			
Initiation			
<i>Re-assessment required after 15 months</i>			
Both:			
1 Patient has focal epilepsy; and			
2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam, and any two of carbamazepine, lamotrigine, and phenytoin sodium (see Note).			
Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.			
Continuation			
Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.			
LAMOTRIGINE			
Tab dispersible 2 mg	55.00	30	Lamictal
Tab dispersible 5 mg	50.00	30	Lamictal
Tab dispersible 25 mg	4.20	56	Logem
Tab dispersible 50 mg	5.11	56	Logem
Tab dispersible 100 mg	6.75	56	Logem
LEVETIRACETAM			
Tab 250 mg	5.84	60	Everet
Tab 500 mg	10.51	60	Everet
Tab 750 mg	16.71	60	Everet
Tab 1,000 mg	21.82	60	Everet
Oral liq 100 mg per ml	44.78	300 ml	Levetiracetam-AFT
Inj 100 mg per ml, 5 ml vial	38.95	10	Levetiracetam-AFT
PHENOBARBITONE			
Tab 15 mg	248.50	500	Noumed Phenobarbitone
Tab 30 mg	398.50	500	Noumed Phenobarbitone
PHENYTOIN			
Tab 50 mg			
PHENYTOIN SODIUM			
Cap 30 mg			
Cap 100 mg			
Oral liq 6 mg per ml			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PREGABALIN			
Note: Pregabalin not to be given in combination with gabapentin			
Cap 25 mg	2.25	56	Lyrica Pregabalin Pfizer
Cap 75 mg	2.65	56	Lyrica Pregabalin Pfizer
Cap 150 mg	4.01	56	Lyrica Pregabalin Pfizer
Cap 300 mg	7.38	56	Lyrica Pregabalin Pfizer
PRIMIDONE			
Tab 250 mg			
SODIUM VALPROATE			
Tab 100 mg			
Tab EC 200 mg			
Tab EC 500 mg			
Oral liq 40 mg per ml			
Inj 100 mg per ml, 4 ml vial	9.98	1	Epilim IV
STIRIPENTOL – Restricted see terms below			
⚡ Cap 250 mg	509.29	60	Diacomit
⚡ Powder for oral liq 250 mg sachet	509.29	60	Diacomit
➔ Restricted (RS1989)			
Initiation			
Paediatric neurologist			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Patient has confirmed diagnosis of Dravet syndrome; and			
2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.			
Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.			
Continuation			
Paediatric neurologist			
Patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.			
TOPIRAMATE			
Tab 25 mg	11.07	60	Arrow-Topiramate
	26.04		Topamax
	11.07		Topiramate Actavis
Tab 50 mg	18.81	60	Arrow-Topiramate
	44.26		Topamax
	18.81		Topiramate Actavis
Tab 100 mg	31.99	60	Arrow-Topiramate
	75.25		Topamax
	31.99		Topiramate Actavis
Tab 200 mg	55.19	60	Arrow-Topiramate
	129.85		Topamax
	55.19		Topiramate Actavis
Cap sprinkle 15 mg	20.84	60	Topamax
Cap sprinkle 25 mg	26.04	60	Topamax
VIGABATRIN – Restricted see terms on the next page			
⚡ Tab 500 mg			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

➔ Restricted (RS1865)

Initiation

Re-assessment required after 15 months

Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
 - 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Continuation

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Antimigraine Preparations

Acute Migraine Treatment

DIHYDROERGOTAMINE MESYLATE

Inj 1 mg per ml, 1 ml ampoule

METOCLOPRAMIDE HYDROCHLORIDE WITH PARACETAMOL

Tab 5 mg with paracetamol 500 mg

RIZATRIPTAN

Tab orodispersible 10 mg – 5% DV Dec-26 to 2029.....	4.10	30	Rizamelt
	2.46	18	Rizatriptan Rising
	4.10	30	Rizatriptan Viatris

SUMATRIPTAN

Tab 50 mg – 1% DV Feb-22 to 2027	14.41	90	Sumagran
Tab 100 mg – 1% DV Feb-22 to 2027	22.68	90	Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen – 5% DV Dec-25 to 2028	29.80	2	Clustran

Prophylaxis of Migraine

PIZOTIFEN

Tab 500 mcg.....	23.21	100	Sandomigran
------------------	-------	-----	-------------

Antinausea and Vertigo Agents

APREPITANT – **Restricted** see terms [on the next page](#)

⚡ Cap 2 x 80 mg and 1 x 125 mg – 5% DV Jan-25 to 2027	21.90	3	Emend Tri-Pack
---	-------	---	-----------------------

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted (RS1154)			
Initiation			
Patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.			
BETAHISTINE DIHYDROCHLORIDE			
Tab 16 mg	3.70	100	Serc
CYCLIZINE HYDROCHLORIDE			
Tab 50 mg – 5% DV Feb-25 to 2027	0.66	10	Nausicalm
CYCLIZINE LACTATE			
Inj 50 mg per ml, 1 ml ampoule	17.86	10	Hameln
DOMPERIDONE			
Tab 10 mg – 5% DV Dec-25 to 2028	3.80	100	Domperidone Viatris
DROPERIDOL			
Inj 2.5 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	28.68	10	Droperidol Medsurge
	43.85		Droperidol Panpharma
GRANISETRON			
Inj 1 mg per ml, 3 ml ampoule – 5% DV Jan-27 to 2029	1.20	1	Deva
	1.71		Granisetron-AFT
<i>(Deva Inj 1 mg per ml, 3 ml ampoule to be delisted 1 January 2027)</i>			
HYOSCINE HYDROBROMIDE			
Inj 400 mcg per ml, 1 ml ampoule			
⚡ Patch 1 mg per 72 hours	88.50	10	Scopolamine Transdermal System Viatris
➔ Restricted (RS1155)			
Initiation			
Any of the following:			
1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or			
2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective; or			
3 For treatment of post-operative nausea and vomiting where cyclizine, droperidol and a 5HT3 antagonist have proven ineffective, are not tolerated or are contraindicated.			
METOCLOPRAMIDE HYDROCHLORIDE			
Tab 10 mg	1.57	100	Metoclopramide Actavis
Oral liq 5 mg per 5 ml			10
Inj 5 mg per ml, 2 ml ampoule – 5% DV Apr-26 to 2028	5.48	10	Medsurge
ONDANSETRON			
Tab 4 mg – 5% DV Dec-25 to 2028	1.95	50	Periset
Tab dispersible 4 mg	0.56	10	Periset ODT
Tab 8 mg – 5% DV Dec-25 to 2028	3.50	50	Periset
Tab dispersible 8 mg	0.90	10	Periset ODT
Inj 2 mg per ml, 2 ml ampoule – 5% DV Feb-26 to 2028	1.54	5	Ondansetron-AFT
Inj 2 mg per ml, 4 ml ampoule – 5% DV Feb-26 to 2028	2.14	5	Ondansetron-AFT

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PROCHLORPERAZINE			
Tab buccal 3 mg			
Tab 5 mg	25.00	250	Nausafix
Inj 12.5 mg per ml, 1 ml ampoule			
Suppos 25 mg			
TROPISETRON			
Inj 1 mg per ml, 2 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			
Antipsychotic Agents			
General			
AMISULPRIDE			
Tab 100 mg – 5% DV Dec-24 to 2027	5.84	30	Sulprix
Tab 200 mg – 5% DV Dec-24 to 2027	14.47	60	Sulprix
Tab 400 mg – 5% DV Dec-24 to 2027	35.06	60	Sulprix
Oral liq 100 mg per ml			
ARIPIRAZOLE			
Tab 5 mg	10.50	30	Aripiprazole Sandoz
Tab 10 mg	10.50	30	Aripiprazole Sandoz
Tab 15 mg	10.50	30	Aripiprazole Sandoz
Tab 20 mg	10.50	30	Aripiprazole Sandoz
Tab 30 mg	10.50	30	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE			
Tab 25 mg	15.62	100	Largactil
Tab 100 mg	36.73	100	Largactil
Oral liq 10 mg per ml			
Oral liq 20 mg per ml			
Inj 25 mg per ml, 2 ml ampoule	30.79	10	Largactil
CLOZAPINE			
Tab 25 mg	6.69	50	Clopine
	13.37	100	Clopine
	6.69	50	Clozaril
	13.37	100	Clozaril
Tab 50 mg	8.67	50	Clopine
	17.33	100	Clopine
Tab 100 mg	17.33	50	Clopine
	34.65	100	Clopine
	17.33	50	Clozaril
	34.65	100	Clozaril
Tab 200 mg	34.65	50	Clopine
	69.30	100	Clopine
Oral liq 50 mg per ml	173.30	100 ml	Versacloz
HALOPERIDOL			
Tab 500 mcg.....	6.23	100	Serenace
Tab 1.5 mg	9.43	100	Serenace
Tab 5 mg	29.72	100	Serenace
Oral liq 2 mg per ml	23.84	100 ml	Serenace
Inj 5 mg per ml, 1 ml ampoule – 5% DV Aug-26 to 2029	12.93	10	Serenace

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LEVOMEPROMAZINE			
Tab 25 mg	16.10	100	Nozinan
Tab 100 mg	41.75	100	Nozinan
LEVOMEPROMAZINE HYDROCHLORIDE			
Inj 25 mg per ml, 1 ml ampoule – 5% DV Dec-25 to 2028	23.26	10	Wockhardt
LITHIUM CARBONATE			
Tab long-acting 400 mg – 5% DV Feb-25 to 2027	82.80	100	Priadel
Cap 250 mg	35.78	100	Douglas
OLANZAPINE			
Tab 2.5 mg	1.40	30	Zypine
Tab 5 mg	1.93	30	Zypine
Tab orodispersible 5 mg	3.93	28	APO-Olanzapine
	8.64	100	Olanzapine ODT Viatris
	2.42	28	Zypine ODT
Tab 10 mg	1.80	28	Zypine
	1.93	30	Zypine
Tab orodispersible 10 mg	4.49	28	APO-Olanzapine
	2.89		Olanzapina Mylan
			Olanzapina Mylan
			Pharma
	10.32	100	Olanzapine ODT Viatris
	2.89	28	Zypine ODT
Inj 10 mg vial			
PERICAZINE			
Tab 2.5 mg			
Tab 10 mg			
QUETIAPINE			
Tab 25 mg – 5% DV Dec-26 to 2029	2.14	90	Quetapel
	0.79	30	Quetiapine Viatris
	13.11	500	Quetiapine Viatris
Tab 100 mg – 5% DV Dec-26 to 2029	5.91	90	Quetapel
Tab 200 mg – 5% DV Dec-26 to 2029	9.93	90	Quetapel
Tab 300 mg – 5% DV Dec-26 to 2029	14.48	90	Quetapel
RISPERIDONE			
Tab 0.5 mg	2.17	60	Risperidone (Teva)
	4.01		Risperidone Sandoz
Tab 1 mg	2.44	60	Risperidone (Teva)
Tab 2 mg	2.72	60	Risperidone (Teva)
	5.38		Risperidone Sandoz
Tab 3 mg	4.50	60	Risperidone (Teva)
Tab 4 mg	6.25	60	Risperdal
			Risperidone (Teva)
Oral liq 1 mg per ml	10.29	30 ml	Risperon
	34.30	100 ml	Risperon
ZIPRASIDONE			
Cap 20 mg	17.90	60	Zusdone
Cap 40 mg	27.41	60	Zusdone
Cap 60 mg	38.39	60	Zusdone
Cap 80 mg	46.55	60	Zusdone

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ZUCLOPENTHIXOL ACETATE			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
ZUCLOPENTHIXOL HYDROCHLORIDE			
Tab 10 mg	31.45	100	Clopixol

Depot Injections

ARIPIPIRAZOLE – **Restricted** see terms [below](#)

↓ Inj 300 mg vial	273.56	1	Abilify Maintena
↓ Inj 400 mg vial	341.96	1	Abilify Maintena

→ **Restricted (RS2058)**

Initiation

Either:

1 Either:

- 1.1 The patient has had an initial Special Authority approval for risperidone depot injection or paliperidone depot injection or olanzapine depot injection; or
- 1.2 All of the following:
 - 1.2.1 The patient has schizophrenia or other psychotic disorder; and
 - 1.2.2 The patient has received treatment with oral atypical antipsychotic agents but has been unable to adhere; and
 - 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or

2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been initiated on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection. (see Note below for the olanzapine Special Authority criteria for new olanzapine depot injection patients prior to 1 April 2024).

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
 - The patient has schizophrenia; and
 - The patient has tried but has not been able to adhere with treatment using oral atypical antipsychotic agents; and
 - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

FLUPENTHIXOL DECANOATE

Inj 20 mg per ml, 1 ml ampoule	13.14	5	Fluanxol
Inj 20 mg per ml, 2 ml ampoule	20.90	5	Fluanxol
Inj 100 mg per ml, 1 ml ampoule	40.87	5	Fluanxol

HALOPERIDOL DECANOATE

Inj 50 mg per ml, 1 ml ampoule	28.39	5	Haldol
Inj 100 mg per ml, 1 ml ampoule	55.90	5	Haldol Concentrate

OLANZAPINE – **Restricted** see terms [below](#)

↓ Inj 210 mg vial	252.00	1	Zyprexa Relprevv
↓ Inj 300 mg vial	414.00	1	Zyprexa Relprevv
↓ Inj 405 mg vial	504.00	1	Zyprexa Relprevv

→ **Restricted (RS2165)**

Initiation

Re-assessment required after 12 months

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

1 Either:

- 1.1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 1.2 All of the following:
 - 1.2.1 The patient has schizophrenia or other psychotic disorder; and
 - 1.2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months; and
- 2 The patient has trialled other funded depot antipsychotics (aripiprazole, risperidone, and paliperidone) unless it is considered clinically inappropriate to use these; and
- 3 The patient continues to have difficulties with adherence on oral antipsychotic treatments; and
- 4 Prescribing clinician has relevant Clinical Director (Mental Health and Addiction services) approval.

Continuation

Re-assessment required after 12 months

The initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE – **Restricted** see terms [below](#)

‡ Inj 25 mg syringe	194.25	1	Invega Sustenna
‡ Inj 50 mg syringe	271.95	1	Invega Sustenna
‡ Inj 75 mg syringe	357.42	1	Invega Sustenna
‡ Inj 100 mg syringe	435.12	1	Invega Sustenna
‡ Inj 150 mg syringe	435.12	1	Invega Sustenna

→ **Restricted (RS2059)**

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 The patient has been unable to adhere to treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE – **Restricted** see terms [below](#)

‡ Inj 175 mg syringe	815.85	1	Invega Trinza
‡ Inj 263 mg syringe	1,072.26	1	Invega Trinza
‡ Inj 350 mg syringe	1,305.36	1	Invega Trinza
‡ Inj 525 mg syringe	1,305.36	1	Invega Trinza

→ **Restricted (RS1932)**

Initiation

Re-assessment required after 12 months

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Continuation

Re-assessment required after 12 months

The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PIPOTHIAZINE PALMITATE – **Restricted**: For continuation only

- ➔ Inj 50 mg per ml, 1 ml ampoule
- ➔ Inj 50 mg per ml, 2 ml ampoule

RISPERIDONE – **Restricted** see terms [below](#)

↓ Inj 25 mg vial	135.98	1	Risperdal Consta
↓ Inj 37.5 mg vial	178.71	1	Risperdal Consta
↓ Inj 50 mg vial	217.56	1	Risperdal Consta

➔ **Restricted (RS2060)**

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 The patient has not been able to adhere to treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE

Inj 200 mg per ml, 1 ml ampoule	19.80	5	Clopixol
Inj 500 mg per ml, 1 ml ampoule			e.g. Clopixol Conc

Anxiolytics

BUSPIRONE HYDROCHLORIDE

Tab 5 mg – 5% DV Dec-24 to 2027	13.95	100	Buspirone Viartis
Tab 10 mg – 5% DV Dec-24 to 2027	12.50	100	Buspirone Viartis

CLONAZEPAM

Tab 500 mcg	5.64	100	Paxam
Tab 2 mg	10.78	100	Paxam

DIAZEPAM

Tab 2 mg	95.00	500	Arrow-Diazepam
Tab 5 mg	115.00	500	Arrow-Diazepam

↓ Oral liq 10 mg per 10 ml

➔ **Restricted (RS2054)**

Initiation

Relevant specialist

Only for use in children where diazepam tablets are not appropriate.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LORAZEPAM			
Tab 1 mg – 5% DV Feb-25 to 2027	10.20	250	Ativan
Tab 2.5 mg – 5% DV Feb-25 to 2027	13.13	100	Ativan
OXAZEPAM			
Tab 10 mg			
Tab 15 mg			

Multiple Sclerosis Treatments

➔ **Restricted** (RS1993)

Initiation – Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide

Any relevant practitioner

Re-assessment required after 12 months

Either:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 – 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
 - 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever ($T > 37.5^{\circ}\text{C}$); and
 - 1.4.5 Either:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
 - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
 - 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory activity is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active approval for ocrelizumab and does not have primary progressive MS.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Continuation – Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

beta-1-beta, natalizumab and teriflunomide

Any relevant practitioner

Patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE – **Restricted** see terms [on the previous page](#)

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

† Cap 120 mg	520.00	14	Tecfidera
† Cap 240 mg	2,000.00	56	Tecfidera

FINGOLIMOD – **Restricted** see terms [on the previous page](#)

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

† Cap 0.5 mg	2,200.00	28	Gilenya
--------------	----------	----	---------

GLATRAMER ACETATE – **Restricted** see terms [on the previous page](#)

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

† Inj 40 mg prefilled syringe	1,500.00	12	Copaxone
-------------------------------	----------	----	----------

INTERFERON BETA-1-ALPHA – **Restricted** see terms [on the previous page](#)

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

† Inj 6 million iu in 0.5 ml syringe	1,170.00	4	Avonex
--------------------------------------	----------	---	--------

INTERFERON BETA-1-BETA – **Restricted** see terms [on the previous page](#)

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

† Inj 8 million iu per ml, 1 ml vial			
--------------------------------------	--	--	--

NATALIZUMAB – **Restricted** see terms [on the previous page](#)

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

† Inj 20 mg per ml, 15 ml vial	1,750.00	1	Tysabri
--------------------------------	----------	---	---------

TERIFLUNOMIDE – **Restricted** see terms [on the previous page](#)

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

† Tab 14 mg	282.81	30	Teriflunomide Aurobindo
	263.96	28	Teriflunomide Sandoz

Multiple Sclerosis Treatments - Other

OCRELIZUMAB – **Restricted** see terms [below](#)

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

† Inj 30 mg per ml, 10 ml vial	8,450.00	1	Ocrevus
† Inj 40 mg per ml, 23 ml vial	16,900.00	1	Ocrevus SC

➔ **Restricted (RS1997)**

Initiation – Multiple Sclerosis - ocrelizumab

Any relevant practitioner

Re-assessment required after 12 months

Either:

1 All of the following:

- 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 1.2 Patient has an EDSS score between 0 – 6.0; and
- 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever ($T > 37.5^{\circ}\text{C}$); and
 - 1.4.5 Either:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Continuation – Multiple Sclerosis - ocrelizumab

Any relevant practitioner

Patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initiation – Primary Progressive Multiple Sclerosis

Any relevant practitioner

Re-assessment required after 12 months

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and
- 3 Patient has no history of relapsing remitting multiple sclerosis.

Continuation – Primary Progressive Multiple Sclerosis

Any relevant practitioner

Patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

Sedatives and Hypnotics

CHLORAL HYDRATE

Oral liq 100 mg per ml

Oral liq 200 mg per ml

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LORMETAZEPAM – Restricted: For continuation only			
➔ Tab 1 mg			
MELATONIN – Restricted see terms below			
↓ Tab modified-release 2 mg – 5% DV Dec-24 to 2027	5.80	30	Vigisom
↓ Tab 3 mg			
Note: Only for use in compounding an oral liquid formulation, for in-hospital use only.			
➔ Restricted (RS1576)			
Initiation – insomnia secondary to neurodevelopmental disorder			
Psychiatrist, paediatrician, neurologist or respiratory specialist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder); and			
2 Behavioural and environmental approaches have been tried or are inappropriate; and			
3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and			
4 Patient is aged 18 years or under.			
Continuation – insomnia secondary to neurodevelopmental disorder			
Psychiatrist, paediatrician, neurologist or respiratory specialist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Patient is aged 18 years or under; and			
2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and			
3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and			
4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.			
Initiation – insomnia where benzodiazepines and zopiclone are contraindicated			
Both:			
1 Patient has insomnia and benzodiazepines and zopiclone are contraindicated; and			
2 For in-hospital use only.			
MIDAZOLAM			
Tab 7.5 mg			
Oral liq 2 mg per ml			
Inj 5 mg per ml, 1 ml plastic ampoule	22.50	10	Midazolam-Pfizer
Inj 1 mg per ml, 5 ml ampoule – 5% DV May-25 to 2027	7.80	10	Midazolam-Baxter
Inj 5 mg per ml, 3 ml ampoule – 5% DV May-25 to 2027	4.75	5	Midazolam-Baxter
PHENOBARBITONE			
Inj 130 mg per ml, 1 ml vial			
Inj 200 mg per ml, 1 ml ampoule			
TEMAZEPAM			
Tab 10 mg	1.40	25	Normison
TRIAZOLAM – Restricted: For continuation only			
➔ Tab 125 mcg			
➔ Tab 250 mcg			
ZOPICLONE			
Tab 7.5 mg – 5% DV Feb-25 to 2027	21.85	500	Zopiclone Actavis

Spinal Muscular Atrophy

NUSINERSEN – Restricted see terms [on the next page](#)

↓ Inj 12 mg per 5 ml vial 120,000.00 1 Spinraza

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ Restricted (RS1938)

Initiation

Re-assessment required after 12 months

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Continuation

Re-assessment required after 12 months

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day), in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

RISDIPLAM – Restricted see terms below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website <https://pharmac.govt.nz/risdiplam>

⚡ Powder for oral soln 750 mcg per ml, 60 mg per bottle..... 14,100.00 80 ml Evrysdi

➔ Restricted (RS1954)

Initiation

Re-assessment required after 12 months

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Continuation

Re-assessment required after 12 months

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day), in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Stimulants / ADHD Treatments			
ATOMOXETINE			
Cap 10 mg.....	43.02	28	APO-Atomoxetine
Cap 18 mg.....	45.57	28	APO-Atomoxetine
Cap 25 mg.....	44.30	28	APO-Atomoxetine
Cap 40 mg.....	46.21	28	APO-Atomoxetine
Cap 60 mg.....	51.31	28	APO-Atomoxetine
Cap 80 mg.....	65.20	28	APO-Atomoxetine
Cap 100 mg.....	65.71	28	APO-Atomoxetine
CAFFEINE			
Tab 100 mg			
DEXAMFETAMINE SULFATE – Restricted see terms below			
↓ Tab 5 mg	29.80	100	Dexamfetamine Aspen Noumed Dexamfetamine
→ Restricted (RS2166)			
Initiation – ADHD			
Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.			
Note: Prescribing practitioner detail is in the relevant approval notice published in the New Zealand Gazette. Approval notices are located through the 'Medicines (controlled drugs) with restrictions under regulation 22 of the Misuse of Drugs Regulations 1977' section of the Medsafe 'Restrictions on the Supply, Prescribing or Administration of Medicines under the Medicines Act 1981 and Misuse of Drugs Regulations 1977' webpage (https://www.medsafe.govt.nz/profs/riss/restrict.asp#MedicinesReg22 as of April 2025).			
Initiation – Narcolepsy			
Neurologist or respiratory specialist			
Patient suffers from narcolepsy.			
LISDEXAMFETAMINE DIMESILATE – Restricted see terms below			
↓ Cap 30 mg.....	60.00	30	Vyvanse
↓ Cap 50 mg.....	60.00	30	Vyvanse
↓ Cap 70 mg.....	60.00	30	Vyvanse
→ Restricted (RS2167)			
Initiation			
Either:			
1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all the following criteria prior to commencing treatment; or			
2 All of the following:			
2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and			
2.2 Diagnosed according to DSM-5 or ICD 11 criteria; and			
2.3 Any of the following:			
2.3.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) for ADHD and has not received sufficient clinical benefit or has experienced intolerable side effects; or			
2.3.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or			
2.3.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate; or			
2.3.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment			

continued...

continued...

- adherence difficulties; or
- 2.3.5 There is significant concern regarding the risk of diversion or abuse of immediate release methylphenidate hydrochloride; or
- 2.3.6 Both:
- 2.3.6.1 Patient would have been prescribed a subsidised formulation of methylphenidate hydrochloride (extended-release) but has been unable to access due to supply issues with methylphenidate hydrochloride (extended-release); and
- 2.3.6.2 Other alternative stimulant presentations (methylphenidate or dexamfetamine) are not appropriate; and
- 2.4 Lisdexamfetamine dimesilate is not to be used in combination with another funded methylphenidate presentation.

Note: Prescribing practitioner detail is in the relevant approval notice published in the New Zealand Gazette. Approval notices are located through the 'Medicines (controlled drugs) with restrictions under regulation 22 of the Misuse of Drugs Regulations 1977' section of the Medsafe 'Restrictions on the Supply, Prescribing or Administration of Medicines under the Medicines Act 1981 and Misuse of Drugs Regulations 1977' webpage (<https://www.medsafe.govt.nz/profs/riss/restrict.asp#MedicinesReg22> as of April 2025).

METHYLPHENIDATE HYDROCHLORIDE – Restricted see terms [below](#)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
⚡ Tab extended-release 18 mg.....	58.96 15.25	30	Concerta Methylphenidate ER - Teva
⚡ Tab extended-release 27 mg.....	65.44 16.25	30	Concerta Methylphenidate ER - Teva
⚡ Tab extended-release 36 mg.....	71.93 21.25	30	Concerta Methylphenidate ER - Teva
⚡ Tab extended-release 54 mg.....	86.24 24.25	30	Concerta Methylphenidate ER - Teva
⚡ Tab immediate-release 5 mg.....	3.20	30	Rubifen
⚡ Tab immediate-release 10 mg.....	4.00 3.00	30	Ritalin Rubifen
⚡ Tab modified-release 18 mg.....	15.25	30	Methylphenidate Sandoz XR
⚡ Tab immediate-release 20 mg.....	7.85	30	Rubifen
⚡ Tab sustained-release 20 mg.....	10.95	30	Rubifen SR
⚡ Tab modified-release 27 mg.....	16.25	30	Methylphenidate Sandoz XR
⚡ Tab modified-release 36 mg.....	21.25	30	Methylphenidate Sandoz XR
⚡ Tab modified-release 54 mg.....	24.25	30	Methylphenidate Sandoz XR
⚡ Cap modified-release 10 mg.....	19.41	30	Ritalin LA
⚡ Cap modified-release 20 mg.....	27.72	30	Ritalin LA
⚡ Cap modified-release 30 mg.....	34.39	30	Ritalin LA
⚡ Cap modified-release 40 mg.....	38.67	30	Ritalin LA

➡ **Restricted (RS2168)**

Initiation – ADHD (immediate-release and sustained-release formulations)

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.

continued...

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$ Per	

continued...

Initiation – Extended-release and modified-release formulations

Both:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Either:
 - 2.1 Patient is taking a currently listed formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
 - 2.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Note: Prescribing practitioner detail is in the relevant approval notice published in the New Zealand Gazette. Approval notices are located through the 'Medicines (controlled drugs) with restrictions under regulation 22 of the Misuse of Drugs Regulations 1977' section of the Medsafe 'Restrictions on the Supply, Prescribing or Administration of Medicines under the Medicines Act 1981 and Misuse of Drugs Regulations 1977' webpage (<https://www.medsafe.govt.nz/profs/riss/restrict.asp#MedicinesReg22> as of April 2025).

Initiation – Narcolepsy (immediate-release and sustained-release formulations)

Neurologist or respiratory specialist

Patient suffers from narcolepsy.

Initiation – Narcolepsy* (extended-release only)

Neurologist or respiratory specialist

Patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Concerta, Ritalin LA or Methylphenidate Sandoz XR.

MODAFINIL – **Restricted** see terms [below](#)

↓ Tab 100 mg – 5% DV May-25 to 2027..... 14.27 30 **Modafinil Max Health**

→ **Restricted (RS2106)**

Initiation – Narcolepsy

Neurologist or respiratory specialist

Either:

- 1 All of the following:
 - 1.1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
 - 1.2 Either:
 - 1.2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 1.2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
 - 1.3 Either:
 - 1.3.1 An effective dose of a listed formulation of methylphenidate or dexamphetamine has been trialed and discontinued because of intolerable side effects; or
 - 1.3.2 Methylphenidate and dexamphetamine are contraindicated; or
- 2 Both:
 - 2.1 Patient meets the Hospital Restriction criteria for methylphenidate hydrochloride for narcolepsy; and
 - 2.2 Patient is unable to access methylphenidate hydrochloride presentations due to an out of stock (see note).

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock of methylphenidate hydrochloride.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE

Tab 5 mg	3.70	84	Ipca-Donepezil
Tab 10 mg	5.50	84	Ipca-Donepezil

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
RIVASTIGMINE – Restricted see terms below			
⚡ Patch 4.6 mg per 24 hour – 5% DV Mar-25 to 2027.....	49.40	30	Rivastigmine Patch BNM 5
⚡ Patch 9.5 mg per 24 hour – 5% DV Mar-25 to 2027.....	49.40	30	Rivastigmine Patch BNM 10

➡ **Restricted (RS2139)**

Initiation

Re-assessment required after 6 months

Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient is contraindicated to or has experienced intolerable side effects from donepezil tablets.

Continuation

Re-assessment required after 12 months

Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE – Restricted see terms [below](#)

⚡ Tab 2 mg with naloxone 0.5 mg – 5% DV May-26 to 2028.....	11.76	28	Buprenorphine Naloxone BNM
⚡ Tab 8 mg with naloxone 2 mg – 5% DV May-26 to 2028.....	26.86	28	Buprenorphine Naloxone BNM

➡ **Restricted (RS1172)**

Initiation – Detoxification

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Prescriber works in an opioid treatment service approved by the Ministry of Health.

Initiation – Maintenance treatment

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Prescriber works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg.....	15.00	30	Zyban
----------------------------------	-------	----	-------

DISULFIRAM

Tab 200 mg	236.40	100	Antabuse
------------------	--------	-----	----------

NALTREXONE HYDROCHLORIDE – Restricted see terms [below](#)

⚡ Tab 50 mg	83.33	30	Naltraccord
-------------------	-------	----	-------------

➡ **Restricted (RS1173)**

Initiation – Alcohol dependence

Both:

- 1 Patient is currently enrolled, or is planned to be enrolled, in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Naltrexone is to be prescribed by, or on the recommendation of, a physician working in an Alcohol and Drug Service.

Initiation – Constipation

For the treatment of opioid-induced constipation.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NICOTINE – Some items restricted see terms below			
Patch 7 mg per 24 hours	19.62	28	Habitrol
Patch 14 mg per 24 hours	21.57	28	Habitrol
Patch 21 mg per 24 hours	24.72	28	Habitrol
↓ Oral spray 1 mg per dose			<i>e.g. Nicorette QuickMist Mouth Spray</i>
Lozenge 1 mg	22.53	216	Habitrol
Lozenge 2 mg	24.68	216	Habitrol
↓ Soln for inhalation 15 mg cartridge			
Gum 2 mg	23.02	204	Habitrol (Fruit) Habitrol (Mint)
Gum 4 mg	25.98	204	Habitrol (Fruit) Habitrol (Mint)

➔ **Restricted (RS1873)**

Initiation

Any of the following:

- 1 For perioperative use in patients who have a 'nil by mouth' instruction; or
- 2 For use within mental health inpatient units; or
- 3 Patient would be admitted to a mental health inpatient unit, but is unable to due to COVID-19 self-isolation requirement; or
- 4 For acute use in agitated patients who are unable to leave the hospital facilities.

VARENICLINE – Restricted see terms [below](#)

↓ Tab 0.5 mg x 11 and 1 mg x 42 – 5% DV Sep-26 to 2028	16.67	53	Champix
	15.99		Pharmacor Varenicline
↓ Tab 1 mg – 5% DV Sep-26 to 2028	17.62	56	Champix
	10.99		Pharmacor Varenicline

(Champix Tab 0.5 mg x 11 and 1 mg x 42 to be delisted 1 September 2026)

(Champix Tab 1 mg to be delisted 1 September 2026)

➔ **Restricted (RS1702)**

Initiation

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline in a 12 month period.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE – **Restricted** see terms [below](#)

⚡ Inj 25 mg vial – 5% DV Apr-25 to 2027	50.05	1	Bendamustine Sandoz
⚡ Inj 100 mg vial – 5% DV Apr-25 to 2027	200.20	1	Bendamustine Sandoz
➡ Restricted (RS2061)			

Initiation – CLL*

All of the following:

- 1 The patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has ECOG performance status 0-2; and
- 3 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a * includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initiation – Indolent, Low-grade lymphomas

Re-assessment required after 9 months

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Continuation – Indolent, Low-grade lymphomas

Re-assessment required after 9 months

Either:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or			
2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.			
Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenström's macroglobulinaemia.			
Initiation – Hodgkin's lymphoma*			
Relevant specialist or medical practitioner on the recommendation of a relevant specialist			
Limited to 6 months treatment			
All of the following:			
1 Patient has Hodgkin's lymphoma requiring treatment; and			
2 Patient has a ECOG performance status of 0-2; and			
3 Patient has received one prior line of chemotherapy; and			
4 Patient's disease relapsed or was refractory following prior chemotherapy; and			
5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m ² twice per cycle, for a maximum of four cycles.			
Note: Indications marked with * are unapproved indications.			
BUSULFAN			
Tab 2 mg89.25	100	Myleran	
Inj 6 mg per ml, 10 ml ampoule			
CARMUSTINE			
Inj 100 mg vial – 5% DV Dec-26 to 2028710.00	1	BiCNU BiCNU S29 Novadoz	
(BiCNU S29 Inj 100 mg vial to be delisted 1 December 2026)			
(Novadoz Inj 100 mg vial to be delisted 1 December 2026)			
CHLORAMBUCIL			
Tab 2 mg			
CYCLOPHOSPHAMIDE			
Tab 50 mg – 5% DV Dec-24 to 2027145.00	50	Cyclonex	
Inj 1 g vial – 5% DV Feb-25 to 202747.46	1	Endoxan	
Inj 2 g vial – 5% DV Feb-25 to 202795.06	1	Endoxan	
IFOSFAMIDE			
Inj 1 g vial96.00	1	Holoxan	
Inj 2 g vial180.00	1	Holoxan	
IFOSFAMIDE WITH MESNA			
Inj 2 g vial with mesna 3 x 100 mg per ml, 4 ml ampoule62.00	1	Ifomid-M	
LOMUSTINE			
Cap 40 mg880.00	20	Medac	
MELPHALAN			
Tab 2 mg			
Inj 50 mg vial – 5% DV Dec-26 to 202946.80	1	Melpha	
THIOTEPA			
Inj 15 mg vial398.00	1	Tepadina	
Inj 100 mg vial1,800.00	1	Tepadina Thiotepa-Reach	

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anthracyclines and Other Cytotoxic Antibiotics			
BLEOMYCIN SULPHATE			
Inj 15,000 iu vial.....	185.16	1	DBL Bleomycin Sulfate
DACTINOMYCIN [ACTINOMYCIN D]			
Inj 0.5 mg vial	255.00	1	Cosmegen
DAUNORUBICIN			
Inj 18.7 mg vial	160.75	1	Pfizer
Inj 20 mg vial	171.93	1	Cerubidine
DOXORUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 5 ml vial			
Inj 2 mg per ml, 25 ml vial.....	75.55	1	Doxorubicin Ebewe
Inj 50 mg vial			
Inj 2 mg per ml, 100 ml vial.....	164.50	1	Adriamycin Doxorubicin Ebewe
EPIRUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 25 ml vial.....	284.36	1	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial.....	284.36	1	Epirubicin Ebewe
IDARUBICIN HYDROCHLORIDE			
Inj 5 mg vial	109.74	1	Zavedos
Inj 10 mg vial	233.64	1	Zavedos
MITOMYCIN C			
Inj 5 mg vial			
Inj 20 mg vial – 5% DV May-26 to 2028.....	1,129.94	1	Teva
MITOZANTRONE			
Inj 2 mg per ml, 10 ml vial.....	170.63	1	Mitozantrone Ebewe
Antimetabolites			
AZACITIDINE – Restricted see terms below			
⚠ Inj 100 mg vial – 5% DV Mar-25 to 2027	50.00	1	Azacitidine Dr Reddy's
➡ Restricted (RS2116)			
Initiation			
<i>Re-assessment required after 12 months</i>			
Both:			
1 Any of the following:			
1.1 The individual has intermediate or high risk MDS based on an internationally recognised scoring system; or			
1.2 The individual has chronic myelomonocytic leukaemia (based on an intermediate or high risk score from an internationally recognised scoring system or 10%-29% marrow blasts without myeloproliferative disorder); or			
1.3 The individual has acute myeloid leukaemia according to World Health Organisation (WHO) Classification; and			
2 The individual has an estimated life expectancy of at least 3 months.			
Continuation			
<i>Re-assessment required after 12 months</i>			
No evidence of disease progression.			
CAPECITABINE			
Tab 150 mg – 5% DV Feb-26 to 2028	10.92	60	Capecitabine Viatris
Tab 500 mg – 5% DV Feb-26 to 2028	50.96	120	Capecitabine Viatris
CLADRIBINE			
Inj 2 mg per ml, 5 ml vial			
Inj 1 mg per ml, 10 ml vial.....	749.96	1	Leustatin

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CYTARABINE			
Inj 20 mg per ml, 5 ml vial.....	472.00	5	Pfizer
Inj 100 mg per ml, 20 ml vial.....	48.80	1	Cytarabine DBL Pfizer
FLUDARABINE PHOSPHATE			
Tab 10 mg	412.00	20	Fludara Oral
Inj 50 mg vial	744.95	5	Fludarabine Ebewe
FLUOROURACIL			
Inj 50 mg per ml, 20 ml vial – 5% DV Dec-24 to 2027	10.51	1	Fluorouracil Accord
Inj 50 mg per ml, 50 ml vial.....	14.72	1	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial – 5% DV Dec-24 to 2027	19.36	1	Fluorouracil Accord
GEMCITABINE HYDROCHLORIDE			
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine), 26.3 ml vial	18.94	1	DBL Gemcitabine
MERCAPTOPURINE			
Tab 50 mg – 5% DV Dec-25 to 2028	19.50	25	Puri-nethol
↓ Oral suspension 20 mg per ml.....	428.00	100 ml	Xaluprine Allmercap
→ Restricted (RS1635)			
Initiation			
Paediatric haematologist or paediatric oncologist			
<i>Re-assessment required after 12 months</i>			
The patient requires a total dose of less than one full 50 mg tablet per day.			
Continuation			
Paediatric haematologist or paediatric oncologist			
<i>Re-assessment required after 12 months</i>			
The patient requires a total dose of less than one full 50 mg tablet per day.			
METHOTREXATE			
Tab 2.5 mg – 5% DV Dec-24 to 2027	7.80	90	Trexate
Tab 10 mg – 5% DV Dec-24 to 2027	26.40	90	Trexate
Inj 2.5 mg per ml, 2 ml vial			
Inj 7.5 mg prefilled syringe – 5% DV Feb-25 to 2027	29.17	1	Methotrexate Pharmascience
Inj 10 mg prefilled syringe – 5% DV Feb-25 to 2027	19.09	1	Methotrexate Sandoz
Inj 15 mg prefilled syringe – 5% DV Feb-25 to 2027	24.53	1	Methotrexate Sandoz
Inj 20 mg prefilled syringe – 5% DV Feb-25 to 2027	16.64	1	Methotrexate Sandoz
Inj 25 mg prefilled syringe – 5% DV Feb-25 to 2027	20.72	1	Methotrexate Sandoz
Inj 30 mg prefilled syringe – 5% DV Feb-25 to 2027	55.00	1	Methotrexate Sandoz
Inj 25 mg per ml, 2 ml vial.....	30.00	5	Methotrexate DBL
Inj 25 mg per ml, 20 ml vial.....	45.00	1	Onco-Vial DBL Methotrexate
Inj 100 mg per ml, 10 ml vial.....	25.00	1	Onco-Vial Methotrexate Ebewe
Inj 100 mg per ml, 50 ml vial.....	67.99	1	Methotrexate Ebewe
PEMETREXED			
Inj 100 mg vial – 5% DV Apr-25 to 2027	8.99	1	Pemetrexed-AFT
Inj 500 mg vial – 5% DV Apr-25 to 2027	29.99	1	Pemetrexed-AFT
THIOGUANINE			
Tab 40 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Other Cytotoxic Agents			
AMSACRINE			
Inj 50 mg per ml, 1.5 ml ampoule			
Inj 75 mg			
ANAGRELIDE HYDROCHLORIDE			
Cap 0.5 mg			
ARSENIC TRIOXIDE			
Inj 1 mg per ml, 10 ml vial.....	4,817.00	10	Phenasen
BORTEZOMIB – Restricted see terms below			
⚡ Inj 3.5 mg vial – 5% DV Sep-26 to 2028	23.99	1	Bortezomib Eugia
	74.93		DBL Bortezomib
<i>(DBL Bortezomib Inj 3.5 mg vial to be delisted 1 September 2026)</i>			
➡ Restricted (RS2169)			
Initiation – plasma cell dyscrasia			
The patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.			
Initiation – Waldenström Macroglobulinaemia			
<i>Re-assessment required after 12 months</i>			
Both:			
1 The patient has Waldenström Macroglobulinaemia/Lymphoplasmacytic Lymphoma requiring treatment; and			
2 The patient has not received prior bortezomib treatment.			
Continuation – Waldenström Macroglobulinaemia			
<i>Re-assessment required after 12 months</i>			
Patient has no evidence of clinical disease progression during bortezomib use.			
DACARBAZINE			
Inj 200 mg vial	72.11	1	DBL Dacarbazine
ETOPOSIDE			
Cap 50 mg	340.73	20	Vepesid
Cap 100 mg	340.73	10	Vepesid
Inj 20 mg per ml, 5 ml vial.....	7.90	1	Rex Medical
ETOPOSIDE (AS PHOSPHATE)			
Inj 100 mg vial	40.00	1	Etopophos
HYDROXYUREA [HYDROXYCARBAMIDE]			
Cap 500 mg	20.72	100	Devatis
IBRUTINIB – Restricted see terms below			
⚡ Tab 140 mg	3,217.00	30	Imbruvica
⚡ Tab 420 mg	9,652.00	30	Imbruvica
➡ Restricted (RS2190)			
Initiation – previously untreated chronic lymphocytic leukaemia in combination with venetoclax			
Either:			
1 Individual is currently on treatment with ibrutinib and/or venetoclax and met all of the following criteria prior to commencing treatment; or			
2 Both:			
2.1 Individual has previously untreated CLL; and			
2.2 Ibrutinib is to be administered at a maximum dose of 420 mg daily for 3 (28 day) cycles as monotherapy, followed by a maximum of 12 (28 day) cycles in combination with venetoclax.			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – chronic lymphocytic leukaemia (CLL)

Re-assessment required after 6 months

All of the following:

- 1 Individual has chronic lymphocytic leukaemia (CLL) requiring therapy; and
- 2 Ibrutinib is to be used as monotherapy; and
- 3 Individual has experienced intolerable side effects, or their disease has relapsed or is refractory following at least one prior line of therapy; and
- 4 Individual has not received ibrutinib monotherapy previously.

Continuation – chronic lymphocytic leukaemia (CLL)

Re-assessment required after 12 months

No evidence of clinical disease progression.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

IRINOTECAN HYDROCHLORIDE

Inj 20 mg per ml, 5 ml vial.....	52.57	1	Accord
Inj 20 mg per ml, 25 ml vial.....	262.85	1	Accord

LENALIDOMIDE (VIATRIS) – **Restricted** see terms [below](#)

↓ Cap 5 mg – 5% DV Feb-25 to 31 Jan 2028.....	76.92	21	Lenalidomide Viatris
↓ Cap 10 mg – 5% DV Feb-25 to 31 Jan 2028.....	50.30	21	Lenalidomide Viatris
↓ Cap 15 mg – 5% DV Feb-25 to 31 Jan 2028.....	62.13	21	Lenalidomide Viatris
↓ Cap 25 mg – 5% DV Feb-25 to 31 Jan 2028.....	65.09	21	Lenalidomide Viatris

→ **Restricted** (RS2044)

Initiation – Plasma cell dyscrasia

Any relevant practitioner

Both:

- 1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient is not refractory to prior lenalidomide use.

Initiation – Myelodysplastic syndrome

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5) associated with a deletion 5q cytogenetic abnormality; and
- 2 Patient has transfusion-dependent anaemia.

Continuation – Myelodysplastic syndrome

Any relevant practitioner

Re-assessment required after 12 months

Both:

- 1 Patient has not needed a transfusion in the last 4 months; and
- 2 No evidence of disease progression.

NIRAPARIB – **Restricted** see terms [below](#)

↓ Tab 100 mg	8,929.84	56	Zejula
	13,393.50	84	Zejula
↓ Cap 100 mg	8,929.84	56	Zejula

(Zejula Cap 100 mg to be delisted 1 October 2026)

→ **Restricted** (RS2027)

Initiation

Re-assessment required after 6 months

All of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1 Patient has advanced high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line** of treatment with platinum-based chemotherapy; and
- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and
- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Either:
 - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen; or
 - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

Continuation

Re-assessment required after 6 months

All of the following:

- 1 No evidence of progressive disease; and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Either:
 - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
 - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: * "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB – **Restricted** see terms [below](#)

⚡ Tab 100 mg	3,701.00	56	Lynparza
⚡ Tab 150 mg	3,701.00	56	Lynparza

➡ **Restricted** (RS1925)

Initiation – Ovarian cancer

Medical oncologist

Re-assessment required after 12 months

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Either:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Continuation – Ovarian cancer

Medical oncologist

Re-assessment required after 12 months

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
 - 5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PEGASPARGASE – **Restricted** see terms [below](#)

↓ Inj 750 iu per ml, 5 ml vial.....3,973.25 1 Oncaspar LYO

→ **Restricted (RS2176)**

Initiation – Newly diagnosed ALL

Limited to 15 months treatment

Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initiation – Relapsed ALL

Limited to 12 months treatment

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initiation – Lymphoma

Limited to 12 months treatment

Patient has lymphoma requiring L-asparaginase containing protocol (e.g. SMILE).

PENTOSTATIN [DEOXYCOFORMYCIN]

Inj 10 mg vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
POMALIDOMIDE – Restricted see terms below			
⚡ Cap 1 mg – 5% DV Aug-24 to 31 Jul 2027	47.45	14	Pomolide
	71.18	21	Pomolide
⚡ Cap 2 mg – 5% DV Aug-24 to 31 Jul 2027	94.90	14	Pomolide
	142.35	21	Pomolide
⚡ Cap 3 mg – 5% DV Aug-24 to 31 Jul 2027	142.35	14	Pomolide
	213.53	21	Pomolide
⚡ Cap 4 mg – 5% DV Aug-24 to 31 Jul 2027	189.81	14	Pomolide
	284.71	21	Pomolide

➡ **Restricted (RS2045)**

Initiation – Relapsed/refractory plasma cell dyscrasia

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 Patient has relapsed or refractory plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient has not received prior funded pomalidomide.

Continuation – Relapsed/refractory plasma cell dyscrasia

Any relevant practitioner

Re-assessment required after 12 months

Patient has no evidence of disease progression.

PROCARBAZINE HYDROCHLORIDE

Cap 50 mg 980.00 50 Natulan

TEMOZOLOMIDE – Restricted see terms [below](#)

⚡ Cap 5 mg	9.13	5	Temaccord
	12.70		Temozolomide (Devatis)
	9.13		Temozolomide Taro
⚡ Cap 20 mg	16.38	5	Temaccord
⚡ Cap 100 mg	35.98	5	Temaccord
⚡ Cap 140 mg	50.12	5	Temaccord
⚡ Cap 250 mg	86.34	5	Temaccord

➡ **Restricted (RS2201)**

Initiation – gliomas

Re-assessment required after 12 months

Patient has a glioma.

Continuation – gliomas

Re-assessment required after 12 months

Treatment remains appropriate and patient is benefitting from treatment.

Initiation – Neuroendocrine tumours

Re-assessment required after 9 months

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Continuation – Neuroendocrine tumours

Re-assessment required after 6 months

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Initiation – ewing's sarcoma

Re-assessment required after 9 months

Patient has relapse or refractory Ewing's sarcoma.

Continuation – ewing's sarcoma

Re-assessment required after 6 months

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Initiation – Neuroblastoma

Re-assessment required after 12 months

Patient has neuroblastoma.

Continuation – Neuroblastoma

Re-assessment required after 12 months

Patient has no evidence of disease progression.

THALIDOMIDE – **Restricted** see terms [below](#)

↓ Cap 50 mg.....	378.00	28	Thalomid
↓ Cap 100 mg.....	756.00	28	Thalomid

→ **Restricted (RS2046)**

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; or
- 2 The patient has erythema nodosum leprosum.

Continuation

Patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen

TRETINOIN

Cap 10 mg.....	479.50	100	Vesanoid
----------------	--------	-----	----------

VENETOCLAX – **Restricted** see terms [below](#)

↓ Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg.....	1,771.86	42	Venclexta
↓ Tab 10 mg.....	13.68	2	Venclexta
↓ Tab 50 mg.....	239.44	7	Venclexta
↓ Tab 100 mg.....	8,209.41	120	Venclexta

→ **Restricted (RS2200)**

Initiation – previously untreated chronic lymphocytic leukaemia in combination with obinutuzumab

Either:

- 1 Individual is currently on treatment with venetoclax and obinutuzumab and met all of the following criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Individual has previously untreated chronic lymphocytic leukaemia; and
 - 2.2 Venetoclax is to be administered with obinutuzumab; and
 - 2.3 Venetoclax is to be used to a maximum dose of 400 mg and for a total of 12 (28 day) cycles.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Initiation – previously untreated chronic lymphocytic leukaemia in combination with ibrutinib

Either:

- 1 Individual is currently on treatment with venetoclax and/or ibrutinib and met all of the following criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Individual has previously untreated chronic lymphocytic leukaemia; and
 - 2.2 Venetoclax is to be administered in combination with ibrutinib; and
 - 2.3 Venetoclax is to be used to a maximum dose of 400 mg and for a total of 12 (28 day) cycles.

Initiation – relapsed/refractory chronic lymphocytic leukaemia

Re-assessment required after 8 months

All of the following:

- 1 Individual has chronic lymphocytic leukaemia requiring treatment; and
- 2 Individual has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 The individual's disease has relapsed; and
- 4 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 5 Individual has an ECOG performance status of 0-2.

Continuation – relapsed/refractory chronic lymphocytic leukaemia

Re-assessment required after 6 months

Both:

- 1 Treatment remains clinically appropriate and the individual is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

Initiation – previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*

Re-assessment required after 6 months

All of the following:

- 1 Individual has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that the individual has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Individual has an ECOG performance status of 0-2.

Continuation – previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*

Re-assessment required after 6 months

No evidence of disease progression.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are unapproved indications

Initiation – previously untreated acute myeloid leukaemia

Re-assessment required after 6 months

Either:

- 1 The individual is currently on treatment with venetoclax and met all remaining special authority criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Individual has previously untreated acute myeloid leukaemia (see note a), according to World Health Organization (WHO) Classification; and
 - 2.2 Venetoclax not to be used in combination with standard intensive remission induction chemotherapy; and
 - 2.3 Venetoclax to be used in combination with azacitidine or low dose cytarabine.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – previously untreated acute myeloid leukaemia

Re-assessment required after 6 months

No evidence of disease progression.

Notes:

- 'Acute myeloid leukaemia' includes myeloid sarcoma*
- Indications marked with * are Unapproved indications

Platinum Compounds

CARBOPLATIN

Inj 10 mg per ml, 45 ml vial – **5% DV Dec-24 to 2027**25.73 1 **Carboplatin Accord**

CISPLATIN

Inj 1 mg per ml, 50 ml vial.....9.45 1 Cisplatin Accord

Inj 1 mg per ml, 100 ml vial – **5% DV Dec-24 to 2027**18.90 1 **Cisplatin Accord**

OXALIPLATIN

Inj 5 mg per ml, 20 ml vial.....33.35 1 Alchemy Oxaliplatin

Protein-Tyrosine Kinase Inhibitors

ALECTINIB – Restricted see terms [below](#)

↓ Cap 150 mg7,935.00 224 Alecensa

→ **Restricted (RS1712)**

Initiation

Re-assessment required after 6 months

All of the following:

- Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- Patient has an ECOG performance score of 0-2.

Continuation

Re-assessment required after 6 months

Both:

- No evidence of progressive disease according to RECIST criteria; and
- The patient is benefitting from and tolerating treatment.

AXITINIB – Restricted see terms [below](#)

↓ Tab 1 mg536.40 28 Inlyta

↓ Tab 5 mg2,682.00 28 Inlyta

→ **Restricted (RS2107)**

Initiation

Re-assessment required after 4 months

All of the following:

- The patient has metastatic renal cell carcinoma; and
- The disease is of predominant clear cell histology; and
- The patient has documented disease progression following one previous line of treatment; and
- The patient has ECOG performance status of 0-2.

Continuation

Re-assessment required after 4 months

No evidence of disease progression..

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CRIZOTINIB – Restricted see terms below			
⚡ Cap 200 mg.....	7,250.00	60	Xalkori
⚡ Cap 250 mg.....	7,250.00	60	Xalkori

➡ **Restricted (RS2144)**

Initiation

Re-assessment required after 6 months

All of the following:

- 1 Individual has locally advanced or metastatic, unresectable, non-squamous non-small cell lung cancer; and
- 2 Either:
 - 2.1 The individual has not received entrectinib; or
 - 2.2 Both:
 - 2.2.1 The individual has received treatment with entrectinib and has discontinued entrectinib due to intolerance; and
 - 2.2.2 The cancer did not progress while the individual was on entrectinib; and
- 3 There is documentation confirming that the patient has a ROS1 rearrangement using an appropriate ROS1 test; and
- 4 Individual has ECOG performance score of 0-3; and
- 5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation

Re-assessment required after 6 months

Both:

- 1 Response to treatment has been determined by comparable radiological assessment following the most recent treatment period; and
- 2 No evidence of disease progression.

DABRAFENIB – Restricted see terms [below](#)

⚡ Cap 50 mg	6,320.86	120	Tafinlar
⚡ Cap 75 mg	9,481.29	120	Tafinlar

➡ **Restricted (RS2145)**

Initiation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 Either:
 - 1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or
 - 1.2 Both:
 - 1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and
 - 1.2.2 Adjuvant treatment with dabrafenib is required; and
- 2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
- 3 Treatment must be adjuvant to complete surgical resection; and
- 4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
- 5 The individual has a confirmed BRAF mutation; and
- 6 Dabrafenib must be administered in combination with trametinib; and
- 7 The individual has ECOG performance score 0-2.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

Any of the following:

- 1 All of the following:
 - 1.1 No evidence of disease recurrence; and
 - 1.2 Dabrafenib must be administered in combination with trametinib; and
 - 1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or
- 2 All of the following:
 - 2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 2.3 The individual meets initiation criteria for dabrafenib for unresectable or metastatic melanoma; or
- 3 All of the following:
 - 3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 3.2 The individual has received a BRAF/MEK inhibitor for unresectable or metastatic melanoma; and
 - 3.3 The individual meets continuation criteria for dabrafenib for unresectable or metastatic melanoma.

Initiation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance score 0-2; and
- 4 The individual has confirmed BRAF mutation; and
- 5 Dabrafenib must be administered in combination with trametinib; and
- 6 Any of the following:
 - 6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or
 - 6.3 All of the following:
 - 6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
 - 6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
 - 6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MEK inhibitor.

Continuation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

Both:

- 1 Any of the following:
 - 1.1 The individual's disease has had a complete response to treatment; or
 - 1.2 The individual's disease has had a partial response to treatment; or
 - 1.3 The individual has stable disease with treatment; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

DASATINIB – Restricted see terms [on the next page](#)

↓ Tab 20 mg – 5% DV Mar-25 to 2027	132.88	60	Dasatinib-Teva
↓ Tab 50 mg – 5% DV Mar-25 to 2027	304.13	60	Dasatinib-Teva
↓ Tab 70 mg – 5% DV Mar-25 to 2027	415.75	60	Dasatinib-Teva

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2055)

Initiation

Haematologist or any relevant practitioner on the recommendation of a haematologist

Re-assessment required after 6 months

Any of the following:

- 1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or
- 2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or
- 3 Both:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Any of the following:
 - 3.2.1 Patient has documented treatment failure* with imatinib; or
 - 3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

Continuation

Haematologist or any relevant practitioner on the recommendation of a haematologist

Re-assessment required after 6 months

Both:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines.

ENTRECTINIB – **Restricted** see terms [below](#)

⚡ Cap 200 mg.....	9,610.00	90	Rozlytrek
-------------------	----------	----	-----------

➔ Restricted (RS2146)

Initiation

Re-assessment required after 6 months

All of the following:

- 1 Individual has locally advanced or metastatic, unresectable, non-squamous non-small cell lung cancer; and
- 2 Either:
 - 2.1 The individual has not received crizotinib; or
 - 2.2 Both:
 - 2.2.1 The individual has received an initial Special Authority approval for crizotinib and has discontinued crizotinib due to intolerance; and
 - 2.2.2 The cancer did not progress while the individual was on crizotinib; and
- 3 There is documentation confirming that the patient has a ROS1 rearrangement using an appropriate ROS1 test; and
- 4 Individual has ECOG performance score of 0-3; and
- 5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation

Re-assessment required after 6 months

Both:

- 1 Response to treatment has been determined by comparable radiological assessment following the most recent treatment period; and
- 2 No evidence of disease progression.

ERLOTINIB – **Restricted** see terms [below](#)

⚡ Tab 100 mg – 5% DV Oct-24 to 2027	280.84	30	Alchemy
⚡ Tab 150 mg – 5% DV Oct-24 to 2027	484.24	30	Alchemy

➔ Restricted (RS2078)

Initiation

Re-assessment required after 4 months

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and			
2 There is documentation confirming that the disease expresses activating mutations of EGFR; and			
3 Any of the following:			
3.1 Patient is treatment naive; or			
3.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or			
3.3 Both:			
3.3.1 The patient has discontinued osimertinib or gefitinib due to intolerance; and			
3.3.2 The cancer did not progress while on osimertinib or gefitinib.			
Continuation			
<i>Re-assessment required after 6 months</i>			
Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.			
GEFITINIB – Restricted see terms below			
↓ Tab 250 mg	918.00	30	Iressa
➔ Restricted (RS2079)			
Initiation			
<i>Re-assessment required after 4 months</i>			
All of the following:			
1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and			
2 Any of the following:			
2.1 Patient is treatment naive; or			
2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or			
2.3 Both:			
2.3.1 The patient has discontinued osimertinib or erlotinib due to intolerance; and			
2.3.2 The cancer did not progress whilst on osimertinib or erlotinib; and			
3 There is documentation confirming that disease expresses activating mutations of EGFR.			
Continuation			
<i>Re-assessment required after 6 months</i>			
Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.			
IMATINIB MESILATE			
Cap 100 mg	44.93	60	Imatinib-Rex
Cap 400 mg	69.76	30	Imatinib-Rex
LAPATINIB – Restricted see terms below			
↓ Tab 250 mg			
➔ Restricted (RS1828)			
Initiation			
For continuation use only.			
Continuation			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and			
2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and			
3 Lapatinib not to be given in combination with trastuzumab; and			
4 Lapatinib to be discontinued at disease progression.			
LENVATINIB – Restricted see terms on the next page			
↓ Cap 4 mg	3,407.40	30	Lenvima
↓ Cap 10 mg	3,407.40	30	Lenvima

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

➔ Restricted (RS2098)

Initiation – thyroid cancer

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
 - 2.2 Either:
 - 2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
 - 2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures; and
 - 2.3 Any of the following:
 - 2.3.1 A lesion without iodine uptake in a RAI scan; or
 - 2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
 - 2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
 - 2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other; and
 - 2.4 Patient has thyroid stimulating hormone (TSH) adequately suppressed; and
 - 2.5 Patient is not a candidate for radiotherapy with curative intent; and
 - 2.6 Surgery is clinically inappropriate; and
 - 2.7 Patient has an ECOG performance status of 0-2.

Continuation – thyroid cancer

Re-assessment required after 6 months

there is no evidence of disease progression.

Initiation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

All of the following:

- 1 Patient has unresectable hepatocellular carcinoma; and
- 2 Patient has preserved liver function (Childs-Pugh A); and
- 3 Transarterial chemoembolisation (TACE) is unsuitable; and
- 4 Patient has an ECOG performance status of 0-2; and
- 5 Either:
 - 5.1 Patient has not received prior systemic therapy for their disease in the palliative setting; or
 - 5.2 Both:
 - 5.2.1 Patient has experienced treatment-limiting toxicity from treatment with atezolizumab with bevacizumab; and
 - 5.2.2 No disease progression since initiation of atezolizumab with bevacizumab.

Continuation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

there is no evidence of disease progression.

Initiation – renal cell carcinoma

Re-assessment required after 4 months

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma; and
 - 1.2 The disease is of predominant clear-cell histology; and
 - 1.3 The patient has documented disease progression following one previous line of treatment; and
 - 1.4 The patient has an ECOG performance status of 0-2; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.5 Lenvatinib is to be used in combination with everolimus; or
- 2 All of the following:
 - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
 - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
 - 2.3 Lenvatinib is to be used in combination with everolimus; and
 - 2.4 There is no evidence of disease progression.

Continuation – renal cell carcinoma

Re-assessment required after 4 months

there is no evidence of disease progression.

MIDOSTAURIN – Restricted see terms [below](#)

↓ Cap 25 mg 10,981.00 56 Rydapt

→ **Restricted (RS2033)**

Initiation

All of the following:

- 1 Patient has a diagnosis of acute myeloid leukaemia; and
- 2 Condition must be FMS tyrosine kinase 3 (FLT3) mutation positive; and
- 3 Patient must not have received a prior line of intensive chemotherapy for acute myeloid leukaemia; and
- 4 Patient is to receive standard intensive chemotherapy in combination with midostaurin only; and
- 5 Midostaurin to be funded for a maximum of 4 cycles.

NILOTINIB – Restricted see terms [below](#)

↓ Cap 150 mg 4,680.00 120 Tasigna
↓ Cap 200 mg 6,532.00 120 Tasigna

→ **Restricted (RS2010)**

Initiation

Haematologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with a tyrosine kinase inhibitor (TKI); or
 - 2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Continuation

Haematologist

Re-assessment required after 6 months

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

OSIMERTINIB – Restricted see terms [on the next page](#)

↓ Tab 40 mg 9,310.00 30 Tagrisso
↓ Tab 80 mg 9,310.00 30 Tagrisso

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2080)

Initiation – NSCLC – first line

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
- 2 Any of the following:
 - 2.1 Patient is treatment naïve; or
 - 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued gefitinib or erlotinib due to intolerance; and
 - 2.3.2 The cancer did not progress while on gefitinib or erlotinib; and
- 3 There is documentation confirming that the cancer expresses activating mutations of EGFR; and
- 4 Patient has an ECOG performance status 0-3; and
- 5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – NSCLC – first line

Re-assessment required after 6 months

response to or stable disease with treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Initiation – NSCLC – second line

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
- 2 Patient has an ECOG performance status 0-3; and
- 3 The patient must have received previous treatment with erlotinib or gefitinib; and
- 4 There is documentation confirming that the cancer expresses T790M mutation of EGFR following progression on or after erlotinib or gefitinib; and
- 5 The treatment must be given as monotherapy; and
- 6 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – NSCLC – second line

Re-assessment required after 6 months

response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

PALBOCICLIB – **Restricted** see terms [below](#)

⚡ Tab 75 mg	1,200.00	21	Palbociclib Pfizer
⚡ Tab 100 mg	1,200.00	21	Palbociclib Pfizer
⚡ Tab 125 mg	1,200.00	21	Palbociclib Pfizer

➔ Restricted (RS2034)

Initiation

Re-assessment required after 6 months

Either:

- 1 All of the following:
 - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
 - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
 - 1.3 Patient has an ECOG performance score of 0-2; and
 - 1.4 Either:
 - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
 - 1.4.2 Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
- 1.4.2.2 Patient has not received prior systemic treatment for metastatic disease; and
- 1.5 Treatment must be used in combination with an endocrine partner; and
- 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has an active Special Authority approval for ribociclib; and
 - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to ribociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
 - 2.3 Treatment must be used in combination with an endocrine partner; and
 - 2.4 There is no evidence of progressive disease since initiation of ribociclib.

Continuation

Re-assessment required after 12 months

Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of palbociclib.

PAZOPANIB – **Restricted** see terms [below](#)

↓ Tab 200 mg – 5% DV May-25 to 2027.....	172.88	30	Pazopanib ADVZ Pazopanib Teva
↓ Tab 400 mg – 5% DV May-25 to 2027.....	464.00	30	Pazopanib Teva

→ **Restricted (RS2089)**

Initiation

Re-assessment required after 3 months

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
 - 1.2 Either:
 - 1.2.1 The patient is treatment naive; or
 - 1.2.2 The patient has only received prior cytokine treatment; and
 - 1.3 The patient has an ECOG performance score of 0-2; and
The patient has intermediate or poor prognosis defined as:
 - 1.4 Any of the following:
 - 1.4.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 1.4.2 Haemoglobin level < lower limit of normal; or
 - 1.4.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 1.4.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 1.4.5 Karnofsky performance score of less than or equal to 70; or
 - 1.4.6 2 or more sites of organ metastasis; or
- 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and
 - 2.2 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on sunitinib; and
 - 2.4 Pazopanib to be used for a maximum of 3 months.

Continuation

Re-assessment required after 3 months

No evidence of disease progression.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

RIBOCICLIB – **Restricted** see terms [below](#)

⚡ Tab 200 mg	1,883.00	21	Kisqali
	3,767.00	42	Kisqali
	5,650.00	63	Kisqali

➡ **Restricted** (RS2131)

Initiation

Re-assessment required after 6 months

Either:

- 1 All of the following:
 - 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
 - 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
 - 1.3 Patient has an ECOG performance score of 0-2; and
 - 1.4 Either:
 - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
 - 1.4.2 Both:
 - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
 - 1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 1.5 Treatment to be used in combination with an endocrine partner; and
 - 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has an active Special Authority approval for palbociclib; and
 - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
 - 2.3 Treatment must be used in combination with an endocrine partner; and
 - 2.4 There is no evidence of progressive disease since initiation of palbociclib.

Continuation

Re-assessment required after 12 months

Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of ribociclib.

RUXOLITINIB – **Restricted** see terms [below](#)

⚡ Tab 5 mg	2,500.00	56	Jakavi
⚡ Tab 10 mg	5,000.00	56	Jakavi
⚡ Tab 15 mg	5,000.00	56	Jakavi
⚡ Tab 20 mg	5,000.00	56	Jakavi

➡ **Restricted** (RS1726)

Initiation

Haematologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and

2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and

3 A maximum dose of 20 mg twice daily is to be given.

Continuation

Relevant specialist or medical practitioner on the recommendation of a Relevant specialist

Re-assessment required after 12 months

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB – **Restricted** see terms [below](#)

↓ Cap 12.5 mg – 5% DV Mar-26 to 2027	103.11	28	Sunitinib Rex
↓ Cap 25 mg – 5% DV Mar-26 to 2027	203.15	28	Sunitinib Rex
↓ Cap 50 mg – 5% DV May-26 to 2027	343.19	28	Sunitinib Rex

→ **Restricted (RS2202)**

Initiation – RCC

Re-assessment required after 4 months

Both:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 The patient has not previously received funded sunitinib.

Continuation – RCC

Re-assessment required after 4 months

No evidence of disease progression.

Initiation – GIST

Re-assessment required after 3 months

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Continuation – GIST

Re-assessment required after 6 months

Both:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Sunitinib is to be discontinued at disease progression.

Note: GIST - It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

TRAMETINIB – **Restricted** see terms [below](#)

↓ Tab 0.5 mg	2,370.32	30	Mekinist
↓ Tab 2 mg	9,481.29	30	Mekinist

→ **Restricted (RS2147)**

Initiation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

1 Either:

1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or

1.2 Both:

1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and

1.2.2 Adjuvant treatment with trametinib is required; and

2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and

3 Treatment must be adjuvant to complete surgical resection; and

4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and

5 The individual has a confirmed BRAF mutation; and

6 Trametinib must be administered in combination with dabrafenib; and

7 The individual has ECOG performance score 0-2.

Notes:

a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition

b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Continuation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

Any of the following:

1 All of the following:

1.1 No evidence of disease recurrence; and

1.2 Trametinib must be administered in combination with dabrafenib; and

1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or

2 All of the following:

2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and

2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and

2.3 The individual meets initiation criteria for trametinib for unresectable or metastatic melanoma; or

3 All of the following:

3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and

3.2 The individual has received a BRAF/MEK inhibitor for unresectable or metastatic melanoma; and

3.3 The individual meets continuation criteria for trametinib for unresectable or metastatic melanoma.

Initiation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

1 The individual has metastatic or unresectable melanoma (excluding uveal melanoma) stage III or IV; and

2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and

3 The individual has ECOG performance score 0-2; and

4 The individual has confirmed BRAF mutation; and

5 Trametinib must be administered in combination with dabrafenib; and

6 Any of the following:

6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or

6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or

6.3 All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
- 6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
- 6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MEK inhibitor.

Continuation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

Both:

- 1 Any of the following:
 - 1.1 The individual's disease has had a complete response to treatment; or
 - 1.2 The individual's disease has had a partial response to treatment; or
 - 1.3 The individual has stable disease with treatment; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Taxanes

DOCETAXEL

Inj 10 mg per ml, 8 ml vial.....	24.91	1	DBL Docetaxel
----------------------------------	-------	---	---------------

PACLITAXEL

Inj 6 mg per ml, 16.7 ml vial.....	19.59	1	Anzatax
Inj 6 mg per ml, 50 ml vial.....	37.89	1	Anzatax

Treatment of Cytotoxic-Induced Side Effects

CALCIUM FOLINATE

Tab 15 mg	135.33	10	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 5 ml vial.....	112.20	5	Eurofolic
Inj 10 mg per ml, 10 ml vial.....	163.35	5	Eurofolic
Inj 10 mg per ml, 30 ml vial			
Inj 10 mg per ml, 35 ml vial			
Inj 10 mg per ml, 100 ml vial.....	139.48	1	Eurofolic

DEXRAZOXANE – **Restricted** see terms [below](#)

↓ Inj 500 mg

e.g. Cardioxane

→ **Restricted (RS1695)**

Initiation

Medical oncologist, paediatric oncologist, haematologist or paediatric haematologist

All of the following:

- 1 Patient is to receive treatment with high dose anthracycline given with curative intent; and
- 2 Based on current treatment plan, patient's cumulative lifetime dose of anthracycline will exceed 250mg/m2 doxorubicin equivalent or greater; and
- 3 Dexrazoxane to be administered only whilst on anthracycline treatment; and
- 4 Either:
 - 4.1 Treatment to be used as a cardioprotectant for a child or young adult; or
 - 4.2 Treatment to be used as a cardioprotectant for secondary malignancy.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MESNA			
Tab 400 mg	314.00	50	Uromitexan
Tab 600 mg	448.50	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule	177.45	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule	407.40	15	Uromitexan

Vinca Alkaloids

VINBLASTINE SULPHATE			
Inj 1 mg per ml, 10 ml vial	270.37	5	Hospira
VINCRIStINE SULPHATE			
Inj 1 mg per ml, 1 ml vial	74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial	102.73	5	DBL Vincristine Sulfate
VINOReLBINE			
Cap 20 mg – 5% DV Feb-26 to 2028	32.10	1	Vinorelbine Te Arai
Cap 30 mg – 5% DV Feb-26 to 2028	42.80	1	Vinorelbine Te Arai
Cap 80 mg – 5% DV Feb-26 to 2028	80.00	1	Vinorelbine Te Arai
Inj 10 mg per ml, 1 ml vial			
Inj 10 mg per ml, 5 ml vial			

Endocrine Therapy

ABIRATERONE ACETATE – Restricted see terms below			
↓ Tab 250 mg	4,276.19	120	Zytiga
→ Restricted (RS1888)			

Initiation

Medical oncologist, radiation oncologist or urologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Continuation

Medical oncologist, radiation oncologist or urologist

Re-assessment required after 6 months

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- The treatment remains appropriate and the patient is benefiting from treatment.

Continuation – pandemic circumstances

Re-assessment required after 6 months

All of the following:

- The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- Abiraterone acetate to be discontinued at progression; and
- No initiation of taxane chemotherapy with abiraterone; and
- The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE

Tab 50 mg 4.18 28 Binarex

FLUTAMIDE

Tab 250 mg 119.50 100 Flutamin

FULVESTRANT – Restricted see terms [below](#)

↓ Inj 50 mg per ml, 5 ml prefilled syringe – 5% DV May-26 to 2028 181.00 2 **Fulvestrant EVER Pharma**

→ **Restricted (RS1732)**

Initiation

Medical oncologist

Re-assessment required after 6 months

All of the following:

- Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- Treatment to be given at a dose of 500 mg monthly following loading doses; and
- Treatment to be discontinued at disease progression.

Continuation

Medical oncologist

Re-assessment required after 6 months

All of the following:

- Treatment remains appropriate and patient is benefitting from treatment; and
- Treatment to be given at a dose of 500 mg monthly; and
- No evidence of disease progression.

OCTREOTIDE

Inj 100 mcg per ml, 1 ml vial 48.50 5 Omega
Inj 50 mcg per ml, 1 ml vial 27.58 5 Omega
Inj 500 mcg per ml, 1 ml vial 113.10 5 Omega
Inj 50 mcg per ml, 1 ml ampoule 27.58 5 Max Health
Inj 100 mcg per ml, 1 ml ampoule 32.71 5 Max Health
Inj 500 mcg per ml, 1 ml ampoule 113.10 5 Max Health

TAMOXIFEN CITRATE

Tab 10 mg – 5% DV Dec-26 to 2029 15.00 60 **Tamoxifen Sandoz**
Tab 20 mg – 5% DV Dec-26 to 2029 5.32 60 **Tamoxifen Sandoz**

Aromatase Inhibitors

ANASTROZOLE

Tab 1 mg 4.39 30 Anatrole

EXEMESTANE

Tab 25 mg 9.86 30 Pfizer Exemestane

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LETROZOLE			
Tab 2.5 mg – 5% DV Dec-24 to 2027	4.67	30	Letrole

Long-acting Somatostatin Analogues

➡ Restricted (RS2100)

Initiation – Malignant bowel obstruction

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has not been successful; and
- 3 Treatment to be given for up to 4 weeks.

Note: Indications marked with * are unapproved indications

Initiation – acromegaly

Re-assessment required after 3 months

All of the following:

- 1 The patient has acromegaly; and
- 2 Either:
 - 2.1 Treatment with surgery and radiotherapy is not suitable or was unsuccessful; or
 - 2.2 Treatment is for an interim period while awaiting the beneficial effects of radiotherapy; and
- 3 Treatment with a dopamine agonist has been unsuccessful.

Continuation – acromegaly

Without reassessment for applications where IGF1 levels have decreased since starting treatment.

Note: In patients with acromegaly, treatment should be discontinued if IGF1 levels have not decreased 3 months after treatment. In patients treated with radiotherapy treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following treatment withdrawal for at least 4 weeks.

Initiation – Other indications

Any of the following:

- 1 VIPomas and glucagonomas - for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:
 - 2.2.1 Surgery has been unsuccessful; or
 - 2.2.2 Patient has metastatic disease after treatment with H2 antagonist or proton pump inhibitors has been unsuccessful; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has not been successful; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Initiation – pre-operative acromegaly

Limited to 12 months treatment

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

Notes: Indications marked with * are unapproved indications

The use of a long-acting somatostatin analogue in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded under Special Authority

LANREOTIDE – **Restricted** see terms [on the previous page](#)

† Inj 60 mg per 0.5 ml, 0.5 ml syringe – 5% DV Aug-25 to 2027	382.77	1	Mytolac
	1,543.79		Somatuline Autogel
† Inj 90 mg per 0.5 ml, 0.5 ml syringe – 5% DV Sep-25 to 2027	562.92	1	Mytolac
	2,054.40		Somatuline Autogel
† Inj 120 mg per 0.5 ml, 0.5 ml syringe – 5% DV Aug-25 to 2027	646.70	1	Mytolac
	2,570.44		Somatuline Autogel

OCTREOTIDE LONG-ACTING – **Restricted** see terms [on the previous page](#)

† Inj depot 10 mg prefilled syringe – 5% DV Dec-24 to 2027	438.40	1	Sandostatin LAR
† Inj depot 20 mg prefilled syringe – 5% DV Dec-24 to 2027	583.70	1	Sandostatin LAR
† Inj depot 30 mg prefilled syringe – 5% DV Dec-24 to 2027	670.80	1	Sandostatin LAR

Imaging Agents

AMINOLEVULINIC ACID HYDROCHLORIDE – **Restricted** see terms [below](#)

↓ Powder for oral soln, 30 mg per ml, 1.5 g vial	4,400.00	1	Gliolan
	44,000.00	10	Gliolan

→ **Restricted (RS1565)**

Initiation – high grade malignant glioma

All of the following:

- 1 Patient has newly diagnosed, untreated, glioblastoma multiforme; and
- 2 Treatment to be used as adjuvant to fluorescence-guided resection; and
- 3 Patient's tumour is amenable to complete resection.

Immunosuppressants

Calcineurin Inhibitors

CICLOSPORIN

Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml	198.13	50 ml	Neoral
Inj 50 mg per ml, 5 ml ampoule	276.30	10	Sandimmun

TACROLIMUS – **Restricted** see terms [below](#)

↓ Cap 0.5 mg	49.60	100	Tacrolimus Sandoz
↓ Cap 0.75 mg	99.30	100	Tacrolimus Sandoz
↓ Cap 1 mg	84.30	100	Tacrolimus Sandoz
↓ Cap 5 mg	248.20	50	Tacrolimus Sandoz
↓ Inj 5 mg per ml, 1 ml ampoule			

→ **Restricted (RS2110)**

Initiation – organ transplant recipients

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 For use in organ transplant recipients; or
- 2 The individual is receiving induction therapy for an organ transplant.

Initiation – non-transplant indications*

Any specialist

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Either:
 - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
 - 2.2 Patient is a child with nephrotic syndrome*.

Note: Indications marked with * are unapproved indications

Fusion Proteins

ETANERCEPT – **Restricted** see terms [below](#)

⚡ Inj 25 mg autoinjector	690.00	4	Enbrel
⚡ Inj 25 mg vial	690.00	4	Enbrel
⚡ Inj 50 mg autoinjector	1,050.00	4	Enbrel
⚡ Inj 50 mg syringe	1,050.00	4	Enbrel

➡ **Restricted (RS2203)**

Initiation – arthritis - polyarticular course juvenile idiopathic

Re-assessment required after 6 months

Any of the following:

- 1 Both:
 - 1.1 Patient has had a Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 At least 5 active joints and at least 3 joints with pain, tenderness or a limited range of motion after a 3-month trial of methotrexate at the maximum tolerated dose, unless contraindicated; or
- 3 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate at the maximum tolerated dose, unless contraindicated; or
- 4 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Continuation – arthritis - polyarticular course juvenile idiopathic

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
- 2 On subsequent reapplications, at least a continuing 30% improvement in active joint count from baseline.

Initiation – arthritis - oligoarticular course juvenile idiopathic

Re-assessment required after 6 months

Any of the following:

- 1 Both:
 - 1.1 Patient has had a Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 At least 2 active joints with pain, tenderness or a limited range of motion, after a 3-month trial of methotrexate at the maximum tolerated dose, unless contraindicated; or
- 3 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate at the maximum tolerated dose, unless contraindicated.

Continuation – arthritis - oligoarticular course juvenile idiopathic

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
- 2 On subsequent reapplications, at least a continuing 30% improvement in active joint count from baseline.

Initiation – arthritis - rheumatoid

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had a Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiologic imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive); and
 - 2.2 Patient has received insufficient benefit from at least 3 months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has received insufficient benefit from at least 3 months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.4 Either:
 - 2.4.1 Patient has received insufficient benefit from at least 3 months of methotrexate in combination with the maximum tolerated dose of ciclosporin, unless contraindicated; or
 - 2.4.2 Patient has received insufficient benefit from at least 3 months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate, unless contraindicated; and
 - 2.5 Either:
 - 2.5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 joints; or
 - 2.5.2 Patient has persistent symptoms of poorly controlled and active disease in at least 4 joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Continuation – arthritis - rheumatoid

Re-assessment required after 2 years

Both:

- 1 Either:
 - 1.1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
 - 1.2 On subsequent reapplications, at least a continuing 30% improvement in active joint count from baseline; and
- 2 Maximum dose 50 mg every 7 days.

Initiation – ankylosing spondylitis

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had a Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.2.1 Patient has experienced intolerable side effects; or
- 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiologic imaging; and
 - 2.4 Disease has not responded adequately to treatment with two or more NSAID (unless contraindicated) while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 BASDAI score of at least 6 on a 10-point scale completed after 3-month exercise trial before ceasing any previous pharmacological treatment and not more than 1 month before the application.

Continuation – ankylosing spondylitis

Re-assessment required after 2 years

Both:

- 1 BASDAI has improved from pre-treatment baseline either by at least 4 points on a 10-point scale, or by at least 50%; and
- 2 Maximum dose 50 mg every 7 days.

Initiation – arthritis - psoriatic

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had a Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has received insufficient benefit from at least 3 months of methotrexate at maximum tolerated dose unless contraindicated; and
 - 2.2 Patient received insufficient benefit from at least 3 months of sulfasalazine or leflunomide at maximum tolerated dose unless contraindicated; and
 - 2.3 Either:
 - 2.3.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 joints; or
 - 2.3.2 Patient has persistent symptoms of poorly controlled and active disease in at least 4 joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.4 Any of the following:
 - 2.4.1 Patient has CRP greater than 15 mg/L measured within one month before the application; or
 - 2.4.2 Patient has an ESR greater than 25 mm per hour measured within one month before the application; or
 - 2.4.3 ESR and CRP not measured as patient is receiving prednisone therapy greater than 5 mg per day received for more than 3 months.

Continuation – arthritis - psoriatic

Re-assessment required after 2 years

Both:

- 1 Either:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
- 1.2 At least a continuing 30% improvement in active joint count from baseline; and

2 Maximum dose 50 mg every 7 days.

Initiation – plaque psoriasis, prior TNF use

Limited to 6 months treatment

All of the following:

- 1 Patient has had a Special Authority approval for adalimumab for plaque psoriasis; and
- 2 Either:
 - 2.1 Patient has experienced intolerable side effects; or
 - 2.2 Patient has received insufficient benefit to meet the renewal criteria for adalimumab for plaque psoriasis; and
- 3 Patient must be reassessed for continuation after 3 doses.

Initiation – plaque psoriasis, treatment-naïve

Limited to 6 months treatment

All of the following:

- 1 Any of the following:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10; or
 - 1.2 Patient has plaque psoriasis of the face, or palm of a hand, or sole of a foot; or
 - 1.3 Patient has localised genital or flexural plaque psoriasis with a DLQI score greater than 10; and
- 2 Patient has received insufficient benefit from (see Note), or has experienced intolerable side effects from, at least 3 of the following at maximum tolerated doses (unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or DLQI assessment has been completed for the most recent prior treatment course within 1 month of stopping that treatment; and
- 4 The most recent PASI or DLQI assessment is within 1 month of before the application.

Note: "Insufficient benefit" is defined as: for whole body plaque psoriasis, a PASI score of greater than 10, for plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. As assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation – plaque psoriasis

Re-assessment required after 2 years

Both:

- 1 Any of the following:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Patient has a PASI score which is reduced by 75% or more, or is sustained at this level, compared with the pre-treatment baseline; or
 - 1.1.2.2 Patient has a DLQI improvement of 5 or more, compared with the pre-treatment baseline; or
 - 1.2 Both:
 - 1.2.1 Patient had plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, compared to the pre-treatment baseline; or
 - 1.2.2.2 Patient has a reduction of 75% or more in the skin area affected, or sustained at this level, compared to the pre-treatment baseline; or
 - 1.3 Both:
 - 1.3.1 Patient had localised genital or flexural plaque psoriasis at the start of treatment; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

1.3.2 Either:

1.3.2.1 Patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, compared to the pre-treatment baseline; or

1.3.2.2 Patient has a DLQI improvement of 5 or more, compared to the pretreatment baseline; and

2 Maximum 50 mg every 7 days.

Initiation – pyoderma gangrenosum*

Both:

1 Patient has received insufficient benefit from 3 months of conventional therapy including a minimum of 3 pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate). Where conventional pharmaceuticals are contraindicated, a 3-month trial has occurred of those that are not contraindicated; and

2 Maximum of 8 doses every 4 months.

Note: Indications marked with * are unapproved indications.

Initiation – Stevens-Johnson syndrome/toxic epidermal necrolysis

Patient has confirmed or suspected Stevens-Johnson syndrome or toxic epidermal necrolysis.

Initiation – Still's disease - adult-onset (AOSD)

Either:

1 Both:

1.1 Patient has had a Special Authority approval for adalimumab or tocilizumab for AOSD; and

1.2 Either:

1.2.1 Patient has experienced intolerable side effects; or

1.2.2 Patient has received insufficient benefit to meet the new renewal criteria from at least a 3-month trial of adalimumab or tocilizumab; or

2 All of the following:

2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and

2.2 Patient has tried and received insufficient benefit from at least 6 months of corticosteroids at a dose of at least 0.5 mg/kg prednisone-equivalents, NSAIDs and methotrexate, unless contraindicated; and

2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initiation – undifferentiated spondyloarthritis

Re-assessment required after 6 months

All of the following:

1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least 4 joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

2 Patient has received insufficient benefit from at least 3 months of each of methotrexate, sulfasalazine, and leflunomide at maximum tolerated doses, unless contraindicated; and

3 Any of the following:

3.1 Patient has a CRP level greater than 15 mg/L measured within one month before the application; or

3.2 Patient has an ESR greater than 25 mm per hour measured within one month before the application; or

3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy greater than 5 mg per day received for more than three months.

Note: Indications marked with * are unapproved indications.

Continuation – undifferentiated spondyloarthritis

Re-assessment required after 2 years

Both:

1 Either:

1.1 Following initial treatment, the patient has experienced at least a 50% decrease in active joint count from baseline; or

1.2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- Maximum 50 mg dose every 7 days.

Monoclonal Antibodies

ABCIXIMAB – **Restricted** see terms [below](#)

↓ Inj 2 mg per ml, 5 ml vial

→ **Restricted** ([RS1202](#))

Initiation

Either:

- For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
- For use in patients undergoing intra-cranial intervention.

ADALIMUMAB (AMGEVITA) – **Restricted** see terms [below](#)

↓ Inj 20 mg per 0.4 ml prefilled syringe..... 190.00 1 Amgevita

↓ Inj 40 mg per 0.8 ml prefilled pen 375.00 2 Amgevita

↓ Inj 40 mg per 0.8 ml prefilled syringe..... 375.00 2 Amgevita

→ **Restricted** ([RS2140](#))

Initiation – Behcet's disease - severe

Any relevant practitioner

Both:

- The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- Either:
 - The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - The patient has severe gastrointestinal, rheumatological and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initiation – Hidradenitis suppurativa

Dermatologist

Re-assessment required after 4 months

All of the following:

- Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- Patient has 3 or more active lesions; and
- The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Continuation – Hidradenitis suppurativa

Any relevant practitioner

Re-assessment required after 2 years

Both:

- The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- The patient has a DLQI improvement of 4 or more from baseline.

Initiation – Plaque psoriasis - severe chronic

Dermatologist

Re-assessment required after 4 months

Either:

- Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or (DLQI) assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Continuation – Plaque psoriasis - severe chronic

Re-assessment required after 2 years

Any of the following:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has experienced a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
- 3 Both:
 - 3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
 - 3.2 Either:
 - 3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

continued...

	Price		Brand or
(ex man.	excl. GST)		Generic
\$	Per		Manufacturer

continued...

Initiation – pyoderma gangrenosum

Dermatologist

Both:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initiation – Crohn's disease - adults

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease - adults

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initiation – Crohn's disease - children

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease - children

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Initiation – Crohn's disease - fistulising

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Continuation – Crohn's disease - fistulising

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initiation – Ocular inflammation - chronic

Any relevant practitioner

Re-assessment required after 4 months

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Continuation – Ocular inflammation - chronic

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initiation – Ocular inflammation - severe

Any relevant practitioner

Re-assessment required after 4 months

Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Continuation – Ocular inflammation - severe

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initiation – ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Continuation – ankylosing spondylitis

Any relevant practitioner

Re-assessment required after 2 years

For applications where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – Arthritis - oligoarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Continuation – Arthritis - oligoarticular course juvenile idiopathic

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - polyarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

continued...

	Price	Brand or
(ex man. excl. GST)	\$	Generic
	Per	Manufacturer

continued...

Continuation – Arthritis - polyarticular course juvenile idiopathic

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - psoriatic

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation – Arthritis - psoriatic

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initiation – Arthritis - rheumatoid

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin (unless contraindicated); or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomid (unless contraindicated) alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Continuation – Arthritis - rheumatoid

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initiation – Still's disease - adult-onset (AOSD)

Rheumatologist

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for (AOSD); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initiation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Continuation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on biologic therapy.

Initiation – undifferentiated spondyloarthritis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulphasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Continuation – undifferentiated spondyloarthritis

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initiation – inflammatory bowel arthritis – axial

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and

continued...

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Continuation – inflammatory bowel arthritis – axial

Any relevant practitioner

Re-assessment required after 2 years

Where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initiation – inflammatory bowel arthritis – peripheral

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulphasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation – inflammatory bowel arthritis – peripheral

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) – Restricted see terms [below](#)

⚡ Inj 20 mg per 0.2 ml prefilled syringe.....	595.50	2	Humira
⚡ Inj 40 mg per 0.4 ml prefilled syringe.....	595.50	2	Humira
⚡ Inj 40 mg per 0.4 ml prefilled pen	595.50	2	HumiraPen

➡ **Restricted (RS1922)**

Initiation – Behcet's disease – severe

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Behcet's disease – severe

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Hidradenitis suppurativa

Dermatologist or Practitioner on the recommendation of a dermatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Continuation – Hidradenitis suppurativa

Dermatologist or Practitioner on the recommendation of a dermatologist

Re-assessment required after 6 months

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initiation – Psoriasis - severe chronic plaque

Dermatologist or Practitioner on the recommendation of a dermatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Psoriasis - severe chronic plaque

Dermatologist or Practitioner on the recommendation of a dermatologist

Re-assessment required after 6 months

Both:

- 1 Either:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Either:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Pyoderma gangrenosum

Dermatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

Continuation – Pyoderma gangrenosum

Dermatologist

Re-assessment required after 6 months

Both:

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initiation – Crohn's disease - adult

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Crohn's disease - adult

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Crohn's disease - children

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Crohn's disease - children

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Crohn's disease - fistulising

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Crohn's disease - fistulising

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Ocular inflammation – chronic

Any relevant practitioner

Re-assessment required after 12 months

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Ocular inflammation – chronic

Any relevant practitioner

Re-assessment required after 12 months

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Ocular inflammation – severe

Any relevant practitioner

Re-assessment required after 12 months

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

response to a change in treatment regimen; or

- 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Ocular inflammation – severe

Any relevant practitioner

Re-assessment required after 12 months

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – ankylosing spondylitis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – ankylosing spondylitis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Arthritis – oligoarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – Arthritis – oligoarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

For patients that demonstrate at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - polyarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Continuation – Arthritis - polyarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

For patients that demonstrate at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - psoriatic

Named specialist or rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Arthritis - psoriatic

Named specialist or rheumatologist

Re-assessment required after 6 months

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Arthritis – rheumatoid

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Continuation – Arthritis – rheumatoid

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initiation – Still's disease – adult-onset (AOSD)

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Continuation – Still's disease – adult-onset (AOSD)

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT – **Restricted** see terms [below](#)

↓ Inj 40 mg per ml, 0.1 ml vial..... 1,250.00 1 Eylea

➔ **Restricted (RS2148)**

Initiation – Wet Age Related Macular Degeneration

Re-assessment required after 3 months

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

bevacizumab; or

1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and

1.3 There is no structural damage to the central fovea of the treated eye; and

1.4 Patient has not previously been treated with ranibizumab or faricimab for longer than 3 months; or

2 Either:

2.1 Patient has current approval to use ranibizumab or faricimab for treatment of wAMD and was found to be intolerant within 3 months; or

2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Continuation – Wet Age Related Macular Degeneration

Re-assessment required after 12 months

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Initiation – Diabetic Macular Oedema

Re-assessment required after 4 months

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 – 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 6 Patient has not previously been treated with faricimab for longer than 3 months.

Continuation – Diabetic Macular Oedema

Re-assessment required after 12 months

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

BASILIXIMAB – Restricted see terms [below](#)

⚡ Inj 20 mg vial2,560.00 1 Simulect

➡ **Restricted (RS1203)**

Initiation

For use in solid organ transplants.

BENRALIZUMAB – Restricted see terms [below](#)

⚡ Inj 30 mg per ml, 1 ml prefilled pen3,539.00 1 Fasenra

➡ **Restricted (RS1920)**

Initiation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 12 months

All of the following:

- 1 Patient must be aged 12 years or older; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5×10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Continuation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 2 years

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

BEVACIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 25 mg per ml, 4 ml vial – 10% DV Aug-25 to 31 Aug 2028	69.00	1	Vegzelma
⚡ Inj 25 mg per ml, 16 ml vial – 10% DV Aug-25 to 31 Aug 2028	276.00	1	Vegzelma

➡ **Restricted (RS2111)**

Initiation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with bevacizumab, and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
 - 2.2 Patient has preserved liver function (Child-Pugh A); and
 - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
 - 2.4 Any of the following:
 - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
- 2.4.3 Both:
 - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
 - 2.4.3.2 No disease progression since initiation of lenvatinib; and
- 2.5 Patient has an ECOG performance status of 0-2; and
- 2.6 To be given in combination with atezolizumab.

Continuation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months
no evidence of disease progression.

Initiation – advanced or metastatic ovarian cancer

Re-assessment required after 4 months
All of the following:

- 1 Either:
 - 1.1 The patient has FIGO Stage IV epithelial ovarian, fallopian tube, or primary peritoneal cancer; or
 - 1.2 Both:
 - 1.2.1 The patient has previously untreated advanced (FIGO Stage IIIB or IIIC) epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
 - 1.2.2 Either:
 - 1.2.2.1 Debulking surgery is inappropriate; or
 - 1.2.2.2 The cancer is sub-optimally debulked (maximum diameter of any gross residual disease greater than 1cm); and
- 2 Bevacizumab to be administered at a maximum dose of 15 mg/kg every three weeks; and
- 3 18 weeks concurrent treatment with chemotherapy is planned.

Continuation – advanced or metastatic ovarian cancer

Re-assessment required after 4 months
no evidence of disease progression.

Initiation – Recurrent Respiratory Papillomatosis

Re-assessment required after 12 months
All of the following:

- 1 Maximum of 6 doses; and
- 2 The patient has recurrent respiratory papillomatosis; and
- 3 The treatment is for intra-lesional administration.

Continuation – Recurrent Respiratory Papillomatosis

Re-assessment required after 12 months
All of the following:

- 1 Maximum of 6 doses; and
- 2 The treatment is for intra-lesional administration; and
- 3 There has been a reduction in surgical treatments or disease regrowth as a result of treatment.

Initiation – Ocular Conditions

- Either:
- 1 Ocular neovascularisation; or
 - 2 Exudative ocular angiopathy.

BEVACIZUMAB (OCULAR) – **Restricted** see terms [below](#)

⚡ Inj 25 mg per ml, 4 ml vial.....	600.00	1	Avastin
⚡ Inj 25 mg per ml, 16 ml vial			

➡ **Restricted (RS2156)**
Initiation – ocular conditions
Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Ocular neovascularisation; or
- 2 Exudative ocular angiopathy.

BRENTUXIMAB VEDOTIN – Restricted see terms [below](#)

↓ Inj 50 mg vial5,275.18 1 Adcetris

➔ **Restricted (RS2183)**

Initiation – CD30 positive systemic anaplastic large-cell lymphoma

Limited to 12 months treatment

Either:

- 1 Patient is currently on treatment with brentuximab vedotin and met all the following criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has CD30 positive systemic anaplastic large-cell lymphoma; and
 - 2.2 Patient must have histological confirmation of CD30 expression; and
 - 2.3 Patient must not have received prior treatment with curative intent chemotherapy for this condition; and
 - 2.4 Treatment must be in combination with cyclophosphamide, anthracycline, and steroids for a maximum of 8 cycles; and
 - 2.5 Brentuximab vedotin is to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Initiation – relapsed/refractory Hodgkin lymphoma

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and
 - 1.1.2 Patient is ineligible for autologous stem cell transplant; or
 - 1.2 Both:
 - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
 - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Continuation – relapsed/refractory Hodgkin lymphoma

Re-assessment required after 9 months

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

Initiation – relapsed/refractory anaplastic large cell lymphoma

Re-assessment required after 9 months

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – relapsed/refractory anaplastic large cell lymphoma

Re-assessment required after 9 months

All of the following:

- 1 Patient has experienced a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

CETUXIMAB – **Restricted** see terms [below](#)

⚡ Inj 5 mg per ml, 20 ml vial.....	364.00	1	Erbixux
⚡ Inj 5 mg per ml, 100 ml vial.....	1,820.00	1	Erbixux

➡ **Restricted (RS2064)**

Initiation – head and neck cancer, locally advanced

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Cisplatin is contraindicated or has resulted in intolerable side effects; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 To be administered in combination with radiation therapy.

Initiation – colorectal cancer, metastatic

Re-assessment required after 6 months

All of the following:

- 1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and
- 2 There is documentation confirming disease is RAS and BRAF wild-type; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Patient has not received prior funded treatment with cetuximab; and
- 5 Either:
 - 5.1 Cetuximab is to be used in combination with chemotherapy; or
 - 5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

Continuation – colorectal cancer, metastatic

Re-assessment required after 6 months

No evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

FARICIMAB – **Restricted** see terms [below](#)

⚡ Inj 120 mg per ml, 0.24 ml vial.....	1,565.00	1	Vabysmo
--	----------	---	---------

➡ **Restricted (RS2149)**

Initiation – Diabetic macular oedema

Re-assessment required after 4 months

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is nonresponsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 – 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 6 Patient has not previously been treated with aflibercept for longer than 3 months.

Continuation – Diabetic macular oedema

Re-assessment required after 12 months

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Initiation – Wet age related macular degeneration

Re-assessment required after 3 months

All of the following:

- 1 Any of the following:
 - 1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.2 Polypoidal choroidal vasculopathy; or
 - 1.3 Choroidal neovascular membrane from causes other than wet AMD; and
- 2 Either:
 - 2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
- 3 There is no structural damage to the central fovea of the treated eye; and
- 4 Patient has not previously been treated with ranibizumab or aflibercept for longer than 3 months.

Continuation – Wet age related macular degeneration

Re-assessment required after 12 months

Both:

- 1 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 2 There is no structural damage to the central fovea of the treated eye.

GEMTUZUMAB OZOGAMICIN – Restricted see terms [below](#)

↓ Inj 5 mg vial 12,973.00 1 Mylotarg

→ **Restricted (RS1923)**

Initiation

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB – Restricted see terms [below](#)

↓ Inj 100 mg 428.00 1 Remicade

→ **Restricted (RS2178)**

Initiation – Graft vs host disease

Patient has steroid-refractory acute graft vs. host disease of the gut.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Initiation – arthritis - rheumatoid

Re-assessment required after 6 months

All of the following:

- 1 Patient has had a Special Authority approval for adalimumab or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Patient has experienced intolerable side effects; or
 - 2.2 Patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; and
- 3 Following initial induction doses, maximum dose 3mg/kg every 8 weeks.

Continuation – arthritis - rheumatoid

Re-assessment required after 2 years

Both:

- 1 Either:
 - 1.1 Following initial treatment, the patient has experienced at least a 50% decrease in active joint count from baseline; or
 - 1.2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline; and
- 2 Maximum dose 3 mg/kg every 8 weeks.

Initiation – ankylosing spondylitis

Re-assessment required after 6 months

All of the following:

- 1 Patient has had a Special Authority approval for adalimumab or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 Patient has experienced intolerable side effects; or
 - 2.2 Patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; and
- 3 Following initial induction doses, maximum dose 5mg/kg every 6-8 weeks.

Continuation – ankylosing spondylitis

Re-assessment required after 2 years

Both:

- 1 BASDAI has improved from pre-treatment baseline either by at least 4 points on a 10-point scale, or by at least 50%; and
- 2 Maximum dose 5 mg/kg every 6-8 weeks.

Initiation – arthritis - psoriatic

Re-assessment required after 6 months

All of the following:

- 1 Patient has had a Special Authority approval for adalimumab or etanercept or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 Patient has experienced intolerable side effects; or
 - 2.2 Patient has received insufficient benefit to meet the renewal criteria for psoriatic arthritis; and
- 3 Following initial induction doses, maximum dose 5mg/kg every 8 weeks.

Continuation – arthritis - psoriatic

Re-assessment required after 2 years

Both:

- 1 Either:
 - 1.1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
 - 1.2 At least a continuing 30% improvement in active joint count from baseline; and
- 2 Maximum dose 5 mg/kg every 8 weeks.

Initiation – ocular inflammation - severe*

Re-assessment required after 4 months

Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

1 Both:

1.1 Patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and

1.2 Either:

1.2.1 Patient has experienced intolerable side effects; or

1.2.2 Patient has received insufficient benefit to meet the renewal criteria for severe ocular inflammation; or

2 Both:

2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and

2.2 Any of the following:

2.2.1 Treatment with high-dose IV corticosteroids followed by high dose oral corticosteroids has been ineffective at controlling symptoms; or

2.2.2 Patient developed new inflammatory symptoms while receiving high dose corticosteroids; or

2.2.3 Patient is aged under 8 years and treatment with high dose oral corticosteroids and other immunosuppressants has been ineffective at controlling symptoms; or

2.2.4 High dose corticosteroids are contraindicated.

Note: Indications marked with * are unapproved indications.

Continuation – ocular inflammation - severe*

Re-assessment required after 2 years

Any of the following:

1 Patient has received a good clinical response following 3 initial doses; or

2 Following each 2 year treatment period, the patient has experienced a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

3 Following each 2 year treatment period, the patient has a sustained corticosteroid sparing effect, allowing reduction in prednisone to < 10mg daily, or corticosteroid drops less than twice daily if under 18 years old.

Notes: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Indications marked with * are unapproved indications.

Initiation – ocular inflammation - chronic*

Re-assessment required after 4 months

Either:

1 Both:

1.1 Patient has had a Special Authority approval for adalimumab for chronic ocular inflammation; and

1.2 Either:

1.2.1 Patient has experienced intolerable side effects; or

1.2.2 Patient has received insufficient benefit to meet the renewal criteria for chronic ocular inflammation; or

2 Both:

2.1 Patient has severe uveitis with a severe risk of vision loss uncontrolled by treatment with corticosteroids and other immunosuppressants; and

2.2 Any of the following:

2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has been ineffective or are contraindicated; or

2.2.2 Patient is under 18 years and treatment with methotrexate has been ineffective, is contraindicated or is not tolerated at a therapeutic dose; or

2.2.3 Patient is under 8 years and treatment with corticosteroids or methotrexate has been ineffective, is contraindicated or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Note: Indications marked with * are unapproved indications.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – ocular inflammation - chronic*

Re-assessment required after 2 years

Any of the following:

- 1 Patient has received a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has experienced a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained corticosteroid sparing effect, allowing reduction in prednisone to < 10mg daily, or corticosteroid drops less than twice daily if under 18 years old.

Notes: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Indications marked with * are unapproved indications.

Initiation – Pulmonary sarcoidosis

Both:

- 1 Patient has life-threatening pulmonary sarcoidosis that is refractory to other treatments; and
- 2 Treatment is to be prescribed by, or has been recommended by, a physician with expertise in the treatment of pulmonary sarcoidosis.

Initiation – Crohn's disease (adults)

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease (adults)

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – Crohn's disease (children)

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease (children)

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complete peri-anal fistula.

Continuation – fistulising Crohn's disease

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – acute fulminant ulcerative colitis

Gastroenterologist

Limited to 6 weeks treatment

Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – fulminant ulcerative colitis

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patients SCCAI is greater than or equal to 4; or
 - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Continuation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 1.2 The PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – plaque psoriasis

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient had a Special Authority approval for adalimumab, etanercept or secukinumab for plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for plaque psoriasis; or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" plaque psoriasis with a PASI score of greater than 10; or
 - 2.1.2 Patient has plaque psoriasis of the face, or palm of a hand or sole of a foot; or
 - 2.1.3 Patient has localised genital or flexural plaque psoriasis with a DLQI score greater than 10; and
 - 2.2 Patient has received insufficient benefit (see Note) or has experienced intolerable side effects from at least 3 of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

- 2.3 A PASI assessment has been completed for the most recent prior treatment course within 1 month of stopping that treatment; and
- 2.4 The most recent PASI assessment is within 1 month before the application.

Note: "Insufficient benefit" is defined as: for whole body plaque psoriasis, a PASI score of greater than 10, for plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation – plaque psoriasis

Re-assessment required after 2 years

Both:

- 1 Any of the following:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" plaque psoriasis at the start of treatment; and
 - 1.1.2 Patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab baseline; or
 - 1.2 Both:
 - 1.2.1 Patient had plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the pre-infliximab baseline; or
 - 1.2.2.2 Patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab baseline; or
 - 1.3 Both:
 - 1.3.1 Patient had localised genital or flexural plaque psoriasis at the start of treatment; and
 - 1.3.2 Either:
 - 1.3.2.1 Patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline; or
 - 1.3.2.2 Patient has a DLQI improvement of 5 or more, as compared to the pre-infliximab baseline; and
- 2 Maximum dose 5 mg/kg every 8 weeks.

Initiation – neurosarcoidosis

Neurologist

Re-assessment required after 18 months

All of the following:

- 1 Biopsy consistent with diagnosis of neurosarcoidosis; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Continuation – neurosarcoidosis

Neurologist

Re-assessment required after 18 months

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.2 There has been a marked reduction in prednisone dose; and
- 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomatology.

Initiation – Behcet disease

All of the following:

- 1 Patient has severe Behcet disease which is significantly impacting their quality of life; and
- 2 Either:
 - 2.1 Patient has severe ocular, neurological and/or vasculitic symptoms and has received insufficient benefit from 1 or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 Patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has received insufficient benefit from 2 or more treatments appropriate for the particular symptom(s); and
- 3 Following initial loading doses, maximum dose 5mg/kg every 8 weeks.

Initiation – pyoderma gangrenosum*

Both:

- 1 Patient has received insufficient benefit from 3 months of conventional therapy including a minimum of 3 pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate). Where conventional pharmaceuticals are contraindicated, a 3-month trial has occurred of those that are not contraindicated; and
- 2 Maximum of 8 doses every 4 months.

Note: Indications marked with * are unapproved indications.

Initiation – Inflammatory bowel arthritis (axial)

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not experienced an adequate response to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Continuation – Inflammatory bowel arthritis (axial)

Re-assessment required after 2 years

Where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initiation – Inflammatory bowel arthritis (peripheral)

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- application; or
- 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation – Inflammatory bowel arthritis (peripheral)

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

Initiation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and
- 2 The individual has received insufficient benefit from use of corticosteroids; and
- 3 Infliximab is to be administered at up to 5mg/kg for up to four doses.

Continuation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

Both:

- 1 The individual has shown clinical improvement and ongoing treatment is required; and
- 2 Infliximab is to be administered at up to 5mg/kg for up to a total of 8 doses.

Note: Indications marked with * are unapproved indications.

INOTUZUMAB OZOGAMICIN – **Restricted** see terms [below](#)

↓ Inj 1 mg vial 14,457.00 1 Responsa

➔ **Restricted (RS2112)**

Initiation

Re-assessment required after 4 months

All of the following:

- 1 Patient has relapsed or refractory CD22-positive B-cell acute lymphoblastic leukaemia/lymphoma, including minimal residual disease; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient has Philadelphia chromosome positive B-Cell ALL; and
 - 3.1.2 Patient has previously received a tyrosine kinase inhibitor; or
 - 3.2 Patient has received one prior line of treatment involving intensive chemotherapy; and
- 4 Treatment is to be administered for a maximum of 3 cycles.

Continuation

Re-assessment required after 4 months

All of the following:

- 1 Patient is not proceeding to a stem cell transplant; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 Either:

- 2.1 Patient has experienced complete disease response; or
- 2.2 Patient has experienced complete remission with incomplete haematological recovery; and

3 Treatment with inotuzumab ozogamicin is to cease after a total duration of 6 cycles.

MEPOLIZUMAB – Restricted see terms [below](#)

⚡ Inj 100 mg prefilled pen	1,638.00	1	Nucala
⚡ Inj 100 mg vial			

➡ **Restricted (RS2024)**

Initiation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 12 months

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5×10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Continuation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 2 years

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – eosinophilic granulomatosis with polyangiitis

Re-assessment required after 12 months

All of the following:

- 1 The patient has eosinophilic granulomatosis with polyangiitis; and
- 2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and
- 3 Either:
 - 3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or
 - 3.2 Corticosteroids are contraindicated.

Continuation – eosinophilic granulomatosis with polyangiitis

Re-assessment required after 12 months

Patient has no evidence of clinical disease progression.

OBINUTUZUMAB – **Restricted** see terms [below](#)

↓ Inj 25 mg per ml, 40 ml vial.....5,910.00 1 Gazyva

→ **Restricted (RS2192)**

Initiation – previously untreated chronic lymphocytic leukaemia in combination with venetoclax

Either:

- 1 Individual is currently on treatment with obinutuzumab and venetoclax and met all of the following criteria prior to commencing treatment; or
- 2 Both:
 - 2.1 Individual has previously untreated chronic lymphocytic leukaemia; and
 - 2.2 Obinutuzumab is to be administered at a maximum cumulative dose of 8,000 mg and in combination with venetoclax for a maximum of 6 (28-day) cycles of treatment.

Initiation

Limited to 6 months treatment

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* greater than or equal to $1.5 \times 10^9/L$ and platelets greater than or equal to $75 \times 10^9/L$

Initiation – follicular / marginal zone lymphoma

Re-assessment required after 9 months

All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Continuation – follicular / marginal zone lymphoma

Re-assessment required after 24 months

All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 150 mg prefilled syringe.....	450.00	1	Xolair
⚡ Inj 150 mg vial	450.00	1	Xolair

➡ **Restricted (RS1652)**

Initiation – severe asthma

Clinical immunologist or respiratory specialist

Re-assessment required after 6 months

All of the following:

- 1 Patient must be aged 6 years or older ; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or formoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Continuation – severe asthma

Respiratory specialist

Re-assessment required after 6 months

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Initiation – severe chronic spontaneous urticaria

Clinical immunologist or dermatologist

Re-assessment required after 6 months

All of the following:

1 Patient must be aged 12 years or older; and

2 Either:

2.1 Both:

2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and

2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; and

3 Any of the following:

3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or

3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or

3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and

4 Either:

4.1 Treatment to be stopped if inadequate response* following 4 doses; or

4.2 Complete response* to 6 doses of omalizumab.

Continuation – severe chronic spontaneous urticaria

Clinical immunologist or dermatologist

Re-assessment required after 6 months

Either:

1 Patient has previously had a complete response* to 6 doses of omalizumab; or

2 Both:

2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and

2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB – **Restricted** see terms [below](#)

⬇ Inj 100 mg per ml, 1 ml vial..... 1,700.00 1 Synagis

➡ **Restricted (RS2193)**

Initiation

Re-assessment required after 12 months

Both:

1 Palivizumab to be administered during the annual RSV season; and

2 Either:

2.1 All of the following:

2.1.1 Infant was born in the last 12 months; and

2.1.2 Infant was born at less than 32 weeks zero days' gestation; and

2.1.3 Infant is to be administered palivizumab within a single calendar year; or

2.2 Both:

2.2.1 Child was born in the last 24 months; and

2.2.2 Any of the following:

2.2.2.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- ventilatory/respiratory support (see Note A) in the community; or
- 2.2.2.2 Both:
- 2.2.2.2.1 Child has haemodynamically significant heart disease; and
- 2.2.2.2.2 Any of the following:
- 2.2.2.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
- 2.2.2.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
- 2.2.2.2.2.3 Child has severe pulmonary hypertension (see Note C); or
- 2.2.2.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
- 2.2.2.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
- 2.2.2.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Continuation

Re-assessment required after 6 months

All of the following:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Child was born in the last 24 months; and
- 3 Any of the following:
 - 3.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
 - 3.2 Both:
 - 3.2.1 Child has haemodynamically significant heart disease; and
 - 3.2.2 Any of the following:
 - 3.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
 - 3.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
 - 3.2.2.3 Child has severe pulmonary hypertension (see Note C); or
 - 3.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
 - 3.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
 - 3.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Notes:

- a) Ventilatory/respiratory support includes those on home oxygen, CPAP/VPAP and those with tracheostomies in situ managed at home
- b) Child requires/will require heart failure medication, and/or child has significant pulmonary hypertension, and/or infant will require surgical palliation/definitive repair within the next 3 months
- c) Mean pulmonary artery pressure more than 25 mmHg
- d) LV Ejection Fraction less than 40%
- e) Inborn errors of immunity include, but are not limited to, IFNAR deficiencies

PERTUZUMAB – Restricted see terms [below](#)

‡ Inj 30 mg per ml, 14 ml vial.....3,927.00 1 Perjeta

→ **Restricted (RS1995)**

Initiation

Re-assessment required after 12 months

All of the following:

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Continuation

Re-assessment required after 12 months

Either:

- 1 Both:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

PERTUZUMAB WITH TRASTUZUMAB – Restricted see terms [below](#)

↓ Inj 600 mg with trastuzumab 600 mg, 10 ml vial	7,707.00	1	Phesgo
↓ Inj 1,200 mg with trastuzumab 600 mg, 15 ml vial	12,894.00	1	Phesgo

➔ **Restricted (RS2152)**

Initiation

Re-assessment required after 12 months

Either:

- 1 Both:
 - 1.1 The individual has received an initial Special Authority approval for intravenous pertuzumab and trastuzumab for metastatic breast cancer; and
 - 1.2 Pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 600 mg pertuzumab with 600 mg trastuzumab every three weeks (or equivalent); or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 Either:
 - 2.2.1 Patient is chemotherapy treatment naïve; or
 - 2.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 2.3 The patient has good performance status (ECOG grade 0-1); and
 - 2.4 Loading dose of pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 1200 mg pertuzumab with 600 mg trastuzumab, respectively; and

continued...

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

- 2.5 Maintenance doses of pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 600 mg pertuzumab with 600 mg trastuzumab every three weeks (or equivalent); and
- 2.6 Pertuzumab with trastuzumab to be discontinued at disease progression.

Continuation

Re-assessment required after 12 months

Either:

- 1 Both:
 - 1.1 The individual has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
 - 2.1 Individual has previously discontinued treatment with pertuzumab with trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Individual has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pertuzumab with trastuzumab.

RANIBIZUMAB – **Restricted** see terms [below](#)

- ⚡ Inj 10 mg per ml, 0.23 ml vial
- ⚡ Inj 10 mg per ml, 0.3 ml vial

➡ **Restricted (RS2151)**

Initiation – Wet Age Related Macular Degeneration

Re-assessment required after 3 months

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with aflibercept or faricimab for longer than 3 months; or
- 2 Patient has current approval to use aflibercept or faricimab for treatment of wAMD and was found to be intolerant within 3 months.

Continuation – Wet Age Related Macular Degeneration

Re-assessment required after 12 months

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

RITUXIMAB (MABTHERA) – **Restricted** see terms [on the next page](#)

⚡ Inj 10 mg per ml, 10 ml vial.....	1,075.50	2	Mabthera
⚡ Inj 10 mg per ml, 50 ml vial.....	2,688.30	1	Mabthera

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

➔ **Restricted (RS2180)**

Initiation – arthritis - rheumatoid - prior TNF inhibitor use

Limited to 4 months treatment

All of the following:

- 1 Patient has had a Special Authority approval for etanercept or adalimumab for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Patient has experienced intolerable side effects; or
 - 2.2 Following at least a 4 month trial of adalimumab or etanercept, the renewal criteria for rheumatoid arthritis were not met; and
- 3 Maximum of two 1000 mg infusions given two weeks apart.

Initiation – arthritis - rheumatoid - TNF inhibitors contraindicated

Limited to 4 months treatment

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had rheumatoid arthritis (either confirmed by radiologic imaging, or the patient is CCP antibody positive); and
- 3 Disease has not responded to at least three months of methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose, unless contraindicated; and
- 4 Disease has not responded to at least 3 months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses), unless contraindicated; and
- 5 Either:
 - 5.1 Disease has not responded to at least 3 months of methotrexate in combination with the maximum tolerated dose of cyclosporin, unless contraindicated; or
 - 5.2 Disease has not responded to at least 3 months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate, unless contraindicated; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least 4 joints from the following: wrist, elbow, knee, ankle, shoulder, or hip; and
- 7 Either:
 - 7.1 Patient has CRP greater than 15 mg/L measured within one month before the application; or
 - 7.2 CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day received for more than 3 months; and
- 8 Maximum of two 1000 mg infusions given two weeks apart.

Continuation – arthritis - rheumatoid - re-treatment for people who have experienced a partial response to rituximab

Re-assessment required after 12 months

All of the following:

- 1 Any of the following:
 - 1.1 Following the initial course of rituximab the patient experienced between a 30% and 50% decrease in active joint count from baseline; or
 - 1.2 Following the second course of rituximab the patient experienced at least a 50% decrease in active joint count from baseline; or
 - 1.3 Following the third and subsequent courses of rituximab, the patient experienced at least a continuing 30% improvement in active joint count from baseline; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Maximum of two 1000 mg infusions given two weeks apart.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – arthritis - rheumatoid - re-treatment for people who experience a response to rituximab

Re-assessment required after 12 months

All of the following:

- 1 Either:
 - 1.1 Following the initial course of rituximab infusions the patient experienced at least a 50% decrease in active joint count from baseline; or
 - 1.2 Following the second and subsequent courses of rituximab, the patient experienced at least a continuing 30% improvement in active joint count from baseline; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Maximum of two 1000 mg infusions per course given two weeks apart.

RITUXIMAB (RIXIMYO) – **Restricted** see terms [below](#)

⚡ Inj 10 mg per ml, 10 ml vial.....	275.33	2	Riximyo
⚡ Inj 10 mg per ml, 50 ml vial.....	688.20	1	Riximyo

➡ **Restricted (RS2204)**

Initiation – haemophilia with inhibitors

Haematologist

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Continuation – haemophilia with inhibitors

Haematologist

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initiation – post-transplant

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Continuation – post-transplant

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initiation – indolent, low-grade lymphomas or hairy cell leukaemia*

Re-assessment required after 9 months

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Continuation – indolent, low-grade lymphomas or hairy cell leukaemia*

Re-assessment required after 12 months

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initiation – aggressive CD20 positive NHL

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Continuation – aggressive CD20 positive NHL

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Initiation – Chronic lymphocytic leukaemia

Re-assessment required after 12 months

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 5 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Continuation – Chronic lymphocytic leukaemia

Re-assessment required after 12 months

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initiation – severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 8 weeks

All of the following:

- 1 Patient has cold haemagglutinin disease*;
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 8 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*;
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initiation – warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 8 weeks

All of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 8 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initiation – immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 8 weeks

All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 8 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – thrombotic thrombocytopenic purpura (TTP)

Haematologist

All of the following:

- 1 The total rituximab dose per cycle would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks; and
- 2 Each treatment cycle at least 6 months apart; and
- 3 Any of the following:
 - 3.1 Patient has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 3.2 Patient has acute idiopathic TTP* with neurological or cardiovascular pathology; or
 - 3.3 Patient has a history of immune-mediated thrombotic thrombocytopenic purpura, currently in clinical remission but has demonstrated a decline in ADAMTS13 levels or equivalent, consistent with high risk of relapse.

Note: Indications marked with * are unapproved indications.

Initiation – pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Continuation – pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initiation – ANCA associated vasculitis

Re-assessment required after 8 weeks

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Continuation – ANCA associated vasculitis

Re-assessment required after 8 weeks

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

Initiation – treatment refractory systemic lupus erythematosus (SLE)

Rheumatologist or nephrologist

All of the following:

- 1 Patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The condition has been refractory to treatment with corticosteroids at a dose of at least 1 mg/kg unless contraindicated; and
- 3 The condition has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Initial treatment maximum of four 1000 mg infusions; and
- 5 Treatment for relapse following initial partial response to rituximab up to a maximum of two 1000 mg infusions every 6 months.

Note: Indications marked with * are unapproved indications.

Initiation – Antibody-mediated organ transplant rejection

Patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initiation – ABO-incompatible organ transplant

Patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initiation – Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)

Re-assessment required after 8 weeks

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with corticosteroids, ciclosporin, and mycophenolate for at least 3 months for each agent has been ineffective, not tolerated, or is contraindicated; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)

Re-assessment required after 8 weeks

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for > 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initiation – Steroid resistant nephrotic syndrome (SRNS)

Re-assessment required after 8 weeks

All of the following:

- 1 Patient is a child with SRNS* and treatment with corticosteroids, ciclosporin and tacrolimus for at least 3 months for each agent has been ineffective, not tolerated, or is contraindicated; and
- 2 Genetic causes of nephrotic syndrome have been excluded; and
- 3 The total rituximab dose per cycle would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – Steroid resistant nephrotic syndrome (SRNS)

Re-assessment required after 8 weeks

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initiation – Neuromyelitis Optica Spectrum Disorder (NMOSD)

All of the following:

- 1 Cumulative dose up to 1500 mg/m² body surface area up to 2000 mg total per cycle; and
- 2 Either:
 - 2.1 Patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms with supporting clinical investigations); or
 - 2.2 All of the following:
 - 2.2.1 Patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 Patient is receiving treatment with mycophenolate unless contraindicated or not tolerated; and
 - 2.2.3 Patient is receiving treatment with corticosteroids unless contraindicated or not tolerated; and
- 3 Each treatment cycle at least 6 months apart.

Note: Indications marked with * are unapproved indications.

Initiation – refractory myasthenia gravis*

Re-assessment required after 2 years

Both:

- 1 Cumulative dose up to 1500 mg/m² body surface area up to 2000 mg total per cycle; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for a minimum period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Note: Indications marked with * are unapproved indications.

Continuation – refractory myasthenia gravis*

Re-assessment required after 2 years

All of the following:

- 1 Cumulative dose up to 1500 mg/m² body surface area up to 2000 mg total per cycle; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 Patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 Patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Note: Indications marked with * are unapproved indications.

Initiation – antisynthetase syndrome

All of the following:

- 1 Patient has severe, immediately life- or organ-threatening disease, including interstitial lung disease; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

2 Either:

- 2.1 Treatment with at least 3 immunosuppressants (oral corticosteroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has been ineffective controlling active disease; or
- 2.2 Rapid treatment is required for life threatening complications; and

3 Maximum of two 1000 mg infusions every 6 months.

Initiation – graft versus host disease

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not been effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initiation – chronic inflammatory demyelinating polyneuropathy (CIPD)*

Neurologist

All of the following:

1 Either:

1.1 Both:

- 1.1.1 Treatment with corticosteroids and intravenous immunoglobulin and/or plasma exchange has been ineffective controlling active disease, is not tolerated, or is contraindicated; and
- 1.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) is not tolerated or has been ineffective controlling active disease. If an immunosuppressant is contraindicated, a trial has occurred of one of those which is not contraindicated (unless all are contraindicated); or

1.2 Rapid treatment is required for life threatening complications; and

2 Cumulative dose up to 1500 mg/m² body surface area up to 2000 mg total per cycle; and

3 Each treatment cycle at least 6 months apart.

Note: Indications marked with * are unapproved indications.

Initiation – anti-NMDA receptor autoimmune encephalitis*

All of the following:

1 Either:

1.1 Both:

- 1.1.1 Treatment with corticosteroids and intravenous immunoglobulin and/or plasma exchange has been ineffective controlling active disease, is not tolerated or is contraindicated; and
- 1.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has been ineffective controlling active disease, is not tolerated or is contraindicated; or

1.2 Rapid treatment is required for life threatening complications; and

2 Cumulative dose up to 1500 mg/m² body surface area up to 2000 mg total per cycle; and

3 Each treatment cycle at least 6 months apart.

Note: Indications marked with * are unapproved indications.

Initiation – CD20+ low grade or follicular B-cell NHL

Re-assessment required after 9 months

Either:

1 Both:

- 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
- 1.2 To be used for a maximum of 6 treatment cycles; or

2 Both:

- 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.2 To be used for a maximum of 6 treatment cycles.

Continuation – CD20+ low grade or follicular B-cell NHL

Re-assessment required after 24 months

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m² every 8 weeks (maximum of 12 cycles).

Initiation – Membranous nephropathy

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; and
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m²; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures that include (unless contraindicated or the patient has experienced intolerable side effects) renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents; and
- 3 The total rituximab dose per cycle would not exceed the equivalent of 375mg/m² of body surface area per week for a total of 4 weeks; and
- 4 Subsequent retreatment only for disease relapse or after partial response.

Note: Indications marked with * are unapproved indications.

Initiation – B-cell acute lymphoblastic leukaemia/lymphoma*

Limited to 2 years treatment

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initiation – desensitisation prior to transplant

Limited to 6 weeks treatment

Both:

- 1 Patient requires desensitisation prior to mismatched allogeneic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m² of body-surface area.

Note: Indications marked with * are unapproved indications.

Initiation – pemphigus*

Dermatologist or relevant specialist

Re-assessment required after 6 months

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.1 Patient has pemphigus; and
- 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Continuation – pemphigus*

Dermatologist or relevant specialist

Re-assessment required after 6 months

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

Initiation – immunoglobulin G4-related disease (IgG4-RD*)

Re-assessment required after 6 weeks

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD*; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or
 - 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

Continuation – immunoglobulin G4-related disease (IgG4-RD*)

Re-assessment required after 12 months

All of the following:

- 1 Either:
 - 1.1 Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
 - 1.2 Patient is receiving maintenance treatment for IgG4-RD*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB – Restricted see terms [below](#)

↓ Inj 150 mg per ml, 1 ml prefilled syringe.....	799.50	1	Cosentyx
	1,599.00	2	Cosentyx

➔ Restricted (RS2182)

Initiation – plaque psoriasis

Re-assessment required after 6 months

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient has "whole body" plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.1.2 Patient has plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.1.3 Patient has localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a DLQI score greater than 10; and
- 1.2 Patient has received insufficient benefit (see Note) or has experienced intolerable side effects from at least 3 of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 1.3 A PASI assessment or DLQI assessment has been completed for the most recent prior treatment course, within 1 month of stopping that treatment; and
- 1.4 The most recent PASI or DLQI assessment is within 1 month before the application; or
- 2 All of the following:
 - 2.1 Patient has had a Special Authority approval for adalimumab, etanercept, or infliximab, for plaque psoriasis; and
 - 2.2 Either:
 - 2.2.1 Patient has experienced intolerable side effects; or
 - 2.2.2 Patient has received insufficient benefit to meet the renewal criteria for plaque psoriasis; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for the most recent prior treatment within 1 month of stopping that treatment; and
 - 2.4 The most recent PASI or DLQI assessment is within 1 month before the application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Insufficient benefit" is defined as: for whole body plaque psoriasis, a PASI score of greater than 10; for plaque psoriasis of the face, hand, foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. As assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation – plaque psoriasis

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient's PASI score has reduced by 75% or more compared to pre-secukinumab baseline; and
 - 1.1.2 Patient has a DLQI improvement of 5 or more compared to pre-secukinumab baseline; or
 - 1.2 Both:
 - 1.2.1 Patient had localised genital or flexural plaque psoriasis at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, compared to the pre-secukinumab baseline; or
 - 1.2.2.2 Patient has a DLQI improvement of 5 or more, compared to pre-secukinumab baseline; and
- 2 Maximum dose 300 mg monthly.

Initiation – ankylosing spondylitis, second-line biologic

Re-assessment required after 3 months

Both:

- 1 Patient has had a Special Authority approval for adalimumab or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 Patient has experienced intolerable side effects; or
 - 2.2 Patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis.

Continuation – ankylosing spondylitis, second-line biologic

Re-assessment required after 6 months

Both:

- 1 BASDAI has improved from the pre-secukinumab baseline either by at least 4 points on a 10-point scale, or by at least 50%, whichever is less; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 Maximum dose 300 mg monthly.

Initiation – arthritis - psoriatic

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had a Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has received insufficient benefit from at least 3 months of methotrexate at a maximum tolerated dose unless contraindicated; and
 - 2.2 Patient has received insufficient benefit from at least 3 months of sulfasalazine or leflunomide at maximum tolerated doses unless contraindicated; and
 - 2.3 Either:
 - 2.3.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 joints; or
 - 2.3.2 Patient has persistent symptoms of poorly controlled and active disease in at least 4 joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.4 Any of the following:
 - 2.4.1 CRP greater than 15 mg/L measured within one month before the application; or
 - 2.4.2 ESR greater than 25 mm per hour measured within one month before the application; or
 - 2.4.3 ESR and CRP not measured as patient is receiving prednisone therapy greater than 5 mg per day received for more than 3 months.

Continuation – arthritis - psoriatic

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
 - 1.2 At least a continuing 30% improvement in active joint count from baseline; and
- 2 Maximum dose 300 mg monthly.

SILTUXIMAB – **Restricted** see terms [below](#)

↓ Inj 100 mg vial	770.57	1	Sylvant
↓ Inj 400 mg vial	3,082.33	1	Sylvant

→ **Restricted (RS1525)**

Initiation

Haematologist or rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Continuation

Haematologist or rheumatologist

Re-assessment required after 12 months

The treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB – **Restricted** see terms [on the next page](#)

↓ Inj 20 mg per ml, 4 ml vial.....	220.00	1	Actemra
↓ Inj 20 mg per ml, 10 ml vial.....	550.00	1	Actemra
↓ Inj 20 mg per ml, 20 ml vial.....	1,100.00	1	Actemra

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

➔ Restricted (RS2125)

Initiation – cytokine release syndrome

Therapy limited to 3 doses

Either:

- 1 Both:
 - 1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
 - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-Cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initiation – previous use

Any relevant practitioner

Limited to 6 months treatment

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initiation – Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Limited to 6 months treatment

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initiation – Rheumatoid Arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initiation – systemic juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initiation – adult-onset Still's disease

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Initiation – polyarticular juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 4 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initiation – idiopathic multicentric Castleman's disease

Haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initiation – moderate to severe COVID-19

Therapy limited to 1 dose

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with baricitinib.

Continuation – Rheumatoid Arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Continuation – systemic juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Continuation – adult-onset Still's disease

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

the patient has a sustained improvement in inflammatory markers and functional status.

Continuation – polyarticular juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Continuation – idiopathic multicentric Castleman's disease

Haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist

Re-assessment required after 12 months

the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

Initiation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and
- 2 The individual has received insufficient benefit from use of corticosteroids; and
- 3 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Continuation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

Both:

- 1 The individual has shown clinical improvement and ongoing treatment is required; and
- 2 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Note: Indications marked with * are unapproved indications.

TRASTUZUMAB (HERZUMA) – **Restricted** see terms [below](#)

↓ Inj 150 mg vial – 5% DV Jun-24 to 31 May 2027	100.00	1	Herzuma
↓ Inj 440 mg vial – 5% DV Jun-24 to 31 May 2027	293.35	1	Herzuma

➔ **Restricted (RS2005)**

Initiation – early breast cancer

Limited to 12 months treatment

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Continuation – early breast cancer*

Re-assessment required after 12 months

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
 - 1.3 Any of the following:
 - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
 - 1.3.3 he cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.4 Either:
 - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 1.4.2 All of the following:
 - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
 - 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

Initiation – metastatic breast cancer

Re-assessment required after 12 months

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – metastatic breast cancer

Re-assessment required after 12 months

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initiation – gastric, gastro-oesophageal junction and oesophageal cancer

Re-assessment required after 12 months

Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an ECOG score of 0-2.

Continuation – gastric, gastro-oesophageal junction and oesophageal cancer

Re-assessment required after 12 months

Both:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB DERUXTECAN – **Restricted** see terms [below](#)

↓ Inj 100 mg per ml, 1 ml vial 2,550.00 1 Enhertu

→ **Restricted (RS2082)**

Initiation

Re-assessment required after 6 months

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Patient has not received prior funded trastuzumab deruxtecan treatment; and
- 6 Treatment to be discontinued at disease progression.

Continuation

Re-assessment required after 6 months

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab deruxtecan; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

TRASTUZUMAB EMTANSINE – **Restricted** see terms [on the next page](#)

↓ Inj 100 mg vial 2,320.00 1 Kadcyla
↓ Inj 160 mg vial 3,712.00 1 Kadcyla

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2083)

Initiation – early breast cancer

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or axillary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initiation – metastatic breast cancer

Re-assessment required after 6 months

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Either:
 - 6.1 Patient has not received prior funded trastuzumab emtansine or trastuzumab deruxtecan treatment; or
 - 6.2 Both:
 - 6.2.1 Patient has discontinued trastuzumab deruxtecan due to intolerance; and
 - 6.2.2 The cancer did not progress while on trastuzumab deruxtecan; and
- 7 Treatment to be discontinued at disease progression.

Continuation – metastatic breast cancer

Re-assessment required after 6 months

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB – Restricted see terms [below](#)

⚡ Inj 130 mg vial	4,162.00	1	Stelara
⚡ Inj 90 mg per ml, 1 ml prefilled syringe	4,162.00	1	Stelara

➔ Restricted (RS1942)

Initiation – Crohn's disease - adults

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.1 Patient has active Crohn's disease; and
- 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Continuation – Crohn's disease - adults

Re-assessment required after 12 months

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initiation – Crohn's disease - children*

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

Continuation – Crohn's disease - children*

Re-assessment required after 12 months

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initiation – ulcerative colitis

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active ulcerative colitis; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.2 Either:

2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or

2.2.2 Both:

2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and

2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Continuation – ulcerative colitis

Re-assessment required after 12 months

Both:

1 Either:

1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or

1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and

2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB – **Restricted** see terms [below](#)

↓ Inj 300 mg vial 3,313.00 1 Entyvio

→ **Restricted (RS1943)**

Initiation – Crohn's disease - adults

Re-assessment required after 6 months

All of the following:

1 Patient has active Crohn's disease; and

2 Any of the following:

2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or

2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or

2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or

2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or

2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and

3 Any of the following:

3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or

3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or

3.3 Immunomodulators and corticosteroids are contraindicated.

Continuation – Crohn's disease - adults

Re-assessment required after 2 years

Both:

1 Any of the following:

1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or

1.2 CDAI score is 150 or less, or HBI is 4 or less; or

1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and

2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Initiation – Crohn's disease - children*

Re-assessment required after 6 months

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Continuation – Crohn's disease - children*

Re-assessment required after 2 years

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initiation – ulcerative colitis

Re-assessment required after 6 months

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Continuation – ulcerative colitis

Re-assessment required after 2 years

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Programmed Cell Death-1 (PD-1) Inhibitors

ATEZOLIZUMAB – **Restricted** see terms [below](#)

↓ Inj 60 mg per ml, 20 ml vial.....9,503.00 1 Tecentriq

→ **Restricted (RS2184)**

Initiation – non-small cell lung cancer second line monotherapy

Medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR, ROS-1 or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – non-small cell lung cancer second line monotherapy

Medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and
- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with atezolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
 - 2.2 Patient has preserved liver function (Child-Pugh A); and
 - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
 - 2.4 Any of the following:
 - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
 - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
 - 2.4.3 Both:
 - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
 - 2.4.3.2 No disease progression since initiation of lenvatinib; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.5 Patient has an ECOG performance status of 0-2; and

2.6 To be given in combination with bevacizumab.

Continuation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

no evidence of disease progression.

DURVALUMAB – Restricted see terms [below](#)

↓ Inj 50 mg per ml, 10 ml vial.....4,700.00 1 Imfinzi

↓ Inj 50 mg per ml, 2.4 ml vial.....1,128.00 1 Imfinzi

→ **Restricted (RS2084)**

Initiation – Non-small cell lung cancer

Re-assessment required after 4 months

All of the following:

- 1 Either:
 - 1.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); or
 - 1.2 Patient has histologically or cytologically documented stage IIb (T1N2a only), locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Continuation – Non-small cell lung cancer

Re-assessment required after 4 months

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

IPILIMUMAB – Restricted see terms [below](#)

↓ Inj 5 mg per ml, 10 ml vial.....5,000.00 1 Yervoy

↓ Inj 5 mg per ml, 40 ml vial.....20,000.00 1 Yervoy

→ **Restricted (RS2195)**

Initiation – stage III or IV resectable melanoma

Either:

- 1 The individual is currently on treatment with ipilimumab for neoadjuvant treatment of resectable stage IIIB, IIIC, IIID or IV melanoma and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.1 The individual has resectable stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note); and
- 2.2 The individual has not received prior funded systemic treatment in the perioperative setting for their stage IIIB, IIIC, IIID or IV melanoma; and
- 2.3 The individual has ECOG performance score 0-2; and
- 2.4 Treatment must be prior to complete surgical resection; and
- 2.5 Neoadjuvant ipilimumab must be administered in combination with nivolumab; and
- 2.6 Ipilimumab to be administered for maximum of two cycles prior to surgical resection.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition.
- b) Disease must be completely resectable and amenable to curative intent surgery, including stage IV disease.

Initiation – renal cell carcinoma

Limited to 4 months treatment

Either:

- 1 The patient is currently on treatment with ipilimumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and
 - 2.2 The patient is treatment naive; and
 - 2.3 The patient has ECOG performance status 0-2; and
 - 2.4 The disease is predominantly of clear cell histology; and
 - 2.5 Any of the following:
 - 2.5.1 The patient has sarcomatoid histology; or
 - 2.5.2 Haemoglobin levels less than the lower limit of normal; or
 - 2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or
 - 2.5.4 Neutrophils greater than the upper limit of normal; or
 - 2.5.5 Platelets greater than the upper limit of normal; or
 - 2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
 - 2.5.7 Karnofsky performance score of less than or equal to 70; and
 - 2.6 Ipilimumab is to be used at a maximum dose of 1 mg/kg for up to four cycles in combination with nivolumab.

NIVOLUMAB – **Restricted** see terms [below](#)

⚡ Inj 10 mg per ml, 4 ml vial.....	1,051.98	1	Opdivo
⚡ Inj 10 mg per ml, 10 ml vial.....	2,629.96	1	Opdivo

➡ **Restricted (RS2196)**

Initiation – stage III or IV resectable melanoma

Re-assessment required after 4 months

Either:

- 1 The individual is currently on treatment with nivolumab for neoadjuvant treatment of resectable stage IIIB, IIIC, IIID or IV melanoma and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The individual has resectable stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note); and
 - 2.2 The individual has not received prior funded systemic treatment in the perioperative setting for their stage IIIB, IIIC, IIID or IV melanoma; and
 - 2.3 The individual has ECOG performance score 0-2; and
 - 2.4 Treatment must be initiated prior to complete surgical resection; and
 - 2.5 Neoadjuvant nivolumab must be administered in combination with ipilimumab; and
 - 2.6 Nivolumab to be administered for maximum of two cycles prior to surgical resection.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – stage III or IV resectable melanoma

Re-assessment required after 4 months

Any of the following:

- 1 All of the following:
 - 1.1 The individual has received funded neoadjuvant treatment with nivolumab in combination with ipilimumab; and
 - 1.2 Adjuvant treatment with nivolumab is required; and
 - 1.3 Treatment must be initiated within 13 weeks of complete surgical resection, unless delay is necessary due to post-surgery recovery; and
 - 1.4 Nivolumab must be administered as monotherapy; and
 - 1.5 Nivolumab to be discontinued at signs of disease recurrence or at completion of 12 months total treatment duration including any systemic neoadjuvant treatment (equivalent to 11 adjuvant cycles at 480 mg every 4 weeks plus initial 2 neoadjuvant treatment cycles); or
- 2 All of the following:
 - 2.1 The individual has received neoadjuvant treatment with nivolumab and ipilimumab; and
 - 2.2 The individual has unresectable or metastatic melanoma (excluding uveal) stage III or IV; and
 - 2.3 The individual meets initial application criteria for nivolumab for unresectable or metastatic melanoma; or
- 3 All of the following:
 - 3.1 The individual has received neoadjuvant treatment with nivolumab and ipilimumab; and
 - 3.2 The individual has received treatment with nivolumab for unresectable or metastatic melanoma; and
 - 3.3 The individual meets the renewal criteria for nivolumab for unresectable or metastatic melanoma.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition.
- b) Disease must be completely resectable and amenable to curative intent surgery, including stage IV disease.

Initiation – unresectable or metastatic melanoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Limited to 4 months treatment

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance 0-2; and
- 4 Either:
 - 4.1 The individual has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 The individual has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the individual was on pembrolizumab; and
- 5 Any of the following:
 - 5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
 - 5.3 All of the following:
 - 5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
 - 5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
 - 5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-L1 inhibitor.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Continuation – unresectable or metastatic melanoma, less than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Both:
 - 1.1 Any of the following:
 - 1.1.1 The individual's disease has had a complete response to treatment; or
 - 1.1.2 The individual's disease has had a partial response to treatment; or
 - 1.1.3 The individual has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; or
- 2 All of the following:
 - 2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 The individual has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Continuation – unresectable or metastatic melanoma, more than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Both:

- 1 The individual has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Any of the following:
 - 2.1.1.1 The individual's disease has had a complete response to treatment; or
 - 2.1.1.2 The individual's disease has had a partial response to treatment; or
 - 2.1.1.3 The individual has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; or
 - 2.2 All of the following:
 - 2.2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 The individual has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

Initiation – renal cell carcinoma, first line

Limited to 4 months treatment

Either:

- 1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and
 - 2.2 The patient is treatment naive; and
 - 2.3 The patient has ECOG performance status 0-2; and
 - 2.4 The disease is predominantly of clear cell histology; and
 - 2.5 Any of the following:
 - 2.5.1 The patient has sarcomatoid histology; or
 - 2.5.2 Haemoglobin levels less than the lower limit of normal; or
 - 2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.5.4 Neutrophils greater than the upper limit of normal; or
- 2.5.5 Platelets greater than the upper limit of normal; or
- 2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
- 2.5.7 Karnofsky performance score of less than or equal to 70; and
- 2.6 Nivolumab is to be used in combination with ipilimumab for the first four treatment cycles at a maximum dose of 3 mg/kg; and
- 2.7 Nivolumab is to be used at a maximum maintenance dose of 240 mg every 2 weeks (or equivalent).

Initiation – renal cell carcinoma, second line

Limited to 4 months treatment

All of the following:

- 1 Patient has metastatic renal-cell carcinoma; and
- 2 The disease is of predominant clear-cell histology; and
- 3 Patient has ECOG performance status 0-2; and
- 4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and
- 5 Patient has not previously received a funded immune checkpoint inhibitor; and
- 6 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

Continuation – renal cell carcinoma

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

PEMBROLIZUMAB – **Restricted** see terms [below](#)

↓ Inj 25 mg per ml, 4 ml vial.....4,680.00 1 Keytruda

➔ **Restricted (RS2185)**

Initiation – stage III or IV resectable melanoma - neoadjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

All of the following:

- 1 The individual has resectable stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note); and
- 2 The individual has not received prior funded systemic treatment in the perioperative setting for their stage IIIB, IIIC, IIID or IV melanoma; and
- 3 Treatment must be prior to complete surgical resection; and
- 4 Pembrolizumab must be administered as monotherapy; and
- 5 The individual has ECOG performance score 0-2; and
- 6 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

Continuation – stage III or IV resectable melanoma - neoadjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Any of the following:

- 1 Both:
 - 1.1 The individual has received neoadjuvant treatment with an immune checkpoint inhibitor; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2 The individual meets initiation criteria for pembrolizumab for stage III or IV resected melanoma – adjuvant; or
- 2 Both:
 - 2.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
 - 2.2 The individual meets continuation criteria for pembrolizumab for stage III or IV resected melanoma – adjuvant; or
- 3 All of the following:
 - 3.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
 - 3.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 3.3 The individual meets initiation criteria for pembrolizumab for unresectable or metastatic melanoma; or
- 4 All of the following:
 - 4.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
 - 4.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma; and
 - 4.3 The individual meets continuation criteria for pembrolizumab for unresectable or metastatic melanoma.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means either 13 weeks after resection (primary or lymphadenectomy) or 13 weeks prior to the scheduled date of the resection (primary or lymphadenectomy)

Initiation – stage III or IV resected melanoma - adjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

All of the following:

- 1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); and
- 2 Adjuvant treatment with pembrolizumab is required; and
- 3 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
- 4 Treatment must be in addition to complete surgical resection; and
- 5 Treatment must be initiated within 13 weeks of complete surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
- 6 Pembrolizumab must be administered as monotherapy; and
- 7 The individual has ECOG performance score 0-2; and
- 8 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Continuation – stage III or IV resected melanoma - adjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Any of the following:

- 1 All of the following:
 - 1.1 No evidence of disease recurrence; and
 - 1.2 Pembrolizumab must be administered as monotherapy; and
 - 1.3 Pembrolizumab to be administered at a fixed dose of 200 mg every three weeks (or equivalent) for a maximum of 12 months total treatment course, including any systemic neoadjuvant treatment; and
 - 1.4 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months total treatment course (equivalent to 18 cycles at a dose of 200 mg every 3 weeks), including any systemic neoadjuvant treatment; or
- 2 All of the following:

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

- 2.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and
- 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2.3 The individual meets initiation criteria for pembrolizumab for unresectable or metastatic melanoma; or
- 3 All of the following:
 - 3.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and
 - 3.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma; and
 - 3.3 The individual meets continuation criteria for pembrolizumab for unresectable or metastatic melanoma.

Initiation – unresectable or metastatic melanoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Limited to 4 months treatment

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance 0-2; and
- 4 Either:
 - 4.1 The individual has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 The individual has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the individual was on nivolumab; and
- 5 Any of the following:
 - 5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
 - 5.3 All of the following:
 - 5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
 - 5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
 - 5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-L1 inhibitor.

Continuation – unresectable or metastatic melanoma, less than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Both:
 - 1.1 Any of the following:
 - 1.1.1 The individual's disease has had a complete response to treatment; or
 - 1.1.2 The individual's disease has had a partial response to treatment; or
 - 1.1.3 The individual has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; or
- 2 All of the following:
 - 2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 The individual has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – unresectable or metastatic melanoma, more than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Both:

- 1 The individual has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 All of the following:
 - 2.1.1 Any of the following:
 - 2.1.1.1 The individual's disease has had a complete response to treatment; or
 - 2.1.1.2 The individual's disease has had a partial response to treatment; or
 - 2.1.1.3 The individual has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
 - 2.1.3 The treatment remains clinically appropriate and the individual is benefitting from the treatment; or
 - 2.2 All of the following:
 - 2.2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 The individual has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initiation – non-small cell lung cancer first-line monotherapy

Medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR, ROS-1 or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Either:
 - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
 - 6.2 Both:
 - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and
 - 6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and
- 7 Patient has an ECOG 0-2; and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – non-small cell lung cancer first-line monotherapy

Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – non-small cell lung cancer first-line combination therapy

Medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR, ROS-1 or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2; and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – non-small cell lung cancer first-line combination therapy

Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – breast cancer, advanced

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
 - 2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- HER2 IHC3+ or ISH+ [including FISH or other technology]); and
- 2.2 Patient is treated with palliative intent; and
- 2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
- 2.4 Patient has received no prior systemic therapy in the palliative setting; and
- 2.5 Patient has an ECOG score of 0–2; and
- 2.6 Pembrolizumab is to be used in combination with chemotherapy; and
- 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – breast cancer, advanced

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period; and
- 4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – head and neck squamous cell carcinoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies; and
 - 2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
 - 2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
 - 2.4 Patient has an ECOG performance score of 0-2; and
 - 2.5 Either:
 - 2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or
 - 2.5.2 Pembrolizumab to be used as monotherapy; and
 - 2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – head and neck squamous cell carcinoma

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – MSI-H/dMMR advanced colorectal cancer

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer; or
 - 2.1.2 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer; and
 - 2.2 Individual is treated with palliative intent; and
 - 2.3 Individual has not previously received funded treatment with pembrolizumab for MSI-H/dMMR advanced colorectal cancer; and
 - 2.4 Individual has an ECOG performance score of 0-2; and
 - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – MSI-H/dMMR advanced colorectal cancer

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 No evidence of disease progression; and
- 2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – Urothelial carcinoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
 - 2.2 Patient has an ECOG performance score of 0-2; and
 - 2.3 Patient has documented disease progression following treatment with chemotherapy; and
 - 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – Urothelial carcinoma

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2 Patient's disease has had a partial response to treatment; and
- 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – relapsed/refractory Hodgkin lymphoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Both:
 - 2.1.1.1 Individual has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and
 - 2.1.1.2 Individual is ineligible for autologous stem cell transplant; or
 - 2.1.2 Individual has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant; and
 - 2.2 Individual has not previously received funded pembrolizumab for relapsed/refractory Hodgkin lymphoma; and
 - 2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Continuation – relapsed/refractory Hodgkin lymphoma

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 Patient has received a partial or complete response to pembrolizumab; and
- 2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Other Immunosuppressants

ANTITHYMOCYTE GLOBULIN (EQUINE)

Inj 50 mg per ml, 5 ml ampoule	4,439.17	5	ATGAM
--------------------------------------	----------	---	-------

ANTITHYMOCYTE GLOBULIN (RABBIT)

Inj 25 mg vial

AZATHIOPRINE

Tab 25 mg – 5% DV Feb-26 to 2028	10.15	60	Azamun
Tab 50 mg – 5% DV Feb-26 to 2028	10.34	100	Azamun
Inj 50 mg vial			
Inj 100 mg vial			

BACILLUS CALMETTE-GUERIN (BCG) – **Restricted** see terms [below](#)

⚡ Inj 2-8 × 10 ⁸ CFU vial	182.45	1	OncoTICE
⚡ Inj 40 mg per ml, vial	182.45	3	SII-Onco-BCG

➡ **Restricted (RS1206)**

Initiation

For use in bladder cancer.

EVEROLIMUS – **Restricted** see terms [on the next page](#)

⚡ Tab 5 mg	4,555.76	30	Afinitor
⚡ Tab 10 mg	6,512.29	30	Afinitor

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2076)

Initiation

Neurologist or oncologist

Re-assessment required after 3 months

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Continuation

Neurologist or oncologist

Re-assessment required after 12 months

All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Initiation – renal cell carcinoma

Re-assessment required after 4 months

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma; and
 - 1.2 The disease is of predominant clear-cell histology; and
 - 1.3 The patient has documented disease progression following one previous line of treatment; and
 - 1.4 The patient has an ECOG performance status of 0-2; and
 - 1.5 Everolimus is to be used in combination with lenvatinib; or
- 2 All of the following:
 - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
 - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
 - 2.3 Everolimus is to be used in combination with lenvatinib; and
 - 2.4 There is no evidence of disease progression.

Continuation – renal cell carcinoma

Re-assessment required after 4 months

there is no evidence of disease progression.

MYCOPHENOLATE MOFETIL

Tab 500 mg	35.90	50	CellCept
Cap 250 mg	35.90	100	CellCept
Powder for oral liq 1 g per 5 ml.....	187.25	165 ml	CellCept
Inj 500 mg vial	133.33	4	CellCept

PICIBANIL

Inj 100 mcg vial

SIROLIMUS – Restricted see terms [below](#)

⚡ Tab 1 mg	749.99	100	Rapamune
⚡ Tab 2 mg	1,499.99	100	Rapamune
⚡ Oral liq 1 mg per ml	449.99	60 ml	Rapamune

➔ Restricted (RS1991)

Initiation

For rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

to calcineurin inhibitor treatment due to any of the following:

- GFR < 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencephalopathy; or
- Significant malignant disease

Initiation – severe non-malignant lymphovascular malformations*

Re-assessment required after 6 months

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Continuation – severe non-malignant lymphovascular malformations*

Re-assessment required after 12 months

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initiation – renal angiomyolipoma(s) associated with tuberous sclerosis complex*

Nephrologist or urologist

Re-assessment required after 6 months

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Continuation – renal angiomyolipoma(s) associated with tuberous sclerosis complex*

Re-assessment required after 12 months

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

continued...

	Price (ex man. excl. GST)		Brand or Generic Manufacturer
	\$	Per	

continued...

Initiation – refractory seizures associated with tuberous sclerosis complex*

Neurologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex*; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, and topiramate. Those who can father children are not required to trial sodium valproate.

Continuation – refractory seizures associated with tuberous sclerosis complex*

Neurologist

Re-assessment required after 12 months

demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – **Restricted** see terms [below](#)

⚡ Tab modified-release 15 mg.....	1,271.00	28	Rinvoq
⚡ Tab modified-release 30 mg.....	2,033.00	28	Rinvoq
⚡ Tab modified-release 45 mg.....	3,049.00	28	Rinvoq

→ **Restricted (RS2120)**

Initiation – Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)

Limited to 6 months treatment

All of the following:

- 1 The individual has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The individual has experienced intolerable side effects with adalimumab and/or etanercept; or
 - 2.2 The individual has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Any of the following:
 - 3.1 Rituximab is not clinically appropriate; or
 - 3.2 The individual is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.3 Both:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

3.3.1 The individual has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and

3.3.2 Either:

3.3.2.1 The individual has experienced intolerable side effects with rituximab; or

3.3.2.2 At four months following the initial course of rituximab the individual has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Continuation – Rheumatoid Arthritis

Re-assessment required after 6 months

Either:

- 1 Following 6 months' initial treatment, the individual has experienced at least a 50% decrease in active joint count from baseline; or
- 2 On subsequent reapplications, the individual has experienced at least a continuing 30% improvement in active joint count from baseline.

Initiation – Atopic dermatitis

Re-assessment required after 6 months

Either:

- 1 Individual is currently on treatment with upadacitinib for atopic dermatitis and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Individual has moderate to severe atopic dermatitis, severity as defined by an Eczema Area and Severity Index (EASI) score of greater than or equal to 16 or a Dermatology Life Quality Index (DLQI) score of greater than or equal to 10; and
 - 2.2 Individual has received insufficient benefit from topical therapy (including topical corticosteroids or topical calcineurin inhibitors) for a 28-day trial within the last 6 months, unless contraindicated to all; and
 - 2.3 Individual has trialed and received insufficient benefit from at least one systemic therapy for a minimum of three months (eg ciclosporin, azathioprine, methotrexate or mycophenolate mofetil), unless contraindicated to all; and
 - 2.4 An EASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.5 The most recent EASI or DQI assessment is no more than 1 month old at the time of application.

Continuation – Atopic dermatitis

Re-assessment required after 12 months

Either:

- 1 Individual has received a 75% or greater reduction in EASI score (EASI 75) as compared to baseline EASI prior to commencing upadacitinib; or
- 2 Individual has received a DLQI improvement of 4 or more as compared to baseline DLQI prior to commencing upadacitinib.

Initiation – Crohn's disease – adult

Re-assessment required after 6 months

Either:

- 1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing treatment; or
- 2 Both:
 - 2.1 Individual has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Individual has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for Crohn's disease; and
- 2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Continuation – Crohn's disease – adult

Re-assessment required after 2 years

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score when the individual was initiated on biologic therapy; or
- 2 HBI score has reduced by 3 points from when individual was initiated on biologic therapy; or
- 3 CDAI score is 150 or less; or
- 4 HBI score is 4 or less; or
- 5 The individual has experienced an adequate response to treatment, but CDAI score cannot be assessed.

Initiation – Crohn's disease – children

Re-assessment required after 6 months

Either:

- 1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing treatment; or
- 2 Both:
 - 2.1 Child has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Child has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Child meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Continuation – Crohn's disease – children

Re-assessment required after 2 years

Any of the following:

- 1 PCDAI score has reduced by 10 points from when the child was initiated on treatment; or
- 2 PCDAI score is 15 or less; or
- 3 The child has experienced an adequate response to treatment, but PCDAI score cannot be assessed.

Note: Indications marked with * are unapproved indications.

Initiation – Ulcerative colitis

Re-assessment required after 6 months

Either:

- 1 Individual is currently on treatment with upadacitinib for ulcerative colitis and met all remaining criteria prior to commencing treatment; or
- 2 Both:
 - 2.1 Individual has active ulcerative colitis; and
 - 2.2 Either:
 - 2.2.1 Individual has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologic therapies for ulcerative colitis are contraindicated.

Continuation – Ulcerative colitis

Re-assessment required after 2 years

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the individual was initiated on treatment; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 PUCAI score has reduced by 10 points or more from the PUCAI score when the individual was initiated on treatment.

	Price (ex man. excl. GST)		Brand or Generic Manufacturer
	\$	Per	

Antiallergy Preparations

Allergic Emergencies

ADRENALINE – **Restricted** see terms [below](#)

↓ Inj 0.15 mg per 0.3 ml auto-injector – 5% DV Dec-25 to 2028.....	85.50	1	Epipen Jr
↓ Inj 0.3 mg per 0.3 ml auto-injector – 5% DV Dec-25 to 2028.....	85.50	1	Epipen

→ **Restricted (RS1944)**

Initiation – anaphylaxis

Either:

- 1 Patient has experienced a previous anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
- 2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner.

ICATIBANT – **Restricted** see terms [below](#)

↓ Inj 10 mg per ml, 3 ml prefilled syringe – 5% DV Oct-26 to 2028	2,668.00	1	Firazyr
	479.00		Icatibant Lupin

(Firazyr Inj 10 mg per ml, 3 ml prefilled syringe to be delisted 1 October 2026)

→ **Restricted (RS1501)**

Initiation

Clinical immunologist or relevant specialist

Re-assessment required after 12 months

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Continuation

Re-assessment required after 12 months

The treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

BEE VENOM – **Restricted** see terms [below](#)

↓ Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent			
↓ Inj 550 mcg vial with diluent			
↓ Initiation kit - 1 vial freeze dried venom with diluent	305.00	1	VENOX
↓ Maintenance Kit - 1 vial freeze dried venom with diluent	305.00	1	VENOX

→ **Restricted (RS1117)**

Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

PAPER WASP VENOM – **Restricted** see terms [below](#)

↓ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent			
↓ Inj 550 mcg vial with diluent			

→ **Restricted (RS1118)**

Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
YELLOW JACKET WASP VENOM – Restricted see terms below			
⚡ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			
➡ Restricted (RS1119)			
Initiation			
Both:			
1 RAST or skin test positive; and			
2 Patient has had severe generalised reaction to the sensitising agent.			
Allergy Prophylactics			
BUDESONIDE			
Nasal spray 50 mcg per dose – 5% DV Feb-25 to 2027	2.59	200 dose	SteroClear
Nasal spray 100 mcg per dose – 5% DV Feb-25 to 2027	2.89	200 dose	SteroClear
FLUTICASONE PROPIONATE			
Metered dose nasal spray 50 mcg per dose – 5% DV Aug-26 to 2028	2.57	120 dose	Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE			
Aqueous nasal spray 0.03%	5.23	15 ml	Univent
SODIUM CROMOGLICATE			
Nasal spray 4%			
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
Tab 10 mg	1.71	100	Zista
Oral liq 1 mg per ml	3.99	200 ml	Histaclear
CHLORPHENIRAMINE MALEATE			
Oral liq 0.4 mg per ml			
Inj 10 mg per ml, 1 ml ampoule			
CYPROHEPTADINE HYDROCHLORIDE			
Tab 4 mg			
FEXOFENADINE HYDROCHLORIDE			
Tab 60 mg			
Tab 120 mg – 5% DV Jul-25 to 2027	3.49	30	Fexaclear
Tab 180 mg – 5% DV Jul-25 to 2027	4.10	30	Fexaclear
LORATADINE			
Tab 10 mg – 5% DV Jun-26 to 2028	1.59	100	Loratadine Noumed
Oral liq 1 mg per ml	1.43	100 ml	Haylor Syrup
PROMETHAZINE HYDROCHLORIDE			
Tab 10 mg – 5% DV Dec-25 to 2028	2.19	100	Allersoothe
Tab 25 mg – 5% DV Dec-25 to 2028	2.69	100	Allersoothe
Oral liq 1 mg per ml	3.39	100 ml	Allersoothe
Inj 25 mg per ml, 1 ml ampoule	52.73	25	XGen
Inj 25 mg per ml, 2 ml ampoule	21.09	5	DBL Promethazine Hydrochloride

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Anticholinergic Agents

IPRATROPIUM BROMIDE

Aerosol inhaler 20 mcg per dose			
Nebuliser soln 250 mcg per ml, 1 ml ampoule			
Nebuliser soln 250 mcg per ml, 2 ml ampoule	11.73	20	Accord
	8.11	10	Ipratropium Viatris
	11.73	20	Univent

Anticholinergic Agents with Beta-Adrenoceptor Agonists

SALBUTAMOL WITH IPRATROPIUM BROMIDE

Aerosol inhaler 100 mcg with ipratropium bromide 20 mcg per dose			
Nebuliser soln 2.5 mg with ipratropium bromide 0.5 mg per 2.5 ml ampoule	96.50	60	Combiprasal
	11.04	20	Duolin
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 3.0 ml vial	16.56	30	Cipla

Long-Acting Muscarinic Agents

GLYCOPYRRONIUM

Note: inhaled glycopyrronium treatment must not be used if the patient is also receiving treatment with subsidised tiotropium or umeclidinium.

Powder for inhalation 50 mcg per dose	61.00	30 dose	Seebri Breezhaler
---	-------	---------	-------------------

TIOTROPIUM BROMIDE

Note: tiotropium treatment must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.

Soln for inhalation 2.5 mcg per dose	50.37	60 dose	Spiriva Respimat
Powder for inhalation 18 mcg per dose	50.37	30 dose	Spiriva

UMECLIDINIUM

Note: Umeclidinium must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.

Powder for inhalation 62.5 mcg per dose	61.50	30 dose	Incruse Ellipta
---	-------	---------	-----------------

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

➔ Restricted (RS2155)

Initiation

Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Note: Combination long acting muscarinic antagonist and long acting beta-2 agonist must not be used if the patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

GLYCOPYRRONIUM WITH INDACATEROL – Restricted see terms [above](#)

† Powder for Inhalation 50 mcg with indacaterol 110 mcg	81.00	30 dose	Ultibro Breezhaler
---	-------	---------	--------------------

TIOTROPIUM BROMIDE WITH OLODATEROL – Restricted see terms [above](#)

† Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg	81.00	60 dose	Spolto Respimat
---	-------	---------	-----------------

UMECLIDINIUM WITH VILANTEROL – Restricted see terms [above](#)

† Powder for inhalation 62.5 mcg with vilanterol 25 mcg	77.00	30 dose	Anoro Ellipta
---	-------	---------	---------------

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

BUDESONIDE WITH GLYCOPYRRONIUM AND EFORMOTEROL – **Restricted** see terms [below](#)

↓ Aerosol inhaler budesonide 160 mcg with glycopyrronium 7.2 mcg and formoterol 5 mcg per dose.....79.15 120 dose Breztri Aerosphere

➔ **Restricted** (RS2085)

Initiation

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

 - 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
 - 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
 - 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
 - 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
 - 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long-acting muscarinic antagonist and long-acting beta-2 agonist – ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler therapy.

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – **Restricted** see terms [below](#)

↓ Powder for inhalation fluticasone furoate 100 mcg with umeclidinium 62.5 mcg and vilanterol 25 mcg.....104.24 30 dose Trelegy Ellipta

➔ **Restricted** (RS2028)

Initiation

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

 - 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
 - 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
 - 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
 - 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
 - 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist – ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Antifibrotics

NINTEDANIB – Restricted see terms [below](#)

↓ Cap 100 mg.....	2,554.00	60	Ofev
↓ Cap 150 mg.....	3,870.00	60	Ofev

→ **Restricted (RS1813)**

Initiation – idiopathic pulmonary fibrosis

Respiratory specialist

Re-assessment required after 12 months

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Continuation – idiopathic pulmonary fibrosis

Respiratory specialist

Re-assessment required after 12 months

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Restricted see terms [below](#)

↓ Tab 267 mg – 5% DV Oct-26 to 2028	1,215.00	90	Esbriet
	145.80		Pirfenidone Sandoz
↓ Tab 801 mg – 5% DV Oct-26 to 2028	3,645.00	90	Esbriet
	437.40		Pirfenidone Sandoz

(Esbriet Tab 267 mg to be delisted 1 October 2026)

(Esbriet Tab 801 mg to be delisted 1 October 2026)

→ **Restricted (RS1814)**

Initiation – idiopathic pulmonary fibrosis

Respiratory specialist

Re-assessment required after 12 months

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Notes); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Continuation – idiopathic pulmonary fibrosis

Respiratory specialist

Re-assessment required after 12 months

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Beta-Adrenoceptor Agonists

SALBUTAMOL

Oral liq 400 mcg per ml – 5% DV May-25 to 2027.....	50.00	150 ml	Ventolin
Inj 500 mcg per ml, 1 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			
Aerosol inhaler, 100 mcg per dose.....	4.18	200 dose	SalAir
	7.45		Ventolin
Nebuliser soln 1 mg per ml, 2.5 ml ampoule	8.96	20	Asthalin
			UK Cipla
Nebuliser soln 2 mg per ml, 2.5 ml ampoule	9.43	20	Asthalin
			UK Cipla

TERBUTALINE SULPHATE

Powder for inhalation 250 mcg per dose			
Inj 0.5 mg per ml, 1 ml ampoule			
Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated.....	22.20	120 dose	Bricanyl Turbuhaler

Decongestants

OXYMETAZOLINE HYDROCHLORIDE

Aqueous nasal spray 0.25 mg per ml	
Aqueous nasal spray 0.5 mg per ml	

PSEUDOEPHEDRINE HYDROCHLORIDE

Tab 60 mg	
-----------	--

SODIUM CHLORIDE

Aqueous nasal spray isotonic	
------------------------------	--

SODIUM CHLORIDE WITH SODIUM BICARBONATE

Soln for nasal irrigation	
---------------------------	--

XYLOMETAZOLINE HYDROCHLORIDE

Aqueous nasal spray 0.05%	
Aqueous nasal spray 0.1%	
Nasal drops 0.05%	
Nasal drops 0.1%	

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler 50 mcg per dose.....	8.54	200 dose	Beclazone 50
	14.01		Qvar
Aerosol inhaler 100 mcg per dose.....	12.50	200 dose	Beclazone 100
	17.52		Qvar
Aerosol inhaler 250 mcg per dose.....	22.67	200 dose	Beclazone 250
BUDESONIDE			
Nebuliser soln 250 mcg per ml, 2 ml ampoule			
Nebuliser soln 500 mcg per ml, 2 ml ampoule			
Powder for inhalation 100 mcg per dose			
Powder for inhalation 200 mcg per dose			
Powder for inhalation 400 mcg per dose			
FLUTICASONE			
Aerosol inhaler 50 mcg per dose.....	7.19	120 dose	Flixotide
Powder for inhalation 50 mcg per dose.....	8.61	60 dose	Flixotide Accuhaler
Powder for inhalation 100 mcg per dose.....	7.81	60 dose	Flixotide Accuhaler
Aerosol inhaler 125 mcg per dose.....	13.60	120 dose	Flixotide
Aerosol inhaler 250 mcg per dose.....	24.62	120 dose	Flixotide
Powder for inhalation 250 mcg per dose.....	11.93	60 dose	Flixotide Accuhaler
Leukotriene Receptor Antagonists			
MONTELUKAST			
Tab 4 mg – 5% DV Dec-25 to 2028	3.10	28	Montelukast Viatrix
Tab 5 mg – 5% DV Dec-25 to 2028	3.10	28	Montelukast Paediatric (Relonchem)
			Montelukast Viatrix
Tab 10 mg – 5% DV Dec-25 to 2028	2.45	28	Montelukast Viatrix
Long-Acting Beta-Adrenoceptor Agonists			
EFORMOTEROL FUMARATE			
Powder for inhalation 12 mcg per dose			
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated (equivalent to eformoterol fumarate 6 mcg metered dose)			
INDACATEROL			
Powder for inhalation 150 mcg per dose.....	61.00	30 dose	Onbrez Breezhaler
Powder for inhalation 300 mcg per dose.....	61.00	30 dose	Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler 25 mcg per dose.....	26.25	120 dose	Serevent
Powder for inhalation 50 mcg per dose.....	26.25	60 dose	Serevent Accuhaler

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

BUDESONIDE WITH EFORMOTEROL

Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg			
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg			
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide with 6 mcg eformoterol fumarate metered dose)	41.50	120 dose	DuoResp Spiromax
Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg	33.74	120 dose	Symbicort Turbuhaler
Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mcg eformoterol fumarate metered dose)	82.50	120 dose	DuoResp Spiromax
Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg	33.74	60 dose	Symbicort Turbuhaler

FLUTICASONE FUROATE WITH VILANTEROL

Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose	Breo Ellipta
--	-------	---------	--------------

FLUTICASONE WITH SALMETEROL

Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.79	120 dose	Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg	33.74	60 dose	Seretide Accuhaler
Aerosol inhaler 125 mcg with salmeterol 25 mcg	32.60	120 dose	Seretide
Powder for inhalation 250 mcg with salmeterol 50 mcg	44.08	60 dose	Seretide Accuhaler

Methylxanthines

AMINOPHYLLINE

Inj 25 mg per ml, 10 ml ampoule	180.00	5	DBL Aminophylline
---------------------------------------	--------	---	-------------------

CAFFEINE CITRATE

Oral liq 20 mg per ml (caffeine 10 mg per ml)	16.91	25 ml	Biomed
Inj 20 mg per ml (caffeine 10 mg per ml), 2.5 ml ampoule	69.70	5	Biomed

THEOPHYLLINE

Tab long-acting 250 mg	25.65	100	Nuelin-SR
Oral liq 80 mg per 15 ml	18.49	500 ml	Nuelin

Mucolytics and Expectorants

DORNASE ALFA – **Restricted** see terms [below](#)

⚡ Nebuliser soln 2.5 mg per 2.5 ml ampoule	250.00	6	Pulmozyme
--	--------	---	-----------

➔ **Restricted (RS1787)**

Initiation – cystic fibrosis

Respiratory physician or paediatrician

Re-assessment required after 12 months

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or
- 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Continuation – cystic fibrosis

Respiratory physician or paediatrician

The treatment remains appropriate and the patient continues to benefit from treatment.

Initiation – significant mucus production

Limited to 4 weeks treatment

Both:

- 1 Patient is an in-patient; and
- 2 The mucus production cannot be cleared by first line chest techniques.

Initiation – pleural emphysema

Limited to 3 days treatment

Both:

- 1 Patient is an in-patient; and
- 2 Patient diagnoses with pleural emphysema.

ELEXACFTOR WITH TEZACFTOR, IVACFTOR AND IVACFTOR – **Restricted** see terms [below](#)

↓ Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg (56) and ivacaftor 75 mg (28).....	27,647.39	84	Trikafta
↓ Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg (56) and ivacaftor 150 mg (28).....	27,647.39	84	Trikafta
↓ Oral granules elexacaftor 80 mg with tezacaftor 40 mg, ivacaftor 60 mg (28) and ivacaftor 59.5mg (28), sachets.....	27,647.39	56	Trikafta
↓ Oral granules elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg (28) and ivacaftor 75 mg (28), sachets.....	27,647.39	56	Trikafta

➔ **Restricted (RS2197)**

Initiation

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 2.2 Patient has a sweat chloride value of at least 60 mmol/L; and
- 3 Either:
 - 3.1 Patient has a heterozygous or homozygous F508del mutation; or
 - 3.2 Patient has a mutation responsive to elexacaftor/tezacaftor/ivacaftor (see note); and
- 4 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 5 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Note: Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information

https://www.accessdata.fda.gov/drugsatfda_docs/label/2026/212273s014lbl.pdf

IVACFTOR – **Restricted** see terms [on the next page](#)

↓ Tab 150 mg	29,386.00	56	Kalydeco
↓ Oral granules 13.4 mg, sachet	29,386.00	56	Kalydeco
↓ Oral granules 25 mg, sachet	29,386.00	56	Kalydeco
↓ Oral granules 50 mg, sachet	29,386.00	56	Kalydeco
↓ Oral granules 75 mg, sachet	29,386.00	56	Kalydeco

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2198)

Initiation

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 2.2 Patients must have a sweat chloride value of at least 60 mmol/L; and
- 3 Patient has a mutation responsive to ivacaftor (see note); and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 The dose of ivacaftor will not exceed one tablet or one sachet twice daily.

Note: Mutations listed in Table 3 of the Food and Drug Administration (FDA) Ivacaftor prescribing information

https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/203188s038lbl.pdf

SODIUM CHLORIDE

Nebuliser soln 7%, 90 ml bottle	25.73	90 ml	Biomed
---------------------------------------	-------	-------	--------

VANZACAFTOR WITH TEZACAFTOR AND DEUTIVACAFTOR – Restricted see terms [below](#)

⚡ Tab vanzacaftor 4 mg with tezacaftor 20 mg and deutivacaftor 50 mg	29,029.76	84	Alyftrek
--	-----------	----	----------

⚡ Tab vanzacaftor 10 mg with tezacaftor 50 mg and deutivacaftor 125 mg	29,029.76	56	Alyftrek
--	-----------	----	----------

➔ Restricted (RS2199)

Initiation

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 2.2 Patient has a sweat chloride value of at least 60 mmol/L; and
- 3 Either:
 - 3.1 Patient has a heterozygous or homozygous F508del mutation; or
 - 3.2 Patient has a mutation responsive to vanzacaftor/tezacaftor/deutivacaftor (see note); and
- 4 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 5 Treatment with vanzacaftor/tezacaftor/deutivacaftor must be given concomitantly with standard therapy for this condition.

Note: Eligible mutations are listed in the in the Food and Drug Administration (FDA) Alyftrek prescribing information

https://www.accessdata.fda.gov/drugsatfda_docs/label/2026/218730s001lbl.pdf

Pulmonary Surfactants

BERACTANT

Soln 200 mg per 8 ml vial

PORACTANT ALFA

Soln 120 mg per 1.5 ml vial	425.00	1	Curosurf
Soln 240 mg per 3 ml vial	695.00	1	Curosurf

Respiratory Stimulants

DOXAPRAM

Inj 20 mg per ml, 5 ml vial

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

Sclerosing Agents

TALC
Powder
Soln (slurry) 100 mg per ml, 50 ml

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anti-Infective Preparations			
Antibacterials			
CHLORAMPHENICOL			
Eye oint 1% – 5% DV Feb-26 to 2028	1.55	5 g	Devatis
Ear drops 0.5%			
Eye drops 0.5% – 5% DV Mar-26 to 2028	1.84	10 ml	Chlorafast
Eye drops 0.5%, single dose			
CIPROFLOXACIN			
Eye drops 0.3% – 5% DV Mar-25 to 2027	10.85	5 ml	Ciprofloxacin Teva
FRAMYCETIN SULPHATE			
Ear/eye drops 0.5%			
GENTAMICIN SULPHATE			
Eye drops 0.3%			
PROPAMIDINE ISETHIONATE			
Eye drops 0.1%			
SODIUM FUSIDATE [FUSIDIC ACID]			
Eye drops 1%	5.29	5 g	Fucithalmic
SULPHACETAMIDE SODIUM			
Eye drops 10%			
TOBRAMYCIN			
Eye oint 0.3%	10.45	3.5 g	Tobrex
Eye drops 0.3%	11.48	5 ml	Tobrex
Antifungals			
NATAMYCIN			
Eye drops 5%			
Antivirals			
ACICLOVIR			
Eye oint 3% – 5% DV Feb-25 to 2027	15.89	4.5 g	ViruPOS
Combination Preparations			
CIPROFLOXACIN WITH HYDROCORTISONE			
Ear drops ciprofloxacin 0.2% with 1% hydrocortisone.....	16.30	10 ml	Ciproxin HC Otic
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml			
DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE			
Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	5.39	3.5 g	Maxitrol
Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	4.50	5 ml	Maxitrol
DEXAMETHASONE WITH TOBRAMYCIN			
Eye drops 0.1% with tobramycin 0.3%	12.64	5 ml	Tobradex

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUMETASONE PIVALATE WITH CLIQUINOL Ear drops 0.02% with cliquinol 1%			
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml	Kenacomb

Anti-Inflammatory Preparations

Corticosteroids

DEXAMETHASONE			
Eye oint 0.1%	5.86	3.5 g	Maxidex
Eye drops 0.1%	4.50	5 ml	Maxidex
↓ Ocular implant 700 mcg.....	1,444.50	1	Ozurdex

→ Restricted (RS1606)

Initiation – Diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

All of the following:

- 1 Patients have diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision; and
- 3 Either:
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Continuation – Diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initiation – Women of child bearing age with diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

All of the following:

- 1 Patients have diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Continuation – Women of child bearing age with diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUOROMETHOLONE			
Eye drops 0.1%	3.09	5 ml	FML
PREDNISOLONE ACETATE			
Eye drops 0.12%			
Eye drops 1%	7.00	5 ml	Pred Forte
	6.92	10 ml	Prednisolone- AFT
PREDNISOLONE SODIUM PHOSPHATE			
Eye drops 0.5%, single dose (preservative free).....	43.26	20 dose	Minims Prednisolone

Non-Steroidal Anti-Inflammatory Drugs

DICLOFENAC SODIUM			
Eye drops 0.1%			
Eye drops 0.1%, single dose – 5% DV Jul-25 to 2027	1.85	10 dose	Diclofenac Devatis
	5.54	30 dose	Diclofenac Devatis
KETOROLAC TROMETAMOL			
Eye drops 0.5%			

Decongestants and Antiallergics

Antiallergic Preparations

LEVOCABASTINE			
Eye drops 0.05%			
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml	Lomide
OLOPATADINE			
Eye drops 0.1% – 5% DV Mar-26 to 2028	3.39	5 ml	Olopatadine Teva
SODIUM CROMOGLICATE			
Eye drops 2% – 5% DV Mar-26 to 2028	2.91	10 ml	Allerfix

Decongestants

NAPHAZOLINE HYDROCHLORIDE			
Eye drops 0.1% – 5% DV Jan-25 to 2027	5.65	15 ml	Albalon

Diagnostic and Surgical Preparations

Diagnostic Dyes

FLUORESCEIN SODIUM			
Eye drops 2%, single dose			
Inj 10%, 5 ml vial	125.00	12	Fluorescite
Ophthalmic strips 1 mg			
FLUORESCEIN SODIUM WITH LIGNOCAINE HYDROCHLORIDE			
Eye drops 0.25% with lignocaine hydrochloride 4%, single dose			
LISSAMINE GREEN			
Ophthalmic strips 1.5 mg			
ROSE BENGAL SODIUM			
Ophthalmic strips 1%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Irrigation Solutions			
MIXED SALT SOLUTION FOR EYE IRRIGATION			
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 15 ml dropper bottle	5.00	15 ml	Balanced Salt Solution
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 250 ml			<i>e.g. Balanced Salt Solution</i>
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml bag			<i>e.g. Balanced Salt Solution</i>
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml bottle.....	10.50	500 ml	Balanced Salt Solution
Ocular Anaesthetics			
OXYBUPROCAINE HYDROCHLORIDE			
Eye drops 0.4%, single dose			
PROXYMETACAINE HYDROCHLORIDE			
Eye drops 0.5%			
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Eye drops 0.5%, single dose			
Eye drops 1%, single dose			
Viscoelastic Substances			
HYPROMELLOSE			
Inj 2%, 1 ml syringe			
Inj 2%, 2 ml syringe			
SODIUM HYALURONATE [HYALURONIC ACID]			
Inj 18 mg per ml, 0.85 ml syringe – 5% DV Mar-26 to 2028	50.00	1	Healon GV Pro
Inj 23 mg per ml, 0.6 ml syringe – 5% DV Mar-26 to 2028	60.00	1	Healon 5
Inj 10 mg per ml, 0.85 ml syringe – 5% DV Mar-26 to 2028	28.50	1	Healon
SODIUM HYALURONATE [HYALURONIC ACID] WITH CHONDROITIN SULPHATE			
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.35 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.4 ml syringe.....	64.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.5 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.55 ml syringe.....	74.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.75 ml syringe.....	67.00	1	Viscoat
Other			
DISODIUM EDETATE			
Inj 150 mg per ml, 20 ml ampoule			
Inj 150 mg per ml, 20 ml vial			
Inj 150 mg per ml, 100 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

RIBOFLAVIN 5-PHOSPHATE
Soln trans epithelial riboflavin
Inj 0.1%
Inj 0.1% plus 20% dextran T500
Inj 0.1% plus hydroxypropyl methylcellulose

Glaucoma Preparations

Beta Blockers

BETAXOLOL
Eye drops 0.25%
Eye drops 0.5%

TIMOLOL
Eye drops 0.25%2.42 5 ml Arrow-Timolol
Eye drops 0.5%2.50 5 ml Arrow-Timolol
➔ Eye drops 0.5%, gel forming – **Restricted:** For continuation only

Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE
Tab 250 mg – **5% DV Sep-25 to 2027**13.96 100 **Medsurge**
Inj 500 mg

BRINZOLAMIDE
Eye drops 1% – **5% DV Dec-24 to 2027**5.11 5 ml **Azopt**

DORZOLAMIDE – **Restricted:** For continuation only
➔ Eye drops 2%

DORZOLAMIDE WITH TIMOLOL
Eye drops 2% with timolol 0.5% – **5% DV Feb-25 to 2027**3.58 5 ml **Dortimopt**

Miotics

ACETYLCHOLINE CHLORIDE
Inj 20 mg vial with diluent

CARBACHOL
Inj 150 mcg vial

PILOCARPINE HYDROCHLORIDE
Eye drops 1%4.26 15 ml Isopto Carpine
Eye drops 2%5.35 15 ml Isopto Carpine
Eye drops 4%7.99 15 ml Isopto Carpine

PILOCARPINE NITRATE
Eye drops 2%, single dose

Prostaglandin Analogues

BIMATOPROST
Eye drops 0.03% – **5% DV Jan-25 to 2027**5.15 3 ml **Lumigan**

LATANOPROST
Eye drops 0.005% – **5% DV Mar-25 to 2027**2.08 2.5 ml **Teva**

LATANOPROST WITH TIMOLOL
Eye drops 0.005% with timolol 0.5%4.95 2.5 ml Arrow - Lattim

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
TRAVOPROST Eye drops 0.004% – 5% DV Dec-24 to 2027	6.80	2.5 ml	Travatan
Sympathomimetics			
APRACLOPIDINE Eye drops 0.5%	19.77	5 ml	Iopidine
BRIMONIDINE TARTRATE Eye drops 0.2% – 5% DV Mar-25 to 2027	5.16	5 ml	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE Eye drops 0.2% with timolol 0.5% – 5% DV Dec-24 to 2027	7.13	5 ml	Combigan
Mydriatics and Cycloplegics			
Anticholinergic Agents			
ATROPINE SULPHATE Eye drops 0.5% Eye drops 1%, single dose Eye drops 1%	18.27	15 ml	Atropt
CYCLOPENTOLATE HYDROCHLORIDE Eye drops 0.5%, single dose Eye drops 1%	68.52	15 ml	Cyclogyl
TROPICAMIDE Eye drops 0.5%	33.89	15 ml	Mydriacyl
Eye drops 0.5%, single dose Eye drops 1%	54.71	15 ml	Mydriacyl
Eye drops 1%, single dose			
Sympathomimetics			
PHENYLEPHRINE HYDROCHLORIDE Eye drops 2.5%, single dose Eye drops 10%, single dose			
Ocular Lubricants			
CARBOMER Ophthalmic gel 0.2%			
CARMELLOSE SODIUM Eye drops 0.5% Eye drops 0.5%, single dose Eye drops 1% Eye drops 1%, single dose			
HYPROMELLOSE Eye drops 0.5%	19.50	15 ml	Methopt
Ophthalmic gel 0.3%			
HYPROMELLOSE WITH DEXTRAN Eye drops 0.3% with dextran 0.1%	2.30	15 ml	Poly-Tears
Eye drops 0.3% with dextran 0.1%, single dose			

SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN Eye oint 42.5% with soft white paraffin 57.3%			
PARAFFIN LIQUID WITH WOOL FAT Eye oint 3% with wool fat 3%	3.63	3.5 g	Poly-Visc
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL Eye drops 0.4% with propylene glycol 0.3%, 10 ml bottle Note: Only for use in compounding an eye drop formulation			
Eye drops 0.4% with propylene glycol 0.3% preservative free, single dose....	10.06	28	Systane Unit Dose
POLYVINYL ALCOHOL WITH POVIDONE Eye drops 1.4% with povidone 0.6%, single dose			
RETINOL PALMITATE Oint 138 mcg per g	3.80	5 g	VitA-POS
SODIUM HYALURONATE [HYALURONIC ACID] Eye drops 1 mg per ml – 5% DV Dec-24 to 2027	13.58	10 ml	Hylo-Fresh

Other Otolological Preparations

ACETIC ACID WITH PROPYLENE GLYCOL Ear drops 2.3% with propylene glycol 2.8%
DOCUSATE SODIUM Ear drops 0.5%

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Agents Used in the Treatment of Poisonings			
Antidotes			
ACETYLCYSTEINE			
Tab eff 200 mg			
Inj 200 mg per ml, 10 ml vial.....	42.99	10	Hikma Acetylcysteine
Inj 200 mg per ml, 10 ml ampoule – 5% DV Apr-25 to 2027	42.99	10	DBL Acetylcysteine
AMYL NITRITE			
Liq 98% in 3 ml capsule			
DIGOXIN IMMUNE FAB			
Inj 38 mg vial			
Inj 40 mg vial			
ETHANOL			
Liq 96%			
ETHANOL WITH GLUCOSE			
Inj 10% with glucose 5%, 500 ml bottle			
ETHANOL, DEHYDRATED			
Inj 100%, 5 ml ampoule			
Inj 96%			
FLUMAZENIL			
Inj 0.1 mg per ml, 5 ml ampoule – 5% DV Dec-24 to 2027	44.00	5	Flumazenil-Baxter
HYDROXOCOBALAMIN			
Inj 5 g vial			
Inj 2.5 g vial			
NALOXONE HYDROCHLORIDE			
Inj 400 mcg per ml, 1 ml ampoule – 5% DV Apr-25 to 2027	13.29	5	DBL Naloxone Hydrochloride
PRALIDOXIME CHLORIDE			
Inj 1 g vial			
PRALIDOXIME IODIDE			
Inj 25 mg per ml, 20 ml ampoule			
SODIUM NITRITE			
Inj 30 mg per ml, 10 ml ampoule			
SODIUM THIOSULFATE			
Inj 250 mg per ml, 100 ml vial			
Inj 250 mg per ml, 10 ml vial			
Inj 250 mg per ml. 50 ml vial			
Inj 500 mg per ml, 10 ml vial			
Inj 500 mg per ml, 20 ml ampoule			
SOYA OIL			
Inj 20%, 500 ml bag			
Inj 20%, 500 ml bottle			

Antitoxins**BOTULISM ANTITOXIN**

Inj 250 ml vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

DIPHTHERIA ANTITOXIN

Inj 10,000 iu vial

Antivenoms

RED BACK SPIDER ANTIVENOM

Inj 500 u vial

SNAKE ANTIVENOM

Inj 50 ml vial

Removal and Elimination

CHARCOAL

Oral liq 200 mg per ml	59.85	250 ml	Carbasorb-X
------------------------------	-------	--------	-------------

DEFERASIROX – **Restricted** see terms [below](#)

⚡ Tab 125 mg dispersible	276.00	28	Exjade
⚡ Tab 250 mg dispersible	552.00	28	Exjade
⚡ Tab 500 mg dispersible	1,105.00	28	Exjade

➡ **Restricted (RS1444)**

Initiation

Haematologist

Re-assessment required after 2 years

All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μ L) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per μ L).

Continuation

Haematologist

Re-assessment required after 2 years

Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels. .

DEFERIPRONE – **Restricted** see terms [below](#)

⚡ Tab 500 mg	533.17	100	Ferriprox
⚡ Oral liq 100 mg per ml	266.59	250 ml	Ferriprox

➡ **Restricted (RS1445)**

Initiation

Patient has been diagnosed with chronic iron overload due to congenital inherited anaemia or acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

Inj 500 mg vial	332.88	10	DBL Desferrioxamine Mesylate for Inj BP
-----------------------	--------	----	--

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DICOBALT EDEATE			
Inj 15 mg per ml, 20 ml ampoule			
DIMERCAPROL			
Inj 50 mg per ml, 2 ml ampoule			
DIMERCAPTOSUCCINIC ACID			
Cap 100 mg			e.g. PCNZ, Optimus Healthcare, Chemet
Cap 200 mg			e.g. PCNZ, Optimus Healthcare, Chemet
SODIUM CALCIUM EDEATE			
Inj 50 mg per ml, 10 ml ampoule			
Inj 200 mg per ml, 2.5 ml ampoule			
Inj 200 mg per ml, 5 ml ampoule			

Antiseptics and Disinfectants

CHLORHEXIDINE			
Soln 0.1%			
Soln 4%			
Soln 5%	15.50	500 ml	healthE
CHLORHEXIDINE WITH CETRIMIDE			
Crm 0.1% with cetrimide 0.5%			
Foaming soln 0.5% with cetrimide 0.5%			
CHLORHEXIDINE WITH ETHANOL			
Soln 0.5% with ethanol 70%			
Soln 2% with ethanol 70%			
Soln 0.5% with ethanol 70%, non-staining (pink) 25 ml	1.55	1	healthE
IODINE WITH ETHANOL			
Soln 1% with ethanol 70%			
ISOPROPYL ALCOHOL			
Soln 70%, 500 ml	5.65	1	healthE
POVIDONE-IODINE			
↓ Vaginal tab 200 mg			
➔ Restricted (RS1354)			
Initiation			
Rectal administration pre-prostate biopsy.			
Oint 10%.....	7.40	65 g	Betadine
Soln 10%.....	4.99	100 ml	Riodine
Soln 5%			
Soln 7.5%			
Soln 10%,	3.83	15 ml	Riodine
	6.99	500 ml	Riodine
Pad 10%			
Swab set 10%			
POVIDONE-IODINE WITH ETHANOL			
Soln 10% with ethanol 30%			
Soln 10% with ethanol 70%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM HYPOCHLORITE			
Soln			
Contrast Media			
Iodinated X-ray Contrast Media			
DIATRIZOATE SODIUM			
Oral liq 370 mg per ml, 10 ml sachet.....	156.12	50	Ioscan
IODISED OIL			
Inj 38% w/w (480 mg per ml), 10 ml ampoule	410.00	1	Lipiodol Ultra Fluid
IODIXANOL			
Inj 270 mg per ml (iodine equivalent), 50 ml bottle.....	305.00	10	Visipaque
Inj 270 mg per ml (iodine equivalent), 100 ml bottle.....	560.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 50 ml bottle.....	310.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 100 ml bottle.....	565.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 200 ml bottle.....	1,130.00	10	Visipaque
IOHEXOL			
Inj 240 mg per ml (iodine equivalent), 50 ml bottle.....	140.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 10 ml bottle.....	109.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 20 ml bottle.....	110.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 50 ml bottle.....	145.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 100 ml bottle.....	240.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 50 ml bottle.....	150.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 100 ml bottle.....	252.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 200 ml bottle.....	522.00	10	Omnipaque
Inj 350 mg per ml, 500 ml bottle	785.00	6	Omnipaque
<i>(Omnipaque Inj 300 mg per ml (iodine equivalent), 20 ml bottle to be delisted 1 October 2026)</i>			
Non-iodinated X-ray Contrast Media			
BARIUM SULPHATE			
Oral liq 400 mg per ml (40% w/v, 30% w/w), bottle	17.39	148 g	Varibar - Thin Liquid
Oral liq 400 mg per ml (40% w/v), bottle	189.15	250 ml	Varibar - Honey
	38.40	240 ml	Varibar - Nectar
	159.05	230 ml	Varibar - Pudding
Grans for oral liq 960 mg per g (96% w/w), 176 g bottle	530.00	24	Vanilla SilQ MD
Grans for oral liq 980 mg per g (98% w/w), 310 g bottle	490.00	24	Vanilla SilQ HD
Oral liq 20.9 mg per ml (2.1% w/v, 2% w/w), 450 ml bottle	97.50	12	Readi-CAT 2
Oral liq 1 mg per ml (0.1% w/v, 0.1% w/w), 450 ml bottle	15.95	1	Neulumex
	191.40	12	Neulumex
Oral liq 400 mg per ml (40% w/v, 30% w/w), 20 ml bottle	52.35	3	Tagitol V
CITRIC ACID WITH SODIUM BICARBONATE			
Powder 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet.....	90.25	50 g	E-Z-Gas II

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Paramagnetic Contrast Media			
GADOBUTROL			
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 5 ml prefilled syringe.....	126.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 7.5 ml prefilled syringe.....	189.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 10 ml prefilled syringe.....	245.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 15 ml vial.....	735.00	10	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 65 ml bottle.....	3,120.00	10	Gadovist 1.0
GADOTERIC ACID			
Inj 279.30 mg per ml, 10 ml prefilled syringe			<i>e.g. Clariscan</i>
Inj 279.30 mg per ml, 10 ml vial			<i>e.g. Clariscan</i>
Inj 279.30 mg per ml, 15 ml prefilled syringe			<i>e.g. Clariscan</i>
Inj 279.30 mg per ml, 20 ml vial			<i>e.g. Clariscan</i>
Inj 279.30 mg per ml, 5 ml vial			<i>e.g. Clariscan</i>
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml prefilled syringe.....	172.00	10	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml bottle.....	25.35	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml prefilled syringe.....	258.00	10	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml prefilled syringe.....	344.00	10	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml bottle.....	14.30	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml bottle.....	28.90	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 5 ml bottle.....	9.10	1	Dotarem
GADOXETATE DISODIUM			
Inj 181.43 mg per ml (equivalent to 0.25 mmol per ml), 10 ml prefilled syringe.....	300.00	1	Primovist
MEGLUMINE GADOPENTETATE			
Inj 469 mg per ml, 10 ml prefilled syringe.....	95.00	5	Magnevist
Inj 469 mg per ml, 10 ml vial.....	185.00	10	Magnevist
MEGLUMINE IOTROXATE			
Inj 105 mg per ml, 100 ml bottle.....	169.15	100 ml	Biliscopein
Ultrasound Contrast Media			
PERFLUTREN			
Inj 1.1 mg per ml, 1.5 ml vial.....	180.00	1	Definity
	720.00	4	Definity
Diagnostic Agents			
ARGININE			
Inj 50 mg per ml, 500 ml bottle			
Inj 100 mg per ml, 300 ml bottle			
HISTAMINE ACID PHOSPHATE			
Nebuliser soln 0.6%, 10 ml vial			
Nebuliser soln 2.5%, 10 ml vial			
Nebuliser soln 5%, 10 ml vial			
MANNITOL			
Powder for inhalation			<i>e.g. Aridol</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METHACHOLINE CHLORIDE			
Powder 100 mg			
SECRETIN PENTAHYDROCHLORIDE			
Inj 100 u vial			
Inj 80 u vial			
Inj 100 u ampoule			
SINCALIDE			
Inj 5 mcg per vial			

Diagnostic Dyes

BONNEY'S BLUE DYE			
Soln			
INDIGO CARMINE			
Inj 4 mg per ml, 5 ml ampoule			
Inj 8 mg per ml, 5 ml ampoule			
INDOCYANINE GREEN			
Inj 25 mg vial			
METHYLTHIONINIUM CHLORIDE [METHYLENE BLUE]			
Inj 5 mg per ml, 10 ml ampoule	259.57	5	Proveblue
PATENT BLUE V			
Inj 2.5%, 2 ml ampoule	440.00	5	Obex Medical
Inj 2.5%, 5 ml prefilled syringe	435.00	5	InterPharma

Irrigation Solutions

CHLORHEXIDINE WITH CETRIMIDE
 ‡ Irrigation soln 0.015% with cetrimide 0.15%, 500 ml bottle

➔ **Restricted (RS1683)**

Initiation

Re-assessment required after 3 months

All of the following:

- 1 Patient has burns that are greater than 30% of total body surface area (BSA); and
- 2 For use in the perioperative preparation and cleansing of large burn areas requiring debridement/skin grafting; and
- 3 The use of 30 ml ampoules is impractical due to the size of the area to be covered.

Continuation

Re-assessment required after 3 months

The treatment remains appropriate for the patient and the patient is benefiting from the treatment.

Irrigation soln 0.015% with cetrimide 0.15%, 100 ml bottle

Irrigation soln 0.015% with cetrimide 0.15%, 30 ml ampoule – **5% DV**

Sep-25 to 2028 29.70 30 **LumaCina**

GLYCINE

Irrigation soln 1.5%, 3,000 ml bag 96.28 4 B Braun

SODIUM CHLORIDE

Irrigation soln 0.9%, 3,000 ml bag 80.00 4 B Braun

Irrigation soln 0.9%, 30 ml ampoule 13.25 20 InterPharma

Irrigation soln 0.9%, 1,000 ml bottle 19.50 10 Baxter Sodium Chloride 0.9%

Irrigation soln 0.9%, 250 ml bottle 24.60 12 Fresenius Kabi

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
WATER			
Irrigation soln, 3,000 ml bag	84.52	4	B Braun
Irrigation soln, 1,000 ml bottle	19.50	10	Baxter Water for Irrigation
Irrigation soln, 250 ml bottle	26.40	12	Fresenius Kabi

Surgical Preparations

BISMUTH SUBNITRATE AND IODOFORM PARAFFIN

Paste

DIMETHYL SULFOXIDE

Soln 50%

Soln 99%

PHENOL

Inj 6%, 10 ml ampoule

PHENOL WITH IOXAGLIC ACID

Inj 12%, 10 ml ampoule

SODIUM HYDROXIDE

Soln 10%

TROMETAMOL

Inj 36 mg per ml, 500 ml bottle

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Cardioplegia Solutions			
ELECTROLYTES			
Inj 15 mmol/l sodium chloride, 9 mmol/l potassium chloride, 1 mmol/l potassium hydrogen 2-ketoglutarate, 4 mmol/l magnesium chloride, 18 mmol/l histidine hydrochloride, 180 mmol/l histidine, 2 mmol/l tryptophan, 30 mmol/l mannitol, 0.015 mmol/l calcium chloride, 1,000 ml bag			<i>e.g. Custodiol-HTK</i>
Inj aspartic acid 10.43 mg per ml, citric acid 0.22476 mg per ml, glutamic acid 11.53 mg per ml, sodium phosphate 0.1725 mg per ml, potassium chloride 2.15211 mg per ml, sodium citrate 1.80768 mg per ml, sodium hydroxide 6.31 mg per ml and trometamol 11.2369 mg per ml, 364 ml bag			<i>e.g. Cardioplegia Enriched Paed. Soln.</i>
Inj aspartic acid 8.481 mg per ml, citric acid 0.8188 mg per ml, glutamic acid 9.375 mg per ml, sodium phosphate 0.6285 mg per ml, potassium chloride 2.5 mg per ml, sodium citrate 6.585 mg per ml, sodium hydroxide 5.133 mg per ml and trometamol 9.097 mg per ml, 527 ml bag			<i>e.g. Cardioplegia Enriched Solution</i>
Inj citric acid 0.07973 mg per ml, sodium phosphate 0.06119 mg per ml, potassium chloride 2.181 mg per ml, sodium chloride 1.788 mg ml, sodium citrate 0.6412 mg per ml and trometamol 5.9 mg per ml, 523 ml bag			<i>e.g. Cardioplegia Base Solution</i>
Inj 110 mmol/l sodium, 16 mmol/l potassium, 1.2 mmol/l calcium, 16 mmol/l magnesium and 160 mmol/l chloride, 1,000 ml bag			<i>e.g. Cardioplegia Solution AHB7832</i>
Inj 143 mmol/l sodium, 16 mmol/l potassium, 16 mmol/l magnesium and 1.2 mmol/l calcium, 1,000 ml bag			<i>e.g. Cardioplegia Electrolyte Solution</i>
MONOSODIUM GLUTAMATE WITH SODIUM ASPARTATE			
Inj 42.68 mg with sodium aspartate 39.48 mg per ml, 250 ml bottle			
MONOSODIUM L-ASPARTATE			
Inj 14 mmol per 10 ml, 10 ml			

Cold Storage Solutions

SODIUM WITH POTASSIUM

Inj 29 mmol/l with potassium 125 mmol/l, 1,000 ml bag

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Extemporaneously Compounded Preparations			
ACETIC ACID Liq			
ALUM Powder BP			
ARACHIS OIL [PEANUT OIL] Liq			
ASCORBIC ACID Powder			
BENZOIN Tincture compound BP			
BISMUTH SUBGALLATE Powder			
BORIC ACID Powder			
CARBOXYMETHYLCELLULOSE Soln 1.5%			
CETRIMIDE Soln 40%			
CHLORHEXIDINE GLUCONATE Soln 20 %			
CHLOROFORM Liq BP			
CITRIC ACID Powder BP			
CLOVE OIL Liq			
COAL TAR Soln BP.....	46.00	200 ml	Midwest
CODEINE PHOSPHATE Powder			
COLLODION FLEXIBLE Liq			
COMPOUND HYDROXYBENZOATE Soln	36.00	100 ml	Midwest
CYSTEAMINE HYDROCHLORIDE Powder			
DISODIUM HYDROGEN PHOSPHATE WITH SODIUM DIHYDROGEN PHOSPHATE Inj 37.46 mg with sodium dihydrogen phosphate 47.7 mg in 1.5 ml ampoule			
DITHRANOL Powder			
GLUCOSE [DEXTROSE] Powder			

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GLYCERIN WITH SODIUM SACCHARIN			
Suspension.....	38.00	473 ml	Ora-Sweet SF
GLYCERIN WITH SUCROSE			
Suspension.....	38.00	473 ml	Ora-Sweet
GLYCEROL			
Liq.....	3.23	500 ml	healthE Glycerol BP Liquid
HYDROCORTISONE			
Powder	49.95	25 g	ABM
LACTOSE			
Powder			
MAGNESIUM HYDROXIDE			
Paste			
MENTHOL			
Crystals			
METHADONE HYDROCHLORIDE			
Powder			
METHYL HYDROXYBENZOATE			
Powder	11.00	25 g	Midwest
METHYLCELLULOSE			
Powder	44.00	100 g	Midwest
Suspension.....	38.00	473 ml	Ora-Plus
<i>(Midwest Powder to be delisted 1 February 2028)</i>			
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN			
Suspension.....	38.00	473 ml	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE			
Suspension.....	38.00	473 ml	Ora-Blend
OLIVE OIL			
Liq			
PARAFFIN			
Liq			
PHENOBARBITONE SODIUM			
Powder			
PHENOL			
Liq			
PILOCARPINE NITRATE			
Powder			
POLYHEXAMETHYLENE BIGUANIDE			
Liq			
POVIDONE K30			
Powder			
SALICYLIC ACID			
Powder			
SILVER NITRATE			
Crystals			
SODIUM BICARBONATE			
Powder BP	13.50	500 g	Midwest

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM CITRATE Powder			
SODIUM METABISULFITE Powder			
STARCH Powder			
SULPHUR Precipitated Sublimed			
SYRUP Liq (pharmaceutical grade).....	25.00	500 ml	Midwest
THEOBROMA OIL Oint			
TRI-SODIUM CITRATE Crystals			
TRICHLORACETIC ACID Grans			
UREA Powder BP			
WOOL FAT Oint, anhydrous			
XANTHAN Gum 1%			
ZINC OXIDE Powder			

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

Food Modules

Carbohydrate

➡ **Restricted (RS1467)**

Initiation – Use as an additive

Any of the following:

- 1 Cystic fibrosis; or
- 2 Chronic kidney disease; or
- 3 Cancer in children; or
- 4 Cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 5 Faltering growth in an infant/child; or
- 6 Bronchopulmonary dysplasia; or
- 7 Premature and post premature infant; or
- 8 Inborn errors of metabolism.

Initiation – Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

CARBOHYDRATE SUPPLEMENT – **Restricted** see terms [above](#)

† Powder 96 g carbohydrate per 100 g, can 6.72 400 g Polycal

Fat

➡ **Restricted (RS1468)**

Initiation – Use as an additive

Any of the following:

- 1 Patient has inborn errors of metabolism; or
- 2 Faltering growth in an infant/child; or
- 3 Bronchopulmonary dysplasia; or
- 4 Fat malabsorption; or
- 5 Lymphangiectasia; or
- 6 Short bowel syndrome; or
- 7 Infants with necrotising enterocolitis; or
- 8 Biliary atresia; or
- 9 For use in a ketogenic diet; or
- 10 Chyle leak; or
- 11 Ascites; or
- 12 Patient has increased energy requirements, and for whom dietary measures have not been successful.

Initiation – Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk. .

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

LONG-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms [above](#)

† Liquid 50 g fat per 100 ml, bottle 15.38 200 ml Calogen (neutral)
Calogen (strawberry)

MEDIUM-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms [above](#)

† Liquid 95 g fat per 100 ml, bottle 37.50 500 ml MCT Oil
† Liquid 50 g fat per 100 ml, 250 ml bottle 143.65 4 Liquigen

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

WALNUT OIL – **Restricted** see terms [on the previous page](#)

† Liq

Protein

➔ **Restricted (RS1469)**

Initiation – Use as an additive

Either:

- 1 Protein losing enteropathy; or
- 2 High protein needs.

Initiation – Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

PROTEIN SUPPLEMENT – **Restricted** see terms [above](#)

† Powder 5 g protein, 0.67 g carbohydrate and 0.6 g fat per 6.6 g, 275 g can			
† Powder 6 g protein per 7 g, can	8.95	227 g	Resource Beneprotein
† Powder 89 g protein, less than 1.5 g carbohydrate and 2 g fat per 100 g, can	13.82	225 g	Protifar

Other Supplements

CARBOHYDRATE AND FAT SUPPLEMENT – **Restricted** see terms [below](#)

† Powder 72.7 g carbohydrate and 22.3 g fat per 100 g, can	71.77	400 g	Duocal Super Soluble Powder
--	-------	-------	-----------------------------

➔ **Restricted (RS1212)**

Initiation

Both:

- 1 Infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 Cystic fibrosis; or
 - 2.2 Cancer in children; or
 - 2.3 Faltering growth; or
 - 2.4 Bronchopulmonary dysplasia; or
 - 2.5 Premature and post premature infants.

HUMAN MILK FORTIFIER

Powder 0.325 g protein, 0.37 g carbohydrate and 0.175 g fat per 1 g sachet	33.48	50	Human Milk Fortifier <i>e.g. FM 85</i>
Powder 0.2 g protein, 0.7 g carbohydrate and 0.02 g fat per 1 g sachet			

Food/Fluid Thickeners

NOTE:

While pre-thickened drinks and supplements have not been included in Section H, Health NZ Hospitals may continue to use such products for patients with dysphagia, provided that:

- use was established prior to 1 July 2013; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- the product has not been specifically considered and excluded by Pharmac; and
- use of the product conforms to any applicable indication restrictions for similar products that are listed in Section H (for example, use of thickened high protein products should be in line with the restriction for high protein oral feed in Section H).

Pharmac intends to make a further decision in relation to pre-thickened drinks and supplements in the future, and will notify of any change to this situation.

CAROB BEAN GUM WITH MAIZE STARCH AND MALTODEXTRIN			
Powder	24.00	380 g	Aptamil Feed Thickener
GUAR GUM			
Powder			<i>e.g. Guarcol</i>
MAIZE STARCH			
Powder	8.29	300 g	Nutilis
MALTODEXTRIN WITH XANTHAN GUM			
Powder			<i>e.g. Instant Thick</i>
MALTODEXTRIN WITH XANTHAN GUM AND ASCORBIC ACID			
Powder			<i>e.g. Easy Thick</i>

Metabolic Products

➔ **Restricted (RS2047)**

Initiation

Either:

- 1 For the dietary management of inherited metabolic disease; or
- 2 Patient has adrenoleukodystrophy.

Supplements for Glutaric Aciduria Type 1

AMINO ACID FORMULA (WITHOUT LYSINE AND LOW TRYPTOPHAN) – **Restricted** see terms [above](#)

⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can		<i>e.g. GA1 Anamix Infant</i>
⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can		<i>e.g. XLYS Low TRY Maxamaid</i>

AMINO ACID FORMULA (WITHOUT LYSINE) – **Restricted** see terms [above](#)

⚡ Powder (neutral) 5 g protein, 5.4 g carbohydrate, 2.3 g fat and 2 g fibre per 18 g sachet.....	750.30	30	GA1 Anamix Junior
⚡ Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30	GA Explore 5
⚡ Powder, 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 3.7 g fibre per 100 g, 400 g can.....	260.00	400 g	GA1 Anamix Infant

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Supplements for Homocystinuria

AMINO ACID FORMULA (WITHOUT METHIONINE) – **Restricted** see terms [on the previous page](#)

† Powder (neutral), 10 g protein, 11.5 g carbohydrate and 4.5 g fat per 36 g sachet.....	750.30	30	HCU Anamix Junior
† Powder, 15 g protein, 3.5 g carbohydrate, 0.55 g fat per 25 g sachet.....	1,048.95	30	HCU Express 15
† Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30	HCU Explore 5
† Powder (neutral) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	480.42	500 g	XMET Maxamum
† Powder (unflavoured) 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	260.00	400 g	HCU Anamix Infant
† Liquid (juicy berries), 20 g protein, 12.63 g carbohydrate and 0.46 g fat per 125 ml bottle.....	1,684.80	30	HCU Lophlex LQ
† Liquid (orange), 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle.....	941.40	36	HCU Anamix Junior LQ

Supplements for MSUD and Short chain enoyl coA hydratase deficiency

AMINO ACID FORMULA (WITHOUT Isoleucine, Leucine and Valine) – **Restricted** see terms [on the previous page](#)

† Powder (neutral) 10 g protein, 11.5 g carbohydrate and 4.5 g fat per 36 g sachet.....	750.00	30	MSUD Anamix Junior
† Powder, 15 g protein, 3.5 g carbohydrate, 0.6 g fat per 25 g sachet.....	1,048.95	30	MSUD Express 15
† Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30	MSUD Explore 5
† Powder (orange) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	454.71	500 g	MSUD Maxamum
† Powder (unflavoured) 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	260.00	400 g	MSUD Anamix Infant
† Powder (unflavoured) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	454.71	500 g	MSUD Maxamum
† Liquid (juicy berries), 20 g protein, 12.63 g carbohydrate and 0.46 g fat per 125 ml pouch.....	1,684.80	30	MSUD Lophlex LQ 20
† Liquid (orange) 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle.....	941.40	36	MSUD Anamix Junior LQ

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Supplements for Phenylketonuria			
AMINO ACID FORMULA (WITHOUT PHENYLALANINE) – Restricted see terms on page 294			
† Tab 8.33 mg	99.00	75	Phlexy 10
† Powder (Berry), 5.0 g protein, 14 g carbohydrate, 0 g fat per 20 g sachet.....	449.28	60	PKU Restore Powder
† Powder (Lemon), 20 g protein, 3.9 g carbohydrate, 0.8 g fat per 34 g sachet.....	883.50	30	PKU Express 20
† Powder (Neutral), 20 g protein, 4.8 g carbohydrate, 0.8 g fat per 34 g sachet.....	883.50	30	PKU Express 20
† Powder (Neutral), 5.0 g protein, 5.2 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	220.88	30	PKU Explore 5
† Powder (Orange), 10 g protein, 9.8 g carbohydrate, 0.4 g fat per 25 g sachet.....	441.75	30	PKU Explore 10
† Powder (Orange), 20 g protein, 3.9 g carbohydrate, 0.8 g fat per 34 g sachet.....	883.50	30	PKU Express 20
† Powder (Orange), 5.0 g protein, 14 g carbohydrate, 0 g fat per 20 g sachet.....	449.28	60	PKU Restore Powder
† Powder (Raspberry), 10 g protein, 9.8 g carbohydrate, 0.4 g fat per 25 g sachet.....	441.75	30	PKU Explore 10
† Powder (Tropical), 20 g protein, 3.9 g carbohydrate, 0.8 g fat per 34 g sachet.....	883.50	30	PKU Express 20
† Powder (berry) 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet.....	936.00	30	PKU Lophlex Powder
† Powder (chocolate) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	393.00	30	PKU Anamix Junior
† Powder (neutral) 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet.....	936.00	30	PKU Lophlex Powder
† Powder (neutral) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	393.00	30	PKU Anamix Junior
† Powder (orange) 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet.....	936.00	30	PKU Lophlex Powder
† Powder (orange) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	393.00	30	PKU Anamix Junior
† Powder (unflavoured), 5 g protein, 4.8 g carbohydrate per 12.5 g sachets	234.00	30	PKU First Spoon
† Powder (vanilla) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	393.00	30	PKU Anamix Junior
† Powder (Neutral), 14.3 g protein, 25 g fat per 100 g, 4 × 400 g can.....	715.16	1,600 g	PKU Start
† Powder (orange) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	320.00	500 g	XP Maxamum
† Powder (unflavoured) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	320.00	500 g	XP Maxamum
† Powder 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	174.72	400 g	PKU Anamix Infant
† Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle	13.10	1	PKU Anamix Junior LQ (Berry) PKU Anamix Junior LQ (Orange)
† Liquid (juicy berries) 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 62.5 ml bottle	939.00	60	PKU Lophlex LQ 10
† Liquid (juicy berries) 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle	936.00	30	PKU Lophlex LQ 20

† Item restricted (see ➡ above); ‡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
† Liquid (juicy orange) 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle	936.00	30	PKU Lophlex LQ 20
† Liquid (juicy tropical) 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 125 ml bottle	936.00	30	PKU Lophlex LQ 20
† Liquid 6.7 g protein, 5.1 g carbohydrate and 2 g fat per 100 ml, 250 ml carton.....	540.00	18	Easiphen Liquid
† Semi-solid 18.3 g protein, 18.5 g carbohydrate and 0.92 g fibre per 100 g, 109 g pot.....	1,123.20	36	PKU Lophlex Sensations 20 (berries)
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME PHENYLALANINE – Restricted see terms on page 294			
† Powder (Neutral), 10 g protein, 0.5 g carbohydrate, 0.6 g fat per 15 g sachet.....	449.28	30	PKU Build 10
† Powder (neutral), 15 g protein, 15 g carbohydrate, 4.5 g fat per 40 g sachet.....	673.92	30	Glytactin Bettermilk
† Powder (unflavoured) 10 g protein, 4 g carbohydrate per 12.5 g sachet	468.00	30	PKU GMPPro Mix-In
† Powder 20 g protein, 1.7 g carbohydrate per 31 g sachet.....	898.56	30	PKU Build 20 Raspberry Lemonade
† Powder 20 g protein, 1.7 g carbohydrate per 32 g sachet.....	898.56	30	PKU Build 20 Smooth
† Powder 20 g protein, 1.7 g carbohydrate per 33 g sachet.....	898.56	30	PKU Build 20 Chocolate
† Powder 20 g protein, 4.9 g carbohydrate per 33.4 g sachet.....	936.00	30	PKU Build 20 Vanilla
† Powder 20 g protein, 6.0 g carbohydrate per 35 g sachet.....	930.00	30	PKU GMPPro Ultra
† Powder 20 g protein, 6.3 g carbohydrate per 35 g sachet.....	930.00	30	PKU GMPPro Ultra Lemonade
† Powder 20 g protein, 6.7 g carbohydrate per 35 g sachet.....	930.00	30	PKU GMPPro Ultra Vanilla
† Liquid (Coffee Mocha), 15 g protein, 3.1 g carbohydrate, 4.6 g fat 250 ml, carton.....	684.45	30	PKU sphere20 Lemon
† Liquid (chocolate), 15 g protein, 22 g carbohydrate, 5.3 g fat per 250 ml, carton.....	684.45	30	PKU sphere20 Chocolate
† Liquid (neutral), 10 g protein, 8.5 g carbohydrate per 250 ml carton.....	280.80	18	PKU sphere20 Red Berry
† Liquid (original), 15 g protein, 22 g carbohydrate, 5.3 g fat per 250 ml, carton.....	684.45	30	PKU sphere20 Vanilla
† Liquid (vanilla), 15 g protein, 3.3 g carbohydrate, 4.6 g fat per 250 ml, carton.....	684.45	30	PKU sphere20 Banana
			PKU Glytactin RTD 15 Lite
			PKU Glytactin RTD 15
			PKU GMPPro LQ
			PKU Glytactin RTD 15
			PKU Glytactin RTD 15 Lite

Protein Free Supplements

PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDRATE, FAT WITH ADDED VITAMINS AND MINERALS –

Restricted see terms [on page 294](#)

† Powder (neutral) nil added protein and 67 g carbohydrate per 100 g, 400 g can.....	49.29	400 g	Energivit
--	-------	-------	-----------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Supplements for Tyrosinaemia			
AMINO ACID FORMULA (WITHOUT PHENYLALANINE AND TYROSINE) – Restricted see terms on page 294			
† Powder (neutral) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	471.00	30	TYR Anamix Junior
† Powder (neutral), 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30	TYR Explore 5
† Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	260.00	400 g	TYR Anamix Infant
† Liquid (orange) 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle.....	941.40	36	TYR Anamix Junior LQ
† Liquid (juicy berries), 20 g protein, 12.75 g carbohydrate and 0.46 g fat and 0 g fibre per 125 ml pouch.....	1,684.80	30	TYR Lophlex LQ 20
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME TYROSINE AND PHENYLALANINE – Restricted see terms on page 294			
† Powder (Red Berry), 20 g protein, 6.3 carbohydrate, 1.6 g fat per 35 g sachet.....	1,398.60	30	TYR Sphere 20
† Powder (Vanilla), 20 g protein, 6.0 g carbohydrate, 1.6 g fat per 35 g sachet.....	1,398.60	30	TYR Sphere 20
X-Linked Adrenoleukodystrophy Products			
GLYCEROL TRIERUCATE – Restricted see terms on page 294			
† Liquid, 1,000 ml bottle			
GLYCEROL TRIOLEATE – Restricted see terms on page 294			
† Liquid, bottle.....	131.80	500 ml	GTO Oil
Supplements for Glycogen Storage Disease			
HIGH AMYLOPECTIN CORN-STARCH – Restricted see terms on page 294			
† Powder 0 g protein, 53 g carbohydrate, 0 g fat per 60 g sachet.....	241.62	30	Glycosade
Supplements for Organic Acidaemias			
AMINO ACID FORMULA (WITHOUT Isoleucine, Methionine, Threonine and Valine) – Restricted see terms on page 294			
† Powder 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	260.00	400 g	MMA/PA Anamix Infant
AMINO ACID FORMULA (WITHOUT METHIONINE, Threonine and Valine) – Restricted see terms on page 294			
† Powder (neutral), 5 g protein, 5.4 g carbohydrate, 2.3 g fat and 2.0 g fibre per 18 g sachet.....	750.30	30	MMA/PA Anamix Junior
† Powder, 15 g protein, 3.4 g carbohydrate, 0.05 g fat per 25 g sachet.....	1,048.95	30	MMA/PA Express 15
† Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30	MMA/PA Explore 5
Single Dose Amino Acids			
ARGININE – Restricted see terms on page 294			
† Powder 1.7 g protein, 1.9 g carbohydrate per 4 g sachet.....	211.45	30	Arginine2000
CITRULLINE – Restricted see terms on page 294			
† Powder 0.8 g protein, 2.9 g carbohydrate per 4 g sachet.....	211.45	30	Citrulline1000
ISOLEUCINE – Restricted see terms on page 294			
† Powder 0.04 g protein, 3.8 g carbohydrate per 4 g sachet.....	141.05	30	Isoleucine50

† Item restricted (see ➡ above); ‡ Item restricted (see ➡ below)
e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LEUCINE – Restricted see terms on page 294			
† Powder 0.08 g protein, 3.7 g carbohydrate per 4 g sachet.....	141.05	30	Leucine100
PHENYLALANINE – Restricted see terms on page 294			
† Powder 0.04 g protein, 3.8 g carbohydrate per 4 g sachet.....	141.05	30	Phenylalanine50
TYROSINE – Restricted see terms on page 294			
† Powder 0.8 g protein, 2.9 g carbohydrate per 4 g sachet.....	211.45	30	Tyrosine1000
VALINE – Restricted see terms on page 294			
† Powder 0.04 g protein, 3.8 g carbohydrate per 4 g sachet.....	141.05	30	Valine50

Other Fat Modified Products

ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCERIDES – **Restricted** see terms [on page 294](#)

† Powder (neutral), 12.5 g protein, 60 g carbohydrate and 16.4 g fat per 100 g sachet	47.01	10	Emsogen
--	-------	----	---------

Essential Amino Acids

ESSENTIAL AMINO ACID FORMULA – **Restricted** see terms [on page 294](#)

† Powder (neutral) 79 g protein per 100 g, 200 g can	313.73	200 g	Essential Amino Acid Mix
--	--------	-------	--------------------------

Specialised Formulas

Diabetic Products

→ **Restricted** (RS1215)

Initiation

Any of the following:

- 1 For patients with type I or type II diabetes suffering weight loss and malnutrition that requires nutritional support; or
- 2 For patients with pancreatic insufficiency; or
- 3 For patients who have, or are expected to, eat little or nothing for 5 days; or
- 4 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or
- 5 For use pre- and post-surgery; or
- 6 For patients being tube-fed; or
- 7 For tube-feeding as a transition from intravenous nutrition.

DIABETIC ORAL FEED 1 KCAL/ML – **Restricted** see terms [above](#)

† Liquid 4.9 g protein, 11.7 g carbohydrate, 3.8 g fat and 2 g fibre per 100 ml, 200 ml bottle	2.25	1	Diasip (strawberry) Diasip (vanilla)
---	------	---	---

LOW-GI ENTERAL FEED 1 KCAL/ML – **Restricted** see terms [above](#)

† Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 500 ml bottle.....	4.65	1	Glucerna Select
† Liquid 4.3 g protein, 11.3 g carbohydrate and 4.2 g fat per 100 ml, 1,000 ml bottle			<i>e.g. Nutrison Advanced Diason</i>

LOW-GI ORAL FEED 1 KCAL/ML – **Restricted** see terms [above](#)

† Liquid 7 g protein, 10.9 g carbohydrate, 2.7 g fat and 2 g fibre per 100 ml, 200 ml bottle	8.40	4	Nutren Diabetes (vanilla)
---	------	---	---------------------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Elemental and Semi-Elemental Products			
➔ Restricted (RS1216)			
Initiation			
Any of the following:			
1 Malabsorption; or			
2 Short bowel syndrome; or			
3 Enterocutaneous fistulas; or			
4 Eosinophilic enteritis (including oesophagitis); or			
5 Inflammatory bowel disease; or			
6 Acute pancreatitis where standard feeds are not tolerated; or			
7 Patients with multiple food allergies requiring enteral feeding.			
AMINO ACID ORAL FEED – Restricted see terms above			
⚡ Powder 11 g protein, 62 g carbohydrate and 1 g fat per sachet, 80 g sachet.....	45.00	10	Vivonex TEN
AMINO ACID ORAL FEED 0.8 KCAL/ML – Restricted see terms above			
⚡ Liquid 2.5 g protein, 11 g carbohydrate and 3.5 g fat per 100 ml, 250 ml carton.....	179.46	18	Elemental 028 Extra (grapefruit) Elemental 028 Extra (pineapple & orange) Elemental 028 Extra (summer fruits)
PEPTIDE-BASED ENTERAL FEED 1 KCAL/ML – Restricted see terms above			
⚡ Liquid 4 g protein, 17.7 g carbohydrate and 1.7 g fat per 100 ml, 500 ml bottle.....	7.47	1	Nutrison Advanced Peptisorb
PEPTIDE-BASED ENTERAL FEED 1.5 KCAL/ML – Restricted see terms above			
⚡ Liquid 6.75 g protein, 18.4 g carbohydrate and 5.5 g fat per 100 ml, 1,000 ml bottle.....	22.39	1	Vital
PEPTIDE-BASED ORAL FEED – Restricted see terms above			
⚡ Powder 13.7 g protein, 62.9 g carbohydrate and 17.5 g fat per 100 g, 400 g can			<i>e.g. Peptamen Junior</i>
⚡ Powder 13.8 g protein, 59 g carbohydrate and 18 g fat per 100 g, 400 g can			<i>e.g. MCT Peptide; MCT Peptide 1+</i>
PEPTIDE-BASED ORAL FEED 1 KCAL/ML – Restricted see terms above			
⚡ Liquid 5 g protein, 16 g carbohydrate and 1.69 g fat per 100 ml, carton.....	4.95	237 ml	Peptamen OS 1.0 (Vanilla)

Fat Modified Products

FAT-MODIFIED FEED – Restricted see terms below			
⚡ Powder 12.8 g protein, 68.6 g carbohydrate and 12.9 g fat per 100 g, can	62.90	400 g	Monogen

➔ **Restricted (RS1470)**

Initiation

Any of the following:

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

Hepatic Products

➔ **Restricted (RS1217)**

Initiation

For children (up to 18 years) who require a liver transplant.

HEPATIC ORAL FEED – **Restricted** see terms [above](#)

⬆ Powder 12 g protein, 56 g carbohydrate and 22 g fat per 100 g, can	93.97	400 g	Heparon Junior
--	-------	-------	----------------

High Calorie Products

➔ **Restricted (RS1317)**

Initiation

Any of the following:

- 1 Patient is fluid volume or rate restricted; or
- 2 Patient requires low electrolyte; or
- 3 Both:
 - 3.1 Any of the following:
 - 3.1.1 Cystic fibrosis; or
 - 3.1.2 Any condition causing malabsorption; or
 - 3.1.3 Faltering growth in an infant/child; or
 - 3.1.4 Increased nutritional requirements; and
 - 3.2 Patient has substantially increased metabolic requirements.

ENTERAL FEED 2 KCAL/ML – **Restricted** see terms [above](#)

⬆ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 500 ml bottle.....	6.82	1	Nutrison Concentrated
⬆ Liquid 8.4 g protein, 21.9 g carbohydrate, 9.1 g fat and 0.5 g fibre per 100 ml, 1,000 ml bottle	13.64	1	Ensure Two Cal HN RTH

ORAL FEED 2 KCAL/ML – **Restricted** see terms [above](#)

⬆ Liquid 8.4 g protein, 22.4 g carbohydrate, 8.9 g fat and 0.8 g fibre per 100 ml, 200 ml bottle	2.34	1	Two Cal HN
--	------	---	------------

High Protein Products

HIGH PROTEIN ENTERAL FEED 1.25 KCAL/ML – **Restricted** see terms [below](#)

⬆ Liquid 6.3 g protein, 14.2 g carbohydrate and 4.9 g fat per 100 ml, bottle	12.00	1,000 ml	Nutrison Protein Plus
--	-------	----------	-----------------------

➔ **Restricted (RS1327)**

Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.1 Patient has liver disease; or
- 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
- 2.3 Patient is fluid restricted; or
- 2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.26 KCAL/ML – **Restricted** see terms [below](#)

⚡ Liquid 10 g protein, 10.4 g carbohydrate and 4.9 g fat per 100 ml, bottle 8.67 500 ml Nutrison Protein Intense
 ➔ **Restricted (RS1327)**

Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
 - 2.1 Patient has liver disease; or
 - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
 - 2.3 Patient is fluid restricted; or
 - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.28 KCAL/ML – **Restricted** see terms [below](#)

⚡ Liquid 6.3 g protein, 14.1 g carbohydrate, 4.9 g fat and 1.5 g fibre per 100 ml, bottle 12.54 1,000 ml Nutrison Protein Plus Multi Fibre
 ➔ **Restricted (RS1327)**

Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
 - 2.1 Patient has liver disease; or
 - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
 - 2.3 Patient is fluid restricted; or
 - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

Infant Formulas

AMINO ACID FORMULA – **Restricted** see terms [below](#)

⚡ Powder 1.95 g protein, 8.1 g carbohydrate and 3.5 g fat per 100 ml, 400 g can		<i>e.g. Neocate</i>
⚡ Powder 13 g protein, 49 g carbohydrate and 23 g fat per 100 g, can 55.61	400 g	Neocate SYNEO
⚡ Powder 13.3 g protein, 56 g carbohydrate and 22 g fat per 100 g, can 55.61	400 g	Neocate Junior Unflavoured
⚡ Powder 13.3 g protein, 57 g carbohydrate and 24.6 g fat per 100 g, can 43.60	400 g	Alfamino
⚡ Powder 13.5 g protein, 52 g carbohydrate and 24.5 g fat per 100 g, can 55.61	400 g	Neocate Gold (Unflavoured)
⚡ Powder 14.8 g protein, 51.4 g carbohydrate and 23 g fat per 100 g, can 55.61	400 g	Neocate Junior Vanilla
⚡ Powder 15 g protein, 56 g carbohydrate and 20 g fat per 100 g, can 43.60	400 g	Alfamino Junior
⚡ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can 65.72	400 g	Elecare LCP (Unflavoured)
⚡ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can 65.72	400 g	Elecare (Unflavoured) Elecare (Vanilla)

➔ **Restricted (RS1867)**

Initiation

Any of the following:

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

- 1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows' milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis; or
- 4 Ultra-short gut; or
- 5 Severe Immune deficiency.

Continuation

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows' milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
- 3 Amino acid formula is required for a nutritional deficit.

Initiation – patients who are currently funded under RS1502 or SA1557

Limited to 3 months treatment

All of the following:

- 1 Patient has a valid initiation or renewal approval for extensively hydrolysed formula (RS1502); and
- 2 Patient is unable to source funded Aptamil powder at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Hospital Restriction RS1502. There is no continuation criteria under this criterion.

ENTERAL LIQUID PEPTIDE FORMULA – **Restricted** see terms [below](#)

↓ Liquid 4.2 g protein, 18.6 g carbohydrate and 6.58 g fat per 100 ml, 500 ml bottle	18.66	1	Nutrini Peptisorb Energy
---	-------	---	--------------------------

→ **Restricted (RS1775)**

Initiation

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure; or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Continuation

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and			
2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula.			
EXTENSIVELY HYDROLYSED FORMULA – Restricted see terms below			
↓ Powder 1.6 g protein, 7.5 g carbohydrate and 3.1 g fat per 100 ml, 900 g can.....	36.20	900 g	Allerpro Syneo 1
↓ Powder 1.6 g protein, 7.8 g carbohydrate and 3.2 g fat per 100 ml, 900 g can.....	36.20	900 g	Allerpro Syneo 2
↓ Powder 14 g protein, 53.4 g carbohydrate and 27.3 g fat per 100 g, can	18.10	450 g	Pepti-Junior
➔ Restricted (RS1502)			
Initiation			
Any of the following:			
1 Both:			
1.1 Cows' milk formula is inappropriate due to severe intolerance or allergy to its protein content; and			
1.2 Either:			
1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or			
1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or			
2 Severe malabsorption; or			
3 Short bowel syndrome; or			
4 Intractable diarrhoea; or			
5 Biliary atresia; or			
6 Cholestatic liver diseases causing malsorption; or			
7 Cystic fibrosis; or			
8 Proven fat malabsorption; or			
9 Severe intestinal motility disorders causing significant malabsorption; or			
10 Intestinal failure; or			
11 For step down from Amino Acid Formula.			
Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.			
Continuation			
Both:			
1 An assessment as to whether the infant can be transitioned to a cows' milk protein or soy infant formula has been undertaken; and			
2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula.			
FRUCTOSE-BASED FORMULA			
Powder 14.6 g protein, 49.7 g carbohydrate and 30.8 g fat per 100 g, 400 g can			e.g. <i>Galactomin 19</i>
LACTOSE-FREE FORMULA			
Powder 1.3 g protein, 7.3 g carbohydrate and 3.5 g fat per 100 ml, 900 g can			e.g. <i>Karicare Aptamil Gold De-Lact</i>
Powder 1.5 g protein, 7.2 g carbohydrate and 3.6 g fat per 100 ml, 900 g can			e.g. <i>S26 Lactose Free</i>
LOW-CALCIUM FORMULA			
Powder 14.8 g protein, 53.7 g carbohydrate and 26.7 g fat per 100 g and tuna fish oil (DHA), can.....	46.18	400 g	Locasol
PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML – Restricted see terms on the next page			
↓ Liquid 2.6 g protein, 10.3 g carbohydrate, 5.4 g fat and 0.6 g fibre per 100 ml, 125 ml bottle	2.80	1	Infatrini

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS1614)

Initiation – Fluid restricted or volume intolerance with faltering growth

Both:

- 1 Either:
 - 1.1 The patient is fluid restricted or volume intolerant; or
 - 1.2 The patient has increased nutritional requirements due to faltering growth; and
- 2 Patient is under 18 months old and weighs less than 8kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialed appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

PRETERM FORMULA – **Restricted** see terms [below](#)

↓ Liquid 2.2 g protein, 8.4 g carbohydrate and 4.4 g fat per 100 ml, bottle	0.75	100 ml	S26 LBW Gold RTF
↓ Liquid 2.3 g protein, 8.6 g carbohydrate and 4.2 g fat per 100 ml, 90 ml bottle			<i>e.g. Pre Nan Gold RTF</i>
↓ Liquid 2.6 g protein, 8.4 g carbohydrate and 3.9 g fat per 100 ml, 70 ml bottle			<i>e.g. Karicare Aptamil Gold+Preterm</i>

➔ Restricted (RS1224)

Initiation

For infants born before 33 weeks' gestation or weighing less than 1.5 kg at birth.

THICKENED FORMULA

Powder 1.8 g protein, 8.1 g carbohydrate and 3.3 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Thickened AR</i>
--	--	--	---

Ketogenic Diet Products

HIGH FAT FORMULA – **Restricted** see terms [below](#)

↓ Powder 14.3 g protein, 2.8 g carbohydrate and 69.2 g fat per 100 g, can	36.92	300 g	Ketocal 4:1 (Unflavoured)
↓ Powder 15.4 g protein, 7.2 g carbohydrate and 68.6 g fat per 100 g, can	36.92	300 g	Ketocal 4:1 (Vanilla)
			Ketocal 3:1 (Unflavoured)

➔ Restricted (RS1225)

Initiation

For patients with intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Paediatric Products

➔ Restricted (RS1473)

Initiation

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 The child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 Any condition causing malabsorption; or
 - 2.3 Faltering growth in an infant/child; or
 - 2.4 Increased nutritional requirements; or
 - 2.5 The child is being transitioned from TPN or tube feeding to oral feeding; or
 - 2.6 The child has eaten, or is expected to eat, little or nothing for 3 days.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PAEDIATRIC ENTERAL FEED 0.76 KCAL/ML – Restricted see terms on the previous page			
† Liquid 2.5 g protein, 12.5 g carbohydrate, 3.3 g fat and 0.7 g fibre per 100 ml, 500 ml bottle	6.27	1	Nutrini Low Energy Multi Fibre RTH
PAEDIATRIC ENTERAL FEED 1 KCAL/ML – Restricted see terms on the previous page			
† Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, 500 ml bottle.....	3.32	1	Pediasure RTH
† Liquid 2.7 g protein, 12.3 g carbohydrate and 4.4 g fat per 100 ml, 500 ml bottle	4.69	1	Nutrini RTH
PAEDIATRIC ENTERAL FEED 1.5 KCAL/ML – Restricted see terms on the previous page			
† Liquid 4.1 g protein, 18.5 g carbohydrate and 6.7 g fat per 100 ml, 500 ml bottle	7.46	1	Nutrini Energy RTH
† Liquid 4.1 g protein, 18.5 g carbohydrate, 6.7 g fat and 0.8 g fibre per 100 ml, 500 ml bottle	7.14	1	Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1 KCAL/ML – Restricted see terms on the previous page			
† Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, 200 ml bottle.....	1.33	1	Pediasure (chocolate) Pediasure (strawberry) Pediasure (vanilla)
PAEDIATRIC ORAL FEED 1.5 KCAL/ML – Restricted see terms on the previous page			
† Liquid 3.4 g protein, 18.8 g carbohydrate and 6.8 g fat per 100 ml, 200 ml bottle	1.90	1	Fortini (Strawberry) Fortini (Vanilla)
† Liquid 4.0 g protein, 18.8 g carbohydrate, 6.8 g fat and 1.5 g fibre per 100 ml, 200 ml bottle	1.90	1	Fortini Multi Fibre (chocolate) Fortini Multi Fibre (strawberry) Fortini Multi Fibre (unflavoured) Fortini Multi Fibre (vanilla)
† Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, 500 ml bottle	8.67	1	Pediasure Plus

Renal Products

LOW ELECTROLYTE ORAL FEED – Restricted see terms below			
‡ Powder 7.5 g protein, 57.6 g carbohydrate and 25.9 g fat per 100 g, can	64.26	400 g	Kindergen
➔ Restricted (RS1227)			
Initiation			
For children (up to 18 years) with acute or chronic kidney disease.			
LOW ELECTROLYTE ORAL FEED 1.8 KCAL/ML			
Liquid 8 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, 220 ml bottle	3.31	1	Nepro HP (strawberry) Nepro HP (vanilla)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LOW ELECTROLYTE ORAL FEED 2 KCAL/ML – Restricted see terms below			
↓ Liquid 3 g protein, 25.5 g carbohydrate and 9.6 g fat per 100 ml, 237 ml bottle			
↓ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 125 ml carton.....	13.72	4	Renilon 7.5 (apricot) Renilon 7.5 (caramel)
↓ Liquid 9.1 g protein, 19 g carbohydrate and 10 g fat per 100 ml, 200 ml bottle.....	13.24	4	Novasource Renal (Vanilla)

→ **Restricted (RS1228)**

Initiation

For patients with acute or chronic kidney disease.

Surgical Products

HIGH ARGININE ORAL FEED 1.4 KCAL/ML – Restricted see terms [below](#)

↓ Liquid 10.4 g protein, 8 g carbohydrate, 4.4 g fat and 0 g fibre per 100 ml, 250 ml carton.....	56.00	10	Impact Advanced Recovery
---	-------	----	-----------------------------

→ **Restricted (RS1231)**

Initiation

Three packs per day for 5 to 7 days prior to major gastrointestinal, head or neck surgery.

PREOPERATIVE CARBOHYDRATE FEED 0.5 KCAL/ML – Restricted see terms [below](#)

↓ Oral liq 0 g protein, 12.6 g carbohydrate and 0 g fat per 100 ml, 200 ml bottle.....	8.64	4	preOp
--	------	---	-------

→ **Restricted (RS1415)**

Initiation

Maximum of 400 ml as part of an Enhanced Recovery After Surgery (ERAS) protocol 2 to 3 hours before major abdominal surgery.

Standard Feeds

→ **Restricted (RS1214)**

Initiation

Any of the following:

For patients with malnutrition, defined as any of the following:

- Any of the following:
 - BMI < 18.5; or
 - Greater than 10% weight loss in the last 3-6 months; or
 - BMI < 20 with greater than 5% weight loss in the last 3-6 months; or
- For patients who have, or are expected to, eat little or nothing for 5 days; or
- For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or
- For use pre- and post-surgery; or
- For patients being tube-fed; or
- For tube-feeding as a transition from intravenous nutrition; or
- For any other condition that meets the community Special Authority criteria.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ENTERAL FEED 1.5 KCAL/ML – Restricted see terms on the previous page			
† Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 1.5 g fibre per 100 ml, 1,000 ml bottle	8.68	1	Nutrison Energy Multi Fibre
† Liquid 6.27 g protein, 20.4 g carbohydrate and 4.9 g fat per 100 ml, 1,000 ml bag	8.68	1	Ensure Plus HN RTH
† Liquid 6.38 g protein, 21.1 g carbohydrate, 4.9 g fat and 1.2 g fibre per 100 ml, 1,000 ml bottle	8.68	1	Jevity HiCal RTH
† Liquid 6 g protein, 18.5 g carbohydrate and 5.8 g fat per 100 ml, 1,000 ml bottle	9.00	1	Nutrison Energy
ENTERAL FEED 1 KCAL/ML – Restricted see terms on the previous page			
† Liquid 4 g protein, 12.3 g carbohydrate, 3.9 g fat and 1.5 g fibre per 100 ml, 1,000 ml bottle	7.21	1	Nutrison Multi Fibre
† Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, 1,000 ml bottle	6.56	1	Osmolite RTH
† Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, 1,000 ml bottle	6.56	1	Jevity RTH
† Liquid 4 g protein, 12.4 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bottle	6.90	1	Nutrison RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Restricted see terms on the previous page			
† Liquid 5.5 g protein, 8.8 g carbohydrate, 2.5 g fat and 1.5 g fibre per 100 ml, 1,000 ml bottle	9.05	1	Nutrison 800 Complete Multi Fibre
HIGH PROTEIN ORAL FEED 2.4 KCAL/ML – Restricted see terms on the previous page			
† Liquid 14.6 g protein, 25.3 g carbohydrate and 9.6 g fat per 100 ml, 125 ml bottle			<i>e.g. Fortisip Compact Protein</i>
ORAL FEED – Restricted see terms on the previous page			
† Powder 15.9 g protein, 57.4 g carbohydrate and 14 g fat per 100 g, can	40.00	800 g 850 g 800 g 850 g	Ensure (Chocolate) Ensure (Chocolate) Ensure (Vanilla) Ensure (Vanilla)
† Powder 23 g protein, 65 g carbohydrate and 2.5 g fat per 100 g, can	15.90	840 g	Sustagen Hospital Formula (Chocolate) Sustagen Hospital Formula (Vanilla)
ORAL FEED 1 KCAL/ML – Restricted see terms on the previous page			
† Liquid 3.8 g protein, 23 g carbohydrate and 12.7 g fibre per 100 ml, 237 ml carton			<i>e.g. Resource Fruit Beverage</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ORAL FEED 1.5 KCAL/ML – Restricted see terms on page 307			
† Liquid 4 g protein and 33.5 g carbohydrate per 100 ml, 200 ml bottle	3.30	200 ml	Fortijuice (Apple) Fortijuice (Orange) Fortijuice (Strawberry)
† Liquid 6 g protein, 18.4 g carbohydrate and 5.8 g fat per 100 ml, 200 ml bottle.....	1.76	1	Fortisip (Banana) Fortisip (Chocolate) Fortisip (Strawberry) Fortisip (Vanilla)
† Liquid 6.25 g protein, 20.2 g carbohydrate and 4.92 g fat per 100 ml, 200 ml bottle	1.56	1	Ensure Plus (Banana) Ensure Plus (Chocolate) Ensure Plus (Fruit of the forest) Ensure Plus (Vanilla)
ORAL FEED WITH FIBRE 1.5 KCAL/ML – Restricted see terms on page 307			
† Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 2.3 g fibre per 100 ml, 200 ml bottle	1.76	1	Fortisip Multi Fibre (chocolate) Fortisip Multi Fibre (strawberry) Fortisip Multi Fibre (vanilla)

Bacterial and Viral Vaccines

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – **Restricted** see terms [below](#)

† Inj 30 IU diphtheria toxoid with 30IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5 ml syringe
– **5% DV Dec-24 to 2027** 0.00 10 **Infanrix IPV**

→ **Restricted** ([RS1387](#))

Initiation

Any of the following:

- 1 A single dose for children up to the age of 7 who have completed primary immunisation; or
- 2 A course of up to four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4 Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE –

Restricted see terms [below](#)

† Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen 10mcg H. influenzae type b with tetanus toxoid 20-40mcg in 0.5ml syringe –
5% DV Dec-24 to 2027 0.00 10 **Infanrix-hexa**

→ **Restricted** ([RS2051](#))

Initiation

Any of the following:

- 1 Up to four doses for children under the age of 10 years for primary immunisation; or
- 2 An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
- 3 An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens; or
- 4 Up to five doses for children under the age of 10 years receiving solid organ transplantation.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Bacterial Vaccines

BACILLUS CALMETTE-GUERIN VACCINE – **Restricted** see terms [below](#)

† Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial with diluent – **5% DV Dec-24 to 2027** 0.00 10 **BCG Vaccine AJV**

→ **Restricted** ([RS1233](#))

Initiation

All of the following:

For infants at increased risk of tuberculosis defined as:

- 1 Living in a house or family with a person with current or past history of TB; and
- 2 Having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

equal to 40 per 100,000 for 6 months or longer; and

3 During their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000.

Note: A list of countries with high rates of TB are available at <http://www.health.govt.nz/tuberculosis> (Search for Downloads) or www.bcgatlas.org/index.php

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – **Restricted** see terms [below](#)

↓ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis
toxoid, 8 mcg pertussis filamentous haemagglutinin and 2.5 mcg
pertactin in 0.5 ml prefilled syringe – **5% DV Dec-24 to 2027** 0.00 10 **Boostrix**

→ **Restricted (RS1790)**

Initiation

Any of the following:

- 1 A single dose for pregnant women in the second or third trimester of each pregnancy; or; or
- 2 A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or; or
- 3 A course of up to four doses is funded for children from age 7 up the age of 18 years inclusive to complete full primary immunisation; or
- 4 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5 A single dose for vaccination of patients aged from 65 years old; or
- 6 A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7 For vaccination of previously unimmunised or partially immunised patients; or
- 8 For revaccination following immunosuppression; or
- 9 For boosting of patients with tetanus-prone wounds.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

HAEMOPHILUS INFLUENZAE TYPE B VACCINE – **Restricted** see terms [below](#)

↓ Inj 10 mcg vial with diluent syringe – **5% DV Dec-24 to 2027** 0.00 1 **Act-HIB**

→ **Restricted (RS1520)**

Initiation

Therapy limited to 1 dose

Any of the following:

- 1 For primary vaccination in children; or
- 2 An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre- or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or
- 3 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

MENINGOCOCCAL (A, C, Y AND W-135) CONJUGATE VACCINE

↓ Inj 10 mcg of each meningococcal polysaccharide conjugated to a total
of approximately 55 mcg of tetanus toxoid carrier per 0.5 ml vial –
5% DV Dec-24 to 2027 0.00 1 **MenQuadfi**

→ **Restricted (RS2019)**

Initiation

Either:

- 1 Any of the following:
 - 1.1 Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with HIV,

continued...

continued...

complement deficiency (acquired or inherited), functional or anatomic asplenia or pre or post solid organ transplant; or

- 1.2 One dose for close contacts of meningococcal cases of any group; or
- 1.3 One dose for person who has previously had meningococcal disease of any group; or
- 1.4 A maximum of two doses for bone marrow transplant patients; or
- 1.5 A maximum of two doses for person pre and post-immunosuppression*; or

2 Both:

2.1 Person is aged between 13 and 25 years, inclusive; and

2.2 Either:

2.2.1 One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or

2.2.2 One dose for individuals who turn 13 years of age while living in boarding school hostels.

Notes: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

‡ Inj 5 mcg of each meningococcal polysaccharide conjugated to a total of approximately 44 mcg of tetanus toxoid carrier in 0.5 ml vial..... 0.00 1 Nimenrix

➔ **Restricted (RS2037)**

Initiation – Children under 12 months of age

Any of the following:

- 1 A maximum of three doses (dependant on age at first dose) for patients pre- and post- splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post- solid organ transplant; or
- 2 A maximum of three doses (dependant on age at first dose) for close contacts of meningococcal cases of any group; or
- 3 A maximum of three doses (dependant on age at first dose) for child who has previously had meningococcal disease of any group; or
- 4 A maximum of three doses (dependant on age at first dose) for bone marrow transplant patients; or
- 5 A maximum of three doses (dependant on age at first dose) for child pre- and post-immunosuppression*.

Notes: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

MENINGOCOCCAL B MULTICOMPONENT VACCINE – Restricted see terms [below](#)

‡ Inj 175 mcg per 0.5 ml prefilled syringe..... 0.00 1 Bexsero
10 Bexsero

➔ **Restricted (RS2141)**

Initiation – Primary immunisation for children up to 59 months of age inclusive

Therapy limited to 3 doses

A primary course of up to three doses (dependent on age at first dose) for previously unvaccinated children up to the age of 59 months inclusive.

Initiation – High-risk individuals 5 years of age or over

Both:

- 1 Person is aged at least 5 years; and
- 2 Any of the following:
 - 2.1 Up to two doses and a booster every five years for patients pre- and post-splenectomy; or
 - 2.2 Up to two doses and a booster every five years for patients with functional or anatomic asplenia, HIV, complement

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- deficiency (acquired or inherited); or
- 2.3 Up to two doses and a booster every five years pre- or post-solid organ transplant; or
- 2.4 Up to two doses for close contacts of meningococcal cases of any group; or
- 2.5 Up to two doses for person who has previously had meningococcal disease of any group; or
- 2.6 Up to two doses for bone marrow transplant patients; or
- 2.7 Up to two doses for person pre- and post-immunosuppression* .

Initiation – Person is aged between 13 and 25 years (inclusive)

Therapy limited to 2 doses

Both:

- 1 Person is aged between 13 and 25 years (inclusive); and
- 2 Either:
 - 2.1 Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or
 - 2.2 Two doses for individuals who turn 13 years of age while living in boarding school hostels.

Note: *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE – Restricted see terms [below](#)

‡ Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5 ml syringe – **5% DV**

Dec-24 to 2027	0.00	1	Prevenar 13
		10	Prevenar 13

→ **Restricted (RS1936)**

Initiation – Primary course for previously unvaccinated children aged under 5 years

Therapy limited to 3 doses

A primary course of three doses for previously unvaccinated children up to the age of 59 months inclusive.

Initiation – High risk individuals who have received PCV10

Therapy limited to 2 doses

Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10.

Initiation – High risk children aged under 5 years

Therapy limited to 4 doses

Both:

- 1 Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high-risk children aged under 5 years; and
- 2 Any of the following:
 - 2.1 on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - 2.2 primary immune deficiencies; or
 - 2.3 HIV infection; or
 - 2.4 renal failure, or nephrotic syndrome; or
 - 2.5 are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - 2.6 cochlear implants or intracranial shunts; or
 - 2.7 cerebrospinal fluid leaks; or
 - 2.8 receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - 2.9 chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or

continued...

continued...

- 2.10 pre term infants, born before 28 weeks gestation; or
- 2.11 cardiac disease, with cyanosis or failure; or
- 2.12 diabetes; or
- 2.13 Down syndrome; or
- 2.14 who are pre-or post-splenectomy, or with functional asplenia.

Initiation – High risk individuals 5 years and over

Therapy limited to 4 doses

Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency.

Initiation – Testing for primary immunodeficiency diseases

For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – **Restricted** see terms [below](#)

‡ Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype) – **5% DV Dec-24 to 2027** 0.00 1 **Pneumovax 23**

➔ **Restricted (RS1587)**

Initiation – High risk patients

Therapy limited to 3 doses

For patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy; or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency.

Initiation – High risk children

Therapy limited to 2 doses

Both:

- 1 Patient is a child under 18 years for (re-)immunisation; and
- 2 Any of the following:
 - 2.1 On immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - 2.2 With primary immune deficiencies; or
 - 2.3 With HIV infection; or
 - 2.4 With renal failure, or nephrotic syndrome; or
 - 2.5 Who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - 2.6 With cochlear implants or intracranial shunts; or
 - 2.7 With cerebrospinal fluid leaks; or
 - 2.8 Receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - 2.9 With chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - 2.10 Pre term infants, born before 28 weeks gestation; or
 - 2.11 With cardiac disease, with cyanosis or failure; or
 - 2.12 With diabetes; or
 - 2.13 With Down syndrome; or
 - 2.14 Who are pre-or post-splenectomy, or with functional asplenia.

Initiation – Testing for primary immunodeficiency diseases

For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

SALMONELLA TYPHI VACCINE – **Restricted** see terms [below](#)

↓ Inj 25 mcg in 0.5 ml syringe

→ **Restricted** ([RS1243](#))

Initiation

For use during typhoid fever outbreaks.

Viral Vaccines

COVID-19 VACCINE

↓ Inj 3 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3 ml, 0.48 ml multi-dose vial; infant vaccine, yellow cap – **5% DV Feb-26**

to 30 Sep 2027 0.00 10 **Comirnaty (LP.8.1)**

→ **Restricted** ([RS2171](#))

Initiation – initial dose

Up to three doses for previously unvaccinated children aged 6 months – 4 years at high risk of severe illness or highly immunocompromised.

Continuation – additional dose

Either:

- 1 One additional dose with the most current variant-matched vaccine every 6 months for highly immunocompromised children aged 6 months to 4 years; or
- 2 One additional dose with the most current variant-matched vaccine every 12 months for children aged 6 months to 4 years old at high risk of severe illness.

↓ Inj 10 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3 ml, 0.48 ml single-dose vial; paediatric vaccine, light blue cap – **5% DV Feb-26 to 30 Sep 2027** 0.00 10 **Comirnaty (LP.8.1)**

→ **Restricted** ([RS2172](#))

Initiation – initial dose

Either:

- 1 One dose for previously unvaccinated children aged 5-11 years old; or
- 2 Up to three doses for immunocompromised children aged 5-11 years old.

Continuation – additional dose

Either:

- 1 One additional dose with the most current variant-matched vaccine every 6 months for highly immunocompromised children aged 5 to 11 years old; or
- 2 One additional dose with the most current variant-matched vaccine up to every 12 months for children aged 5 to 11 years old at high-risk of severe illness.

↓ Inj 30 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3 ml, pre-filled syringe; adult dose – **5% DV Feb-26 to 30 Sep 2027** 0.00 10 **Comirnaty (LP.8.1)**

→ **Restricted** ([RS2173](#))

Initiation – initial dose

Any of the following:

- 1 One dose for previously unvaccinated people aged 12-15 years and over 30 years old; or
- 2 Two doses for previously unvaccinated people aged 16-29 years old; or
- 3 Up to three doses for previously unvaccinated immunocompromised people from 12 years old; or
- 4 Up to four doses for people at risk of severe illness aged from 12-29 years.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – additional dose

Both:

- 1 One additional dose with the most current variant-matched vaccine every 6 months, additional dose to be given at least 6 months after last dose; and
- 2 Any of the following:
 - 2.1 Previously vaccinated people aged 30 years and over; or
 - 2.2 Previously vaccinated immunocompromised people from 12 years; or
 - 2.3 Previously vaccinated people at high-risk of severe illness from 12 years.

HEPATITIS A VACCINE – **Restricted** see terms [below](#)

⚡ Inj 720 ELISA units in 0.5 ml syringe – 5% DV Dec-24 to 2027	0.00	1	Havrix Junior
⚡ Inj 1440 ELISA units in 1 ml syringe – 5% DV Dec-24 to 2027	0.00	1	Havrix 1440
➡ Restricted (RS1638)			

Initiation

Any of the following:

- 1 Two vaccinations for use in transplant patients; or
- 2 Two vaccinations for use in children with chronic liver disease; or
- 3 One dose of vaccine for close contacts of known hepatitis A cases.

HEPATITIS B RECOMBINANT VACCINE

⚡ Inj 10 mcg per 0.5 ml prefilled syringe – 5% DV Dec-24 to 2027	0.00	1	Engerix-B
➡ Restricted (RS2049)			

Initiation

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4 For HIV positive patients; or
- 5 For hepatitis C positive patients; or
- 6 For patients following non-consensual sexual intercourse; or
- 7 For patients prior to planned immunosuppression for greater than 28 days; or
- 8 For patients following immunosuppression; or
- 9 For solid organ transplant patients; or
- 10 For post-haematopoietic stem cell transplant (HSCT) patients; or
- 11 Following needle stick injury.

⚡ Inj 20 mcg per 1 ml prefilled syringe – 5% DV Dec-24 to 2027	0.00	1	Engerix-B
➡ Restricted (RS2174)			

Initiation

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4 For HIV positive patients; or
- 5 For hepatitis C positive patients; or
- 6 For patients following non-consensual sexual intercourse; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 7 For patients prior to planned immunosuppression for greater than 28 days; or
- 8 For patients following immunosuppression; or
- 9 For solid organ transplant patients; or
- 10 For post-haematopoietic stem cell transplant (HSCT) patients; or
- 11 Following needle stick injury; or
- 12 For chronic kidney disease (CKD) stage 4 or 5 patients; or
- 13 For liver or kidney transplant patients.

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] – **Restricted** see terms [below](#)

↓ Inj 270 mcg in 0.5 ml syringe – **5% DV Dec-24 to 2027** 0.00 10 **Gardasil 9**

→ **Restricted (RS2038)**

Initiation – Children aged 14 years and under

Therapy limited to 2 doses

Children aged 14 years and under.

Initiation – other conditions

Either:

- 1 Up to 3 doses for people aged 15 to 26 years inclusive; or
- 2 Both:
 - 2.1 People aged 9 to 26 years inclusive; and
 - 2.2 Any of the following:
 - 2.2.1 Up to 3 doses for confirmed HIV infection; or
 - 2.2.2 Up to 3 doses people with a transplant (including stem cell); or
 - 2.2.3 Up to 4 doses for Post chemotherapy.

Initiation – Recurrent Respiratory Papillomatosis

All of the following:

- 1 Either:
 - 1.1 Maximum of two doses for children aged 14 years and under; or
 - 1.2 Maximum of three doses for people aged 15 years and over; and
- 2 The person has recurrent respiratory papillomatosis; and
- 3 The person has not previously had an HPV vaccine.

INFLUENZA VACCINE

↓ Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)..... 120.00 10 **Influvac Tetra**
(2026 formulation)

→ **Restricted (RS2013)**

Initiation – People over 65

The patient is 65 years of age or over.

Initiation – cardiovascular disease

Any of the following:

- 1 Ischaemic heart disease; or
- 2 Congestive heart failure; or
- 3 Rheumatic heart disease; or
- 4 Congenital heart disease; or
- 5 Cerebro-vascular disease.

Note: hypertension and/or dyslipidaemia without evidence of end-organ disease is excluded from funding.

Initiation – chronic respiratory disease

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Asthma, if on a regular preventative therapy; or
- 2 Other chronic respiratory disease with impaired lung function.

Note: asthma not requiring regular preventative therapy is excluded from funding.

Initiation – Other conditions

Either:

- 1 Any of the following:
 - 1.1 Diabetes; or
 - 1.2 chronic renal disease; or
 - 1.3 Any cancer, excluding basal and squamous skin cancers if not invasive; or
 - 1.4 Autoimmune disease; or
 - 1.5 Immune suppression or immune deficiency; or
 - 1.6 HIV; or
 - 1.7 Transplant recipient; or
 - 1.8 Neuromuscular and CNS diseases/ disorders; or
 - 1.9 Haemoglobinopathies; or
 - 1.10 Is a child on long term aspirin; or
 - 1.11 Has a cochlear implant; or
 - 1.12 Errors of metabolism at risk of major metabolic decompensation; or
 - 1.13 Pre and post splenectomy; or
 - 1.14 Down syndrome; or
 - 1.15 Is pregnant; or
 - 1.16 Is a child 4 years of age or under (inclusive) who has been hospitalised for respiratory illness or has a history of significant respiratory illness; or
- 2 Patients in a long-stay inpatient mental health care unit or who are compulsorily detained long-term in a forensic unit within a Public Hospital.

Initiation – Serious mental health conditions or addiction

Any of the following:

- 1 schizophrenia; or
- 2 major depressive disorder; or
- 3 bipolar disorder; or
- 4 schizoaffective disorder; or
- 5 person is currently accessing secondary or tertiary mental health and addiction services.

MEASLES, MUMPS AND RUBELLA VACCINE – **Restricted** see terms [below](#)

⚠ Injection, measles virus 1,000 CCID50, mumps virus 5,012 CCID50, Rubella virus 1,000 CCID50; prefilled syringe/ampoule of diluent 0.5 ml – 5% DV Dec-24 to 2027	0.00	10	Priorix
➡ Restricted (RS1487)			

Initiation – first dose prior to 12 months

Therapy limited to 3 doses

Any of the following:

- 1 For primary vaccination in children; or
- 2 For revaccination following immunosuppression; or
- 3 For any individual susceptible to measles, mumps or rubella.

Initiation – first dose after 12 months

Therapy limited to 2 doses

Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 For primary vaccination in children; or
- 2 For revaccination following immunosuppression; or
- 3 For any individual susceptible to measles, mumps or rubella.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

POLIOMYELITIS VACCINE – **Restricted** see terms [below](#)

↓ Inj 80 D-antigen units in 0.5 ml syringe – **5% DV Dec-24 to 2027** 0.00 1 **IPOL**
 ➔ **Restricted (RS1398)**

Initiation

Therapy limited to 3 doses

Either:

- 1 For partially vaccinated or previously unvaccinated individuals; or
- 2 For revaccination following immunosuppression.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

RABIES VACCINE

Inj 2.5 IU vial with diluent

ROTAVIRUS ORAL VACCINE – **Restricted** see terms [below](#)

↓ Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose,
 prefilled oral applicator – **5% DV Dec-24 to 2027** 0.00 10 **Rotarix**
 ↓ Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose,
 squeezable tube (PVC free) 0.00 10 Rotarix

➔ **Restricted (RS1590)**

Initiation

Therapy limited to 2 doses

Both:

- 1 First dose to be administered in infants aged under 14 weeks of age; and
- 2 No vaccination being administered to children aged 24 weeks or over.

VARICELLA VACCINE [CHICKENPOX VACCINE]

↓ Inj 2000 PFU prefilled syringe plus vial – **5% DV Dec-24 to 2027** 0.00 10 **Varilrix**

➔ **Restricted (RS1591)**

Initiation – primary vaccinations

Therapy limited to 1 dose

Either:

- 1 Any infant born on or after 1 April 2016; or
- 2 For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox).

Initiation – other conditions

Therapy limited to 2 doses

Any of the following:

- 1 Any of the following:
 for non-immune patients:
 1.1 With chronic liver disease who may in future be candidates for transplantation; or
 1.2 With deteriorating renal function before transplantation; or
 1.3 Prior to solid organ transplant; or
 1.4 Prior to any elective immunosuppression*; or
 1.5 For post exposure prophylaxis who are immune competent inpatients; or
- 2 For patients at least 2 years after bone marrow transplantation, on advice of their specialist; or

continued...

VACCINES

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

- 3 For patients at least 6 months after completion of chemotherapy, on advice of their specialist; or
- 4 For HIV positive patients non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
- 5 For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
- 6 For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
- 7 For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

Note: * immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] – **Restricted** see terms [below](#)

⚡ Inj 50 mcg per 0.5 ml vial plus vial.....	0.00	1	Shingrix
		10	Shingrix

➡ **Restricted** ([RS2039](#))

Initiation – people aged 18 years and over (Shingrix)

Therapy limited to 2 doses

Any of the following:

- 1 Pre- and post-haematopoietic stem cell transplant or cellular therapy; or
- 2 Pre- or post-solid organ transplant; or
- 3 Haematological malignancies; or
- 4 People living with poorly controlled HIV infection; or
- 5 Planned or receiving disease modifying anti-rheumatic drugs (DMARDs – targeted synthetic, biologic, or conventional synthetic) for polymyalgia rheumatica, systemic lupus erythematosus or rheumatoid arthritis; or
- 6 End stage kidney disease (CKD 4 or 5);; or
- 7 Primary immunodeficiency.

Diagnostic Agents

TUBERCULIN PPD [MANTOUX] TEST

Inj 5 TU per 0.1 ml, 1 ml vial – 5% DV Dec-24 to 2027	0.00	1	Tubersol
--	------	---	-----------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Optional Pharmaceuticals			
NOTE:			
In addition to the products expressly listed here in Part III: Optional Pharmaceuticals, a range of hospital medical devices are listed in an addendum to Part III which is available at schedule.pharmac.govt.nz . The Optional Pharmaceuticals listed in the addendum are deemed to be listed in Part III, and the Rules of the Pharmaceutical Schedule applying to products listed in Part III apply to them.			
BETA-HCG LOW SENSITIVITY URINE TEST KIT			
Note: For use in abortion services only.			
Midstream.....	17.47	1 test	CheckTop
BLOOD GLUCOSE DIAGNOSTIC TEST METER			
1 meter with 50 lancets, a lancing device, and 10 diagnostic test strips	20.00	1	CareSens N Premier
	10.00		Caresens N
			Caresens N POP
BLOOD GLUCOSE DIAGNOSTIC TEST STRIP			
Blood glucose test strips.....	10.56	50 test	CareSens N
Test strips	10.56	50 test	CareSens PRO
BLOOD KETONE DIAGNOSTIC TEST STRIP			
Test strips	15.50	10 strip	KetoSens
DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER			
Meter with 50 lancets, a lancing device, and 10 blood glucose diagnostic test strips	20.00	1	CareSens Dual
MASK FOR SPACER DEVICE			
Small.....	2.70	1	e-chamber Mask
PEAK FLOW METER			
Low Range	9.54	1	Mini-Wright AFS Low Range
Normal Range	9.54	1	Mini-Wright Standard
PREGNANCY TEST - HCG URINE			
Cassette – 5% DV Mar-25 to 2027	16.00	40 test	David One Step Cassette Pregnancy Test
SODIUM NITROPRUSSIDE			
Test strip.....	22.00	50 strip	Ketostix
SPACER DEVICE			
220 ml (single patient)	3.65	1	e-chamber Turbo
510 ml (single patient)	5.95	1	e-chamber La Grande
800 ml.....	6.50	1	Volumatic

- Symbols -

Xaluprine	153
8-methoxypsoralen	71

- A -

A-Scabies	68
Abacavir sulphate	105
Abacavir sulphate with lamivudine	105
Abacavir/lamivudine Viatris	105
Abciximab	183
Abilify Maintena	137
Abiraterone acetate	174
Acarbose	9
Accarb	9
Acetazolamide	278
Acetic acid Extemporaneously Compounded Preparations	289
Genito-Urinary	74
Acetic acid with hydroxyquinoline, glycerol and ricinoleic acid	74
Acetic acid with propylene glycol	280
Acetylcholine chloride	278
Acetylcysteine	281
Aciclovir Infections	108
Sensory	274
Aciclovir-Baxter	108
Acid Citrate Dextrose A	36
Acidex	5
Acipimox	54
Acitretin	71
Act-HIB	311
Actemra	233
Actinomycin D	152
Adalimumab (Amgevita)	183
Adalimumab (Humira - alternative brand)	192
Adapalene	68
Adcetris	203
Adenosine	46
Adenosine Baxter	46
Adrenaline Cardiovascular	54
Respiratory	263
Adriamycin	152
Adsine	46
Advantan	70
Advate	35
Adynovate	35
Aerrane	122
Afinitor	256
Aflibercept	199

Agents Affecting the Renin-Angiotensin System	44
Agents for Parkinsonism and Related Disorders	121
Agents Used in the Treatment of Poisonings	281
Ajmaline	46
Albalon	276
Albendazole	102
Alchemy Caspofungin	100
Alchemy Oxaliplatin	161
Alchemy Oxybutynin	77
Aldara	72
Aldurazyme	19
Alecensa	161
Alectinib	161
Alendronate sodium	115
Alendronate sodium with colecalciferol	115
Alfacalcidol	28
Alfamino	302
Alfamino Junior	302
Alfentanil	126
Alglucosidase alfa	17
Allegron	129
Allerfix	276
Allerpro Syneo 1	304
Allerpro Syneo 2	304
Allersoothe	264
Allmercap	153
Allopurinol	117
Almarytm	47
Alpha tocopheryl	28
Alpha tocopheryl acetate	29
Alpha-Adrenoceptor Blockers	46
Alphamox 125	93
Alphamox 250	93
Alprolix	34
Alprostadil	55
Alprostadil hydrochloride	55
Alteplase	39
Alum	289
Aluminium chloride	32
Aluminium hydroxide	5
Aluminium hydroxide with magnesium hydroxide and simeticone	5
Alyftrek	272
Amantadine hydrochloride	121
Ambrisentan	56
Ambrisentan Viatris	56
Amdipharm	93
Amethocaine Nervous	125

Sensory	277
Amgevita	183
Amikacin	89
Amiloride hydrochloride	50
Amiloride hydrochloride with furosemide	50
Amiloride hydrochloride with hydrochlorothiazide	50
Aminolevulinic acid hydrochloride	177
Aminophylline	270
Amiodarone hydrochloride	46
Amisulpride	135
Amitriptyline	128
Amlodipine	48
Amorolfine	67
Amoxicillin	93
Amoxicillin with clavulanic acid	93
Amoxiclav Devatis Forte	93
Amoxycillin Sandoz	93
Amphotericin B Alimentary	26
Infections	98
Amphotericin Liposomal SUN	98
Amsacrine	154
Amyl nitrite	281
Anabolic Agents	78
Anaesthetics	122
Anagrelide hydrochloride	154
Analgesics	125
Anastrozole	175
Anatrole	175
Androgen Agonists and Antagonists	78
Anoro Ellipta	265
Antabuse	148
Antacids and Antiflatulents	5
Anti-Infective Agents	74
Anti-Infective Preparations Dermatological	67
Sensory	274
Anti-Inflammatory Preparations	275
Antiacne Preparations	68
Antiallergy Preparations	263
Antianaemics	30
Antiarrhythmics	46
Antibacterials	89
Anticholinergic Agents	265
Anticholinesterases	115
Antidepressants	128
Antidiarrhoeals and Intestinal Anti-Inflammatory Agents	5
Antiepilepsy Drugs	130
Antifibrinolytics, Haemostatics and	

Local Sclerosants	32	Arrow-Bendrofluazide	51	Azathioprine	256
Antifibrotics	267	Arrow-Brimonidine	279	Azilect	122
Antifungals	97	Arrow-Diazepam	139	Azithromycin	91
Antihypotensives	47	Arrow-Fluoxetine	130	Azopt	278
Antimigraine Preparations	133	Arrow-Norfloxacin	95	Aztreonam	96
Antimycobacterials	101	Arrow-Ornidazole	103	- B -	
Antinausea and Vertigo Agents	133	Arrow-Quinapril 10	44	Bacillus calmette-guerin (BCG)	256
Antiparasitics	102	Arrow-Quinapril 20	44	Bacillus calmette-guerin vaccine	310
Antipruritic Preparations	68	Arrow-Quinapril 5	44	Baclofen	118
Antipsychotic Agents	135	Arrow-Roxithromycin	93	Bacterial and Viral Vaccines	310
Antiretrovirals	104	Arrow-Timolol	278	Bacterial Vaccines	310
Antirheumatoid Agents	115	Arrow-Topiramate	132	Balanced Salt Solution	277
Antiseptics and Disinfectants	283	Arrow-Tramadol	128	Barium sulphate	284
Antispasmodics and Other Agents		Arsenic trioxide	154	Barrier Creams and Emollients	68
Altering Gut Motility	7	Artemether with lumefantrine	103	Basiliximab	200
Antithrombotics	36	Artesunate	103	BCG Vaccine AJV	310
Antithymocyte globulin		Articaine hydrochloride	123	Beclazone 100	269
(equine)	256	Articaine hydrochloride with adrenaline	123	Beclazone 250	269
Antithymocyte globulin (rabbit)	256	Asacol	6	Beclazone 50	269
Antulcerants	7	Ascorbic acid		Beclomethasone dipropionate	269
Antivirals	107	Alimentary	28	Bedaquiline	101
Anxiolytics	139	Extemporaneously Compounded Preparations	289	Bee venom	263
Anzatax	173	Aspen Adrenaline	54	Bendamustine hydrochloride	150
Apidra	10	Aspirin		Bendamustine Sandoz	150
Apidra Solostar	10	Blood	37	Bendrofluazide	51
APO Clomipramine	128	Nervous	125	Bendroflumethiazide [Bendrofluazide]	51
APO-Atomoxetine	145	Asthalin	268	Benralizumab	200
Apo-Candesartan	45	Atazanavir sulphate	106	Benzathine benzylpenicillin	93
APO-Candesartan HCTZ		Atazanavir Viatrix	106	Benzatropine mesylate	121
16/12.5	45	Atenolol	47	Benzbromaron AL 100	117
APO-Candesartan HCTZ		Atenolol Viatrix	47	Benzbromarone	117
32/12.5	45	Atenolol-AFT	47	Benzetacil	93
APO-Olanzapine	136	Atezolizumab	244	Benzocaine	123
APO-Ursodeoxycholic Acid	14	ATGAM	256	Benzocaine with tetracaine hydrochloride	123
Apomorphine hydrochloride	121	Ativan	140	Benzoin	289
Apraclonidine	279	Atomoxetine	145	Benzoyl peroxide	68
Aprepitant	133	Atorvastatin	52	Benzotrop	121
Apresoline	55	Atovaquone with proguanil hydrochloride	103	Benzydamine hydrochloride	25
Aprotinin	32	Atracurium besylate	118	Benzydamine hydrochloride with cetylpyridinium chloride	26
Aptamil Feed Thickener	294	Atropine sulphate		Benzylpenicillin sodium [Penicillin G]	93
Aqueous cream	69	Cardiovascular	46	Beractant	272
Arachis oil [Peanut oil]	289	Sensory	279	Besponsa	213
Aratac	46	Atropt	279	Beta Cream	70
Arava	115	Augmentin	93	Beta Ointment	70
Arginine		Aurorix	129	Beta Scalp	72
Alimentary	18	Avastin	202	Beta-Adrenoceptor Agonists	268
Various	285	Avelox	94	Beta-Adrenoceptor Blockers	47
Arginine2000	298	Avonex	141	Beta-hCG low sensitivity urine test kit	321
Argipressin [Vasopressin]	88	Axitinib	161	Betadine	283
Aripiprazole	135, 137	Azacididine	152	Betahistine dihydrochloride	134
Aripiprazole Sandoz	135	Azacididine Dr Reddy's	152		
Aristocort	70	Azactam	96		
Arrotex-Prazosin S29	46	Azamun	256		
Arrow - Clopid	37				
Arrow - Lattim	278				
Arrow-Amitriptyline	128				

7

324

Cefoxitin.....	90	Choriogonadotropin alfa.....	83	Clostridium botulinum type A toxin.....	118
Ceftaroline fosamil.....	91	Ciclopirox olamine.....	67	Clotrimazole	
Ceftazidime.....	90	Ciclosporin.....	177	Dermatological.....	67
Ceftazidime Kabi.....	90	Cidofovir.....	108	Genito-Urinary.....	74
Ceftazidime with avibactam.....	90	Cilicaine VK.....	94	Clove oil.....	289
Ceftriaxone.....	91	Cimetidine.....	8	Clozapine.....	135
Ceftriaxone-AFT.....	91	Cinacalcet.....	78	Clozaril.....	135
Cefuroxime.....	90	Cinacalcet Devatis.....	78	Clustran.....	133
Cefuroxime Devatis.....	90	Cinchocaine hydrochloride with hydrocortisone.....	7	Co-trimoxazole.....	97
Celapram.....	129	Cipla.....	265	Coal tar.....	289
Celebrex.....	119	Ciprofloxacin		Coal tar with salicylic acid and sulphur.....	71
Celecoxib.....	119	Infections.....	94	Cocaine hydrochloride.....	124
Celiprolol.....	48	Sensory.....	274	Cocaine hydrochloride with adrenaline.....	124
CellCept.....	257	Ciprofloxacin Kabi.....	94	Codeine phosphate	
Celostea.....	119	Ciprofloxacin Teva.....	274	Extemporaneously Compounded Preparations.....	289
Centrally-Acting Agents.....	50	Ciprofloxacin with hydrocortisone.....	274	Nervous.....	126
Cerazette.....	75	Ciproxin HC Otic.....	274	Coenzyme Q10.....	18
Cerobact.....	93	Cisplatin.....	161	Colchicine.....	117
Cerubidine.....	152	Cisplatin Accord.....	161	Colchicine Indoco.....	117
Cetirizine hydrochloride.....	264	Citalopram hydrobromide.....	129	Colecalciferol.....	28
Cetomacrogol.....	69	Citanest.....	125	Colestimethate.....	96
Cetomacrogol Cream AFT.....	69	Citrate sodium.....	36	Colestipol hydrochloride.....	53
Cetomacrogol with glycerol.....	69	Citric acid.....	289	Colestyramine.....	53
Cetomacrogol-AFT.....	69	Citric acid with magnesium carbonate hydrate and sodium picosulfate.....	15	Colestyramine - Mylan.....	53
Cetrimide.....	289	Citric acid with sodium bicarbonate.....	284	Colgout.....	117
Cetuximab.....	204	Citrulline 1000.....	298	Colifoam.....	6
Champix.....	149	Cladribine.....	152	Colistin sulphomethate [Colestimethate].....	96
Charcoal.....	282	Clarithromycin.....	92	Collodion flexible.....	289
CheckTop.....	321	Clexane.....	36	Colloidal bismuth subcitrate.....	9
Chemotherapeutic Agents.....	150	Clexane Forte.....	36	Colofac.....	7
Chickenpox vaccine.....	319	Clindamycin.....	96	Colomycin.....	96
Chlorafast.....	274	Clinicians.....	28	Colony-Stimulating Factors.....	39
Chloral hydrate.....	142	Clinicians Multivit & Mineral Boost.....	26	Coloxyl.....	15
Chlorambucil.....	151	Clinicians Renal Vit.....	27	Combigan.....	279
Chloramphenicol		Clobazam.....	130	Combiprasal.....	265
Infections.....	96	Clobetasol propionate.....	70, 72	Comirnaty (LP.8.1).....	315
Sensory.....	274	Clobetasone butyrate.....	70	Compound electrolytes.....	41, 43
Chlorhexidine.....	283	Clofazimine.....	101	Compound electrolytes with glucose [Dextrose].....	41, 43
Chlorhexidine gluconate		Clomazol		Compound hydroxybenzoate.....	289
Alimentary.....	26	Dermatological.....	67	Compound sodium lactate [Hartmann's solution].....	41
Extemporaneously Compounded Preparations.....	289	Genito-Urinary.....	74	Concerta.....	146
Genito-Urinary.....	74	Clomifene citrate.....	82	Condylina.....	72
Chlorhexidine with cetrimide.....	283, 286	Clomipramine hydrochloride.....	128	Contraceptives.....	74
Chlorhexidine with ethanol.....	283	Clonazepam.....	130, 139	Contrast Media.....	284
Chloroform.....	289	Clonidine.....	50	Copaxone.....	141
Chloroquine phosphate.....	103	Clonidine hydrochloride.....	50	Copper.....	23
Chlorothiazide.....	51	Clonidine Teva.....	50	Copper chloride.....	23
Chlorpheniramine maleate.....	264	Clopidogrel.....	37	Corticotrelin (ovine).....	82
Chlorpromazine hydrochloride.....	135	Clopine.....	135		
Chlortalidone [Chlorthalidone].....	51	Clopixol.....	137, 139		
Chlortalidone.....	51				
Choice 380 7med Nsha Silver/copper Short.....	75				
Cholestyramine.....	53				

Corticosteroids		Daunorubicin	152	Dexamethasone with	
Dermatological	70	David One Step Cassette Pregnancy		tobramycin	274
Hormone Preparations	79	Test	321	Dexamfetamine sulfate	145
Cosentyx	231	DBL Acetylcysteine	281	Dexmedetomidine	122
Cosmegen	152	DBL Adrenaline	54	Dexmedetomidine Viatrix	122
Coversyl	44	DBL Amikacin	89	Dexmethsone	80
COVID-19 vaccine	315	DBL Aminophylline	270	Dextrazoxane	173
Creon 10000	14	DBL Bleomycin Sulfate	152	Dextrose	
Creon 25000	14	DBL Bortezomib	154	Alimentary	9
Creon Micro	14	DBL Cefotaxime	90	Blood	41, 43
Crizotinib	162	DBL Dacarbazine	154	Extemporaneously Compounded	
Crotamiton	68	DBL Desferrioxamine Mesylate for Inj		Preparations	289
Crystadern	67	BP	282	Dextrose with sodium citrate and	
Cu 375 Standard	75	DBL Docetaxel	173	citric acid [Acid Citrate Dextrose	
Curam Duo 500/125	93	DBL Ergometrine	75	A]	36
Curosurf	272	DBL Gemcitabine	153	DHC Continus	126
Cvite	28	DBL Gentamicin	89	Diabetes	9
Cyclizine hydrochloride	134	DBL Leucovorin Calcium	173	Diacomit	132
Cyclizine lactate	134	DBL Methotrexate Onco-Vial	153	Diagnostic Agents	
Cyclogyl	279	DBL Naloxone Hydrochloride	281	Vaccines	320
Cyclonex	151	DBL Pethidine Hydrochloride	128	Various	285
Cyclopentolate hydrochloride	279	DBL Promethazine		Diagnostic and Surgical	
Cyclophosphamide	151	Hydrochloride	264	Preparations	276
Cycloserine	101	DBL Vincristine Sulfate	174	Diamide Relief	5
Cymevene	108	Decongestants	268	Diasip (strawberry)	299
Cyproheptadine hydrochloride	264	Decongestants and		Diasip (vanilla)	299
Cyproterone acetate	78	Antiallergics	276	Diatrizoate sodium	284
Cyproterone acetate with		Decozol	26	Diazepam	130, 139
ethinyloestradiol	74	Deferasirox	282	Diazoxide	
Cystadane	18	Deferiprone	282	Alimentary	9
Cysteamine hydrochloride	289	Defibrotide	36	Cardiovascular	55
Cytarabine	153	Definity	285	Dichlorobenzyl alcohol with	
Cytotec	7	Demeclocycline hydrochloride	95	amylmetacresol	26
- D -		Denosumab	116	Diclofenac Devatis	276
D-Penamamine	115	Deolate	101	Diclofenac Sandoz	119
Dabigatran	36	Deoxycoformycin	157	Diclofenac sodium	
Dabrafenib	162	Depo-Medrol	80	Musculoskeletal	119
Dacarbazine	154	Depo-Provera	75	Sensory	276
Dactinomycin [Actinomycin D]	152	Depo-Testosterone	78	Dicobalt edetate	283
Daivobet	71	Deprim	97	Diflucan	98
Daivonex	71	Dermol	70, 72	Diflucortolone valerate	70
Dalacin C	96	Desferrioxamine mesilate	282	Digestives Including Enzymes	14
Danaparoid	36	Desflurane	122	Digoxin	47
Dantrium	118	Desmopressin	88	Digoxin immune Fab	281
Dantrium IV	118	Desmopressin acetate	88	Dihydrocodeine tartrate	126
Dantrolene	118	Desmopressin-PH&T	88	Dihydroergotamine mesylate	133
Daonil	11	Desogestrel	75	Diltiazem CD Clinect	49
Dapa-Tabs	51	Dexamethasone		Diltiazem hydrochloride	49
Dapsone	101	Hormone Preparations	80	Dimercaprol	283
Daptomycin	96	Sensory	275	Dimercaptosuccinic acid	283
Daptomycin Dr Reddy's	96	Dexamethasone phosphate	80	Dimethicone	67-68
Daptomycin-AFT	96	Dexamethasone with framycetin and		Dimethyl fumarate	141
Darunavir	106	gramicidin	274	Dimethyl sulfoxide	287
Darunavir Viatrix	106	Dexamethasone with neomycin		Dinoprostone	75
Dasatinib	163	sulphate and polymyxin B		Dipentum	7
Dasatinib-Teva	163	sulphate	274	Diphemanil metilsulfate	72

Diphenoxylate hydrochloride with atropine sulphate.....	5	DP-Captopril.....	44	Emend Tri-Pack.....	133
Diphtheria antitoxin.....	282	Dr Reddy's Omeprazole.....	8	Emicizumab.....	33
Diphtheria, tetanus and pertussis vaccine.....	311	Drofate.....	48	EMLA.....	125
Diphtheria, tetanus, pertussis and polio vaccine.....	310	Droperidol.....	134	Empagliflozin.....	13
Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine.....	310	Droperidol Panpharma.....	134	Empagliflozin with metformin hydrochloride.....	13
Diprosone.....	70	Drugs Affecting Bone Metabolism.....	115	Emsogen.....	299
Dipyridamole.....	37	Dual blood glucose and blood ketone diagnostic test meter.....	321	Emtricitabine.....	105
Dipyridamole - Strides.....	37	Dulaglutide.....	11	Emtricitabine with tenofovir disoproxil.....	110
Disodium edetate.....	277	Dulcolax SP Drop.....	16	Emtriva.....	105
Disodium hydrogen phosphate with sodium dihydrogen phosphate.....	289	Duocal Super Soluble Powder.....	293	Emulsifying ointment.....	69
Disopyramide phosphate.....	47	Duolin.....	265	Enalapril maleate.....	44
Disulfiram.....	148	DuoResp Spiromax.....	270	Enbrel.....	178
Dithranol.....	289	Duovisc.....	277	Endocrine Therapy.....	174
Diuretics.....	50	Duride.....	54	Endoxan.....	151
Dobutamine.....	54	Durvalumab.....	245	Energivit.....	297
Dobutamine-hameln.....	54	Dynastat.....	120	Engerix-B.....	316
Docetaxel.....	173	Dysport.....	118	Enhertu.....	239
Docusate sodium		- E -		Enlifax XR.....	129
Alimentary.....	15	e-chamber La Grande.....	321	Enoxaparin sodium.....	36
Sensory.....	280	e-chamber Mask.....	321	Enstilar.....	71
Docusate sodium with sennosides.....	15	e-chamber Turbo.....	321	Ensure (Chocolate).....	308
Dolutegravir.....	107	E-Mycin.....	93	Ensure (Vanilla).....	308
Dolutegravir with lamivudine.....	107	E-Z-Gas II.....	284	Ensure Plus (Banana).....	309
Domperidone.....	134	Easiphen Liquid.....	297	Ensure Plus (Chocolate).....	309
Domperidone Viatris.....	134	Econazole nitrate.....	67	Ensure Plus (Fruit of the forest).....	309
Donepezil hydrochloride.....	147	Edrophonium chloride.....	115	Ensure Plus (Vanilla).....	309
Dopamine Basi.....	54	Efavirenz.....	104	Ensure Plus HN RTH.....	308
Dopamine hydrochloride.....	54	Efavirenz Milpharm.....	104	Ensure Two Cal HN RTH.....	301
Dornase alfa.....	270	Efavirenz with emtricitabine and tenofovir disoproxil.....	105	Entacapone.....	121
Dortimopt.....	278	Eformoterol fumarate.....	269	Entacapone Viatris.....	121
Dorzolamide.....	278	Eformoterol fumarate dihydrate.....	269	Entecavir.....	107
Dorzolamide with timolol.....	278	Eltrenonacog alfa [Recombinant factor IX].....	34	Entecavir (Rex).....	107
Dostinex.....	81	Efudix.....	72	Entrectinib.....	164
Dosulepin [Dothiepin] hydrochloride.....	129	Elaprase.....	19	Entresto 24/26.....	45
Dosulepin Viatris.....	129	Elecare (Unflavoured).....	302	Entresto 49/51.....	45
Dotarem.....	285	Elecare (Vanilla).....	302	Entresto 97/103.....	45
Dothiepin.....	129	Elecare LCP (Unflavoured).....	302	Entyvio.....	242
Dovato.....	107	Electral.....	43	Enzymes.....	117
Doxapram.....	272	Electrolytes.....	288	Ephedrine.....	54
Doxazosin.....	46	Elelyso.....	22	Ephedrine Juno.....	54
Doxazosin Clinect.....	46	Elemental 028 Extra (grapefruit).....	300	Epilim IV.....	132
Doxepin hydrochloride.....	129	Elemental 028 Extra (pineapple & orange).....	300	Epipen.....	263
Doxine.....	95	Elemental 028 Extra (summer fruits).....	300	Epipen Jr.....	263
Doxorubicin Ebewe.....	152	Elaxacaftor with tezacaftor, ivacaftor and ivacaftor.....	271	Epirubicin Ebewe.....	152
Doxorubicin hydrochloride.....	152	Elidel.....	71	Epirubicin hydrochloride.....	152
Doxycycline.....	95	Elocon.....	70	Eplerenone.....	50
DP Lotn HC.....	70	Elocon Alcohol Free.....	70	Epoetin alfa.....	30
		Eltrombopag.....	32	Epoetin beta.....	31
				Epoprostenol.....	62
				Eptacog alfa [Recombinant factor VIIa].....	34
				Eptifibatide.....	37
				Eptifibatide Viatris.....	37

Erbix	204	fraction	34	Fludarabine Ebewe	153
Ergometrine maleate	75	Famotidine	8	Fludarabine phosphate	153
Eritrox	93	Famotidine Hovid MY	8	Fludrocortisone acetate	80
Erlotinib	164	Faricimab	204	Fluids and Electrolytes	40
Ertapenem	89	Fasenra	200	Flumazenil	281
Erythrocin IV	93	Fatty Emulsion Cream (Evara)	69	Flumazenil-Baxter	281
Erythromycin (as ethylsuccinate)	93	Febuxostat	117	Flumetasone pivalate with clioquinol	275
Erythromycin (as lactobionate)	93	Febuxostat (Teva)	117	Fluocortolone caproate with fluocortolone pivalate and cinchocaine	7
Erythromycin (as stearate)	93	FEIBA NF	34	Fluorescein sodium	276
Esbriet	267	Felo 10 ER	49	Fluorescein sodium with lignocaine hydrochloride	276
Escitalopram	129	Felo 5 ER	49	Fluorescein sodium	276
Esmolol hydrochloride	48	Felodipine	49	Fluorometholone	276
Essential Amino Acid Mix	299	Fentanyl	127	Fluorouracil	153
Estradiol TDP Mylan	81	Fentanyl Sandoz	127	Fluorouracil Accord	153
Estradot	81	Ferinject	24	Fluorouracil sodium	72
Estrogel	81	Ferric derisomaltose	24	Fluox	130
Etanercept	178	Ferric subsulfate	33	Fluoxetine hydrochloride	130
Ethambutol hydrochloride	101	Feriprox	282	Flupenthixol decanoate	137
Ethanol	281	Ferro-F-Tabs	24	Flutamide	175
Ethanol with glucose	281	Ferro-Liquid	24	Flutamin	175
Ethanol, dehydrated	281	Ferro-tab	24	Fluticasone	269
Ethics Aspirin	125	Ferrosig	24	Fluticasone furoate with umeclidinium and vilanterol	266
Ethics Aspirin EC	37	Ferrous fumarate	24	Fluticasone furoate with vilanterol	270
Ethinylestradiol with desogestrel	74	Ferrous fumarate with folic acid	24	Fluticasone propionate	264
Ethinylestradiol with levonorgestrel	74	Ferrous gluconate with ascorbic acid	24	Fluticasone with salmeterol	270
Ethinylestradiol with norethisterone	74	Ferrous sulfate	24	Flynn	90
Ethosuximide	130	Ferrous sulfate with ascorbic acid	24	FML	276
Ethyl chloride	124	Fexaclear	264	Foban	67
Etomidate	122	Fexofenadine hydrochloride	264	Folic acid	31
Etopophos	154	Filgrastim	40	Folic Acid multichem	31
Etoposide	154	Finasteride	76	Folic Acid Viatrix	31
Etoposide (as phosphate)	154	Fingolimod	141	Fondaparinux sodium	36
Etoricoxib	119	Firazyr	263	Food Modules	292
Etravirine	105	Flagyl	103	Food/Fluid Thickeners	293
Eugia	55	Flagyl-S	103	Forteo	116
Eurofolic	173	Flamazine	67	Fortijuice (Apple)	309
Evara	68-69	Flecainide acetate	47	Fortijuice (Orange)	309
Evara Emulsifying Ointment	69	Flecainide BNM	47	Fortijuice (Strawberry)	309
EVARA White Soft Paraffin	69	Flecainide Controlled Release Teva	47	Fortini (Strawberry)	306
Everet	131	Fleet Phosphate Enema	16	Fortini (Vanilla)	306
Everolimus	256	Flixonase Hayfever & Allergy	264	Fortini Multi Fibre (chocolate)	306
Evrysdi	144	Flixotide	269	Fortini Multi Fibre (strawberry)	306
Exemestane	175	Flixotide Accuhaler	269	Fortini Multi Fibre (unflavoured)	306
Exjade	282	Florinef	80	Fortisip (Banana)	309
Extemporaneously Compounded Preparations	289	Fluanxol	137	Fortisip (Chocolate)	309
Eylea	199	Fluazazole	98	Fortisip (Strawberry)	309
Ezetimibe	53	Flucil	94	Fortisip (Vanilla)	309
Ezetimibe Sandoz	53	Flucloxacillin	94	Fortisip Multi Fibre (chocolate)	309
Ezetimibe with simvastatin	53	Flucloxin	94		
- F -		Fluconazole	98		
Factor eight inhibitor bypassing		Fluconazole-Baxter	98		
		Flucytosine	101		
		Fludara Oral	153		

Fortisip Multi Fibre (strawberry)	309	Glizide.....	11	HCU Anamix Infant.....	295
Fortisip Multi Fibre (vanilla)	309	Glucagen Hypokit	9	HCU Anamix Junior.....	295
Fosamax.....	115	Glucagon hydrochloride.....	9	HCU Anamix Junior LQ	295
Fosamax Plus.....	115	Glucerna Select.....	299	HCU Explore 5.....	295
Foscarnet sodium.....	108	Glucose [Dextrose] Alimentary	9	HCU Express 15.....	295
Fosfomycin.....	96	Blood.....	41	HCU Lophlex LQ	295
Framycetin sulphate	274	Extemporaneously Compounded Preparations	289	Healon	277
Fresofol 1% MCT/LCT.....	123	Glucose with potassium chloride	41	Healon 5.....	277
Frusemide.....	50	Glucose with potassium chloride and sodium chloride	41	Healon GV Pro	277
Fucidin.....	97	Glucose with sodium chloride.....	41	healthE Calamine Aqueous.....	68
Fucithalmic.....	274	Glucose with sucrose and fructose.....	9	healthE Dimethicone 10%.....	68
Fulvestrant.....	175	Glycerin with sodium saccharin.....	290	healthE Dimethicone 4% Lotion	67
Fulvestrant EVER Pharma	175	Glycerin with sucrose	290	healthE Dimethicone 5%.....	68
Fungilin.....	26	Glycerol		healthE Glycerol BP Liquid.....	290
Furosemide [Frusemide].....	50	Alimentary	16	healthE Urea Cream.....	70
Furosemide-AFT.....	50	Extemporaneously Compounded Preparations	290	Hemibra	33
Fusidic acid Dermatological.....	67, 71	Glycerol with paraffin.....	69	Heparin sodium	37
Infections.....	97	Glyceryl trinitrate Alimentary	7	Heparinised saline	37
Sensory.....	274	Cardiovascular	54	Heparon Junior.....	301
- G -		Glycine.....	286	Hepatitis A vaccine	316
GA Explore 5.....	294	Glycoprep Orange	15	Hepatitis B recombinant vaccine	316
GA1 Anamix Infant	294	Glycopyrronium	265	Herzuma	237
GA1 Anamix Junior.....	294	Glycopyrronium bromide	7	Hikma	46
Gabapentin.....	130	Glycopyrronium with indacaterol	265	Hikma Acetylcysteine	281
Gacet.....	126	Glycopyrronium-AFT	7	Hiprex	96
Gadobutrol.....	285	Glycosade.....	298	Histaclear.....	264
Gadoteric acid	285	Glytactin Bettermilk.....	297	Histamine acid phosphate	285
Gadovist 1.0	285	Gonadorelin	82	Holoxan	151
Gadoxetate disodium.....	285	Gonadron.....	78	Hormone Replacement Therapy	81
Galsulfase.....	19	Goserelin	83	HPV	317
Galvumet.....	11	Granisetron	134	Humalog Mix 25.....	10
Galvus	11	Granisetron-AFT.....	134	Humalog Mix 50.....	10
Ganciclovir.....	108	GTO Oil	298	Human Milk Fortifier	293
Gardasil 9	317	- H -		Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV]	317
Gastrodenol.....	9	Habitrol	149	Humatin	89
Gazyva	215	Habitrol (Fruit).....	149	Humira	192
Gefitinib	165	Habitrol (Mint).....	149	HumiraPen.....	192
Gelatine, succinylated	43	Haem arginate	19	Hyaluronic acid Alimentary.....	26
Gelofusine.....	43	Haemophilus influenzae type B vaccine	311	Sensory.....	277, 280
Gemcitabine Hydrochloride	153	Haldol	137	Hyaluronic acid with lidocaine [lignocaine]	26
Gemtuzumab ozogamicin.....	205	Haldol Concentrate.....	137	Hyaluronidase.....	117
Gentamicin sulphate Infections.....	89	Haloperidol	135	Hydralazine hydrochloride.....	55
Sensory.....	274	Haloperidol decanoate.....	137	Hydralock.....	68
Gestrinone.....	82	Hartmann's solution	41	Hydrapres.....	55
Gilenya	141	Harvoni.....	108	Hydrocortisone Dermatological.....	70
Ginet.....	74	Havrix 1440	316	Extemporaneously Compounded Preparations	290
Glatiramer acetate	141	Havrix Junior.....	316	Hormone Preparations.....	80
Glaucoma Preparations.....	278	Haylor Syrup.....	264	Hydrocortisone acetate.....	6
Glecaprevir with pibrentasvir	108			Hydrocortisone acetate with	
Glibenclamide.....	11				
Gliclazide.....	11				
Gliolan	177				
Glipizide.....	11				

pramoxine hydrochloride	6	Indapamide.....	51	IPCA-Frusemide	50
Hydrocortisone and paraffin liquid and lanolin	70	Indigo carmine.....	286	Ipca-Hydroxychloroquine.....	115
Hydrocortisone butyrate	70, 72	Indinavir.....	106	IPCA-Metoprolol	48
Hydrocortisone with miconazole.....	71	Indocyanine green.....	286	IPCA-Propranolol.....	48
Hydrocortisone with natamycin and neomycin	71	Indometacin [Indomethacin]	119	Ipilimumab	245
Hydrogen peroxide	67	Indomethacin.....	119	IPOLE	319
Hydroxocobalamin		Infanrix IPV	310	Ipratropium bromide	264-265
Alimentary	28	Infanrix-hexa.....	310	Ipratropium Viatrix	265
Various.....	281	Infatrin.....	304	Iressa.....	165
Hydroxocobalamin Panpharma	28	Infliximab	205	Irinotecan hydrochloride	155
hydroxycarbamide	154	Influenza vaccine.....	317	Iron (as ferric carboxymaltose).....	24
Hydroxychloroquine sulphate	115	Influvac Tetra		Iron (as sucrose).....	24
Hydroxyurea		(2026 formulation)	317	Iron polymaltose.....	24
[hydroxycarbamide].....	154	Inhaled Corticosteroids.....	269	Irrigation Solutions.....	286
Hygroton	51	Inlyta.....	161	Isentress	107
Hylo-Fresh	280	Inotuzumab ozogamicin.....	213	Isentress HD	107
Hyoscine butylbromide	7	Inresa.....	25	Ismo 20.....	54
Hyoscine Butylbromide		Inspra.....	50	Ismo 40 Retard	54
(Adiramedita)	7	Instillagel Lido	124	Isoflurane.....	122
Hyoscine hydrobromide	134	Insulin aspart	10	Isoleucine50	298
Hyperuricaemia and Antigout	117	Insulin aspart with insulin aspart protamine.....	10	Isoniazid	102
HypoPak Glucose.....	9	Insulin degludec with insulin aspart.....	10	Isoniazid Teva	102
Hypromellose.....	277, 279	Insulin glargine	10	Isoniazid with rifampicin.....	102
Hypromellose with dextran	279	Insulin glulisine	10	Isoprenaline [Isoproterenol]	55
- I -		Insulin isophane.....	10	Isopropyl alcohol.....	283
Ibiamox.....	93	Insulin isopro.....	10	Isoproterenol.....	55
Ibrutinib.....	154	Insulin lispro with insulin lispro protamine.....	10	Isoptin	49
Ibuprofen	119	Insulin lispro.....	10	Isoptin SR.....	49
Ibuprofen SR BNM	119	Insulin lispro with insulin lispro protamine.....	10	Isopto Carpine	278
Icatibant.....	263	Insulin neutral	10	Isosorbide mononitrate.....	54
Icatibant Lupin	263	Insulin neutral with insulin isophane.....	10	Isotretinoin	68
Idarubicin hydrochloride	152	Intelence	105	Ispaghula (psyllium) husk	15
Idarucizumab	34	Interferon alfa-2b.....	112	Isradipine.....	49
Idursulfase	19	Interferon beta-1-alpha.....	141	Itch-Soothe	68
Ifomid-M.....	151	Interferon beta-1-beta.....	141	Itraconazole.....	98
Ifosfamide.....	151	Interferon gamma	112	Itraconazole Crescent.....	98
Ifosfamide with mesna.....	151	Intra-uterine device.....	75	Itrazole.....	98
Ilomedin.....	64	Invanz	89	Ivabradine.....	47
Iloprost.....	64	Invega Sustenna.....	138	Ivacaftor	271
Imaging Agents.....	177	Invega Trinza.....	138	Ivermectin	102
Imatinib mesilate.....	165	Inza	120	- J -	
Imatinib-Rex	165	Iodine	87	Jadelle	75
Imbruvica	154	Iodine with ethanol.....	283	Jakavi	170
Imfinzi	245	Iodised oil	284	Jardiamet.....	13
Imipenem with cilastatin	89	Iodixanol	284	Jardiance.....	13
Imipenem+Cilastatin RBX	89	Iohexol	284	Jaydess	75
Imipramine Crescent	129	Iopidine	279	Jevity HiCal RTH	308
Imipramine hydrochloride	129	Ioscan	284	Jevity RTH	308
Imiquimod.....	72	Ipca-Allopurinol.....	117	Jinarc.....	51
Immune Modulators.....	112	Ipca-Bisoprolol.....	47	- K -	
Immunosuppressants	177	Ipca-Ciprofloxacin.....	94	Kadcyla.....	239
Impact Advanced Recovery.....	307	Ipca-Donepezil.....	147	Kalydeco.....	271
Incruse Ellipta.....	265	Ipca-Enalapril.....	44	Kenacomb	275
Indacaterol.....	269	Ipca-Escitalopram.....	129	Kenacort-A 10.....	80
				Kenacort-A 40.....	80
				Kenalog in Orabase.....	26

Ketalar.....	123	Lenvatinib.....	165	Lo-Oralcon 20 ED.....	74
Ketamine.....	123	Lenvima.....	165	Local Preparations for Anal and Rectal Disorders.....	7
Ketocal 3:1 (Unflavoured).....	305	Letemovir.....	108	Locasol.....	304
Ketocal 4:1 (Unflavoured).....	305	Letrole.....	176	Locoid.....	70, 72
Ketocal 4:1 (Vanilla).....	305	Letrozole.....	176	Locoid Crelo.....	70
Ketoconazole.....		Leucine100.....	299	Locoid Lipocream.....	70
Dermatological.....	67	Leukotriene Receptor Antagonists.....	269	Lodoxamide.....	276
Infections.....	97	Leuprorelin acetate.....	83	Logem.....	131
Ketoprofen.....	119	Leustatin.....	152	Lomide.....	276
Ketorolac trometamol.....	276	Levetiracetam.....	131	Lomustine.....	151
KetoSens.....	321	Levetiracetam-AFT.....	131	Long-Acting Beta-Adrenoceptor Agonists.....	269
Ketostix.....	321	Levocabastine.....	276	Loniten.....	56
Keytruda.....	249	Levocarnitine.....	20	Loperamide hydrochloride.....	5
Kindergen.....	306	Levodopa with benserazide.....	122	Lopinavir with ritonavir.....	107
Kisqali.....	170	Levodopa with carbidopa.....	122	Lopinavir/Ritonavir Mylan.....	107
Klacid.....	92	Levodopa with carbidopa and entacapone.....	122	Loratadine.....	264
Klacid IV.....	92	Levomepromazine.....	136	Lorazepam.....	130, 140
Klaricid.....	92	Levomepromazine hydrochloride.....	136	Lormetazepam.....	143
Kogenate FS.....	35	Levonorgestrel.....	75	Lorstat.....	52
Konakion MM.....	35	Levonorgestrel -1 (Lupin).....	75	Losartan & Hydrochlorothiazide (lpca).....	45
Konakion MM Paediatric.....	35	Levosimendan.....	54	Losartan Actavis.....	45
Konsyl-D.....	15	Levothyroxine.....	87	Losartan potassium.....	45
Kuvan.....	21	Lidocaine [Lignocaine].....	124	Losartan potassium with hydrochlorothiazide.....	45
- L -		Lidocaine [Lignocaine] hydrochloride.....	124	Lovir.....	108
L-ornithine L-aspartate.....	9	Lidocaine [Lignocaine] hydrochloride with adrenaline.....	124	Loxamine.....	130
Labetalol.....	48	Lidocaine [Lignocaine] hydrochloride with adrenaline and tetracaine hydrochloride.....	124	Lucrin Depot 1-month.....	83
Lacosamide.....	131	Lidocaine [Lignocaine] hydrochloride with phenylephrine hydrochloride.....	125	Lucrin Depot 3-month.....	83
Lactose.....	290	Lidocaine [Lignocaine] with prilocaine.....	125	LumaCina.....	286
Lactulose.....	16	Lidocaine-Baxter.....	124	Lumigan.....	278
Laevolac.....	16	lignocaine.....		Lynparza.....	156
Lamictal.....	131	Alimentary.....	26	Lyrica.....	132
Lamivudine.....	106-107	Nervous.....	124-125	Lysine acetylsalicylate [Lysine aspirin].....	38
Lamivudine Viatris.....	106	Lincomycin.....	96	Lysine aspirin.....	38
Lamivudine/Zidovudine Viatris.....	106	Linezolid.....	96	- M -	
Lamotrigine.....	131	Linezolid Kabi.....	96	m-Eslon.....	127
Lanoxin.....	47	Lioresal Intrathecal.....	118	Mabthera.....	220
Lanoxin PG.....	47	Liothyronine sodium.....	87	Macrobid.....	97
Lanreotide.....	177	Lipid-Modifying Agents.....	52	Macrogl 3350 with ascorbic acid, potassium chloride, sodium chloride and citric acid with magnesium carbonate hydrate and sodium picosulfate.....	15
Lansoprazole.....	8	Lipiodol Ultra Fluid.....	284	Macrogl 3350 with potassium chloride and sodium chloride.....	15
Lansoprazole (Viatris).....	8	Liquigen.....	292	Macrogl 3350 with potassium chloride and sodium chloride with/ without sodium sulfate, sodium ascorbate, ascorbic acid.....	15
Lantus.....	10	Liraglutide.....	12	Macrogl 3350 with potassium chloride, sodium bicarbonate and sodium chloride.....	16
Lantus SoloStar.....	10	Lisdexamfetamine dimesilate.....	145		
Lanzol Relief.....	8	Lisinopril.....	44		
Lapatinib.....	165	Lissamine green.....	276		
Largactil.....	135	Lithium carbonate.....	136		
Laronidase.....	19	LMX4.....	124		
Lasix.....	50				
Latanoprost.....	278				
Latanoprost with timolol.....	278				
Lax-Suppositories.....	16				
Lax-suppositories Glycerol.....	16				
Laxatives.....	15				
Ledipasvir with sofosbuvir.....	108				
Leflunomide.....	115				
Lenalidomide (Viatris).....	155				
Lenalidomide Viatris.....	155				

Madopar 125	122	Melphalan	151	Methylprednisolone (as sodium succinate)	80
Madopar 250	122	Meningococcal (A, C, Y and W-135) conjugate vaccine	311	Methylprednisolone aceponate	70
Madopar 62.5	122	Meningococcal B multicomponent vaccine	312	Methylprednisolone acetate	80
Madopar HBS	122	MenQuadfi	311	Methylthionium chloride [Methylene blue]	286
Madopar Rapid	122	Menthol	290	Methylxanthines	270
Mafenide acetate	67	Mepivacaine hydrochloride	125	Metoclopramide Actavis 10	134
Magnesium amino acid chelate	24	Mepivacaine hydrochloride with adrenaline	125	Metoclopramide hydrochloride	134
Magnesium chloride	24	Mepolizumab	214	Metoclopramide hydrochloride with paracetamol	133
Magnesium hydroxide		Mercaptopurine	153	Metolazone	51
Alimentary	24	Meropenem	90	Metoprolol IV Mylan	48
Extemporaneously Compounded Preparations	290	Meropenem-AFT	90	Metoprolol IV Viatris	48
Magnesium oxide	25	Mesalazine	6	Metoprolol succinate	48
Magnesium oxide with magnesium aspartate, magnesium amino acid chelate and magnesium		Mesna	174	Metoprolol tartrate	48
citrate	25	Mestinon	115	Metronidamed	103
Magnesium sulphate	25	Metabolic Disorder Agents	17	Metronidazole	
Magnevist	285	Metabolic Products	294	Dermatological	67
Malarone	103	Metalyse	39	Infections	103
Malarone Junior	103	Metaraminol	55	Metyrapone	82
Malathion [Maldison]	68	Metformin hydrochloride	11	Mexiletine hydrochloride	47
Maldison	68	Metformin Viatris	11	Miacalcin	78
Mannitol		Methacholine chloride	286	Mianserin hydrochloride	129
Cardiovascular	50	Methadone BNM	127	Micolette	16
Various	285	Methadone hydrochloride		Miconazole	26
Mantoux	320	Extemporaneously Compounded Preparations	290	Miconazole nitrate	
Maprotiline hydrochloride	129	Nervous	127	Dermatological	67
Marcaïn	123	Methenamine (Hexamine) hippurate	96	Genito-Urinary	74
Marcaïn Heavy	124	Methohexital sodium	123	Micreme	74
Marcaïn Isobaric	123	Methopt	279	Micreme H	71
Marcaïn with Adrenaline	123	Methotrexate	153	Microlut	75
Marevan	37	Methotrexate DBL Onco-Vial	153	Midazolam	143
Marine Blue Lotion SPF 50+	72	Methotrexate Ebewe	153	Midazolam-Baxter	143
Mask for spacer device	321	Methotrexate Pharmascience	153	Midazolam-Pfizer	143
Maviret	108	Methotrexate Sandoz	153	Midodrine	47
Maxidex	275	Methoxsalen		Midostaurin	167
Maxitrol	274	[8-methoxypsoralen]	71	Mifepristone	75
MCT Oil	292	Methoxyflurane	125	Milrinone	56
Measles, mumps and rubella vaccine	318	Methyl aminolevulinate hydrochloride	72	Milrinone-Baxter	56
Mebedazole	103	Methyl hydroxybenzoate	290	Minerals	23
Mebeverine hydrochloride	7	Methylcellulose	290	Mini-Wright AFS Low Range	321
Medac	151	Methylcellulose with glycerin and sodium saccharin	290	Mini-Wright Standard	321
Medrol	80	Methylcellulose with glycerin and sucrose	290	Minidiab	11
Medroxyprogesterone	82	Methyldopa	50	Minims Prednisolone	276
Medroxyprogesterone acetate		Methylphenidate Viatris	50	Minirin	88
Genito-Urinary	75	Methylene blue	286	Minirin Melt	88
Hormone Preparations	81	Methylnaltrexone bromide	16	Minocycline	95
Mefenamic acid	120	Methylphenidate ER - Teva	146	Minoxidil	56
Mefloquine	103	Methylphenidate hydrochloride	146	Mirena	75
Meglumine gadopentetate	285	Methylphenidate Sandoz XR	146	Miro-Amoxicillin	93
Meglumine iotroxate	285			Mirtazapine	129
Mekinist	171			Misoprostol	7
Melatonin	143			Mitomycin C	152
Melpha	151			Mitozantrone	152
				Mitozantrone Ebewe	152

Mivacurium chloride.....	118	Mydratics and Cycloplegics	279	Nimotop	49
Mixed salt solution for eye irrigation.....	277	Mylan (24 hr release).....	49	Nintedanib	267
MMA/PA Anamix Infant	298	Mylan Clomiphen.....	82	Niraparib	155
MMA/PA Anamix Junior.....	298	Mylan Italy (24 hr release)	49	Nirmatrelvir with ritonavir	111
MMA/PA Explore 5	298	Myleran	151	Nitazoxanide.....	103
MMA/PA Express 15	298	Myloc CR	48	Nitisinone.....	20
Moclobemide	129	Mylotarg.....	205	Nitisinone LogixX Pharma	20
Modafinil	147	Myozyme	17	Nitrates	54
Modafinil Max Health	147	Mytolac	177	Nitroderm TTS 10.....	54
Molaxole	16	- N -		Nitroderm TTS 5.....	54
Mometasone furoate.....	70	Nadolol	48	Nitrofurantoin.....	97
Monofer	24	Nadolol BNM	48	Nitrofurantoin Clinect	97
Monogen.....	300	Naglazyme.....	19	Nitrolingual Pump Spray.....	54
Monosodium glutamate with sodium aspartate.....	288	Naloxone hydrochloride.....	281	Nivestim	40
Monosodium l-aspartate.....	288	Naltraccord	148	Nivolumab.....	246
Montelukast	269	Naltrexone hydrochloride.....	148	Nodia	5
Montelukast Paediatric (Relonchem).....	269	Naphazoline hydrochloride.....	276	Noflam 250	120
Montelukast Viatrix	269	Naprosyn SR 1000	120	Noflam 500	120
Morotocog alfa [Recombinant factor VIII]	35	Naprosyn SR 750	120	Non-Steroidal Anti-Inflammatory Drugs.....	119
Morphine hydrochloride	127	Naproxen	120	Nonacog gamma, [Recombinant factor IX]	35
Morphine sulphate	127	Natalizumab.....	141	Noradrenaline.....	55
Morphine tartrate	127	Natamycin.....	274	Norethisterone Genito-Urinary.....	75
Motetis	121	Natulan	158	Hormone Preparations.....	82
Mouth and Throat.....	25	Nausafix.....	135	Norethisterone with mestranol.....	74
Movapo.....	121	Nausicalm.....	134	Norflex	118
Movicol	16	Nefopam hydrochloride	126	Norflex	95
Moxifloxacin.....	94	Neo-Mercazole	87	Noriday	75
Moxifloxacin Kabi.....	94	Neocate Gold (Unflavoured).....	302	Noriday 28	75
Moximed	94	Neocate Junior Unflavoured	302	Normison	143
Mozobil	39	Neocate Junior Vanilla.....	302	Nortriptyline hydrochloride.....	129
MSUD Anamix Infant.....	295	Neocate SYNEO.....	302	Norvir	107
MSUD Anamix Junior	295	Neoral	177	Noumed Dexamfetamine.....	145
MSUD Anamix Junior LQ	295	Neostigmine metilsulfate	115	Noumed Isoniazid.....	102
MSUD Explore 5	295	Neostigmine metilsulfate with glycopyrronium bromide	115	Noumed Paracetamol.....	126
MSUD Express 15	295	Neosyneprine HCL	55	Noumed Pethidine	128
MSUD Lophlex LQ 20.....	295	Nepro HP (strawberry).....	306	Noumed Phenobarbitone.....	131
MSUD Maxamum	295	Nepro HP (vanilla)	306	Novadoz	151
Mucolytics and Expectorants.....	270	Neulumex	284	Novasource Renal (Vanilla).....	307
Multiple Sclerosis Treatments	140	Neupogen	40	Novatretin	71
Multivitamin and mineral supplement.....	26	NeuroTabs.....	24	NovoMix 30 FlexPen.....	10
Multivitamin renal.....	27	Nevirapine	105	NovoRapid FlexPen.....	10
Multivitamins.....	27	Nevirapine Viatrix	105	NovoSeven RT	34
Mupirocin.....	67	Nicardipine hydrochloride	49	Nozinan	136
Muscle Relaxants and Related Agents	118	Nicorandil.....	56	Nucala	214
Mvite	27	Nicotine	149	Nuelin	270
Myambutol	101	Nifedipine.....	49	Nuelin-SR	270
Mycobutin	102	Nifuran	97	Nupentin	130
MycosNail.....	67	Nilotinib.....	167	Nusinersen	143
Mycophenolate mofetil.....	257	Nilstat		Nutilis.....	294
Mydracyl	279	Alimentary	26	Nutren Diabetes (vanilla).....	299
		Genito-Urinary.....	74	Nutrini Energy Multi Fibre	306
		Infections.....	98	Nutrini Energy RTH	306
		Nimenrix	312	Nutrini Low Energy Multi Fibre	
		Nimodipine.....	49		
		Nimorhage	49		

RTH.....	306	Olopatadine.....	276	Oxytocin.....	75
Nutrini Peptisorb Energy.....	303	Olopatadine Teva.....	276	Oxytocin BNM.....	75
Nutrini RTH.....	306	Olsalazine.....	7	Oxytocin with ergometrine maleate.....	75
Nutrison 800 Complete Multi Fibre.....	308	Omalizumab.....	216	Ozurdex.....	275
Nutrison Advanced Peptisorb.....	300	Omeprazole.....	8	- P -	
Nutrison Concentrated.....	301	Omeprazole actavis 10.....	8	Pacifen.....	118
Nutrison Energy.....	308	Omeprazole actavis 20.....	8	Pacimol.....	126
Nutrison Energy Multi Fibre.....	308	Omeprazole actavis 40.....	8	Paclitaxel.....	173
Nutrison Multi Fibre.....	308	Omeprazole Teva.....	8	Padagis.....	72
Nutrison Protein Intense.....	302	Omezol IV.....	8	Paediatric Seravit.....	27
Nutrison Protein Plus.....	301	Omnipaque.....	284	Palbociclib.....	168
Nutrison Protein Plus Multi Fibre.....	302	Omnitrope.....	83	Palbociclib Pfizer.....	168
Nutrison RTH.....	308	Onbrez Breezhaler.....	269	Paliperidone.....	138
Nyefax Retard.....	49	Oncaspar LYO.....	157	Paliperidone palmitate.....	138
Nystatin.....		OncoTICE.....	256	Palivizumab.....	217
Alimentary.....	26	Ondansetron.....	134	Pamidronate disodium.....	115
Dermatological.....	67	Ondansetron-AFT.....	134	Pamisol.....	115
Genito-Urinary.....	74	One-Alpha.....	28	Pamol.....	126
Infections.....	98	Opdivo.....	246	Pancreatic enzyme.....	14
- O -		Optional Pharmaceuticals.....	321	Pancuronium bromide.....	118
Obinutuzumab.....	215	Ora-Blend.....	290	Pantoprazole.....	8
Obstetric Preparations.....	75	Ora-Blend SF.....	290	Panzop Relief.....	8
Ocrelizumab.....	141	Ora-Plus.....	290	Papaverine hydrochloride.....	56
Ocrevus.....	141	Ora-Sweet.....	290	Paper wasp venom.....	263
Ocrevus SC.....	141	Ora-Sweet SF.....	290	Para-aminosalicylic Acid.....	102
Octasa.....	6	Oralcon 30 ED.....	74	Paracetamol.....	126
Octocog alfa [Recombinant factor VIII] (Advate).....	35	Oramorph.....	127	Paracetamol (Ethics).....	126
Octocog alfa [Recombinant factor VIII] (Kogenate FS).....	35	Oramorph CDC S29.....	127	Paracetamol Kabi.....	126
Octreotide.....	175	Oratane.....	68	Paracetamol with codeine.....	128
Octreotide Long-Acting.....	177	Ornidazole.....	103	Paraffin	
Ocular Lubricants.....	279	Orphenadrine citrate.....	118	Alimentary.....	15
Oestradiol.....	81-82	Oruvail SR.....	119	Dermatological.....	69
Oestradiol valerate.....	81	Oseltamivir.....	111	Extemporaneously Compounded Preparations.....	290
Oestradiol with norethisterone acetate.....	81	Osimertinib.....	167	Paraffin liquid with soft white paraffin.....	280
Oestriol		Osmolite RTH.....	308	Paraffin liquid with wool fat.....	280
Genito-Urinary.....	76	Other Cardiac Agents.....	54	Paraffin with wool fat.....	69
Hormone Preparations.....	82	Other Endocrine Agents.....	81	Paraldehyde.....	130
Oestrogens.....	76	Other Oestrogen Preparations.....	82	Parecoxib.....	120
Oestrogens (conjugated equine).....	81	Other Otological Preparations.....	280	Paromomycin.....	89
Oestrogens with medroxyprogesterone acetate.....	81	Other Progesterone Preparations.....	82	Paroxetine.....	130
Ofev.....	267	Other Skin Preparations.....	72	Paser.....	102
Oil in water emulsion.....	69	Ovestin		Patent blue V.....	286
Oily phenol [Phenol oily].....	7	Genito-Urinary.....	76	Paxam.....	139
Olanzapina Mylan.....	136	Hormone Preparations.....	82	Paxlovid.....	111
Olanzapina Mylan Pharma.....	136	Oxaliplatin.....	161	Pazopanib.....	169
Olanzapine.....	136-137	Oxandrolone.....	78	Pazopanib ADVZ.....	169
Olanzapine ODT Viatris.....	136	Oxazepam.....	140	Pazopanib Teva.....	169
Olaparib.....	156	Oxpentifylline.....	56	Peak flow meter.....	321
Olive oil.....	290	Oxybuprocaine hydrochloride.....	277	Peanut oil.....	289
		Oxybutynin.....	77	Pedialyte.....	43
		Oxycodone Amneal.....	128	Pediasure (chocolate).....	306
		Oxycodone hydrochloride.....	128	Pediasure (strawberry).....	306
		Oxycodone Sandoz.....	128	Pediasure (vanilla).....	306
		Oxymetazoline hydrochloride.....	268		

Pediasure Plus	306	Phentolamine mesylate	46	PKU Lophlex Sensations	
Pediasure RTH	306	Phenylalanine50	299	20 (berries)	297
Pegaspargase	157	Phenylephrine hydrochloride		PKU Restore Powder	296
Pegasys	112	Cardiovascular	55	PKU sphere20 Banana	297
Pegasys (S29)	112	Sensory	279	PKU sphere20 Chocolate	297
Pegfilgrastim	40	Phenytoin	131	PKU sphere20 Lemon	297
Pegylated interferon alfa-2a	112	Phenytoin sodium	130–131	PKU sphere20 Red Berry	297
Pembrolizumab	249	Phesgo	219	PKU sphere20 Vanilla	297
Pemetrexed	153	Phlexy 10	296	PKU Start	296
Pemetrexed-AFT	153	Phosphorus	43	Plasma-Lyte 148	41
Penicillamine	115	Phytomenadione	35	Plasma-Lyte 148 & 5% Glucose	41
Penicillin G	93	Picibanil	257	Plendil ER	49
Penicillin V	94	Pilocarpine hydrochloride	278	Plenvu	15
Pentacarinat	103	Pilocarpine nitrate		Plerixafor	39
Pentagastrin	82	Extemporaneously Compounded		Plerixafor-AFT	39
Pentamidine isethionate	103	Preparations	290	Pneumococcal (PCV13) conjugate	
Pentasa	6	Sensory	278	vaccine	313
Penthrox	125	Pimafulcort	71	Pneumococcal (PPV23)	
Pentostatit [Deoxycofomycin]	157	Pimecrolimus	71	polysaccharide vaccine	314
Pentoxifylline [Oxpentifylline]	56	Pine tar with trolamine laurilsulfate		Pneumovax 23	314
Peptamen OS 1.0 (Vanilla)	300	and fluorescein	71	Podophyllotoxin	72
Pepti-Junior	304	Pinetarsol	71	Polidocanol	33
Perflutren	285	Pioglitazone	11	Poliomyelitis vaccine	319
Perhexiline maleate	49	Piperacillin with tazobactam	94	Poloxamer	15
Pericyazine	136	Pipothiazine palmitate	139	Poly-Tears	279
Perindopril	44	PipTaz-AFT	94	Poly-Visc	280
Periset	134	Pirfenidone	267	Polycal	292
Periset ODT	134	Pirfenidone Sandoz	267	Polyethylene glycol 400 and	
Perjeta	218	Pituitary and Hypothalamic		propylene glycol	280
Permethrin	68	Hormones and Analogues	82	Polyflush	42
Perrigo	72	Pivmecillinam	97	Polyhexamethylene biguanide	290
Pertuzumab	218	Pizotifen	133	Polyvinyl alcohol with povidone	280
Pertuzumab with trastuzumab	219	PKU Anamix Infant	296	Pomalidomide	158
Peteha	102	PKU Anamix Junior	296	Pomolide	158
Pethidine hydrochloride	128	PKU Anamix Junior LQ (Berry)	296	Poractant alfa	272
Pevaryl	67	PKU Anamix Junior LQ		Posaconazole	98
Pexsig	49	(Orange)	296	Posaconazole Juno	98
Pfizer Exemestane	175	PKU Build 10	297	Potassium chloride	41, 43
Pharmacor Varenicline	149	PKU Build 20 Chocolate	297	Potassium chloride with sodium	
Pheburane	21	PKU Build 20 Raspberry		chloride	42
Phenasen	154	Lemonade	297	Potassium citrate	76
Phenelzine sulphate	129	PKU Build 20 Smooth	297	Potassium dihydrogen	
Phenindione	37	PKU Build 20 Vanilla	297	phosphate	42
Phenobarbitone	131, 143	PKU Explore 10	296	Potassium iodate	
Phenobarbitone sodium	290	PKU Explore 5	296	Alimentary	24
Phenol		PKU Express 20	296	Hormone Preparations	87
Extemporaneously Compounded		PKU First Spoon	296	Potassium iodate with iodine	24
Preparations	290	PKU Glytactin RTD 15	297	Potassium perchlorate	87
Various	287	PKU Glytactin RTD 15 Lite	297	Potassium permanganate	72
Phenol oily	7	PKU GMPro LQ	297	Povidone K30	290
Phenol with ioxaglic acid	287	PKU GMPro Mix-In	297	Povidone-iodine	283
Phenothrin	68	PKU GMPro Ultra Lemonade	297	Povidone-iodine with ethanol	283
Phenoxybenzamine		PKU GMPro Ultra Vanilla	297	Pradaxa	36
hydrochloride	46	PKU Lophlex LQ 10	296	Pralidoxime chloride	281
Phenoxyethylpenicillin [Penicillin		PKU Lophlex LQ 20	296–297	Pralidoxime iodide	281
V]	94	PKU Lophlex Powder	296	Pramipexole hydrochloride	122

Pravastatin.....	52	Provera.....	81	Resource Beneprotein.....	293
Praxbind.....	34	Provera HD.....	82	Respiratory Stimulants.....	272
Praziquantel.....	103	Proxymetacaine hydrochloride.....	277	Retinol.....	27
Prazosin.....	46	Pseudoephedrine.....		Retinol Palmitate.....	280
Prazosin Mylan.....	46	hydrochloride.....	268	ReTrieve.....	68
Pred Forte.....	276	Psoriasis and Eczema.....		Retrovir.....	106
Prednisolone.....	80	Preparations.....	71	Retrovir IV.....	106
Prednisolone acetate.....	276	PTU.....	87	Revolade.....	32
Prednisolone hexanoate with.....		Pulmonary Surfactants.....	272	Ribociclib.....	170
cinchocaine hydrochloride.....	7	Pulmozyme.....	270	Riboflavin.....	20
Prednisolone sodium.....		Puri-nethol.....	153	Riboflavin 5-phosphate.....	278
phosphate.....	276	Pyrazinamide.....	102	Ricit.....	76
Prednisolone- AFT.....	276	Pyridostigmine bromide.....	115	Ricovir.....	108
Prednisone.....	80	Pyridoxal-5-phosphate.....	20	Rifabutin.....	102
Prednisone Clinect.....	80	Pyridoxine hydrochloride.....	28	Rifadin.....	102
Pregabalin.....	132	Pyridoxine multichem.....	28	Rifamazid.....	102
Pregabalin Pfizer.....	132	Pyrimethamine.....	104	Rifampicin.....	102
Pregnancy test - hCG urine.....	321	- Q -.....		Rifaximin.....	9
preOp.....	307	Questran Light (Neon).....	53	Rifinah.....	102
Presolol.....	48	Quetapel.....	136	Rilutek.....	121
Prevenar 13.....	313	Quetiapine.....	136	Riluzole.....	121
Prevymis.....	108	Quetiapine Viatrix.....	136	Ringer's solution.....	42
Priadel.....	136	Quinapril.....	44	Rinvoq.....	259
Prilocaine hydrochloride.....	125	Quinine dihydrochloride.....	104	Riodine.....	283
Prilocaine hydrochloride with.....		Qvar.....	269	Risdiplam.....	144
felypressin.....	125	- R -.....		Risedronate Sandoz.....	115
Primaquine.....	103	RA-Morph.....	127	Risedronate sodium.....	115
Primidone.....	132	Rabies vaccine.....	319	Risperdal.....	136
Primolut N.....	82	Raltegravir potassium.....	107	Risperdal Consta.....	139
Primovist.....	285	Ramipex.....	122	Risperidone.....	136, 139
Priorix.....	318	Ramipril.....	44	Risperidone (Teva).....	136
Probenecid.....	118	Ramipril Mylan Generics.....	44	Risperidone Sandoz.....	136
Procaine penicillin.....	94	Ramipril Viatrix.....	44	Risperon.....	136
Procarbazine hydrochloride.....	158	Ranbaxy-Cefaclor.....	90	Ritalin.....	146
Prochlorperazine.....	135	Ranibizumab.....	220	Ritalin LA.....	146
Proctosedyl.....	7	Ranitidine.....	8	Ritonavir.....	107
Procyclidine hydrochloride.....	121	Rapamune.....	257	Rituximab (mabthera).....	220
Progesterone.....	76	Rasagiline.....	122	Rituximab (riximyo).....	222
Proglicem.....	9	Rasburicase.....	118	Rivaroxaban.....	37
Proglycem.....	9	Readi-CAT 2.....	284	Rivastigmine.....	148
Prodynova.....	81	Reandron 1000.....	78	Rivastigmine Patch BNM 10.....	148
Prolia.....	116	Recombinant factor IX.....	34-35	Rivastigmine Patch BNM 5.....	148
Promethazine hydrochloride.....	264	Recombinant factor VIIa.....	34	Riximyo.....	222
Propafenone hydrochloride.....	47	Recombinant factor VIII.....	35	RIXUBIS.....	35
Propamide isethionate.....	274	Rectogesic.....	7	Rizamelt.....	133
Propofol.....	123	Red back spider antivenom.....	282	Rizatriptan.....	133
Propranolol.....	48	Redipred.....	80	Rizatriptan Rising.....	133
Propranolol LA Lupin.....	48	Relenza Rotadisk.....	111	Rizatriptan Viatrix.....	133
Propylthiouracil.....	87	Relistor.....	16	Rocuronium bromide.....	118
Prostin E2.....	75	Remdesivir.....	112	Roma.....	43
Prostin VR.....	55	Remicade.....	205	Ropin.....	122
Protamine sulphate.....	37	Remifentanil.....	128	Ropinrole hydrochloride.....	122
Protifar.....	293	Remifentanil-AFT.....	128	Ropivacaine hydrochloride.....	125
Protionamide.....	102	Renilon 7.5 (apricot).....	307	Ropivacaine Kabi.....	125
Protirelin.....	88	Renilon 7.5 (caramel).....	307	Rose bengal sodium.....	276
Proveblue.....	286	Resonium A.....	43	Rosemont.....	128

Rosuvastatin.....	52	Shingles vaccine.....	320	Sodium cromoglicate	
Rosuvastatin - Sandoz	52	Shingrix.....	320	Alimentary.....	7
Rosuvastatin Mylan.....	52	SII-Onco-BCG.....	256	Respiratory.....	264
Rosuvastatin Viatriis.....	52	Sildenafil.....	61	Sensory.....	276
Rotarix.....	319	Siltuximab.....	233	Sodium dihydrogen phosphate	
Rotavirus oral vaccine.....	319	Silver diamine fluoride.....	26	[Sodium acid phosphate].....	42
Roxithromycin.....	93	Silver nitrate		Sodium fluoride.....	24
Rozlytrek.....	164	Dermatological.....	72	Sodium fusidate [Fusidic acid]	
Rubifen.....	146	Extemporaneously Compounded		Dermatological.....	67
Rubifen SR.....	146	Preparations.....	290	Infections.....	97
Rugby.....	25	Simdax.....	54	Sensory.....	274
Rurioctocog alfa pegol [Recombinant		Simeticone.....	5	Sodium hyaluronate [Hyaluronic acid]	
factor VIII].....	35	Simulect.....	200	Alimentary.....	26
Ruxolitinib.....	170	Simvastatin.....	53	Sensory.....	277, 280
Rydapt.....	167	Simvastatin Mylan.....	53	Sodium hyaluronate [Hyaluronic acid]	
Ryzodeg 70/30 Penfill.....	10	Simvastatin Viatriis.....	53	with chondroitin sulphate.....	277
- S -		Sincalide.....	286	Sodium hydroxide.....	287
S26 LBW Gold RTF.....	305	Sinemet.....	122	Sodium hypochlorite.....	284
Sacubitril with valsartan.....	45	Sinemet CR.....	122	Sodium metabisulfite.....	291
SalAir.....	268	Sintetica Baclofen Intrathecal.....	118	Sodium nitrite.....	281
Salazopyrin.....	7	Sirolimus.....	257	Sodium nitroprusside	
Salazopyrin EN.....	7	Sirturo.....	101	Cardiovascular.....	56
Salbutamol.....	268	Siterone.....	78	Optional Pharmaceuticals.....	321
Salbutamol with ipratropium		Slow-Lopresor.....	48	Sodium phenylbutyrate.....	21
bromide.....	265	Snake antivenom.....	282	Sodium phosphate with phosphoric	
Salicylic acid.....	290	Sodibic.....	43	acid.....	16
Salmeterol.....	269	Sodium acetate.....	42	Sodium picosulfate.....	16
Salmonella typhi vaccine.....	315	Sodium acid phosphate.....	42	Sodium polystyrene sulphonate.....	43
Sandimmun.....	177	Sodium alginate with magnesium		Sodium stibogluconate.....	104
Sandomigran.....	133	alginate.....	5	Sodium tetradecyl sulphate.....	34
Sandostatin LAR.....	177	Sodium alginate with sodium		Sodium thiosulfate.....	281
Sapropterin Dihydrochloride.....	21	bicarbonate and calcium		Sodium valproate.....	132
Scalp Preparations.....	72	carbonate.....	5	Sodium with potassium.....	288
Scandonest 3%.....	125	Sodium aurothiomalate.....	115	Solax.....	15
Scheriproct.....	7	Sodium benzoate.....	21	Solifenacin succinate.....	77
Sclerosing Agents.....	273	Sodium bicarbonate		Solifenacin succinate Max	
Scopolamine Transdermal System		Blood.....	42-43	Health.....	77
Viatriis.....	134	Extemporaneously Compounded		Solu-Cortef.....	80
Sebizole.....	67	Preparations.....	290	Solu-Medrol.....	80
Secretin pentahydrochloride.....	286	Sodium calcium edetate.....	283	Solu-Medrol Act-O-Vial.....	80
Secukinumab.....	231	Sodium chloride		Somatropin.....	83
Sedatives and Hypnotics.....	142	Blood.....	42-43	Somatuline Autogel.....	177
Seebri Breezhaler.....	265	Respiratory.....	268, 272	Sotalol.....	48
Selegiline hydrochloride.....	122	Various.....	286	Soya oil.....	281
Selenium.....	25	Sodium chloride with sodium		Spacer device.....	321
Sennosides.....	16	bicarbonate.....	268	Span-K.....	43
Serc.....	134	Sodium citrate		Spazmol.....	7
Serenace.....	135	Alimentary.....	5	Specialised Formulas.....	299
Seretide.....	270	Extemporaneously Compounded		Spinal Muscular Atrophy.....	143
Seretide Accuhaler.....	270	Preparations.....	291	Spinraza.....	143
Serevent.....	269	Sodium citrate with sodium chloride		Spiocto Respimat.....	265
Serevent Accuhaler.....	269	and potassium chloride.....	37	Spiractin.....	51
Sertraline.....	130	Sodium citrate with sodium lauryl		Spiramycin.....	104
Setrona.....	130	sulphoacetate.....	16	Spiriva.....	265
Sevoflurane.....	123	Sodium citro-tartrate.....	76	Spiriva Respimat.....	265
Sevredol.....	127				

Spironolactone.....	51	Talc.....	273	Tetracosactide [Tetracosactrin].....	82
Stalevo.....	122	Taliglucerase alfa.....	22	Tetracosactrin.....	82
Standard Feeds.....	307	Tambocor.....	47	Tetracycline.....	95
Staphlex.....	94	Tambocor German.....	47	Tetracycline AA.....	95
Starch.....	291	Tamoxifen citrate.....	175	Teva Lisinopril.....	44
Stavudine.....	106	Tamoxifen Sandoz.....	175	Thalidomide.....	159
Stelara.....	240	Tamsulosin hydrochloride.....	76	Thalomid.....	159
Sterculia with frangula.....	15	Tamsulosin-Rex.....	76	Theobroma oil.....	291
SteroClear.....	264	Tasigna.....	167	Theophylline.....	270
Stesolid.....	130	Tasmar.....	122	Thiamine hydrochloride.....	28
Stimulants / ADHD Treatments.....	145	Taurine.....	23	Thiamine multichem.....	28
Stiripentol.....	132	TCu 380 Plus Normal.....	75	Thiamine Sterop.....	28
Streptomycin sulphate.....	89	Tecentriq.....	244	Thiamine-REX.....	28
Stromectol.....	102	Tecfidera.....	141	Thioguanine.....	153
Sucralfate.....	9	TEEVIR.....	105	Thiopental [Thiopentone] sodium.....	123
Sucrose.....	126	TEEVIR (India).....	105	Thiopentone.....	123
Sugammadex.....	118	Tegretol.....	130	Thiotepa.....	151
Sugammadex BNM.....	118	Tegretol CR.....	130	Thiotepa-Reach.....	151
Sulfadiazine silver.....	67	Teicoplanin.....	97	Thrombin.....	34
Sulfadiazine sodium.....	97	Temaccord.....	158	Thyroid and Antithyroid Preparations.....	87
Sulfasalazine.....	7	Temazepam.....	143	Thyrotropin alfa.....	82
Sulindac.....	120	Temozolomide.....	158	Ticagrelor.....	38
Sulphacetamide sodium.....	274	Temozolomide (Devatis).....	158	Ticagrelor Mylan.....	38
Sulphur.....	291	Temozolomide Taro.....	158	Ticagrelor Sandoz.....	38
Sulprix.....	135	Tenecteplase.....	39	Ticagrelor Sandoz S29.....	38
Sumagran.....	133	Tenofovir disoproxil.....	108	Ticarcillin with clavulanic acid.....	94
Sumatriptan.....	133	Tenofovir Disoproxil Emtricitabine Mylan.....	110	Ticlopidine.....	39
Sunitinib.....	171	Tenofovir Disoproxil Emtricitabine Viatr.....	110	Tigecycline.....	95
Sunitinib Rex.....	171	Tenofovir Disoproxil Viatris.....	108	Tilcotil.....	120
Sunscreen, proprietary.....	72	Tenofovir Efavirenz Emtricitabine(Mylan).....	105	Tillomed.....	103
Suprane.....	122	Tenoxicam.....	120	Timolol.....	278
Surgical Preparations.....	287	Tensipine MR10.....	49	Tiotropium bromide.....	265
Sustagen Hospital Formula (Chocolate).....	308	Tepadina.....	151	Tiotropium bromide with olodaterol.....	265
Sustagen Hospital Formula (Vanilla).....	308	Terazosin.....	46	Tivicay.....	107
Suxamethonium chloride.....	118	Terbinafine.....	101	TMP.....	97
Sylvant.....	233	Terbutaline.....	76	Tobradex.....	274
Symbicort Turbuhaler.....	270	Terbutaline sulphate.....	268	Tobramycin Infections.....	89
Symmetrel.....	121	Terflunomide.....	141	Sensory.....	274
Sympathomimetics.....	54	Terflunomide Aurobindo.....	141	Tobramycin (Viatris).....	89
Synacthen.....	82	Terflunomide Sandoz.....	141	Tobramycin BNM.....	89
Synacthen Depot.....	82	Teriparatide.....	116	Tobrex.....	274
Synagis.....	217	Teriparatide - Teva.....	116	Tocilizumab.....	233
Synermox.....	93	Terlipressin.....	88	Tofranil.....	129
Syntometrine.....	75	Terlipressin Ever Pharma.....	88	Tolcapone.....	122
Syrup.....	291	Testogel.....	78	Tolvaptan.....	51
Systane Unit Dose.....	280	Testosterone.....	78	Topamax.....	132
- T -					
Tacrolimus Dermatological.....	72	Testosterone cypionate.....	78	Topamine.....	26
Oncology.....	177	Testosterone esters.....	78	Topical Products for Joint and Muscular Pain.....	120
Tacrolimus Sandoz.....	177	Testosterone undecanoate.....	78	Topiramate.....	132
Tafinlar.....	162	Tetrabenazine.....	121	Topiramate Actavis.....	132
Tagitol V.....	284	Tetracaine [Amethocaine] hydrochloride Nervous.....	125		
Tagrisso.....	167	Sensory.....	277		

Torbay	55	TYR Anamix Junior.....	298	Vasodilators.....	55
Tramadol hydrochloride.....	128	TYR Anamix Junior LQ.....	298	Vasopressin.....	88
Tramal 100	128	TYR Explore 5	298	Vasopressin Agents.....	88
Tramal 50	128	TYR Lophlex LQ 20	298	Vasorex	48
Tramal SR 100	128	TYR Sphere 20	298	Vebulis.....	64
Tramal SR 150	128	Tyrosine1000.....	299	Vecure	118
Tramal SR 200	128	Tysabri.....	141	Vecuronium bromide	118
Trametinib.....	171	- U -			
Trandate	48	UK Cipla	268	Vedafil.....	61
Tranexamic acid	34	UK Synacthen.....	82	Vedolizumab.....	242
Tranexamic-AFT	34	Ultibro Breezhaler.....	265	Vegzelma.....	201
Tranylcypromine sulphate	129	Ultraproct	7	Veklury.....	112
Trastuzumab (Herzuma).....	237	Umeclidinium	265	Veletri	62
Trastuzumab deruxtecan.....	239	Umeclidinium with vilanterol	265	Venclexta.....	159
Trastuzumab emtansine.....	239	Univent	264-265	Venetoclax.....	159
Travatan	279	Upadacitinib.....	259	Venlafaxine.....	129
Travoprost	279	Ural	76	Venofer	24
Treatments for Dementia.....	147	Urea		VENOX.....	263
Treatments for Substance		Dermatological	70	Ventolin.....	268
Dependence	148	Extemporaneously Compounded		Vepesid.....	154
Trelegy Ellipta.....	266	Preparations	291	Verapamil hydrochloride.....	49
Tretinoin		Urex Forte.....	50	Vermox.....	103
Dermatological.....	68	UroFos.....	96	Versacloz.....	135
Oncology.....	159	Urokinase	39	Vesanoid.....	159
Trexate	153	Urologicals.....	76	Vexazone.....	11
Tri-sodium citrate.....	291	Uromitexan	174	Vfend	99
Triamcinolone acetoneide		Ursodeoxycholic acid.....	14	Victoza.....	12
Alimentary.....	26	Ursosan	14	Vigabatrin	132
Dermatological.....	70	Ustekinumab.....	240	Vigisom.....	143
Hormone Preparations.....	80	Utrogestan.....	76	Vildagliptin	11
Triamcinolone acetonide with		- V -			
gramicidin, neomycin and		Vabysmo.....	204	Vildagliptin with metformin	
nystatin	275	Vaclovir.....	109	hydrochloride	11
Triamcinolone acetonide with		Valaciclovir	109	Vimpat.....	131
neomycin sulphate, gramicidin		Valaciclovir Mylan.....	109	Vinblastine sulphate	174
and nystatin	71	Valaciclovir Viatriis	109	Vincristine sulphate	174
Triamcinolone hexacetoneide.....	80	Valganciclovir	109	Vinorelbine.....	174
Triazolam.....	143	Valganciclovir Viatriis	109	Vinorelbine Te Arai.....	174
Trichloroacetic acid.....	291	Valine50.....	299	Viral Vaccines.....	315
Trientine.....	23	Valni Retard.....	49	Viramune Suspension	105
Trientine Waymade	23	Vancomycin	97	ViruPOS.....	274
Trikafta.....	271	Vancomycin Viatriis.....	97	Viscoat.....	277
Trimethoprim	97	Vanilla SilQ HD	284	Visipaque.....	284
Trimethoprim with		Vanilla SilQ MD	284	Vit.D3.....	28
sulphamethoxazole		Vanzacaftor with tezacaftor and		VitA-POS.....	280
[Co-trimoxazole]	97	deutivacaftor	272	Vital.....	300
Trisul.....	97	Varenicline.....	149	Vitamin B complex.....	28
Trometamol	287	Varbar - Honey	284	Vitamin B6 25	28
Tropicamide.....	279	Varbar - Nectar	284	Vitamins.....	26
Tropisetron	135	Varbar - Pudding.....	284	Vivonex TEN.....	300
Trulicity.....	11	Varbar - Thin Liquid	284	Voltaren	119
Tryzan.....	44	Varicella vaccine [Chickenpox		Voltaren D.....	119
Tuberculin PPD [Mantoux] test.....	320	vaccine]	319	Voltaren SR.....	119
Tubersol.....	320	Varicella zoster vaccine [Shingles		Volumatic.....	321
Two Cal HN	301	vaccine]	320	Voriconazole.....	99
TYR Anamix Infant	298	Varilrix.....	319	Vttack.....	99
				Vytorin.....	53
				Vyvanse.....	145

- W -	
Warfarin sodium	37
Wart Preparations.....	72
Water	
Blood.....	43
Various.....	287
Wool fat	
Dermatological.....	70
Extemporaneously Compounded	
Preparations	291
- X -	
Xalkori.....	162
Xanthan	291
Xarelto	37
XGen	264
Xgeva	116
Xifaxan.....	9
XMET Maxamum.....	295
Xolair	216
XP Maxamum.....	296
Xylocaine.....	124
Xylocaine Viscous	124
Xylometazoline hydrochloride.....	268
Xyntha	35
- Y -	
Yellow jacket wasp venom	264
Yervoy	245
- Z -	
Zanamivir.....	111
Zarontin	130
Zavedos.....	152
Zavicefta.....	90
Zeffix.....	107
Zejula.....	155
Zematop	72
Zetlam.....	107
Ziagen.....	105
Zidovudine	106
Zidovudine with lamivudine	106
Ziextenzo	40
Zimyebe.....	53
Zinc	
Alimentary	25
Dermatological.....	68
Zinc and castor oil	68
Zinc chloride	25
Zinc oxide	291
Zinc sulphate	25
Zinc with wool fat	69
Zincaps.....	25
Zinfoo	91
Ziprasidone.....	136
Zista.....	264
Zithromax.....	91
Zo-Rub HP.....	125
Zo-Rub Osteo.....	120
Zoladex.....	83
Zoledronic acid	
Hormone Preparations.....	79
Musculoskeletal	115
Zoledronic Acid Injection Mylan.....	79
Zoledronic acid Viatris	
Hormone Preparations.....	79
Musculoskeletal	115
Zopiclone	143
Zopiclone Actavis	143
Zostrix.....	120
Zostrix HP	125
Zuclopenthixol acetate.....	137
Zuclopenthixol decanoate.....	139
Zuclopenthixol hydrochloride.....	137
Zusdone.....	136
Zyban.....	148
Zypine.....	136
Zypine ODT	136
Zyprexa Relprevv	137
Zytiga.....	174
Zyvox.....	96

















