Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER	PATIENT:
Name:	Name:
Ward:	NHI:
Pazopanib	
INITIATION Re-assessment required after 3 months Prerequisites (tick boxes where appropriate)	
and	The patient has metastatic renal cell carcinoma
or	O The patient is treatment naive
or	O The patient has only received prior cytokine treatment
	O The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance and
	O The cancer did not progress whilst on sunitinib
and and and	The patient has good performance status (WHO/ECOG grade 0-2) The disease is of predominant clear cell histology
an	C Lactate dehydrogenase level > 1.5 times upper limit of normal
an	
an	Corrected serum calcium level > 10 mg/dL (2.5 mmol/L) d Interval of < 1 year from original diagnosis to the start of systemic therapy
an	
an	
CONTINUATION Re-assessment required after 3 months Prerequisites (tick boxes where appropriate)	
and	No evidence of disease progression
	The treatment remains appropriate and the patient is benefiting from treatment
Note: Pazopanib treatment should be stopped if disease progresses. Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.	