# SA2032 - Somatropin

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

# Somatropin

[	Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 w	
or	life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device)	
	Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using standards of Tanner and Davies (1985)	g the
	and A current bone age is < 14 years (female patients) or < 16 years (male patients) and	
	Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In chi are 5 years or older, GH testing with sex steroid priming is required	dren w
	and If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is	

Renewal — growth hormone deficiency in children			
Current approval Number (if known): Applications only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months.			
Prerequisites(tick boxes where appropriate)			
A current bone age is 14 years or under (female patients) or 16 years or under (male patients)			
Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985)			
and Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months and			
No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred and			
No malignancy has developed since starting growth hormone			
Initial application — Turner syndrome Applications only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months.			

Prerequisites(tick boxes where appropriate)

The patient has a post-natal genotype confirming Turner Syndrome
and Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985)
A current bone age is < 14 years

Enquiries	to	Ministry	of	Health
0800 855	06	6		

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:		
Reg No:	First Names:			
Name:				
Address:	DOB:	Address:		
Address:				
Fax Number:		Fax Number:		
Somatropin - continued				
Renewal — Turner syndrome				
Prerequisites(tick boxes where appropriate)	t or endocrinologist. Approvals valid for 12 months.			
<ul> <li>Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts)</li> <li>and</li> <li>Height velocity is greater than or equal to 2 cm per year, calculated over six months</li> <li>and</li> <li>A current bone age is 14 years or under</li> <li>and</li> <li>No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred</li> <li>and</li> <li>No malignancy has developed since starting growth hormone</li> </ul>				
Initial application — short stature without grow Applications only from a paediatric endocrinologis Prerequisites(tick boxes where appropriate)	vth hormone deficiency st or endocrinologist. Approvals valid for 9 months.			
The patient's height is more than 3 delay and	3 standard deviations below the mean for age or for b	one age if there is marked growth acceleration or		
Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985) and				
A current bone age is < 14 years or under (female patients) or < 16 years (male patients)				
The patient does not have severe medications known to impair heigh	chronic disease (including malignancy or recognized nt velocity	severe skeletal dysplasia) and is not receiving		
Renewal — short stature without growth horm	one deficiency			
Current approval Number (if known):				
Applications only from a paediatric endocrinologis <b>Prerequisites</b> (tick boxes where appropriate)	t or endocrinologist. Approvals valid for 12 months.			
Height velocity is greater than or e 12 months using the standards of and	equal to 50th percentile (adjusted for bone age/pubert Tanner and Davies (1985)	al status if appropriate) as calculated over 6 to		
	equal to 2 cm per year as calculated over six months			
	under (female patients) or 16 years or under (male patients)	atients)		
	patient's specialist considers is likely to be attributab	le to growth hormone treatment has occurred		

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

# Somatropin - continued

#### Initial application — short stature due to chronic renal insufficiency

Applications only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months.

Prerequisites(tick boxes where appropriate)

and		The patient's height is more than 2 standard deviations below the mean
		Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985)
and		A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients)
and		
Ĺ		The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease
and		The patient is under the supervision of a specialist with expertise in renal medicine
and		
	or	The patient has a GFR less than or equal to 30 ml/min/1.73m <sup>2</sup> as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73m <sup>2</sup> ) in a child who may or may not be receiving dialysis
	U	The patient has received a renal transplant and has received < 5mg/ m <sup>2</sup> /day of prednisone or equivalent for at least 6 months.

### Renewal — short stature due to chronic renal insufficiency

Current approval Number (if known):.....

Applications only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months.

Prerequisites(tick boxes where appropriate)

	and	Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985)
		Height velocity is greater than or equal to 2 cm per year as calculated over six months
	and	A current bone age is 14 years or under (female patients) or 16 years or under (male patients)
	and	No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred
	and	No malignancy has developed after growth hormone therapy was commenced
	and	The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results
	and	The patient has not received renal transplantation since starting growth hormone treatment
á	and	If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

# Somatropin - continued

Initial application — Prader-Willi syndrome Applications only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months. Prerequisites(tick boxes where appropriate)				
and	The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria			
and	The patient is aged six months or older			
and	A current bone age is < 14 years (female patients) or < 16 years (male patients)			
	Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon			
and				
	and			
	There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months			
	or The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation			

## Renewal — Prader-Willi syndrome

Current approval Number (if known):.....

Applications only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months. **Prerequisites**(tick boxes where appropriate)

	Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985)
and and	Height velocity is greater than or equal to 2 cm per year as calculated over six months
and	A current bone age is 14 years or under (female patients) or 16 years or under (male patients)
and	No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred
and	No malignancy has developed after growth hormone therapy was commenced
	The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

# Somatropin - continued

<b>Initial application — adults and adolescents</b> Applications only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months.
Prerequisites (tick boxes where appropriate)
The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour)
and
The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses
and
The patient has severe growth hormone deficiency (see notes)
and The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex
and
The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®)
Note: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.
Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with
isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where
an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre. The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the
mean normal value for age and sex; and
Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.
At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid
and levothyroxine.

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
<b>-</b>		

Somatropin - continued

newar	— adults and adolescents
rrent a	pproval Number (if known):
•	ns only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months. sites(tick boxes where appropriate)
	The patient has been treated with somatropin for < 12 months and There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline and Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex
	and The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients
or	The patient has been treated with somatropin for more than 12 months and The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors)
	and Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors) and
	The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients
or	The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication
	The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses and
	The patient has severe growth hormone deficiency (see notes) and The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex
	and The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire fo adult growth hormone deficiency (QoL-AGHDA®)

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre. The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients. At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.