SA2487 - Infliximab

Crohn's disease (adults) - Initial application	
Crohn's disease (adults) - Renewal	
Crohn's disease (children) - Initial application	
Crohn's disease (children) - Renewal	
Graft vs host disease - Initial application	
Pulmonary sarcoidosis - Initial application	
Acute fulminant ulcerative colitis - Initial application	
Ankylosing spondylitis - Initial application	
Ankylosing spondylitis - Renewal	
Chronic ocular inflammation - Initial application	
Chronic ocular inflammation - Renewal	
istulising Crohn's disease - Initial application	
istulising Crohn's disease - Renewal	
ulminant ulcerative colitis - Renewal	
mmune checkpoint inhibitor toxicity in malignancy* - Initial application	1
mmune checkpoint inhibitor toxicity in malignancy* - Renewal	1
nflammatory bowel arthritis – axial - Initial application	
nflammatory bowel arthritis – axial - Renewal	
nflammatory bowel arthritis - peripheral - Initial application	
nflammatory bowel arthritis – peripheral - Renewal	
leurosarcoidosis - Initial application	
leurosarcoidosis - Renewal	
laque psoriasis - Initial application	
laque psoriasis - Renewal	
revious use - Initial application	
soriatic arthritis - Initial application	
soriatic arthritis - Renewal	
voderma gangrenosum - Initial application	
yoderma gangrenosum - Renewal	
heumatoid arthritis - Initial application	
Iheumatoid arthritis - Renewal	
evere Behcet's disease - Initial application	
evere Behcet's disease - Renewal	
Severe ocular inflammation - Initial application	
Severe ocular inflammation - Renewal	
JIcerative colitis - Initial application	
JIcerative colitis - Initial application	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab

Applicati	ions f	tion — Crohn's disease (adults) rom any relevant practitioner. Approvals valid for 6 months. (tick boxes where appropriate)
an	d	Patient has active Crohn's disease
	0	Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10
	0	Patient has extensive small intestine disease affecting more than 50 cm of the small intestine
	0	Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection
		Patient has an ileostomy or colostomy, and has intestinal inflammation
an	d	Patient has tried but has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids
Renewal	I — C	rohn's disease (adults)
	•••	/al Number (if known):
••		om any relevant practitioner. Approvals valid for 2 years. (tick boxes where appropriate)
	0	CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab
		CDAI score is 150 or less, or HBI is 4 or less
	0	The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed
an	d	Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — Crohn's disease (children) Applications from any relevant practitioner. Approvals valid for 6 months.
Prerequisites(tick boxes where appropriate)
Paediatric patient has active Crohn's disease and
Patient has a PCDAI score of greater than or equal to 30
Patient has extensive small intestine disease
and Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids
Renewal — Crohn's disease (children)
Current approval Number (if known):
Applications from any relevant practitioner. Approvals valid for 2 years.
Prerequisites(tick boxes where appropriate)
PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab
PCDAI score is 15 or less
or The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed
and Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019
Initial application — Graft vs host disease Applications from any relevant practitioner. Approvals valid without further renewal unless notified. Prerequisites(tick box where appropriate)
Patient has steroid-refractory acute graft vs. host disease of the gut
Initial application — Pulmonary sarcoidosis Applications from any relevant practitioner. Approvals valid without further renewal unless notified. Prerequisites(tick box where appropriate)
Patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — acute fulminant ulcerative colitis Applications only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks. Prerequisites(tick boxes where appropriate)
Patient has acute, fulminant ulcerative colitis and Treatment with intravenous or high dose oral corticosteroids has not been successful
Initial application — ankylosing spondylitis Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months. Prerequisites(tick boxes where appropriate)
The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis and
The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept or
Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumal and/or etanercept for ankylosing spondylitis

Renewal — ankylosing spondylitis

and

and

Current approval Number	' (if	known):
-------------------------	-------	---------

Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. **Prerequisites**(tick boxes where appropriate)

> Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less

Physician considers that the patient has benefited from treatment and that continued treatment is appropriate

Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

	n any relevant practitioner. Approvals valid for 4 months. ck boxes where appropriate)
and	The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation
	The patient has experienced intolerable side effects from adalimumab
	or The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation
r	
and	Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss
	Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective or
	Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose or
	Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate
	and

Renewal — chronic ocular inflammation

Current approval Number (if known): Applications from any relevant practitioner. Approvals valid for 12 months. Prerequisites (tick boxes where appropriate)			
	or The patient has had a good clinical response following 3 initial doses		
	Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema)		
	Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old		
	Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.		

Enquiries to Ministry of	Health
0800 855 066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:
Infliximab - continued		

Initial application — fistulising Crohn's disease Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)		
	and	Patient has confirmed Crohn's disease
		Patient has one or more complex externally draining enterocutaneous fistula(e)
		Patient has one or more rectovaginal fistula(e)
		Patent has complex peri-anal fistula
Ren	ewal –	– fistulising Crohn's disease
Appl	ication	proval Number (if known): as only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 2 years. ites(tick boxes where appropriate)
		The number of open draining fistulae have decreased from baseline by at least 50%
		There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain
	and [Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019
Арр	lication	lication — neurosarcoidosis ns only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months. ites (tick boxes where appropriate)
	and	Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team
	and	Patient has CNS involvement
	and	Patient has steroid-refractory disease
		IV cyclophosphamide has been tried
		Treatment with IV cyclophosphamide is clinically inappropriate

Enquiries to Ministry of Health 0800 855 066

APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:				
Reg No:	First Names:	First Names:				
Name:	Surname:	Surname:				
Address:	DOB:	Address:				
	Address:					
Fax Number:		Fax Number:				
Infliximab - continued						
Renewal — neurosarcoidosis						
Current approval Number (if known):						
Applications only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months.						
Prerequisites(tick boxes where appropriate)						
A withdrawal period has been tried and the patient has relapsed						
A withdrawal period has been considered but would not be clinically appropriate						
and There has been a marked re	duction in prednisone dose					
or	provement in MRI appearances					
Marked improvement i	n other symptomology					

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Applic	Initial application — plaque psoriasis Applications only from a dermatologist or any relevant practitioner on the recommendation of a dermatologist. Approvals valid for 3 months. Prerequisites(tick boxes where appropriate)			
		The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis and		
			or	Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab
				Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis
	or			
			or or	 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis. Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10
		and [and		Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin
		and		A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course
		[The most recent PASI assessment is no more than 1 month old at the time of initiation
while s face, h severe	still o nand, e, and	n treat foot, g I for th	tmen genit ne fa	sponse" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably it but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the ital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very ce, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed n treatment but no longer than 1 month following cessation of the most recent prior treatment.

Enquiries	to Ministry	of Health
0800 855	066	

eg No:	mp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:	
	No: First Names: First Names:			
ame:		Surname:	Surname:	
ddress:		DOB:	Address:	
		Address:		
ax Number:			Fax Number:	
nfliximab - co	ontinued			
Renewal — plac	que psoriasis			
	and Following each prior i	vals valid for 6 months. ody" severe chronic plaque psoriasis at the start of tre nfliximab treatment course the patient has a PASI sco I, when compared with the pre-infliximab treatment ba	ore which is reduced by 75% or more, or is	
or	Patient had severe ch	nronic plaque psoriasis of the face, or palm of a hand	or sole of a foot at the start of treatment	
		prior infliximab treatment course the patient has a rea na, thickness and scaling, to slight or better, or sustain e values		
		prior infliximab treatment course the patient has a re- stained at this level, as compared to the pre-infliximab		
or	Patient had severe ch	nronic localised genital or flexural plaque psoriasis at t	he start of treatment	

Enquiries to Ministry of Health 0800 855 066

APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — previous use Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate)						
and	Patient was being treated with infliximab prior to 1 February 2019					
	Rheumatoid arthritis					
	Ankylosing spondylitis					
	Psoriatic arthritis or					
	Severe ocular inflammation					
	Chronic ocular inflammation or					
	Crohn's disease (adults) or					
	Crohn's disease (children)					
	Fistulising Crohn's disease or					
	Severe fulminant ulcerative colitis					
	Severe ulcerative colitis					
	Plaque psoriasis or					
	Neurosarcoidosis or					
	Severe Behcet's disease					
Initial appli	cation — psoriatic arthritis					
Applications only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months. Prerequisites(tick boxes where appropriate)						

and		The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis			
	or		The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab		
	01		Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis		

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

	1aD - continued
Renewa	al — psoriatic arthritis
Current	approval Number (if known):
	tions only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. uisites(tick boxes where appropriate)
	Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician
	The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician
a	nd Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks
Applica	pplication — rheumatoid arthritis ations only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months. uisites(tick boxes where appropriate)
a	The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis
	The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept or
	Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept
a	nd Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance
Renewa	al — rheumatoid arthritis
Current	approval Number (if known):
	tions only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months. uisites(tick boxes where appropriate)
a	Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance
	Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician
	The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician
a	nd Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — severe Behcet's disease Applications from any relevant practitioner. Approvals valid for 4 months.				
Prerequisites (tick boxes where appropriate)				
The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes) and				
The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes)				
The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes)				
and The patient is experiencing significant loss of quality of life				
Note: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7. Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.				
Renewal — severe Behcet's disease				
Current approval Number (if known):				
Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites (tick boxes where appropriate)				
Patient has had a good clinical response to initial treatment with measurably improved quality of life and				
Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks				
Renewal — fulminant ulcerative colitis				
Current approval Number (if known):				
Applications from any relevant practitioner. Approvals valid for 2 years. Prerequisites(tick boxes where appropriate)				
Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months and				
Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019				

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Initial application — severe ocular inflammation Applications from any relevant practitioner. Approvals valid for 4 months. Prerequisites(tick boxes where appropriate)								
		and	The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation and					
			or	The patient has experienced intolerable side effects from adalimumab				
				The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation				
	or							
		and		Patient has severe, vision-threatening ocular inflammation requiring rapid control				
				Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms				
			or	Patient developed new inflammatory symptoms while receiving high dose steroids				
			or	Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms				

Renewal — severe ocular inflammation				
Current approval Number (if known):				
Applications from any relevant practitioner. Approvals valid for 12 months. Prerequisites (tick boxes where appropriate)				
The patient has had a good clinical response following 3 initial doses				
Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema)				
or Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old				
Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.				

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Applic	cations	ation — ulcerative colitis from any relevant practitioner. Approvals valid for 6 months. s (tick boxes where appropriate)		
	Patient has active ulcerative colitis			
	c	Patients SCCAI is greater than or equal to 4		
		Patients PUCAI score is greater than or equal to 20		
	and	Patient has tried but has experienced an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids		
Renev	wal — u	ulcerative colitis		
Currer	nt appro	oval Number (if known):		
		from any relevant practitioner. Approvals valid for 2 years.		
Preree	quisite	s(tick boxes where appropriate)		
	c	The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on infliximab		
	and	Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019		
Applic	cations	ation — pyoderma gangrenosum only from a dermatologist. Approvals valid for 4 months. s (tick boxes where appropriate)		
	and	Patient has pyoderma gangrenosum*		
		Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response		
	and	A maximum of 8 doses		

Note: Note: Indications marked with * are unapproved indications.

Enquiries	to Mini	istry of	Health
0800 855	066	-	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:				
Reg No:	First Names:	First Names:				
Name:	Surname:	Surname:				
Address:	DOB:	Address:				
	Address:					
Fax Number:		Fax Number:				
Infliximab - continued						
Renewal — pyoderma gangrenosum						
Current approval Number (if known):						
	ner on the recommendation of a dermatologist. Appr	ovals valid for 4 months.				
Prerequisites(tick boxes where appropriate)						
Patient has shown clinical improve	ement					
Patient continues to require treatm	lent					
A maximum of 8 doses						
Initial application — inflammatory bowel arthri Applications from any relevant practitioner. Appro						
Prerequisites(tick boxes where appropriate)						
	lcerative colitis or active Crohn's disease					
And Patient has had axial inflammatory	pain for six months or more					
and Patient is unable to take NSAIDs and Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI and Patient's disease has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a						
				physiotherapist and		o months of an exclose regime supervised by a
					on a 0 - 10 scale completed after the 3 month exer	rcise trial, but prior to ceasing any previous
Renewal — inflammatory bowel arthritis – axial						
Current approval Number (if known):						
Applications from any relevant practitioner. Approvals valid for 2 years.						

Prerequisites(tick box where appropriate)

Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less

Enquiries	to Ministry	of Health
0800 855	066	

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:
Reg No:	First Names:	First Names:
Name:	Surname:	Surname:
Address:	DOB:	Address:
	Address:	
Fax Number:		Fax Number:

Infliximab - continued

Applications from any relevant practitioner. Approvals valid for 6 months. Prerequisites(tick boxes where appropriate) Patient has a diagnosis of active ulcerative colitis or active Crohn's disease and Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular	
and Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular	
Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular	
and	
Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated do (unless contraindicated)	e
and Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated) and	
Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application or	
Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application	
or ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and h done so for more than three months	3
Renewal — inflammatory bowel arthritis – peripheral	
Current approval Number (if known):	
Applications from any relevant practitioner. Approvals valid for 2 years. Prerequisites (tick boxes where appropriate)	
Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically signific response to treatment in the opinion of the physician	Int
or Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician	
Initial application — immune checkpoint inhibitor toxicity in malignancy*	_
Applications from any relevant practitioner. Approvals valid for 4 months. Prerequisites (tick boxes where appropriate)	
The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy	
and The individual has received insufficient benefit from use of corticosteroids	
and	

Enquiries to Ministry of Health 0800 855 066

APPLICATION FOR SUBSIDY BY SPECIAL AUTHORITY

APPLICANT (stamp or sticker acceptable)	PATIENT NHI:	REFERRER Reg No:		
Reg No:	First Names:	First Names:		
Name:	Surname:	Surname:		
Address:	DOB:	Address:		
	Address:			
Fax Number:		Fax Number:		
Infliximab - continued	Infliximab - continued			
Renewal — immune checkpoint inhibitor toxicity in malignancy*				
Current approval Number (if known):				
Applications from any relevant practitioner. Approvals valid for 4 months.				
Prerequisites(tick boxes where appropriate)				
The individual has shown clinical ir	nprovement and ongoing treatment is required			
	up to 5mg/kg for up to a total of 8 doses			

Note: Indications marked with * are unapproved indications.