Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER		PATIENT:
Name:		Name:
Ward:		NHI:
Epoetin alfa		
INITIATION – chronic rena Prerequisites (tick boxes v		
_	hronic renal failure	
and Haemoglob	oin is less than or equal to 100g/L	
and O	Patient does not have diabetes mellitus Glomerular filtration rate is less than or equal to 30	Oml/min
or O and	Patient has diabetes mellitus	
	Glomerular filtration rate is less than or equal to 4	5ml/min
and Patient is o	n haemodialysis or peritoneal dialysis	
INITIATION – myelodyspla Re-assessment required af Prerequisites (tick boxes v	fter 2 months	
O Patient has	a confirmed diagnosis of myelodysplasia (MDS)	
\sim	mptomatic anaemia with haemoglobin < 100g/L an	d is red cell transfusion-dependent
O Patient has syndrome (ne WHO classification-based prognostic scoring system for myelodysplastic
Other cause	es of anaemia such as B12 and folate deficiency h	ave been excluded
	a serum epoetin level of < 500 IU/L	
	um necessary dose of epoetin would be used and v	vill not exceed 80,000 iu per week
CONTINUATION – myeloc Re-assessment required af Prerequisites (tick boxes v	fter 12 months	
O The patient	's transfusion requirement continues to be reduced	with epoetin treatment
	tion to acute myeloid leukaemia has not occurred	
and	um necessary dose of epoetin would be used and v	vill not exceed 80,000 iu per week

I confirm that the above details are correct:

Signed: Date:

HOSPITAL MEDICINES LIST RESTRICTIONS CHECKLIST

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER	PATIENT:		
Name:	Name:		
Ward:	NHI:		
Epoetin alfa - continued			
INITIATION – all other indications Prerequisites (tick box where appropriate)			
O Prescribed by, or recommended by a haematologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.			
O For use in patients where blood transfusion is not a viable treatment Note: Indications marked with * are unapproved indications	alternative		