Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

Name:
NHI:
tioner, or in accordance with a protocol or guideline that has been
r than wet AMD
or severe posterior uveitis following treatment with bevacizumab y despite three intraocular injections of bevacizumab four weeks reated eye for longer than 3 months
ment of wAMD and was found to be intolerant to ranibizumab within ment with ranibizumab for wAMD and disease was stable while on
tioner, or in accordance with a protocol or guideline that has been re eye
t t

I confirm that the above details are correct:

Signed: Date:

HOSPITAL MEDICINES LIST RESTRICTIONS CHECKLIST

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER	PATIENT:
Name:	Name:
Ward:	NHI:
Aflibercept - continued	
INITIATION – Diabetic Macular Oedema Re-assessment required after 4 months Prerequisites (tick boxes where appropriate) O Prescribed by, or recommended by an ophthalmologist or nurse praceed endorsed by the Health NZ Hospital. And O Patient has centre involving diabetic macular oedema (DMO) and O Patient's disease is non responsive to 4 doses of intravitreal beand O Patient has reduced visual acuity between 6/9 – 6/36 with functionand O Patient has DMO within central OCT (ocular coherence tomogrand O There is no centre-involving sub-retinal fibrosis or foveal atroptions.	evacizumab when administered 4-6 weekly etional awareness of reduction in vision raphy) subfield > 350 micrometers
CONTINUATION – Diabetic Macular Oedema Re-assessment required after 12 months Prerequisites (tick boxes where appropriate)	
Prescribed by, or recommended by an ophthalmologist or nurse praceendorsed by the Health NZ Hospital.	etitioner, or in accordance with a protocol or guideline that has been
and Patient's vision is 6/36 or better on the Snellen visual acuity so and There is no centre-involving sub-retinal fibrosis or foveal atrop and	