Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER		PATIENT:
Name:		Name:
Ward:		NHI:
Pneumococc	cal (PCV13) conjugate vaccine	
Re-assessment Prerequisites (t	Primary course for previously unvaccinated children aged un t required after 3 doses (tick box where appropriate) nary course of three doses for previously unvaccinated children	
INITIATION – Hi Re-assessment Prerequisites (t	High risk individuals who have received PCV10 t required after 2 doses (tick box where appropriate)	onths and under 18 years) who have previously received two doses of the
Re-assessment Prerequisites (t	High risk children aged under 5 years t required after 4 doses (tick boxes where appropriate)	
and or	O Primary immune deficiencies O HIV infection O Renal failure, or nephrotic syndrome O Are immune-suppressed following organ transplantation O Cochlear implants or intracranial shunts O Cerebrospinal fluid leaks	cinate when there is expected to be a sufficient immune response (including haematopoietic stem cell transplant) s, and who are on an equivalent daily dosage of prednisone of 2 mg/kg g on a total daily dosage of 20 mg or greater

I confirm that the above details are correct:		
Cianadi	Deter	

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER	PATIENT:			
Name:	Name:			
Ward:	NHI:			
Pneumococcal (PCV13) conjugate vaccine - continued				
INITIATION – High risk individuals 5 years and over Re-assessment required after 4 doses Prerequisites (tick box where appropriate) Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency				
INITIATION – Testing for primary immunodeficiency diseases Prerequisites (tick box where appropriate) O For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician				
Note: Please refer to the Immunisation Handbook for the appropriate schedu	lle for catch up programmes			