HOSPITAL MEDICINES LIST RESTRICTIONS CHECKLIST

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER	PATIENT:
Name:	Name:
Ward:	NHI:
Bendamustine hydro	chloride
Bendamustine hydro INITIATION – CLL* Prerequisites (tick boxes of and O Patient has and O Bendamust) Note: Indication marked wit (SLL). INITIATION – Indolent, Lo Re-assessment required at Prerequisites (tick boxes of O The patient and O The patient of O The pati	chloride where appropriate) has chronic lymphocytic leukaemia requiring treatment ECOG performance status 0-2 ine is to be administered at a maximum dose of 100 mg/m ² on days 1 and 2 every 4 weeks for a maximum of 6 cycles th a * includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma w-grade lymphomas ter 9 months
or and or and and and or and or Bend	Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles The patient has not received prior bendamustine therapy Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+) Patient has had a rituximab treatment-free interval of 12 months or more amustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients

Signed: Date:

HOSPITAL MEDICINES LIST RESTRICTIONS CHECKLIST

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIE	BER		PATIENT:	
Name:			Name:	
Ward:			NHI:	
Bendam	ustin	e hy	drochloride - continued	
			lolent, Low-grade lymphomas ed after 9 months	
Prerequisites (tick boxes where appropriate)				
	O Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine and O Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles			
or	(and	O	Patients have not received a bendamustine regimen within the last 12 months O Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+) and O Patient has had a rituximab treatment-free interval of 12 months or more	
			O Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients	

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenström's macroglobulinaemia.

INITIATION – Hodgkin's lymphoma*

Re-assessment required after 6 months **Prerequisites** (tick boxes where appropriate)

Prerequisites (tick boxes where appropriate) Patient has Hodgkin's lymphoma requiring treatment and Patient has a ECOG performance status of 0-2 and Patient has received one prior line of chemotherapy and Patient's disease relapsed or was refractory following prior chemotherapy and Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles Note: Indications marked with * are unapproved indications.