#### HOSPITAL MEDICINES LIST RESTRICTIONS CHECKLIST

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER	PATIENT:
Name:	Name:
Ward:	NHI:

### **Epoetin beta**

0	Patient in	chronic renal failure
and O and	Haemoglo	bin is less than or equal to 100g/L
	O	Patient does not have diabetes mellitus
	and	Glomerular filtration rate is less than or equal to 30ml/min
o	r	
	and	Patient has diabetes mellitus
		Glomerular filtration rate is less than or equal to 45ml/min

#### INITIATION – myelodysplasia\*

Re-assessment	t required	after 1	12 months	;

Prerequisites (tick boxes where appropriate)

and	Patient has a confirmed diagnosis of myelodysplasia (MDS)
and	Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent
	Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS)
and	Other causes of anaemia such as B12 and folate deficiency have been excluded
and O and	Patient has a serum epoetin level of < 500 IU/L
0	The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week

## CONTINUATION – myelodysplasia\*

Re-assessment required after 2 months
Prerequisites (tick boxes where appropriate)
O
The patient's transfusion requirement continues to be reduced with epoetin treatment
and
O
Transformation to acute myeloid leukaemia has not occurred
and
O
The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week

# HOSPITAL MEDICINES LIST RESTRICTIONS CHECKLIST

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

Name: Name:				
Ward: NHI:				
Epoetin beta - continued				
INITIATION – all other indications Prerequisites (tick boxes where appropriate)				
Haematologist				
O For use in patients where blood transfusion is not a viable treatment alternative				
O *Note: Indications marked with * are unapproved indications				

I confirm that the above details are correct: