## HOSPITAL MEDICINES LIST RESTRICTIONS CHECKLIST

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER	PATIENT:
Name:	Name:
Ward:	NHI:

## Nintedanib

Re-as	ssessi	men	diopathic pulmonary fibrosis t required after 12 months (tick boxes where appropriate)	
( and			cribed by, or recommended by a respiratory specialist, or in accordance with a protocol or guideline that has been endorsed by the Health ospital.	
	( and	С	Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist	
	( and	C	Forced vital capacity is between 50% and 90% predicted	
	and		Nintedanib is to be discontinued at disease progression (See Note)	
	and		Nintedanib is not to be used in combination with subsidised pirfenidone	
		or	<ul> <li>O The patient has not previously received treatment with pirfenidone</li> <li>O Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance</li> </ul>	
		or	<ul> <li>Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone)</li> </ul>	
		_		
CONTINUATION – idiopathic pulmonary fibrosis Re-assessment required after 12 months Prerequisites (tick boxes where appropriate)				
and			bribed by, or recommended by a respiratory specialist, or in accordance with a protocol or guideline that has been endorsed by the Health ospital.	
	( and	С	Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment	
	(and	C	Nintedanib is not to be used in combination with subsidised pirfenidone	
	(	$\mathcal{C}$	Nintedanib is to be discontinued at disease progression (See Note)	

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.