Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER PA	ATIENT:		
Name: N	ame:		
Ward: N	H:		
COVID-19 vaccine			
INITIATION – initial dose			
Prerequisites (tick boxes where appropriate)			
One dose for previously unvaccinated people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to three doses for immunocompromised people aged 12-15 years of Up to Up to three doses for immunocompromised people aged 12-15 years of Up to Up t			
or O Up to two doses for previously unvaccinated people 16-29 years old or			
O Up to four doses for people aged 16-29 at high risk of severe illness. One dose for previously unvaccinated people aged 30 and older.	ss		
INITIATION – additional dose Prerequisites (tick box where appropriate) One additional dose every 6 months for people aged 30 years and over, additional dose is given at least 6 months after last dose			
CONTINUATION – additional dose Prerequisites (tick box where appropriate) One additional dose every 6 months for people aged 30 years and over	additional dose is given at least 6 months after last dose		
	,		

I confirm that the above details are correct:

0:	D - 1 - 1	
Zigneg.	i jate:	
Oigilica.	 Duic.	