Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER	PATIENT:
Name:	Name:
Ward:	NHI:
Dabrafenib	
NZ Hospital. The individual is currently on treator The individual has restor The individual has restor Adjuvant treator and The individual has not received and Treatment must be adjuvant and Treatment must be initiated note b) and The individual has a confirm and	relevant practitioner, or in accordance with a protocol or guideline that has been endorsed by the Health trent with dabrafenib and trametinib and met all remaining criteria prior to commencing treatment sected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a) nas received neoadjuvant treatment with a PD-1/PD-L1 inhibitor ment with dabrafenib is required ived prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma to complete surgical resection within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see need BRAF mutation stered in combination with trametinib
Note:	
a) Stage IIIB, IIIC, IIID or IV melanoma defined a	as per American Joint Committee on Cancer (AJCC) 8th Edition
b) Initiating treatment within 13 weeks of comple	te surgical resection means 13 weeks after resection (primary or lymphadenectomy)

C:	D-1	
Signed.	Date:	
Oigilica.	 Daic.	

HOSPITAL MEDICINES LIST RESTRICTIONS CHECKLIST

August 2025

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRI	BER	PATIENT:
Name:		Name:
Ward:		. NHI:
Dabrafei	nib - continued	
	JATION – stage III or IV resected melanoma - adjuvant sment required after 4 months	
	sites (tick boxes where appropriate)	
	Prescribed by, or recommended by any relevant practitioner, or in NZ Hospital.	accordance with a protocol or guideline that has been endorsed by the Health
	No evidence of disease recurrence and Dabrafenib must be administered in combination with and Treatment to be discontinued at signs of disease recu any systemic neoadjuvant treatment	trametinib rrence or at completion of 12 months' total treatment course, including
The individual has received adjuvant treatment with a BRAF/MEK inhibitor and The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV and The individual meets initiation criteria for dabrafenib for unresectable or metastatic melanoma		ma (excluding uveal) stage III or IV
	The individual has received adjuvant treatment with a and The individual has received a BRAF/MEK inhibitor for and The individual meets continuation criteria for dabrafen	unresectable or metastatic melanoma

HOSPITAL MEDICINES LIST RESTRICTIONS CHECKLIST

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

	,g,g,	
PRESCRIBER		PATIENT:
Name:		Name:
Ward:		NHI:
Dabrafenib	- continued	
INITIATION – u Re-assessmen Prerequisites	unresectable or metastatic melanoma nt required after 4 months (tick boxes where appropriate) cribed by, or recommended by any relevant practitioner, or in acclospital. The individual is currently on treatment with dabrafenib and tra The individual has metastatic or unresectable melanoma Baseline measurement of overall tumour burden is docured. The individual has ECOG performance score 0-2 The individual has confirmed BRAF mutation Dabrafenib must be administered in combination with trained. The individual has been diagnosed in the metastate or The individual did not receive treatment in the adjuinant and The individual received treatment in the adjuinant and The individual did not experience disease received and	mented clinically and radiologically metinib tic or unresectable stage III or IV setting evant setting with a BRAF/MEK inhibitor
Re-assessmen Prerequisites Presc	ON – unresectable or metastatic melanoma at required after 4 months (tick boxes where appropriate) cribed by, or recommended by any relevant practitioner, or in accessive to the complete response	
and	O The individual has stable disease with treatment Response to treatment in target lesions has been determined by treatment period	by comparable radiologic assessment following the most recent