

September 2025

Volume 13

Editor:

Kaye Wilson, Doris Chong, Sophie Molloy
email: enquiry@pharmac.govt.nz
Telephone +64 4 460 4990
Level 9, 40 Mercer Street
PO Box 10 254 Wellington 6143

Freephone Information Line

0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule, Section H for Hospital Pharmaceuticals (link to PDF copy) emailed to your nominated email address each month by subscribing at schedule.pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX.
XML version of the Schedule available from schedule.pharmac.govt.nz/pub/HML

Programmers

Anrik Drenth

email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency



ISSN 1179-3708 pdf

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

Part I

General Rules

4

Part II

Alimentary Tract and Metabolism

5

Blood and Blood Forming Organs

30

Cardiovascular System

44

Dermatologicals

67

Genito-Urinary System

73

Hormone Preparations

77

Infections

88

Musculoskeletal System

113

Nervous System

120

Oncology Agents and Immunosuppressants

148

Respiratory System and Allergies

264

Sensory Organs

274

Various

281

Extemporaneous Compounds (ECPs)

289

Special Foods

292

Vaccines

310

Part III

Optional Pharmaceuticals

321

Index

322

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at <https://pharmac.govt.nz/about>.

Glossary

Units of Measure

gram	g	microgram.....	mcg	millimole.....	mmol
kilogram.....	kg	milligram.....	mg	unit.....	u
international unit.....	iu	millilitre.....	ml		

Abbreviations

application	app	enteric coated.....	EC	solution.....	soln
capsule	cap	granules.....	grans	suppository.....	suppos
cream.....	crm	injection	inj	tablet.....	tab
dispersible	disp	liquid	liq	tincture.....	tinc
effervescent.....	eff	lotion	lotn		
emulsion	emul	ointment.....	oint		

HSS Hospital Supply Status

Guide to Section H listings

Example

ANATOMICAL HEADING			
	Price (ex man. Excl. GST) \$	Per	Brand or Generic Manufacturer
THERAPEUTIC HEADING			
Generic name listed by therapeutic group and subgroup	CHEMICAL A - Restricted see terms below ⚡ Presentation A.....10.00	100	Brand A
	➡ Restricted Only for use in children under 12 years of age		Brand or manufacturer's name
Indicates only presentation B1 is Restricted	CHEMICAL B - Some items restricted see terms below ⚡ Presentation B1.....1,589,00 Presentation B2 ➡ Restricted Oncologist or haematologist	1	Brand B1 e.g. Brand B2
From 1 January 2012 to 30 June 2014, at least 99% of the total volume of this item purchased must be Brand C	CHEMICAL C Presentation C - -1% DV Limit Jan-12 to 201415.00	28	Brand C
	CHEMICAL D - Restricted see terms below ⚡ Presentation D - -1% DV Limit Mar-13 to 201438.65	500	Brand D
Standard national price excluding GST	➡ Restricted <i>Limited to five weeks' treatment</i> Either: 1 For the prophylaxis of venous thromboembolism following a total hip replacement; or 2 For the prophylaxis of venous thromboembolism following a total knee replacement.		Quantity the Price applies to
Form and strength	CHEMICAL E Presentation E		e.g. Brand E
			Not a contracted product
⚡ Item restricted (see above); ⚡ Item restricted (see below) Products with Hospital Supply Status (HSS) are in bold			

PART I: GENERAL RULES

General Rules for Section H of the Pharmaceutical Schedule are included in Section A.

Read the [General Rules](https://pharmac.govt.nz/section-a) : <https://pharmac.govt.nz/section-a>.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antacids and Antiflatulents			
Antacids and Reflux Barrier Agents			
ALUMINIUM HYDROXIDE WITH MAGNESIUM HYDROXIDE AND SIMETICONE			
Tab 200 mg with magnesium hydroxide 200 mg and simeticone 20 mg			<i>e.g. Mylanta</i>
Oral liq 400 mg with magnesium hydroxide 400 mg and simeticone 30 mg per 5 ml			<i>e.g. Mylanta Double Strength</i>
SIMETICONE			
Oral drops 100 mg per ml			
Oral drops 20 mg per 0.3 ml			
Oral drops 40 mg per ml			
SODIUM ALGINATE WITH MAGNESIUM ALGINATE			
Powder for oral soln 225 mg with magnesium alginate 87.5 mg, sachet			<i>e.g. Gaviscon Infant</i>
SODIUM ALGINATE WITH SODIUM BICARBONATE AND CALCIUM CARBONATE			
Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg			<i>e.g. Gaviscon Extra Strength</i>
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml.....	7.50	500 ml	Acidex
SODIUM CITRATE			
Oral liq 8.8% (300 mmol/l).....	25.00	90 ml	Biomed
Phosphate Binding Agents			
ALUMINIUM HYDROXIDE			
Tab 600 mg			
CALCIUM CARBONATE – Restricted see terms below			
↓ Oral liq 250 mg per ml (100 mg elemental per ml).....	47.30	473 ml	Calcium carbonate PAI
	39.00	500 ml	Roxane
➔ Restricted (RS1698)			
Initiation			
Only when prescribed for patients unable to swallow calcium carbonate tablets or where calcium carbonate tablets are inappropriate..			
Antidiarrhoeals and Intestinal Anti-Inflammatory Agents			
Antipropulsives			
DIPHENOXYLATE HYDROCHLORIDE WITH ATROPINE SULPHATE			
Tab 2.5 mg with atropine sulphate 25 mcg			
LOPERAMIDE HYDROCHLORIDE			
Tab 2 mg.....	10.75	400	Nodia
Cap 2 mg – 5% DV Feb-26 to 2028	12.00	400	Diamide Relief
Rectal and Colonic Anti-Inflammatories			
BUDESONIDE – Restricted see terms on the next page			
↓ Cap modified-release 3 mg – 5% DV Dec-25 to 2028	33.38	90	Budesonide Te Arai

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS1723)

Initiation – Crohn's disease

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or
 - 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initiation – Collagenous and lymphocytic colitis (microscopic colitis)

Patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initiation – Gut Graft versus Host disease

Patient has gut Graft versus Host disease following allogenic bone marrow transplantation.

Initiation – non-cirrhotic autoimmune hepatitis

Re-assessment required after 6 months

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indications marked with * are unapproved indications.

Continuation – non-cirrhotic autoimmune hepatitis

Re-assessment required after 6 months

Treatment remains appropriate and the patient is benefitting from the treatment.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC free (14 applications)	57.09	15 g	Colifoam
---	-------	------	----------

HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE

Topical Aerosol foam, 1% with pramoxine hydrochloride 1%

MESALAZINE

Tab EC 400 mg	49.50	100	Asacol
Tab long-acting 500 mg	56.10	100	Pentasa
Tab 800 mg	85.50	90	Asacol
Tab 1,600 mg	85.50	60	Asacol
Modified release granules 1 g	118.10	100 g	Pentasa
Suppos 500 mg	22.80	20	Asacol
Suppos 1 g	50.96	28	Pentasa
Enema 1 g per 100 ml	41.30	7	Pentasa

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
OLSALAZINE			
Tab 500 mg	93.37	100	Dipentum
Cap 250 mg	53.00	100	Dipentum
SODIUM CROMOGLICATE			
Cap 100 mg			
SULFASALAZINE			
Tab 500 mg	19.49	100	Salazopyrin
Tab EC 500 mg	20.54	100	Salazopyrin EN

Local Preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

CINCHOCAINE HYDROCHLORIDE WITH HYDROCORTISONE			
Oint 5 mg with hydrocortisone 5 mg per g	15.00	30 g	Proctosedyl
Suppos 5 mg with hydrocortisone 5 mg per g	9.90	12	Proctosedyl
FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE			
Oint 950 mcg with fluocortolone pivalate 920 mcg and cinchocaine hydrochloride 5 mg per g	13.05	30 g	Ultraproct
Suppos 630 mcg with fluocortolone pivalate 610 mcg and cinchocaine hydrochloride 1 mg	8.61	12	Ultraproct

Management of Anal Fissures

GLYCERYL TRINITRATE			
Oint 0.2%	22.00	30 g	Rectogesic

Rectal Sclerosants

OILY PHENOL [PHENOL OILY]			
Inj 5%, 5 ml vial			

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule	19.00	5	Robinul
HYOSCINE BUTYLBROMIDE			
Tab 10 mg – 5% DV Apr-25 to 2027	2.25	20	Hyoscine Butylbromide (Adiramédica)
Inj 20 mg, 1 ml ampoule – 5% DV Dec-23 to 2026	1.91	1	Spazmol
MEBEVERINE HYDROCHLORIDE			
Tab 135 mg – 5% DV Dec-23 to 2026	8.50	90	Colofac

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL			
Tab 200 mcg	47.73	120	Cytotec

Price
(ex man. excl. GST)
\$ Per Brand or
Generic
Manufacturer

H2 Antagonists

CIMETIDINE

Tab 200 mg
Tab 400 mg

FAMOTIDINE

Tab 20 mg
Tab 40 mg 10.27 100 Famotidine Hovid MY
Inj 10 mg per ml, 2 ml vial
Inj 10 mg per ml, 4 ml vial

RANITIDINE – **Restricted** see terms [below](#)

⚡ Tab 150 mg
⚡ Tab 300 mg
⚡ Inj 25 mg per ml, 2 ml ampoule

➡ **Restricted** ([RS1703](#))

Initiation

Either:

- 1 For continuation use; or
- 2 Routine prevention of allergic reactions..

Proton Pump Inhibitors

LANSOPRAZOLE

Cap 15 mg – **5% DV Feb-25 to 2027** 4.04 100 **Lanzol Relief**
Cap 30 mg – **5% DV Feb-25 to 2027** 5.43 100 **Lanzol Relief**

OMEPRAZOLE

⚡ Tab dispersible 10 mg

➡ **Restricted** ([RS1027](#))

Initiation

Only for use in tube-fed patients.

⚡ Tab dispersible 20 mg

➡ **Restricted** ([RS1027](#))

Initiation

Only for use in tube-fed patients.

Cap 10 mg – **5% DV Mar-24 to 2026** 2.06 90 Omeprazole Teva
Omeprazole actavis 10
Cap 20 mg – **5% DV Mar-24 to 2026** 2.02 90 Omeprazole Teva
Omeprazole actavis 20
Cap 40 mg – **5% DV Mar-24 to 2026** 3.18 90 Omeprazole Teva
Omeprazole actavis 40
Powder for oral liq..... 42.50 5 g Midwest
Inj 40 mg ampoule with diluent..... 37.38 5 Dr Reddy's Omeprazole
Inj 40 mg vial 11.95 5 Omezol IV

PANTOPRAZOLE

Tab EC 20 mg 1.99 90 Panzop Relief
Tab EC 40 mg 2.74 90 Panzop Relief
Inj 40 mg vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Site Protective Agents

COLLOIDAL BISMUTH SUBCITRATE

Tab 120 mg 14.51 50 Gastrodenol

SUCRALFATE

Tab 1 g

Bile and Liver Therapy

L-ORNITHINE L-ASPARTATE – **Restricted** see terms [below](#)

↓ Grans for oral liquid 3 g

→ **Restricted (RS1261)**

Initiation

For patients with chronic hepatic encephalopathy who have not responded to treatment with, or are intolerant to lactulose, or where lactulose is contraindicated.

RIFAXIMIN – **Restricted** see terms [below](#)

↓ Tab 550 mg – 5% DV **Dec-24 to 2027** 625.00 56 **Xifaxan**

→ **Restricted (RS1416)**

Initiation

For patients with hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Diabetes

Alpha Glucosidase Inhibitors

ACARBOSE

Tab 50 mg – 5% DV **Feb-25 to 2027** 11.20 90 **Accarb**

Tab 100 mg – 5% DV **Feb-25 to 2027** 17.38 90 **Accarb**

Hyperglycaemic Agents

DIAZOXIDE – **Restricted** see terms [below](#)

↓ Cap 25 mg 110.00 100 Proglidem

↓ Cap 100 mg 280.00 100 Proglidem

↓ Oral liq 50 mg per ml 620.00 30 ml Proglycem

→ **Restricted (RS1028)**

Initiation

For patients with confirmed hypoglycaemia caused by hyperinsulinism.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit 32.00 1 Glucagen Hypokit

GLUCOSE [DEXTROSE]

Tab 1.5 g

Tab 3.1 g

Tab 4 g

Oral soln 15 g per 80 ml sachet 70.00 50 HypoPak Glucose

Gel 40%

GLUCOSE WITH SUCROSE AND FRUCTOSE

Gel 19.7% with sucrose 35% and fructose 19.7%, 18 g sachet

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Insulin - Intermediate-Acting Preparations			
INSULIN ASPART WITH INSULIN ASPART PROTAMINE			
Inj insulin aspart 30% with insulin aspart protamine 70%, 100 u per ml, 3 ml prefilled pen	52.15	5	NovoMix 30 FlexPen
INSULIN DEGLUDEC WITH INSULIN ASPART			
Inj degludec 70 u with insulin aspart 30 u, 100 u per ml, 3 ml.....	80.00	5	Ryzodeg 70/30 Penfill
INSULIN ISOPHANE			
Inj insulin human 100 u per ml, 10 ml vial			
Inj insulin human 100 u per ml, 3 ml cartridge			
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj insulin lispro 25% with insulin lispro protamine 75%, 100 u per ml, 3 ml cartridge.....	42.66	5	Humalog Mix 25
Inj insulin lispro 50% with insulin lispro protamine 50%, 100 u per ml, 3 ml cartridge.....	42.66	5	Humalog Mix 50
INSULIN NEUTRAL WITH INSULIN ISOPHANE			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 10 ml vial			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 40% with insulin isophane 60%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 50% with insulin isophane 50%, 100 u per ml, 3 ml cartridge			
Insulin - Long-Acting Preparations			
INSULIN GLARGINE			
Inj 100 u per ml, 3 ml disposable pen.....	94.50	5	Lantus SoloStar
Inj 100 u per ml, 3 ml cartridge.....	94.50	5	Lantus
Inj 100 u per ml, 10 ml vial.....	63.00	1	Lantus
Insulin - Rapid-Acting Preparations			
INSULIN ASPART			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
Inj 100 u per ml, 3 ml syringe	51.19	5	NovoRapid FlexPen
INSULIN GLULISINE			
Inj 100 u per ml, 10 ml vial.....	27.03	1	Apidra
Inj 100 u per ml, 3 ml cartridge	46.07	5	Apidra
Inj 100 u per ml, 3 ml disposable pen.....	46.07	5	Apidra Solostar
INSULIN LISPRO			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
Insulin - Short-Acting Preparations			
INSULIN NEUTRAL			
Inj human 100 u per ml, 10 ml vial			
Inj human 100 u per ml, 3 ml cartridge			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
Tab 5 mg	7.50	100	Daonil
GLICLAZIDE			
Tab 80 mg – 5% DV Feb-24 to 2026	20.10	500	Glizide
GLIPIZIDE			
Tab 5 mg – 5% DV Mar-25 to 2027	6.86	100	Minidiab
METFORMIN HYDROCHLORIDE			
Tab immediate-release 500 mg – 1% DV Mar-23 to 2027	14.74	1,000	Metformin Viatris
Tab immediate-release 850 mg – 1% DV Aug-23 to 2027	11.28	500	Metformin Viatris
PIOGLITAZONE			
Tab 15 mg – 5% DV Dec-24 to 2027	6.15	90	Vexazone
Tab 30 mg – 5% DV Dec-24 to 2027	7.25	90	Vexazone
Tab 45 mg – 5% DV Dec-24 to 2027	12.00	90	Vexazone
VILDAGLIPTIN			
Tab 50 mg	35.00	60	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE			
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	Galvumet

GLP-1 Agonists

DULAGLUTIDE – **Restricted** see terms [below](#)

Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.

↓ Inj 1.5 mg per 0.5 ml prefilled pen 115.23 4 Trulicity

→ **Restricted (RS2135)**

Initiation

Either:

- 1 For continuation use; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and
 - 2.3 Any of the following:
 - 2.3.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack,

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin / empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

LIRAGLUTIDE – **Restricted** see terms [below](#)

Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.

⚡ Inj 6 mg per ml, 3 ml prefilled pen383.72 3 Victoza

➡ **Restricted** (RS2136)

Initiation

Either:

- 1 For continuation use; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and
 - 2.3 Any of the following:
 - 2.3.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause identified.
- c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin / empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

SGLT2 Inhibitors

➡ **Restricted** (RS2069)

Initiation – heart failure reduced ejection fraction

All of the following:

- 1 Patient has heart failure; and
- 2 Patient is in NYHA functional class II or III or IV; and

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

3 Either:

- 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or
- 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initiation – Type 2 Diabetes

Any of the following:

- 1 For continuation use; or
- 2 Patient has previously had an initial approval for a GLP-1 agonist; or
- 3 All of the following:
 - 3.1 Patient has type 2 diabetes; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 3.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 3.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 3.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 3.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 3.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause.
- c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride) for the treatment of heart failure.

EMPAGLIFLOZIN – **Restricted** see terms [on the previous page](#)

† Tab 10 mg	58.56	30	Jardiance
† Tab 25 mg	58.56	30	Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – **Restricted** see terms [on the previous page](#)

† Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
† Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
† Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
† Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Digestives Including Enzymes

PANCREATIC ENZYME

Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph Eur U)	34.93	20 g	Creon Micro
Powder pancreatin 60.12 mg (3,600 Ph. Eur. u/amylase, 5,000 Ph. Eur. u/lipase and 200 Ph. Eur. u/protease)			

URSODEOXYCHOLIC ACID – Restricted see terms [below](#)

⚡ Cap 250 mg – 5% DV Feb-24 to 2026	33.95	100	Ursosan
---	-------	-----	---------

➡ Restricted (RS2103)

Initiation – Alagille syndrome or progressive familial intrahepatic cholestasis

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initiation – Chronic severe drug induced cholestatic liver injury

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initiation – Primary biliary cholangitis

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initiation – Pregnancy

Patient diagnosed with cholestasis of pregnancy.

Initiation – Haematological transplant

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogeneic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initiation – Total parenteral nutrition induced cholestasis

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by TPN; and
- 2 Liver function has not improved with modifying the TPN composition.

Initiation – prevention of sinusoidal obstruction syndrome

The individual has leukaemia/lymphoma and requires prophylaxis for medications/therapies with a high risk of sinusoidal obstruction syndrome.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Laxatives			
Bowel-Cleansing Preparations			
CITRIC ACID WITH MAGNESIUM CARBONATE HYDRATE AND SODIUM PICOSULFATE			
Powder for oral soln 12 g with magnesium carbonate hydrate 7.4 g and sodium picosulfate 10 mg per sachet			<i>e.g. PicoPrep Orange</i>
MACROGOL 3350 WITH ASCORBIC ACID, POTASSIUM CHLORIDE, SODIUM CHLORIDE AND CITRIC ACID WITH MAGNESIUM CARBONATE HYDRATE AND SODIUM PICOSULFATE			
Powder for oral soln 52.9 g with ascorbic acid 6 g, potassium chloride 740 mg, sodium chloride 2.6 g and sodium sulphate 5.6 g per sachet (1) and powder for oral soln citric acid 12 g with magnesium carbonate hydrate 7.4 g and sodium picosulfate 10 mg per sachet (2)			<i>e.g. Prepkit Orange</i>
MACROGOL 3350 WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE			
Powder for oral soln 755.68 mg with potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 70 g sachet – 5% DV Feb-25 to 2027	16.10	3	Glycoprep Orange
	64.32	12	Glycoprep Orange
Powder for oral soln 755.68 mg with potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 210 g sachet			<i>e.g. Glycoprep Orange</i>
MACROGOL 3350 WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE WITH/WITHOUT SODIUM SULFATE, SODIUM ASCORBATE, ASCORBIC ACID			
Powd for oral soln 100g with potassium chloride 1g, sodium chloride 2g and sodium sulfate 9g per sach(1), powd for oral soln 40g with potassium chloride 1.2g and sodium chloride 3.2g per sach(1) and powd for oral soln ascorbic acid 7.54g and sodium ascorbate 48.11g per sach(1) – 5% DV Oct-23 to 2026	18.52	3	Plenvu
Bulk-Forming Agents			
ISPAGHULA (PSYLLIUM) HUSK			
Powder for oral soln	22.10	500 g	Konsyl-D
STERCULIA WITH FRANGULA – Restricted: For continuation only			
➡ Powder for oral soln			
Faecal Softeners			
DOCUSATE SODIUM			
Tab 50 mg – 5% DV Feb-24 to 2026	3.20	100	Coloxyl
Tab 120 mg – 5% DV Feb-24 to 2026	4.98	100	Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES			
Tab 50 mg with sennosides 8 mg	3.50	200	Laxsol
PARAFFIN			
Oral liquid 1 mg per ml			
Enema 133 ml			
POLOXAMER			
Oral drops 10% – 5% DV Feb-24 to 2026	4.17	30 ml	Coloxyl

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Opioid Receptor Antagonists - Peripheral			
METHYLNALTREXONE BROMIDE – Restricted see terms below			
⚡ Inj 12 mg per 0.6 ml vial	36.00	1	Relistor
	246.00	7	Relistor
➔ Restricted (RS2057)			
Initiation – Opioid induced constipation			
Both:			
1 The patient is receiving palliative care; and			
2 Either:			
2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or			
2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.			
Initiation – Opioid induced constipation outside of palliative care			
<i>Limited to 14 days treatment</i>			
All of the following:			
1 Individual has opioid induced constipation; and			
2 Oral and rectal treatments for opioid induced constipation, including bowel-cleansing preparations, are ineffective or inappropriate; and			
3 Mechanical bowel obstruction has been excluded.			
Osmotic Laxatives			
GLYCEROL			
Suppos 2.8/4.0 g – 5% DV Feb-26 to 2028	12.39	20	Lax-suppositories Glycerol
Note: DV limit applies to glycerol suppository presentations.			
LACTULOSE			
Oral liq 10 g per 15 ml – 5% DV Feb-26 to 2028	6.16	500 ml	Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE			
Powder for oral soln 6.563 g with potassium chloride 23.3 mg, sodium bicarbonate 89.3 mg and sodium chloride 175.4 mg			
Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg.....	8.50	30	APO Health Macrogol Molaxole Movicol
	10.15		
	12.19		
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE			
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml – 5% DV Feb-26 to 2028	36.89	50	Micolette
SODIUM PHOSPHATE WITH PHOSPHORIC ACID			
Oral liq 16.4% with phosphoric acid 25.14%			
Enema 10% with phosphoric acid 6.58%	3.70	1	Fleet Phosphate Enema
Stimulant Laxatives			
BISACODYL			
Tab 5 mg	5.80	200	Bisacodyl Viatris
Suppos 10 mg – 5% DV Feb-25 to 2027	4.14	10	Lax-Suppositories
SENNOSIDES			
Tab 7.5 mg			
SODIUM PICOSULFATE – Restricted see terms on the next page			
⚡ Oral soln 7.5 mg per ml	7.40	30 ml	Dulcolax SP Drop

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

➔ **Restricted (RS1843)**

Initiation

Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation.

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA – **Restricted** see terms [below](#)

↓ Inj 50 mg vial 1,142.60 1 Myozyme

➔ **Restricted (RS1793)**

Initiation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Continuation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

ARGININE

- Tab 1,000 mg
- Cap 500 mg
- Powder
- Inj 500 mg per ml, 10 ml vial
- Inj 600 mg per ml, 25 ml vial

BETAININE – **Restricted** see terms [below](#)

↓ Powder for oral soln.....575.00 180 g Cystadane

→ **Restricted** (RS1794)

Initiation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Continuation

Metabolic physician

Re-assessment required after 12 months

The treatment remains appropriate and the patient is benefiting from treatment.

BIOTIN – **Restricted** see terms [below](#)

- ↓ Cap 50 mg
- ↓ Cap 100 mg
- ↓ Inj 10 mg per ml, 5 ml vial

→ **Restricted** (RS1330)

Metabolic physician or metabolic disorders dietitian

CARGLUMIC ACID – **Restricted** see terms [below](#)

- ↓ Tab disp 200 mg

→ **Restricted** (RS1831)

Initiation

Metabolic physician

For the acute in-patient treatment of organic acidaemias as an alternative to haemofiltration.

COENZYME Q10 – **Restricted** see terms [below](#)

- ↓ Cap 120 mg
- ↓ Cap 160 mg

→ **Restricted** (RS1832)

Initiation

Metabolic physician

Re-assessment required after 6 months

The patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Continuation

Metabolic physician

Re-assessment required after 24 months

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GALSULFASE – Restricted see terms below			
⚡ Inj 1 mg per ml, 5 ml vial.....	2,234.00	1	Naglazyme
➔ Restricted (RS1795)			
Initiation			
Metabolic physician			
<i>Re-assessment required after 12 months</i>			
Both:			
1 The patient has been diagnosed with mucopolysaccharidosis VI; and			
2 Either:			
2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency confirmed by either enzyme activity assay in leukocytes or skin fibroblasts; or			
2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.			
Continuation			
Metabolic physician			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and			
2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and			
3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and			
4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.			
HAEM ARGINATE			
Inj 25 mg per ml, 10 ml ampoule			
IDURSULFASE – Restricted see terms below			
⚡ Inj 2 mg per ml, 3 ml vial.....	4,608.30	1	Elaprase
➔ Restricted (RS1546)			
Initiation			
Metabolic physician			
<i>Limited to 24 weeks treatment</i>			
All of the following:			
1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and			
2 Either:			
2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or			
2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and			
3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and			
4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and			
5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.			
LARONIDASE – Restricted see terms below			
⚡ Inj 100 U per ml, 5 ml vial.....	1,335.16	1	Aldurazyme
➔ Restricted (RS1607)			
Initiation			
Metabolic physician			
<i>Limited to 24 weeks treatment</i>			
All of the following:			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysaccharidosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE – **Restricted** see terms [below](#)

- ⚡ Tab 500 mg
- ⚡ Cap 250 mg
- ⚡ Cap 500 mg
- ⚡ Oral liq 500 mg per 10 ml
- ⚡ Oral soln 1,000 mg per 10 ml
- ⚡ Oral soln 1,100 mg per 15 ml
- ⚡ Inj 200 mg per ml, 5 ml vial

➡ **Restricted** (RS1035)

Neurologist, metabolic physician or metabolic disorders dietitian

PYRIDOXAL-5-PHOSPHATE – **Restricted** see terms [below](#)

- ⚡ Tab 50 mg

➡ **Restricted** (RS1331)

Neurologist, metabolic physician or metabolic disorders dietitian

RIBOFLAVIN – **Restricted** see terms [below](#)

- ⚡ Tab 100 mg
- ⚡ Cap 100 mg

➡ **Restricted** (RS1833)

Initiation

Metabolic physician or neurologist

Re-assessment required after 6 months

The patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Continuation

Metabolic physician or neurologist

Re-assessment required after 24 months

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE – **Restricted** see terms [below](#)

- ⚡ Tab soluble 100 mg 1,452.70 30 Kuvan

➡ **Restricted** (RS1796)

Initiation

Metabolic physician

Re-assessment required after 1 month

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Continuation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE

Cap 500 mg

Powder

Soln 100 mg per ml

Inj 20%, 10 ml ampoule

SODIUM PHENYLBUTYRATE – **Some items restricted** see terms [below](#)

Tab 500 mg

↓ Grans 483 mg per g.....2,016.00 174 g Pheburane

Oral liq 250 mg per ml

Inj 200 mg per ml, 10 ml ampoule

→ **Restricted (RS1797)**

Initiation

Metabolic physician

Re-assessment required after 12 months

For the chronic management of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Continuation

Metabolic physician

Re-assessment required after 12 months

The treatment remains appropriate and the patient is benefiting from treatment.

TALIGLUCERASE ALFA – **Restricted** see terms [on the next page](#)

↓ Inj 200 unit vial.....1,072.00 1 Eleyso

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

➔ Restricted (RS1897)

Initiation

Metabolic physician

Re-assessment required after 12 months

All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Continuation

Metabolic physician or any relevant practitioner on the recommendation of a metabolic physician

Re-assessment required after 3 years

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3 RRadiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

TAURINE – **Restricted** see terms [below](#)

⚡ Cap 500 mg

⚡ Cap 1,000 mg

⚡ Powder

➔ Restricted (RS1834)

Initiation

Metabolic physician

Re-assessment required after 6 months

The patient has a suspected specific mitochondrial disorder that may respond to taurine supplementation.

Continuation

Metabolic physician

Re-assessment required after 24 months

Both:

- 1 The patient has a confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

TRIENTINE – **Restricted** see terms [below](#)

↓ Cap 250 mg 2,022.00 100 Trientine Waymade

→ **Restricted** ([RS2026](#))

Initiation

All of the following:

- 1 Patient has confirmed Wilson disease; and
- 2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
- 3 Treatment with zinc has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

Minerals

Calcium

CALCIUM CARBONATE

Tab 1.25 g (500 mg elemental) – **5% DV Feb-24 to 2026** 7.28 250 **Calci-Tab 500**

Tab eff 1.25 g (500 mg elemental)

Tab eff 1.75 g (1 g elemental)

Copper

→ **Restricted** ([RS1928](#))

Initiation – Moderate to severe burns

Limited to 3 months treatment

Both:

- 1 Patient has been hospitalised with moderate to severe burns; and
- 2 Treatment is recommended by a National Burns Unit specialist.

COPPER – **Restricted** see terms [above](#)

† Tab 2.5 mg, chelated

COPPER CHLORIDE – **Restricted** see terms [above](#)

† Inj 0.4 mg per ml, 10 ml vial

Fluoride

SODIUM FLUORIDE

Tab 1.1 mg (0.5 mg elemental)

Iodine

POTASSIUM IODATE

Tab 253 mcg (150 mcg elemental iodine) – **5% DV Feb-24 to 2026** 5.99 90 **NeuroTabs**

POTASSIUM IODATE WITH IODINE

Oral liq 10% with iodine 5%

Iron

FERROUS FUMARATE

Tab 200 mg (65 mg elemental) – **5% DV Feb-25 to 2027** 3.49 100 **Ferro-tab**

ALIMENTARY TRACT AND METABOLISM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FERROUS FUMARATE WITH FOLIC ACID			
Tab 310 mg (100 mg elemental) with folic acid 350 mcg – 5% DV Dec-24 to 2027	5.98	100	Ferro-F-Tabs
FERROUS GLUCONATE WITH ASCORBIC ACID			
Tab 170 mg (20 mg elemental) with ascorbic acid 40 mg			
FERROUS SULFATE			
➔ Tab long-acting 325 mg (105 mg elemental) – Restricted: For continuation only.....	2.55	30	Ferrograd
Oral liq 30 mg (6 mg elemental) per ml – 5% DV Feb-26 to 2028	13.10	500 ml	Ferodan
	10.25	250 ml	Ferro-Liquid
<i>(Ferrograd Tab long-acting 325 mg (105 mg elemental) to be delisted 1 March 2026)</i>			
<i>(Ferodan Oral liq 30 mg (6 mg elemental) per ml to be delisted 1 February 2026)</i>			
FERROUS SULFATE WITH ASCORBIC ACID			
Tab long-acting 325 mg (105 mg elemental) with ascorbic acid 500 mg			
IRON (AS FERRIC CARBOXYMALTOSE) – Restricted see terms below			
⚡ Inj 50 mg per ml, 10 ml vial.....	150.00	1	Ferinject
➔ Restricted (RS1417)			
Initiation			
Treatment with oral iron has proven ineffective or is clinically inappropriate.			
IRON (AS SUCROSE)			
Inj 20 mg per ml, 5 ml ampoule	100.00	5	Venofer
IRON POLYMALTOSE			
Inj 50 mg per ml, 2 ml ampoule	37.95	5	Ferrosig

Magnesium

MAGNESIUM AMINO ACID CHELATE			
Cap 750 mg (150 mg elemental)			
MAGNESIUM CHLORIDE			
Inj 1 mmol per 1 ml, 100 ml bag			
MAGNESIUM HYDROXIDE			
Tab 311 mg (130 mg elemental)			
Suspension 8%			
MAGNESIUM OXIDE			
Cap 663 mg (400 mg elemental)			
Cap 696 mg (420 mg elemental)			
MAGNESIUM OXIDE WITH MAGNESIUM ASPARTATE, MAGNESIUM AMINO ACID CHELATE AND MAGNESIUM CITRATE			
Cap 500 mg with magnesium aspartate 100 mg, magnesium amino acid chelate 100 mg and magnesium citrate 100 mg (360 mg elemental magnesium)			
MAGNESIUM SULPHATE			
Inj 100 mg per ml, 40 ml bag			
Inj 0.4 mmol per ml, 250 ml bag			
Inj 2 mmol per ml, 10 ml ampoule	75.06	10	Inresa
Inj 2 mmol per ml, 5 ml ampoule – 5% DV Jun-24 to 2026	37.53	10	Martindale
Inj 100 mg per ml, 50 ml bag			

↑ Item restricted (see ➔ above); ⚡ Item restricted (see ➔ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Selenium

SELENIUM – **Restricted** see terms [below](#)

↓ Oral liq 150 mcg per 3 drops

*e.g. Clinicians selenium
oral drops*

↓ Inj 300 mcg per ml, 1 ml ampoule

→ **Restricted (RS1929)**

Initiation – Moderate to severe burns

Limited to 3 months treatment

Both:

- 1 Patient has been hospitalised with moderate to severe burns; and
- 2 Treatment is recommended by a National Burns Unit specialist.

Zinc

ZINC

Oral liq 5 mg per 5 drops

ZINC CHLORIDE

Inj 5.3 mg per ml (5.1 mg per ml elemental), 2 ml ampoule

ZINC SULPHATE

Cap 137.4 mg (50 mg elemental) 11.00 100 Zincaps

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

Soln 0.15%

Spray 0.15%

Spray 0.3%

BENZYDAMINE HYDROCHLORIDE WITH CETYLPYRIDINIUM CHLORIDE

Lozenge 3 mg with cetylpyridinium chloride

CARBOXYMETHYLCELLULOSE

Oral spray

CARMELLOSE SODIUM WITH PECTIN AND GELATINE

Paste

Powder

CHLORHEXIDINE GLUCONATE

Mouthwash 0.2% – **5% DV Jan-25 to 2027** 3.99 200 ml **healthE**

DICHLOROBENZYL ALCOHOL WITH AMYLMETACRESOL

Lozenge 1.2 mg with amylmetacresol 0.6 mg

TRIAMCINOLONE ACETONIDE

Paste 0.1% – **5% DV Feb-24 to 2026** 5.49 5 g **Kenalog in Orabase**

Oropharyngeal Anti-Infectives

AMPHOTERICIN B

Lozenge 10 mg 5.86 20 Fungilin

MICONAZOLE

Oral gel 20 mg per g – **5% DV Feb-25 to 2027** 5.19 40 g **Decozol**

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NYSTATIN			
Oral liquid 100,000 u per ml – 5% DV Feb-24 to 2026	2.22	24 ml	Nilstat

Other Oral Agents

HYALURONIC ACID WITH LIDOCAINE [LIGNOCAINE]

Inj 20 mg per ml

SODIUM HYALURONATE [HYALURONIC ACID] – **Restricted** see terms [below](#)

↓ Inj 20 mg per ml, 1 ml syringe

→ **Restricted (RS1175)**

Otolaryngologist

Vitamins

Multivitamin Preparations

MULTIVITAMIN AND MINERAL SUPPLEMENT – **Restricted** see terms [below](#)

↓ Cap..... 23.35 180 Clinicians Multivit & Mineral Boost

→ **Restricted (RS1498)**

Initiation

Limited to 3 months treatment

Both:

- 1 Patient was admitted to hospital with burns; and
- 2 Any of the following:
 - 2.1 Burn size is greater than 15% of total body surface area (BSA) for all types of burns; or
 - 2.2 Burn size is greater than 10% of BSA for mid-dermal or deep dermal burns; or
 - 2.3 Nutritional status prior to admission or dietary intake is poor.

MULTIVITAMIN RENAL – **Restricted** see terms [below](#)

↓ Cap..... 7.28 30 Clinicians Renal Vit

→ **Restricted (RS1499)**

Initiation

Either:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73m² body surface area (BSA).

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MULTIVITAMINS			
Tab (BPC cap strength).....	18.50	1,000	Mvite
↓ cap vitamin A 2500 u, betacarotene 3 mg, cholecalciferol 11 mcg, alpha tocopherol 150 u, phytomenadione 150 mcg, folic acid 0.2 mg, ascorbic acid 100 mg, thiamine 1.5 mg, pantothenic acid 12 mg, riboflavin 1.7 mg, niacin 20 mg, pyridoxine hydrochloride 1.9 mg, cyanocobalamin 3 mcg, zinc 7.5 mg and biotin 100 mcg			e.g. Vitabdeck
→ Restricted (RS1620)			
Initiation			
Any of the following:			
1 Patient has cystic fibrosis with pancreatic insufficiency; or			
2 Patient is an infant or child with liver disease or short gut syndrome; or			
3 Patient has severe malabsorption syndrome.			
↓ Powder vitamin A 3200 mcg with vitamin D 100 mcg, vitamin E 54.2 mg, vitamin C 400 mg, vitamin K1 108 mcg thiamine 3.2 mg, riboflavin 4.4 mg, niacin 41 mg, vitamin B6 3.6 mg, folic acid 600 mcg, vitamin B12 9 mcg, biotin 120 mcg, pantothenic acid 24 mg, choline 1250 mg and inositol 700 mg	74.88	200 g	Paediatric Seravit
→ Restricted (RS1178)			
Initiation			
Patient has inborn errors of metabolism.			
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg and glucose 1000 mg, 5 ml ampoule (1)			e.g. Pabrinex IV
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg, 2 ml ampoule (1)			
Inj thiamine hydrochloride 500 mg with riboflavin 8 mg and pyridoxine hydrochloride 100 mg, 10 ml ampoule (1) and inj ascorbic acid 1000 mg with nicotinamide 320 mg and glucose 2000 mg, 10 ml ampoule (1)			
Vitamin A			
RETINOL			
Tab 10,000 iu			
Cap 25,000 iu			
Oral liq 150,000 iu per ml			
Oral liq 666.7 mcg per 2 drops, 10 ml			
Oral liq 5,000 iu per drop, 30 ml			
Vitamin B			
HYDROXOCOBALAMIN			
Inj 1 mg per ml, 1 ml ampoule – 5% DV Jul-25 to 2027	3.95	3	Hydroxocobalamin Panpharma
PYRIDOXINE HYDROCHLORIDE			
Tab 25 mg – 5% DV Feb-24 to 2026	3.43	90	Vitamin B6 25
Tab 50 mg	23.45	500	Pyridoxine multichem
Inj 100 mg per ml, 2 ml vial			
Inj 100 mg per ml, 1 ml ampoule			
Inj 100 mg per ml, 30 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
THIAMINE HYDROCHLORIDE			
Tab 50 mg	4.65	100	Thiamine multichem
Tab 100 mg			
Inj 100 mg per ml, 1 ml vial			<i>e.g. Benerva</i>
Inj 100 mg per ml, 2 ml vial			
Inj 125 mg per ml, 2 ml vial			
VITAMIN B COMPLEX			
Tab strong, BPC	11.25	500	Bplex

Vitamin C

ASCORBIC ACID			
Tab 100 mg	12.50	500	Cvite
Tab chewable 250 mg			

Vitamin D

ALFACALCIDOL			
Cap 0.25 mcg	26.32	100	One-Alpha
Cap 1 mcg	87.98	100	One-Alpha
Oral drops 2 mcg per ml	60.68	20 ml	One-Alpha
CALCITRIOL			
Cap 0.25 mcg	7.89	100	Calcitriol XL
			Calcitriol-AFT
Cap 0.5 mcg	13.68	100	Calcitriol XL
			Calcitriol-AFT
Oral liq 1 mcg per ml			
Inj 1 mcg per ml, 1 ml ampoule			
COLECALCIFEROL			
Cap 1.25 mg (50,000 iu) – 5% DV Jun-24 to 2026	3.65	12	Vit.D3
Oral liq 188 mcg per ml (7,500 iu per ml)	9.00	5 ml	Clinicians

Vitamin E

ALPHA TOCOPHERYL – **Restricted** see terms [below](#)

↓ Oral liq 156 u per ml

→ **Restricted (RS1632)**

Initiation – Cystic fibrosis

Both:

1 Cystic fibrosis patient; and

2 Either:

2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or

2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

Initiation – Osteoradionecrosis

For the treatment of osteoradionecrosis.

Initiation – Other indications

All of the following:

1 Infant or child with liver disease or short gut syndrome; and

continued...

	Price	Brand or
	(ex man. excl. GST)	Generic
	\$	Manufacturer
	Per	

continued...

- 2 Requires vitamin supplementation; and
- 3 Either:
 - 3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
 - 3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

ALPHA TOCOPHERYL ACETATE – **Restricted** see terms [below](#)

- ↓ Cap 100 u
- ↓ Cap 500 u
- ↓ Oral liq 156 u per ml

➡ **Restricted (RS1176)**

Initiation – Cystic fibrosis

- Both:
- 1 Cystic fibrosis patient; and
 - 2 Either:
 - 2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
 - 2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

Initiation – Osteoradionecrosis

For the treatment of osteoradionecrosis.

Initiation – Other indications

- All of the following:
- 1 Infant or child with liver disease or short gut syndrome; and
 - 2 Requires vitamin supplementation; and
 - 3 Either:
 - 3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
 - 3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Antianaemics

Hypoplastic and Haemolytic

EPOETIN ALFA – **Restricted** see terms [below](#)

⚡ Inj 1,000 iu in 0.5 ml syringe.....	250.00	6	Binocrit
⚡ Inj 2,000 iu in 1 ml syringe.....	100.00	6	Binocrit
⚡ Inj 3,000 iu in 0.3 ml syringe.....	150.00	6	Binocrit
⚡ Inj 4,000 iu in 0.4 ml syringe.....	96.50	6	Binocrit
⚡ Inj 5,000 iu in 0.5 ml syringe.....	125.00	6	Binocrit
⚡ Inj 6,000 iu in 0.6 ml syringe.....	145.00	6	Binocrit
⚡ Inj 8,000 iu in 0.8 ml syringe.....	175.00	6	Binocrit
⚡ Inj 10,000 iu in 1 ml syringe.....	197.50	6	Binocrit
⚡ Inj 40,000 iu in 1 ml syringe.....	250.00	1	Binocrit

➡ **Restricted (RS1660)**

Initiation – chronic renal failure

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initiation – myelodysplasia*

Re-assessment required after 2 months

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Continuation – myelodysplasia*

Re-assessment required after 12 months

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with epoetin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Initiation – all other indications

Haematologist

For use in patients where blood transfusion is not a viable treatment alternative.

Note: Indications marked with * are unapproved indications

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

EPOETIN BETA – Restricted see terms [below](#)

Note: Epoetin beta is considered a Discretionary Variance Pharmaceutical for epoetin alfa.

- ↓ Inj 2,000 iu in 0.3 ml syringe
- ↓ Inj 3,000 iu in 0.3 ml syringe
- ↓ Inj 4,000 iu in 0.3 ml syringe
- ↓ Inj 5,000 iu in 0.3 ml syringe
- ↓ Inj 6,000 iu in 0.3 ml syringe
- ↓ Inj 10,000 iu in 0.6 ml syringe

→ **Restricted (RS1661)**

Initiation – chronic renal failure

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initiation – myelodysplasia*

Re-assessment required after 12 months

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Continuation – myelodysplasia*

Re-assessment required after 2 months

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with epoetin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Initiation – all other indications

All of the following:

- 1 Haematologist; and
- 2 For use in patients where blood transfusion is not a viable treatment alternative; and
- 3 *Note: Indications marked with * are unapproved indications.

Megaloblastic

FOLIC ACID

Tab 0.8 mg	26.60	1,000	Folic Acid multichem
Tab 5 mg – 1% DV Mar-23 to 2027	5.82	100	Folic Acid Viatrix
Oral liq 50 mcg per ml	31.77	25 ml	Biomed
Inj 5 mg per ml, 10 ml vial			

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

Antifibrinolytics, Haemostatics and Local Sclerosants

ALUMINIUM CHLORIDE – **Restricted** see terms [below](#)

↓ Topical soln 20% w/v

e.g. Driclor

→ **Restricted** (RS1500)

Initiation

For use as a haemostatis agent.

APROTININ – **Restricted** see terms [below](#)

↓ Inj 10,000 kIU per ml (equivalent to 200 mg per ml), 50 ml vial

→ **Restricted** (RS1332)

Initiation

Cardiac anaesthetist

Either:

- 1 Paediatric patient undergoing cardiopulmonary bypass procedure; or
- 2 Adult patient undergoing cardiac surgical procedure where the significant risk of massive bleeding outweighs the potential adverse effects of the drug.

ELTROMBOPAG – **Restricted** see terms [below](#)

↓ Tab 25 mg 1,550.00

28 Revolade

↓ Tab 50 mg 3,100.00

28 Revolade

→ **Restricted** (RS1648)

Initiation – idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

Re-assessment required after 6 weeks

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initiation – idiopathic thrombocytopenic purpura - preparation for splenectomy

Haematologist

Limited to 6 weeks treatment

The patient requires eltrombopag treatment as preparation for splenectomy.

Continuation – idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

Re-assessment required after 12 months

The patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre

Initiation – idiopathic thrombocytopenic purpura contraindicated to splenectomy

Haematologist

Re-assessment required after 3 months

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Continuation – idiopathic thrombocytopenic purpura contraindicated to splenectomy

Haematologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Initiation – severe aplastic anaemia

Haematologist

Re-assessment required after 3 months

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Continuation – severe aplastic anaemia

Haematologist

Re-assessment required after 12 months

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB – Restricted see terms below

⚡ Inj 30 mg in 1 ml vial.....	3,570.00	1	Hemlibra
⚡ Inj 60 mg in 0.4 ml vial.....	7,138.00	1	Hemlibra
⚡ Inj 105 mg in 0.7 ml vial.....	12,492.00	1	Hemlibra
⚡ Inj 150 mg in 1 ml vial.....	17,846.00	1	Hemlibra

➡ Restricted (RS1998)

Initiation – Severe Haemophilia A with or without FVIII inhibitors

Haematologist

Both:

- 1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
- 2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

FERRIC SUBSULFATE

Gel 25.9%

Soln 500 ml

POLIDOCANOL

Inj 0.5%, 30 ml vial

BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM TETRADECYL SULPHATE			
Inj 3%, 2 ml ampoule			
THROMBIN			
Powder			
TRANEXAMIC ACID			
Tab 500 mg	10.45	60	Mercury Pharma
Inj 100 mg per ml, 5 ml ampoule – 5% DV Mar-25 to 2027	5.39	5	Tranexamic-AFT
Inj 100 mg per ml, 10 ml ampoule – 5% DV Mar-25 to 2027	7.99	5	Tranexamic-AFT

Anticoagulant Reversal Agents

IDARUCIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 50 mg per ml, 50 ml vial.....4,250.00 2 Praxbind

➡ **Restricted (RS1535)**

Initiation

For the reversal of the anticoagulant effects of dabigatran when required in situations of life-threatening or uncontrolled bleeding, or for emergency surgery or urgent procedures.

Blood Factors

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – **Restricted** see terms [below](#)

⚡ Inj 250 iu vial.....612.50 1 Alprolix
 ⚡ Inj 500 iu vial.....1,225.00 1 Alprolix
 ⚡ Inj 1,000 iu vial.....2,450.00 1 Alprolix
 ⚡ Inj 2,000 iu vial.....4,900.00 1 Alprolix
 ⚡ Inj 3,000 iu vial.....7,350.00 1 Alprolix
 ⚡ Inj 4,000 iu vial.....9,800.00 1 Alprolix

➡ **Restricted (RS1684)**

Initiation

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] – **Restricted** see terms [below](#)

⚡ Inj 1 mg syringe.....1,178.30 1 NovoSeven RT
 ⚡ Inj 2 mg syringe.....2,356.60 1 NovoSeven RT
 ⚡ Inj 5 mg syringe.....5,891.50 1 NovoSeven RT
 ⚡ Inj 8 mg syringe.....9,426.40 1 NovoSeven RT

➡ **Restricted (RS1704)**

Initiation

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

FACTOR EIGHT INHIBITOR BYPASSING FRACTION – **Restricted** see terms [below](#)

⚡ Inj 500 U1,315.00 1 FEIBA NF
 ⚡ Inj 1,000 U2,630.00 1 FEIBA NF
 ⚡ Inj 2,500 U6,575.00 1 FEIBA NF

➡ **Restricted (RS1705)**

Initiation

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – Restricted see terms below			
⚡ Inj 250 iu prefilled syringe.....	287.50	1	Xyntha
⚡ Inj 500 iu prefilled syringe.....	575.00	1	Xyntha
⚡ Inj 1,000 iu prefilled syringe.....	1,150.00	1	Xyntha
⚡ Inj 2,000 iu prefilled syringe.....	2,300.00	1	Xyntha
⚡ Inj 3,000 iu prefilled syringe.....	3,450.00	1	Xyntha
➔ Restricted (RS1706)			
Initiation			
For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.			
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – Restricted see terms below			
⚡ Inj 1,000 iu vial.....	870.00	1	RIXUBIS
⚡ Inj 2,000 iu vial.....	1,740.00	1	RIXUBIS
⚡ Inj 3,000 iu vial.....	2,610.00	1	RIXUBIS
➔ Restricted (RS1679)			
Initiation			
For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.			
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – Restricted see terms below			
⚡ Inj 500 iu vial.....	420.00	1	Advate
⚡ Inj 1,000 iu vial.....	840.00	1	Advate
⚡ Inj 2,000 iu vial.....	1,680.00	1	Advate
⚡ Inj 3,000 iu vial.....	2,520.00	1	Advate
➔ Restricted (RS1707)			
Initiation			
For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.			
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) – Restricted see terms below			
⚡ Inj 250 iu vial.....	237.50	1	Kogenate FS
⚡ Inj 500 iu vial.....	475.00	1	Kogenate FS
⚡ Inj 1,000 iu vial.....	950.00	1	Kogenate FS
⚡ Inj 2,000 iu vial.....	1,900.00	1	Kogenate FS
⚡ Inj 3,000 iu vial.....	2,850.00	1	Kogenate FS
➔ Restricted (RS1708)			
Initiation			
For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.			
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] – Restricted see terms below			
⚡ Inj 1,000 iu vial.....	1,200.00	1	Adynovate
⚡ Inj 2,000 iu vial.....	2,400.00	1	Adynovate
➔ Restricted (RS1682)			
Initiation			
For patients with haemophilia A receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.			
Vitamin K			
PHYTOMENADIONE			
Inj 2 mg in 0.2 ml ampoule	8.00	5	Konakion MM Paediatric
Inj 10 mg per ml, 1 ml ampoule	9.21	5	Konakion MM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antithrombotics			
Anticoagulants			
BIVALIRUDIN – Restricted see terms below			
⚡ Inj 250 mg vial			
➡ Restricted (RS1181)			
Initiation			
Either:			
1 For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance; or			
2 For use in patients undergoing endovascular procedures.			
CITRATE SODIUM			
Inj 4% (200 mg per 5 ml), 5 ml ampoule			
Inj 46.7% (1.4 g per 3 ml), 3 ml syringe			
Inj 46.7% (2.36 g per 5 ml), 5 ml ampoule			
DABIGATRAN			
Cap 75 mg – 5% DV Jul-24 to 2026	27.99	60	Pradaxa
Cap 110 mg – 5% DV Jul-24 to 2026	27.99	60	Pradaxa
Cap 150 mg – 5% DV Jul-24 to 2026	27.99	60	Pradaxa
DANAPAROID – Restricted see terms below			
⚡ Inj 750 u in 0.6 ml ampoule			
➡ Restricted (RS1182)			
Initiation			
For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance.			
DEFIBROTIDE – Restricted see terms below			
⚡ Inj 80 mg per ml, 2.5 ml ampoule			
➡ Restricted (RS1183)			
Initiation			
Haematologist			
Patient has moderate or severe sinusoidal obstruction syndrome as a result of chemotherapy or regimen-related toxicities.			
DEXTROSE WITH SODIUM CITRATE AND CITRIC ACID [ACID CITRATE DEXTROSE A]			
Inj 24.5 mg with sodium citrate 22 mg and citric acid 7.3 mg per ml, 100 ml bag			
ENOXAPARIN SODIUM			
Inj 20 mg in 0.2 ml syringe – 5% DV Feb-25 to 2027	21.90	10	Clexane
Inj 40 mg in 0.4 ml ampoule			
Inj 40 mg in 0.4 ml syringe – 5% DV Feb-25 to 2027	29.74	10	Clexane
Inj 60 mg in 0.6 ml syringe – 5% DV Feb-25 to 2027	42.47	10	Clexane
Inj 80 mg in 0.8 ml syringe – 5% DV Feb-25 to 2027	56.62	10	Clexane
Inj 100 mg in 1 ml syringe – 5% DV Feb-25 to 2027	70.91	10	Clexane
Inj 120 mg in 0.8 ml syringe – 5% DV Feb-25 to 2027	88.11	10	Clexane Forte
Inj 150 mg in 1 ml syringe – 5% DV Feb-25 to 2027	100.70	10	Clexane Forte
FONDAPARINUX SODIUM – Restricted see terms below			
⚡ Inj 2.5 mg in 0.5 ml syringe			
⚡ Inj 7.5 mg in 0.6 ml syringe			
➡ Restricted (RS1184)			
Initiation			
For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
HEPARIN SODIUM			
Inj 5,000 iu per ml, 5 ml vial.....	83.00	10	Heparin Sodium Panpharma
Inj 100 iu per ml, 250 ml bag			
Inj 1,000 iu per ml, 1 ml ampoule	362.98	50	Hospira
Inj 1,000 iu per ml, 5 ml ampoule	127.44	50	Pfizer
	25.49	10	Wockhardt
	103.70		Wockhardt PSF
Inj 5,000 iu in 0.2 ml ampoule			
Inj 5,000 iu per ml, 1 ml ampoule	70.33	5	Hospira
Inj 1,000 iu per ml, 10 ml vial.....	127.44	25	Pfizer
<i>(Wockhardt Inj 1,000 iu per ml, 5 ml ampoule to be delisted 1 January 2026)</i>			
<i>(Wockhardt PSF Inj 1,000 iu per ml, 5 ml ampoule to be delisted 1 January 2026)</i>			
<i>(Pfizer Inj 1,000 iu per ml, 10 ml vial to be delisted 1 January 2026)</i>			
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml ampoule	96.91	50	Pfizer
Inj 100 iu per ml, 2 ml ampoule			
Inj 100 iu per ml, 5 ml ampoule			
PHENINDIONE			
Tab 10 mg			
Tab 25 mg			
Tab 50 mg			
PROTAMINE SULPHATE			
Inj 10 mg per ml, 5 ml ampoule			
RIVAROXABAN			
Tab 10 mg – 5% DV Dec-23 to 2026	15.60	30	Xarelto
Tab 15 mg – 5% DV Dec-23 to 2026	14.56	28	Xarelto
Tab 20 mg – 5% DV Dec-23 to 2026	14.56	28	Xarelto
SODIUM CITRATE WITH SODIUM CHLORIDE AND POTASSIUM CHLORIDE			
Inj 4.2 mg with sodium chloride 5.7 mg and potassium chloride 74.6 mcg per ml, 5,000 ml bag			
WARFARIN SODIUM			
Tab 1 mg	7.50	100	Marevan
Tab 2 mg			
Tab 3 mg	12.00	100	Marevan
Tab 5 mg	13.50	100	Marevan
Antiplatelets			
ASPIRIN			
Tab 100 mg – 5% DV Jun-24 to 2026	12.65	990	Ethics Aspirin EC
Suppos 300 mg			
CLOPIDOGREL			
Tab 75 mg – 5% DV Dec-25 to 2028	5.07	84	Arrow - Clopid
DIPYRIDAMOLE – Restricted: For continuation only			
➡ Tab 25 mg			
➡ Tab long-acting 150 mg.....	13.93	60	Pytazen SR
➡ Cap modified-release 200 mg	55.13	60	Dipyridamole - Strides
➡ Inj 5 mg per ml, 2 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
EPTIFIBATIDE – Restricted see terms below			
↓ Inj 2 mg per ml, 10 ml vial.....	180.38	1	Eptifibatide Viatris
↓ Inj 750 mcg per ml, 100 ml vial.....	526.50	1	Eptifibatide Viatris

➔ **Restricted (RS1759)**

Initiation

Any of the following:

- 1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
- 2 For use in patients with definite or strongly suspected intra-coronary thrombus on coronary angiography; or
- 3 For use in patients undergoing intra-cranial intervention.

LYSINE ACETYSALICYLATE [LYSINE ASPRIN] – Restricted see terms [below](#)

↓ Inj 500 mg

e.g. Aspegic

➔ **Restricted (RS1689)**

Initiation

Both:

- 1 For use when an immediate antiplatelet effect is required prior to an urgent interventional neuro-radiology or interventional cardiology procedure; and
- 2 Administration of oral aspirin would delay the procedure.

TICAGRELOR – Restricted see terms [below](#)

↓ Tab 90 mg – **5% DV Dec-24 to 2027**.....20.35

56

Ticagrelor Sandoz

➔ **Restricted (RS1774)**

Initiation

Restricted to treatment of acute coronary syndromes specifically for patients who have recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome, and in whom fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initiation – thrombosis prevention neurological stenting

Re-assessment required after 12 months

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent..

Continuation – thrombosis prevention neurological stenting

Re-assessment required after 12 months

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Initiation – Percutaneous coronary intervention with stent deployment

Limited to 12 months treatment

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

3 Patient is clopidogrel-allergic**.

Initiation – Stent thrombosis

Patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Initiation – Myocardial infarction

Limited to 1 week treatment

For short term use while in hospital following ST-elevated myocardial infarction.

Notes: Indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment

TICLOPIDINE

Tab 250 mg

Fibrinolytic Agents

ALTEPLASE

Inj 2 mg vial
Inj 10 mg vial
Inj 50 mg vial

TENECTEPLASE

Inj 50 mg vial

UROKINASE

Inj 5,000 iu vial
Inj 10,000 iu vial
Inj 50,000 iu vial
Inj 100,000 iu vial
Inj 250,000 iu vial
Inj 500,000 iu vial

Colony-Stimulating Factors

Drugs Used to Mobilise Stem Cells

PLERIXAFOR – **Restricted** see terms [below](#)

↓ Inj 20 mg per ml, 1.2 ml vial.....8,740.00 1 Mozobil

→ **Restricted (RS1536)**

Initiation – Autologous stem cell transplant

Haematologist

Limited to 3 days treatment

All of the following:

- 1 Patient is to undergo stem cell transplantation; and
- 2 Patient has not had a previous unsuccessful mobilisation attempt with plerixafor; and
- 3 Any of the following:

3.1 Both:

3.1.1 Patient is undergoing G-CSF mobilisation; and

3.1.2 Either:

3.1.2.1 Has a suboptimal peripheral blood CD34 count of less than or equal to $10 \times 10^6/L$ on day 5 after 4 days of G-CSF treatment; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 3.1.2.2 Efforts to collect $> 1 \times 10^6$ CD34 cells/kg have failed after one apheresis procedure; or
- 3.2 Both:
- 3.2.1 Patient is undergoing chemotherapy and G-CSF mobilisation; and
- 3.2.2 Any of the following:
- 3.2.2.1 Both:
- 3.2.2.1.1 Has rising white blood cell counts of $> 5 \times 10^9/L$; and
- 3.2.2.1.2 Has a suboptimal peripheral blood CD34 count of less than or equal to $10 \times 10^6/L$; or
- 3.2.2.2 Efforts to collect $> 1 \times 10^6$ CD34 cells/kg have failed after one apheresis procedure; or
- 3.2.2.3 The peripheral blood CD34 cell counts are decreasing before the target has been received; or
- 3.3 A previous mobilisation attempt with G-CSF or G-CSF plus chemotherapy has failed.

Granulocyte Colony-Stimulating Factors

FILGRASTIM – **Restricted** see terms [below](#)

‡ Inj 300 mcg in 0.5 ml prefilled syringe – 5% DV Dec-24 to 2027	86.60	10	Nivestim
‡ Inj 300 mcg in 1 ml vial	520.00	4	Neupogen
‡ Inj 480 mcg in 0.5 ml prefilled syringe – 5% DV Dec-24 to 2027	133.72	10	Nivestim

→ **Restricted (RS1188)**

Haematologist or oncologist

PEGFILGRASTIM – **Restricted** see terms [below](#)

‡ Inj 6 mg per 0.6 ml syringe – 5% DV Feb-26 to 2028	69.50	1	Ziextenzo
--	-------	---	-----------

→ **Restricted (RS1743)**

Initiation

For prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*).

Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines

Fluids and Electrolytes

Intravenous Administration

CALCIUM CHLORIDE

Inj 100 mg per ml, 10 ml vial			
Inj 100 mg per ml, 50 ml syringe			e.g. Baxter

CALCIUM GLUCONATE

Inj 10%, 10 ml ampoule			e.g. Max Health
------------------------	--	--	-----------------

COMPOUND ELECTROLYTES

Inj sodium 140 mmol/l, potassium 5 mmol/l, magnesium 1.5 mmol/l, chloride 98 mmol/l, acetate 27 mmol/l, gluconate 23 mmol/l, 500 ml bag	62.82	18	Plasma-Lyte 148
Inj sodium 140 mmol/l, potassium 5 mmol/l, magnesium 1.5 mmol/l, chloride 98 mmol/l, acetate 27 mmol/l, gluconate 23 mmol/l, 1,000 ml bag	30.72	12	Plasma-Lyte 148

COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]

Inj sodium 140 mmol/l, 5 mmol/l potassium, 1.5 mmol/l magnesium, 98 mmol/l chloride, 27 mmol/l acetate and 23 mmol/l gluconate, glucose 23 mmol/l (5%), 1,000 ml bag	239.04	12	Plasma-Lyte 148 & 5% Glucose
--	--------	----	------------------------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
COMPOUND SODIUM LACTATE [HARTMANN'S SOLUTION]			
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bicarbonate 29 mmol/l, chloride 111 mmol/l, 500 ml bag	27.90	18	Baxter
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bicarbonate 29 mmol/l, chloride 111 mmol/l, 1,000 ml bag	19.32	12	Baxter
GLUCOSE [DEXTROSE]			
Inj 5%, 1,000 ml bag.....	52.00	10	Fresenius Kabi
Inj 5%, 100 ml bag.....	95.00	50	Fresenius Kabi
Inj 5%, 250 ml bag.....	61.50	30	Fresenius Kabi
Inj 5%, 50 ml bag.....	162.00	60	Baxter Glucose 5%
Inj 5%, 500 ml bag.....	66.00	20	Fresenius Kabi
Inj 10%, 1,000 ml bag.....	162.00	12	Baxter Glucose 10%
Inj 10%, 500 ml bag.....	126.00	18	Baxter Glucose 10%
Inj 50%, 10 ml ampoule – 5% DV Feb-24 to 2026	34.75	5	Biomed
Inj 50%, 500 ml bag.....	423.00	18	Baxter Glucose 50%
Inj 50%, 90 ml bottle – 5% DV Feb-24 to 2026	17.50	1	Biomed
GLUCOSE WITH POTASSIUM CHLORIDE			
Inj 10% glucose with 20 mmol/l potassium chloride, 500 ml bag			
GLUCOSE WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE			
Inj 2.5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 3,000 ml bag			
Inj 10% glucose with potassium chloride 10 mmol/l and sodium chloride 15 mmol/l, 500 ml bag			
Inj 4% glucose with potassium chloride 20 mmol/l and sodium chloride 0.18%, 1,000 ml bag.....	240.36	12	Baxter
Inj 5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 1,000 ml bag.....	189.00	12	Baxter
Inj 5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.9%, 1,000 ml bag.....	334.08	12	Baxter
GLUCOSE WITH SODIUM CHLORIDE			
Inj glucose 2.5% with sodium chloride 0.45%, 500 ml bag.....	318.78	18	Baxter
Inj 4% glucose and sodium chloride 0.18%, 1,000 ml bag.....	192.96	12	Baxter
Inj 5% glucose and sodium chloride 0.45%, 1,000 ml bag.....	192.84	12	Baxter
Inj 5% glucose and sodium chloride 0.9%, 1,000 ml bag.....	204.84	12	Baxter
POTASSIUM CHLORIDE			
Inj 75 mg (1 mmol) per ml, 10 ml ampoule			
Inj 225 mg (3 mmol) per ml, 20 ml ampoule			
POTASSIUM CHLORIDE WITH SODIUM CHLORIDE			
Inj 10 mmol potassium chloride with 0.29% sodium chloride, 100 ml bag	563.52	48	Baxter
Inj 20 mmol potassium chloride with 0.9% sodium chloride, 1,000 ml bag.....	192.72	12	Baxter
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 1,000 ml bag.....	299.40	12	Baxter
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 100 ml bag	912.96	48	Baxter
POTASSIUM DIHYDROGEN PHOSPHATE			
Inj 1 mmol per ml, 10 ml ampoule	174.57	10	Hospira
RINGER'S SOLUTION			
Inj sodium 147 mmol/l with potassium 4 mmol/l, calcium 2.2 mmol/l, chloride 156 mmol/l, 1,000 ml bag.....	227.52	12	Baxter
SODIUM ACETATE			
Inj 4 mmol per ml, 20 ml ampoule			

BLOOD AND BLOOD FORMING ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM BICARBONATE			
Inj 8.4%, 10 ml vial			
Inj 8.4%, 50 ml vial	24.70	1	Biomed
Inj 8.4%, 100 ml vial	25.31	1	Biomed
SODIUM CHLORIDE			
Inj 0.9%, 5 ml ampoule – 5% DV Feb-26 to 2028	4.12	20	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – 5% DV Feb-26 to 2028	7.50	50	Fresenius Kabi
⚡ Inj 0.9%, 3 ml syringe, non-sterile pack	12.00	30	BD PosiFlush
➡ Restricted (RS1297)			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⚡ Inj 0.9%, 5 ml syringe, non-sterile pack	12.00	30	BD PosiFlush
➡ Restricted (RS1297)			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⚡ Inj 0.9%, 10 ml syringe, non-sterile pack	11.70	30	BD PosiFlush
➡ Restricted (RS1297)			
Initiation			
For use in flushing of in-situ vascular access devices only.			
Inj 0.9%, 20 ml ampoule – 5% DV Feb-26 to 2028	5.20	20	Fresenius Kabi
Inj 23.4% (4 mmol/ml), 20 ml ampoule	40.15	5	Biomed
Inj 0.45%, 500 ml bag	84.42	18	Baxter
Inj 3%, 1,000 ml bag	165.84	12	Baxter
Inj 0.9%, 50 ml bag	124.20	60	Baxter
	147.75	75	Baxter-Viaflo
Inj 0.9%, 100 ml bag	88.80	48	Baxter
	105.60	60	Baxter-Viaflo
Inj 0.9%, 250 ml bag	50.40	24	Baxter
Inj 0.9%, 500 ml bag	27.54	18	Baxter
Inj 0.9%, 1,000 ml bag	18.96	12	Baxter
Inj 1.8%, 500 ml bottle			
SODIUM DIHYDROGEN PHOSPHATE [SODIUM ACID PHOSPHATE]			
Inj 1 mmol per ml, 20 ml ampoule	59.10	5	Biomed
WATER			
Inj 10 ml ampoule	7.60	50	Fresenius Kabi
			Multichem
Inj 20 ml ampoule	5.00	20	Fresenius Kabi
Inj 250 ml bag			
Inj 500 ml bag			
Inj, 1,000 ml bag	24.12	12	Baxter

Oral Administration

CALCIUM POLYSTYRENE SULPHONATE			
Powder	169.85	300 g	Calcium Resonium
COMPOUND ELECTROLYTES			
Powder for oral soln – 5% DV Dec-25 to 2028	9.50	50	Electral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]			
Soln with electrolytes	6.53	1	Hydralyte - Lemonade
PHOSPHORUS			
Tab eff 500 mg (16 mmol)			

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
POTASSIUM CHLORIDE			
Tab eff 548 mg (14 mmol) with chloride 285 mg (8 mmol)			
Tab long-acting 600 mg (8 mmol) – 5% DV Feb-26 to 2028	16.15	200	Span-K
Oral liq 2 mmol per ml			
SODIUM BICARBONATE			
Cap 840 mg	8.52	100	Sodibic
SODIUM CHLORIDE			
Tab 600 mg			
Oral liq 2 mmol/ml			
SODIUM POLYSTYRENE SULPHONATE			
Powder	84.65	454 g	Resonium A
Plasma Volume Expanders			
GELATINE, SUCCINYLATED			
Inj 4%, 500 ml bag	139.10	10	Gelofusine

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Agents Affecting the Renin-Angiotensin System			
ACE Inhibitors			
CAPTOPRIL			
↓ Oral liq 5 mg per ml – 5% DV Apr-24 to 2026	86.00	100 ml	DP-Captopril
➔ Restricted (RS1263)			
Initiation			
Any of the following:			
1 For use in children under 12 years of age; or			
2 For use in tube-fed patients; or			
3 For management of rebound transient hypertension following cardiac surgery.			
ENALAPRIL MALEATE			
Tab 5 mg	1.75	90	Acetec
Tab 10 mg	1.97	90	Acetec
Tab 20 mg	2.35	90	Acetec
LISINOPRIL			
Tab 5 mg	11.07	90	Teva Lisinopril
Tab 10 mg	11.67	90	Teva Lisinopril
Tab 20 mg	14.69	90	Teva Lisinopril
PERINDOPRIL			
Tab 2 mg – 5% DV Dec-24 to 2027	1.79	30	Coversyl
Tab 4 mg – 5% DV Dec-24 to 2027	2.44	30	Coversyl
Tab 8 mg – 5% DV Dec-24 to 2027	3.94	30	Coversyl
QUINAPRIL			
Tab 5 mg – 5% DV Mar-25 to 2027	10.24	90	Arrow-Quinapril 5
Tab 10 mg – 5% DV Mar-25 to 2027	12.51	90	Arrow-Quinapril 10
Tab 20 mg – 5% DV Mar-25 to 2027	14.38	90	Arrow-Quinapril 20
RAMIPRIL			
Cap 1.25 mg – 5% DV Feb-25 to 2027	17.25	90	Tryzan
Cap 2.5 mg – 5% DV Feb-25 to 2027	16.50	90	Tryzan
Cap 5 mg – 5% DV Feb-25 to 2027	16.88	90	Tryzan
Cap 10 mg – 5% DV Feb-25 to 2027	17.63	90	Tryzan
Angiotensin II Antagonists			
CANDESARTAN CILEXETIL			
Tab 4 mg – 5% DV Feb-25 to 2027	2.68	90	Candestar
Tab 8 mg – 5% DV Feb-25 to 2027	2.67	90	Candestar
Tab 16 mg – 5% DV Feb-25 to 2027	4.22	90	Candestar
Tab 32 mg – 5% DV Feb-25 to 2027	5.24	90	Candestar
LOSARTAN POTASSIUM			
Tab 12.5 mg – 5% DV Mar-24 to 2026	2.00	84	Losartan Actavis
Tab 25 mg – 5% DV Mar-24 to 2026	2.29	84	Losartan Actavis
Tab 50 mg – 5% DV Mar-24 to 2026	2.86	84	Losartan Actavis
Tab 100 mg – 5% DV Mar-24 to 2026	4.57	84	Losartan Actavis

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Angiotensin II Antagonists with Diuretics			
CANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDE			
Tab 16 mg with hydrochlorothiazide 12.5 mg.....	4.10	30	APO-Candesartan HCTZ 16/12.5
Tab 32 mg with hydrochlorothiazide 12.5 mg.....	5.25	30	APO-Candesartan HCTZ 32/12.5
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE			
Tab 50 mg with hydrochlorothiazide 12.5 mg.....	4.00	30	Arrow-Losartan & Hydrochlorothiazide
Angiotensin II Antagonists with Neprilysin Inhibitors			
SACUBITRIL WITH VALSARTAN – Restricted see terms below			
↓ Tab 24.3 mg with valsartan 25.7 mg	190.00	56	Entresto 24/26
↓ Tab 48.6 mg with valsartan 51.4 mg	190.00	56	Entresto 49/51
↓ Tab 97.2 mg with valsartan 102.8 mg	190.00	56	Entresto 97/103
→ Restricted (RS2014)			
Initiation			
All of the following:			
1 Patient has heart failure; and			
2 Any of the following:			
2.1 Patient is in NYHA/WHO functional class II; or			
2.2 Patient is in NYHA/WHO functional class III; or			
2.3 Patient is in NYHA/WHO functional class IV; and			
3 Either:			
3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or			
3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and			
4 Patient is receiving concomitant optimal standard chronic heart failure treatments.			
Alpha-Adrenoceptor Blockers			
DOXAZOSIN			
Tab 2 mg	17.35	500	Doxazosin Clinect
Tab 4 mg	20.94	500	Doxazosin Clinect
PHENOXYBENZAMINE HYDROCHLORIDE			
Cap 10 mg			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
PHENTOLAMINE MESYLATE			
Inj 5 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 1 ml ampoule			
PRAZOSIN			
Tab 1 mg	5.53	100	Arrotex-Prazosin S29
Tab 2 mg	7.00	100	Arrotex-Prazosin S29
Tab 5 mg	11.70	100	Arrotex-Prazosin S29
Cap 1 mg	15.40	100	Prazosin Mylan
Cap 2 mg	15.58	100	Prazosin Mylan
Cap 5 mg	23.32	100	Prazosin Mylan

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

TERAZOSIN – **Restricted**: For continuation only

➔ Tab 1 mg

Antiarrhythmics

ADENOSINE

Inj 3 mg per ml, 2 ml vial – 5% DV Dec-24 to 2027	34.50	5	Adsine
⚡ Inj 3 mg per ml, 10 ml vial – 5% DV Dec-24 to 2027	100.00	5	Adenosine Baxter
➔ Restricted (RS1266)			

Initiation

For use in cardiac catheterisation, electrophysiology and MRI.

AJMALINE – **Restricted** see terms [below](#)

⚡ Inj 5 mg per ml, 10 ml ampoule

➔ **Restricted** (RS1001)

Cardiologist

AMIODARONE HYDROCHLORIDE

Tab 100 mg – 5% DV Feb-26 to 2028	4.95	30	Aratac
Tab 200 mg – 5% DV Feb-26 to 2028	5.86	30	Aratac
Inj 50 mg per ml, 3 ml ampoule – 5% DV Feb-26 to 2028	17.96	10	Max Health

ATROPINE SULPHATE

Inj 600 mcg per ml, 1 ml ampoule – 5% DV Feb-25 to 2027	16.10	10	Hikma Juno Martindale
---	-------	----	-----------------------------

(Juno Inj 600 mcg per ml, 1 ml ampoule to be delisted 1 October 2025)

DIGOXIN

Tab 62.5 mcg – 5% DV Feb-26 to 2028	8.58	240	Lanoxin PG
Tab 250 mcg – 5% DV Feb-26 to 2028	18.75	240	Lanoxin
Oral liq 50 mcg per ml			
Inj 250 mcg per ml, 2 ml vial			

DISOPYRAMIDE PHOSPHATE

Cap 100 mg

FLECAINIDE ACETATE

Tab 50 mg – 5% DV Dec-23 to 2026	19.95	60	Flecainide BNM
Cap long-acting 100 mg – 5% DV Aug-23 to 2026	35.78	90	Flecainide Controlled Release Teva
Cap long-acting 200 mg – 5% DV Aug-23 to 2026	54.28	90	Flecainide Controlled Release Teva
Inj 10 mg per ml, 15 ml ampoule	102.79 108.16	5	Almarytm Tambacor Tambacor German

IVABRADINE – **Restricted** see terms [below](#)

⚡ Tab 5 mg

➔ **Restricted** (RS1566)

Initiation

Both:

- 1 Patient is indicated for computed tomography coronary angiography; and
- 2 Either:
 - 2.1 Patient has a heart rate of greater than 70 beats per minute while taking a maximally tolerated dose of beta blocker; or
 - 2.2 Patient is unable to tolerate beta blockers.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MEXILETINE HYDROCHLORIDE			
Cap 150 mg	162.00	100	Teva
Cap 250 mg	202.00	100	Teva
PROPAFENONE HYDROCHLORIDE			
Tab 150 mg			
Antihypotensives			
MIDODRINE – Restricted see terms below			
⬇ Tab 2.5 mg – 5% DV Feb-25 to 2027	36.68	100	MAR-Midodrine Midodrine Medsurge
⬇ Tab 5 mg – 5% DV Feb-25 to 2027	58.88	100	MAR-Midodrine Midodrine Medsurge
(MAR-Midodrine Tab 2.5 mg to be delisted 1 October 2025)			
(MAR-Midodrine Tab 5 mg to be delisted 1 October 2025)			
➡ Restricted (RS1427)			
Initiation			
Patient has disabling orthostatic hypotension not due to drugs.			
Beta-Adrenoceptor Blockers			
ATENOLOL			
Tab 50 mg – 5% DV Feb-25 to 2027	11.00	500	Viatrix
Tab 100 mg – 5% DV Feb-25 to 2027	18.50	500	Atenolol Viatrix
Oral liq 5 mg per ml	49.85	300 ml	Atenolol-AFT
BISOPROLOL FUMARATE			
Tab 2.5 mg – 5% DV Apr-24 to 2026	1.36	90	Ipca-Bisoprolol
Tab 5 mg – 5% DV Apr-24 to 2026	1.91	90	Ipca-Bisoprolol
Tab 10 mg – 5% DV Apr-24 to 2026	2.71	90	Ipca-Bisoprolol
CARVEDILOL			
Tab 6.25 mg	2.24	60	Carvedilol Sandoz
Tab 12.5 mg	2.30	60	Carvedilol Sandoz
Tab 25 mg	2.95	60	Carvedilol Sandoz
CELIPROLOL – Restricted: For continuation only			
➡ Tab 200 mg			
ESMOLOL HYDROCHLORIDE			
Inj 10 mg per ml, 10 ml vial			
LABELTALOL			
Tab 50 mg			
Tab 100 mg	49.54	100	Biocon
	14.50		Trandate
Tab 200 mg	42.07	100	Presolol
	27.00		Trandate
Inj 5 mg per ml, 20 ml ampoule			
METOPROLOL SUCCINATE			
Tab long-acting 23.75 mg – 5% DV Apr-24 to 2026	4.20	90	Myloc CR
Tab long-acting 47.5 mg – 5% DV Apr-24 to 2026	3.65	90	Myloc CR
Tab long-acting 95 mg – 5% DV Apr-24 to 2026	5.24	90	Myloc CR
Tab long-acting 190 mg – 5% DV Apr-24 to 2026	9.76	90	Myloc CR

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METOPROLOL TARTRATE			
Tab 50 mg – 1% DV Mar-22 to 2027	5.66	100	IPCA-Metoprolol
Tab 100 mg – 1% DV Mar-22 to 2027	7.55	60	IPCA-Metoprolol
Tab long-acting 200 mg.....	23.40	28	Slow-Lopresor
Inj 1 mg per ml, 5 ml vial.....	26.50	5	Metoprolol IV Mylan Metoprolol IV Viatris
NADOLOL			
Tab 40 mg – 1% DV Mar-22 to 2027	19.19	100	Nadolol BNM
Tab 80 mg – 1% DV Mar-22 to 2027	30.39	100	Nadolol BNM
PROPRANOLOL			
Tab 10 mg – 1% DV Mar-22 to 2027	7.04	100	Drofate
Tab 40 mg – 1% DV Mar-22 to 2027	8.75	100	IPCA-Propranolol
Cap long-acting 160 mg	18.17	100	Cardinol LA
Oral liq 4 mg per ml			<i>e.g. Hikma-Propranolol</i>
Inj 1 mg per ml, 1 ml ampoule			
SOTALOL			
Tab 80 mg – 5% DV Feb-26 to 2028	40.00	500	Mylan
	22.50	300	Sotalol Viatris
Tab 160 mg – 5% DV Feb-26 to 2028	20.00	100	Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMLODIPINE			
Tab 2.5 mg – 5% DV Feb-24 to 2026	1.45	90	Vasorex
Tab 5 mg – 5% DV Feb-24 to 2026	1.21	90	Vasorex
Tab 10 mg – 5% DV Feb-24 to 2026	1.31	90	Vasorex
FELODIPINE			
Tab long-acting 2.5 mg – 5% DV Feb-25 to 2027	2.18	30	Plendil ER
Tab long-acting 5 mg – 5% DV Feb-25 to 2027	6.57	90	Felo 5 ER
Tab long-acting 10 mg – 5% DV Feb-25 to 2027	6.95	90	Felo 10 ER

ISRADIPINE

Tab 2.5 mg

Cap 2.5 mg

NICARDIPINE HYDROCHLORIDE – **Restricted** see terms [below](#)

⚡ Inj 2.5 mg per ml, 10 ml vial

➡ **Restricted (RS1699)**

Initiation

Anaesthetist, intensivist, cardiologist or paediatric cardiologist

Any of the following:

- 1 Patient has hypertension requiring urgent treatment with an intravenous agent; or
- 2 Patient has excessive ventricular afterload; or
- 3 Patient is awaiting or undergoing cardiac surgery using cardiopulmonary bypass.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NIFEDIPINE			
Tab long-acting 10 mg.....	19.42	56	Tensipine MR10
Tab long-acting 20 mg.....	17.72	100	Nyefax Retard
Tab long-acting 30 mg.....	34.10	100	Mylan (24 hr release)
	4.78	14	Mylan Italy (24 hr release)
Tab long-acting 60 mg.....	52.81	100	Mylan (24 hr release)
Cap 5 mg			
NIMODIPINE			
Tab 30 mg – 5% DV Feb-26 to 2028	350.00	100	Nimotop
Inj 0.2 mg per ml, 50 ml vial – 5% DV Feb-26 to 2028	337.50	5	Nimotop
Other Calcium Channel Blockers			
DILTIAZEM HYDROCHLORIDE			
Tab 30 mg			
Cap long-acting 120 mg – 5% DV Dec-25 to 2028	65.35	500	Diltiazem CD Clinect
Cap long-acting 180 mg – 1% DV Mar-22 to 2027	7.00	30	Cardizem CD
Cap long-acting 240 mg – 1% DV Mar-22 to 2027	9.30	30	Cardizem CD
Inj 5 mg per ml, 5 ml vial			
PERHEXILINE MALEATE			
Tab 100 mg	62.90	100	Pexsig
VERAPAMIL HYDROCHLORIDE			
Tab 40 mg	7.01	100	Isoptin
Tab 80 mg	11.74	100	Isoptin
Tab long-acting 120 mg.....	36.02	100	Isoptin SR
Tab long-acting 240 mg.....	15.12	30	Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule	25.00	5	Isoptin
Centrally-Acting Agents			
CLONIDINE			
Patch 2.5 mg, 100 mcg per day – 5% DV Feb-24 to 2026	11.70	4	Mylan
Patch 5 mg, 200 mcg per day – 5% DV Feb-24 to 2026	12.80	4	Mylan
Patch 7.5 mg, 300 mcg per day – 5% DV Feb-24 to 2026	17.90	4	Mylan
CLONIDINE HYDROCHLORIDE			
Tab 25 mcg – 5% DV Feb-26 to 2028	29.74	112	Clonidine Teva
Tab 150 mcg – 5% DV Feb-25 to 2027	40.41	100	Catapres
Inj 150 mcg per ml, 1 ml ampoule – 5% DV Jan-25 to 2027	14.10	5	Catapres
METHYLDOPA			
Tab 250 mg	15.10	100	Methyldopa Viatris

Diuretics

Loop Diuretics

BUMETANIDE			
Tab 1 mg	16.36	100	Burinex
Inj 500 mcg per ml, 4 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FUROSEMIDE [FRUSEMIDE]			
Tab 40 mg – 5% DV Feb-25 to 2027	12.80	1,000	IPCA-Frusemide
Tab 500 mg	25.00	50	Urex Forte
Oral liq 10 mg per ml	11.20	30 ml	Lasix
Inj 10 mg per ml, 2 ml ampoule	2.40	5	Furosemide-Baxter
Inj 10 mg per ml, 25 ml ampoule	60.65	6	Lasix

Osmotic Diuretics

MANNITOL			
Inj 10%, 1,000 ml bag	882.84	12	Baxter
Inj 20%, 500 ml bag	1,296.00	18	Baxter

Potassium Sparing Combination Diuretics

AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE			
Tab 5 mg with furosemide 40 mg			
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE			
Tab 5 mg with hydrochlorothiazide 50 mg			

Potassium Sparing Diuretics

AMILORIDE HYDROCHLORIDE			
Tab 5 mg			
Oral liq 1 mg per ml	35.40	25 ml	Biomed
EPLERENONE – Restricted see terms below			
⚡ Tab 25 mg – 5% DV Dec-24 to 2027	15.84	30	Inspra
⚡ Tab 50 mg – 5% DV Dec-24 to 2027	25.00	30	Inspra
➡ Restricted (RS1640)			

Initiation

- Both:
- 1 Patient has heart failure with ejection fraction less than 40%; and
 - 2 Either:
 - 2.1 Patient is intolerant to optimal dosing of spironolactone; or
 - 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

SPIRONOLACTONE			
Tab 25 mg	3.68	100	Spiractin
Tab 100 mg	10.65	100	Spiractin
Oral liq 5 mg per ml	35.70	25 ml	Biomed

Thiazide and Related Diuretics

BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]			
Tab 2.5 mg – 5% DV Mar-24 to 2026	51.50	500	Arrow-Bendrofluzide
Tab 5 mg – 5% DV Mar-24 to 2026	61.00	500	Arrow-Bendrofluzide
CHLOROTHIAZIDE			
Oral liq 50 mg per ml	30.67	25 ml	Biomed
CHLORTALIDONE [CHLORTHALIDONE]			
Tab 25 mg – 5% DV Feb-26 to 2028	6.95	50	Hygroton
INDAPAMIDE			
Tab 2.5 mg – 5% DV Feb-24 to 2026	16.00	90	Dapa-Tabs

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

METOLAZONE
Tab 5 mg

Vasopressin receptor antagonists

TOLVAPTAN – **Restricted** see terms [below](#)

↓ Tab 15 mg	873.50	28	Jinarc
↓ Tab 30 mg	873.50	28	Jinarc
↓ Tab 45 mg + 15 mg	1,747.00	56	Jinarc
↓ Tab 60 mg + 30 mg	1,747.00	56	Jinarc
↓ Tab 90 mg + 30 mg	1,747.00	56	Jinarc

→ **Restricted (RS1930)**

Initiation – autosomal dominant polycystic kidney disease

Renal physician or any relevant practitioner on the recommendation of a renal physician

Re-assessment required after 12 months

All of the following:

- 1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
- 2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 mL/min/1.73 m² at treatment initiation; and
- 3 Either:
 - 3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one-year; or
 - 3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five-year period.

Continuation – autosomal dominant polycystic kidney disease

Renal physician or any relevant practitioner on the recommendation of a renal physician

Re-assessment required after 12 months

Both:

- 1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m²; and
- 2 Patient has not undergone a kidney transplant.

Lipid-Modifying Agents

Fibrates

BEZAFIBRATE

Tab 200 mg – 5% DV Mar-25 to 2027	22.65	90	Bezalip
Tab long-acting 400 mg – 5% DV Mar-25 to 2027	21.54	30	Bezalip Retard

HMG CoA Reductase Inhibitors (Statins)

ATORVASTATIN

Tab 10 mg – 5% DV Dec-24 to 2027	0.31	30	Lorstat
	5.16	500	Lorstat
Tab 20 mg – 5% DV Dec-24 to 2027	8.12	500	Lorstat
Tab 40 mg – 5% DV Dec-24 to 2027	13.79	500	Lorstat
Tab 80 mg – 5% DV Dec-24 to 2027	25.39	500	Lorstat

PRAVASTATIN

Tab 10 mg			
Tab 20 mg – 5% DV May-24 to 2026	7.16	100	Clinect
Tab 40 mg – 5% DV May-24 to 2026	12.25	100	Clinect

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ROSUVASTATIN – Restricted see terms below			
⚡ Tab 5 mg – 5% DV Oct-24 to 2026	1.29	30	Rosuvastatin Viatris
⚡ Tab 10 mg – 5% DV Oct-24 to 2026	1.69	30	Rosuvastatin Viatris
⚡ Tab 20 mg – 5% DV Apr-24 to 2026	2.71	30	Rosuvastatin Viatris
⚡ Tab 40 mg – 5% DV Apr-24 to 2026	4.55	30	Rosuvastatin Viatris

➡ **Restricted (RS1868)**

Initiation – cardiovascular disease risk

Either:

- 1 Both:
 - 1.1 Patient is considered to be at risk of cardiovascular disease; and
 - 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:
 - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
 - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initiation – familial hypercholesterolemia

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initiation – established cardiovascular disease

Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initiation – recurrent major cardiovascular events

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIMVASTATIN

Tab 10 mg – 5% DV Mar-24 to 2026	1.68	90	Simvastatin Mylan Simvastatin Viatris
Tab 20 mg – 5% DV Mar-24 to 2026	2.54	90	Simvastatin Viatris
Tab 40 mg – 5% DV Jun-24 to 2026	4.11	90	Simvastatin Viatris
Tab 80 mg – 5% DV Jun-24 to 2026	8.81	90	Simvastatin Viatris

Resins

CHOLESTYRAMINE

Powder for oral liq 4 g

COLESTIPOL HYDROCHLORIDE

Grans for oral liq 5 g

COLESTYRAMINE

Powder for oral suspension 4 g sachet	61.50	50	Colestyramine - Mylan
---	-------	----	-----------------------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Selective Cholesterol Absorption Inhibitors

EZETIMIBE			
Tab 10 mg – 5% DV Dec-23 to 2026	1.76	30	Ezetimibe Sandoz
EZETIMIBE WITH SIMVASTATIN			
Tab 10 mg with simvastatin 10 mg	5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	Zimybe
Tab 10 mg with simvastatin 40 mg	7.15	30	Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	Zimybe

Other Lipid-Modifying Agents

ACIPIMOX
Cap 250 mg

Nitrates

GLYCERYL TRINITRATE			
Inj 1 mg per ml, 5 ml ampoule			
Inj 1 mg per ml, 10 ml ampoule			
Inj 1 mg per ml, 50 ml vial			
Inj 5 mg per ml, 10 ml ampoule	118.00	5	Hospira
Oral pump spray, 400 mcg per dose	7.48	250 dose	Nitrolingual Pump Spray
Patch 25 mg, 5 mg per day	15.73	30	Nitroderm TTS 5
Patch 50 mg, 10 mg per day	18.62	30	Nitroderm TTS 10
ISOSORBIDE MONONITRATE			
Tab 20 mg – 5% DV Feb-24 to 2026	22.49	100	Ismo 20
Tab long-acting 40 mg – 5% DV Feb-24 to 2026	9.80	30	Ismo 40 Retard
Tab long-acting 60 mg – 5% DV Feb-24 to 2026	13.50	90	Duride

Other Cardiac Agents

LEVOSIMENDAN – Restricted see terms below			
⚠ Inj 2.5 mg per ml, 5 ml vial – 5% DV Nov-24 to 2027	509.60	1	Simdax
⚠ Inj 2.5 mg per ml, 10 ml vial			

➡ **Restricted (RS1007)**
Initiation – Heart transplant
Either:

- 1 For use as a bridge to heart transplant, in patients who have been accepted for transplant; or
- 2 For the treatment of heart failure following heart transplant.

Initiation – Heart failure
Cardiologist or intensivist
For the treatment of severe acute decompensated heart failure that is non-responsive to dobutamine.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Sympathomimetics			
ADRENALINE			
Inj 1 in 1,000, 1 ml ampoule	4.98	5	Aspen Adrenaline
	13.27		DBL Adrenaline
	25.30	10	Hameln
Inj 1 in 1,000, 30 ml vial			
Inj 1 in 10,000, 10 ml ampoule	49.00	10	Aspen Adrenaline
	27.00	5	Hospira
Inj 1 in 10,000, 10 ml syringe			
<i>(Hameln Inj 1 in 1,000, 1 ml ampoule to be delisted 1 October 2025)</i>			
DOBUTAMINE			
Inj 12.5 mg per ml, 20 ml ampoule – 5% DV Dec-24 to 2027	61.13	5	Dobutamine-hameln
DOPAMINE HYDROCHLORIDE			
Inj 40 mg per ml, 5 ml ampoule – 5% DV Feb-25 to 2027	46.38	10	Dopamine Basi Max Health Ltd
EPHEDRINE			
Inj 3 mg per ml, 10 ml syringe – 5% DV Aug-25 to 2026	142.00	10	Ephedrine Aguettant Ephedrine Juno
		10	Max Health
Inj 30 mg per ml, 1 ml ampoule – 5% DV Feb-24 to 2026	34.31		
ISOPRENALINE [ISOPROTERENOL]			
Inj 200 mcg per ml, 1 ml ampoule			
Inj 200 mcg per ml, 5 ml ampoule			
METARAMINOL			
Inj 0.5 mg per ml, 10 ml syringe			
Inj 0.5 mg per ml, 20 ml syringe			
Inj 0.5 mg per ml, 5 ml syringe			
Inj 1 mg per ml, 1 ml ampoule			
Inj 1 mg per ml, 10 ml syringe			
Inj 10 mg per ml, 1 ml ampoule – 5% DV Feb-24 to 2026	53.00	10	Torbay
NORADRENALINE			
Inj 0.06 mg per ml, 100 ml bag			
Inj 0.06 mg per ml, 50 ml syringe			
Inj 0.1 mg per ml, 100 ml bag			
Inj 0.1 mg per ml, 50 ml syringe			
Inj 0.12 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 50 ml syringe			
Inj 0.16 mg per ml, 50 ml syringe			
Inj 1 mg per ml, 100 ml bag			
Inj 1 mg per ml, 4 ml ampoule – 5% DV Apr-26 to 2028	45.00	10	Noradrenaline BNM
	32.78		Noradrenaline Medsurge
<i>(Noradrenaline BNM Inj 1 mg per ml, 4 ml ampoule to be delisted 1 April 2026)</i>			
PHENYLEPHRINE HYDROCHLORIDE			
Inj 10 mg per ml, 1 ml ampoule	310.42	25	Neosynephrine HCL

Vasodilators

ALPROSTADIL – **Restricted** see terms [on the next page](#)

⚡ Inj 10 mcg vial

⚡ Inj 20 mcg vial

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted (RS1992)			
Initiation			
Both:			
1 Patient has erectile dysfunction; and			
2 Patient is to receive a penile Doppler ultrasonography.			
ALPROSTADIL HYDROCHLORIDE			
Inj 500 mcg per ml, 1 ml ampoule	2,030.33	5	Prostin VR
DIAZOXIDE			
Inj 15 mg per ml, 20 ml ampoule			
HYDRALAZINE HYDROCHLORIDE			
↓ Tab 25 mg			
➔ Restricted (RS1008)			
Initiation			
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure, in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.			
Inj 20 mg ampoule	25.90	5	Apresoline
MILRINONE			
Inj 1 mg per ml, 10 ml ampoule – 5% DV Dec-24 to 2027	68.00	10	Milrinone-Baxter
MINOXIDIL			
Tab 10 mg	78.40	100	Loniten
NICORANDIL			
Tab 10 mg – 5% DV Feb-26 to 2028	27.81	60	Max Health
Tab 20 mg – 5% DV Feb-26 to 2028	35.12	60	Max Health
PAPAVERINE HYDROCHLORIDE			
Inj 30 mg per ml, 1 ml vial			
Inj 12 mg per ml, 10 ml ampoule	257.12	5	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg			
SODIUM NITROPRUSSIDE			
Inj 50 mg vial			

Endothelin Receptor Antagonists

AMBRISENTAN – Restricted see terms below			
↓ Tab 5 mg – 5% DV Dec-23 to 2026	200.00	30	Ambrisentan Viatris
↓ Tab 10 mg – 5% DV Dec-23 to 2026	200.00	30	Ambrisentan Viatris
➔ Restricted (RS2121)			
Initiation – PAH monotherapy			
Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist			
<i>Limited to 6 months treatment</i>			
All of the following:			
1 Patient has pulmonary arterial hypertension (PAH); and			
2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and			

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
 - 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH monotherapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH dual therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient has tried bosentan (either as PAH monotherapy, or PAH dual therapy with sildenafil) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; or
 - 5.2.2 Patient has experienced intolerable side effects on bosentan; or
 - 5.2.3 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
 - 5.2.4 Patient is presenting in NYHA/WHO functional class III or IV, and would benefit from initial dual therapy in the opinion of the treating clinician and has an absolute or relative contraindication to bosentan (eg. due to current liver disease or use of a combined oral contraceptive).

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Ambrisentan is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

5.2.2 Both:

5.2.2.1 Patient is presenting in NYHA/WHO functional class IV; and

5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or

5.2.3 Both:

5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**;

5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

The patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: [2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH](#)

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN – **Restricted** see terms [below](#)

⚡ Tab 62.5 mg – 5% DV Jan-25 to 2027 100.00

60

Bosentan Dr Reddy's

⚡ Tab 125 mg – 5% DV Jan-25 to 2027 100.00

60

Bosentan Dr Reddy's

➡ **Restricted (RS1982)**

Initiation – PAH monotherapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH)*; and

2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and

3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and

4 Any of the following:

4.1 All of the following:

4.1.1 PAH has been confirmed by right heart catheterisation; and

4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and

4.1.5 Any of the following:

4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or

4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**;

4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or

4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:

- 5.1 Bosentan is to be used as PAH monotherapy; and
- 5.2 Any of the following:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute contraindication to sildenafil; or
 - 5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as part of PAH dual therapy; and
 - 5.2 Either:
 - 5.2.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool**; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy.

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Bosentan is to be used as part of PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

Patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**.

Notes: † The European Respiratory Journal Guidelines can be found here: [2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH](#)

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – **Restricted** see terms [on the next page](#)

⚡ Tab 25 mg – 5% DV Dec-24 to 2027	0.72	4	Vedafil
⚡ Tab 50 mg – 5% DV Dec-24 to 2027	1.45	4	Vedafil
⚡ Tab 100 mg – 5% DV Dec-24 to 2027	11.22	12	Vedafil
⚡ Inj 0.8 mg per ml, 12.5 ml vial			

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

➔ **Restricted (RS1983)**

Initiation – tablets Raynaud's Phenomenon

All of the following:

- 1 Patient has Raynaud's phenomenon; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (proper body insulation, avoidance of cold exposure, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient has persisting severe symptoms despite treatment with calcium channel blockers and nitrates (unless contraindicated or not tolerated).

Initiation – tablets Pulmonary arterial hypertension

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH is confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
 - 4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or at least 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

Initiation – tablets other conditions

Any of the following:

- 1 For use in weaning patients from inhaled nitric oxide; or
- 2 For perioperative use in cardiac surgery patients; or
- 3 For use in intensive care as an alternative to nitric oxide; or
- 4 For use in the treatment of erectile dysfunction secondary to spinal cord injury in patients being treated in a spinal unit.

Initiation – injection

Both:

- 1 For use in the treatment of pulmonary hypertension in infants or children being treated in paediatric intensive care units and neonatal intensive care units when the enteral route is not accessible; and
- 2 Any of the following:
 - 2.1 For perioperative use following cardiac surgery; or
 - 2.2 For use in persistent pulmonary hypertension of the newborn (PPHN); or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.3 For use in congenital diaphragmatic hernia.

Notes: † The European Respiratory Journal Guidelines can be found here: [2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH](#)

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Prostacyclin Analogues

EPOPROSTENOL – **Restricted** see terms [below](#)

⚡ Inj 500 mcg vial.....	36.61	1	Veletri
⚡ Inj 1.5 mg vial	73.21	1	Veletri

➔ **Restricted (RS1984)**

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † ; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 All of the following:
 - 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Patient is presenting in NYHA/WHO functional class IV; and
 - 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist,

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...
cardiologist or rheumatologist
Limited to 6 months treatment
All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † ; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**;
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Epoprostenol is to be used as PAH triple therapy; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is on the lung transplant list; or
 - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
 - 5.2.3 Both:
 - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
 - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

Patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool.

Notes: † The European Respiratory Journal Guidelines can be found here: [2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH](#)

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST

Inj 50 mcg in 0.5 ml ampoule.....	380.00	5	Ilomedin
⚡ Nebuliser soln 10 mcg per ml, 2 ml – 5% DV Dec-25 to 2028	166.53	30	Veblis

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

➔ Restricted (RS1985)

Initiation – PAH monotherapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) †; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool*; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
 - 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
 - 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
- 5 Both:
 - 5.1 Iloprost is to be used as PAH monotherapy; and
 - 5.2 Either:
 - 5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
 - 5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
- 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † ; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and
- 5 All of the following:
 - 5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
 - 5.2 Either:
 - 5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or
 - 5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and
 - 5.3 Either:
 - 5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; or
 - 5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH); and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
- 3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
- 4 Any of the following:
 - 4.1 All of the following:
 - 4.1.1 PAH has been confirmed by right heart catheterisation; and
 - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
 - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
 - 4.1.5 Any of the following:
 - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH (see note below for link to these guidelines) † ; or
 - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
 - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

developmental lung disorders including severe chronic neonatal lung disease; or

- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and

5 Both:

- 5.1 Iloprost is to be used as PAH triple therapy; and

- 5.2 Any of the following:

5.2.1 Patient is on the lung transplant list; or

5.2.2 Patient is presenting in NYHA/WHO functional class IV; or

5.2.3 Both:

5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and

5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

Patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool.

Notes: † The European Respiratory Journal Guidelines can be found here: [2022 ECS/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension PAH](#)

** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anti-Infective Preparations			
Antibacterials			
HYDROGEN PEROXIDE			
Crm 1% – 5% DV Jan-26 to 2028	4.89	15 g	Crystaderm
Soln 3% (10 vol)			
MAFENIDE ACETATE – Restricted see terms below			
↓ Crm 8.5%			
↓ Powder 5%			
➔ Restricted (RS1299)			
Initiation			
For the treatment of burns patients.			
MUPIROCIN			
Oint 2%			
SODIUM FUSIDATE [FUSIDIC ACID]			
Crm 2% – 5% DV Feb-25 to 2027	1.69	5 g	Foban
Oint 2% – 5% DV Feb-25 to 2027	1.69	5 g	Foban
SULFADIAZINE SILVER			
Crm 1%.....	10.80	50 g	Flamazine
Antifungals			
AMOROLFINE			
Nail soln 5% – 5% DV Feb-24 to 2026	21.87	5 ml	MycoNail
CICLOPIROX OLAMINE			
Nail soln 8%			
➔ Soln 1% – Restricted: For continuation only			
CLOTRIMAZOLE			
Crm 1%.....	1.10	20 g	Clomazol
➔ Soln 1% – Restricted: For continuation only			
ECONAZOLE NITRATE			
Crm 1% – 5% DV Jun-25 to 2027	8.04	20 g	Pevaryl
Foaming soln 1%			
KETOCONAZOLE			
Shampoo 2% – 5% DV May-24 to 2026	4.09	100 ml	Sebizole
METRONIDAZOLE			
Gel 0.75%			
MICONAZOLE NITRATE			
Crm 2% – 5% DV May-24 to 2026	0.90	15 g	Multichem
➔ Lotn 2% – Restricted: For continuation only			
Tinc 2%			
NYSTATIN			
Crm 100,000 u per g			
Antiparasitics			
DIMETHICONE			
Lotn 4%	4.25	200 ml	healthE Dimethicone 4% Lotion

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MALATHION [MALDISON] Lotn 0.5% Shampoo 1%			
PERMETHRIN Lotn 5% – 5% DV Feb-24 to 2026	4.28	30 ml	A-Scabies
PHENOTHRIN Shampoo 0.5%			

Antiacne Preparations

ADAPALENE Crm 0.1% Gel 0.1%			
BENZOYL PEROXIDE Soln 5%			
ISOTRETINOIN Cap 5 mg – 5% DV Dec-24 to 2027	11.26	60	Oratane
Cap 10 mg – 5% DV Dec-24 to 2027	18.75	120	Oratane
Cap 20 mg – 5% DV Dec-24 to 2027	26.73	120	Oratane
TRETINOIN Crm 0.05% – 5% DV Feb-25 to 2027	16.82	50 g	ReTrieve

Antipruritic Preparations

CALAMINE Crm, aqueous, BP – 5% DV Apr-25 to 2027	3.45	100 g	healthE Calamine Aqueous
CROTAMITON Crm 10% – 5% DV Feb-25 to 2027	3.49	20 g	Itch-Soothe

Barrier Creams and Emollients

Barrier Creams

DIMETHICONE Crm 10% pump bottle	4.52	460 g	healthE Dimethicone 10%
Crm 5% pump bottle	4.30	460 g	healthE Dimethicone 5%
Crm 5% tube	1.47	100 g	healthE Dimethicone 5%
ZINC Crm			<i>e.g. Zinc Cream (Orion-) ;Zinc Cream (PSM)</i>
Oint Paste			<i>e.g. Zinc oxide (PSM)</i>
ZINC AND CASTOR OIL Crm	1.63	20 g	Orion
Oint	4.25	500 g	Evvara
Note: DV limit applies to the pack sizes of greater than 30 g.			
Oint, BP	1.26	20 g	healthE
Note: DV limit applies to the pack sizes of 30 g or less.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ZINC WITH WOOL FAT			
Crm zinc 15.25% with wool fat 4%			<i>e.g. Sudocrem</i>
Emollients			
AQUEOUS CREAM			
Crm 100 g – 5% DV Mar-25 to 2027	1.25	100 g	Evvara
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm 500 g – 5% DV Mar-25 to 2027	1.65	500 g	Evvara
Note: DV limit applies to the pack sizes of greater than 100 g.			
CETOMACROGOL			
Crm BP, 100 g – 5% DV Jun-25 to 2027	0.99	100 g	Cetomacrogol Cream
			AFT
Crm BP, 500 g – 5% DV Feb-25 to 2027	2.29	500 g	Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL			
Crm 90% with glycerol 10% – 5% DV Dec-25 to 2028	1.92	460 g	Evvara
	3.25	920 g	Evvara
Note: DV limit applies to the pack sizes of greater than 100 g.			
Crm 90% with glycerol 10%,	1.65	100 g	healthE
Note: DV limit applies to the pack sizes of 100 g or less.			
EMULSIFYING OINTMENT			
Oint BP – 5% DV Feb-24 to 2026	2.30	100 g	Jaychem
Note: DV limit applies to pack sizes of less than 200 g.			
Oint BP, 500 g – 5% DV May-24 to 2026	3.13	500 g	Evvara Emulsifying Ointment
Note: DV limit applies to pack sizes of greater than 200 g.			
GLYCEROL WITH PARAFFIN			
Crm glycerol 10% with white soft paraffin 5% and liquid paraffin 10%			<i>e.g. QV cream</i>
OIL IN WATER EMULSION			
Crm, 100 g – 5% DV Apr-25 to 2027	1.43	100 g	Fatty Emulsion Cream (Evvara)
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm, 500 g – 5% DV Apr-25 to 2027	2.10	500 g	Fatty Emulsion Cream (Evvara)
Note: DV limit applies to the pack sizes of greater than 100 g.			
PARAFFIN			
Oint liquid paraffin 50% with white soft paraffin 50%.....	1.84	100 g	White Soft Liquid Paraffin AFT
Note: DV limit applies to the pack sizes of 100 g or less.			
White soft.....	0.79	10 g	healthE
Note: DV limit applies to pack sizes of 30 g or less, and to both white soft paraffin and yellow soft paraffin.			
White soft, – 5% DV Jun-24 to 2026	4.74	450 g	EVARA White Soft Paraffin
Note: DV limit applies to the pack sizes of 500 g or less and greater than 30 g.			
Yellow soft			
Lotn liquid paraffin 85%			<i>e.g. QV Bath Oil</i>
PARAFFIN WITH WOOL FAT			
Lotn liquid paraffin 15.9% with wool fat 0.6%			<i>e.g. AlphaKeri;BK ;DP; Hydroderm Lotn</i>
Lotn liquid paraffin 91.7% with wool fat 3%			<i>e.g. Alpha Keri Bath Oil</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
UREA			
Crm 10%.....	1.37	100 g	healthE Urea Cream
WOOL FAT			
Crm			
Corticosteroids			
BETAMETHASONE DIPROPIONATE			
Crm 0.05% – 5% DV Jul-24 to 2026	36.00	50 g	Diprosone
Note: DV limit applies to the pack sizes of greater than 30 g.			
Oint 0.05% – 5% DV Jul-24 to 2026	36.00	50 g	Diprosone
Note: DV limit applies to the pack sizes of greater than 30 g.			
BETAMETHASONE VALERATE			
Crm 0.1% – 5% DV Feb-25 to 2027	5.85	50 g	Beta Cream
Oint 0.1% – 5% DV Feb-25 to 2027	7.90	50 g	Beta Ointment
Lotn 0.1% – 5% DV May-25 to 2027	30.00	50 ml	Betnovate
CLOBETASOL PROPIONATE			
Crm 0.05% – 5% DV Feb-26 to 2028	3.75	30 g	Dermol
Oint 0.05% – 5% DV Feb-26 to 2028	3.68	30 g	Dermol
CLOBETASONE BUTYRATE			
Crm 0.05%			
DIFLUCORTOLONE VALERATE – Restricted: For continuation only			
➡ Crm 0.1%			
➡ Fatty oint 0.1%			
HYDROCORTISONE			
Crm 1%, 30 g.....	1.78	30 g	Ethics
Note: DV limit applies to the pack sizes of less than or equal to 100 g.			
Crm 1%, 500 g – 5% DV Feb-26 to 2028	20.40	500 g	Noumed
Note: DV limit applies to the pack sizes of greater than 100 g.			
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – 5% DV Jun-24 to 2026	12.83	250 ml	DP Lotn HC
HYDROCORTISONE BUTYRATE			
Crm 0.1%.....	4.85	100 g	Locoid Lipocream
Oint 0.1%.....	10.28	100 g	Locoid
Milky emul 0.1%	12.33	100 ml	Locoid Crelo
METHYLPREDNISOLONE ACEPONATE			
Crm 0.1% – 5% DV Feb-24 to 2026	4.95	15 g	Advantan
Oint 0.1% – 5% DV Feb-24 to 2026	4.95	15 g	Advantan
MOMETASONE FUROATE			
Crm 0.1% – 5% DV Feb-25 to 2027	2.25	15 g	Elocon Alcohol Free
	3.50	50 g	Elocon Alcohol Free
Oint 0.1% – 5% DV Feb-25 to 2027	2.25	15 g	Elocon
	3.50	50 g	Elocon
Lotn 0.1% – 5% DV Feb-25 to 2027	4.99	30 ml	Elocon
TRIAMCINOLONE ACETONIDE			
Crm 0.02% – 5% DV Feb-24 to 2026	6.49	100 g	Aristocort
Oint 0.02% – 5% DV Feb-24 to 2026	6.54	100 g	Aristocort

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

Corticosteroids with Anti-Infective Agents

BETAMETHASONE VALERATE WITH CLIOQUINOL – **Restricted** see terms [below](#)

↓ Crm 0.1% with clioquinol 3%

→ **Restricted (RS1125)**

Initiation

Either:

- 1 For the treatment of intertrigo; or
- 2 For continuation use.

BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID]

Crm 0.1% with sodium fusidate (fusidic acid) 2%

HYDROCORTISONE WITH MICONAZOLE

Crm 1% with miconazole nitrate 2% – **5% DV Feb-25 to 2027** 2.85 15 g **Micreme H**

HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN

Oint 1% with natamycin 1% and neomycin sulphate 0.5%..... 4.34 15 g Pimafucort

TRIAMCINOLONE ACETONIDE WITH NEOMYCIN SULPHATE, GRAMICIDIN AND NYSTATIN

Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and
gramicidin 250 mcg per g

Psoriasis and Eczema Preparations

ACITRETIN

Cap 10 mg – **5% DV Jul-24 to 2026** 26.20 60 **Novatretin**

Cap 25 mg – **5% DV Jul-24 to 2026** 57.37 60 **Novatretin**

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per g 59.95 60 g Enstilar

Gel 500 mcg with calcipotriol 50 mcg per g – **5% DV Dec-24 to 2027** 40.92 60 g **Daivobet**

Oint 500 mcg with calcipotriol 50 mcg per g – **5% DV Dec-24 to 2027** 14.31 30 g **Daivobet**

CALCIPOTRIOL

Oint 50 mcg per g 40.00 120 g Daivonex

COAL TAR WITH SALICYLIC ACID AND SULPHUR

Oint 12% with salicylic acid 2% and sulphur 4%

METHOXSALEN [8-METHOXYPsorALEN]

Tab 10 mg

Lotn 1.2%

PIMECROLIMUS – **Restricted** see terms [below](#)

↓ Crm 1% – **5% DV Feb-24 to 2026** 33.00 15 g **Elidel**

→ **Restricted (RS1781)**

Initiation

Dermatologist, paediatrician or ophthalmologist

Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN

Soln 2.3% with trolamine laurilsulfate and fluorescein sodium – **5% DV
Feb-24 to 2026** 5.41 500 ml **Pinetarsol**

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

POTASSIUM PERMANGANATE

Tab 400 mg
Crystals

TACROLIMUS

⚠ Oint 0.1% – 5% DV Dec-23 to 202633.00 30 g **Zematop**
➡ **Restricted (RS1859)**

Initiation

Dermatologist or paediatrician

Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

BETAMETHASONE VALERATE

Scalp app 0.1% – 5% DV Feb-25 to 202712.95 100 ml **Beta Scalp**

CLOBETASOL PROPIONATE

Scalp app 0.05% – 5% DV Feb-26 to 20286.90 30 ml **Dermol**

HYDROCORTISONE BUTYRATE

Scalp lotn 0.1%.....6.57 100 ml Locoid

Wart Preparations

PODOPHYLLOTOXIN

Soln 0.5%33.60 3.5 ml Condylina

SILVER NITRATE

Sticks with applicator

Other Skin Preparations

DIPHENYL METILSULFATE

Powder 2%

IMIQUIMOD

Crm 5%, 250 mg sachet21.72 24 Perrigo

SUNSCREEN, PROPRIETARY

Lotn.....6.50 200 g Marine Blue Lotion SPF 50+

Antineoplastics

FLUOROURACIL SODIUM

Crm 5% – 5% DV Dec-24 to 20275.56 20 g **Efudix**

METHYL AMINOLEVULINATE HYDROCHLORIDE – **Restricted** see terms [below](#)

⚠ Crm 16%

➡ **Restricted (RS1127)**

Dermatologist or plastic surgeon

Wound Management Products

CALCIUM GLUCONATE

Gel 2.5%e.g. Orion

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anti-Infective Agents			
ACETIC ACID			
Soln 3%			
Soln 5%			
ACETIC ACID WITH HYDROXYQUINOLINE, GLYCEROL AND RICINOLEIC ACID			
Jelly 0.94% with hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator			
CHLORHEXIDINE GLUCONATE			
Crm 1%			
Lotn 1%			
CLOTTRIMAZOLE			
Vaginal crm 1% with applicator	3.50	35 g	Clomazol
Vaginal crm 2% with applicator	3.85	20 g	Clomazol
MICONAZOLE NITRATE			
Vaginal crm 2% with applicator	6.89	40 g	Micreme
NYSTATIN			
Vaginal crm 100,000 u per 5 g with applicator(s) – 5% DV Feb-24 to 2026	5.70	75 g	Nilstat
Contraceptives			
Antiandrogen Oral Contraceptives			
CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL			
Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tablets – 5% DV Feb-24 to 2026	5.08	168	Ginet
Combined Oral Contraceptives			
ETHINYLOESTRADIOL WITH DESOGESTREL			
Tab 20 mcg with desogestrel 150 mcg			
Tab 30 mcg with desogestrel 150 mcg			
ETHINYLOESTRADIOL WITH LEVONORGESTREL			
Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	1.50	84	Lo-Oralcon 20 ED
Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	1.50	84	Oralcon 30 ED
Tab 20 mcg with levonorgestrel 100 mcg			
Tab 30 mcg with levonorgestrel 150 mcg			
ETHINYLOESTRADIOL WITH NORETHISTERONE			
Tab 35 mcg with norethisterone 1 mg			
Tab 35 mcg with norethisterone 1 mg and 7 inert tab	12.25	84	Alyacen Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg			
NORETHISTERONE WITH MESTRANOL			
Tab 1 mg with mestranol 50 mcg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Contraceptive Devices			
INTRA-UTERINE DEVICE			
IUD 29.1 mm length x 23.2 mm width	29.80	1	Choice 380 7med Nsha Silver/copper Short
IUD 33.6 mm length x 29.9 mm width	26.80	1	TCu 380 Plus Normal
IUD 35.5 mm length x 19.6 mm width	33.00	1	Cu 375 Standard
Emergency Contraception			
LEVONORGESTREL			
Tab 1.5 mg	1.75	1	Levonorgestrel BNM
Progestogen-Only Contraceptives			
DESOGESTREL			
Tab 75 mcg.....	24.50	84	Cerazette
LEVONORGESTREL			
Tab 30 mcg.....	22.00	112	Microlut
Intra-uterine device 52 mg.....	269.50	1	Mirena
Intra-uterine device 13.5 mg.....	215.60	1	Jaydess
Subdermal implant (2 x 75 mg rods) – 5% DV Apr-25 to 2026	106.92	2	Jadelle
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe	10.56	1	Depo-Provera
NORETHISTERONE			
Tab 350 mcg.....	12.25	84	Norethindrone - CDC Noriday Noriday 28
Obstetric Preparations			
Antiprogestogens			
MIFEPRISTONE			
Tab 200 mg			
Oxytocics			
CARBOPROST TROMETAMOL			
Inj 250 mcg per ml, 1 ml ampoule			
DINOPROSTONE			
Pessaries 10 mg			
Vaginal gel 1 mg in 3 g	65.39	1	Prostin E2
Vaginal gel 2 mg in 3 g	82.33	1	Prostin E2
ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule	160.00	5	DBL Ergometrine
OXYTOCIN			
Inj 5 iu per ml, 1 ml ampoule	4.98	5	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	5.98	5	Oxytocin BNM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

OXYTOCIN WITH ERGOMETRINE MALEATE

Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	41.47	5	Syntometrine
--	-------	---	--------------

Tocolytics

PROGESTERONE

Cap 100 mg.....	14.85	30	Utrogestan
-----------------	-------	----	------------

TERBUTALINE – **Restricted** see terms [below](#)

↓ Inj 500 mcg ampoule

→ **Restricted (RS1130)**

Obstetrician

Oestrogens

OESTRIOL

Crm 1 mg per g with applicator – 5% DV Feb-24 to 2026	6.95	15 g	Ovestin
Pessaries 500 mcg – 5% DV Feb-24 to 2026	7.55	15	Ovestin

Urologicals

5-Alpha Reductase Inhibitors

FINASTERIDE – **Restricted** see terms [below](#)

↓ Tab 5 mg – **5% DV Dec-23 to 2026** 4.79 100 Ricit

→ **Restricted (RS1131)**

Initiation

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoceptor Blockers

TAMSULOSIN HYDROCHLORIDE – **Restricted** see terms [below](#)

↓ Cap 400 mcg – **5% DV Feb-26 to 2028** 28.56 100 Tamsulosin-Rex

→ **Restricted (RS1132)**

Initiation

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Urinary Alkalisers

POTASSIUM CITRATE – **Restricted** see terms [below](#)

↓ Oral liq 3 mmol per ml 37.49 200 ml Biomed

→ **Restricted (RS1133)**

Initiation

Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

SODIUM CITRO-TARTRATE			
Grans eff 4 g sachets – 5% DV Feb-24 to 2026	3.50	28	Ural

Urinary Antispasmodics

OXYBUTYNIN			
Tab 5 mg	5.42	100	Alchemy Oxybutynin
Oral liq 5 mg per 5 ml			

SOLIFENACIN SUCCINATE			
Tab 5 mg – 5% DV Jun-25 to 2027	3.15	30	Solifenacin Viatris
	1.95		Solifenacin succinate
			Max Health
Tab 10 mg – 5% DV Jun-25 to 2027	3.53	30	Solifenacin succinate
			Max Health

(Solifenacin Viatris Tab 5 mg to be delisted 1 November 2025)

	Price (ex man. excl. GST)		Brand or Generic Manufacturer
	\$	Per	

Anabolic Agents

OXANDROLONE

↓ Tab 2.5 mg

➔ **Restricted (RS1302)**

Initiation

For the treatment of burns patients.

Androgen Agonists and Antagonists

CYPROTERONE ACETATE

Tab 50 mg – 5% DV Jul-25 to 2027	17.05	50	Siterone
Tab 100 mg – 5% DV Jul-25 to 2027	31.00	50	Siterone

TESTOSTERONE

Gel (transdermal) 16.2 mg per g, 88 g – 5% DV Apr-25 to 2027	52.00	60	Testogel
--	-------	----	-----------------

TESTOSTERONE CIPIONATE

Inj 100 mg per ml, 10 ml vial.....	85.00	1	Depo-Testosterone
------------------------------------	-------	---	--------------------------

TESTOSTERONE ESTERS

Inj testosterone decanoate 100 mg, testosterone isocaproate 60 mg,
testosterone phenylpropionate 60 mg and testosterone propionate
30 mg per ml, 1 ml ampoule

TESTOSTERONE UNDECANOATE

➔ Cap 40 mg – **Restricted:** For continuation only

Inj 250 mg per ml, 4 ml vial.....	86.00	1	Reandron 1000
-----------------------------------	-------	---	----------------------

Calcium Homeostasis

CALCITONIN

Inj 100 iu per ml, 1 ml ampoule	121.00	5	Miacalcic
---------------------------------------	--------	---	------------------

CINACALCET – **Restricted** see terms [below](#)

↓ Tab 30 mg – 5% DV Dec-24 to 2027	25.24	28	Cinacalcet Devatis
↓ Tab 60 mg – 5% DV Dec-24 to 2027	50.47	28	Cinacalcet Devatis

➔ **Restricted (RS1931)**

Initiation – parathyroid carcinoma or calciphylaxis

Nephrologist or endocrinologist

Re-assessment required after 6 months

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

thiosulfate.

Continuation – parathyroid carcinoma or calciphylaxis

Nephrologist or endocrinologist

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initiation – primary hyperparathyroidism

All of the following:

- 1 Patient has primary hyperparathyroidism; and
- 2 Either:
 - 2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
 - 2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
- 3 Surgery is not feasible or has failed; and
- 4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initiation – secondary or tertiary hyperparathyroidism

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
 - 1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
- 2 Patient is on renal replacement therapy; and
- 3 Any of the following:
 - 3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or
 - 3.2 Parathyroid tissue is surgically inaccessible; or
 - 3.3 Parathyroid surgery is not feasible.

Continuation – secondary or tertiary hyperparathyroidism

Re-assessment required after 12 months

Either:

- 1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
- 2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial – 5% DV Dec-24 to 2027	15.65	1	Zoledronic Acid Injection Mylan Zoledronic acid Viatrix
--	-------	---	---

Corticosteroids

BETAMETHASONE

- Tab 500 mcg
Inj 4 mg per ml, 1 ml ampoule

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

- Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml ampoule

DEXAMETHASONE

Tab 0.5 mg – 5% DV Feb-25 to 2027	1.80	30	Dexamethsone
Tab 4 mg – 5% DV Feb-25 to 2027	3.18	30	Dexamethsone
Oral liq 1 mg per ml	53.86	25 ml	Biomed

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DEXAMETHASONE PHOSPHATE			
Inj 4 mg per ml, 1 ml ampoule	7.86	10	Hameln
Inj 4 mg per ml, 2 ml ampoule	13.10	10	Hameln
FLUDROCORTISONE ACETATE			
Tab 100 mcg – 5% DV Dec-25 to 2028	8.05	100	Florinef
HYDROCORTISONE			
Tab 5 mg	8.10	100	Douglas
Tab 20 mg	20.32	100	Douglas
Inj 100 mg vial – 5% DV Dec-24 to 2027	3.96	1	Solu-Cortef
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Tab 4 mg	112.00	100	Medrol
Tab 100 mg	223.10	20	Medrol
Inj 40 mg vial	22.30	1	Solu-Medrol Act-O-Vial
Inj 125 mg vial	34.10	1	Solu-Medrol Act-O-Vial
Inj 500 mg vial	43.01	1	Solu-Medrol Act-O-Vial
Inj 1 g vial	52.54	1	Solu-Medrol
METHYLPREDNISOLONE ACETATE			
Inj 40 mg per ml, 1 ml vial.....	47.06	5	Depo-Medrol
PREDNISOLONE			
Oral liq 5 mg per ml – 5% DV Dec-24 to 2027	6.00	30 ml	Redipred
Enema 200 mcg per ml, 100 ml			
PREDNISONE			
Tab 1 mg	18.58	500	Prednisone Clinect
Tab 2.5 mg	21.04	500	Prednisone Clinect
Tab 5 mg	19.30	500	Prednisone Clinect
Tab 20 mg	50.51	500	Prednisone Clinect
TRIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule – 10% DV Feb-24 to 2026	21.42	5	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule – 5% DV Feb-24 to 2026	52.63	5	Kenacort-A 40
TRIAMCINOLONE HEXACETONIDE			
Inj 20 mg per ml, 1 ml vial			

Price
(ex man. excl. GST)
\$ Per Brand or
Generic
Manufacturer

Hormone Replacement Therapy

Oestrogens

OESTRADIOL

Tab 1 mg			
Gel (transdermal) 0.06% (750 mcg/actuation) – 5% DV Nov-24 to 31 Oct 2027	14.25	80 g	Estrogel
Patch 25 mcg per day – 5% DV Dec-25 to 2027	8.89	8	Estradiol TDP Mylan
	16.23		Estradot
	21.35		Lyllana
Patch 50 mcg per day – 5% DV Dec-25 to 2027	9.26	8	Estradiol TDP Mylan
	15.79		Estradot
	21.55		Lyllana
Patch 75 mcg per day – 5% DV Dec-25 to 2027	10.33	8	Estradiol TDP Mylan
	16.53		Estradot
	22.37		Lyllana
Patch 100 mcg per day – 5% DV Dec-25 to 2027	10.59	8	Estradiol TDP Mylan
	16.18		Estradot
	22.77		Lyllana

(Lyllana Patch 25 mcg per day to be delisted 1 December 2025)
(Lyllana Patch 50 mcg per day to be delisted 1 December 2025)
(Lyllana Patch 75 mcg per day to be delisted 1 December 2025)
(Lyllana Patch 100 mcg per day to be delisted 1 December 2025)

OESTRADIOL VALERATE

Tab 1 mg – 5% DV Dec-25 to 2028	12.36	84	Progynova
Tab 2 mg – 5% DV Dec-25 to 2028	12.36	84	Progynova

OESTROGENS (CONJUGATED EQUINE)

Tab 300 mcg
Tab 625 mcg

Progestogen and Oestrogen Combined Preparations

OESTRADIOL WITH NORETHISTERONE ACETATE

Tab 1 mg with 0.5 mg norethisterone acetate
Tab 2 mg with 1 mg norethisterone acetate
Tab 2 mg with 1 mg norethisterone acetate (10), and tab 2 mg oestradiol (12) and tab 1 mg oestradiol (6)

OESTROGENS WITH MEDROXYPROGESTERONE ACETATE

Tab 625 mcg conjugated equine with 2.5 mg medroxyprogesterone acetate
Tab 625 mcg conjugated equine with 5 mg medroxyprogesterone acetate

Progestogens

MEDROXYPROGESTERONE ACETATE

Tab 2.5 mg	6.56	30	Provera
Tab 5 mg	20.13	100	Provera
Tab 10 mg	10.28	30	Provera

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Other Endocrine Agents

CABERGOLINE – Restricted see terms below			
↓ Tab 0.5 mg	4.43	2	Dostinex
	17.94	8	Dostinex

→ **Restricted (RS1855)**

Initiation

Any of the following:

- 1 Inhibition of lactation; or
- 2 Patient has hyperprolactinemia; or
- 3 Patient has acromegaly.

Note: Indication marked with * is an unapproved indication.

CLOMIFENE CITRATE			
Tab 50 mg	29.84	10	Mylan Clomiphene
GESTRINONE			
Cap 2.5 mg			
METYRAPONE			
Cap 250 mg			
PENTAGASTRIN			
Inj 250 mcg per ml, 2 ml ampoule			

Other Oestrogen Preparations

OESTRADIOL			
Implant 50 mg			
OESTRIOL			
Tab 2 mg – 5% DV Feb-24 to 2026	7.70	30	Ovestin

Other Progestogen Preparations

MEDROXYPROGESTERONE			
Tab 100 mg	133.57	100	Provera HD
NORETHISTERONE			
Tab 5 mg	5.49	30	Primolut N

Pituitary and Hypothalamic Hormones and Analogues

CORTICORELIN (OVINE)			
Inj 100 mcg vial			
THYROTROPIN ALFA			
Inj 900 mcg vial			

Adrenocorticotropic Hormones

TETRACOSACTIDE [TETRACOSACTRIN]			
Inj 250 mcg per ml, 1 ml ampoule	86.25	1	Synacthen
			UK Synacthen
Inj 1 mg per ml, 1 ml ampoule	690.00	1	Synacthen Depot

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

GnRH Agonists and Antagonists

BUSERELIN

Inj 1 mg per ml, 5.5 ml vial

GONADORELIN

Inj 100 mcg vial

GOSERELIN

Implant 3.6 mg, syringe – **5% DV Apr-24 to 2026**66.48

Implant 10.8 mg, syringe – **5% DV Apr-24 to 2026**138.23

1

Zoladex

1

Zoladex

LEUPRORELIN ACETATE

Inj 3.75 mg prefilled dual chamber syringe.....221.60

Inj 11.25 mg prefilled dual chamber syringe.....591.68

1

Lucrin Depot 1-month

1

Lucrin Depot 3-month

Gonadotrophins

CHORIOGONADOTROPIN ALFA

Inj 250 mcg in 0.5 ml syringe

Growth Hormone

SOMATROPIN – **Restricted** see terms [below](#)

⚠ Inj 5 mg cartridge – **5% DV Feb-25 to 2027**80.21

⚠ Inj 10 mg cartridge – **5% DV Feb-25 to 2027**80.21

⚠ Inj 15 mg cartridge – **5% DV Feb-25 to 2027**139.50

1

Omnitrope

1

Omnitrope

1

Omnitrope

➡ **Restricted (RS1826)**

Initiation – growth hormone deficiency in children

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age; and adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Continuation – growth hormone deficiency in children

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and

continued...

Price			Brand or
(ex man.	excl.	GST)	Generic
\$		Per	Manufacturer

continued...

- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initiation – Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Continuation – Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initiation – short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Continuation – short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 Current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initiation – short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of an endocrinologist or paediatric

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73 m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73 m²) in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m² /day of prednisone or equivalent for at least 6 months.

Continuation – short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of an endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initiation – Prader-Willi syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

- 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Continuation – Prader-Willi syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initiation – adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until it is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Continuation – adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in the Quality of Life Assessment defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have increased to within $\pm 1SD$ of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin does not exceed 0.7 mg per day for male patients, or 1 mg per day for female patients; or

continued...

HORMONE PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 All of the following:

- 2.1 The patient has been treated with somatropin for more than 12 months; and
- 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
- 2.3 Serum IGF-I levels have continued to be maintained within ± 1 SD of the mean of the normal range for age and sex (other than for obvious external factors); and
- 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or

3 All of the following:

- 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
- 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3.3 The patient has severe growth hormone deficiency (see notes); and
- 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Thyroid and Antithyroid Preparations

CARBIMAZOLE

Tab 5 mg – 5% DV Dec-25 to 2028 7.56 100 Neo-Mercazole

IODINE

Soln BP 50 mg per ml

LEVOTHYROXINE

Tab 25 mcg

Tab 50 mcg

Tab 100 mcg

LIOTHYRONINE SODIUM

↓ Tab 20 mcg

➔ Restricted (RS1301)

Initiation

For a maximum of 14 days' treatment in patients with thyroid cancer who are due to receive radioiodine therapy.

Inj 10 mcg vial

Inj 20 mcg vial

Inj 100 mcg vial

POTASSIUM IODATE

Tab 170 mg

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
POTASSIUM PERCHLORATE Cap 200 mg			
PROPYLTHIOURACIL – Restricted see terms below			
⬇ Tab 50 mg	35.00	100	PTU
➡ Restricted (RS1276)			
Initiation			
Both:			
1 The patient has hyperthyroidism; and			
2 The patient is intolerant of carbimazole or carbimazole is contraindicated.			
PROTIRELIN Inj 100 mcg per ml, 2 ml ampoule			

Vasopressin Agents

ARGIPRESSIN [VASOPRESSIN] Inj 20 u per ml, 1 ml ampoule			
DESMOPRESSIN Wafer 120 mcg	47.00	30	Minirin Melt
DESMOPRESSIN ACETATE Tab 100 mcg.....	25.00	30	Minirin
Tab 200 mcg.....	54.45	30	Minirin
Inj 4 mcg per ml, 1 ml ampoule			
Inj 15 mcg per ml, 1 ml ampoule			
Nasal drops 100 mcg per ml			
Nasal spray 10 mcg per dose, 6 ml – 5% DV Apr-25 to 2026	34.95	60	Desmopressin-PH&T
TERLIPRESSIN Inj 0.2 mg per ml, 5 ml vial – 5% DV Feb-25 to 2027	110.00	5	Terlipressin Ever Pharma

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antibacterials			
Aminoglycosides			
AMIKACIN – Restricted see terms below			
⚡ Inj 5 mg per ml, 10 ml syringe			
⚡ Inj 5 mg per ml, 5 ml syringe			
⚡ Inj 15 mg per ml, 5 ml syringe			
⚡ Inj 250 mg per ml, 2 ml vial – 5% DV Dec-24 to 2027	169.97	5	DBL Amikacin
➡ Restricted (RS1041)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule	95.00	5	DBL Gentamicin
Inj 10 mg per ml, 2 ml ampoule			
Inj 40 mg per ml, 2 ml ampoule	36.70	5	Cidomycin P/Free
	18.38	10	Gentamicin Amdipharm
	91.90	50	Gentamicin Noridem
	18.38	10	Pfizer
<i>(Cidomycin P/Free Inj 40 mg per ml, 2 ml ampoule to be delisted 1 December 2025)</i>			
<i>(Gentamicin Amdipharm Inj 40 mg per ml, 2 ml ampoule to be delisted 1 December 2025)</i>			
<i>(Gentamicin Noridem Inj 40 mg per ml, 2 ml ampoule to be delisted 1 December 2025)</i>			
PAROMOMYCIN – Restricted see terms below			
⚡ Cap 250 mg	126.00	16	Humatin
➡ Restricted (RS1603)			
Clinical microbiologist, infectious disease specialist or gastroenterologist			
STREPTOMYCIN SULPHATE – Restricted see terms below			
⚡ Inj 400 mg per ml, 2.5 ml ampoule			
➡ Restricted (RS1043)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
TOBRAMYCIN			
⚡ Powder			
➡ Restricted (RS1475)			
Initiation			
For addition to orthopaedic bone cement.			
⚡ Inj 40 mg per ml, 2 ml vial – 5% DV Dec-24 to 2027	15.50	5	Tobramycin (Viatris)
➡ Restricted (RS1044)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
⚡ Inj 100 mg per ml, 5 ml vial			
➡ Restricted (RS1044)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
⚡ Solution for inhalation 60 mg per ml, 5 ml – 5% DV Dec-23 to 2026	395.00	56 dose	Tobramycin BNM
➡ Restricted (RS1435)			
Initiation			
Patient has cystic fibrosis.			
Carbapenems			
ERTAPENEM – Restricted see terms on the next page			
⚡ Inj 1 g vial	70.00	1	Invanz

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted (RS1045)			
Clinical microbiologist or infectious disease specialist			
IMPENEM WITH CILASTATIN – Restricted see terms below			
↓ Inj 500 mg with 500 mg cilastatin vial	60.00	1	Imipenem+Cilastatin RBX
➔ Restricted (RS1046)			
Clinical microbiologist or infectious disease specialist			
MEROPENEM – Restricted see terms below			
↓ Inj 500 mg vial – 5% DV Jun-24 to 2026	33.48	10	Meropenem-AFT
↓ Inj 1 g vial – 5% DV Jun-24 to 2026	44.97	10	Meropenem-AFT
➔ Restricted (RS1047)			
Clinical microbiologist or infectious disease specialist			
Cephalosporins and Cephamycins - 1st Generation			
CEFALEXIN			
Cap 250 mg	3.85	20	Cephalexin ABM
Cap 500 mg	5.85	20	Cephalexin ABM
Grans for oral liq 25 mg per ml	7.88	100 ml	Flynn
Grans for oral liq 50 mg per ml	11.75	100 ml	Cefalexin Sandoz
	10.38		Flynn
CEFAZOLIN			
Inj 500 mg vial – 5% DV Mar-24 to 2026	3.39	5	Cefazolin-AFT
Inj 1 g vial – 5% DV Mar-24 to 2026	3.59	5	Cefazolin-AFT
Inj 2 g vial – 5% DV Mar-24 to 2026	7.09	5	Cefazolin-AFT
Cephalosporins and Cephamycins - 2nd Generation			
CEFACLOR			
Cap 250 mg – 5% DV Feb-26 to 2028	29.73	100	Ranbaxy-Cefaclor
Grans for oral liq 25 mg per ml – 5% DV Feb-26 to 2028	5.83	100 ml	Ranbaxy-Cefaclor
CEFOXITIN			
Inj 1 g vial			
CEFUROXIME			
Tab 250 mg			
Inj 750 mg vial – 5% DV Jun-24 to 2026	8.16	10	Cefuroxime Devatis
Inj 1.5 g vial – 5% DV Jun-24 to 2026	13.01	10	Cefuroxime Devatis
Cephalosporins and Cephamycins - 3rd Generation			
CEFOTAXIME			
Inj 500 mg vial	1.90	1	Cefotaxime Sandoz
Inj 1 g vial – 5% DV Dec-23 to 2026	38.98	10	DBL Cefotaxime
CEFTAZIDIME – Restricted see terms below			
↓ Inj 1 g vial – 5% DV Dec-23 to 2026	25.80	10	Ceftazidime Kabi
➔ Restricted (RS1048)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
CEFTAZIDIME WITH AVIBACTAM – Restricted see terms on the next page			
↓ Inj ceftazidime 2,000 mg with avibactam 500 mg, vial	2,250.00	10	Zavicefta

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2104)

Initiation

Both:

- 1 Prescribed by, or recommended by a clinical microbiologist or infectious disease specialist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital; and
- 2 Either:
 - 2.1 Proven infection with a carbapenem-resistant micro-organism, based on microbiology report; or
 - 2.2 Probable infection with a carbapenem-resistant micro-organism, based on assessment by a clinical microbiologist or infectious disease specialist.

CEFTRIAXONE

Inj 500 mg vial – 5% DV Feb-26 to 2028	0.94	1	Ceftriaxone-AFT
Inj 1 g vial – 5% DV Feb-26 to 2028	3.49	5	Ceftriaxone-AFT
Inj 2 g vial – 5% DV Feb-26 to 2028	8.15	5	Ceftriaxone-AFT

Cephalosporins and Cephamycins - 4th Generation

CEFEPIME – Restricted see terms [below](#)

⚡ Inj 1 g vial – 5% DV Dec-24 to 2027	3.19	1	Cefepime-AFT
⚡ Inj 2 g vial – 5% DV Dec-24 to 2027	4.99	1	Cefepime-AFT

➔ Restricted (RS1049)

Clinical microbiologist or infectious disease specialist

Cephalosporins and Cephamycins - 5th Generation

CEFTAROLINE FOSAMIL – Restricted see terms [below](#)

⚡ Inj 600 mg vial	1,834.25	10	Zinforo
-------------------------	----------	----	---------

➔ Restricted (RS1446)

Initiation – multi-resistant organism salvage therapy

Clinical microbiologist or infectious disease specialist

Either:

- 1 for patients where alternative therapies have failed; or
- 2 for patients who have a contraindication or hypersensitivity to standard current therapies.

Macrolides

AZITHROMYCIN – Restricted see terms [below](#)

⚡ Tab 250 mg			
⚡ Tab 500 mg – 5% DV Jan-26 to 2027	2.80	2	Zithromax
⚡ Grans for oral liq 200 mg per 5 ml (40 mg per ml).....	16.97	15 ml	Zithromax

➔ Restricted (RS1598)

Initiation – bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections

Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: Indications marked with * are unapproved indications

Initiation – non-cystic fibrosis bronchiectasis*

Respiratory specialist or paediatrician

Re-assessment required after 12 months

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis will be subsidised in the community.

Continuation – non-cystic fibrosis bronchiectasis*

Respiratory specialist or paediatrician

Re-assessment required after 12 months

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

Note: Indications marked with * are unapproved indications. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis will be subsidised in the community.

Initiation – other indications

Re-assessment required after 5 days

For any other condition.

Continuation – other indications

Re-assessment required after 5 days

For any other condition.

CLARITHROMYCIN – Restricted see terms [below](#)

↓ Tab 250 mg – 1% DV Feb-22 to 2027	8.53	14	Klacid
	7.31	12	Klaricid
↓ Tab 500 mg – 1% DV Feb-22 to 2027	14.58	14	Klacid
↓ Grans for oral liq 50 mg per ml	192.00	50 ml	Klacid
↓ Inj 500 mg vial – 5% DV Jul-24 to 2026	9.10	1	Klacid IV

→ Restricted (RS1709)

Initiation – Tab 250 mg and oral liquid

Any of the following:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents; or
- 3 Helicobacter pylori eradication; or
- 4 Prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated.

Initiation – Tab 500 mg

Helicobacter pylori eradication.

Initiation – Infusion

Any of the following:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents; or
- 3 Community-acquired pneumonia.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ERYTHROMYCIN (AS ETHYLSUCCINATE)			
Tab 400 mg	35.82	100	E-Mycin
Grans for oral liq 200 mg per 5 ml	6.53	100 ml	E-Mycin
Grans for oral liq 400 mg per 5 ml	9.41	100 ml	E-Mycin
ERYTHROMYCIN (AS LACTOBIONATE)			
Inj 1 g vial – 5% DV Dec-25 to 2028	10.00	1	Erythrocin IV
ERYTHROMYCIN (AS STEARATE) – Restricted: For continuation only			
➡ Tab 250 mg			
➡ Tab 500 mg			
ROXITHROMYCIN – Some items restricted see terms below			
⚡ Tab dispersible 50 mg			
Tab 150 mg – 5% DV Aug-23 to 2026	13.19	50	Arrow-Roxithromycin
Tab 300 mg – 5% DV Aug-23 to 2026	25.00	50	Arrow-Roxithromycin
➡ Restricted (RS1569)			
Initiation			
Only for use in patients under 12 years of age.			
Penicillins			
AMOXICILLIN			
Cap 250 mg – 5% DV Feb-26 to 2028	54.00	500	Miro-Amoxicillin
Cap 500 mg	41.00	500	Miro-Amoxicillin
Grans for oral liq 125 mg per 5 ml – 5% DV Feb-24 to 2026	2.22	100 ml	Alphamox 125
Grans for oral liq 250 mg per 5 ml – 5% DV Feb-24 to 2026	2.81	100 ml	Alphamox 250
Inj 250 mg vial	15.97	10	Ibiamox
Inj 500 mg vial	17.43	10	Ibiamox
Inj 1 g vial	21.64	10	Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg – 5% DV Feb-24 to 2026	1.59	10	Curam Duo 500/125
Grans for oral liq 25 mg with clavulanic acid 6.25 mg per ml – 5% DV May-25 to 2027	8.50	100 ml	Augmentin
Grans for oral liq 50 mg with clavulanic acid 12.5 mg per ml – 5% DV Jun-25 to 2027	5.61	100 ml	Amoxiclav Devatis Forte
Inj 500 mg with clavulanic acid 100 mg vial – 5% DV Sep-25 to 2027	22.48	10	Synermox
Inj 1,000 mg with clavulanic acid 200 mg vial – 5% DV Sep-25 to 2027	26.90	10	Cerobact
	29.61		Synermox
BENZATHINE BENZYL PENICILLIN			
Inj 900 mg (1.2 million units) vial	43.24	1	Benzetacil
Inj 900 mg (1.2 million units) in 2.3 ml syringe	432.37	10	Bicillin LA
BENZYL PENICILLIN SODIUM [PENICILLIN G]			
Inj 600 mg (1 million units) vial – 5% DV Feb-24 to 2026	16.50	10	Sandoz
FLUCLOXACILLIN			
Cap 250 mg – 5% DV Aug-25 to 2027	22.58	250	Staphlex
Cap 500 mg – 5% DV Aug-25 to 2027	72.71	500	Staphlex
Grans for oral liq 25 mg per ml – 5% DV Feb-25 to 2027	4.89	100 ml	AFT
Grans for oral liq 50 mg per ml – 5% DV Feb-25 to 2027	5.89	100 ml	AFT
Inj 250 mg vial – 5% DV Jul-24 to 2026	42.60	10	Flucloxin
Inj 500 mg vial – 5% DV Jul-24 to 2026	45.63	10	Flucloxin
Inj 1 g vial – 5% DV Feb-24 to 2026	6.00	5	Flucil

↑ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PHENOXYMETHYLPENICILLIN [PENICILLIN V]			
Cap 250 mg – 5% DV Feb-25 to 2027	7.68	50	Cilicaine VK
Cap 500 mg – 5% DV Feb-25 to 2027	13.72	50	Cilicaine VK
Grans for oral liq 125 mg per 5 ml – 5% DV Feb-26 to 2028	5.75	100 ml	AFT
Grans for oral liq 250 mg per 5 ml – 5% DV Feb-26 to 2028	5.89	100 ml	AFT

PIPERACILLIN WITH TAZOBACTAM – Restricted see terms [below](#)

↓ Inj 4 g with tazobactam 0.5 g vial – **5% DV Dec-25 to 2028**.....3.15

→ **Restricted (RS1053)**

Clinical microbiologist, infectious disease specialist or respiratory specialist

PROCAINE PENICILLIN

Inj 1.5 g in 3.4 ml syringe

TICARCILLIN WITH CLAVULANIC ACID – Restricted see terms [below](#)

↓ Inj 3 g with clavulanic acid 0.1 mg vial

→ **Restricted (RS1054)**

Clinical microbiologist, infectious disease specialist or respiratory specialist

Quinolones

CIPROFLOXACIN – Restricted see terms [below](#)

↓ Tab 250 mg – **5% DV Nov-24 to 2026**.....1.95

↓ Tab 500 mg – **5% DV Nov-24 to 2026**.....3.10

↓ Tab 750 mg – **5% DV Dec-24 to 2026**.....4.80

↓ Oral liq 50 mg per ml

↓ Oral liq 100 mg per ml

↓ Inj 2 mg per ml, 100 ml bag

↓ Inj 2 mg per ml, 100 ml bottle166.50

→ **Restricted (RS1055)**

Clinical microbiologist or infectious disease specialist

MOXIFLOXACIN – Restricted see terms [below](#)

↓ Tab 400 mg42.00

↓ Inj 1.6 mg per ml, 250 ml bottle – **5% DV Feb-24 to 2026**.....413.40

→ **Restricted (RS2129)**

Initiation – Mycobacterium infection

Infectious disease specialist, clinical microbiologist or respiratory specialist

Any of the following:

1 Both:

1.1 Active tuberculosis; and

1.2 Any of the following:

1.2.1 Documented resistance to one or more first-line medications; or

1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or

1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or

1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or

1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;
or

2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated; or

3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – Pneumonia

Infectious disease specialist or clinical microbiologist

Either:

- 1 Immunocompromised patient with pneumonia that is unresponsive to first-line treatment; or
- 2 Pneumococcal pneumonia or other invasive pneumococcal disease highly resistant to other antibiotics.

Initiation – Penetrating eye injury

Ophthalmologist

Five days treatment for patients requiring prophylaxis following a penetrating eye injury.

Initiation – Mycoplasma genitalium

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initiation – severe delayed beta-lactam allergy

Infectious disease specialist or clinical microbiologist

Individual has a history of severe delayed beta-lactam allergy.

NORFLOXACIN

Tab 400 mg	245.00	100	Arrow-Norfloxacin
------------------	--------	-----	-------------------

Tetracyclines

DEMECLOCYCLINE HYDROCHLORIDE

Tab 150 mg
Cap 150 mg
Cap 300 mg

DOXYCYCLINE

➔ Tab 50 mg – **Restricted:** For continuation only

Tab 100 mg	64.43	500	Doxine
Inj 5 mg per ml, 20 ml vial			

MINOCYCLINE

Tab 50 mg

➔ Cap 100 mg – **Restricted:** For continuation only

TETRACYCLINE

Tab 250 mg	68.44	28	Accord
Cap 500 mg			

TIGECYCLINE – **Restricted** see terms [below](#)

⚡ Inj 50 mg vial

➔ **Restricted (RS1059)**

Clinical microbiologist or infectious disease specialist

Other Antibacterials

AZTREONAM – **Restricted** see terms [below](#)

⚡ Inj 1 g vial	364.92	10	Azactam
----------------------	--------	----	---------

➔ **Restricted (RS1277)**

Clinical microbiologist or infectious disease specialist

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CHLORAMPHENICOL – Restricted see terms below			
↓ Inj 1 g vial			
→ Restricted (RS1277)			
Clinical microbiologist or infectious disease specialist			
CLINDAMYCIN – Restricted see terms below			
↓ Cap 150 mg – 5% DV Dec-24 to 2027	4.94	24	Dalacin C
↓ Oral liq 15 mg per ml			
↓ Inj 150 mg per ml, 4 ml ampoule	35.10	10	Hameln
→ Restricted (RS1061)			
Clinical microbiologist or infectious disease specialist			
COLISTIN SULPHOMETHATE [COLESTIMETHATE] – Restricted see terms below			
↓ Inj 2 million iu, 10 ml vial	216.67	10	Colomycin
→ Restricted (RS1062)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
DAPTOMYCIN – Restricted see terms below			
↓ Inj 500 mg vial	115.36	1	Daptomycin Dr Reddy's
→ Restricted (RS1063)			
Clinical microbiologist or infectious disease specialist			
FOSFOMYCIN – Restricted see terms below			
↓ Powder for oral solution, 3 g sachet – 5% DV Apr-25 to 2027	18.70	1	UroFos
→ Restricted (RS1315)			
Clinical microbiologist or infectious disease specialist			
LINCOMYCIN – Restricted see terms below			
↓ Inj 300 mg per ml, 2 ml vial			
→ Restricted (RS1065)			
Clinical microbiologist or infectious disease specialist			
LINEZOLID – Restricted see terms below			
↓ Tab 600 mg – 5% DV Dec-24 to 2027	194.60	10	Zyvox
↓ Oral liq 20 mg per ml	1,879.00	150 ml	Zyvox
↓ Inj 2 mg per ml, 300 ml bottle – 5% DV Dec-24 to 2027	155.00	10	Linezolid Kabi
→ Restricted (RS1066)			
Clinical microbiologist or infectious disease specialist			
METHENAMINE (HEXAMINE) HIPPURATE			
Tab 1 g	19.95	100	Hiprex
NITROFURANTOIN			
Tab 50 mg – 5% DV Dec-24 to 2027	22.20	100	Nifuran
Tab 100 mg	37.50	100	Nifuran
Cap modified-release 100 mg – 5% DV Dec-23 to 2026	81.20	100	Macrobid
PIVMECILLINAM – Restricted see terms below			
↓ Tab 200 mg			
→ Restricted (RS1322)			
Clinical microbiologist or infectious disease specialist			
SODIUM FUSIDATE [FUSIDIC ACID] – Restricted see terms below			
↓ Tab 250 mg	135.70	36	Fucidin
→ Restricted (RS1064)			
Clinical microbiologist or infectious disease specialist			
SULFADIAZINE SODIUM – Restricted see terms on the next page			
↓ Tab 500 mg			<i>e.g. Sulfadiazin-Heyl; Wockhardt</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted (RS1067)			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
TEICOPLANIN – Restricted see terms below			
⚡ Inj 400 mg vial – 5% DV Apr-25 to 2027	38.85	1	Teicoplanin Medsurge
➔ Restricted (RS1068)			
Clinical microbiologist or infectious disease specialist			
TRIMETHOPRIM			
Tab 100 mg			
Tab 300 mg – 5% DV Feb-25 to 2027	27.83	50	TMP
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]			
Tab 80 mg with sulphamethoxazole 400 mg – 5% DV Feb-25 to 2027	115.74	500	Trisul
Oral liq 8 mg with sulphamethoxazole 40 mg per ml – 5% DV Aug-25 to 2028	4.95	100 ml	Deprim
Inj 16 mg with sulphamethoxazole 80 mg per ml, 5 ml ampoule			
VANCOMYCIN – Restricted see terms below			
⚡ Inj 500 mg vial – 5% DV Feb-24 to 2026	3.38	1	Mylan
➔ Restricted (RS1069)			
Clinical microbiologist or infectious disease specialist			
Antifungals			
Imidazoles			
KETOCONAZOLE			
⚡ Tab 200 mg			
➔ Restricted (RS1410)			
Oncologist			
Polyene Antimycotics			
AMPHOTERICIN B			
⚡ Inj (liposomal) 50 mg vial	3,450.00	10	AmBisome
➔ Restricted (RS1071)			
Initiation			
Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist			
Either:			
1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or			
2 Both:			
2.1 Possible invasive fungal infection; and			
2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.			
⚡ Inj 50 mg vial			
➔ Restricted (RS1316)			
Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist			
NYSTATIN			
Tab 500,000 u	17.09	50	Nilstat
Cap 500,000 u	15.47	50	Nilstat

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Triazoles			
FLUCONAZOLE – Restricted see terms below			
↓ Cap 50 mg – 5% DV Dec-23 to 2026.....	4.10	28	Mylan
↓ Cap 150 mg – 5% DV Dec-23 to 2026.....	0.45	1	Mylan
↓ Cap 200 mg – 5% DV Dec-23 to 2026.....	8.90	28	Mylan
↓ Oral liquid 50 mg per 5 ml	129.02	35 ml	Diflucan
↓ Inj 2 mg per ml, 50 ml vial.....	11.20	1	Fluconazole-Baxter
↓ Inj 2 mg per ml, 100 ml vial.....	5.20	1	Fluconazole-Baxter
→ Restricted (RS1072)			
Consultant			
ITRACONAZOLE – Restricted see terms below			
↓ Cap 100 mg	27.32	60	Itracap
	6.83	15	Itraconazole Crescent Itrazole
↓ Oral liquid 10 mg per ml			
<i>(Itracap Cap 100 mg to be delisted 1 December 2025)</i>			
→ Restricted (RS1073)			
Clinical immunologist, clinical microbiologist, dermatologist or infectious disease specialist			
POSACONAZOLE – Restricted see terms below			
↓ Tab modified-release 100 mg – 5% DV Dec-25 to 2028	123.60	24	Posaconazole Juno
↓ Oral liq 40 mg per ml – 5% DV Dec-25 to 2028	308.26	105 ml	Devatis
→ Restricted (RS2052)			
Initiation			
Haematologist or infectious disease specialist			
<i>Re-assessment required after 6 weeks</i>			
Both:			
1 Either:			
1.1 Patient has acute myeloid leukaemia; or			
1.2 Patient is planned to receive a stem cell transplant and is at high risk for aspergillus infection; and			
2 Patient is to be treated with high dose remission induction therapy or re-induction therapy.			
Continuation			
Haematologist or infectious disease specialist			
<i>Re-assessment required after 6 weeks</i>			
Both:			
1 Patient has previously received posaconazole prophylaxis during remission induction therapy; and			
2 Any of the following:			
2.1 Patient is to be treated with high dose remission re-induction therapy; or			
2.2 Patient is to be treated with high dose consolidation therapy; or			
2.3 Patient is receiving a high risk stem cell transplant.			
Initiation – Invasive fungal infection prophylaxis			
Any relevant practitioner			
<i>Re-assessment required after 6 months</i>			
Both:			
1 The patient is at risk of invasive fungal infection; and			
2 Either:			
2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Continuation – Invasive fungal infection prophylaxis

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

VORICONAZOLE – **Restricted** see terms [below](#)

⚡ Tab 50 mg – 5% DV Aug-25 to 2028	71.00	56	Vttack
⚡ Tab 200 mg – 5% DV Aug-25 to 2028	263.00	56	Vttack
⚡ Powder for oral suspension 40 mg per ml.....	1,523.22	70 ml	Vfend
⚡ Inj 200 mg vial – 5% DV Dec-25 to 2028	16.89	1	AFT

➡ **Restricted (RS2053)**

Initiation – Proven or probable aspergillus infection

Clinical microbiologist, haematologist or infectious disease specialist

Both:

- 1 Patient is immunocompromised; and
- 2 Patient has proven or probable invasive aspergillus infection.

Initiation – Possible aspergillus infection

Clinical microbiologist, haematologist or infectious disease specialist

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient has possible invasive aspergillus infection; and
- 3 A multidisciplinary team (including an infectious disease physician) considers the treatment to be appropriate.

Initiation – Resistant candidiasis infections and other moulds

Clinical microbiologist, haematologist or infectious disease specialist

All of the following:

- 1 Patient is immunocompromised; and
- 2 Either:
 - 2.1 Patient has fluconazole resistant candidiasis; or
 - 2.2 Patient has mould strain such as *Fusarium* spp. and *Scedosporium* spp; and
- 3 A multidisciplinary team (including an infectious disease physician or clinical microbiologist) considers the treatment to be appropriate.

Initiation – Invasive fungal infection prophylaxis

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:

continued...

	Price (ex man. excl. GST)		Brand or Generic Manufacturer
	\$	Per	

continued...

- 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
- 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Continuation – Invasive fungal infection prophylaxis

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 The patient is at risk of invasive fungal infection; and
- 2 Either:
 - 2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
 - 2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Other Antifungals

CASPOFUNGIN – **Restricted** see terms [below](#)

↓ Inj 50 mg vial	110.00	1	Alchemy Caspofungin
↓ Inj 70 mg vial	135.00	1	Alchemy Caspofungin

→ **Restricted (RS1076)**

Initiation

Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist

Either:

- 1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
- 2 Both:
 - 2.1 Possible invasive fungal infection; and
 - 2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

FLUCYTOSINE – **Restricted** see terms [below](#)

↓ Tab 500 mg
↓ Cap 500 mg

→ **Restricted (RS1279)**

Clinical microbiologist or infectious disease specialist

TERBINAFINE

Tab 250 mg – 5% DV Feb-24 to 2026	8.97	84	Deolatte
---	------	----	----------

Antimycobacterials

Antileprotics

CLOFAZIMINE – **Restricted** see terms [below](#)

↓ Cap 50 mg

→ **Restricted (RS1077)**

Clinical microbiologist, dermatologist or infectious disease specialist

DAPSONE – **Restricted** see terms [on the next page](#)

↓ Tab 25 mg	268.50	100	Dapsone
↓ Tab 100 mg	329.50	100	Dapsone

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted (RS1078) Clinical microbiologist, dermatologist or infectious disease specialist			
Antituberculotics			
BEDAQUILINE – Restricted see terms below			
⚡ Tab 100 mg	3,084.51	24	Sirturo
➔ Restricted (RS1977) Initiation – multi-drug resistant tuberculosis <i>Limited to 6 months treatment</i> Both: 1 The person has multi-drug resistant tuberculosis (MDR-TB); and 2 Ministry of Health's Tuberculosis Clinical Network has reviewed the individual case and recommends bedaquiline as part of the treatment regimen.			
CYCLOSERINE – Restricted see terms below			
⚡ Cap 250 mg			
➔ Restricted (RS1079) Clinical microbiologist, infectious disease specialist or respiratory specialist			
ETHAMBUTOL HYDROCHLORIDE – Restricted see terms below			
⚡ Tab 100 mg			
⚡ Tab 400 mg	49.34	56	Myambutol
➔ Restricted (RS1080) Clinical microbiologist, infectious disease specialist or respiratory specialist			
ISONIAZID – Restricted see terms below			
⚡ Tab 100 mg – 5% DV May-25 to 2027	94.50	100	Isoniazid Teva
	327.41		Noumed Isoniazid
➔ Restricted (RS1281) Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician			
ISONIAZID WITH RIFAMPICIN – Restricted see terms below			
⚡ Tab 100 mg with rifampicin 150 mg – 5% DV Feb-25 to 2027	89.82	100	Rifinah
⚡ Tab 150 mg with rifampicin 300 mg – 5% DV Feb-25 to 2027	179.13	100	Rifinah
➔ Restricted (RS1282) Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician			
PARA-AMINOSALICYLIC ACID – Restricted see terms below			
⚡ Grans for oral liq 4 g	280.00	30	Paser
➔ Restricted (RS1083) Clinical microbiologist, infectious disease specialist or respiratory specialist			
PROTIONAMIDE – Restricted see terms below			
⚡ Tab 250 mg	305.00	100	Peteha
➔ Restricted (RS1084) Clinical microbiologist, infectious disease specialist or respiratory specialist			
PYRAZINAMIDE – Restricted see terms below			
⚡ Tab 500 mg			
➔ Restricted (RS1085) Clinical microbiologist, infectious disease specialist or respiratory specialist			
RIFABUTIN – Restricted see terms below			
⚡ Cap 150 mg	353.71	30	Mycobutin
➔ Restricted (RS1086) Clinical microbiologist, gastroenterologist, infectious disease specialist or respiratory specialist			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
RIFAMPICIN – Restricted see terms below			
↓ Cap 150 mg – 5% DV Dec-23 to 2026	58.54	100	Rifadin
↓ Cap 300 mg – 5% DV Dec-23 to 2026	122.06	100	Rifadin
↓ Oral liq 100 mg per 5 ml – 5% DV Dec-23 to 2026	12.60	60 ml	Rifadin
↓ Inj 600 mg vial – 5% DV Dec-23 to 2026	134.98	1	Rifadin

→ **Restricted (RS1087)**

Clinical microbiologist, dermatologist, internal medicine physician, paediatrician or public health physician

Antiparasitics

Anthelmintics

ALBENDAZOLE – **Restricted** see terms [below](#)

↓ Tab 200 mg

↓ Tab 400 mg

→ **Restricted (RS1088)**

Clinical microbiologist or infectious disease specialist

IVERMECTIN – **Restricted** see terms [below](#)

↓ Tab 3 mg 17.20 4 Stromectol

→ **Restricted (RS1283)**

Clinical microbiologist, dermatologist or infectious disease specialist

MEBENDAZOLE

Tab 100 mg – **5% DV Dec-24 to 2027** 5.18 6 **Vermox**

Oral liq 100 mg per 5 ml

PRAZIQUANTEL

Tab 600 mg

Antiprotozoals

ARTEMETHER WITH LUMEFANTRINE – **Restricted** see terms [below](#)

↓ Tab 20 mg with lumefantrine 120 mg

→ **Restricted (RS1090)**

Clinical microbiologist or infectious disease specialist

ARTESUNATE – **Restricted** see terms [below](#)

↓ Inj 60 mg vial

→ **Restricted (RS1091)**

Clinical microbiologist or infectious disease specialist

ATOVAQUONE WITH PROGUANIL HYDROCHLORIDE – **Restricted** see terms [below](#)

↓ Tab 62.5 mg with proguanil hydrochloride 25 mg..... 27.20 12 Malarone Junior

↓ Tab 250 mg with proguanil hydrochloride 100 mg..... 69.50 12 Malarone

→ **Restricted (RS1092)**

Clinical microbiologist or infectious disease specialist

CHLOROQUINE PHOSPHATE – **Restricted** see terms [below](#)

↓ Tab 250 mg

→ **Restricted (RS1093)**

Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist

MEFLOQUINE – **Restricted** see terms [below](#)

↓ Tab 250 mg

→ **Restricted (RS1094)**

Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METRONIDAZOLE			
Tab 200 mg – 5% DV Mar-25 to 2027	25.86	250	Metronidamed
Tab 400 mg – 5% DV Mar-25 to 2027	4.29	21	Metronidamed
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	Flagyl-S
Inj 5 mg per ml, 100 ml bag – 5% DV Dec-23 to 2026	18.00	10	Baxter
Suppos 500 mg	24.48	10	Flagyl
NITAZOXANIDE – Restricted see terms below			
↓ Tab 500 mg			
↓ Oral liq 100 mg per 5 ml			
➔ Restricted (RS1095)			
Clinical microbiologist or infectious disease specialist			
ORNIDAZOLE			
Tab 500 mg – 5% DV Mar-25 to 2027	36.52	10	Arrow-Ornidazole
PENTAMIDINE ISETHIONATE – Restricted see terms below			
↓ Inj 300 mg vial	216.00	5	Pentacarinat
➔ Restricted (RS1096)			
Clinical microbiologist or infectious disease specialist			
PRIMAQUINE – Restricted see terms below			
↓ Tab 15 mg			
↓ Tab 7.5 mg			
➔ Restricted (RS1097)			
Clinical microbiologist or infectious disease specialist			
PYRIMETHAMINE – Restricted see terms below			
↓ Tab 25 mg			
➔ Restricted (RS1098)			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
QUININE DIHYDROCHLORIDE – Restricted see terms below			
↓ Inj 60 mg per ml, 10 ml ampoule			
↓ Inj 300 mg per ml, 2 ml vial			
➔ Restricted (RS1099)			
Clinical microbiologist or infectious disease specialist			
SODIUM STIBOGLUCONATE – Restricted see terms below			
↓ Inj 100 mg per ml, 1 ml vial			
➔ Restricted (RS1100)			
Clinical microbiologist or infectious disease specialist			
SPIRAMYCIN – Restricted see terms below			
↓ Tab 500 mg			
➔ Restricted (RS1101)			
Maternal-foetal medicine specialist			

Antiretrovirals

Non-Nucleoside Reverse Transcriptase Inhibitors

➔ **Restricted** ([RS1898](#))

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (<https://www.ashm.org.au/hiv/hiv-management/pep/>).

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

EFAVIRENZ – **Restricted** see terms [on the previous page](#)

➔ Tab 600 mg – **Restricted:** For continuation only65.38 30 Efavirenz Milpharm
 † Oral liq 30 mg per ml

(Efavirenz Milpharm Tab 600 mg to be delisted 1 November 2026)

ETRAVIRINE – **Restricted** see terms [on the previous page](#)

† Tab 200 mg770.00 60 Intelence

NEVIRAPINE – **Restricted** see terms [on the previous page](#)

† Tab 200 mg – **5% DV Feb-25 to 2027**198.25 60 **Nevirapine Viatris**
 † Oral suspension 10 mg per ml.....203.55 240 ml Viramune Suspension

Nucleoside Reverse Transcriptase Inhibitors

➔ **Restricted (RS1899)**

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
prophylaxis is required; or			
2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.			
Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).			
Initiation – Percutaneous exposure			
Patient has percutaneous exposure to blood known to be HIV positive.			
ABACAVIR SULPHATE – Restricted see terms on the previous page			
† Tab 300 mg	180.00	60	Ziagen
† Oral liq 20 mg per ml			
ABACAVIR SULPHATE WITH LAMIVUDINE – Restricted see terms on the previous page			
† Tab 600 mg with lamivudine 300 mg – 5% DV Feb-26 to 2028	35.00	30	Abacavir/lamivudine Viatris
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPROXIL – Restricted see terms on the previous page			
† Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate)	106.88	30	Viatris
† Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a fumarate)	106.88	30	TEEVIR Triovir
EMTRICITABINE – Restricted see terms on the previous page			
† Cap 200 mg	307.20	30	Emtriva
LAMIVUDINE – Restricted see terms on the previous page			
† Tab 150 mg – 5% DV Feb-24 to 2026	98.00	60	Lamivudine Viatris
† Oral liq 10 mg per ml			
STAVUDINE – Restricted see terms on the previous page			
† Cap 30 mg			
† Cap 40 mg			
† Powder for oral soln 1 mg per ml			
ZIDOVUDINE [AZT] – Restricted see terms on the previous page			
† Cap 100 mg	152.25	100	Retrovir
† Oral liq 10 mg per ml	30.45	200 ml	Retrovir
† Inj 10 mg per ml, 20 ml vial	750.00	5	Retrovir IV
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Restricted see terms on the previous page			
† Tab 300 mg with lamivudine 150 mg	92.40	60	Lamivudine/Zidovudine Viatris

Protease Inhibitors

➔ **Restricted (RS1900)**

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
Initiation – Post-exposure prophylaxis following exposure to HIV			
Both:			
1 Treatment course to be initiated within 72 hours post exposure; and			
2 Any of the following:			
2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or			
2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or			
2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or			
2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.			
Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).			
Initiation – Percutaneous exposure			
Patient has percutaneous exposure to blood known to be HIV positive.			
ATAZANAVIR SULPHATE – Restricted see terms on the previous page			
† Cap 150 mg – 5% DV Feb-26 to 2028	85.00	60	Atazanavir Mylan
	102.50		Atazanavir Viatris
† Cap 200 mg – 5% DV Feb-26 to 2028	152.30	60	Atazanavir Viatris
<i>(Atazanavir Mylan Cap 150 mg to be delisted 1 November 2025)</i>			
DARUNAVIR – Restricted see terms on the previous page			
† Tab 400 mg – 5% DV Feb-24 to 2026	150.00	60	Darunavir Viatris
† Tab 600 mg – 5% DV Feb-24 to 2026	225.00	60	Darunavir Viatris
INDINAVIR – Restricted see terms on the previous page			
† Cap 200 mg			
† Cap 400 mg			
LOPINAVIR WITH RITONAVIR – Restricted see terms on the previous page			
† Tab 100 mg with ritonavir 25 mg	150.00	60	Lopinavir/Ritonavir Mylan
† Tab 200 mg with ritonavir 50 mg – 5% DV Feb-25 to 2027	875.00	120	Lopinavir/Ritonavir Mylan
RITONAVIR – Restricted see terms on the previous page			
† Tab 100 mg	43.31	30	Norvir

Strand Transfer Inhibitors

➔ Restricted (RS1901)

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 Any of the following:

- 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
- 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
- 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
- 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (<https://www.ashm.org.au/hiv/hiv-management/pep/>).

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

DOLUTEGRAVIR – Restricted see terms [on the previous page](#)

† Tab 50 mg	1,090.00	30	Tivicay
-------------------	----------	----	---------

DOLUTEGRAVIR WITH LAMIVUDINE – Restricted see terms [on the previous page](#)

† Tab 50 mg with lamivudine 300 mg.....	1,090.00	30	Dovato
---	----------	----	--------

RALTEGRAVIR POTASSIUM – Restricted see terms [on the previous page](#)

† Tab 400 mg	1,090.00	60	Isentress
† Tab 600 mg	1,090.00	60	Isentress HD

Antivirals

Hepatitis B

ENTECAVIR

Tab 0.5 mg – 5% DV Mar-24 to 2026	12.04	30	Entecavir (Rex)
---	-------	----	-----------------

LAMIVUDINE

Tab 100 mg – 5% DV Feb-24 to 2026	12.06	28	Zetlam
Oral liq 5 mg per ml	270.00	240 ml	Zeffix

TENOFOVIR DISOPROXIL

Tab 245 mg (300 mg as a maleate) – 5% DV Dec-25 to 2028	13.80	30	Tenofovir Disoproxil Viatris
Tab 245 mg (300 mg as a fumarate)	13.80	30	Ricovir

Hepatitis C

GLECAPREVIR WITH PIBRENTASVIR

Note: the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website <https://www.pharmac.govt.nz/maviret>.

Tab 100 mg with pibrentasvir 40 mg	24,750.00	84	Maviret
--	-----------	----	---------

LEDIPASVIR WITH SOFOSBUVIR – Restricted see terms [below](#)

† Tab 90 mg with sofosbuvir 400 mg.....	24,363.46	28	Harvoni
---	-----------	----	---------

➔ **Restricted (RS1528)**

Note: Only for use in patients with approval by the Hepatitis C Treatment Panel (HepCTP). Applications will be considered by HepCTP at its regular meetings and approved subject to eligibility according to the Access Criteria (set out in Section B of the Pharmaceutical Schedule).

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Herpesviridae			
ACICLOVIR			
Tab dispersible 200 mg – 5% DV Feb-26 to 2028	2.05	25	Lovir
Tab dispersible 400 mg – 5% DV Feb-26 to 2028	7.55	56	Lovir
Tab dispersible 800 mg – 5% DV Feb-26 to 2028	7.43	35	Lovir
Inj 250 mg vial – 5% DV Feb-25 to 2027	13.75	5	Aciclovir-Baxter
CIDOFOVIR – Restricted see terms below			
↓ Inj 75 mg per ml, 5 ml vial			
➔ Restricted (RS1108)			
Clinical microbiologist, infectious disease specialist, otolaryngologist or oral surgeon			
FOSCARNET SODIUM – Restricted see terms below			
↓ Inj 24 mg per ml, 250 ml bottle			
➔ Restricted (RS1109)			
Clinical microbiologist or infectious disease specialist			
GANCICLOVIR – Restricted see terms below			
↓ Inj 500 mg vial	380.00	5	Cymevene
➔ Restricted (RS1110)			
Clinical microbiologist or infectious disease specialist			
VALACICLOVIR			
Tab 500 mg – 5% DV Feb-25 to 2027	9.64	30	Vaclovir
Tab 1,000 mg – 5% DV Feb-25 to 2027	17.78	30	Vaclovir
VALGANCICLOVIR – Restricted see terms below			
↓ Tab 450 mg – 5% DV Feb-25 to 2027	140.89	60	Valganciclovir Viatris
➔ Restricted (RS2137)			
Initiation – Transplant cytomegalovirus prophylaxis			
<i>Re-assessment required after 3 months</i>			
Patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.			
Continuation – Transplant cytomegalovirus prophylaxis			
<i>Re-assessment required after 3 months</i>			
Either:			
1 Both:			
1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and			
1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or			
2 Both:			
2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and			
2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.			
Initiation – Lung transplant cytomegalovirus prophylaxis			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Patient has undergone a lung transplant; and			
2 Either:			
2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or			
2.2 The recipient is cytomegalovirus positive; and			
3 Patient has a high risk of CMV disease.			

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – Lung transplant cytomegalovirus prophylaxis

Re-assessment required after 12 months

All of the following:

- 1 Patient has undergone a lung re-transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initiation – Cytomegalovirus in immunocompromised patients

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – **Restricted** see terms [below](#)

⚡ Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a maleate) – 5% DV Dec-25 to 2028	13.45	30	Tenofovir Disoproxil Emtricitabine Viatr
---	-------	----	---

➡ **Restricted (RS1902)**

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

Initiation – Prevention of maternal transmission

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

Initiation – Pre-exposure prophylaxis

Re-assessment required after 24 months

Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines (<https://ashm.org.au/HIV/PrEP/>)

continued...

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)
e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price		Brand or
(ex man. excl. GST)			Generic
\$	Per		Manufacturer

continued...

Continuation – Pre-exposure prophylaxis

Re-assessment required after 24 months

Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines (<https://ashm.org.au/HIV/PrEP/>)

Influenza

OSELTAMIVIR – **Restricted** see terms [below](#)

Note: The restriction on the use of oseltamivir to hospitalised patients means that supply into the community for a new course is not permitted. Supply of a part original pack on discharge where initiated as a hospital inpatient is permitted.

↓ Tab 75 mg

↓ Powder for oral suspension 6 mg per ml

→ **Restricted** ([RS1307](#))

Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a Health NZ Hospital approved infections control plan.

ZANAMIVIR

Note: The restriction on the use of zanamivir to hospitalised patients means that supply into the community for a new course is not permitted. Supply of a part original pack on discharge where initiated as a hospital inpatient is permitted.

↓ Powder for inhalation 5 mg 37.38 20 dose Relenza Rotadisk

→ **Restricted** ([RS1369](#))

Initiation

Either:

- 1 Only for hospitalised patient with known or suspected influenza; or
- 2 For prophylaxis of influenza in hospitalised patients as part of a Health NZ Hospital approved infections control plan.

COVID-19 Treatments

NIRMATRELVIR WITH RITONAVIR – **Restricted** see terms [below](#)

↓ Tab 150 mg with ritonavir 100 mg 0.00 30 Paxlovid

→ **Restricted** ([RS1894](#))

Initiation

Only if patient meets access criteria (as per <https://pharmac.govt.nz/covid-oral-antivirals>). Note the supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

REMEDESIVIR

Note: For patients meeting access criteria for oral antiviral treatments (as on Pharmac's website).

Inj 100 mg vial 615.23 1 Veklury

Immune Modulators

INTERFERON ALFA-2B

Inj 18 m iu, 1.2 ml multidose pen

Inj 30 m iu, 1.2 ml multidose pen

Inj 60 m iu, 1.2 ml multidose pen

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

INTERFERON GAMMA – Restricted see terms [below](#)

‡ Inj 100 mcg in 0.5 ml vial

➔ **Restricted (RS1113)**

Initiation

Patient has chronic granulomatous disease and requires interferon gamma.

PEGYLATED INTERFERON ALFA-2A – Restricted see terms [below](#)

‡ Inj 135 mcg prefilled syringe.....	887.35	1	Pegasys (S29)
‡ Inj 180 mcg prefilled syringe.....	748.50	4	Pegasys
	1,355.71		Pegasys

➔ **Restricted (RS1827)**

Initiation – Chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant

Limited to 48 weeks treatment

Any of the following:

- 1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
- 2 Patient has chronic hepatitis C and is co-infected with HIV; or
- 3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant.

Notes: Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.

Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml.

Continuation – Chronic hepatitis C - genotype 1 infection

Gastroenterologist, infectious disease specialist or general physician

Re-assessment required after 48 weeks

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir.

Initiation – Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir.

Initiation – Chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV

Limited to 6 months treatment

Patient has chronic hepatitis C, genotype 2 or 3 infection.

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

Initiation – Hepatitis B

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naïve; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log₁₀ IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 Serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (greater than or equal to Metavir Stage F2 or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon.

Initiation – myeloproliferative disorder or cutaneous T cell lymphoma

Re-assessment required after 12 months

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Continuation – myeloproliferative disorder or cutaneous T cell lymphoma

Re-assessment required after 12 months

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications

Initiation – ocular surface squamous neoplasia

Ophthalmologist

Re-assessment required after 12 months

Patient has ocular surface squamous neoplasia*.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – ocular surface squamous neoplasia

Ophthalmologist

Re-assessment required after 12 months

The treatment remains appropriate and patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initiation – post-allogenic bone marrow transplant

Re-assessment required after 3 months

Patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Continuation – post-allogenic bone marrow transplant

Re-assessment required after 3 months

Patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

Anticholinesterases

EDROPHONIUM CHLORIDE – **Restricted** see terms [below](#)

↓ Inj 10 mg per ml, 15 ml vial

↓ Inj 10 mg per ml, 1 ml ampoule

→ **Restricted (RS1015)**

Initiation

For the diagnosis of myasthenia gravis.

NEOSTIGMINE METILSULFATE

Inj 2.5 mg per ml, 1 ml ampoule – **5% DV Feb-25 to 2027** 48.25 10 **Max Health**

NEOSTIGMINE METILSULFATE WITH GLYCOPYRRONIUM BROMIDE

Inj 2.5 mg with glycopyrronium bromide 0.5 mg per ml, 1 ml ampoule –
5% DV Feb-26 to 2028 26.13 10 **Max Health**

PYRIDOSTIGMINE BROMIDE

Tab 60 mg 50.28 100 Mestinon

Antirheumatoid Agents

HYDROXYCHLOROQUINE SULPHATE

Tab 200 mg – **5% DV May-25 to 2027** 7.80 100 **Ipca-
Hydroxychloroquine**

LEFLUNOMIDE

Tab 10 mg – **5% DV Dec-23 to 2026** 6.00 30 **Arava**

Tab 20 mg – **5% DV Dec-23 to 2026** 6.00 30 **Arava**

PENICILLAMINE

Tab 125 mg 67.23 100 D-Penamine

Tab 250 mg 110.12 100 D-Penamine

SODIUM AUROTHIOMALATE

Inj 10 mg in 0.5 ml ampoule

Inj 20 mg in 0.5 ml ampoule

Inj 50 mg in 0.5 ml ampoule

Drugs Affecting Bone Metabolism

Bisphosphonates

ALENDRONATE SODIUM

Tab 70 mg – **5% DV Jul-24 to 2026** 3.10 4 **Fosamax**

ALENDRONATE SODIUM WITH COLECALCIFEROL

Tab 70 mg with colecalciferol 5,600 iu – **5% DV Jul-24 to 2026** 1.99 4 **Fosamax Plus**

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial 32.49 1 Pamisol

Inj 6 mg per ml, 10 ml vial 88.11 1 Pamisol

Inj 9 mg per ml, 10 ml vial 94.34 1 Pamisol

RISEDRONATE SODIUM

Tab 35 mg – **5% DV Feb-26 to 2028** 3.00 4 **Risedronate Sandoz**

ZOLEDRONIC ACID

Inj 5 mg per 100 ml, bag – **5% DV Feb-26 to 2028** 19.45 1 **Zoledronic Acid Viatrix**

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

Other Drugs Affecting Bone Metabolism

DENOSUMAB – **Restricted** see terms [below](#)

Note: Denosumab inj 60 mg per 1 ml pre-filled syringe is Medsafe approved for use in osteoporosis. Denosumab inj 120 mg per 1.7 ml vial is Medsafe approved for use in hypercalcaemia of malignancy.

⚡ Inj 120 mg per 1.7 ml vial	500.00	1	Xgeva
⚡ Inj 60 mg per 1 ml pre-filled syringe.....	250.00	1	Prolia

➔ **Restricted** ([RS2097](#))

Initiation – Osteoporosis

All of the following:

- 1 The patient has established osteoporosis; and
- 2 Any of the following:
 - 2.1 History of one significant osteoporotic fracture demonstrated radiologically, with a documented T-Score less than or equal to -2.5, that incorporates BMD measured using dual-energy x-ray absorptiometry (DEXA); or
 - 2.2 History of one significant osteoporotic fracture, demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of logistical, technical or pathophysiological reasons; or
 - 2.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 2.4 Documented T-Score less than or equal to -3.0; or
 - 2.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm that incorporates BMD measured using DEXA; and
- 3 Any of the following:
 - 3.1 Bisphosphonates are contraindicated because the patient's creatinine clearance or eGFR is less than 35 mL/min; or
 - 3.2 The patient has experienced at least two symptomatic new fractures or a BMD loss greater than 2% per year, after at least 12 months' continuous therapy with a funded antiresorptive agent; or
 - 3.3 Bisphosphonates result in intolerable side effects; or
 - 3.4 Intravenous bisphosphonates cannot be administered due to logistical or technical reasons.

Initiation – Hypercalcaemia

Both:

- 1 Patient has hypercalcaemia of malignancy; and
- 2 Patient has severe renal impairment.

RALOXIFENE – **Restricted** see terms [below](#)

⚡ Tab 60 mg	53.76	28	Evista
-------------------	-------	----	--------

➔ **Restricted** ([RS1666](#))

Initiation

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score greater than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause - Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

TERIPARATIDE – **Restricted** see terms [below](#)

↓ Inj 250 mcg per ml, 2.4 ml	195.00	1	Teriparatide - Teva
------------------------------------	--------	---	---------------------

➔ **Restricted (RS1143)****Initiation***Limited to 18 months treatment*

All of the following:

- The patient has severe, established osteoporosis; and
- The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- The patient has had two or more fractures due to minimal trauma; and
- The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- Antiresorptive agents and their adequate doses for the purposes of this restriction are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Enzymes

HYALURONIDASE

Inj 1,500 iu ampoule

Hyperuricaemia and Antigout

ALLOPURINOL

Tab 100 mg – 5% DV Jun-24 to 2026	17.99	1,000	Ipca-Allopurinol
Tab 300 mg – 5% DV Jun-24 to 2026	22.50	500	Ipca-Allopurinol

BENZBROMARONE – **Restricted:** For continuation only

➔ Tab 50 mg			
➔ Tab 100 mg	45.00	100	Benzbromaron AL 100

MUSCULOSKELETAL SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
COLCHICINE			
Tab 500 mcg.....	6.00	100	Colgout
FEBUXOSTAT – Restricted see terms below			
⚡ Tab 80 mg – 5% DV Jun-24 to 2026	4.73	28	Febuxostat (Teva)
⚡ Tab 120 mg – 5% DV Jun-24 to 2026	11.78	28	Febuxostat (Teva)

➡ **Restricted (RS1844)**

Initiation – Gout

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout..

Initiation – Tumour lysis syndrome

Haematologist or oncologist

Re-assessment required after 6 weeks

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Continuation – Tumour lysis syndrome

Haematologist or oncologist

Re-assessment required after 6 weeks

The treatment remains appropriate and patient is benefitting from treatment.

PROBENECID

Tab 500 mg

RASBURICASE – Restricted see terms [below](#)

⚡ Inj 1.5 mg vial

➡ **Restricted (RS1016)**

Haematologist

Muscle Relaxants and Related Agents

ATRAURIUM BESYLATE

Inj 10 mg per ml, 2.5 ml ampoule – 5% DV Jun-25 to 2026 7.69

Inj 10 mg per ml, 5 ml ampoule – 5% DV Jun-25 to 2026 9.86

BACLOFEN

Tab 10 mg – 5% DV Dec-24 to 2027 3.70

Oral liq 1 mg per ml

Inj 0.05 mg per ml, 1 ml ampoule 11.55

Inj 2 mg per ml, 5 ml ampoule – 5% DV Mar-25 to 2027 490.91

CLOSTRIDIUM BOTULINUM TYPE A TOXIN

Inj 100 u vial 467.50

Inj 300 u vial 388.50

Inj 500 u vial 1,295.00

Medsurge

Medsurge

Pacifen

Lioresal Intrathecal

**Sintetica Baclofen
Intrathecal**

Botox

Dysport

Dysport

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DANTROLENE			
Cap 25 mg.....	145.77	100	Dantrium
Cap 50 mg.....	77.00	100	Dantrium
Inj 20 mg vial	1,143.74	6	Dantrium IV
MIVACURIUM CHLORIDE			
Inj 2 mg per ml, 10 ml ampoule			
ORPHENADRINE CITRATE			
Tab 100 mg – 5% DV Feb-25 to 2027	23.25	100	Norflex
PANCURONIUM BROMIDE			
Inj 2 mg per ml, 2 ml ampoule			
ROCURONIUM BROMIDE			
Inj 10 mg per ml, 5 ml ampoule	37.06	10	Hameln
SUXAMETHONIUM CHLORIDE			
Inj 50 mg per ml, 2 ml ampoule – 5% DV Feb-24 to 2026	35.40	10	Martindale
VECURONIUM BROMIDE			
Inj 10 mg vial – 5% DV Apr-25 to 2027	380.00	10	Vecure

Reversers of Neuromuscular Blockade

SUGAMMADEX – **Restricted** see terms [below](#)

⚠ Inj 100 mg per ml, 2 ml vial – 5% DV Dec-24 to 2027	80.64	10	Sugammadex BNM
⚠ Inj 100 mg per ml, 5 ml vial – 5% DV Dec-24 to 2027	201.60	10	Sugammadex BNM

➔ **Restricted (RS1370)**

Initiation

Any of the following:

- 1 Patient requires reversal of profound neuromuscular blockade following rapid sequence induction that has been undertaken using rocuronium (i.e. suxamethonium is contraindicated or undesirable); or
- 2 Severe neuromuscular degenerative disease where the use of neuromuscular blockade is required; or
- 3 Patient has an unexpectedly difficult airway that cannot be intubated and requires a rapid reversal of anaesthesia and neuromuscular blockade; or
- 4 The duration of the patient's surgery is unexpectedly short; or
- 5 Neostigmine or a neostigmine/anticholinergic combination is contraindicated (for example the patient has ischaemic heart disease, morbid obesity or COPD); or
- 6 Patient has a partial residual block after conventional reversal.

Non-Steroidal Anti-Inflammatory Drugs

CELECOXIB			
Cap 100 mg – 5% DV Feb-26 to 2028	3.60	60	Celebrex
	3.45		Celecoxib Pfizer
Cap 200 mg	3.20	30	Celecoxib Pfizer

(Celecoxib Pfizer Cap 100 mg to be delisted 1 February 2026)

MUSCULOSKELETAL SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DICLOFENAC SODIUM			
Tab EC 25 mg – 5% DV Feb-25 to 2027	2.19	50	Diclofenac Sandoz
Tab 50 mg dispersible	1.50	20	Voltaren D
Tab EC 50 mg – 5% DV Feb-25 to 2027	2.19	50	Diclofenac Sandoz
Tab long-acting 75 mg – 5% DV Aug-25 to 2028	10.00	100	Voltaren SR
Inj 25 mg per ml, 3 ml ampoule	13.20	5	Voltaren
Suppos 12.5 mg	2.04	10	Voltaren
Suppos 25 mg	2.44	10	Voltaren
Suppos 50 mg	4.22	10	Voltaren
Suppos 100 mg	7.00	10	Voltaren
ETORICOXIB – Restricted see terms below			
⚡ Tab 30 mg			
⚡ Tab 60 mg			
⚡ Tab 90 mg			
⚡ Tab 120 mg			
➡ Restricted (RS1592)			
Initiation			
For in-vivo investigation of allergy only.			
IBUPROFEN			
Tab 200 mg - 1,000 tablet pack – 1% DV Feb-21 to 2026	21.40	1,000	Relieve
➡ Tab 400 mg – Restricted: For continuation only			
➡ Tab 600 mg – Restricted: For continuation only			
Tab long-acting 800 mg – 5% DV Apr-25 to 2027	3.65	30	Ibuprofen SR BNM
Oral liq 20 mg per ml – 5% DV Apr-25 to 2027	2.85	200 ml	Ethics
Inj 5 mg per ml, 2 ml ampoule			
Inj 10 mg per ml, 2 ml vial			
INDOMETACIN [INDOMETHACIN]			
Cap 25 mg			
Cap 50 mg			
Cap long-acting 75 mg			
Inj 1 mg vial			
Suppos 100 mg			
KETOPROFEN			
Cap long-acting 200 mg	12.07	28	Oruvail SR
MEFENAMIC ACID – Restricted: For continuation only			
➡ Cap 250 mg			
NAPROXEN			
Tab 250 mg – 5% DV Feb-25 to 2027	39.23	500	Noflam 250
Tab 500 mg – 5% DV Feb-25 to 2027	34.45	250	Noflam 500
Tab long-acting 750 mg – 5% DV Feb-25 to 2027	10.40	28	Naprosyn SR 750
Tab long-acting 1 g – 5% DV Feb-25 to 2027	11.50	28	Naprosyn SR 1000
PARECOXIB			
Inj 40 mg vial – 5% DV Dec-24 to 2027	46.00	10	Dynastat
SULINDAC			
Tab 100 mg			
Tab 200 mg			
TENOXICAM			
Tab 20 mg – 5% DV Feb-26 to 2028	23.50	100	Tilcotil
Inj 20 mg vial	9.95	1	AFT

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Topical Products for Joint and Muscular Pain

CAPSAICIN – Restricted see terms below			
↓ Crm 0.025%.....	9.75	45 g	Zo-Rub Osteo Zostrix

➡ **Restricted (RS1309)**

Initiation

Patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Agents for Parkinsonism and Related Disorders

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE – Restricted see terms below			
⚠ Tab 50 mg – 5% DV Feb-25 to 2027	117.00	56	Rilutek
➡ Restricted (RS1351)			

Initiation
Neurologist or respiratory specialist
Re-assessment required after 6 months
All of the following:
1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
3 The patient has not undergone a tracheostomy; and
4 The patient has not experienced respiratory failure; and
5 Any of the following:
5.1 The patient is ambulatory; or
5.2 The patient is able to use upper limbs; or
5.3 The patient is able to swallow.

Continuation
Re-assessment required after 18 months
All of the following:
1 The patient has not undergone a tracheostomy; and
2 The patient has not experienced respiratory failure; and
3 Any of the following:
3.1 The patient is ambulatory; or
3.2 The patient is able to use upper limbs; or
3.3 The patient is able to swallow.

TETRABENAZINE			
Tab 25 mg – 5% DV Feb-26 to 2028	126.02	112	Motetis

Anticholinergics

BENZATROPINE MESYLATE			
Tab 2 mg	10.99	60	Benztrop
Inj 1 mg per ml, 2 ml ampoule	95.00	5	Phebra
PROCYCLIDINE HYDROCHLORIDE			
Tab 5 mg			

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE			
Cap 100 mg	38.24	60	Symmetrel
APOMORPHINE HYDROCHLORIDE			
Inj 10 mg per ml, 2 ml ampoule	59.50	5	Movapo
Inj 10 mg per ml, 5 ml ampoule	121.84	5	Movapo
BROMOCRIPTINE			
Cap 5 mg			
ENTACAPONE			
Tab 200 mg – 5% DV Jul-25 to 2027	13.73	100	Entacapone Viatris

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LEVODOPA WITH BENSERAZIDE			
Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	Madopar Rapid
Cap 50 mg with benserazide 12.5 mg	13.75	100	Madopar 62.5
Cap 100 mg with benserazide 25 mg	15.80	100	Madopar 125
Cap long-acting 100 mg with benserazide 25 mg	22.85	100	Madopar HBS
Cap 200 mg with benserazide 50 mg	26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
Tab 100 mg with carbidopa 25 mg – 5% DV Feb-25 to 2027	26.49	100	Sinemet
Tab long-acting 100 mg with carbidopa 25 mg			
Tab long-acting 200 mg with carbidopa 50 mg – 5% DV Feb-25 to 2027	44.99	100	Sinemet CR
Tab 250 mg with carbidopa 25 mg – 5% DV Feb-25 to 2027	39.49	100	Sinemet
LEVODOPA WITH CARBIDOPA AND ENTACAPONE			
Tab 50 mg with carbidopa 12.5 mg and entacapone 200 mg – 5% DV Jul-25 to 2027	27.01	100	Stalevo
Tab 100 mg with carbidopa 25 mg and entacapone 200 mg – 5% DV Jul-25 to 2027	34.18	100	Stalevo
Tab 150 mg with carbidopa 37.5 mg and entacapone 200 mg – 5% DV Jul-25 to 2027	44.96	100	Stalevo
Tab 200 mg with carbidopa 50 mg and entacapone 200 mg – 5% DV Jul-25 to 2027	51.23	100	Stalevo
PRAMIPEXOLE HYDROCHLORIDE			
Tab 0.25 mg – 5% DV Dec-25 to 2028	5.23	100	Ramipex
Tab 1 mg – 5% DV Dec-25 to 2028	17.73	100	Ramipex
RASAGILINE			
Tab 1 mg	53.50	30	Azilect
ROPINIROLE HYDROCHLORIDE			
Tab 0.25 mg	4.05	84	Ropin
Tab 1 mg	4.95	84	Ropin
Tab 2 mg	6.48	84	Ropin
Tab 5 mg	14.50	84	Ropin
SELEGILINE HYDROCHLORIDE – Restricted: For continuation only			
➔ Tab 5 mg			
TOLCAPONE			
Tab 100 mg	152.38	100	Tasmar

Anaesthetics

General Anaesthetics

DESFLURANE			
Soln for inhalation 100%, 240 ml bottle	1,350.00	6	Suprane
DEXMEDETOMIDINE			
Inj 100 mcg per ml, 2 ml vial – 5% DV May-24 to 2026	42.00	5	Dexmedetomidine Viatris
ETOMIDATE			
Inj 2 mg per ml, 10 ml ampoule			
ISOFLURANE			
Soln for inhalation 100%, 250 ml bottle	2,730.00	6	Aerrane

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
KETAMINE			
Inj 1 mg per ml, 100 ml bag	146.00	5	Biomed
Inj 10 mg per ml, 10 ml syringe	76.00	5	Biomed
Inj 100 mg per ml, 2 ml vial	36.23	5	Ketalar Ketamine-Baxter
<i>(Ketamine-Baxter Inj 100 mg per ml, 2 ml vial to be delisted 1 November 2025)</i>			
METHOHEXITAL SODIUM			
Inj 10 mg per ml, 50 ml vial			
PROPOFOL			
Inj 10 mg per ml, 20 ml ampoule – 5% DV Feb-26 to 2028	5.75	5	Fresofol 1% MCT/LCT
Inj 10 mg per ml, 50 ml vial – 5% DV Feb-26 to 2028	27.50	10	Fresofol 1% MCT/LCT
Inj 10 mg per ml, 100 ml vial – 5% DV Feb-26 to 2028	39.90	10	Fresofol 1% MCT/LCT
SEVOFLURANE			
Soln for inhalation 100%, 250 ml bottle	930.00	6	Baxter
THIOPENTAL [THIOPENTONE] SODIUM			
Inj 500 mg ampoule			
Local Anaesthetics			
ARTICAINE HYDROCHLORIDE			
Inj 1%			
ARTICAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 4% with adrenaline 1:100,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:100,000, 1.8 ml dental cartridge			
Inj 4% with adrenaline 1:100,000, 2.2 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:200,000 1.8 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 2.2 ml dental cartridge			
BENZOCAINE			
Gel 20%			
BENZOCAINE WITH TETRACAINE HYDROCHLORIDE			
Gel 18% with tetracaine hydrochloride 2%			<i>e.g. ZAP Topical Anaesthetic Gel</i>
BUPIVACAINE HYDROCHLORIDE			
Inj 5 mg per ml, 4 ml ampoule – 5% DV Feb-24 to 2026	62.50	5	Marcaïn Isobaric
Inj 2.5 mg per ml, 20 ml ampoule			
Inj 2.5 mg per ml, 20 ml ampoule sterile pack – 5% DV Feb-24 to 2026	28.00	5	Marcaïn
Inj 5 mg per ml, 10 ml ampoule sterile pack	16.20	5	Marcaïn
Inj 5 mg per ml, 20 ml ampoule			
Inj 5 mg per ml, 20 ml ampoule sterile pack	16.56	5	Marcaïn
Inj 1.25 mg per ml, 100 ml bag			
Inj 1.25 mg per ml, 200 ml bag			
Inj 2.5 mg per ml, 100 ml bag	150.00	5	Marcaïn
Inj 2.5 mg per ml, 200 ml bag			
Inj 1.25 mg per ml, 500 ml bag			
<i>(Marcaïn Inj 2.5 mg per ml, 100 ml bag to be delisted 1 November 2025)</i>			
BUPIVACAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 2.5 mg per ml with adrenaline 1:200,000, 10 ml ampoule			
Inj 2.5 mg per ml with adrenaline 1:400,000, 20 ml vial	94.50	5	Marcaïn with Adrenaline
Inj 5 mg per ml with adrenaline 1:200,000, 20 ml vial	80.50	5	Marcaïn with Adrenaline

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
BUPIVACAINE HYDROCHLORIDE WITH FENTANYL			
Inj 0.625 mg with fentanyl 2 mcg per ml, 100 ml bag			
Inj 0.625 mg with fentanyl 2 mcg per ml, 200 ml bag	165.00	5	Biomed
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml bag – 5% DV Feb-26 to 2028	135.00	5	Bupafen
Inj 1.25 mg with fentanyl 2 mcg per ml, 200 ml bag – 5% DV Feb-26 to 2028	138.00	5	Bupafen
Inj 1.25 mg with fentanyl 2 mcg per ml, 50 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 20 ml syringe.....	57.35	5	Biomed
BUPIVACAINE HYDROCHLORIDE WITH GLUCOSE			
Inj 0.5% with glucose 8%, 4 ml ampoule – 5% DV Dec-25 to 2028	21.40	5	Marcaïn Heavy
COCAINE HYDROCHLORIDE			
Paste 5%			
Soln 15%, 2 ml syringe			
Soln 4%, 2 ml syringe.....	30.77	1	Biomed
COCAINE HYDROCHLORIDE WITH ADRENALINE			
Paste 15% with adrenaline 0.06%			
Paste 25% with adrenaline 0.06%			
ETHYL CHLORIDE			
Spray 100%			
LIDOCAINE [LIGNOCAINE]			
Crm 4%.....	5.40	5 g	LMX4
	27.00	30 g	LMX4
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Gel 2%.....	4.87	20 g	Orion
Soln 4%			
Spray 10% – 5% DV Feb-26 to 2028	82.90	50 ml	Xylocaine
Oral (gel) soln 2%.....	44.00	200 ml	Mucosoothe
Inj 1%, 20 ml ampoule, sterile pack			
Inj 2%, 20 ml ampoule, sterile pack			
Inj 1%, 5 ml ampoule	15.00	25	Lidocaine-Baxter
Inj 1%, 20 ml vial	19.50	5	Lidocaine-Baxter
Inj 2%, 5 ml ampoule	27.50	25	Lidocaine-Baxter
Inj 2%, 20 ml vial	14.00	5	Lidocaine-Baxter
Inj 10%, 5 ml ampoule			
Gel 2%, 11 ml urethral syringe – 5% DV Feb-26 to 2028	65.45	10	Instillagel Lido
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE			
Inj 1% with adrenaline 1:100,000, 20 ml vial			
Inj 1% with adrenaline 1:100,000, 5 ml ampoule – 5% DV Dec-25 to 2028	32.00	10	Xylocaine
Inj 1% with adrenaline 1:200,000, 20 ml vial	50.00	5	Xylocaine
Inj 2% with adrenaline 1:100,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 2.2 ml dental cartridge			
Inj 2% with adrenaline 1:200,000, 20 ml vial	60.00	5	Xylocaine
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE AND TETRACAINE HYDROCHLORIDE			
Soln 4% with adrenaline 0.1% and tetracaine hydrochloride 0.5%, 5 ml syringe.....	20.50	1	Topicaïne

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH PHENYLEPHRINE HYDROCHLORIDE			
Nasal spray 5% with phenylephrine hydrochloride 0.5%			
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE			
Crn 2.5% with prilocaine 2.5%.....	45.00	30 g	EMLA
Patch 25 mcg with prilocaine 25 mcg	115.00	20	EMLA
Crn 2.5% with prilocaine 2.5%, 5 g.....	45.00	5	EMLA
MEPIVACAINE HYDROCHLORIDE			
Inj 3%, 1.8 ml dental cartridge	43.60	50	Scandonest 3%
Inj 3%, 2.2 ml dental cartridge	43.60	50	Scandonest 3%
MEPIVACAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 2% with adrenaline 1:100,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:100,000, 2.2 ml dental cartridge			
PRILOCAINE HYDROCHLORIDE			
Inj 0.5%, 50 ml vial	100.00	5	Citanest
Inj 2%, 5 ml ampoule			
PRILOCAINE HYDROCHLORIDE WITH FELYPRESSIN			
Inj 3% with felypressin 0.03 iu per ml, 1.8 ml dental cartridge			
Inj 3% with felypressin 0.03 iu per ml, 2.2 ml dental cartridge			
ROPIVACAINE HYDROCHLORIDE			
Inj 2 mg per ml, 10 ml ampoule – 5% DV Feb-24 to 2026	9.80	5	Ropivacaine Kabi
Inj 2 mg per ml, 20 ml ampoule – 5% DV Feb-24 to 2026	10.25	5	Ropivacaine Kabi
Inj 2 mg per ml, 100 ml bag – 5% DV Feb-24 to 2026	32.85	5	Ropivacaine Kabi
Inj 2 mg per ml, 200 ml bag – 5% DV Feb-24 to 2026	43.40	5	Ropivacaine Kabi
Inj 7.5 mg per ml, 10 ml ampoule – 5% DV Feb-24 to 2026	11.00	5	Ropivacaine Kabi
Inj 7.5 mg per ml, 20 ml ampoule – 5% DV Feb-24 to 2026	13.50	5	Ropivacaine Kabi
Inj 10 mg per ml, 10 ml ampoule – 5% DV Feb-24 to 2026	11.75	5	Ropivacaine Kabi
Inj 10 mg per ml, 20 ml ampoule – 5% DV Feb-24 to 2026	17.60	5	Ropivacaine Kabi
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Gel 4%			

Analgesics

Non-Opioid Analgesics

ASPIRIN			
Tab dispersible 300 mg – 5% DV May-24 to 2026	5.65	100	Ethics Aspirin
CAPSAICIN – Restricted see terms below			
‡ Crn 0.075%.....	11.95	45 g	Zo-Rub HP Zostrix HP
➡ Restricted (RS1145)			
Initiation			
For post-herpetic neuralgia or diabetic peripheral neuropathy.			
METHOXYFLURANE – Restricted see terms below			
‡ Soln for inhalation 99.9%, 3 ml bottle			
➡ Restricted (RS1292)			
Initiation			
Both:			
1 Patient is undergoing a painful procedure with an expected duration of less than one hour; and			
2 Only to be used under supervision by a medical practitioner or nurse who is trained in the use of methoxyflurane.			

‡ Item restricted (see ➡ above); ‡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NEFOPAM HYDROCHLORIDE			
Tab 30 mg			
PARACETAMOL – Some items restricted see terms below			
Tab soluble 500 mg			
Tab 500 mg - blister pack - 1,000 tablet pack – 1% DV Feb-22 to 2026	19.75	1,000	Pacimol
Tab 500 mg - blister pack - 12 tablet pack			
Tab 500 mg - blister pack - 20 tablet pack			
Tab 500 mg - bottle pack – 1% DV Feb-22 to 2026	17.92	1,000	Noumed Paracetamol
Oral liq 120 mg per 5 ml	3.98	200 ml	Paracetamol (Ethics)
Oral liq 250 mg per 5 ml	3.35	200 ml	Pamol
↓ Inj 10 mg per ml, 100 ml vial – 5% DV Feb-26 to 2028	15.30	10	Paracetamol Kabi
Suppos 25 mg			
Suppos 50 mg			
Suppos 125 mg – 5% DV Feb-24 to 2026	4.29	10	Gacet
Suppos 250 mg – 5% DV Feb-24 to 2026	5.39	10	Gacet
Suppos 500 mg – 5% DV Feb-24 to 2026	16.55	50	Gacet

→ **Restricted (RS1146)**

Initiation

Intravenous paracetamol is only to be used where other routes are unavailable or impractical, or where there is reduced absorption. The need for IV paracetamol must be re-assessed every 24 hours.

SUCROSE

Oral liq 25%	14.61	25 ml	Biomed
↓ Oral liq 66.7% (preservative free)			

→ **Restricted (RS1763)**

Initiation

For use in neonatal patients only.

Opioid Analgesics

ALFENTANIL

Inj 0.5 mg per ml, 2 ml ampoule – 5% DV Feb-24 to 2026	8.99	5	Medsurge
---	------	---	-----------------

CODEINE PHOSPHATE

Tab 15 mg – 5% DV Dec-25 to 2028	5.82	100	Noumed
Tab 30 mg – 5% DV Dec-25 to 2028	6.88	100	Noumed
Tab 60 mg – 5% DV Dec-25 to 2028	13.89	100	Noumed

DIHYDROCODEINE TARTRATE

Tab long-acting 60 mg – 5% DV Feb-26 to 2028	9.20	60	DHC Continus
---	------	----	---------------------

FENTANYL

Inj 10 mcg per ml, 10 ml syringe – 5% DV Feb-25 to 2027	44.50	5	Biomed Fentanyl
Inj 50 mcg per ml, 2 ml ampoule – 5% DV May-25 to 2027	4.25	10	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule – 5% DV May-25 to 2027	9.41	10	Boucher and Muir
Inj 10 mcg per ml, 100 ml bag – 5% DV Feb-24 to 2026	114.25	5	Biomed
Inj 20 mcg per ml, 50 ml syringe – 5% DV Feb-25 to 2027	136.50	5	Biomed
Inj 20 mcg per ml, 100 ml bag			
Patch 12 mcg per hour – 5% DV May-25 to 2027	6.02	5	Fentanyl Sandoz
Patch 12.5 mcg per hour	6.02	5	Fentanyl Sandoz
Patch 25 mcg per hour – 5% DV Dec-24 to 2027	6.91	5	Fentanyl Sandoz
Patch 50 mcg per hour – 5% DV Dec-24 to 2027	9.28	5	Fentanyl Sandoz
Patch 75 mcg per hour – 5% DV Dec-24 to 2027	15.50	5	Fentanyl Sandoz
Patch 100 mcg per hour – 5% DV Dec-24 to 2027	16.37	5	Fentanyl Sandoz

(Fentanyl Sandoz Patch 12.5 mcg per hour to be delisted 1 January 2026)

NERVOUS SYSTEM

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METHADONE HYDROCHLORIDE			
Tab 5 mg	1.45	10	Methadone BNM
Oral liq 2 mg per ml – 5% DV Feb-25 to 2027	7.80	200 ml	Biodone
Oral liq 5 mg per ml – 5% DV Feb-25 to 2027	7.80	200 ml	Biodone Forte
Oral liq 10 mg per ml – 5% DV Feb-25 to 2027	9.65	200 ml	Biodone Extra Forte
Inj 10 mg per ml, 1 ml vial.....	72.99	10	AFT
MORPHINE HYDROCHLORIDE			
Oral liq 1 mg per ml	19.00	200 ml	RA-Morph
Oral liq 2 mg per ml	23.55	200 ml	RA-Morph
Oral liq 5 mg per ml	28.20	200 ml	RA-Morph
Oral liq 10 mg per ml	40.25	200 ml	RA-Morph
MORPHINE SULPHATE			
Tab immediate-release 10 mg.....	2.80	10	Sevredol
Tab immediate-release 20 mg.....	5.52	10	Sevredol
Cap long-acting 10 mg	3.00	10	m-Eslon
Cap long-acting 30 mg	4.30	10	m-Eslon
Cap long-acting 60 mg	9.00	10	m-Eslon
Cap long-acting 100 mg	10.50	10	m-Eslon
Oral liq 2 mg per ml	42.56	300 ml	Oramorph
	29.80	100 ml	Oramorph CDC S29
	16.31		Wockhardt
Inj 1 mg per ml, 100 ml bag – 5% DV Feb-24 to 2026	114.25	5	Biomed
Inj 1 mg per ml, 10 ml syringe – 5% DV Feb-24 to 2026	27.25	5	Biomed
Inj 1 mg per ml, 50 ml syringe – 5% DV Feb-24 to 2026	63.75	5	Biomed
Inj 1 mg per ml, 2 ml syringe			
Inj 5 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	5.96	5	Medsurge
Inj 10 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	4.99	5	Medsurge
Inj 10 mg per ml, 100 mg cassette			
Inj 10 mg per ml, 100 ml bag			
Inj 15 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	6.93	5	Medsurge
Inj 30 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	7.28	5	Medsurge
Inj 200 mcg in 0.4 ml syringe			
Inj 300 mcg in 0.3 ml syringe			
MORPHINE TARTRATE			
Inj 80 mg per ml, 1.5 ml ampoule			
OXYCODONE HYDROCHLORIDE			
Tab controlled-release 5 mg – 5% DV Dec-24 to 2027	2.49	20	Oxycodone Sandoz
Tab immediate-release 5 mg.....	13.77	100	Oxycodone Amneal
Tab controlled-release 10 mg – 5% DV Dec-24 to 2027	2.49	20	Oxycodone Sandoz
Tab immediate-release 10 mg.....	18.77	100	Oxycodone Amneal
Tab controlled-release 20 mg – 5% DV Dec-24 to 2027	3.41	20	Oxycodone Sandoz
Tab immediate-release 20 mg.....	26.77	100	Oxycodone Amneal
Tab controlled-release 40 mg – 5% DV Dec-24 to 2027	6.67	20	Oxycodone Sandoz
Tab controlled-release 80 mg – 5% DV Dec-24 to 2027	12.99	20	Oxycodone Sandoz
Oral liq 1 mg per ml	37.08	250 ml	Oxycodone Lucis S29
Inj 1 mg per ml, 100 ml bag			
Inj 10 mg per ml, 1 ml ampoule – 5% DV Dec-24 to 2027	4.37	5	Hameln
Inj 10 mg per ml, 2 ml ampoule – 5% DV Dec-24 to 2027	8.62	5	Hameln
Inj 50 mg per ml, 1 ml ampoule – 5% DV Dec-24 to 2027	14.90	5	Hameln

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PARACETAMOL WITH CODEINE			
Tab paracetamol 500 mg with codeine phosphate 8 mg – 5% DV Feb-26 to 2028	31.95	1,000	Paracetamol + Codeine (Relieve)
PETHIDINE HYDROCHLORIDE			
Tab 50 mg – 5% DV Feb-26 to 2028	8.68	10	Noumed Pethidine
Inj 5 mg per ml, 10 ml syringe			
Inj 5 mg per ml, 100 ml bag			
Inj 10 mg per ml, 100 ml bag			
Inj 10 mg per ml, 50 ml syringe			
Inj 50 mg per ml, 1 ml ampoule	29.88	5	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule	30.72	5	DBL Pethidine Hydrochloride
REMIFENTANIL			
Inj 1 mg vial – 5% DV Feb-24 to 2026	14.95	5	Remifentanil-AFT
Inj 2 mg vial – 5% DV Feb-24 to 2026	20.95	5	Remifentanil-AFT
TRAMADOL HYDROCHLORIDE			
Tab sustained-release 100 mg – 5% DV May-24 to 2026	1.95	20	Tramal SR 100
Tab sustained-release 150 mg – 5% DV May-24 to 2026	2.95	20	Tramal SR 150
Tab sustained-release 200 mg – 5% DV May-24 to 2026	3.80	20	Tramal SR 200
Cap 50 mg – 5% DV Jan-24 to 2026	3.33	100	Arrow-Tramadol
Oral soln 10 mg per ml			
Inj 10 mg per ml, 100 ml bag			
Inj 50 mg per ml, 1 ml ampoule – 5% DV May-24 to 2026	10.00	5	Tramal 50
Inj 50 mg per ml, 2 ml ampoule – 5% DV May-24 to 2026	9.00	5	Tramal 100
Antidepressants			
Cyclic and Related Agents			
AMITRIPTYLINE			
Tab 10 mg – 5% DV Mar-24 to 2026	2.99	100	Arrow-Amitriptyline
Tab 25 mg – 5% DV Mar-24 to 2026	1.99	100	Arrow-Amitriptyline
Tab 50 mg – 5% DV Mar-24 to 2026	3.14	100	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE			
Tab 25 mg – 5% DV Jul-25 to 2027	16.99	50	APO Clomipramine
Cap 10 mg	35.50	28	Clomipramine Teva
<i>(Clomipramine Teva Cap 10 mg to be delisted 1 April 2026)</i>			
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Restricted: For continuation only			
➔ Tab 75 mg	3.85	30	Dosulepin Viatrix
➔ Cap 25 mg	7.83	50	Dosulepin Viatrix
DOXEPIN HYDROCHLORIDE – Restricted: For continuation only			
➔ Cap 10 mg			
➔ Cap 25 mg			
➔ Cap 50 mg			
IMIPRAMINE HYDROCHLORIDE			
Tab 10 mg	5.48	50	Tofranil
	6.58	60	Tofranil
Tab 25 mg	4.93	28	Imipramine Crescent
	8.80	50	Tofranil

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MAPROTILINE HYDROCHLORIDE – Restricted: For continuation only			
➡ Tab 25 mg			
➡ Tab 75 mg			
MIANSERIN HYDROCHLORIDE – Restricted: For continuation only			
➡ Tab 30 mg			
NORTRIPTYLINE HYDROCHLORIDE			
Tab 10 mg	2.24	50	Allegron
	2.46	100	Norpress
Tab 25 mg	2.95	50	Allegron
	6.29	180	Norpress
Monoamine-Oxidase Inhibitors - Non-Selective			
PHENELZINE SULPHATE			
Tab 15 mg			
TRANLYCYPROMINE SULPHATE			
Tab 10 mg			
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE			
Tab 150 mg – 5% DV Feb-25 to 2027	23.60	60	Aurorix
Tab 300 mg – 5% DV Feb-25 to 2027	38.50	60	Aurorix
Other Antidepressants			
MIRTAZAPINE			
Tab 30 mg – 5% DV Jan-26 to 2028	2.34	30	Noumed
Tab 45 mg – 5% DV Jan-26 to 2028	3.10	30	Noumed
VENLAFAXINE			
Cap 37.5 mg	8.29	84	Enlafax XR
Cap 75 mg	10.32	84	Enlafax XR
Cap 150 mg	13.95	84	Enlafax XR
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE			
Tab 20 mg	2.86	84	Celapram
ESCITALOPRAM			
Tab 10 mg – 5% DV Apr-24 to 2026	0.79	28	Ipca-Escitalopram
Tab 20 mg – 5% DV Apr-24 to 2026	1.49	28	Ipca-Escitalopram
FLUOXETINE HYDROCHLORIDE			
Tab dispersible 20 mg, scored	2.50	28	Fluox
Cap 20 mg	3.13	90	Arrow-Fluoxetine
PAROXETINE			
Tab 20 mg – 5% DV Feb-26 to 2028	4.98	90	Loxamine
	1.66	30	Paxtine
SERTRALINE			
Tab 50 mg	0.99	30	Setrona
Tab 100 mg	1.74	30	Setrona

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antiepilepsy Drugs			
Agents for the Control of Status Epilepticus			
CLONAZEPAM			
Inj 1 mg per ml, 1 ml ampoule			
DIAZEPAM			
Inj 5 mg per ml, 2 ml ampoule	27.92	5	Hospira
Rectal tubes 5 mg.....	54.58	5	Stesolid
Rectal tubes 10 mg			
LORAZEPAM			
Inj 2 mg vial			
Inj 4 mg per ml, 1 ml vial			
PARALDEHYDE			
Soln 97%			
Inj 5 ml ampoule			
PHENYTOIN SODIUM			
Inj 50 mg per ml, 2 ml ampoule	104.58	5	Hospira
Inj 50 mg per ml, 5 ml ampoule	154.01	5	Hospira
<i>(Hospira Inj 50 mg per ml, 2 ml ampoule to be delisted 1 February 2026)</i>			
Control of Epilepsy			
CARBAMAZEPINE			
Tab 200 mg	14.53	100	Tegretol
Tab long-acting 200 mg.....	16.98	100	Tegretol CR
Tab 400 mg	34.58	100	Tegretol
Tab long-acting 400 mg.....	39.17	100	Tegretol CR
Oral liq 20 mg per ml	26.37	250 ml	Tegretol
CLOBAZAM			
Tab 10 mg			
CLONAZEPAM			
Oral drops 2.5 mg per ml			
ETHOSUXIMIDE			
Cap 250 mg	140.88	100	Zarontin
Oral liq 50 mg per ml	56.35	200 ml	Zarontin
GABAPENTIN			
Note: Gabapentin not to be given in combination with pregabalin			
Cap 100 mg – 1% DV Feb-22 to 2027	6.45	100	Nupentin
Cap 300 mg – 1% DV Feb-22 to 2027	8.45	100	Nupentin
Cap 400 mg – 1% DV Feb-22 to 2027	10.26	100	Nupentin
LACOSAMIDE – Restricted see terms on the next page			
⚡ Tab 50 mg	25.04	14	Vimpat
⚡ Tab 100 mg	50.06	14	Vimpat
	200.24	56	Vimpat
⚡ Tab 150 mg	75.10	14	Vimpat
	300.40	56	Vimpat
⚡ Tab 200 mg	400.55	56	Vimpat
⚡ Inj 10 mg per ml, 20 ml vial			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
➔ Restricted (RS1988)			
Initiation			
<i>Re-assessment required after 15 months</i>			
Both:			
1 Patient has focal epilepsy; and			
2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam, and any two of carbamazepine, lamotrigine, and phenytoin sodium (see Note).			
Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.			
Continuation			
Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.			
LAMOTRIGINE			
Tab dispersible 2 mg	55.00	30	Lamictal
Tab dispersible 5 mg	50.00	30	Lamictal
Tab dispersible 25 mg	4.20	56	Logem
Tab dispersible 50 mg	5.11	56	Logem
Tab dispersible 100 mg	6.75	56	Logem
LEVETIRACETAM			
Tab 250 mg	5.84	60	Everet
Tab 500 mg	10.51	60	Everet
Tab 750 mg	16.71	60	Everet
Tab 1,000 mg	21.82	60	Everet
Oral liq 100 mg per ml	44.78	300 ml	Levetiracetam-AFT
Inj 100 mg per ml, 5 ml vial.....	38.95	10	Levetiracetam-AFT
PHENOBARBITONE			
Tab 15 mg	248.50	500	Noumed Phenobarbitone
Tab 30 mg	398.50	500	Noumed Phenobarbitone
PHENYTOIN			
Tab 50 mg			
PHENYTOIN SODIUM			
Cap 30 mg			
Cap 100 mg			
Oral liq 6 mg per ml			
PREGABALIN			
Note: Pregabalin not to be given in combination with gabapentin			
Cap 25 mg	2.25	56	Lyrica Pregabalin Pfizer
Cap 75 mg	2.65	56	Lyrica Pregabalin Pfizer
Cap 150 mg	4.01	56	Lyrica Pregabalin Pfizer
Cap 300 mg	7.38	56	Lyrica Pregabalin Pfizer
PRIMIDONE			
Tab 250 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM VALPROATE			
Tab 100 mg			
Tab EC 200 mg			
Tab EC 500 mg			
Oral liq 40 mg per ml			
Inj 100 mg per ml, 4 ml vial.....	9.98	1	Epilim IV
STIRIPENTOL – Restricted see terms below			
↓ Cap 250 mg.....	509.29	60	Diacomit
↓ Powder for oral liq 250 mg sachet.....	509.29	60	Diacomit
➔ Restricted (RS1989)			
Initiation			
Paediatric neurologist			
<i>Re-assessment required after 6 months</i>			
Both:			
1 Patient has confirmed diagnosis of Dravet syndrome; and			
2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.			
Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.			
Continuation			
Paediatric neurologist			
Patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.			
TOPIRAMATE			
Tab 25 mg	11.07	60	Arrow-Topiramate
	26.04		Topamax
	11.07		Topiramate Actavis
Tab 50 mg	18.81	60	Arrow-Topiramate
	44.26		Topamax
	18.81		Topiramate Actavis
Tab 100 mg	31.99	60	Arrow-Topiramate
	75.25		Topamax
	31.99		Topiramate Actavis
Tab 200 mg	55.19	60	Arrow-Topiramate
	129.85		Topamax
	55.19		Topiramate Actavis
Cap sprinkle 15 mg.....	20.84	60	Topamax
Cap sprinkle 25 mg.....	26.04	60	Topamax
VIGABATRIN – Restricted see terms below			
↓ Tab 500 mg			
↓ Powder for oral soln 500 mg per sachet.....	71.58	60	Sabril
➔ Restricted (RS1865)			
Initiation			
<i>Re-assessment required after 15 months</i>			
Both:			
1 Any of the following:			
1.1 Patient has infantile spasms; or			
1.2 Both:			
1.2.1 Patient has epilepsy; and			

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

1.2.2 Either:

1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or

1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or

1.3 Patient has tuberous sclerosis complex; and

2 Either:

2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or

2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Continuation

Both:

1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and

2 Either:

2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or

2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Antimigraine Preparations

Acute Migraine Treatment

DIHYDROERGOTAMINE MESYLATE

Inj 1 mg per ml, 1 ml ampoule

METOCLOPRAMIDE HYDROCHLORIDE WITH PARACETAMOL

Tab 5 mg with paracetamol 500 mg

RIZATRIPTAN

Tab orodispersible 10 mg – 5% DV Feb-24 to 2026 4.84 30 Rizamelt

SUMATRIPTAN

Tab 50 mg – 1% DV Feb-22 to 2027 14.41 90 Sumagran

Tab 100 mg – 1% DV Feb-22 to 2027 22.68 90 Sumagran

Inj 12 mg per ml, 0.5 ml prefilled pen – 5% DV Dec-25 to 2028 29.80 2 Clustran

Prophylaxis of Migraine

PIZOTIFEN

Tab 500 mcg 23.21 100 Sandomigran

Antinausea and Vertigo Agents

APREPITANT – **Restricted** see terms [below](#)

⚡ Cap 2 x 80 mg and 1 x 125 mg – 5% DV Jan-25 to 2027 21.90 3 Emend Tri-Pack

➡ **Restricted (RS1154)**

Initiation

Patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

Tab 16 mg – 5% DV Dec-23 to 2026 3.70 100 Serc

CYCLIZINE HYDROCHLORIDE

Tab 50 mg – 5% DV Feb-25 to 2027 0.66 10 Nausicalm

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CYCLIZINE LACTATE			
Inj 50 mg per ml, 1 ml ampoule	16.36	10	Hameln
DOMPERIDONE			
Tab 10 mg – 5% DV Dec-25 to 2028	3.80	100	Domperidone Viatris
DROPERIDOL			
Inj 2.5 mg per ml, 1 ml ampoule – 5% DV Feb-26 to 2028	28.68	10	Droperidol Medsurge
	43.85		Droperidol Panpharma
<i>(Droperidol Panpharma Inj 2.5 mg per ml, 1 ml ampoule to be delisted 1 February 2026)</i>			
GRANISETRON			
Inj 1 mg per ml, 3 ml ampoule – 5% DV Feb-24 to 2026	1.20	1	Deva
HYOSCINE HYDROBROMIDE			
Inj 400 mcg per ml, 1 ml ampoule			
↓ Patch 1 mg per 72 hours	88.50	10	Scopolamine Transdermal System Viatris
➔ Restricted (RS1155)			
Initiation			
Any of the following:			
1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or			
2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective; or			
3 For treatment of post-operative nausea and vomiting where cyclizine, droperidol and a 5HT3 antagonist have proven ineffective, are not tolerated or are contraindicated.			
METOCLOPRAMIDE HYDROCHLORIDE			
Tab 10 mg – 5% DV Mar-24 to 2026	1.57	100	Metoclopramide Actavis 10
Oral liq 5 mg per 5 ml			
Inj 5 mg per ml, 2 ml ampoule	7.00	10	Baxter
ONDANSETRON			
Tab 4 mg – 5% DV Dec-25 to 2028	1.95	50	Periset
Tab dispersible 4 mg – 5% DV Mar-24 to 2026	0.56	10	Periset ODT
Tab 8 mg – 5% DV Dec-25 to 2028	3.50	50	Periset
Tab dispersible 8 mg – 5% DV Mar-24 to 2026	0.90	10	Periset ODT
Inj 2 mg per ml, 2 ml ampoule – 5% DV Feb-26 to 2028	1.54	5	Ondansetron-AFT
Inj 2 mg per ml, 4 ml ampoule – 5% DV Feb-26 to 2028	2.14	5	Ondansetron-AFT
PROCHLORPERAZINE			
Tab buccal 3 mg			
Tab 5 mg – 5% DV Mar-24 to 2026	25.00	250	Nausafix
Inj 12.5 mg per ml, 1 ml ampoule			
Suppos 25 mg			
TROPISETRON			
Inj 1 mg per ml, 2 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Antipsychotic Agents			
General			
AMISULPRIDE			
Tab 100 mg – 5% DV Dec-24 to 2027	5.84	30	Sulprix
Tab 200 mg – 5% DV Dec-24 to 2027	14.47	60	Sulprix
Tab 400 mg – 5% DV Dec-24 to 2027	35.06	60	Sulprix
Oral liq 100 mg per ml			
ARIPIPAZOLE			
Tab 5 mg	10.50	30	Aripiprazole Sandoz
Tab 10 mg	10.50	30	Aripiprazole Sandoz
Tab 15 mg	10.50	30	Aripiprazole Sandoz
Tab 20 mg	10.50	30	Aripiprazole Sandoz
Tab 30 mg	10.50	30	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE			
Tab 25 mg	15.62	100	Largactil
Tab 100 mg	36.73	100	Largactil
Oral liq 10 mg per ml			
Oral liq 20 mg per ml			
Inj 25 mg per ml, 2 ml ampoule	30.79	10	Largactil
CLOZAPINE			
Tab 25 mg	6.69	50	Clopine
	13.37	100	Clopine
	6.69	50	Clozaril
	13.37	100	Clozaril
Tab 50 mg	8.67	50	Clopine
	17.33	100	Clopine
Tab 100 mg	17.33	50	Clopine
	34.65	100	Clopine
	17.33	50	Clozaril
	34.65	100	Clozaril
Tab 200 mg	34.65	50	Clopine
	69.30	100	Clopine
Oral liq 50 mg per ml	173.30	100 ml	Versacloz
HALOPERIDOL			
Tab 500 mcg	6.23	100	Serenace
Tab 1.5 mg	9.43	100	Serenace
Tab 5 mg	29.72	100	Serenace
Oral liq 2 mg per ml	23.84	100 ml	Serenace
Inj 5 mg per ml, 1ml ampoule	21.55	10	Serenace
LEVOMEPROMAZINE			
Tab 25 mg	16.10	100	Nozinan
Tab 100 mg	41.75	100	Nozinan
LEVOMEPROMAZINE HYDROCHLORIDE			
Inj 25 mg per ml, 1 ml ampoule – 5% DV Dec-25 to 2028	23.26	10	Wockhardt
LITHIUM CARBONATE			
Tab long-acting 400 mg – 5% DV Feb-25 to 2027	82.80	100	Priadel
Cap 250 mg	35.78	100	Douglas

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
OLANZAPINE			
Tab 2.5 mg – 5% DV Aug-24 to 2026.....	1.40	30	Zypine
Tab 5 mg – 5% DV Aug-24 to 2026.....	1.93	30	Zypine
Tab orodispersible 5 mg – 5% DV Feb-24 to 2026.....	2.42	28	Zypine ODT
Tab 10 mg – 5% DV Aug-24 to 2026.....	1.93	30	Zypine
Tab orodispersible 10 mg – 5% DV Feb-24 to 2026.....	2.89	28	Zypine ODT
Inj 10 mg vial			
PERICYAZINE			
Tab 2.5 mg			
Tab 10 mg			
QUETIAPINE			
Tab 25 mg – 5% DV Feb-24 to 2026.....	2.36	90	Quetapel
	0.79	30	Quetiapine Viatris
	13.11	500	Quetiapine Viatris
Tab 100 mg – 5% DV Feb-24 to 2026.....	6.40	90	Quetapel
Tab 200 mg – 5% DV Feb-24 to 2026.....	10.97	90	Quetapel
Tab 300 mg – 5% DV Feb-24 to 2026.....	15.83	90	Quetapel
RISPERIDONE			
Tab 0.5 mg – 5% DV Mar-24 to 2026.....	2.17	60	Risperidone (Teva)
Tab 1 mg – 5% DV Mar-24 to 2026.....	2.44	60	Risperidone (Teva)
Tab 2 mg – 5% DV Mar-24 to 2026.....	2.72	60	Risperidone (Teva)
Tab 3 mg – 5% DV Mar-24 to 2026.....	4.50	60	Risperidone (Teva)
Tab 4 mg – 5% DV Mar-24 to 2026.....	6.25	60	Risperdal
			Risperidone (Teva)
Oral liq 1 mg per ml – 5% DV Mar-24 to 2026.....	10.29	30 ml	Risperon
ZIPRASIDONE			
Cap 20 mg.....	17.90	60	Zusdone
Cap 40 mg.....	27.41	60	Zusdone
Cap 60 mg.....	38.39	60	Zusdone
Cap 80 mg.....	46.55	60	Zusdone
ZUCLOPENTHIXOL ACETATE			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
ZUCLOPENTHIXOL HYDROCHLORIDE			
Tab 10 mg.....	31.45	100	Clopixol

Depot Injections

ARIPIPRAZOLE – Restricted see terms [below](#)

↓ Inj 300 mg vial	273.56	1	Abilify Maintena
↓ Inj 400 mg vial	341.96	1	Abilify Maintena

➔ **Restricted (RS2058)**

Initiation

Either:

1 Either:

- 1.1 The patient has had an initial Special Authority approval for risperidone depot injection or paliperidone depot injection or olanzapine depot injection; or
- 1.2 All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.2.1 The patient has schizophrenia or other psychotic disorder; and
 - 1.2.2 The patient has received treatment with oral atypical antipsychotic agents but has been unable to adhere; and
 - 1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or
- 2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been initiated on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection. (see Note below for the olanzapine Special Authority criteria for new olanzapine depot injection patients prior to 1 April 2024).

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
 - The patient has schizophrenia; and
 - The patient has tried but has not been able to adhere with treatment using oral atypical antipsychotic agents; and
 - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

FLUPENTHIXOL DECANOATE

Inj 20 mg per ml, 1 ml ampoule	13.14	5	Fluanxol
Inj 20 mg per ml, 2 ml ampoule	20.90	5	Fluanxol
Inj 100 mg per ml, 1 ml ampoule	40.87	5	Fluanxol

HALOPERIDOL DECANOATE

Inj 50 mg per ml, 1 ml ampoule	28.39	5	Haldol
Inj 100 mg per ml, 1 ml ampoule	55.90	5	Haldol Concentrate

OLANZAPINE – **Restricted:** For continuation only

➡ Inj 210 mg vial	252.00	1	Zyprexa Relprevv
➡ Inj 300 mg vial	414.00	1	Zyprexa Relprevv
➡ Inj 405 mg vial	504.00	1	Zyprexa Relprevv

➡ **Restricted (RS2018)**

Continuation

Re-assessment required after 12 months

The initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE – **Restricted** see terms [below](#)

⚡ Inj 25 mg syringe	194.25	1	Invega Sustenna
⚡ Inj 50 mg syringe	271.95	1	Invega Sustenna
⚡ Inj 75 mg syringe	357.42	1	Invega Sustenna
⚡ Inj 100 mg syringe	435.12	1	Invega Sustenna
⚡ Inj 150 mg syringe	435.12	1	Invega Sustenna

➡ **Restricted (RS2059)**

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

- 2.1 The patient has schizophrenia or other psychotic disorder; and
- 2.2 The patient has been unable to adhere to treatment using oral atypical antipsychotic agents; and
- 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE – Restricted see terms [below](#)

↓ Inj 175 mg syringe	815.85	1	Invega Trinza
↓ Inj 263 mg syringe	1,072.26	1	Invega Trinza
↓ Inj 350 mg syringe	1,305.36	1	Invega Trinza
↓ Inj 525 mg syringe	1,305.36	1	Invega Trinza

→ **Restricted (RS1932)**

Initiation

Re-assessment required after 12 months

Both:

- 1 The patient has schizophrenia; and
- 2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Continuation

Re-assessment required after 12 months

The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PIPOTHIAZINE PALMITATE – Restricted: For continuation only

- Inj 50 mg per ml, 1 ml ampoule
- Inj 50 mg per ml, 2 ml ampoule

RISPERIDONE – Restricted see terms [below](#)

↓ Inj 25 mg vial	135.98	1	Risperdal Consta
↓ Inj 37.5 mg vial	178.71	1	Risperdal Consta
↓ Inj 50 mg vial	217.56	1	Risperdal Consta

→ **Restricted (RS2060)**

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 The patient has not been able to adhere to treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE

Inj 200 mg per ml, 1 ml ampoule	19.80	5	Clopixol
Inj 500 mg per ml, 1 ml ampoule			e.g. Clopixol Conc

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anxiolytics			
BUSPIRONE HYDROCHLORIDE			
Tab 5 mg – 5% DV Dec-24 to 2027	13.95	100	Buspirone Viatris
Tab 10 mg – 5% DV Dec-24 to 2027	12.50	100	Buspirone Viatris
CLONAZEPAM			
Tab 500 mcg	5.64	100	Paxam
Tab 2 mg	10.78	100	Paxam
DIAZEPAM			
Tab 2 mg – 5% DV Mar-24 to 2026	95.00	500	Arrow-Diazepam
Tab 5 mg – 5% DV Mar-24 to 2026	115.00	500	Arrow-Diazepam
↓ Oral liq 10 mg per 10 ml			
➔ Restricted (RS2054)			
Initiation			
Relevant specialist			
Only for use in children where diazepam tablets are not appropriate.			
LORAZEPAM			
Tab 1 mg – 5% DV Feb-25 to 2027	10.20	250	Ativan
Tab 2.5 mg – 5% DV Feb-25 to 2027	13.13	100	Ativan
OXAZEPAM			
Tab 10 mg			
Tab 15 mg			

Multiple Sclerosis Treatments

➔ **Restricted (RS1993)**

Initiation – Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide

Any relevant practitioner

Re-assessment required after 12 months

Either:

1 All of the following:

- 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 1.2 Patient has an EDSS score between 0 – 6.0; and
- 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
fever ($T > 37.5^{\circ}\text{C}$); and			
1.4.5 Either:			
1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or			
1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and			
1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and			
1.6 Any of the following:			
1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or			
1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or			
1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or			
1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or			
1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or			
2 Patient has an active approval for ocrelizumab and does not have primary progressive MS.			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
Continuation – Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide			
Any relevant practitioner			
Patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
DIMETHYL FUMARATE – Restricted see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
† Cap 120 mg	520.00	14	Tecfidera
† Cap 240 mg	2,000.00	56	Tecfidera
FINGOLIMOD – Restricted see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
† Cap 0.5 mg	2,200.00	28	Gilenya
GLATIRAMER ACETATE – Restricted see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
† Inj 40 mg prefilled syringe	1,137.48	12	Copaxone
INTERFERON BETA-1-ALPHA – Restricted see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
† Inj 6 million iu in 0.5 ml syringe	1,170.00	4	Avonex
INTERFERON BETA-1-BETA – Restricted see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
† Inj 8 million iu per ml, 1 ml vial			
NATALIZUMAB – Restricted see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
† Inj 20 mg per ml, 15 ml vial	1,750.00	1	Tysabri
TERIFLUNOMIDE – Restricted see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
† Tab 14 mg – 5% DV Apr-25 to 2026	263.96	28	Teriflunomide Sandoz

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

Multiple Sclerosis Treatments - Other

OCRELIZUMAB – **Restricted** see terms [below](#)

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

⚡ Inj 30 mg per ml, 10 ml vial.....9,346.00 1 Ocrevus

➡ **Restricted (RS1997)**

Initiation – Multiple Sclerosis - ocrelizumab

Any relevant practitioner

Re-assessment required after 12 months

Either:

- 1 All of the following:
 - 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
 - 1.2 Patient has an EDSS score between 0 – 6.0; and
 - 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
 - 1.4 All of the following:
 - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever ($T > 37.5^{\circ}\text{C}$); and
 - 1.4.5 Either:
 - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtzle Functional System scores by at least 1 point; or
 - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
 - 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
 - 1.6 Any of the following:
 - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or
- 2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Continuation – Multiple Sclerosis - ocrelizumab

Any relevant practitioner

Patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Initiation – Primary Progressive Multiple Sclerosis

Any relevant practitioner

Re-assessment required after 12 months

All of the following:

- 1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and
- 2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and
- 3 Patient has no history of relapsing remitting multiple sclerosis.

Continuation – Primary Progressive Multiple Sclerosis

Any relevant practitioner

Patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

Sedatives and Hypnotics

CHLORAL HYDRATE

Oral liq 100 mg per ml

Oral liq 200 mg per ml

LORMETAZEPAM – **Restricted:** For continuation only

➔ Tab 1 mg

MELATONIN – **Restricted** see terms [below](#)

↓ Tab modified-release 2 mg – **5% DV Dec-24 to 2027** 5.80 30 **Vigisom**

↓ Tab 3 mg

Note: Only for use in compounding an oral liquid formulation, for in-hospital use only.

➔ **Restricted (RS1576)**

Initiation – insomnia secondary to neurodevelopmental disorder

Psychiatrist, paediatrician, neurologist or respiratory specialist

Re-assessment required after 12 months

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder); and
- 2 Behavioural and environmental approaches have been tried or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under.

Continuation – insomnia secondary to neurodevelopmental disorder

Psychiatrist, paediatrician, neurologist or respiratory specialist

Re-assessment required after 12 months

All of the following:

- 1 Patient is aged 18 years or under; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Initiation – insomnia where benzodiazepines and zopiclone are contraindicated

Both:

- 1 Patient has insomnia and benzodiazepines and zopiclone are contraindicated; and
- 2 For in-hospital use only.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MIDAZOLAM			
Tab 7.5 mg			
Oral liq 2 mg per ml			
Inj 5 mg per ml, 1 ml plastic ampoule	22.50	10	Midazolam-Pfizer
Inj 1 mg per ml, 5 ml ampoule – 5% DV May-25 to 2027	7.80	10	Midazolam-Baxter
Inj 5 mg per ml, 3 ml ampoule – 5% DV May-25 to 2027	4.75	5	Midazolam-Baxter
PHENOBARBITONE			
Inj 130 mg per ml, 1 ml vial			
Inj 200 mg per ml, 1 ml ampoule			
TEMAZEPAM			
Tab 10 mg – 5% DV Feb-24 to 2026	1.40	25	Normison
TRIAZOLAM – Restricted: For continuation only			
➡ Tab 125 mcg			
➡ Tab 250 mcg			
ZOPICLONE			
Tab 7.5 mg – 5% DV Feb-25 to 2027	21.85	500	Zopiclone Actavis

Spinal Muscular Atrophy

NUSINERSEN – Restricted see terms [below](#)

⚡ Inj 12 mg per 5 ml vial 120,000.00 1 Spinraza

➡ **Restricted (RS1938)**

Initiation

Re-assessment required after 12 months

All of the following:

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Continuation

Re-assessment required after 12 months

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day), in the absence of a potentially reversible cause while being treated with nusinersen; and
- 3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

RISDIPLAM – Restricted see terms [below](#)

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website <https://pharmac.govt.nz/risdiplam>

⚡ Powder for oral soln 750 mcg per ml, 60 mg per bottle..... 14,100.00 80 ml Evrysdi

➡ **Restricted (RS1954)**

Initiation

Re-assessment required after 12 months

All of the following:

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

- 1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
- 2 Patient is 18 years of age or under; and
- 3 Either:
 - 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
 - 3.2 Both:
 - 3.2.1 Patient is pre-symptomatic; and
 - 3.2.2 Patient has three or less copies of SMN2.

Continuation

Re-assessment required after 12 months

All of the following:

- 1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
- 2 Patient does not require invasive permanent ventilation (at least 16 hours per day), in the absence of a potentially reversible cause while being treated with risdiplam; and
- 3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

Stimulants / ADHD Treatments

ATOMOXETINE

Cap 10 mg – 5% DV Aug-24 to 2026	43.02	28	APO-Atomoxetine
Cap 18 mg – 5% DV Aug-24 to 2026	45.57	28	APO-Atomoxetine
Cap 25 mg – 5% DV Aug-24 to 2026	44.30	28	APO-Atomoxetine
Cap 40 mg – 5% DV Aug-24 to 2026	46.21	28	APO-Atomoxetine
Cap 60 mg – 5% DV Aug-24 to 2026	51.31	28	APO-Atomoxetine
Cap 80 mg – 5% DV Aug-24 to 2026	65.20	28	APO-Atomoxetine
Cap 100 mg – 5% DV Aug-24 to 2026	65.71	28	APO-Atomoxetine

CAFFEINE

Tab 100 mg

DEXAMFETAMINE SULFATE – **Restricted** see terms [below](#)

↓ Tab 5 mg29.80 100 Noumed Dexamfetamine

→ **Restricted (RS2071)**

Initiation – ADHD

Paediatrician or psychiatrist

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.

Initiation – Narcolepsy

Neurologist or respiratory specialist

Patient suffers from narcolepsy.

LISDEXAMFETAMINE DIMESILATE – **Restricted** see terms [below](#)

↓ Cap 30 mg	60.00	30	Vyvanse
↓ Cap 50 mg	60.00	30	Vyvanse
↓ Cap 70 mg	60.00	30	Vyvanse

→ **Restricted (RS2070)**

Initiation

Paediatrician or psychiatrist

Either:

- 1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all remaining criteria prior to commencing treatment; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 All of the following:

2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and

2.2 Diagnosed according to DSM-V or ICD 11 criteria; and

2.3 Any of the following:

- 2.3.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) and has not received sufficient benefit or has experienced intolerable side effects; or
- 2.3.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
- 2.3.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate; or
- 2.3.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
- 2.3.5 There is significant concern regarding the risk of diversion or abuse of immediate release methylphenidate hydrochloride; or
- 2.3.6 Both:
 - 2.3.6.1 Patient would have been prescribed a subsidised formulation of methylphenidate hydrochloride (extended-release) but has been unable to access due to supply issues with methylphenidate hydrochloride (extended-release); and
 - 2.3.6.2 Other alternative stimulant presentations (methylphenidate or dexamfetamine) are not appropriate; and

2.4 Lisdexamfetamine dimesilate is not to be used in combination with another funded methylphenidate presentation.

METHYLPHENIDATE HYDROCHLORIDE – Restricted see terms [below](#)

↓ Tab extended-release 18 mg.....	58.96	30	Concerta
	7.75		Methylphenidate ER - Teva
↓ Tab extended-release 27 mg.....	65.44	30	Concerta
	11.45		Methylphenidate ER - Teva
↓ Tab extended-release 36 mg.....	71.93	30	Concerta
	15.50		Methylphenidate ER - Teva
↓ Tab extended-release 54 mg.....	86.24	30	Concerta
	22.25		Methylphenidate ER - Teva
↓ Tab immediate-release 5 mg.....	3.20	30	Rubifen
↓ Tab immediate-release 10 mg.....	4.00	30	Ritalin
	3.00		Rubifen
↓ Tab immediate-release 20 mg.....	7.85	30	Rubifen
↓ Tab sustained-release 20 mg.....	10.95	30	Rubifen SR
↓ Cap modified-release 10 mg.....	19.41	30	Ritalin LA
↓ Cap modified-release 20 mg.....	27.72	30	Ritalin LA
↓ Cap modified-release 30 mg.....	34.39	30	Ritalin LA
↓ Cap modified-release 40 mg.....	38.67	30	Ritalin LA

→ **Restricted (RS2105)**

Initiation – ADHD (immediate-release and sustained-release formulations)

Paediatrician or psychiatrist

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.

continued...

	Price (ex man. excl. GST)		Brand or Generic Manufacturer
	\$	Per	

continued...

Initiation – Narcolepsy (immediate-release and sustained-release formulations)

Neurologist or respiratory specialist

Patient suffers from narcolepsy.

Initiation – Extended-release and modified-release formulations

Paediatrician or psychiatrist

Both:

- 1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria; and
- 2 Either:
 - 2.1 Patient is taking a currently listed formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 2.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Initiation – Narcolepsy* (extended-release only)

Neurologist or respiratory specialist

Patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Concerta or Ritalin LA.

MODAFINIL – **Restricted** see terms [below](#)

↓ Tab 100 mg – 5% DV May-25 to 2027 14.27 30 **Modafinil Max Health**

→ **Restricted (RS2106)**

Initiation – Narcolepsy

Neurologist or respiratory specialist

Either:

- 1 All of the following:
 - 1.1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
 - 1.2 Either:
 - 1.2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 1.2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
 - 1.3 Either:
 - 1.3.1 An effective dose of a listed formulation of methylphenidate or dexamphetamine has been trialed and discontinued because of intolerable side effects; or
 - 1.3.2 Methylphenidate and dexamphetamine are contraindicated; or
- 2 Both:
 - 2.1 Patient meets the Hospital Restriction criteria for methylphenidate hydrochloride for narcolepsy; and
 - 2.2 Patient is unable to access methylphenidate hydrochloride presentations due to an out of stock (see note).

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock of methylphenidate hydrochloride.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE

Tab 5 mg – 5% DV Jun-24 to 2026	3.70	84	Ipca-Donepezil
Tab 10 mg – 5% DV Jun-24 to 2026	5.50	84	Ipca-Donepezil

RIVASTIGMINE – **Restricted** see terms [on the next page](#)

↓ Patch 4.6 mg per 24 hour – 5% DV Mar-25 to 2027	49.40	30	Rivastigmine Patch BNM 5
↓ Patch 9.5 mg per 24 hour – 5% DV Mar-25 to 2027	49.40	30	Rivastigmine Patch BNM 10

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2139)

Initiation

Re-assessment required after 6 months

Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient is contraindicated to or has experienced intolerable side effects from donepezil tablets.

Continuation

Re-assessment required after 12 months

Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE – **Restricted** see terms [below](#)

⚡ Tab 2 mg with naloxone 0.5 mg	11.76	28	Buprenorphine Naloxone BNM
⚡ Tab 8 mg with naloxone 2 mg	34.00	28	Buprenorphine Naloxone BNM

➔ Restricted (RS1172)

Initiation – Detoxification

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Prescriber works in an opioid treatment service approved by the Ministry of Health.

Initiation – Maintenance treatment

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Prescriber works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg – 5% DV May-24 to 2026	15.00	30	Zyban
---	-------	----	--------------

DISULFIRAM

Tab 200 mg	236.40	100	Antabuse
------------------	--------	-----	----------

NALTREXONE HYDROCHLORIDE – **Restricted** see terms [below](#)

⚡ Tab 50 mg – 5% DV Dec-23 to 2026	83.33	30	Naltraccord
---	-------	----	--------------------

➔ Restricted (RS1173)

Initiation – Alcohol dependence

Both:

- 1 Patient is currently enrolled, or is planned to be enrolled, in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Naltrexone is to be prescribed by, or on the recommendation of, a physician working in an Alcohol and Drug Service.

Initiation – Constipation

For the treatment of opioid-induced constipation.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
NICOTINE – Some items restricted see terms below			
Patch 7 mg per 24 hours	19.62	28	Habitrol
Patch 14 mg per 24 hours	21.57	28	Habitrol
Patch 21 mg per 24 hours	24.72	28	Habitrol
↓ Oral spray 1 mg per dose			<i>e.g. Nicorette QuickMist Mouth Spray</i>
Lozenge 1 mg	22.53	216	Habitrol
Lozenge 2 mg	24.68	216	Habitrol
↓ Soln for inhalation 15 mg cartridge			
Gum 2 mg	23.02	204	Habitrol (Fruit) Habitrol (Mint)
Gum 4 mg	25.98	204	Habitrol (Fruit) Habitrol (Mint)

➔ **Restricted (RS1873)**

Initiation

Any of the following:

- 1 For perioperative use in patients who have a 'nil by mouth' instruction; or
- 2 For use within mental health inpatient units; or
- 3 Patient would be admitted to a mental health inpatient unit, but is unable to due to COVID-19 self-isolation requirement; or
- 4 For acute use in agitated patients who are unable to leave the hospital facilities.

VARENICLINE – Restricted see terms [below](#)

↓ Tab 0.5 mg x 11 and 1 mg x 42	16.67	53	Champix
↓ Tab 1 mg	17.62	56	Champix

➔ **Restricted (RS1702)**

Initiation

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline in a 12 month period.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE – **Restricted** see terms [below](#)

⚡ Inj 25 mg vial – 5% DV Apr-25 to 2027	50.05	1	Bendamustine Sandoz
⚡ Inj 100 mg vial – 5% DV Apr-25 to 2027	200.20	1	Bendamustine Sandoz
➡ Restricted (RS2061)			

Initiation – CLL*

All of the following:

- 1 The patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has ECOG performance status 0-2; and
- 3 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a * includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initiation – Indolent, Low-grade lymphomas

Re-assessment required after 9 months

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Continuation – Indolent, Low-grade lymphomas

Re-assessment required after 9 months

Either:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or

2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenström's macroglobulinaemia.

Initiation – Hodgkin's lymphoma*

Relevant specialist or medical practitioner on the recommendation of a relevant specialist

Limited to 6 months treatment

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m² twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN

Tab 2 mg	89.25	100	Myleran
Inj 6 mg per ml, 10 ml ampoule			

CARMUSTINE

Inj 100 mg vial	710.00	1	BiCNU BiCNU S29 Novadoz
-----------------------	--------	---	-------------------------------

CHLORAMBUCIL

Tab 2 mg

CYCLOPHOSPHAMIDE

Tab 50 mg – 5% DV Dec-24 to 2027	145.00	50	Cyclonex
Inj 1 g vial – 5% DV Feb-25 to 2027	47.46	1	Endoxan
Inj 2 g vial – 5% DV Feb-25 to 2027	95.06	1	Endoxan

IFOSFAMIDE

Inj 1 g vial	96.00	1	Holoxan
Inj 2 g vial	180.00	1	Holoxan

LOMUSTINE

Cap 40 mg	880.00	20	Medac
-----------------	--------	----	-------

MELPHALAN

Tab 2 mg			
Inj 50 mg vial – 5% DV Dec-23 to 2026	48.25	1	Melpha

THIOTEPA

Inj 15 mg vial – 5% DV Apr-24 to 2026	398.00	1	Tepadina
Inj 100 mg vial – 5% DV Apr-24 to 2026	1,800.00	1	Tepadina

Anthracyclines and Other Cytotoxic Antibiotics

BLEOMYCIN SULPHATE

Inj 15,000 iu vial.....	185.16	1	DBL Bleomycin Sulfate
-------------------------	--------	---	-----------------------

DACTINOMYCIN [ACTINOMYCIN D]

Inj 0.5 mg vial	255.00	1	Cosmegen
-----------------------	--------	---	----------

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DAUNORUBICIN			
Inj 18.7 mg vial	171.93	1	Pfizer
Inj 2 mg per ml, 10 ml vial.....	171.93	1	Pfizer
<i>(Pfizer Inj 2 mg per ml, 10 ml vial to be delisted 1 January 2026)</i>			
DOXORUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 5 ml vial.....	11.50	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial.....	23.00	1	Doxorubicin Ebewe
Inj 50 mg vial.....	69.99	1	Doxorubicin Ebewe
Inj 2 mg per ml, 50 ml vial.....	23.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial.....	69.99	1	Doxorubicin Ebewe
EPIRUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 5 ml vial.....	25.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial.....	30.00	1	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial.....	99.99	1	Epirubicin Ebewe
IDARUBICIN HYDROCHLORIDE			
Inj 5 mg vial	109.74	1	Zavedos
Inj 10 mg vial	233.64	1	Zavedos
MITOMYCIN C			
Inj 5 mg vial			
Inj 20 mg vial	1,250.00	1	Teva
MITOZANTRONE			
Inj 2 mg per ml, 10 ml vial.....	97.50	1	Mitozantrone Ebewe
Antimetabolites			
AZACITIDINE – Restricted see terms below			
⚡ Inj 100 mg vial – 5% DV Mar-25 to 2027	50.00	1	Azacitidine Dr Reddy's
➡ Restricted (RS2116)			
Initiation			
<i>Re-assessment required after 12 months</i>			
Both:			
1 Any of the following:			
1.1 The individual has intermediate or high risk MDS based on an internationally recognised scoring system; or			
1.2 The individual has chronic myelomonocytic leukaemia (based on an intermediate or high risk score from an internationally recognised scoring system or 10%-29% marrow blasts without myeloproliferative disorder); or			
1.3 The individual has acute myeloid leukaemia according to World Health Organisation (WHO) Classification; and			
2 The individual has an estimated life expectancy of at least 3 months.			
Continuation			
<i>Re-assessment required after 12 months</i>			
No evidence of disease progression.			
CAPECITABINE			
Tab 150 mg – 5% DV Feb-26 to 2028	10.92	60	Capecitabine Viatris
Tab 500 mg – 5% DV Feb-26 to 2028	50.96	120	Capecitabine Viatris
CLADRIBINE			
Inj 2 mg per ml, 5 ml vial.....			
Inj 1 mg per ml, 10 ml vial.....	749.96	1	Leustatin
CYTARABINE			
Inj 20 mg per ml, 5 ml vial.....	472.00	5	Pfizer
Inj 100 mg per ml, 20 ml vial.....	48.80	1	Cytarabine DBL Pfizer

⚡ Item restricted (see ➡ above); ⚡ Item restricted (see ➡ below)
e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUDARABINE PHOSPHATE			
Tab 10 mg	412.00	20	Fludara Oral
Inj 50 mg vial	634.00	5	Fludarabine Ebewe
	126.80	1	Fludarabine Sagent
<i>(Fludarabine Sagent Inj 50 mg vial to be delisted 1 November 2025)</i>			
FLUOROURACIL			
Inj 50 mg per ml, 20 ml vial – 5% DV Dec-24 to 2027	10.51	1	Fluorouracil Accord
Inj 50 mg per ml, 50 ml vial	14.72	1	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial – 5% DV Dec-24 to 2027	19.36	1	Fluorouracil Accord
GEMCITABINE HYDROCHLORIDE			
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine), 26.3 ml vial – 5% DV Jun-24 to 2026	18.94	1	DBL Gemcitabine
MERCAPTOPURINE			
Tab 50 mg – 5% DV Dec-25 to 2028	19.50	25	Puri-nethol
↓ Oral suspension 20 mg per ml	428.00	100 ml	Xaluprine Allmercap
→ Restricted (RS1635)			
Initiation			
Paediatric haematologist or paediatric oncologist			
<i>Re-assessment required after 12 months</i>			
The patient requires a total dose of less than one full 50 mg tablet per day.			
Continuation			
Paediatric haematologist or paediatric oncologist			
<i>Re-assessment required after 12 months</i>			
The patient requires a total dose of less than one full 50 mg tablet per day.			
METHOTREXATE			
Tab 2.5 mg – 5% DV Dec-24 to 2027	7.80	90	Trexate
Tab 10 mg – 5% DV Dec-24 to 2027	26.40	90	Trexate
Inj 2.5 mg per ml, 2 ml vial			
Inj 7.5 mg prefilled syringe – 5% DV Feb-25 to 2027	29.17	1	Methotrexate Sandoz
Inj 10 mg prefilled syringe – 5% DV Feb-25 to 2027	19.09	1	Methotrexate Sandoz
Inj 15 mg prefilled syringe – 5% DV Feb-25 to 2027	24.53	1	Methotrexate Sandoz
Inj 20 mg prefilled syringe – 5% DV Feb-25 to 2027	16.64	1	Methotrexate Sandoz
Inj 25 mg prefilled syringe – 5% DV Feb-25 to 2027	20.72	1	Methotrexate Sandoz
Inj 30 mg prefilled syringe – 5% DV Feb-25 to 2027	55.00	1	Methotrexate Sandoz
Inj 25 mg per ml, 2 ml vial	30.00	5	Methotrexate DBL
			Onco-Vial
Inj 25 mg per ml, 20 ml vial	45.00	1	DBL Methotrexate
			Onco-Vial
Inj 100 mg per ml, 10 ml vial	25.00	1	Methotrexate Ebewe
Inj 100 mg per ml, 50 ml vial – 5% DV Dec-23 to 2026	67.99	1	Methotrexate Ebewe
PEMETREXED			
Inj 100 mg vial – 5% DV Apr-25 to 2027	8.99	1	Pemetrexed-AFT
Inj 500 mg vial – 5% DV Apr-25 to 2027	29.99	1	Pemetrexed-AFT
THIOGUANINE			
Tab 40 mg			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Other Cytotoxic Agents			
AMSACRINE			
Inj 50 mg per ml, 1.5 ml ampoule			
Inj 75 mg			
ANAGRELIDE HYDROCHLORIDE			
Cap 0.5 mg			
ARSENIC TRIOXIDE			
Inj 1 mg per ml, 10 ml vial.....	4,817.00	10	Phenasen
BORTEZOMIB – Restricted see terms below			
⚡ Inj 3.5 mg vial	74.93	1	DBL Bortezomib
➡ Restricted (RS2043)			
Initiation – plasma cell dyscrasia			
The patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.			
DACARBAZINE			
Inj 200 mg vial	72.11	1	DBL Dacarbazine
ETOPOSIDE			
Cap 50 mg	340.73	20	Vepesid
Cap 100 mg	340.73	10	Vepesid
Inj 20 mg per ml, 5 ml vial	7.90	1	Rex Medical
ETOPOSIDE (AS PHOSPHATE)			
Inj 100 mg vial	40.00	1	Etopophos
HYDROXYUREA [HYDROXYCARBAMIDE]			
Cap 500 mg – 5% DV Dec-23 to 2026.....	20.72	100	Devatis
IBRUTINIB – Restricted see terms below			
⚡ Tab 140 mg	3,217.00	30	Imbruvica
⚡ Tab 420 mg	9,652.00	30	Imbruvica
➡ Restricted (RS2117)			
Initiation – chronic lymphocytic leukaemia (CLL)			
<i>Re-assessment required after 6 months</i>			
All of the following:			
1 Individual has chronic lymphocytic leukaemia (CLL) requiring therapy; and			
2 Individual has not previously received funded ibrutinib; and			
3 Ibrutinib is to be used as monotherapy; and			
4 Any of the following:			
4.1 Both:			
4.1.1 There is documentation confirming that the individual has 17p deletion or TP53 mutation; and			
4.1.2 Individual has experienced intolerable side effects with venetoclax monotherapy; or			
4.2 All of the following:			
4.2.1 Individual has received at least one prior immunochemotherapy for CLL; and			
4.2.2 Individual's CLL has relapsed; and			
4.2.3 Individual has experienced intolerable side effects with venetoclax in combination with rituximab regimen; or			
4.3 Individual's CLL is refractory to or has relapsed following a venetoclax regimen.			
Continuation – chronic lymphocytic leukaemia (CLL)			
<i>Re-assessment required after 12 months</i>			
No evidence of clinical disease progression.			
Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
IRINOTECAN HYDROCHLORIDE			
Inj 20 mg per ml, 5 ml vial.....	52.57	1	Accord
Inj 20 mg per ml, 25 ml vial.....	262.85	1	Accord
LENALIDOMIDE (VIATRIS) – Restricted see terms below			
↓ Cap 5 mg – 5% DV Feb-25 to 31 Jan 2028.....	76.92	21	Lenalidomide Viatris
↓ Cap 10 mg – 5% DV Feb-25 to 31 Jan 2028.....	50.30	21	Lenalidomide Viatris
↓ Cap 15 mg – 5% DV Feb-25 to 31 Jan 2028.....	62.13	21	Lenalidomide Viatris
↓ Cap 25 mg – 5% DV Feb-25 to 31 Jan 2028.....	65.09	21	Lenalidomide Viatris

→ **Restricted (RS2044)**

Initiation – Plasma cell dyscrasia

Any relevant practitioner

Both:

- 1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
- 2 Patient is not refractory to prior lenalidomide use.

Initiation – Myelodysplastic syndrome

Any relevant practitioner

Re-assessment required after 6 months

Both:

- 1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5) associated with a deletion 5q cytogenetic abnormality; and
- 2 Patient has transfusion-dependent anaemia.

Continuation – Myelodysplastic syndrome

Any relevant practitioner

Re-assessment required after 12 months

Both:

- 1 Patient has not needed a transfusion in the last 4 months; and
- 2 No evidence of disease progression.

NIRAPARIB – Restricted see terms below

↓ Tab 100 mg	13,393.50	84	Zejula
↓ Cap 100 mg	8,929.84	56	Zejula

→ **Restricted (RS2027)**

Initiation

Re-assessment required after 6 months

All of the following:

- 1 Patient has advanced high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 Patient has received at least one line** of treatment with platinum-based chemotherapy; and
- 3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and
- 4 Patient has not previously received funded treatment with a PARP inhibitor; and
- 5 Either:
 - 5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen; or
 - 5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and
- 6 Treatment to be administered as maintenance treatment; and
- 7 Treatment not to be administered in combination with other chemotherapy.

Continuation

Re-assessment required after 6 months

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 No evidence of progressive disease; and
- 2 Treatment to be administered as maintenance treatment; and
- 3 Treatment not to be administered in combination with other chemotherapy; and
- 4 Either:
 - 4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
 - 4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: * "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB – **Restricted** see terms [below](#)

⚡ Tab 100 mg	3,701.00	56	Lynparza
⚡ Tab 150 mg	3,701.00	56	Lynparza

➡ **Restricted (RS1925)**

Initiation – Ovarian cancer

Medical oncologist

Re-assessment required after 12 months

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Either:
 - 3.1 All of the following:
 - 3.1.1 Patient has newly diagnosed, advanced disease; and
 - 3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 3.2 All of the following:
 - 3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
 - 3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
 - 3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
 - 3.2.4 Patient has not previously received funded olaparib treatment; and
- 4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 5 Treatment to be administered as maintenance treatment; and
- 6 Treatment not to be administered in combination with other chemotherapy.

Continuation – Ovarian cancer

Medical oncologist

Re-assessment required after 12 months

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

5 Either:

5.1 Both:

5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and

5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or

5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PEGASPARGASE – **Restricted** see terms [below](#)

⚡ Inj 750 iu per ml, 5 ml vial.....3,973.25 1 Oncaspar LYO

➔ **Restricted (RS1788)**

Initiation – Newly diagnosed ALL

Limited to 12 months treatment

Both:

1 The patient has newly diagnosed acute lymphoblastic leukaemia; and

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initiation – Relapsed ALL

Limited to 12 months treatment

Both:

1 The patient has relapsed acute lymphoblastic leukaemia; and

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initiation – Lymphoma

Limited to 12 months treatment

Patient has lymphoma requiring L-asparaginase containing protocol (e.g. SMILE).

PENTOSTATIN [DEOXYCOFORMYCIN]

Inj 10 mg vial

POMALIDOMIDE – **Restricted** see terms [below](#)

⚡ Cap 1 mg – 5% DV Aug-24 to 31 Jul 202747.45 14 **Pomolide**

71.18 21 **Pomolide**

⚡ Cap 2 mg – 5% DV Aug-24 to 31 Jul 202794.90 14 **Pomolide**

142.35 21 **Pomolide**

⚡ Cap 3 mg – 5% DV Aug-24 to 31 Jul 2027142.35 14 **Pomolide**

213.53 21 **Pomolide**

⚡ Cap 4 mg – 5% DV Aug-24 to 31 Jul 2027189.81 14 **Pomolide**

284.71 21 **Pomolide**

➔ **Restricted (RS2045)**

Initiation – Relapsed/refractory plasma cell dyscrasia

Any relevant practitioner

Re-assessment required after 6 months

Both:

1 Patient has relapsed or refractory plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and

2 Patient has not received prior funded pomalidomide.

Continuation – Relapsed/refractory plasma cell dyscrasia

Any relevant practitioner

Re-assessment required after 12 months

Patient has no evidence of disease progression.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PROCARBAZINE HYDROCHLORIDE			
Cap 50 mg	980.00	50	Natulan
TEMOZOLOMIDE – Restricted see terms below			
⚡ Cap 5 mg	9.13	5	Temaccord
⚡ Cap 20 mg	16.38	5	Temaccord
⚡ Cap 100 mg	35.98	5	Temaccord
⚡ Cap 140 mg	50.12	5	Temaccord
⚡ Cap 250 mg	86.34	5	Temaccord

➡ **Restricted (RS1994)**

Initiation – gliomas

Re-assessment required after 12 months

Patient has a glioma.

Continuation – gliomas

Re-assessment required after 12 months

Treatment remains appropriate and patient is benefitting from treatment.

Initiation – Neuroendocrine tumours

Re-assessment required after 9 months

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Continuation – Neuroendocrine tumours

Re-assessment required after 6 months

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Initiation – ewing's sarcoma

Re-assessment required after 9 months

Patient has relapse or refractory Ewing's sarcoma.

Continuation – ewing's sarcoma

Re-assessment required after 6 months

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not funded for the treatment of relapsed high grade glioma.

THALIDOMIDE – **Restricted** see terms [below](#)

⚡ Cap 50 mg	378.00	28	Thalomid
⚡ Cap 100 mg	756.00	28	Thalomid

➡ **Restricted (RS2046)**

Initiation

Re-assessment required after 12 months

Either:

- 1 The patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; or
- 2 The patient has erythema nodosum leprosum.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation

Patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen

TRETINOIN

Cap 10 mg	479.50	100	Vesanoid
-----------------	--------	-----	----------

VENETOCLAX – **Restricted** see terms [below](#)

↓ Tab 14 x 10 mg, 7 x 50 mg, 21 x 100 mg	1,771.86	42	Venclexta
↓ Tab 10 mg	13.68	2	Venclexta
↓ Tab 50 mg	239.44	7	Venclexta
↓ Tab 100 mg	8,209.41	120	Venclexta

→ **Restricted (RS2118)**

Initiation – relapsed/refractory chronic lymphocytic leukaemia

Re-assessment required after 7 months

All of the following:

- 1 Individual has chronic lymphocytic leukaemia requiring treatment; and
- 2 Individual has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Individual has not previously received funded venetoclax; and
- 4 The individual's disease has relapsed; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Individual has an ECOG performance status of 0-2.

Continuation – relapsed/refractory chronic lymphocytic leukaemia

Re-assessment required after 6 months

Both:

- 1 Treatment remains clinically appropriate and the individual is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initiation – previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*

Re-assessment required after 6 months

All of the following:

- 1 Individual has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that the individual has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Individual has an ECOG performance status of 0-2.

Continuation – previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*

Re-assessment required after 6 months

No evidence of disease progression.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are unapproved indications

Initiation – previously untreated acute myeloid leukaemia

Re-assessment required after 6 months

Either:

- 1 The individual is currently on treatment with venetoclax and met all remaining special authority criteria prior to commencing treatment; or
- 2 All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.1 Individual has previously untreated acute myeloid leukaemia (see note a), according to World Health Organization (WHO) Classification; and
- 2.2 Venetoclax not to be used in combination with standard intensive remission induction chemotherapy; and
- 2.3 Venetoclax to be used in combination with azacitidine or low dose cytarabine.

Continuation – previously untreated acute myeloid leukaemia

Re-assessment required after 6 months

No evidence of disease progression.

Notes:

- a) 'Acute myeloid leukaemia' includes myeloid sarcoma*
- b) Indications marked with * are Unapproved indications

Platinum Compounds

CARBOPLATIN

Inj 10 mg per ml, 45 ml vial – 5% DV Dec-24 to 2027	25.73	1	Carboplatin Accord DBL Carboplatin
---	-------	---	--

CISPLATIN

Inj 1 mg per ml, 50 ml vial	9.45	1	Cisplatin Accord
Inj 1 mg per ml, 100 ml vial – 5% DV Dec-24 to 2027	18.90	1	Cisplatin Accord

OXALIPLATIN

Inj 5 mg per ml, 20 ml vial	33.35	1	Alchemy Oxaliplatin
-----------------------------------	-------	---	---------------------

Protein-Tyrosine Kinase Inhibitors

ALECTINIB – Restricted see terms [below](#)

⚡ Cap 150 mg	7,935.00	224	Alecensa
--------------------	----------	-----	----------

➡ **Restricted (RS1712)**

Initiation

Re-assessment required after 6 months

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- 3 Patient has an ECOG performance score of 0-2.

Continuation

Re-assessment required after 6 months

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

AXITINIB – Restricted see terms [below](#)

⚡ Tab 1 mg	536.40	28	Inlyta
⚡ Tab 5 mg	2,682.00	28	Inlyta

➡ **Restricted (RS2107)**

Initiation

Re-assessment required after 4 months

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 The disease is of predominant clear cell histology; and
- 3 The patient has documented disease progression following one previous line of treatment; and
- 4 The patient has ECOG performance status of 0-2.

Continuation

Re-assessment required after 4 months

No evidence of disease progression..

CRIZOTINIB – **Restricted** see terms [below](#)

⬇ Cap 200 mg	7,250.00	60	Xalkori
⬇ Cap 250 mg	7,250.00	60	Xalkori

➡ **Restricted (RS2108)**

Initiation

Re-assessment required after 6 months

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has a ROS1 rearrangement using an appropriate ROS1 test; and
- 3 Patient has ECOG performance score of 0-3; and
- 4 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation

Re-assessment required after 6 months

Both:

- 1 Response to treatment has been determined by comparable radiological assessment following the most recent treatment period; and
- 2 No evidence of disease progression..

DABRAFENIB – **Restricted** see terms [below](#)

⬇ Cap 50 mg	6,320.86	120	Tafinlar
⬇ Cap 75 mg	9,481.29	120	Tafinlar

➡ **Restricted (RS2130)**

Initiation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

Either:

- 1 The individual is currently on treatment with dabrafenib and trametinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or
 - 2.1.2 Both:
 - 2.1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and
 - 2.1.2.2 Adjuvant treatment with dabrafenib is required; and
 - 2.2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
 - 2.3 Treatment must be adjuvant to complete surgical resection; and
 - 2.4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
 - 2.5 The individual has a confirmed BRAF mutation; and
 - 2.6 Dabrafenib must be administered in combination with trametinib; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.7 The individual has ECOG performance score 0-2.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Continuation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

Any of the following:

- 1 All of the following:
 - 1.1 No evidence of disease recurrence; and
 - 1.2 Dabrafenib must be administered in combination with trametinib; and
 - 1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or
- 2 All of the following:
 - 2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 2.3 The individual meets initiation criteria for dabrafenib for unresectable or metastatic melanoma; or
- 3 All of the following:
 - 3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 3.2 The individual has received a BRAF/MEK inhibitor for unresectable or metastatic melanoma; and
 - 3.3 The individual meets continuation criteria for dabrafenib for unresectable or metastatic melanoma.

Initiation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

Either:

- 1 The individual is currently on treatment with dabrafenib and trametinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 2.2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.3 The individual has ECOG performance score 0-2; and
 - 2.4 The individual has confirmed BRAF mutation; and
 - 2.5 Dabrafenib must be administered in combination with trametinib; and
 - 2.6 Any of the following:
 - 2.6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 2.6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or
 - 2.6.3 All of the following:
 - 2.6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
 - 2.6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
 - 2.6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MEK inhibitor.

Continuation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

1 Any of the following:

- 1.1 The individual's disease has had a complete response to treatment; or
- 1.2 The individual's disease has had a partial response to treatment; or
- 1.3 The individual has stable disease with treatment; and

2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

DASATINIB – Restricted see terms [below](#)

↓ Tab 20 mg – 5% DV Mar-25 to 2027	132.88	60	Dasatinib-Teva
↓ Tab 50 mg – 5% DV Mar-25 to 2027	304.13	60	Dasatinib-Teva
↓ Tab 70 mg – 5% DV Mar-25 to 2027	415.75	60	Dasatinib-Teva

→ **Restricted (RS2055)**

Initiation

Haematologist or any relevant practitioner on the recommendation of a haematologist

Re-assessment required after 6 months

Any of the following:

- 1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or
- 2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or
- 3 Both:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Any of the following:
 - 3.2.1 Patient has documented treatment failure* with imatinib; or
 - 3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

Continuation

Haematologist or any relevant practitioner on the recommendation of a haematologist

Re-assessment required after 6 months

Both:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines.

ERLOTINIB – Restricted see terms [below](#)

↓ Tab 100 mg – 5% DV Oct-24 to 2027	280.84	30	Alchemy
↓ Tab 150 mg – 5% DV Oct-24 to 2027	484.24	30	Alchemy

→ **Restricted (RS2078)**

Initiation

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR; and
- 3 Any of the following:
 - 3.1 Patient is treatment naive; or
 - 3.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 3.3 Both:
 - 3.3.1 The patient has discontinued osimertinib or gefitinib due to intolerance; and
 - 3.3.2 The cancer did not progress while on osimertinib or gefitinib.

Continuation

Re-assessment required after 6 months

Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

GEFITINIB – **Restricted** see terms [below](#)

⚡ Tab 250 mg918.00 30 Iressa

➡ **Restricted (RS2079)**

Initiation

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Any of the following:
 - 2.1 Patient is treatment naive; or
 - 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued osimertinib or erlotinib due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on osimertinib or erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR.

Continuation

Re-assessment required after 6 months

Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

IMATINIB MESILATE

Cap 100 mg – **5% DV Dec-23 to 2026**44.93 60 **Imatinib-Rex**

Cap 400 mg – **5% DV Dec-23 to 2026**69.76 30 **Imatinib-Rex**

LAPATINIB – **Restricted** see terms [below](#)

⚡ Tab 250 mg

➡ **Restricted (RS1828)**

Initiation

For continuation use only.

Continuation

Re-assessment required after 12 months

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

LENVATINIB – **Restricted** see terms [below](#)

⚡ Cap 4 mg3,407.40 30 Lenvima

⚡ Cap 10 mg3,407.40 30 Lenvima

➡ **Restricted (RS2098)**

Initiation – thyroid cancer

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
 - 2.2 Either:
 - 2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
 - 2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures; and

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

- 2.3 Any of the following:
 - 2.3.1 A lesion without iodine uptake in a RAI scan; or
 - 2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
 - 2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
 - 2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other; and
- 2.4 Patient has thyroid stimulating hormone (TSH) adequately suppressed; and
- 2.5 Patient is not a candidate for radiotherapy with curative intent; and
- 2.6 Surgery is clinically inappropriate; and
- 2.7 Patient has an ECOG performance status of 0-2.

Continuation – thyroid cancer

Re-assessment required after 6 months

there is no evidence of disease progression.

Initiation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

All of the following:

- 1 Patient has unresectable hepatocellular carcinoma; and
- 2 Patient has preserved liver function (Childs-Pugh A); and
- 3 Transarterial chemoembolisation (TACE) is unsuitable; and
- 4 Patient has an ECOG performance status of 0-2; and
- 5 Either:
 - 5.1 Patient has not received prior systemic therapy for their disease in the palliative setting; or
 - 5.2 Both:
 - 5.2.1 Patient has experienced treatment-limiting toxicity from treatment with atezolizumab with bevacizumab; and
 - 5.2.2 No disease progression since initiation of atezolizumab with bevacizumab.

Continuation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

there is no evidence of disease progression.

Initiation – renal cell carcinoma

Re-assessment required after 4 months

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma; and
 - 1.2 The disease is of predominant clear-cell histology; and
 - 1.3 The patient has documented disease progression following one previous line of treatment; and
 - 1.4 The patient has an ECOG performance status of 0-2; and
 - 1.5 Lenvatinib is to be used in combination with everolimus; or
- 2 All of the following:
 - 2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
 - 2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
 - 2.3 Lenvatinib is to be used in combination with everolimus; and
 - 2.4 There is no evidence of disease progression.

Continuation – renal cell carcinoma

Re-assessment required after 4 months

there is no evidence of disease progression.

MIDOSTAURIN – Restricted see terms [on the next page](#)

⚡ Cap 25 mg	10,981.00	56	Rydapt
-------------------	-----------	----	--------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2033)

Initiation

All of the following:

- 1 Patient has a diagnosis of acute myeloid leukaemia; and
- 2 Condition must be FMS tyrosine kinase 3 (FLT3) mutation positive; and
- 3 Patient must not have received a prior line of intensive chemotherapy for acute myeloid leukaemia; and
- 4 Patient is to receive standard intensive chemotherapy in combination with midostaurin only; and
- 5 Midostaurin to be funded for a maximum of 4 cycles.

NILOTINIB – **Restricted** see terms [below](#)

⚡ Cap 150 mg	4,680.00	120	Tasigna
⚡ Cap 200 mg	6,532.00	120	Tasigna

➔ Restricted (RS2010)

Initiation

Haematologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with a tyrosine kinase inhibitor (TKI); or
 - 2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Continuation

Haematologist

Re-assessment required after 6 months

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

OSIMERTINIB – **Restricted** see terms [below](#)

⚡ Tab 40 mg	9,310.00	30	Tagrisso
⚡ Tab 80 mg	9,310.00	30	Tagrisso

➔ Restricted (RS2080)

Initiation – NSCLC – first line

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
- 2 Any of the following:
 - 2.1 Patient is treatment naïve; or
 - 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued gefitinib or erlotinib due to intolerance; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.3.2 The cancer did not progress while on gefitinib or erlotinib; and

3 There is documentation confirming that the cancer expresses activating mutations of EGFR; and

4 Patient has an ECOG performance status 0-3; and

5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – NSCLC – first line*Re-assessment required after 6 months*

response to or stable disease with treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Initiation – NSCLC – second line*Re-assessment required after 4 months*

All of the following:

1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and

2 Patient has an ECOG performance status 0-3; and

3 The patient must have received previous treatment with erlotinib or gefitinib; and

4 There is documentation confirming that the cancer expresses T790M mutation of EGFR following progression on or after erlotinib or gefitinib; and

5 The treatment must be given as monotherapy; and

6 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – NSCLC – second line*Re-assessment required after 6 months*

response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

PALBOCICLIB – Restricted see terms [below](#)

↓ Tab 75 mg	4,000.00	21	Ibrance
	1,200.00		Palbociclib Pfizer
↓ Tab 100 mg	4,000.00	21	Ibrance
	1,200.00		Palbociclib Pfizer
↓ Tab 125 mg	4,000.00	21	Ibrance
	1,200.00		Palbociclib Pfizer

*(Ibrance Tab 75 mg to be delisted 1 December 2025)**(Ibrance Tab 100 mg to be delisted 1 December 2025)**(Ibrance Tab 125 mg to be delisted 1 December 2025)*➔ **Restricted (RS2034)****Initiation***Re-assessment required after 6 months*

Either:

1 All of the following:

1.1 Patient has unresectable locally advanced or metastatic breast cancer; and

1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and

1.3 Patient has an ECOG performance score of 0-2; and

1.4 Either:

1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or

1.4.2 Both:

1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and

1.4.2.2 Patient has not received prior systemic treatment for metastatic disease; and

1.5 Treatment must be used in combination with an endocrine partner; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has an active Special Authority approval for ribociclib; and
 - 2.2 Patient has experienced a grade 3 or 4 adverse reaction to ribociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
 - 2.3 Treatment must be used in combination with an endocrine partner; and
 - 2.4 There is no evidence of progressive disease since initiation of ribociclib.

Continuation

Re-assessment required after 12 months

Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of palbociclib.

PAZOPANIB – **Restricted** see terms [below](#)

⚡ Tab 200 mg – 5% DV May-25 to 2027	172.88	30	Pazopanib Teva
⚡ Tab 400 mg – 5% DV May-25 to 2027	464.00	30	Pazopanib Teva

➡ **Restricted (RS2089)**

Initiation

Re-assessment required after 3 months

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and
 - 1.2 Either:
 - 1.2.1 The patient is treatment naive; or
 - 1.2.2 The patient has only received prior cytokine treatment; and
 - 1.3 The patient has an ECOG performance score of 0-2; and
The patient has intermediate or poor prognosis defined as:
 - 1.4 Any of the following:
 - 1.4.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 1.4.2 Haemoglobin level < lower limit of normal; or
 - 1.4.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 1.4.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 1.4.5 Karnofsky performance score of less than or equal to 70; or
 - 1.4.6 2 or more sites of organ metastasis; or
- 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and
 - 2.2 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on sunitinib; and
 - 2.4 Pazopanib to be used for a maximum of 3 months.

Continuation

Re-assessment required after 3 months

No evidence of disease progression.

RIBOCICLIB – **Restricted** see terms [below](#)

⚡ Tab 200 mg	1,883.00	21	Kisqali
	3,767.00	42	Kisqali
	5,650.00	63	Kisqali

➡ **Restricted (RS2131)**

Initiation

Re-assessment required after 6 months

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

1 All of the following:

- 1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 1.3 Patient has an ECOG performance score of 0-2; and
- 1.4 Either:
 - 1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
 - 1.4.2 Both:
 - 1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
 - 1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
- 1.5 Treatment to be used in combination with an endocrine partner; and
- 1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or

2 All of the following:

- 2.1 Patient has an active Special Authority approval for palbociclib; and
- 2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
- 2.3 Treatment must be used in combination with an endocrine partner; and
- 2.4 There is no evidence of progressive disease since initiation of palbociclib.

Continuation

Re-assessment required after 12 months

Both:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 There is no evidence of progressive disease since initiation of ribociclib.

RUXOLITINIB – **Restricted** see terms [below](#)

↓ Tab 5 mg	2,500.00	56	Jakavi
↓ Tab 10 mg	5,000.00	56	Jakavi
↓ Tab 15 mg	5,000.00	56	Jakavi
↓ Tab 20 mg	5,000.00	56	Jakavi

→ **Restricted (RS1726)**

Initiation

Haematologist

Re-assessment required after 12 months

All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation

Relevant specialist or medical practitioner on the recommendation of a Relevant specialist

Re-assessment required after 12 months

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB – **Restricted** see terms [below](#)

⚡ Cap 12.5 mg	208.38	28	Sunitinib Pfizer
⚡ Cap 25 mg	416.77	28	Sunitinib Pfizer
⚡ Cap 50 mg	694.62	28	Sunitinib Pfizer

➡ **Restricted (RS2109)**

Initiation – RCC

Re-assessment required after 4 months

Both:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 The patient has not previously received funded sunitinib.

Continuation – RCC

Re-assessment required after 4 months

No evidence of disease progression.

Initiation – GIST

Re-assessment required after 3 months

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Continuation – GIST

Re-assessment required after 6 months

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non-measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Continuation – GIST pandemic circumstances

Re-assessment required after 6 months

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Note: GIST - It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
TRAMETINIB – Restricted see terms below			
⬇ Tab 0.5 mg	2,370.32	30	Mekinist
⬇ Tab 2 mg	9,481.29	30	Mekinist

➔ **Restricted (RS2132)**

Initiation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

Either:

- 1 The individual is currently on treatment with dabrafenib and trametinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or
 - 2.1.2 Both:
 - 2.1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and
 - 2.1.2.2 Adjuvant treatment with trametinib is required; and
 - 2.2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
 - 2.3 Treatment must be adjuvant to complete surgical resection; and
 - 2.4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
 - 2.5 The individual has a confirmed BRAF mutation; and
 - 2.6 Trametinib must be administered in combination with dabrafenib; and
 - 2.7 The individual has ECOG performance score 0-2.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Continuation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

Any of the following:

- 1 All of the following:
 - 1.1 No evidence of disease recurrence; and
 - 1.2 Trametinib must be administered in combination with dabrafenib; and
 - 1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or
- 2 All of the following:
 - 2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 2.3 The individual meets initiation criteria for trametinib for unresectable or metastatic melanoma; or
- 3 All of the following:
 - 3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
 - 3.2 The individual has received a BRAF/MEK inhibitor for unresectable or metastatic melanoma; and
 - 3.3 The individual meets continuation criteria for trametinib for unresectable or metastatic melanoma.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

Either:

- 1 The individual is currently on treatment with dabrafenib and trametinib and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 2.2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.3 The individual has ECOG performance score 0-2; and
 - 2.4 The individual has confirmed BRAF mutation; and
 - 2.5 Trametinib must be administered in combination with dabrafenib; and
 - 2.6 Any of the following:
 - 2.6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 2.6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or
 - 2.6.3 All of the following:
 - 2.6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
 - 2.6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
 - 2.6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MEK inhibitor.

Continuation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

Both:

- 1 Any of the following:
 - 1.1 The individual's disease has had a complete response to treatment; or
 - 1.2 The individual's disease has had a partial response to treatment; or
 - 1.3 The individual has stable disease with treatment; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Taxanes

DOCETAXEL

Inj 10 mg per ml, 8 ml vial – 5% DV Dec-23 to 2026	24.91	1	DBL Docetaxel
--	-------	---	---------------

PACLITAXEL

Inj 6 mg per ml, 16.7 ml vial – 5% DV Aug-24 to 2026	19.59	1	Anzatax
Inj 6 mg per ml, 50 ml vial – 5% DV Aug-24 to 2026	37.89	1	Anzatax

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Treatment of Cytotoxic-Induced Side Effects

CALCIUM FOLINATE

Tab 15 mg	135.33	10	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 5 ml ampoule	18.25	5	Calcium Folate Ebewe
Inj 10 mg per ml, 5 ml vial.....	7.28	1	Calcium Folate Sandoz
	112.20	5	Eurofolic
Inj 10 mg per ml, 10 ml vial.....	9.49	1	Calcium Folate Sandoz
	163.35	5	Eurofolic
Inj 10 mg per ml, 30 ml vial.....	22.51	1	Calcium Folate Ebewe
Inj 10 mg per ml, 35 ml vial.....	25.14	1	Calcium Folate Sandoz
Inj 10 mg per ml, 100 ml vial.....	72.00	1	Calcium Folate Sandoz
	139.48		Eurofolic

(Calcium Folate Ebewe Inj 10 mg per ml, 5 ml ampoule to be delisted 1 November 2025)

(Calcium Folate Sandoz Inj 10 mg per ml, 5 ml vial to be delisted 1 November 2025)

(Calcium Folate Sandoz Inj 10 mg per ml, 10 ml vial to be delisted 1 November 2025)

(Calcium Folate Ebewe Inj 10 mg per ml, 30 ml vial to be delisted 1 November 2025)

(Calcium Folate Sandoz Inj 10 mg per ml, 35 ml vial to be delisted 1 November 2025)

(Calcium Folate Sandoz Inj 10 mg per ml, 100 ml vial to be delisted 1 November 2025)

DEXRAZOXANE – **Restricted** see terms [below](#)

↓ Inj 500 mg

e.g. Cardioxane

➔ **Restricted (RS1695)**

Initiation

Medical oncologist, paediatric oncologist, haematologist or paediatric haematologist

All of the following:

- 1 Patient is to receive treatment with high dose anthracycline given with curative intent; and
- 2 Based on current treatment plan, patient's cumulative lifetime dose of anthracycline will exceed 250mg/m² doxorubicin equivalent or greater; and
- 3 Dexrazoxane to be administered only whilst on anthracycline treatment; and
- 4 Either:
 - 4.1 Treatment to be used as a cardioprotectant for a child or young adult; or
 - 4.2 Treatment to be used as a cardioprotectant for secondary malignancy.

MESNA

Tab 400 mg	314.00	50	Uromitexan
Tab 600 mg	448.50	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule	177.45	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule	407.40	15	Uromitexan

Vinca Alkaloids

VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial.....	270.37	5	Hospira
----------------------------------	--------	---	---------

VINCISTINE SULPHATE

Inj 1 mg per ml, 1 ml vial.....	74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial.....	102.73	5	DBL Vincristine Sulfate

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
VINORELBINE			
Cap 20 mg – 5% DV Feb-26 to 2028	32.10	1	Vinorelbine Te Arai
Cap 30 mg – 5% DV Feb-26 to 2028	42.80	1	Vinorelbine Te Arai
Cap 80 mg – 5% DV Feb-26 to 2028	80.00	1	Vinorelbine Te Arai
Inj 10 mg per ml, 1 ml vial			
Inj 10 mg per ml, 5 ml vial			

Endocrine Therapy

ABIRATERONE ACETATE – Restricted see terms [below](#)

⚡ Tab 250 mg4,276.19 120 Zytiga

➡ **Restricted (RS1888)**

Initiation

Medical oncologist, radiation oncologist or urologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Continuation

Medical oncologist, radiation oncologist or urologist

Re-assessment required after 6 months

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Continuation – pandemic circumstances

Re-assessment required after 6 months

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE

Tab 50 mg – **5% DV Dec-23 to 2026**.....4.18 28 **Binarex**

FLUTAMIDE

Tab 250 mg119.50 100 Flutamin

FULVESTRANT – Restricted see terms [on the next page](#)

⚡ Inj 50 mg per ml, 5 ml prefilled syringe.....1,068.00 2 Faslodex

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ **Restricted (RS1732)**

Initiation

Medical oncologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Continuation

Medical oncologist

Re-assessment required after 6 months

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 No evidence of disease progression.

OCTREOTIDE

Inj 100 mcg per ml, 1 ml vial.....	48.50	5	Omega
Inj 50 mcg per ml, 1 ml vial.....	27.58	5	Omega
Inj 500 mcg per ml, 1 ml vial.....	113.10	5	Omega
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	Max Health
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	Max Health
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	Max Health

TAMOXIFEN CITRATE

Tab 10 mg – 5% DV Dec-23 to 2026	15.00	60	Tamoxifen Sandoz
Tab 20 mg – 5% DV Dec-23 to 2026	5.32	60	Tamoxifen Sandoz

Aromatase Inhibitors

ANASTROZOLE

Tab 1 mg – 5% DV Dec-23 to 2026	4.39	30	Anatrole
--	------	----	-----------------

EXEMESTANE

Tab 25 mg – 5% DV Nov-23 to 2026	9.86	30	Pfizer Exemestane
---	------	----	--------------------------

LETROZOLE

Tab 2.5 mg – 5% DV Dec-24 to 2027	4.36	28	Accord
	4.67	30	Letrole

Long-acting Somatostatin Analogues

➔ **Restricted (RS2100)**

Initiation – Malignant bowel obstruction

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has not been successful; and
- 3 Treatment to be given for up to 4 weeks.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: Indications marked with * are unapproved indications

Initiation – acromegaly

Re-assessment required after 3 months

All of the following:

- 1 The patient has acromegaly; and
- 2 Either:
 - 2.1 Treatment with surgery and radiotherapy is not suitable or was unsuccessful; or
 - 2.2 Treatment is for an interim period while awaiting the beneficial effects of radiotherapy; and
- 3 Treatment with a dopamine agonist has been unsuccessful.

Continuation – acromegaly

Without reassessment for applications where IGF1 levels have decreased since starting treatment.

Note: In patients with acromegaly, treatment should be discontinued if IGF1 levels have no decreased 3 months after treatment.

In patients treated with radiotherapy treatment should be withdrawn every 2 years, for 1 month, for assessment of remission.

Treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following treatment withdrawal for at least 4 weeks.

Initiation – Other indications

Any of the following:

- 1 VIPomas and glucagonomas - for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:
 - 2.2.1 Surgery has been unsuccessful; or
 - 2.2.2 Patient has metastatic disease after treatment with H2 antagonist or proton pump inhibitors has been unsuccessful; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has not been successful; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Initiation – pre-operative acromegaly

Limited to 12 months treatment

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

Notes: Indications marked with * are unapproved indications

The use of a long-acting somatostatin analogue in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded under Special Authority

LANREOTIDE – **Restricted** see terms [on the previous page](#)

† Inj 60 mg per 0.5 ml, 0.5 ml syringe – 5% DV Aug-25 to 2027	382.77	1	Mytolac
† Inj 90 mg per 0.5 ml, 0.5 ml syringe – 5% DV Sep-25 to 2027	562.92	1	Mytolac
† Inj 120 mg per 0.5 ml, 0.5 ml syringe – 5% DV Aug-25 to 2027	646.70	1	Mytolac

OCTREOTIDE LONG-ACTING – **Restricted** see terms [on the previous page](#)

† Inj depot 10 mg prefilled syringe – 5% DV Dec-24 to 2027	438.40	1	Sandostatin LAR
† Inj depot 20 mg prefilled syringe – 5% DV Dec-24 to 2027	583.70	1	Sandostatin LAR
† Inj depot 30 mg prefilled syringe – 5% DV Dec-24 to 2027	670.80	1	Sandostatin LAR

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Imaging Agents

AMINOLEVULINIC ACID HYDROCHLORIDE – **Restricted** see terms [below](#)

↓ Powder for oral soln, 30 mg per ml, 1.5 g vial	4,400.00	1	Gliolan
	44,000.00	10	Gliolan

→ **Restricted (RS1565)**

Initiation – high grade malignant glioma

All of the following:

- 1 Patient has newly diagnosed, untreated, glioblastoma multiforme; and
- 2 Treatment to be used as adjuvant to fluorescence-guided resection; and
- 3 Patient's tumour is amenable to complete resection.

Immunosuppressants

Calcineurin Inhibitors

CICLOSPORIN

Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml	198.13	50 ml	Neoral
Inj 50 mg per ml, 5 ml ampoule	276.30	10	Sandimmun

TACROLIMUS – **Restricted** see terms [below](#)

↓ Cap 0.5 mg	49.60	100	Tacrolimus Sandoz
↓ Cap 0.75 mg	99.30	100	Tacrolimus Sandoz
↓ Cap 1 mg	84.30	100	Tacrolimus Sandoz
↓ Cap 5 mg	248.20	50	Tacrolimus Sandoz
↓ Inj 5 mg per ml, 1 ml ampoule			

→ **Restricted (RS2110)**

Initiation – organ transplant recipients

Either:

- 1 For use in organ transplant recipients; or
- 2 The individual is receiving induction therapy for an organ transplant.

Initiation – non-transplant indications*

Any specialist

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Either:
 - 2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
 - 2.2 Patient is a child with nephrotic syndrome*.

Note: Indications marked with * are unapproved indications

Fusion Proteins

ETANERCEPT – **Restricted** see terms [on the next page](#)

↓ Inj 25 mg autoinjector	690.00	4	Enbrel
↓ Inj 25 mg vial	690.00	4	Enbrel
↓ Inj 50 mg autoinjector	1,050.00	4	Enbrel
↓ Inj 50 mg syringe	1,050.00	4	Enbrel

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2062)

Initiation – polyarticular course juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Continuation – polyarticular course juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 6 months

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – oligoarticular course juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Continuation – oligoarticular course juvenile idiopathic arthritis

Rheumatologist or named specialist

Re-assessment required after 6 months

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - rheumatoid

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – Arthritis - rheumatoid

Any relevant practitioner

Re-assessment required after 2 years

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation – ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of starting treatment.

Average normal chest expansion corrected for age and gender:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Age	Male	Female
18-24	7.0 cm	5.5 cm
25-34	7.5 cm	5.5 cm
35-44	6.5 cm	4.5 cm
45-54	6.0 cm	5.0 cm
55-64	5.5 cm	4.0 cm
65-74	4.0 cm	4.0 cm
75+	3.0 cm	2.5 cm

Continuation – ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation – psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- Both:
 - The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - Either:
 - The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- All of the following:
 - Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - Either:
 - Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - Any of the following:
 - Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Continuation – psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and

2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation – severe chronic plaque psoriasis, prior TNF use

Dermatologist

Limited to 4 months treatment

All of the following:

1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and

2 Either:

- 2.1 The patient has experienced intolerable side effects from adalimumab; or
- 2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; and

3 Patient must be reassessed for continuation after 3 doses.

Initiation – severe chronic plaque psoriasis, treatment-naïve

Dermatologist

Limited to 4 months treatment

All of the following:

1 Any of the following:

- 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
- 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
- 1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and

2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation – severe chronic plaque psoriasis

Re-assessment required after 6 months

Both:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

1 Any of the following:

1.1 Both:

1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

1.1.2 Either:

1.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-etanercept treatment baseline value; or

1.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or

1.2 Both:

1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

1.2.2 Either:

1.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or

1.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-etanercept treatment baseline value; or

1.3 Both:

1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and

1.3.2 Either:

1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or

1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing etanercept; and

2 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initiation – pyoderma gangrenosum

Dermatologist

All of the following:

1 Patient has pyoderma gangrenosum*; and

2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and

3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Continuation – pyoderma gangrenosum

Dermatologist

All of the following:

1 Patient has shown clinical improvement; and

2 Patient continues to require treatment; and

3 A maximum of 8 doses.

Initiation – adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

Either:

1 Both:

1.1 Either:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Continuation – adult-onset Still's disease

Rheumatologist

Re-assessment required after 6 months

The patient has a sustained improvement in inflammatory markers and functional status.

Initiation – undifferentiated spondyloarthritis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Continuation – undifferentiated spondyloarthritis

Rheumatologist or medical practitioner on the recommendation of a Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Monoclonal Antibodies

ABCIXIMAB – **Restricted** see terms [below](#)

↓ Inj 2 mg per ml, 5 ml vial

→ **Restricted (RS1202)**

Initiation

Either:

- 1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
- 2 For use in patients undergoing intra-cranial intervention.

ADALIMUMAB (AMGEVITA) – **Restricted** see terms [below](#)

↓ Inj 20 mg per 0.4 ml prefilled syringe – 5% DV Oct-22 to 31 Jul 2026.....190.00 1 **Amgevita**

↓ Inj 40 mg per 0.8 ml prefilled pen – 5% DV Oct-22 to 31 Jul 2026375.00 2 **Amgevita**

↓ Inj 40 mg per 0.8 ml prefilled syringe – 5% DV Oct-22 to 31 Jul 2026.....375.00 2 **Amgevita**

→ **Restricted (RS2140)**

Initiation – Behcet's disease - severe

Any relevant practitioner

Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initiation – Hidradenitis suppurativa

Dermatologist

Re-assessment required after 4 months

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Continuation – Hidradenitis suppurativa

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – Plaque psoriasis - severe chronic

Dermatologist

Re-assessment required after 4 months

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or (DLQI) assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Continuation – Plaque psoriasis - severe chronic

Re-assessment required after 2 years

Any of the following:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has experienced a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
- 3 Both:
 - 3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
 - 3.2 Either:

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

- 3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
- 3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

Initiation – pyoderma gangrenosum

Dermatologist

Both:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initiation – Crohn's disease - adults

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease - adults

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initiation – Crohn's disease - children

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease - children

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initiation – Crohn's disease - fistulising

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Continuation – Crohn's disease - fistulising

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initiation – Ocular inflammation - chronic

Any relevant practitioner

Re-assessment required after 4 months

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Continuation – Ocular inflammation - chronic

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Initiation – Ocular inflammation - severe

Any relevant practitioner

Re-assessment required after 4 months

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Continuation – Ocular inflammation - severe

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initiation – ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – ankylosing spondylitis

Any relevant practitioner

Re-assessment required after 2 years

For applications where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initiation – Arthritis - oligoarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Continuation – Arthritis - oligoarticular course juvenile idiopathic

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - polyarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

- 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Continuation – Arthritis - polyarticular course juvenile idiopathic

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - psoriatic

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation – Arthritis - psoriatic

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – Arthritis - rheumatoid

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin (unless contraindicated); or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomid (unless contraindicated) alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Continuation – Arthritis - rheumatoid

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initiation – Still's disease - adult-onset (AOSD)

Rheumatologist

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for (AOSD); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initiation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 20; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Continuation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or
- 2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on biologic therapy.

Initiation – undifferentiated spondyloarthritis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulphasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Continuation – undifferentiated spondyloarthritis

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

continued...

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

Initiation – inflammatory bowel arthritis – axial

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Continuation – inflammatory bowel arthritis – axial

Any relevant practitioner

Re-assessment required after 2 years

Where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initiation – inflammatory bowel arthritis – peripheral

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulphasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation – inflammatory bowel arthritis – peripheral

Any relevant practitioner

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) – **Restricted** see terms [below](#)

⚡ Inj 20 mg per 0.2 ml prefilled syringe.....	595.50	2	Humira
⚡ Inj 40 mg per 0.4 ml prefilled syringe.....	595.50	2	Humira
⚡ Inj 40 mg per 0.4 ml prefilled pen	595.50	2	HumiraPen

➡ **Restricted (RS1922)**

Initiation – Behcet's disease – severe

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

1 Either:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

2 Patient has received a maximum of 6 months treatment with Amgevita; and

3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Behcet's disease – severe

Any relevant practitioner

Re-assessment required after 6 months

Both:

1 The patient has had a good clinical response to treatment with measurably improved quality of life; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Hidradenitis suppurativa

Dermatologist or Practitioner on the recommendation of a dermatologist

Re-assessment required after 6 months

All of the following:

1 Either:

1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

2 Patient has received a maximum of 6 months treatment with Amgevita; and

3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Continuation – Hidradenitis suppurativa

Dermatologist or Practitioner on the recommendation of a dermatologist

Re-assessment required after 6 months

All of the following:

1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and

2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and

3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initiation – Psoriasis - severe chronic plaque

Dermatologist or Practitioner on the recommendation of a dermatologist

Re-assessment required after 6 months

All of the following:

1 Either:

1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

2 Patient has received a maximum of 6 months treatment with Amgevita; and

3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – Psoriasis - severe chronic plaque

Dermatologist or Practitioner on the recommendation of a dermatologist

Re-assessment required after 6 months

Both:

1 Either:

1.1 Both:

1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

1.1.2 Either:

1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or

1.2 Both:

1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

1.2.2 Either:

1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or

1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Pyoderma gangrenosum

Dermatologist

Re-assessment required after 6 months

All of the following:

1 Either:

1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

2 Patient has received a maximum of 6 months treatment with Amgevita; and

3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

4 A maximum of 8 doses.

Continuation – Pyoderma gangrenosum

Dermatologist

Re-assessment required after 6 months

Both:

1 The patient has demonstrated clinical improvement and continues to require treatment; and

2 A maximum of 8 doses.

Initiation – Crohn's disease - adult

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

All of the following:

1 Any of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Crohn's disease - adult

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Crohn's disease - children

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Crohn's disease - children

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Crohn's disease - fistulising

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

All of the following:

- 1 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Crohn's disease - fistulising

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Ocular inflammation – chronic

Any relevant practitioner

Re-assessment required after 12 months

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Ocular inflammation – chronic

Any relevant practitioner

Re-assessment required after 12 months

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Ocular inflammation – severe

Any relevant practitioner

Re-assessment required after 12 months

All of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

1 Any of the following:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and

2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Ocular inflammation – severe

Any relevant practitioner

Re-assessment required after 12 months

Both:

1 Any of the following:

- 1.1 The patient has had a good clinical response following 3 initial doses; or
- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – ankylosing spondylitis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

1 Either:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and

2 Patient has received a maximum of 6 months treatment with Amgevita; and

3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – ankylosing spondylitis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Both:

1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Arthritis – oligoarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

All of the following:

1 Either:

- 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Continuation – Arthritis – oligoarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

For patients that demonstrate at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - polyarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Continuation – Arthritis - polyarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

For patients that demonstrate at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - psoriatic

Named specialist or rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Arthritis - psoriatic

Named specialist or rheumatologist

Re-assessment required after 6 months

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

Initiation – Arthritis – rheumatoid

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Continuation – Arthritis – rheumatoid

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Either:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initiation – Still's disease – adult-onset (AOSD)

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Continuation – Still's disease – adult-onset (AOSD)

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT – Restricted see terms [below](#)

↓ Inj 40 mg per ml, 0.1 ml vial..... 1,250.00 1 Eylea

➔ **Restricted (RS1872)**

Initiation – Wet Age Related Macular Degeneration

Ophthalmologist or nurse practitioner

Re-assessment required after 3 months

Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Continuation – Wet Age Related Macular Degeneration

Ophthalmologist or nurse practitioner
Re-assessment required after 12 months
All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Initiation – Diabetic Macular Oedema

Ophthalmologist or nurse practitioner
Re-assessment required after 4 months
All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 – 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Continuation – Diabetic Macular Oedema

Ophthalmologist or nurse practitioner
Re-assessment required after 12 months
All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with aflibercept, patient has retrialled with at least one injection of bevacizumab and had no response.

BASILIXIMAB – Restricted see terms below

⚡ Inj 20 mg vial2,560.00 1 Simulect

➡ Restricted (RS1203)

Initiation

For use in solid organ transplants.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

BENRALIZUMAB – Restricted see terms [below](#)

↓ Inj 30 mg per ml, 1 ml prefilled pen 3,539.00 1 Fasenra

→ **Restricted (RS1920)**

Initiation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 12 months

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5×10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Continuation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 2 years

- Both:
- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
 - 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

BEVACIZUMAB – Restricted see terms [below](#)

↓ Inj 25 mg per ml, 4 ml vial – **10% DV Aug-25 to 31 Aug 2028** 69.00 1 **Vegzelma**

↓ Inj 25 mg per ml, 16 ml vial – **10% DV Aug-25 to 31 Aug 2028** 276.00 1 **Vegzelma**

→ **Restricted (RS2111)**

Initiation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

Either:

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- 1 Patient is currently on treatment with bevacizumab, and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
 - 2.2 Patient has preserved liver function (Child-Pugh A); and
 - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
 - 2.4 Any of the following:
 - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
 - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
 - 2.4.3 Both:
 - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
 - 2.4.3.2 No disease progression since initiation of lenvatinib; and
 - 2.5 Patient has an ECOG performance status of 0-2; and
 - 2.6 To be given in combination with atezolizumab.

Continuation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

no evidence of disease progression.

Initiation – advanced or metastatic ovarian cancer

Re-assessment required after 4 months

All of the following:

- 1 Either:
 - 1.1 The patient has FIGO Stage IV epithelial ovarian, fallopian tube, or primary peritoneal cancer; or
 - 1.2 Both:
 - 1.2.1 The patient has previously untreated advanced (FIGO Stage IIIB or IIIC) epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
 - 1.2.2 Either:
 - 1.2.2.1 Debulking surgery is inappropriate; or
 - 1.2.2.2 The cancer is sub-optimally debulked (maximum diameter of any gross residual disease greater than 1cm); and
- 2 Bevacizumab to be administered at a maximum dose of 15 mg/kg every three weeks; and
- 3 18 weeks concurrent treatment with chemotherapy is planned.

Continuation – advanced or metastatic ovarian cancer

Re-assessment required after 4 months

no evidence of disease progression.

Initiation – Recurrent Respiratory Papillomatosis

Re-assessment required after 12 months

All of the following:

- 1 Maximum of 6 doses; and
- 2 The patient has recurrent respiratory papillomatosis; and
- 3 The treatment is for intra-lesional administration.

Continuation – Recurrent Respiratory Papillomatosis

Re-assessment required after 12 months

All of the following:

- 1 Maximum of 6 doses; and
- 2 The treatment is for intra-lesional administration; and
- 3 There has been a reduction in surgical treatments or disease regrowth as a result of treatment.

Initiation – Ocular Conditions

Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Ocular neovascularisation; or
- 2 Exudative ocular angiopathy.

BEVACIZUMAB (OCULAR) – **Restricted** see terms [below](#)

↓ Inj 25 mg per ml, 4 ml vial

↓ Inj 25 mg per ml, 16 ml vial

→ **Restricted (RS1691)**

Initiation – Recurrent Respiratory Papillomatosis

Otolaryngologist

Re-assessment required after 12 months

All of the following:

- 1 Maximum of 6 doses; and
- 2 The patient has recurrent respiratory papillomatosis; and
- 3 The treatment is for intra-lesional administration.

Continuation – Recurrent Respiratory Papillomatosis

Otolaryngologist

Re-assessment required after 12 months

All of the following:

- 1 Maximum of 6 doses; and
- 2 The treatment is for intra-lesional administration; and
- 3 There has been a reduction in surgical treatments or disease regrowth as a result of treatment.

Initiation – ocular conditions

Either:

- 1 Ocular neovascularisation; or
- 2 Exudative ocular angiopathy.

BRENTUXIMAB VEDOTIN – **Restricted** see terms [below](#)

↓ Inj 50 mg vial 5,275.18 1 Adcetris

→ **Restricted (RS2002)**

Initiation – relapsed/refractory Hodgkin lymphoma

Re-assessment required after 6 months

All of the following:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and
 - 1.1.2 Patient is ineligible for autologous stem cell transplant; or
 - 1.2 Both:
 - 1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and
 - 1.2.2 Patient has previously undergone autologous stem cell transplant; and
- 2 Patient has not previously received funded brentuximab vedotin; and
- 3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Continuation – relapsed/refractory Hodgkin lymphoma

Re-assessment required after 9 months

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

Initiation – anaplastic large cell lymphoma

Re-assessment required after 9 months

All of the following:

- 1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and
- 2 Patient has an ECOG performance status of 0-1; and
- 3 Patient has not previously received brentuximab vedotin; and
- 4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and
- 5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Continuation – anaplastic large cell lymphoma

Re-assessment required after 9 months

All of the following:

- 1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and
- 2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and
- 3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

CETUXIMAB – Restricted see terms [below](#)

⚡ Inj 5 mg per ml, 20 ml vial.....	364.00	1	Erbix
⚡ Inj 5 mg per ml, 100 ml vial.....	1,820.00	1	Erbix

➡ Restricted (RS2064)

Initiation – head and neck cancer, locally advanced

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Cisplatin is contraindicated or has resulted in intolerable side effects; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 To be administered in combination with radiation therapy.

Initiation – colorectal cancer, metastatic

Re-assessment required after 6 months

All of the following:

- 1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and
- 2 There is documentation confirming disease is RAS and BRAF wild-type; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Patient has not received prior funded treatment with cetuximab; and
- 5 Either:
 - 5.1 Cetuximab is to be used in combination with chemotherapy; or
 - 5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

Continuation – colorectal cancer, metastatic

Re-assessment required after 6 months

No evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

GEMTUZUMAB OZOGAMICIN – Restricted see terms [below](#)

⚡ Inj 5 mg vial	12,973.00	1	Myotarg
-----------------------	-----------	---	---------

➡ Restricted (RS1923)

Initiation

All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB – **Restricted** see terms [below](#)

↓ Inj 100 mg.....428.00 1 Remicade

→ **Restricted (RS2124)**

Initiation – Graft vs host disease

Patient has steroid-refractory acute graft vs. host disease of the gut.

Initiation – rheumatoid arthritis

Rheumatologist

Re-assessment required after 4 months

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Continuation – rheumatoid arthritis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initiation – ankylosing spondylitis

Rheumatologist

Re-assessment required after 3 months

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

adalimumab and/or etanercept for ankylosing spondylitis.

Continuation – ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initiation – psoriatic arthritis

Rheumatologist

Re-assessment required after 4 months

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Continuation – psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initiation – severe ocular inflammation

Re-assessment required after 4 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – severe ocular inflammation

Re-assessment required after 12 months

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initiation – chronic ocular inflammation

Re-assessment required after 4 months

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Continuation – chronic ocular inflammation

Re-assessment required after 12 months

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initiation – Pulmonary sarcoidosis

Both:

- 1 Patient has life-threatening pulmonary sarcoidosis that is refractory to other treatments; and
- 2 Treatment is to be prescribed by, or has been recommended by, a physician with expertise in the treatment of pulmonary sarcoidosis.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Initiation – Crohn's disease (adults)

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease (adults)

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – Crohn's disease (children)

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease (children)

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

Initiation – fistulising Crohn's disease

Gastroenterologist

Re-assessment required after 6 months

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complete peri-anal fistula.

Continuation – fistulising Crohn's disease

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – acute fulminant ulcerative colitis

Gastroenterologist

Limited to 6 weeks treatment

Both:

- 1 Patient has acute, fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Continuation – fulminant ulcerative colitis

Any relevant practitioner

Re-assessment required after 2 years

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Either:
 - 2.1 Patients SCCAI is greater than or equal to 4; or
 - 2.2 Patients PUCAI score is greater than or equal to 20; and
- 3 Patient has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 2 years

Both:

1 Either:

- 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
- 1.2 The PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and

2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – plaque psoriasis

Dermatologist

Re-assessment required after 3 doses

Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or

2 All of the following:

- 2.1 Any of the following:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, cyclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

Continuation – plaque psoriasis

Re-assessment required after 3 doses

Both:

1 Any of the following:

1.1 Both:

- 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or

1.2 Both:

- 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; or

1.3 Both:

- 1.3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
- 1.3.2 Either:
 - 1.3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing infliximab; and

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initiation – neurosarcoidosis

Neurologist

Re-assessment required after 18 months

All of the following:

- 1 Biopsy consistent with diagnosis of neurosarcoidosis; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Continuation – neurosarcoidosis

Neurologist

Re-assessment required after 18 months

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

Initiation – severe Behcet's disease

Re-assessment required after 4 months

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes:

- a) Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.
- b) Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Continuation – severe Behcet's disease

Re-assessment required after 6 months

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initiation – pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has pyoderma gangrenosum*;
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Continuation – pyoderma gangrenosum

Dermatologist

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initiation – Inflammatory bowel arthritis (axial)

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has had axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
- 5 Patient has not experienced an adequate response to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 Patient has a BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$ Per	Manufacturer

continued...

previous pharmacological treatment .

Continuation – Inflammatory bowel arthritis (axial)

Re-assessment required after 2 years

Where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initiation – Inflammatory bowel arthritis (peripheral)

Re-assessment required after 6 months

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
- 4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation – Inflammatory bowel arthritis (peripheral)

Re-assessment required after 2 years

Either:

- 1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

Initiation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and
- 2 The individual has received insufficient benefit from use of corticosteroids; and
- 3 Infliximab is to be administered at up to 5mg/kg for up to four doses.

Continuation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

Both:

- 1 The individual has shown clinical improvement and ongoing treatment is required; and
- 2 Infliximab is to be administered at up to 5mg/kg for up to a total of 8 doses.

Note: Indications marked with * are unapproved indications.

INOTUZUMAB OZOGAMICIN – **Restricted** see terms [below](#)

↓ Inj 1 mg vial 14,457.00 1 Besponsa

➔ **Restricted (RS2112)**

Initiation

Re-assessment required after 4 months

All of the following:

continued...

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has relapsed or refractory CD22-positive B-cell acute lymphoblastic leukaemia/lymphoma, including minimal residual disease; and
- 2 Patient has ECOG performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient has Philadelphia chromosome positive B-Cell ALL; and
 - 3.1.2 Patient has previously received a tyrosine kinase inhibitor; or
 - 3.2 Patient has received one prior line of treatment involving intensive chemotherapy; and
- 4 Treatment is to be administered for a maximum of 3 cycles.

Continuation

Re-assessment required after 4 months

All of the following:

- 1 Patient is not proceeding to a stem cell transplant; and
- 2 Either:
 - 2.1 Patient has experienced complete disease response; or
 - 2.2 Patient has experienced complete remission with incomplete haematological recovery; and
- 3 Treatment with inotuzumab ozogamicin is to cease after a total duration of 6 cycles.

MEPOLIZUMAB – Restricted see terms below

⚡ Inj 100 mg prefilled pen	1,638.00	1	Nucala
⚡ Inj 100 mg vial			

➡ Restricted (RS2024)

Initiation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 12 months

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5×10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab; and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

9.2 Both:

9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and

9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Continuation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 2 years

Both:

1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and

2 Either:

2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or

2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initiation – eosinophilic granulomatosis with polyangiitis

Re-assessment required after 12 months

All of the following:

1 The patient has eosinophilic granulomatosis with polyangiitis; and

2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and

3 Either:

3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or

3.2 Corticosteroids are contraindicated.

Continuation – eosinophilic granulomatosis with polyangiitis

Re-assessment required after 12 months

Patient has no evidence of clinical disease progression.

OBINUTUZUMAB – **Restricted** see terms [below](#)

↓ Inj 25 mg per ml, 40 ml vial.....5,910.00 1 Gazyva

➔ **Restricted (RS1919)**

Initiation

Haematologist

Limited to 6 months treatment

All of the following:

1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and

2 The patient is obinutuzumab treatment naive; and

3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and

4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and

5 Patient has good performance status; and

6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

* greater than or equal to $1.5 \times 10^9/L$ and platelets greater than or equal to $75 \times 10^9/L$

Initiation – follicular / marginal zone lymphoma

Re-assessment required after 9 months

All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Continuation – follicular / marginal zone lymphoma

Re-assessment required after 24 months

All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB – Restricted see terms [below](#)

⚡ Inj 150 mg prefilled syringe.....	450.00	1	Xolair
⚡ Inj 150 mg vial	450.00	1	Xolair

➡ **Restricted (RS1652)**

Initiation – severe asthma

Clinical immunologist or respiratory specialist

Re-assessment required after 6 months

All of the following:

- 1 Patient must be aged 6 years or older ; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or formoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – severe asthma

Respiratory specialist

Re-assessment required after 6 months

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Initiation – severe chronic spontaneous urticaria

Clinical immunologist or dermatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; and
 - 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
 - 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Continuation – severe chronic spontaneous urticaria

Clinical immunologist or dermatologist

Re-assessment required after 6 months

Either:

- 1 Patient has previously had a complete response* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB – **Restricted** see terms [below](#)

↓ Inj 100 mg per ml, 1 ml vial.....1,700.00 1 Synagis

➔ **Restricted (RS2081)**

Initiation

Re-assessment required after 6 months

Both:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Infant was born in the last 12 months; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.1.2 Infant was born at less than 32 weeks zero days' gestation; or
- 2.2 Both:
 - 2.2.1 Child was born in the last 24 months; and
 - 2.2.2 Any of the following:
 - 2.2.2.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
 - 2.2.2.2 Both:
 - 2.2.2.2.1 Child has haemodynamically significant heart disease; and
 - 2.2.2.2.2 Any of the following:
 - 2.2.2.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
 - 2.2.2.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
 - 2.2.2.2.2.3 Child has severe pulmonary hypertension (see Note C); or
 - 2.2.2.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
 - 2.2.2.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
 - 2.2.2.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Continuation

Re-assessment required after 6 months

All of the following:

- 1 Palivizumab to be administered during the annual RSV season; and
- 2 Child was born in the last 24 months; and
- 3 Any of the following:
 - 3.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
 - 3.2 Both:
 - 3.2.1 Child has haemodynamically significant heart disease; and
 - 3.2.2 Any of the following:
 - 3.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
 - 3.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
 - 3.2.2.3 Child has severe pulmonary hypertension (see Note C); or
 - 3.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
 - 3.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
 - 3.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Notes:

- a) Ventilatory/respiratory support includes those on home oxygen, CPAP/VPAP and those with tracheostomies in situ managed at home
- b) Child requires/will require heart failure medication, and/or child has significant pulmonary hypertension, and/or infant will require surgical palliation/definitive repair within the next 3 months
- c) Mean pulmonary artery pressure more than 25 mmHg
- d) LV Ejection Fraction less than 40%
- e) Inborn errors of immunity include, but are not limited to, IFNAR deficiencies

PERTUZUMAB – Restricted see terms [on the next page](#)

⚡ Inj 30 mg per ml, 14 ml vial.....3,927.00 1 Perjeta

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

➔ Restricted (RS1995)

Initiation

Re-assessment required after 12 months

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naive; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Continuation

Re-assessment required after 12 months

Either:

- 1 Both:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

RANIBIZUMAB – **Restricted** see terms [below](#)

⬇ Inj 10 mg per ml, 0.23 ml vial

⬇ Inj 10 mg per ml, 0.3 ml vial

➔ Restricted (RS1870)

Initiation – Wet Age Related Macular Degeneration

Ophthalmologist or nurse practitioner

Re-assessment required after 3 months

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with aflibercept for longer than 3 months; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2 Patient has current approval to use aflibercept for treatment of wAMD and was found to be intolerant to aflibercept within 3 months.

Continuation – Wet Age Related Macular Degeneration

Ophthalmologist or nurse practitioner

Re-assessment required after 12 months

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

RITUXIMAB (MABTHERA) – **Restricted** see terms [below](#)

⚡ Inj 10 mg per ml, 10 ml vial.....	1,075.50	2	Mabthera
⚡ Inj 10 mg per ml, 50 ml vial.....	2,688.30	1	Mabthera

➡ **Restricted (RS1785)**

Initiation – rheumatoid arthritis - prior TNF inhibitor use

Rheumatologist

Limited to 4 months treatment

All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initiation – rheumatoid arthritis - TNF inhibitors contraindicated

Rheumatologist

Limited to 4 months treatment

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of cyclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

6 Either:

- 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
- 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

7 Either:

- 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and

8 Either:

- 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and

9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Continuation – rheumatoid arthritis - re-treatment in 'partial responders' to rituximab

Rheumatologist

Re-assessment required after 4 months

All of the following:

1 Any of the following:

- 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and

2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and

3 Either:

- 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and

4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Continuation – rheumatoid arthritis - re-treatment in 'responders' to rituximab

Rheumatologist

Re-assessment required after 4 months

All of the following:

1 Either:

- 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and

2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and

3 Either:

- 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and

4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) – **Restricted** see terms [on the next page](#)

⚡ Inj 10 mg per ml, 10 ml vial.....	275.33	2	Riximyo
⚡ Inj 10 mg per ml, 50 ml vial.....	688.20	1	Riximyo

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

➔ Restricted (RS2133)

Initiation – haemophilia with inhibitors

Haematologist

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Continuation – haemophilia with inhibitors

Haematologist

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initiation – post-transplant

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Continuation – post-transplant

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initiation – indolent, low-grade lymphomas or hairy cell leukaemia*

Re-assessment required after 9 months

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Continuation – indolent, low-grade lymphomas or hairy cell leukaemia*

Re-assessment required after 12 months

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initiation – aggressive CD20 positive NHL

Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Continuation – aggressive CD20 positive NHL

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Initiation – Chronic lymphocytic leukaemia

Re-assessment required after 12 months

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Continuation – Chronic lymphocytic leukaemia

Re-assessment required after 12 months

Both:

- 1 Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1.1 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; or
- 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initiation – severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 8 weeks

All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 8 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initiation – warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 8 weeks

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Continuation – warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 8 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initiation – immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 8 weeks

All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 8 weeks

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initiation – thrombotic thrombocytopenic purpura (TTP)

Haematologist

Re-assessment required after 8 weeks

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

persistent thrombocytopenia despite plasma exchange; or

2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Continuation – thrombotic thrombocytopenic purpura (TTP)

Haematologist

Re-assessment required after 8 weeks

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initiation – pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Continuation – pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initiation – ANCA associated vasculitis

Re-assessment required after 8 weeks

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Continuation – ANCA associated vasculitis

Re-assessment required after 8 weeks

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Initiation – treatment refractory systemic lupus erythematosus (SLE)

Rheumatologist or nephrologist

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Continuation – treatment refractory systemic lupus erythematosus (SLE)

Rheumatologist or nephrologist

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initiation – Antibody-mediated organ transplant rejection

Patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initiation – ABO-incompatible organ transplant

Patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initiation – Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)

Nephrologist

Re-assessment required after 8 weeks

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with a * are unapproved indications.

Continuation – Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)

Nephrologist

Re-assessment required after 8 weeks

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for > 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with a * are unapproved indications.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Initiation – Steroid resistant nephrotic syndrome (SRNS)

Nephrologist

Re-assessment required after 8 weeks

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with a * are unapproved indications.

Continuation – Steroid resistant nephrotic syndrome (SRNS)

Nephrologist

Re-assessment required after 8 weeks

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with a * are unapproved indications.

Initiation – Neuromyelitis Optica Spectrum Disorder (NMOSD)

Re-assessment required after 6 months

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m² administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patient is receiving treatment with corticosteroids.

Continuation – Neuromyelitis Optica Spectrum Disorder (NMOSD)

Re-assessment required after 2 years

All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m² administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initiation – Severe Refractory Myasthenia Gravis

Neurologist

Re-assessment required after 2 years

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
- 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Continuation – Severe Refractory Myasthenia Gravis

Neurologist

Re-assessment required after 2 years

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initiation – Severe antisynthetase syndrome

Re-assessment required after 12 months

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not been effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000 mg infusions of rituximab.

Continuation – Severe antisynthetase syndrome

Re-assessment required after 12 months

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000 mg infusions of rituximab given two weeks apart.

Initiation – graft versus host disease

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not been effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

Initiation – severe chronic inflammatory demyelinating polyneuropathy

Neurologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Continuation – severe chronic inflammatory demyelinating polyneuropathy

Neurologist or medical practitioner on the recommendation of a Neurologist

Re-assessment required after 6 months

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initiation – anti-NMDA receptor autoimmune encephalitis

Neurologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Continuation – anti-NMDA receptor autoimmune encephalitis

Neurologist

Re-assessment required after 6 months

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

Initiation – CD20+ low grade or follicular B-cell NHL

Re-assessment required after 9 months

Either:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Continuation – CD20+ low grade or follicular B-cell NHL

Re-assessment required after 24 months

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m² every 8 weeks (maximum of 12 cycles).

Initiation – Membranous nephropathy

Re-assessment required after 6 weeks

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m²; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m² of body surface area per week for a total of 4 weeks.

Continuation – Membranous nephropathy

Re-assessment required after 6 weeks

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initiation – B-cell acute lymphoblastic leukaemia/lymphoma*

Limited to 2 years treatment

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initiation – desensitisation prior to transplant

Limited to 6 weeks treatment

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m² of body-surface area.

Note: Indications marked with * are unapproved indications.

Initiation – pemphigus*

Dermatologist or relevant specialist

Re-assessment required after 6 months

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Continuation – pemphigus*

Dermatologist or relevant specialist

Re-assessment required after 6 months

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

Initiation – immunoglobulin G4-related disease (IgG4-RD*)

Re-assessment required after 6 weeks

All of the following:

- 1 Patient has confirmed diagnosis of IgG4-RD*; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or
 - 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and
- 3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – immunoglobulin G4-related disease (IgG4-RD*)

Re-assessment required after 12 months

All of the following:

- 1 Either:
 - 1.1 Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
 - 1.2 Patient is receiving maintenance treatment for IgG4-RD*; and
- 2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and
- 3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB – **Restricted** see terms [below](#)

↓ Inj 150 mg per ml, 1 ml prefilled syringe.....	799.50	1	Cosentyx
	1,599.00	2	Cosentyx

➔ **Restricted (RS2119)**

Initiation – severe chronic plaque psoriasis, second-line biologic

Dermatologist

Re-assessment required after 4 months

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Continuation – severe chronic plaque psoriasis, second-line biologic

Dermatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initiation – severe chronic plaque psoriasis, first-line biologic

Dermatologist

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
 - 1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and

- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand, foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation – severe chronic plaque psoriasis, first-line biologic

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Either:
 - 1.1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
 - 1.2.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initiation – ankylosing spondylitis, second-line biologic

Rheumatologist

Re-assessment required after 3 months

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Continuation – ankylosing spondylitis, second-line biologic

Rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 300 mg monthly.

Initiation – psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation – psoriatic arthritis

Rheumatologist

Re-assessment required after 6 months

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB – **Restricted** see terms [below](#)

⚡ Inj 100 mg vial	770.57	1	Sylvant
⚡ Inj 400 mg vial	3,082.33	1	Sylvant

➔ **Restricted (RS1525)**

Initiation

Haematologist or rheumatologist

Re-assessment required after 6 months

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Continuation

Haematologist or rheumatologist

Re-assessment required after 12 months

The treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 20 mg per ml, 4 ml vial.....	220.00	1	Actemra
⚡ Inj 20 mg per ml, 10 ml vial.....	550.00	1	Actemra
⚡ Inj 20 mg per ml, 20 ml vial.....	1,100.00	1	Actemra

➡ **Restricted (RS2125)**

Initiation – cytokine release syndrome

Therapy limited to 3 doses

Either:

- 1 Both:
 - 1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
 - 2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-Cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initiation – previous use

Any relevant practitioner

Limited to 6 months treatment

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initiation – Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Limited to 6 months treatment

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
- 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initiation – Rheumatoid Arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initiation – systemic juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initiation – adult-onset Still's disease

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or

1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or

2 All of the following:

2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and

2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and

2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initiation – polyarticular juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 4 months

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and

1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or

2 All of the following:

2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and

2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and

2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

2.4 Any of the following:

2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or

2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or

2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initiation – idiopathic multicentric Castleman's disease

Haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist

Re-assessment required after 6 months

All of the following:

1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and

2 Treatment with an adequate trial of corticosteroids has proven ineffective; and

3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initiation – moderate to severe COVID-19

Therapy limited to 1 dose

All of the following:

1 Patient has confirmed (or probable) COVID-19; and

2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and

3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and

4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and

5 Tocilizumab is not to be administered in combination with baricitinib.

continued...

	Price	Brand or
(ex man. excl. GST)		Generic
\$	Per	Manufacturer

continued...

Continuation – Rheumatoid Arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Continuation – systemic juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Continuation – adult-onset Still's disease

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

the patient has a sustained improvement in inflammatory markers and functional status.

Continuation – polyarticular juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Continuation – idiopathic multicentric Castleman's disease

Haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist

Re-assessment required after 12 months

the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

Initiation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and
- 2 The individual has received insufficient benefit from use of corticosteroids; and
- 3 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Continuation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

Both:

- 1 The individual has shown clinical improvement and ongoing treatment is required; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Note: Indications marked with * are unapproved indications.

TRASTUZUMAB (HERZUMA) – **Restricted** see terms [below](#)

† Inj 150 mg vial – 5% DV Jun-24 to 31 May 2027	100.00	1	Herzuma
† Inj 440 mg vial – 5% DV Jun-24 to 31 May 2027	293.35	1	Herzuma

→ **Restricted** (RS2005)

Initiation – early breast cancer

Limited to 12 months treatment

Both:

- 1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Continuation – early breast cancer*

Re-assessment required after 12 months

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
 - 1.3 Any of the following:
 - 1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or
 - 1.3.3 the cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.4 Either:
 - 1.4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 1.4.2 All of the following:
 - 1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 1.4.2.3 The patient has good performance status (ECOG grade 0-1); and
 - 1.5 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

Initiation – metastatic breast cancer

Re-assessment required after 12 months

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and

3 Either:

- 3.1 Trastuzumab will not be given in combination with pertuzumab; or
- 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab to be discontinued at disease progression.

Continuation – metastatic breast cancer*Re-assessment required after 12 months*

Either:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
 - 1.3 Trastuzumab to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with trastuzumab.

Initiation – gastric, gastro-oesophageal junction and oesophageal cancer*Re-assessment required after 12 months*

Both:

- 1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and
- 2 Patient has an ECOG score of 0-2.

Continuation – gastric, gastro-oesophageal junction and oesophageal cancer*Re-assessment required after 12 months*

Both:

- 1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 2 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB DERUXTECAN – Restricted see terms [below](#)

↓ Inj 100 mg per ml, 1 ml vial.....2,550.00 1 Enhertu

➔ **Restricted (RS2082)****Initiation***Re-assessment required after 6 months*

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Patient has not received prior funded trastuzumab deruxtecan treatment; and
- 6 Treatment to be discontinued at disease progression.

Continuation

Re-assessment required after 6 months

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab deruxtecan; and
- 2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

TRASTUZUMAB EMTANSINE – **Restricted** see terms [below](#)

⚡ Inj 100 mg vial	2,320.00	1	Kadcyla
⚡ Inj 160 mg vial	3,712.00	1	Kadcyla

➡ **Restricted (RS2083)**

Initiation – early breast cancer

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or axillary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initiation – metastatic breast cancer

Re-assessment required after 6 months

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Either:
 - 6.1 Patient has not received prior funded trastuzumab emtansine or trastuzumab deruxtecan treatment; or
 - 6.2 Both:
 - 6.2.1 Patient has discontinued trastuzumab deruxtecan due to intolerance; and
 - 6.2.2 The cancer did not progress while on trastuzumab deruxtecan; and
- 7 Treatment to be discontinued at disease progression.

Continuation – metastatic breast cancer

Re-assessment required after 6 months

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB – **Restricted** see terms [below](#)

↓ Inj 130 mg vial	4,162.00	1	Stelara
↓ Inj 90 mg per ml, 1 ml prefilled syringe.....	4,162.00	1	Stelara

→ **Restricted (RS1942)**

Initiation – Crohn's disease - adults

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Continuation – Crohn's disease - adults

Re-assessment required after 12 months

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initiation – Crohn's disease - children*

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Continuation – Crohn's disease - children*

Re-assessment required after 12 months

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initiation – ulcerative colitis

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
- 2 Both:
 - 2.1 Patient has active ulcerative colitis; and
 - 2.2 Either:
 - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Continuation – ulcerative colitis

Re-assessment required after 12 months

Both:

- 1 Either:
 - 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
 - 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and
- 2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 300 mg vial3,313.00 1 Entyvio

➡ **Restricted (RS1943)**

Initiation – Crohn's disease - adults

Re-assessment required after 6 months

All of the following:

- 1 Patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Any of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
- 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
- 3.3 Immunomodulators and corticosteroids are contraindicated.

Continuation – Crohn's disease - adults

Re-assessment required after 2 years

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
 - 1.2 CDAI score is 150 or less, or HBI is 4 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initiation – Crohn's disease - children*

Re-assessment required after 6 months

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.3 Patient has extensive small intestine disease; and
- 3 Any of the following:
 - 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
 - 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
 - 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Continuation – Crohn's disease - children*

Re-assessment required after 2 years

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initiation – ulcerative colitis

Re-assessment required after 6 months

All of the following:

- 1 Patient has active ulcerative colitis; and
- 2 Any of the following:
 - 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
 - 2.2 Patient has a SCCAI score is greater than or equal to 4; or
 - 2.3 Patient's PUCAI score is greater than or equal to 20*; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

3 Any of the following:

- 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
- 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
- 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Continuation – ulcerative colitis

Re-assessment required after 2 years

Both:

1 Either:

- 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
- 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and
- 2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Programmed Cell Death-1 (PD-1) Inhibitors

ATEZOLIZUMAB – **Restricted** see terms [below](#)

‡ Inj 60 mg per ml, 20 ml vial.....9,503.00 1 Tecentriq

→ **Restricted (RS2099)**

Initiation – non-small cell lung cancer second line monotherapy

Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic non-small cell lung cancer; and
- 2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 4 Patient has an ECOG 0-2; and
- 5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
- 6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – non-small cell lung cancer second line monotherapy

Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with atezolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and
 - 2.2 Patient has preserved liver function (Child-Pugh A); and
 - 2.3 Transarterial chemoembolisation (TACE) is unsuitable; and
 - 2.4 Any of the following:
 - 2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or
 - 2.4.2 Patient received funded lenvatinib before 1 March 2025; or
 - 2.4.3 Both:
 - 2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and
 - 2.4.3.2 No disease progression since initiation of lenvatinib; and
 - 2.5 Patient has an ECOG performance status of 0-2; and
 - 2.6 To be given in combination with bevacizumab.

Continuation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

no evidence of disease progression.

DURVALUMAB – Restricted see terms [below](#)

↓ Inj 50 mg per ml, 10 ml vial.....	4,700.00	1	Imfinzi
↓ Inj 50 mg per ml, 2.4 ml vial.....	1,128.00	1	Imfinzi

➔ **Restricted (RS2084)**

Initiation – Non-small cell lung cancer

Re-assessment required after 4 months

All of the following:

- 1 Either:
 - 1.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); or
 - 1.2 Patient has histologically or cytologically documented stage IIb (T1N2a only), locally advanced, unresectable non-small cell lung cancer (NSCLC); and
- 2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
- 3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
- 4 Patient has a ECOG performance status of 0 or 1; and
- 5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 7 Either:
 - 7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 8 Treatment with durvalumab to cease upon signs of disease progression.

Continuation – Non-small cell lung cancer

Re-assessment required after 4 months

All of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

IPILIMUMAB – Restricted see terms [below](#)

⚡ Inj 5 mg per ml, 10 ml vial.....	5,000.00	1	Yervoy
⚡ Inj 5 mg per ml, 40 ml vial.....	20,000.00	1	Yervoy

➔ **Restricted (RS2115)**

Initiation – renal cell carcinoma

Limited to 4 months treatment

- Either:
- 1 The patient is currently on treatment with ipilimumab and met all remaining criteria prior to commencing treatment; or
 - 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and
 - 2.2 The patient is treatment naive; and
 - 2.3 The patient has ECOG performance status 0-2; and
 - 2.4 The disease is predominantly of clear cell histology; and
 - 2.5 Any of the following:
 - 2.5.1 The patient has sarcomatoid histology; or
 - 2.5.2 Haemoglobin levels less than the lower limit of normal; or
 - 2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or
 - 2.5.4 Neutrophils greater than the upper limit of normal; or
 - 2.5.5 Platelets greater than the upper limit of normal; or
 - 2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
 - 2.5.7 Karnofsky performance score of less than or equal to 70; and
 - 2.6 Ipilimumab is to be used at a maximum dose of 1 mg/kg for up to four cycles in combination with nivolumab.

NIVOLUMAB – Restricted see terms [below](#)

⚡ Inj 10 mg per ml, 4 ml vial.....	1,051.98	1	Opdivo
⚡ Inj 10 mg per ml, 10 ml vial.....	2,629.96	1	Opdivo

➔ **Restricted (RS2126)**

Initiation – unresectable or metastatic melanoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Limited to 4 months treatment

- All of the following:
- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 3 The individual has ECOG performance 0-2; and
 - 4 Either:
 - 4.1 The individual has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 The individual has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the individual was on pembrolizumab; and
 - 5 Any of the following:

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
- 5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
- 5.3 All of the following:
 - 5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
 - 5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
 - 5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-L1 inhibitor.

Continuation – unresectable or metastatic melanoma, less than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Both:
 - 1.1 Any of the following:
 - 1.1.1 The individual's disease has had a complete response to treatment; or
 - 1.1.2 The individual's disease has had a partial response to treatment; or
 - 1.1.3 The individual has stable disease; and
 - 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; or
- 2 All of the following:
 - 2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 The individual has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Continuation – unresectable or metastatic melanoma, more than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Both:

- 1 The individual has been on treatment for more than 24 months; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Any of the following:
 - 2.1.1.1 The individual's disease has had a complete response to treatment; or
 - 2.1.1.2 The individual's disease has had a partial response to treatment; or
 - 2.1.1.3 The individual has stable disease; and
 - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; or
 - 2.2 All of the following:
 - 2.2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2.2 The individual has signs of disease progression; and
 - 2.2.3 Disease has not progressed during previous treatment with nivolumab.

Initiation – renal cell carcinoma, first line

Limited to 4 months treatment

Either:

- 1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The patient has metastatic renal cell carcinoma; and

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.2 The patient is treatment naive; and
- 2.3 The patient has ECOG performance status 0-2; and
- 2.4 The disease is predominantly of clear cell histology; and
- 2.5 Any of the following:
 - 2.5.1 The patient has sarcomatoid histology; or
 - 2.5.2 Haemoglobin levels less than the lower limit of normal; or
 - 2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or
 - 2.5.4 Neutrophils greater than the upper limit of normal; or
 - 2.5.5 Platelets greater than the upper limit of normal; or
 - 2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
 - 2.5.7 Karnofsky performance score of less than or equal to 70; and
- 2.6 Nivolumab is to be used in combination with ipilimumab for the first four treatment cycles at a maximum dose of 3 mg/kg; and
- 2.7 Nivolumab is to be used at a maximum maintenance dose of 240 mg every 2 weeks (or equivalent).

Initiation – renal cell carcinoma, second line

Limited to 4 months treatment

All of the following:

- 1 Patient has metastatic renal-cell carcinoma; and
- 2 The disease is of predominant clear-cell histology; and
- 3 Patient has ECOG performance status 0-2; and
- 4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and
- 5 Patient has not previously received a funded immune checkpoint inhibitor; and
- 6 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

Continuation – renal cell carcinoma

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

PEMBROLIZUMAB – **Restricted** see terms [below](#)

⚡ Inj 25 mg per ml, 4 ml vial.....4,680.00 1 Keytruda

➡ **Restricted (RS2134)**

Initiation – stage III or IV resectable melanoma - neoadjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 The individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The individual has resectable stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note); and
 - 2.2 The individual has not received prior funded systemic treatment in the perioperative setting for their stage IIIB, IIIC, IIID or IV melanoma; and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 2.3 Treatment must be prior to complete surgical resection; and
- 2.4 Pembrolizumab must be administered as monotherapy; and
- 2.5 The individual has ECOG performance score 0-2; and
- 2.6 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

Continuation – stage III or IV resectable melanoma - neoadjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Any of the following:

- 1 Both:
 - 1.1 The individual has received neoadjuvant treatment with an immune checkpoint inhibitor; and
 - 1.2 The individual meets initiation criteria for pembrolizumab for stage III or IV resected melanoma – adjuvant; or
- 2 Both:
 - 2.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
 - 2.2 The individual meets continuation criteria for pembrolizumab for stage III or IV resected melanoma – adjuvant; or
- 3 All of the following:
 - 3.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
 - 3.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 3.3 The individual meets initiation criteria for pembrolizumab for unresectable or metastatic melanoma; or
- 4 All of the following:
 - 4.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
 - 4.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma; and
 - 4.3 The individual meets continuation criteria for pembrolizumab for unresectable or metastatic melanoma.

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
- b) Initiating treatment within 13 weeks of complete surgical resection means either 13 weeks after resection (primary or lymphadenectomy) or 13 weeks prior to the scheduled date of the resection (primary or lymphadenectomy)

Initiation – stage III or IV resected melanoma - adjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 The individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); and
 - 2.2 Adjuvant treatment with pembrolizumab is required; and
 - 2.3 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
 - 2.4 Treatment must be in addition to complete surgical resection; and
 - 2.5 Treatment must be initiated within 13 weeks of complete surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
 - 2.6 Pembrolizumab must be administered as monotherapy; and
 - 2.7 The individual has ECOG performance score 0-2; and
 - 2.8 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

Notes:

- a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

- b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Continuation – stage III or IV resected melanoma - adjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Any of the following:

- 1 All of the following:
 - 1.1 No evidence of disease recurrence; and
 - 1.2 Pembrolizumab must be administered as monotherapy; and
 - 1.3 Pembrolizumab to be administered at a fixed dose of 200 mg every three weeks (or equivalent) for a maximum of 12 months total treatment course, including any systemic neoadjuvant treatment; and
 - 1.4 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months total treatment course (equivalent to 18 cycles at a dose of 200 mg every 3 weeks), including any systemic neoadjuvant treatment; or
- 2 All of the following:
 - 2.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and
 - 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
 - 2.3 The individual meets initiation criteria for pembrolizumab for unresectable or metastatic melanoma; or
- 3 All of the following:
 - 3.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and
 - 3.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma; and
 - 3.3 The individual meets continuation criteria for pembrolizumab for unresectable or metastatic melanoma.

Initiation – unresectable or metastatic melanoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Limited to 4 months treatment

All of the following:

- 1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
- 3 The individual has ECOG performance 0-2; and
- 4 Either:
 - 4.1 The individual has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 The individual has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the individual was on nivolumab; and
- 5 Any of the following:
 - 5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
 - 5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
 - 5.3 All of the following:
 - 5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
 - 5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
 - 5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-L1 inhibitor.

Continuation – unresectable or metastatic melanoma, less than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

continued...

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

continued...

1 Both:

1.1 Any of the following:

- 1.1.1 The individual's disease has had a complete response to treatment; or
- 1.1.2 The individual's disease has had a partial response to treatment; or
- 1.1.3 The individual has stable disease; and

1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; or

2 All of the following:

- 2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
- 2.2 The individual has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Continuation – unresectable or metastatic melanoma, more than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Both:

1 The individual has been on treatment for more than 24 months; and

2 Either:

2.1 All of the following:

2.1.1 Any of the following:

- 2.1.1.1 The individual's disease has had a complete response to treatment; or
- 2.1.1.2 The individual's disease has had a partial response to treatment; or
- 2.1.1.3 The individual has stable disease; and

2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and

2.1.3 The treatment remains clinically appropriate and the individual is benefitting from the treatment; or

2.2 All of the following:

2.2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and

2.2.2 The individual has signs of disease progression; and

2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initiation – non-small cell lung cancer first-line monotherapy

Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 Patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used as monotherapy; and
- 6 Either:
 - 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
 - 6.2 Both:
 - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and

continued...

	Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
	Per	

continued...

6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and

- 7 Patient has an ECOG 0-2; and
- 8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – non-small cell lung cancer first-line monotherapy

Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – non-small cell lung cancer first-line combination therapy

Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
- 2 The patient has not had chemotherapy for their disease in the palliative setting; and
- 3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
- 4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR or ALK tyrosine kinase unless not possible to ascertain; and
- 5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
- 6 Patient has an ECOG 0-2; and
- 7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
- 8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – non-small cell lung cancer first-line combination therapy

Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
- 3 No evidence of disease progression; and
- 4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
- 5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and

continued...

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

continued...

- 6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – breast cancer, advanced

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 6 months

Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
 - 2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); and
 - 2.2 Patient is treated with palliative intent; and
 - 2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
 - 2.4 Patient has received no prior systemic therapy in the palliative setting; and
 - 2.5 Patient has an ECOG score of 0–2; and
 - 2.6 Pembrolizumab is to be used in combination with chemotherapy; and
 - 2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – breast cancer, advanced

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period; and
- 4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – head and neck squamous cell carcinoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies; and
 - 2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
 - 2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
 - 2.4 Patient has an ECOG performance score of 0-2; and
 - 2.5 Either:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or
- 2.5.2 Pembrolizumab to be used as monotherapy; and
- 2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – head and neck squamous cell carcinoma

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 Any of the following:
 - 1.1 Patient's disease has had a complete response to treatment; or
 - 1.2 Patient's disease has had a partial response to treatment; or
 - 1.3 Patient has stable disease; and
- 2 No evidence of disease progression; and
- 3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – MSI-H/dMMR advanced colorectal cancer

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer; or
 - 2.1.2 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer; and
 - 2.2 Individual is treated with palliative intent; and
 - 2.3 Individual has not previously received funded treatment with pembrolizumab for MSI-H/dMMR advanced colorectal cancer; and
 - 2.4 Individual has an ECOG performance score of 0-2; and
 - 2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
 - 2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – MSI-H/dMMR advanced colorectal cancer

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

- 1 No evidence of disease progression; and
- 2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
- 3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – Urothelial carcinoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

- 1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or

continued...

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

continued...

2 All of the following:

- 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
- 2.2 Patient has an ECOG performance score of 0-2; and
- 2.3 Patient has documented disease progression following treatment with chemotherapy; and
- 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – Urothelial carcinoma

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

1 Any of the following:

- 1.1 Patient's disease has had a complete response to treatment; or
- 1.2 Patient's disease has had a partial response to treatment; or
- 1.3 Patient has stable disease; and

2 No evidence of disease progression; and

3 Pembrolizumab is to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent); and

4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – relapsed/refractory Hodgkin lymphoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or

2 All of the following:

2.1 Either:

2.1.1 Both:

2.1.1.1 Individual has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and

2.1.1.2 Individual is ineligible for autologous stem cell transplant; or

2.1.2 Individual has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant; and

2.2 Individual has not previously received funded pembrolizumab for relapsed/refractory Hodgkin lymphoma; and

2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Continuation – relapsed/refractory Hodgkin lymphoma

Any relevant practitioner

Re-assessment required after 6 months

Both:

1 Patient has received a partial or complete response to pembrolizumab; and

2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Other Immunosuppressants

ANTITHYMOCYTE GLOBULIN (EQUINE)

Inj 50 mg per ml, 5 ml ampoule 4,439.17 5 ATGAM

ANTITHYMOCYTE GLOBULIN (RABBIT)

Inj 25 mg vial

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
AZATHIOPRINE			
Tab 25 mg – 5% DV Feb-26 to 2028	10.15	60	Azamun
Tab 50 mg – 5% DV Feb-26 to 2028	10.34	100	Azamun
Inj 50 mg vial			
Inj 100 mg vial			
BACILLUS CALMETTE-GUERIN (BCG) – Restricted see terms below			
¶ Inj 2-8 × 10 ⁸ CFU vial	149.37	1	OncoTICE
¶ Inj 40 mg per ml, vial	182.45	3	SII-Onco-BCG
➔ Restricted (RS1206)			
Initiation			
For use in bladder cancer.			
EVEROLIMUS – Restricted see terms below			
¶ Tab 5 mg	4,555.76	30	Afinitor
¶ Tab 10 mg	6,512.29	30	Afinitor
➔ Restricted (RS2076)			
Initiation			
Neurologist or oncologist			
<i>Re-assessment required after 3 months</i>			
Both:			
1 Patient has tuberous sclerosis; and			
2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.			
Continuation			
Neurologist or oncologist			
<i>Re-assessment required after 12 months</i>			
All of the following:			
1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and			
2 The treatment remains appropriate and the patient is benefiting from treatment; and			
3 Everolimus to be discontinued at progression of SEGAs.			
Initiation – renal cell carcinoma			
<i>Re-assessment required after 4 months</i>			
Either:			
1 All of the following:			
1.1 The patient has metastatic renal cell carcinoma; and			
1.2 The disease is of predominant clear-cell histology; and			
1.3 The patient has documented disease progression following one previous line of treatment; and			
1.4 The patient has an ECOG performance status of 0-2; and			
1.5 Everolimus is to be used in combination with lenvatinib; or			
2 All of the following:			
2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and			
2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and			
2.3 Everolimus is to be used in combination with lenvatinib; and			
2.4 There is no evidence of disease progression.			
Continuation – renal cell carcinoma			
<i>Re-assessment required after 4 months</i>			
there is no evidence of disease progression.			
MYCOPHENOLATE MOFETIL			
Tab 500 mg	35.90	50	CellCept
Cap 250 mg	35.90	100	CellCept
Powder for oral liq 1 g per 5 ml.....	187.25	165 ml	CellCept
Inj 500 mg vial	133.33	4	CellCept

¶ Item restricted (see ➔ above); ¶ Item restricted (see ➔ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
PICIBANIL			
Inj 100 mcg vial			
SIROLIMUS – Restricted see terms below			
↓ Tab 1 mg	749.99	100	Rapamune
↓ Tab 2 mg	1,499.99	100	Rapamune
↓ Oral liq 1 mg per ml	449.99	60 ml	Rapamune
→ Restricted (RS1991)			

Initiation

For rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR < 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencephalopathy; or
- Significant malignant disease

Initiation – severe non-malignant lymphovascular malformations*

Re-assessment required after 6 months

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Continuation – severe non-malignant lymphovascular malformations*

Re-assessment required after 12 months

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initiation – renal angiomyolipoma(s) associated with tuberous sclerosis complex*

Nephrologist or urologist

Re-assessment required after 6 months

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

continued...

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

Continuation – renal angiomyolipoma(s) associated with tuberous sclerosis complex*

Re-assessment required after 12 months

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initiation – refractory seizures associated with tuberous sclerosis complex*

Neurologist

Re-assessment required after 6 months

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex*; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialed and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, and topiramate. Those who can father children are not required to trial sodium valproate.

Continuation – refractory seizures associated with tuberous sclerosis complex*

Neurologist

Re-assessment required after 12 months

demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – **Restricted** see terms [below](#)

⚡ Tab modified-release 15 mg.....	1,271.00	28	Rinvoq
⚡ Tab modified-release 30 mg.....	2,033.00	28	Rinvoq
⚡ Tab modified-release 45 mg.....	3,049.00	28	Rinvoq

➡ **Restricted (RS2120)**

Initiation – Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)

Limited to 6 months treatment

All of the following:

continued...

	Price	Brand or
	(ex man. excl. GST)	Generic
	\$	Manufacturer
	Per	

continued...

- 1 The individual has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The individual has experienced intolerable side effects with adalimumab and/or etanercept; or
 - 2.2 The individual has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Any of the following:
 - 3.1 Rituximab is not clinically appropriate; or
 - 3.2 The individual is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.3 Both:
 - 3.3.1 The individual has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.3.2 Either:
 - 3.3.2.1 The individual has experienced intolerable side effects with rituximab; or
 - 3.3.2.2 At four months following the initial course of rituximab the individual has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Continuation – Rheumatoid Arthritis

Re-assessment required after 6 months

Either:

- 1 Following 6 months' initial treatment, the individual has experienced at least a 50% decrease in active joint count from baseline; or
- 2 On subsequent reapplications, the individual has experienced at least a continuing 30% improvement in active joint count from baseline.

Initiation – Atopic dermatitis

Re-assessment required after 6 months

Either:

- 1 Individual is currently on treatment with upadacitinib for atopic dermatitis and met all remaining criteria prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Individual has moderate to severe atopic dermatitis, severity as defined by an Eczema Area and Severity Index (EASI) score of greater than or equal to 16 or a Dermatology Life Quality Index (DLQI) score of greater than or equal to 10; and
 - 2.2 Individual has received insufficient benefit from topical therapy (including topical corticosteroids or topical calcineurin inhibitors) for a 28-day trial within the last 6 months, unless contraindicated to all; and
 - 2.3 Individual has trialled and received insufficient benefit from at least one systemic therapy for a minimum of three months (eg ciclosporin, azathioprine, methotrexate or mycophenolate mofetil), unless contraindicated to all; and
 - 2.4 An EASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.5 The most recent EASI or DQLI assessment is no more than 1 month old at the time of application.

Continuation – Atopic dermatitis

Re-assessment required after 12 months

Either:

- 1 Individual has received a 75% or greater reduction in EASI score (EASI 75) as compared to baseline EASI prior to commencing upadacitinib; or
- 2 Individual has received a DLQI improvement of 4 or more as compared to baseline DLQI prior to commencing upadacitinib.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Initiation – Crohn's disease – adult

Re-assessment required after 6 months

Either:

- 1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing treatment; or
- 2 Both:
 - 2.1 Individual has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Individual has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Continuation – Crohn's disease – adult

Re-assessment required after 2 years

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score when the individual was initiated on biologic therapy; or
- 2 HBI score has reduced by 3 points from when individual was initiated on biologic therapy; or
- 3 CDAI score is 150 or less; or
- 4 HBI score is 4 or less; or
- 5 The individual has experienced an adequate response to treatment, but CDAI score cannot be assessed.

Initiation – Crohn's disease – children

Re-assessment required after 6 months

Either:

- 1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing treatment; or
- 2 Both:
 - 2.1 Child has active Crohn's disease; and
 - 2.2 Either:
 - 2.2.1 Child has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Child meets the initiation criteria for prior biologic therapies for Crohn's disease; and
 - 2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Continuation – Crohn's disease – children

Re-assessment required after 2 years

Any of the following:

- 1 PCDAI score has reduced by 10 points from when the child was initiated on treatment; or
- 2 PCDAI score is 15 or less; or
- 3 The child has experienced an adequate response to treatment, but PCDAI score cannot be assessed.

Note: Indications marked with * are unapproved indications.

Initiation – Ulcerative colitis

Re-assessment required after 6 months

Either:

- 1 Individual is currently on treatment with upadacitinib for ulcerative colitis and met all remaining criteria prior to commencing treatment; or
- 2 Both:

continued...

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

- 2.1 Individual has active ulcerative colitis; and
- 2.2 Either:
 - 2.2.1 Individual has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
 - 2.2.2 Both:
 - 2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
 - 2.2.2.2 Other biologic therapies for ulcerative colitis are contraindicated.

Continuation – Ulcerative colitis

Re-assessment required after 2 years

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the individual was initiated on treatment; or
- 2 PUCAI score has reduced by 10 points or more from the PUCAI score when the individual was initiated on treatment.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Antiallergy Preparations

Allergic Emergencies

ADRENALINE – **Restricted** see terms [below](#)

⚡ Inj 0.15 mg per 0.3 ml auto-injector – 5% DV Dec-25 to 2028.....	85.50	1	Epipen Jr
⚡ Inj 0.3 mg per 0.3 ml auto-injector – 5% DV Dec-25 to 2028.....	85.50	1	Epipen

➡ **Restricted (RS1944)**

Initiation – anaphylaxis

Either:

- 1 Patient has experienced a previous anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or
- 2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner.

ICATIBANT – **Restricted** see terms [below](#)

⚡ Inj 10 mg per ml, 3 ml prefilled syringe	2,668.00	1	Firazyr
--	----------	---	---------

➡ **Restricted (RS1501)**

Initiation

Clinical immunologist or relevant specialist

Re-assessment required after 12 months

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Continuation

Re-assessment required after 12 months

The treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

BEE VENOM – **Restricted** see terms [below](#)

⚡ Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			
⚡ Initiation kit - 1 vial freeze dried venom with diluent	305.00	1	VENOX
⚡ Maintenance Kit - 1 vial freeze dried venom with diluent	305.00	1	VENOX

➡ **Restricted (RS1117)**

Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

PAPER WASP VENOM – **Restricted** see terms [below](#)

⚡ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			

➡ **Restricted (RS1118)**

Initiation

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

YELLOW JACKET WASP VENOM – **Restricted** see terms [on the next page](#)

⚡ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent			
⚡ Inj 550 mcg vial with diluent			

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

➔ **Restricted (RS1119)****Initiation**

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Allergy Prophylactics**BUDESONIDE**Nasal spray 50 mcg per dose – **5% DV Feb-25 to 2027** 2.59 200 dose **SteroClear**Nasal spray 100 mcg per dose – **5% DV Feb-25 to 2027** 2.89 200 dose **SteroClear****FLUTICASONE PROPIONATE**Metered dose nasal spray 50 mcg per dose – **5% DV Feb-26 to 2028** 2.57 120 dose **Flixonase Hayfever & Allergy****IPRATROPIUM BROMIDE**

Aqueous nasal spray 0.03% 5.23 15 ml Univent

SODIUM CROMOGLICATE

Nasal spray 4%

Antihistamines**CETIRIZINE HYDROCHLORIDE**Tab 10 mg – **5% DV Sep-23 to 2026** 1.71 100 **Zista**

Oral liq 1 mg per ml 3.99 200 ml Histaclear

CHLORPHENIRAMINE MALEATE

Oral liq 0.4 mg per ml

Inj 10 mg per ml, 1 ml ampoule

CYPROHEPTADINE HYDROCHLORIDE

Tab 4 mg

FEXOFENADINE HYDROCHLORIDE

Tab 60 mg

Tab 120 mg – **5% DV Jul-25 to 2027** 3.49 30 **Fexaclear**Tab 180 mg – **5% DV Jul-25 to 2027** 4.10 30 **Fexaclear****LORATADINE**

Tab 10 mg 1.78 100 Lorafix

Oral liq 1 mg per ml 1.43 100 ml Haylor Syrup

PROMETHAZINE HYDROCHLORIDETab 10 mg – **5% DV Dec-25 to 2028** 2.19 100 **Allersoothe**Tab 25 mg – **5% DV Dec-25 to 2028** 2.69 100 **Allersoothe**

Oral liq 1 mg per ml 3.39 100 ml Allersoothe

Inj 25 mg per ml, 2 ml ampoule 21.09 5 Hospira

Anticholinergic Agents**IPRATROPIUM BROMIDE**

Aerosol inhaler 20 mcg per dose

Nebuliser soln 250 mcg per ml, 1 ml ampoule

Nebuliser soln 250 mcg per ml, 2 ml ampoule 11.73 20 Accord
Univent

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Anticholinergic Agents with Beta-Adrenoceptor Agonists

SALBUTAMOL WITH IPRATROPIUM BROMIDE

Aerosol inhaler 100 mcg with ipratropium bromide 20 mcg per dose

Nebuliser soln 2.5 mg with ipratropium bromide 0.5 mg per 2.5 ml

ampoule.....	11.04	20	Duolin
--------------	-------	----	--------

Long-Acting Muscarinic Agents

GLYCOPYRRONIUM

Note: inhaled glycopyrronium treatment must not be used if the patient is also receiving treatment with subsidised tiotropium or umeclidinium.

Powder for inhalation 50 mcg per dose	61.00	30 dose	Seebri Breezhaler
---	-------	---------	-------------------

TIOTROPIUM BROMIDE

Note: tiotropium treatment must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.

Soln for inhalation 2.5 mcg per dose	50.37	60 dose	Spiriva Respimat
--	-------	---------	------------------

Powder for inhalation 18 mcg per dose	50.37	30 dose	Spiriva
---	-------	---------	---------

UMECLIDINIUM

Note: Umeclidinium must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.

Powder for inhalation 62.5 mcg per dose	61.50	30 dose	Incruse Ellipta
---	-------	---------	-----------------

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

➡ Restricted (RS1518)

Initiation

Re-assessment required after 2 years

- Both:
- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
 - 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Continuation

Re-assessment required after 2 years

- Both:
- 1 Patient is compliant with the medication; and
 - 2 Patient has experienced improved COPD symptom control (prescriber determined).

Note: Combination long acting muscarinic antagonist and long acting beta-2 agonist must not be used if the patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

GLYCOPYRRONIUM WITH INDACATEROL – **Restricted** see terms [above](#)

† Powder for Inhalation 50 mcg with indacaterol 110 mcg	81.00	30 dose	Ultibro Breezhaler
---	-------	---------	--------------------

TIOTROPIUM BROMIDE WITH OLODATEROL – **Restricted** see terms [above](#)

† Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg	81.00	60 dose	Spiolto Respimat
---	-------	---------	------------------

UMECLIDINIUM WITH VILANTEROL – **Restricted** see terms [above](#)

† Powder for inhalation 62.5 mcg with vilanterol 25 mcg	77.00	30 dose	Anoro Ellipta
---	-------	---------	---------------

Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

BUDESONIDE WITH GLYCOPYRRONIUM AND EFORMOTEROL – **Restricted** see terms [on the next page](#)

† Aerosol inhaler budesonide 160 mcg with glycopyrronium 7.2 mcg and formoterol 5 mcg per dose.....	79.15	120 dose	Breztri Aerosphere
---	-------	----------	--------------------

	Price		Brand or
	(ex man. excl. GST)		Generic
	\$	Per	Manufacturer

➔ **Restricted (RS2085)****Initiation**

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

 - 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
 - 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
 - 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
 - 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
 - 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long-acting muscarinic antagonist and long-acting beta-2 agonist – ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler therapy.

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – **Restricted** see terms [below](#)

↓ Powder for inhalation fluticasone furoate 100 mcg with umeclidinium			
62.5 mcg and vilanterol 25 mcg.....	104.24	30 dose	Trelegy Ellipta

➔ **Restricted (RS2028)****Initiation**

Both:

- 1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and
 - 2.1.2 Any of the following:

Clinical criteria:

 - 2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or
 - 2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or
 - 2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or
 - 2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3×10^9 cells/L in the previous 12 months; or
 - 2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist – ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

AntifibroticsNINTEDANIB – **Restricted** see terms [below](#)

↓ Cap 100 mg.....	2,554.00	60	Ofev
↓ Cap 150 mg.....	3,870.00	60	Ofev

➔ **Restricted (RS1813)****Initiation – idiopathic pulmonary fibrosis**

Respiratory specialist

Re-assessment required after 12 months

All of the following:

continued...

Products with Hospital Supply Status (HSS) are in **bold**

Expiry date of HSS period is 30 June of the year indicated unless otherwise stated.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and

2 Forced vital capacity is between 50% and 90% predicted; and

3 Nintedanib is to be discontinued at disease progression (See Note); and

4 Nintedanib is not to be used in combination with subsidised pirfenidone; and

5 Any of the following:

5.1 The patient has not previously received treatment with pirfenidone; or

5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or

5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Continuation – idiopathic pulmonary fibrosis

Respiratory specialist
Re-assessment required after 12 months
All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and

2 Nintedanib is not to be used in combination with subsidised pirfenidone; and

3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – **Restricted** see terms [below](#)

⚡ Tab 267 mg	1,215.00	90	Esbriet
⚡ Tab 801 mg	3,645.00	90	Esbriet

➡ **Restricted (RS1814)**

Initiation – idiopathic pulmonary fibrosis

Respiratory specialist
Re-assessment required after 12 months
All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and

2 Forced vital capacity is between 50% and 90% predicted; and

3 Pirfenidone is to be discontinued at disease progression (See Notes); and

4 Pirfenidone is not to be used in combination with subsidised nintedanib; and

5 Any of the following:

5.1 The patient has not previously received treatment with nintedanib; or

5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or

5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Continuation – idiopathic pulmonary fibrosis

Respiratory specialist
Re-assessment required after 12 months
All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and

2 Pirfenidone is not to be used in combination with subsidised nintedanib; and

3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Oral liq 400 mcg per ml – 5% DV May-25 to 2027	50.00	150 ml	Ventolin
Inj 500 mcg per ml, 1 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			
Aerosol inhaler, 100 mcg per dose	4.18	200 dose	SalAir
	6.80		Ventolin
Nebuliser soln 1 mg per ml, 2.5 ml ampoule	8.96	20	Asthalin
			UK Cipla
Nebuliser soln 2 mg per ml, 2.5 ml ampoule	9.43	20	Asthalin
			UK Cipla
TERBUTALINE SULPHATE			
Powder for inhalation 250 mcg per dose			
Inj 0.5 mg per ml, 1 ml ampoule			
Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated	22.20	120 dose	Bricanyl Turbuhaler
Decongestants			
OXYMETAZOLINE HYDROCHLORIDE			
Aqueous nasal spray 0.25 mg per ml			
Aqueous nasal spray 0.5 mg per ml			
PSEUDOEPHEDRINE HYDROCHLORIDE			
Tab 60 mg			
SODIUM CHLORIDE			
Aqueous nasal spray isotonic			
SODIUM CHLORIDE WITH SODIUM BICARBONATE			
Soln for nasal irrigation			
XYLOMETAZOLINE HYDROCHLORIDE			
Aqueous nasal spray 0.05%			
Aqueous nasal spray 0.1%			
Nasal drops 0.05%			
Nasal drops 0.1%			
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler 50 mcg per dose	8.54	200 dose	Beclazone 50
	14.01		Qvar
Aerosol inhaler 100 mcg per dose	12.50	200 dose	Beclazone 100
	17.52		Qvar
Aerosol inhaler 250 mcg per dose	22.67	200 dose	Beclazone 250
BUDESONIDE			
Nebuliser soln 250 mcg per ml, 2 ml ampoule			
Nebuliser soln 500 mcg per ml, 2 ml ampoule			
Powder for inhalation 100 mcg per dose			
Powder for inhalation 200 mcg per dose			
Powder for inhalation 400 mcg per dose			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUTICASONE			
Aerosol inhaler 50 mcg per dose.....	7.19	120 dose	Flixotide
Powder for inhalation 50 mcg per dose.....	8.61	60 dose	Flixotide Accuhaler
Powder for inhalation 100 mcg per dose.....	7.81	60 dose	Flixotide Accuhaler
Aerosol inhaler 125 mcg per dose.....	13.60	120 dose	Flixotide
Aerosol inhaler 250 mcg per dose.....	24.62	120 dose	Flixotide
Powder for inhalation 250 mcg per dose.....	11.93	60 dose	Flixotide Accuhaler

Leukotriene Receptor Antagonists

MONTELUKAST			
Tab 4 mg – 5% DV Dec-25 to 2028.....	3.10	28	Montelukast Viatris
Tab 5 mg – 5% DV Dec-25 to 2028.....	3.10	28	Montelukast Viatris
Tab 10 mg – 5% DV Dec-25 to 2028.....	2.45	28	Montelukast Viatris

Long-Acting Beta-Adrenoceptor Agonists

EFORMOTEROL FUMARATE			
Powder for inhalation 12 mcg per dose			
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated (equivalent to eformoterol fumarate 6 mcg metered dose)			
INDACATEROL			
Powder for inhalation 150 mcg per dose.....	61.00	30 dose	Onbrez Breezhaler
Powder for inhalation 300 mcg per dose.....	61.00	30 dose	Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler 25 mcg per dose.....	26.25	120 dose	Serevent
Powder for inhalation 50 mcg per dose.....	26.25	60 dose	Serevent Accuhaler

Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

BUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg			
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg			
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide with 6 mcg eformoterol fumarate metered dose).....	41.50	120 dose	DuoResp Spiromax
Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg.....	33.74	120 dose	Symbicort Turbuhaler
Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mcg eformoterol fumarate metered dose).....	82.50	120 dose	DuoResp Spiromax
Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg.....	33.74	60 dose	Symbicort Turbuhaler
FLUTICASONE FURATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg.....	44.08	30 dose	Breo Ellipta
FLUTICASONE WITH SALMETEROL			
Aerosol inhaler 50 mcg with salmeterol 25 mcg.....	25.79	120 dose	Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg.....	33.74	60 dose	Seretide Accuhaler
Aerosol inhaler 125 mcg with salmeterol 25 mcg.....	32.60	120 dose	Seretide
Powder for inhalation 250 mcg with salmeterol 50 mcg.....	44.08	60 dose	Seretide Accuhaler

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Methylxanthines			
AMINOPHYLLINE			
Inj 25 mg per ml, 10 ml ampoule	180.00	5	DBL Aminophylline
CAFFEINE CITRATE			
Oral liq 20 mg per ml (caffeine 10 mg per ml)	16.91	25 ml	Biomed
Inj 20 mg per ml (caffeine 10 mg per ml), 2.5 ml ampoule	69.70	5	Biomed
THEOPHYLLINE			
Tab long-acting 250 mg.....	25.65	100	Nuelin-SR
Oral liq 80 mg per 15 ml.....	18.49	500 ml	Nuelin

Mucolytics and Expectorants

DORNASE ALFA – **Restricted** see terms [below](#)

↓ Nebuliser soln 2.5 mg per 2.5 ml ampoule250.00 6 Pulmozyme

→ **Restricted (RS1787)**

Initiation – cystic fibrosis

Respiratory physician or paediatrician

Re-assessment required after 12 months

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Continuation – cystic fibrosis

Respiratory physician or paediatrician

The treatment remains appropriate and the patient continues to benefit from treatment.

Initiation – significant mucus production

Limited to 4 weeks treatment

Both:

- 1 Patient is an in-patient; and
- 2 The mucus production cannot be cleared by first line chest techniques.

Initiation – pleural emphyema

Limited to 3 days treatment

Both:

- 1 Patient is an in-patient; and
- 2 Patient diagnoses with pleural emphyema.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR – **Restricted** see terms [on the next page](#)

↓ Tab elexacafator 50 mg with tezacafator 25 mg, ivacafator 37.5 mg (56) and
ivacafator 75 mg (28).....27,647.39 84 Trikafta

↓ Tab elexacafator 100 mg with tezacafator 50 mg, ivacafator 75 mg (56) and
ivacafator 150 mg (28).....27,647.39 84 Trikafta

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2114)

Initiation

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Patient is 6 years of age or older; and
- 3 Either:
 - 3.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
 - 3.2 Patient has a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Either:
 - 4.1 Patient has a heterozygous or homozygous F508del mutation; or
 - 4.2 Patient has a G551D mutation or other mutation responsive in vitro to elexacaftor/tezacaftor/ivacaftor (see note a); and
- 5 The treatment must be the sole funded CFTR modulator therapy for this condition; and
- 6 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Notes:

- a) Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information
<https://nctr-crs.fda.gov/fdalabel/services/spl/set-ids/f354423a-85c2-41c3-a9db-0f3aee135d8d/spl-doc>

IVACAFTOR – Restricted see terms below

⚡ Tab 150 mg	29,386.00	56	Kalydeco
⚡ Oral granules 50 mg, sachet	29,386.00	56	Kalydeco
⚡ Oral granules 75 mg, sachet	29,386.00	56	Kalydeco

➔ Restricted (RS1818)

Initiation

Respiratory specialist or paediatrician

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

Nebuliser soln 7%, 90 ml bottle	25.73	90 ml	Biomed
---------------------------------------	-------	-------	--------

Pulmonary Surfactants

BERACTANT

Soln 200 mg per 8 ml vial

PORACTANT ALFA

Soln 120 mg per 1.5 ml vial	425.00	1	Curosurf
Soln 240 mg per 3 ml vial	695.00	1	Curosurf

⚡ Item restricted (see ➔ above); ⚡ Item restricted (see ➔ below)

e.g. Brand indicates brand example only. It is not a contracted product.

Price	Brand or
(ex man. excl. GST)	Generic
\$	Manufacturer
Per	

Respiratory Stimulants

DOXAPRAM

Inj 20 mg per ml, 5 ml vial

Sclerosing Agents

TALC

Powder

Soln (slurry) 100 mg per ml, 50 ml

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Anti-Infective Preparations			
Antibacterials			
CHLORAMPHENICOL			
Eye oint 1% – 5% DV Feb-26 to 2028	1.55	5 g	Devatis
Ear drops 0.5%			
Eye drops 0.5%	1.45	10 ml	Chlorsig
Eye drops 0.5%, single dose			
CIPROFLOXACIN			
Eye drops 0.3% – 5% DV Mar-25 to 2027	10.85	5 ml	Ciprofloxacin Teva
FRAMYCETIN SULPHATE			
Ear/eye drops 0.5%			
GENTAMICIN SULPHATE			
Eye drops 0.3%			
PROPAMIDINE ISETHIONATE			
Eye drops 0.1%			
SODIUM FUSIDATE [FUSIDIC ACID]			
Eye drops 1%	5.29	5 g	Fucithalmic
SULPHACETAMIDE SODIUM			
Eye drops 10%			
TOBRAMYCIN			
Eye oint 0.3%	10.45	3.5 g	Tobrex
Eye drops 0.3%	11.48	5 ml	Tobrex
Antifungals			
NATAMYCIN			
Eye drops 5%			
Antivirals			
ACICLOVIR			
Eye oint 3% – 5% DV Feb-25 to 2027	15.89	4.5 g	ViruPOS
Combination Preparations			
CIPROFLOXACIN WITH HYDROCORTISONE			
Ear drops ciprofloxacin 0.2% with 1% hydrocortisone.....	16.30	10 ml	Ciproxin HC Otic
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml			
DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE			
Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	5.39	3.5 g	Maxitrol
Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	4.50	5 ml	Maxitrol
DEXAMETHASONE WITH TOBRAMYCIN			
Eye drops 0.1% with tobramycin 0.3%	12.64	5 ml	Tobradex

↑ Item restricted (see ➡ above); ↓ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUMETASONE PIVALATE WITH CLIQUINOL Ear drops 0.02% with cliquinol 1%			
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml	Kenacomb

Anti-Inflammatory Preparations

Corticosteroids

DEXAMETHASONE			
Eye oint 0.1%	5.86	3.5 g	Maxidex
Eye drops 0.1%	4.50	5 ml	Maxidex
↓ Ocular implant 700 mcg.....	1,444.50	1	Ozurdex

→ Restricted (RS1606)

Initiation – Diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

All of the following:

- 1 Patients have diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision; and
- 3 Either:
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Continuation – Diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initiation – Women of child bearing age with diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

All of the following:

- 1 Patients have diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Continuation – Women of child bearing age with diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
FLUOROMETHOLONE			
Eye drops 0.1%	3.09	5 ml	FML
PREDNISOLONE ACETATE			
Eye drops 0.12%			
Eye drops 1%	7.00	5 ml	Pred Forte
	6.92	10 ml	Prednisolone- AFT
PREDNISOLONE SODIUM PHOSPHATE			
Eye drops 0.5%, single dose (preservative free).....	43.26	20 dose	Minims Prednisolone

Non-Steroidal Anti-Inflammatory Drugs

DICLOFENAC SODIUM			
Eye drops 0.1%			
Eye drops 0.1%, single dose – 5% DV Jul-25 to 2027	1.85	10 dose	Diclofenac Devatis
	5.54	30 dose	Diclofenac Devatis
KETOROLAC TROMETAMOL			
Eye drops 0.5%			

Decongestants and Antiallergics

Antiallergic Preparations

LEVOCABASTINE			
Eye drops 0.05%			
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml	Lomide
OLOPATADINE			
Eye drops 0.1%	2.17	5 ml	Olopatadine Teva
SODIUM CROMOGLICATE			
Eye drops 2%	2.62	10 ml	Allerfix

Decongestants

NAPHAZOLINE HYDROCHLORIDE			
Eye drops 0.1% – 5% DV Jan-25 to 2027	5.65	15 ml	Albalon

Diagnostic and Surgical Preparations

Diagnostic Dyes

FLUORESCEIN SODIUM			
Eye drops 2%, single dose			
Inj 10%, 5 ml vial	125.00	12	Fluorescite
Ophthalmic strips 1 mg			
FLUORESCEIN SODIUM WITH LIGNOCAINE HYDROCHLORIDE			
Eye drops 0.25% with lignocaine hydrochloride 4%, single dose			
LISSAMINE GREEN			
Ophthalmic strips 1.5 mg			
ROSE BENGAL SODIUM			
Ophthalmic strips 1%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Irrigation Solutions			
MIXED SALT SOLUTION FOR EYE IRRIGATION			
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 15 ml dropper bottle	5.00	15 ml	Balanced Salt Solution
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 250 ml			<i>e.g. Balanced Salt Solution</i>
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml bag			<i>e.g. Balanced Salt Solution</i>
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml bottle.....	10.50	500 ml	Balanced Salt Solution
Ocular Anaesthetics			
OXYBUPROCAINE HYDROCHLORIDE			
Eye drops 0.4%, single dose			
PROXYMETACAINE HYDROCHLORIDE			
Eye drops 0.5%			
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Eye drops 0.5%, single dose			
Eye drops 1%, single dose			
Viscoelastic Substances			
HYPROMELLOSE			
Inj 2%, 1 ml syringe			
Inj 2%, 2 ml syringe			
SODIUM HYALURONATE [HYALURONIC ACID]			
Inj 14 mg per ml, 0.85 ml syringe	50.00	1	Healon GV
Inj 18 mg per ml, 0.85 ml syringe	50.00	1	Healon GV Pro
Inj 23 mg per ml, 0.6 ml syringe	60.00	1	Healon 5
Inj 10 mg per ml, 0.85 ml syringe	28.50	1	Healon
SODIUM HYALURONATE [HYALURONIC ACID] WITH CHONDROITIN SULPHATE			
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.35 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.4 ml syringe	64.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.5 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.55 ml syringe	74.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.75 ml syringe.....	67.00	1	Viscoat
Other			
DISODIUM EDETATE			
Inj 150 mg per ml, 20 ml ampoule			
Inj 150 mg per ml, 20 ml vial			
Inj 150 mg per ml, 100 ml vial			

SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

RIBOFLAVIN 5-PHOSPHATE
Soln trans epithelial riboflavin
Inj 0.1%
Inj 0.1% plus 20% dextran T500
Inj 0.1% plus hydroxypropyl methylcellulose

Glaucoma Preparations

Beta Blockers

BETAXOLOL			
Eye drops 0.25%	11.80	5 ml	Betoptic S
Eye drops 0.5%	7.50	5 ml	Betoptic
<i>(Betoptic S Eye drops 0.25% to be delisted 1 December 2025)</i>			
<i>(Betoptic Eye drops 0.5% to be delisted 1 December 2025)</i>			
TIMOLOL			
Eye drops 0.25% – 5% DV Mar-24 to 2026	2.42	5 ml	Arrow-Timolol
Eye drops 0.5% – 5% DV Mar-24 to 2026	2.50	5 ml	Arrow-Timolol
➡ Eye drops 0.5%, gel forming – Restricted: For continuation only			

Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE			
Tab 250 mg – 5% DV Sep-25 to 2027	13.96	100	Medsurge
Inj 500 mg			
BRINZOLAMIDE			
Eye drops 1% – 5% DV Dec-24 to 2027	5.11	5 ml	Azopt
DORZOLAMIDE – Restricted: For continuation only			
➡ Eye drops 2%			
DORZOLAMIDE WITH TIMOLOL			
Eye drops 2% with timolol 0.5% – 5% DV Feb-25 to 2027	3.58	5 ml	Dortimopt

Miotics

ACETYLCHOLINE CHLORIDE			
Inj 20 mg vial with diluent			
CARBACHOL			
Inj 150 mcg vial			
PILOCARPINE HYDROCHLORIDE			
Eye drops 1%	4.26	15 ml	Isopto Carpine
Eye drops 2%	5.35	15 ml	Isopto Carpine
Eye drops 4%	7.99	15 ml	Isopto Carpine
PILOCARPINE NITRATE			
Eye drops 2%, single dose			

Prostaglandin Analogues

BIMATOPROST			
Eye drops 0.03% – 5% DV Jan-25 to 2027	5.15	3 ml	Lumigan
LATANOPROST			
Eye drops 0.005% – 5% DV Mar-25 to 2027	2.08	2.5 ml	Teva

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LATANOPROST WITH TIMOLOL			
Eye drops 0.005% with timolol 0.5% – 5% DV Mar-24 to 2026	4.95	2.5 ml	Arrow - Lattim
TRAVOPROST			
Eye drops 0.004% – 5% DV Dec-24 to 2027	6.80	2.5 ml	Travatan

Sympathomimetics

APRACLOPIDINE			
Eye drops 0.5%	19.77	5 ml	Iopidine
BRIMONIDINE TARTRATE			
Eye drops 0.2% – 5% DV Mar-25 to 2027	5.16	5 ml	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE			
Eye drops 0.2% with timolol 0.5% – 5% DV Dec-24 to 2027	7.13	5 ml	Combigan

Mydriatics and Cycloplegics

Anticholinergic Agents

ATROPINE SULPHATE			
Eye drops 0.5%			
Eye drops 1%, single dose			
Eye drops 1% – 5% DV Feb-24 to 2026	18.27	15 ml	Atropt
CYCLOPENTOLATE HYDROCHLORIDE			
Eye drops 0.5%, single dose			
Eye drops 1%	25.16	15 ml	Cyclogyl
Eye drops 1%, single dose			
TROPICAMIDE			
Eye drops 0.5%	20.52	15 ml	Mydriacyl
Eye drops 0.5%, single dose			
Eye drops 1%	24.82	15 ml	Mydriacyl
Eye drops 1%, single dose			

Sympathomimetics

PHENYLEPHRINE HYDROCHLORIDE			
Eye drops 2.5%, single dose			
Eye drops 10%, single dose			

Ocular Lubricants

CARBOMER			
Ophthalmic gel 0.2%			
CARMELLOSE SODIUM WITH PECTIN AND GELATINE			
Eye drops 0.5%			
Eye drops 0.5%, single dose			
Eye drops 1%			
Eye drops 1%, single dose			
HYPROMELLOSE			
Eye drops 0.5%	19.50	15 ml	Methopt
Ophthalmic gel 0.3%			

SENSORY ORGANS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
HYPROMELLOSE WITH DEXTRAN			
Eye drops 0.3% with dextran 0.1%.....	2.30	15 ml	Poly-Tears
Eye drops 0.3% with dextran 0.1%, single dose			
PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN			
Eye oint 42.5% with soft white paraffin 57.3%			
PARAFFIN LIQUID WITH WOOL FAT			
Eye oint 3% with wool fat 3%	3.63	3.5 g	Poly-Visc
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL			
Eye drops 0.4% with propylene glycol 0.3%, 10 ml bottle			
Note: Only for use in compounding an eye drop formulation			
Eye drops 0.4% with propylene glycol 0.3% preservative free, single dose....	10.78	30	Systane Unit Dose
POLYVINYL ALCOHOL WITH POVIDONE			
Eye drops 1.4% with povidone 0.6%, single dose			
RETINOL PALMITATE			
Oint 138 mcg per g	3.80	5 g	VitA-POS
SODIUM HYALURONATE [HYALURONIC ACID]			
Eye drops 1 mg per ml – 5% DV Dec-24 to 2027	13.58	10 ml	Hylo-Fresh

Other Otological Preparations

ACETIC ACID WITH PROPYLENE GLYCOL			
Ear drops 2.3% with propylene glycol 2.8%			
DOCUSATE SODIUM			
Ear drops 0.5%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Agents Used in the Treatment of Poisonings			
Antidotes			
ACETYLCYSTEINE			
Tab eff 200 mg			
Inj 200 mg per ml, 10 ml vial.....	42.99	10	Hikma Acetylcysteine
Inj 200 mg per ml, 10 ml ampoule – 5% DV Apr-25 to 2027	42.99	10	DBL Acetylcysteine
	52.88		Martindale Pharma
<i>(Martindale Pharma Inj 200 mg per ml, 10 ml ampoule to be delisted 1 November 2025)</i>			
AMYL NITRITE			
Liq 98% in 3 ml capsule			
DIGOXIN IMMUNE FAB			
Inj 38 mg vial			
Inj 40 mg vial			
ETHANOL			
Liq 96%			
ETHANOL WITH GLUCOSE			
Inj 10% with glucose 5%, 500 ml bottle			
ETHANOL, DEHYDRATED			
Inj 100%, 5 ml ampoule			
Inj 96%			
FLUMAZENIL			
Inj 0.1 mg per ml, 5 ml ampoule – 5% DV Dec-24 to 2027	44.00	5	Flumazenil-Baxter
HYDROXOCOBALAMIN			
Inj 5 g vial			
Inj 2.5 g vial			
NALOXONE HYDROCHLORIDE			
Inj 400 mcg per ml, 1 ml ampoule – 5% DV Apr-25 to 2027	13.29	5	DBL Naloxone Hydrochloride
PRALIDOXIME CHLORIDE			
Inj 1 g vial			
PRALIDOXIME IODIDE			
Inj 25 mg per ml, 20 ml ampoule			
SODIUM NITRITE			
Inj 30 mg per ml, 10 ml ampoule			
SODIUM THIOSULFATE			
Inj 250 mg per ml, 100 ml vial			
Inj 250 mg per ml, 10 ml vial			
Inj 250 mg per ml. 50 ml vial			
Inj 500 mg per ml, 10 ml vial			
Inj 500 mg per ml, 20 ml ampoule			
SOYA OIL			
Inj 20%, 500 ml bag			
Inj 20%, 500 ml bottle			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Antitoxins

BOTULISM ANTITOXIN

Inj 250 ml vial

DIPHThERIA ANTITOXIN

Inj 10,000 iu vial

Antivenoms

RED BACK SPIDER ANTIVENOM

Inj 500 u vial

SNAKE ANTIVENOM

Inj 50 ml vial

Removal and Elimination

CHARCOAL

Oral liq 200 mg per ml 59.85 250 ml Carbasorb-X

DEFERASIROX – **Restricted** see terms [below](#)

⚡ Tab 125 mg dispersible 276.00 28 Exjade

⚡ Tab 250 mg dispersible 552.00 28 Exjade

⚡ Tab 500 mg dispersible 1,105.00 28 Exjade

➡ **Restricted (RS1444)**

Initiation

Haematologist

Re-assessment required after 2 years

All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per µL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per µL).

Continuation

Haematologist

Re-assessment required after 2 years

Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels. .

DEFERIPRONE – **Restricted** see terms [below](#)

⚡ Tab 500 mg 533.17 100 Ferriprox

⚡ Oral liq 100 mg per ml 266.59 250 ml Ferriprox

➡ **Restricted (RS1445)**

Initiation

Patient has been diagnosed with chronic iron overload due to congenital inherited anaemia or acquired red cell aplasia.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
DEFERRIOXAMINE MESILATE			
Inj 500 mg vial	332.88	10	DBL Desferrioxamine Mesylate for Inj BP
DICOBALT EDETATE			
Inj 15 mg per ml, 20 ml ampoule			
DIMERCAPROL			
Inj 50 mg per ml, 2 ml ampoule			
DIMERCAPTOSUCCINIC ACID			
Cap 100 mg			e.g. PCNZ, Optimus Healthcare, Chemet
Cap 200 mg			e.g. PCNZ, Optimus Healthcare, Chemet
SODIUM CALCIUM EDETATE			
Inj 50 mg per ml, 10 ml ampoule			
Inj 200 mg per ml, 2.5 ml ampoule			
Inj 200 mg per ml, 5 ml ampoule			

Antiseptics and Disinfectants

CHLORHEXIDINE			
Soln 0.1%			
Soln 4%			
Soln 5%	15.50	500 ml	healthE
CHLORHEXIDINE WITH CETRIMIDE			
Crm 0.1% with cetrimide 0.5%			
Foaming soln 0.5% with cetrimide 0.5%			
CHLORHEXIDINE WITH ETHANOL			
Soln 0.5% with ethanol 70%			
Soln 2% with ethanol 70%			
Soln 0.5% with ethanol 70%, non-staining (pink) 25 ml	1.55	1	healthE
IODINE WITH ETHANOL			
Soln 1% with ethanol 70%			
ISOPROPYL ALCOHOL			
Soln 70%, 500 ml	5.65	1	healthE
POVIDONE-IODINE			
↓ Vaginal tab 200 mg			
➔ Restricted (RS1354)			
Initiation			
Rectal administration pre-prostate biopsy.			
Oint 10%	7.40	65 g	Betadine
Soln 10%	4.99	100 ml	Riodine
Soln 5%			
Soln 7.5%			
Soln 10%,	3.83	15 ml	Riodine
	6.99	500 ml	Riodine
Pad 10%			
Swab set 10%			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
POVIDONE-IODINE WITH ETHANOL			
Soln 10% with ethanol 30%			
Soln 10% with ethanol 70%			
SODIUM HYPOCHLORITE			
Soln			

Contrast Media

Iodinated X-ray Contrast Media

DIATRIZOATE MEGLUMINE WITH SODIUM AMIDOTRIZOATE			
Oral liq 660 mg per ml with sodium amidotrizoate 100 mg per ml, 100 ml bottle.....	30.00	100 ml	Gastrografin
Inj 260 mg with sodium amidotrizoate 40 mg per ml, 250 ml bottle.....	120.00	1	Urografin
DIATRIZOATE SODIUM			
Oral liq 370 mg per ml, 10 ml sachet.....	156.12	50	Ioscan
IODISED OIL			
Inj 38% w/w (480 mg per ml), 10 ml ampoule	410.00	1	Lipiodol Ultra Fluid
IODIXANOL			
Inj 270 mg per ml (iodine equivalent), 50 ml bottle.....	275.00	10	Visipaque
Inj 270 mg per ml (iodine equivalent), 100 ml bottle.....	505.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 50 ml bottle.....	280.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 100 ml bottle.....	510.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 200 ml bottle.....	1,020.00	10	Visipaque
IOHEXOL			
Inj 240 mg per ml (iodine equivalent), 50 ml bottle.....	117.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 20 ml bottle.....	110.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 50 ml bottle.....	121.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 100 ml bottle.....	200.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 50 ml bottle.....	125.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 100 ml bottle.....	210.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 200 ml bottle.....	420.00	10	Omnipaque
Inj 350 mg per ml, 500 ml bottle	655.00	6	Omnipaque

Non-iodinated X-ray Contrast Media

BARIUM SULPHATE			
Oral liq 400 mg per ml (40% w/v, 30% w/w), bottle	17.39	148 g	Varibar - Thin Liquid
Oral liq 400 mg per ml (40% w/v), bottle	189.15	250 ml	Varibar - Honey
	38.40	240 ml	Varibar - Nectar
	159.05	230 ml	Varibar - Pudding
Grans for oral liq 960 mg per g (96% w/w), 176 g bottle	530.00	24	Vanilla SilQ MD
Grans for oral liq 980 mg per g (98% w/w), 310 g bottle	490.00	24	Vanilla SilQ HD
Oral liq 20.9 mg per ml (2.1% w/v, 2% w/w), 450 ml bottle	97.50	12	Readi-CAT 2
Oral liq 1 mg per ml (0.1% w/v, 0.1% w/w), 450 ml bottle	15.95	1	Neulumex
	191.40	12	Neulumex
Oral liq 400 mg per ml (40% w/v, 30% w/w), 20 ml bottle	52.35	3	Tagitol V
CITRIC ACID WITH SODIUM BICARBONATE			
Powder 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet.....	90.25	50 g	E-Z-Gas II

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Paramagnetic Contrast Media

GADOBUTROL			
Inj 1 mmol per ml, 15 ml vial			
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 5 ml prefilled syringe.....	126.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 7.5 ml prefilled syringe.....	189.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 15 ml prefilled syringe.....	735.00	10	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 65 ml bottle.....	3,120.00	10	Gadovist 1.0
GADOTERIC ACID			
Inj 279.30 mg per ml, 10 ml prefilled syringe			<i>e.g. Clariscan</i>
Inj 279.30 mg per ml, 10 ml vial			<i>e.g. Clariscan</i>
Inj 279.30 mg per ml, 15 ml prefilled syringe			<i>e.g. Clariscan</i>
Inj 279.30 mg per ml, 20 ml vial			<i>e.g. Clariscan</i>
Inj 279.30 mg per ml, 5 ml vial			<i>e.g. Clariscan</i>
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml prefilled syringe.....	172.00	10	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml bottle	25.35	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml prefilled syringe.....	258.00	10	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml prefilled syringe.....	344.00	10	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml bottle	14.30	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml bottle	28.90	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 5 ml bottle	9.10	1	Dotarem
GADOXETATE DISODIUM			
Inj 181.43 mg per ml (equivalent to 0.25 mmol per ml), 10 ml prefilled syringe.....	300.00	1	Primovist
MEGLUMINE GADOPENTETATE			
Inj 469 mg per ml, 10 ml prefilled syringe.....	95.00	5	Magnevist
Inj 469 mg per ml, 10 ml vial.....	185.00	10	Magnevist
MEGLUMINE IOTROXATE			
Inj 105 mg per ml, 100 ml bottle	169.15	100 ml	Biliscopin

Ultrasound Contrast Media

PERFLUTREN			
Inj 1.1 mg per ml, 1.5 ml vial.....	180.00	1	Definity
	720.00	4	Definity

Diagnostic Agents

ARGININE			
Inj 50 mg per ml, 500 ml bottle			
Inj 100 mg per ml, 300 ml bottle			
HISTAMINE ACID PHOSPHATE			
Nebuliser soln 0.6%, 10 ml vial			
Nebuliser soln 2.5%, 10 ml vial			
Nebuliser soln 5%, 10 ml vial			
MANNITOL			
Powder for inhalation			<i>e.g. Aridol</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
METHACHOLINE CHLORIDE Powder 100 mg			
SECRETIN PENTAHYDROCHLORIDE Inj 100 u vial Inj 80 u vial Inj 100 u ampoule			
SINCALIDE Inj 5 mcg per vial			

Diagnostic Dyes

BONNEY'S BLUE DYE Soln			
INDIGO CARMINE Inj 4 mg per ml, 5 ml ampoule Inj 8 mg per ml, 5 ml ampoule			
INDOCYANINE GREEN Inj 25 mg vial			
METHYLTHIONINIUM CHLORIDE [METHYLENE BLUE] Inj 5 mg per ml, 10 ml ampoule	259.57	5	Proveblue
PATENT BLUE V Inj 2.5%, 2 ml ampoule	440.00	5	Obex Medical
Inj 2.5%, 5 ml prefilled syringe.....	420.00	5	InterPharma

Irrigation Solutions

CHLORHEXIDINE WITH CETRIMIDE
 ‡ Irrigation soln 0.015% with cetrimide 0.15%, 500 ml bottle

➔ **Restricted (RS1683)**

Initiation

Re-assessment required after 3 months

All of the following:

- 1 Patient has burns that are greater than 30% of total body surface area (BSA); and
- 2 For use in the perioperative preparation and cleansing of large burn areas requiring debridement/skin grafting; and
- 3 The use of 30 ml ampoules is impractical due to the size of the area to be covered.

Continuation

Re-assessment required after 3 months

The treatment remains appropriate for the patient and the patient is benefiting from the treatment.

Irrigation soln 0.015% with cetrimide 0.15%, 100 ml bottle

Irrigation soln 0.015% with cetrimide 0.15%, 30 ml ampoule – **5% DV**

Sep-25 to 202829.70 30 **LumaCina**

GLYCINE

Irrigation soln 1.5%, 3,000 ml bag96.28 4 B Braun

SODIUM CHLORIDE

Irrigation soln 0.9%, 3,000 ml bag80.00 4 B Braun

Irrigation soln 0.9%, 30 ml ampoule12.50 20 InterPharma

Irrigation soln 0.9%, 1,000 ml bottle19.50 10 Baxter Sodium Chloride
0.9%

Irrigation soln 0.9%, 250 ml bottle21.60 12 Fresenius Kabi

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
WATER			
Irrigation soln, 3,000 ml bag	84.52	4	B Braun
Irrigation soln, 1,000 ml bottle	19.50	10	Baxter Water for Irrigation
Irrigation soln, 250 ml bottle	21.60	12	Fresenius Kabi

Surgical Preparations

BISMUTH SUBNITRATE AND IODOFORM PARAFFIN

Paste

DIMETHYL SULFOXIDE

Soln 50%

Soln 99%

PHENOL

Inj 6%, 10 ml ampoule

PHENOL WITH IOXAGLIC ACID

Inj 12%, 10 ml ampoule

SODIUM HYDROXIDE

Soln 10%

TROMETAMOL

Inj 36 mg per ml, 500 ml bottle

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Cardioplegia Solutions			
ELECTROLYTES			
Inj 15 mmol/l sodium chloride, 9 mmol/l potassium chloride, 1 mmol/l potassium hydrogen 2-ketoglutarate, 4 mmol/l magnesium chloride, 18 mmol/l histidine hydrochloride, 180 mmol/l histidine, 2 mmol/l tryptophan, 30 mmol/l mannitol, 0.015 mmol/l calcium chloride, 1,000 ml bag			<i>e.g. Custodiol-HTK</i>
Inj aspartic acid 10.43 mg per ml, citric acid 0.22476 mg per ml, glutamic acid 11.53 mg per ml, sodium phosphate 0.1725 mg per ml, potassium chloride 2.15211 mg per ml, sodium citrate 1.80768 mg per ml, sodium hydroxide 6.31 mg per ml and trometamol 11.2369 mg per ml, 364 ml bag			<i>e.g. Cardioplegia Enriched Paed. Soln.</i>
Inj aspartic acid 8.481 mg per ml, citric acid 0.8188 mg per ml, glutamic acid 9.375 mg per ml, sodium phosphate 0.6285 mg per ml, potassium chloride 2.5 mg per ml, sodium citrate 6.585 mg per ml, sodium hydroxide 5.133 mg per ml and trometamol 9.097 mg per ml, 527 ml bag			<i>e.g. Cardioplegia Enriched Solution</i>
Inj citric acid 0.07973 mg per ml, sodium phosphate 0.06119 mg per ml, potassium chloride 2.181 mg per ml, sodium chloride 1.788 mg ml, sodium citrate 0.6412 mg per ml and trometamol 5.9 mg per ml, 523 ml bag			<i>e.g. Cardioplegia Base Solution</i>
Inj 110 mmol/l sodium, 16 mmol/l potassium, 1.2 mmol/l calcium, 16 mmol/l magnesium and 160 mmol/l chloride, 1,000 ml bag			<i>e.g. Cardioplegia Solution AHB7832</i>
Inj 143 mmol/l sodium, 16 mmol/l potassium, 16 mmol/l magnesium and 1.2 mmol/l calcium, 1,000 ml bag			<i>e.g. Cardioplegia Electrolyte Solution</i>
MONOSODIUM GLUTAMATE WITH SODIUM ASPARTATE			
Inj 42.68 mg with sodium aspartate 39.48 mg per ml, 250 ml bottle			
MONOSODIUM L-ASPARTATE			
Inj 14 mmol per 10 ml, 10 ml			

Cold Storage Solutions

SODIUM WITH POTASSIUM

Inj 29 mmol/l with potassium 125 mmol/l, 1,000 ml bag

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Extemporaneously Compounded Preparations			
ACETIC ACID Liq			
ALUM Powder BP			
ARACHIS OIL [PEANUT OIL] Liq			
ASCORBIC ACID Powder			
BENZOIN Tincture compound BP			
BISMUTH SUBGALLATE Powder			
BORIC ACID Powder			
CARBOXYMETHYLCELLULOSE Soln 1.5%			
CETRIMIDE Soln 40%			
CHLORHEXIDINE GLUCONATE Soln 20 %			
CHLOROFORM Liq BP			
CITRIC ACID Powder BP			
CLOVE OIL Liq			
COAL TAR Soln BP.....	36.25	200 ml	Midwest
CODEINE PHOSPHATE Powder			
COLLODION FLEXIBLE Liq			
COMPOUND HYDROXYBENZOATE Soln	30.00	100 ml	Midwest
CYSTEAMINE HYDROCHLORIDE Powder			
DISODIUM HYDROGEN PHOSPHATE WITH SODIUM DIHYDROGEN PHOSPHATE Inj 37.46 mg with sodium dihydrogen phosphate 47.7 mg in 1.5 ml ampoule			
DITHRANOL Powder			
GLUCOSE [DEXTROSE] Powder			

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
GLYCERIN WITH SODIUM SACCHARIN			
Suspension.....	30.95	473 ml	Ora-Sweet SF
GLYCERIN WITH SUCROSE			
Suspension.....	30.95	473 ml	Ora-Sweet
GLYCEROL			
Liq.....	3.23	500 ml	healthE Glycerol BP Liquid
HYDROCORTISONE			
Powder	49.95	25 g	ABM
LACTOSE			
Powder			
MAGNESIUM HYDROXIDE			
Paste			
MENTHOL			
Crystals			
METHADONE HYDROCHLORIDE			
Powder			
METHYL HYDROXYBENZOATE			
Powder	8.98	25 g	Midwest
METHYLCELLULOSE			
Powder	36.95	100 g	Midwest
Suspension.....	30.95	473 ml	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN			
Suspension.....	30.95	473 ml	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE			
Suspension.....	30.95	473 ml	Ora-Blend
OLIVE OIL			
Liq			
PARAFFIN			
Liq			
PHENOBARBITONE SODIUM			
Powder			
PHENOL			
Liq			
PILOCARPINE NITRATE			
Powder			
POLYHEXAMETHYLENE BIGUANIDE			
Liq			
POVIDONE K30			
Powder			
SALICYLIC ACID			
Powder			
SILVER NITRATE			
Crystals			
SODIUM BICARBONATE			
Powder BP	10.05	500 g	Midwest

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SODIUM CITRATE Powder			
SODIUM METABISULFITE Powder			
STARCH Powder			
SULPHUR Precipitated Sublimed			
SYRUP Liq (pharmaceutical grade).....	14.95	500 ml	Midwest
THEOBROMA OIL Oint			
TRI-SODIUM CITRATE Crystals			
TRICHLORACETIC ACID Grans			
UREA Powder BP			
WOOL FAT Oint, anhydrous			
XANTHAN Gum 1%			
ZINC OXIDE Powder			

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

Food Modules

Carbohydrate

➡ **Restricted (RS1467)**

Initiation – Use as an additive

Any of the following:

- 1 Cystic fibrosis; or
- 2 Chronic kidney disease; or
- 3 Cancer in children; or
- 4 Cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 5 Faltering growth in an infant/child; or
- 6 Bronchopulmonary dysplasia; or
- 7 Premature and post premature infant; or
- 8 Inborn errors of metabolism.

Initiation – Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.
Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

CARBOHYDRATE SUPPLEMENT – **Restricted** see terms [above](#)

† Powder 96 g carbohydrate per 100 g, can	6.72	400 g	Polycal
---	------	-------	---------

Fat

➡ **Restricted (RS1468)**

Initiation – Use as an additive

Any of the following:

- 1 Patient has inborn errors of metabolism; or
- 2 Faltering growth in an infant/child; or
- 3 Bronchopulmonary dysplasia; or
- 4 Fat malabsorption; or
- 5 Lymphangiectasia; or
- 6 Short bowel syndrome; or
- 7 Infants with necrotising enterocolitis; or
- 8 Biliary atresia; or
- 9 For use in a ketogenic diet; or
- 10 Chyle leak; or
- 11 Ascites; or
- 12 Patient has increased energy requirements, and for whom dietary measures have not been successful.

Initiation – Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk. .
Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

LONG-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms [above](#)

† Liquid 50 g fat per 100 ml, bottle	15.38	200 ml	Calogen (neutral)
	38.44	500 ml	Calogen (neutral)
	15.38	200 ml	Calogen (strawberry)

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
MEDIUM-CHAIN TRIGLYCERIDE SUPPLEMENT – Restricted see terms on the previous page			
† Liquid 95 g fat per 100 ml, bottle	37.50	500 ml	MCT Oil
† Liquid 50 g fat per 100 ml, 250 ml bottle	143.65	4	Liquigen
WALNUT OIL – Restricted see terms on the previous page			
† Liq			

Protein

→ **Restricted (RS1469)**

Initiation – Use as an additive

Either:

- 1 Protein losing enteropathy; or
- 2 High protein needs.

Initiation – Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

PROTEIN SUPPLEMENT – Restricted see terms [above](#)

† Powder 5 g protein, 0.67 g carbohydrate and 0.6 g fat per 6.6 g, 275 g can			
† Powder 6 g protein per 7 g, can	8.95	227 g	Resource Beneprotein
† Powder 89 g protein, less than 1.5 g carbohydrate and 2 g fat per 100 g, can.....	13.82	225 g	Protifar

Other Supplements

CARBOHYDRATE AND FAT SUPPLEMENT – Restricted see terms [below](#)

† Powder 72.7 g carbohydrate and 22.3 g fat per 100 g, can	71.77	400 g	Duocal Super Soluble Powder
--	-------	-------	-----------------------------

→ **Restricted (RS1212)**

Initiation

Both:

- 1 Infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 Cystic fibrosis; or
 - 2.2 Cancer in children; or
 - 2.3 Faltering growth; or
 - 2.4 Bronchopulmonary dysplasia; or
 - 2.5 Premature and post premature infants.

HUMAN MILK FORTIFIER

Powder 0.325 g protein, 0.37 g carbohydrate and 0.175 g fat per 1 g sachet	33.48	50	Human Milk Fortifier <i>e.g. FM 85</i>
Powder 0.2 g protein, 0.7 g carbohydrate and 0.02 g fat per 1 g sachet			

Food/Fluid Thickeners

NOTE:

continued...

Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
------------------------------------	-----	-------------------------------------

continued...

While pre-thickened drinks and supplements have not been included in Section H, Health NZ Hospitals may continue to use such products for patients with dysphagia, provided that:

- use was established prior to 1 July 2013; and
- the product has not been specifically considered and excluded by Pharmac; and
- use of the product conforms to any applicable indication restrictions for similar products that are listed in Section H (for example, use of thickened high protein products should be in line with the restriction for high protein oral feed in Section H).

Pharmac intends to make a further decision in relation to pre-thickened drinks and supplements in the future, and will notify of any change to this situation.

CAROB BEAN GUM WITH MAIZE STARCH AND MALTODEXTRIN		
Powder	24.00	380 g Aptamil Feed Thickener
GUAR GUM		
Powder		e.g. Guarcol
MAIZE STARCH		
Powder	8.29	300 g Nutilis
MALTODEXTRIN WITH XANTHAN GUM		
Powder		e.g. Instant Thick
MALTODEXTRIN WITH XANTHAN GUM AND ASCORBIC ACID		
Powder		e.g. Easy Thick

Metabolic Products

➡ Restricted (RS2047)

Initiation

Either:

- 1 For the dietary management of inherited metabolic disease; or
- 2 Patient has adrenoleukodystrophy.

Supplements for Glutaric Aciduria Type 1

AMINO ACID FORMULA (WITHOUT LYSINE AND LOW TRYPTOPHAN) – Restricted see terms above		
⚡ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can		e.g. GA1 Anamix Infant
⚡ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can		e.g. XLYS Low TRY Maxamaid
AMINO ACID FORMULA (WITHOUT LYSINE) – Restricted see terms above		
⚡ Powder (neutral) 5 g protein, 5.4 g carbohydrate, 2.3 g fat and 2 g fibre per 18 g sachet.....	750.30	30 GA1 Anamix Junior
⚡ Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30 GA Explore 5
⚡ Powder, 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 3.7 g fibre per 100 g, 400 g can.....	260.00	400 g GA1 Anamix Infant

Price (ex man. excl. GST)	Brand or Generic Manufacturer
\$ Per	

Supplements for Homocystinuria

AMINO ACID FORMULA (WITHOUT METHIONINE) – **Restricted** see terms [on the previous page](#)

† Powder (neutral), 10 g protein, 11.5 g carbohydrate and 4.5 g fat per 36 g sachet.....	750.30	30	HCU Anamix Junior
† Powder, 15 g protein, 3.5 g carbohydrate, 0.55 g fat per 25 g sachet.....	1,048.95	30	HCU Express 15
† Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30	HCU Explore 5
† Powder (neutral) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	480.42	500 g	XMET Maxamum
† Powder (unflavoured) 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	260.00	400 g	HCU Anamix Infant
† Liquid (juicy berries), 20 g protein, 9.3 g carbohydrate, 0.44 g fat and 0.44 g fibre per 125 ml bottle.....	1,684.80	30	HCU Lophlex LQ
† Liquid (orange), 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle.....	941.40	36	HCU Anamix Junior LQ

Supplements for MSUD and Short chain enoyl coA hydratase deficiency

AMINO ACID FORMULA (WITHOUT Isoleucine, Leucine and Valine) – **Restricted** see terms [on the previous page](#)

† Powder (neutral) 10 g protein, 11.5 g carbohydrate and 4.5 g fat per 36 g sachet.....	750.00	30	MSUD Anamix Junior
† Powder, 15 g protein, 3.5 g carbohydrate, 0.6 g fat per 25 g sachet.....	1,048.95	30	MSUD Express 15
† Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30	MSUD Explore 5
† Powder (orange) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	454.71	500 g	MSUD Maxamum
† Powder (unflavoured) 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	260.00	400 g	MSUD Anamix Infant
† Powder (unflavoured) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	454.71	500 g	MSUD Maxamum
† Liquid (juicy berries), 20 g protein, 8.8 g carbohydrate, 0.44 g fat and 0.5 g fibre per 125 ml pouch.....	1,684.80	30	MSUD Lophlex LQ 20
† Liquid (orange) 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle.....	941.40	36	MSUD Anamix Junior LQ

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Supplements for Phenylketonuria			
AMINO ACID FORMULA (WITHOUT PHENYLALANINE) – Restricted see terms on page 294			
† Tab 8.33 mg	99.00	75	Phlexy 10
† Powder (Berry), 5.0 g protein, 14 g carbohydrate, 0 g fat per 20 g sachet.....	449.28	60	PKU Restore Powder
† Powder (Lemon), 20 g protein, 3.9 g carbohydrate, 0.8 g fat per 34 g sachet.....	883.50	30	PKU Express 20
† Powder (Neutral), 20 g protein, 4.8 g carbohydrate, 0.8 g fat per 34 g sachet.....	883.50	30	PKU Express 20
† Powder (Neutral), 5.0 g protein, 5.2 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	220.88	30	PKU Explore 5
† Powder (Orange), 10 g protein, 9.8 g carbohydrate, 0.4 g fat per 25 g sachet.....	441.75	30	PKU Explore 10
† Powder (Orange), 20 g protein, 3.9 g carbohydrate, 0.8 g fat per 34 g sachet.....	883.50	30	PKU Express 20
† Powder (Orange), 5.0 g protein, 14 g carbohydrate, 0 g fat per 20 g sachet.....	449.28	60	PKU Restore Powder
† Powder (Raspberry), 10 g protein, 9.8 g carbohydrate, 0.4 g fat per 25 g sachet.....	441.75	30	PKU Explore 10
† Powder (Tropical), 20 g protein, 3.9 g carbohydrate, 0.8 g fat per 34 g sachet.....	883.50	30	PKU Express 20
† Powder (berry) 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet.....	936.00	30	PKU Lophlex Powder
† Powder (chocolate) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	393.00	30	PKU Anamix Junior
† Powder (neutral) 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet.....	936.00	30	PKU Lophlex Powder
† Powder (neutral) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	393.00	30	PKU Anamix Junior
† Powder (orange) 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet.....	936.00	30	PKU Lophlex Powder
† Powder (orange) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	393.00	30	PKU Anamix Junior
† Powder (unflavoured), 5 g protein, 4.8 g carbohydrate per 12.5 g sachets	234.00	30	PKU First Spoon
† Powder (vanilla) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	393.00	30	PKU Anamix Junior
† Powder (Neutral), 14.3 g protein, 25 g fat per 100 g, 4 × 400 g can.....	715.16	1,600 g	PKU Start
† Powder (orange) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	320.00	500 g	XP Maxamum
† Powder (unflavoured) 39 g protein and 34 g carbohydrate per 100 g, 500 g can.....	320.00	500 g	XP Maxamum
† Powder 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	174.72	400 g	PKU Anamix Infant
† Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle	13.10	1	PKU Anamix Junior LQ (Berry) PKU Anamix Junior LQ (Orange)
† Liquid (juicy berries) 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 62.5 ml bottle	939.00	60	PKU Lophlex LQ 10
† Liquid (juicy berries) 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle	936.00	30	PKU Lophlex LQ 20

† Item restricted (see ➡ above); ‡ Item restricted (see ➡ below)

e.g. *Brand* indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
† Liquid (juicy orange) 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle	936.00	30	PKU Lophlex LQ 20
† Liquid (juicy tropical) 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 125 ml bottle	936.00	30	PKU Lophlex LQ 20
† Liquid 6.7 g protein, 5.1 g carbohydrate and 2 g fat per 100 ml, 250 ml carton.....	540.00	18	Easiphen Liquid
† Semi-solid 18.3 g protein, 18.5 g carbohydrate and 0.92 g fibre per 100 g, 109 g pot.....	1,123.20	36	PKU Lophlex Sensations 20 (berries)
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME PHENYLALANINE – Restricted see terms on page 294			
† Powder (Neutral), 10 g protein, 0.5 g carbohydrate, 0.6 g fat per 15 g sachet.....	449.28	30	PKU Build 10
† Powder (neutral), 15 g protein, 15 g carbohydrate, 4.5 g fat per 40 g sachet.....	673.92	30	Glytactin Bettermilk
† Powder (unflavoured) 10 g protein, 4 g carbohydrate per 12.5 g sachet	468.00	30	PKU GMPPro Mix-In
† Powder 20 g protein, 1.7 g carbohydrate per 31 g sachet.....	898.56	30	PKU Build 20 Raspberry Lemonade
† Powder 20 g protein, 1.7 g carbohydrate per 32 g sachet.....	898.56	30	PKU Build 20 Smooth
† Powder 20 g protein, 1.7 g carbohydrate per 33 g sachet.....	898.56	30	PKU Build 20 Chocolate
† Powder 20 g protein, 4.9 g carbohydrate per 33.4 g sachet.....	936.00	30	PKU Build 20 Vanilla
† Powder 20 g protein, 6.0 g carbohydrate per 35 g sachet.....	930.00	30	PKU GMPPro Ultra
† Powder 20 g protein, 6.3 g carbohydrate per 35 g sachet.....	930.00	30	PKU GMPPro Ultra Lemonade
† Powder 20 g protein, 6.7 g carbohydrate per 35 g sachet.....	930.00	30	PKU GMPPro Ultra Vanilla
† Powder 20 g protein, 6.7 g carbohydrate per 35 g sachet.....	930.00	30	PKU sphere20 Lemon
† Powder 20 g protein, 6.3 g carbohydrate per 35 g sachet.....	930.00	30	PKU sphere20 Chocolate
† Powder 20 g protein, 6.7 g carbohydrate per 35 g sachet.....	930.00	30	PKU sphere20 Red Berry
† Powder 20 g protein, 6.7 g carbohydrate per 35 g sachet.....	930.00	30	PKU sphere20 Vanilla
† Liquid (Coffee Mocha), 15 g protein, 3.1 g carbohydrate, 4.6 g fat 250 ml, carton.....	684.45	30	PKU sphere20 Banana
† Liquid (chocolate), 15 g protein, 22 g carbohydrate, 5.3 g fat per 250 ml, carton.....	684.45	30	PKU Glytactin RTD 15 Lite
† Liquid (neutral), 10 g protein, 8.5 g carbohydrate per 250 ml carton.....	280.80	18	PKU Glytactin RTD 15
† Liquid (original), 15 g protein, 22 g carbohydrate, 5.3 g fat per 250 ml, carton.....	684.45	30	PKU GMPPro LQ
† Liquid (vanilla), 15 g protein, 3.3 g carbohydrate, 4.6 g fat per 250 ml, carton.....	684.45	30	PKU Glytactin RTD 15
		30	PKU Glytactin RTD 15 Lite

Protein Free Supplements

PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDRATE, FAT WITH ADDED VITAMINS AND MINERALS –

Restricted see terms [on page 294](#)

† Powder (neutral) nil added protein and 67 g carbohydrate per 100 g, 400 g can.....	49.29	400 g	Energivit
--	-------	-------	-----------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Supplements for Tyrosinaemia			
AMINO ACID FORMULA (WITHOUT PHENYLALANINE AND TYROSINE) – Restricted see terms on page 294			
† Powder (neutral) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet.....	471.00	30	TYR Anamix Junior
† Powder (neutral), 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30	TYR Explore 5
† Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	260.00	400 g	TYR Anamix Infant
† Liquid (orange) 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle.....	941.40	36	TYR Anamix Junior LQ
† Liquid (juicy berries), 20 g protein, 12.75 g carbohydrate and 0.46 g fat and 0 g fibre per 125 ml pouch.....	1,684.80	30	TYR Lophlex LQ 20
† Liquid (juicy berries), 20 g protein, 8.8 g carbohydrate, 0.44 g fat and 0.5 g fibre per 125 ml pouch.....	1,684.80	30	TYR Lophlex LQ 20
<i>(TYR Lophlex LQ 20 Liquid (juicy berries), 20 g protein, 8.8 g carbohydrate, 0.44 g fat and 0.5 g fibre per 125 ml pouch to be delisted 1 December 2025)</i>			
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME TYROSINE AND PHENYLALANINE – Restricted see terms on page 294			
† Powder (Red Berry), 20 g protein, 6.3 carbohydrate, 1.6 g fat per 35 g sachet.....	1,398.60	30	TYR Sphere 20
† Powder (Vanilla), 20 g protein, 6.0 g carbohydrate, 1.6 g fat per 35 g sachet.....	1,398.60	30	TYR Sphere 20
X-Linked Adrenoleukodystrophy Products			
GLYCEROL TRIERUCATE – Restricted see terms on page 294			
† Liquid, 1,000 ml bottle			
GLYCEROL TRIOLEATE – Restricted see terms on page 294			
† Liquid, bottle.....	131.80	500 ml	GTO Oil
Supplements for Glycogen Storage Disease			
HIGH AMYLOPECTIN CORN-STARCH – Restricted see terms on page 294			
† Powder 0 g protein, 53 g carbohydrate, 0 g fat per 60 g sachet.....	241.62	30	Glycosade
Supplements for Organic Acidaemias			
AMINO ACID FORMULA (WITHOUT ISOLEUCINE, METHIONINE, THREONINE AND VALINE) – Restricted see terms on page 294			
† Powder 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can.....	260.00	400 g	MMA/PA Anamix Infant
AMINO ACID FORMULA (WITHOUT METHIONINE, THREONINE AND VALINE) – Restricted see terms on page 294			
† Powder (neutral), 5 g protein, 5.4 g carbohydrate, 2.3 g fat and 2.0 g fibre per 18 g sachet.....	750.30	30	MMA/PA Anamix Junior
† Powder, 15 g protein, 3.4 g carbohydrate, 0.05 g fat per 25 g sachet.....	1,048.95	30	MMA/PA Express 15
† Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet.....	349.65	30	MMA/PA Explore 5
Single Dose Amino Acids			
ARGININE – Restricted see terms on page 294			
† Powder 1.7 g protein, 1.9 g carbohydrate per 4 g sachet.....	211.45	30	Arginine2000

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
CITRULLINE – Restricted see terms on page 294			
† Powder 0.8 g protein, 2.9 g carbohydrate per 4 g sachet.....	211.45	30	Citrulline1000
ISOLEUCINE – Restricted see terms on page 294			
† Powder 0.04 g protein, 3.8 g carbohydrate per 4 g sachet.....	141.05	30	Isoleucine50
LEUCINE – Restricted see terms on page 294			
† Powder 0.08 g protein, 3.7 g carbohydrate per 4 g sachet.....	141.05	30	Leucine100
PHENYLALANINE – Restricted see terms on page 294			
† Powder 0.04 g protein, 3.8 g carbohydrate per 4 g sachet.....	141.05	30	Phenylalanine50
TYROSINE – Restricted see terms on page 294			
† Powder 0.8 g protein, 2.9 g carbohydrate per 4 g sachet.....	211.45	30	Tyrosine1000
VALINE – Restricted see terms on page 294			
† Powder 0.04 g protein, 3.8 g carbohydrate per 4 g sachet.....	141.05	30	Valine50

Other Fat Modified Products

ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCERIDES – Restricted see terms [on page 294](#)

† Powder (neutral), 12.5 g protein, 60 g carbohydrate and 16.4 g fat per 100 g sachet.....	47.01	10	Emsogen
---	-------	----	---------

Essential Amino Acids

ESSENTIAL AMINO ACID FORMULA – Restricted see terms [on page 294](#)

† Powder (neutral) 79 g protein per 100 g, 200 g can.....	313.73	200 g	Essential Amino Acid Mix
---	--------	-------	--------------------------

Specialised Formulas

Diabetic Products

➔ **Restricted (RS1215)**

Initiation

Any of the following:

- 1 For patients with type I or type II diabetes suffering weight loss and malnutrition that requires nutritional support; or
- 2 For patients with pancreatic insufficiency; or
- 3 For patients who have, or are expected to, eat little or nothing for 5 days; or
- 4 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or
- 5 For use pre- and post-surgery; or
- 6 For patients being tube-fed; or
- 7 For tube-feeding as a transition from intravenous nutrition.

DIABETIC ORAL FEED 1 KCAL/ML – Restricted see terms [above](#)

† Liquid 4.9 g protein, 11.7 g carbohydrate, 3.8 g fat and 2 g fibre per 100 ml, 200 ml bottle	2.25	1	Diasip (strawberry) Diasip (vanilla)
---	------	---	---

LOW-GI ENTERAL FEED 1 KCAL/ML – Restricted see terms [above](#)

† Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 500 ml bottle.....	4.65	1	Glucerna Select
† Liquid 4.3 g protein, 11.3 g carbohydrate and 4.2 g fat per 100 ml, 1,000 ml bottle			<i>e.g. Nutrison Advanced Diasion</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LOW-GI ORAL FEED 1 KCAL/ML – Restricted see terms on the previous page			
† Liquid 7 g protein, 10.9 g carbohydrate, 2.7 g fat and 2 g fibre per 100 ml, 200 ml bottle	2.10	1	Nutren Diabetes (vanilla)

Elemental and Semi-Elemental Products

➔ Restricted (RS1216)

Initiation

Any of the following:

- 1 Malabsorption; or
- 2 Short bowel syndrome; or
- 3 Enterocutaneous fistulas; or
- 4 Eosinophilic enteritis (including oesophagitis); or
- 5 Inflammatory bowel disease; or
- 6 Acute pancreatitis where standard feeds are not tolerated; or
- 7 Patients with multiple food allergies requiring enteral feeding.

AMINO ACID ORAL FEED – Restricted

 see terms [above](#)

† Powder 11 g protein, 62 g carbohydrate and 1 g fat per sachet, 80 g sachet.....	4.50	1	Vivonex TEN
---	------	---	-------------

AMINO ACID ORAL FEED 0.8 KCAL/ML – Restricted

 see terms [above](#)

† Liquid 2.5 g protein, 11 g carbohydrate and 3.5 g fat per 100 ml, 250 ml carton.....	179.46	18	Elemental 028 Extra (grapefruit) Elemental 028 Extra (pineapple & orange) Elemental 028 Extra (summer fruits)
--	--------	----	---

PEPTIDE-BASED ENTERAL FEED 1 KCAL/ML – Restricted

 see terms [above](#)

† Liquid 4 g protein, 17.7 g carbohydrate and 1.7 g fat per 100 ml, 500 ml bottle.....	7.47	1	Nutrison Advanced Peptisorb
--	------	---	-----------------------------

PEPTIDE-BASED ENTERAL FEED 1.5 KCAL/ML – Restricted

 see terms [above](#)

† Liquid 6.75 g protein, 18.4 g carbohydrate and 5.5 g fat per 100 ml, 1,000 ml bottle	22.39	1	Vital
--	-------	---	-------

PEPTIDE-BASED ORAL FEED – Restricted

 see terms [above](#)

† Powder 13.7 g protein, 62.9 g carbohydrate and 17.5 g fat per 100 g, 400 g can			<i>e.g. Peptamen Junior</i>
† Powder 13.8 g protein, 59 g carbohydrate and 18 g fat per 100 g, 400 g can			<i>e.g. MCT Peptide; MCT Peptide 1+</i>

PEPTIDE-BASED ORAL FEED 1 KCAL/ML – Restricted

 see terms [above](#)

† Liquid 5 g protein, 16 g carbohydrate and 1.69 g fat per 100 ml, carton	4.95	237 ml	Peptamen OS 1.0 (Vanilla)
---	------	--------	---------------------------

Fat Modified Products

FAT-MODIFIED FEED – Restricted

 see terms [on the next page](#)

† Powder 12.8 g protein, 68.6 g carbohydrate and 12.9 g fat per 100 g, can	62.90	400 g	Monogen
---	-------	-------	---------

	Price (ex man. excl. GST)		Brand or Generic Manufacturer
	\$	Per	

➔ Restricted (RS1470)

Initiation

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

Hepatic Products

➔ Restricted (RS1217)

Initiation

For children (up to 18 years) who require a liver transplant.

HEPATIC ORAL FEED – **Restricted** see terms [above](#)

⬆ Powder 12 g protein, 56 g carbohydrate and 22 g fat per 100 g, can	93.97	400 g	Heparon Junior
--	-------	-------	----------------

High Calorie Products

➔ Restricted (RS1317)

Initiation

Any of the following:

- 1 Patient is fluid volume or rate restricted; or
- 2 Patient requires low electrolyte; or
- 3 Both:
 - 3.1 Any of the following:
 - 3.1.1 Cystic fibrosis; or
 - 3.1.2 Any condition causing malabsorption; or
 - 3.1.3 Faltering growth in an infant/child; or
 - 3.1.4 Increased nutritional requirements; and
 - 3.2 Patient has substantially increased metabolic requirements.

ENTERAL FEED 2 KCAL/ML – **Restricted** see terms [above](#)

⬆ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 500 ml bottle.....	6.82	1	Nutrison Concentrated
⬆ Liquid 8.4 g protein, 21.9 g carbohydrate, 9.1 g fat and 0.5 g fibre per 100 ml, 1,000 ml bottle	13.64	1	Ensure Two Cal HN RTH

ORAL FEED 2 KCAL/ML – **Restricted** see terms [above](#)

⬆ Liquid 8.4 g protein, 22.4 g carbohydrate, 8.9 g fat and 0.8 g fibre per 100 ml, 200 ml bottle	2.34	1	Two Cal HN
--	------	---	------------

High Protein Products

HIGH PROTEIN ENTERAL FEED 1.25 KCAL/ML – **Restricted** see terms [below](#)

⬇ Liquid 6.3 g protein, 14.2 g carbohydrate and 4.9 g fat per 100 ml, bottle	12.00	1,000 ml	Nutrison Protein Plus
--	-------	----------	-----------------------

➔ Restricted (RS1327)

Initiation

Both:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
 - 2.1 Patient has liver disease; or
 - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
 - 2.3 Patient is fluid restricted; or
 - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.26 KCAL/ML – **Restricted** see terms [below](#)

⚡ Liquid 10 g protein, 10.4 g carbohydrate and 4.9 g fat per 100 ml, bottle 8.67 500 ml Nutrison Protein Intense

➔ **Restricted** (RS1327)

Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
 - 2.1 Patient has liver disease; or
 - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
 - 2.3 Patient is fluid restricted; or
 - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.28 KCAL/ML – **Restricted** see terms [below](#)

⚡ Liquid 6.3 g protein, 14.1 g carbohydrate, 4.9 g fat and 1.5 g fibre per 100 ml, bottle 12.54 1,000 ml Nutrison Protein Plus Multi Fibre

➔ **Restricted** (RS1327)

Initiation

Both:

- 1 The patient has a high protein requirement; and
- 2 Any of the following:
 - 2.1 Patient has liver disease; or
 - 2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
 - 2.3 Patient is fluid restricted; or
 - 2.4 Patient's needs cannot be more appropriately met using high calorie product.

Infant Formulas

AMINO ACID FORMULA – **Restricted** see terms [on the next page](#)

⚡ Powder 1.95 g protein, 8.1 g carbohydrate and 3.5 g fat per 100 ml, 400 g can		<i>e.g. Neocate</i>
⚡ Powder 13 g protein, 49 g carbohydrate and 23 g fat per 100 g, can55.61	400 g	Neocate SYNEO
⚡ Powder 13.3 g protein, 56 g carbohydrate and 22 g fat per 100 g, can55.61	400 g	Neocate Junior Unflavoured
⚡ Powder 13.3 g protein, 57 g carbohydrate and 24.6 g fat per 100 g, can43.60	400 g	Alfamino
⚡ Powder 13.5 g protein, 52 g carbohydrate and 24.5 g fat per 100 g, can55.61	400 g	Neocate Gold (Unflavoured)
⚡ Powder 14.8 g protein, 51.4 g carbohydrate and 23 g fat per 100 g, can55.61	400 g	Neocate Junior Vanilla
⚡ Powder 15 g protein, 56 g carbohydrate and 20 g fat per 100 g, can43.60	400 g	Alfamino Junior
⚡ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can.....65.72	400 g	Elecare LCP (Unflavoured)
⚡ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can.....65.72	400 g	Elecare (Unflavoured) Elecare (Vanilla)

Price (ex man. excl. GST) \$	Brand or Generic Manufacturer
Per	

➔ Restricted (RS1867)

Initiation

Any of the following:

- 1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
- 2 History of anaphylaxis to cows' milk protein formula or dairy products; or
- 3 Eosinophilic oesophagitis; or
- 4 Ultra-short gut; or
- 5 Severe Immune deficiency.

Continuation

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows' milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
- 3 Amino acid formula is required for a nutritional deficit.

Initiation – patients who are currently funded under RS1502 or SA1557

Limited to 3 months treatment

All of the following:

- 1 Patient has a valid initiation or renewal approval for extensively hydrolysed formula (RS1502); and
- 2 Patient is unable to source funded Aptamil powder at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Hospital Restriction RS1502. There is no continuation criteria under this criterion.

ENTERAL LIQUID PEPTIDE FORMULA – **Restricted** see terms [below](#)

↓ Liquid 4.2 g protein, 18.6 g carbohydrate and 6.58 g fat per 100 ml, 500 ml bottle	18.66	1	Nutrini Peptisorb Energy
---	-------	---	--------------------------

➔ Restricted (RS1775)

Initiation

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure; or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

Note: A reasonable trial is defined as a 2-4 week trial.

Continuation

Both:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula.

EXTENSIVELY HYDROLYSED FORMULA – **Restricted** see terms [below](#)

↓ Powder 1.6 g protein, 7.5 g carbohydrate and 3.1 g fat per 100 ml, 900 g can.....	36.20	900 g	Allerpro Syneo 1
↓ Powder 1.6 g protein, 7.8 g carbohydrate and 3.2 g fat per 100 ml, 900 g can.....	36.20	900 g	Allerpro Syneo 2
↓ Powder 14 g protein, 53.4 g carbohydrate and 27.3 g fat per 100 g, can	18.10	450 g	Pepti-Junior

➔ **Restricted (RS1502)**

Initiation

Any of the following:

- 1 Both:
 - 1.1 Cows' milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 For step down from Amino Acid Formula.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Continuation

Both:

- 1 An assessment as to whether the infant can be transitioned to a cows' milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula.

FRUCTOSE-BASED FORMULA

Powder 14.6 g protein, 49.7 g carbohydrate and 30.8 g fat per 100 g, 400 g can			<i>e.g. Galactomin 19</i>
--	--	--	---------------------------

LACTOSE-FREE FORMULA

Powder 1.3 g protein, 7.3 g carbohydrate and 3.5 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Gold De-Lact</i>
Powder 1.5 g protein, 7.2 g carbohydrate and 3.6 g fat per 100 ml, 900 g can			<i>e.g. S26 Lactose Free</i>

LOW-CALCIUM FORMULA

Powder 14.8 g protein, 53.7 g carbohydrate and 26.7 g fat per 100 g and tuna fish oil (DHA), can.....	46.18	400 g	Locasol
---	-------	-------	---------

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML – Restricted see terms [below](#)

↓ Liquid 2.6 g protein, 10.3 g carbohydrate, 5.4 g fat and 0.6 g fibre per 100 ml, 125 ml bottle	2.80	1	Infatrini
--	------	---	-----------

➔ **Restricted (RS1614)**

Initiation – Fluid restricted or volume intolerance with faltering growth

Both:

- 1 Either:
 - 1.1 The patient is fluid restricted or volume intolerant; or
 - 1.2 The patient has increased nutritional requirements due to faltering growth; and
- 2 Patient is under 18 months old and weighs less than 8kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

PRETERM FORMULA – Restricted see terms [below](#)

↓ Liquid 2.2 g protein, 8.4 g carbohydrate and 4.4 g fat per 100 ml, bottle	0.75	100 ml	S26 LBW Gold RTF
↓ Liquid 2.3 g protein, 8.6 g carbohydrate and 4.2 g fat per 100 ml, 90 ml bottle			<i>e.g. Pre Nan Gold RTF</i>
↓ Liquid 2.6 g protein, 8.4 g carbohydrate and 3.9 g fat per 100 ml, 70 ml bottle			<i>e.g. Karicare Aptamil Gold+Preterm</i>

➔ **Restricted (RS1224)**

Initiation

For infants born before 33 weeks' gestation or weighing less than 1.5 kg at birth.

THICKENED FORMULA

Powder 1.8 g protein, 8.1 g carbohydrate and 3.3 g fat per 100 ml, 900 g can			<i>e.g. Karicare Aptamil Thickened AR</i>
--	--	--	---

Ketogenic Diet Products

HIGH FAT FORMULA – Restricted see terms [below](#)

↓ Powder 14.3 g protein, 2.8 g carbohydrate and 69.2 g fat per 100 g, can	36.92	300 g	Ketocal 4:1 (Unflavoured)
↓ Powder 15.4 g protein, 7.2 g carbohydrate and 68.6 g fat per 100 g, can	36.92	300 g	Ketocal 4:1 (Vanilla)
			Ketocal 3:1 (Unflavoured)

➔ **Restricted (RS1225)**

Initiation

For patients with intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Paediatric Products

➔ **Restricted (RS1473)**

Initiation

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
continued...			
2.1 The child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or			
2.2 Any condition causing malabsorption; or			
2.3 Faltering growth in an infant/child; or			
2.4 Increased nutritional requirements; or			
2.5 The child is being transitioned from TPN or tube feeding to oral feeding; or			
2.6 The child has eaten, or is expected to eat, little or nothing for 3 days.			
PAEDIATRIC ENTERAL FEED 0.76 KCAL/ML – Restricted see terms on the previous page			
† Liquid 2.5 g protein, 12.5 g carbohydrate, 3.3 g fat and 0.7 g fibre per 100 ml, 500 ml bottle	6.27	1	Nutrini Low Energy Multi Fibre RTH
PAEDIATRIC ENTERAL FEED 1 KCAL/ML – Restricted see terms on the previous page			
† Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, 500 ml bottle.....	3.32	1	Pediasure RTH
† Liquid 2.7 g protein, 12.3 g carbohydrate and 4.4 g fat per 100 ml, 500 ml bottle	4.69	1	Nutrini RTH
PAEDIATRIC ENTERAL FEED 1.5 KCAL/ML – Restricted see terms on the previous page			
† Liquid 4.1 g protein, 18.5 g carbohydrate and 6.7 g fat per 100 ml, 500 ml bottle	7.46	1	Nutrini Energy RTH
† Liquid 4.1 g protein, 18.5 g carbohydrate, 6.7 g fat and 0.8 g fibre per 100 ml, 500 ml bottle	7.14	1	Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1 KCAL/ML – Restricted see terms on the previous page			
† Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, 200 ml bottle.....	1.33	1	Pediasure (chocolate) Pediasure (strawberry) Pediasure (vanilla)
† Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, 250 ml can.....	1.66	1	Pediasure (vanilla)
PAEDIATRIC ORAL FEED 1.5 KCAL/ML – Restricted see terms on the previous page			
† Liquid 3.4 g protein, 18.8 g carbohydrate and 6.8 g fat per 100 ml, 200 ml bottle	1.90	1	Fortini (Strawberry) Fortini (Vanilla)
† Liquid 4.0 g protein, 18.8 g carbohydrate, 6.8 g fat and 1.5 g fibre per 100 ml, 200 ml bottle	1.90	1	Fortini Multi Fibre (chocolate) Fortini Multi Fibre (strawberry) Fortini Multi Fibre (unflavoured) Fortini Multi Fibre (vanilla)
† Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, 500 ml bottle	8.67	1	Pediasure Plus

Renal Products

LOW ELECTROLYTE ORAL FEED – **Restricted** see terms [below](#)

↓ Powder 7.5 g protein, 57.6 g carbohydrate and 25.9 g fat per 100 g, can 64.26

400 g

Kindergen

→ **Restricted** (RS1227)

Initiation

For children (up to 18 years) with acute or chronic kidney disease.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
LOW ELECTROLYTE ORAL FEED 1.8 KCAL/ML			
Liquid 8 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, 220 ml bottle	3.31	1	Nepro HP (strawberry) Nepro HP (vanilla)
LOW ELECTROLYTE ORAL FEED 2 KCAL/ML – Restricted see terms below			
⬇ Liquid 3 g protein, 25.5 g carbohydrate and 9.6 g fat per 100 ml, 237 ml bottle			
⬇ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 125 ml carton.....	13.72	4	Renilon 7.5 (apricot) Renilon 7.5 (caramel)
⬇ Liquid 9.1 g protein, 19 g carbohydrate and 10 g fat per 100 ml, 200 ml bottle.....	13.24	4	Novasource Renal (Vanilla)

➔ **Restricted (RS1228)**

Initiation

For patients with acute or chronic kidney disease.

Surgical Products

HIGH ARGININE ORAL FEED 1.4 KCAL/ML – **Restricted see terms [below](#)**

⬇ Liquid 10.4 g protein, 8 g carbohydrate, 4.4 g fat and 0 g fibre per 100 ml, 250 ml carton	56.00	10	Impact Advanced Recovery
--	-------	----	-----------------------------

➔ **Restricted (RS1231)**

Initiation

Three packs per day for 5 to 7 days prior to major gastrointestinal, head or neck surgery.

PREOPERATIVE CARBOHYDRATE FEED 0.5 KCAL/ML – **Restricted see terms [below](#)**

⬇ Oral liq 0 g protein, 12.6 g carbohydrate and 0 g fat per 100 ml, 200 ml bottle.....	8.64	4	preOp
--	------	---	-------

➔ **Restricted (RS1415)**

Initiation

Maximum of 400 ml as part of an Enhanced Recovery After Surgery (ERAS) protocol 2 to 3 hours before major abdominal surgery.

Standard Feeds

➔ **Restricted (RS1214)**

Initiation

Any of the following:

For patients with malnutrition, defined as any of the following:

1 Any of the following:

1.1 BMI < 18.5; or

1.2 Greater than 10% weight loss in the last 3-6 months; or

1.3 BMI < 20 with greater than 5% weight loss in the last 3-6 months; or

2 For patients who have, or are expected to, eat little or nothing for 5 days; or

3 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or

4 For use pre- and post-surgery; or

5 For patients being tube-fed; or

6 For tube-feeding as a transition from intravenous nutrition; or

7 For any other condition that meets the community Special Authority criteria.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ENTERAL FEED 1.5 KCAL/ML – Restricted see terms on the previous page			
† Liquid 6 g protein, 18.3 g carbohydrate and 5.8 g fat per 100 ml, 1,000 ml bottle	9.00	1	Nutrison Energy
† Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 1.5 g fibre per 100 ml, 1,000 ml bottle	8.68	1	Nutrison Energy Multi Fibre
† Liquid 6.27 g protein, 20.4 g carbohydrate and 4.9 g fat per 100 ml, 1,000 ml bag	8.68	1	Ensure Plus HN RTH
† Liquid 6.38 g protein, 21.1 g carbohydrate, 4.9 g fat and 1.2 g fibre per 100 ml, 1,000 ml bottle	8.68	1	Jevity HiCal RTH
† Liquid 6 g protein, 18.5 g carbohydrate and 5.8 g fat per 100 ml, 1,000 ml bottle	9.00	1	Nutrison Energy
† Liquid 6.25 g protein, 20 g carbohydrate and 5 g fat per 100 ml, 250 ml can....	2.17	1	Ensure Plus HN
<i>(Nutrison Energy Liquid 6 g protein, 18.3 g carbohydrate and 5.8 g fat per 100 ml, 1,000 ml bottle to be delisted 1 January 2026)</i>			
ENTERAL FEED 1 KCAL/ML – Restricted see terms on the previous page			
† Liquid 4 g protein, 12.3 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bottle	6.90	1	Nutrison RTH
† Liquid 4 g protein, 12.3 g carbohydrate, 3.9 g fat and 1.5 g fibre per 100 ml, 1,000 ml bottle	7.21	1	Nutrison Multi Fibre
† Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, 1,000 ml bottle	6.56	1	Osmolite RTH
† Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, 1,000 ml bottle	6.56	1	Jevity RTH
† Liquid 4 g protein, 12.4 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bottle	6.90	1	Nutrison RTH
<i>(Nutrison RTH Liquid 4 g protein, 12.3 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bottle to be delisted 1 January 2026)</i>			
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Restricted see terms on the previous page			
† Liquid 5.5 g protein, 8.8 g carbohydrate, 2.5 g fat and 1.5 g fibre per 100 ml, 1,000 ml bottle	9.05	1	Nutrison 800 Complete Multi Fibre
HIGH PROTEIN ORAL FEED 2.4 KCAL/ML – Restricted see terms on the previous page			
† Liquid 14.6 g protein, 25.3 g carbohydrate and 9.6 g fat per 100 ml, 125 ml bottle			<i>e.g. Fortisip Compact Protein</i>
ORAL FEED – Restricted see terms on the previous page			
† Powder 15.9 g protein, 57.4 g carbohydrate and 14 g fat per 100 g, can	26.00	850 g	Ensure (Chocolate) Ensure (Vanilla)
† Powder 23 g protein, 65 g carbohydrate and 2.5 g fat per 100 g, can	15.90	840 g	Sustagen Hospital Formula (Chocolate) Sustagen Hospital Formula (Vanilla)
ORAL FEED 1 KCAL/ML – Restricted see terms on the previous page			
† Liquid 3.8 g protein, 23 g carbohydrate and 12.7 g fibre per 100 ml, 237 ml carton			<i>e.g. Resource Fruit Beverage</i>

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
ORAL FEED 1.5 KCAL/ML – Restricted see terms on page 307			
† Liquid 4 g protein and 33.5 g carbohydrate per 100 ml, 200 ml bottle	3.30	200 ml	Fortijuice (Apple) Fortijuice (Orange) Fortijuice (Strawberry)
† Liquid 6 g protein, 18.4 g carbohydrate and 5.8 g fat per 100 ml, 200 ml bottle.....	1.76	1	Fortisip (Banana) Fortisip (Chocolate) Fortisip (Strawberry) Fortisip (Vanilla)
† Liquid 6.25 g protein, 20.2 g carbohydrate and 4.92 g fat per 100 ml, 200 ml bottle	1.56	1	Ensure Plus (Banana) Ensure Plus (Chocolate) Ensure Plus (Fruit of the forest) Ensure Plus (Vanilla)
† Liquid 5.5 g protein, 21.1 g carbohydrate and 4.81 g fat per 100 ml, 237 ml can	1.65	1	Ensure Plus (Vanilla)
ORAL FEED WITH FIBRE 1.5 KCAL/ML – Restricted see terms on page 307			
† Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 2.3 g fibre per 100 ml, 200 ml bottle	1.76	1	Fortisip Multi Fibre (chocolate) Fortisip Multi Fibre (strawberry) Fortisip Multi Fibre (vanilla)

Bacterial and Viral Vaccines

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – **Restricted** see terms [below](#)

† Inj 30 IU diphtheria toxoid with 30IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5 ml syringe – 5% DV Dec-24 to 2027	0.00	10	Infanrix IPV
--	------	----	---------------------

➔ **Restricted** ([RS1387](#))

Initiation

Any of the following:

- 1 A single dose for children up to the age of 7 who have completed primary immunisation; or
- 2 A course of up to four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4 Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE –

Restricted see terms [below](#)

† Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen 10mcg H. influenzae type b with tetanus toxoid 20-40mcg in 0.5ml syringe – 5% DV Dec-24 to 2027	0.00	10	Infanrix-hexa
--	------	----	----------------------

➔ **Restricted** ([RS2051](#))

Initiation

Any of the following:

- 1 Up to four doses for children under the age of 10 years for primary immunisation; or
- 2 An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
- 3 An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens; or
- 4 Up to five doses for children under the age of 10 years receiving solid organ transplantation.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Bacterial Vaccines

BACILLUS CALMETTE-GUERIN VACCINE – **Restricted** see terms [below](#)

† Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial with diluent – 5% DV Dec-24 to 2027	0.00	10	BCG Vaccine AJV
---	------	----	------------------------

➔ **Restricted** ([RS1233](#))

Initiation

All of the following:

For infants at increased risk of tuberculosis defined as:

- 1 Living in a house or family with a person with current or past history of TB; and
- 2 Having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

equal to 40 per 100,000 for 6 months or longer; and

3 During their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000.

Note: A list of countries with high rates of TB are available at <http://www.health.govt.nz/tuberculosis> (Search for Downloads) or www.bcgatlas.org/index.php

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – **Restricted** see terms [below](#)

↓ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis
toxoid, 8 mcg pertussis filamentous haemagglutinin and 2.5 mcg
pertactin in 0.5 ml prefilled syringe – **5% DV Dec-24 to 2027** 0.00 10 **Boostrix**

→ **Restricted (RS1790)**

Initiation

Any of the following:

- 1 A single dose for pregnant women in the second or third trimester of each pregnancy; or; or
- 2 A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or; or
- 3 A course of up to four doses is funded for children from age 7 up the age of 18 years inclusive to complete full primary immunisation; or
- 4 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5 A single dose for vaccination of patients aged from 65 years old; or
- 6 A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7 For vaccination of previously unimmunised or partially immunised patients; or
- 8 For revaccination following immunosuppression; or
- 9 For boosting of patients with tetanus-prone wounds.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

HAEMOPHILUS INFLUENZAE TYPE B VACCINE – **Restricted** see terms [below](#)

↓ Inj 10 mcg vial with diluent syringe – **5% DV Dec-24 to 2027** 0.00 1 **Act-HIB**

→ **Restricted (RS1520)**

Initiation

Therapy limited to 1 dose

Any of the following:

- 1 For primary vaccination in children; or
- 2 An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre- or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or
- 3 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

MENINGOCOCCAL (A, C, Y AND W-135) CONJUGATE VACCINE

↓ Inj 10 mcg of each meningococcal polysaccharide conjugated to a total
of approximately 55 mcg of tetanus toxoid carrier per 0.5 ml vial –
5% DV Dec-24 to 2027 0.00 1 **MenQuadfi**

→ **Restricted (RS2019)**

Initiation

Either:

- 1 Any of the following:
 - 1.1 Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with HIV,

continued...

continued...

complement deficiency (acquired or inherited), functional or anatomic asplenia or pre or post solid organ transplant; or

- 1.2 One dose for close contacts of meningococcal cases of any group; or
- 1.3 One dose for person who has previously had meningococcal disease of any group; or
- 1.4 A maximum of two doses for bone marrow transplant patients; or
- 1.5 A maximum of two doses for person pre and post-immunosuppression*; or

2 Both:

2.1 Person is aged between 13 and 25 years, inclusive; and

2.2 Either:

2.2.1 One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or

2.2.2 One dose for individuals who turn 13 years of age while living in boarding school hostels.

Notes: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

‡ Inj 5 mcg of each meningococcal polysaccharide conjugated to a total of approximately 44 mcg of tetanus toxoid carrier in 0.5 ml vial..... 0.00 1 Nimenrix

➔ **Restricted (RS2037)**

Initiation – Children under 12 months of age

Any of the following:

- 1 A maximum of three doses (dependant on age at first dose) for patients pre- and post- splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post- solid organ transplant; or
- 2 A maximum of three doses (dependant on age at first dose) for close contacts of meningococcal cases of any group; or
- 3 A maximum of three doses (dependant on age at first dose) for child who has previously had meningococcal disease of any group; or
- 4 A maximum of three doses (dependant on age at first dose) for bone marrow transplant patients; or
- 5 A maximum of three doses (dependant on age at first dose) for child pre- and post-immunosuppression*.

Notes: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

MENINGOCOCCAL B MULTICOMPONENT VACCINE – Restricted see terms [below](#)

‡ Inj 175 mcg per 0.5 ml prefilled syringe..... 0.00 1 Bexsero
10 Bexsero

➔ **Restricted (RS2141)**

Initiation – Primary immunisation for children up to 59 months of age inclusive

Therapy limited to 3 doses

A primary course of up to three doses (dependent on age at first dose) for previously unvaccinated children up to the age of 59 months inclusive.

Initiation – High-risk individuals 5 years of age or over

Both:

- 1 Person is aged at least 5 years; and
- 2 Any of the following:
 - 2.1 Up to two doses and a booster every five years for patients pre- and post-splenectomy; or
 - 2.2 Up to two doses and a booster every five years for patients with functional or anatomic asplenia, HIV, complement

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- deficiency (acquired or inherited); or
- 2.3 Up to two doses and a booster every five years pre- or post-solid organ transplant; or
- 2.4 Up to two doses for close contacts of meningococcal cases of any group; or
- 2.5 Up to two doses for person who has previously had meningococcal disease of any group; or
- 2.6 Up to two doses for bone marrow transplant patients; or
- 2.7 Up to two doses for person pre- and post-immunosuppression* .

Initiation – Person is aged between 13 and 25 years (inclusive)

Therapy limited to 2 doses

Both:

- 1 Person is aged between 13 and 25 years (inclusive); and
- 2 Either:
 - 2.1 Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or
 - 2.2 Two doses for individuals who turn 13 years of age while living in boarding school hostels.

Note: *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE – Restricted see terms [below](#)

‡ Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5 ml syringe – **5% DV**

Dec-24 to 2027	0.00	1	Prevenar 13
		10	Prevenar 13

→ **Restricted (RS1936)**

Initiation – Primary course for previously unvaccinated children aged under 5 years

Therapy limited to 3 doses

A primary course of three doses for previously unvaccinated children up to the age of 59 months inclusive.

Initiation – High risk individuals who have received PCV10

Therapy limited to 2 doses

Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10.

Initiation – High risk children aged under 5 years

Therapy limited to 4 doses

Both:

- 1 Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high-risk children aged under 5 years; and
- 2 Any of the following:
 - 2.1 on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - 2.2 primary immune deficiencies; or
 - 2.3 HIV infection; or
 - 2.4 renal failure, or nephrotic syndrome; or
 - 2.5 are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - 2.6 cochlear implants or intracranial shunts; or
 - 2.7 cerebrospinal fluid leaks; or
 - 2.8 receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - 2.9 chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or

continued...

continued...

- 2.10 pre term infants, born before 28 weeks gestation; or
- 2.11 cardiac disease, with cyanosis or failure; or
- 2.12 diabetes; or
- 2.13 Down syndrome; or
- 2.14 who are pre-or post-splenectomy, or with functional asplenia.

Initiation – High risk individuals 5 years and over

Therapy limited to 4 doses

Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency.

Initiation – Testing for primary immunodeficiency diseases

For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – **Restricted** see terms [below](#)

‡ Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype) – **5% DV Dec-24 to 2027** 0.00 1 **Pneumovax 23**

➔ **Restricted (RS1587)**

Initiation – High risk patients

Therapy limited to 3 doses

For patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy; or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency.

Initiation – High risk children

Therapy limited to 2 doses

Both:

- 1 Patient is a child under 18 years for (re-)immunisation; and
- 2 Any of the following:
 - 2.1 On immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - 2.2 With primary immune deficiencies; or
 - 2.3 With HIV infection; or
 - 2.4 With renal failure, or nephrotic syndrome; or
 - 2.5 Who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - 2.6 With cochlear implants or intracranial shunts; or
 - 2.7 With cerebrospinal fluid leaks; or
 - 2.8 Receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - 2.9 With chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - 2.10 Pre term infants, born before 28 weeks gestation; or
 - 2.11 With cardiac disease, with cyanosis or failure; or
 - 2.12 With diabetes; or
 - 2.13 With Down syndrome; or
 - 2.14 Who are pre-or post-splenectomy, or with functional asplenia.

Initiation – Testing for primary immunodeficiency diseases

For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
SALMONELLA TYPHI VACCINE – Restricted see terms below			
↓ Inj 25 mcg in 0.5 ml syringe			
→ Restricted (RS1243)			
Initiation			
For use during typhoid fever outbreaks.			
Viral Vaccines			
COVID-19 VACCINE			
↓ Inj 3 mcg brettevameran per 0.3 ml, 0.48 ml vial; infant vaccine, yellow cap.....	0.00	10	Comirnaty Omicron (JN.1)
→ Restricted (RS2042)			
Initiation – initial dose			
Up to three doses for previously unvaccinated children aged 6 months – 4 years at high risk of severe illness.			
↓ Inj 10 mcg brettevameran per 0.3 ml, 0.48 ml vial; paediatric vaccine, light blue cap	0.00	10	Comirnaty Omicron (JN.1)
→ Restricted (RS2041)			
Initiation – initial dose			
Either:			
1 One dose for previously unvaccinated children aged 5-11 years old; or			
2 Up to three doses for immunocompromised children aged 5-11 years old.			
↓ Inj 30 mcg brettevameran per 0.3 ml, 0.48 ml vial; adult vaccine, light grey cap	0.00	10	Comirnaty Omicron (JN.1)
→ Restricted (RS2040)			
Initiation – initial dose			
Any of the following:			
1 One dose for previously unvaccinated people aged 12-15 years old; or			
2 Up to three doses for immunocompromised people aged 12-15 years old; or			
3 Up to two doses for previously unvaccinated people 16-29 years old; or			
4 Up to four doses for people aged 16-29 at high risk of severe illness; or			
5 One dose for previously unvaccinated people aged 30 and older.			
Initiation – additional dose			
One additional dose every 6 months for people aged 30 years and over, additional dose is given at least 6 months after last dose.			
Continuation – additional dose			
One additional dose every 6 months for people aged 30 years and over, additional dose is given at least 6 months after last dose.			
HEPATITIS A VACCINE – Restricted see terms below			
↓ Inj 720 ELISA units in 0.5 ml syringe – 5% DV Dec-24 to 2027	0.00	1	Havrix Junior
↓ Inj 1440 ELISA units in 1 ml syringe – 5% DV Dec-24 to 2027	0.00	1	Havrix 1440
→ Restricted (RS1638)			
Initiation			
Any of the following:			
1 Two vaccinations for use in transplant patients; or			
2 Two vaccinations for use in children with chronic liver disease; or			
3 One dose of vaccine for close contacts of known hepatitis A cases.			
HEPATITIS B RECOMBINANT VACCINE			
↓ Inj 10 mcg per 0.5 ml prefilled syringe – 5% DV Dec-24 to 2027	0.00	1	Engerix-B

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ Restricted (RS2049)

Initiation

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4 For HIV positive patients; or
- 5 For hepatitis C positive patients; or
- 6 For patients following non-consensual sexual intercourse; or
- 7 For patients prior to planned immunosuppression for greater than 28 days; or
- 8 For patients following immunosuppression; or
- 9 For solid organ transplant patients; or
- 10 For post-haematopoietic stem cell transplant (HSCT) patients; or
- 11 Following needle stick injury.

⚡ Inj 20 mcg per 1 ml prefilled syringe – 5% DV Dec-24 to 2027 0.00 1 **Engerix-B**

➔ Restricted (RS2050)

Initiation

Any of the following:

- 1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
- 2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
- 3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
- 4 For HIV positive patients; or
- 5 For hepatitis C positive patients; or
- 6 For patients following non-consensual sexual intercourse; or
- 7 For patients prior to planned immunosuppression for greater than 28 days; or
- 8 For patients following immunosuppression; or
- 9 For solid organ transplant patients; or
- 10 For post-haematopoietic stem cell transplant (HSCT) patients; or
- 11 Following needle stick injury; or
- 12 For dialysis patients; or
- 13 For liver or kidney transplant patients.

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] – **Restricted** see terms [below](#)

⚡ Inj 270 mcg in 0.5 ml syringe – 5% DV Dec-24 to 2027 0.00 10 **Gardasil 9**

➔ Restricted (RS2038)

Initiation – Children aged 14 years and under

Therapy limited to 2 doses

Children aged 14 years and under.

Initiation – other conditions

Either:

- 1 Up to 3 doses for people aged 15 to 26 years inclusive; or
- 2 Both:
 - 2.1 People aged 9 to 26 years inclusive; and
 - 2.2 Any of the following:
 - 2.2.1 Up to 3 doses for confirmed HIV infection; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

2.2.2 Up to 3 doses people with a transplant (including stem cell); or

2.2.3 Up to 4 doses for Post chemotherapy.

Initiation – Recurrent Respiratory Papillomatosis

All of the following:

- 1 Either:
 - 1.1 Maximum of two doses for children aged 14 years and under; or
 - 1.2 Maximum of three doses for people aged 15 years and over; and
- 2 The person has recurrent respiratory papillomatosis; and
- 3 The person has not previously had an HPV vaccine.

INFLUENZA VACCINE

↓ Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine).....	120.00	10	Influvac Tetra (2025 formulation)
--	--------	----	--------------------------------------

→ Restricted (RS2013)

Initiation – People over 65

The patient is 65 years of age or over.

Initiation – cardiovascular disease

Any of the following:

- 1 Ischaemic heart disease; or
- 2 Congestive heart failure; or
- 3 Rheumatic heart disease; or
- 4 Congenital heart disease; or
- 5 Cerebro-vascular disease.

Note: hypertension and/or dyslipidaemia without evidence of end-organ disease is excluded from funding.

Initiation – chronic respiratory disease

Either:

- 1 Asthma, if on a regular preventative therapy; or
- 2 Other chronic respiratory disease with impaired lung function.

Note: asthma not requiring regular preventative therapy is excluded from funding.

Initiation – Other conditions

Either:

- 1 Any of the following:
 - 1.1 Diabetes; or
 - 1.2 chronic renal disease; or
 - 1.3 Any cancer, excluding basal and squamous skin cancers if not invasive; or
 - 1.4 Autoimmune disease; or
 - 1.5 Immune suppression or immune deficiency; or
 - 1.6 HIV; or
 - 1.7 Transplant recipient; or
 - 1.8 Neuromuscular and CNS diseases/ disorders; or
 - 1.9 Haemoglobinopathies; or
 - 1.10 Is a child on long term aspirin; or
 - 1.11 Has a cochlear implant; or
 - 1.12 Errors of metabolism at risk of major metabolic decompensation; or
 - 1.13 Pre and post splenectomy; or
 - 1.14 Down syndrome; or
 - 1.15 Is pregnant; or
 - 1.16 Is a child 4 years of age or under (inclusive) who has been hospitalised for respiratory illness or has a history of

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

significant respiratory illness; or

- 2 Patients in a long-stay inpatient mental health care unit or who are compulsorily detained long-term in a forensic unit within a Public Hospital.

Initiation – Serious mental health conditions or addiction

Any of the following:

- 1 schizophrenia; or
- 2 major depressive disorder; or
- 3 bipolar disorder; or
- 4 schizoaffective disorder; or
- 5 person is currently accessing secondary or tertiary mental health and addiction services.

MEASLES, MUMPS AND RUBELLA VACCINE – **Restricted** see terms [below](#)

⚡ Injection, measles virus 1,000 CCID50, mumps virus 5,012 CCID50, Rubella virus 1,000 CCID50; prefilled syringe/ampoule of diluent 0.5 ml – 5% DV Dec-24 to 2027	0.00	10	Priorix
--	------	----	----------------

➡ **Restricted (RS1487)**

Initiation – first dose prior to 12 months

Therapy limited to 3 doses

Any of the following:

- 1 For primary vaccination in children; or
- 2 For revaccination following immunosuppression; or
- 3 For any individual susceptible to measles, mumps or rubella.

Initiation – first dose after 12 months

Therapy limited to 2 doses

Any of the following:

- 1 For primary vaccination in children; or
- 2 For revaccination following immunosuppression; or
- 3 For any individual susceptible to measles, mumps or rubella.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

POLIOMYELITIS VACCINE – **Restricted** see terms [below](#)

⚡ Inj 80 D-antigen units in 0.5 ml syringe – 5% DV Dec-24 to 2027	0.00	1	IPOL
---	------	---	-------------

➡ **Restricted (RS1398)**

Initiation

Therapy limited to 3 doses

Either:

- 1 For partially vaccinated or previously unvaccinated individuals; or
- 2 For revaccination following immunosuppression.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

RABIES VACCINE

Inj 2.5 IU vial with diluent

ROTAVIRUS ORAL VACCINE – **Restricted** see terms [on the next page](#)

⚡ Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator – 5% DV Dec-24 to 2027	0.00	10	Rotarix
⚡ Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube	0.00	10	Rotarix
⚡ Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube (PVC free)	0.00	10	Rotarix

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

➔ **Restricted (RS1590)**

Initiation

Therapy limited to 2 doses

Both:

- 1 First dose to be administered in infants aged under 14 weeks of age; and
- 2 No vaccination being administered to children aged 24 weeks or over.

VARICELLA VACCINE [CHICKENPOX VACCINE]

↓ Inj 2000 PFU prefilled syringe plus vial – **5% DV Dec-24 to 2027** 0.00 10 **Varilrix**

➔ **Restricted (RS1591)**

Initiation – primary vaccinations

Therapy limited to 1 dose

Either:

- 1 Any infant born on or after 1 April 2016; or
- 2 For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox).

Initiation – other conditions

Therapy limited to 2 doses

Any of the following:

- 1 Any of the following:
 - for non-immune patients:
 - 1.1 With chronic liver disease who may in future be candidates for transplantation; or
 - 1.2 With deteriorating renal function before transplantation; or
 - 1.3 Prior to solid organ transplant; or
 - 1.4 Prior to any elective immunosuppression*; or
 - 1.5 For post exposure prophylaxis who are immune competent inpatients; or
 - 2 For patients at least 2 years after bone marrow transplantation, on advice of their specialist; or
 - 3 For patients at least 6 months after completion of chemotherapy, on advice of their specialist; or
 - 4 For HIV positive patients non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or
 - 5 For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or
 - 6 For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or
 - 7 For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

Note: * immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] – Restricted see terms below

↓ Inj 50 mcg per 0.5 ml vial plus vial..... 0.00 1 **Shingrix**
10 **Shingrix**

➔ **Restricted (RS2039)**

Initiation – people aged 18 years and over (Shingrix)

Therapy limited to 2 doses

Any of the following:

- 1 Pre- and post-haematopoietic stem cell transplant or cellular therapy; or
- 2 Pre- or post-solid organ transplant; or

continued...

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
--	------------------------------------	-----	-------------------------------------

continued...

- 3 Haematological malignancies; or
- 4 People living with poorly controlled HIV infection; or
- 5 Planned or receiving disease modifying anti-rheumatic drugs (DMARDs – targeted synthetic, biologic, or conventional synthetic) for polymyalgia rheumatica, systemic lupus erythematosus or rheumatoid arthritis; or
- 6 End stage kidney disease (CKD 4 or 5);; or
- 7 Primary immunodeficiency.

Diagnostic Agents

TUBERCULIN PPD [MANTOUX] TEST			
Inj 5 TU per 0.1 ml, 1 ml vial – 5% DV Dec-24 to 2027	0.00	1	Tubersol

	Price (ex man. excl. GST) \$	Per	Brand or Generic Manufacturer
Optional Pharmaceuticals			
NOTE:			
In addition to the products expressly listed here in Part III: Optional Pharmaceuticals, a range of hospital medical devices are listed in an addendum to Part III which is available at schedule.pharmac.govt.nz . The Optional Pharmaceuticals listed in the addendum are deemed to be listed in Part III, and the Rules of the Pharmaceutical Schedule applying to products listed in Part III apply to them.			
BETA-HCG LOW SENSITIVITY URINE TEST KIT			
Note: For use in abortion services only.			
Midstream.....	16.28	1 test	CheckTop
BLOOD GLUCOSE DIAGNOSTIC TEST METER			
1 meter with 50 lancets, a lancing device, and 10 diagnostic test strips	20.00	1	CareSens N Premier
	10.00		Caresens N
			Caresens N POP
BLOOD GLUCOSE DIAGNOSTIC TEST STRIP			
Blood glucose test strips.....	10.56	50 test	CareSens N
Test strips	10.56	50 test	CareSens PRO
BLOOD KETONE DIAGNOSTIC TEST STRIP			
Test strips	15.50	10 strip	KetoSens
DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER			
Meter with 50 lancets, a lancing device, and 10 blood glucose diagnostic test strips	20.00	1	CareSens Dual
MASK FOR SPACER DEVICE			
Small.....	2.70	1	e-chamber Mask
PEAK FLOW METER			
Low Range	9.54	1	Mini-Wright AFS Low Range
Normal Range	9.54	1	Mini-Wright Standard
PREGNANCY TEST - HCG URINE			
Cassette – 5% DV Mar-25 to 2027	16.00	40 test	David One Step Cassette Pregnancy Test
SODIUM NITROPRUSSIDE			
Test strip.....	22.00	50 strip	Ketostix
SPACER DEVICE			
220 ml (single patient)	3.65	1	e-chamber Turbo
510 ml (single patient)	5.95	1	e-chamber La Grande
800 ml.....	6.50	1	Volumatic

- Symbols -

Xaluprine 151
8-methoxypsoralen 71

- A -

A-Scabies 68
Abacavir sulphate 104
Abacavir sulphate with
 lamivudine 104
Abacavir/lamivudine Viatris 104
Abciximab 183
Abilify Maintena 135
Abiraterone acetate 172
Acarbose 9
Accarb 9
Acetazolamide 278
Acetec 44
Acetic acid
 Extemporaneously Compounded
 Preparations 289
 Genito-Urinary 73
Acetic acid with hydroxyquinoline,
 glycerol and ricinoleic acid 73
Acetic acid with propylene
 glycol 280
Acetylcholine chloride 278
Acetylcysteine 281
Aciclovir
 Infections 107
 Sensory 274
Aciclovir-Baxter 107
Acid Citrate Dextrose A 36
Acidex 5
Acipimox 53
Acitretin 71
Act-HIB 311
Actemra 236
Actinomycin D 149
Adalimumab (Amgevita) 183
Adalimumab (Humira - alternative
 brand) 192
Adapalene 68
Adcetris 203
Adenosine 46
Adenosine Baxter 46
Adrenaline
 Cardiovascular 54
 Respiratory 264
Adsine 46
Advantan 70
Advate 35
Adynovate 35
Aerrane 121
Afinitor 258
Aflibercept 199

Agents Affecting the
 Renin-Angiotensin System 44
Agents for Parkinsonism and Related
 Disorders 120
Agents Used in the Treatment of
 Poisonings 281
Ajmaline 46
Albalon 276
Albendazole 101
Alchemy Caspofungin 99
Alchemy Oxaliplatin 158
Alchemy Oxybutynin 76
Aldurazyme 19
Alecensa 158
Alectinib 158
Alendronate sodium 113
Alendronate sodium with
 colecalciferol 113
Alfacalcidol 28
Alfamino 302
Alfamino Junior 302
Alfentanil 125
Alglucosidase alfa 17
Allegron 128
Allerfix 276
Allerpro Syneo 1 304
Allerpro Syneo 2 304
Allersoothe 265
Allmercap 151
Allopurinol 115
Almarytm 46
Alpha tocopheryl 28
Alpha tocopheryl acetate 29
Alpha-Adrenoceptor Blockers 45
Alphamox 125 92
Alphamox 250 92
Alprolix 34
Alprostadi 54
Alprostadi hydrochloride 55
Alteplase 39
Alum 289
Aluminium chloride 32
Aluminium hydroxide 5
Aluminium hydroxide with
 magnesium hydroxide and
 simeticone 5
Alyacen 73
Amantadine hydrochloride 120
AmBisome 96
Ambrisentan 55
Ambrisentan Viatris 55
Amethocaine
 Nervous 124
 Sensory 277

Amgevita 183
Amikacin 88
Amiloride hydrochloride 50
Amiloride hydrochloride with
 furosemide 50
Amiloride hydrochloride with
 hydrochlorothiazide 50
Aminolevulinic acid
 hydrochloride 175
Aminophylline 271
Amiodarone hydrochloride 46
Amisulpride 134
Amitriptyline 127
Amlodipine 48
Amorolfine 67
Amoxicillin 92
Amoxicillin with clavulanic acid 92
Amoxiclav Devatis Forte 92
Amphotericin B
 Alimentary 25
 Infections 96
Amsacrine 152
Amyl nitrite 281
Anabolic Agents 77
Anaesthetics 121
Anagrelide hydrochloride 152
Analgesics 124
Anastrozole 173
Anatrole 173
Androgen Agonists and
 Antagonists 77
Anoro Ellipta 266
Antabuse 146
Antacids and Antiflatulents 5
Anti-Infective Agents 73
Anti-Infective Preparations
 Dermatological 67
 Sensory 274
Anti-Inflammatory Preparations 275
Antiacne Preparations 68
Antiallergic Preparations 264
Antianaemics 30
Antiarrhythmics 46
Antibacterials 88
Anticholinergic Agents 265
Anticholinesterases 113
Antidepressants 127
Antidiarrhoeals and Intestinal
 Anti-Inflammatory Agents 5
Antiepilepsy Drugs 129
Antifibrinolytics, Haemostatics and
 Local Sclerosants 32
Antifibrotics 267
Antifungals 96

Antihypotensives	47	Hydrochlorothiazide	45	Azopt	278
Antimigraine Preparations	132	Arrow-Norfloxacin	94	AZT	104
Antimycobacterials	99	Arrow-Ornidazole	102	Aztreonam	94
Antinausea and Vertigo Agents	132	Arrow-Quinapril 10.....	44	- B -	
Antiparasitics	101	Arrow-Quinapril 20.....	44	Bacillus calmette-guerin (BCG)	258
Antipruritic Preparations	68	Arrow-Quinapril 5.....	44	Bacillus calmette-guerin vaccine	310
Antipsychotic Agents	134	Arrow-Roxithromycin	92	Baclofen	116
Antiretrovirals.....	102	Arrow-Timolol	278	Bacterial and Viral Vaccines.....	310
Antirheumatoid Agents	113	Arrow-Topiramate.....	131	Bacterial Vaccines	310
Antiseptics and Disinfectants.....	283	Arrow-Tramadol.....	127	Balanced Salt Solution	277
Antispasmodics and Other Agents		Arsenic trioxide	152	Barium sulphate.....	284
Altering Gut Motility	7	Artemether with lumefantrine.....	101	Barrier Creams and Emollients.....	68
Antithrombotics.....	36	Artesunate	101	Basiliximab	200
Antithymocyte globulin		Articaine hydrochloride	122	BCG Vaccine AJV	310
(equine)	257	Articaine hydrochloride with adrenaline.....	122	BD PosiFlush.....	42
Antithymocyte globulin (rabbit).....	257	Asacol.....	6	Beclazone 100	269
Antiulcerants	7	Ascorbic acid		Beclazone 250	269
Antivirals	106	Alimentary	28	Beclazone 50	269
Anxiolytics.....	138	Extemporaneously Compounded Preparations	289	Beclomethasone dipropionate	269
Anzatax.....	170	Aspen Adrenaline	54	Bedaquiline.....	100
Apidra	10	Aspirin		Bee venom	264
Apidra Solostar	10	Blood.....	37	Bendamustine hydrochloride	148
APO Clomipramine.....	127	Nervous.....	124	Bendamustine Sandoz	148
APO Health Macrogol.....	16	Asthalin.....	269	Bendrofluazide.....	50
APO-Atomoxetine.....	143	Atazanavir Mylan	105	Bendroflumethiazide [Bendrofluazide].....	50
APO-Candesartan HCTZ		Atazanavir sulphate	105	Benralizumab.....	201
16/12.5.....	45	Atazanavir Viatrix.....	105	Benzathine benzylpenicillin	92
APO-Candesartan HCTZ		Atenolol.....	47	Benzatropine mesylate	120
32/12.5.....	45	Atenolol Viatrix.....	47	Benzbromaron AL 100.....	115
Apomorphine hydrochloride.....	120	Atenolol-AFT.....	47	Benzbromarone	115
Apraclonidine.....	279	Atazolizumab	246	Benzetacil	92
Aprepitant	132	ATGAM	257	Benzocaine.....	122
Apresoline.....	55	Ativan.....	138	Benzocaine with tetracaine hydrochloride	122
Aprotinin	32	Atomoxetine.....	143	Benzoil	289
Aptamil Feed Thickener.....	294	Atorvastatin.....	51	Benzoyl peroxide	68
Aqueous cream	69	Atovaquone with proguanil hydrochloride	101	Benzotrop	120
Arachis oil [Peanut oil].....	289	Atracurium besylate	116	Benzydamine hydrochloride	25
Aratac	46	Atropine sulphate		Benzydamine hydrochloride with cetylpyridinium chloride	25
Arava	113	Cardiovascular	46	Benzylpenicillin sodium [Penicillin G].....	92
Arginine		Sensory.....	279	Beractant	272
Alimentary	18	Atropt	279	Besponsa.....	213
Various.....	285	Augmentin	92	Beta Cream	70
Arginine2000	298	Aurorix	128	Beta Ointment.....	70
Argipressin [Vasopressin].....	87	Avelox.....	93	Beta Scalp	72
Aripiprazole.....	134-135	Avonex.....	139	Beta-Adrenoceptor Agonists.....	269
Aripiprazole Sandoz	134	Axitinib	158	Beta-Adrenoceptor Blockers.....	47
Aristocort	70	Azaciditidine	150	Beta-hCG low sensitivity urine test kit.....	321
Arrotex-Prazosin S29	45	Azaciditidine Dr Reddy's	150	Betadine	283
Arrow - Clopid.....	37	Azactam	94	Betahistine dihydrochloride	132
Arrow - Lattim	279	Azamun	258	Betaine	18
Arrow-Amitriptyline	127	Azathioprine.....	258		
Arrow-Bendrofluazide	50	Azilect	121		
Arrow-Brimonidine	279	Azithromycin.....	90		
Arrow-Diazepam	138				
Arrow-Fluoxetine	128				
Arrow-Losartan &					

Betamethasone	78	Botulism antitoxin	282	Dermatological	72
Betamethasone dipropionate.....	70	Bplex.....	28	Calcium Homeostasis.....	77
Betamethasone dipropionate with calcipotriol.....	71	Brentuximab vedotin.....	203	Calcium polystyrene sulphonate.....	42
Betamethasone sodium phosphate with betamethasone acetate.....	78	Breo Ellipta	270	Calcium Resonium	42
Betamethasone valerate.....	70, 72	Brevinor 1/28	73	Calogen (neutral).....	292
Betamethasone valerate with clioquinol.....	71	Breztri Aerosphere.....	266	Calogen (strawberry).....	292
Betamethasone valerate with sodium fusidate [Fusidic acid].....	71	Bricanyl Turbuhaler	269	Candesartan cilexetil	44
Betaxolol	278	Brimonidine tartrate	279	Candesartan cilexetil with hydrochlorothiazide	45
Betnovate	70	Brimonidine tartrate with timolol maleate.....	279	Candestar	44
Betoptic.....	278	Brinzolamide.....	278	Capecitabine.....	150
Betoptic S	278	Bromocriptine	120	Capecitabine Viatrix.....	150
Bevacizumab	201	Budesonide		Capsaicin	
Bevacizumab (Ocular).....	203	Alimentary.....	5	Musculoskeletal	119
Bexsero	312	Respiratory.....	265, 269	Nervous.....	124
Bezafibrate	51	Budesonide Te Arai	5	Captopril	44
Bezalip.....	51	Budesonide with eformoterol	270	Carbachol	278
Bezalip Retard.....	51	Budesonide with glycopyrronium and eformoterol	266	Carbamazepine	129
Bicalutamide	172	Bumetanide	49	Carbasorb-X	282
Bicillin LA	92	Bupafen	123	Carbimazole	86
BiCNU.....	149	Bupivacaine hydrochloride	122	Carbomer.....	279
BiCNU S29	149	Bupivacaine hydrochloride with adrenaline	122	Carboplatin	158
Bile and Liver Therapy.....	9	Bupivacaine hydrochloride with fentanyl.....	123	Carboplatin Accord.....	158
Biliscopin	285	Bupivacaine hydrochloride with glucose	123	Carboprost trometamol.....	74
Bimatoprost	278	Buprenorphine Naloxone BNM.....	146	Carboxymethylcellulose Alimentary	25
Binarex	172	Buprenorphine with naloxone	146	Extemporaneously Compounded Preparations	289
Binocrit	30	Bupropion hydrochloride.....	146	Cardinol LA	48
Biocon	47	Burinex	49	Cardizem CD.....	49
Biodone	126	Buserelin.....	82	CareSens Dual	321
Biodone Extra Forte.....	126	Buspirone hydrochloride.....	138	CareSens N.....	321
Biodone Forte.....	126	Buspirone Viatrix	138	CareSens N POP	321
Biotin.....	18	Busulfan.....	149	CareSens N Premier	321
Bisacodyl	16			CareSens PRO.....	321
Bisacodyl Viatrix	16	- C -		Carglumic acid.....	18
Bismuth subgallate	289	Cabergoline	81	Carbomethoxymethylcellulose sodium with pectin and gelatine	
Bismuth subnitrate and iodoform paraffin.....	287	Caffeine	143	Alimentary	25
Bisoprolol fumarate.....	47	Caffeine citrate	271	Sensory.....	279
Bivalirudin.....	36	Calamine	68	Carmustine	149
Bleomycin sulphate	149	Calci-Tab 500	23	Carvedilol.....	47
Blood glucose diagnostic test meter	321	Calcipotriol.....	71	Carvedilol Sandoz	47
Blood glucose diagnostic test strip.....	321	Calcitonin.....	77	Casporfungin	99
Blood ketone diagnostic test strip.....	321	Calcitriol.....	28	Catapres	49
Bonney's blue dye	286	Calcitriol XL	28	Cefaclor	89
Boostrix.....	311	Calcitriol-AFT.....	28	Cefalexin.....	89
Boric acid.....	289	Calcium carbonate.....	5, 23	Cefalexin Sandoz	89
Bortezomib	152	Calcium carbonate PAI.....	5	Cefazolin.....	89
Bosentan	58	Calcium Channel Blockers	48	Cefazolin-AFT.....	89
Bosentan Dr Reddy's.....	58	Calcium chloride	40	Cefepime	90
Botox	116	Calcium folinate	171	Cefepime-AFT	90
		Calcium Folate Ebewe	171	Cefotaxime	89
		Calcium Folate Sandoz.....	171	Cefotaxime Sandoz	89
		Calcium gluconate		Cefoxitin.....	89
		Blood.....	40	Ceftaroline fosamil.....	90

Ceftazidime.....	89	Ciclosporin.....	175	toxin.....	116
Ceftazidime Kabi.....	89	Cidofovir.....	107	Clotrimazole.....	
Ceftazidime with avibactam.....	89	Cidomycin P/Free.....	88	Dermatological.....	67
Ceftriaxone.....	90	Cilicaine VK.....	93	Genito-Urinary.....	73
Ceftriaxone-AFT.....	90	Cimetidine.....	8	Clove oil.....	289
Cefuroxime.....	89	Cinacalcet.....	77	Clozapine.....	134
Cefuroxime Devatis.....	89	Cinacalcet Devatis.....	77	Clozaril.....	134
Celapram.....	128	Cinchocaine hydrochloride with		Clustran.....	132
Celebrex.....	117	hydrocortisone.....	7	Co-trimoxazole.....	96
Celecoxib.....	117	Ciprofloxacin.....		Coal tar.....	289
Celecoxib Pfizer.....	117	Infections.....	93	Coal tar with salicylic acid and	
Celiprolol.....	47	Sensory.....	274	sulphur.....	71
CellCept.....	258	Ciprofloxacin Kabi.....	93	Cocaine hydrochloride.....	123
Centrally-Acting Agents.....	49	Ciprofloxacin Teva.....	274	Cocaine hydrochloride with	
Cephalexin ABM.....	89	Ciprofloxacin with		adrenaline.....	123
Cerazette.....	74	hydrocortisone.....	274	Codeine phosphate.....	
Cerobact.....	92	Ciproxin HC Otic.....	274	Extemporaneously Compounded	
Cetirizine hydrochloride.....	265	Cisplatin.....	158	Preparations.....	289
Cetomacrogol.....	69	Cisplatin Accord.....	158	Nervous.....	125
Cetomacrogol Cream AFT.....	69	Citalopram hydrobromide.....	128	Coenzyme Q10.....	18
Cetomacrogol with glycerol.....	69	Citanest.....	124	Colchicine.....	116
Cetomacrogol-AFT.....	69	Citrate sodium.....	36	Colecalciferol.....	28
Cetrimide.....	289	Citric acid.....	289	Colestimethate.....	95
Cetuximab.....	204	Citric acid with magnesium carbonate		Colestipol hydrochloride.....	52
Champix.....	147	hydrate and sodium		Colestyramine.....	52
Charcoal.....	282	picosulfate.....	15	Colestyramine - Mylan.....	52
CheckTop.....	321	Citric acid with sodium		Colgout.....	116
Chemotherapeutic Agents.....	148	bicarbonate.....	284	Colifoam.....	6
Chickenpox vaccine.....	319	Citrulline 1000.....	299	Colistin sulphomethate	
Chloral hydrate.....	141	Cladribine.....	150	[Colestimethate].....	95
Chlorambucil.....	149	Clarithromycin.....	91	Collodion flexible.....	289
Chloramphenicol.....		Clexane.....	36	Colloidal bismuth subcitrate.....	9
Infections.....	95	Clexane Forte.....	36	Colofac.....	7
Sensory.....	274	Clindamycin.....	95	Colomycin.....	95
Chlorhexidine.....	283	Clinicians.....	28	Colony-Stimulating Factors.....	39
Chlorhexidine gluconate		Clinicians Multivit & Mineral		Coloxyl.....	15
Alimentary.....	25	Boost.....	26	Combigan.....	279
Extemporaneously Compounded		Clinicians Renal Vit.....	26	Comirnaty Omicron (JN.1).....	315
Preparations.....	289	Clobazam.....	129	Compound electrolytes.....	40, 42
Genito-Urinary.....	73	Clobetasol propionate.....	70, 72	Compound electrolytes with glucose	
Chlorhexidine with		Clobetasone butyrate.....	70	[Dextrose].....	40, 42
cetrimide.....	283, 286	Clofazimine.....	99	Compound hydroxybenzoate.....	289
Chlorhexidine with ethanol.....	283	Clomazol.....		Compound sodium lactate	
Chloroform.....	289	Dermatological.....	67	[Hartmann's solution].....	41
Chloroquine phosphate.....	101	Genito-Urinary.....	73	Concerta.....	144
Chlorothiazide.....	50	Clomifene citrate.....	81	Condyline.....	72
Chlorpheniramine maleate.....	265	Clomipramine hydrochloride.....	127	Contraceptives.....	73
Chlorpromazine hydrochloride.....	134	Clomipramine Teva.....	127	Contrast Media.....	284
Chlorsig.....	274	Clonazepam.....	129, 138	Copaxone.....	139
Chlortalidone [Chlorthalidone].....	50	Clonidine.....	49	Copper.....	23
Chlorthalidone.....	50	Clonidine hydrochloride.....	49	Copper chloride.....	23
Choice 380 7med Nsha Silver/copper		Clonidine Teva.....	49	Corticorelin (ovine).....	81
Short.....	74	Clopidogrel.....	37	Corticosteroids.....	
Cholestyramine.....	52	Clopine.....	134	Dermatological.....	70
Choriogonadotropin alfa.....	82	Clopirol.....	135, 137	Hormone Preparations.....	78
Ciclopirox olamine.....	67	Clostridium botulinum type A		Cosentyx.....	233

Cosmegen	149	DBL Amikacin	88	Dexrazoxane	171
Coversyl	44	DBL Aminophylline	271	Dextrose	
COVID-19 vaccine	315	DBL Bleomycin Sulfate	149	Alimentary	9
Creon 10000	14	DBL Bortezomib	152	Blood	40–42
Creon 25000	14	DBL Carboplatin	158	Extemporaneously Compounded	
Creon Micro	14	DBL Cefotaxime	89	Preparations	289
Crizotinib	159	DBL Dacarbazine	152	Dextrose with sodium citrate and	
Crotamiton	68	DBL Desferrioxamine Mesylate for Inj		citric acid [Acid Citrate Dextrose	
Crystaderm	67	BP	283	A]	36
Cu 375 Standard	74	DBL Docetaxel	170	DHC Continus	125
Curam Duo 500/125	92	DBL Ergometrine	74	Diabetes	9
Curosurf	272	DBL Gemcitabine	151	Diacomit	131
Cvite	28	DBL Gentamicin	88	Diagnostic Agents	
Cyclizine hydrochloride	132	DBL Leucovorin Calcium	171	Vaccines	320
Cyclizine lactate	133	DBL Methotrexate Onco-Vial	151	Various	285
Cyclogyl	279	DBL Naloxone Hydrochloride	281	Diagnostic and Surgical	
Cyclonex	149	DBL Pethidine Hydrochloride	127	Preparations	276
Cyclopentolate hydrochloride	279	DBL Vincristine Sulfate	171	Diamide Relief	5
Cyclophosphamide	149	Decongestants	269	Diasip (strawberry)	299
Cycloserine	100	Decongestants and		Diasip (vanilla)	299
Cymevene	107	Antiallergics	276	Diatrizoate meglumine with sodium	
Cyproheptadine hydrochloride	265	Decozol	25	amidotrizoate	284
Cyproterone acetate	77	Deferasirox	282	Diatrizoate sodium	284
Cyproterone acetate with		Deferiprone	282	Diazepam	129, 138
ethinyloestradiol	73	Defibrotide	36	Diazoxide	
Cystadane	18	Definity	285	Alimentary	9
Cysteamine hydrochloride	289	Demeclocycline hydrochloride	94	Cardiovascular	55
Cytarabine	150	Denosumab	114	Dichlorobenzyl alcohol with	
Cytotec	7	Deolate	99	amylmetacresol	25
- D -					
D-Penamine	113	Deoxycoformycin	155	Diclofenac Devatis	276
Dabigatran	36	Depo-Medrol	79	Diclofenac Sandoz	118
Dabrafenib	159	Depo-Provera	74	Diclofenac sodium	
Dacarbazine	152	Depo-Testosterone	77	Musculoskeletal	118
Dactinomycin [Actinomycin D]	149	Deprim	96	Sensory	276
Daivobet	71	Dermol	70, 72	Dicobalt edetate	283
Daivonex	71	Desferrioxamine mesilate	283	Diffucan	97
Dalacin C	95	Desflurane	121	Diffucortolone valerate	70
Danaparoid	36	Desmopressin	87	Digestives Including Enzymes	14
Dantrium	117	Desmopressin acetate	87	Digoxin	46
Dantrium IV	117	Desmopressin-PH&T	87	Digoxin immune Fab	281
Dantrolene	117	Desogestrel	74	Dihydrocodeine tartrate	125
Daonil	11	Dexamethasone		Dihydroergotamine mesylate	132
Dapa-Tabs	50	Hormone Preparations	78	Diltiazem CD Clinect	49
Dapsone	99	Sensory	275	Diltiazem hydrochloride	49
Daptomycin	95	Dexamethasone phosphate	79	Dimercaprol	283
Daptomycin Dr Reddy's	95	Dexamethasone with framycetin and		Dimercaptosuccinic acid	283
Darunavir	105	gramicidin	274	Dimethicone	67–68
Darunavir Viatrix	105	Dexamethasone with neomycin		Dimethyl fumarate	139
Dasatinib	161	sulphate and polymyxin B		Dimethyl sulfoxide	287
Dasatinib-Teva	161	sulphate	274	Dinoprostone	74
Daurorubicin	150	Dexamethasone with		Dipentum	7
David One Step Cassette Pregnancy		tobramycin	274	Diphenamil metilsulfate	72
Test	321	Dexamfetamine sulfate	143	Diphenoxylate hydrochloride with	
DBL Acetylcysteine	281	Dexmedetomidine	121	atropine sulphate	5
DBL Adrenaline	54	Dexmedetomidine Viatrix	121	Diphtheria antitoxin	282
		Dexmethsone	78	Diphtheria, tetanus and pertussis	

vaccine 311	Droperidol Panpharma 133	Empagliflozin with metformin hydrochloride 13
Diphtheria, tetanus, pertussis and polio vaccine 310	Drugs Affecting Bone Metabolism 113	Emsogen 299
Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine 310	Dual blood glucose and blood ketone diagnostic test meter 321	Emtricitabine 104
Diprosone 70	Dulaglutide 11	Emtricitabine with tenofovir disoproxil 108
Dipyridamole 37	Dulcolax SP Drop 16	Emtriva 104
Dipyridamole - Strides 37	Duocal Super Soluble Powder 293	Emulsifying ointment 69
Disodium edetate 277	Duolin 266	Enalapril maleate 44
Disodium hydrogen phosphate with sodium dihydrogen phosphate 289	DuoResp Spiromax 270	Enbrel 175
Disopyramide phosphate 46	Duovisc 277	Endocrine Therapy 172
Disulfiram 146	Duride 53	Endoxan 149
Dithranol 289	Durvalumab 247	Energivit 297
Diuretics 49	Dynastat 118	Engerix-B 315-316
Dobutamine 54	Dysport 116	Enhertu 241
Dobutamine-hameln 54	- E -	Enlax XR 128
Docetaxel 170	e-chamber La Grande 321	Enoxaparin sodium 36
Docusate sodium	e-chamber Mask 321	Enstilar 71
Alimentary 15	e-chamber Turbo 321	Ensure (Chocolate) 308
Sensory 280	E-Mycin 92	Ensure (Vanilla) 308
Docusate sodium with sennosides 15	E-Z-Gas II 284	Ensure Plus (Banana) 309
Dolutegravir 106	Easiphen Liquid 297	Ensure Plus (Chocolate) 309
Dolutegravir with lamivudine 106	Econazole nitrate 67	Ensure Plus (Fruit of the forest) 309
Domperidone 133	Edrophonium chloride 113	Ensure Plus (Vanilla) 309
Domperidone Viatrix 133	Efavirenz 103	Ensure Plus HN 308
Donepezil hydrochloride 145	Efavirenz Milpharm 103	Ensure Plus HN RTH 308
Dopamine Basi 54	Efavirenz with emtricitabine and tenofovir disoproxil 104	Ensure Two Cal HN RTH 301
Dopamine hydrochloride 54	Eformoterol fumarate 270	Entacapone 120
Dornase alfa 271	Eformoterol fumarate dihydrate 270	Entacapone Viatrix 120
Dortimopt 278	Eftrenonacog alfa [Recombinant factor IX] 34	Entecavir 106
Dorzolamide 278	Efudix 72	Entecavir (Rex) 106
Dorzolamide with timolol 278	Elaprase 19	Entresto 24/26 45
Dostinex 81	Elecare (Unflavoured) 302	Entresto 49/51 45
Dosulepin [Dothiepin] hydrochloride 127	Elecare (Vanilla) 302	Entresto 97/103 45
Dosulepin Viatrix 127	Elecare LCP (Unflavoured) 302	Entyvio 244
Dotarem 285	Electral 42	Enzymes 115
Dothiepin 127	Electrolytes 288	Ephedrine 54
Dovato 106	Elelyso 21	Ephedrine Aguetant 54
Doxapram 273	Elemental 028 Extra (grapefruit) 300	Ephedrine Juno 54
Doxazosin 45	Elemental 028 Extra (pineapple & orange) 300	Epilim IV 131
Doxazosin Clinect 45	Elemental 028 Extra (summer fruits) 300	Epipen 264
Doxepin hydrochloride 127	Elexacafter with tezacaftor, ivacaftor and ivacaftor 271	Epipen Jr 264
Doxine 94	Elidel 71	Epirubicin Ebewe 150
Doxorubicin Ebewe 150	Elocon 70	Epirubicin hydrochloride 150
Doxorubicin hydrochloride 150	Elocon Alcohol Free 70	Eplerenone 50
Doxycycline 94	Eltrombopag 32	Epoetin alfa 30
DP Lotn HC 70	Emend Tri-Pack 132	Epoetin beta 31
DP-Captopril 44	Emicizumab 33	Epoprostenol 62
Dr Reddy's Omeprazole 8	EMLA 124	Eptacog alfa [Recombinant factor VIIa] 34
Drofate 48	Empagliflozin 13	Eptifibatide 38
Droperidol 133		Eptifibatide Viatrix 38
		Eributux 204
		Ergometrine maleate 74
		Erlotinib 161

Ertapenem	88	Faslodex	172	Fluids and Electrolytes	40
Erythrocin IV	92	Fatty Emulsion Cream (Evara)	69	Flumazenil	281
Erythromycin (as ethylsuccinate)	92	Febuxostat	116	Flumazenil-Baxter	281
Erythromycin (as lactobionate)	92	Febuxostat (Teva)	116	Flumetasone pivalate with clioquinol	275
Erythromycin (as stearate)	92	FEIBA NF	34	Fluocortolone caproate with fluocortolone pivalate and cinchocaine	7
Esbriet	268	Felo 10 ER	48	Fluorescein sodium	276
Escitalopram	128	Felo 5 ER	48	Fluorescein sodium with lignocaine hydrochloride	276
Esmolol hydrochloride	47	Felodipine	48	Fluorescein sodium with lignocaine hydrochloride	276
Essential Amino Acid Mix	299	Fentanyl	125	Fluorometholone	276
Estradiol TDP Mylan	80	Fentanyl Sandoz	125	Fluorouracil	151
Estradot	80	Ferinject	24	Fluorouracil Accord	151
Estrogeol	80	Ferodan	24	Fluorouracil sodium	72
Etanercept	175	Ferric subsulfate	33	Fluox	128
Ethambutol hydrochloride	100	Ferriprox	282	Fluoxetine hydrochloride	128
Ethanol	281	Ferro-F-Tabs	24	Flupenthixol decanoate	136
Ethanol with glucose	281	Ferro-Liquid	24	Flutamide	172
Ethanol, dehydrated	281	Ferro-tab	23	Flutamin	172
Ethics Aspirin	124	Ferrograd	24	Fluticasone	270
Ethics Aspirin EC	37	Ferrosig	24	Fluticasone furoate with umecclidinium and vilanterol	267
Ethinylestradiol with desogestrel	73	Ferrous fumarate	23	Fluticasone furoate with vilanterol	270
Ethinylestradiol with levonorgestrel	73	Ferrous fumarate with folic acid	24	Fluticasone propionate	265
Ethinylestradiol with norethisterone	73	Ferrous gluconate with ascorbic acid	24	Fluticasone with salmeterol	270
Ethosuximide	129	Ferrous sulfate	24	Flynn	89
Ethyl chloride	123	Ferrous sulfate with ascorbic acid	24	FML	276
Etomidate	121	Fexaclear	265	Foban	67
Etopophos	152	Fexofenadine hydrochloride	265	Folic acid	31
Etoposide	152	Filgrastim	40	Folic Acid multicchem	31
Etoposide (as phosphate)	152	Finasteride	75	Folic Acid Viatrix	31
Etoricoxib	118	Fingolimod	139	Fondaparinux sodium	36
Etravirine	103	Flagyl	102	Food Modules	292
Eurofolic	171	Flagyl-S	102	Food/Fluid Thickeners	293
Evara	68-69	Flamazine	67	Fortijuice (Apple)	309
Evara Emulsifying Ointment	69	Flecainide acetate	46	Fortijuice (Orange)	309
EVARA White Soft Paraffin	69	Flecainide BNM	46	Fortijuice (Strawberry)	309
Everet	130	Flecainide Controlled Release Teva	46	Fortini (Strawberry)	306
Everolimus	258	Fleet Phosphate Enema	16	Fortini (Vanilla)	306
Evista	114	Flixonase Hayfever & Allergy	265	Fortini Multi Fibre (chocolate)	306
Evrysdi	142	Flixotide	270	Fortini Multi Fibre (strawberry)	306
Exemestane	173	Flixotide Accuhaler	270	Fortini Multi Fibre (unflavoured)	306
Exjade	282	Florinef	79	Fortini Multi Fibre (vanilla)	306
Extemporaneously Compounded Preparations	289	Fluanxol	136	Fortisip (Banana)	309
Eylea	199	Flucil	92	Fortisip (Chocolate)	309
Ezetimibe	53	Flucloxacillin	92	Fortisip (Strawberry)	309
Ezetimibe Sandoz	53	Flucloxacillin	92	Fortisip (Vanilla)	309
Ezetimibe with simvastatin	53	Fluconazole	97	Fortisip Multi Fibre (chocolate)	309
- F -					
Factor eight inhibitor bypassing fraction	34	Fluconazole-Baxter	97	Fortisip Multi Fibre (strawberry)	309
Famotidine	8	Flucytosine	99	Fortisip Multi Fibre (vanilla)	309
Famotidine Hovid MY	8	Fludara Oral	151	Fosamax	113
Fasenra	201	Fludarabine Ebewe	151		
		Fludarabine phosphate	151		
		Fludarabine Sagent	151		
		Fludrocortisone acetate	79		

Fosamax Plus.....	113	Glucagon hydrochloride.....	9	HCU Lophlex LQ	295
Foscarnet sodium.....	107	Glucerna Select	299	Healon	277
Fosfomycin.....	95	Glucose [Dextrose]		Healon 5	277
Framycetin sulphate.....	274	Alimentary	9	Healon GV	277
Fresofol 1% MCT/LCT.....	122	Blood.....	41	Healon GV Pro	277
Frusemide.....	50	Extemporaneously Compounded		healthE Calamine Aqueous.....	68
Fucidin	95	Preparations	289	healthE Dimethicone 10%.....	68
Fucithalmic	274	Glucose with potassium chloride.....	41	healthE Dimethicone 4% Lotion	67
Fulvestrant.....	172	Glucose with potassium chloride and		healthE Dimethicone 5%.....	68
Fungilin.....	25	sodium chloride.....	41	healthE Glycerol BP Liquid.....	290
Furosemide [Frusemide].....	50	Glucose with sodium chloride.....	41	healthE Urea Cream.....	70
Furosemide-Baxter	50	Glucose with sucrose and		Hemilibra	33
Fusidic acid		fructose.....	9	Heparin sodium	37
Dermatological.....	67, 71	Glycerin with sodium saccharin	290	Heparin Sodium Panpharma	37
Infections.....	95	Glycerin with sucrose	290	Heparinised saline	37
Sensory.....	274	Glycerol		Heparon Junior	301
- G -		Alimentary	16	Hepatitis A vaccine	315
GA Explore 5	294	Extemporaneously Compounded		Hepatitis B recombinant	
GA1 Anamix Infant	294	Preparations	290	vaccine	315
GA1 Anamix Junior.....	294	Glycerol with paraffin	69	Herzuma	240
Gabapentin.....	129	Glycerol trinitrate		Hikma	46
Gacet.....	125	Alimentary	7	Hikma Acetylcysteine	281
Gadobutrol.....	285	Cardiovascular	53	Hiprex	95
Gadoteric acid	285	Glycine.....	286	Histaclear.....	265
Gadovist 1.0	285	Glycoprep Orange	15	Histamine acid phosphate	285
Gadoxetate disodium.....	285	Glycopyrronium	266	Holoxan	149
Galsulfase.....	19	Glycopyrronium bromide	7	Hormone Replacement Therapy	80
Galvumet.....	11	Glycopyrronium with		HPV	316
Galvus	11	indacaterol	266	Humalog Mix 25.....	10
Ganciclovir.....	107	Glycosade.....	298	Humalog Mix 50.....	10
Gardasil 9	316	Glytactin Bettermilk.....	297	Human Milk Fortifier	293
Gastrodenol	9	Gonadorelin	82	Human papillomavirus (6, 11, 16, 18,	
Gastrografin.....	284	Goserelin	82	31, 33, 45, 52 and 58) vaccine	
Gazyva	215	Granisetron	133	[HPV]	316
Gefitinib	162	GTO Oil	298	Humatin	88
Gelatine, succinylated	43	- H -		Humira	192
Gelofusine	43	Habitrol	147	HumiraPen.....	192
Gemcitabine Hydrochloride	151	Habitrol (Fruit).....	147	Hyaluronic acid	
Gemtuzumab ozogamicin.....	204	Habitrol (Mint).....	147	Alimentary.....	26
Gentamicin Amdipharm	88	Haem arginate.....	19	Sensory.....	277, 280
Gentamicin Noridem.....	88	Haemophilus influenzae type B		Hyaluronic acid with lidocaine	
Gentamicin sulphate		vaccine	311	[lignocaine]	26
Infections.....	88	Haldol	136	Hyaluronidase	115
Sensory.....	274	Haldol Concentrate.....	136	Hydralazine hydrochloride.....	55
Gestrinone.....	81	Haloperidol	134	Hydralyte - Lemonade	42
Gilenya	139	Haloperidol decanoate.....	136	Hydrocortisone	
Ginet.....	73	Hartmann's solution	41	Dermatological.....	70
Glatiramer acetate	139	Harvoni	106	Extemporaneously Compounded	
Glaucoma Preparations.....	278	Havrix 1440	315	Preparations	290
Glecaprevir with pibrentasvir	106	Havrix Junior.....	315	Hormone Preparations.....	79
Glibenclamide.....	11	Haylor Syrup.....	265	Hydrocortisone acetate.....	6
Gliclazide	11	HCU Anamix Infant.....	295	Hydrocortisone acetate with	
Glolan	175	HCU Anamix Junior	295	pramoxine hydrochloride	6
Glipizide.....	11	HCU Anamix Junior LQ	295	Hydrocortisone and paraffin liquid	
Glizide.....	11	HCU Explore 5.....	295	and lanolin	70
Glucagen Hypokit	9	HCU Express 15.....	295	Hydrocortisone butyrate	70, 72

Hydrocortisone with miconazole.....	71	Infanrix IPV.....	310	Irinotecan hydrochloride.....	153
Hydrocortisone with natamycin and neomycin.....	71	Infanrix-hexa.....	310	Iron (as ferric carboxymaltose).....	24
Hydrogen peroxide.....	67	Infatrin.....	305	Iron (as sucrose).....	24
Hydroxocobalamin.....		Infliximab.....	205	Iron polymaltose.....	24
Alimentary.....	27	Influenza vaccine.....	317	Irrigation Solutions.....	286
Various.....	281	Influvac Tetra (2025 formulation).....	317	Isentress.....	106
Hydroxocobalamin Panpharma.....	27	Inhaled Corticosteroids.....	269	Isentress HD.....	106
hydroxycarbamide.....	152	Inlyta.....	158	Ismo 20.....	53
Hydroxychloroquine sulphate.....	113	Inotuzumab ozogamicin.....	213	Ismo 40 Retard.....	53
Hydroxyurea.....		Inresa.....	24	Isoflurane.....	121
[hydroxycarbamide].....	152	Inspra.....	50	Isoleucine50.....	299
Hygroton.....	50	Instillagel Lido.....	123	Isoniazid.....	100
Hylo-Fresh.....	280	Insulin aspart.....	10	Isoniazid Teva.....	100
Hyoscine butylbromide.....	7	Insulin aspart with insulin aspart protamine.....	10	Isoniazid with rifampicin.....	100
Hyoscine Butylbromide (Adiramedita).....	7	Insulin degludec with insulin aspart.....	10	Isoprenaline [Isoproterenol].....	54
Hyoscine hydrobromide.....	133	Insulin glargine.....	10	Isopropyl alcohol.....	283
Hyperuricaemia and Antigout.....	115	Insulin glulisine.....	10	Isoproterenol.....	54
HypoPak Glucose.....	9	Insulin isophane.....	10	Isoptin.....	49
Hypromellose.....	277, 279	Insulin lispro.....	10	Isoptin SR.....	49
Hypromellose with dextran.....	280	Insulin lispro with insulin lispro protamine.....	10	Isopto Carpine.....	278
- I -.....		Insulin neutral.....	10	Isosorbide mononitrate.....	53
Ibiamox.....	92	Insulin neutral with insulin isophane.....	10	Isotretinoin.....	68
Ibrance.....	165	Intelence.....	103	Ispaghula (psyllium) husk.....	15
Ibrutinib.....	152	Interferon alfa-2b.....	109	Isradipine.....	48
Ibuprofen.....	118	Interferon beta-1-alpha.....	139	Itch-Soothe.....	68
Ibuprofen SR BNM.....	118	Interferon beta-1-beta.....	139	Itracap.....	97
Icatibant.....	264	Interferon gamma.....	110	Itraconazole.....	97
Idarubicin hydrochloride.....	150	Intra-uterine device.....	74	Itraconazole Cresent.....	97
Idarucizumab.....	34	Invanz.....	88	Itrazole.....	97
Idursulfase.....	19	Invega Sustenna.....	136	Ivabradine.....	46
Ifosfamide.....	149	Invega Trinza.....	137	Ivacaftor.....	272
Ilomedin.....	63	Iodine.....	86	Ivermectin.....	101
Iloprost.....	63	Iodine with ethanol.....	283	- J -.....	
Imaging Agents.....	175	Iodised oil.....	284	Jadelle.....	74
Imatinib mesilate.....	162	Iodixanol.....	284	Jakavi.....	167
Imatinib-Rex.....	162	Iohexol.....	284	Jardiamet.....	13
Imbruvica.....	152	Iopidine.....	279	Jardiance.....	13
Imfinzi.....	247	Ioscan.....	284	Jaydess.....	74
Imipenem with cilastatin.....	89	Ipc-a-Allopurinol.....	115	Jevity HiCal RTH.....	308
Imipenem+Cilastatin RBX.....	89	Ipc-a-Bisoprolol.....	47	Jevity RTH.....	308
Imipramine Crescent.....	127	Ipc-a-Ciprofloxacin.....	93	Jinarc.....	51
Imipramine hydrochloride.....	127	Ipc-a-Donepezil.....	145	Juno.....	46
Imiquimod.....	72	Ipc-a-Escitalopram.....	128	- K -.....	
Immune Modulators.....	109	IPCA-Frusemide.....	50	Kadcyla.....	242
Immunosuppressants.....	175	Ipc-a-Hydroxychloroquine.....	113	Kalydeco.....	272
Impact Advanced Recovery.....	307	Ipc-a-Metoprolol.....	48	Kenacomb.....	275
Incruse Ellipta.....	266	IPCA-Propranolol.....	48	Kenacort-A 10.....	79
Indacaterol.....	270	Ipilimumab.....	248	Kenacort-A 40.....	79
Indapamide.....	50	IPOL.....	318	Kenalog in Orabase.....	25
Indigo carmine.....	286	Ipratropium bromide.....	265	Ketalar.....	122
Indinavir.....	105	Iressa.....	162	Ketamine.....	122
Indocyanine green.....	286			Ketamine-Baxter.....	122
Indometacin [Indomethacin].....	118			Ketocal 3:1 (Unflavoured).....	305
Indomethacin.....	118			Ketocal 4:1 (Unflavoured).....	305
				Ketocal 4:1 (Vanilla).....	305

Ketoconazole		Leukotriene Receptor		Locoid Lipocream	70
Dermatological	67	Antagonists	270	Lodoxamide	276
Infections	96	Leuporelin acetate	82	Logem	130
Ketoprofen	118	Leustatin	150	Lomide	276
Ketorolac trometamol	276	Levetiracetam	130	Lomustine	149
KetoSens	321	Levetiracetam-AFT	130	Long-Acting Beta-Adrenoceptor	
Ketostix	321	Levocabastine	276	Agonists	270
Keytruda	250	Levocarnitine	20	Loniten	55
Kindergen	306	Levodopa with benserazide	121	Loperamide hydrochloride	5
Kisqali	166	Levodopa with carbidopa	121	Lopinavir with ritonavir	105
Klacid	91	Levodopa with carbidopa and		Lopinavir/Ritonavir Mylan	105
Klacid IV	91	entacapone	121	Lorafax	265
Klaricid	91	Levomepromazine	134	Loratadine	265
Kogenate FS	35	Levomepromazine		Lorazepam	129, 138
Konakion MM	35	hydrochloride	134	Lormetazepam	141
Konakion MM Paediatric	35	Levonorgestrel	74	Lorstat	51
Konsyl-D	15	Levonorgestrel BNM	74	Losartan Actavis	44
Kuvan	20	Levosimendan	53	Losartan potassium	44
- L -		Levothyroxine	86	Losartan potassium with	
L-ornithine L-aspartate	9	Lidocaine [Lignocaine]	123	hydrochlorothiazide	45
Labetalol	47	Lidocaine [Lignocaine]		Lovir	107
Lacosamide	129	hydrochloride	123	Loxamine	128
Lactose	290	Lidocaine [Lignocaine] hydrochloride		Lucrin Depot 1-month	82
Lactulose	16	with adrenaline	123	Lucrin Depot 3-month	82
Laevolac	16	Lidocaine [Lignocaine] hydrochloride		LumaCina	286
Lamictal	130	with adrenaline and tetracaine		Lumigan	278
Lamivudine	104, 106	hydrochloride	123	Lyllana	80
Lamivudine Viatris	104	Lidocaine [Lignocaine] hydrochloride		Lynparza	154
Lamivudine/Zidovudine Viatris	104	with phenylephrine		Lyrica	130
Lamotrigine	130	hydrochloride	124	Lysine acetylsalicylate [Lysine	
Lanoxin	46	Lidocaine [Lignocaine] with		asprin]	38
Lanoxin PG	46	prilocaine	124	Lysine asprin	38
Lanreotide	174	Lidocaine-Baxter	123	- M -	
Lansoprazole	8	lignocaine		m-Eslon	126
Lantus	10	Alimentary	26	Mabthera	220
Lantus SoloStar	10	Nervous	123-124	Macrobid	95
Lanzol Relief	8	Lincomycin	95	Macrogl 3350 with ascorbic acid,	
Lapatinib	162	Linezolid	95	potassium chloride, sodium	
Largactil	134	Linezolid Kabi	95	chloride and citric acid with	
Laronidase	19	Lioresal Intrathecal	116	magnesium carbonate hydrate	
Lasix	50	Liothyronine sodium	86	and sodium picosulfate	15
Latanoprost	278	Lipid-Modifying Agents	51	Macrogl 3350 with potassium	
Latanoprost with timolol	279	Lipiodol Ultra Fluid	284	chloride and sodium chloride	15
Lax-Suppositories	16	Liquigen	293	Macrogl 3350 with potassium	
Lax-suppositories Glycerol	16	Liraglutide	12	chloride and sodium chloride with/	
Laxatives	15	Lisdexamfetamine dimesilate	143	without sodium sulfate, sodium	
Laxsol	15	Lisinopril	44	ascorbate, ascorbic acid	15
Ledipasvir with sofosbuvir	106	Lissamine green	276	Macrogl 3350 with potassium	
Leflunomide	113	Lithium carbonate	134	chloride, sodium bicarbonate and	
Lenalidomide (Viatris)	153	LMX4	123	sodium chloride	16
Lenalidomide Viatris	153	Lo-Oralcon 20 ED	73	Madopar 125	121
Lenvatinib	162	Local Preparations for Anal and		Madopar 250	121
Lenvima	162	Rectal Disorders	7	Madopar 62.5	121
Letrole	173	Locasol	304	Madopar HBS	121
Letrozole	173	Locoid	70, 72	Madopar Rapid	121
Leucine100	299	Locoid Crelo	70	Mafenide acetate	67

Magnesium amino acid chelate.....	24	vaccine.....	312	Metoclopramide Actavis 10.....	133
Magnesium chloride.....	24	MenQuadfi.....	311	Metoclopramide hydrochloride.....	133
Magnesium hydroxide		Menthol.....	290	Metoclopramide hydrochloride with	
Alimentary.....	24	Mepivacaine hydrochloride.....	124	paracetamol.....	132
Extemporaneously Compounded		Mepivacaine hydrochloride with		Metolazone.....	51
Preparations.....	290	adrenaline.....	124	Metoprolol IV Mylan.....	48
Magnesium oxide.....	24	Mepolizumab.....	214	Metoprolol IV Viatris.....	48
Magnesium oxide with magnesium		Mercaptopurine.....	151	Metoprolol succinate.....	47
aspartate, magnesium amino acid		Meropenem.....	89	Metoprolol tartrate.....	48
chelate and magnesium		Meropenem-AFT.....	89	Metronidamed.....	102
citrate.....	24	Mesalazine.....	6	Metronidazole	
Magnesium sulphate.....	24	Mesna.....	171	Dermatological.....	67
Magnevist.....	285	Mestinon.....	113	Infections.....	102
Malarone.....	101	Metabolic Disorder Agents.....	17	Metyrapone.....	81
Malarone Junior.....	101	Metabolic Products.....	294	Mexiletine hydrochloride.....	47
Malathion [Maldison].....	68	Metaraminol.....	54	Miacalcic.....	77
Maldison.....	68	Metformin hydrochloride.....	11	Mianserin hydrochloride.....	128
Mannitol		Metformin Viatris.....	11	Micolette.....	16
Cardiovascular.....	50	Methacholine chloride.....	286	Miconazole.....	25
Various.....	285	Methadone BNM.....	126	Miconazole nitrate	
Mantoux.....	320	Methadone hydrochloride		Dermatological.....	67
Maprotiline hydrochloride.....	128	Extemporaneously Compounded		Genito-Urinary.....	73
MAR-Midodrine.....	47	Preparations.....	290	Micreme.....	73
Marcaïn.....	122	Nervous.....	126	Micreme H.....	71
Marcaïn Heavy.....	123	Methenamine (Hexamine)		Microlut.....	74
Marcaïn Isobaric.....	122	hippurate.....	95	Midazolam.....	142
Marcaïn with Adrenaline.....	122	Methohexital sodium.....	122	Midazolam-Baxter.....	142
Marevan.....	37	Methopt.....	279	Midazolam-Pfizer.....	142
Marine Blue Lotion SPF 50+.....	72	Methotrexate.....	151	Midodrine.....	47
Martindale Pharma.....	281	Methotrexate DBL Onco-Vial.....	151	Midostaurin.....	163
Mask for spacer device.....	321	Methotrexate Ebewe.....	151	Mifepristone.....	74
Maviret.....	106	Methotrexate Sandoz.....	151	Milrinone.....	55
Maxidex.....	275	Methoxsalen		Milrinone-Baxter.....	55
Maxitrol.....	274	[8-methoxypsoralen].....	71	Minerals.....	23
MCT Oil.....	293	Methoxyflurane.....	124	Mini-Wright AFS Low Range.....	321
Measles, mumps and rubella		Methyl aminolevulinate		Mini-Wright Standard.....	321
vaccine.....	318	hydrochloride.....	72	Minidiab.....	11
Mebendazole.....	101	Methyl hydroxybenzoate.....	290	Minims Prednisolone.....	276
Mebeverine hydrochloride.....	7	Methylcellulose.....	290	Minirin.....	87
Medac.....	149	Methylcellulose with glycerin and		Minirin Melt.....	87
Medrol.....	79	sodium saccharin.....	290	Minocycline.....	94
Medroxyprogesterone.....	81	Methylcellulose with glycerin and		Minoxidil.....	55
Medroxyprogesterone acetate		sucrose.....	290	Mirena.....	74
Genito-Urinary.....	74	Methylidopa.....	49	Miro-Amoxicillin.....	92
Hormone Preparations.....	80	Methylidopa Viatris.....	49	Mirtazapine.....	128
Mefenamic acid.....	118	Methylene blue.....	286	Misoprostol.....	7
Mefloquine.....	101	Methylnaltrexone bromide.....	16	Mitomycin C.....	150
Meglumine gadopentetate.....	285	Methylphenidate ER - Teva.....	144	Mitozantrone.....	150
Meglumine iotroxate.....	285	Methylphenidate hydrochloride.....	144	Mitozantrone Ebewe.....	150
Mekinist.....	169	Methylprednisolone (as sodium		Mivacurium chloride.....	117
Melatonin.....	141	succinate).....	79	Mixed salt solution for eye	
Melpha.....	149	Methylprednisolone aceponate.....	70	irrigation.....	277
Melphalan.....	149	Methylprednisolone acetate.....	79	MMA/PA Anamix Infant.....	298
Meningococcal (A, C, Y and W-135)		Methylthionium chloride [Methylene		MMA/PA Anamix Junior.....	298
conjugate vaccine.....	311	blue].....	286	MMA/PA Explore 5.....	298
Meningococcal B multicomponent		Methylxanthines.....	271	MMA/PA Express 15.....	298

Moclobemide	128	Nadolol	48	Nivolumab	248
Modafinil	145	Nadolol BNM	48	Nodia	5
Modafinil Max Health	145	Naglazyme	19	Noflam 250	118
Molaxole	16	Naloxone hydrochloride	281	Noflam 500	118
Mometasone furoate	70	Naltraccord	146	Non-Steroidal Anti-Inflammatory Drugs	117
Monogen	300	Naltrexone hydrochloride	146	Nonacog gamma, [Recombinant factor IX]	35
Monosodium glutamate with sodium aspartate	288	Naprosyn SR 1000	118	Noradrenaline	54
Monosodium l-aspartate	288	Naprosyn SR 750	118	Noradrenaline BNM	54
Montelukast	270	Naproxen	118	Norethindrone - CDC	74
Montelukast Viatris	270	Natalizumab	139	Norethisterone	
Moroctocog alfa [Recombinant factor VIII]	35	Natamycin	274	Genito-Urinary	74
Morphine hydrochloride	126	Natulan	156	Hormone Preparations	81
Morphine sulphate	126	Nausafix	133	Norethisterone with mestranol	73
Morphine tartrate	126	Nausicalm	132	Norflex	117
Motetis	120	Nefopam hydrochloride	125	Norflexacin	94
Mouth and Throat	25	Neo-Mercazole	86	Noriday	74
Movapo	120	Neocate Gold (Unflavoured)	302	Noriday 28	74
Movicol	16	Neocate Junior Unflavoured	302	Normison	142
Moxifloxacin	93	Neocate Junior Vanilla	302	Norpress	128
Moxifloxacin Kabi	93	Neocate SYNEO	302	Nortriptyline hydrochloride	128
Mozobil	39	Neoral	175	Norvir	105
MSUD Anamix Infant	295	Neostigmine metilsulfate	113	Noumed Dexamfetamine	143
MSUD Anamix Junior	295	Neostigmine metilsulfate with glycopyrronium bromide	113	Noumed Isoniazid	100
MSUD Anamix Junior LQ	295	Neosyneprine HCL	54	Noumed Paracetamol	125
MSUD Explore 5	295	Nepro HP (strawberry)	307	Noumed Pethidine	127
MSUD Express 15	295	Nepro HP (vanilla)	307	Noumed Phenobarbitone	130
MSUD Lophlex LQ 20	295	Neulumex	284	Novadoz	149
MSUD Maxamum	295	Neupogen	40	Novasource Renal (Vanilla)	307
Mucolytics and Expectorants	271	NeuroTabs	23	Novatretin	71
Mucosoothe	123	Nevirapine	103	NovoMix 30 FlexPen	10
Multiple Sclerosis Treatments	138	Nevirapine Viatris	103	NovoRapid FlexPen	10
Multivitamin and mineral supplement	26	Nicardipine hydrochloride	48	NovoSeven RT	34
Multivitamin renal	26	Nicorandil	55	Nozinan	134
Multivitamins	27	Nicotine	147	Nucala	214
Mupirocin	67	Nifedipine	49	Nuelin	271
Muscle Relaxants and Related Agents	116	Nifuran	95	Nuelin-SR	271
Mvite	27	Nilotinib	164	Nupentin	129
Myambutol	100	Nilstat		Nusinersen	142
Mycobutin	100	Alimentary	26	Nutilis	294
MycosNail	67	Genito-Urinary	73	Nutren Diabetes (vanilla)	300
Mycophenolate mofetil	258	Infections	96	Nutrini Energy Multi Fibre	306
Mydriacyl	279	Nimenrix	312	Nutrini Energy RTH	306
Mydriatics and Cycloplegics	279	Nimodipine	49	Nutrini Low Energy Multi Fibre RTH	306
Mylan (24 hr release)	49	Nimetop	49	Nutrini Peptisorb Energy	303
Mylan Clomiphen	81	Nintedanib	267	Nutrini RTH	306
Mylan Italy (24 hr release)	49	Niraparib	153	Nutrison 800 Complete Multi Fibre	308
Myleran	149	Nirmatrelvir with ritonavir	109	Nutrison Advanced Peptisorb	300
Myloc CR	47	Nitazoxanide	102	Nutrison Concentrated	301
Mylotarg	204	Nitrates	53	Nutrison Energy	308
Myozyme	17	Nitroderm TTS 10	53	Nutrison Energy Multi Fibre	308
Mytolac	174	Nitroderm TTS 5	53	Nutrison Multi Fibre	308
		Nitrofurantoin	95	Nutrison Protein Intense	302
		Nitrolingual Pump Spray	53		
		Nivestim	40		

Nutrison Protein Plus.....	301	Ondansetron-AFT.....	133	Pamisol.....	113
Nutrison Protein Plus Multi Fibre.....	302	One-Alpha.....	28	Pamol.....	125
Nutrison RTH.....	308	Opdivo.....	248	Pancreatic enzyme.....	14
Nyefax Retard.....	49	Optional Pharmaceuticals.....	321	Pancuronium bromide.....	117
Nystatin.....		Ora-Blend.....	290	Pantoprazole.....	8
Alimentary.....	26	Ora-Blend SF.....	290	Panzop Relief.....	8
Dermatological.....	67	Ora-Plus.....	290	Papaverine hydrochloride.....	55
Genito-Urinary.....	73	Ora-Sweet.....	290	Paper wasp venom.....	264
Infections.....	96	Ora-Sweet SF.....	290	Para-aminosalicylic Acid.....	100
- O -		Oralcon 30 ED.....	73	Paracetamol.....	125
Obinutuzumab.....	215	Oramorph.....	126	Paracetamol (Ethics).....	125
Obstetric Preparations.....	74	Oramorph CDC S29.....	126	Paracetamol Kabi.....	125
Ocrelizumab.....	140	Oratane.....	68	Paracetamol with codeine.....	127
Ocrevus.....	140	Ornidazole.....	102	Paraffin.....	
Octocog alfa [Recombinant factor VIII] (Advate).....	35	Orphenadrine citrate.....	117	Alimentary.....	15
Octocog alfa [Recombinant factor VIII] (Kogenate FS).....	35	Oruvail SR.....	118	Dermatological.....	69
Octreotide.....	173	Oseltamivir.....	109	Extemporaneously Compounded Preparations.....	290
Octreotide Long-Acting.....	174	Osimertinib.....	164	Paraffin liquid with soft white paraffin.....	280
Ocular Lubricants.....	279	Osmolite RTH.....	308	Paraffin liquid with wool fat.....	280
Oestradiol.....	80-81	Other Cardiac Agents.....	53	Paraffin with wool fat.....	69
Oestradiol valerate.....	80	Other Endocrine Agents.....	81	Paraldehyde.....	129
Oestradiol with norethisterone acetate.....	80	Other Oestrogen Preparations.....	81	Parecoxib.....	118
Oestriol.....		Other Otolological Preparations.....	280	Paromomycin.....	88
Genito-Urinary.....	75	Other Progestogen Preparations.....	81	Paroxetine.....	128
Hormone Preparations.....	81	Other Skin Preparations.....	72	Paser.....	100
Oestrogens.....	75	Ovestin.....		Patent blue V.....	286
Oestrogens (conjugated equine).....	80	Genito-Urinary.....	75	Paxam.....	138
Oestrogens with medroxyprogesterone acetate.....	80	Hormone Preparations.....	81	Paxlovid.....	109
Ofev.....	267	Oxaliplatin.....	158	Paxtine.....	128
Oil in water emulsion.....	69	Oxandrolone.....	77	Pazopanib.....	166
Oily phenol [Phenol oily].....	7	Oxazepam.....	138	Pazopanib Teva.....	166
Olanzapine.....	135-136	Oxpentifylline.....	55	Peak flow meter.....	321
Olaparib.....	154	Oxybuprocaine hydrochloride.....	277	Peanut oil.....	289
Olive oil.....	290	Oxybutynin.....	76	Pediasure (chocolate).....	306
Olopatadine.....	276	Oxycodone Amneal.....	126	Pediasure (strawberry).....	306
Olopatadine Teva.....	276	Oxycodone hydrochloride.....	126	Pediasure (vanilla).....	306
Olsalazine.....	7	Oxycodone Lucis S29.....	126	Pediasure Plus.....	306
Omaliuzumab.....	216	Oxycodone Sandoz.....	126	Pediasure RTH.....	306
Omeprazole.....	8	Oxymetazoline hydrochloride.....	269	Pegaspargase.....	155
Omeprazole actavis 10.....	8	Oxytocin.....	74	Pegasys.....	110
Omeprazole actavis 20.....	8	Oxytocin BNM.....	74	Pegasys (S29).....	110
Omeprazole actavis 40.....	8	Oxytocin with ergometrine maleate.....	75	Pegfilgrastim.....	40
Omeprazole Teva.....	8	Ozurdex.....	275	Pegylated interferon alfa-2a.....	110
Omezol IV.....	8	- P -		Pembrolizumab.....	250
Omnipaque.....	284	Pacifen.....	116	Pemetrexed.....	151
Omnitrope.....	82	Pacimol.....	125	Pemetrexed-AFT.....	151
Onbrez Breezhaler.....	270	Paclitaxel.....	170	Penicillamine.....	113
Oncaspar LYO.....	155	Paediatric Seravit.....	27	Penicillin G.....	92
OncoTICE.....	258	Palbociclib.....	165	Penicillin V.....	93
Ondansetron.....	133	Palbociclib Pfizer.....	165	Pentacarinat.....	102
		Paliperidone.....	136	Pentagastrin.....	81
		Paliperidone palmitate.....	137	Pentamidine isethionate.....	102
		Palivizumab.....	217	Pentasa.....	6
		Pamidronate disodium.....	113	Pentostatin [Deoxycofomycin].....	155

Pentoxifylline [Oxpentifylline].....	55	Pioglitazone.....	11	Poly-Tears.....	280
Peptamen OS 1.0 (Vanilla).....	300	Piperacillin with tazobactam.....	93	Poly-Visc.....	280
Pepti-Junior.....	304	Pipthiazine palmitate.....	137	Polycal.....	292
Perflutren.....	285	PipTaz-AFT.....	93	Polyethylene glycol 400 and propylene glycol.....	280
Perhexiline maleate.....	49	Pirfenidone.....	268	Polyhexamethylene biguanide.....	290
Pericyazine.....	135	Pituitary and Hypothalamic Hormones and Analogues.....	81	Polyvinyl alcohol with povidone.....	280
Perindopril.....	44	Pivmecillinam.....	95	Pomalidomide.....	155
Periset.....	133	Pizotifen.....	132	Pomolide.....	155
Periset ODT.....	133	PKU Anamix Infant.....	296	Poractant alfa.....	272
Perjeta.....	218	PKU Anamix Junior.....	296	Posaconazole.....	97
Permethrin.....	68	PKU Anamix Junior LQ (Berry).....	296	Posaconazole Juno.....	97
Perrigo.....	72	PKU Anamix Junior LQ (Orange).....	296	Potassium chloride.....	41, 43
Perzumab.....	218	PKU Build 10.....	297	Potassium chloride with sodium chloride.....	41
Peteha.....	100	PKU Build 20 Chocolate.....	297	Potassium citrate.....	75
Pethidine hydrochloride.....	127	PKU Build 20 Raspberry Lemonade.....	297	Potassium dihydrogen phosphate.....	41
Pevaryl.....	67	PKU Build 20 Smooth.....	297	Potassium iodate Alimentary.....	23
Pexsig.....	49	PKU Build 20 Vanilla.....	297	Hormone Preparations.....	86
Pfizer Exemestane.....	173	PKU Explore 10.....	296	Potassium iodate with iodine.....	23
Pheburane.....	21	PKU Explore 5.....	296	Potassium perchlorate.....	87
Phenaseen.....	152	PKU Express 20.....	296	Potassium permanganate.....	72
Phenelzine sulphate.....	128	PKU First Spoon.....	296	Povidone K30.....	290
Phenindione.....	37	PKU Glytactin RTD 15.....	297	Povidone-iodine.....	283
Phenobarbitone.....	130, 142	PKU Glytactin RTD 15 Lite.....	297	Povidone-iodine with ethanol.....	284
Phenobarbitone sodium.....	290	PKU GMPro LQ.....	297	Pradaxa.....	36
Phenol		PKU GMPro Mix-In.....	297	Pralidoxime chloride.....	281
Extemporaneously Compounded Preparations.....	290	PKU GMPro Ultra Lemonade.....	297	Pralidoxime iodide.....	281
Various.....	287	PKU GMPro Ultra Vanilla.....	297	Pramipexole hydrochloride.....	121
Phenol oily.....	7	PKU Lophlex LQ 10.....	296	Pravastatin.....	51
Phenol with ioxaglic acid.....	287	PKU Lophlex LQ 20.....	296-297	Praxbind.....	34
Phenothrin.....	68	PKU Lophlex Powder.....	296	Praziquantel.....	101
Phenoxybenzamine hydrochloride.....	45	PKU Lophlex Sensations 20 (berries).....	297	Prazosin.....	45
Phenoxyethylpenicillin [Penicillin V].....	93	PKU Restore Powder.....	296	Prazosin Mylan.....	45
Phentolamine mesylate.....	45	PKU sphere20 Banana.....	297	Pred Forte.....	276
Phenylalanine50.....	299	PKU sphere20 Chocolate.....	297	Prednisolone.....	79
Phenylephrine hydrochloride Cardiovascular.....	54	PKU sphere20 Lemon.....	297	Prednisolone acetate.....	276
Sensory.....	279	PKU sphere20 Red Berry.....	297	Prednisolone sodium phosphate.....	276
Phenytoin.....	130	PKU sphere20 Vanilla.....	297	Prednisolone- AFT.....	276
Phenytoin sodium.....	129-130	PKU Start.....	296	Prednisone.....	79
Phlexy 10.....	296	Plasma-Lyte 148.....	40	Prednisone Clinect.....	79
Phosphorus.....	42	Plasma-Lyte 148 & 5% Glucose.....	40	Pregabalin.....	130
Phytomenadione.....	35	Plendil ER.....	48	Pregabalin Pfizer.....	130
Picibanil.....	259	Plenvu.....	15	Pregnancy test - hCG urine.....	321
Pilocarpine hydrochloride.....	278	Plerixafor.....	39	preOp.....	307
Pilocarpine nitrate		Pneumococcal (PCV13) conjugate vaccine.....	313	Presolol.....	47
Extemporaneously Compounded Preparations.....	290	Pneumococcal (PPV23) polysaccharide vaccine.....	314	Prevenar 13.....	313
Sensory.....	278	Pneumovax 23.....	314	Priadel.....	134
Pimafucort.....	71	Podophyllotoxin.....	72	Prilocaine hydrochloride.....	124
Pimecrolimus.....	71	Polidocanol.....	33	Prilocaine hydrochloride with felypressin.....	124
Pine tar with trolamine laurilsulfate and fluorescein.....	71	Poliomyelitis vaccine.....	318	Primaquine.....	102
Pinetarsol.....	71	Poloxamer.....	15	Primidone.....	130

Primolut N.....	81	Raltegravir potassium.....	106	Risperon.....	135
Primovist.....	285	Ramipex.....	121	Ritalin.....	144
Priorix.....	318	Ramipril.....	44	Ritalin LA.....	144
Probenecid.....	116	Ranbaxy-Cefaclor.....	89	Ritonavir.....	105
Procaine penicillin.....	93	Ranibizumab.....	219	Rituximab (mabthera).....	220
Procarbazine hydrochloride.....	156	Ranitidine.....	8	Rituximab (riximyo).....	221
Prochlorperazine.....	133	Rapamune.....	259	Rivaroxaban.....	37
Proctosedyl.....	7	Rasagiline.....	121	Rivastigmine.....	145
Procyclidine hydrochloride.....	120	Rasburicase.....	116	Rivastigmine Patch BNM 10.....	145
Progesterone.....	75	Readi-CAT 2.....	284	Rivastigmine Patch BNM 5.....	145
Proglcem.....	9	Reandron 1000.....	77	Riximyo.....	221
Proglycem.....	9	Recombinant factor IX.....	34-35	RIXUBIS.....	35
Prodynova.....	80	Recombinant factor VIIa.....	34	Rizamelt.....	132
Prolia.....	114	Recombinant factor VIII.....	35	Rizatriptan.....	132
Promethazine hydrochloride.....	265	Rectogesic.....	7	Robinul.....	7
Propafenone hydrochloride.....	47	Red back spider antivenom.....	282	Rocuronium bromide.....	117
Propamide isethionate.....	274	Redipred.....	79	Ropin.....	121
Propofol.....	122	Relenza Rotadisk.....	109	Ropinrole hydrochloride.....	121
Propranolol.....	48	Relistor.....	16	Ropivacaine hydrochloride.....	124
Propylthiouracil.....	87	Remdesivir.....	109	Ropivacaine Kabi.....	124
Prostin E2.....	74	Remicade.....	205	Rose bengal sodium.....	276
Prostin VR.....	55	Remifentanil.....	127	Rosuvastatin.....	52
Protamine sulphate.....	37	Remifentanil-AFT.....	127	Rosuvastatin Viatrix.....	52
Protifar.....	293	Renilon 7.5 (apricot).....	307	Rotarix.....	318
Protonamide.....	100	Renilon 7.5 (caramel).....	307	Rotavirus oral vaccine.....	318
Protirelin.....	87	Resonium A.....	43	Roxithromycin.....	92
Proveblue.....	286	Resource Beneprotein.....	293	Rubifen.....	144
Provera.....	80	Respiratory Stimulants.....	273	Rubifen SR.....	144
Provera HD.....	81	Retinol.....	27	Ruriotocog alfa pegol [Recombinant factor VIII].....	35
Proxymetacaine hydrochloride.....	277	Retinol Palmitate.....	280	Ruxolitinib.....	167
Pseudoephedrine hydrochloride.....	269	ReTrieve.....	68	Rydapt.....	163
Psoriasis and Eczema Preparations.....	71	Retrovir.....	104	Ryzodeg 70/30 Penfill.....	10
PTU.....	87	Retrovir IV.....	104	- S -	
Pulmonary Surfactants.....	272	Revolade.....	32	S26 LBW Gold RTF.....	305
Pulmozyme.....	271	Ribociclib.....	166	Sabril.....	131
Puri-nethol.....	151	Riboflavin.....	20	Sacubitril with valsartan.....	45
Pyrazinamide.....	100	Riboflavin 5-phosphate.....	278	SalAir.....	269
Pyridostigmine bromide.....	113	Ricit.....	75	Salazopyrin.....	7
Pyridoxal-5-phosphate.....	20	Ricovir.....	106	Salazopyrin EN.....	7
Pyridoxine hydrochloride.....	27	Rifabutin.....	100	Salbutamol.....	269
Pyridoxine multichem.....	27	Rifadin.....	101	Salbutamol with ipratropium bromide.....	266
Pyrimethamine.....	102	Rifampicin.....	101	Salicylic acid.....	290
Pytazen SR.....	37	Rifaximin.....	9	Salmeterol.....	270
- Q -		Rifinah.....	100	Salmonella typhi vaccine.....	315
Quetapel.....	135	Rilutek.....	120	Sandimmun.....	175
Quetiapine.....	135	Riluzole.....	120	Sandomigran.....	132
Quetiapine Viatrix.....	135	Ringer's solution.....	41	Sandostatin LAR.....	174
Quinapril.....	44	Rinvoq.....	260	Sapropterin Dihydrochloride.....	20
Quinine dihydrochloride.....	102	Riodine.....	283	Scalp Preparations.....	72
Qvar.....	269	Risdiplam.....	142	Scandonest 3%.....	124
- R -		Risedronate Sandoz.....	113	Sclerosing Agents.....	273
RA-Morph.....	126	Risedronate sodium.....	113	Scopolamine Transdermal System Viatrix.....	133
Rabies vaccine.....	318	Risperdal.....	135	Sebizole.....	67
Raloxifene.....	114	Risperdal Consta.....	137		
		Risperidone.....	135, 137		
		Risperidone (Teva).....	135		

Secretin pentahydrochloride.....	286	Sodium chloride	Sotalol.....	48
Secukinumab.....	233	Blood.....	Sotalol Viatris.....	48
Sedatives and Hypnotics.....	141	Respiratory.....	Soya oil.....	281
Seebri Breezhaler.....	266	Various.....	Spacer device.....	321
Selegiline hydrochloride.....	121	Sodium chloride with sodium	Span-K.....	43
Selenium.....	25	bicarbonate.....	Spazmol.....	7
Sennosides.....	16	Sodium citrate	Specialised Formulas.....	299
Serc.....	132	Alimentary.....	Spinal Muscular Atrophy.....	142
Serenace.....	134	Extemporaneously Compounded	Spinraza.....	142
Seretide.....	270	Preparations.....	Spiolto Respimat.....	266
Seretide Accuhaler.....	270	Sodium citrate with sodium chloride	Spiractin.....	50
Serevent.....	270	and potassium chloride.....	Spiramycin.....	102
Serevent Accuhaler.....	270	Sodium citrate with sodium lauryl	Spiriva.....	266
Sertraline.....	128	sulphoacetate.....	Spiriva Respimat.....	266
Setrona.....	128	Sodium citro-tartrate.....	Spironolactone.....	50
Sevoflurane.....	122	Sodium cromoglicate	Stalevo.....	121
Sevredol.....	126	Alimentary.....	Standard Feeds.....	307
Shingles vaccine.....	319	Respiratory.....	Staphlex.....	92
Shingrix.....	319	Sensory.....	Starch.....	291
SII-Onco-BCG.....	258	Sodium dihydrogen phosphate	Stavudine.....	104
Sildenafil.....	60	[Sodium acid phosphate].....	Stelara.....	243
Siltuximab.....	235	Sodium fluoride.....	Sterculia with frangula.....	15
Silver nitrate		Sodium fusidate [Fusidic acid]	SteroClear.....	265
Dermatological.....	72	Dermatological.....	Stesolid.....	129
Extemporaneously Compounded		Infections.....	Stimulants / ADHD Treatments.....	143
Preparations.....	290	Sensory.....	Stiripentol.....	131
Simdax.....	53	Sodium hyaluronate [Hyaluronic acid]	Streptomycin sulphate.....	88
Simeticone.....	5	Alimentary.....	Stromectol.....	101
Simulect.....	200	Sensory.....	Sucalfate.....	9
Simvastatin.....	52	Sodium hyaluronate [Hyaluronic acid]	Sucrose.....	125
Simvastatin Mylan.....	52	with chondroitin sulphate.....	Sugammadex.....	117
Simvastatin Viatris.....	52	Sodium hydroxide.....	Sugammadex BNM.....	117
Sincalide.....	286	Sodium hypochlorite.....	Sulfadiazine silver.....	67
Sinemet.....	121	Sodium metabisulfite.....	Sulfadiazine sodium.....	95
Sinemet CR.....	121	Sodium nitrite.....	Sulfasalazine.....	7
Sintetica Baclofen Intrathecal.....	116	Sodium nitroprusside	Sulindac.....	118
Sinolimus.....	259	Cardiovascular.....	Sulphacetamide sodium.....	274
Sirturo.....	100	Optional Pharmaceuticals.....	Sulphur.....	291
Siterone.....	77	Sodium phenylbutyrate.....	Sulprix.....	134
Slow-Lopresor.....	48	Sodium phosphate with phosphoric	Sumagran.....	132
Snake antivenom.....	282	acid.....	Sumatriptan.....	132
Sodibic.....	43	Sodium picosulfate.....	Sunitinib.....	168
Sodium acetate.....	41	Sodium polystyrene sulphonate.....	Sunitinib Pfizer.....	168
Sodium acid phosphate.....	42	Sodium stibogluconate.....	Sunscreen, proprietary.....	72
Sodium alginate with magnesium		Sodium tetradecyl sulphate.....	Suprane.....	121
alginate.....	5	Sodium thiosulfate.....	Surgical Preparations.....	287
Sodium alginate with sodium		Sodium valproate.....	Sustagen Hospital Formula	
bicarbonate and calcium		Sodium with potassium.....	(Chocolate).....	308
carbonate.....	5	Solifenacin succinate.....	Sustagen Hospital Formula	
Sodium aurothiomalate.....	113	Solifenacin succinate Max	(Vanilla).....	308
Sodium benzoate.....	21	Health.....	Suxamethonium chloride.....	117
Sodium bicarbonate		Solifenacin Viatris.....	Sylvant.....	235
Blood.....	42-43	Solu-Cortef.....	Symbicort Turbuhaler.....	270
Extemporaneously Compounded		Solu-Medrol.....	Symmetrel.....	120
Preparations.....	290	Solu-Medrol Act-O-Vial.....	Sympathomimetics.....	54
Sodium calcium edetate.....	283	Somatropin.....	Synacthen.....	81

Synacthen Depot.....	81	Testosterone cipionate.....	77	Tramadol hydrochloride.....	127
Synagis.....	217	Testosterone esters.....	77	Tramal 100.....	127
Synermox.....	92	Testosterone undecanoate.....	77	Tramal 50.....	127
Syntometrine.....	75	Tetrabenazine.....	120	Tramal SR 100.....	127
Syrup.....	291	Tetracaine [Amethocaine] hydrochloride		Tramal SR 150.....	127
Systane Unit Dose.....	280	Nervous.....	124	Tramal SR 200.....	127
- T -		Sensory.....	277	Trametinib.....	169
Tacrolimus		Tetracosactide [Tetracosactrin].....	81	Trandate.....	47
Dermatological.....	72	Tetracosactrin.....	81	Tranexamic acid.....	34
Oncology.....	175	Tetracycline.....	94	Tranexamic-AFT.....	34
Tacrolimus Sandoz.....	175	Teva Lisinopril.....	44	Tranylcypromine sulphate.....	128
Tafinlar.....	159	Thalidomide.....	156	Trastuzumab (Herzuma).....	240
Tagitol V.....	284	Thalomid.....	156	Trastuzumab deruxitecan.....	241
Tagrisso.....	164	Theobroma oil.....	291	Trastuzumab emtansine.....	242
Talc.....	273	Theophylline.....	271	Travatan.....	279
Taliglucerase alfa.....	21	Thiamine hydrochloride.....	28	Travoprost.....	279
Tambocor.....	46	Thiamine multichem.....	28	Treatments for Dementia.....	145
Tambocor German.....	46	Thioguanine.....	151	Treatments for Substance	
Tamoxifen citrate.....	173	Thiopental [Thiopentone]		Dependence.....	146
Tamoxifen Sandoz.....	173	sodium.....	122	Trelegy Ellipta.....	267
Tamsulosin hydrochloride.....	75	Thiopentone.....	122	Tretinoin	
Tamsulosin-Rex.....	75	Thiotepa.....	149	Dermatological.....	68
Tasigna.....	164	Thrombin.....	34	Oncology.....	157
Tasmar.....	121	Thyroid and Antithyroid		Trexate.....	151
Taurine.....	22	Preparations.....	86	Tri-sodium citrate.....	291
TCu 380 Plus Normal.....	74	Thyrotropin alfa.....	81	Triamcinolone acetonide	
Tecentriq.....	246	Ticagrelor.....	38	Alimentary.....	25
Tecfidera.....	139	Ticagrelor Sandoz.....	38	Dermatological.....	70
TEEVIR.....	104	Ticarcillin with clavulanic acid.....	93	Hormone Preparations.....	79
Tegretol.....	129	Ticlopidine.....	39	Triamcinolone acetonide with	
Tegretol CR.....	129	Tigecycline.....	94	gramicidin, neomycin and	
Teicoplanin.....	96	Tilcotil.....	118	nystatin.....	275
Temacord.....	156	Timolol.....	278	Triamcinolone acetonide with	
Temazepam.....	142	Tiotropium bromide.....	266	neomycin sulphate, gramicidin	
Temozolomide.....	156	Tiotropium bromide with		and nystatin.....	71
Temozolomide Taro.....	156	olodaterol.....	266	Triamcinolone hexacetone.....	79
Tenecteplase.....	39	Tivicay.....	106	Triazolam.....	142
Tenofovir disoproxil.....	106	TMP.....	96	Trichloroacetic acid.....	291
Tenofovir Disoproxil Emtricitabine		Tobradex.....	274	Trientine.....	23
Viatr.....	108	Tobramycin		Trientine Waymade.....	23
Tenofovir Disoproxil Viatris.....	106	Infections.....	88	Trikafta.....	271
Tenoxicam.....	118	Sensory.....	274	Trimethoprim.....	96
Tensipine MR10.....	49	Tobramycin (Viatris).....	88	Trimethoprim with	
Tepadina.....	149	Tobramycin BNM.....	88	sulphamethoxazole	
Terazosin.....	46	Tobrex.....	274	[Co-trimoxazole].....	96
Terbinafine.....	99	Tocilizumab.....	236	Triovir.....	104
Terbutaline.....	75	Tofranol.....	127	Trisul.....	96
Terbutaline sulphate.....	269	Tolcapone.....	121	Trometamol.....	287
Teriflunomide.....	139	Tolvaptan.....	51	Tropicamide.....	279
Teriflunomide Sandoz.....	139	Topamax.....	131	Tropisetron.....	133
Teriparatide.....	115	Topicaline.....	123	Trulicity.....	11
Teriparatide - Teva.....	115	Topical Products for Joint and		Tryzan.....	44
Terlipressin.....	87	Muscular Pain.....	119	Tuberculin PPD [Mantoux] test.....	320
Terlipressin Ever Pharma.....	87	Topiramate.....	131	Tubersol.....	320
Testogel.....	77	Topiramate Actavis.....	131	Two Cal HN.....	301
Testosterone.....	77	Torbay.....	54	TYR Anamix Infant.....	298

TYR Anamix Junior.....	298	Vecuronium bromide	117	Wool fat	
TYR Anamix Junior LQ.....	298	Vedafil.....	60	Dermatological.....	70
TYR Explore 5.....	298	Vedolizumab.....	244	Extemporaneously Compounded	
TYR Lophlex LQ 20.....	298	Vegzelma.....	201	Preparations	291
TYR Sphere 20.....	298	Veklury.....	109	- X -	
Tyrosine1000.....	299	Veletri	62	Xalkori.....	159
Tysabri.....	139	Venclexta.....	157	Xanthan	291
- U -		Venetoclax.....	157	Xarelto	37
UK Cipla	269	Venlafaxine.....	128	Xgeva	114
UK Synacthen.....	81	Venofer	24	Xifaxan.....	9
Ultibro Breezhaler.....	266	VENOX.....	264	XMET Maxamum.....	295
Ultraproct.....	7	Ventolin.....	269	Xolair	216
Umeclidinium.....	266	Vepesid.....	152	XP Maxamum	296
Umeclidinium with vilanterol	266	Verapamil hydrochloride.....	49	Xylocaine.....	123
Univent	265	Vermox.....	101	Xylometazoline hydrochloride.....	269
Upadacitinib.....	260	Versacloz.....	134	Xyntha	35
Ural	76	Vesanoid.....	157	- Y -	
Urea		Vexazone.....	11	Yellow jacket wasp venom	264
Dermatological.....	70	Vfend.....	98	Yervoy	248
Extemporaneously Compounded		Victoza	12	- Z -	
Preparations	291	Vigabatrin	131	Zanamivir.....	109
Urex Forte.....	50	Vigisom	141	Zarontin	129
UroFos.....	95	Vildagliptin.....	11	Zavedos.....	150
Urografin.....	284	Vildagliptin with metformin		Zavicefta.....	89
Urokinase	39	hydrochloride	11	Zeffix.....	106
Urologicals.....	75	Vimpat	129	Zejula	153
Uromitexan.....	171	Vinblastine sulphate	171	Zematop	72
Ursodeoxycholic acid.....	14	Vincristine sulphate	171	Zetlam.....	106
Ursosan.....	14	Vinorelbine.....	172	Ziagen.....	104
Ustekinumab.....	243	Vinorelbine Te Arai.....	172	Zidovudine [AZT].....	104
Utrogestan.....	75	Viral Vaccines.....	315	Zidovudine [AZT] with	
- V -		Viramune Suspension	103	lamivudine	104
Vaclovir.....	107	Virupos.....	274	Ziextenzo.....	40
Valaciclovir	107	Viscoat.....	277	Zimybe.....	53
Valganciclovir	107	Visipaque.....	284	Zinc	
Valganciclovir Viatris	107	Vit.D3.....	28	Alimentary.....	25
Valine50.....	299	VitA-POS.....	280	Dermatological.....	68
Vancomycin.....	96	Vital.....	300	Zinc and castor oil	68
Vanilla SilQ HD.....	284	Vitamin B complex.....	28	Zinc chloride	25
Vanilla SilQ MD	284	Vitamin B6 25	27	Zinc oxide	291
Varenicline.....	147	Vitamins.....	26	Zinc sulphate	25
Varibar - Honey.....	284	Vivonex TEN.....	300	Zinc with wool fat.....	69
Varibar - Nectar.....	284	Voltaren	118	Zincaps.....	25
Varibar - Pudding.....	284	Voltaren D.....	118	Zinfo.....	90
Varibar - Thin Liquid	284	Voltaren SR	118	Ziprasidone.....	135
Varicella vaccine [Chickenpox		Volumatic.....	321	Zista	265
vaccine]	319	Voriconazole.....	98	Zithromax.....	90
Varicella zoster vaccine [Shingles		Vttack.....	98	Zo-Rub HP.....	124
vaccine]	319	Vyvanse.....	143	Zo-Rub Osteo.....	119
Varilrix.....	319	- W -		Zoladex.....	82
Vasodilators.....	54	Warfarin sodium.....	37	Zoledronic acid	
Vasopressin.....	87	Wart Preparations.....	72	Hormone Preparations.....	78
Vasopressin Agents.....	87	Water		Musculoskeletal	113
Vasorex	48	Blood.....	42	Zoledronic Acid Injection Mylan.....	78
Vebulis.....	63	Various.....	287	Zoledronic acid Viatris	
Vecure.....	117	White Soft Liquid Paraffin AFT	69	Hormone Preparations.....	78

Musculoskeletal	113
Zopiclone	142
Zopiclone Actavis	142
Zostrix	119
Zostrix HP	124
Zuclopenthixol acetate	135
Zuclopenthixol decanoate	137
Zuclopenthixol hydrochloride	135
Zusdone	135
Zyban	146
Zypine	135
Zypine ODT	135
Zyprexa Relprevv	136
Zytiga	172
Zyvox	95















