Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRES	CRIBER	PATIENT:					
Name):						
Ward:		NHI:					
Dexa	Dexamethasone						
Re-a	ssessmer equisites	Diabetic macular oedema It required after 12 months (tick boxes where appropriate) Cribed by, or recommended by an ophthalmologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ object. Patients have diabetic macular oedema with pseudophakic lens Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision Patient's disease has progressed despite 3 injections with bevacizumab Patient is unsuitable or contraindicated to treatment with anti-VEGF agents Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year					
CONTINUATION – Diabetic macular oedema Re-assessment required after 12 months Prerequisites (tick boxes where appropriate) Orescribed by, or recommended by an ophthalmologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital. and Orea Patient's vision is stable or has improved (prescriber determined) and Orea Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum							
		of 3 implants per eye per year					
Re-a	ssessmer equisites	Women of child bearing age with diabetic macular oedema at required after 12 months (tick boxes where appropriate) cribed by, or recommended by an ophthalmologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ oital.					
and	and on an analysis of an analys	Patients have diabetic macular oedema Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision Patient is of child bearing potential and has not yet completed a family Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year					

I confirm that the above details are correct:

Cianad.	Doto.	
Siurieu.	 Date.	

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRES	CRIBER		PATIENT:					
Name	e:		Name:					
Ward			NHI:					
Dexamethasone - continued								
Re-a	ssessment (tequisites (t	ON – Women of child bearing age with diabetic macular oedema nt required after 12 months to (tick boxes where appropriate) scribed by, or recommended by an ophthalmologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ poital.						
	and F	Patient's vision is stable or has improved (prescriber determine Patient is of child bearing potential and has not yet completed a Dexamethasone implants are to be administered not more freq of 3 implants per eye per year						