HOSPITAL MEDICINES LIST
RESTRICTIONS CHECKLIST

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER

Name: .....

Ward: .....

PATIENT:

Name: .....

NHI: .....

Faricimab

INITIATION – Diabetic macular oedema
Re-assessment required after 4 months
Prerequisites (tick boxes where appropriate)

- ○ Patient has centre involving diabetic macular oedema (DMO)

and

- ○ Patient's disease is nonresponsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly

and

- ○ Patient has reduced visual acuity between 6/9 – 6/36 with functional awareness of reduction in vision

and

- ○ Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers

and

- ○ There is no centre-involving sub-retinal fibrosis or foveal atrophy

and

- ○ Patient has not previously been treated with aflibercept for longer than 3 months

CONTINUATION – Diabetic macular oedema
Re-assessment required after 12 months
Prerequisites (tick boxes where appropriate)

- ○ There is stability or two lines of Snellen visual acuity gain

and

- ○ There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid)

and

- ○ Patient's vision is 6/36 or better on the Snellen visual acuity score

and

- ○ There is no centre-involving sub-retinal fibrosis or foveal atrophy

INITIATION – Wet age related macular degeneration
Re-assessment required after 3 months
Prerequisites (tick boxes where appropriate)

- ○ Wet age-related macular degeneration (wet AMD)

or

- ○ Polypoidal choroidal vasculopathy

or

- ○ Choroidal neovascular membrane from causes other than wet AMD

and

- ○ The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab

or

- ○ There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart

and

- ○ There is no structural damage to the central fovea of the treated eye

and

- ○ Patient has not previously been treated with ranibizumab or aflibercept for longer than 3 months

I confirm that the above details are correct:

Signed: ..... Date: .....

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to Section H of the Pharmaceutical Schedule. For community funding, see the Special Authority Criteria.

PRESCRIBER

Name: .....

Ward: .....

PATIENT:

Name: .....

NHI: .....

Faricimab *- continued*

CONTINUATION – Wet age related macular degeneration
Re-assessment required after 12 months
Prerequisites (tick boxes where appropriate)

- ❍ Patient's vision is 6/36 or better on the Snellen visual acuity score

and

- ❍ There is no structural damage to the central fovea of the treated eye

I confirm that the above details are correct:

Signed: ..... Date: .....