

RS1826 - Somatropin

Prader-Willi syndrome - INITIATION	5
Prader-Willi syndrome - CONTINUATION	5
Turner syndrome - INITIATION	2
Turner syndrome - CONTINUATION	3
Adults and adolescents - INITIATION	6
Adults and adolescents - CONTINUATION	7
Growth hormone deficiency in children - INITIATION	2
Growth hormone deficiency in children - CONTINUATION	2
Short stature due to chronic renal insufficiency - INITIATION	4
Short stature due to chronic renal insufficiency - CONTINUATION	4
Short stature without growth hormone deficiency - INITIATION	3
Short stature without growth hormone deficiency - CONTINUATION	3

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PRESCRIBER

Name: Name:

Ward: NHI:

Somatropin

INITIATION – growth hormone deficiency in children

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with $\text{GH} < 5 \text{ mcg/l}$ on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose $< 2 \text{ mmol/l}$ using a laboratory device)

or

Height velocity $< 25\text{th percentile}$ for age; and adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985)

and

A current bone age is < 14 years (female patients) or < 16 years (male patients)

and

Peak growth hormone value of $< 5.0 \text{ mcg per litre}$ in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required

and

If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate

and

Appropriate imaging of the pituitary gland has been obtained

CONTINUATION – growth hormone deficiency in children

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

A current bone age is 14 years or under (female patients) or 16 years or under (male patients)

and

Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davies (1985)

and

Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months

and

No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred

and

No malignancy has developed since starting growth hormone

INITIATION – Turner syndrome

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

The patient has a post-natal genotype confirming Turner Syndrome

and

Height velocity is $< 25\text{th percentile}$ over 6-12 months using the standards of Tanner and Davies (1985)

and

A current bone age is < 14 years

I confirm that the above details are correct:

Signed: Date:

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PRESCRIBER

Name: Name:

Ward: NHI:

Somatropin - continued

CONTINUATION – Turner syndrome

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

Height velocity greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts)

and

Height velocity is greater than or equal to 2 cm per year, calculated over six months

and

A current bone age is 14 years or under

and

No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred

and

No malignancy has developed since starting growth hormone

INITIATION – short stature without growth hormone deficiency

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay

and

Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985)

and

A current bone age is < 14 years (female patients) or < 16 years (male patients)

and

The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity

CONTINUATION – short stature without growth hormone deficiency

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985)

and

Height velocity is greater than or equal to 2 cm per year as calculated over six months

and

Current bone age is 14 years or under (female patients) or 16 years or under (male patients)

and

No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred

I confirm that the above details are correct:

Signed: Date:

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PRESCRIBER

Name: Name:

Ward: NHI:

Somatropin - continued

INITIATION – short stature due to chronic renal insufficiency

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist, paediatric endocrinologist or renal physician on the recommendation of a endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.
and

The patient's height is more than 2 standard deviations below the mean
and

Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985)
and

A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients)
and

The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease
and

The patient is under the supervision of a specialist with expertise in renal medicine
and

The patient has a GFR less than or equal to $30 \text{ ml/min}/1.73 \text{ m}^2$ as measured by the Schwartz method ($\text{Height(cm)}/\text{plasma creatinine (umol/l)} \times 40 = \text{corrected GFR (ml/min}/1.73 \text{ m}^2)$ in a child who may or may not be receiving dialysis
or

The patient has received a renal transplant and has received < $5\text{mg}/\text{m}^2/\text{day}$ of prednisone or equivalent for at least 6 months

CONTINUATION – short stature due to chronic renal insufficiency

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist, paediatric endocrinologist or renal physician on the recommendation of a endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.
and

Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985)
and

Height velocity is greater than or equal to 2 cm per year as calculated over six months
and

A current bone age is 14 years or under (female patients) or 16 years or under (male patients)
and

No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred
and

No malignancy has developed after growth hormone therapy was commenced
and

The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results
and

The patient has not received renal transplantation since starting growth hormone treatment
and

If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria

I confirm that the above details are correct:

Signed: Date:

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PRESCRIBER

Name: Name:

Ward: NHI:

Somatropin - continued

INITIATION – Prader-Willi syndrome

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria

and

The patient is aged six months or older

and

A current bone age is < 14 years (female patients) or < 16 years (male patients)

and

Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon

and

The patient is aged two years or older

and

There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months

or

The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation

CONTINUATION – Prader-Willi syndrome

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985)

and

Height velocity is greater than or equal to 2 cm per year as calculated over six months

and

A current bone age is 14 years or under (female patients) or 16 years or under (male patients)

and

No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred

and

No malignancy has developed after growth hormone therapy was commenced

and

The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months

I confirm that the above details are correct:

Signed: Date:

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PREScriBER

Name: Name:

Ward: NHI:

Somatropin - continued

INITIATION – adults and adolescents

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour)

and

The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses

and

The patient has severe growth hormone deficiency (see notes)

and

The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex

and

The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®)

Note: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre. The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until it is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

I confirm that the above details are correct:

Signed: Date:

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PRESCRIBER

Name: Name:

Ward: NHI:

Somatropin - continued

CONTINUATION – adults and adolescents

Re-assessment required after 12 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by an endocrinologist or paediatric endocrinologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

The patient has been treated with somatropin for < 12 months
and
 There has been an improvement in the Quality of Life Assessment defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline
and
 Serum IGF-I levels have increased to within $\pm 1SD$ of the mean of the normal range for age and sex
and
 The dose of somatropin does not exceed 0.7 mg per day for male patients, or 1 mg per day for female patients

or

The patient has been treated with somatropin for more than 12 months
and
 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors)
and
 Serum IGF-I levels have continued to be maintained within $\pm 1SD$ of the mean of the normal range for age and sex (other than for obvious external factors)
and
 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients

or

The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication
and
 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses
and
 The patient has severe growth hormone deficiency (see notes)
and
 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex
and
 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®)

Note: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre. The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

I confirm that the above details are correct:

Signed: Date: