

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PRESCRIBER

Name: Name:

Ward: NHI:

Iloprost

INITIATION – PAH monotherapy

Re-assessment required after 6 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

Patient has pulmonary arterial hypertension (PAH)

and

PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications

and

PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV

and

PAH has been confirmed by right heart catheterisation

and

A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)

and

A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg

and

A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵)

and

PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH

or

Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**

or

Patient has PAH other than idiopathic / heritable or drug-associated type

or

Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease

or

Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures

and

Iloprost is to be used as PAH monotherapy

and

Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan)

or

Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists

I confirm that the above details are correct:

Signed: Date:

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PRESCRIBER

Name: Name:

Ward: NHI:

Iloprost - *continued*

INITIATION – PAH dual therapy

Re-assessment required after 6 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

Patient has pulmonary arterial hypertension (PAH)

and

PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications

and

PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV

and

PAH has been confirmed by right heart catheterisation

and

A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)

and

A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg

and

A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵)

and

PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH

or

Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**

or

Patient has PAH other than idiopathic / heritable or drug-associated type

or

Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease

or

Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures

and

Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist

and

Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil

or

Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist

and

Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**

or

Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy

I confirm that the above details are correct:

Signed: Date:

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PRESCRIBER

Name: Name:

Ward: NHI:

Iloprost - continued

INITIATION – PAH triple therapy

Re-assessment required after 6 months

Prerequisites (tick boxes where appropriate)

Prescribed by, or recommended by a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

Patient has pulmonary arterial hypertension (PAH)

and

PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications

and

PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV

and

PAH has been confirmed by right heart catheterisation

and

A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair)

and

A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg

and

A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm⁻⁵)

and

PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH

or

Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**

or

Patient has PAH other than idiopathic / heritable or drug-associated type

or

Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease

or

Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures

and

Iloprost is to be used as PAH triple therapy

and

Patient is on the lung transplant list

or

Patient is presenting in NYHA/WHO functional class IV

or

Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**

and

Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario

I confirm that the above details are correct:

Signed: Date:

Use this checklist to determine if a patient meets the restrictions for funding in the **hospital setting**. For more details, refer to [Section H](#) of the Pharmaceutical Schedule. For community funding, see the [Special Authority Criteria](#).

PREScriBER

Name:

Ward:

PATIENT:

Name:

NHI:

Iloprost - *continued*

CONTINUATION

Re-assessment required after 2 years

Prerequisites (tick box where appropriate)

Prescribed by, or recommended by a respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital.

and

Patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

I confirm that the above details are correct:

Signed: Date: