May 2026
Volume 14

Editor:
Kaye Wilson, Doris Chong, Teresa Tay
email: enquiry@pharmac.govt.nz
Telephone +64 4 460 4990
Level 9, 40 Mercer Street
PO Box 10 254 Wellington 6143

Freephone Information Line
0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule, Section H for Hospital Pharmaceuticals (link to PDF copy) emailed to your nominated email address each month by subscribing at schedule.pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and TEX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/HML

Programmers

Anrik Drenth

email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency

ISSN 1179-3708 pdf

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

Introducing Pharmac 2

Part I General Rules 4

Part II
Alimentary Tract and Metabolism 5
Blood and Blood Forming Organs 30
Cardiovascular System 45
Dermatologicals 68
Genito-Urinary System 75
Hormone Preparations 79
Infections 90
Musculoskeletal System 116
Nervous System 122
Oncology Agents and Immunosuppressants 151
Respiratory System and Allergies 264
Sensory Organs 275
Various 282
Extemporaneous Compounds (ECPs) 290
Special Foods 293
Vaccines 311

Part III Optional Pharmaceuticals 323

Index 324

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.
Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://pharmac.govt.nz/about.

Glossary

Units of Measure

gram ..... g	microgram..... mcg	millimole..... mmol
kilogram..... kg	milligram ..... mg	unit..... u
international unit ..... iu	millilitre..... ml	

Abbreviations

application ..... app	enteric coated..... EC	solution .....soln
capsule ..... cap	granules..... grans	suppository .....suppos
cream..... crm	injection .....inj	tablet..... tab
dispersible .....disp	liquid .....liq	tincture.....tinc
effervescent..... eff	lotion..... lotn	
emulsion ..... emul	ointment..... oint	

HSS Hospital Supply Status

Guide to Section H listings

Example



General Rules for Section H of the Pharmaceutical Schedule are included in Section A.

Read the General Rules : https://pharmac.govt.nz/section-a.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Antacids and Antiflatulents

Antacids and Reflux Barrier Agents

ALUMINIUM HYDROXIDE WITH MAGNESIUM HYDROXIDE AND SIMETICONE

Tab 200 mg with magnesium hydroxide 200 mg and simeticone 20 mg — *e.g. Mylanta*

Oral liq 400 mg with magnesium hydroxide 400 mg and simeticone 30 mg per 5 ml — *e.g. Mylanta Double Strength*

SIMETICONE

Oral drops 100 mg per ml

Oral drops 20 mg per 0.3 ml

Oral drops 40 mg per ml

SODIUM ALGINATE WITH MAGNESIUM ALGINATE

Powder for oral soln 225 mg with magnesium alginate 87.5 mg, sachet — *e.g. Gaviscon Infant*

SODIUM ALGINATE WITH SODIUM BICARBONATE AND CALCIUM CARBONATE

Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg — *e.g. Gaviscon Extra Strength*

Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml — 7.50 — 500 ml — Acidex

SODIUM CITRATE

Oral liq 8.8% (300 mmol/l) — 25.00 — 90 ml — Biomed

Phosphate Binding Agents

ALUMINIUM HYDROXIDE

Tab 600 mg

CALCIUM CARBONATE – **Restricted** see terms below

⬇ Oral liq 250 mg per ml (100 mg elemental per ml) — 47.30 — 473 ml — Calcium carbonate PAI

39.00 — 500 ml — Roxane

➥ **Restricted** (RS1698)

Initiation

Only when prescribed for patients unable to swallow calcium carbonate tablets or where calcium carbonate tablets are inappropriate..

Antidiarrhoeals and Intestinal Anti-Inflammatory Agents

Antipropulsives

DIPHENOXYLATE HYDROCHLORIDE WITH ATROPINE SULPHATE

Tab 2.5 mg with atropine sulphate 25 mcg

LOPERAMIDE HYDROCHLORIDE

Tab 2 mg — 10.75 — 400 — Nodia

Cap 2 mg – **5% DV Feb-26 to 2028** — 12.00 — 400 — **Diamide Relief**

Rectal and Colonic Anti-Inflammatories

BUDESONIDE – **Restricted** see terms on the next page

⬇ Cap modified-release 3 mg – **5% DV Dec-25 to 2028** — 33.38 — 90 — **Budesonide Te Arai**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1723)

Initiation – Crohn's disease

Both:

1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
2 Any of the following:
- 2.1 Diabetes; or
- 2.2 Cushingoid habitus; or
- 2.3 Osteoporosis where there is significant risk of fracture; or
- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initiation – Collagenous and lymphocytic colitis (microscopic colitis)

Patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initiation – Gut Graft versus Host disease

Patient has gut Graft versus Host disease following allogenic bone marrow transplantation.

Initiation – non-cirrhotic autoimmune hepatitis

Re-assessment required after 6 months

All of the following:

1 Patient has autoimmune hepatitis*; and
2 Patient does not have cirrhosis; and
3 Any of the following:
- 3.1 Diabetes; or
- 3.2 Cushingoid habitus; or
- 3.3 Osteoporosis where there is significant risk of fracture; or
- 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
- 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indications marked with * are unapproved indications.

Continuation – non-cirrhotic autoimmune hepatitis

Re-assessment required after 6 months

Treatment remains appropriate and the patient is benefitting from the treatment.

HYDROCORTISONE ACETATE			
Rectal foam 10%, CFC free (14 applications)	57.09	15 g	Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE			
Topical Aerosol foam, 1% with pramoxine hydrochloride 1%			
MESALAZINE			
Tab EC 400 mg	49.50	100	Asacol
	71.00	90	Octasa
Tab long-acting 500 mg	56.10	100	Pentasa
Tab 800 mg	85.50	90	Asacol
Tab 1,600 mg	85.50	60	Asacol
Modified release granules 1 g	118.10	100 g	Pentasa
Suppos 500 mg	22.80	20	Asacol
Suppos 1 g	50.96	28	Pentasa
Enema 1 g per 100 ml	41.30	7	Pentasa

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
OLSALAZINE			
Tab 500 mg	93.37	100	Dipentum
Cap 250 mg	53.00	100	Dipentum
SODIUM CROMOGLICATE			
Cap 100 mg			
SULFASALAZINE			
Tab 500 mg	19.49	100	Salazopyrin
Tab EC 500 mg	20.54	100	Salazopyrin EN

Local Preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

CINCHOCAINE HYDROCHLORIDE WITH HYDROCORTISONE			
Oint 5 mg with hydrocortisone 5 mg per g	15.00	30 g	Proctosedyl
Suppos 5 mg with hydrocortisone 5 mg per g	9.90	12	Proctosedyl
FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE			
Oint 950 mcg with fluocortolone pivalate 920 mcg and cinchocaine hydrochloride 5 mg per g	13.05	30 g	Ultraproct
Suppos 630 mcg with fluocortolone pivalate 610 mcg and cinchocaine hydrochloride 1 mg	8.61	12	Ultraproct
PREDNISOLONE HEXANOATE WITH CINCHOCAINE HYDROCHORIDE			
Suppos 1.3 mg with cinchocaine hydrochloride 1 mg per g	8.61	12	Scheriproct

Management of Anal Fissures

GLYCERYL TRINITRATE			
Oint 0.2%	22.00	30 g	Rectogesic

Rectal Sclerosants

OILY PHENOL [PHENOL OILY]			
Inj 5%, 5 ml vial			

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – **5% DV Jul-26 to 2028**	11.99	10	**Glycopyrronium-AFT**
	19.00	5	Robinul
(Robinul Inj 200 mcg per ml, 1 ml ampoule to be delisted 1 July 2026)			
HYOSCINE BUTYLBROMIDE			
Tab 10 mg – **5% DV Apr-25 to 2027**	2.25	20	**Hyoscine Butylbromide (Adiramedica)**
Inj 20 mg, 1 ml ampoule – **5% DV Dec-23 to 2026**	1.91	5	**Spazmol**
MEBEVERINE HYDROCHLORIDE			
Tab 135 mg – **5% DV Dec-23 to 2026**	8.50	90	**Colofac**

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL			
Tab 200 mcg	47.73	120	Cytotec

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

H2 Antagonists

CIMETIDINE
- Tab 200 mg
- Tab 400 mg

FAMOTIDINE
- Tab 20 mg
- Tab 40 mg ..... 10.27 — 100 — Famotidine Hovid MY
- Inj 10 mg per ml, 2 ml vial
- Inj 10 mg per ml, 4 ml vial

RANITIDINE – **Restricted** see terms below
- ⬇ Tab 150 mg
- ⬇ Tab 300 mg
- ⬇ Inj 25 mg per ml, 2 ml ampoule

➥ **Restricted** (RS1703)
Initiation
Either:
1 For continuation use; or
2 Routine prevention of allergic reactions..

Proton Pump Inhibitors

LANSOPRAZOLE
- Cap 15 mg – **5% DV Feb-25 to 2027** ..... 1.21 — 30 — **Lanzol Relief**
- 4.04 — 100 — **Lanzol Relief**
- Cap 30 mg – **5% DV Feb-25 to 2027** ..... 5.43 — 100 — **Lanzol Relief**

OMEPRAZOLE
- ⬇ Tab dispersible 10 mg

➥ **Restricted** (RS1027)
Initiation
Only for use in tube-fed patients.
- ⬇ Tab dispersible 20 mg

➥ **Restricted** (RS1027)
Initiation
Only for use in tube-fed patients.
- Cap 10 mg – **5% DV Mar-24 to 2026** ..... 2.06 — 90 — Omeprazole Teva **Omeprazole actavis 10**
- Cap 20 mg – **5% DV Mar-24 to 2026** ..... 2.02 — 90 — Omeprazole Teva **Omeprazole actavis 20**
- Cap 40 mg – **5% DV Mar-24 to 2026** ..... 3.18 — 90 — Omeprazole Teva **Omeprazole actavis 40**
- Powder for oral liq ..... 52.00 — 5 g — Midwest
- Inf 40 mg vial – **5% DV Aug-26 to 2029** ..... 14.50 — 5 — **Omezol IV**
- Inj 40 mg ampoule with diluent ..... 37.38 — 5 — Dr Reddy's Omeprazole

PANTOPRAZOLE
- Tab EC 20 mg – **5% DV May-26 to 2028** ..... 1.81 — 90 — **Panzop Relief**
- Tab EC 40 mg – **5% DV May-26 to 2028** ..... 2.70 — 90 — **Panzop Relief**
- Inj 40 mg vial

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Site Protective Agents

COLLOIDAL BISMUTH SUBCITRATE
Tab 120 mg ..... 14.51 | 50 | Gastrodenol

SUCRALFATE
Tab 1 g

Bile and Liver Therapy

L-ORNITHINE L-ASPARTATE – **Restricted** see terms below
⬇ Grans for oral liquid 3 g
➥ **Restricted** (RS1261)
Initiation
For patients with chronic hepatic encephalopathy who have not responded to treatment with, or are intolerant to lactulose, or where lactulose is contraindicated.

RIFAXIMIN – **Restricted** see terms below
⬇ Tab 550 mg – **5% DV Dec-24 to 2027** ..... 625.00 | 56 | **Xifaxan**
➥ **Restricted** (RS1416)
Initiation
For patients with hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Diabetes

Alpha Glucosidase Inhibitors

ACARBOSE
Tab 50 mg – **5% DV Feb-25 to 2027** ..... 11.20 | 90 | **Accarb**
Tab 100 mg – **5% DV Feb-25 to 2027** ..... 17.38 | 90 | **Accarb**

Hyperglycaemic Agents

DIAZOXIDE – **Restricted** see terms below
⬇ Cap 25 mg ..... 110.00 | 100 | Proglicem
⬇ Cap 100 mg ..... 280.00 | 100 | Proglicem
⬇ Oral liq 50 mg per ml ..... 620.00 | 30 ml | Proglycem
➥ **Restricted** (RS1028)
Initiation
For patients with confirmed hypoglycaemia caused by hyperinsulinism.

GLUCAGON HYDROCHLORIDE
Inj 1 mg syringe kit ..... 32.00 | 1 | Glucagen Hypokit

GLUCOSE [DEXTROSE]
Tab 1.5 g
Tab 3.1 g
Tab 4 g
Oral soln 15 g per 80 ml sachet ..... 70.00 | 50 | HypoPak Glucose
Gel 40%

GLUCOSE WITH SUCROSE AND FRUCTOSE
Gel 19.7% with sucrose 35% and fructose 19.7%, 18 g sachet

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Insulin - Intermediate-Acting Preparations			
INSULIN ASPART WITH INSULIN ASPART PROTAMINE			
Inj insulin aspart 30% with insulin aspart protamine 70%, 100 u per ml, 3 ml prefilled pen	52.15	5	NovoMix 30 FlexPen
INSULIN DEGLUDEC WITH INSULIN ASPART			
Inj degludec 70 u with insulin aspart 30 u, 100 u per ml, 3 ml	80.00	5	Ryzodeg 70/30 Penfill
INSULIN ISOPHANE			
Inj insulin human 100 u per ml, 10 ml vial			
Inj insulin human 100 u per ml, 3 ml cartridge			
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj insulin lispro 25% with insulin lispro protamine 75%, 100 u per ml, 3 ml cartridge	42.66	5	Humalog Mix 25
Inj insulin lispro 50% with insulin lispro protamine 50%, 100 u per ml, 3 ml cartridge	42.66	5	Humalog Mix 50
INSULIN NEUTRAL WITH INSULIN ISOPHANE			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 10 ml vial			
Inj insulin neutral 30% with insulin isophane 70%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 40% with insulin isophane 60%, 100 u per ml, 3 ml cartridge			
Inj insulin neutral 50% with insulin isophane 50%, 100 u per ml, 3 ml cartridge			
Insulin - Long-Acting Preparations			
INSULIN GLARGINE			
Inj 100 u per ml, 3 ml disposable pen	94.50	5	Lantus SoloStar
Inj 100 u per ml, 3 ml cartridge	94.50	5	Lantus
Inj 100 u per ml, 10 ml vial	63.00	1	Lantus
Insulin - Rapid-Acting Preparations			
INSULIN ASPART			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
Inj 100 u per ml, 3 ml syringe	51.19	5	NovoRapid FlexPen
INSULIN GLULISINE			
Inj 100 u per ml, 10 ml vial	27.03	1	Apidra
Inj 100 u per ml, 3 ml cartridge	46.07	5	Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	Apidra Solostar
INSULIN LISPRO			
Inj 100 u per ml, 10 ml vial			
Inj 100 u per ml, 3 ml cartridge			
Insulin - Short-Acting Preparations			
INSULIN NEUTRAL			
Inj human 100 u per ml, 10 ml vial			
Inj human 100 u per ml, 3 ml cartridge			

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE			
Tab 5 mg	7.50	100	Daonil
GLICLAZIDE			
Tab 80 mg – **5% DV Feb-24 to 2026**	20.10	500	**Glizide**
GLIPIZIDE			
Tab 5 mg – **5% DV Mar-25 to 2027**	6.86	100	**Minidiab**
METFORMIN HYDROCHLORIDE			
Tab immediate-release 500 mg – **1% DV Mar-23 to 2027**	14.74	1,000	**Metformin Viatris**
Tab immediate-release 850 mg – **1% DV Aug-23 to 2027**	11.28	500	**Metformin Viatris**
PIOGLITAZONE			
Tab 15 mg – **5% DV Dec-24 to 2027**	6.15	90	**Vexazone**
Tab 30 mg – **5% DV Dec-24 to 2027**	7.25	90	**Vexazone**
Tab 45 mg – **5% DV Dec-24 to 2027**	12.00	90	**Vexazone**
VILDAGLIPTIN			
Tab 50 mg	35.00	60	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE			
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	Galvumet
GLP-1 Agonists			
DULAGLUTIDE – **Restricted** see terms below			
Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.			
⬇ Inj 1.5 mg per 0.5 ml prefilled pen	115.23	4	Trulicity

➥ **Restricted** (RS2135)

Initiation

Either:

1 For continuation use; or

2 All of the following:

2.1 Patient has type 2 diabetes; and

2.2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and

2.3 Any of the following:

2.3.1 Patient is Māori or any Pacific ethnicity*; or

2.3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or

2.3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or

2.3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or

2.3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack,

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause identified.

c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin / empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

LIRAGLUTIDE – **Restricted** see terms below

Note: Not to be given in combination with another funded GLP-1 agonist or empagliflozin / empagliflozin with metformin hydrochloride unless receiving empagliflozin / empagliflozin with metformin hydrochloride for the treatment of heart failure.

⬇ Inj 6 mg per ml, 3 ml prefilled pen ..... 383.72 3 Victoza

➥ **Restricted** (RS2136)

Initiation

Either:

1 For continuation use; or

2 All of the following:

2.1 Patient has type 2 diabetes; and

2.2 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of all of the following funded blood glucose lowering agents for a period of least 6 months, where clinically appropriate: empagliflozin, metformin, and vildagliptin; and

2.3 Any of the following:

2.3.1 Patient is Māori or any Pacific ethnicity*; or

2.3.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or

2.3.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or

2.3.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or

2.3.5 Patient has diabetic kidney disease (see note b)*.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause identified.

c) Funded GLP-1a treatment is not to be given in combination with funded (empagliflozin / empagliflozin with metformin hydrochloride) unless receiving funded (empagliflozin or empagliflozin in combination with metformin hydrochloride) for the treatment of heart failure.

SGLT2 Inhibitors

➥ **Restricted** (RS2069)

Initiation – heart failure reduced ejection fraction

All of the following:

1 Patient has heart failure; and

2 Patient is in NYHA functional class II or III or IV; and

continued…

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3 Either:
- 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%; or
- 3.2 An ECHO is not reasonably practicable, and in the opinion of the treating practitioner the patient would benefit from treatment; and

4 Patient is receiving concomitant optimal standard funded chronic heart failure treatment.

Initiation – Type 2 Diabetes

Any of the following:

1 For continuation use; or

2 Patient has previously had an initial approval for a GLP-1 agonist; or

3 All of the following:
- 3.1 Patient has type 2 diabetes; and
- 3.2 Any of the following:
  - 3.2.1 Patient is Māori or any Pacific ethnicity*; or
  - 3.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
  - 3.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
  - 3.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
  - 3.2.5 Patient has diabetic kidney disease (see note b)*; and
- 3.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

c) Funded [empagliflozin / empagliflozin with metformin hydrochloride] treatment is not to be given in combination with a funded GLP-1 unless receiving (empagliflozin / empagliflozin with metformin hydrochloride] for the treatment of heart failure.

EMPAGLIFLOZIN – **Restricted** see terms on the previous page

⬆	Tab 10 mg	58.56	30	Jardiance
⬆	Tab 25 mg	58.56	30	Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – **Restricted** see terms on the previous page

⬆	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
⬆	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
⬆	Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
⬆	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Digestives Including Enzymes

PANCREATIC ENZYME

Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	Creon 10000
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph Eur U)	34.93	20 g	Creon Micro
Powder pancreatin 60.12 mg (3,600 Ph. Eur. u/amylase, 5,000 Ph. Eur. u/lipase and 200 Ph. Eur. u/protease)			

URSODEOXYCHOLIC ACID – **Restricted** see terms below

⬇ Cap 250 mg – **5% DV Feb-24 to 2026**	33.95	100	**Ursosan**

➥ **Restricted** (RS2103)

Initiation – Alagille syndrome or progressive familial intrahepatic cholestasis

Either:

1 Patient has been diagnosed with Alagille syndrome; or
2 Patient has progressive familial intrahepatic cholestasis.

Initiation – Chronic severe drug induced cholestatic liver injury

All of the following:

1 Patient has chronic severe drug induced cholestatic liver injury; and
2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initiation – Primary biliary cholangitis

Both:

1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative by liver biopsy; and
2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initiation – Pregnancy

Patient diagnosed with cholestasis of pregnancy.

Initiation – Haematological transplant

Both:

1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
2 Treatment for up to 13 weeks.

Initiation – Total parenteral nutrition induced cholestasis

Both:

1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by TPN; and
2 Liver function has not improved with modifying the TPN composition.

Initiation – prevention of sinusoidal obstruction syndrome

The individual has leukaemia/lymphoma and requires prophylaxis for medications/therapies with a high risk of sinusoidal obstruction syndrome.

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Laxatives

Bowel-Cleansing Preparations

CITRIC ACID WITH MAGNESIUM CARBONATE HYDRATE AND SODIUM PICOSULFATE
- Powder for oral soln 12 g with magnesium carbonate hydrate 7.4 g and sodium picosulfate 10 mg per sachet — *e.g. PicoPrep Orange*

MACROGOL 3350 WITH ASCORBIC ACID, POTASSIUM CHLORIDE, SODIUM CHLORIDE AND CITRIC ACID WITH MAGNESIUM CARBONATE HYDRATE AND SODIUM PICOSULFATE
- Powder for oral soln 52.9 g with ascorbic acid 6 g, potassium chloride 740 mg, sodium chloride 2.6 g and sodium sulphate 5.6 g per sachet (1) and powder for oral soln citric acid 12 g with magnesium carbonate hydrate 7.4 g and sodium picosulfate 10 mg per sachet (2) — *e.g. Prepkit Orange*

MACROGOL 3350 WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE
- Powder for oral soln 755.68 mg with potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 70 g sachet – **5% DV Feb-25 to 2027** — 16.10 — 3 — **Glycoprep Orange**
- 64.32 — 12 — **Glycoprep Orange**
- Powder for oral soln 755.68 mg with potassium chloride 10.55 mg, sodium chloride 37.33 mg and sodium sulphate 80.62 mg per g, 210 g sachet — *e.g. Glycoprep Orange*

MACROGOL 3350 WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE WITH/WITHOUT SODIUM SULFATE, SODIUM ASCORBATE, ASCORBIC ACID
- Powd for oral soln 100g with potassium chloride 1g, sodium chloride 2g and sodium sulfate 9g per sach(1), powd for oral soln 40g with potassium chloride 1.2g and sodium chloride 3.2g per sach(1) and powd for oral soln ascorbic acid 7.54g and sodium ascorbate 48.11g per sach(1) – **5% DV Oct-23 to 2026** — 18.52 — 3 — **Plenvu**

Bulk-Forming Agents

ISPAGHULA (PSYLLIUM) HUSK
- Powder for oral soln — 22.10 — 500 g — Konsyl-D

STERCULIA WITH FRANGULA – **Restricted:** For continuation only
- ➥ Powder for oral soln

Faecal Softeners

DOCUSATE SODIUM
- Tab 50 mg – **5% DV Feb-24 to 2026** — 3.20 — 100 — **Coloxyl**
- Tab 120 mg – **5% DV Feb-24 to 2026** — 4.98 — 100 — **Coloxyl**

DOCUSATE SODIUM WITH SENNOSIDES
- Tab 50 mg with sennosides 8 mg – **5% DV May-26 to 2028** — 1.50 — 100 — **Solax**

PARAFFIN
- Oral liquid 1 mg per ml
- Enema 133 ml

POLOXAMER
- Oral drops 10% – **5% DV Feb-24 to 2026** — 4.17 — 30 ml — **Coloxyl**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE – **Restricted** see terms below

⬇ Inj 12 mg per 0.6 ml vial ..... 36.00 — 1 — Relistor
246.00 — 7 — Relistor

➥ **Restricted** (RS2057)

Initiation – Opioid induced constipation

Both:

1 The patient is receiving palliative care; and
2 Either:
  2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Initiation – Opioid induced constipation outside of palliative care

Limited to 14 days treatment

All of the following:

1 Individual has opioid induced constipation; and
2 Oral and rectal treatments for opioid induced constipation, including bowel-cleansing preparations, are ineffective or inappropriate; and
3 Mechanical bowel obstruction has been excluded.

Osmotic Laxatives

GLYCEROL

Suppos 2.8/4.0 g – **5% DV Feb-26 to 2028** ..... 12.39 — 20 — **Lax-suppositories Glycerol**

Note: DV limit applies to glycerol suppository presentations.

LACTULOSE

Oral liq 10 g per 15 ml – **5% DV Feb-26 to 2028** ..... 6.16 — 500 ml — **Laevolac**

MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE

Powder for oral soln 6.563 g with potassium chloride 23.3 mg, sodium bicarbonate 89.3 mg and sodium chloride 175.4 mg

Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg ..... 10.15 — 30 — Molaxole
12.19 — Movicol

SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE

Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml – **5% DV Feb-26 to 2028** ..... 36.89 — 50 — **Micolette**

SODIUM PHOSPHATE WITH PHOSPHORIC ACID

Oral liq 16.4% with phosphoric acid 25.14%

Enema 10% with phosphoric acid 6.58% ..... 3.70 — 1 — Fleet Phosphate Enema

Stimulant Laxatives

BISACODYL

Tab 5 mg – **5% DV Jul-26 to 2028** ..... 10.00 — 200 — Bisacodyl Viatris
6.28 — **Bisacodyl-AFT**

Suppos 10 mg – **5% DV Feb-25 to 2027** ..... 4.14 — 10 — **Lax-Suppositories**

(Bisacodyl Viatris Tab 5 mg to be delisted 1 July 2026)

SENNOSIDES

Tab 7.5 mg

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
SODIUM PICOSULFATE – **Restricted** see terms below			
⬇ Oral soln 7.5 mg per ml	7.40	30 ml	Dulcolax SP Drop

➥ **Restricted** (RS1843)

Initiation

Both:

1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
2 The patient would otherwise require a high-volume bowel cleansing preparation.

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA – **Restricted** see terms below			
⬇ Inj 50 mg vial	1,142.60	1	Myozyme

➥ **Restricted** (RS1793)

Initiation

Metabolic physician

Re-assessment required after 12 months

All of the following:

1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
2 Any of the following:
  2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
  2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
  2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
  2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Continuation

Metabolic physician

Re-assessment required after 12 months

All of the following:

1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
7 There is no evidence of new or progressive cardiomyopathy.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

ARGININE
- Tab 1,000 mg
- Cap 500 mg
- Powder
- Inj 500 mg per ml, 10 ml vial
- Inj 600 mg per ml, 25 ml vial

BETAINE – **Restricted** see terms below
⬇ Powder for oral soln.....575.00 180 g Cystadane
➥ **Restricted** (RS1794)
Initiation
Metabolic physician
Re-assessment required after 12 months
All of the following:
1 The patient has a confirmed diagnosis of homocystinuria; and
2 Any of the following:
2.1 A cystathionine beta-synthase (CBS) deficiency; or
2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
2.3 A disorder of intracellular cobalamin metabolism; and
3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.
Continuation
Metabolic physician
Re-assessment required after 12 months
The treatment remains appropriate and the patient is benefiting from treatment.

BIOTIN – **Restricted** see terms below
⬇ Cap 50 mg
⬇ Cap 100 mg
⬇ Inj 10 mg per ml, 5 ml vial
⬇ Inj 5 mg per ml, 1 ml ampoule
➥ **Restricted** (RS1330)
Metabolic physician or metabolic disorders dietitian

CARGLUMIC ACID – **Restricted** see terms below
⬇ Tab disp 200 mg
➥ **Restricted** (RS1831)
Initiation
Metabolic physician
For the acute in-patient treatment of organic acidaemias as an alternative to haemofiltration.

COENZYME Q10 – **Restricted** see terms below
⬇ Cap 120 mg
⬇ Cap 160 mg
➥ **Restricted** (RS1832)
Initiation
Metabolic physician
Re-assessment required after 6 months
The patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.
Continuation
Metabolic physician
Re-assessment required after 24 months
Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE – **Restricted** see terms below

⬇ Inj 1 mg per ml, 5 ml vial ..... 2,234.00 1 Naglazyme

➥ **Restricted** (RS1795)

Initiation

Metabolic physician

Re-assessment required after 12 months

Both:

1 The patient has been diagnosed with mucopolysaccharidosis VI; and
2 Either:
  2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency confirmed by either enzyme activity assay in leukocytes or skin fibroblasts; or
  2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Continuation

Metabolic physician

Re-assessment required after 12 months

All of the following:

1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

HAEM ARGINATE

Inj 25 mg per ml, 10 ml ampoule

IDURSULFASE – **Restricted** see terms below

⬇ Inj 2 mg per ml, 3 ml vial ..... 4,608.30 1 Elaprase

➥ **Restricted** (RS1546)

Initiation

Metabolic physician

Limited to 24 weeks treatment

All of the following:

1 The patient has been diagnosed with Hunter Syndrome (mucopolysacchardosis II); and
2 Either:
  2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE – **Restricted** see terms on the next page

⬇ Inj 100 U per ml, 5 ml vial ..... 1,335.16 1 Aldurazyme

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1607)
Initiation
Metabolic physician
Limited to 24 weeks treatment
All of the following:

1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
2 Either:
  2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE – **Restricted** see terms below

⬇	Tab 500 mg			
⬇	Cap 250 mg			
⬇	Cap 500 mg			
⬇	Oral liq 500 mg per 10 ml			
⬇	Oral soln 1,000 mg per 10 ml			
⬇	Oral soln 1,100 mg per 15 ml			
⬇	Inj 200 mg per ml, 5 ml vial			

➥ **Restricted** (RS1035)
Neurologist, metabolic physician or metabolic disorders dietitian

NITISINONE – **Restricted** see terms below

⬇	Cap 2 mg – **5% DV Jul-26 to 2028**	676.00	60	**Nitisinone LogixX Pharma**
⬇	Cap 5 mg – **5% DV Jul-26 to 2028**	1,302.00	60	**Nitisinone LogixX Pharma**
⬇	Cap 10 mg – **5% DV Jul-26 to 2028**	1,704.00	60	**Nitisinone LogixX Pharma**

➥ **Restricted** (RS2164)
Initiation
Patient requires nitisinone for the management of inherited metabolic disorders.

PYRIDOXAL-5-PHOSPHATE – **Restricted** see terms below

⬇	Tab 50 mg			

➥ **Restricted** (RS1331)
Neurologist, metabolic physician or metabolic disorders dietitian

RIBOFLAVIN – **Restricted** see terms below

⬇	Tab 100 mg			
⬇	Cap 100 mg			

➥ **Restricted** (RS1833)
Initiation
Metabolic physician or neurologist
Re-assessment required after 6 months
The patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation

Metabolic physician or neurologist

Re-assessment required after 24 months

Both:

1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE – **Restricted** see terms below

⬇ Tab soluble 100 mg ..... 1,452.70 30 Kuvan

➥ **Restricted** (RS1796)

Initiation

Metabolic physician

Re-assessment required after 1 month

All of the following:

1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
4 Sapropterin to be used alone or in combination with PKU dietary management; and
5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Continuation

Metabolic physician

Re-assessment required after 12 months

All of the following:

1 Either:
  1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
  1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
2 Any of the following:
  2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
4 Sapropterin to be used alone or in combination with PKU dietary management; and
5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE

Cap 500 mg
Powder
Soln 100 mg per ml
Inj 20%, 10 ml ampoule

SODIUM PHENYLBUTYRATE – **Some items restricted** see terms on the next page

Tab 500 mg

⬇ Grans 483 mg per g ..... 2,016.00 174 g Pheburane

Oral liq 250 mg per ml
Inj 200 mg per ml, 10 ml ampoule

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1797)

Initiation

Metabolic physician

Re-assessment required after 12 months

For the chronic management of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Continuation

Metabolic physician

Re-assessment required after 12 months

The treatment remains appropriate and the patient is benefiting from treatment.

TALIGLUCERASE ALFA – **Restricted** see terms below

⬇	Inj 200 unit vial	1,072.00	1	Elelyso

➥ **Restricted** (RS1897)

Initiation

Metabolic physician

Re-assessment required after 12 months

All of the following:

1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
3 Any of the following:
    3.1 Patient has haematological complications of Gaucher disease; or
    3.2 Patient has skeletal complications of Gaucher disease; or
    3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
    3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
    3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Continuation

Metabolic physician or any relevant practitioner on the recommendation of a metabolic physician

Re-assessment required after 3 years

All of the following:

1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
3 RRadiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

TAURINE – **Restricted** see terms below
- ⬇ Cap 500 mg
- ⬇ Cap 1,000 mg
- ⬇ Powder

➥ **Restricted** (RS1834)
Initiation
Metabolic physician
Re-assessment required after 6 months
The patient has a suspected specific mitochondrial disorder that may respond to taurine supplementation.
Continuation
Metabolic physician
Re-assessment required after 24 months
Both:
1 The patient has a confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
2 The treatment remains appropriate and the patient is benefiting from treatment.

TRIENTINE – **Restricted** see terms below
⬇ Cap 250 mg ..... 2,022.00 100 Trientine Waymade

➥ **Restricted** (RS2026)
Initiation
All of the following:
1 Patient has confirmed Wilson disease; and
2 Treatment with D-penicillamine has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit; and
3 Treatment with zinc has been trialled and discontinued because the person has experienced intolerable side effects or has not received sufficient benefit, or zinc is considered clinically inappropriate as the person has symptomatic liver disease and requires copper chelation.

Minerals

Calcium

CALCIUM CARBONATE
Tab 1.25 g (500 mg elemental) – **5% DV Feb-24 to 2026** ..... 7.28 250 **Calci-Tab 500**
Tab eff 1.25 g (500 mg elemental)
Tab eff 1.75 g (1 g elemental)

Copper

➥ **Restricted** (RS1928)
Initiation – Moderate to severe burns
Limited to 3 months treatment
Both:
1 Patient has been hospitalised with moderate to severe burns; and
2 Treatment is recommended by a National Burns Unit specialist.

COPPER – **Restricted** see terms above
⬆ Tab 2.5 mg, chelated

COPPER CHLORIDE – **Restricted** see terms above
⬆ Inj 0.4 mg per ml, 10 ml vial

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Fluoride

SODIUM FLUORIDE
Tab 1.1 mg (0.5 mg elemental)

Iodine

POTASSIUM IODATE
Tab 253 mcg (150 mcg elemental iodine) – **5% DV Feb-24 to 2026** ..... 5.99 — 90 — **NeuroTabs**

POTASSIUM IODATE WITH IODINE
Oral liq 10% with iodine 5%

Iron

FERRIC DERISOMALTOSE – **Restricted** see terms below
⬇ Inj 500 mg per 5 ml vial ..... 249.99 — 1 — Monofer
➥ **Restricted** (RS2175)
Initiation
Patient had previously developed iron-infusion related hypophosphataemia or other severe adverse reaction.

FERROUS FUMARATE
Tab 200 mg (65.7 mg elemental) – **5% DV Feb-25 to 2027** ..... 3.49 — 100 — **Ferro-tab**

FERROUS FUMARATE WITH FOLIC ACID
Tab 310 mg (100 mg elemental) with folic acid 350 mcg – **5% DV Dec-24 to 2027** ..... 5.98 — 100 — **Ferro-F-Tabs**

FERROUS GLUCONATE WITH ASCORBIC ACID
Tab 170 mg (20 mg elemental) with ascorbic acid 40 mg

FERROUS SULFATE
Oral liq 30 mg (6 mg elemental) per ml – **5% DV Feb-26 to 2028** ..... 10.25 — 250 ml — **Ferro-Liquid**

FERROUS SULFATE WITH ASCORBIC ACID
Tab long-acting 325 mg (105 mg elemental) with ascorbic acid 500 mg

IRON (AS FERRIC CARBOXYMALTOSE) – **Restricted** see terms below
⬇ Inj 50 mg per ml, 10 ml vial ..... 150.00 — 1 — Ferinject
➥ **Restricted** (RS1417)
Initiation
Treatment with oral iron has proven ineffective or is clinically inappropriate.

IRON (AS SUCROSE)
Inj 20 mg per ml, 5 ml ampoule ..... 100.00 — 5 — Venofer
(Venofer Inj 20 mg per ml, 5 ml ampoule to be delisted 1 September 2026)

IRON POLYMALTOSE
Inj 50 mg per ml, 2 ml ampoule ..... 41.75 — 5 — Ferrosig

Magnesium

MAGNESIUM AMINO ACID CHELATE
Cap 750 mg (150 mg elemental)

MAGNESIUM CHLORIDE
Inj 1 mmol per 1 ml, 100 ml bag

MAGNESIUM HYDROXIDE
Tab 311 mg (130 mg elemental)
Suspension 8%

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
MAGNESIUM OXIDE			
Cap 663 mg (400 mg elemental)			
Cap 696 mg (420 mg elemental)			
MAGNESIUM OXIDE WITH MAGNESIUM ASPARTATE, MAGNESIUM AMINO ACID CHELATE AND MAGNESIUM CITRATE			
Cap 500 mg with magnesium aspartate 100 mg, magnesium amino acid chelate 100 mg and magnesium citrate 100 mg (345 mg elemental magnesium)			
Cap 500 mg with magnesium aspartate 100 mg, magnesium amino acid chelate 100 mg and magnesium citrate 100 mg (360 mg elemental magnesium)			
(Any Cap 500 mg with magnesium aspartate 100 mg, magnesium amino acid chelate 100 mg and magnesium citrate 100 mg (360 mg elemental magnesium) to be delisted 1 November 2026)			
MAGNESIUM SULPHATE			
Inj 100 mg per ml, 40 ml bag			
Inj 0.4 mmol per ml, 250 ml bag			
Inj 2 mmol per ml, 10 ml ampoule	75.06	10	Inresa
Inj 2 mmol per ml, 5 ml ampoule – **5% DV Jun-24 to 2026**	37.53	10	**Martindale**
Inj 100 mg per ml, 50 ml bag			

Selenium

SELENIUM – **Restricted** see terms below

⬇ Oral liq 150 mcg per 3 drops — *e.g. Clinicians selenium oral drops*

⬇ Inj 300 mcg per ml, 1 ml ampoule

➥ **Restricted** (RS1929)

Initiation – Moderate to severe burns

Limited to 3 months treatment

Both:

1 Patient has been hospitalised with moderate to severe burns; and
2 Treatment is recommended by a National Burns Unit specialist.

Zinc

ZINC			
Oral liq 5 mg per 5 drops			
ZINC CHLORIDE			
Inj 5.3 mg per ml (5.1 mg per ml elemental), 2 ml ampoule			
ZINC SULPHATE			
Cap 220 mg (50 mg elemental)	29.14	100	Rugby
Cap 137.4 mg (50 mg elemental)	11.00	100	Zincaps

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE
- Soln 0.15%
- Spray 0.15%
- Spray 0.3%

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
BENZYDAMINE HYDROCHLORIDE WITH CETYLPYRIDINIUM CHLORIDE			
Lozenge 3 mg with cetylpyridinium chloride			
CARBOXYMETHYLCELLULOSE			
Oral spray			
CARMELLOSE SODIUM WITH PECTIN AND GELATINE			
Paste			
Powder			
CHLORHEXIDINE GLUCONATE			
Mouthwash 0.2% – **5% DV Jan-25 to 2027**	3.99	200 ml	**healthE**
DICHLOROBENZYL ALCOHOL WITH AMYLMETACRESOL			
Lozenge 1.2 mg with amylmetacresol 0.6 mg			
TRIAMCINOLONE ACETONIDE			
Paste 0.1% – **5% DV Feb-24 to 2026**	5.49	5 g	**Kenalog in Orabase**

Oropharyngeal Anti-Infectives

AMPHOTERICIN B			
Lozenge 10 mg	5.86	20	Fungilin
MICONAZOLE			
Oral gel 20 mg per g – **5% DV Feb-25 to 2027**	5.19	40 g	**Decozol**
NYSTATIN			
Oral liquid 100,000 u per ml – **5% DV Feb-24 to 2026**	2.22	24 ml	**Nilstat**

Other Oral Agents

HYALURONIC ACID WITH LIDOCAINE [LIGNOCAINE]			
Inj 20 mg per ml			
SILVER DIAMINE FLUORIDE			
Oral application 38%	139.00	5 ml	Topamine

SODIUM HYALURONATE [HYALURONIC ACID] – **Restricted** see terms below

⬇ Inj 20 mg per ml, 1 ml syringe

➥ **Restricted** (RS1175)

Otolaryngologist

Vitamins

Multivitamin Preparations

MULTIVITAMIN AND MINERAL SUPPLEMENT – **Restricted** see terms below			
⬇ Cap	23.35	180	Clinicians Multivit & Mineral Boost

➥ **Restricted** (RS1498)

Initiation

Limited to 3 months treatment

Both:

1 Patient was admitted to hospital with burns; and

2 Any of the following:

2.1 Burn size is greater than 15% of total body surface area (BSA) for all types of burns; or

2.2 Burn size is greater than 10% of BSA for mid-dermal or deep dermal burns; or

2.3 Nutritional status prior to admission or dietary intake is poor.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
MULTIVITAMIN RENAL – **Restricted** see terms below			
⬇ Cap	7.28	30	Clinicians Renal Vit

➥ **Restricted** (RS1499)

Initiation

Either:

1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or

2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73m^2 body surface area (BSA).

MULTIVITAMINS			
Tab (BPC cap strength)	24.00	1,000	Mvite
⬇ cap vitamin A 2500 u, betacarotene 3 mg, cholecalciferol 11 mcg, alpha tocopherol 150 u, phytomenadione 150 mcg, folic acid 0.2 mg, ascorbic acid 100 mg, thiamine 1.5 mg, pantothenic acid 12 mg, riboflavin 1.7 mg, niacin 20 mg, pyridoxine hydrochloride 1.9 mg, cyanocobalamin 3 mg, zinc 7.5 mg and biotin 100 mcg			*e.g. Vitabdeck*

➥ **Restricted** (RS1620)

Initiation

Any of the following:

1 Patient has cystic fibrosis with pancreatic insufficiency; or

2 Patient is an infant or child with liver disease or short gut syndrome; or

3 Patient has severe malabsorption syndrome.

⬇ Powder vitamin A 3200 mcg with vitamin D 100 mcg, vitamin E 54.2 mg, vitamin C 400 mg, vitamin K1 108 mcg thiamine 3.2 mg, riboflavin 4.4 mg, niacin 41 mg, vitamin B6 3.6 mg, folic acid 600 mcg, vitamin B12 9 mcg, biotin 120 mcg, pantothenic acid 24 mg, choline 1250 mg and inositol 700 mg	74.88	200 g	Paediatric Seravit

➥ **Restricted** (RS1178)

Initiation

Patient has inborn errors of metabolism.

Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg and glucose 1000 mg, 5 ml ampoule (1)			*e.g. Pabrinex IV*
Inj thiamine hydrochloride 250 mg with riboflavin 4 mg and pyridoxine hydrochloride 50 mg, 5 ml ampoule (1) and inj ascorbic acid 500 mg with nicotinamide 160 mg, 2 ml ampoule (1)			
Inj thiamine hydrochloride 500 mg with riboflavin 8 mg and pyridoxine hydrochloride 100 mg, 10 ml ampoule (1) and inj ascorbic acid 1000 mg with nicotinamide 320 mg and glucose 2000 mg, 10 ml ampoule (1)			

Vitamin A

RETINOL

Tab 10,000 iu

Cap 25,000 iu

Oral liq 150,000 iu per ml

Oral liq 666.7 mcg per 2 drops, 10 ml

Oral liq 5,000 iu per drop, 30 ml

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Vitamin B			
HYDROXOCOBALAMIN			
Inj 1 mg per ml, 1 ml ampoule – **5% DV Jul-25 to 2027**	3.95	3	**Hydroxocobalamin Panpharma**
PYRIDOXINE HYDROCHLORIDE			
Tab 25 mg – **5% DV Feb-24 to 2026**	3.43	90	**Vitamin B6 25**
Tab 50 mg	23.45	500	Pyridoxine multichem
Inj 100 mg per ml, 2 ml vial			
Inj 100 mg per ml, 1 ml ampoule			
Inj 100 mg per ml, 30 ml vial			
THIAMINE HYDROCHLORIDE			
Tab 50 mg	4.65	100	Thiamine multichem
Tab 100 mg			
Inj 100 mg per ml, 1 ml vial			*e.g. Benerva*
Inj 50 mg per ml, 2 ml ampoule – **5% DV Sep-26 to 2028**	49.95	10	**Thiamine Sterop**
Inj 100 mg per ml, 2 ml vial			
Inj 125 mg per ml, 2 ml vial			
VITAMIN B COMPLEX			
Tab strong, BPC	11.25	500	Bplex
Vitamin C			
ASCORBIC ACID			
Tab 100 mg – **5% DV Mar-26 to 2028**	16.00	500	**Cvite**
Tab chewable 250 mg			
Vitamin D			
ALFACALCIDOL			
Cap 0.25 mcg	26.32	100	One-Alpha
Cap 1 mcg	87.98	100	One-Alpha
Oral drops 2 mcg per ml	60.68	20 ml	One-Alpha
CALCITRIOL			
Cap 0.25 mcg	7.89	100	Calcitriol XL Calcitriol-AFT
Cap 0.5 mcg	13.68	100	Calcitriol XL Calcitriol-AFT
Oral liq 1 mcg per ml			
Inj 1 mcg per ml, 1 ml ampoule			
COLECALCIFEROL			
Cap 1.25 mg (50,000 iu) – **5% DV Jun-24 to 2026**	3.65	12	**Vit.D3**
Oral liq 188 mcg per ml (7,500 iu per ml)	9.00	5 ml	Clinicians
Vitamin E			
ALPHA TOCOPHERYL – **Restricted** see terms below			
⬇ Oral liq 156 u per ml			
➥ **Restricted** (RS1632)			
Initiation – Cystic fibrosis			
Both:			

continued…

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Cystic fibrosis patient; and
2 Either:
  2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
  2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

Initiation – Osteoradionecrosis
For the treatment of osteoradionecrosis.
Initiation – Other indications
All of the following:

1 Infant or child with liver disease or short gut syndrome; and
2 Requires vitamin supplementation; and
3 Either:
  3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
  3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

ALPHA TOCOPHERYL ACETATE – **Restricted** see terms below
⬇ Cap 100 u
⬇ Cap 500 u
⬇ Oral liq 156 u per ml
➥ **Restricted** (RS1176)
Initiation – Cystic fibrosis
Both:

1 Cystic fibrosis patient; and
2 Either:
  2.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck); or
  2.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for the patient.

Initiation – Osteoradionecrosis
For the treatment of osteoradionecrosis.
Initiation – Other indications
All of the following:

1 Infant or child with liver disease or short gut syndrome; and
2 Requires vitamin supplementation; and
3 Either:
  3.1 Patient has tried and failed the other available funded fat soluble vitamin A,D,E,K supplements (Vitabdeck); or
  3.2 The other available funded fat soluble vitamin A,D,E,K supplement (Vitabdeck) is contraindicated or clinically inappropriate for patient.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

EPOETIN ALFA – **Restricted** see terms below

⬇	Inj 1,000 iu in 0.5 ml syringe	250.00	6	Binocrit
⬇	inj 2,000 iu in 1 ml syringe	100.00	6	Binocrit
⬇	Inj 3,000 iu in 0.3 ml syringe	150.00	6	Binocrit
⬇	Inj 4,000 iu in 0.4 ml syringe	96.50	6	Binocrit
⬇	Inj 5,000 iu in 0.5 ml syringe	125.00	6	Binocrit
⬇	Inj 6,000 iu in 0.6 ml syringe	145.00	6	Binocrit
⬇	Inj 8,000 iu in 0.8 ml syringe	175.00	6	Binocrit
⬇	Inj 10,000 iu in 1 ml syringe	197.50	6	Binocrit
⬇	Inj 40,000 iu in 1 ml syringe	250.00	1	Binocrit

➥ **Restricted** (RS1660)

Initiation – chronic renal failure

All of the following:

1 Patient in chronic renal failure; and
2 Haemoglobin is less than or equal to 100g/L; and
3 Any of the following:
  3.1 Both:
    3.1.1 Patient does not have diabetes mellitus; and
    3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  3.2 Both:
    3.2.1 Patient has diabetes mellitus; and
    3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  3.3 Patient is on haemodialysis or peritoneal dialysis.

Initiation – myelodysplasia*

Re-assessment required after 2 months

All of the following:

1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
5 Patient has a serum epoetin level of < 500 IU/L; and
6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Continuation – myelodysplasia*

Re-assessment required after 12 months

All of the following:

1 The patient's transfusion requirement continues to be reduced with epoetin treatment; and
2 Transformation to acute myeloid leukaemia has not occurred; and
3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Initiation – all other indications

Haematologist

For use in patients where blood transfusion is not a viable treatment alternative.

Note: Indications marked with * are unapproved indications

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

EPOETIN BETA – **Restricted** see terms below

Note: Epoetin beta is considered a Discretionary Variance Pharmaceutical for epoetin alfa.

⬇ Inj 2,000 iu in 0.3 ml syringe
⬇ Inj 3,000 iu in 0.3 ml syringe
⬇ Inj 4,000 iu in 0.3 ml syringe
⬇ Inj 5,000 iu in 0.3 ml syringe
⬇ Inj 6,000 iu in 0.3 ml syringe
⬇ Inj 10,000 iu in 0.6 ml syringe

➥ **Restricted** (RS1661)

Initiation – chronic renal failure

All of the following:

1 Patient in chronic renal failure; and
2 Haemoglobin is less than or equal to 100g/L; and
3 Any of the following:
  3.1 Both:
    3.1.1 Patient does not have diabetes mellitus; and
    3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  3.2 Both:
    3.2.1 Patient has diabetes mellitus; and
    3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  3.3 Patient is on haemodialysis or peritoneal dialysis.

Initiation – myelodysplasia*

Re-assessment required after 12 months

All of the following:

1 Patient has a confirmed diagnosis of myelodysplasia (MDS); and
2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
3 Patient has very low, low or intermediate risk MDS based on the WHO classification-based prognostic scoring system for myelodysplastic syndrome (WPSS); and
4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
5 Patient has a serum epoetin level of < 500 IU/L; and
6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Continuation – myelodysplasia*

Re-assessment required after 2 months

All of the following:

1 The patient's transfusion requirement continues to be reduced with epoetin treatment; and
2 Transformation to acute myeloid leukaemia has not occurred; and
3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Initiation – all other indications

All of the following:

1 Haematologist; and
2 For use in patients where blood transfusion is not a viable treatment alternative; and
3 *Note: Indications marked with * are unapproved indications.

Megaloblastic

FOLIC ACID

Tab 0.8 mg	26.60	1,000	Folic Acid multichem
Tab 5 mg – **1% DV Mar-23 to 2027**	5.82	100	**Folic Acid Viatris**
Oral liq 50 mcg per ml	31.77	25 ml	Biomed
Inj 5 mg per ml, 10 ml vial			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Antifibrinolytics, Haemostatics and Local Sclerosants

ALUMINIUM CHLORIDE – **Restricted** see terms below

⬇ Topical soln 20% w/v *e.g. Driclor*

➥ **Restricted** (RS1500)

Initiation

For use as a haemostatis agent.

APROTININ – **Restricted** see terms below

⬇ Inj 10,000 kIU per ml (equivalent to 200 mg per ml), 50 ml vial

➥ **Restricted** (RS1332)

Initiation

Cardiac anaesthetist

Either:

1 Paediatric patient undergoing cardiopulmonary bypass procedure; or
2 Adult patient undergoing cardiac surgical procedure where the significant risk of massive bleeding outweighs the potential adverse effects of the drug.

ELTROMBOPAG – **Restricted** see terms below

⬇ Tab 25 mg ..... 1,550.00 28 Revolade

⬇ Tab 50 mg ..... 3,100.00 28 Revolade

➥ **Restricted** (RS1648)

Initiation – idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

Re-assessment required after 6 weeks

All of the following:

1 Patient has had a splenectomy; and
2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
3 Any of the following:
  3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
  3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initiation – idiopathic thrombocytopenic purpura - preparation for splenectomy

Haematologist

Limited to 6 weeks treatment

The patient requires eltrombopag treatment as preparation for splenectomy.

Continuation – idiopathic thrombocytopenic purpura - post-splenectomy

Haematologist

Re-assessment required after 12 months

The patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre

Initiation – idiopathic thrombocytopenic purpura contraindicated to splenectomy

Haematologist

Re-assessment required after 3 months

All of the following:

1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
3 Either:
3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Continuation – idiopathic thrombocytopenic purpura contraindicated to splenectomy
Haematologist
Re-assessment required after 12 months
All of the following:

1 The patient's significant contraindication to splenectomy remains; and
2 The patient has obtained a response from treatment during the initial approval period; and
3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
4 Further treatment with eltrombopag is required to maintain response.

Initiation – severe aplastic anaemia
Haematologist
Re-assessment required after 3 months
Both:

1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
2 Either:
2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Continuation – severe aplastic anaemia
Haematologist
Re-assessment required after 12 months
Both:

1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB – **Restricted** see terms below

⬇	Inj 30 mg in 1 ml vial	3,570.00	1	Hemlibra
⬇	Inj 60 mg in 0.4 ml vial	7,138.00	1	Hemlibra
⬇	Inj 105 mg in 0.7 ml vial	12,492.00	1	Hemlibra
⬇	Inj 150 mg in 1 ml vial	17,846.00	1	Hemlibra

➥ **Restricted** (RS1998)

Initiation – Severe Haemophilia A with or without FVIII inhibitors
Haematologist
Both:

1 Patient has severe congenital haemophilia A with a severe bleeding phenotype (endogenous factor VIII activity less than or equal to 2%); and
2 Emicizumab is to be administered at a dose of no greater than 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

FERRIC SUBSULFATE
Gel 25.9%
Soln 500 ml

POLIDOCANOL
Inj 0.5%, 30 ml vial

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
SODIUM TETRADECYL SULPHATE			
Inj 3%, 2 ml ampoule			
THROMBIN			
Powder			
TRANEXAMIC ACID			
Tab 500 mg – **5% DV May-26 to 2028**	9.93	60	**Mercury Pharma**
Inj 100 mg per ml, 5 ml ampoule – **5% DV Mar-25 to 2027**	5.39	5	**Tranexamic-AFT**
Inj 100 mg per ml, 10 ml ampoule – **5% DV Mar-25 to 2027**	7.99	5	**Tranexamic-AFT**

Anticoagulant Reversal Agents

IDARUCIZUMAB – **Restricted** see terms below

⬇ Inj 50 mg per ml, 50 ml vial	4,250.00	2	Praxbind

➥ **Restricted** (RS1535)

Initiation

For the reversal of the anticoagulant effects of dabigatran when required in situations of life-threatening or uncontrolled bleeding, or for emergency surgery or urgent procedures.

Blood Factors

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – **Restricted** see terms below

⬇ Inj 250 iu vial	612.50	1	Alprolix
⬇ Inj 500 iu vial	1,225.00	1	Alprolix
⬇ Inj 1,000 iu vial	2,450.00	1	Alprolix
⬇ Inj 2,000 iu vial	4,900.00	1	Alprolix
⬇ Inj 3,000 iu vial	7,350.00	1	Alprolix
⬇ Inj 4,000 iu vial	9,800.00	1	Alprolix

➥ **Restricted** (RS1684)

Initiation

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] – **Restricted** see terms below

⬇ Inj 1 mg syringe	1,178.30	1	NovoSeven RT
⬇ Inj 2 mg syringe	2,356.60	1	NovoSeven RT
⬇ Inj 5 mg syringe	5,891.50	1	NovoSeven RT
⬇ Inj 8 mg syringe	9,426.40	1	NovoSeven RT

➥ **Restricted** (RS1704)

Initiation

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

FACTOR EIGHT INHIBITOR BYPASSING FRACTION – **Restricted** see terms below

⬇ Inj 500 U	1,315.00	1	FEIBA NF
⬇ Inj 1,000 U	2,630.00	1	FEIBA NF
⬇ Inj 2,500 U	6,575.00	1	FEIBA NF

➥ **Restricted** (RS1705)

Initiation

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – **Restricted** see terms below			
⬇ Inj 250 iu prefilled syringe	287.50	1	Xyntha
⬇ Inj 500 iu prefilled syringe	575.00	1	Xyntha
⬇ Inj 1,000 iu prefilled syringe	1,150.00	1	Xyntha
⬇ Inj 2,000 iu prefilled syringe	2,300.00	1	Xyntha
⬇ Inj 3,000 iu prefilled syringe	3,450.00	1	Xyntha

➥ **Restricted** (RS1706)

Initiation

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

NONACOG GAMMA, [RECOMBINANT FACTOR IX] – **Restricted** see terms below			
⬇ Inj 1,000 iu vial	870.00	1	RIXUBIS
⬇ Inj 2,000 iu vial	1,740.00	1	RIXUBIS
⬇ Inj 3,000 iu vial	2,610.00	1	RIXUBIS

➥ **Restricted** (RS1679)

Initiation

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – **Restricted** see terms below			
⬇ Inj 500 iu vial	420.00	1	Advate
⬇ Inj 1,000 iu vial	840.00	1	Advate
⬇ Inj 2,000 iu vial	1,680.00	1	Advate
⬇ Inj 3,000 iu vial	2,520.00	1	Advate

➥ **Restricted** (RS1707)

Initiation

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) – **Restricted** see terms below			
⬇ Inj 250 iu vial	237.50	1	Kogenate FS
⬇ Inj 500 iu vial	475.00	1	Kogenate FS
⬇ Inj 1,000 iu vial	950.00	1	Kogenate FS
⬇ Inj 2,000 iu vial	1,900.00	1	Kogenate FS
⬇ Inj 3,000 iu vial	2,850.00	1	Kogenate FS

➥ **Restricted** (RS1708)

Initiation

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] – **Restricted** see terms below			
⬇ Inj 1,000 iu vial	1,200.00	1	Adynovate
⬇ Inj 2,000 iu vial	2,400.00	1	Adynovate

➥ **Restricted** (RS1682)

Initiation

For patients with haemophilia A receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Vitamin K

PHYTOMENADIONE			
Inj 2 mg in 0.2 ml ampoule	8.00	5	Konakion MM Paediatric
Inj 10 mg per ml, 1 ml ampoule	9.21	5	Konakion MM

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Antithrombotics

Anticoagulants

BIVALIRUDIN – **Restricted** see terms below
⬇ Inj 250 mg vial
➥ **Restricted** (RS1181)
Initiation
Either:
1 For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance; or
2 For use in patients undergoing endovascular procedures.

CITRATE SODIUM
Inj 4% (200 mg per 5 ml), 5 ml ampoule
Inj 46.7% (1.4 g per 3 ml), 3 ml syringe
Inj 46.7% (2.36 g per 5 ml), 5 ml ampoule

DABIGATRAN

Cap 75 mg – **5% DV Jul-24 to 2026**	27.99	60	**Pradaxa**
Cap 110 mg – **5% DV Jul-24 to 2026**	27.99	60	**Pradaxa**
Cap 150 mg – **5% DV Jul-24 to 2026**	27.99	60	**Pradaxa**

DANAPAROID – **Restricted** see terms below
⬇ Inj 750 u in 0.6 ml ampoule
➥ **Restricted** (RS1182)
Initiation
For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance.

DEFIBROTIDE – **Restricted** see terms below
⬇ Inj 80 mg per ml, 2.5 ml ampoule
➥ **Restricted** (RS1183)
Initiation
Haematologist
Patient has moderate or severe sinusoidal obstruction syndrome as a result of chemotherapy or regimen-related toxicities.

DEXTROSE WITH SODIUM CITRATE AND CITRIC ACID [ACID CITRATE DEXTROSE A]
Inj 24.5 mg with sodium citrate 22 mg and citric acid 7.3 mg per ml, 100 ml bag

ENOXAPARIN SODIUM

Inj 20 mg in 0.2 ml syringe – **5% DV Feb-25 to 2027**	21.90	10	**Clexane**
Inj 40 mg in 0.4 ml ampoule			
Inj 40 mg in 0.4 ml syringe – **5% DV Feb-25 to 2027**	29.74	10	**Clexane**
Inj 60 mg in 0.6 ml syringe – **5% DV Feb-25 to 2027**	42.47	10	**Clexane**
Inj 80 mg in 0.8 ml syringe – **5% DV Feb-25 to 2027**	56.62	10	**Clexane**
Inj 100 mg in 1 ml syringe – **5% DV Feb-25 to 2027**	70.91	10	**Clexane**
Inj 120 mg in 0.8 ml syringe – **5% DV Feb-25 to 2027**	88.11	10	**Clexane Forte**
Inj 150 mg in 1 ml syringe – **5% DV Feb-25 to 2027**	100.70	10	**Clexane Forte**

FONDAPARINUX SODIUM – **Restricted** see terms below
⬇ Inj 2.5 mg in 0.5 ml syringe
⬇ Inj 7.5 mg in 0.6 ml syringe
➥ **Restricted** (RS1184)
Initiation
For use in heparin-induced thrombocytopaenia, heparin resistance or heparin intolerance.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
HEPARIN SODIUM			
Inj 10 iu per ml, 5 ml ampoule (flushing solution)	19.38	10	Wockhardt
Inj 100 iu per ml, 250 ml bag			
Inj 1,000 iu per ml, 1 ml ampoule	362.98	50	Hospira
Inj 1,000 iu per ml, 5 ml ampoule	164.40	50	Pfizer
Inj 5,000 iu in 0.2 ml ampoule			
Inj 5,000 iu per ml, 1 ml ampoule	70.33	5	Hospira
Inj 5,000 iu per ml, 5 ml ampoule – **5% DV May-26 to 2028**	406.15	50	**Pfizer**
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml ampoule	96.91	50	Pfizer
Inj 100 iu per ml, 2 ml ampoule			
Inj 100 iu per ml, 5 ml ampoule			
PHENINDIONE			
Tab 10 mg			
Tab 25 mg			
Tab 50 mg			
PROTAMINE SULPHATE			
Inj 10 mg per ml, 5 ml ampoule			
RIVAROXABAN			
Tab 10 mg – **5% DV Dec-23 to 2026**	15.60	30	**Xarelto**
Tab 15 mg – **5% DV Dec-23 to 2026**	14.56	28	**Xarelto**
Tab 20 mg – **5% DV Dec-23 to 2026**	14.56	28	**Xarelto**
SODIUM CITRATE WITH SODIUM CHLORIDE AND POTASSIUM CHLORIDE			
Inj 4.2 mg with sodium chloride 5.7 mg and potassium chloride 74.6 mcg per ml, 5,000 ml bag			
WARFARIN SODIUM			
Tab 1 mg	7.50	100	Marevan
Tab 2 mg			
Tab 3 mg	12.00	100	Marevan
Tab 5 mg	13.50	100	Marevan

Antiplatelets

ASPIRIN			
Tab 100 mg – **5% DV Jun-24 to 2026**	12.65	990	**Ethics Aspirin EC**
Suppos 300 mg			
CLOPIDOGREL			
Tab 75 mg – **5% DV Dec-25 to 2028**	5.07	84	**Arrow - Clopid**
DIPYRIDAMOLE – **Restricted:** For continuation only			
➥ Tab 25 mg			
➥ Cap modified-release 200 mg	55.13	60	Dipyridamole - Strides
➥ Inj 5 mg per ml, 2 ml ampoule			
EPTIFIBATIDE – **Restricted** see terms below			
⬇ Inj 2 mg per ml, 10 ml vial	180.38	1	Eptifibatide Viatris
⬇ Inj 750 mcg per ml, 100 ml vial	526.50	1	Eptifibatide Viatris

➥ **Restricted** (RS1759)

Initiation

Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
2 For use in patients with definite or strongly suspected intra-coronary thrombus on coronary angiography; or
3 For use in patients undergoing intra-cranial intervention.

LYSINE ACETYLSALICYLATE [LYSINE ASPIRIN] – **Restricted** see terms below

⬇ Inj 500 mg *e.g. Aspegic*

➥ **Restricted** (RS1689)

Initiation

Both:

1 For use when an immediate antiplatelet effect is required prior to an urgent interventional neuro-radiology or interventional cardiology procedure; and
2 Administration of oral aspirin would delay the procedure.

TICAGRELOR – **Restricted** see terms below

⬇ Tab 90 mg – **5% DV Dec-24 to 2027**	23.85	56	Ticagrelor Mylan
	20.35		**Ticagrelor Sandoz**
			Ticagrelor Sandoz S29

➥ **Restricted** (RS2142)

Initiation

Restricted to treatment of acute coronary syndromes specifically for patients who have recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome, and in whom fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initiation – thrombosis prevention neurological stenting

Re-assessment required after 12 months

Both:

1 Either:
  1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
  1.2 Patient is about to have a neurological stenting procedure performed*; and
2 Either:
  2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
  2.2 Either:
    2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
    2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent..

Continuation – thrombosis prevention neurological stenting

Re-assessment required after 12 months

Both:

1 Patient is continuing to benefit from treatment; and
2 Treatment continues to be clinically appropriate.

Initiation – Percutaneous coronary intervention with stent deployment

Limited to 12 months treatment

All of the following:

1 Patient has undergone percutaneous coronary intervention; and
2 Patient has had a stent deployed in the previous 4 weeks; and
3 Patient is clopidogrel-allergic**.

Initiation – Stent thrombosis

Patient has experienced cardiac stent thrombosis whilst on clopidogrel.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – Myocardial infarction

Limited to 1 week treatment

For short term use while in hospital following ST-elevated myocardial infarction.

Initiation – acute minor stroke or high-risk transient ischemic attack (TIA)*

All of the following:

1 Patient has been diagnosed with a minor stroke (NIHSS† score 3 or less), high-risk TIA (ABCD2 score 4 or more) or Crescendo TIA; and

2 Either:

2.1 Patient is expected to be a poor metaboliser of clopidogrel, with documented clinical rationale; or

2.2 Patient is allergic to clopidogrel**; and

3 Ticagrelor to be prescribed for a maximum of 21 days following minor stroke or TIA.

Continuation – subsequent minor stroke or high-risk transient ischemic attack

Re-assessment required after 1 month

Patient has been diagnosed with a minor stroke (NIHSS score 3 or less), high-risk transient ischemic attack (ABCD2 score 4 or more) or Crescendo TIA.

Notes: Indications marked with * are unapproved indications.

Note:** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment

Note:NIHSS† National Institutes of Health Stroke Scale.

TICLOPIDINE

Tab 250 mg

Fibrinolytic Agents

ALTEPLASE

Inj 2 mg vial

Inj 10 mg vial

Inj 50 mg vial

TENECTEPLASE

Inj 50 mg vial

UROKINASE

Inj 5,000 iu vial

Inj 10,000 iu vial

Inj 50,000 iu vial

Inj 100,000 iu vial

Inj 250,000 iu vial

Inj 500,000 iu vial

Colony-Stimulating Factors

Drugs Used to Mobilise Stem Cells

PLERIXAFOR – **Restricted** see terms below

⬇	Inj 20 mg per ml, 1.2 ml vial – **5% DV Oct-26 to 2028**	8,740.00	1	Mozobil
		869.00		**Plerixafor-AFT**

(Mozobil Inj 20 mg per ml, 1.2 ml vial to be delisted 1 October 2026)

➥ **Restricted** (RS2157)

Initiation – Stem cell transplant

Haematologist

Limited to 3 days treatment

All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Either:
- 1.1 Patient is to undergo stem cell transplantation; or
- 1.2 Patient is a donor for stem cell transplantation; and

2 Patient has not had more than one previous unsuccessful mobilisation attempt with plerixafor; and

3 Any of the following:
- 3.1 Both:
  - 3.1.1 Patient is undergoing G-CSF mobilisation; and
  - 3.1.2 Either:
    - 3.1.2.1 Has a suboptimal peripheral blood CD34 count of less than or equal to 20 x 10^6/L on day 5 after 4 days of G-CSF treatment; or
    - 3.1.2.2 Efforts to collect > 1 x 10^6 CD34 cells/kg have failed after one apheresis procedure; or
- 3.2 Both:
  - 3.2.1 Patient is undergoing chemotherapy and G-CSF mobilisation; and
  - 3.2.2 Any of the following:
    - 3.2.2.1 Both:
      - 3.2.2.1.1 Has rising white blood cell counts of > 2 x 10^9/L; and
      - 3.2.2.1.2 Has a suboptimal peripheral blood CD34 count of less than or equal to 20 x 10^6/L; or
    - 3.2.2.2 Efforts to collect > 1 x 10^6 CD34 cells/kg have failed after one apheresis procedure; or
    - 3.2.2.3 The peripheral blood CD34 cell counts are decreasing before the target has been received; or
- 3.3 A previous mobilisation attempt with G-CSF or G-CSF plus chemotherapy has failed.

Granulocyte Colony-Stimulating Factors

FILGRASTIM – **Restricted** see terms below

⬇	Inj 300 mcg in 0.5 ml prefilled syringe – **5% DV Dec-24 to 2027**	86.60	10	**Nivestim**
⬇	Inj 300 mcg in 1 ml vial	520.00	4	Neupogen
⬇	Inj 480 mcg in 0.5 ml prefilled syringe – **5% DV Dec-24 to 2027**	133.72	10	**Nivestim**

➥ **Restricted** (RS1188)
Haematologist or oncologist

PEGFILGRASTIM – **Restricted** see terms below

⬇	Inj 6 mg per 0.6 ml syringe – **5% DV Feb-26 to 2028**	69.50	1	**Ziextenzo**

➥ **Restricted** (RS1743)
Initiation
For prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*).
Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines

Fluids and Electrolytes

Intravenous Administration

CALCIUM CHLORIDE
Inj 100 mg per ml, 10 ml vial
Inj 100 mg per ml, 50 ml syringe — *e.g. Baxter*

CALCIUM GLUCONATE
Inj 10%, 10 ml ampoule — *e.g. Max Health*

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
COMPOUND ELECTROLYTES			
Inj sodium 140 mmol/l, potassium 5 mmol/l, magnesium 1.5 mmol/l, chloride 98 mmol/l, acetate 27 mmol/l, gluconate 23 mmol/l, 500 ml bag	62.82	18	Plasma-Lyte 148
Inj sodium 140 mmol/l, potassium 5 mmol/l, magnesium 1.5 mmol/l, chloride 98 mmol/l, acetate 27 mmol/l, gluconate 23 mmol/l, 1,000 ml bag	30.72	12	Plasma-Lyte 148
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]			
Inj sodium 140 mmol/l, 5 mmol/l potassium, 1.5 mmol/l magnesium, 98 mmol/l chloride, 27 mmol/l acetate and 23 mmol/l gluconate, glucose 23 mmol/l (5%), 1,000 ml bag	239.04	12	Plasma-Lyte 148 & 5% Glucose
COMPOUND SODIUM LACTATE [HARTMANN'S SOLUTION]			
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bicarbonate 29 mmol/l, chloride 111 mmol/l, 500 ml bag	27.90	18	Baxter
Inj sodium 131 mmol/l with potassium 5 mmol/l, calcium 2 mmol/l, bicarbonate 29 mmol/l, chloride 111 mmol/l, 1,000 ml bag	19.32	12	Baxter
GLUCOSE [DEXTROSE]			
Inj 5%, 1,000 ml bag	53.10	10	Fresenius Kabi
Inj 5%, 100 ml bag	97.00	50	Fresenius Kabi
Inj 5%, 250 ml bag	63.00	30	Fresenius Kabi
Inj 5%, 50 ml bag	162.00	60	Baxter Glucose 5%
Inj 5%, 500 ml bag	67.40	20	Fresenius Kabi
Inj 10%, 1,000 ml bag	162.00	12	Baxter Glucose 10%
Inj 10%, 500 ml bag	126.00	18	Baxter Glucose 10%
Inj 50%, 10 ml ampoule – **5% DV Feb-24 to 2026**	34.75	5	**Biomed**
Inj 50%, 500 ml bag	423.00	18	Baxter Glucose 50%
Inj 50%, 90 ml bottle – **5% DV Feb-24 to 2026**	17.50	1	**Biomed**
GLUCOSE WITH POTASSIUM CHLORIDE			
Inj 10% glucose with 20 mmol/l potassium chloride, 500 ml bag			
GLUCOSE WITH POTASSIUM CHLORIDE AND SODIUM CHLORIDE			
Inj 2.5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 3,000 ml bag			
Inj 10% glucose with potassium chloride 10 mmol/l and sodium chloride 15 mmol/l, 500 ml bag			
Inj 4% glucose with potassium chloride 20 mmol/l and sodium chloride 0.18%, 1,000 ml bag	240.36	12	Baxter
Inj 5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.45%, 1,000 ml bag	189.00	12	Baxter
Inj 5% glucose with potassium chloride 20 mmol/l and sodium chloride 0.9%, 1,000 ml bag	334.08	12	Baxter
GLUCOSE WITH SODIUM CHLORIDE			
Inj glucose 2.5% with sodium chloride 0.45%, 500 ml bag	318.78	18	Baxter
Inj 4% glucose and sodium chloride 0.18%, 1,000 ml bag	192.96	12	Baxter
Inj 5% glucose and sodium chloride 0.45%, 1,000 ml bag	192.84	12	Baxter
Inj 5% glucose and sodium chloride 0.9%, 1,000 ml bag	204.84	12	Baxter
POTASSIUM CHLORIDE			
Inj 75 mg (1 mmol) per ml, 10 ml ampoule			
Inj 225 mg (3 mmol) per ml, 20 ml ampoule			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
POTASSIUM CHLORIDE WITH SODIUM CHLORIDE			
Inj 10 mmol potassium chloride with 0.29% sodium chloride, 100 ml bag	563.52	48	Baxter
Inj 20 mmol potassium chloride with 0.9% sodium chloride, 1,000 ml bag	192.72	12	Baxter
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 1,000 ml bag	299.40	12	Baxter
Inj 40 mmol potassium chloride with 0.9% sodium chloride, 100 ml bag	912.96	48	Baxter
POTASSIUM DIHYDROGEN PHOSPHATE			
Inj 1 mmol per ml, 10 ml ampoule	174.57	10	Hospira
RINGER'S SOLUTION			
Inj sodium 147 mmol/l with potassium 4 mmol/l, calcium 2.2 mmol/l, chloride 156 mmol/l, 1,000 ml bag	227.52	12	Baxter
SODIUM ACETATE			
Inj 4 mmol per ml, 20 ml ampoule			
SODIUM BICARBONATE			
Inj 8.4%, 10 ml vial			
Inj 8.4%, 50 ml vial	24.70	1	Biomed
Inj 8.4%, 100 ml vial	25.31	1	Biomed
SODIUM CHLORIDE			
Inj 0.9%, 5 ml ampoule – **5% DV Feb-26 to 2028**	4.12	20	**Fresenius Kabi**
Inj 0.9%, 10 ml ampoule – **5% DV Feb-26 to 2028**	7.50	50	**Fresenius Kabi**
⬇ Inj 0.9%, 3 ml, prefilled syringe, non-sterile pack – **5% DV Jul-26 to 2028**	12.00	30	BD PosiFlush
	12.60		**Polyflush**
➥ **Restricted** (RS1297)			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⬇ Inj 0.9%, 5 ml, prefilled syringe, non-sterile pack – **5% DV Jul-26 to 2028**	12.00	30	BD PosiFlush
	12.60		**Polyflush**
➥ **Restricted** (RS1297)			
Initiation			
For use in flushing of in-situ vascular access devices only.			
⬇ Inj 0.9%, 10 ml, prefilled syringe, non-sterile pack – **5% DV Jul-26 to 2028**	11.70	30	BD PosiFlush
	12.60		**Polyflush**
➥ **Restricted** (RS1297)			
Initiation			
For use in flushing of in-situ vascular access devices only.			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Inj 0.9%, 20 ml ampoule – **5% DV Feb-26 to 2028**	5.20	20	**Fresenius Kabi**
Inj 23.4% (4 mmol/ml), 20 ml ampoule	40.15	5	Biomed
Inj 0.45%, 500 ml bag	84.42	18	Baxter
Inj 3%, 1,000 ml bag	165.84	12	Baxter
Inj 0.9%, 50 ml bag	124.20	60	Baxter
	147.75	75	Baxter-Viaflo
Inj 0.9%, 100 ml bag	88.80	48	Baxter
	105.60	60	Baxter-Viaflo
Inj 0.9%, 250 ml bag	50.40	24	Baxter
Inj 0.9%, 500 ml bag	27.54	18	Baxter
Inj 0.9%, 1,000 ml bag	18.96	12	Baxter
Inj 1.8%, 500 ml bottle			
(BD PosiFlush Inj 0.9%, 3 ml, prefilled syringe, non-sterile pack to be delisted 1 July 2026)			
(BD PosiFlush Inj 0.9%, 5 ml, prefilled syringe, non-sterile pack to be delisted 1 July 2026)			
(BD PosiFlush Inj 0.9%, 10 ml, prefilled syringe, non-sterile pack to be delisted 1 July 2026)			
SODIUM DIHYDROGEN PHOSPHATE [SODIUM ACID PHOSPHATE]			
Inj 1 mmol per ml, 20 ml ampoule	59.10	5	Biomed
WATER			
Inj 10 ml ampoule	7.60	50	Fresenius Kabi Multichem
Inj 20 ml ampoule	5.00	20	Fresenius Kabi
Inj 250 ml bag			
Inj 500 ml bag			
Inj, 1,000 ml bag	24.12	12	Baxter

Oral Administration

CALCIUM POLYSTYRENE SULPHONATE			
Powder	169.85	300 g	Calcium Resonium Roma
COMPOUND ELECTROLYTES			
Powder for oral soln – **5% DV Dec-25 to 2028**	9.50	50	**Electral**
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]			
Soln with electrolytes	6.53	1	Hydralyte - Lemonade
Soln with electrolytes (2 × 500 ml) – **5% DV Jul-26 to 2028**	8.45	1	**Pedialyte**
(Hydralyte - Lemonade Soln with electrolytes to be delisted 1 July 2026)			
PHOSPHORUS			
Tab eff 500 mg (16 mmol)			
POTASSIUM CHLORIDE			
Tab eff 548 mg (14 mmol) with chloride 285 mg (8 mmol)			
Tab long-acting 600 mg (8 mmol) – **5% DV Feb-26 to 2028**	16.15	200	**Span-K**
Oral liq 2 mmol per ml			
SODIUM BICARBONATE			
Cap 840 mg	8.52	100	Sodibic
SODIUM CHLORIDE			
Tab 600 mg			
Oral liq 2 mmol/ml			
SODIUM POLYSTYRENE SULPHONATE			
Powder	84.65	454 g	Resonium A

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Plasma Volume Expanders			
GELATINE, SUCCINYLATED			
Inj 4%, 500 ml bag	139.10	10	Gelofusine

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL

⬇ Oral liq 5 mg per ml – **5% DV Apr-24 to 2026**	86.00	100 ml	**DP-Captopril**

➥ **Restricted** (RS1263)
Initiation
Any of the following:
1 For use in children under 12 years of age; or
2 For use in tube-fed patients; or
3 For management of rebound transient hypertension following cardiac surgery.

ENALAPRIL MALEATE

Tab 5 mg – **5% DV Jul-26 to 2028**	4.25	90	Acetec
	1.40		**Ipca-Enalapril**
Tab 10 mg – **5% DV Jul-26 to 2028**	5.50	90	Acetec
	1.58		**Ipca-Enalapril**
Tab 20 mg – **5% DV Jul-26 to 2028**	6.50	90	Acetec
	2.00		**Ipca-Enalapril**

(Acetec Tab 5 mg to be delisted 1 July 2026)
(Acetec Tab 10 mg to be delisted 1 July 2026)
(Acetec Tab 20 mg to be delisted 1 July 2026)

LISINOPRIL

Tab 5 mg – **5% DV Mar-26 to 2028**	12.00	90	**Teva Lisinopril**
Tab 10 mg – **5% DV Mar-26 to 2028**	12.00	90	**Teva Lisinopril**
Tab 20 mg – **5% DV Mar-26 to 2028**	16.00	90	**Teva Lisinopril**

PERINDOPRIL

Tab 2 mg – **5% DV Dec-24 to 2027**	1.79	30	**Coversyl**
Tab 4 mg – **5% DV Dec-24 to 2027**	2.44	30	**Coversyl**
Tab 8 mg – **5% DV Dec-24 to 2027**	3.94	30	**Coversyl**

QUINAPRIL

Tab 5 mg – **5% DV Mar-25 to 2027**	10.24	90	**Arrow-Quinapril 5**
Tab 10 mg – **5% DV Mar-25 to 2027**	12.51	90	**Arrow-Quinapril 10**
Tab 20 mg – **5% DV Mar-25 to 2027**	14.38	90	**Arrow-Quinapril 20**

RAMIPRIL

Tab 1.25 mg	5.75	30	Ramipril Viatris
Tab 5 mg	5.25	28	Ramipril Mylan Generics
Tab 10 mg	5.48	28	Ramipril Mylan Generics
Cap 1.25 mg – **5% DV Feb-25 to 2027**	17.25	90	**Tryzan**
Cap 2.5 mg – **5% DV Feb-25 to 2027**	16.50	90	**Tryzan**
Cap 5 mg – **5% DV Feb-25 to 2027**	16.88	90	**Tryzan**
Cap 10 mg – **5% DV Feb-25 to 2027**	17.63	90	**Tryzan**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Angiotensin II Antagonists			
CANDESARTAN CILEXETIL			
Tab 4 mg – **5% DV Feb-25 to 2027**	2.68	90	Candesartan Viatris Candesartan Viatris Sante **Candestar**
Tab 8 mg – **5% DV Feb-25 to 2027**	2.67	90	Candesartan Viatris **Candestar**
Tab 16 mg – **5% DV Feb-25 to 2027**	4.22	90	Candesartan Viatris **Candestar**
Tab 32 mg – **5% DV Feb-25 to 2027**	5.24	90	Candesartan Viatris **Candestar**
LOSARTAN POTASSIUM			
Tab 12.5 mg – **5% DV Mar-24 to 2026**	2.00	84	**Losartan Actavis**
Tab 25 mg – **5% DV Mar-24 to 2026**	2.29	84	**Losartan Actavis**
Tab 50 mg – **5% DV Mar-24 to 2026**	2.86	84	**Losartan Actavis**
Tab 100 mg – **5% DV Mar-24 to 2026**	4.57	84	**Losartan Actavis**
Angiotensin II Antagonists with Diuretics			
CANDESARTAN CILEXETIL WITH HYDROCHLOROTHIAZIDE			
Tab 16 mg with hydrochlorothiazide 12.5 mg	4.10	30	APO-Candesartan HCTZ 16/12.5
Tab 32 mg with hydrochlorothiazide 12.5 mg	5.25	30	APO-Candesartan HCTZ 32/12.5
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE			
Tab 50 mg with hydrochlorothiazide 12.5 mg – **5% DV Jul-26 to 2028**	4.31	30	Arrow-Losartan & Hydrochlorothiazide
	7.25	90	**Losartan & Hydrochlorothiazide (Ipca)**

(Arrow-Losartan & Hydrochlorothiazide Tab 50 mg with hydrochlorothiazide 12.5 mg to be delisted 1 July 2026)

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN – **Restricted** see terms below

⬇ Tab 24.3 mg with valsartan 25.7 mg	190.00	56	Entresto 24/26
⬇ Tab 48.6 mg with valsartan 51.4 mg	190.00	56	Entresto 49/51
⬇ Tab 97.2 mg with valsartan 102.8 mg	190.00	56	Entresto 97/103

➥ **Restricted** (RS2014)

Initiation

All of the following:

1 Patient has heart failure; and

2 Any of the following:

2.1 Patient is in NYHA/WHO functional class II; or

2.2 Patient is in NYHA/WHO functional class III; or

2.3 Patient is in NYHA/WHO functional class IV; and

3 Either:

3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or

continued…

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and

4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Alpha-Adrenoceptor Blockers

DOXAZOSIN

Tab 2 mg	17.35	500	Doxazosin Clinect
Tab 4 mg	20.94	500	Doxazosin Clinect

PHENOXYBENZAMINE HYDROCHLORIDE

Cap 10 mg

Inj 50 mg per ml, 1 ml ampoule

Inj 50 mg per ml, 2 ml ampoule

PHENTOLAMINE MESYLATE

Inj 5 mg per ml, 1 ml ampoule

Inj 10 mg per ml, 1 ml ampoule

PRAZOSIN

Tab 1 mg	5.53	100	Arrotex-Prazosin S29
Tab 2 mg	7.00	100	Arrotex-Prazosin S29
Tab 5 mg	11.70	100	Arrotex-Prazosin S29
Cap 1 mg	15.40	100	Prazosin Mylan
Cap 2 mg	15.58	100	Prazosin Mylan
Cap 5 mg	23.32	100	Prazosin Mylan

TERAZOSIN – **Restricted:** For continuation only

➥ Tab 1 mg

Antiarrhythmics

ADENOSINE

Inj 3 mg per ml, 2 ml vial – **5% DV Dec-24 to 2027**	34.50	5	**Adsine**
⬇ Inj 3 mg per ml, 10 ml vial – **5% DV Dec-24 to 2027**	100.00	5	**Adenosine Baxter**

➥ **Restricted** (RS2158)

Initiation

For use in cardiac catheterisation, myocardial perfusion scans, electrophysiology and MRI.

AJMALINE – **Restricted** see terms below

⬇ Inj 5 mg per ml, 10 ml ampoule

➥ **Restricted** (RS1001)

Cardiologist

AMIODARONE HYDROCHLORIDE

Tab 100 mg – **5% DV Feb-26 to 2028**	4.95	30	**Aratac**
Tab 200 mg – **5% DV Feb-26 to 2028**	5.86	30	**Aratac**
Inj 50 mg per ml, 3 ml ampoule – **5% DV Feb-26 to 2028**	17.96	10	**Max Health**

ATROPINE SULPHATE

Inj 600 mcg per ml, 1 ml ampoule – **5% DV Feb-25 to 2027**	16.10	10	Hikma **Martindale**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
DIGOXIN			
Tab 62.5 mcg – **5% DV Feb-26 to 2028**	8.58	240	**Lanoxin PG**
Tab 250 mcg – **5% DV Feb-26 to 2028**	18.75	240	**Lanoxin**
Oral liq 50 mcg per ml			
Inj 250 mcg per ml, 2 ml vial			
DISOPYRAMIDE PHOSPHATE			
Cap 100 mg			
FLECAINIDE ACETATE			
Tab 50 mg – **5% DV Dec-23 to 2026**	19.95	60	**Flecainide BNM**
Cap long-acting 100 mg – **5% DV Aug-23 to 2026**	35.78	90	**Flecainide Controlled Release Teva**
Cap long-acting 200 mg – **5% DV Aug-23 to 2026**	54.28	90	**Flecainide Controlled Release Teva**
Inj 10 mg per ml, 15 ml ampoule	102.79	5	Almarytm
	108.16		Tambocor
			Tambocor German

IVABRADINE – **Restricted** see terms below

⬇ Tab 5 mg

➥ **Restricted** (RS1566)

Initiation

Both:

1 Patient is indicated for computed tomography coronary angiography; and

2 Either:

2.1 Patient has a heart rate of greater than 70 beats per minute while taking a maximally tolerated dose of beta blocker; or

2.2 Patient is unable to tolerate beta blockers.

MEXILETINE HYDROCHLORIDE			
Cap 150 mg	162.00	100	Teva
Cap 250 mg	202.00	100	Teva
PROPAFENONE HYDROCHLORIDE			
Tab 150 mg			

Antihypotensives

MIDODRINE – **Restricted** see terms below			
⬇ Tab 2.5 mg – **5% DV Feb-25 to 2027**	36.68	100	**Midodrine Medsurge**
⬇ Tab 5 mg – **5% DV Feb-25 to 2027**	58.88	100	**Midodrine Medsurge**

➥ **Restricted** (RS1427)

Initiation

Patient has disabling orthostatic hypotension not due to drugs.

Beta-Adrenoceptor Blockers

ATENOLOL			
Tab 50 mg – **5% DV Feb-25 to 2027**	11.00	500	**Viatris**
Tab 100 mg – **5% DV Feb-25 to 2027**	18.50	500	**Atenolol Viatris**
Oral liq 5 mg per ml	49.85	300 ml	Atenolol-AFT
BISOPROLOL FUMARATE			
Tab 2.5 mg – **5% DV Apr-24 to 2026**	1.36	90	**Ipca-Bisoprolol**
Tab 5 mg – **5% DV Apr-24 to 2026**	1.91	90	**Ipca-Bisoprolol**
Tab 10 mg – **5% DV Apr-24 to 2026**	2.71	90	**Ipca-Bisoprolol**

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
CARVEDILOL			
Tab 6.25 mg – **5% DV Aug-26 to 2029**	1.97	60	**Carvedilol Sandoz**
Tab 12.5 mg – **5% DV Aug-26 to 2029**	2.03	60	**Carvedilol Sandoz**
Tab 25 mg – **5% DV Aug-26 to 2029**	2.46	60	**Carvedilol Sandoz**
CELIPROLOL – **Restricted:** For continuation only			
➥ Tab 200 mg			
ESMOLOL HYDROCHLORIDE			
Inj 10 mg per ml, 10 ml vial			
LABETALOL			
Tab 50 mg			
Tab 100 mg	49.54	100	Biocon
	14.50		Trandate
Tab 200 mg	42.07	100	Presolol
	27.00		Trandate
Inj 5 mg per ml, 20 ml ampoule			
METOPROLOL SUCCINATE			
Tab long-acting 23.75 mg – **5% DV Apr-24 to 2026**	4.20	90	**Myloc CR**
Tab long-acting 47.5 mg – **5% DV Apr-24 to 2026**	3.65	90	**Myloc CR**
Tab long-acting 95 mg – **5% DV Apr-24 to 2026**	5.24	90	**Myloc CR**
Tab long-acting 190 mg – **5% DV Apr-24 to 2026**	9.76	90	**Myloc CR**
METOPROLOL TARTRATE			
Tab 50 mg – **1% DV Mar-22 to 2027**	5.66	100	**IPCA-Metoprolol**
Tab 100 mg – **1% DV Mar-22 to 2027**	7.55	60	**IPCA-Metoprolol**
Tab long-acting 200 mg	23.40	28	Slow-Lopresor
Inj 1 mg per ml, 5 ml vial	26.50	5	Metoprolol IV Mylan Metoprolol IV Viatris
NADOLOL			
Tab 40 mg – **1% DV Mar-22 to 2027**	19.19	100	**Nadolol BNM**
Tab 80 mg – **1% DV Mar-22 to 2027**	30.39	100	**Nadolol BNM**
PROPRANOLOL			
Tab 10 mg – **1% DV Mar-22 to 2027**	7.04	100	**Drofate**
Tab 40 mg – **1% DV Mar-22 to 2027**	8.75	100	**IPCA-Propranolol**
Cap long-acting 160 mg	18.17	100	Cardinol LA
Oral liq 4 mg per ml			*e.g. Hikma-Propranolol*
Inj 1 mg per ml, 1 ml ampoule			
SOTALOL			
Tab 80 mg – **5% DV Feb-26 to 2028**	40.00	500	**Mylan**
Tab 160 mg – **5% DV Feb-26 to 2028**	20.00	100	**Mylan**

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMLODIPINE			
Tab 2.5 mg – **5% DV Feb-24 to 2026**	1.45	90	**Vasorex**
Tab 5 mg – **5% DV Feb-24 to 2026**	1.21	90	**Vasorex**
Tab 10 mg – **5% DV Feb-24 to 2026**	1.31	90	**Vasorex**
FELODIPINE			
Tab long-acting 2.5 mg – **5% DV Feb-25 to 2027**	2.18	30	**Plendil ER**
Tab long-acting 5 mg – **5% DV Feb-25 to 2027**	6.57	90	**Felo 5 ER**
Tab long-acting 10 mg – **5% DV Feb-25 to 2027**	6.95	90	**Felo 10 ER**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
ISRADIPINE			
Tab 2.5 mg			
Cap 2.5 mg			
NICARDIPINE HYDROCHLORIDE – **Restricted** see terms below			
⬇ Inj 2.5 mg per ml, 10 ml vial			
➥ **Restricted** (RS1699)			
Initiation			
Anaesthetist, intensivist, cardiologist or paediatric cardiologist			
Any of the following:			
1 Patient has hypertension requiring urgent treatment with an intravenous agent; or			
2 Patient has excessive ventricular afterload; or			
3 Patient is awaiting or undergoing cardiac surgery using cardiopulmonary bypass.			
NIFEDIPINE			
Tab long-acting 10 mg	19.42	56	Tensipine MR10
Tab long-acting 20 mg	17.72	100	Nyefax Retard
	9.92	56	Valni Retard
Tab long-acting 30 mg	34.10	100	Mylan (24 hr release)
	4.78	14	Mylan Italy (24 hr release)
Tab long-acting 60 mg	52.81	100	Mylan (24 hr release)
Cap 5 mg			
NIMODIPINE			
Tab 30 mg – **5% DV Feb-26 to 2028**	350.00	100	**Nimotop**
Inj 0.2 mg per ml, 50 ml vial – **5% DV Feb-26 to 2028**	337.50	5	**Nimotop**

Other Calcium Channel Blockers

DILTIAZEM HYDROCHLORIDE			
Tab 30 mg			
Cap long-acting 120 mg – **5% DV Dec-25 to 2028**	65.35	500	**Diltiazem CD Clinect**
Cap long-acting 180 mg – **1% DV Mar-22 to 2027**	7.00	30	**Cardizem CD**
Cap long-acting 240 mg – **1% DV Mar-22 to 2027**	9.30	30	**Cardizem CD**
Inj 5 mg per ml, 5 ml vial			
PERHEXILINE MALEATE			
Tab 100 mg	62.90	100	Pexsig
VERAPAMIL HYDROCHLORIDE			
Tab 40 mg	7.01	100	Isoptin
Tab 80 mg	11.74	100	Isoptin
Tab long-acting 120 mg	36.02	100	Isoptin SR
Tab long-acting 240 mg	15.12	30	Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule	25.00	5	Isoptin

Centrally-Acting Agents

CLONIDINE			
Patch 2.5 mg, 100 mcg per day – **5% DV Feb-24 to 2026**	11.70	4	**Mylan**
Patch 5 mg, 200 mcg per day – **5% DV Feb-24 to 2026**	12.80	4	**Mylan**
Patch 7.5 mg, 300 mcg per day – **5% DV Feb-24 to 2026**	17.90	4	**Mylan**

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
CLONIDINE HYDROCHLORIDE			
Tab 25 mcg – **5% DV Feb-26 to 2028**	29.74	112	**Clonidine Teva**
Tab 150 mcg – **5% DV Feb-25 to 2027**	40.41	100	**Catapres**
Inj 150 mcg per ml, 1 ml ampoule – **5% DV Jan-25 to 2027**	14.10	5	**Catapres**
METHYLDOPA			
Tab 250 mg	15.10	100	Methyldopa Viatris

Diuretics

Loop Diuretics

BUMETANIDE			
Tab 1 mg	16.36	100	Burinex
Inj 500 mcg per ml, 4 ml vial			
FUROSEMIDE [FRUSEMIDE]			
Tab 40 mg – **5% DV Feb-25 to 2027**	12.80	1,000	**IPCA-Frusemide**
Tab 500 mg	25.00	50	Urex Forte
Oral liq 10 mg per ml	11.20	30 ml	Lasix
Inj 10 mg per ml, 2 ml ampoule – **5% DV Aug-26 to 2028**	3.97	10	**Furosemide-AFT**
	2.40	5	Furosemide-Baxter
Inj 10 mg per ml, 25 ml ampoule	60.65	6	Lasix

(Furosemide-Baxter Inj 10 mg per ml, 2 ml ampoule to be delisted 1 August 2026)

Osmotic Diuretics

MANNITOL			
Inj 10%, 1,000 ml bag	882.84	12	Baxter
Inj 20%, 500 ml bag	1,296.00	18	Baxter

Potassium Sparing Combination Diuretics

AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE
Tab 5 mg with furosemide 40 mg

AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE
Tab 5 mg with hydrochlorothiazide 50 mg

Potassium Sparing Diuretics

AMILORIDE HYDROCHLORIDE			
Tab 5 mg			
Oral liq 1 mg per ml	35.40	25 ml	Biomed
EPLERENONE – **Restricted** see terms below			
⬇ Tab 25 mg – **5% DV Dec-24 to 2027**	15.84	30	**Inspra**
⬇ Tab 50 mg – **5% DV Dec-24 to 2027**	25.00	30	**Inspra**

➥ **Restricted** (RS1640)

Initiation

Both:

1 Patient has heart failure with ejection fraction less than 40%; and

2 Either:

2.1 Patient is intolerant to optimal dosing of spironolactone; or

2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
SPIRONOLACTONE			
Tab 25 mg – **5% DV Mar-26 to 2028**	4.20	100	**Spiractin**
Tab 100 mg – **5% DV Mar-26 to 2028**	11.40	100	**Spiractin**
Oral liq 5 mg per ml	35.70	25 ml	Biomed

Thiazide and Related Diuretics

BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]			
Tab 2.5 mg – **5% DV Mar-24 to 2026**	51.50	500	**Arrow-Bendrofluazide**
Tab 5 mg – **5% DV Mar-24 to 2026**	61.00	500	**Arrow-Bendrofluazide**
CHLOROTHIAZIDE			
Oral liq 50 mg per ml	30.67	25 ml	Biomed
CHLORTALIDONE [CHLORTHALIDONE]			
Tab 25 mg – **5% DV Feb-26 to 2028**	6.95	50	**Hygroton**
INDAPAMIDE			
Tab 2.5 mg – **5% DV Feb-24 to 2026**	16.00	90	**Dapa-Tabs**
METOLAZONE			
Tab 5 mg			

Vasopressin receptor antagonists

TOLVAPTAN – **Restricted** see terms below

⬇ Tab 15 mg	873.50	28	Jinarc
⬇ Tab 30 mg	873.50	28	Jinarc
⬇ Tab 45 mg + 15 mg	1,747.00	56	Jinarc
⬇ Tab 60 mg + 30 mg	1,747.00	56	Jinarc
⬇ Tab 90 mg + 30 mg	1,747.00	56	Jinarc

➥ **Restricted** (RS1930)

Initiation – autosomal dominant polycystic kidney disease

Renal physician or any relevant practitioner on the recommendation of a renal physician

Re-assessment required after 12 months

All of the following:

1 Patient has a confirmed diagnosis of autosomal dominant polycystic kidney disease; and
2 Patient has an estimated glomerular filtration rate (eGFR) of greater than or equal to 25 ml/min/1.73 m^2 at treatment initiation; and
3 Either:
  3.1 Patient's disease is rapidly progressing, with a decline in eGFR of greater than or equal to 5 mL/min/1.73 m^2 within one-year; or
  3.2 Patient's disease is rapidly progressing, with an average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m^2 per year over a five-year period.

Continuation – autosomal dominant polycystic kidney disease

Renal physician or any relevant practitioner on the recommendation of a renal physician

Re-assessment required after 12 months

Both:

1 Patient has not developed end-stage renal disease, defined as an eGFR of less than 15 mL/min/1.73 m^2; and
2 Patient has not undergone a kidney transplant.

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Lipid-Modifying Agents

Fibrates

BEZAFIBRATE

Tab 200 mg – **5% DV Mar-25 to 2027**	22.65	90	**Bezalip**
Tab long-acting 400 mg – **5% DV Mar-25 to 2027**	21.54	30	**Bezalip Retard**

HMG CoA Reductase Inhibitors (Statins)

ATORVASTATIN

Tab 10 mg – **5% DV Dec-24 to 2027**	0.31	30	**Lorstat**
	5.16	500	**Lorstat**
Tab 20 mg – **5% DV Dec-24 to 2027**	8.12	500	**Lorstat**
Tab 40 mg – **5% DV Dec-24 to 2027**	13.79	500	**Lorstat**
Tab 80 mg – **5% DV Dec-24 to 2027**	1.52	30	**Lorstat**
	25.39	500	**Lorstat**

PRAVASTATIN

Tab 10 mg			
Tab 20 mg – **5% DV May-24 to 2026**	7.16	100	**Clinect**
Tab 40 mg – **5% DV May-24 to 2026**	12.25	100	**Clinect**

ROSUVASTATIN – **Restricted** see terms below

⬇ Tab 5 mg – **5% DV Oct-24 to 2026**	4.21	30	Rosuvastatin - Sandoz
	1.29		**Rosuvastatin Viatris**
⬇ Tab 10 mg – **5% DV Oct-24 to 2026**	4.21	30	Rosuvastatin - Sandoz
	1.69		**Rosuvastatin Viatris**
⬇ Tab 20 mg – **5% DV Apr-24 to 2026**	4.21	30	Rosuvastatin - Sandoz
	2.71		**Rosuvastatin Viatris**
⬇ Tab 40 mg – **5% DV Apr-24 to 2026**	4.55	30	Rosuvastatin - Sandoz
			Rosuvastatin Viatris

➥ **Restricted** (RS1868)

Initiation – cardiovascular disease risk

Either:

1 Both:
- 1.1 Patient is considered to be at risk of cardiovascular disease; and
- 1.2 Patient is Māori or any Pacific ethnicity; or

2 Both:
- 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
- 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initiation – familial hypercholesterolemia

Both:

1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initiation – established cardiovascular disease

Both:

1 Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.1 Patient has proven coronary artery disease (CAD); or
1.2 Patient has proven peripheral artery disease (PAD); or
1.3 Patient has experienced an ischaemic stroke; and

2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initiation – recurrent major cardiovascular events

Both:

1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIMVASTATIN

Tab 10 mg – **5% DV Mar-24 to 2026**	1.68	90	**Simvastatin Mylan** Simvastatin Viatris
Tab 20 mg – **5% DV Mar-24 to 2026**	2.54	90	**Simvastatin Viatris**
Tab 40 mg – **5% DV Jun-24 to 2026**	4.11	90	**Simvastatin Viatris**
Tab 80 mg – **5% DV Jun-24 to 2026**	8.81	90	**Simvastatin Viatris**

Resins

CHOLESTYRAMINE

Powder for oral liq 4 g

COLESTIPOL HYDROCHLORIDE

Grans for oral liq 5 g

COLESTYRAMINE

Powder for oral suspension 4 g sachet	61.50	50	Colestyramine - Mylan

Selective Cholesterol Absorption Inhibitors

EZETIMIBE

Tab 10 mg – **5% DV Dec-23 to 2026**	1.76	30	**Ezetimibe Sandoz**

EZETIMIBE WITH SIMVASTATIN

Tab 10 mg with simvastatin 10 mg	11.86	30	Zimybe
Tab 10 mg with simvastatin 20 mg	12.55	30	Zimybe
Tab 10 mg with simvastatin 40 mg	11.55	30	Vytorin Zimybe
Tab 10 mg with simvastatin 80 mg	14.27	30	Zimybe

Other Lipid-Modifying Agents

ACIPIMOX

Cap 250 mg

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Nitrates

GLYCERYL TRINITRATE

Inj 1 mg per ml, 5 ml ampoule			
Inj 1 mg per ml, 10 ml ampoule			
Inj 1 mg per ml, 50 ml vial			
Inj 5 mg per ml, 10 ml ampoule	118.00	5	Hospira
Oral pump spray, 400 mcg per dose	7.48	250 dose	Nitrolingual Pump Spray
Patch 25 mg, 5 mg per day	15.73	30	Nitroderm TTS 5
Patch 50 mg, 10 mg per day	18.62	30	Nitroderm TTS 10

ISOSORBIDE MONONITRATE

Tab 20 mg – **5% DV Feb-24 to 2026**	22.49	100	**Ismo 20**
Tab long-acting 40 mg – **5% DV Feb-24 to 2026**	9.80	30	**Ismo 40 Retard**
Tab long-acting 60 mg – **5% DV Feb-24 to 2026**	13.50	90	**Duride**

Other Cardiac Agents

LEVOSIMENDAN – **Restricted** see terms below

⬇ Inj 2.5 mg per ml, 5 ml vial – **5% DV Nov-24 to 2027**	509.60	1	**Simdax**
⬇ Inj 2.5 mg per ml, 10 ml vial			

➥ **Restricted** (RS1007)

Initiation – Heart transplant

Either:

1 For use as a bridge to heart transplant, in patients who have been accepted for transplant; or

2 For the treatment of heart failure following heart transplant.

Initiation – Heart failure

Cardiologist or intensivist

For the treatment of severe acute decompensated heart failure that is non-responsive to dobutamine.

Sympathomimetics

ADRENALINE

Inj 1 in 1,000, 1 ml ampoule	4.98	5	Aspen Adrenaline
	17.78		DBL Adrenaline
Inj 1 in 1,000, 30 ml vial			
Inj 1 in 10,000, 10 ml ampoule	49.00	10	Aspen Adrenaline
	36.18	5	DBL Adrenaline
Inj 1 in 10,000, 10 ml syringe			

DOBUTAMINE

Inj 12.5 mg per ml, 20 ml ampoule – **5% DV Dec-24 to 2027**	61.13	5	**Dobutamine-hameln**

DOPAMINE HYDROCHLORIDE

Inj 40 mg per ml, 5 ml ampoule – **5% DV Feb-25 to 2027**	46.38	10	Dopamine Basi
			Max Health Ltd

EPHEDRINE

Inj 3 mg per ml, 10 ml syringe – **5% DV Aug-25 to 2026**	142.00	10	Ephedrine Aguettant
			Ephedrine Juno
Inj 30 mg per ml, 1 ml ampoule – **5% DV Feb-24 to 2026**	34.31	10	**Max Health**

ISOPRENALINE [ISOPROTERENOL]

Inj 200 mcg per ml, 1 ml ampoule			
Inj 200 mcg per ml, 5 ml ampoule			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
METARAMINOL			
Inj 0.5 mg per ml, 10 ml syringe			
Inj 0.5 mg per ml, 20 ml syringe			
Inj 0.5 mg per ml, 5 ml syringe			
Inj 1 mg per ml, 1 ml ampoule			
Inj 1 mg per ml, 10 ml syringe			
Inj 10 mg per ml, 1 ml ampoule – **5% DV Feb-24 to 2026**	53.00	10	**Torbay**
NORADRENALINE			
Inj 0.06 mg per ml, 100 ml bag			
Inj 0.06 mg per ml, 50 ml syringe			
Inj 0.1 mg per ml, 100 ml bag			
Inj 0.1 mg per ml, 50 ml syringe			
Inj 0.12 mg per ml, 100 ml bag			
Inj 0.12 mg per ml, 50 ml syringe			
Inj 0.16 mg per ml, 50 ml syringe			
Inj 1 mg per ml, 100 ml bag			
Inj 1 mg per ml, 4 ml ampoule – **5% DV Apr-26 to 2028**	32.78	10	**Noradrenaline Medsurge**
PHENYLEPHRINE HYDROCHLORIDE			
Inj 10 mg per ml, 1 ml ampoule	310.42	25	Neosynephrine HCL

Vasodilators

ALPROSTADIL – **Restricted** see terms below

⬇ Inj 10 mcg vial

⬇ Inj 20 mcg vial

➥ **Restricted** (RS1992)

Initiation

Both:

1 Patient has erectile dysfunction; and

2 Patient is to receive a penile Doppler ultrasonography.

ALPROSTADIL HYDROCHLORIDE			
Inj 500 mcg per ml, 1 ml ampoule	2,030.33	5	Prostin VR
DIAZOXIDE			
Inj 15 mg per ml, 20 ml ampoule			
HYDRALAZINE HYDROCHLORIDE			

⬇ Tab 25 mg

➥ **Restricted** (RS1008)

Initiation

Either:

1 For the treatment of refractory hypertension; or

2 For the treatment of heart failure, in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

Inj 20 mg ampoule	25.90	5	Apresoline Hydrapres
Inj 20 mg per ml, 1 ml vial	283.00	25	Eugia
MILRINONE			
Inj 1 mg per ml, 10 ml ampoule – **5% DV Dec-24 to 2027**	68.00	10	**Milrinone-Baxter**
MINOXIDIL			
Tab 10 mg	78.40	100	Loniten

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
NICORANDIL			
Tab 10 mg – **5% DV Feb-26 to 2028**	27.81	60	**Max Health**
Tab 20 mg – **5% DV Feb-26 to 2028**	35.12	60	**Max Health**
PAPAVERINE HYDROCHLORIDE			
Inj 30 mg per ml, 1 ml vial			
Inj 12 mg per ml, 10 ml ampoule	257.12	5	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg			
SODIUM NITROPRUSSIDE			
Inj 50 mg vial			

Endothelin Receptor Antagonists

AMBRISENTAN – **Restricted** see terms below

⬇ Tab 5 mg – **5% DV Dec-23 to 2026**	200.00	30	**Ambrisentan Viatris**
⬇ Tab 10 mg – **5% DV Dec-23 to 2026**	200.00	30	**Ambrisentan Viatris**

➥ **Restricted** (RS2159)

Initiation – PAH monotherapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH); and
2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
4 Any of the following:
  4.1 All of the following:
    4.1.1 PAH has been confirmed by right heart catheterisation; and
    4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
    4.1.5 Any of the following:
      4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
5 Both:
  5.1 Ambrisentan is to be used as PAH monotherapy; and
  5.2 Any of the following:
    5.2.1 Patient has experienced intolerable side effects with both sildenafil and bosentan; or
    5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH); and
2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
4 Any of the following:
  4.1 All of the following:
    4.1.1 PAH has been confirmed by right heart catheterisation; and
    4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
    4.1.5 Any of the following:
      4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
  4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
5 Both:
  5.1 Ambrisentan is to be used as PAH dual therapy; and
  5.2 Any of the following:
    5.2.1 Patient has tried bosentan (either as PAH monotherapy, or PAH dual therapy with sildenafil) for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; or
    5.2.2 Patient has experienced intolerable side effects on bosentan; or
    5.2.3 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
    5.2.4 Patient is presenting in NYHA/WHO functional class III or IV, and would benefit from initial dual therapy in the opinion of the treating clinician and has an absolute or relative contraindication to bosentan (eg. due to current liver disease or use of a combined oral contraceptive).

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH); and
2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
4 Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

4.1 All of the following:
4.1.1 PAH has been confirmed by right heart catheterisation; and
4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
4.1.5 Any of the following:
4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including chronic neonatal lung disease; or
4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:
5.1 Ambrisentan is to be used as PAH triple therapy; and
5.2 Any of the following:
5.2.1 Patient is on the lung transplant list; or
5.2.2 Both:
5.2.2.1 Patient is presenting in NYHA/WHO functional class IV; and
5.2.2.2 Patient has an absolute or relative contraindication to bosentan (e.g. due to current use of a combined oral contraceptive or liver disease); or
5.2.3 Both:
5.2.3.1 Patient has tried PAH dual therapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; and
5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation
Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist
Re-assessment required after 2 years
The patient is continuing to derive benefit from ambrisentan treatment according to a validated PAH risk stratification tool**.
Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

BOSENTAN – **Restricted** see terms below

⬇	Tab 62.5 mg – **5% DV Jan-25 to 2027**	100.00	60	**Bosentan Dr Reddy's**
⬇	Tab 125 mg – **5% DV Jan-25 to 2027**	100.00	60	**Bosentan Dr Reddy's**

➥ **Restricted** (RS2160)
Initiation – PAH monotherapy
Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist
Limited to 6 months treatment
All of the following:

1 Patient has pulmonary arterial hypertension (PAH)*; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
4 Any of the following:
  4.1 All of the following:
    4.1.1 PAH has been confirmed by right heart catheterisation; and
    4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
    4.1.5 Any of the following:
      4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and
5 Both:
  5.1 Bosentan is to be used as PAH monotherapy; and
  5.2 Any of the following:
    5.2.1 Patient has experienced intolerable side effects on sildenafil; or
    5.2.2 Patient has an absolute contraindication to sildenafil; or
    5.2.3 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease.

Initiation – PAH dual therapy
Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist
Limited to 6 months treatment
All of the following:
1 Patient has pulmonary arterial hypertension (PAH)*; and
2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
4 Any of the following:
  4.1 All of the following:
    4.1.1 PAH has been confirmed by right heart catheterisation; and
    4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
    4.1.5 Any of the following:
      4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

developmental lung disorders including severe chronic neonatal lung disease; or

4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:

5.1 Bosentan is to be used as part of PAH dual therapy; and

5.2 Either:

5.2.1 Patient has tried a PAH monotherapy (sildenafil) for at least three months and has experienced an inadequate therapeutic response to treatment according to a validated risk stratification tool**; or

5.2.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would likely benefit from initial dual therapy.

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH)*; and

2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and

3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and

4 Any of the following:

4.1 All of the following:

4.1.1 PAH has been confirmed by right heart catheterisation; and

4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

4.1.4 Pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and

4.1.5 Any of the following:

4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or

4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or

4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or

4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or

4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:

5.1 Bosentan is to be used as part of PAH triple therapy; and

5.2 Any of the following:

5.2.1 Patient is on the lung transplant list; or

5.2.2 Patient is presenting in NYHA/WHO functional class IV; or

5.2.3 Both:

5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and

5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist,

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…
cardiologist or rheumatologist
Re-assessment required after 2 years
Patient is continuing to derive benefit from bosentan treatment according to a validated PAH risk stratification tool**.
Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – **Restricted** see terms below

⬇	Tab 25 mg – **5% DV Dec-24 to 2027**	0.72	4	**Vedafil**
⬇	Tab 50 mg – **5% DV Dec-24 to 2027**	1.45	4	**Vedafil**
⬇	Tab 100 mg – **5% DV Dec-24 to 2027**	11.22	12	**Vedafil**
⬇	Inj 0.8 mg per ml, 12.5 ml vial			

➥ **Restricted** (RS2161)

Initiation – tablets Raynaud's Phenomenon

All of the following:

1 Patient has Raynaud's phenomenon; and
2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
3 Patient is following lifestyle management (proper body insulation, avoidance of cold exposure, smoking cessation support, avoidance of sympathomimetic drugs); and
4 Patient has persisting severe symptoms despite treatment with calcium channel blockers and nitrates (unless contraindicated or not tolerated).

Initiation – tablets Pulmonary arterial hypertension

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist
All of the following:

1 Patient has pulmonary arterial hypertension (PAH)*; and
2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
4 Any of the following:
  4.1 All of the following:
    4.1.1 PAH is confirmed by right heart catheterisation; and
    4.1.2 A mean pulmonary artery pressure (PAPm) of greater than 20 mmHg; and
    4.1.3 A pulmonary capillary wedge pressure (PCWP) that is less than or equal to 15 mmHg; and
    4.1.4 Pulmonary vascular resistance (PVR) of at least 2 Wood Units or at least 160 International Units (dyn s cm^{-5}); and
    4.1.5 Any of the following:
      4.1.5.1 PAH is non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – tablets other conditions

Any of the following:

1 For use in weaning patients from inhaled nitric oxide; or
2 For perioperative use in cardiac surgery patients; or
3 For use in intensive care as an alternative to nitric oxide; or
4 For use in the treatment of erectile dysfunction secondary to spinal cord injury in patients being treated in a spinal unit.

Initiation – injection

Both:

1 For use in the treatment of pulmonary hypertension in infants or children being treated in paediatric intensive care units and neonatal intensive care units when the enteral route is not accessible; and
2 Any of the following:
  2.1 For perioperative use following cardiac surgery; or
  2.2 For use in persistent pulmonary hypertension of the newborn (PPHN); or
  2.3 For use in congenital diaphragmatic hernia.

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

Prostacyclin Analogues

EPOPROSTENOL – **Restricted** see terms below

⬇	Inj 500 mcg vial	36.61	1	Veletri
⬇	Inj 1.5 mg vial	73.21	1	Veletri

➥ **Restricted** (RS2162)

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH); and
2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and
4 Any of the following:
  4.1 All of the following:
    4.1.1 PAH has been confirmed by right heart catheterisation; and
    4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
    4.1.5 Any of the following:
      4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 All of the following:
- 5.1 Epoprostenol is to be used as part of PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and
- 5.2 Patient is presenting in NYHA/WHO functional class IV; and
- 5.3 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool.

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH); and

2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and

3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class III or IV; and

4 Any of the following:
- 4.1 All of the following:
  - 4.1.1 PAH has been confirmed by right heart catheterisation; and
  - 4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
  - 4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
  - 4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
  - 4.1.5 Any of the following:
    - 4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
    - 4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
    - 4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
- 4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
- 4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures; and

5 Both:
- 5.1 Epoprostenol is to be used as PAH triple therapy; and
- 5.2 Any of the following:
  - 5.2.1 Patient is on the lung transplant list; or
  - 5.2.2 Patient is presenting in NYHA/WHO functional class IV; or
  - 5.2.3 Both:
    - 5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool; and
    - 5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

Patient is continuing to derive benefit from epoprostenol treatment according to a validated PAH risk stratification tool.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

ILOPROST

Inj 50 mcg in 0.5 ml ampoule	380.00	5	Ilomedin
⬇ Nebuliser soln 10 mcg per ml, 2 ml – **5% DV Dec-25 to 2028**	166.53	30	**Vebulis**

➥ **Restricted** (RS2163)

Initiation – PAH monotherapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH); and
2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and
4 Any of the following:
  4.1 All of the following:
    4.1.1 PAH has been confirmed by right heart catheterisation; and
    4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and
    4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and
    4.1.5 Any of the following:
      4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or
      4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or
      4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or
  4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or
  4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and
5 Both:
  5.1 Iloprost is to be used as PAH monotherapy; and
  5.2 Either:
    5.2.1 Patient has experienced intolerable side effects on sildenafil and both the funded endothelin receptor antagonists (i.e. both bosentan and ambrisentan); or
    5.2.2 Patient has an absolute contraindication to sildenafil and an absolute or relative contraindication to endothelin receptor antagonists.

Initiation – PAH dual therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH); and
2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and
3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

4 Any of the following:

4.1 All of the following:

4.1.1 PAH has been confirmed by right heart catheterisation; and

4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and

4.1.5 Any of the following:

4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or

4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or

4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or

4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or

4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and

5 All of the following:

5.1 Iloprost is to be used as PAH dual therapy with either sildenafil or an endothelin receptor antagonist; and

5.2 Either:

5.2.1 Patient has an absolute contraindication to or has experienced intolerable side effects on sildenafil; or

5.2.2 Patient has an absolute or relative contraindication to or experienced intolerable side effects with a funded endothelin receptor antagonist; and

5.3 Either:

5.3.1 Patient has tried a PAH monotherapy for at least three months and remains in an unacceptable risk category according to a validated risk stratification tool**; or

5.3.2 Patient is presenting in NYHA/WHO functional class III or IV, and in the opinion of the treating clinician would benefit from initial dual therapy.

Initiation – PAH triple therapy

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Limited to 6 months treatment

All of the following:

1 Patient has pulmonary arterial hypertension (PAH); and

2 PAH is in Group 1, 4 or 5 of the WHO (Venice 2003) clinical classifications; and

3 PAH is in New York Heart Association/World Health Organization (NYHA/WHO) functional class II, III or IV; and

4 Any of the following:

4.1 All of the following:

4.1.1 PAH has been confirmed by right heart catheterisation; and

4.1.2 A mean pulmonary artery pressure (PAPm) greater than 20 mmHg (unless peri Fontan repair); and

4.1.3 A pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

4.1.4 A pulmonary vascular resistance greater than 2 Wood Units or greater than 160 International Units (dyn s cm^{-5}); and

4.1.5 Any of the following:

4.1.5.1 PAH has been demonstrated to be non-responsive in vasoreactivity assessment using iloprost or nitric oxide, as defined in the 2022 ECS/ERS Guidelines for PAH; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

4.1.5.2 Patient has not experienced an acceptable response to calcium antagonist treatment, according to a validated risk stratification tool**; or

4.1.5.3 Patient has PAH other than idiopathic / heritable or drug-associated type; or

4.2 Patient is a child with PAH secondary to congenital heart disease or PAH due to idiopathic, congenital or developmental lung disorders including severe chronic neonatal lung disease; or

4.3 Patient has palliated single ventricle congenital heart disease and elevated pulmonary pressures or a major complication of the Fontan circulation requiring the minimising of pulmonary/venous filling pressures ; and

5 Both:

5.1 Iloprost is to be used as PAH triple therapy; and

5.2 Any of the following:

5.2.1 Patient is on the lung transplant list; or

5.2.2 Patient is presenting in NYHA/WHO functional class IV; or

5.2.3 Both:

5.2.3.1 Patient has tried PAH dual therapy for at least three months and has not experienced an acceptable response to treatment according to a validated risk stratification tool**; and

5.2.3.2 Patient does not have major life-threatening comorbidities and triple therapy is not being used in a palliative scenario.

Continuation

Respiratory specialist, cardiologist, rheumatologist or any relevant practitioner on the recommendation of a respiratory specialist, cardiologist or rheumatologist

Re-assessment required after 2 years

Patient is continuing to derive benefit from iloprost treatment according to a validated PAH risk stratification tool.

Note: ** the requirement to use a validated risk stratification tool to determine insufficient response applies to adults. Determining insufficient response in children does not require use of a validated PAH risk stratification tool, where currently no such validated tools exist for PAH risk stratification in children.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Anti-Infective Preparations

Antibacterials

HYDROGEN PEROXIDE
- Crm 1% – **5% DV Jan-26 to 2028** — 4.89 — 15 g — **Crystaderm**
- Soln 3% (10 vol)

MAFENIDE ACETATE – **Restricted** see terms below
- ⬇ Crm 8.5%
- ⬇ Powder 5%

➥ **Restricted** (RS1299)

Initiation

For the treatment of burns patients.

MUPIROCIN
- Oint 2%

SODIUM FUSIDATE [FUSIDIC ACID]
- Crm 2% – **5% DV Feb-25 to 2027** — 1.69 — 5 g — **Foban**
- Oint 2% – **5% DV Feb-25 to 2027** — 1.69 — 5 g — **Foban**

SULFADIAZINE SILVER
- Crm 1% — 10.80 — 50 g — Flamazine

Antifungals

AMOROLFINE
- Nail soln 5% – **5% DV Feb-24 to 2026** — 21.87 — 5 ml — **MycoNail**

CICLOPIROX OLAMINE
- Nail soln 8%
- ➥ Soln 1% – **Restricted:** For continuation only

CLOTRIMAZOLE
- Crm 1% – **5% DV Jul-26 to 2028** — 1.15 — 20 g — **Clomazol**
- ➥ Soln 1% – **Restricted:** For continuation only

ECONAZOLE NITRATE
- Crm 1% – **5% DV Jun-25 to 2027** — 8.04 — 20 g — **Pevaryl**
- Foaming soln 1%

KETOCONAZOLE
- Shampoo 2% – **5% DV May-24 to 2026** — 4.09 — 100 ml — **Sebizole**

METRONIDAZOLE
- Gel 0.75%

MICONAZOLE NITRATE
- Crm 2% – **5% DV May-24 to 2026** — 0.90 — 15 g — **Multichem**
- ➥ Lotn 2% – **Restricted:** For continuation only
- Tinc 2%

NYSTATIN
- Crm 100,000 u per g

Antiparasitics

DIMETHICONE
- Lotn 4% – **5% DV Jun-26 to 2028** — 4.60 — 200 ml — **healthE Dimethicone 4% Lotion**

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
MALATHION [MALDISON]			
Lotn 0.5%			
Shampoo 1%			
PERMETHRIN			
Lotn 5% – **5% DV Feb-24 to 2026**	4.28	30 ml	**A-Scabies**
PHENOTHRIN			
Shampoo 0.5%			

Antiacne Preparations

ADAPALENE			
Crm 0.1%			
Gel 0.1%			
BENZOYL PEROXIDE			
Soln 5%			
ISOTRETINOIN			
Cap 5 mg – **5% DV Dec-24 to 2027**	11.26	60	**Oratane**
Cap 10 mg – **5% DV Dec-24 to 2027**	18.75	120	**Oratane**
Cap 20 mg – **5% DV Dec-24 to 2027**	26.73	120	**Oratane**
TRETINOIN			
Crm 0.05% – **5% DV Feb-25 to 2027**	16.82	50 g	**ReTrieve**

Antipruritic Preparations

CALAMINE			
Crm, aqueous, BP – **5% DV Apr-25 to 2027**	3.45	100 g	**healthE Calamine Aqueous**
CROTAMITON			
Crm 10% – **5% DV Feb-25 to 2027**	3.49	20 g	**Itch-Soothe**

Barrier Creams and Emollients

Barrier Creams

DIMETHICONE			
Crm 10% pump bottle	4.52	460 g	healthE Dimethicone 10%
Crm 5% pump bottle – **5% DV Jun-26 to 2028**	4.14	460 g	**Hydralock**
	4.30		healthE Dimethicone 5%
Crm 5% tube – **5% DV Jun-26 to 2028**	1.52	100 g	**healthE Dimethicone 5%**
(healthE Dimethicone 5% Crm 5% pump bottle to be delisted 1 June 2026)			
ZINC			
Crm			*e.g. Zinc Cream (Orion-) ;Zinc Cream (PSM)*
Oint			*e.g. Zinc oxide (PSM)*
Paste			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
ZINC AND CASTOR OIL			
Crm	1.63	20 g	Orion
Oint	4.25	500 g	Evara
Note: DV limit applies to the pack sizes of greater than 30 g.			
Oint, BP	1.26	20 g	healthE
Note: DV limit applies to the pack sizes of 30 g or less.			
ZINC WITH WOOL FAT			
Crm zinc 15.25% with wool fat 4%			*e.g. Sudocrem*

Emollients

AQUEOUS CREAM			
Crm 100 g – **5% DV Mar-25 to 2027**	1.25	100 g	**Evara**
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm 500 g – **5% DV Mar-25 to 2027**	1.65	500 g	**Evara**
Note: DV limit applies to the pack sizes of greater than 100 g.			
CETOMACROGOL			
Crm BP, 100 g – **5% DV Jun-25 to 2027**	0.99	100 g	**Cetomacrogol Cream AFT**
Crm BP, 500 g – **5% DV Feb-25 to 2027**	2.29	500 g	**Cetomacrogol-AFT**
CETOMACROGOL WITH GLYCEROL			
Crm 90% with glycerol 10% – **5% DV Dec-25 to 2028**	1.92	460 g	**Evara**
	3.25	920 g	**Evara**
Note: DV limit applies to the pack sizes of greater than 100 g.			
Crm 90% with glycerol 10%,	1.65	100 g	healthE
Note: DV limit applies to the pack sizes of 100 g or less.			
EMULSIFYING OINTMENT			
Oint BP – **5% DV Feb-24 to 2026**	2.30	100 g	**Jaychem**
Note: DV limit applies to pack sizes of less than 200 g.			
Oint BP, 500 g – **5% DV May-24 to 2026**	3.13	500 g	**Evara Emulsifying Ointment**
Note: DV limit applies to pack sizes of greater than 200 g.			
GLYCEROL WITH PARAFFIN			
Crm glycerol 10% with white soft paraffin 5% and liquid paraffin 10%			*e.g. QV cream*
OIL IN WATER EMULSION			
Crm, 100 g – **5% DV Apr-25 to 2027**	1.43	100 g	**Fatty Emulsion Cream (Evara)**
Note: DV limit applies to the pack sizes of 100 g or less.			
Crm, 500 g – **5% DV Apr-25 to 2027**	2.10	500 g	**Fatty Emulsion Cream (Evara)**
Note: DV limit applies to the pack sizes of greater than 100 g.			

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
PARAFFIN			
Oint liquid paraffin 50% with white soft paraffin 50% – **5% DV Aug-26 to 2028**	1.84	100 g	White Soft Liquid Paraffin AFT
	1.78		**healthE**
Note: DV limit applies to the pack sizes of 100 g or less.			
White soft	0.79	10 g	healthE
Note: DV limit applies to pack sizes of 30 g or less, and to both white soft paraffin and yellow soft paraffin.			
White soft, – **5% DV Jun-24 to 2026**	4.74	450 g	**EVARA White Soft Paraffin**
Note: DV limit applies to the pack sizes of 500 g or less and greater than 30 g.			
Yellow soft			
Lotn liquid paraffin 85%			*e.g QV Bath Oil*
(White Soft Liquid Paraffin AFT Oint liquid paraffin 50% with white soft paraffin 50% to be delisted 1 August 2026)			
PARAFFIN WITH WOOL FAT			
Lotn liquid paraffin 15.9% with wool fat 0.6%			*e.g. AlphaKeri;BK ;DP; Hydroderm Lotn*
Lotn liquid paraffin 91.7% with wool fat 3%			*e.g. Alpha Keri Bath Oil*
UREA			
Crm 10%	1.37	100 g	healthE Urea Cream
WOOL FAT			
Crm			

Corticosteroids

BETAMETHASONE DIPROPIONATE			
Crm 0.05% – **5% DV Jul-24 to 2026**	36.00	50 g	**Diprosone**
Note: DV limit applies to the pack sizes of greater than 30 g.			
Oint 0.05% – **5% DV Jul-24 to 2026**	36.00	50 g	**Diprosone**
Note: DV limit applies to the pack sizes of greater than 30 g.			
BETAMETHASONE VALERATE			
Crm 0.1% – **5% DV Feb-25 to 2027**	5.85	50 g	**Beta Cream**
Oint 0.1% – **5% DV Feb-25 to 2027**	7.90	50 g	**Beta Ointment**
Lotn 0.1% – **5% DV May-25 to 2027**	30.00	50 ml	**Betnovate**
CLOBETASOL PROPIONATE			
Crm 0.05% – **5% DV Feb-26 to 2028**	3.75	30 g	**Dermol**
Oint 0.05% – **5% DV Feb-26 to 2028**	3.68	30 g	**Dermol**
CLOBETASONE BUTYRATE			
Crm 0.05%			
DIFLUCORTOLONE VALERATE – **Restricted:** For continuation only			
➥ Crm 0.1%			
➥ Fatty oint 0.1%			
HYDROCORTISONE			
Crm 1%, 30 g – **5% DV Jul-26 to 2028**	1.75	30 g	**Ethics**
Note: DV limit applies to the pack sizes of less than or equal to 100 g.			
Crm 1%, 500 g – **5% DV Feb-26 to 2028**	20.40	500 g	**Noumed**
Note: DV limit applies to the pack sizes of greater than 100 g.			
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – **5% DV Jun-24 to 2026**	12.83	250 ml	**DP Lotn HC**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
HYDROCORTISONE BUTYRATE			
Crm 0.1%	4.85	100 g	Locoid Lipocream
Oint 0.1%	10.28	100 g	Locoid
Milky emul 0.1%	12.33	100 ml	Locoid Crelo
METHYLPREDNISOLONE ACEPONATE			
Crm 0.1% – **5% DV Feb-24 to 2026**	4.95	15 g	**Advantan**
Oint 0.1% – **5% DV Feb-24 to 2026**	4.95	15 g	**Advantan**
MOMETASONE FUROATE			
Crm 0.1% – **5% DV Feb-25 to 2027**	2.25	15 g	**Elocon Alcohol Free**
	3.50	50 g	**Elocon Alcohol Free**
Oint 0.1% – **5% DV Feb-25 to 2027**	2.25	15 g	**Elocon**
	3.50	50 g	**Elocon**
Lotn 0.1% – **5% DV Feb-25 to 2027**	4.99	30 ml	**Elocon**
TRIAMCINOLONE ACETONIDE			
Crm 0.02% – **5% DV Feb-24 to 2026**	6.49	100 g	**Aristocort**
Oint 0.02% – **5% DV Feb-24 to 2026**	6.54	100 g	**Aristocort**

Corticosteroids with Anti-Infective Agents

BETAMETHASONE VALERATE WITH CLIOQUINOL – **Restricted** see terms below

⬇ Crm 0.1% with clioquiniol 3%

➥ **Restricted** (RS1125)

Initiation

Either:

1 For the treatment of intertrigo; or

2 For continuation use.

BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID]

Crm 0.1% with sodium fusidate (fusidic acid) 2%

HYDROCORTISONE WITH MICONAZOLE			
Crm 1% with miconazole nitrate 2% – **5% DV Feb-25 to 2027**	2.85	15 g	**Micreme H**
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN			
Oint 1% with natamycin 1% and neomycin sulphate 0.5%	4.34	15 g	Pimafucort

TRIAMCINOLONE ACETONIDE WITH NEOMYCIN SULPHATE, GRAMICIDIN AND NYSTATIN

Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g

Psoriasis and Eczema Preparations

ACITRETIN			
Cap 10 mg – **5% DV Jul-24 to 2026**	26.20	60	**Novatretin**
Cap 25 mg – **5% DV Jul-24 to 2026**	57.37	60	**Novatretin**
BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g – **5% DV Dec-24 to 2027**	40.92	60 g	**Daivobet**
Oint 500 mcg with calcipotriol 50 mcg per g – **5% DV Dec-24 to 2027**	14.31	30 g	**Daivobet**
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g	Daivonex

COAL TAR WITH SALICYLIC ACID AND SULPHUR

Oint 12% with salicylic acid 2% and sulphur 4%

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
METHOXSALEN [8-METHOXYPSORALEN]			
Tab 10 mg			
Lotn 1.2%			
PIMECROLIMUS – **Restricted** see terms below			
⬇ Crm 1% – **5% DV Feb-24 to 2026**	33.00	15 g	**Elidel**
➥ **Restricted** (RS1781)			
Initiation			
Dermatologist, paediatrician or ophthalmologist			
Both:			
1 Patient has atopic dermatitis on the eyelid; and			
2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.			
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN			
Soln 2.3% with trolamine laurilsulfate and fluorescein sodium – **5% DV Feb-24 to 2026**	5.41	500 ml	**Pinetarsol**
POTASSIUM PERMANGANATE			
Tab 400 mg			
Crystals			
TACROLIMUS			
⬇ Oint 0.1% – **5% DV Dec-23 to 2026**	33.00	30 g	**Zematop**
➥ **Restricted** (RS1859)			
Initiation			
Dermatologist or paediatrician			
Both:			
1 Patient has atopic dermatitis on the face; and			
2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.			

Scalp Preparations

BETAMETHASONE VALERATE			
Scalp app 0.1% – **5% DV Feb-25 to 2027**	12.95	100 ml	**Beta Scalp**
CLOBETASOL PROPIONATE			
Scalp app 0.05% – **5% DV Feb-26 to 2028**	6.90	30 ml	**Dermol**
HYDROCORTISONE BUTYRATE			
Scalp lotn 0.1%	6.57	100 ml	Locoid

Wart Preparations

PODOPHYLLOTOXIN			
Soln 0.5%	33.60	3.5 ml	Condyline
SILVER NITRATE			
Sticks with applicator			

Other Skin Preparations

DIPHEMANIL METILSULFATE			
Powder 2%			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
IMIQUIMOD			
Crm 5%, 250 mg sachet	34.50	12	Aldara
	21.72	24	Padagis
			Perrigo
SUNSCREEN, PROPRIETARY			
Lotn	6.50	200 g	Marine Blue Lotion SPF 50+

Antineoplastics

FLUOROURACIL SODIUM			
Crm 5% – **5% DV Dec-24 to 2027**	5.56	20 g	**Efudix**
METHYL AMINOLEVULINATE HYDROCHLORIDE – **Restricted** see terms below			
⬇ Crm 16%			

➥ **Restricted** (RS1127)
Dermatologist or plastic surgeon

Wound Management Products

CALCIUM GLUCONATE			
Gel 2.5%			*e.g. Orion*

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Anti-Infective Agents

ACETIC ACID
- Soln 3%
- Soln 5%

ACETIC ACID WITH HYDROXYQUINOLINE, GLYCEROL AND RICINOLEIC ACID
- Jelly 0.94% with hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator

CHLORHEXIDINE GLUCONATE
- Crm 1%
- Lotn 1%

CLOTRIMAZOLE

Vaginal crm 1% with applicator – **5% DV Jul-26 to 2028**	4.20	35 g	**Clomazol**
Vaginal crm 2% with applicator – **5% DV Jul-26 to 2028**	4.60	20 g	**Clomazol**

MICONAZOLE NITRATE

Vaginal crm 2% with applicator	6.89	40 g	Micreme

NYSTATIN

Vaginal crm 100,000 u per 5 g with applicator(s) – **5% DV Feb-24 to 2026**	5.70	75 g	**Nilstat**

Contraceptives

Antiandrogen Oral Contraceptives

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tablets – **5% DV Feb-24 to 2026**	5.08	168	**Ginet**

Combined Oral Contraceptives

ETHINYLOESTRADIOL WITH DESOGESTREL
- Tab 20 mcg with desogestrel 150 mcg
- Tab 30 mcg with desogestrel 150 mcg

ETHINYLOESTRADIOL WITH LEVONORGESTREL

Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets – **5% DV Apr-26 to 2028**	2.00	84	**Lo-Oralcon 20 ED**
Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets – **5% DV Apr-26 to 2028**	2.30	84	**Oralcon 30 ED**

- Tab 20 mcg with levonorgestrel 100 mcg
- Tab 30 mcg with levonorgestrel 150 mcg

ETHINYLOESTRADIOL WITH NORETHISTERONE
- Tab 35 mcg with norethisterone 1 mg

Tab 35 mcg with norethisterone 1 mg and 7 inert tab	14.09	84	Brevinor 1/28

- Tab 35 mcg with norethisterone 500 mcg

NORETHISTERONE WITH MESTRANOL
- Tab 1 mg with mestranol 50 mcg

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Contraceptive Devices			
INTRA-UTERINE DEVICE			
IUD 29.1 mm length × 23.2 mm width	29.80	1	Choice 380 7med Nsha Silver/copper Short
IUD 33.6 mm length × 29.9 mm width	26.80	1	TCu 380 Plus Normal
IUD 35.5 mm length × 19.6 mm width	33.00	1	Cu 375 Standard
Emergency Contraception			
LEVONORGESTREL			
Tab 1.5 mg – **5% DV Jun-26 to 2028**	1.31	1	**Levonorgestrel -1 (Lupin)**
	1.75		Levonorgestrel BNM
(Levonorgestrel BNM Tab 1.5 mg to be delisted 1 June 2026)			
Progestogen-Only Contraceptives			
DESOGESTREL			
Tab 75 mcg	24.50	84	Cerazette
LEVONORGESTREL			
Tab 30 mcg	22.00	112	Microlut
Intra-uterine device 52 mg	269.50	1	Mirena
Intra-uterine device 13.5 mg	215.60	1	Jaydess
Subdermal implant (2 × 75 mg rods) – **5% DV Apr-25 to 2026**	106.92	2	**Jadelle**
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe	10.56	1	Depo-Provera
NORETHISTERONE			
Tab 350 mcg	12.25	84	Noriday
	13.23		Noriday 28
Obstetric Preparations			
Antiprogestogens			
MIFEPRISTONE			
Tab 200 mg			
Oxytocics			
CARBOPROST TROMETAMOL			
Inj 250 mcg per ml, 1 ml ampoule			
DINOPROSTONE			
Pessaries 10 mg			
Vaginal gel 1 mg in 3 g	65.39	1	Prostin E2
Vaginal gel 2 mg in 3 g	82.33	1	Prostin E2
ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule	160.00	5	DBL Ergometrine
OXYTOCIN			
Inj 5 iu per ml, 1 ml ampoule – **5% DV Mar-26 to 2028**	5.98	5	**Oxytocin BNM**
Inj 10 iu per ml, 1 ml ampoule – **5% DV Mar-26 to 2028**	7.18	5	**Oxytocin BNM**

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
OXYTOCIN WITH ERGOMETRINE MALEATE			
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoule – **5% DV Feb-26 to 2028**	41.47	5	**Syntometrine**

Tocolytics

PROGESTERONE			
Cap 100 mg	14.85	30	Utrogestan

TERBUTALINE – **Restricted** see terms below
⬇ Inj 500 mcg ampoule
➥ **Restricted** (RS1130)
Obstetrician

Oestrogens

OESTRIOL			
Crm 1 mg per g with applicator – **5% DV Feb-24 to 2026**	6.95	15 g	**Ovestin**
Pessaries 500 mcg – **5% DV Feb-24 to 2026**	7.55	15	**Ovestin**

Urologicals

5-Alpha Reductase Inhibitors

FINASTERIDE – **Restricted** see terms below
⬇ Tab 5 mg – **5% DV Dec-23 to 2026** 4.79 100 **Ricit**
➥ **Restricted** (RS1131)
Initiation
Both:
1 Patient has symptomatic benign prostatic hyperplasia; and
2 Either:
2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoceptor Blockers

TAMSULOSIN HYDROCHLORIDE – **Restricted** see terms below
⬇ Cap 400 mcg – **5% DV Feb-26 to 2028** 28.56 100 **Tamsulosin-Rex**
➥ **Restricted** (RS1132)
Initiation
Both:
1 Patient has symptomatic benign prostatic hyperplasia; and
2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Urinary Alkalisers

POTASSIUM CITRATE – **Restricted** see terms below
⬇ Oral liq 3 mmol per ml 37.49 200 ml Biomed
➥ **Restricted** (RS1133)
Initiation
Both:
1 The patient has recurrent calcium oxalate urolithiasis; and
2 The patient has had more than two renal calculi in the two years prior to the application.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
SODIUM CITRO-TARTRATE			
Grans eff 4 g sachets – **5% DV Feb-24 to 2026**	3.50	28	**Ural**
Urinary Antispasmodics			
OXYBUTYNIN			
Tab 5 mg	5.42	100	Alchemy Oxybutynin
Oral liq 5 mg per 5 ml			
SOLIFENACIN SUCCINATE			
Tab 5 mg – **5% DV Jun-25 to 2027**	1.95	30	**Solifenacin succinate Max Health**
Tab 10 mg – **5% DV Jun-25 to 2027**	3.53	30	**Solifenacin succinate Max Health**

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Anabolic Agents

OXANDROLONE
⬇ Tab 2.5 mg
➥ **Restricted** (RS1302)
Initiation
For the treatment of burns patients.

Androgen Agonists and Antagonists

CYPROTERONE ACETATE
Tab 50 mg – **5% DV Jul-25 to 2027** ..... 17.05 — 50 — **Siterone**
Tab 100 mg – **5% DV Jul-25 to 2027** ..... 31.00 — 50 — **Siterone**

TESTOSTERONE
Gel (transdermal) 16.2 mg per g, 88 g – **5% DV Apr-25 to 2027** ..... 52.00 — 60 — **Testogel**

TESTOSTERONE CIPIONATE
Inj 100 mg per ml, 10 ml vial ..... 97.75 — 1 — Depo-Testosterone

TESTOSTERONE ESTERS
Inj testosterone decanoate 100 mg, testosterone isocarproate 60 mg, testosterone phenylpropionate 60 mg and testosterone propionate 30 mg per ml, 1 ml ampoule

TESTOSTERONE UNDECANOATE
➥ Cap 40 mg – **Restricted:** For continuation only
Inj 250 mg per ml, 4 ml vial ..... 86.00 — 1 — Reandron 1000

Calcium Homeostasis

CALCITONIN
Inj 100 iu per ml, 1 ml ampoule ..... 121.00 — 5 — Miacalcic

CINACALCET – **Restricted** see terms below
⬇ Tab 30 mg – **5% DV Dec-24 to 2027** ..... 25.24 — 28 — **Cinacalet Devatis**
⬇ Tab 60 mg – **5% DV Dec-24 to 2027** ..... 50.47 — 28 — **Cinacalet Devatis**
➥ **Restricted** (RS1931)
Initiation – parathyroid carcinoma or calciphylaxis
Nephrologist or endocrinologist
Re-assessment required after 6 months
Either:

1 All of the following:
- 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
- 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
- 1.3 The patient is symptomatic; or

2 All of the following:
- 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
- 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
- 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

thiosulfate.

Continuation – parathyroid carcinoma or calciphylaxis

Nephrologist or endocrinologist

Both:

1 The patient's serum calcium level has fallen to < 3mmol/L; and
2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

Initiation – primary hyperparathyroidism

All of the following:

1 Patient has primary hyperparathyroidism; and
2 Either:
  2.1 Patient has hypercalcaemia of more than 3 mmol/L with or without symptoms; or
  2.2 Patient has hypercalcaemia of more than 2.85 mmol/L with symptoms; and
3 Surgery is not feasible or has failed; and
4 Patient has other comorbidities, severe bone pain, or calciphylaxis.

Initiation – secondary or tertiary hyperparathyroidism

Re-assessment required after 6 months

All of the following:

1 Either:
  1.1 Patient has tertiary hyperparathyroidism and markedly elevated parathyroid hormone (PTH) with hypercalcaemia; or
  1.2 Patient has symptomatic secondary hyperparathyroidism and elevated PTH; and
2 Patient is on renal replacement therapy; and
3 Any of the following:
  3.1 Residual parathyroid tissue has not been localised despite repeat unsuccessful parathyroid explorations; or
  3.2 Parathyroid tissue is surgically inaccessible; or
  3.3 Parathyroid surgery is not feasible.

Continuation – secondary or tertiary hyperparathyroidism

Re-assessment required after 12 months

Either:

1 The patient has had a kidney transplant, and following a treatment free interval of at least 12 weeks a clinically acceptable parathyroid hormone (PTH) level to support ongoing cessation of treatment has not been reached; or
2 The patient has not received a kidney transplant and trial of withdrawal of cinacalcet is clinically inappropriate.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial – **5% DV Dec-24 to 2027**	15.65	1	Zoledronic Acid Injection Mylan **Zoledronic acid Viatris**

(Zoledronic Acid Injection Mylan Inj 4 mg per 5 ml, vial to be delisted 1 November 2026)

Corticosteroids

BETAMETHASONE

Tab 500 mcg

Inj 4 mg per ml, 1 ml ampoule

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml ampoule

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
DEXAMETHASONE			
Tab 0.5 mg – **5% DV Feb-25 to 2027**	1.80	30	**Dexmethsone**
Tab 4 mg – **5% DV Feb-25 to 2027**	3.18	30	**Dexmethsone**
Oral liq 1 mg per ml	53.86	25 ml	Biomed
DEXAMETHASONE PHOSPHATE			
Inj 4 mg per ml, 1 ml ampoule – **5% DV Mar-26 to 2028**	6.88	10	**Dexamethasone Medsurge**
Inj 4 mg per ml, 2 ml ampoule – **5% DV Mar-26 to 2028**	10.98	10	**Dexamethasone Medsurge**
FLUDROCORTISONE ACETATE			
Tab 100 mcg – **5% DV Dec-25 to 2028**	8.05	100	**Florinef**
HYDROCORTISONE			
Tab 5 mg	8.10	100	Douglas
Tab 20 mg	20.32	100	Douglas
Inj 100 mg vial – **5% DV Dec-24 to 2027**	3.96	1	**Solu-Cortef**
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Tab 4 mg	112.00	100	Medrol
Tab 100 mg	223.10	20	Medrol
Inj 40 mg vial	22.30	1	Solu-Medrol Act-O-Vial
Inj 125 mg vial	34.10	1	Solu-Medrol Act-O-Vial
Inj 500 mg vial	43.01	1	Solu-Medrol Act-O-Vial
Inj 1 g vial	52.54	1	Solu-Medrol
METHYLPREDNISOLONE ACETATE			
Inj 40 mg per ml, 1 ml vial	47.06	5	Depo-Medrol
PREDNISOLONE			
Oral liq 5 mg per ml – **5% DV Dec-24 to 2027**	6.00	30 ml	**Redipred**
Enema 200 mcg per ml, 100 ml			
PREDNISONE			
Tab 1 mg	18.58	500	Prednisone Clinect
Tab 2.5 mg	21.04	500	Prednisone Clinect
Tab 5 mg	19.30	500	Prednisone Clinect
Tab 20 mg	50.51	500	Prednisone Clinect
TRIAMCINOLONE ACETONIDE			
Inj 10 mg per ml, 1 ml ampoule – **10% DV Feb-24 to 2026**	21.42	5	**Kenacort-A 10**
Inj 40 mg per ml, 1 ml ampoule – **5% DV Feb-24 to 2026**	52.63	5	**Kenacort-A 40**
TRIAMCINOLONE HEXACETONIDE			
Inj 20 mg per ml, 1 ml vial			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Hormone Replacement Therapy

Oestrogens

OESTRADIOL

Tab 1 mg

Gel (transdermal) 0.06% (750 mcg/actuation) – **5% DV Nov-24 to 31 Oct 2027**

	14.25	80 g	**Estrogel**
Patch 25 mcg per day – **5% DV Dec-25 to 2027**	8.89	8	**Estradiol TDP Mylan**
	16.23		Estradot
Patch 50 mcg per day – **5% DV Dec-25 to 2027**	9.26	8	**Estradiol TDP Mylan**
	15.79		Estradot
Patch 75 mcg per day – **5% DV Dec-25 to 2027**	10.33	8	**Estradiol TDP Mylan**
	16.53		Estradot
Patch 100 mcg per day – **5% DV Dec-25 to 2027**	10.59	8	**Estradiol TDP Mylan**
	16.18		Estradot

OESTRADIOL VALERATE

Tab 1 mg – **5% DV Dec-25 to 2028**	12.36	84	**Progynova**
Tab 2 mg – **5% DV Dec-25 to 2028**	12.36	84	**Progynova**

OESTROGENS (CONJUGATED EQUINE)

Tab 300 mcg

Tab 625 mcg

Progestogen and Oestrogen Combined Preparations

OESTRADIOL WITH NORETHISTERONE ACETATE

Tab 1 mg with 0.5 mg norethisterone acetate

Tab 2 mg with 1 mg norethisterone acetate

Tab 2 mg with 1 mg norethisterone acetate (10), and tab 2 mg oestradiol (12) and tab 1 mg oestradiol (6)

OESTROGENS WITH MEDROXYPROGESTERONE ACETATE

Tab 625 mcg conjugated equine with 2.5 mg medroxyprogesterone acetate

Tab 625 mcg conjugated equine with 5 mg medroxyprogesterone acetate

Progestogens

MEDROXYPROGESTERONE ACETATE

Tab 2.5 mg	7.54	30	Provera
Tab 5 mg	23.15	100	Provera
Tab 10 mg	11.10	30	Provera

Other Endocrine Agents

CABERGOLINE – **Restricted** see terms below

⬇ Tab 0.5 mg	4.43	2	Dostinex
	17.94	8	Dostinex

➥ **Restricted** (RS1855)

Initiation

Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
continued…			
1 Inhibition of lactation; or			
2 Patient has hyperprolactinemia; or			
3 Patient has acromegaly.			
Note: Indication marked with * is an unapproved indication.			
CLOMIFENE CITRATE			
Tab 50 mg	29.84	10	Mylan Clomiphen
GESTRINONE			
Cap 2.5 mg			
METYRAPONE			
Cap 250 mg			
PENTAGASTRIN			
Inj 250 mcg per ml, 2 ml ampoule			

Other Oestrogen Preparations

OESTRADIOL			
Implant 50 mg			
OESTRIOL			
Tab 2 mg – **5% DV Feb-24 to 2026**	7.70	30	**Ovestin**

Other Progestogen Preparations

MEDROXYPROGESTERONE			
Tab 100 mg	153.60	100	Provera HD
NORETHISTERONE			
Tab 5 mg	5.49	30	Primolut N

Pituitary and Hypothalamic Hormones and Analogues

CORTICORELIN (OVINE)			
Inj 100 mcg vial			
THYROTROPIN ALFA			
Inj 900 mcg vial			

Adrenocorticotropic Hormones

TETRACOSACTIDE [TETRACOSACTRIN]			
Inj 250 mcg per ml, 1 ml ampoule	86.25	1	Synacthen UK Synacthen
Inj 1 mg per ml, 1 ml ampoule	690.00	1	Synacthen Depot

GnRH Agonists and Antagonists

BUSERELIN			
Inj 1 mg per ml, 5.5 ml vial			
GONADORELIN			
Inj 100 mcg vial			
GOSERELIN			
Implant 3.6 mg, syringe – **5% DV Apr-24 to 2026**	66.48	1	**Zoladex**
Implant 10.8 mg, syringe – **5% DV Apr-24 to 2026**	138.23	1	**Zoladex**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
LEUPRORELIN ACETATE			
Inj 3.75 mg prefilled dual chamber syringe	221.60	1	Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe	591.68	1	Lucrin Depot 3-month

Gonadotrophins

CHORIOGONADOTROPIN ALFA
Inj 250 mcg in 0.5 ml syringe

Growth Hormone

SOMATROPIN – **Restricted** see terms below

⬇	Inj 5 mg cartridge – **5% DV Feb-25 to 2027**	80.21	1	**Omnitrope** Omnitrope AU
⬇	Inj 10 mg cartridge – **5% DV Feb-25 to 2027**	80.21	1	**Omnitrope**
⬇	Inj 15 mg cartridge – **5% DV Feb-25 to 2027**	139.50	1	**Omnitrope**

➥ **Restricted** (RS1826)

Initiation – growth hormone deficiency in children
Endocrinologist or paediatric endocrinologist
Re-assessment required after 12 months
Either:

1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
2 All of the following:

2.1 Height velocity < 25th percentile for age; and adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
2.5 Appropriate imaging of the pituitary gland has been obtained.

Continuation – growth hormone deficiency in children
Endocrinologist or paediatric endocrinologist
Re-assessment required after 12 months
All of the following:

1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
5 No malignancy has developed since starting growth hormone.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

1 The patient has a post-natal genotype confirming Turner Syndrome; and
2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
3 A current bone age is < 14 years.

Continuation – Turner syndrome

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

1 Height velocity greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
3 A current bone age is 14 years or under; and
4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
5 No malignancy has developed since starting growth hormone.

Initiation – short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Continuation – short stature without growth hormone deficiency

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
3 Current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initiation – short stature due to chronic renal insufficiency

Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of a endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

1 The patient's height is more than 2 standard deviations below the mean; and
2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
5 The patient is under the supervision of a specialist with expertise in renal medicine; and
6 Either:
  6.1 The patient has a GFR less than or equal to 30 ml/min/1.73 m^2 as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l)) × 40 = corrected GFR (ml/min/1.73 m^2) in a child who may or may not be receiving dialysis; or
  6.2 The patient has received a renal transplant and has received < 5mg/ m^2 /day of prednisone or equivalent for at least 6 months.

Continuation – short stature due to chronic renal insufficiency
Endocrinologist, paediatric endocrinologist or renal physician on the recommendation of a endocrinologist or paediatric endocrinologist
Re-assessment required after 12 months
All of the following:

1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
5 No malignancy has developed after growth hormone therapy was commenced; and
6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
7 The patient has not received renal transplantation since starting growth hormone treatment; and
8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initiation – Prader-Willi syndrome
Endocrinologist or paediatric endocrinologist
Re-assessment required after 12 months
All of the following:

1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
2 The patient is aged six months or older; and
3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
5 Either:
  5.1 Both:
    5.1.1 The patient is aged two years or older; and
    5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
  5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Continuation – Prader-Willi syndrome
Endocrinologist or paediatric endocrinologist
Re-assessment required after 12 months
All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
4 No serious adverse effect that the patient's specialist con siders is likely to be attributable to growth hormone treatment has occurred; and
5 No malignancy has developed after growth hormone therapy was commenced; and
6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initiation – adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

All of the following:

1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
3 The patient has severe growth hormone deficiency (see notes); and
4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until it is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Continuation – adults and adolescents

Endocrinologist or paediatric endocrinologist

Re-assessment required after 12 months

Any of the following:

1 All of the following:
  1.1 The patient has been treated with somatropin for < 12 months; and
  1.2 There has been an improvement in the Quality of Life Assessment defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  1.3 Serum IGF-I levels have increased to within ±1SD of the mean of the normal range for age and sex; and
  1.4 The dose of somatropin does not exceed 0.7 mg per day for male patients, or 1 mg per day for female patients; or
2 All of the following:
  2.1 The patient has been treated with somatropin for more than 12 months; and
  2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or

3 All of the following:

3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and

3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and

3.3 The patient has severe growth hormone deficiency (see notes); and

3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and

3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

The dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Thyroid and Antithyroid Preparations

CARBIMAZOLE

Tab 5 mg – **5% DV Dec-25 to 2028** ..... 7.56 100 **Neo-Mercazole**

IODINE

Soln BP 50 mg per ml

LEVOTHYROXINE

Tab 25 mcg

Tab 50 mcg

Tab 100 mcg

LIOTHYRONINE SODIUM

⬇ Tab 20 mcg

➥ **Restricted** (RS1301)

Initiation

For a maximum of 14 days' treatment in patients with thyroid cancer who are due to receive radioiodine therapy.

Inj 10 mcg vial

Inj 20 mcg vial

Inj 100 mcg vial

POTASSIUM IODATE

Tab 170 mg

POTASSIUM PERCHLORATE

Cap 200 mg

PROPYLTHIOURACIL – **Restricted** see terms on the next page

⬇ Tab 50 mg ..... 35.00 100 PTU

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1276)
Initiation
Both:

1 The patient has hyperthyroidism; and
2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

PROTIRELIN
Inj 100 mcg per ml, 2 ml ampoule

Vasopressin Agents

ARGIPRESSIN [VASOPRESSIN]
Inj 20 u per ml, 1 ml ampoule

DESMOPRESSIN
Wafer 120 mcg ..... 47.00 — 30 — Minirin Melt

DESMOPRESSIN ACETATE
Tab 100 mcg ..... 25.00 — 30 — Minirin
Tab 200 mcg ..... 54.45 — 30 — Minirin
Inj 4 mcg per ml, 1 ml ampoule
Inj 15 mcg per ml, 1 ml ampoule
Nasal drops 100 mcg per ml
Nasal spray 10 mcg per dose, 6 ml – **5% DV Apr-25 to 2026** ..... 34.95 — 60 — **Desmopressin-PH&T**

TERLIPRESSIN
Inj 0.2 mg per ml, 5 ml vial – **5% DV Feb-25 to 2027** ..... 110.00 — 5 — **Terlipressin Ever Pharma**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Antibacterials

Aminoglycosides

AMIKACIN – **Restricted** see terms below

⬇ Inj 5 mg per ml, 10 ml syringe

⬇ Inj 5 mg per ml, 5 ml syringe

⬇ Inj 15 mg per ml, 5 ml syringe

⬇ Inj 250 mg per ml, 2 ml vial – **5% DV Dec-24 to 2027** ..... 169.97 — 5 — **DBL Amikacin**

➥ **Restricted** (RS1041)

Clinical microbiologist, infectious disease specialist or respiratory specialist

GENTAMICIN SULPHATE

Inj 10 mg per ml, 1 ml ampoule ..... 102.60 — 5 — DBL Gentamicin

Inj 10 mg per ml, 2 ml ampoule

Inj 40 mg per ml, 2 ml ampoule ..... 18.38 — 10 — Pfizer

PAROMOMYCIN – **Restricted** see terms below

⬇ Cap 250 mg ..... 126.00 — 16 — Humatin

➥ **Restricted** (RS1603)

Clinical microbiologist, infectious disease specialist or gastroenterologist

STREPTOMYCIN SULPHATE – **Restricted** see terms below

⬇ Inj 400 mg per ml, 2.5 ml ampoule

➥ **Restricted** (RS1043)

Clinical microbiologist, infectious disease specialist or respiratory specialist

TOBRAMYCIN

⬇ Powder

➥ **Restricted** (RS1475)

Initiation

For addition to orthopaedic bone cement.

⬇ Inj 40 mg per ml, 2 ml vial – **5% DV Dec-24 to 2027** ..... 15.50 — 5 — **Tobramycin (Viatris)**

➥ **Restricted** (RS1044)

Clinical microbiologist, infectious disease specialist or respiratory specialist

⬇ Inj 100 mg per ml, 5 ml vial

➥ **Restricted** (RS1044)

Clinical microbiologist, infectious disease specialist or respiratory specialist

⬇ Solution for inhalation 60 mg per ml, 5 ml – **5% DV Dec-23 to 2026** ..... 395.00 — 56 dose — **Tobramycin BNM**

➥ **Restricted** (RS1435)

Initiation

Patient has cystic fibrosis.

Carbapenems

ERTAPENEM – **Restricted** see terms below

⬇ Inj 1 g vial ..... 70.00 — 1 — Invanz

➥ **Restricted** (RS1045)

Clinical microbiologist or infectious disease specialist

IMIPENEM WITH CILASTATIN – **Restricted** see terms on the next page

⬇ Inj 500 mg with 500 mg cilastatin vial ..... 60.00 — 1 — Imipenem+Cilastatin RBX

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
➥ **Restricted** (RS1046)			
Clinical microbiologist or infectious disease specialist			
MEROPENEM – **Restricted** see terms below			
⬇ Inj 500 mg vial – **5% DV Jun-24 to 2026**	33.48	10	**Meropenem-AFT**
⬇ Inj 1 g vial – **5% DV Jun-24 to 2026**	44.97	10	**Meropenem-AFT**
➥ **Restricted** (RS1047)			
Clinical microbiologist or infectious disease specialist			

Cephalosporins and Cephamycins - 1st Generation

CEFALEXIN			
Cap 250 mg – **5% DV Jul-26 to 2028**	3.90	20	**Cefalexin Lupin**
	3.85		Cephalexin ABM
Cap 500 mg – **5% DV Jul-26 to 2028**	3.33	20	**Cefalexin Sandoz**
	5.85		Cephalexin ABM
Grans for oral liq 25 mg per ml – **5% DV Dec-26 to 2028**	5.41	100 ml	**Cefalexin Sandoz**
	7.88		Flynn
Grans for oral liq 50 mg per ml – **5% DV Dec-26 to 2028**	7.54	100 ml	**Cefalexin Sandoz**
	10.38		Flynn
(Cephalexin ABM Cap 250 mg to be delisted 1 July 2026)			
(Cephalexin ABM Cap 500 mg to be delisted 1 July 2026)			
(Flynn Grans for oral liq 25 mg per ml to be delisted 1 December 2026)			
(Flynn Grans for oral liq 50 mg per ml to be delisted 1 December 2026)			
CEFAZOLIN			
Inj 500 mg vial – **5% DV Mar-24 to 2026**	3.39	5	**Cefazolin-AFT**
Inj 1 g vial – **5% DV Mar-24 to 2026**	3.59	5	**Cefazolin-AFT**
Inj 2 g vial – **5% DV Mar-24 to 2026**	7.09	5	**Cefazolin-AFT**

Cephalosporins and Cephamycins - 2nd Generation

CEFACLOR			
Cap 250 mg – **5% DV Feb-26 to 2028**	29.73	100	**Ranbaxy-Cefaclor**
Grans for oral liq 25 mg per ml – **5% DV Feb-26 to 2028**	5.83	100 ml	**Ranbaxy-Cefaclor**
CEFOXITIN			
Inj 1 g vial			
CEFUROXIME			
Tab 250 mg			
Inj 750 mg vial – **5% DV Jun-24 to 2026**	8.16	10	**Cefuroxime Devatis**
Inj 1.5 g vial – **5% DV Jun-24 to 2026**	13.01	10	**Cefuroxime Devatis**

Cephalosporins and Cephamycins - 3rd Generation

CEFOTAXIME			
Inj 500 mg vial	1.90	1	Cefotaxime Sandoz
Inj 1 g vial – **5% DV Dec-23 to 2026**	38.98	10	**DBL Cefotaxime**
CEFTAZIDIME – **Restricted** see terms below			
⬇ Inj 1 g vial – **5% DV Dec-23 to 2026**	25.80	10	**Ceftazidime Kabi**
➥ **Restricted** (RS1048)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
CEFTAZIDIME WITH AVIBACTAM – **Restricted** see terms on the next page			
⬇ Inj ceftazidime 2,000 mg with avibactam 500 mg, vial	2,250.00	10	Zavicefta

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS2104)

Initiation

Both:

1 Prescribed by, or recommended by a clinical microbiologist or infectious disease specialist, or in accordance with a protocol or guideline that has been endorsed by the Health NZ Hospital; and

2 Either:

2.1 Proven infection with a carbapenem-resistant micro-organism, based on microbiology report; or

2.2 Probable infection with a carbapenem-resistant micro-organism, based on assessment by a clinical microbiologist or infectious disease specialist..

CEFTRIAXONE

Inj 500 mg vial – **5% DV Feb-26 to 2028**	0.94	1	**Ceftriaxone-AFT**
Inj 1 g vial – **5% DV Feb-26 to 2028**	3.49	5	**Ceftriaxone-AFT**
Inj 2 g vial – **5% DV Feb-26 to 2028**	8.15	5	**Ceftriaxone-AFT**

Cephalosporins and Cephamycins - 4th Generation

CEFEPIME – **Restricted** see terms below

⬇ Inj 1 g vial – **5% DV Dec-24 to 2027**	3.19	1	**Cefepime-AFT**
⬇ Inj 2 g vial – **5% DV Dec-24 to 2027**	4.99	1	**Cefepime-AFT**

➥ **Restricted** (RS1049)

Clinical microbiologist or infectious disease specialist

Cephalosporins and Cephamycins - 5th Generation

CEFTAROLINE FOSAMIL – **Restricted** see terms below

⬇ Inj 600 mg vial	1,834.25	10	Zinforo

➥ **Restricted** (RS1446)

Initiation – multi-resistant organisn salvage therapy

Clinical microbiologist or infectious disease specialist

Either:

1 for patients where alternative therapies have failed; or

2 for patients who have a contraindication or hypersensitivity to standard current therapies.

Macrolides

AZITHROMYCIN – **Restricted** see terms below

⬇ Tab 250 mg			
⬇ Tab 500 mg – **5% DV Jan-26 to 2027**	2.80	2	**Zithromax**
⬇ Grans for oral liq 200 mg per 5 ml (40 mg per ml)	16.97	15 ml	Zithromax

➥ **Restricted** (RS1598)

Initiation – bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections

Any of the following:

1 Patient has received a lung transplant, stem cell transplant or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or

2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or

3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas related gram negative organisms*; or

4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – non-cystic fibrosis bronchiectasis*

Respiratory specialist or paediatrician

Re-assessment required after 12 months

All of the following:

1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
2 Patient is aged 18 and under; and
3 Either:
  3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis will be subsidised in the community.

Continuation – non-cystic fibrosis bronchiectasis*

Respiratory specialist or paediatrician

Re-assessment required after 12 months

All of the following:

1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

Note: Indications marked with * are unapproved indications. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis will be subsidised in the community.

Initiation – other indications

Re-assessment required after 5 days

For any other condition.

Continuation – other indications

Re-assessment required after 5 days

For any other condition.

CLARITHROMYCIN – **Restricted** see terms below

⬇	Tab 250 mg – **1% DV Feb-22 to 2027**	8.53	14	**Klacid**
		7.31	12	Klaricid
⬇	Tab 500 mg – **1% DV Feb-22 to 2027**	14.58	14	**Klacid**
⬇	Grans for oral liq 50 mg per ml	192.00	50 ml	Klacid
⬇	Inj 500 mg vial – **5% DV Jul-24 to 2026**	9.10	1	**Klacid IV**

➥ **Restricted** (RS1709)

Initiation – Tab 250 mg and oral liquid

Any of the following:

1 Atypical mycobacterial infection; or
2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents; or
3 Helicobacter pylori eradication; or
4 Prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated.

Initiation – Tab 500 mg

Helicobacter pylori eradication.

Initiation – Infusion

Any of the following:

1 Atypical mycobacterial infection; or
2 Mycobacterium tuberculosis infection where there is drug resistance or intolerance to standard pharmaceutical agents; or
3 Community-acquired pneumonia.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
ERYTHROMYCIN (AS ETHYLSUCCINATE)			
Tab 400 mg	35.82	100	E-Mycin
Grans for oral liq 200 mg per 5 ml	6.53	100 ml	E-Mycin
Grans for oral liq 400 mg per 5 ml	9.41	100 ml	E-Mycin
ERYTHROMYCIN (AS LACTOBIONATE)			
Inj 1 g vial – **5% DV Dec-25 to 2028**	10.00	1	**Erythrocin IV**
ERYTHROMYCIN (AS STEARATE) – **Restricted:** For continuation only			
➥ Tab 250 mg			
➥ Tab 500 mg			
ROXITHROMYCIN – **Some items restricted** see terms below			
⬇ Tab dispersible 50 mg			
Tab 150 mg – **5% DV Aug-23 to 2026**	13.19	50	**Arrow-Roxithromycin**
Tab 300 mg – **5% DV Aug-23 to 2026**	25.00	50	**Arrow-Roxithromycin**
➥ **Restricted** (RS1569)			
Initiation			
Only for use in patients under 12 years of age.			

Penicillins

AMOXICILLIN			
Cap 250 mg – **5% DV Feb-26 to 2028**	54.00	500	**Miro-Amoxicillin**
Cap 500 mg – **5% DV Nov-26 to 2028**	1.14	20	**Amoxycillin Sandoz**
	41.00	500	Miro-Amoxicillin
Grans for oral liq 125 mg per 5 ml – **5% DV Feb-24 to 2026**	2.22	100 ml	**Alphamox 125**
Grans for oral liq 250 mg per 5 ml – **5% DV Feb-24 to 2026**	2.81	100 ml	**Alphamox 250**
Inj 250 mg vial	15.97	10	Ibiamox
Inj 500 mg vial	17.43	10	Ibiamox
Inj 1 g vial	21.64	10	Ibiamox
(Miro-Amoxicillin Cap 500 mg to be delisted 1 November 2026)			
AMOXICILLIN WITH CLAVULANIC ACID			
Tab 500 mg with clavulanic acid 125 mg – **5% DV Feb-24 to 2026**	1.59	10	**Curam Duo 500/125**
Grans for oral liq 25 mg with clavulanic acid 6.25 mg per ml – **5% DV May-25 to 2027**	8.50	100 ml	**Augmentin**
Grans for oral liq 50 mg with clavulanic acid 12.5 mg per ml – **5% DV Jun-25 to 2027**	5.61	100 ml	**Amoxiclav Devatis Forte**
Inj 500 mg with clavulanic acid 100 mg vial – **5% DV Sep-25 to 2027**	22.48	10	**Synermox**
Inj 1,000 mg with clavulanic acid 200 mg vial – **5% DV Sep-25 to 2027**	26.90	10	Cerobact
	29.61		**Synermox**
BENZATHINE BENZYLPENICILLIN			
Inj 900 mg (1.2 million units) vial	43.24	1	Benzetacil
Inj 900 mg (1.2 million units) in 2.3 ml syringe	432.37	10	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]			
Inj 600 mg (1 million units) vial – **5% DV Feb-24 to 2026**	16.50	10	**Sandoz**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
FLUCLOXACILLIN			
Cap 250 mg – **5% DV Aug-25 to 2027**	22.58	250	**Staphlex**
Cap 500 mg – **5% DV Aug-25 to 2027**	72.71	500	**Staphlex**
Grans for oral liq 25 mg per ml – **5% DV Feb-25 to 2027**	4.89	100 ml	**AFT**
Grans for oral liq 50 mg per ml – **5% DV Feb-25 to 2027**	5.89	100 ml	**AFT**
Inj 250 mg vial – **5% DV Jul-24 to 2026**	42.60	10	**Flucloxin**
Inj 500 mg vial – **5% DV Jul-24 to 2026**	45.63	10	**Flucloxin**
Inj 1 g vial – **5% DV Feb-24 to 2026**	6.00	5	**Flucil**
PHENOXYMETHYLPENICILLIN [PENICILLIN V]			
Cap 250 mg – **5% DV Feb-25 to 2027**	7.68	50	**Cilicaine VK**
Cap 500 mg – **5% DV Feb-25 to 2027**	13.72	50	**Cilicaine VK**
Grans for oral liq 125 mg per 5 ml – **5% DV Feb-26 to 2028**	5.75	100 ml	**AFT**
Grans for oral liq 250 mg per 5 ml – **5% DV Feb-26 to 2028**	5.89	100 ml	**AFT**
PIPERACILLIN WITH TAZOBACTAM – **Restricted** see terms below			
⬇ Inj 4 g with tazobactam 0.5 g vial – **5% DV Dec-25 to 2028**	3.15	1	**PipTaz-AFT**
➥ **Restricted** (RS1053)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PROCAINE PENICILLIN			
Inj 1.5 g in 3.4 ml syringe			
TICARCILLIN WITH CLAVULANIC ACID – **Restricted** see terms below			
⬇ Inj 3 g with clavulanic acid 0.1 mg vial			
➥ **Restricted** (RS1054)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			

Quinolones

CIPROFLOXACIN – **Restricted** see terms below			
⬇ Tab 250 mg – **5% DV Nov-24 to 2026**	1.95	28	**Ipca-Ciprofloxacin**
⬇ Tab 500 mg – **5% DV Nov-24 to 2026**	3.10	28	**Ipca-Ciprofloxacin**
⬇ Tab 750 mg – **5% DV Dec-24 to 2026**	4.80	28	**Ipca-Ciprofloxacin**
⬇ Oral liq 50 mg per ml			
⬇ Oral liq 100 mg per ml			
⬇ Inj 2 mg per ml, 100 ml bag			
⬇ Inj 2 mg per ml, 100 ml bottle	166.50	10	Ciprofloxacin Kabi
➥ **Restricted** (RS1055)			
Clinical microbiologist or infectious disease specialist			
MOXIFLOXACIN – **Restricted** see terms below			
⬇ Tab 400 mg	42.00	5	Avelox
⬇ Inj 1.6 mg per ml, 250 ml bottle – **5% DV Feb-24 to 2026**	413.40	10	**Moxifloxacin Kabi**
➥ **Restricted** (RS2129)			

Initiation – Mycobacterium infection

Infectious disease specialist, clinical microbiologist or respiratory specialist

Any of the following:

1 Both:

1.1 Active tuberculosis; and

1.2 Any of the following:

1.2.1 Documented resistance to one or more first-line medications; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or

2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated; or
3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Initiation – Pneumonia
Infectious disease specialist or clinical microbiologist
Either:

1 Immunocompromised patient with pneumonia that is unresponsive to first-line treatment; or
2 Pneumococcal pneumonia or other invasive pneumococcal disease highly resistant to other antibiotics.

Initiation – Penetrating eye injury
Ophthalmologist
Five days treatment for patients requiring prophylaxis following a penetrating eye injury.

Initiation – Mycoplasma genitalium
All of the following:

1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium and is symptomatic; and
2 Either:
2.1 Has tried and failed to clear infection using azithromycin; or
2.2 Has laboratory confirmed azithromycin resistance; and
3 Treatment is only for 7 days.

Initiation – severe delayed beta-lactam allergy
Infectious disease specialist or clinical microbiologist
Individual has a history of severe delayed beta-lactam allergy.

NORFLOXACIN
Tab 400 mg ..... 245.00 100 Arrow-Norfloxacin

Tetracyclines

DEMECLOCYCLINE HYDROCHLORIDE
Tab 150 mg
Cap 150 mg
Cap 300 mg

DOXYCYCLINE
➥ Tab 50 mg – **Restricted:** For continuation only
Tab 100 mg ..... 64.43 500 Doxine
Inj 5 mg per ml, 20 ml vial

MINOCYCLINE
Tab 50 mg
➥ Cap 100 mg – **Restricted:** For continuation only

TETRACYCLINE
Tab 250 mg ..... 68.44 28 Accord
Cap 500 mg

TIGECYCLINE – **Restricted** see terms below
⬇ Inj 50 mg vial
➥ **Restricted** (RS1059)
Clinical microbiologist or infectious disease specialist

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Other Antibacterials			
AZTREONAM – **Restricted** see terms below			
⬇ Inj 1 g vial	364.92	10	Azactam
➥ **Restricted** (RS1277)			
Clinical microbiologist or infectious disease specialist			
CHLORAMPHENICOL – **Restricted** see terms below			
⬇ Inj 1 g vial			
➥ **Restricted** (RS1277)			
Clinical microbiologist or infectious disease specialist			
CLINDAMYCIN – **Restricted** see terms below			
⬇ Cap 150 mg – **5% DV Dec-24 to 2027**	4.94	24	**Dalacin C**
⬇ Oral liq 15 mg per ml			
⬇ Inj 150 mg per ml, 4 ml ampoule – **5% DV Mar-26 to 2028**	48.78	10	**Dalacin C**
➥ **Restricted** (RS1061)			
Clinical microbiologist or infectious disease specialist			
COLISTIN SULPHOMETHATE [COLESTIMETHATE] – **Restricted** see terms below			
⬇ Inj 2 million iu, 10 ml vial	216.67	10	Colomycin
➥ **Restricted** (RS1062)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
DAPTOMYCIN – **Restricted** see terms below			
⬇ Inj 500 mg vial – **5% DV Oct-26 to 2028**	115.36 44.90	1	Daptomycin Dr Reddy's **Daptomycin-AFT**
(Daptomycin Dr Reddy's Inj 500 mg vial to be delisted 1 October 2026)			
➥ **Restricted** (RS1063)			
Clinical microbiologist or infectious disease specialist			
FOSFOMYCIN – **Restricted** see terms below			
⬇ Powder for oral solution, 3 g sachet – **5% DV Apr-25 to 2027**	18.70	1	**UroFos**
➥ **Restricted** (RS1315)			
Clinical microbiologist or infectious disease specialist			
LINCOMYCIN – **Restricted** see terms below			
⬇ Inj 300 mg per ml, 2 ml vial			
➥ **Restricted** (RS1065)			
Clinical microbiologist or infectious disease specialist			
LINEZOLID – **Restricted** see terms below			
⬇ Tab 600 mg – **5% DV Dec-24 to 2027**	194.60	10	**Zyvox**
⬇ Oral liq 20 mg per ml	1,879.00	150 ml	Zyvox
⬇ Inj 2 mg per ml, 300 ml bottle – **5% DV Dec-24 to 2027**	155.00	10	**Linezolid Kabi**
➥ **Restricted** (RS1066)			
Clinical microbiologist or infectious disease specialist			
METHENAMINE (HEXAMINE) HIPPURATE			
Tab 1 g	19.95	100	Hiprex
NITROFURANTOIN			
Tab 50 mg – **5% DV Dec-24 to 2027**	22.20	100	**Nifuran** Nitrofurantoin Clinect
Tab 100 mg	37.50	100	Nifuran
Cap modified-release 100 mg – **5% DV Dec-23 to 2026**	81.20	100	**Macrobid**
(Nifuran Tab 50 mg to be delisted 1 November 2026)			
(Nifuran Tab 100 mg to be delisted 1 November 2026)			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
PIVMECILLINAM – **Restricted** see terms below			
⬇ Tab 200 mg			
➥ **Restricted** (RS1322)			
Clinical microbiologist or infectious disease specialist			
SODIUM FUSIDATE [FUSIDIC ACID] – **Restricted** see terms below			
⬇ Tab 250 mg	135.70	36	Fucidin
➥ **Restricted** (RS1064)			
Clinical microbiologist or infectious disease specialist			
SULFADIAZINE SODIUM – **Restricted** see terms below			
⬇ Tab 500 mg			*e.g. Sulfadiazin-Heyl; Wockhardt*
➥ **Restricted** (RS1067)			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
TEICOPLANIN – **Restricted** see terms below			
⬇ Inj 400 mg vial – **5% DV Apr-25 to 2027**	38.85	1	**Teicoplanin Medsurge**
➥ **Restricted** (RS1068)			
Clinical microbiologist or infectious disease specialist			
TRIMETHOPRIM			
Tab 100 mg			
Tab 300 mg – **5% DV Feb-25 to 2027**	27.83	50	**TMP**
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]			
Tab 80 mg with sulphamethoxazole 400 mg – **5% DV Feb-25 to 2027**	115.74	500	**Trisul**
Oral liq 8 mg with sulphamethoxazole 40 mg per ml – **5% DV Aug-25 to 2028**	4.95	100 ml	**Deprim**
Inj 16 mg with sulphamethoxazole 80 mg per ml, 5 ml ampoule			
VANCOMYCIN – **Restricted** see terms below			
⬇ Inj 500 mg vial – **5% DV Dec-25 to 2026**	3.38	1	**Vancomycin Viatris**
➥ **Restricted** (RS1069)			
Clinical microbiologist or infectious disease specialist			

Antifungals

Imidazoles

KETOCONAZOLE			
⬇ Tab 200 mg			
➥ **Restricted** (RS1410)			
Oncologist			

Polyene Antimycotics

AMPHOTERICIN B			
⬇ Inj (liposomal) 50 mg vial – **5% DV Apr-26 to 2028**	125.00	1	**Amphotericin Liposomal SUN**
➥ **Restricted** (RS1071)			
Initiation			
Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist			
Either:			

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
2 Both:
2.1 Possible invasive fungal infection; and
2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

⬇ Inj 50 mg vial
➥ **Restricted** (RS1316)
Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist

NYSTATIN

Tab 500,000 u	17.09	50	Nilstat
Cap 500,000 u	15.47	50	Nilstat

Triazoles

FLUCONAZOLE – **Restricted** see terms below

⬇ Cap 50 mg – **5% DV Dec-23 to 2026**	4.10	28	**Mylan**
⬇ Cap 150 mg – **5% DV Dec-23 to 2026**	0.45	1	Flucazole **Mylan**
⬇ Cap 200 mg – **5% DV Dec-23 to 2026**	8.90	28	**Mylan**
⬇ Oral liquid 50 mg per 5 ml	129.02	35 ml	Diflucan
⬇ Inj 2 mg per ml, 50 ml vial	11.20	1	Fluconazole-Baxter
⬇ Inj 2 mg per ml, 100 ml vial	5.20	1	Fluconazole-Baxter

➥ **Restricted** (RS1072)
Consultant

ITRACONAZOLE – **Restricted** see terms below

⬇ Cap 100 mg	6.83	15	Itraconazole Cresent Itrazole

⬇ Oral liquid 10 mg per ml
➥ **Restricted** (RS1073)
Clinical immunologist, clinical microbiologist, dermatologist or infectious disease specialist

POSACONAZOLE – **Restricted** see terms below

⬇ Tab modified-release 100 mg – **5% DV Dec-25 to 2028**	123.60	24	**Posaconazole Juno**
⬇ Oral liq 40 mg per ml – **5% DV Dec-25 to 2028**	308.26	105 ml	**Devatis**

➥ **Restricted** (RS2052)
Initiation
Haematologist or infectious disease specialist
Re-assessment required after 6 weeks
Both:

1 Either:
1.1 Patient has acute myeloid leukaemia; or
1.2 Patient is planned to receive a stem cell transplant and is at high risk for aspergillus infection; and
2 Patient is to be treated with high dose remission induction therapy or re-induction therapy.

Continuation
Haematologist or infectious disease specialist
Re-assessment required after 6 weeks
Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient has previously received posaconazole prophylaxis during remission induction therapy; and
2 Any of the following:
  2.1 Patient is to be treated with high dose remission re-induction therapy; or
  2.2 Patient is to be treated with high dose consolidation therapy; or
  2.3 Patient is receiving a high risk stem cell transplant.

Initiation – Invasive fungal infection prophylaxis
Any relevant practitioner
Re-assessment required after 6 months
Both:

1 The patient is at risk of invasive fungal infection; and
2 Either:
  2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Continuation – Invasive fungal infection prophylaxis
Any relevant practitioner
Re-assessment required after 6 months
Both:

1 The patient is at risk of invasive fungal infection; and
2 Either:
  2.1 Posaconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  2.2 Prescribing posaconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

VORICONAZOLE – **Restricted** see terms below

⬇	Tab 50 mg – **5% DV Aug-25 to 2028**	71.00	56	**Vttack**
⬇	Tab 200 mg – **5% DV Aug-25 to 2028**	263.00	56	**Vttack**
⬇	Powder for oral suspension 40 mg per ml	1,523.22	70 ml	Vfend
⬇	Inj 200 mg vial – **5% DV Dec-25 to 2028**	16.89	1	**AFT**

➥ **Restricted** (RS2053)

Initiation – Proven or probable aspergillus infection
Clinical microbiologist, haematologist or infectious disease specialist
Both:

1 Patient is immunocompromised; and
2 Patient has proven or probable invasive aspergillus infection.

Initiation – Possible aspergillus infection
Clinical microbiologist, haematologist or infectious disease specialist
All of the following:

1 Patient is immunocompromised; and
2 Patient has possible invasive aspergillus infection; and
3 A multidisciplinary team (including an infectious disease physician) considers the treatment to be appropriate.

Initiation – Resistant candidiasis infections and other moulds
Clinical microbiologist, haematologist or infectious disease specialist
All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient is immunocompromised; and
2 Either:
  2.1 Patient has fluconazole resistant candidiasis; or
  2.2 Patient has mould strain such as Fusarium spp. and Scedosporium spp; and
3 A multidisciplinary team (including an infectious disease physician or clinical microbiologist) considers the treatment to be appropriate.

Initiation – Invasive fungal infection prophylaxis
Any relevant practitioner
Re-assessment required after 6 months
Both:
1 The patient is at risk of invasive fungal infection; and
2 Either:
  2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Continuation – Invasive fungal infection prophylaxis
Any relevant practitioner
Re-assessment required after 6 months
Both:
1 The patient is at risk of invasive fungal infection; and
2 Either:
  2.1 Voriconazole is prescribed by, or recommended by a haematologist, transplant physician, infectious disease specialist, paediatric haematologist or paediatric oncologist; or
  2.2 Prescribing voriconazole is in accordance with a protocol or guideline that has been endorsed by the Health New Zealand - Te Whatu Ora Hospital in the specific settings where there is a greater than 10% risk of invasive fungal infection (IFI).

Other Antifungals

CASPOFUNGIN – **Restricted** see terms below

⬇	Inj 50 mg vial – **5% DV Mar-26 to 2028**	110.00	1	**Alchemy Caspofungin**
⬇	Inj 70 mg vial – **5% DV Mar-26 to 2028**	135.00	1	**Alchemy Caspofungin**

➥ **Restricted** (RS1076)
Initiation
Clinical microbiologist, haematologist, infectious disease specialist, oncologist, respiratory specialist or transplant specialist
Either:
1 Proven or probable invasive fungal infection, to be prescribed under an established protocol; or
2 Both:
  2.1 Possible invasive fungal infection; and
  2.2 A multidisciplinary team (including an infectious disease physician or a clinical microbiologist) considers the treatment to be appropriate.

FLUCYTOSINE – **Restricted** see terms below
⬇ Tab 500 mg
⬇ Cap 500 mg
➥ **Restricted** (RS1279)
Clinical microbiologist or infectious disease specialist

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
TERBINAFINE			
Tab 250 mg – **5% DV Feb-24 to 2026**	8.97	84	**Deolate**

Antimycobacterials

Antileprotics

CLOFAZIMINE – **Restricted** see terms below

⬇ Cap 50 mg

➥ **Restricted** (RS1077)

Clinical microbiologist, dermatologist or infectious disease specialist

DAPSONE – **Restricted** see terms below

⬇ Tab 25 mg	268.50	100	Dapsone
⬇ Tab 100 mg	329.50	100	Dapsone

➥ **Restricted** (RS1078)

Clinical microbiologist, dermatologist or infectious disease specialist

Antituberculotics

BEDAQUILINE – **Restricted** see terms below

⬇ Tab 100 mg	3,084.51	24	Sirturo

➥ **Restricted** (RS1977)

Initiation – multi-drug resistant tuberculosis

Limited to 6 months treatment

Both:

1 The person has multi-drug resistant tuberculosis (MDR-TB); and
2 Ministry of Health's Tuberculosis Clinical Network has reviewed the individual case and recommends bedaquiline as part of the treatment regimen.

CYCLOSERINE – **Restricted** see terms below

⬇ Cap 250 mg

➥ **Restricted** (RS1079)

Clinical microbiologist, infectious disease specialist or respiratory specialist

ETHAMBUTOL HYDROCHLORIDE – **Restricted** see terms below

⬇ Tab 100 mg			
⬇ Tab 400 mg	49.34	56	Myambutol

➥ **Restricted** (RS1080)

Clinical microbiologist, infectious disease specialist or respiratory specialist

ISONIAZID – **Restricted** see terms below

⬇ Tab 100 mg – **5% DV May-25 to 2027**	94.50	100	Isoniazid Teva
	327.41		**Noumed Isoniazid**

➥ **Restricted** (RS1281)

Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician

ISONIAZID WITH RIFAMPICIN – **Restricted** see terms below

⬇ Tab 100 mg with rifampicin 150 mg – **5% DV Feb-25 to 2027**	89.82	100	**Rifinah**
⬇ Tab 150 mg with rifampicin 300 mg – **5% DV Feb-25 to 2027**	179.13	100	**Rifinah**
⬇ Cap 100 mg with rifampicin 150 mg	199.00	100	Rifamazid

➥ **Restricted** (RS1282)

Clinical microbiologist, dermatologist, paediatrician, public health physician or internal medicine physician

PARA-AMINOSALICYLIC ACID – **Restricted** see terms on the next page

⬇ Grans for oral liq 4 g	280.00	30	Paser

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
➥ **Restricted** (RS1083)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PROTIONAMIDE – **Restricted** see terms below			
⬇ Tab 250 mg	305.00	100	Peteha
➥ **Restricted** (RS1084)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
PYRAZINAMIDE – **Restricted** see terms below			
⬇ Tab 500 mg			
➥ **Restricted** (RS1085)			
Clinical microbiologist, infectious disease specialist or respiratory specialist			
RIFABUTIN – **Restricted** see terms below			
⬇ Cap 150 mg	353.71	30	Mycobutin
➥ **Restricted** (RS1086)			
Clinical microbiologist, gastroenterologist, infectious disease specialist or respiratory specialist			
RIFAMPICIN – **Restricted** see terms below			
⬇ Cap 150 mg – **5% DV Dec-23 to 2026**	58.54	100	**Rifadin**
⬇ Cap 300 mg – **5% DV Dec-23 to 2026**	122.06	100	**Rifadin**
⬇ Oral liq 100 mg per 5 ml – **5% DV Dec-23 to 2026**	12.60	60 ml	**Rifadin**
⬇ Inj 600 mg vial – **5% DV Dec-23 to 2026**	134.98	1	**Rifadin**
➥ **Restricted** (RS1087)			
Clinical microbiologist, dermatologist, internal medicine physician, paediatrician or public health physician			

Antiparasitics

Anthelmintics

ALBENDAZOLE – **Restricted** see terms below			
⬇ Tab 200 mg			
⬇ Tab 400 mg			
➥ **Restricted** (RS1088)			
Clinical microbiologist or infectious disease specialist			
IVERMECTIN – **Restricted** see terms below			
⬇ Tab 3 mg	17.20	4	Stromectol
➥ **Restricted** (RS1283)			
Clinical microbiologist, dermatologist or infectious disease specialist			
MEBENDAZOLE			
Tab 100 mg – **5% DV Dec-24 to 2027**	5.18	6	**Vermox**
Oral liq 100 mg per 5 ml			
PRAZIQUANTEL			
Tab 600 mg			

Antiprotozoals

ARTEMETHER WITH LUMEFANTRINE – **Restricted** see terms below			
⬇ Tab 20 mg with lumefantrine 120 mg			
➥ **Restricted** (RS1090)			
Clinical microbiologist or infectious disease specialist			
ARTESUNATE – **Restricted** see terms on the next page			
⬇ Inj 60 mg vial			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
➥ **Restricted** (RS1091)			
Clinical microbiologist or infectious disease specialist			
ATOVAQUONE WITH PROGUANIL HYDROCHLORIDE – **Restricted** see terms below			
⬇ Tab 62.5 mg with proguanil hydrochloride 25 mg	29.50	12	Malarone Junior
⬇ Tab 250 mg with proguanil hydrochloride 100 mg	72.00	12	Malarone
➥ **Restricted** (RS1092)			
Clinical microbiologist or infectious disease specialist			
CHLOROQUINE PHOSPHATE – **Restricted** see terms below			
⬇ Tab 250 mg			
➥ **Restricted** (RS1093)			
Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist			
MEFLOQUINE – **Restricted** see terms below			
⬇ Tab 250 mg			
➥ **Restricted** (RS1094)			
Clinical microbiologist, dermatologist, infectious disease specialist or rheumatologist			
METRONIDAZOLE			
Tab 200 mg – **5% DV Mar-25 to 2026**	25.86	250	**Metronidamed**
Tab 400 mg – **5% DV Mar-25 to 2026**	4.29	21	**Metronidamed**
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	Flagyl-S
Inj 5 mg per ml, 100 ml bag – **5% DV Dec-23 to 2026**	18.00	10	**Baxter**
Suppos 500 mg	24.48	10	Flagyl
NITAZOXANIDE – **Restricted** see terms below			
⬇ Tab 500 mg			
⬇ Oral liq 100 mg per 5 ml			
➥ **Restricted** (RS1095)			
Clinical microbiologist or infectious disease specialist			
ORNIDAZOLE			
Tab 500 mg – **5% DV Mar-25 to 2027**	36.52	10	**Arrow-Ornidazole**
PENTAMIDINE ISETHIONATE – **Restricted** see terms below			
⬇ Inj 300 mg vial	216.00	5	Pentacarinat
	638.69		Tillomed
➥ **Restricted** (RS1096)			
Clinical microbiologist or infectious disease specialist			
PRIMAQUINE – **Restricted** see terms below			
⬇ Tab 15 mg			
⬇ Tab 7.5 mg			
➥ **Restricted** (RS1097)			
Clinical microbiologist or infectious disease specialist			
PYRIMETHAMINE – **Restricted** see terms below			
⬇ Tab 25 mg			
➥ **Restricted** (RS1098)			
Clinical microbiologist, infectious disease specialist or maternal-foetal medicine specialist			
QUININE DIHYDROCHLORIDE – **Restricted** see terms below			
⬇ Inj 60 mg per ml, 10 ml ampoule			
⬇ Inj 300 mg per ml, 2 ml vial			
➥ **Restricted** (RS1099)			
Clinical microbiologist or infectious disease specialist			
SODIUM STIBOGLUCONATE – **Restricted** see terms on the next page			
⬇ Inj 100 mg per ml, 1 ml vial			

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1100)
Clinical microbiologist or infectious disease specialist

SPIRAMYCIN – **Restricted** see terms below
⬇ Tab 500 mg
➥ **Restricted** (RS1101)
Maternal-foetal medicine specialist

Antiretrovirals

Non-Nucleoside Reverse Transcriptase Inhibitors

➥ **Restricted** (RS1898)
Initiation – Confirmed HIV
Patient has confirmed HIV infection.
Initiation – Prevention of maternal transmission
Either:
1 Prevention of maternal foetal transmission; or
2 Treatment of the newborn for up to eight weeks.
Initiation – Post-exposure prophylaxis following exposure to HIV
Both:
1 Treatment course to be initiated within 72 hours post exposure; and
2 Any of the following:
2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).
Initiation – Percutaneous exposure
Patient has percutaneous exposure to blood known to be HIV positive.

EFAVIRENZ – **Restricted** see terms above

➥ Tab 600 mg – **Restricted:** For continuation only	65.38	30	Efavirenz Milpharm
⬆ Oral liq 30 mg per ml			

(Efavirenz Milpharm Tab 600 mg to be delisted 1 November 2026)

ETRAVIRINE – **Restricted** see terms above

⬆ Tab 200 mg	770.00	60	Intelence

NEVIRAPINE – **Restricted** see terms above

⬆ Tab 200 mg – **5% DV Feb-25 to 2027**	198.25	60	**Nevirapine Viatris**
⬆ Oral suspension 10 mg per ml	203.55	240 ml	Viramune Suspension

Nucleoside Reverse Transcriptase Inhibitors

➥ **Restricted** (RS1899)
Initiation – Confirmed HIV
Patient has confirmed HIV infection.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – Prevention of maternal transmission

Either:

1 Prevention of maternal foetal transmission; or
2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following exposure to HIV

Both:

1 Treatment course to be initiated within 72 hours post exposure; and
2 Any of the following:
  2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

ABACAVIR SULPHATE – **Restricted** see terms on the previous page

⬆ Tab 300 mg	180.00	60	Ziagen
⬆ Oral liq 20 mg per ml			

ABACAVIR SULPHATE WITH LAMIVUDINE – **Restricted** see terms on the previous page

⬆ Tab 600 mg with lamivudine 300 mg – **5% DV Feb-26 to 2028**	35.00	30	**Abacavir/lamivudine Viatris**

EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPROXIL – **Restricted** see terms on the previous page

⬆ Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate)	106.88	30	Viatris
⬆ Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a fumarate)	106.88	30	TEEVIR

EMTRICITABINE – **Restricted** see terms on the previous page

⬆ Cap 200 mg	307.20	30	Emtriva

LAMIVUDINE – **Restricted** see terms on the previous page

⬆ Tab 150 mg – **5% DV Feb-24 to 2026**	98.00	60	**Lamivudine Viatris**
⬆ Oral liq 10 mg per ml			

STAVUDINE – **Restricted** see terms on the previous page

⬆ Cap 30 mg			
⬆ Cap 40 mg			
⬆ Powder for oral soln 1 mg per ml			

ZIDOVUDINE – **Restricted** see terms on the previous page

⬆ Cap 100 mg	152.25	100	Retrovir
⬆ Oral liq 10 mg per ml	30.45	200 ml	Retrovir
⬆ Inj 10 mg per ml, 20 ml vial	750.00	5	Retrovir IV

ZIDOVUDINE WITH LAMIVUDINE – **Restricted** see terms on the previous page

⬆ Tab 300 mg with lamivudine 150 mg	92.40	60	Lamivudine/Zidovudine Viatris

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Protease Inhibitors

➡ **Restricted** (RS1900)
Initiation – Confirmed HIV
Patient has confirmed HIV infection.
Initiation – Prevention of maternal transmission
Either:

1 Prevention of maternal foetal transmission; or
2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following exposure to HIV
Both:

1 Treatment course to be initiated within 72 hours post exposure; and
2 Any of the following:
  2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).
Initiation – Percutaneous exposure
Patient has percutaneous exposure to blood known to be HIV positive.

ATAZANAVIR SULPHATE – **Restricted** see terms above

⬆	Cap 150 mg – **5% DV Feb-26 to 2028**	102.50	60	**Atazanavir Viatris**
⬆	Cap 200 mg – **5% DV Feb-26 to 2028**	152.30	60	**Atazanavir Viatris**

DARUNAVIR – **Restricted** see terms above

⬆	Tab 400 mg – **5% DV Feb-24 to 2026**	150.00	60	**Darunavir Viatris**
⬆	Tab 600 mg – **5% DV Feb-24 to 2026**	225.00	60	**Darunavir Viatris**

INDINAVIR – **Restricted** see terms above

⬆	Cap 200 mg			
⬆	Cap 400 mg			

LOPINAVIR WITH RITONAVIR – **Restricted** see terms above

⬆	Tab 100 mg with ritonavir 25 mg			
⬆	Tab 200 mg with ritonavir 50 mg – **5% DV Feb-25 to 2027**	875.00	120	**Lopinavir/Ritonavir Mylan**
⬆	Oral liq 80 mg per ml with ritonavir 20 mg per ml			

RITONAVIR – **Restricted** see terms above

⬆	Tab 100 mg	43.31	30	Norvir

Strand Transfer Inhibitors

➡ **Restricted** (RS1901)
Initiation – Confirmed HIV
Patient has confirmed HIV infection.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – Prevention of maternal transmission

Either:

1 Prevention of maternal foetal transmission; or
2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following exposure to HIV

Both:

1 Treatment course to be initiated within 72 hours post exposure; and
2 Any of the following:
  2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
  2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
  2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

DOLUTEGRAVIR – **Restricted** see terms on the previous page			
⬆ Tab 50 mg	1,090.00	30	Tivicay
DOLUTEGRAVIR WITH LAMIVUDINE – **Restricted** see terms on the previous page			
⬆ Tab 50 mg with lamivudine 300 mg	1,090.00	30	Dovato
RALTEGRAVIR POTASSIUM – **Restricted** see terms on the previous page			
⬆ Tab 400 mg	1,090.00	60	Isentress
⬆ Tab 600 mg	1,090.00	60	Isentress HD

Antivirals

Hepatitis B

ENTECAVIR			
Tab 0.5 mg – **5% DV Mar-24 to 2026**	12.04	30	**Entecavir (Rex)**
LAMIVUDINE			
Tab 100 mg – **5% DV Feb-24 to 2026**	12.06	28	**Zetlam**
Oral liq 5 mg per ml	270.00	240 ml	Zeffix
TENOFOVIR DISOPROXIL			
Tab 245 mg (300 mg as a maleate) – **5% DV Dec-25 to 2028**	13.80	30	**Tenofovir Disoproxil Viatris**
Tab 245 mg (300 mg as a fumarate)	13.80	30	Ricovir

Hepatitis C

GLECAPREVIR WITH PIBRENTASVIR			
Note: the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://www.pharmac.govt.nz/maviret.			
Tab 100 mg with pibrentasvir 40 mg	24,750.00	84	Maviret
LEDIPASVIR WITH SOFOSBUVIR – **Restricted** see terms on the next page			
⬇ Tab 90 mg with sofosbuvir 400 mg	24,363.46	28	Harvoni

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1528)
Note: Only for use in patients with approval by the Hepatitis C Treatment Panel (HepCTP). Applications will be considered by HepCTP at its regular meetings and approved subject to eligibility according to the Access Criteria (set out in Section B of the Pharmaceutical Schedule).

Herpesviridae

ACICLOVIR

Tab dispersible 200 mg – **5% DV Feb-26 to 2028**	2.05	25	**Lovir**
Tab dispersible 400 mg – **5% DV Feb-26 to 2028**	7.55	56	**Lovir**
Tab dispersible 800 mg – **5% DV Feb-26 to 2028**	7.43	35	**Lovir**
Inj 250 mg vial – **5% DV Feb-25 to 2027**	13.75	5	Aciclovir Injection DBL **Aciclovir-Baxter**

CIDOFOVIR – **Restricted** see terms below
⬇ Inj 75 mg per ml, 5 ml vial
➥ **Restricted** (RS1108)
Clinical microbiologist, infectious disease specialist, otolaryngologist or oral surgeon

FOSCARNET SODIUM – **Restricted** see terms below
⬇ Inj 24 mg per ml, 250 ml bottle
➥ **Restricted** (RS1109)
Clinical microbiologist or infectious disease specialist

GANCICLOVIR – **Restricted** see terms below

⬇ Inj 500 mg vial	380.00	5	Cymevene

➥ **Restricted** (RS1110)
Clinical microbiologist or infectious disease specialist

LETERMOVIR – **Restricted** see terms below

⬇ Tab 240 mg	6,664.00	28	Prevymis

➥ **Restricted** (RS2189)
Initiation – CMV prophylaxis post HSCT
Limited to 4 months treatment
All of the following:
1 Patient has undergone an allogeneic haematopoietic stem cell transplant; and
2 The patient has confirmed presence of cytomegalovirus-specific antibodies; and
3 Treatment to commence within 28 days of an allogeneic haematopoietic stem cell transplant; and
4 Maximum treatment duration of 100 days post-transplant.

Continuation – CMV prophylaxis second or subsequent HSCT
Re-assessment required after 4 months
All of the following:
1 Patient has undergone an allogeneic haematopoietic stem cell transplant; and
2 The patient has confirmed presence of cytomegalovirus-specific antibodies; and
3 Treatment to commence within 28 days of an allogeneic haematopoietic stem cell transplant; and
4 Maximum treatment duration of 100 days post-transplant.

Initiation – CMV prophylaxis in severe immunosuppression*
Clinical microbiologist or infectious disease specialist
Re-assessment required after 6 months
Both:
1 Patient has severe immunosuppression requiring prophylaxis of CMV; and
2 Either:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.1 Patient is contraindicated to all other funded CMV active oral antiviral agents; or
2.2 Patients CMV is resistant to all other funded CMV active oral antiviral agents.

Continuation – CMV prophylaxis in severe immunosuppression*
Clinical microbiologist or infectious disease specialist
Re-assessment required after 6 months
Both:

1 Patient has severe immunosuppression requiring prophylaxis of CMV; and
2 Either:
2.1 Patient is contraindicated to all other funded CMV active oral antiviral agents; or
2.2 Patients CMV is resistant to all other funded CMV active oral antiviral agents.

Note: Indications marked with * are unapproved indications.

VALACICLOVIR

Tab 500 mg – **5% DV Feb-25 to 2027**	9.64	30	**Vaclovir**
Tab 1,000 mg – **5% DV Feb-25 to 2027**	17.78	30	**Vaclovir**
	12.45	21	Valaciclovir Mylan Valaciclovir Viatris

VALGANCICLOVIR – **Restricted** see terms below

⬇ Tab 450 mg – **5% DV Feb-25 to 2027**	140.89	60	**Valganciclovir Viatris**

➥ **Restricted** (RS2137)
Initiation – Transplant cytomegalovirus prophylaxis
Re-assessment required after 3 months
Patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.
Continuation – Transplant cytomegalovirus prophylaxis
Re-assessment required after 3 months
Either:

1 Both:
1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
2 Both:
2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initiation – Lung transplant cytomegalovirus prophylaxis
Re-assessment required after 12 months
All of the following:

1 Patient has undergone a lung transplant; and
2 Either:
2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
2.2 The recipient is cytomegalovirus positive; and
3 Patient has a high risk of CMV disease.

Continuation – Lung transplant cytomegalovirus prophylaxis
Re-assessment required after 12 months
All of the following:

1 Patient has undergone a lung re-transplant; and
2 Either:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
2.2 The recipient is cytomegalovirus positive; and

3 Patient has a high risk of CMV disease.

Initiation – Cytomegalovirus in immunocompromised patients

Both:

1 Patient is immunocompromised; and
2 Any of the following:
2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
2.3 Patient has cytomegalovirus retinitis.

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – **Restricted** see terms below

⬇ Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a maleate) – **5% DV Dec-25 to 2028**	13.45	30	**Tenofovir Disoproxil Emtricitabine Viatr**
⬇ Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a fumarate)	13.45	30	Tenofovir Disoproxil Emtricitabine Mylan

➥ **Restricted** (RS1902)

Initiation – Confirmed HIV

Patient has confirmed HIV infection.

Initiation – Prevention of maternal transmission

Either:

1 Prevention of maternal foetal transmission; or
2 Treatment of the newborn for up to eight weeks.

Initiation – Post-exposure prophylaxis following non-occupational exposure to HIV

Both:

1 Treatment course to be initiated within 72 hours post exposure; and
2 Any of the following:
2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initiation – Percutaneous exposure

Patient has percutaneous exposure to blood known to be HIV positive.

Initiation – Pre-exposure prophylaxis

Re-assessment required after 24 months

Both:

1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines (https://ashm.org.au/HIV/PrEP/)

Continuation – Pre-exposure prophylaxis

Re-assessment required after 24 months

Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines (https://ashm.org.au/HIV/PrEP/)

Influenza

OSELTAMIVIR – **Restricted** see terms below

Note: The restriction on the use of oseltamivir to hospitalised patients means that supply into the community for a new course is not permitted. Supply of a part original pack on discharge where initiated as a hospital inpatient is permitted.

⬇ Tab 75 mg
⬇ Powder for oral suspension 6 mg per ml
➥ **Restricted** (RS1307)

Initiation

Either:

1 Only for hospitalised patient with known or suspected influenza; or
2 For prophylaxis of influenza in hospitalised patients as part of a Health NZ Hospital approved infections control plan.

ZANAMIVIR

Note: The restriction on the use of zanamivir to hospitalised patients means that supply into the community for a new course is not permitted. Supply of a part original pack on discharge where initiated as a hospital inpatient is permitted.

⬇ Powder for inhalation 5 mg ..... 37.38 20 dose Relenza Rotadisk
➥ **Restricted** (RS1369)

Initiation

Either:

1 Only for hospitalised patient with known or suspected influenza; or
2 For prophylaxis of influenza in hospitalised patients as part of a Health NZ Hospital approved infections control plan.

COVID-19 Treatments

NIRMATRELVIR WITH RITONAVIR – **Restricted** see terms below

⬇ Tab 150 mg with ritonavir 100 mg ..... 1,274.00 30 Paxlovid
➥ **Restricted** (RS1894)

Initiation

Only if patient meets access criteria (as per https://pharmac.govt.nz/covid-oral-antivirals). Note the supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

REMDESIVIR

Note: For patients meeting access criteria for oral antiviral treatments (as on Pharmac's website).

Inj 100 mg vial ..... 615.23 1 Veklury

Immune Modulators

INTERFERON ALFA-2B

Inj 18 m iu, 1.2 ml multidose pen
Inj 30 m iu, 1.2 ml multidose pen
Inj 60 m iu, 1.2 ml multidose pen

INTERFERON GAMMA – **Restricted** see terms on the next page

⬇ Inj 100 mcg in 0.5 ml vial

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1113)

Initiation

Patient has chronic granulomatous disease and requires interferon gamma.

PEGYLATED INTERFERON ALFA-2A – **Restricted** see terms below

⬇	Inj 135 mcg prefilled syringe	887.35	1	Pegasys (S29)
⬇	Inj 180 mcg prefilled syringe	1,074.79	4	Pegasys
		1,355.71		Pegasys

➥ **Restricted** (RS1827)

Initiation – Chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant

Limited to 48 weeks treatment

Any of the following:

1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
2 Patient has chronic hepatitis C and is co-infected with HIV; or
3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant.

Notes: Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml.

Continuation – Chronic hepatitis C - genotype 1 infection

Gastroenterologist, infectious disease specialist or general physician

Re-assessment required after 48 weeks

All of the following:

1 Patient has chronic hepatitis C, genotype 1; and
2 Patient has had previous treatment with pegylated interferon and ribavirin; and
3 Either:
  3.1 Patient has responder relapsed; or
  3.2 Patient was a partial responder; and
4 Patient is to be treated in combination with boceprevir.

Initiation – Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

1 Patient has chronic hepatitis C, genotype 1; and
2 Patient has had previous treatment with pegylated interferon and ribavirin; and
3 Any of the following:
  3.1 Patient has responder relapsed; or
  3.2 Patient was a partial responder; or
  3.3 Patient received interferon treatment prior to 2004; and
4 Patient is to be treated in combination with boceprevir.

Initiation – Chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV

Limited to 6 months treatment

Patient has chronic hepatitis C, genotype 2 or 3 infection.

Initiation – Hepatitis B

Gastroenterologist, infectious disease specialist or general physician

Limited to 48 weeks treatment

All of the following:

1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 Patient is Hepatitis B treatment-naive; and
3 ALT > 2 times Upper Limit of Normal; and
4 HBV DNA < 10 log10 IU/ml; and
5 Either:
5.1 HBeAg positive; or
5.2 Serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (greater than or equal to Metavir Stage F2 or moderate fibrosis); and
6 Compensated liver disease; and
7 No continuing alcohol abuse or intravenous drug use; and
8 Not co-infected with HCV, HIV or HDV; and
9 Neither ALT nor AST > 10 times upper limit of normal; and
10 No history of hypersensitivity or contraindications to pegylated interferon.

Initiation – myeloproliferative disorder or cutaneous T cell lymphoma
Re-assessment required after 12 months
Any of the following:
1 Patient has a cutaneous T cell lymphoma*; or
2 All of the following:
2.1 Patient has a myeloproliferative disorder*; and
2.2 Patient is intolerant of hydroxyurea; and
2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
3 Both:
3.1 Patient has a myeloproliferative disorder; and
3.2 Patient is pregnant, planning pregnancy or lactating.

Continuation – myeloproliferative disorder or cutaneous T cell lymphoma
Re-assessment required after 12 months
All of the following:
1 No evidence of disease progression; and
2 The treatment remains appropriate and patient is benefitting from treatment; and
3 Either:
3.1 Patient has a cutaneous T cell lymphoma*; or
3.2 Both:
3.2.1 Patient has a myeloproliferative disorder*; and
3.2.2 Either:
3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications

Initiation – ocular surface squamous neoplasia
Ophthalmologist
Re-assessment required after 12 months
Patient has ocular surface squamous neoplasia*.

Continuation – ocular surface squamous neoplasia
Ophthalmologist
Re-assessment required after 12 months
The treatment remains appropriate and patient is benefitting from treatment.
Note: Indications marked with * are unapproved indications

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – post-allogenic bone marrow transplant

Re-assessment required after 3 months

Patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Continuation – post-allogenic bone marrow transplant

Re-assessment required after 3 months

Patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Anticholinesterases

EDROPHONIUM CHLORIDE – **Restricted** see terms below
⬇ Inj 10 mg per ml, 15 ml vial
⬇ Inj 10 mg per ml, 1 ml ampoule
➥ **Restricted** (RS1015)
Initiation
For the diagnosis of myasthenia gravis.

NEOSTIGMINE METILSULFATE
Inj 2.5 mg per ml, 1 ml ampoule – **5% DV Feb-25 to 2027** ..... 48.25 — 10 — **Max Health**

NEOSTIGMINE METILSULFATE WITH GLYCOPYRRONIUM BROMIDE
Inj 2.5 mg with glycopyrronium bromide 0.5 mg per ml, 1 ml ampoule – **5% DV Feb-26 to 2028** ..... 26.13 — 10 — **Max Health**

PYRIDOSTIGMINE BROMIDE
Tab 60 mg ..... 50.28 — 100 — Mestinon

Antirheumatoid Agents

HYDROXYCHLOROQUINE SULPHATE
Tab 200 mg – **5% DV May-25 to 2027** ..... 7.80 — 100 — **Ipca-Hydroxychloroquine**

LEFLUNOMIDE
Tab 10 mg – **5% DV Dec-23 to 2026** ..... 6.00 — 30 — **Arava**
Tab 20 mg – **5% DV Dec-23 to 2026** ..... 6.00 — 30 — **Arava**

PENICILLAMINE
Tab 125 mg ..... 67.23 — 100 — D-Penamine
Tab 250 mg ..... 110.12 — 100 — D-Penamine

SODIUM AUROTHIOMALATE
Inj 10 mg in 0.5 ml ampoule
Inj 20 mg in 0.5 ml ampoule
Inj 50 mg in 0.5 ml ampoule

Drugs Affecting Bone Metabolism

Bisphosphonates

ALENDRONATE SODIUM
Tab 70 mg – **5% DV Jul-24 to 2026** ..... 3.10 — 4 — **Fosamax**

ALENDRONATE SODIUM WITH COLECALCIFEROL
Tab 70 mg with colecalciferol 5,600 iu – **5% DV Jul-24 to 2026** ..... 1.99 — 4 — **Fosamax Plus**

PAMIDRONATE DISODIUM
Inj 3 mg per ml, 10 ml vial ..... 32.49 — 1 — Pamisol
Inj 6 mg per ml, 10 ml vial ..... 88.11 — 1 — Pamisol
Inj 9 mg per ml, 10 ml vial ..... 94.34 — 1 — Pamisol

RISEDRONATE SODIUM
Tab 35 mg – **5% DV Feb-26 to 2028** ..... 3.00 — 4 — **Risedronate Sandoz**

ZOLEDRONIC ACID
Inj 5 mg per 100 ml, bag – **5% DV Feb-26 to 2028** ..... 19.45 — 1 — **Zoledronic Acid Viatris**

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Other Drugs Affecting Bone Metabolism

DENOSUMAB – **Restricted** see terms below

Note: Denosumab inj 60 mg per 1 ml pre-filled syringe is Medsafe approved for use in osteoporosis. Denosumab inj 120 mg per 1.7 ml vial is Medsafe approved for use in hypercalcaemia of malignancy.

⬇	Inj 120 mg per 1.7 ml vial	375.00	1	Xgeva
⬇	Inj 60 mg per 1 ml prefilled syringe	187.50	1	Prolia

➥ **Restricted** (RS2097)

Initiation – Osteoporosis

All of the following:

1 The patient has established osteoporosis; and
2 Any of the following:
  2.1 History of one significant osteoporotic fracture demonstrated radiologically, with a documented T-Score less than or equal to -2.5, that incorporates BMD measured using dual-energy x-ray absorptiometry (DEXA); or
  2.2 History of one significant osteoporotic fracture, demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of logistical, technical or pathophysiological reasons; or
  2.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  2.4 Documented T-Score less than or equal to -3.0; or
  2.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm that incorporates BMD measured using DEXA; and
3 Any of the following:
  3.1 Bisphosphonates are contraindicated because the patient's creatinine clearance or eGFR is less than 35 mL/min; or
  3.2 The patient has experienced at least two symptomatic new fractures or a BMD loss greater than 2% per year, after at least 12 months' continuous therapy with a funded antiresorptive agent; or
  3.3 Bisphosphonates result in intolerable side effects; or
  3.4 Intravenous bisphosphonates cannot be administered due to logistical or technical reasons.

Initiation – Hypercalcaemia

Both:

1 Patient has hypercalcaemia of malignancy; and
2 Patient has severe renal impairment.

TERIPARATIDE – **Restricted** see terms below

⬇	Inj 250 mcg per ml, 2.4 ml	490.00	1	Forteo
		200.27		Teriparatide - Teva

➥ **Restricted** (RS1143)

Initiation

Limited to 18 months treatment

All of the following:

1 The patient has severe, established osteoporosis; and
2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
3 The patient has had two or more fractures due to minimal trauma; and
4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
b) Antiresorptive agents and their adequate doses for the purposes of this restriction are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Enzymes

HYALURONIDASE

Inj 1,500 iu ampoule

Hyperuricaemia and Antigout

ALLOPURINOL

Tab 100 mg – **5% DV Jun-24 to 2026**	17.99	1,000	**Ipca-Allopurinol**
Tab 300 mg – **5% DV Jun-24 to 2026**	22.50	500	**Ipca-Allopurinol**

BENZBROMARONE – **Restricted:** For continuation only

➥ Tab 50 mg			
➥ Tab 100 mg	45.00	100	Benzbromaron AL 100

COLCHICINE

Tab 500 mcg	6.30	100	Colgout

FEBUXOSTAT – **Restricted** see terms below

⬇ Tab 80 mg – **5% DV Jun-24 to 2026**	4.73	28	**Febuxostat (Teva)**
⬇ Tab 120 mg – **5% DV Jun-24 to 2026**	11.78	28	**Febuxostat (Teva)**

➥ **Restricted** (RS1844)

Initiation – Gout

Both:

1 Patient has been diagnosed with gout; and

2 Any of the following:

2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or

2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout..

Initiation – Tumour lysis syndrome

Haematologist or oncologist

Re-assessment required after 6 weeks

Both:

1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and

2 Patient has a documented history of allopurinol intolerance.

Continuation – Tumour lysis syndrome

Haematologist or oncologist

Re-assessment required after 6 weeks

The treatment remains appropriate and patient is benefitting from treatment.

PROBENECID

Tab 500 mg

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
RASBURICASE – **Restricted** see terms below			
⬇ Inj 1.5 mg vial			
➥ **Restricted** (RS1016)			
Haematologist			

Muscle Relaxants and Related Agents

ATRACURIUM BESYLATE			
Inj 10 mg per ml, 2.5 ml ampoule – **5% DV Jun-25 to 2026**	7.69	5	**Medsurge**
Inj 10 mg per ml, 5 ml ampoule – **5% DV Jun-25 to 2026**	9.86	5	**Medsurge**
BACLOFEN			
Tab 10 mg – **5% DV Dec-24 to 2027**	3.70	100	**Pacifen**
Oral liq 1 mg per ml			
Inj 0.05 mg per ml, 1 ml ampoule	11.55	1	Lioresal Intrathecal
Inj 2 mg per ml, 5 ml ampoule – **5% DV Mar-25 to 2027**	490.91	10	**Sintetica Baclofen Intrathecal**
CLOSTRIDIUM BOTULINUM TYPE A TOXIN			
Inj 100 u vial	467.50	1	Botox
Inj 300 u vial	388.50	1	Dysport
Inj 500 u vial	1,295.00	2	Dysport
DANTROLENE			
Cap 25 mg	145.77	100	Dantrium
Cap 50 mg	77.00	100	Dantrium
Inj 20 mg vial	1,143.74	6	Dantrium IV
MIVACURIUM CHLORIDE			
Inj 2 mg per ml, 10 ml ampoule			
ORPHENADRINE CITRATE			
Tab 100 mg – **5% DV Feb-25 to 2027**	23.25	100	**Norflex**
PANCURONIUM BROMIDE			
Inj 2 mg per ml, 2 ml ampoule			
ROCURONIUM BROMIDE			
Inj 10 mg per ml, 5 ml vial – **5% DV May-26 to 2028**	28.96	10	**Medsurge**
SUXAMETHONIUM CHLORIDE			
Inj 50 mg per ml, 2 ml ampoule – **5% DV Feb-24 to 2026**	35.40	10	**Martindale**
VECURONIUM BROMIDE			
Inj 10 mg vial – **5% DV Apr-25 to 2027**	380.00	10	**Vecure**

Reversers of Neuromuscular Blockade

SUGAMMADEX – **Restricted** see terms below			
⬇ Inj 100 mg per ml, 2 ml vial – **5% DV Dec-24 to 2027**	80.64	10	**Sugammadex BNM**
⬇ Inj 100 mg per ml, 5 ml vial – **5% DV Dec-24 to 2027**	201.60	10	**Sugammadex BNM**
➥ **Restricted** (RS1370)			

Initiation

Any of the following:

1 Patient requires reversal of profound neuromuscular blockade following rapid sequence induction that has been undertaken using rocuronium (i.e. suxamethonium is contraindicated or undesirable); or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 Severe neuromuscular degenerative disease where the use of neuromuscular blockade is required; or
3 Patient has an unexpectedly difficult airway that cannot be intubated and requires a rapid reversal of anaesthesia and neuromuscular blockade; or
4 The duration of the patient's surgery is unexpectedly short; or
5 Neostigmine or a neostigmine/anticholinergic combination is contraindicated (for example the patient has ischaemic heart disease, morbid obesity or COPD); or
6 Patient has a partial residual block after conventional reversal.

Non-Steroidal Anti-Inflammatory Drugs

CELECOXIB			
Cap 100 mg – **5% DV Feb-26 to 2028**	3.60	60	**Celebrex**
Cap 200 mg – **5% DV Jul-26 to 2028**	3.20	30	Celecoxib Pfizer
	2.55		**Celostea**
(Celecoxib Pfizer Cap 200 mg to be delisted 1 July 2026)			
DICLOFENAC SODIUM			
Tab EC 25 mg – **5% DV Feb-25 to 2027**	2.19	50	**Diclofenac Sandoz**
Tab 50 mg dispersible	1.50	20	Voltaren D
Tab EC 50 mg – **5% DV Feb-25 to 2027**	2.19	50	**Diclofenac Sandoz**
Tab long-acting 75 mg – **5% DV Aug-25 to 2028**	10.00	100	**Voltaren SR**
Inj 25 mg per ml, 3 ml ampoule	13.20	5	Voltaren
Suppos 12.5 mg	2.04	10	Voltaren
Suppos 25 mg	2.44	10	Voltaren
Suppos 50 mg	4.22	10	Voltaren
Suppos 100 mg	7.00	10	Voltaren

ETORICOXIB – **Restricted** see terms below

⬇ Tab 30 mg
⬇ Tab 60 mg
⬇ Tab 90 mg
⬇ Tab 120 mg

➥ **Restricted** (RS1592)

Initiation

For in-vivo investigation of allergy only.

IBUPROFEN			
Tab 200 mg - 1,000 tablet pack – **1% DV Feb-21 to 2026**	21.40	1,000	**Relieve**
➥ Tab 400 mg – **Restricted:** For continuation only			
➥ Tab 600 mg – **Restricted:** For continuation only			
Tab long-acting 800 mg – **5% DV Apr-25 to 2027**	3.65	30	**Ibuprofen SR BNM**
Oral liq 20 mg per ml – **5% DV Apr-25 to 2027**	2.85	200 ml	**Ethics**
Inj 5 mg per ml, 2 ml ampoule			
Inj 10 mg per ml, 2 ml vial			
INDOMETACIN [INDOMETHACIN]			
Cap 25 mg			
Cap 50 mg			
Cap long-acting 75 mg			
Inj 1 mg vial			
Suppos 100 mg			
KETOPROFEN			
Cap long-acting 200 mg	12.07	28	Oruvail SR
(Oruvail SR Cap long-acting 200 mg to be delisted 1 October 2026)			

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
MEFENAMIC ACID – **Restricted:** For continuation only			
➥ Cap 250 mg			
NAPROXEN			
Tab 250 mg – **5% DV Feb-25 to 2027**	3.92	50	Inza
	7.06	90	**Noflam 250**
	39.23	500	**Noflam 250**
Tab 500 mg – **5% DV Feb-25 to 2027**	6.89	50	Inza
	34.45	250	**Noflam 500**
Tab long-acting 750 mg – **5% DV Feb-25 to 2027**	10.40	28	**Naprosyn SR 750**
Tab long-acting 1 g – **5% DV Feb-25 to 2027**	11.50	28	**Naprosyn SR 1000**
PARECOXIB			
Inj 40 mg vial – **5% DV Dec-24 to 2027**	46.00	10	**Dynastat** Parecoxib Juno
SULINDAC			
Tab 100 mg			
Tab 200 mg			
TENOXICAM			
Tab 20 mg – **5% DV Feb-26 to 2028**	23.50	100	**Tilcotil**
Inj 20 mg vial	9.95	1	AFT

Topical Products for Joint and Muscular Pain

CAPSAICIN – **Restricted** see terms below			
⬇ Crm 0.025%	9.75	45 g	Zo-Rub Osteo Zostrix

➥ **Restricted** (RS1309)

Initiation

Patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE – **Restricted** see terms below

⬇ Tab 50 mg – **5% DV Feb-25 to 2027** ..... 117.00 56 **Rilutek**

➥ **Restricted** (RS1351)

Initiation

Neurologist or respiratory specialist

Re-assessment required after 6 months

All of the following:

1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
3 The patient has not undergone a tracheostomy; and
4 The patient has not experienced respiratory failure; and
5 Any of the following:
  5.1 The patient is ambulatory; or
  5.2 The patient is able to use upper limbs; or
  5.3 The patient is able to swallow.

Continuation

Re-assessment required after 18 months

All of the following:

1 The patient has not undergone a tracheostomy; and
2 The patient has not experienced respiratory failure; and
3 Any of the following:
  3.1 The patient is ambulatory; or
  3.2 The patient is able to use upper limbs; or
  3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg – **5% DV Feb-26 to 2028** ..... 126.02 112 **Motetis**

Anticholinergics

BENZATROPINE MESYLATE

Tab 2 mg ..... 10.99 60 Benztrop
Inj 1 mg per ml, 2 ml ampoule ..... 95.00 5 Phebra

PROCYCLIDINE HYDROCHLORIDE

Tab 5 mg

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE

Cap 100 mg ..... 38.24 60 Symmetrel

APOMORPHINE HYDROCHLORIDE

Inj 10 mg per ml, 2 ml ampoule ..... 59.50 5 Movapo
Inj 10 mg per ml, 5 ml ampoule ..... 121.84 5 Movapo

BROMOCRIPTINE

Cap 5 mg

ENTACAPONE

Tab 200 mg – **5% DV Jul-25 to 2027** ..... 13.73 100 **Entacapone Viatris**

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
LEVODOPA WITH BENSERAZIDE			
Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	Madopar Rapid
Cap 50 mg with benserazide 12.5 mg	13.75	100	Madopar 62.5
Cap 100 mg with benserazide 25 mg	15.80	100	Madopar 125
Cap long-acting 100 mg with benserazide 25 mg	22.85	100	Madopar HBS
Cap 200 mg with benserazide 50 mg	26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
Tab 100 mg with carbidopa 25 mg – **5% DV Feb-25 to 2027**	26.49	100	**Sinemet**
Tab long-acting 100 mg with carbipoda 25 mg			
Tab long-acting 200 mg with carbidopa 50 mg – **5% DV Feb-25 to 2027**	44.99	100	**Sinemet CR**
Tab 250 mg with carbidopa 25 mg – **5% DV Feb-25 to 2027**	39.49	100	**Sinemet**
LEVODOPA WITH CARBIDOPA AND ENTACAPONE			
Tab 50 mg with carbidopa 12.5 mg and entacapone 200 mg – **5% DV Jul-25 to 2027**	27.01	100	**Stalevo**
Tab 100 mg with carbidopa 25 mg and entacapone 200 mg – **5% DV Jul-25 to 2027**	34.18	100	**Stalevo**
Tab 150 mg with carbidopa 37.5 mg and entacapone 200 mg – **5% DV Jul-25 to 2027**	44.96	100	**Stalevo**
Tab 200 mg with carbidopa 50 mg and entacapone 200 mg – **5% DV Jul-25 to 2027**	51.23	100	**Stalevo**
PRAMIPEXOLE HYDROCHLORIDE			
Tab 0.25 mg – **5% DV Dec-25 to 2028**	5.23	100	**Ramipex**
Tab 1 mg – **5% DV Dec-25 to 2028**	17.73	100	**Ramipex**
RASAGILINE			
Tab 1 mg	53.50	30	Azilect
ROPINIROLE HYDROCHLORIDE			
Tab 0.25 mg	8.83	84	Ropin
Tab 1 mg	10.09	84	Ropin
Tab 2 mg	12.29	84	Ropin
Tab 5 mg	25.94	84	Ropin
SELEGILINE HYDROCHLORIDE – **Restricted:** For continuation only			
➥ Tab 5 mg			
TOLCAPONE			
Tab 100 mg	152.38	100	Tasmar

Anaesthetics

General Anaesthetics

DESFLURANE			
Soln for inhalation 100%, 240 ml bottle	1,350.00	6	Suprane
DEXMEDETOMIDINE			
Inj 100 mcg per ml, 2 ml vial – **5% DV May-24 to 2026**	42.00	5	**Dexmedetomidine Viatris**
ETOMIDATE			
Inj 2 mg per ml, 10 ml ampoule			
ISOFLURANE			
Soln for inhalation 100%, 250 ml bottle	2,730.00	6	Aerrane

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
KETAMINE			
Inj 1 mg per ml, 100 ml bag	146.00	5	Biomed
Inj 10 mg per ml, 10 ml syringe	76.00	5	Biomed
Inj 100 mg per ml, 2 ml vial	36.23	5	Ketalar
	91.98		Ketalar
METHOHEXITAL SODIUM			
Inj 10 mg per ml, 50 ml vial			
PROPOFOL			
Inj 10 mg per ml, 20 ml ampoule – **5% DV Feb-26 to 2028**	5.75	5	**Fresofol 1% MCT/LCT**
Inj 10 mg per ml, 50 ml vial – **5% DV Feb-26 to 2028**	27.50	10	**Fresofol 1% MCT/LCT**
Inj 10 mg per ml, 100 ml vial – **5% DV Feb-26 to 2028**	39.90	10	**Fresofol 1% MCT/LCT**
SEVOFLURANE			
Soln for inhalation 100%, 250 ml bottle	930.00	6	Baxter
THIOPENTAL [THIOPENTONE] SODIUM			
Inj 500 mg ampoule			

Local Anaesthetics

ARTICAINE HYDROCHLORIDE			
Inj 1%			
ARTICAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 4% with adrenaline 1:100,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:100,000, 1.8 ml dental cartridge			
Inj 4% with adrenaline 1:100,000, 2.2 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 1.7 ml dental cartridge			
Inj 4% with adrenaline 1:200,000 1.8 ml dental cartridge			
Inj 4% with adrenaline 1:200,000, 2.2 ml dental cartridge			
BENZOCAINE			
Gel 20%			
BENZOCAINE WITH TETRACAINE HYDROCHLORIDE			
Gel 18% with tetracaine hydrochloride 2%			*e.g. ZAP Topical Anaesthetic Gel*
BUPIVACAINE HYDROCHLORIDE			
Inj 5 mg per ml, 4 ml ampoule – **5% DV Feb-24 to 2026**	62.50	5	**Marcain Isobaric**
Inj 2.5 mg per ml, 20 ml ampoule			
Inj 2.5 mg per ml, 20 ml ampoule sterile pack – **5% DV Feb-24 to 2026**	28.00	5	**Marcain**
Inj 5 mg per ml, 10 ml ampoule sterile pack	16.20	5	Marcain
Inj 5 mg per ml, 20 ml ampoule			
Inj 5 mg per ml, 20 ml ampoule sterile pack	16.56	5	Marcain
Inj 1.25 mg per ml, 100 ml bag			
Inj 1.25 mg per ml, 200 ml bag			
Inj 2.5 mg per ml, 200 ml bag			
Inj 1.25 mg per ml, 500 ml bag			
BUPIVACAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 2.5 mg per ml with adrenaline 1:200,000, 10 ml ampoule			
Inj 2.5 mg per ml with adrenaline 1:400,000, 20 ml vial – **5% DV Aug-26 to 2029**	57.00	5	**Marcain with Adrenaline**
Inj 5 mg per ml with adrenaline 1:200,000, 20 ml vial – **5% DV Aug-26 to 2029**	61.68	5	**Marcain with Adrenaline**

⬆ Item restricted (see ➡ above); ⬇ Item restricted (see ➡ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
BUPIVACAINE HYDROCHLORIDE WITH FENTANYL			
Inj 0.625 mg with fentanyl 2 mcg per ml, 100 ml bag			
Inj 0.625 mg with fentanyl 2 mcg per ml, 200 ml bag	165.00	5	Biomed
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 100 ml bag – **5% DV Feb-26 to 2028**	135.00	5	**Bupafen**
Inj 1.25 mg with fentanyl 2 mcg per ml, 200 ml bag – **5% DV Feb-26 to 2028**	138.00	5	**Bupafen**
Inj 1.25 mg with fentanyl 2 mcg per ml, 50 ml syringe			
Inj 1.25 mg with fentanyl 2 mcg per ml, 20 ml syringe	57.35	5	Biomed
BUPIVACAINE HYDROCHLORIDE WITH GLUCOSE			
Inj 0.5% with glucose 8%, 4 ml ampoule – **5% DV Dec-25 to 2028**	21.40	5	**Marcain Heavy**
COCAINE HYDROCHLORIDE			
Paste 5%			
Soln 15%, 2 ml syringe			
Soln 4%, 2 ml syringe	30.77	1	Biomed
COCAINE HYDROCHLORIDE WITH ADRENALINE			
Paste 15% with adrenaline 0.06%			
Paste 25% with adrenaline 0.06%			
ETHYL CHLORIDE			
Spray 100%			
LIDOCAINE [LIGNOCAINE]			
Crm 4%	7.60	5 g	LMX4
	30.00	30 g	LMX4
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Gel 2%	4.87	20 g	Orion
Soln 4%			
Spray 10% – **5% DV Feb-26 to 2028**	82.90	50 ml	**Xylocaine**
Oral (gel) soln 2% – **5% DV Apr-26 to 2028**	30.80	200 ml	**Xylocaine Viscous**
Inj 1%, 20 ml ampoule, sterile pack			
Inj 2%, 20 ml ampoule, sterile pack			
Inj 1%, 5 ml ampoule	15.00	25	Lidocaine-Baxter
Inj 1%, 20 ml vial	19.50	5	Lidocaine-Baxter
Inj 2%, 5 ml ampoule	27.50	25	Lidocaine-Baxter
Inj 2%, 20 ml vial	14.00	5	Lidocaine-Baxter
Inj 10%, 5 ml ampoule			
Gel 2%, 11 ml urethral syringe – **5% DV Feb-26 to 2028**	65.45	10	**Instillagel Lido**
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE			
Inj 1% with adrenaline 1:100,000, 20 ml vial			
Inj 1% with adrenaline 1:100,000, 5 ml ampoule – **5% DV Dec-25 to 2028**	32.00	10	**Xylocaine**
Inj 1% with adrenaline 1:200,000, 20 ml vial	50.00	5	Xylocaine
Inj 2% with adrenaline 1:100,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.7 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:80,000, 2.2 ml dental cartridge			
Inj 2% with adrenaline 1:200,000, 20 ml vial	60.00	5	Xylocaine
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH ADRENALINE AND TETRACAINE HYDROCHLORIDE			
Soln 4% with adrenaline 0.1% and tetracaine hydrochloride 0.5%, 5 ml syringe	20.50	1	Topicaine

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE WITH PHENYLEPHRINE HYDROCHLORIDE			
Nasal spray 5% with phenylephrine hydrochloride 0.5%			
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE			
Crm 2.5% with prilocaine 2.5%	45.00	30 g	EMLA
Patch 25 mcg with prilocaine 25 mcg	115.00	20	EMLA
Crm 2.5% with prilocaine 2.5%, 5 g	45.00	5	EMLA
MEPIVACAINE HYDROCHLORIDE			
Inj 3%, 1.8 ml dental cartridge	43.60	50	Scandonest 3%
Inj 3%, 2.2 ml dental cartridge	43.60	50	Scandonest 3%
MEPIVACAINE HYDROCHLORIDE WITH ADRENALINE			
Inj 2% with adrenaline 1:100,000, 1.8 ml dental cartridge			
Inj 2% with adrenaline 1:100,000, 2.2 ml dental cartridge			
PRILOCAINE HYDROCHLORIDE			
Inj 0.5%, 50 ml vial	100.00	5	Citanest
Inj 2%, 5 ml ampoule			
PRILOCAINE HYDROCHLORIDE WITH FELYPRESSIN			
Inj 3% with felypressin 0.03 iu per ml, 1.8 ml dental cartridge			
Inj 3% with felypressin 0.03 iu per ml, 2.2 ml dental cartridge			
ROPIVACAINE HYDROCHLORIDE			
Inj 2 mg per ml, 10 ml ampoule – **5% DV Feb-24 to 2026**	9.80	5	**Ropivacaine Kabi**
Inj 2 mg per ml, 20 ml ampoule – **5% DV Feb-24 to 2026**	10.25	5	**Ropivacaine Kabi**
Inj 2 mg per ml, 100 ml bag – **5% DV Feb-24 to 2026**	32.85	5	**Ropivacaine Kabi**
Inj 2 mg per ml, 200 ml bag – **5% DV Feb-24 to 2026**	43.40	5	**Ropivacaine Kabi**
Inj 7.5 mg per ml, 10 ml ampoule – **5% DV Feb-24 to 2026**	11.00	5	**Ropivacaine Kabi**
Inj 7.5 mg per ml, 20 ml ampoule – **5% DV Feb-24 to 2026**	13.50	5	**Ropivacaine Kabi**
Inj 10 mg per ml, 10 ml ampoule – **5% DV Feb-24 to 2026**	11.75	5	**Ropivacaine Kabi**
Inj 10 mg per ml, 20 ml ampoule – **5% DV Feb-24 to 2026**	17.60	5	**Ropivacaine Kabi**
TETRACAINE [AMETHOCAINE] HYDROCHLORIDE			
Gel 4%			

Analgesics

Non-Opioid Analgesics

ASPIRIN			
Tab dispersible 300 mg – **5% DV May-24 to 2026**	5.65	100	**Ethics Aspirin**
CAPSAICIN – **Restricted** see terms below			
⬇ Crm 0.075%	11.95	45 g	Zo-Rub HP Zostrix HP

➥ **Restricted** (RS1145)

Initiation

For post-herpetic neuralgia or diabetic peripheral neuropathy.

METHOXYFLURANE – **Restricted** see terms below			
⬇ Soln for inhalation 99.9 mg per g, 3 ml bottle	276.00	10	Penthrox
⬇ Soln for inhalation 999.9 mg per g, 3 ml bottle with inhaler device and activated carbon chamber	54.00	1	Penthrox

➥ **Restricted** (RS1292)

Initiation

Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient is undergoing a painful procedure with an expected duration of less than one hour; and
2 Only to be used under supervision by a medical practitioner or nurse who is trained in the use of methoxyflurane.

NEFOPAM HYDROCHLORIDE

Tab 30 mg			

PARACETAMOL – **Some items restricted** see terms below

Tab soluble 500 mg			
Tab 500 mg - blister pack - 1,000 tablet pack – **1% DV Feb-22 to 2026**	19.75	1,000	**Pacimol**
Tab 500 mg - blister pack - 12 tablet pack			
Tab 500 mg - blister pack - 20 tablet pack			
Tab 500 mg - bottle pack – **1% DV Feb-22 to 2026**	17.92	1,000	**Noumed Paracetamol**
Oral liq 120 mg per 5 ml	3.98	200 ml	Paracetamol (Ethics)
Oral liq 250 mg per 5 ml – **5% DV Jul-26 to 2028**	3.18	200 ml	**Pamol**
⬇ Inj 10 mg per ml, 100 ml vial – **5% DV Feb-26 to 2028**	15.30	10	**Paracetamol Kabi**
Suppos 25 mg			
Suppos 50 mg			
Suppos 125 mg – **5% DV Feb-24 to 2026**	4.29	10	**Gacet**
Suppos 250 mg – **5% DV Feb-24 to 2026**	5.39	10	**Gacet**
Suppos 500 mg – **5% DV Feb-24 to 2026**	16.55	50	**Gacet**

➥ **Restricted** (RS1146)

Initiation

Intravenous paracetamol is only to be used where other routes are unavailable or impractical, or where there is reduced absorption. The need for IV paracetamol must be re-assessed every 24 hours.

SUCROSE

Oral liq 25%	14.61	25 ml	Biomed
⬇ Oral liq 66.7% (preservative free)			

➥ **Restricted** (RS1763)

Initiation

For use in neonatal patients only.

Opioid Analgesics

ALFENTANIL

Inj 0.5 mg per ml, 2 ml ampoule – **5% DV Feb-24 to 2026**	8.99	5	**Medsurge**

CODEINE PHOSPHATE

Tab 15 mg – **5% DV Dec-25 to 2028**	5.82	100	**Noumed**
Tab 30 mg – **5% DV Dec-25 to 2028**	6.88	100	**Noumed**
Tab 60 mg – **5% DV Dec-25 to 2028**	13.89	100	**Noumed**

DIHYDROCODEINE TARTRATE

Tab long-acting 60 mg – **5% DV Feb-26 to 2028**	9.20	60	**DHC Continus**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
FENTANYL			
Inj 10 mcg per ml, 10 ml syringe – **5% DV Feb-25 to 2027**	44.50	5	**Biomed Fentanyl**
Inj 50 mcg per ml, 2 ml ampoule – **5% DV May-25 to 2027**	4.25	10	**Boucher and Muir**
Inj 50 mcg per ml, 10 ml ampoule – **5% DV May-25 to 2027**	9.41	10	**Boucher and Muir**
Inj 10 mcg per ml, 100 ml bag – **5% DV Feb-24 to 2026**	114.25	5	**Biomed**
Inj 20 mcg per ml, 50 ml syringe – **5% DV Feb-25 to 2027**	136.50	5	**Biomed**
Inj 20 mcg per ml, 100 ml bag			
Patch 12 mcg per hour – **5% DV May-25 to 2027**	6.02	5	**Fentanyl Sandoz**
Patch 25 mcg per hour – **5% DV Dec-24 to 2027**	6.91	5	**Fentanyl Sandoz**
Patch 50 mcg per hour – **5% DV Dec-24 to 2027**	9.28	5	**Fentanyl Sandoz**
Patch 75 mcg per hour – **5% DV Dec-24 to 2027**	15.50	5	**Fentanyl Sandoz**
Patch 100 mcg per hour – **5% DV Dec-24 to 2027**	16.37	5	**Fentanyl Sandoz**
METHADONE HYDROCHLORIDE			
Tab 5 mg – **5% DV Sep-26 to 2028**	1.38	10	**Methadone BNM**
Oral liq 2 mg per ml – **5% DV Feb-25 to 2027**	7.80	200 ml	**Biodone**
Oral liq 5 mg per ml – **5% DV Feb-25 to 2027**	7.80	200 ml	**Biodone Forte**
Oral liq 10 mg per ml – **5% DV Feb-25 to 2027**	9.65	200 ml	**Biodone Extra Forte**
Inj 10 mg per ml, 1 ml vial	72.99	10	AFT
MORPHINE HYDROCHLORIDE			
Oral liq 1 mg per ml	19.00	200 ml	RA-Morph
Oral liq 2 mg per ml	23.55	200 ml	RA-Morph
Oral liq 5 mg per ml	28.20	200 ml	RA-Morph
Oral liq 10 mg per ml	40.25	200 ml	RA-Morph
MORPHINE SULPHATE			
Tab immediate-release 10 mg	2.80	10	Sevredol
Tab immediate-release 20 mg	5.52	10	Sevredol
Cap long-acting 10 mg – **5% DV Jul-26 to 2028**	4.10	10	**m-Eslon**
Cap long-acting 30 mg – **5% DV Jul-26 to 2028**	6.05	10	**m-Eslon**
Cap long-acting 60 mg – **5% DV Jul-26 to 2028**	12.10	10	**m-Eslon**
Cap long-acting 100 mg – **5% DV Jul-26 to 2028**	14.50	10	**m-Eslon**
Oral liq 2 mg per ml	42.56	300 ml	Oramorph
	29.80	100 ml	Oramorph CDC S29
	16.31		Wockhardt
Inj 1 mg per ml, 100 ml bag – **5% DV Feb-24 to 2026**	114.25	5	**Biomed**
Inj 1 mg per ml, 10 ml syringe – **5% DV Feb-24 to 2026**	27.25	5	**Biomed**
Inj 1 mg per ml, 50 ml syringe – **5% DV Feb-24 to 2026**	63.75	5	**Biomed**
Inj 1 mg per ml, 2 ml syringe			
Inj 5 mg per ml, 1 ml ampoule – **5% DV Feb-26 to 2028**	5.96	5	**Medsurge**
Inj 10 mg per ml, 1 ml ampoule – **5% DV Feb-26 to 2028**	4.99	5	**Medsurge**
Inj 10 mg per ml, 100 mg cassette			
Inj 10 mg per ml, 100 ml bag			
Inj 15 mg per ml, 1 ml ampoule – **5% DV Feb-26 to 2028**	6.93	5	**Medsurge**
Inj 30 mg per ml, 1 ml ampoule – **5% DV Feb-26 to 2028**	7.28	5	**Medsurge**
Inj 200 mcg in 0.4 ml syringe			
Inj 300 mcg in 0.3 ml syringe			
MORPHINE TARTRATE			
Inj 80 mg per ml, 1.5 ml ampoule			

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
OXYCODONE HYDROCHLORIDE			
Tab controlled-release 5 mg – **5% DV Dec-24 to 2027**	2.49	20	**Oxycodone Sandoz**
Tab immediate-release 5 mg	13.77	100	Oxycodone Amneal
Tab controlled-release 10 mg – **5% DV Dec-24 to 2027**	2.49	20	**Oxycodone Sandoz**
Tab immediate-release 10 mg	18.77	100	Oxycodone Amneal
Tab controlled-release 20 mg – **5% DV Dec-24 to 2027**	3.41	20	**Oxycodone Sandoz**
Tab immediate-release 20 mg	26.77	100	Oxycodone Amneal
Tab controlled-release 40 mg – **5% DV Dec-24 to 2027**	6.67	20	**Oxycodone Sandoz**
Tab controlled-release 80 mg – **5% DV Dec-24 to 2027**	12.99	20	**Oxycodone Sandoz**
Oral liq 1 mg per ml	37.08	250 ml	Oxycodone Lucis S29 Rosemont
Inj 1 mg per ml, 100 ml bag			
Inj 10 mg per ml, 1 ml ampoule – **5% DV Dec-24 to 2027**	4.37	5	**Hameln**
Inj 10 mg per ml, 2 ml ampoule – **5% DV Dec-24 to 2027**	8.62	5	**Hameln**
Inj 50 mg per ml, 1 ml ampoule – **5% DV Dec-24 to 2027**	14.90	5	**Hameln**
(Oxycodone Lucis S29 Oral liq 1 mg per ml to be delisted 1 June 2026)			
PARACETAMOL WITH CODEINE			
Tab paracetamol 500 mg with codeine phosphate 8 mg – **5% DV Feb-26 to 2028**	31.95	1,000	**Paracetamol + Codeine (Relieve)**
PETHIDINE HYDROCHLORIDE			
Tab 50 mg – **5% DV Feb-26 to 2028**	8.68	10	**Noumed Pethidine**
Inj 5 mg per ml, 10 ml syringe			
Inj 5 mg per ml, 100 ml bag			
Inj 10 mg per ml, 100 ml bag			
Inj 10 mg per ml, 50 ml syringe			
Inj 50 mg per ml, 1 ml ampoule	29.88	5	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule	30.72	5	DBL Pethidine Hydrochloride
REMIFENTANIL			
Inj 1 mg vial – **5% DV Feb-24 to 2026**	14.95	5	**Remifentanil-AFT**
Inj 2 mg vial – **5% DV Feb-24 to 2026**	20.95	5	**Remifentanil-AFT**
TRAMADOL HYDROCHLORIDE			
Tab sustained-release 100 mg – **5% DV May-24 to 2026**	1.95	20	**Tramal SR 100**
Tab sustained-release 150 mg – **5% DV May-24 to 2026**	2.95	20	**Tramal SR 150**
Tab sustained-release 200 mg – **5% DV May-24 to 2026**	3.80	20	**Tramal SR 200**
Cap 50 mg – **5% DV Jan-24 to 2026**	3.33	100	**Arrow-Tramadol**
Oral soln 10 mg per ml			
Inj 10 mg per ml, 100 ml bag			
Inj 50 mg per ml, 1 ml ampoule – **5% DV May-24 to 2026**	10.00	5	**Tramal 50**
Inj 50 mg per ml, 2 ml ampoule – **5% DV May-24 to 2026**	9.00	5	**Tramal 100**

Antidepressants

Cyclic and Related Agents

AMITRIPTYLINE			
Tab 10 mg – **5% DV Mar-24 to 2026**	2.99	100	**Arrow-Amitriptyline**
Tab 25 mg – **5% DV Mar-24 to 2026**	1.99	100	**Arrow-Amitriptyline**
Tab 50 mg – **5% DV Mar-24 to 2026**	3.14	100	**Arrow-Amitriptyline**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
CLOMIPRAMINE HYDROCHLORIDE			
Tab 25 mg – **5% DV Jul-25 to 2027**	16.99	50	**APO Clomipramine**
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – **Restricted:** For continuation only			
➥ Tab 75 mg	3.85	30	Dosulepin Viatris
➥ Cap 25 mg	7.83	50	Dosulepin Viatris
DOXEPIN HYDROCHLORIDE – **Restricted:** For continuation only			
➥ Cap 10 mg			
➥ Cap 25 mg			
➥ Cap 50 mg			
IMIPRAMINE HYDROCHLORIDE			
Tab 10 mg	5.48	50	Tofranil
	6.58	60	Tofranil
Tab 25 mg	4.93	28	Imipramine Crescent
	8.80	50	Tofranil
MAPROTILINE HYDROCHLORIDE – **Restricted:** For continuation only			
➥ Tab 25 mg			
➥ Tab 75 mg			
MIANSERIN HYDROCHLORIDE – **Restricted:** For continuation only			
➥ Tab 30 mg			
NORTRIPTYLINE HYDROCHLORIDE			
Tab 10 mg	2.24	50	Allegron
Tab 25 mg	2.95	50	Allegron

Monoamine-Oxidase Inhibitors - Non-Selective

PHENELZINE SULPHATE
Tab 15 mg

TRANYLCYPROMINE SULPHATE
Tab 10 mg

Monoamine-Oxidase Type A Inhibitors

MOCLOBEMIDE			
Tab 150 mg – **5% DV Feb-25 to 2027**	23.60	60	**Aurorix**
Tab 300 mg – **5% DV Feb-25 to 2027**	38.50	60	**Aurorix**

Other Antidepressants

MIRTAZAPINE			
Tab 30 mg – **5% DV Jan-26 to 2028**	2.34	30	**Noumed**
Tab 45 mg – **5% DV Jan-26 to 2028**	3.10	30	**Noumed**
VENLAFAXINE			
Cap 37.5 mg	8.29	84	Enlafax XR
Cap 75 mg	10.32	84	Enlafax XR
Cap 150 mg	13.95	84	Enlafax XR

Selective Serotonin Reuptake Inhibitors

CITALOPRAM HYDROBROMIDE			
Tab 20 mg – **5% DV May-26 to 2028**	3.55	84	**Celapram**

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
ESCITALOPRAM			
Tab 10 mg – **5% DV Apr-24 to 2026**	0.79	28	**Ipca-Escitalopram**
Tab 20 mg – **5% DV Apr-24 to 2026**	1.49	28	**Ipca-Escitalopram**
FLUOXETINE HYDROCHLORIDE			
Tab dispersible 20 mg, scored – **5% DV Mar-26 to 2028**	2.37	28	**Fluox**
Cap 20 mg – **5% DV Mar-26 to 2028**	3.50	90	**Arrow-Fluoxetine**
PAROXETINE			
Tab 20 mg – **5% DV Feb-26 to 2028**	4.98	90	**Loxamine**
SERTRALINE			
Tab 50 mg – **5% DV Apr-26 to 2028**	1.24	30	**Setrona**
Tab 100 mg – **5% DV Apr-26 to 2028**	2.00	30	**Setrona**

Antiepilepsy Drugs

Agents for the Control of Status Epilepticus

CLONAZEPAM			
Inj 1 mg per ml, 1 ml ampoule			
DIAZEPAM			
Inj 5 mg per ml, 2 ml ampoule	27.92	5	Hospira
Rectal tubes 5 mg	54.58	5	Stesolid
Rectal tubes 10 mg			
LORAZEPAM			
Inj 2 mg vial			
Inj 4 mg per ml, 1 ml vial			
PARALDEHYDE			
Soln 97%			
Inj 5 ml ampoule			
PHENYTOIN SODIUM			
Inj 50 mg per ml, 5 ml ampoule	154.01	5	Hospira

Control of Epilepsy

CARBAMAZEPINE			
Tab 200 mg	14.53	100	Tegretol
Tab long-acting 200 mg	16.98	100	Tegretol CR
Tab 400 mg	34.58	100	Tegretol
Tab long-acting 400 mg	39.17	100	Tegretol CR
Oral liq 20 mg per ml	26.37	250 ml	Tegretol
CLOBAZAM			
Tab 10 mg			
CLONAZEPAM			
Oral drops 2.5 mg per ml			
ETHOSUXIMIDE			
Cap 250 mg	140.88	100	Zarontin
Oral liq 50 mg per ml	56.35	200 ml	Zarontin

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
GABAPENTIN			
Note: Gabapentin not to be given in combination with pregabalin			
Cap 100 mg – **1% DV Feb-22 to 2027**	6.45	100	**Nupentin**
Cap 300 mg – **1% DV Feb-22 to 2027**	8.45	100	**Nupentin**
Cap 400 mg – **1% DV Feb-22 to 2027**	10.26	100	**Nupentin**
LACOSAMIDE – **Restricted** see terms below			
⬇ Tab 50 mg	25.04	14	Vimpat
⬇ Tab 100 mg	50.06	14	Vimpat
	200.24	56	Vimpat
⬇ Tab 150 mg	75.10	14	Vimpat
	300.40	56	Vimpat
⬇ Tab 200 mg	400.55	56	Vimpat
⬇ Inj 10 mg per ml, 20 ml vial			

➥ **Restricted** (RS1988)

Initiation

Re-assessment required after 15 months

Both:

1 Patient has focal epilepsy; and

2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam, and any two of carbamazepine, lamotrigine, and phenytoin sodium (see Note).

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, or topiramate. Those who can father children are not required to trial sodium valproate.

Continuation

Patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

LAMOTRIGINE			
Tab dispersible 2 mg	55.00	30	Lamictal
Tab dispersible 5 mg	50.00	30	Lamictal
Tab dispersible 25 mg	4.20	56	Logem
Tab dispersible 50 mg	5.11	56	Logem
Tab dispersible 100 mg	6.75	56	Logem
LEVETIRACETAM			
Tab 250 mg	5.84	60	Everet
Tab 500 mg	10.51	60	Everet
Tab 750 mg	16.71	60	Everet
Tab 1,000 mg	21.82	60	Everet
Oral liq 100 mg per ml	44.78	300 ml	Levetiracetam-AFT
Inj 100 mg per ml, 5 ml vial	38.95	10	Levetiracetam-AFT
PHENOBARBITONE			
Tab 15 mg	248.50	500	Noumed Phenobarbitone
Tab 30 mg	398.50	500	Noumed Phenobarbitone
PHENYTOIN			
Tab 50 mg			
PHENYTOIN SODIUM			
Cap 30 mg			
Cap 100 mg			
Oral liq 6 mg per ml			

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
PREGABALIN			
Note: Pregabalin not to be given in combination with gabapentin			
Cap 25 mg	2.25	56	Lyrica
			Pregabalin Pfizer
Cap 75 mg	2.65	56	Lyrica
			Pregabalin Pfizer
Cap 150 mg	4.01	56	Lyrica
			Pregabalin Pfizer
Cap 300 mg	7.38	56	Lyrica
			Pregabalin Pfizer
PRIMIDONE			
Tab 250 mg			
SODIUM VALPROATE			
Tab 100 mg			
Tab EC 200 mg			
Tab EC 500 mg			
Oral liq 40 mg per ml			
Inj 100 mg per ml, 4 ml vial	9.98	1	Epilim IV
STIRIPENTOL – **Restricted** see terms below			
⬇ Cap 250 mg	509.29	60	Diacomit
⬇ Powder for oral liq 250 mg sachet	509.29	60	Diacomit

➥ **Restricted** (RS1989)

Initiation

Paediatric neurologist

Re-assessment required after 6 months

Both:

1 Patient has confirmed diagnosis of Dravet syndrome; and

2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Note: Those of childbearing potential are not required to trial sodium valproate or topiramate. Those who can father children are not required to trial sodium valproate.

Continuation

Paediatric neurologist

Patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE			
Tab 25 mg	11.07	60	Arrow-Topiramate
	26.04		Topamax
	11.07		Topiramate Actavis
Tab 50 mg	18.81	60	Arrow-Topiramate
	44.26		Topamax
	18.81		Topiramate Actavis
Tab 100 mg	31.99	60	Arrow-Topiramate
	75.25		Topamax
	31.99		Topiramate Actavis
Tab 200 mg	55.19	60	Arrow-Topiramate
	129.85		Topamax
	55.19		Topiramate Actavis
Cap sprinkle 15 mg	20.84	60	Topamax
Cap sprinkle 25 mg	26.04	60	Topamax
VIGABATRIN – **Restricted** see terms on the next page			
⬇ Tab 500 mg			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1865)

Initiation

Re-assessment required after 15 months

Both:

1 Any of the following:
- 1.1 Patient has infantile spasms; or
- 1.2 Both:
  - 1.2.1 Patient has epilepsy; and
  - 1.2.2 Either:
    - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
    - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and

2 Either:
- 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
- 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Continuation

Both:

1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and

2 Either:
- 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
- 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields.

Antimigraine Preparations

Acute Migraine Treatment

DIHYDROERGOTAMINE MESYLATE

Inj 1 mg per ml, 1 ml ampoule			

METOCLOPRAMIDE HYDROCHLORIDE WITH PARACETAMOL

Tab 5 mg with paracetamol 500 mg			

RIZATRIPTAN

Tab orodispersible 10 mg – **5% DV Feb-24 to 2026**	4.84	30	**Rizamelt**

SUMATRIPTAN

Tab 50 mg – **1% DV Feb-22 to 2027**	14.41	90	**Sumagran**
Tab 100 mg – **1% DV Feb-22 to 2027**	22.68	90	**Sumagran**
Inj 12 mg per ml, 0.5 ml prefilled pen – **5% DV Dec-25 to 2028**	29.80	2	**Clustran**

Prophylaxis of Migraine

PIZOTIFEN

Tab 500 mcg	23.21	100	Sandomigran

Antinausea and Vertigo Agents

APREPITANT – **Restricted** see terms on the next page

⬇ Cap 2 × 80 mg and 1 × 125 mg – **5% DV Jan-25 to 2027**	21.90	3	**Emend Tri-Pack**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
➥ **Restricted** (RS1154)			
Initiation			
Patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.			
BETAHISTINE DIHYDROCHLORIDE			
Tab 16 mg – **5% DV Dec-23 to 2026**	3.70	100	**Serc**
CYCLIZINE HYDROCHLORIDE			
Tab 50 mg – **5% DV Feb-25 to 2027**	0.66	10	**Nausicalm**
CYCLIZINE LACTATE			
Inj 50 mg per ml, 1 ml ampoule	16.36	10	Hameln
DOMPERIDONE			
Tab 10 mg – **5% DV Dec-25 to 2028**	3.80	100	**Domperidone Viatris**
DROPERIDOL			
Inj 2.5 mg per ml, 1 ml ampoule – **5% DV Feb-26 to 2028**	28.68	10	**Droperidol Medsurge**
GRANISETRON			
Inj 1 mg per ml, 3 ml ampoule – **5% DV Feb-24 to 2026**	1.20	1	**Deva**
HYOSCINE HYDROBROMIDE			
Inj 400 mcg per ml, 1 ml ampoule			
⬇ Patch 1 mg per 72 hours	88.50	10	Scopolamine Transdermal System Viatris
➥ **Restricted** (RS1155)			
Initiation			
Any of the following:			
1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or			
2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective; or			
3 For treatment of post-operative nausea and vomiting where cyclizine, droperidol and a 5HT3 antagonist have proven ineffective, are not tolerated or are contraindicated.			
METOCLOPRAMIDE HYDROCHLORIDE			
Tab 10 mg – **5% DV Mar-24 to 2026**	1.57	100	**Metoclopramide Actavis 10**
Oral liq 5 mg per 5 ml			
Inj 5 mg per ml, 2 ml ampoule – **5% DV Apr-26 to 2028**	5.48	10	**Medsurge**
ONDANSETRON			
Tab 4 mg – **5% DV Dec-25 to 2028**	1.95	50	**Periset**
Tab dispersible 4 mg – **5% DV Mar-24 to 2026**	0.56	10	**Periset ODT**
Tab 8 mg – **5% DV Dec-25 to 2028**	3.50	50	**Periset**
Tab dispersible 8 mg – **5% DV Mar-24 to 2026**	0.90	10	**Periset ODT**
Inj 2 mg per ml, 2 ml ampoule – **5% DV Feb-26 to 2028**	1.54	5	**Ondansetron-AFT**
Inj 2 mg per ml, 4 ml ampoule – **5% DV Feb-26 to 2028**	2.14	5	**Ondansetron-AFT**
PROCHLORPERAZINE			
Tab buccal 3 mg			
Tab 5 mg – **5% DV Mar-24 to 2026**	25.00	250	**Nausafix**
Inj 12.5 mg per ml, 1 ml ampoule			
Suppos 25 mg			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
TROPISETRON			
Inj 1 mg per ml, 2 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			

Antipsychotic Agents

General

AMISULPRIDE			
Tab 100 mg – **5% DV Dec-24 to 2027**	5.84	30	**Sulprix**
Tab 200 mg – **5% DV Dec-24 to 2027**	14.47	60	**Sulprix**
Tab 400 mg – **5% DV Dec-24 to 2027**	35.06	60	**Sulprix**
Oral liq 100 mg per ml			
ARIPIPRAZOLE			
Tab 5 mg	10.50	30	Aripiprazole Sandoz
Tab 10 mg	10.50	30	Aripiprazole Sandoz
Tab 15 mg	10.50	30	Aripiprazole Sandoz
Tab 20 mg	10.50	30	Aripiprazole Sandoz
Tab 30 mg	10.50	30	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE			
Tab 25 mg	15.62	100	Largactil
Tab 100 mg	36.73	100	Largactil
Oral liq 10 mg per ml			
Oral liq 20 mg per ml			
Inj 25 mg per ml, 2 ml ampoule	30.79	10	Largactil
CLOZAPINE			
Tab 25 mg	6.69	50	Clopine
	13.37	100	Clopine
	6.69	50	Clozaril
	13.37	100	Clozaril
Tab 50 mg	8.67	50	Clopine
	17.33	100	Clopine
Tab 100 mg	17.33	50	Clopine
	34.65	100	Clopine
	17.33	50	Clozaril
	34.65	100	Clozaril
Tab 200 mg	34.65	50	Clopine
	69.30	100	Clopine
Oral liq 50 mg per ml	173.30	100 ml	Versacloz
HALOPERIDOL			
Tab 500 mcg	6.23	100	Serenace
Tab 1.5 mg	9.43	100	Serenace
Tab 5 mg	29.72	100	Serenace
Oral liq 2 mg per ml	23.84	100 ml	Serenace
Inj 5 mg per ml, 1 ml ampoule – **5% DV Aug-26 to 2029**	12.93	10	**Serenace**
LEVOMEPROMAZINE			
Tab 25 mg	16.10	100	Nozinan
Tab 100 mg	41.75	100	Nozinan
LEVOMEPROMAZINE HYDROCHLORIDE			
Inj 25 mg per ml, 1 ml ampoule – **5% DV Dec-25 to 2028**	23.26	10	**Wockhardt**

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
LITHIUM CARBONATE			
Tab long-acting 400 mg – **5% DV Feb-25 to 2027**	82.80	100	**Priadel**
Cap 250 mg	35.78	100	Douglas
OLANZAPINE			
Tab 2.5 mg – **5% DV Aug-24 to 2026**	1.40	30	**Zypine**
Tab 5 mg – **5% DV Aug-24 to 2026**	1.93	30	**Zypine**
Tab orodispersible 5 mg – **5% DV Feb-24 to 2026**	3.93	28	APO-Olanzapine
	8.64	100	Olanzapine ODT Viatris
	2.42	28	**Zypine ODT**
Tab 10 mg – **5% DV Aug-24 to 2026**	1.80	28	**Zypine**
	1.93	30	**Zypine**
Tab orodispersible 10 mg – **5% DV Feb-24 to 2026**	4.49	28	APO-Olanzapine
	2.89		Olanzapina Mylan
			Olanzapina Mylan Pharma
	10.32	100	Olanzapine ODT Viatris
	2.89	28	**Zypine ODT**
Inj 10 mg vial			
PERICYAZINE			
Tab 2.5 mg			
Tab 10 mg			
QUETIAPINE			
Tab 25 mg – **5% DV Feb-24 to 2026**	2.36	90	**Quetapel**
	0.79	30	Quetiapine Viatris
	13.11	500	Quetiapine Viatris
Tab 100 mg – **5% DV Feb-24 to 2026**	6.40	90	**Quetapel**
Tab 200 mg – **5% DV Feb-24 to 2026**	10.97	90	**Quetapel**
Tab 300 mg – **5% DV Feb-24 to 2026**	15.83	90	**Quetapel**
RISPERIDONE			
Tab 0.5 mg – **5% DV Mar-24 to 2026**	2.17	60	**Risperidone (Teva)**
Tab 1 mg – **5% DV Mar-24 to 2026**	2.44	60	**Risperidone (Teva)**
Tab 2 mg – **5% DV Mar-24 to 2026**	2.72	60	**Risperidone (Teva)**
Tab 3 mg – **5% DV Mar-24 to 2026**	4.50	60	**Risperidone (Teva)**
Tab 4 mg – **5% DV Mar-24 to 2026**	6.25	60	Risperdal
			Risperidone (Teva)
Oral liq 1 mg per ml – **5% DV Mar-24 to 2026**	10.29	30 ml	**Risperon**
	34.30	100 ml	**Risperon**
ZIPRASIDONE			
Cap 20 mg	17.90	60	Zusdone
Cap 40 mg	27.41	60	Zusdone
Cap 60 mg	38.39	60	Zusdone
Cap 80 mg	46.55	60	Zusdone
ZUCLOPENTHIXOL ACETATE			
Inj 50 mg per ml, 1 ml ampoule			
Inj 50 mg per ml, 2 ml ampoule			
ZUCLOPENTHIXOL HYDROCHLORIDE			
Tab 10 mg	31.45	100	Clopixol

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Depot Injections

ARIPIPRAZOLE – **Restricted** see terms below

⬇ Inj 300 mg vial	273.56	1	Abilify Maintena
⬇ Inj 400 mg vial	341.96	1	Abilify Maintena

➥ **Restricted** (RS2058)

Initiation

Either:

1 Either:

1.1 The patient has had an initial Special Authority approval for risperidone depot injection or paliperidone depot injection or olanzapine depot injection; or

1.2 All of the following:

1.2.1 The patient has schizophrenia or other psychotic disorder; and

1.2.2 The patient has received treatment with oral atypical antipsychotic agents but has been unable to adhere; and

1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months; or

2 Patient has been unable to access olanzapine depot injection due to supply issues with olanzapine depot injection, or otherwise would have been initiated on olanzapine depot injection but has been unable to due to supply issues with olanzapine depot injection. (see Note below for the olanzapine Special Authority criteria for new olanzapine depot injection patients prior to 1 April 2024).

Notes: The Olanzapine depot injection Special Authority criteria that apply to criterion 2 in this Aripiprazole Special Authority application are as follows:

- The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- All of the following:
  - The patient has schizophrenia; and
  - The patient has tried but has not been able to adhere with treatment using oral atypical antipsychotic agents; and
  - The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

FLUPENTHIXOL DECANOATE

Inj 20 mg per ml, 1 ml ampoule	13.14	5	Fluanxol
Inj 20 mg per ml, 2 ml ampoule	20.90	5	Fluanxol
Inj 100 mg per ml, 1 ml ampoule	40.87	5	Fluanxol

HALOPERIDOL DECANOATE

Inj 50 mg per ml, 1 ml ampoule	28.39	5	Haldol
Inj 100 mg per ml, 1 ml ampoule	55.90	5	Haldol Concentrate

OLANZAPINE – **Restricted** see terms below

⬇ Inj 210 mg vial	252.00	1	Zyprexa Relprevv
⬇ Inj 300 mg vial	414.00	1	Zyprexa Relprevv
⬇ Inj 405 mg vial	504.00	1	Zyprexa Relprevv

➥ **Restricted** (RS2165)

Initiation

Re-assessment required after 12 months

All of the following:

1 Either:

1.1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.2 All of the following:
1.2.1 The patient has schizophrenia or other psychotic disorder; and
1.2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
1.2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months; and

2 The patient has trialled other funded depot antipsychotics (aripiprazole, risperidone, and paliperidone) unless it is considered clinically inappropriate to use these; and
3 The patient continues to have difficulties with adherence on oral antipsychotic treatments; and
4 Prescribing clinician has relevant Clinical Director (Mental Health and Addiction services) approval.

Continuation

Re-assessment required after 12 months

The initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE – **Restricted** see terms below

⬇	Inj 25 mg syringe	194.25	1	Invega Sustenna
⬇	Inj 50 mg syringe	271.95	1	Invega Sustenna
⬇	Inj 75 mg syringe	357.42	1	Invega Sustenna
⬇	Inj 100 mg syringe	435.12	1	Invega Sustenna
⬇	Inj 150 mg syringe	435.12	1	Invega Sustenna

➥ **Restricted** (RS2059)

Initiation

Re-assessment required after 12 months

Either:

1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or
2 All of the following:
2.1 The patient has schizophrenia or other psychotic disorder; and
2.2 The patient has been unable to adhere to treatment using oral atypical antipsychotic agents; and
2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE PALMITATE – **Restricted** see terms below

⬇	Inj 175 mg syringe	815.85	1	Invega Trinza
⬇	Inj 263 mg syringe	1,072.26	1	Invega Trinza
⬇	Inj 350 mg syringe	1,305.36	1	Invega Trinza
⬇	Inj 525 mg syringe	1,305.36	1	Invega Trinza

➥ **Restricted** (RS1932)

Initiation

Re-assessment required after 12 months

Both:

1 The patient has schizophrenia; and
2 The patient has had an initial Special Authority approval for paliperidone once-monthly depot injection.

Continuation

Re-assessment required after 12 months

The initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
PIPOTHIAZINE PALMITATE – **Restricted:** For continuation only			
➥ Inj 50 mg per ml, 1 ml ampoule			
➥ Inj 50 mg per ml, 2 ml ampoule			
RISPERIDONE – **Restricted** see terms below			
⬇ Inj 25 mg vial	135.98	1	Risperdal Consta
⬇ Inj 37.5 mg vial	178.71	1	Risperdal Consta
⬇ Inj 50 mg vial	217.56	1	Risperdal Consta

➥ **Restricted** (RS2060)

Initiation

Re-assessment required after 12 months

Either:

1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection or aripiprazole depot injection; or

2 All of the following:

2.1 The patient has schizophrenia or other psychotic disorder; and

2.2 The patient has not been able to adhere to treatment using oral atypical antipsychotic agents; and

2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Continuation

Re-assessment required after 12 months

The initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE			
Inj 200 mg per ml, 1 ml ampoule	19.80	5	Clopixol
Inj 500 mg per ml, 1 ml ampoule			*e.g. Clopixol Conc*

Anxiolytics

BUSPIRONE HYDROCHLORIDE			
Tab 5 mg – **5% DV Dec-24 to 2027**	13.95	100	**Buspirone Viatris**
Tab 10 mg – **5% DV Dec-24 to 2027**	12.50	100	**Buspirone Viatris**
CLONAZEPAM			
Tab 500 mcg	5.64	100	Paxam
Tab 2 mg	10.78	100	Paxam
DIAZEPAM			
Tab 2 mg – **5% DV Mar-24 to 2026**	95.00	500	**Arrow-Diazepam**
Tab 5 mg – **5% DV Mar-24 to 2026**	115.00	500	**Arrow-Diazepam**
⬇ Oral liq 10 mg per 10 ml			

➥ **Restricted** (RS2054)

Initiation

Relevant specialist

Only for use in children where diazepam tablets are not appropriate.

LORAZEPAM			
Tab 1 mg – **5% DV Feb-25 to 2027**	10.20	250	**Ativan**
Tab 2.5 mg – **5% DV Feb-25 to 2027**	13.13	100	**Ativan**
OXAZEPAM			
Tab 10 mg			
Tab 15 mg			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Multiple Sclerosis Treatments

➡ **Restricted** (RS1993)

Initiation – Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide

Any relevant practitioner

Re-assessment required after 12 months

Either:

1 All of the following:
- 1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 1.2 Patient has an EDSS score between 0 – 6.0; and
- 1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 1.4 All of the following:
  - 1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
  - 1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
  - 1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
  - 1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
  - 1.4.5 Either:
    - 1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
    - 1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 1.6 Any of the following:
  - 1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
  - 1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
  - 1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
  - 1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
  - 1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or

2 Patient has an active approval for ocrelizumab and does not have primary progressive MS.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Continuation – Multiple Sclerosis - dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab and teriflunomide

Any relevant practitioner

Patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE – **Restricted** see terms above

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

⬆	Cap 120 mg	520.00	14	Tecfidera
⬆	Cap 240 mg	2,000.00	56	Tecfidera

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
FINGOLIMOD – **Restricted** see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
⬆ Cap 0.5 mg	2,200.00	28	Gilenya
GLATIRAMER ACETATE – **Restricted** see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
⬆ Inj 40 mg prefilled syringe	1,500.00	12	Copaxone
INTERFERON BETA-1-ALPHA – **Restricted** see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
⬆ Inj 6 million iu in 0.5 ml syringe	1,170.00	4	Avonex
INTERFERON BETA-1-BETA – **Restricted** see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
⬆ Inj 8 million iu per ml, 1 ml vial			
NATALIZUMAB – **Restricted** see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
⬆ Inj 20 mg per ml, 15 ml vial	1,750.00	1	Tysabri
TERIFLUNOMIDE – **Restricted** see terms on the previous page			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
⬆ Tab 14 mg – **5% DV Apr-25 to 2026**	263.96	28	**Teriflunomide Sandoz**

Multiple Sclerosis Treatments - Other

OCRELIZUMAB – **Restricted** see terms below			
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.			
⬇ Inj 30 mg per ml, 10 ml vial	8,450.00	1	Ocrevus
⬇ Inj 40 mg per ml, 23 ml vial	16,900.00	1	Ocrevus SC

➥ **Restricted** (RS1997)

Initiation – Multiple Sclerosis - ocrelizumab

Any relevant practitioner

Re-assessment required after 12 months

Either:

1 All of the following:

1.1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and

1.2 Patient has an EDSS score between 0 – 6.0; and

1.3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and

1.4 All of the following:

1.4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and

1.4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and

1.4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and

1.4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and

1.4.5 Either:

1.4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze

continued…

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Functional System scores by at least 1 point; or

1.4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and

1.5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and

1.6 Any of the following:

1.6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or

1.6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or

1.6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or

1.6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or

1.6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan; or

2 Patient has an active Special Authority approval for either dimethyl fumarate, fingolimod, glatiramer acetate, interferon beta-1-alpha, interferon beta-1-beta, natalizumab or teriflunomide.

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Continuation – Multiple Sclerosis - ocrelizumab

Any relevant practitioner

Patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use unilateral or bilateral aids at any time in the last six months (ie the patient has walked 100 metres or more with or without aids in the last six months).

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Initiation – Primary Progressive Multiple Sclerosis

Any relevant practitioner

Re-assessment required after 12 months

All of the following:

1 Diagnosis of primary progressive multiple sclerosis (PPMS) meets the 2017 McDonald criteria and has been confirmed by a neurologist; and

2 Patient has an EDSS 2.0 (score equal to or greater than 2 on pyramidal functions) to EDSS 6.5; and

3 Patient has no history of relapsing remitting multiple sclerosis.

Continuation – Primary Progressive Multiple Sclerosis

Any relevant practitioner

Patient has had an EDSS score of less than or equal to 6.5 at any time in the last six months (ie patient has walked 20 metres with bilateral assistance/aids, without rest in the last six months).

Sedatives and Hypnotics

CHLORAL HYDRATE

Oral liq 100 mg per ml

Oral liq 200 mg per ml

LORMETAZEPAM – **Restricted:** For continuation only

➥ Tab 1 mg

MELATONIN – **Restricted** see terms below

⬇ Tab modified-release 2 mg – **5% DV Dec-24 to 2027** ..... 5.80 30 **Vigisom**

⬇ Tab 3 mg

Note: Only for use in compounding an oral liquid formulation, for in-hospital use only.

➥ **Restricted** (RS1576)

Initiation – insomnia secondary to neurodevelopmental disorder

Psychiatrist, paediatrician, neurologist or respiratory specialist

Re-assessment required after 12 months

All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder); and
2 Behavioural and environmental approaches have been tried or are inappropriate; and
3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
4 Patient is aged 18 years or under.

Continuation – insomnia secondary to neurodevelopmental disorder
Psychiatrist, paediatrician, neurologist or respiratory specialist
Re-assessment required after 12 months
All of the following:

1 Patient is aged 18 years or under; and
2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Initiation – insomnia where benzodiazepines and zopiclone are contraindicated
Both:

1 Patient has insomnia and benzodiazepines and zopiclone are contraindicated; and
2 For in-hospital use only.

MIDAZOLAM			
Tab 7.5 mg			
Oral liq 2 mg per ml			
Inj 5 mg per ml, 1 ml plastic ampoule	22.50	10	Midazolam-Pfizer
Inj 1 mg per ml, 5 ml ampoule – **5% DV May-25 to 2027**	7.80	10	**Midazolam-Baxter**
Inj 5 mg per ml, 3 ml ampoule – **5% DV May-25 to 2027**	4.75	5	**Midazolam-Baxter**
PHENOBARBITONE			
Inj 130 mg per ml, 1 ml vial			
Inj 200 mg per ml, 1 ml ampoule			
TEMAZEPAM			
Tab 10 mg – **5% DV Feb-24 to 2026**	1.40	25	**Normison**
TRIAZOLAM – **Restricted:** For continuation only			
➥ Tab 125 mcg			
➥ Tab 250 mcg			
ZOPICLONE			
Tab 7.5 mg – **5% DV Feb-25 to 2027**	21.85	500	**Zopiclone Actavis**

Spinal Muscular Atrophy

NUSINERSEN – **Restricted** see terms below			
⬇ Inj 12 mg per 5 ml vial	120,000.00	1	Spinraza

➥ **Restricted** (RS1938)
Initiation
Re-assessment required after 12 months
All of the following:

1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
2 Patient is 18 years of age or under; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3 Either:
- 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
- 3.2 Both:
  - 3.2.1 Patient is pre-symptomatic; and
  - 3.2.2 Patient has three or less copies of SMN2.

Continuation

Re-assessment required after 12 months

All of the following:

1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
2 Patient does not require invasive permanent ventilation (at least 16 hours per day), in the absence of a potentially reversible cause while being treated with nusinersen; and
3 Nusinersen not to be administered in combination other SMA disease modifying treatments or gene therapy.

RISDIPLAM – **Restricted** see terms below

Note: the supply of risdiplam is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/risdiplam

⬇	Powder for oral soln 750 mcg per ml, 60 mg per bottle	14,100.00	80 ml	Evrysdi

➥ **Restricted** (RS1954)

Initiation

Re-assessment required after 12 months

All of the following:

1 Patient has genetic documentation of homozygous SMN1 gene deletion, homozygous SMN1 point mutation, or compound heterozygous mutation; and
2 Patient is 18 years of age or under; and
3 Either:
- 3.1 Patient has experienced the defined signs and symptoms of SMA type I, II or IIIa prior to three years of age; or
- 3.2 Both:
  - 3.2.1 Patient is pre-symptomatic; and
  - 3.2.2 Patient has three or less copies of SMN2.

Continuation

Re-assessment required after 12 months

All of the following:

1 There has been demonstrated maintenance of motor milestone function since treatment initiation; and
2 Patient does not require invasive permanent ventilation (at least 16 hours per day), in the absence of a potentially reversible cause while being treated with risdiplam; and
3 Risdiplam not to be administered in combination other SMA disease modifying treatments or gene therapy.

Stimulants / ADHD Treatments

ATOMOXETINE

Cap 10 mg – **5% DV Aug-24 to 2026**	43.02	28	**APO-Atomoxetine**
Cap 18 mg – **5% DV Aug-24 to 2026**	45.57	28	**APO-Atomoxetine**
Cap 25 mg – **5% DV Aug-24 to 2026**	44.30	28	**APO-Atomoxetine**
Cap 40 mg – **5% DV Aug-24 to 2026**	46.21	28	**APO-Atomoxetine**
Cap 60 mg – **5% DV Aug-24 to 2026**	51.31	28	**APO-Atomoxetine**
Cap 80 mg – **5% DV Aug-24 to 2026**	65.20	28	**APO-Atomoxetine**
Cap 100 mg – **5% DV Aug-24 to 2026**	65.71	28	**APO-Atomoxetine**

CAFFEINE

Tab 100 mg

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
DEXAMFETAMINE SULFATE – **Restricted** see terms below			
⬇ Tab 5 mg	29.80	100	Dexamfetamine Aspen Noumed Dexamfetamine

➥ **Restricted** (RS2166)

Initiation – ADHD

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.

Note: Prescribing practitioner detail is in the relevant approval notice published in the New Zealand Gazette. Approval notices are located through the 'Medicines (controlled drugs) with restrictions under regulation 22 of the Misuse of Drugs Regulations 1977' section of the Medsafe 'Restrictions on the Supply, Prescribing or Administration of Medicines under the Medicines Act 1981 and Misuse of Drugs Regulations 1977' webpage (https://www.medsafe.govt.nz/profs/riss/restrict.asp#MedicinesReg22 as of April 2025).

Initiation – Narcolepsy

Neurologist or respiratory specialist

Patient suffers from narcolepsy.

LISDEXAMFETAMINE DIMESILATE – **Restricted** see terms below			
⬇ Cap 30 mg	60.00	30	Vyvanse
⬇ Cap 50 mg	60.00	30	Vyvanse
⬇ Cap 70 mg	60.00	30	Vyvanse

➥ **Restricted** (RS2167)

Initiation

Either:

1 Patient is currently on treatment with lisdexamfetamine dimesilate and met all the following criteria prior to commencing treatment; or

2 All of the following:

2.1 ADHD (Attention Deficit and Hyperactivity Disorder); and

2.2 Diagnosed according to DSM-5 or ICD 11 criteria; and

2.3 Any of the following:

2.3.1 Patient is taking a currently subsidised formulation of atomoxetine or methylphenidate hydrochloride (extended-release) for ADHD and has not received sufficient clinical benefit or has experienced intolerable side effects; or

2.3.2 Patient is taking a currently subsidised formulation of dexamfetamine sulfate (immediate-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or

2.3.3 There is significant concern regarding the risk of diversion or abuse of immediate release dexamfetamine sulfate; or

2.3.4 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained release) which has not been effective due to significant administration and/or treatment adherence difficulties; or

2.3.5 There is significant concern regarding the risk of diversion or abuse of immediate release methylphenidate hydrochloride; or

2.3.6 Both:

2.3.6.1 Patient would have been prescribed a subsidised formulation of methylphenidate hydrochloride (extended-release) but has been unable to access due to supply issues with methylphenidate hydrochloride (extended-release); and

2.3.6.2 Other alternative stimulant presentations (methylphenidate or dexamfetamine) are not appropriate; and

2.4 Lisdexamfetamine dimesilate is not to be used in combination with another funded methylphenidate presentation.

Note: Prescribing practitioner detail is in the relevant approval notice published in the New Zealand Gazette. Approval notices are located through the 'Medicines (controlled drugs) with restrictions under regulation 22 of the Misuse of Drugs Regulations 1977' section of the Medsafe 'Restrictions on the Supply, Prescribing or Administration of Medicines under the Medicines Act 1981 and Misuse of Drugs Regulations 1977' webpage (https://www.medsafe.govt.nz/profs/riss/restrict.asp#MedicinesReg22 as of April 2025).

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
METHYLPHENIDATE HYDROCHLORIDE – **Restricted** see terms below			
⬇ Tab extended-release 18 mg	58.96	30	Concerta
	15.25		Methylphenidate ER - Teva
⬇ Tab extended-release 27 mg	65.44	30	Concerta
	16.25		Methylphenidate ER - Teva
⬇ Tab extended-release 36 mg	71.93	30	Concerta
	21.25		Methylphenidate ER - Teva
⬇ Tab extended-release 54 mg	86.24	30	Concerta
	24.25		Methylphenidate ER - Teva
⬇ Tab immediate-release 5 mg	3.20	30	Rubifen
⬇ Tab immediate-release 10 mg	4.00	30	Ritalin
	3.00		Rubifen
⬇ Tab modified-release 18 mg	15.25	30	Methylphenidate Sandoz XR
⬇ Tab immediate-release 20 mg	7.85	30	Rubifen
⬇ Tab sustained-release 20 mg	10.95	30	Rubifen SR
⬇ Tab modified-release 27 mg	16.25	30	Methylphenidate Sandoz XR
⬇ Tab modified-release 36 mg	21.25	30	Methylphenidate Sandoz XR
⬇ Tab modified-release 54 mg	24.25	30	Methylphenidate Sandoz XR
⬇ Cap modified-release 10 mg	19.41	30	Ritalin LA
⬇ Cap modified-release 20 mg	27.72	30	Ritalin LA
⬇ Cap modified-release 30 mg	34.39	30	Ritalin LA
⬇ Cap modified-release 40 mg	38.67	30	Ritalin LA

➥ **Restricted** (RS2168)

Initiation – ADHD (immediate-release and sustained-release formulations)

Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria.

Initiation – Extended-release and modified-release formulations

Both:

1 Patient has ADHD (Attention Deficit and Hyperactivity Disorder), diagnosed according to DSM-IV or ICD 10 criteria; and
2 Either:
  2.1 Patient is taking a currently listed formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or treatment adherence difficulties; or
  2.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Note: Prescribing practitioner detail is in the relevant approval notice published in the New Zealand Gazette. Approval notices are located through the 'Medicines (controlled drugs) with restrictions under regulation 22 of the Misuse of Drugs Regulations 1977' section of the Medsafe 'Restrictions on the Supply, Prescribing or Administration of Medicines under the Medicines Act 1981 and Misuse of Drugs Regulations 1977' webpage (https://www.medsafe.govt.nz/profs/riss/restrict.asp#MedicinesReg22 as of April 2025).

Initiation – Narcolepsy (immediate-release and sustained-release formulations)

Neurologist or respiratory specialist

Patient suffers from narcolepsy.

Initiation – Narcolepsy* (extended-release only)

Neurologist or respiratory specialist

Patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Concerta, Ritalin LA or Methylphenidate Sandoz XR.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
MODAFINIL – **Restricted** see terms below			
⬇ Tab 100 mg – **5% DV May-25 to 2027**	14.27	30	**Modafinil Max Health**

➥ **Restricted** (RS2106)

Initiation – Narcolepsy

Neurologist or respiratory specialist

Either:

1 All of the following:
- 1.1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 1.2 Either:
  - 1.2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 1.2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 1.3 Either:
  - 1.3.1 An effective dose of a listed formulation of methylphenidate or dexamphetamine has been trialled and discontinued because of intolerable side effects; or
  - 1.3.2 Methylphenidate and dexamphetamine are contraindicated; or

2 Both:
- 2.1 Patient meets the Hospital Restriction criteria for methylphenidate hydrochloride for narcolepsy; and
- 2.2 Patient is unable to access methylphenidate hydrochloride presentations due to an out of stock (see note).

Note: Criterion 2 is to permit short-term funding to cover an out-of-stock of methylphenidate hydrochloride.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
Tab 5 mg – **5% DV Jun-24 to 2026**	3.70	84	**Ipca-Donepezil**
Tab 10 mg – **5% DV Jun-24 to 2026**	5.50	84	**Ipca-Donepezil**
RIVASTIGMINE – **Restricted** see terms below			
⬇ Patch 4.6 mg per 24 hour – **5% DV Mar-25 to 2027**	49.40	30	**Rivastigmine Patch BNM 5**
⬇ Patch 9.5 mg per 24 hour – **5% DV Mar-25 to 2027**	49.40	30	**Rivastigmine Patch BNM 10**

➥ **Restricted** (RS2139)

Initiation

Re-assessment required after 6 months

Both:

1 The patient has been diagnosed with dementia; and

2 The patient is contraindicated to or has experienced intolerable side effects from donepezil tablets.

Continuation

Re-assessment required after 12 months

Both:

1 The treatment remains appropriate; and

2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE – **Restricted** see terms on the next page			
⬇ Tab 2 mg with naloxone 0.5 mg – **5% DV May-26 to 2028**	11.76	28	**Buprenorphine Naloxone BNM**
⬇ Tab 8 mg with naloxone 2 mg – **5% DV May-26 to 2028**	26.86	28	**Buprenorphine Naloxone BNM**

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

		Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1172)
Initiation – Detoxification
All of the following:
1 Patient is opioid dependent; and
2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
3 Prescriber works in an opioid treatment service approved by the Ministry of Health.
Initiation – Maintenance treatment
All of the following:
1 Patient is opioid dependent; and
2 Patient will not be receiving methadone; and
3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
4 Prescriber works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE
Tab modified-release 150 mg – **5% DV May-24 to 2026** ... 15.00 | 30 | **Zyban**

DISULFIRAM
Tab 200 mg ... 236.40 | 100 | Antabuse

NALTREXONE HYDROCHLORIDE – **Restricted** see terms below
⬇ Tab 50 mg – **5% DV Dec-23 to 2026** ... 83.33 | 30 | **Naltraccord**
➥ **Restricted** (RS1173)
Initiation – Alcohol dependence
Both:
1 Patient is currently enrolled, or is planned to be enrolled, in a recognised comprehensive treatment programme for alcohol dependence; and
2 Naltrexone is to be prescribed by, or on the recommendation of, a physician working in an Alcohol and Drug Service.
Initiation – Constipation
For the treatment of opioid-induced constipation.

NICOTINE – **Some items restricted** see terms below
Patch 7 mg per 24 hours ... 19.62 | 28 | Habitrol
Patch 14 mg per 24 hours ... 21.57 | 28 | Habitrol
Patch 21 mg per 24 hours ... 24.72 | 28 | Habitrol
⬇ Oral spray 1 mg per dose | | | *e.g. Nicorette QuickMist Mouth Spray*
Lozenge 1 mg ... 22.53 | 216 | Habitrol
Lozenge 2 mg ... 24.68 | 216 | Habitrol
⬇ Soln for inhalation 15 mg cartridge
Gum 2 mg ... 23.02 | 204 | Habitrol (Fruit)
Habitrol (Mint)
Gum 4 mg ... 25.98 | 204 | Habitrol (Fruit)
Habitrol (Mint)

➥ **Restricted** (RS1873)
Initiation
Any of the following:
1 For perioperative use in patients who have a 'nil by mouth' instruction; or
2 For use within mental health inpatient units; or
3 Patient would be admitted to a mental health inpatient unit, but is unable to due to COVID-19 self-isolation requirement; or
4 For acute use in agitated patients who are unable to leave the hospital facilities.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
VARENICLINE – **Restricted** see terms below			
⬇ Tab 0.5 mg x 11 and 1 mg x 42 – **5% DV Sep-26 to 2028**	16.67	53	Champix
	15.99		**Pharmacor Varenicline**
⬇ Tab 1 mg – **5% DV Sep-26 to 2028**	17.62	56	Champix
	10.99		**Pharmacor Varenicline**

(Champix Tab 0.5 mg x 11 and 1 mg x 42 to be delisted 1 September 2026)
(Champix Tab 1 mg to be delisted 1 September 2026)

➥ **Restricted** (RS1702)

Initiation

All of the following:

1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
3 Either:
  3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
6 The patient is not pregnant; and
7 The patient will not be prescribed more than 12 weeks' funded varenicline in a 12 month period.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE – **Restricted** see terms below

⬇	Inj 25 mg vial – **5% DV Apr-25 to 2027**	50.05	1	**Bendamustine Sandoz**
⬇	Inj 100 mg vial – **5% DV Apr-25 to 2027**	200.20	1	**Bendamustine Sandoz**

➥ **Restricted** (RS2061)

Initiation – CLL*

All of the following:

1 The patient has chronic lymphocytic leukaemia requiring treatment; and
2 Patient has ECOG performance status 0-2; and
3 Bendamustine is to be administered at a maximum dose of 100 mg/m^2 on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: Indication marked with a * includes indications that are unapproved. 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL).

Initiation – Indolent, Low-grade lymphomas

Re-assessment required after 9 months

All of the following:

1 The patient has indolent low grade NHL requiring treatment; and
2 Patient has ECOG performance status of 0-2; and
3 Any of the following:
  3.1 Both:
    3.1.1 Patient is treatment naive; and
    3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  3.2 Both:
    3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
    3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
  3.3 All of the following:
    3.3.1 The patient has not received prior bendamustine therapy; and
    3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
    3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
  3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Continuation – Indolent, Low-grade lymphomas

Re-assessment required after 9 months

Either:

1 Both:
  1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
  1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
2 Both:
  2.1 Patients have not received a bendamustine regimen within the last 12 months; and
  2.2 Either:
    2.2.1 Both:
      2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or

2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenström's macroglobulinaemia.

Initiation – Hodgkin's lymphoma*

Relevant specialist or medical practitioner on the recommendation of a relevant specialist

Limited to 6 months treatment

All of the following:

1 Patient has Hodgkin's lymphoma requiring treatment; and
2 Patient has a ECOG performance status of 0-2; and
3 Patient has received one prior line of chemotherapy; and
4 Patient's disease relapsed or was refractory following prior chemotherapy; and
5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN			
Tab 2 mg	89.25	100	Myleran
Inj 6 mg per ml, 10 ml ampoule			
CARMUSTINE			
Inj 100 mg vial	710.00	1	BiCNU BiCNU S29 Novadoz
CHLORAMBUCIL			
Tab 2 mg			
CYCLOPHOSPHAMIDE			
Tab 50 mg – **5% DV Dec-24 to 2027**	145.00	50	**Cyclonex**
Inj 1 g vial – **5% DV Feb-25 to 2027**	47.46	1	**Endoxan**
Inj 2 g vial – **5% DV Feb-25 to 2027**	95.06	1	**Endoxan**
IFOSFAMIDE			
Inj 1 g vial	96.00	1	Holoxan
Inj 2 g vial	180.00	1	Holoxan
LOMUSTINE			
Cap 40 mg	880.00	20	Medac
MELPHALAN			
Tab 2 mg			
Inj 50 mg vial – **5% DV Dec-23 to 2026**	48.25	1	**Melpha**
THIOTEPA			
Inj 15 mg vial – **5% DV Apr-24 to 2026**	398.00	1	**Tepadina**
Inj 100 mg vial – **5% DV Apr-24 to 2026**	1,800.00	1	**Tepadina** Thiotepa-Reach

Anthracyclines and Other Cytotoxic Antibiotics

BLEOMYCIN SULPHATE			
Inj 15,000 iu vial	185.16	1	DBL Bleomycin Sulfate
DACTINOMYCIN [ACTINOMYCIN D]			
Inj 0.5 mg vial	255.00	1	Cosmegen

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
DAUNORUBICIN			
Inj 18.7 mg vial	160.75	1	Pfizer
Inj 20 mg vial	171.93	1	Cerubidine
DOXORUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 5 ml vial			
Inj 2 mg per ml, 25 ml vial	75.55	1	Doxorubicin Ebewe
Inj 50 mg vial			
Inj 2 mg per ml, 100 ml vial	164.50	1	Doxorubicin Ebewe
EPIRUBICIN HYDROCHLORIDE			
Inj 2 mg per ml, 25 ml vial	284.36	1	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial	284.36	1	Epirubicin Ebewe
IDARUBICIN HYDROCHLORIDE			
Inj 5 mg vial	109.74	1	Zavedos
Inj 10 mg vial	233.64	1	Zavedos
MITOMYCIN C			
Inj 5 mg vial			
Inj 20 mg vial – **5% DV May-26 to 2028**	1,129.94	1	**Teva**
MITOZANTRONE			
Inj 2 mg per ml, 10 ml vial	170.63	1	Mitozantrone Ebewe

Antimetabolites

AZACITIDINE – **Restricted** see terms below

⬇ Inj 100 mg vial – **5% DV Mar-25 to 2027** 50.00 | 1 | **Azacitidine Dr Reddy's**

➥ **Restricted** (RS2116)

Initiation

Re-assessment required after 12 months

Both:

1 Any of the following:

1.1 The individual has intermediate or high risk MDS based on an internationally recognised scoring system; or

1.2 The individual has chronic myelomonocytic leukaemia (based on an intermediate or high risk score from an internationally recognised scoring system or 10%-29% marrow blasts without myeloproliferative disorder); or

1.3 The individual has acute myeloid leukaemia according to World Health Organisation (WHO) Classification; and

2 The individual has an estimated life expectancy of at least 3 months.

Continuation

Re-assessment required after 12 months

No evidence of disease progression.

CAPECITABINE			
Tab 150 mg – **5% DV Feb-26 to 2028**	10.92	60	**Capecitabine Viatris**
Tab 500 mg – **5% DV Feb-26 to 2028**	50.96	120	**Capecitabine Viatris**
CLADRIBINE			
Inj 2 mg per ml, 5 ml vial			
Inj 1 mg per ml, 10 ml vial	749.96	1	Leustatin
CYTARABINE			
Inj 20 mg per ml, 5 ml vial	472.00	5	Pfizer
Inj 100 mg per ml, 20 ml vial	48.80	1	Cytarabine DBL Pfizer
FLUDARABINE PHOSPHATE			
Tab 10 mg	412.00	20	Fludara Oral
Inj 50 mg vial	744.95	5	Fludarabine Ebewe

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
FLUOROURACIL			
Inj 50 mg per ml, 20 ml vial – **5% DV Dec-24 to 2027**	10.51	1	**Fluorouracil Accord**
Inj 50 mg per ml, 50 ml vial	14.72	1	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial – **5% DV Dec-24 to 2027**	19.36	1	**Fluorouracil Accord**
GEMCITABINE HYDROCHLORIDE			
Inj 43.3 mg per ml (equivalent to 38 mg per ml gemcitabine), 26.3 ml vial – **5% DV Jun-24 to 2026**	18.94	1	**DBL Gemcitabine**
MERCAPTOPURINE			
Tab 50 mg – **5% DV Dec-25 to 2028**	19.50	25	**Puri-nethol**
⬇ Oral suspension 20 mg per ml	428.00	100 ml	Xaluprine Allmercap

➥ **Restricted** (RS1635)
Initiation
Paediatric haematologist or paediatric oncologist
Re-assessment required after 12 months
The patient requires a total dose of less than one full 50 mg tablet per day.
Continuation
Paediatric haematologist or paediatric oncologist
Re-assessment required after 12 months
The patient requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE			
Tab 2.5 mg – **5% DV Dec-24 to 2027**	7.80	90	**Trexate**
Tab 10 mg – **5% DV Dec-24 to 2027**	26.40	90	**Trexate**
Inj 2.5 mg per ml, 2 ml vial			
Inj 7.5 mg prefilled syringe – **5% DV Feb-25 to 2027**	29.17	1	**Methotrexate Sandoz**
Inj 10 mg prefilled syringe – **5% DV Feb-25 to 2027**	19.09	1	**Methotrexate Sandoz**
Inj 15 mg prefilled syringe – **5% DV Feb-25 to 2027**	24.53	1	**Methotrexate Sandoz**
Inj 20 mg prefilled syringe – **5% DV Feb-25 to 2027**	16.64	1	**Methotrexate Sandoz**
Inj 25 mg prefilled syringe – **5% DV Feb-25 to 2027**	20.72	1	**Methotrexate Sandoz**
Inj 30 mg prefilled syringe – **5% DV Feb-25 to 2027**	55.00	1	**Methotrexate Sandoz**
Inj 25 mg per ml, 2 ml vial	30.00	5	Methotrexate DBL Onco-Vial
Inj 25 mg per ml, 20 ml vial	45.00	1	DBL Methotrexate Onco-Vial
Inj 100 mg per ml, 10 ml vial	25.00	1	Methotrexate Ebewe
Inj 100 mg per ml, 50 ml vial – **5% DV Dec-23 to 2026**	67.99	1	**Methotrexate Ebewe**
PEMETREXED			
Inj 100 mg vial – **5% DV Apr-25 to 2027**	8.99	1	**Pemetrexed-AFT**
Inj 500 mg vial – **5% DV Apr-25 to 2027**	29.99	1	**Pemetrexed-AFT**
THIOGUANINE			
Tab 40 mg			

Other Cytotoxic Agents

AMSACRINE			
Inj 50 mg per ml, 1.5 ml ampoule			
Inj 75 mg			
ANAGRELIDE HYDROCHLORIDE			
Cap 0.5 mg			
ARSENIC TRIOXIDE			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	Phenasen

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
BORTEZOMIB – **Restricted** see terms below			
⬇ Inj 3.5 mg vial – **5% DV Sep-26 to 2028**	23.99	1	**Bortezomib Eugia**
	74.93		DBL Bortezomib

(DBL Bortezomib Inj 3.5 mg vial to be delisted 1 September 2026)

➥ **Restricted** (RS2169)

Initiation – plasma cell dyscrasia

The patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment.

Initiation – Waldenström Macroglobulinaemia

Re-assessment required after 12 months

Both:

1 The patient has Waldenström Macroglobulinaemia/Lymphoplasmacytic Lymphoma requiring treatment; and
2 The patient has not received prior bortezomib treatment.

Continuation – Waldenström Macroglobulinaemia

Re-assessment required after 12 months

Patient has no evidence of clinical disease progression during bortezomib use.

DACARBAZINE			
Inj 200 mg vial	72.11	1	DBL Dacarbazine
ETOPOSIDE			
Cap 50 mg	340.73	20	Vepesid
Cap 100 mg	340.73	10	Vepesid
Inj 20 mg per ml, 5 ml vial	7.90	1	Rex Medical
ETOPOSIDE (AS PHOSPHATE)			
Inj 100 mg vial	40.00	1	Etopophos
HYDROXYUREA [HYDROXYCARBAMIDE]			
Cap 500 mg – **5% DV Dec-23 to 2026**	20.72	100	**Devatis**
IBRUTINIB – **Restricted** see terms below			
⬇ Tab 140 mg	3,217.00	30	Imbruvica
⬇ Tab 420 mg	9,652.00	30	Imbruvica

➥ **Restricted** (RS2190)

Initiation – previously untreated chronic lymphocytic leukaemia in combination with venetoclax

Either:

1 Individual is currently on treatment with ibrutinib and/or venetoclax and met all of the following criteria prior to commencing treatment; or
2 Both:
  2.1 Individual has previously untreated CLL; and
  2.2 Ibrutinib is to be administered at a maximum dose of 420 mg daily for 3 (28 day) cycles as monotherapy, followed by a maximum of 12 (28 day) cycles in combination with venetoclax.

Initiation – chronic lymphocytic leukaemia (CLL)

Re-assessment required after 6 months

All of the following:

1 Individual has chronic lymphocytic leukaemia (CLL) requiring therapy; and
2 Ibrutinib is to be used as monotherapy; and
3 Individual has experienced intolerable side effects, or their disease has relapsed or is refractory following at least one prior line of therapy; and
4 Individual has not received ibrutinib monotherapy previously.

Continuation – chronic lymphocytic leukaemia (CLL)

Re-assessment required after 12 months

No evidence of clinical disease progression.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
IRINOTECAN HYDROCHLORIDE			
Inj 20 mg per ml, 5 ml vial	52.57	1	Accord
Inj 20 mg per ml, 25 ml vial	262.85	1	Accord
LENALIDOMIDE (VIATRIS) – **Restricted** see terms below			
⬇ Cap 5 mg – **5% DV Feb-25 to 31 Jan 2028**	76.92	21	**Lenalidomide Viatris**
⬇ Cap 10 mg – **5% DV Feb-25 to 31 Jan 2028**	50.30	21	**Lenalidomide Viatris**
⬇ Cap 15 mg – **5% DV Feb-25 to 31 Jan 2028**	62.13	21	**Lenalidomide Viatris**
⬇ Cap 25 mg – **5% DV Feb-25 to 31 Jan 2028**	65.09	21	**Lenalidomide Viatris**

➥ **Restricted** (RS2044)

Initiation – Plasma cell dyscrasia

Any relevant practitioner

Both:

1 Patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; and
2 Patient is not refractory to prior lenalidomide use.

Initiation – Myelodysplastic syndrome

Any relevant practitioner

Re-assessment required after 6 months

Both:

1 Patient has low or intermediate-1 risk myelodysplastic syndrome (based on IPSS or an IPSS-R score of less than 3.5) associated with a deletion 5q cytogenetic abnormality; and
2 Patient has transfusion-dependent anaemia.

Continuation – Myelodysplastic syndrome

Any relevant practitioner

Re-assessment required after 12 months

Both:

1 Patient has not needed a transfusion in the last 4 months; and
2 No evidence of disease progression.

NIRAPARIB – **Restricted** see terms below			
⬇ Tab 100 mg	8,929.84	56	Zejula
	13,393.50	84	Zejula
⬇ Cap 100 mg	8,929.84	56	Zejula

➥ **Restricted** (RS2027)

Initiation

Re-assessment required after 6 months

All of the following:

1 Patient has advanced high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
2 Patient has received at least one line** of treatment with platinum-based chemotherapy; and
3 Patient has experienced a partial or complete response to the preceding treatment with platinum-based chemotherapy; and
4 Patient has not previously received funded treatment with a PARP inhibitor; and
5 Either:

5.1 Treatment will be commenced within 12 weeks of the patient's last dose of the preceding platinum-based regimen; or
5.2 Patient commenced treatment with niraparib prior to 1 May 2024; and

6 Treatment to be administered as maintenance treatment; and
7 Treatment not to be administered in combination with other chemotherapy.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation

Re-assessment required after 6 months

All of the following:

1 No evidence of progressive disease; and
2 Treatment to be administered as maintenance treatment; and
3 Treatment not to be administered in combination with other chemotherapy; and
4 Either:
  4.1 Treatment with niraparib to cease after a total duration of 36 months from commencement; or
  4.2 Treatment with niraparib is being used in the second-line or later maintenance setting.

Notes: * "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments

OLAPARIB – **Restricted** see terms below

⬇	Tab 100 mg	3,701.00	56	Lynparza
⬇	Tab 150 mg	3,701.00	56	Lynparza

➥ **Restricted** (RS1925)

Initiation – Ovarian cancer

Medical oncologist

Re-assessment required after 12 months

All of the following:

1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
3 Either:
  3.1 All of the following:
    3.1.1 Patient has newly diagnosed, advanced disease; and
    3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
    3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
  3.2 All of the following:
    3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
    3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
    3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
    3.2.4 Patient has not previously received funded olaparib treatment; and
4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
5 Treatment to be administered as maintenance treatment; and
6 Treatment not to be administered in combination with other chemotherapy.

Continuation – Ovarian cancer

Medical oncologist

Re-assessment required after 12 months

All of the following:

1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
2 Either:
  2.1 No evidence of progressive disease; or
  2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

clinician's opinion; and

3 Treatment to be administered as maintenance treatment; and
4 Treatment not to be administered in combination with other chemotherapy; and
5 Either:
5.1 Both:
5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component. **A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PEGASPARGASE – **Restricted** see terms below

⬇	Inj 750 iu per ml, 5 ml vial	3,973.25	1	Oncaspar LYO

➥ **Restricted** (RS2176)

Initiation – Newly diagnosed ALL

Limited to 15 months treatment

Both:

1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initiation – Relapsed ALL

Limited to 12 months treatment

Both:

1 The patient has relapsed acute lymphoblastic leukaemia; and
2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initiation – Lymphoma

Limited to 12 months treatment

Patient has lymphoma requiring L-asparaginase containing protocol (e.g. SMILE).

PENTOSTATIN [DEOXYCOFORMYCIN]

	Inj 10 mg vial			

POMALIDOMIDE – **Restricted** see terms below

⬇	Cap 1 mg – **5% DV Aug-24 to 31 Jul 2027**	47.45	14	**Pomolide**
		71.18	21	**Pomolide**
⬇	Cap 2 mg – **5% DV Aug-24 to 31 Jul 2027**	94.90	14	**Pomolide**
		142.35	21	**Pomolide**
⬇	Cap 3 mg – **5% DV Aug-24 to 31 Jul 2027**	142.35	14	**Pomolide**
		213.53	21	**Pomolide**
⬇	Cap 4 mg – **5% DV Aug-24 to 31 Jul 2027**	189.81	14	**Pomolide**
		284.71	21	**Pomolide**

➥ **Restricted** (RS2045)

Initiation – Relapsed/refractory plasma cell dyscrasia

Any relevant practitioner

Re-assessment required after 6 months

Both:

1 Patient has relapsed or refractory plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

treatment; and

2 Patient has not received prior funded pomalidomide.

Continuation – Relapsed/refractory plasma cell dyscrasia

Any relevant practitioner

Re-assessment required after 12 months

Patient has no evidence of disease progression.

PROCARBAZINE HYDROCHLORIDE

Cap 50 mg	980.00	50	Natulan

TEMOZOLOMIDE – **Restricted** see terms below

⬇ Cap 5 mg	9.13	5	Temaccord Temozolomide Taro
⬇ Cap 20 mg	16.38	5	Temaccord
⬇ Cap 100 mg	35.98	5	Temaccord
⬇ Cap 140 mg	50.12	5	Temaccord
⬇ Cap 250 mg	86.34	5	Temaccord

➥ **Restricted** (RS2170)

Initiation – gliomas

Re-assessment required after 12 months

Patient has a glioma.

Continuation – gliomas

Re-assessment required after 12 months

Treatment remains appropriate and patient is benefitting from treatment.

Initiation – Neuroendocrine tumours

Re-assessment required after 9 months

All of the following:

1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
2 Temozolomide is to be given in combination with capecitabine; and
3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m^2 per day; and
4 Temozolomide to be discontinued at disease progression.

Continuation – Neuroendocrine tumours

Re-assessment required after 6 months

Both:

1 No evidence of disease progression; and
2 The treatment remains appropriate and the patient is benefitting from treatment.

Initiation – ewing's sarcoma

Re-assessment required after 9 months

Patient has relapse or refractory Ewing's sarcoma.

Continuation – ewing's sarcoma

Re-assessment required after 6 months

Both:

1 No evidence of disease progression; and
2 The treatment remains appropriate and the patient is benefitting from treatment.

Initiation – Neuroblastoma

Re-assessment required after 12 months

Patient has neuroblastoma.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation – Neuroblastoma

Re-assessment required after 12 months

Patient has no evidence of disease progression.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not funded for the treatment of relapsed high grade glioma.

THALIDOMIDE – **Restricted** see terms below

⬇	Cap 50 mg	378.00	28	Thalomid
⬇	Cap 100 mg	756.00	28	Thalomid

➥ **Restricted** (RS2046)

Initiation

Re-assessment required after 12 months

Either:

1 The patient has plasma cell dyscrasia, not including Waldenström macroglobulinaemia, requiring treatment; or
2 The patient has erythema nodosum leprosum.

Continuation

Patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen

TRETINOIN

	Cap 10 mg	479.50	100	Vesanoid

VENETOCLAX – **Restricted** see terms below

⬇	Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42	Venclexta
⬇	Tab 10 mg	13.68	2	Venclexta
⬇	Tab 50 mg	239.44	7	Venclexta
⬇	Tab 100 mg	8,209.41	120	Venclexta

➥ **Restricted** (RS2191)

Initiation – previously untreated chronic lymphocytic leukaemia in combination with obinutuzumab

Either:

1 Individual is currently on treatment with venetoclax and obinutuzumab and met all of the following criteria prior to commencing treatment; or
2 All of the following:
  2.1 Individual has previously untreated chronic lymphocytic leukaemia; and
  2.2 Venetoclax is to be administered with obinutuzumab; and
  2.3 Venetoclax is to be used to a maximum dose of 400 mg and for a total of 12 (28 day) cycles*.

Note: *maximum number of cycles refers to 12 cycles of full dose venetoclax, in addition to the initial 5-week dose ramp-up period.

Initiation – previously untreated chronic lymphocytic leukaemia in combination with ibrutinib

Either:

1 Individual is currently on treatment with venetoclax and/or ibrutinib and met all of the following criteria prior to commencing treatment; or
2 All of the following:
  2.1 Individual has previously untreated chronic lymphocytic leukaemia; and
  2.2 Venetoclax is to be administered in combination with ibrutinib; and
  2.3 Venetoclax is to be used to a maximum dose of 400 mg and for a total of 12 (28 day) cycles*.

Note: *maximum number of cycles refers to 12 cycles of full dose venetoclax, in addition to the initial 5-week dose ramp-up

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

period.

Initiation – relapsed/refractory chronic lymphocytic leukaemia

Re-assessment required after 8 months

All of the following:

1 Individual has chronic lymphocytic leukaemia requiring treatment; and
2 Individual has received at least one prior therapy for chronic lymphocytic leukaemia; and
3 The individual's disease has relapsed; and
4 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
5 Individual has an ECOG performance status of 0-2.

Continuation – relapsed/refractory chronic lymphocytic leukaemia

Re-assessment required after 6 months

Both:

1 Treatment remains clinically appropriate and the individual is benefitting from and tolerating treatment; and
2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL) and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

Initiation – previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*

Re-assessment required after 6 months

All of the following:

1 Individual has previously untreated chronic lymphocytic leukaemia; and
2 There is documentation confirming that the individual has 17p deletion by FISH testing or TP53 mutation by sequencing; and
3 Individual has an ECOG performance status of 0-2.

Continuation – previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*

Re-assessment required after 6 months

No evidence of disease progression.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are unapproved indications

Initiation – previously untreated acute myeloid leukaemia

Re-assessment required after 6 months

Either:

1 The individual is currently on treatment with venetoclax and met all remaining special authority criteria prior to commencing treatment; or
2 All of the following:
  2.1 Individual has previously untreated acute myeloid leukaemia (see note a), according to World Health Organization (WHO) Classification; and
  2.2 Venetoclax not to be used in combination with standard intensive remission induction chemotherapy; and
  2.3 Venetoclax to be used in combination with azacitidine or low dose cytarabine.

Continuation – previously untreated acute myeloid leukaemia

Re-assessment required after 6 months

No evidence of disease progression.

Notes:

a) 'Acute myeloid leukaemia' includes myeloid sarcoma*
b) Indications marked with * are Unapproved indications

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Platinum Compounds			
CARBOPLATIN			
Inj 10 mg per ml, 45 ml vial – **5% DV Dec-24 to 2027**	25.73	1	**Carboplatin Accord**
CISPLATIN			
Inj 1 mg per ml, 50 ml vial	9.45	1	Cisplatin Accord
Inj 1 mg per ml, 100 ml vial – **5% DV Dec-24 to 2027**	18.90	1	**Cisplatin Accord**
OXALIPLATIN			
Inj 5 mg per ml, 20 ml vial	33.35	1	Alchemy Oxaliplatin
Protein-Tyrosine Kinase Inhibitors			
ALECTINIB – **Restricted** see terms below			
⬇ Cap 150 mg	7,935.00	224	Alecensa

➥ **Restricted** (RS1712)

Initiation

Re-assessment required after 6 months

All of the following:

1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
3 Patient has an ECOG performance score of 0-2.

Continuation

Re-assessment required after 6 months

Both:

1 No evidence of progressive disease according to RECIST criteria; and
2 The patient is benefitting from and tolerating treatment.

AXITINIB – **Restricted** see terms below			
⬇ Tab 1 mg	536.40	28	Inlyta
⬇ Tab 5 mg	2,682.00	28	Inlyta

➥ **Restricted** (RS2107)

Initiation

Re-assessment required after 4 months

All of the following:

1 The patient has metastatic renal cell carcinoma; and
2 The disease is of predominant clear cell histology; and
3 The patient has documented disease progression following one previous line of treatment; and
4 The patient has ECOG performance status of 0-2.

Continuation

Re-assessment required after 4 months

No evidence of disease progression..

CRIZOTINIB – **Restricted** see terms below			
⬇ Cap 200 mg	7,250.00	60	Xalkori
⬇ Cap 250 mg	7,250.00	60	Xalkori

➥ **Restricted** (RS2144)

Initiation

Re-assessment required after 6 months

All of the following:

continued…

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $		Brand or Generic Manufacturer

continued…

1 Individual has locally advanced or metastatic, unresectable, non-squamous non-small cell lung cancer; and
2 Either:
  2.1 The individual has not received entrectinib; or
  2.2 Both:
    2.2.1 The individual has received treatment with entrectinib and has discontinued entrectinib due to intolerance; and
    2.2.2 The cancer did not progress while the individual was on entrectinib; and
3 There is documentation confirming that the patient has a ROS1 rearrangement using an appropriate ROS1 test; and
4 Individual has ECOG performance score of 0-3; and
5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation

Re-assessment required after 6 months

Both:

1 Response to treatment has been determined by comparable radiological assessment following the most recent treatment period; and
2 No evidence of disease progression.

DABRAFENIB – **Restricted** see terms below

⬇	Cap 50 mg	6,320.86	120	Tafinlar
⬇	Cap 75 mg	9,481.29	120	Tafinlar

➥ **Restricted** (RS2145)

Initiation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

1 Either:
  1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or
  1.2 Both:
    1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and
    1.2.2 Adjuvant treatment with dabrafenib is required; and
2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
3 Treatment must be adjuvant to complete surgical resection; and
4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
5 The individual has a confirmed BRAF mutation; and
6 Dabrafenib must be administered in combination with trametinib; and
7 The individual has ECOG performance score 0-2.

Notes:

a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Continuation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

Any of the following:

1 All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.1 No evidence of disease recurrence; and
1.2 Dabrafenib must be administered in combination with trametinib; and
1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or

2 All of the following:
2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
2.3 The individual meets initiation criteria for dabrafenib for unresectable or metastatic melanoma; or

3 All of the following:
3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
3.2 The individual has received a BRAF/MEK inhibitor for unresectable or metastatic melanoma; and
3.3 The individual meets continuation criteria for dabrafenib for unresectable or metastatic melanoma.

Initiation – unresectable or metastatic melanoma
Any relevant practitioner
Re-assessment required after 4 months
All of the following:
1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
3 The individual has ECOG performance score 0-2; and
4 The individual has confirmed BRAF mutation; and
5 Dabrafenib must be administered in combination with trametinib; and
6 Any of the following:
6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or
6.3 All of the following:
6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MEK inhibitor.

Continuation – unresectable or metastatic melanoma
Any relevant practitioner
Re-assessment required after 4 months
Both:
1 Any of the following:
1.1 The individual's disease has had a complete response to treatment; or
1.2 The individual's disease has had a partial response to treatment; or
1.3 The individual has stable disease with treatment; and
2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

DASATINIB – **Restricted** see terms below

⬇	Tab 20 mg – **5% DV Mar-25 to 2027**	132.88	60	**Dasatinib-Teva**
⬇	Tab 50 mg – **5% DV Mar-25 to 2027**	304.13	60	**Dasatinib-Teva**
⬇	Tab 70 mg – **5% DV Mar-25 to 2027**	415.75	60	**Dasatinib-Teva**

➥ **Restricted** (RS2055)
Initiation
Haematologist or any relevant practitioner on the recommendation of a haematologist
Re-assessment required after 6 months
Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; or
2 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); or
3 Both:
  3.1 The patient has a diagnosis of CML in chronic phase; and
  3.2 Any of the following:
    3.2.1 Patient has documented treatment failure* with imatinib; or
    3.2.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
    3.2.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system.

Continuation

Haematologist or any relevant practitioner on the recommendation of a haematologist

Re-assessment required after 6 months

Both:

1 Lack of treatment failure while on dasatinib*; and
2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines.

ENTRECTINIB – **Restricted** see terms below

⬇	Cap 200 mg	9,610.00	90	Rozlytrek

➥ **Restricted** (RS2146)

Initiation

Re-assessment required after 6 months

All of the following:

1 Individual has locally advanced or metastatic, unresectable, non-squamous non-small cell lung cancer; and
2 Either:
  2.1 The individual has not received crizotinib; or
  2.2 Both:
    2.2.1 The individual has received an initial Special Authority approval for crizotinib and has discontinued crizotinib due to intolerance; and
    2.2.2 The cancer did not progress while the individual was on crizotinib; and
3 There is documentation confirming that the patient has a ROS1 rearrangement using an appropriate ROS1 test; and
4 Individual has ECOG performance score of 0-3; and
5 Baseline measurement of overall tumour burden is documented clinically and radiologicallyy.

Continuation

Re-assessment required after 6 months

Both:

1 Response to treatment has been determined by comparable radiological assessment following the most recent treatment period; and
2 No evidence of disease progression.

ERLOTINIB – **Restricted** see terms below

⬇	Tab 100 mg – **5% DV Oct-24 to 2027**	280.84	30	**Alchemy**
⬇	Tab 150 mg – **5% DV Oct-24 to 2027**	484.24	30	**Alchemy**

➥ **Restricted** (RS2078)

Initiation

Re-assessment required after 4 months

All of the following:

1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
2 There is documentation confirming that the disease expresses activating mutations of EGFR; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3 Any of the following:
- 3.1 Patient is treatment naive; or
- 3.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
- 3.3 Both:
  - 3.3.1 The patient has discontinued osimertinib or getitinib due to intolerance; and
  - 3.3.2 The cancer did not progress while on osimertinib or gefitinib.

Continuation

Re-assessment required after 6 months

Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB – **Restricted** see terms below

⬇ Tab 250 mg	918.00	30	Iressa

➥ **Restricted** (RS2079)

Initiation

Re-assessment required after 4 months

All of the following:

1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and

2 Any of the following:
- 2.1 Patient is treatment naive; or
- 2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
- 2.3 Both:
  - 2.3.1 The patient has discontinued osimertinib or erlotinib due to intolerance; and
  - 2.3.2 The cancer did not progress whilst on osimertinib or erlotinib; and

3 There is documentation confirming that disease expresses activating mutations of EGFR.

Continuation

Re-assessment required after 6 months

Radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

IMATINIB MESILATE

Cap 100 mg – **5% DV Dec-23 to 2026**	44.93	60	**Imatinib-Rex**
Cap 400 mg – **5% DV Dec-23 to 2026**	69.76	30	**Imatinib-Rex**

LAPATINIB – **Restricted** see terms below

⬇ Tab 250 mg

➥ **Restricted** (RS1828)

Initiation

For continuation use only.

Continuation

Re-assessment required after 12 months

All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and

3 Lapatinib not to be given in combination with trastuzumab; and

4 Lapatinib to be discontinued at disease progression.

LENVATINIB – **Restricted** see terms below

⬇ Cap 4 mg	3,407.40	30	Lenvima
⬇ Cap 10 mg	3,407.40	30	Lenvima

➥ **Restricted** (RS2098)

Initiation – thyroid cancer

Re-assessment required after 6 months

Either:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient is currently on treatment with lenvatinib and met all remaining criteria prior to commencing treatment; or
2 All of the following:
  2.1 The patient has locally advanced or metastatic differentiated thyroid cancer; and
  2.2 Either:
    2.2.1 Patient must have symptomatic progressive disease prior to treatment; or
    2.2.2 Patient must progressive disease at critical anatomical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures; and
  2.3 Any of the following:
    2.3.1 A lesion without iodine uptake in a RAI scan; or
    2.3.2 Receiving cumulative RAI greater than or equal to 600 mCi; or
    2.3.3 Experiencing disease progression after a RAI treatment within 12 months; or
    2.3.4 Experiencing disease progression after two RAI treatments administered within 12 months of each other; and
  2.4 Patient has thyroid stimulating hormone (TSH) adequately supressed; and
  2.5 Patient is not a candidate for radiotherapy with curative intent; and
  2.6 Surgery is clinically inappropriate; and
  2.7 Patient has an ECOG performance status of 0-2.

Continuation – thyroid cancer

Re-assessment required after 6 months

there is no evidence of disease progression.

Initiation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

All of the following:

1 Patient has unresectable hepatocellular carcinoma; and
2 Patient has preserved liver function (Childs-Pugh A); and
3 Transarterial chemoembolisation (TACE) is unsuitable; and
4 Patient has an ECOG performance status of 0-2; and
5 Either:
  5.1 Patient has not received prior systemic therapy for their disease in the palliative setting; or
  5.2 Both:
    5.2.1 Patient has experienced treatment-limiting toxicity from treatment with atezolizumab with bevacizumab; and
    5.2.2 No disease progression since initiation of atezolizumab with bevacizumab.

Continuation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

there is no evidence of disease progression.

Initiation – renal cell carcinoma

Re-assessment required after 4 months

Either:

1 All of the following:
  1.1 The patient has metastatic renal cell carcinoma; and
  1.2 The disease is of predominant clear-cell histology; and
  1.3 The patient has documented disease progression following one previous line of treatment; and
  1.4 The patient has an ECOG performance status of 0-2; and
  1.5 Lenvatinib is to be used in combination with everolimus; or
2 All of the following:
  2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
2.3 Lenvatinib is to be used in combination with everolimus; and
2.4 There is no evidence of disease progression.

Continuation – renal cell carcinoma
Re-assessment required after 4 months
there is no evidence of disease progression.

MIDOSTAURIN – **Restricted** see terms below

⬇	Cap 25 mg	10,981.00	56	Rydapt

➥ **Restricted** (RS2033)
Initiation
All of the following:
1 Patient has a diagnosis of acute myeloid leukaemia; and
2 Condition must be FMS tyrosine kinase 3 (FLT3) mutation positive; and
3 Patient must not have received a prior line of intensive chemotherapy for acute myeloid leukaemia; and
4 Patient is to receive standard intensive chemotherapy in combination with midostaurin only; and
5 Midostaurin to be funded for a maximum of 4 cycles.

NILOTINIB – **Restricted** see terms below

⬇	Cap 150 mg	4,680.00	120	Tasigna
⬇	Cap 200 mg	6,532.00	120	Tasigna

➥ **Restricted** (RS2010)
Initiation
Haematologist
Re-assessment required after 6 months
All of the following:
1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, high risk chronic phase, or in chronic phase; and
2 Either:
2.1 Patient has documented CML treatment failure* with a tyrosine kinase inhibitor (TKI); or
2.2 Patient has experienced treatment limiting toxicity with a tyrosine kinase inhibitor (TKI) precluding further treatment; and
3 Maximum nilotinib dose of 800 mg/day; and
4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.
Continuation
Haematologist
Re-assessment required after 6 months
All of the following:
1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
3 Maximum nilotinib dose of 800 mg/day; and
4 Subsidised for use as monotherapy only.

OSIMERTINIB – **Restricted** see terms below

⬇	Tab 40 mg	9,310.00	30	Tagrisso
⬇	Tab 80 mg	9,310.00	30	Tagrisso

➥ **Restricted** (RS2080)
Initiation – NSCLC – first line
Re-assessment required after 4 months
All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
2 Any of the following:
  2.1 Patient is treatment naïve; or
  2.2 Patient has received prior treatment in the adjuvant setting and/or while awaiting EGFR results; or
  2.3 Both:
    2.3.1 The patient has discontinued gefitinib or erlotinib due to intolerance; and
    2.3.2 The cancer did not progress while on gefitinib or erlotinib; and
3 There is documentation confirming that the cancer expresses activating mutations of EGFR; and
4 Patient has an ECOG performance status 0-3; and
5 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – NSCLC – first line

Re-assessment required after 6 months

response to or stable disease with treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Initiation – NSCLC – second line

Re-assessment required after 4 months

All of the following:

1 Patient has locally advanced or metastatic, incurable, non-squamous non-small cell lung cancer (NSCLC); and
2 Patient has an ECOG performance status 0-3; and
3 The patient must have received previous treatment with erlotinib or gefitinib; and
4 There is documentation confirming that the cancer expresses T790M mutation of EGFR following progression on or after erlotinib or gefitinib; and
5 The treatment must be given as monotherapy; and
6 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – NSCLC – second line

Re-assessment required after 6 months

response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

PALBOCICLIB – **Restricted** see terms below

⬇	Tab 75 mg	1,200.00	21	Palbociclib Pfizer
⬇	Tab 100 mg	1,200.00	21	Palbociclib Pfizer
⬇	Tab 125 mg	1,200.00	21	Palbociclib Pfizer

➥ **Restricted** (RS2034)

Initiation

Re-assessment required after 6 months

Either:

1 All of the following:
  1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
  1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
  1.3 Patient has an ECOG performance score of 0-2; and
  1.4 Either:
    1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
    1.4.2 Both:
      1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
      1.4.2.2 Patient has not received prior systemic treatment for metastatic disease; and
  1.5 Treatment must be used in combination with an endocrine partner; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or

2 All of the following:

2.1 Patient has an active Special Authority approval for ribociclib; and

2.2 Patient has experienced a grade 3 or 4 adverse reaction to ribociclib that cannot be managed by dose reductions and requires treatment discontinuation; and

2.3 Treatment must be used in combination with an endocrine partner; and

2.4 There is no evidence of progressive disease since initiation of ribociclib.

Continuation

Re-assessment required after 12 months

Both:

1 Treatment must be used in combination with an endocrine partner; and

2 There is no evidence of progressive disease since initiation of palbociclib.

PAZOPANIB – **Restricted** see terms below

⬇	Tab 200 mg – **5% DV May-25 to 2027**	172.88	30	Pazopanib ADVZ **Pazopanib Teva**
⬇	Tab 400 mg – **5% DV May-25 to 2027**	464.00	30	**Pazopanib Teva**

➥ **Restricted** (RS2089)

Initiation

Re-assessment required after 3 months

Either:

1 All of the following:

1.1 The patient has metastatic renal cell carcinoma of predominantly clear cell histology; and

1.2 Either:

1.2.1 The patient is treatment naive; or

1.2.2 The patient has only received prior cytokine treatment; and

1.3 The patient has an ECOG performance score of 0-2; and
The patient has intermediate or poor prognosis defined as:

1.4 Any of the following:

1.4.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or

1.4.2 Haemoglobin level < lower limit of normal; or

1.4.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or

1.4.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or

1.4.5 Karnofsky performance score of less than or equal to 70; or

1.4.6 2 or more sites of organ metastasis; or

2 All of the following:

2.1 The patient has metastatic renal cell carcinoma; and

2.2 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and

2.3 The cancer did not progress whilst on sunitinib; and

2.4 Pazopanib to be used for a maximum of 3 months.

Continuation

Re-assessment required after 3 months

No evidence of disease progression.

RIBOCICLIB – **Restricted** see terms on the next page

⬇	Tab 200 mg	1,883.00	21	Kisqali
		3,767.00	42	Kisqali
		5,650.00	63	Kisqali

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS2131)
Initiation
Re-assessment required after 6 months
Either:
1 All of the following:
1.1 Patient has unresectable locally advanced or metastatic breast cancer; and
1.2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
1.3 Patient has an ECOG performance score of 0-2; and
1.4 Either:
1.4.1 Disease has relapsed or progressed during prior endocrine therapy; or
1.4.2 Both:
1.4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal or without menstrual-potential state; and
1.4.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
1.5 Treatment to be used in combination with an endocrine partner; and
1.6 Patient has not received prior funded treatment with a CDK4/6 inhibitor; or
2 All of the following:
2.1 Patient has an active Special Authority approval for palbociclib; and
2.2 Patient has experienced a grade 3 or 4 adverse reaction to palbociclib that cannot be managed by dose reductions and requires treatment discontinuation; and
2.3 Treatment must be used in combination with an endocrine partner; and
2.4 There is no evidence of progressive disease since initiation of palbociclib.
Continuation
Re-assessment required after 12 months
Both:
1 Treatment must be used in combination with an endocrine partner; and
2 There is no evidence of progressive disease since initiation of ribociclib.

RUXOLITINIB – **Restricted** see terms below

⬇	Tab 5 mg	2,500.00	56	Jakavi
⬇	Tab 10 mg	5,000.00	56	Jakavi
⬇	Tab 15 mg	5,000.00	56	Jakavi
⬇	Tab 20 mg	5,000.00	56	Jakavi

➥ **Restricted** (RS1726)
Initiation
Haematologist
Re-assessment required after 12 months
All of the following:
1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
2 Either:
2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
2.2 Both:
2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy;

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

and

3 A maximum dose of 20 mg twice daily is to be given.

Continuation

Relevant specialist or medical practitioner on the recommendation of a Relevant specialist

Re-assessment required after 12 months

Both:

1 The treatment remains appropriate and the patient is benefiting from treatment; and
2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB – **Restricted** see terms below

⬇ Cap 12.5 mg – **5% DV Mar-26 to 2027**	103.11	28	**Sunitinib Rex**
⬇ Cap 25 mg – **5% DV Mar-26 to 2027**	203.15	28	**Sunitinib Rex**
⬇ Cap 50 mg – **5% DV May-26 to 2027**	343.19	28	**Sunitinib Rex**

➥ **Restricted** (RS2109)

Initiation – RCC

Re-assessment required after 4 months

Both:

1 The patient has metastatic renal cell carcinoma; and
2 The patient has not previously received funded sunitinib.

Continuation – RCC

Re-assessment required after 4 months

No evidence of disease progression.

Initiation – GIST

Re-assessment required after 3 months

Both:

1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
2 Either:
  2.1 The patient's disease has progressed following treatment with imatinib; or
  2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Continuation – GIST

Re-assessment required after 6 months

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

1 Any of the following:
  1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non-measurable disease); or
  1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
2 The treatment remains appropriate and the patient is benefiting from treatment.

Continuation – GIST pandemic circumstances

Re-assessment required after 6 months

All of the following:

1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and

continued…

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3 Sunitinib is to be discontinued at progression; and
4 The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Note: GIST - It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

TRAMETINIB – **Restricted** see terms below

⬇	Tab 0.5 mg	2,370.32	30	Mekinist
⬇	Tab 2 mg	9,481.29	30	Mekinist

➥ **Restricted** (RS2147)

Initiation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

1 Either:
  1.1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); or
  1.2 Both:
    1.2.1 The individual has received neoadjuvant treatment with a PD-1/PD-L1 inhibitor; and
    1.2.2 Adjuvant treatment with trametinib is required; and
2 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
3 Treatment must be adjuvant to complete surgical resection; and
4 Treatment must be initiated within 13 weeks of surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
5 The individual has a confirmed BRAF mutation; and
6 Trametinib must be administered in combination with dabrafenib; and
7 The individual has ECOG performance score 0-2.

Notes:

a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Continuation – stage III or IV resected melanoma - adjuvant

Any relevant practitioner

Re-assessment required after 4 months

Any of the following:

1 All of the following:
  1.1 No evidence of disease recurrence; and
  1.2 Trametinib must be administered in combination with dabrafenib; and
  1.3 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months' total treatment course, including any systemic neoadjuvant treatment; or
2 All of the following:
  2.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
  2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
  2.3 The individual meets initiation criteria for trametinib for unresectable or metastatic melanoma; or
3 All of the following:
  3.1 The individual has received adjuvant treatment with a BRAF/MEK inhibitor; and
  3.2 The individual has received a BRAF/MEK inhibitor for unresectable or metastatic melanoma; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3.3 The individual meets continuation criteria for trametinib for unresectable or metastatic melanoma.

Initiation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

1 The individual has metastatic or unresectable melanoma (excluding uveal melanoma) stage III or IV; and
2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
3 The individual has ECOG performance score 0-2; and
4 The individual has confirmed BRAF mutation; and
5 Trametinib must be administered in combination with dabrafenib; and
6 Any of the following:
  6.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
  6.2 The individual did not receive treatment in the adjuvant setting with a BRAF/MEK inhibitor; or
  6.3 All of the following:
    6.3.1 The individual received treatment in the adjuvant setting with a BRAF/MEK inhibitor; and
    6.3.2 The individual did not experience disease recurrence while on treatment with that BRAF/MEK inhibitor; and
    6.3.3 The individual did not experience disease recurrence within six months of completing adjuvant treatment with a BRAF/MEK inhibitor.

Continuation – unresectable or metastatic melanoma

Any relevant practitioner

Re-assessment required after 4 months

Both:

1 Any of the following:
  1.1 The individual's disease has had a complete response to treatment; or
  1.2 The individual's disease has had a partial response to treatment; or
  1.3 The individual has stable disease with treatment; and
2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period.

Taxanes

DOCETAXEL			
Inj 10 mg per ml, 8 ml vial – **5% DV Dec-23 to 2026**	24.91	1	**DBL Docetaxel**
PACLITAXEL			
Inj 6 mg per ml, 16.7 ml vial – **5% DV Aug-24 to 2026**	19.59	1	**Anzatax**
Inj 6 mg per ml, 50 ml vial – **5% DV Aug-24 to 2026**	37.89	1	**Anzatax**

Treatment of Cytotoxic-Induced Side Effects

CALCIUM FOLINATE			
Tab 15 mg	135.33	10	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml ampoule			
Inj 10 mg per ml, 5 ml vial	112.20	5	Eurofolic
Inj 10 mg per ml, 10 ml vial	163.35	5	Eurofolic
Inj 10 mg per ml, 30 ml vial			
Inj 10 mg per ml, 35 ml vial			
Inj 10 mg per ml, 100 ml vial	139.48	1	Eurofolic
DEXRAZOXANE – **Restricted** see terms on the next page			
⬇ Inj 500 mg			*e.g. Cardioxane*

⬆ Item restricted (see ➡ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1695)
Initiation
Medical oncologist, paediatric oncologist, haematologist or paediatric haematologist
All of the following:

1 Patient is to receive treatment with high dose anthracycline given with curative intent; and
2 Based on current treatment plan, patient's cumulative lifetime dose of anthracycline will exceed 250mg/m2 doxorubicin equivalent or greater; and
3 Dexrazoxane to be administered only whilst on anthracycline treatment; and
4 Either:
  4.1 Treatment to be used as a cardioprotectant for a child or young adult; or
  4.2 Treatment to be used as a cardioprotectant for secondary malignancy.

MESNA

Tab 400 mg	314.00	50	Uromitexan
Tab 600 mg	448.50	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule	177.45	15	Uromitexan
Inj 100 mg per ml, 10 ml ampoule	407.40	15	Uromitexan

Vinca Alkaloids

VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial	270.37	5	Hospira

VINCRISTINE SULPHATE

Inj 1 mg per ml, 1 ml vial	74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial	102.73	5	DBL Vincristine Sulfate

VINORELBINE

Cap 20 mg – **5% DV Feb-26 to 2028**	32.10	1	**Vinorelbine Te Arai**
Cap 30 mg – **5% DV Feb-26 to 2028**	42.80	1	**Vinorelbine Te Arai**
Cap 80 mg – **5% DV Feb-26 to 2028**	80.00	1	**Vinorelbine Te Arai**
Inj 10 mg per ml, 1 ml vial			
Inj 10 mg per ml, 5 ml vial			

Endocrine Therapy

ABIRATERONE ACETATE – **Restricted** see terms below

⬇ Tab 250 mg	4,276.19	120	Zytiga

➥ **Restricted** (RS1888)
Initiation
Medical oncologist, radiation oncologist or urologist
Re-assessment required after 6 months
All of the following:

1 Patient has prostate cancer; and
2 Patient has metastases; and
3 Patient's disease is castration resistant; and
4 Either:
  4.1 All of the following:
    4.1.1 Patient is symptomatic; and
    4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

4.1.3 Patient has ECOG performance score of 0-1; and
4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
4.2 All of the following:
4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
4.2.2 Patient has ECOG performance score of 0-2; and
4.2.3 Patient has not had prior treatment with abiraterone.

Continuation
Medical oncologist, radiation oncologist or urologist
Re-assessment required after 6 months
All of the following:
1 Significant decrease in serum PSA from baseline; and
2 No evidence of clinical disease progression; and
3 No initiation of taxane chemotherapy with abiraterone; and
4 The treatment remains appropriate and the patient is benefiting from treatment.

Continuation – pandemic circumstances
Re-assessment required after 6 months
All of the following:
1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
2 Abiraterone acetate to be discontinued at progression; and
3 No initiation of taxane chemotherapy with abiraterone; and
4 The regular renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE
Tab 50 mg – **5% DV Dec-23 to 2026** ..... 4.18 — 28 — **Binarex**

FLUTAMIDE
Tab 250 mg ..... 119.50 — 100 — Flutamin

FULVESTRANT – **Restricted** see terms below
⬇ Inj 50 mg per ml, 5 ml prefilled syringe – **5% DV May-26 to 2028** ..... 181.00 — 2 — **Fulvestrant EVER Pharma**

➥ **Restricted** (RS1732)
Initiation
Medical oncologist
Re-assessment required after 6 months
All of the following:
1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
4 Treatment to be discontinued at disease progression.

Continuation
Medical oncologist
Re-assessment required after 6 months
All of the following:
1 Treatment remains appropriate and patient is benefitting from treatment; and
2 Treatment to be given at a dose of 500 mg monthly; and
3 No evidence of disease progression.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
OCTREOTIDE			
Inj 100 mcg per ml, 1 ml vial	48.50	5	Omega
Inj 50 mcg per ml, 1 ml vial	27.58	5	Omega
Inj 500 mcg per ml, 1 ml vial	113.10	5	Omega
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	Max Health
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	Max Health
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	Max Health
TAMOXIFEN CITRATE			
Tab 10 mg – **5% DV Dec-23 to 2026**	15.00	60	**Tamoxifen Sandoz**
Tab 20 mg – **5% DV Dec-23 to 2026**	5.32	60	**Tamoxifen Sandoz**

Aromatase Inhibitors

ANASTROZOLE			
Tab 1 mg – **5% DV Dec-23 to 2026**	4.39	30	**Anatrole**
EXEMESTANE			
Tab 25 mg – **5% DV Nov-23 to 2026**	9.86	30	**Pfizer Exemestane**
LETROZOLE			
Tab 2.5 mg – **5% DV Dec-24 to 2027**	4.36	28	Accord
	4.67	30	**Letrole**

(Accord Tab 2.5 mg to be delisted 1 July 2026)

Long-acting Somatostatin Analogues

➡ **Restricted** (RS2100)

Initiation – Malignant bowel obstruction

All of the following:

1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has not been successful; and
3 Treatment to be given for up to 4 weeks.

Note: Indications marked with * are unapproved indications

Initiation – acromegaly

Re-assessment required after 3 months

All of the following:

1 The patient has acromegaly; and
2 Either:
  2.1 Treatment with surgery and radiotherapy is not suitable or was unsuccessful; or
  2.2 Treatment is for an interim period while awaiting the beneficial effects of radiotherapy; and
3 Treatment with a dopamine agonist has been unsuccessful.

Continuation – acromegaly

Without reassessment for applications where IGF1 levels have decreased since starting treatment.

Note: In patients with acromegaly, treatment should be discontinued if IGF1 levels have no decreased 3 months after treatment. In patients treated with radiotherapy treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following treatment withdrawal for at least 4 weeks.

Initiation – Other indications

Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 VIPomas and glucagonomas - for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
2 Both:
  2.1 Gastrinoma; and
  2.2 Either:
    2.2.1 Surgery has been unsuccessful; or
    2.2.2 Patient has metastatic disease after treatment with H2 antagonist or proton pump inhibitors has been unsuccessful; or
3 Both:
  3.1 Insulinomas; and
  3.2 Surgery is contraindicated or has not been successful; or
4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
5 Both:
  5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
  5.2 Disabling symptoms not controlled by maximal medical therapy.

Initiation – pre-operative acromegaly
Limited to 12 months treatment
All of the following:
1 Patient has acromegaly; and
2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
3 Patient is scheduled to undergo pituitary surgery in the next six months.

Notes: Indications marked with * are unapproved indications
The use of a long-acting somatostatin analogue in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded under Special Authority

LANREOTIDE – **Restricted** see terms on the previous page

⬆	Inj 60 mg per 0.5 ml, 0.5 ml syringe – **5% DV Aug-25 to 2027**	382.77	1	**Mytolac**
		1,543.79		Somatuline Autogel
⬆	Inj 90 mg per 0.5 ml, 0.5 ml syringe – **5% DV Sep-25 to 2027**	562.92	1	**Mytolac**
		2,054.40		Somatuline Autogel
⬆	Inj 120 mg per 0.5 ml, 0.5 ml syringe – **5% DV Aug-25 to 2027**	646.70	1	**Mytolac**
		2,570.44		Somatuline Autogel

OCTREOTIDE LONG-ACTING – **Restricted** see terms on the previous page

⬆	Inj depot 10 mg prefilled syringe – **5% DV Dec-24 to 2027**	438.40	1	**Sandostatin LAR**
⬆	Inj depot 20 mg prefilled syringe – **5% DV Dec-24 to 2027**	583.70	1	**Sandostatin LAR**
⬆	Inj depot 30 mg prefilled syringe – **5% DV Dec-24 to 2027**	670.80	1	**Sandostatin LAR**

Imaging Agents

AMINOLEVULINIC ACID HYDROCHLORIDE – **Restricted** see terms below

⬇	Powder for oral soln, 30 mg per ml, 1.5 g vial	4,400.00	1	Gliolan
		44,000.00	10	Gliolan

➥ **Restricted** (RS1565)
Initiation – high grade malignant glioma
All of the following:
1 Patient has newly diagnosed, untreated, glioblastoma multiforme; and
2 Treatment to be used as adjuvant to fluorescence-guided resection; and
3 Patient's tumour is amenable to complete resection.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Immunosuppressants

Calcineurin Inhibitors

CICLOSPORIN

Cap 25 mg	44.63	50	Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml	198.13	50 ml	Neoral
Inj 50 mg per ml, 5 ml ampoule	276.30	10	Sandimmun

TACROLIMUS – **Restricted** see terms below

⬇ Cap 0.5 mg	49.60	100	Tacrolimus Sandoz
⬇ Cap 0.75 mg	99.30	100	Tacrolimus Sandoz
⬇ Cap 1 mg	84.30	100	Tacrolimus Sandoz
⬇ Cap 5 mg	248.20	50	Tacrolimus Sandoz
⬇ Inj 5 mg per ml, 1 ml ampoule			

➥ **Restricted** (RS2110)

Initiation – organ transplant recipients

Either:

1 For use in organ transplant recipients; or
2 The individual is receiving induction therapy for an organ transplant.

Initiation – non-transplant indications*

Any specialist

Both:

1 Patient requires long-term systemic immunosuppression; and
2 Either:
  2.1 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response; or
  2.2 Patient is a child with nephrotic syndrome*.

Note: Indications marked with * are unapproved indications

Fusion Proteins

ETANERCEPT – **Restricted** see terms below

⬇ Inj 25 mg autoinjector	690.00	4	Enbrel
⬇ Inj 25 mg vial	690.00	4	Enbrel
⬇ Inj 50 mg autoinjector	1,050.00	4	Enbrel
⬇ Inj 50 mg syringe	1,050.00	4	Enbrel

➥ **Restricted** (RS2177)

Initiation – arthritis - polyarticular course juvenile idiopathic

Re-assessment required after 6 months

Any of the following:

1 Both:
  1.1 Patient has had a Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  1.2 Either:
    1.2.1 Patient has experienced intolerable side effects; or
    1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 At least 5 active joints and at least 3 joints with pain, tenderness or a limited range of motion after a 3-month trial of methotrexate at the maximum tolerated dose, unless contraindicated; or
3 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate at the maximum tolerated dose, unless contraindicated; or
4 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Continuation – arthritis - polyarticular course juvenile idiopathic
Re-assessment required after 2 years
Either:

1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
2 On subsequent reapplications, at least a continuing 30% improvement in active joint count from baseline.

Initiation – arthritis - oligoarticular course juvenile idiopathic
Re-assessment required after 6 months
Any of the following:

1 Both:
1.1 Patient has had a Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
1.2 Either:
1.2.1 Patient has experienced intolerable side effects; or
1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
2 At least 2 active joints with pain, tenderness or a limited range of motion, after a 3-month trial of methotrexate at the maximum tolerated dose, unless contraindicated; or
3 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate at the maximum tolerated dose, unless contraindicated.

Continuation – arthritis - oligoarticular course juvenile idiopathic
Re-assessment required after 2 years
Either:

1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
2 On subsequent reapplications, at least a continuing 30% improvement in active joint count from baseline.

Initiation – arthritis - rheumatoid
Re-assessment required after 6 months
Either:

1 Both:
1.1 Patient has had a Special Authority approval for adalimumab for rheumatoid arthritis; and
1.2 Either:
1.2.1 Patient has experienced intolerable side effects; or
1.2.2 Patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
2 All of the following:
2.1 Patient has had rheumatoid arthritis (either confirmed by radiologic imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive); and
2.2 Patient has received insufficient benefit from at least 3 months of methotrexate at a maximum tolerated dose (unless contraindicated); and
2.3 Patient has received insufficient benefit from at least 3 months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
2.4 Either:
2.4.1 Patient has received insufficient benefit from at least 3 months of methotrexate in combination with the maximum tolerated dose of ciclosporin, unless contraindicated; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.4.2 Patient has received insufficient benefit from at least 3 months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate, unless contraindicated; and

2.5 Either:

2.5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 joints; or

2.5.2 Patient has persistent symptoms of poorly controlled and active disease in at least 4 joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Continuation – arthritis - rheumatoid

Re-assessment required after 2 years

Both:

1 Either:

1.1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or

1.2 On subsequent reapplications, at least a continuing 30% improvement in active joint count from baseline; and

2 Maximum dose 50 mg every 7 days.

Initiation – ankylosing spondylitis

Re-assessment required after 6 months

Either:

1 Both:

1.1 Patient has had a Special Authority approval for adalimumab for ankylosing spondylitis; and

1.2 Either:

1.2.1 Patient has experienced intolerable side effects; or

1.2.2 Patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or

2 All of the following:

2.1 Patient has a confirmed diagnosis of ankylosing spondylitis; and

2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and

2.3 Patient has bilateral sacroiliitis demonstrated by radiologic imaging; and

2.4 Disease has not responded adequately to treatment with two or more NSAID (unless contraindicated) while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and

2.5 Either:

2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and

2.6 BASDAI score of at least 6 on a 10-point scale completed after 3-month exercise trial before ceasing any previous pharmacological treatment and not more than 1 month before the application.

Continuation – ankylosing spondylitis

Re-assessment required after 2 years

Both:

1 BASDAI has improved from pre-treatment baseline either by at least 4 points on a 10-point scale, or by at least 50%; and

2 Maximum dose 50 mg every 7 days.

Initiation – arthritis - psoriatic

Re-assessment required after 6 months

Either:

1 Both:

1.1 Patient has had a Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and

1.2 Either:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.2.1 Patient has experienced intolerable side effects; or
1.2.2 Patient has received insufficient benefit to meet the renewal criteria for psoriatic arthritis; or

2 All of the following:
2.1 Patient has received insufficient benefit from at least 3 months of methotrexate at maximum tolerated dose unless contraindicated; and
2.2 Patient received insufficient benefit from at least 3 months of sulfasalazine or leflunomide at maximum tolerated dose unless contraindicated; and
2.3 Either:
2.3.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 joints; or
2.3.2 Patient has persistent symptoms of poorly controlled and active disease in at least 4 joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
2.4 Any of the following:
2.4.1 Patient has CRP greater than 15 mg/L measured within one month before the application; or
2.4.2 Patient has an ESR greater than 25 mm per hour measured within one month before the application; or
2.4.3 ESR and CRP not measured as patient is receiving prednisone therapy greater than 5 mg per day received for more than 3 months.

Continuation – arthritis - psoriatic
Re-assessment required after 2 years
Both:
1 Either:
1.1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
1.2 At least a continuing 30% improvement in active joint count from baseline; and
2 Maximum dose 50 mg every 7 days.

Initiation – plaque psoriasis, prior TNF use
Limited to 6 months treatment
All of the following:
1 Patient has had a Special Authority approval for adalimumab for plaque psoriasis; and
2 Either:
2.1 Patient has experienced intolerable side effects; or
2.2 Patient has received insufficient benefit to meet the renewal criteria for adalimumab for plaque psoriasis; and
3 Patient must be reassessed for continuation after 3 doses.

Initiation – plaque psoriasis, treatment-naive
Limited to 6 months treatment
All of the following:
1 Any of the following:
1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10; or
1.2 Patient has plaque psoriasis of the face, or palm of a hand, or sole of a foot; or
1.3 Patient has localised genital or flexural plaque psoriasis with a DLQI score greater than 10; and
2 Patient has received insufficient benefit from (see Note), or has experienced intolerable side effects from, at least 3 of the following at maximum tolerated doses (unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
3 A PASI assessment or DLQI assessment has been completed for the most recent prior treatment course within 1 month of stopping that treatment; and
4 The most recent PASI or DLQI assessment is within 1 month of before the applicattion.

Note: "Insufficient benefit" is defined as: for whole body plaque psoriasis, a PASI score of greater than 10, for plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face,

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

palm of a hand or sole of a foot. As assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation – plaque psoriasis

Re-assessment required after 2 years

Both:

1 Any of the following:
- 1.1 Both:
  - 1.1.1 Patient had "whole body" plaque psoriasis at the start of treatment; and
  - 1.1.2 Either:
    - 1.1.2.1 Patient has a PASI score which is reduced by 75% or more, or is sustained at this level, compared with the pre-treatment baseline; or
    - 1.1.2.2 Patient has a DLQI improvement of 5 or more, compared with the pre-treatment baseline; or
- 1.2 Both:
  - 1.2.1 Patient had plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 1.2.2 Either:
    - 1.2.2.1 Patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, compared to the pre-treatment baseline; or
    - 1.2.2.2 Patient has a reduction of 75% or more in the skin area affected, or sustained at this level, compared to the pre-treatment baseline; or
- 1.3 Both:
  - 1.3.1 Patient had localised genital or flexural plaque psoriasis at the start of treatment; and
  - 1.3.2 Either:
    - 1.3.2.1 Patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, compared to the pre-treatment baseline; or
    - 1.3.2.2 Patient has a DLQI improvement of 5 or more, compared to the pretreatment baseline; and

2 Maximum 50 mg every 7 days.

Initiation – pyoderma gangrenosum*

Both:

1 Patient has received insufficient benefit from 3 months of conventional therapy including a minimum of 3 pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate). Where conventional pharmaceuticals are contraindicated, a 3-month trial has occurred of those that are not contraindicated; and

2 Maximum of 8 doses every 4 months.

Note: Indications marked with * are unapproved indications.

Initiation – Still's disease - adult-onset (AOSD)

Either:

1 Both:
- 1.1 Patient has had a Special Authority approval for adalimumab or tocilizumab for AOSD; and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects; or
  - 1.2.2 Patient has received insufficient benefit to meet the new renewal criteria from at least a 3-month trial of adalimumab or tocilizumab; or

2 All of the following:
- 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
- 2.2 Patient has tried and received insufficient benefit from at least 6 months of corticosteroids at a dose of at least 0.5 mg/kg prednisone-equivalents, NSAIDs and methotrexate, unless contraindicated; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – undifferentiated spondyloarthritis

Re-assessment required after 6 months

All of the following:

1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least 4 joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
2 Patient has received insufficient benefit from at least 3 months of each of methotrexate, sulfasalazine, and leflunomide at maximum tolerated doses, unless contraindicated; and
3 Any of the following:
  3.1 Patient has a CRP level greater than 15 mg/L measured within one month before the application; or
  3.2 Patient has an ESR greater than 25 mm per hour measured within one month before the application; or
  3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy greater than 5 mg per day received for more than three months.

Note: Indications marked with * are unapproved indications.

Continuation – undifferentiated spondyloarthritis

Re-assessment required after 2 years

Both:

1 Either:
  1.1 Following initial treatment, the patient has experienced at least a 50% decrease in active joint count from baseline; or
  1.2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline; and
2 Maximum 50 mg dose every 7 days.

Monoclonal Antibodies

ABCIXIMAB – **Restricted** see terms below

⬇ Inj 2 mg per ml, 5 ml vial

➥ **Restricted** (RS1202)

Initiation

Either:

1 For use in patients with acute coronary syndromes undergoing percutaneous coronary intervention; or
2 For use in patients undergoing intra-cranial intervention.

ADALIMUMAB (AMGEVITA) – **Restricted** see terms below

⬇	Inj 20 mg per 0.4 ml prefilled syringe – **5% DV Oct-22 to 31 Jul 2026**	190.00	1	**Amgevita**
⬇	Inj 40 mg per 0.8 ml prefilled pen – **5% DV Oct-22 to 31 Jul 2026**	375.00	2	**Amgevita**
⬇	Inj 40 mg per 0.8 ml prefilled syringe – **5% DV Oct-22 to 31 Jul 2026**	375.00	2	**Amgevita**

➥ **Restricted** (RS2140)

Initiation – Behcet's disease - severe

Any relevant practitioner

Both:

1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
2 Either:
  2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
  2.2 The patient has severe gastrointestinal, rheumatological and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – Hidradenitis suppurativa

Dermatologist

Re-assessment required after 4 months

All of the following:

1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
3 Patient has 3 or more active lesions; and
4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Continuation – Hidradenitis suppurativa

Any relevant practitioner

Re-assessment required after 2 years

Both:

1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
2 The patient has a DLQI improvement of 4 or more from baseline.

Initiation – Plaque psoriasis - severe chronic

Dermatologist

Re-assessment required after 4 months

Either:

1 Both:
  1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
  1.2 Either:
    1.2.1 Patient has experienced intolerable side effects; or
    1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
2 All of the following:
  2.1 Any of the following:
    2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
    2.1.3 Patient has severe chronic localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a Dermatology Life Quality Index (DLQI) score greater than 10; and
  2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  2.3 A PASI assessment or (DLQI) assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Continuation – Plaque psoriasis - severe chronic

Re-assessment required after 2 years

Any of the following:

1 Both:
  1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
  1.2 Either:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.2.1 The patient has experienced a 75% or more reduction in PASI score, or is sustained at this level, when compared with the pre-treatment baseline value; or
1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or

2 Both:
2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
2.2 Either:
2.2.1 The patient has experienced a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
2.2.2 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or

3 Both:
3.1 Patient had severe chronic localised genital or flexural plaque psoriasis at the start of treatment; and
3.2 Either:
3.2.1 The patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; or
3.2.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing adalimumab.

Initiation – pyoderma gangrenosum
Dermatologist
Both:
1 Patient has pyoderma gangrenosum*; and
2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initiation – Crohn's disease - adults
Any relevant practitioner
Re-assessment required after 6 months
All of the following:
1 Patient has severe active Crohn's disease; and
2 Any of the following:
2.1 Patient has a CDAI score of greater than or equal to 300 or HBI score of greater than or equal to 10; or
2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease - adults
Any relevant practitioner
Re-assessment required after 2 years
Any of the following:
1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced 3 points, from when the patient was initiated on adalimumab; or
2 CDAI score is 150 or less, or HBI is 4 or less; or
3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – Crohn's disease - children

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

1 Paediatric patient has active Crohn's disease; and
2 Either:
  2.1 Patient has a PCDAI score of greater than or equal to 30; or
  2.2 Patient has extensive small intestine disease; and
3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease - children

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
2 PCDAI score is 15 or less; or
3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initiation – Crohn's disease - fistulising

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

1 Patient has confirmed Crohn's disease; and
2 Any of the following:
  2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  2.2 Patient has one or more rectovaginal fistula(e); or
  2.3 Patient has complex peri-anal fistula; and
3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Continuation – Crohn's disease - fistulising

Any relevant practitioner

Re-assessment required after 2 years

Either:

1 The number of open draining fistulae have decreased from baseline by at least 50%; or
2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initiation – Ocular inflammation - chronic

Any relevant practitioner

Re-assessment required after 4 months

Either:

1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
2 Both:
  2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  2.2 Any of the following:
    2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
    2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Continuation – Ocular inflammation - chronic

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

1 The patient has had a good clinical response following 12 weeks' initial treatment; or
2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initiation – Ocular inflammation - severe

Any relevant practitioner

Re-assessment required after 4 months

Either:

1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
2 Both:
  2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  2.2 Any of the following:
    2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Continuation – Ocular inflammation - severe

Any relevant practitioner

Re-assessment required after 2 years

Any of the following:

1 The patient has had a good clinical response following 3 initial doses; or
2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initiation – ankylosing spondylitis

Rheumatologist

Re-assessment required after 6 months

Either:

1 Both:
  1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  1.2 Either:
    1.2.1 The patient has experienced intolerable side effects; or
    1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
2 All of the following:
  2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
2.5 Either:

2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and

2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Continuation – ankylosing spondylitis
Any relevant practitioner
Re-assessment required after 2 years
For applications where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.
Initiation – Arthritis - oligoarticular course juvenile idiopathic
Named specialist or rheumatologist
Re-assessment required after 6 months
Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
1.2 Either:

1.2.1 Patient has experienced intolerable side effects; or
1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or

2 All of the following:

2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
2.3 Either:

2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Continuation – Arthritis - oligoarticular course juvenile idiopathic
Any relevant practitioner
Re-assessment required after 2 years
Either:

1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – Arthritis - polyarticular course juvenile idiopathic

Named specialist or rheumatologist

Re-assessment required after 6 months

Either:

1 Both:
- 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects; or
  - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or

2 All of the following:
- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
  - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Continuation – Arthritis - polyarticular course juvenile idiopathic

Any relevant practitioner

Re-assessment required after 2 years

Either:

1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or

2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - psoriatic

Rheumatologist

Re-assessment required after 6 months

Either:

1 Both:
- 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects; or
  - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for psoriatic arthritis; or

2 All of the following:
- 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
- 2.4 Either:
  - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
  - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.5 Any of the following:
- 2.5.1 Patient has CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.5.2 Patient has an elevated ESR greater than 25 mm per hour; or
- 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation – Arthritis - psoriatic

Any relevant practitioner

Re-assessment required after 2 years

Either:

1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or

2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initiation – Arthritis - rheumatoid

Rheumatologist

Re-assessment required after 6 months

Either:

1 Both:
- 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects; or
  - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or

2 All of the following:
- 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
- 2.5 Either:
  - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin (unless contraindicated); or
  - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomid (unless contraindicated) alone or in combination with methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Continuation – Arthritis - rheumatoid

Any relevant practitioner

Re-assessment required after 2 years

Either:

1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

significant response to treatment in the opinion of the physician; or

2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initiation – Still's disease - adult-onset (AOSD)

Rheumatologist

Either:

1 Both:
- 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for (AOSD); and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
  - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or

2 All of the following:
- 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
- 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initiation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

1 Patient has active ulcerative colitis; and

2 Either:
- 2.1 Patient's SCCAI score is greater than or equal to 4; or
- 2.2 Patient's PUCAI score is greater than or equal to 20; and

3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids; and

4 Surgery (or further surgery) is considered to be clinically inappropriate.

Continuation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 2 years

Either:

1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on biologic therapy; or

2 The PUCAI score has reduced by 10 points or more from the PUCAI score when the patient was initiated on biologic therapy.

Initiation – undifferentiated spondyloarthiritis

Rheumatologist

Re-assessment required after 6 months

All of the following:

1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

2 Patient has tried and not responded to at least three months of each of methotrexate, sulphasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and

3 Any of the following:
- 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or

3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Continuation – undifferentiated spondyloarthiritis

Any relevant practitioner

Re-assessment required after 2 years

Either:

1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initiation – inflammatory bowel arthritis – axial

Rheumatologist

Re-assessment required after 6 months

All of the following:

1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and

2 Patient has axial inflammatory pain for six months or more; and

3 Patient is unable to take NSAIDs; and

4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and

5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and

6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Continuation – inflammatory bowel arthritis – axial

Any relevant practitioner

Re-assessment required after 2 years

Where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initiation – inflammatory bowel arthritis – peripheral

Rheumatologist

Re-assessment required after 6 months

All of the following:

1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and

2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and

3 Patient has tried and not experienced a response to at least three months of methotrexate, or azathioprine at a maximum tolerated dose (unless contraindicated); and

4 Patient has tried and not experienced a response to at least three months of sulphasalazine at a maximum tolerated dose (unless contraindicated); and

5 Any of the following:

5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or

5.2 Patient has an ESR greater than 25 mm per hour; or

5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation – inflammatory bowel arthritis – peripheral
Any relevant practitioner
Re-assessment required after 2 years
Either:

1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
2 Patient demonstrates at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) – **Restricted** see terms below

⬇	Inj 20 mg per 0.2 ml prefilled syringe	595.50	2	Humira
⬇	Inj 40 mg per 0.4 ml prefilled syringe	595.50	2	Humira
⬇	Inj 40 mg per 0.4 ml prefilled pen	595.50	2	HumiraPen

➥ **Restricted** (RS1922)
Initiation – Behcet's disease – severe
Any relevant practitioner
Re-assessment required after 6 months
All of the following:

1 Either:
  1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
2 Patient has received a maximum of 6 months treatment with Amgevita; and
3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Behcet's disease – severe
Any relevant practitioner
Re-assessment required after 6 months
Both:

1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Hidradenitis suppurativa
Dermatologist or Practitioner on the recommendation of a dermatologist
Re-assessment required after 6 months
All of the following:

1 Either:
  1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
2 Patient has received a maximum of 6 months treatment with Amgevita; and
3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Continuation – Hidradenitis suppurativa
Dermatologist or Practitioner on the recommendation of a dermatologist
Re-assessment required after 6 months
All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initiation – Psoriasis - severe chronic plaque

Dermatologist or Practitioner on the recommendation of a dermatologist
Re-assessment required after 6 months
All of the following:

1 Either:
  1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
2 Patient has received a maximum of 6 months treatment with Amgevita; and
3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Psoriasis - severe chronic plaque

Dermatologist or Practitioner on the recommendation of a dermatologist
Re-assessment required after 6 months
Both:

1 Either:
  1.1 Both:
    1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    1.1.2 Either:
      1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
      1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  1.2 Both:
    1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    1.2.2 Either:
      1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Pyoderma gangrenosum

Dermatologist
Re-assessment required after 6 months
All of the following:

1 Either:
  1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

2 Patient has received a maximum of 6 months treatment with Amgevita; and

3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

4 A maximum of 8 doses.

Continuation – Pyoderma gangrenosum

Dermatologist

Re-assessment required after 6 months

Both:

1 The patient has demonstrated clinical improvement and continues to require treatment; and

2 A maximum of 8 doses.

Initiation – Crohn's disease - adult

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

All of the following:

1 Any of the following:

1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or

1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or

1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and

2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Crohn's disease - adult

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

Both:

1 Any of the following:

1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or

1.2 CDAI score is 150 or less; or

1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Crohn's disease - children

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

All of the following:

1 Any of the following:

1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or

1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or

1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and

2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Crohn's disease - children

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

Both:

1 Any of the following:
   1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
   1.2 PCDAI score is 15 or less; or
   1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Crohn's disease - fistulising

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

All of the following:

1 Any of the following:
   1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
   1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
   1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Crohn's disease - fistulising

Gastroenterologist or Practitioner on the recommendation of a gastroenterologist

Re-assessment required after 6 months

Both:

1 Either:
   1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
   1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Ocular inflammation – chronic

Any relevant practitioner

Re-assessment required after 12 months

All of the following:

1 Any of the following:
   1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
   1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
   1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation – Ocular inflammation – chronic

Any relevant practitioner

Re-assessment required after 12 months

Both:

1 Any of the following:
   1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
   1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
   1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Ocular inflammation – severe

Any relevant practitioner

Re-assessment required after 12 months

All of the following:

1 Any of the following:
   1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
   1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
   1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Ocular inflammation – severe

Any relevant practitioner

Re-assessment required after 12 months

Both:

1 Any of the following:
   1.1 The patient has had a good clinical response following 3 initial doses; or
   1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
   1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – ankylosing spondylitis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

1 Either:
   1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
   1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
2 Patient has received a maximum of 6 months treatment with Amgevita; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – ankylosing spondylitis
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Re-assessment required after 6 months
Both:

1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Arthritis – oligoarticular course juvenile idiopathic
Named specialist or rheumatologist
Re-assessment required after 6 months
All of the following:

1 Either:
  1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
2 Patient has received a maximum of 6 months treatment with Amgevita; and
3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Continuation – Arthritis – oligoarticular course juvenile idiopathic
Named specialist or rheumatologist
Re-assessment required after 6 months
For patients that demonstrate at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - polyarticular course juvenile idiopathic
Named specialist or rheumatologist
Re-assessment required after 6 months
All of the following:

1 Either:
  1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
  1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
2 Patient has received a maximum of 6 months treatment with Amgevita; and
3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Continuation – Arthritis - polyarticular course juvenile idiopathic
Named specialist or rheumatologist
Re-assessment required after 6 months
For patients that demonstrate at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initiation – Arthritis - psoriatic
Named specialist or rheumatologist
Re-assessment required after 6 months
All of the following:

1 Either:
  1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

treatment; or

1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

2 Patient has received a maximum of 6 months treatment with Amgevita; and

3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Continuation – Arthritis - psoriatic

Named specialist or rheumatologist

Re-assessment required after 6 months

Both:

1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initiation – Arthritis – rheumatoid

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

1 Either:

1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

2 Patient has received a maximum of 6 months treatment with Amgevita; and

3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and

4 Either:

4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or

4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Continuation – Arthritis – rheumatoid

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

Both:

1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and

2 Either:

2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or

2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initiation – Still's disease – adult-onset (AOSD)

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 6 months

All of the following:

1 Either:

1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or

1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 Patient has received a maximum of 6 months treatment with Amgevita; and
3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Continuation – Still's disease – adult-onset (AOSD)
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Re-assessment required after 6 months
the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT – **Restricted** see terms below
⬇ Inj 40 mg per ml, 0.1 ml vial ..... 1,250.00 1 Eylea
➥ **Restricted** (RS2148)
Initiation – Wet Age Related Macular Degeneration
Re-assessment required after 3 months
Either:
1 All of the following:
1.1 Any of the following:
1.1.1 Wet age-related macular degeneration (wet AMD); or
1.1.2 Polypoidal choroidal vasculopathy; or
1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
1.2 Either:
1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
1.3 There is no structural damage to the central fovea of the treated eye; and
1.4 Patient has not previously been treated with ranibizumab or faricimab for longer than 3 months; or
2 Either:
2.1 Patient has current approval to use ranibizumab or faricimab for treatment of wAMD and was found to be intolerant within 3 months; or
2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Continuation – Wet Age Related Macular Degeneration
Re-assessment required after 12 months
All of the following:
1 Documented benefit must be demonstrated to continue; and
2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
3 There is no structural damage to the central fovea of the treated eye.

Initiation – Diabetic Macular Oedema
Re-assessment required after 4 months
All of the following:
1 Patient has centre involving diabetic macular oedema (DMO); and
2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
3 Patient has reduced visual acuity between 6/9 – 6/36 with functional awareness of reduction in vision; and
4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
5 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
6 Patient has not previously been treated with faricimab for longer than 3 months.

Continuation – Diabetic Macular Oedema
Re-assessment required after 12 months
All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 There is stability or two lines of Snellen visual acuity gain; and
2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
4 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

BASILIXIMAB – **Restricted** see terms below

⬇ Inj 20 mg vial ..... 2,560.00 | 1 | Simulect

➥ **Restricted** (RS1203)

Initiation

For use in solid organ transplants.

BENRALIZUMAB – **Restricted** see terms below

⬇ Inj 30 mg per ml, 1 ml prefilled pen ..... 3,539.00 | 1 | Fasenra

➥ **Restricted** (RS1920)

Initiation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 12 months

All of the following:

1 Patient must be aged 12 years or older; and
2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
4 Patient has a blood eosinophil count of greater than 0.5 × 10^9 cells/L in the last 12 months; and
5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
6 Either:
  6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
  6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
7 Treatment is not to be used in combination with subsidised mepolizumab; and
8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
9 Either:
  9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
  9.2 Both:
    9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
    9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Continuation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 2 years

Both:

1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 Either:

2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with benralizumab; or

2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

BEVACIZUMAB – **Restricted** see terms below

⬇	Inj 25 mg per ml, 4 ml vial – **10% DV Aug-25 to 31 Aug 2028**	69.00	1	**Vegzelma**
⬇	Inj 25 mg per ml, 16 ml vial – **10% DV Aug-25 to 31 Aug 2028**	276.00	1	**Vegzelma**

➥ **Restricted** (RS2111)

Initiation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

Either:

1 Patient is currently on treatment with bevacizumab, and met all remaining criteria prior to commencing treatment; or

2 All of the following:

2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and

2.2 Patient has preserved liver function (Child-Pugh A); and

2.3 Transarterial chemoembolisation (TACE) is unsuitable; and

2.4 Any of the following:

2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or

2.4.2 Patient received funded lenvatinib before 1 March 2025; or

2.4.3 Both:

2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and

2.4.3.2 No disease progression since initiation of lenvatinib; and

2.5 Patient has an ECOG performance status of 0-2; and

2.6 To be given in combination with atezolizumab.

Continuation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

no evidence of disease progression.

Initiation – advanced or metastatic ovarian cancer

Re-assessment required after 4 months

All of the following:

1 Either:

1.1 The patient has FIGO Stage IV epithelial ovarian, fallopian tube, or primary peritoneal cancer; or

1.2 Both:

1.2.1 The patient has previously untreated advanced (FIGO Stage IIIB or IIIC) epithelial ovarian, fallopian tube, or primary peritoneal cancer; and

1.2.2 Either:

1.2.2.1 Debulking surgery is inappropriate; or

1.2.2.2 The cancer is sub-optimally debulked (maximum diameter of any gross residual disease greater than 1cm); and

2 Bevacizumab to be administered at a maximum dose of 15 mg/kg every three weeks; and

3 18 weeks concurrent treatment with chemotherapy is planned.

Continuation – advanced or metastatic ovarian cancer

Re-assessment required after 4 months

no evidence of disease progression.

Initiation – Recurrent Respiratory Papillomatosis

Re-assessment required after 12 months

All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Maximum of 6 doses; and
2 The patient has recurrent respiratory papillomatosis; and
3 The treatment is for intra-lesional administration.

Continuation – Recurrent Respiratory Papillomatosis
Re-assessment required after 12 months
All of the following:

1 Maximum of 6 doses; and
2 The treatment is for intra-lesional administration; and
3 There has been a reduction in surgical treatments or disease regrowth as a result of treatment.

Initiation – Ocular Conditions
Either:

1 Ocular neovascularisation; or
2 Exudative ocular angiopathy.

BEVACIZUMAB (OCULAR) – **Restricted** see terms below

⬇	Inj 25 mg per ml, 4 ml vial	600.00	1	Avastin
⬇	Inj 25 mg per ml, 16 ml vial			

➥ **Restricted** (RS2156)
Initiation – ocular conditions
Either:

1 Ocular neovascularisation; or
2 Exudative ocular angiopathy.

BRENTUXIMAB VEDOTIN – **Restricted** see terms below

⬇	Inj 50 mg vial	5,275.18	1	Adcetris

➥ **Restricted** (RS2183)
Initiation – CD30 positive systemic anaplastic large-cell lymphoma
Limited to 12 months treatment
Either:

1 Patient is currently on treatment with brentuximab vedotin and met all the following criteria prior to commencing treatment; or
2 All of the following:
  2.1 Patient has CD30 positive systemic anaplastic large-cell lymphoma; and
  2.2 Patient must have histological confirmation of CD30 expression; and
  2.3 Patient must not have received prior treatment with curative intent chemotherapy for this condition; and
  2.4 Treatment must be in combination with cyclophosphamide, anthracycline, and steroids for a maximum of 8 cycles; and
  2.5 Brentuximab vedotin is to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Initiation – relapsed/refractory Hodgkin lymphoma
Re-assessment required after 6 months
All of the following:

1 Either:
  1.1 Both:
    1.1.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma after two or more lines of chemotherapy; and
    1.1.2 Patient is ineligible for autologous stem cell transplant; or
  1.2 Both:
    1.2.1 Patient has relapsed/refractory CD30-positive Hodgkin lymphoma; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.2.2 Patient has previously undergone autologous stem cell transplant; and

2 Patient has not previously received funded brentuximab vedotin; and

3 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and

4 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Continuation – relapsed/refractory Hodgkin lymphoma

Re-assessment required after 9 months

All of the following:

1 Patient has achieved a partial or complete response to brentuximab vedotin after 6 treatment cycles; and

2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and

3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

Initiation – relapsed/refractory anaplastic large cell lymphoma

Re-assessment required after 9 months

All of the following:

1 Patient has relapsed/refractory CD30-positive systemic anaplastic large cell lymphoma; and

2 Patient has an ECOG performance status of 0-1; and

3 Patient has not previously received brentuximab vedotin; and

4 Response to brentuximab vedotin treatment is to be reviewed after a maximum of 6 treatment cycles; and

5 Brentuximab vedotin to be administered at doses no greater than 1.8 mg/kg every 3 weeks.

Continuation – relapsed/refractory anaplastic large cell lymphoma

Re-assessment required after 9 months

All of the following:

1 Patient has experienced a partial or complete response to brentuximab vedotin after 6 treatment cycles; and

2 Treatment remains clinically appropriate and the patient is benefitting from treatment and treatment is being tolerated; and

3 Patient is to receive a maximum of 16 total cycles of brentuximab vedotin treatment.

CETUXIMAB – **Restricted** see terms below

⬇	Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
⬇	Inj 5 mg per ml, 100 ml vial	1,820.00	1	Erbitux

➥ **Restricted** (RS2064)

Initiation – head and neck cancer, locally advanced

All of the following:

1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and

2 Cisplatin is contraindicated or has resulted in intolerable side effects; and

3 Patient has an ECOG performance score of 0-2; and

4 To be administered in combination with radiation therapy.

Initiation – colorectal cancer, metastatic

Re-assessment required after 6 months

All of the following:

1 Patient has metastatic colorectal cancer located on the left side of the colon (see Note); and

2 There is documentation confirming disease is RAS and BRAF wild-type; and

3 Patient has an ECOG performance score of 0-2; and

4 Patient has not received prior funded treatment with cetuximab; and

5 Either:

5.1 Cetuximab is to be used in combination with chemotherapy; or

5.2 Chemotherapy is determined to not be in the best interest of the patient based on clinician assessment.

Continuation – colorectal cancer, metastatic

Re-assessment required after 6 months

No evidence of disease progression.

Note: Left-sided colorectal cancer comprises of the distal one-third of the transverse colon, the splenic flexure, the descending colon, the sigmoid colon, or the rectum.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
FARICIMAB – **Restricted** see terms below			
⬇ Inj 120 mg per ml, 0.24 ml vial	1,565.00	1	Vabysmo

➥ **Restricted** (RS2149)

Initiation – Diabetic macular oedema

Re-assessment required after 4 months

All of the following:

1 Patient has centre involving diabetic macular oedema (DMO); and
2 Patient's disease is nonresponsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
3 Patient has reduced visual acuity between 6/9 – 6/36 with functional awareness of reduction in vision; and
4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
5 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
6 Patient has not previously been treated with aflibercept for longer than 3 months.

Continuation – Diabetic macular oedema

Re-assessment required after 12 months

All of the following:

1 There is stability or two lines of Snellen visual acuity gain; and
2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
4 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Initiation – Wet age related macular degeneration

Re-assessment required after 3 months

All of the following:

1 Any of the following:
  1.1 Wet age-related macular degeneration (wet AMD); or
  1.2 Polypoidal choroidal vasculopathy; or
  1.3 Choroidal neovascular membrane from causes other than wet AMD; and
2 Either:
  2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
  2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
3 There is no structural damage to the central fovea of the treated eye; and
4 Patient has not previously been treated with ranibizumab or aflibercept for longer than 3 months.

Continuation – Wet age related macular degeneration

Re-assessment required after 12 months

Both:

1 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
2 There is no structural damage to the central fovea of the treated eye.

GEMTUZUMAB OZOGAMICIN – **Restricted** see terms below			
⬇ Inj 5 mg vial	12,973.00	1	Mylotarg

➥ **Restricted** (RS1923)

Initiation

All of the following:

1 Patient has not received prior chemotherapy for this condition; and
2 Patient has de novo CD33-positive acute myeloid leukaemia; and
3 Patient does not have acute promyelocytic leukaemia; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
5 Patient is being treated with curative intent; and
6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m^2 body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB – **Restricted** see terms below

⬇ Inj 100 mg ..... 428.00 1 Remicade

➥ **Restricted** (RS2178)

Initiation – Graft vs host disease

Patient has steroid-refractory acute graft vs. host disease of the gut.

Initiation – arthritis - rheumatoid

Re-assessment required after 6 months

All of the following:

1 Patient has had a Special Authority approval for adalimumab or etanercept for rheumatoid arthritis; and
2 Either:
  2.1 Patient has experienced intolerable side effects; or
  2.2 Patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; and
3 Following initial induction doses, maximum dose 3mg/kg every 8 weeks.

Continuation – arthritis - rheumatoid

Re-assessment required after 2 years

Both:

1 Either:
  1.1 Following initial treatment, the patient has experienced at least a 50% decrease in active joint count from baseline; or
  1.2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline; and
2 Maximum dose 3 mg/kg every 8 weeks.

Initiation – ankylosing spondylitis

Re-assessment required after 6 months

All of the following:

1 Patient has had a Special Authority approval for adalimumab or etanercept for ankylosing spondylitis; and
2 Either:
  2.1 Patient has experienced intolerable side effects; or
  2.2 Patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; and
3 Following initial induction doses, maximum dose 5mg/kg every 6-8 weeks.

Continuation – ankylosing spondylitis

Re-assessment required after 2 years

Both:

1 BASDAI has improved from pre-treatment baseline either by at least 4 points on a 10-point scale, or by at least 50%; and
2 Maximum dose 5 mg/kg every 6-8 weeks.

Initiation – arthritis - psoriatic

Re-assessment required after 6 months

All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient has had a Special Authority approval for adalimumab or etanercept or secukinumab for psoriatic arthritis; and
2 Either:
  2.1 Patient has experienced intolerable side effects; or
  2.2 Patient has received insufficient benefit to meet the renewal criteria for psoriatic arthritis; and
3 Following initial induction doses, maximum dose 5mg/kg every 8 weeks.

Continuation – arthritis - psoriatic

Re-assessment required after 2 years

Both:

1 Either:
  1.1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
  1.2 At least a continuing 30% improvement in active joint count from baseline; and
2 Maximum dose 5 mg/kg every 8 weeks.

Initiation – ocular inflammation - severe*

Re-assessment required after 4 months

Either:

1 Both:
  1.1 Patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  1.2 Either:
    1.2.1 Patient has experienced intolerable side effects; or
    1.2.2 Patient has received insufficient benefit to meet the renewal criteria for severe ocular inflammation; or
2 Both:
  2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  2.2 Any of the following:
    2.2.1 Treatment with high-dose IV corticosteroids followed by high dose oral corticosteroids has been ineffective at controlling symptoms; or
    2.2.2 Patient developed new inflammatory symptoms while receiving high dose corticosteroids; or
    2.2.3 Patient is aged under 8 years and treatment with high dose oral corticosteroids and other immunosuppressants has been ineffective at controlling symptoms; or
    2.2.4 High dose corticosteroids are contraindicated.

Note: Indications marked with * are unapproved indications.

Continuation – ocular inflammation - severe*

Re-assessment required after 2 years

Any of the following:

1 Patient has received a good clinical response following 3 initial doses; or
2 Following each 2 year treatment period, the patient has experienced a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
3 Following each 2 year treatment period, the patient has a sustained corticosteroid sparing effect, allowing reduction in prednisone to < 10mg daily, or corticosteroid drops less than twice daily if under 18 years old.

Notes: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.
Indications marked with * are unapproved indications.

Initiation – ocular inflammation - chronic*

Re-assessment required after 4 months

Either:

1 Both:
  1.1 Patient has had a Special Authority approval for adalimumab for chronic ocular inflammation; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.2 Either:

1.2.1 Patient has experienced intolerable side effects; or
1.2.2 Patient has received insufficient benefit to meet the renewal criteria for chronic ocular inflammation; or

2 Both:

2.1 Patient has severe uveitis with a severe risk of vision loss uncontrolled by treatment with corticosteroids and other immunosuppressants; and
2.2 Any of the following:

2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has been ineffective or are contraindicated; or
2.2.2 Patient is under 18 years and treatment with methotrexate has been ineffective, is contraindicated or is not tolerated at a therapeutic dose; or
2.2.3 Patient is under 8 years and treatment with corticosteroids or methotrexate has been ineffective, is contraindicated or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Note: Indications marked with * are unapproved indications.

Continuation – ocular inflammation - chronic*

Re-assessment required after 2 years

Any of the following:

1 Patient has received a good clinical response following 3 initial doses; or
2 Following each 2 year treatment period, the patient has experienced a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
3 Following each 2 year treatment period, the patient has a sustained corticosteroid sparing effect, allowing reduction in prednisone to < 10mg daily, or corticosteroid drops less than twice daily if under 18 years old.

Notes: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.
Indications marked with * are unapproved indications.

Initiation – Pulmonary sarcoidosis

Both:

1 Patient has life-threatening pulmonary sarcoidosis that is refractory to other treatments; and
2 Treatment is to be prescribed by, or has been recommended by, a physician with expertise in the treatment of pulmonary sarcoidosis.

Initiation – Crohn's disease (adults)

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

1 Patient has active Crohn's disease; and
2 Any of the following:

2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and

3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation – Crohn's disease (adults)
Any relevant practitioner
Re-assessment required after 2 years
Both:

1 Any of the following:
  1.1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on infliximab; or
  1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  1.3 The patient has demonstrated an adequate response to treatment but CDAI score and/or HBI score cannot be assessed; and
2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – Crohn's disease (children)
Any relevant practitioner
Re-assessment required after 6 months
All of the following:

1 Paediatric patient has active Crohn's disease; and
2 Either:
  2.1 Patient has a PCDAI score of greater than or equal to 30; or
  2.2 Patient has extensive small intestine disease; and
3 Patient has tried but experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and corticosteroids.

Continuation – Crohn's disease (children)
Any relevant practitioner
Re-assessment required after 2 years
Both:

1 Any of the following:
  1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  1.2 PCDAI score is 15 or less; or
  1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – fistulising Crohn's disease
Gastroenterologist
Re-assessment required after 6 months
Both:

1 Patient has confirmed Crohn's disease; and
2 Any of the following:
  2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  2.2 Patient has one or more rectovaginal fistula(e); or
  2.3 Patient has complete peri-anal fistula.

Continuation – fistulising Crohn's disease
Any relevant practitioner
Re-assessment required after 2 years
Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Either:

1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or

1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and

2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – acute fulminant ulcerative colitis

Gastroenterologist

Limited to 6 weeks treatment

Both:

1 Patient has acute, fulminant ulcerative colitis; and

2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Continuation – fulminant ulcerative colitis

Any relevant practitioner

Re-assessment required after 2 years

Both:

1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and

2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

Initiation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

1 Patient has active ulcerative colitis; and

2 Either:

2.1 Patients SCCAI is greater than or equal to 4; or

2.2 Patients PUCAI score is greater than or equal to 20; and

3 Patient has experienced an inadequate response to, or intolerable side effects from, prior therapy with immunomodulators and systemic corticosteroids.

Continuation – ulcerative colitis

Any relevant practitioner

Re-assessment required after 2 years

Both:

1 Either:

1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or

1.2 The PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and

2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – plaque psoriasis

Re-assessment required after 6 months

Either:

1 Both:
- 1.1 Patient had a Special Authority approval for adalimumab, etanercept or secukinumab for plaque psoriasis; and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects; or
  - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for plaque psoriasis; or

2 All of the following:
- 2.1 Any of the following:
  - 2.1.1 Patient has "whole body" plaque psoriasis with a PASI score of greater than 10; or
  - 2.1.2 Patient has plaque psoriasis of the face, or palm of a hand or sole of a foot; or
  - 2.1.3 Patient has localised genital or flexural plaque psoriasis with a DLQI score greater than 10; and
- 2.2 Patient has received insufficient benefit (see Note) or has experienced intolerable side effects from at least 3 of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for the most recent prior treatment course within 1 month of stopping that treatment; and
- 2.4 The most recent PASI assessment is within 1 month before the application.

Note: "Insufficient benefit" is defined as: for whole body plaque psoriasis, a PASI score of greater than 10, for plaque psoriasis of the face, hand, foot, genital or flexural areas at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation – plaque psoriasis

Re-assessment required after 2 years

Both:

1 Any of the following:
- 1.1 Both:
  - 1.1.1 Patient had "whole body" plaque psoriasis at the start of treatment; and
  - 1.1.2 Patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab baseline; or
- 1.2 Both:
  - 1.2.1 Patient had plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 1.2.2 Either:
    - 1.2.2.1 Patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the pre-infliximab baseline; or
    - 1.2.2.2 Patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab baseline; or
- 1.3 Both:
  - 1.3.1 Patient had localised genital or flexural plaque psoriasis at the start of treatment; and
  - 1.3.2 Either:
    - 1.3.2.1 Patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline; or
    - 1.3.2.2 Patient has a DLQI improvement of 5 or more, as compared to the pre-infliximab baseline; and

2 Maximum dose 5 mg/kg every 8 weeks.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – neurosarcoidosis

Neurologist

Re-assessment required after 18 months

All of the following:

1 Biopsy consistent with diagnosis of neurosarcoidosis; and
2 Patient has CNS involvement; and
3 Patient has steroid-refractory disease; and
4 Either:
  4.1 IV cyclophosphamide has been tried; or
  4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Continuation – neurosarcoidosis

Neurologist

Re-assessment required after 18 months

Either:

1 A withdrawal period has been tried and the patient has relapsed; or
2 All of the following:
  2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  2.2 There has been a marked reduction in prednisone dose; and
  2.3 Either:
    2.3.1 There has been an improvement in MRI appearances; or
    2.3.2 Marked improvement in other symptomology.

Initiation – Behcet disease

All of the following:

1 Patient has severe Behcet disease which is significantly impacting their quality of life; and
2 Either:
  2.1 Patient has severe ocular, neurological and/or vasculitic symptoms and has received insufficient benefit from 1 or more treatment(s) appropriate for the particular symptom(s); or
  2.2 Patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has received insufficient benefit from 2 or more treatments appropriate for the particular symptom(s); and
3 Following initial loading doses, maximum dose 5mg/kg every 8 weeks.

Initiation – pyoderma gangrenosum*

Both:

1 Patient has received insufficient benefit from 3 months of conventional therapy including a minimum of 3 pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate). Where conventional pharmaceuticals are contraindicated, a 3-month trial has occurred of those that are not contraindicated; and
2 Maximum of 8 doses every 4 months.

Note: Indications marked with * are unapproved indications.

Initiation – Inflammatory bowel arthritis (axial)

Re-assessment required after 6 months

All of the following:

1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
2 Patient has had axial inflammatory pain for six months or more; and
3 Patient is unable to take NSAIDs; and
4 Patient has unequivocal sacroiliitis demonstrated by radiological imaging or MRI; and
5 Patient has not experienced an adequate response to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

6 Patient has a BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment .

Continuation – Inflammatory bowel arthritis (axial)

Re-assessment required after 2 years

Where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10-point scale, or an improvement in BASDAI of 50%, whichever is less.

Initiation – Inflammatory bowel arthritis (peripheral)

Re-assessment required after 6 months

All of the following:

1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
3 Patient has tried and not experienced a response to at least three months of methotrexate or azathioprine at a maximum tolerated dose (unless contraindicated); and
4 Patient has tried and not experienced a response to at least three months of sulfasalazine at a maximum tolerated dose (unless contraindicated); and
5 Any of the following:
  5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Continuation – Inflammatory bowel arthritis (peripheral)

Re-assessment required after 2 years

Either:

1 Following initial treatment, patient has experienced at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
2 Patient has experienced at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

Initiation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and
2 The individual has received insufficient benefit from use of corticosteroids; and
3 Infliximab is to be administered at up to 5mg/kg for up to four doses.

Continuation – immune checkpoint inhibitor toxicity in malignancy*

Any relevant practitioner

Re-assessment required after 4 months

Both:

1 The individual has shown clinical improvement and ongoing treatment is required; and
2 Infliximab is to be administered at up to 5mg/kg for up to a total of 8 doses.

Note: Indications marked with * are unapproved indications.

INOTUZUMAB OZOGAMICIN – **Restricted** see terms on the next page

⬇	Inj 1 mg vial	14,457.00	1	Besponsa

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS2112)

Initiation

Re-assessment required after 4 months

All of the following:

1 Patient has relapsed or refractory CD22-positive B-cell acute lymphoblastic leukaemia/lymphoma, including minimal residual disease; and
2 Patient has ECOG performance status of 0-2; and
3 Either:
  3.1 Both:
    3.1.1 Patient has Philadelphia chromosome positive B-Cell ALL; and
    3.1.2 Patient has previously received a tyrosine kinase inhibitor; or
  3.2 Patient has received one prior line of treatment involving intensive chemotherapy; and
4 Treatment is to be administered for a maximum of 3 cycles.

Continuation

Re-assessment required after 4 months

All of the following:

1 Patient is not proceeding to a stem cell transplant; and
2 Either:
  2.1 Patient has experienced complete disease response; or
  2.2 Patient has experienced complete remission with incomplete haematological recovery; and
3 Treatment with inotuzumab ozogamicin is to cease after a total duration of 6 cycles.

MEPOLIZUMAB – **Restricted** see terms below

⬇	Inj 100 mg prefilled pen	1,638.00	1	Nucala
⬇	Inj 100 mg vial			

➥ **Restricted** (RS2024)

Initiation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 12 months

All of the following:

1 Patient must be aged 12 years or older; and
2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
4 Patient has a blood eosinophil count of greater than 0.5 × 10^9 cells/L in the last 12 months; and
5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
6 Either:
  6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
  6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
7 Treatment is not to be used in combination with subsidised benralizumab; and
8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

the first dose to assess response to treatment; and

9 Either:

9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or

9.2 Both:

9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and

9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Continuation – Severe eosinophilic asthma

Respiratory physician or clinical immunologist

Re-assessment required after 2 years

Both:

1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and

2 Either:

2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or

2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

Initiation – eosinophilic granulomatosis with polyangiitis

Re-assessment required after 12 months

All of the following:

1 The patient has eosinophilic granulomatosis with polyangiitis; and

2 The patient has trialled and not received adequate benefit from at least one of the following for at least three months (unless contraindicated to all): azathioprine, cyclophosphamide, leflunomide, methotrexate, mycophenolate, or rituximab; and

3 Either:

3.1 The patient has trialled prednisone for a minimum of three months and is unable to maintain disease control at doses below 7.5 mg per day; or

3.2 Corticosteroids are contraindicated.

Continuation – eosinophilic granulomatosis with polyangiitis

Re-assessment required after 12 months

Patient has no evidence of clinical disease progression.

OBINUTUZUMAB – **Restricted** see terms below

⬇	Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva

➥ **Restricted** (RS2192)

Initiation – previously untreated chronic lymphocytic leukaemia in combination with venetoclax

Either:

1 Individual is currently on treatment with obinutuzumab and venetoclax and met all of the following criteria prior to commencing treatment; or

2 Both:

2.1 Individual has previously untreated chronic lymphocytic leukaemia; and

2.2 Obinutuzumab is to be administered at a maximum cumulative dose of 8,000 mg and in combination with venetoclax for a maximum of 6 (28-day) cycles of treatment.

Initiation

Limited to 6 months treatment

All of the following:

1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and

2 The patient is obinutuzumab treatment naive; and

3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

(CIRS) or reduced renal function (creatinine clearance < 70mL/min); and

4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and

5 Patient has good performance status; and

6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* greater than or equal to 1.5×10^9/L and platelets greater than or equal to 75×10^9/L

Initiation – follicular / marginal zone lymphoma

Re-assessment required after 9 months

All of the following:

1 Either:

1.1 Patient has follicular lymphoma; or

1.2 Patient has marginal zone lymphoma; and

2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*; and

3 Patient has an ECOG performance status of 0-2; and

4 Patient has been previously treated with no more than four chemotherapy regimens; and

5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Continuation – follicular / marginal zone lymphoma

Re-assessment required after 24 months

All of the following:

1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and

2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and

3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB – **Restricted** see terms below

⬇	Inj 150 mg prefilled syringe	450.00	1	Xolair
⬇	Inj 150 mg vial	450.00	1	Xolair

➥ **Restricted** (RS1652)

Initiation – severe asthma

Clinical immunologist or respiratory specialist

Re-assessment required after 6 months

All of the following:

1 Patient must be aged 6 years or older ; and

2 Patient has a diagnosis of severe asthma; and

3 Past or current evidence of atopy, documented by skin prick testing or RAST; and

4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and

5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and

6 Either:

6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

12 months, unless contraindicated or not tolerated; or

6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and

7 Patient has an Asthma Control Test (ACT) score of 10 or less; and

8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Continuation – severe asthma

Respiratory specialist

Re-assessment required after 6 months

Both:

1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and

2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Initiation – severe chronic spontaneous urticaria

Clinical immunologist or dermatologist

Re-assessment required after 6 months

All of the following:

1 Patient must be aged 12 years or older; and

2 Either:

2.1 Both:

2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and

2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; and

3 Any of the following:

3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or

3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or

3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and

4 Either:

4.1 Treatment to be stopped if inadequate response* following 4 doses; or

4.2 Complete response* to 6 doses of omalizumab.

Continuation – severe chronic spontaneous urticaria

Clinical immunologist or dermatologist

Re-assessment required after 6 months

Either:

1 Patient has previously had a complete response* to 6 doses of omalizumab; or

2 Both:

2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and

2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB – **Restricted** see terms below

⬇ Inj 100 mg per ml, 1 ml vial ..... 1,700.00 1 Synagis

➥ **Restricted** (RS2193)

Initiation

Re-assessment required after 12 months

Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Palivizumab to be administered during the annual RSV season; and
2 Either:
2.1 All of the following:
2.1.1 Infant was born in the last 12 months; and
2.1.2 Infant was born at less than 32 weeks zero days' gestation; and
2.1.3 Infant is to be administered palivizumab within a single calendar year; or
2.2 Both:
2.2.1 Child was born in the last 24 months; and
2.2.2 Any of the following:
2.2.2.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
2.2.2.2 Both:
2.2.2.2.1 Child has haemodynamically significant heart disease; and
2.2.2.2.2 Any of the following:
2.2.2.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
2.2.2.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
2.2.2.2.2.3 Child has severe pulmonary hypertension (see Note C); or
2.2.2.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
2.2.2.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
2.2.2.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Continuation

Re-assessment required after 6 months

All of the following:

1 Palivizumab to be administered during the annual RSV season; and
2 Child was born in the last 24 months; and
3 Any of the following:
3.1 Child has severe lung, airway, neurological or neuromuscular disease that requires ongoing ventilatory/respiratory support (see Note A) in the community; or
3.2 Both:
3.2.1 Child has haemodynamically significant heart disease; and
3.2.2 Any of the following:
3.2.2.1 Child has unoperated simple congenital heart disease with significant left to right shunt (see Note B); or
3.2.2.2 Child has unoperated or surgically palliated complex congenital heart disease; or
3.2.2.3 Child has severe pulmonary hypertension (see Note C); or
3.2.2.4 Child has moderate or severe left ventricular (LV) failure (see Note D); or
3.3 Child has severe combined immune deficiency, confirmed by an immunologist, but has not received a stem cell transplant; or
3.4 Child has inborn errors of immunity (see Note E) that increase susceptibility to life-threatening viral respiratory infections, confirmed by an immunologist.

Notes:

a) Ventilatory/respiratory support includes those on home oxygen, CPAP/VPAP and those with tracheostomies in situ managed at home

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

b) Child requires/will require heart failure medication, and/or child has significant pulmonary hypertension, and/or infant will require surgical palliation/definitive repair within the next 3 months
c) Mean pulmonary artery pressure more than 25 mmHg
d) LV Ejection Fraction less than 40%
e) Inborn errors of immunity include, but are not limited to, IFNAR deficiencies

PERTUZUMAB – **Restricted** see terms below

⬇	Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta

➥ **Restricted** (RS1995)

Initiation

Re-assessment required after 12 months

All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
2 Either:
  2.1 Patient is chemotherapy treatment naive; or
  2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
3 The patient has good performance status (ECOG grade 0-1); and
4 Pertuzumab to be administered in combination with trastuzumab; and
5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
6 Pertuzumab to be discontinued at disease progression.

Continuation

Re-assessment required after 12 months

Either:

1 Both:
  1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or
2 All of the following:
  2.1 Patient has previously discontinued treatment with pertuzumab and trastuzumab for reasons other than severe toxicity or disease progression; and
  2.2 Patient has signs of disease progression; and
  2.3 Disease has not progressed during previous treatment with pertuzumab and trastuzumab.

PERTUZUMAB WITH TRASTUZUMAB – **Restricted** see terms below

⬇	Inj 600 mg with trastuzumab 600 mg, 10 ml vial	7,707.00	1	Phesgo
⬇	Inj 1,200 mg with trastuzumab 600 mg, 15 ml vial	12,894.00	1	Phesgo

➥ **Restricted** (RS2152)

Initiation

Re-assessment required after 12 months

Either:

1 Both:
  1.1 The individual has received an initial Special Authority approval for intravenous pertuzumab and trastuzumab for metastatic breast cancer; and
  1.2 Pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 600 mg pertuzumab with 600 mg trastuzumab every three weeks (or equivalent); or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 All of the following:
- 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2.2 Either:
  - 2.2.1 Patient is chemotherapy treatment naïve; or
  - 2.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 2.3 The patient has good performance status (ECOG grade 0-1); and
- 2.4 Loading dose of pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 1200 mg pertuzumab with 600 mg trastuzumab, respectively; and
- 2.5 Maintenance doses of pertuzumab with trastuzumab to be administered subcutaneously at a maximum dose of 600 mg pertuzumab with 600 mg trastuzumab every three weeks (or equivalent); and
- 2.6 Pertuzumab with trastuzumab to be discontinued at disease progression.

Continuation

Re-assessment required after 12 months

Either:

1 Both:
- 1.1 The individual has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 1.2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab; or

2 All of the following:
- 2.1 Individual has previously discontinued treatment with pertuzumab with trastuzumab for reasons other than severe toxicity or disease progression; and
- 2.2 Individual has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with pertuzumab with trastuzumab.

RANIBIZUMAB – **Restricted** see terms below

⬇ Inj 10 mg per ml, 0.23 ml vial

⬇ Inj 10 mg per ml, 0.3 ml vial

➥ **Restricted** (RS2151)

Initiation – Wet Age Related Macular Degeneration

Re-assessment required after 3 months

Either:

1 All of the following:
- 1.1 Any of the following:
  - 1.1.1 Wet age-related macular degeneration (wet AMD); or
  - 1.1.2 Polypoidal choroidal vasculopathy; or
  - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
- 1.2 Either:
  - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
  - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
- 1.3 There is no structural damage to the central fovea of the treated eye; and
- 1.4 Patient has not previously been treated with aflibercept or faricimab for longer than 3 months; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 Patient has current approval to use aflibercept or faricimab for treatment of wAMD and was found to be intolerant within 3 months.

Continuation – Wet Age Related Macular Degeneration

Re-assessment required after 12 months

All of the following:

1 Documented benefit must be demonstrated to continue; and
2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
3 There is no structural damage to the central fovea of the treated eye.

RITUXIMAB (MABTHERA) – **Restricted** see terms below

⬇	Inj 10 mg per ml, 10 ml vial	1,075.50	2	Mabthera
⬇	Inj 10 mg per ml, 50 ml vial	2,688.30	1	Mabthera

➥ **Restricted** (RS2180)

Initiation – arthritis - rheumatoid - prior TNF inhibitor use

Limited to 4 months treatment

All of the following:

1 Patient has had a Special Authority approval for etanercept or adalimumab for rheumatoid arthritis; and
2 Either:
  2.1 Patient has experienced intolerable side effects; or
  2.2 Following at least a 4 month trial of adalimumab or etanercept, the renewal criteria for rheumatoid arthritis were not met; and
3 Maximum of two 1000 mg infusions given two weeks apart.

Initiation – arthritis - rheumatoid - TNF inhibitors contraindicated

Limited to 4 months treatment

All of the following:

1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
2 Patient has had rheumatoid arthritis (either confirmed by radiologic imaging, or the patient is CCP antibody positive); and
3 Disease has not responded to at least three months of methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose, unless contraindicated; and
4 Disease has not responded to at least 3 months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses), unless contraindicated; and
5 Either:
  5.1 Disease has not responded to at least 3 months of methotrexate in combination with the maximum tolerated dose of cyclosporin, unless contraindicated; or
  5.2 Disease has not responded to at least 3 months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate, unless contraindicated; and
6 Either:
  6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 joints; or
  6.2 Patient has persistent symptoms of poorly controlled and active disease in at least 4 joints from the following: wrist, elbow, knee, ankle, shoulder, or hip; and
7 Either:
  7.1 Patient has CRP greater than 15 mg/L measured within one month before the application; or
  7.2 CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day received for more than 3 months; and
8 Maximum of two 1000 mg infusions given two weeks apart.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation – arthritis - rheumatoid - re-treatment for people who have experienced a partial response to rituximab

Re-assessment required after 12 months

All of the following:

1 Any of the following:
  1.1 Following the initial course of rituximab the patient experienced between a 30% and 50% decrease in active joint count from baseline; or
  1.2 Following the second course of rituximab the patient experienced at least a 50% decrease in active joint count from baseline; or
  1.3 Following the third and subsequent courses of rituximab, the patient experienced at least a continuing 30% improvement in active joint count from baseline; and
2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
3 Maximum of two 1000 mg infusions given two weeks apart.

Continuation – arthritis - rheumatoid - re-treatment for people who experience a response to rituximab

Re-assessment required after 12 months

All of the following:

1 Either:
  1.1 Following the initial course of rituximab infusions the patient experienced at least a 50% decrease in active joint count from baseline; or
  1.2 Following the second and subsequent courses of rituximab, the patient experienced at least a continuing 30% improvement in active joint count from baseline; and
2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
3 Maximum of two 1000 mg infusions per course given two weeks apart.

RITUXIMAB (RIXIMYO) – **Restricted** see terms below

⬇	Inj 10 mg per ml, 10 ml vial	275.33	2	Riximyo
⬇	Inj 10 mg per ml, 50 ml vial	688.20	1	Riximyo

➥ **Restricted** (RS2194)

Initiation – haemophilia with inhibitors

Haematologist

Any of the following:

1 Patient has mild congenital haemophilia complicated by inhibitors; or
2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
3 Patient has acquired haemophilia.

Continuation – haemophilia with inhibitors

Haematologist

All of the following:

1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
2 An initial response lasting at least 12 months was demonstrated; and
3 Patient now requires repeat treatment.

Initiation – post-transplant

Both:

1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Continuation – post-transplant

All of the following:

1 The patient has had a rituximab treatment-free interval of 12 months or more; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initiation – indolent, low-grade lymphomas or hairy cell leukaemia*

Re-assessment required after 9 months

Either:

1 Both:
  1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
  1.2 To be used for a maximum of 6 treatment cycles; or
2 Both:
  2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
  2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Continuation – indolent, low-grade lymphomas or hairy cell leukaemia*

Re-assessment required after 12 months

All of the following:

1 The patient has had a rituximab treatment-free interval of 12 months or more; and
2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initiation – aggressive CD20 positive NHL

Either:

1 All of the following:
  1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  1.3 To be used for a maximum of 8 treatment cycles; or
2 Both:
  2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Continuation – aggressive CD20 positive NHL

All of the following:

1 The patient has had a rituximab treatment-free interval of 12 months or more; and
2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia.

Initiation – Chronic lymphocytic leukaemia

Re-assessment required after 12 months

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
2 Any of the following:
  2.1 The patient is rituximab treatment naive; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.2 Either:

2.2.1 The patient is chemotherapy treatment naive; or
2.2.2 Both:

2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or

2.3 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; and

3 The patient has good performance status; and
4 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
5 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Continuation – Chronic lymphocytic leukaemia

Re-assessment required after 12 months

Both:

1 Either:

1.1 The patient's disease has relapsed and rituximab treatment is to be used in combination with funded venetoclax; or
1.2 All of the following:

1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
1.2.3 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustin; and

2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initiation – severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 8 weeks

All of the following:

1 Patient has cold haemagglutinin disease*; and
2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
3 The total rituximab dose used would not exceed the equivalent of 375 mg/m^2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation – severe cold haemagglutinin disease (CHAD)

Haematologist

Re-assessment required after 8 weeks

Either:

1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m^2 weekly for 4 weeks) is now planned; or

2 All of the following:

2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and

2.2 An initial response lasting at least 12 months was demonstrated; and

2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initiation – warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 8 weeks

All of the following:

1 Patient has warm autoimmune haemolytic anaemia*; and

2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m^2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – warm autoimmune haemolytic anaemia (warm AIHA)

Haematologist

Re-assessment required after 8 weeks

Either:

1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m^2 weekly for 4 weeks) is now planned; or

2 All of the following:

2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and

2.2 An initial response lasting at least 12 months was demonstrated; and

2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initiation – immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 8 weeks

All of the following:

1 Either:

1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or

1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and

2 Any of the following:

2.1 Treatment with steroids and splenectomy have been ineffective; or

2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or

2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m^2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – immune thrombocytopenic purpura (ITP)

Haematologist

Re-assessment required after 8 weeks

Either:

1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m^2 weekly for 4 weeks) is now planned; or

2 All of the following:

2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and

2.2 An initial response lasting at least 12 months was demonstrated; and

2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initiation – thrombotic thrombocytopenic purpura (TTP)

Haematologist

All of the following:

1 The total rituximab dose per cycle would not exceed the equivalent of 375 mg/m^2 of body surface area per week for a total of 4 weeks; and

2 Each treatment cycle at least 6 months apart; and

3 Either:

3.1 Patient has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or

3.2 Patient has acute idiopathic TTP* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Initiation – pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Continuation – pure red cell aplasia (PRCA)

Haematologist

Re-assessment required after 6 weeks

Patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initiation – ANCA associated vasculitis

Re-assessment required after 8 weeks

All of the following:

1 Patient has been diagnosed with ANCA associated vasculitis*; and

2 The total rituximab dose would not exceed the equivalent of 375 mg/m^2 of body-surface area per week for a total of 4 weeks; and

3 Any of the following:

3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or

3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3.3 Cyclophosphamide and methotrexate are contraindicated; or
3.4 Patient is a female of child-bearing potential; or
3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Continuation – ANCA associated vasculitis

Re-assessment required after 8 weeks

All of the following:

1 Patient has been diagnosed with ANCA associated vasculitis*; and
2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
3 The total rituximab dose would not exceed the equivalent of 375 mg/m^2 of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initiation – treatment refractory systemic lupus erythematosus (SLE)

Rheumatologist or nephrologist

All of the following:

1 Patient has severe, immediately life- or organ-threatening SLE*; and
2 The condition has been refractory to treatment with corticosteroids at a dose of at least 1 mg/kg unless contraindicated; and
3 The condition has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
4 Initial treatment maximum of four 1000 mg infusions; and
5 Treatment for relapse following initial partial response to rituximab up to a maximum of two 1000 mg infusions every 6 months.

Note: Indications marked with * are unapproved indications.

Initiation – Antibody-mediated organ transplant rejection

Patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initiation – ABO-incompatible organ transplant

Patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initiation – Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)

Re-assessment required after 8 weeks

All of the following:

1 Patient is a child with SDNS* or FRNS*; and
2 Treatment with corticosteroids, ciclosporin, and mycophenolate for at least 3 months for each agent has been ineffective, not tolerated, or is contraindicated; and
3 The total rituximab dose used would not exceed the equivalent of 375 mg/m^2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)

Re-assessment required after 8 weeks

All of the following:

1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
2 Treatment with rituximab was previously successful and has demonstrated sustained response for > 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
3 The total rituximab dose used would not exceed the equivalent of 375 mg/m^2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – Steroid resistant nephrotic syndrome (SRNS)

Re-assessment required after 8 weeks

All of the following:

1 Patient is a child with SRNS* and treatment with corticosteroids, ciclosporin and tacrolimus for at least 3 months for each agent has been ineffective, not tolerated, or is contraindicated; and
2 Genetic causes of nephrotic syndrome have been excluded; and
3 The total rituximab dose per cycle would not exceed the equivalent of 375 mg/m^2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Continuation – Steroid resistant nephrotic syndrome (SRNS)

Re-assessment required after 8 weeks

All of the following:

1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
3 The total rituximab dose used would not exceed the equivalent of 375 mg/m^2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initiation – Neuromyelitis Optica Spectrum Disorder (NMOSD)

All of the following:

1 Cumulative dose up to 1500 mg/m^2 body surface area up to 2000 mg total per cycle; and
2 Either:
  2.1 Patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms with supporting clinical investigations); or
  2.2 All of the following:
    2.2.1 Patient has experienced a breakthrough attack of NMOSD; and
    2.2.2 Patient is receiving treatment with mycophenolate unless contraindicated or not tolerated; and
    2.2.3 Patient is receiving treatment with corticosteroids unless contraindicated or not tolerated; and
3 Each treatment cycle at least 6 months apart.

Note: Indications marked with * are unapproved indications.

Initiation – refractory myasthenia gravis*

Re-assessment required after 2 years

Both:

1 Cumulative dose up to 1500 mg/m^2 body surface area up to 2000 mg total per cycle; and
2 Either:
  2.1 Treatment with corticosteroids and at least one other immunosuppressant for a minimum period of 12 months has been ineffective; or
  2.2 Both:
    2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Note: Indications marked with * are unapproved indications.

Continuation – refractory myasthenia gravis*

Re-assessment required after 2 years

All of the following:

1 Cumulative dose up to 1500 mg/m^2 body surface area up to 2000 mg total per cycle; and
2 An initial response lasting at least 12 months was demonstrated; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3 Either:
- 3.1 Patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
- 3.2 Both:
  - 3.2.1 Patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
  - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Note: Indications marked with * are unapproved indications.

Initiation – antisynthetase syndrome

All of the following:
1 Patient has severe, immediately life- or organ-threatening disease, including interstitial lung disease; and
2 Either:
- 2.1 Treatment with at least 3 immunosuppressants (oral corticosteroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has been ineffective controlling active disease; or
- 2.2 Rapid treatment is required for life threatening complications; and

3 Maximum of two 1000 mg infusions every 6 months.

Initiation – graft versus host disease

All of the following:
1 Patient has refractory graft versus host disease following transplant; and
2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
3 The total rituximab dose used would not exceed the equivalent of 375 mg/m^2 of body surface area per week for a total of 4 weeks.

Initiation – chronic inflammatory demyelinating polyneuropathy (CIPD)*

Neurologist

All of the following:
1 Either:
- 1.1 Both:
  - 1.1.1 Treatment with corticosteroids and intravenous immunoglobulin and/or plasma exchange has been ineffective controlling active disease, is not tolerated, or is contraindicated; and
  - 1.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) is not tolerated or has been ineffective controlling active disease. If an immunosuppressant is contraindicated, a trial has occurred of one of those which is not contraindicated (unless all are contraindicated); or
- 1.2 Rapid treatment is required for life threatening complications; and

2 Cumulative dose up to 1500 mg/m^2 body surface area up to 2000 mg total per cycle; and
3 Each treatment cycle at least 6 months apart.

Note: Indications marked with * are unapproved indications.

Initiation – anti-NMDA receptor autoimmune encephalitis*

All of the following:
1 Either:
- 1.1 Both:
  - 1.1.1 Treatment with corticosteroids and intravenous immunoglobulin and/or plasma exchange has been ineffective controlling active disease, is not tolerated or is contraindicated; and
  - 1.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has been ineffective controlling active disease, is not tolerated or is contraindicated; or
- 1.2 Rapid treatment is required for life threatening complications; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 Cumulative dose up to 1500 mg/m^2 body surface area up to 2000 mg total per cycle; and
3 Each treatment cycle at least 6 months apart.

Note: Indications marked with * are unapproved indications.

Initiation – CD20+ low grade or follicular B-cell NHL

Re-assessment required after 9 months

Either:

1 Both:
  1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  1.2 To be used for a maximum of 6 treatment cycles; or
2 Both:
  2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
  2.2 To be used for a maximum of 6 treatment cycles.

Continuation – CD20+ low grade or follicular B-cell NHL

Re-assessment required after 24 months

Both:

1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m^2 every 8 weeks (maximum of 12 cycles).

Initiation – Membranous nephropathy

All of the following:

1 Either:
  1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
  1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m^2; and
2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures that include (unless contraindicated or the patient has experienced intolerable side effects) renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents; and
3 The total rituximab dose per cycle would not exceed the equivalent of 375mg/m^2 of body surface area per week for a total of 4 weeks; and
4 Subsequent retreatment only for disease relapse or after partial response.

Note: Indications marked with * are unapproved indications.

Initiation – B-cell acute lymphoblastic leukaemia/lymphoma*

Limited to 2 years treatment

All of the following:

1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
3 The total rituximab dose would not exceed the equivalent of 375 mg/m^2 per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initiation – desensitisation prior to transplant

Limited to 6 weeks treatment

Both:

1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
2 Patient would receive no more than two doses at 375 mg/m^2 of body-surface area.

Note: Indications marked with * are unapproved indications.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – pemiphigus*

Dermatologist or relevant specialist

Re-assessment required after 6 months

Either:

1 All of the following:
- 1.1 Patient has severe rapidly progressive pemphigus; and
- 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
- 1.3 Any of the following:
  - 1.3.1 Skin involvement is at least 5% body surface area; or
  - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
  - 1.3.3 Involvement of two or more mucosal sites; or

2 Both:
- 2.1 Patient has pemphigus; and
- 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Continuation – pemiphigus*

Dermatologist or relevant specialist

Re-assessment required after 6 months

Both:

1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and

2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

Initiation – immunoglobulin G4-related disease (IgG4-RD*)

Re-assessment required after 6 weeks

All of the following:

1 Patient has confirmed diagnosis of IgG4-RD*; and

2 Either:
- 2.1 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs for at least 3 months has been ineffective in lowering corticosteroid dose below 5 mg per day (prednisone equivalent) without relapse; or
- 2.2 Treatment with corticosteroids and/or disease modifying anti-rheumatic drugs is contraindicated or associated with evidence of toxicity or intolerance; and

3 Total rituximab dose used should not exceed a maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

Continuation – immunoglobulin G4-related disease (IgG4-RD*)

Re-assessment required after 12 months

All of the following:

1 Either:
- 1.1 Treatment with rituximab for IgG4-RD* was previously successful and patient's disease has demonstrated sustained response, but the condition has relapsed; or
- 1.2 Patient is receiving maintenance treatment for IgG4-RD*; and

2 Rituximab re-treatment not to be given within 6 months of previous course of treatment; and

3 Maximum of two 1000 mg infusions of rituximab given two weeks apart.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB – **Restricted** see terms on the next page

⬇	Inj 150 mg per ml, 1 ml prefilled syringe	799.50	1	Cosentyx
		1,599.00	2	Cosentyx

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS2182)

Initiation – plaque psoriasis

Re-assessment required after 6 months

Either:

1 All of the following:
- 1.1 Any of the following:
  - 1.1.1 Patient has "whole body" plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.1.2 Patient has plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; or
  - 1.1.3 Patient has localised genital or flexural plaque psoriasis where the plaques or lesions have been present for at least 6 months from the time of initial diagnosis, and with a DLQI score greater than 10; and
- 1.2 Patient has received insufficient benefit (see Note) or has experienced intolerable side effects from at least 3 of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 1.3 A PASI assessment or DLQI assessment has been completed for the most recent prior treatment course, within 1 month of stopping that treatment; and
- 1.4 The most recent PASI or DQLI assessment is within 1 month before the application; or

2 All of the following:
- 2.1 Patient has had a Special Authority approval for adalimumab, etanercept, or infliximab, for plaque psoriasis; and
- 2.2 Either:
  - 2.2.1 Patient has experienced intolerable side effects; or
  - 2.2.2 Patient has received insufficient benefit to meet the renewal criteria for plaque psoriasis; and
- 2.3 A PASI assessment or DLQI assessment has been completed for the most recent prior treatment within 1 month of stopping that treatment; and
- 2.4 The most recent PASI or DQLI assessment is within 1 month before the application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Insufficient benefit" is defined as: for whole body plaque psoriasis, a PASI score of greater than 10; for plaque psoriasis of the face, hand, foot, genital or flexural areas, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and for the face, palm of a hand or sole of a foot the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. As assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Continuation – plaque psoriasis

Re-assessment required after 6 months

Both:

1 Either:
- 1.1 Both:
  - 1.1.1 Patient's PASI score has reduced by 75% or more compared to pre-secukinumab baseline; and
  - 1.1.2 Patient has a DLQI improvement of 5 or more compared to pre-secukinumab baseline; or
- 1.2 Both:
  - 1.2.1 Patient had localised genital or flexural plaque psoriasis at the start of treatment; and
  - 1.2.2 Either:
    - 1.2.2.1 Patient has experienced a reduction of 75% or more in the skin area affected, or sustained at this level, compared to the pre-secukinumab baseline; or
    - 1.2.2.2 Patient has a DLQI improvement of 5 or more, compared to pre-secukinumab baseline; and

2 Maximum dose 300 mg monthly.

Initiation – ankylosing spondylitis, second-line biologic

Re-assessment required after 3 months

Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient has had a Special Authority approval for adalimumab or etanercept for ankylosing spondylitis; and
2 Either:
2.1 Patient has experienced intolerable side effects; or
2.2 Patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis.

Continuation – ankylosing spondylitis, second-line biologic
Re-assessment required after 6 months
Both:
1 BASDAI has improved from the pre-secukinumab baseline either by at least 4 points on a 10-point scale, or by at least 50%, whichever is less; and
2 Maximum dose 300 mg monthly.

Initiation – arthritis - psoriatic
Re-assessment required after 6 months
Either:
1 Both:
1.1 Patient has had a Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
1.2 Either:
1.2.1 Patient has experienced intolerable side effects; or
1.2.2 Patient has received insufficient benefit to meet the renewal criteria for psoriatic arthritis; or
2 All of the following:
2.1 Patient has received insufficient benefit from at least 3 months of methotrexate at a maximum tolerated dose unless contraindicated; and
2.2 Patient has received insufficient benefit from at least 3 months of sulfasalazine or leflunomide at maximum tolerated doses unless contraindicated; and
2.3 Either:
2.3.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 joints; or
2.3.2 Patient has persistent symptoms of poorly controlled and active disease in at least 4 joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
2.4 Any of the following:
2.4.1 CRP greater than 15 mg/L measured within one month before the application; or
2.4.2 ESR greater than 25 mm per hour measured within one month before the application; or
2.4.3 ESR and CRP not measured as patient is receiving prednisone therapy greater than 5 mg per day received for more than 3 months.

Continuation – arthritis - psoriatic
Re-assessment required after 6 months
Both:
1 Either:
1.1 Following initial treatment, at least a 50% decrease in active joint count from baseline; or
1.2 At least a continuing 30% improvement in active joint count from baseline; and
2 Maximum dose 300 mg monthly.

SILTUXIMAB – **Restricted** see terms below

⬇	Inj 100 mg vial	770.57	1	Sylvant
⬇	Inj 400 mg vial	3,082.33	1	Sylvant

➥ **Restricted** (RS1525)
Initiation
Haematologist or rheumatologist
Re-assessment required after 6 months
All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Continuation
Haematologist or rheumatologist
Re-assessment required after 12 months
The treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB – **Restricted** see terms below

⬇	Inj 20 mg per ml, 4 ml vial	220.00	1	Actemra
⬇	Inj 20 mg per ml, 10 ml vial	550.00	1	Actemra
⬇	Inj 20 mg per ml, 20 ml vial	1,100.00	1	Actemra

➥ **Restricted** (RS2125)

Initiation – cytokine release syndrome
Therapy limited to 3 doses
Either:
1 Both:
1.1 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
1.2 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
2 All of the following:
2.1 The patient is enrolled in the Malaghan Institute of Medical Research ENABLE trial programme; and
2.2 The patient has developed CRS or Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) following CAR T-Cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS or ICANS for CAR T-cell therapy at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initiation – previous use
Any relevant practitioner
Limited to 6 months treatment
Both:
1 Patient was being treated with tocilizumab prior to 1 February 2019; and
2 Any of the following:
2.1 rheumatoid arthritis; or
2.2 systemic juvenile idiopathic arthritis; or
2.3 adult-onset Still's disease; or
2.4 polyarticular juvenile idiopathic arthritis; or
2.5 idiopathic multicentric Castleman's disease.

Initiation – Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Limited to 6 months treatment
All of the following:
1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
2 Either:
2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
3 Either:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
3.2 Both:
3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
3.2.2 Either:
3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initiation – Rheumatoid Arthritis
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Re-assessment required after 6 months
All of the following:
1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
2 Tocilizumab is to be used as monotherapy; and
3 Either:
3.1 Treatment with methotrexate is contraindicated; or
3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
4 Either:
4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
5 Either:
5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
6 Either:
6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initiation – systemic juvenile idiopathic arthritis
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Re-assessment required after 6 months
Both:
1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initiation – adult-onset Still's disease
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Re-assessment required after 6 months
Either:
1 Both:
1.1 Either:
1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and

1.2 Either:

1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or

1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or

2 All of the following:

2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and

2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and

2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initiation – polyarticular juvenile idiopathic arthritis

Rheumatologist or Practitioner on the recommendation of a rheumatologist

Re-assessment required after 4 months

Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and

1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or

2 All of the following:

2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and

2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and

2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

2.4 Any of the following:

2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or

2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or

2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initiation – idiopathic multicentric Castleman's disease

Haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist

Re-assessment required after 6 months

All of the following:

1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and

2 Treatment with an adequate trial of corticosteroids has proven ineffective; and

3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initiation – moderate to severe COVID-19

Therapy limited to 1 dose

All of the following:

1 Patient has confirmed (or probable) COVID-19; and

2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and

3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and

4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and

5 Tocilizumab is not to be administered in combination with barcitinib.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation – Rheumatoid Arthritis
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Re-assessment required after 6 months
Either:

1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Continuation – systemic juvenile idiopathic arthritis
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Re-assessment required after 6 months
Either:

1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Continuation – adult-onset Still's disease
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Re-assessment required after 6 months
the patient has a sustained improvement in inflammatory markers and functional status.

Continuation – polyarticular juvenile idiopathic arthritis
Rheumatologist or Practitioner on the recommendation of a rheumatologist
Re-assessment required after 6 months
Both:

1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
2 Either:
  2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Continuation – idiopathic multicentric Castleman's disease
Haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist
Re-assessment required after 12 months
the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

Initiation – immune checkpoint inhibitor toxicity in malignancy*
Any relevant practitioner
Re-assessment required after 4 months
All of the following:

1 The individual requires treatment for moderate to severe autoimmune toxicity following immune checkpoint inhibitor treatment for malignancy; and
2 The individual has received insufficient benefit from use of corticosteroids; and
3 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Continuation – immune checkpoint inhibitor toxicity in malignancy*
Any relevant practitioner
Re-assessment required after 4 months
Both:

1 The individual has shown clinical improvement and ongoing treatment is required; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 Tocilizumab is to be administered at a maximum dose of 8 mg/kg fortnightly.

Note: Indications marked with * are unapproved indications.

TRASTUZUMAB (HERZUMA) – **Restricted** see terms below

⬇ Inj 150 mg vial – **5% DV Jun-24 to 31 May 2027** ..... 100.00 1 **Herzuma**

⬇ Inj 440 mg vial – **5% DV Jun-24 to 31 May 2027** ..... 293.35 1 **Herzuma**

➥ **Restricted** (RS2005)

Initiation – early breast cancer

Limited to 12 months treatment

Both:

1 The patient has early breast cancer expressing HER-2 IHC 3+ or ISH + (including FISH or other current technology); and

2 Maximum cumulative dose of 106 mg/kg (12 months' treatment).

Continuation – early breast cancer*

Re-assessment required after 12 months

Either:

1 All of the following:

1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

1.2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and

1.3 Any of the following:

1.3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or

1.3.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; or

1.3.3 he cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and

1.4 Either:

1.4.1 Trastuzumab will not be given in combination with pertuzumab; or

1.4.2 All of the following:

1.4.2.1 Trastuzumab to be administered in combination with pertuzumab; and

1.4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and

1.4.2.3 The patient has good performance status (ECOG grade 0-1); and

1.5 Trastuzumab to be discontinued at disease progression; or

2 All of the following:

2.1 Patient has previously discontinued treatment with trastuzumab in the metastatic setting for reasons other than severe toxicity or disease progression; and

2.2 Patient has signs of disease progression; and

2.3 Disease has not progressed during previous treatment with trastuzumab.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer

Initiation – metastatic breast cancer

Re-assessment required after 12 months

All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

2 Either:

2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.2 The patient discontinued lapatinib within 3 months due to intolerable side effects and the cancer did not progress whilst on lapatinib; and

3 Either:

3.1 Trastuzumab will not be given in combination with pertuzumab; or

3.2 All of the following:

3.2.1 Trastuzumab to be administered in combination with pertuzumab; and

3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and

3.2.3 The patient has good performance status (ECOG grade 0-1); and

4 Trastuzumab to be discontinued at disease progression.

Continuation – metastatic breast cancer

Re-assessment required after 12 months

Either:

1 All of the following:

1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

1.2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and

1.3 Trastuzumab to be discontinued at disease progression; or

2 All of the following:

2.1 Patient has previously discontinued treatment with trastuzumab for reasons other than severe toxicity or disease progression; and

2.2 Patient has signs of disease progression; and

2.3 Disease has not progressed during previous treatment with trastuzumab.

Initiation – gastric, gastro-oesophageal junction and oesophageal cancer

Re-assessment required after 12 months

Both:

1 The patient has locally advanced or metastatic gastric, gastro-oesophageal junction or oesophageal cancer expressing HER-2 IHC 2+ FISH+ or IHC3+ (or other current technology); and

2 Patient has an ECOG score of 0-2.

Continuation – gastric, gastro-oesophageal junction and oesophageal cancer

Re-assessment required after 12 months

Both:

1 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and

2 Trastuzumab to be discontinued at disease progression.

TRASTUZUMAB DERUXTECAN – **Restricted** see terms below

⬇	Inj 100 mg per ml, 1 ml vial	2,550.00	1	Enhertu

➥ **Restricted** (RS2082)

Initiation

Re-assessment required after 6 months

All of the following:

1 Patient has metastatic breast cancer expressing HER-2 IHC3+ or ISH+ (including FISH or other current technology); and

2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and

3 Either:

3.1 The patient has received prior therapy for metastatic disease; or

3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

4 Patient has a good performance status (ECOG 0-1); and
5 Patient has not received prior funded trastuzumab deruxtecan treatment; and
6 Treatment to be discontinued at disease progression.

Continuation

Re-assessment required after 6 months

Both:

1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab deruxtecan; and
2 Treatment to be discontinued at disease progression.

Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

TRASTUZUMAB EMTANSINE – **Restricted** see terms below

⬇	Inj 100 mg vial	2,320.00	1	Kadcyla
⬇	Inj 160 mg vial	3,712.00	1	Kadcyla

➥ **Restricted** (RS2083)

Initiation – early breast cancer

All of the following:

1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
2 Documentation of pathological invasive residual disease in the breast and/or axiliary lymph nodes following completion of surgery; and
3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
4 Disease has not progressed during neoadjuvant therapy; and
5 Patient has left ventricular ejection fraction of 45% or greater; and
6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
7 Trastuzumab emtansine to be discontinued at disease progression; and
8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initiation – metastatic breast cancer

Re-assessment required after 6 months

All of the following:

1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
3 Either:
  3.1 The patient has received prior therapy for metastatic disease*; or
  3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
4 Patient has a good performance status (ECOG 0-1); and
5 Either:
  5.1 Patient does not have symptomatic brain metastases; or
  5.2 Patient has brain metastases and has received prior local CNS therapy; and
6 Either:
  6.1 Patient has not received prior funded trastuzumab emtansine or trastuzumab deruxtecan treatment; or
  6.2 Both:
    6.2.1 Patient has discontinued trastuzumab deruxtecan due to intolerance; and
    6.2.2 The cancer did not progress while on trastuzumab deruxtecan; and
7 Treatment to be discontinued at disease progression.

Continuation – metastatic breast cancer

Re-assessment required after 6 months

Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

USTEKINUMAB – **Restricted** see terms below

⬇	Inj 130 mg vial 4,162.00	1	Stelara
⬇	Inj 90 mg per ml, 1 ml prefilled syringe 4,162.00	1	Stelara

➥ **Restricted** (RS1942)

Initiation – Crohn's disease - adults

Re-assessment required after 6 months

Either:

1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
2 Both:
  2.1 Patient has active Crohn's disease; and
  2.2 Either:
    2.2.1 Patient has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    2.2.2 Both:
      2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Continuation – Crohn's disease - adults

Re-assessment required after 12 months

Both:

1 Any of the following:
  1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
  1.2 CDAI score is 150 or less, or HBI is 4 or less; or
  1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
2 Ustekinumab to be administered at a dose no greater than 90 mg every 8 weeks.

Initiation – Crohn's disease - children*

Re-assessment required after 6 months

Either:

1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or
2 Both:
  2.1 Patient has active Crohn's disease; and
  2.2 Either:
    2.2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
    2.2.2 Both:
      2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for Crohn's disease; and
      2.2.2.2 Other biologics for Crohn's disease are contraindicated.

Note: Indication marked with * is an unapproved indication.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation – Crohn's disease - children*

Re-assessment required after 12 months

Both:

1 Any of the following:
- 1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
- 1.2 PCDAI score is 15 or less; or
- 1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and

2 Ustekinumab to administered at a dose no greater than 90 mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initiation – ulcerative colitis

Re-assessment required after 6 months

Either:

1 Patient is currently on treatment with ustekinumab commenced prior to 1 February 2023 and met all remaining criteria (criterion 2) below at the time of commencing treatment; or

2 Both:
- 2.1 Patient has active ulcerative colitis; and
- 2.2 Either:
  - 2.2.1 Patient has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
  - 2.2.2 Both:
    - 2.2.2.1 Patient meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
    - 2.2.2.2 Other biologics for ulcerative colitis are contraindicated.

Continuation – ulcerative colitis

Re-assessment required after 12 months

Both:

1 Either:
- 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
- 1.2 PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy*; and

2 Ustekinumab will be used at a dose no greater than 90 mg intravenously every 8 weeks.

Note: Criterion marked with * is for an unapproved indication.

VEDOLIZUMAB – **Restricted** see terms below

⬇	Inj 300 mg vial	3,313.00	1	Entyvio

➥ **Restricted** (RS1943)

Initiation – Crohn's disease - adults

Re-assessment required after 6 months

All of the following:

1 Patient has active Crohn's disease; and

2 Any of the following:
- 2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
- 2.2 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
- 2.3 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
- 2.4 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
- 2.5 Patient has an ileostomy or colostomy, and has intestinal inflammation; and

3 Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
3.3 Immunomodulators and corticosteroids are contraindicated.

Continuation – Crohn's disease - adults
Re-assessment required after 2 years
Both:
1 Any of the following:
1.1 CDAI score has reduced by 100 points, or HBI score has reduced by 3 points, from when the patient was initiated on biologic therapy; or
1.2 CDAI score is 150 or less, or HBI is 4 or less; or
1.3 The patient has experienced an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed; and
2 Vedolizumab to administered at a dose no greater than 300 mg every 8 weeks.

Initiation – Crohn's disease - children*
Re-assessment required after 6 months
All of the following:
1 Paediatric patient has active Crohn's disease; and
2 Any of the following:
2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
2.2 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
2.3 Patient has extensive small intestine disease; and
3 Any of the following:
3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Continuation – Crohn's disease - children*
Re-assessment required after 2 years
Both:
1 Any of the following:
1.1 PCDAI score has reduced by 10 points from when the patient was initiated on biologic therapy; or
1.2 PCDAI score is 15 or less; or
1.3 The patient has experienced an adequate response to treatment, but CDAI score cannot be assessed; and
2 Vedolizumab to administered at a dose no greater than 300mg every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Initiation – ulcerative colitis
Re-assessment required after 6 months
All of the following:
1 Patient has active ulcerative colitis; and
2 Any of the following:
2.1 Patient has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria (unless contraindicated); or
2.2 Patient has a SCCAI score is greater than or equal to 4; or
2.3 Patient's PUCAI score is greater than or equal to 20*; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3 Any of the following:
- 3.1 Patient has tried but experienced an inadequate response to (including lack of initial response and/or loss of initial response) from prior therapy with immunomodulators and corticosteroids; or
- 3.2 Patient has experienced intolerable side effects from immunomodulators and corticosteroids; or
- 3.3 Immunomodulators and corticosteroids are contraindicated.

Note: Indication marked with * is an unapproved indication.

Continuation – ulcerative colitis

Re-assessment required after 2 years

Both:

1 Either:
- 1.1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on biologic therapy; or
- 1.2 The PUCAI score has reduced by 10 points or more from the PUCAI score since initiation on biologic therapy *; and

2 Vedolizumab will be used at a dose no greater than 300 mg intravenously every 8 weeks.

Note: Indication marked with * is an unapproved indication.

Programmed Cell Death-1 (PD-1) Inhibitors

ATEZOLIZUMAB – **Restricted** see terms below

⬇ Inj 60 mg per ml, 20 ml vial ..... 9,503.00 1 Tecentriq

➥ **Restricted** (RS2184)

Initiation – non-small cell lung cancer second line monotherapy

Medical oncologist

Re-assessment required after 4 months

All of the following:

1 Patient has locally advanced or metastatic non-small cell lung cancer; and
2 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
3 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR, ROS-1 or ALK tyrosine kinase unless not possible to ascertain; and
4 Patient has an ECOG 0-2; and
5 Patient has documented disease progression following treatment with at least two cycles of platinum-based chemotherapy; and
6 Atezolizumab is to be used as monotherapy at a dose of 1200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
7 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – non-small cell lung cancer second line monotherapy

Medical oncologist

Re-assessment required after 4 months

All of the following:

1 Any of the following:
- 1.1 Patient's disease has had a complete response to treatment; or
- 1.2 Patient's disease has had a partial response to treatment; or
- 1.3 Patient has stable disease; and

2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
3 No evidence of disease progression; and
4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
5 Atezolizumab to be used at a maximum dose of 1200 mg every three weeks (or equivalent); and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

6 Treatment with atezolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

Either:

1 Patient is currently on treatment with atezolizumab and met all remaining criteria prior to commencing treatment; or

2 All of the following:

2.1 Patient has locally advanced or metastatic, unresectable hepatocellular carcinoma; and

2.2 Patient has preserved liver function (Child-Pugh A); and

2.3 Transarterial chemoembolisation (TACE) is unsuitable; and

2.4 Any of the following:

2.4.1 Patient has not received prior systemic therapy for the treatment of hepatocellular carcinoma; or

2.4.2 Patient received funded lenvatinib before 1 March 2025; or

2.4.3 Both:

2.4.3.1 Patient has experienced treatment-limiting toxicity from treatment with lenvatinib; and

2.4.3.2 No disease progression since initiation of lenvatinib; and

2.5 Patient has an ECOG performance status of 0-2; and

2.6 To be given in combination with bevacizumab.

Continuation – unresectable hepatocellular carcinoma

Re-assessment required after 6 months

no evidence of disease progression.

DURVALUMAB – **Restricted** see terms below

⬇	Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
⬇	Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	Imfinzi

➥ **Restricted** (RS2084)

Initiation – Non-small cell lung cancer

Re-assessment required after 4 months

All of the following:

1 Either:

1.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); or

1.2 Patient has histologically or cytologically documented stage IIb (T1N2a only), locally advanced, unresectable non-small cell lung cancer (NSCLC); and

2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and

3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and

4 Patient has a ECOG performance status of 0 or 1; and

5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and

6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and

7 Either:

7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or

7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and

8 Treatment with durvalumab to cease upon signs of disease progression.

Continuation – Non-small cell lung cancer

Re-assessment required after 4 months

All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
2 Either:
  2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
  2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
3 Treatment with durvalumab to cease upon signs of disease progression; and
4 Total continuous treatment duration must not exceed 12 months.

IPILIMUMAB – **Restricted** see terms below

⬇	Inj 5 mg per ml, 10 ml vial	5,000.00	1	Yervoy
⬇	Inj 5 mg per ml, 40 ml vial	20,000.00	1	Yervoy

➥ **Restricted** (RS2195)

Initiation – stage III or IV resectable melanoma

Either:

1 The individual is currently on treatment with ipilimumab for neoadjuvant treatment of resectable stage IIIB, IIIC, IIID or IV melanoma and met all remaining criteria prior to commencing treatment; or
2 All of the following:
  2.1 The individual has resectable stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note); and
  2.2 The individual has not received prior funded systemic treatment in the perioperative setting for their stage IIIB, IIIC, IIID or IV melanoma; and
  2.3 The individual has ECOG performance score 0-2; and
  2.4 Treatment must be prior to complete surgical resection; and
  2.5 Neoadjuvant ipilimumab must be administered in combination with nivolumab; and
  2.6 Ipilimumab to be administered for maximum of two cycles prior to surgical resection.

Notes:

a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition.
b) Disease must be completely resectable and amenable to curative intent surgery, including stage IV disease.

Initiation – renal cell carcinoma

Limited to 4 months treatment

Either:

1 The patient is currently on treatment with ipilimumab and met all remaining criteria prior to commencing treatment; or
2 All of the following:
  2.1 The patient has metastatic renal cell carcinoma; and
  2.2 The patient is treatment naive; and
  2.3 The patient has ECOG performance status 0-2; and
  2.4 The disease is predominantly of clear cell histology; and
  2.5 Any of the following:
    2.5.1 The patient has sarcomatoid histology; or
    2.5.2 Haemoglobin levels less than the lower limit of normal; or
    2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or
    2.5.4 Neutrophils greater than the upper limit of normal; or
    2.5.5 Platelets greater than the upper limit of normal; or
    2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
    2.5.7 Karnofsky performance score of less than or equal to 70; and
  2.6 Ipilimumab is to be used at a maximum dose of 1 mg/kg for up to four cycles in combination with nivolumab.

NIVOLUMAB – **Restricted** see terms on the next page

⬇	Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
⬇	Inj 10 mg per ml, 10 ml vial	2,629.96	1	Opdivo

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS2196)

Initiation – stage III or IV resectable melanoma

Re-assessment required after 4 months

Either:

1 The individual is currently on treatment with nivolumab for neoadjuvant treatment of resectable stage IIIB, IIIC, IIID or IV melanoma and met all remaining criteria prior to commencing treatment; or

2 All of the following:

2.1 The individual has resectable stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note); and

2.2 The individual has not received prior funded systemic treatment in the perioperative setting for their stage IIIB, IIIC, IIID or IV melanoma; and

2.3 The individual has ECOG performance score 0-2; and

2.4 Treatment must be initiated prior to complete surgical resection; and

2.5 Neoadjuvant nivolumab must be administered in combination with ipilimumab; and

2.6 Nivolumab to be administered for maximum of two cycles prior to surgical resection.

Continuation – stage III or IV resectable melanoma

Re-assessment required after 4 months

Any of the following:

1 All of the following:

1.1 The individual has received funded neoadjuvant treatment with nivolumab in combination with ipilimumab; and

1.2 Adjuvant treatment with nivolumab is required; and

1.3 Treatment must be initiated within 13 weeks of complete surgical resection, unless delay is necessary due to post-surgery recovery; and

1.4 Nivolumab must be administered as monotherapy; and

1.5 Nivolumab to be discontinued at signs of disease recurrence or at completion of 12 months total treatment duration including any systemic neoadjuvant treatment (equivalent to 11 adjuvant cycles at 480 mg every 4 weeks plus initial 2 neoadjuvant treatment cycles); or

2 All of the following:

2.1 The individual has received neoadjuvant treatment with nivolumab and ipilimumab; and

2.2 The individual has unresectable or metastatic melanoma (excluding uveal) stage III or IV; and

2.3 The individual meets initial application criteria for nivolumab for unresectable or metastatic melanoma; or

3 All of the following:

3.1 The individual has received neoadjuvant treatment with nivolumab and ipilimumab; and

3.2 The individual has received treatment with nivolumab for unresectable or metastatic melanoma; and

3.3 The individual meets the renewal criteria for nivolumab for unresectable or metastatic melanoma.

Notes:

a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition.

b) Disease must be completely resectable and amenable to curative intent surgery, including stage IV disease.

Initiation – unresectable or metastatic melanoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Limited to 4 months treatment

All of the following:

1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and

2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and

3 The individual has ECOG performance 0-2; and

4 Either:

4.1 The individual has not received funded pembrolizumab; or

4.2 Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

4.2.1 The individual has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
4.2.2 The cancer did not progress while the individual was on pembrolizumab; and

5 Any of the following:
5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
5.3 All of the following:
5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-L1 inhibitor.

Continuation – unresectable or metastatic melanoma, less than 24 months on treatment
Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist
Re-assessment required after 4 months
Either:

1 Both:
1.1 Any of the following:
1.1.1 The individual's disease has had a complete response to treatment; or
1.1.2 The individual's disease has had a partial response to treatment; or
1.1.3 The individual has stable disease; and
1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; or

2 All of the following:
2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
2.2 The individual has signs of disease progression; and
2.3 Disease has not progressed during previous treatment with nivolumab.

Continuation – unresectable or metastatic melanoma, more than 24 months on treatment
Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist
Re-assessment required after 4 months
Both:

1 The individual has been on treatment for more than 24 months; and
2 Either:
2.1 Both:
2.1.1 Any of the following:
2.1.1.1 The individual's disease has had a complete response to treatment; or
2.1.1.2 The individual's disease has had a partial response to treatment; or
2.1.1.3 The individual has stable disease; and
2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; or
2.2 All of the following:
2.2.1 The individual has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
2.2.2 The individual has signs of disease progression; and
2.2.3 Disease has not progressed during previous treatment with nivolumab.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – renal cell carcinoma, first line

Limited to 4 months treatment

Either:

1 Patient is currently on treatment with nivolumab and met all remaining criteria prior to commencing treatment; or
2 All of the following:
  2.1 The patient has metastatic renal cell carcinoma; and
  2.2 The patient is treatment naive; and
  2.3 The patient has ECOG performance status 0-2; and
  2.4 The disease is predominantly of clear cell histology; and
  2.5 Any of the following:
    2.5.1 The patient has sarcomatoid histology; or
    2.5.2 Haemoglobin levels less than the lower limit of normal; or
    2.5.3 Corrected serum calcium level greater than 10 mg/dL (2.5 mmol/L); or
    2.5.4 Neutrophils greater than the upper limit of normal; or
    2.5.5 Platelets greater than the upper limit of normal; or
    2.5.6 Interval of less than 1 year from original diagnosis to the start of systemic therapy; or
    2.5.7 Karnofsky performance score of less than or equal to 70; and
  2.6 Nivolumab is to be used in combination with ipilimumab for the first four treatment cycles at a maximum dose of 3 mg/kg; and
  2.7 Nivolumab is to be used at a maximum maintenance dose of 240 mg every 2 weeks (or equivalent).

Initiation – renal cell carcinoma, second line

Limited to 4 months treatment

All of the following:

1 Patient has metastatic renal-cell carcinoma; and
2 The disease is of predominant clear-cell histology; and
3 Patient has ECOG performance status 0-2; and
4 Patient has documented disease progression following one or two previous regimens of antiangiogenic therapy; and
5 Patient has not previously received a funded immune checkpoint inhibitor; and
6 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

Continuation – renal cell carcinoma

Re-assessment required after 4 months

All of the following:

1 Any of the following:
  1.1 Patient's disease has had a complete response to treatment; or
  1.2 Patient's disease has had a partial response to treatment; or
  1.3 Patient has stable disease; and
2 No evidence of disease progression; and
3 Nivolumab is to be used as monotherapy at a maximum dose of 240 mg every 2 weeks (or equivalent) and discontinued at disease progression.

PEMBROLIZUMAB – **Restricted** see terms below

⬇	Inj 25 mg per ml, 4 ml vial	4,680.00	1	Keytruda

➥ **Restricted** (RS2185)

Initiation – stage III or IV resectable melanoma - neoadjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 The individual has resectable stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note); and
2 The individual has not received prior funded systemic treatment in the perioperative setting for their stage IIIB, IIIC, IIID or IV melanoma; and
3 Treatment must be prior to complete surgical resection; and
4 Pembrolizumab must be administered as monotherapy; and
5 The individual has ECOG performance score 0-2; and
6 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

Continuation – stage III or IV resectable melanoma - neoadjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Any of the following:

1 Both:
  1.1 The individual has received neoadjuvant treatment with an immune checkpoint inhibitor; and
  1.2 The individual meets initiation criteria for pembrolizumab for stage III or IV resected melanoma – adjuvant; or
2 Both:
  2.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
  2.2 The individual meets continuation criteria for pembrolizumab for stage III or IV resected melanoma – adjuvant; or
3 All of the following:
  3.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
  3.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
  3.3 The individual meets initiation criteria for pembrolizumab for unresectable or metastatic melanoma; or
4 All of the following:
  4.1 The individual has received neoadjuvant and adjuvant treatment with an immune checkpoint inhibitor; and
  4.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma; and
  4.3 The individual meets continuation criteria for pembrolizumab for unresectable or metastatic melanoma.

Notes:

a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition
b) Initiating treatment within 13 weeks of complete surgical resection means either 13 weeks after resection (primary or lymphadenectomy) or 13 weeks prior to the scheduled date of the resection (primary or lymphadenectomy)

Initiation – stage III or IV resected melanoma - adjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

All of the following:

1 The individual has resected stage IIIB, IIIC, IIID or IV melanoma (excluding uveal) (see note a); and
2 Adjuvant treatment with pembrolizumab is required; and
3 The individual has not received prior funded systemic treatment in the adjuvant setting for stage IIIB, IIIC, IIID or IV melanoma; and
4 Treatment must be in addition to complete surgical resection; and
5 Treatment must be initiated within 13 weeks of complete surgical resection, unless delay is necessary due to post-surgery recovery (see note b); and
6 Pembrolizumab must be administered as monotherapy; and
7 The individual has ECOG performance score 0-2; and
8 Pembrolizumab to be administered at a fixed dose of 200 mg every 3 weeks (or equivalent).

Notes:

a) Stage IIIB, IIIC, IIID or IV melanoma defined as per American Joint Committee on Cancer (AJCC) 8th Edition

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

b) Initiating treatment within 13 weeks of complete surgical resection means 13 weeks after resection (primary or lymphadenectomy)

Continuation – stage III or IV resected melanoma - adjuvant

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Any of the following:

1 All of the following:
- 1.1 No evidence of disease recurrence; and
- 1.2 Pembrolizumab must be administered as monotherapy; and
- 1.3 Pembrolizumab to be administered at a fixed dose of 200 mg every three weeks (or equivalent) for a maximum of 12 months total treatment course, including any systemic neoadjuvant treatment; and
- 1.4 Treatment to be discontinued at signs of disease recurrence or at completion of 12 months total treatment course (equivalent to 18 cycles at a dose of 200 mg every 3 weeks), including any systemic neoadjuvant treatment; or

2 All of the following:
- 2.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and
- 2.2 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2.3 The individual meets initiation criteria for pembrolizumab for unresectable or metastatic melanoma; or

3 All of the following:
- 3.1 The individual has received adjuvant treatment with an immune checkpoint inhibitor; and
- 3.2 The individual has received treatment with an immune checkpoint inhibitor for unresectable or metastatic melanoma; and
- 3.3 The individual meets continuation criteria for pembrolizumab for unresectable or metastatic melanoma.

Initiation – unresectable or metastatic melanoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Limited to 4 months treatment

All of the following:

1 The individual has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and

2 Baseline measurement of overall tumour burden is documented clinically and radiologically; and

3 The individual has ECOG performance 0-2; and

4 Either:
- 4.1 The individual has not received funded nivolumab; or
- 4.2 Both:
  - 4.2.1 The individual has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
  - 4.2.2 The cancer did not progress while the individual was on nivolumab; and

5 Any of the following:
- 5.1 The individual has been diagnosed in the metastatic or unresectable stage III or IV setting; or
- 5.2 The individual did not receive treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; or
- 5.3 All of the following:
  - 5.3.1 The individual received treatment in the perioperative setting with a PD-1/PD-L1 inhibitor; and
  - 5.3.2 The individual did not experience disease recurrence while on treatment with that PD-1/PD-L1 inhibitor; and
  - 5.3.3 The individual did not experience disease recurrence within six months of completing perioperative treatment with a PD-1/PD-L1 inhibitor.

Continuation – unresectable or metastatic melanoma, less than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Both:
- 1.1 Any of the following:
  - 1.1.1 The individual's disease has had a complete response to treatment; or
  - 1.1.2 The individual's disease has had a partial response to treatment; or
  - 1.1.3 The individual has stable disease; and
- 1.2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; or

2 All of the following:
- 2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
- 2.2 The individual has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Continuation – unresectable or metastatic melanoma, more than 24 months on treatment

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Both:

1 The individual has been on treatment for more than 24 months; and

2 Either:
- 2.1 All of the following:
  - 2.1.1 Any of the following:
    - 2.1.1.1 The individual's disease has had a complete response to treatment; or
    - 2.1.1.2 The individual's disease has had a partial response to treatment; or
    - 2.1.1.3 The individual has stable disease; and
  - 2.1.2 Response to treatment in target lesions has been determined by comparable radiologic or clinical assessment following the most recent treatment period; and
  - 2.1.3 The treatment remains clinically appropriate and the individual is benefitting from the treatment; or
- 2.2 All of the following:
  - 2.2.1 The individual has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
  - 2.2.2 The individual has signs of disease progression; and
  - 2.2.3 Disease has not progressed during previous treatment with pembrolizumab.

Initiation – non-small cell lung cancer first-line monotherapy

Medical oncologist

Re-assessment required after 4 months

All of the following:

1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and

2 Patient has not had chemotherapy for their disease in the palliative setting; and

3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and

4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR, ROS-1 or ALK tyrosine kinase unless not possible to ascertain; and

5 Pembrolizumab to be used as monotherapy; and

6 Either:
- 6.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 50% as determined by a validated test unless not possible to ascertain; or
- 6.2 Both:
  - 6.2.1 There is documentation confirming the disease expresses PD-L1 at a level greater than or equal to 1% as determined by a validated test unless not possible to ascertain; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

6.2.2 Chemotherapy is determined to be not in the best interest of the patient based on clinician assessment; and

7 Patient has an ECOG 0-2; and
8 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
9 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – non-small cell lung cancer first-line monotherapy
Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist
Re-assessment required after 4 months
All of the following:

1 Any of the following:
  1.1 Patient's disease has had a complete response to treatment; or
  1.2 Patient's disease has had a partial response to treatment; or
  1.3 Patient has stable disease; and
2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
3 No evidence of disease progression; and
4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – non-small cell lung cancer first-line combination therapy
Medical oncologist
Re-assessment required after 4 months
All of the following:

1 Patient has locally advanced or metastatic, unresectable, non-small cell lung cancer; and
2 The patient has not had chemotherapy for their disease in the palliative setting; and
3 Patient has not received prior funded treatment with an immune checkpoint inhibitor for NSCLC; and
4 For patients with non-squamous histology there is documentation confirming that the disease does not express activating mutations of EGFR, ROS-1 or ALK tyrosine kinase unless not possible to ascertain; and
5 Pembrolizumab to be used in combination with platinum-based chemotherapy; and
6 Patient has an ECOG 0-2; and
7 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks; and
8 Baseline measurement of overall tumour burden is documented clinically and radiologically.

Continuation – non-small cell lung cancer first-line combination therapy
Medical oncologist or any relevant practitioner on the recommendation of a medical oncologist
Re-assessment required after 4 months
All of the following:

1 Any of the following:
  1.1 Patient's disease has had a complete response to treatment; or
  1.2 Patient's disease has had a partial response to treatment; or
  1.3 Patient has stable disease; and
2 Response to treatment in target lesions has been determined by comparable radiologic assessment following the most recent treatment period; and
3 No evidence of disease progression; and
4 The treatment remains clinically appropriate and patient is benefitting from treatment; and
5 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

6 Treatment with pembrolizumab to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – breast cancer, advanced

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 6 months

Either:

1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
2 All of the following:
  2.1 Either:
    2.1.1 Patient has recurrent or de novo unresectable, inoperable locally advanced triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); or
    2.1.2 Patient has recurrent or de novo metastatic triple-negative breast cancer (that does not express ER, PR or HER2 IHC3+ or ISH+ [including FISH or other technology]); and
  2.2 Patient is treated with palliative intent; and
  2.3 Patient's cancer has confirmed PD-L1 Combined Positive Score (CPS) is greater than or equal to 10; and
  2.4 Patient has received no prior systemic therapy in the palliative setting; and
  2.5 Patient has an ECOG score of 0–2; and
  2.6 Pembrolizumab is to be used in combination with chemotherapy; and
  2.7 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
  2.8 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – breast cancer, advanced

Any relevant practitioner

Re-assessment required after 6 months

All of the following:

1 Any of the following:
  1.1 Patient's disease has had a complete response to treatment; or
  1.2 Patient's disease has had a partial response to treatment; or
  1.3 Patient has stable disease; and
2 No evidence of disease progression; and
3 Response to treatment in target lesions has been determined by a comparable radiologic assessment following the most recent treatment period; and
4 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
5 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – head and neck squamous cell carcinoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
2 All of the following:
  2.1 Patient has recurrent or metastatic head and neck squamous cell carcinoma of mucosal origin (excluding nasopharyngeal carcinoma) that is incurable by local therapies; and
  2.2 Patient has not received prior systemic therapy in the recurrent or metastatic setting; and
  2.3 Patient has a positive PD-L1 combined positive score (CPS) of greater than or equal to 1; and
  2.4 Patient has an ECOG performance score of 0-2; and
  2.5 Either:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.5.1 Pembrolizumab to be used in combination with platinum-based chemotherapy; or
2.5.2 Pembrolizumab to be used as monotherapy; and
2.6 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – head and neck squamous cell carcinoma
Any relevant practitioner
Re-assessment required after 4 months
All of the following:
1 Any of the following:
1.1 Patient's disease has had a complete response to treatment; or
1.2 Patient's disease has had a partial response to treatment; or
1.3 Patient has stable disease; and
2 No evidence of disease progression; and
3 Pembrolizumab is to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – MSI-H/dMMR advanced colorectal cancer
Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist
Re-assessment required after 4 months
Either:
1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or
2 All of the following:
2.1 Either:
2.1.1 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) metastatic colorectal cancer; or
2.1.2 Individual has deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) unresectable colorectal cancer; and
2.2 Individual is treated with palliative intent; and
2.3 Individual has not previously received funded treatment with pembrolizumab for MSI-H/dMMR advanced colorectal cancer; and
2.4 Individual has an ECOG performance score of 0-2; and
2.5 Baseline measurement of overall tumour burden is documented clinically and radiologically; and
2.6 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – MSI-H/dMMR advanced colorectal cancer
Any relevant practitioner
Re-assessment required after 4 months
All of the following:
1 No evidence of disease progression; and
2 Pembrolizumab to be used at a maximum dose of 200 mg every three weeks (or equivalent); and
3 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – Urothelial carcinoma
Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist
Re-assessment required after 4 months
Either:
1 Patient is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 All of the following:
- 2.1 Patient has inoperable locally advanced (T4) or metastatic urothelial carcinoma; and
- 2.2 Patient has an ECOG performance score of 0-2; and
- 2.3 Patient has documented disease progression following treatment with chemotherapy; and
- 2.4 Pembrolizumab to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent) for a maximum of 16 weeks.

Continuation – Urothelial carcinoma

Any relevant practitioner

Re-assessment required after 4 months

All of the following:

1 Any of the following:
- 1.1 Patient's disease has had a complete response to treatment; or
- 1.2 Patient's disease has had a partial response to treatment; or
- 1.3 Patient has stable disease; and

2 No evidence of disease progression; and

3 Pembrolizumab is to be used as monotherapy at a maximum dose of 200 mg every three weeks (or equivalent); and

4 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Initiation – relapsed/refractory Hodgkin lymphoma

Relevant specialist or any relevant practitioner on the recommendation of a relevant specialist

Re-assessment required after 4 months

Either:

1 Individual is currently on treatment with pembrolizumab and met all remaining criteria prior to commencing treatment; or

2 All of the following:
- 2.1 Either:
  - 2.1.1 Both:
    - 2.1.1.1 Individual has relapsed/refractory Hodgkin lymphoma after two or more lines of chemotherapy; and
    - 2.1.1.2 Individual is ineligible for autologous stem cell transplant; or
  - 2.1.2 Individual has relapsed/refractory Hodgkin lymphoma and has previously undergone an autologous stem cell transplant; and
- 2.2 Individual has not previously received funded pembrolizumab for relapsed/refractory Hodgkin lymphoma; and
- 2.3 Pembrolizumab to be administered at doses no greater than 200 mg once every 3 weeks.

Continuation – relapsed/refractory Hodgkin lymphoma

Any relevant practitioner

Re-assessment required after 6 months

Both:

1 Patient has received a partial or complete response to pembrolizumab; and

2 Treatment with pembrolizumab is to cease after a total duration of 24 months from commencement (or equivalent of 35 cycles dosed every 3 weeks).

Other Immunosuppressants

ANTITHYMOCYTE GLOBULIN (EQUINE)

Inj 50 mg per ml, 5 ml ampoule	4,439.17	5	ATGAM

ANTITHYMOCYTE GLOBULIN (RABBIT)

Inj 25 mg vial

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
AZATHIOPRINE			
Tab 25 mg – **5% DV Feb-26 to 2028**	10.15	60	**Azamun**
Tab 50 mg – **5% DV Feb-26 to 2028**	10.34	100	**Azamun**
Inj 50 mg vial			
Inj 100 mg vial			
BACILLUS CALMETTE-GUERIN (BCG) – **Restricted** see terms below			
⬇ Inj 2-8 × 10^8 CFU vial	149.37	1	OncoTICE
⬇ Inj 40 mg per ml, vial	182.45	3	SII-Onco-BCG

➥ **Restricted** (RS1206)

Initiation

For use in bladder cancer.

EVEROLIMUS – **Restricted** see terms below			
⬇ Tab 5 mg	4,555.76	30	Afinitor
⬇ Tab 10 mg	6,512.29	30	Afinitor

➥ **Restricted** (RS2076)

Initiation

Neurologist or oncologist

Re-assessment required after 3 months

Both:

1 Patient has tuberous sclerosis; and
2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Continuation

Neurologist or oncologist

Re-assessment required after 12 months

All of the following:

1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
2 The treatment remains appropriate and the patient is benefiting from treatment; and
3 Everolimus to be discontinued at progression of SEGAs.

Initiation – renal cell carcinoma

Re-assessment required after 4 months

Either:

1 All of the following:
  1.1 The patient has metastatic renal cell carcinoma; and
  1.2 The disease is of predominant clear-cell histology; and
  1.3 The patient has documented disease progression following one previous line of treatment; and
  1.4 The patient has an ECOG performance status of 0-2; and
  1.5 Everolimus is to be used in combination with lenvatinib; or
2 All of the following:
  2.1 Patient has received funded treatment with nivolumab for the second line treatment of metastatic renal cell carcinoma; and
  2.2 Patient has experienced treatment limiting toxicity from treatment with nivolumab; and
  2.3 Everolimus is to be used in combination with lenvatinib; and
  2.4 There is no evidence of disease progression.

Continuation – renal cell carcinoma

Re-assessment required after 4 months

there is no evidence of disease progression.

MYCOPHENOLATE MOFETIL			
Tab 500 mg	35.90	50	CellCept
Cap 250 mg	35.90	100	CellCept
Powder for oral liq 1 g per 5 ml	187.25	165 ml	CellCept
Inj 500 mg vial	133.33	4	CellCept

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
PICIBANIL			
Inj 100 mcg vial			
SIROLIMUS – **Restricted** see terms below			
⬇ Tab 1 mg	749.99	100	Rapamune
⬇ Tab 2 mg	1,499.99	100	Rapamune
⬇ Oral liq 1 mg per ml	449.99	60 ml	Rapamune

➥ **Restricted** (RS1991)

Initiation

For rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR < 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencepthalopathy; or
- Significant malignant disease

Initiation – severe non-malignant lymphovascular malformations*

Re-assessment required after 6 months

All of the following:

1 Patient has severe non-malignant lymphovascular malformation*; and
2 Any of the following:
  2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
  2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
  2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Continuation – severe non-malignant lymphovascular malformations*

Re-assessment required after 12 months

All of the following:

1 Either:
  1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
  1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
2 No evidence of progressive disease; and
3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initiation – renal angiomyolipoma(s) associated with tuberous sclerosis complex*

Nephrologist or urologist

Re-assessment required after 6 months

Both:

1 Patient has tuberous sclerosis complex*; and
2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Continuation – renal angiomyolipoma(s) associated with tuberous sclerosis complex*

Re-assessment required after 12 months

All of the following:

1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
2 Demonstrated stabilisation or improvement in renal function; and
3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initiation – refractory seizures associated with tuberous sclerosis complex*

Neurologist

Re-assessment required after 6 months

All of the following:

1 Patient has epilepsy with a background of documented tuberous sclerosis complex*; and
2 Either:
  2.1 Both:
    2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
    2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
  2.2 Both:
    2.2.1 Vigabatrin is contraindicated; and
    2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
3 Seizures have a significant impact on quality of life; and
4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Those of childbearing potential are not required to trial phenytoin sodium, sodium valproate, and topiramate. Those who can father children are not required to trial sodium valproate.

Continuation – refractory seizures associated with tuberous sclerosis complex*

Neurologist

Re-assessment required after 12 months

demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – **Restricted** see terms below

⬇	Tab modified-release 15 mg	1,271.00	28	Rinvoq
⬇	Tab modified-release 30 mg	2,033.00	28	Rinvoq
⬇	Tab modified-release 45 mg	3,049.00	28	Rinvoq

➥ **Restricted** (RS2120)

Initiation – Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)

Limited to 6 months treatment

All of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 The individual has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
2 Either:
  2.1 The individual has experienced intolerable side effects with adalimumab and/or etanercept; or
  2.2 The individual has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
3 Any of the following:
  3.1 Rituximab is not clinically appropriate; or
  3.2 The individual is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  3.3 Both:
    3.3.1 The individual has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
    3.3.2 Either:
      3.3.2.1 The individual has experienced intolerable side effects with rituximab; or
      3.3.2.2 At four months following the initial course of rituximab the individual has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Continuation – Rheumatoid Arthritis
Re-assessment required after 6 months
Either:

1 Following 6 months' initial treatment, the individual has experienced at least a 50% decrease in active joint count from baseline; or
2 On subsequent reapplications, the individual has experienced at least a continuing 30% improvement in active joint count from baseline.

Initiation – Atopic dermatitis
Re-assessment required after 6 months
Either:

1 Individual is currently on treatment with upadacitinib for atopic dermatitis and met all remaining criteria prior to commencing treatment; or
2 All of the following:
  2.1 Individual has moderate to severe atopic dermatitis, severity as defined by an Eczema Area and Severity Index (EASI) score of greater than or equal to 16 or a Dermatology Life Quality Index (DLQI) score of greater than or equal to 10; and
  2.2 Individual has received insufficient benefit from topical therapy (including topical corticosteroids or topical calcineurin inhibitors) for a 28-day trial within the last 6 months, unless contraindicated to all; and
  2.3 Individual has trialled and received insufficient benefit from at least one systemic therapy for a minimum of three months (eg ciclosporin, azathioprine, methotrexate or mycophenolate mofetil), unless contraindicated to all; and
  2.4 An EASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  2.5 The most recent EASI or DQLI assessment is no more than 1 month old at the time of application.

Continuation – Atopic dermatitis
Re-assessment required after 12 months
Either:

1 Individual has received a 75% or greater reduction in EASI score (EASI 75) as compared to baseline EASI prior to commencing upadacitinib; or
2 Individual has received a DLQI improvement of 4 or more as compared to baseline DLQI prior to commencing upadacitinib.

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

Initiation – Crohn's disease – adult

Re-assessment required after 6 months

Either:

1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing treatment; or

2 Both:

2.1 Individual has active Crohn's disease; and

2.2 Either:

2.2.1 Individual has had an initial approval for prior biologic therapy and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or

2.2.2 Both:

2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for Crohn's disease; and

2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Continuation – Crohn's disease – adult

Re-assessment required after 2 years

Any of the following:

1 CDAI score has reduced by 100 points from the CDAI score when the individual was initiated on biologic therapy; or

2 HBI score has reduced by 3 points from when individual was initiated on biologic therapy; or

3 CDAI score is 150 or less; or

4 HBI score is 4 or less; or

5 The individual has experienced an adequate response to treatment, but CDAI score cannot be assessed.

Initiation – Crohn's disease – children

Re-assessment required after 6 months

Either:

1 Individual is currently on treatment with upadacitinib for Crohn's disease and met all remaining criteria prior to commencing treatment; or

2 Both:

2.1 Child has active Crohn's disease; and

2.2 Either:

2.2.1 Child has had an initial approval for prior biologic therapy for Crohn's disease and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or

2.2.2 Both:

2.2.2.1 Child meets the initiation criteria for prior biologic therapies for Crohn's disease; and

2.2.2.2 Other biologic therapies for Crohn's disease are contraindicated.

Continuation – Crohn's disease – children

Re-assessment required after 2 years

Any of the following:

1 PCDAI score has reduced by 10 points from when the child was initiated on treatment; or

2 PCDAI score is 15 or less; or

3 The child has experienced an adequate response to treatment, but PCDAI score cannot be assessed.

Note: Indications marked with * are unapproved indications.

Initiation – Ulcerative colitis

Re-assessment required after 6 months

Either:

1 Individual is currently on treatment with upadacitinib for ulcerative colitis and met all remaining criteria prior to commencing treatment; or

2 Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.1 Individual has active ulcerative colitis; and
2.2 Either:
2.2.1 Individual has had an initial approval for prior biologic therapy for ulcerative colitis and has experienced intolerable side effects or insufficient benefit to meet renewal criteria; or
2.2.2 Both:
2.2.2.1 Individual meets the initiation criteria for prior biologic therapies for ulcerative colitis; and
2.2.2.2 Other biologic therapies for ulcerative colitis are contraindicated.

Continuation – Ulcerative colitis
Re-assessment required after 2 years
Either:
1 The SCCAI score has reduced by 2 points or more from the SCCAI score when the individual was initiated on treatment; or
2 PUCAI score has reduced by 10 points or more from the PUCAI score when the individual was initiated on treatment.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Antiallergy Preparations

Allergic Emergencies

ADRENALINE – **Restricted** see terms below

⬇ Inj 0.15 mg per 0.3 ml auto-injector – **5% DV Dec-25 to 2028**	85.50	1	**Epipen Jr**
⬇ Inj 0.3 mg per 0.3 ml auto-injector – **5% DV Dec-25 to 2028**	85.50	1	**Epipen**

➥ **Restricted** (RS1944)

Initiation – anaphylaxis

Either:

1 Patient has experienced a previous anaphylactic reaction which has resulted in presentation to a hospital or emergency department; or

2 Patient has been assessed to be at significant risk of anaphylaxis by a relevant practitioner.

ICATIBANT – **Restricted** see terms below

⬇ Inj 10 mg per ml, 3 ml prefilled syringe – **5% DV Oct-26 to 2028**	2,668.00	1	Firazyr
	479.00		**Icatibant Lupin**

(Firazyr Inj 10 mg per ml, 3 ml prefilled syringe to be delisted 1 October 2026)

➥ **Restricted** (RS1501)

Initiation

Clinical immunologist or relevant specialist

Re-assessment required after 12 months

Both:

1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and

2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Continuation

Re-assessment required after 12 months

The treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

BEE VENOM – **Restricted** see terms below

⬇ Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent			
⬇ Inj 550 mcg vial with diluent			
⬇ Initiation kit - 1 vial freeze dried venom with diluent	305.00	1	VENOX
⬇ Maintenance Kit - 1 vial freeze dried venom with diluent	305.00	1	VENOX

➥ **Restricted** (RS1117)

Initiation

Both:

1 RAST or skin test positive; and

2 Patient has had severe generalised reaction to the sensitising agent.

PAPER WASP VENOM – **Restricted** see terms below

⬇ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent			
⬇ Inj 550 mcg vial with diluent			

➥ **Restricted** (RS1118)

Initiation

Both:

1 RAST or skin test positive; and

2 Patient has had severe generalised reaction to the sensitising agent.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

YELLOW JACKET WASP VENOM – **Restricted** see terms below
⬇ Treatment kit - 6 vials 120 mcg freeze dried venom, with diluent
⬇ Inj 550 mcg vial with diluent
➥ **Restricted** (RS1119)
Initiation
Both:
1 RAST or skin test positive; and
2 Patient has had severe generalised reaction to the sensitising agent.

Allergy Prophylactics

BUDESONIDE			
Nasal spray 50 mcg per dose – **5% DV Feb-25 to 2027**	2.59	200 dose	**SteroClear**
Nasal spray 100 mcg per dose – **5% DV Feb-25 to 2027**	2.89	200 dose	**SteroClear**
FLUTICASONE PROPIONATE			
Metered dose nasal spray 50 mcg per dose – **5% DV Aug-26 to 2028**	2.57	120 dose	**Flixonase Hayfever & Allergy**
IPRATROPIUM BROMIDE			
Aqueous nasal spray 0.03%	5.23	15 ml	Univent
SODIUM CROMOGLICATE			
Nasal spray 4%			

Antihistamines

CETIRIZINE HYDROCHLORIDE			
Tab 10 mg – **5% DV Sep-23 to 2026**	1.71	100	**Zista**
Oral liq 1 mg per ml	3.99	200 ml	Histaclear
CHLORPHENIRAMINE MALEATE			
Oral liq 0.4 mg per ml			
Inj 10 mg per ml, 1 ml ampoule			
CYPROHEPTADINE HYDROCHLORIDE			
Tab 4 mg			
FEXOFENADINE HYDROCHLORIDE			
Tab 60 mg			
Tab 120 mg – **5% DV Jul-25 to 2027**	3.49	30	**Fexaclear**
Tab 180 mg – **5% DV Jul-25 to 2027**	4.10	30	**Fexaclear**
LORATADINE			
Tab 10 mg – **5% DV Jun-26 to 2028**	6.02	100	Lorafix
	1.59		**Loratadine Noumed**
Oral liq 1 mg per ml	1.43	100 ml	Haylor Syrup
(Lorafix Tab 10 mg to be delisted 1 June 2026)			
PROMETHAZINE HYDROCHLORIDE			
Tab 10 mg – **5% DV Dec-25 to 2028**	2.19	100	**Allersoothe**
Tab 25 mg – **5% DV Dec-25 to 2028**	2.69	100	**Allersoothe**
Oral liq 1 mg per ml	3.39	100 ml	Allersoothe
Inj 25 mg per ml, 2 ml ampoule	21.09	5	Hospira

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Anticholinergic Agents			
IPRATROPIUM BROMIDE			
Aerosol inhaler 20 mcg per dose			
Nebuliser soln 250 mcg per ml, 1 ml ampoule			
Nebuliser soln 250 mcg per ml, 2 ml ampoule	11.73	20	Accord
	8.11	10	Ipratropium Viatris
	11.73	20	Univent
Anticholinergic Agents with Beta-Adrenoceptor Agonists			
SALBUTAMOL WITH IPRATROPIUM BROMIDE			
Aerosol inhaler 100 mcg with ipratropium bromide 20 mcg per dose			
Nebuliser soln 2.5 mg with ipratropium bromide 0.5 mg per 2.5 ml ampoule	96.50	60	Combiprasal
	11.04	20	Duolin
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 3.0 ml vial	16.56	30	Cipla
Long-Acting Muscarinic Agents			
GLYCOPYRRONIUM			
Note: inhaled glycopyrronium treatment must not be used if the patient is also receiving treatment with subsidised tiotropium or umeclidinium.			
Powder for inhalation 50 mcg per dose	61.00	30 dose	Seebri Breezhaler
TIOTROPIUM BROMIDE			
Note: tiotropium treatment must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.			
Soln for inhalation 2.5 mcg per dose	50.37	60 dose	Spiriva Respimat
Powder for inhalation 18 mcg per dose	50.37	30 dose	Spiriva
UMECLIDINIUM			
Note: Umeclidinium must not be used if the patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.			
Powder for inhalation 62.5 mcg per dose	61.50	30 dose	Incruse Ellipta
Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists			
➦ **Restricted** (RS2155)			
Initiation			
Both:			
1 Patient has been stabilised on a long acting muscarinic antagonist; and			
2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.			
Note: Combination long acting muscarinic antagonist and long acting beta-2 agonist must not be used if the patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.			
GLYCOPYRRONIUM WITH INDACATEROL – **Restricted** see terms above			
⬆ Powder for Inhalation 50 mcg with indacaterol 110 mcg	81.00	30 dose	Ultibro Breezhaler
TIOTROPIUM BROMIDE WITH OLODATEROL – **Restricted** see terms above			
⬆ Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg	81.00	60 dose	Spiolto Respimat
UMECLIDINIUM WITH VILANTEROL – **Restricted** see terms above			
⬆ Powder for inhalation 62.5 mcg with vilanterol 25 mcg	77.00	30 dose	Anoro Ellipta

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Inhaled Corticosteroid with Long-Acting Muscarinic Antagonist and Beta Agonist

BUDESONIDE WITH GLYCOPYRRONIUM AND EFORMOTEROL – **Restricted** see terms below

⬇ Aerosol inhaler budesonide 160 mcg with glycopyrronium 7.2 mcg and formoterol 5 mcg per dose..... 79.15 120 dose Breztri Aerosphere

➥ **Restricted** (RS2085)

Initiation

Both:

1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and

2 Either:

2.1 Both:

2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and

2.1.2 Any of the following:

Clinical criteria:

2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or

2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or

2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or

2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3 × 10^9 cells/L in the previous 12 months; or

2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long-acting muscarinic antagonist and long-acting beta-2 agonist – ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler therapy.

FLUTICASONE FUROATE WITH UMECLIDINIUM AND VILANTEROL – **Restricted** see terms below

⬇ Powder for inhalation fluticasone furoate 100 mcg with umeclidinium 62.5 mcg and vilanterol 25 mcg..... 104.24 30 dose Trelegy Ellipta

➥ **Restricted** (RS2028)

Initiation

Both:

1 Patient has a diagnosis of COPD confirmed by spirometry or spirometry has been attempted and technically acceptable results are not possible; and

2 Either:

2.1 Both:

2.1.1 Patient is currently receiving an inhaled corticosteroid with long acting beta-2 agonist (ICS/LABA) or a long acting muscarinic antagonist with long acting beta-2 agonist (LAMA/LABA); and

2.1.2 Any of the following:

Clinical criteria:

2.1.2.1 Patient has a COPD Assessment Test (CAT) score greater than 10; or

2.1.2.2 Patient has had 2 or more exacerbations in the previous 12 months; or

2.1.2.3 Patient has had one exacerbation requiring hospitalisation in the previous 12 months; or

2.1.2.4 Patient has had an eosinophil count greater than or equal to 0.3 × 10^9 cells/L in the previous 12 months; or

2.2 Patient is currently receiving multiple inhaler triple therapy (inhaled corticosteroid with long acting muscarinic antagonist and long acting beta-2 agonist – ICS/LAMA/LABA) and met at least one of the clinical criteria above prior to commencing multiple inhaler triple therapy.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Antifibrotics

NINTEDANIB – **Restricted** see terms below

⬇ Cap 100 mg	2,554.00	60	Ofev
⬇ Cap 150 mg	3,870.00	60	Ofev

➥ **Restricted** (RS1813)

Initiation – idiopathic pulmonary fibrosis

Respiratory specialist

Re-assessment required after 12 months

All of the following:

1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
2 Forced vital capacity is between 50% and 90% predicted; and
3 Nintedanib is to be discontinued at disease progression (See Note); and
4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
5 Any of the following:
  5.1 The patient has not previously received treatment with pirfenidone; or
  5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Continuation – idiopathic pulmonary fibrosis

Respiratory specialist

Re-assessment required after 12 months

All of the following:

1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – **Restricted** see terms below

⬇ Tab 267 mg – **5% DV Oct-26 to 2028**	1,215.00	90	Esbriet
	145.80		**Pirfenidone Sandoz**
⬇ Tab 801 mg – **5% DV Oct-26 to 2028**	3,645.00	90	Esbriet
	437.40		**Pirfenidone Sandoz**

(Esbriet Tab 267 mg to be delisted 1 October 2026)

(Esbriet Tab 801 mg to be delisted 1 October 2026)

➥ **Restricted** (RS1814)

Initiation – idiopathic pulmonary fibrosis

Respiratory specialist

Re-assessment required after 12 months

All of the following:

1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
2 Forced vital capacity is between 50% and 90% predicted; and
3 Pirfenidone is to be discontinued at disease progression (See Notes); and
4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
5 Any of the following:
  5.1 The patient has not previously received treatment with nintedanib; or
  5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Continuation – idiopathic pulmonary fibrosis

Respiratory specialist

Re-assessment required after 12 months

All of the following:

1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Beta-Adrenoceptor Agonists

SALBUTAMOL			
Oral liq 400 mcg per ml – **5% DV May-25 to 2027**	50.00	150 ml	**Ventolin**
Inj 500 mcg per ml, 1 ml ampoule			
Inj 1 mg per ml, 5 ml ampoule			
Aerosol inhaler, 100 mcg per dose	4.18	200 dose	SalAir
	7.45		Ventolin
Nebuliser soln 1 mg per ml, 2.5 ml ampoule	8.96	20	Asthalin
			UK Cipla
Nebuliser soln 2 mg per ml, 2.5 ml ampoule	9.43	20	Asthalin
			UK Cipla
TERBUTALINE SULPHATE			
Powder for inhalation 250 mcg per dose			
Inj 0.5 mg per ml, 1 ml ampoule			
Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated	22.20	120 dose	Bricanyl Turbuhaler

Decongestants

OXYMETAZOLINE HYDROCHLORIDE
- Aqueous nasal spray 0.25 mg per ml
- Aqueous nasal spray 0.5 mg per ml

PSEUDOEPHEDRINE HYDROCHLORIDE
- Tab 60 mg

SODIUM CHLORIDE
- Aqueous nasal spray isotonic

SODIUM CHLORIDE WITH SODIUM BICARBONATE
- Soln for nasal irrigation

XYLOMETAZOLINE HYDROCHLORIDE
- Aqueous nasal spray 0.05%
- Aqueous nasal spray 0.1%
- Nasal drops 0.05%
- Nasal drops 0.1%

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler 50 mcg per dose	8.54	200 dose	Beclazone 50
	14.01		Qvar
Aerosol inhaler 100 mcg per dose	12.50	200 dose	Beclazone 100
	17.52		Qvar
Aerosol inhaler 250 mcg per dose	22.67	200 dose	Beclazone 250
BUDESONIDE			
Nebuliser soln 250 mcg per ml, 2 ml ampoule			
Nebuliser soln 500 mcg per ml, 2 ml ampoule			
Powder for inhalation 100 mcg per dose			
Powder for inhalation 200 mcg per dose			
Powder for inhalation 400 mcg per dose			
FLUTICASONE			
Aerosol inhaler 50 mcg per dose	7.19	120 dose	Flixotide
Powder for inhalation 50 mcg per dose	8.61	60 dose	Flixotide Accuhaler
Powder for inhalation 100 mcg per dose	7.81	60 dose	Flixotide Accuhaler
Aerosol inhaler 125 mcg per dose	13.60	120 dose	Flixotide
Aerosol inhaler 250 mcg per dose	24.62	120 dose	Flixotide
Powder for inhalation 250 mcg per dose	11.93	60 dose	Flixotide Accuhaler
Leukotriene Receptor Antagonists			
MONTELUKAST			
Tab 4 mg – **5% DV Dec-25 to 2028**	3.10	28	**Montelukast Viatris**
Tab 5 mg – **5% DV Dec-25 to 2028**	3.10	28	**Montelukast Viatris**
			Relonchem
Tab 10 mg – **5% DV Dec-25 to 2028**	2.45	28	**Montelukast Viatris**
Long-Acting Beta-Adrenoceptor Agonists			
EFORMOTEROL FUMARATE			
Powder for inhalation 12 mcg per dose			
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated (equivalent to eformoterol fumarate 6 mcg metered dose)			
INDACATEROL			
Powder for inhalation 150 mcg per dose	61.00	30 dose	Onbrez Breezhaler
Powder for inhalation 300 mcg per dose	61.00	30 dose	Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler 25 mcg per dose	26.25	120 dose	Serevent
Powder for inhalation 50 mcg per dose	26.25	60 dose	Serevent Accuhaler

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

BUDESONIDE WITH EFORMOTEROL

Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide with 6 mcg eformoterol fumarate metered dose)	41.50	120 dose	DuoResp Spiromax
Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg	33.74	120 dose	Symbicort Turbuhaler
Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mcg eformoterol fumarate metered dose)	82.50	120 dose	DuoResp Spiromax
Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg	33.74	60 dose	Symbicort Turbuhaler

FLUTICASONE FUROATE WITH VILANTEROL

Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose	Breo Ellipta

FLUTICASONE WITH SALMETEROL

Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.79	120 dose	Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg	33.74	60 dose	Seretide Accuhaler
Aerosol inhaler 125 mcg with salmeterol 25 mcg	32.60	120 dose	Seretide
Powder for inhalation 250 mcg with salmeterol 50 mcg	44.08	60 dose	Seretide Accuhaler

Methylxanthines

AMINOPHYLLINE

Inj 25 mg per ml, 10 ml ampoule	180.00	5	DBL Aminophylline

CAFFEINE CITRATE

Oral liq 20 mg per ml (caffeine 10 mg per ml)	16.91	25 ml	Biomed
Inj 20 mg per ml (caffeine 10 mg per ml), 2.5 ml ampoule	69.70	5	Biomed

THEOPHYLLINE

Tab long-acting 250 mg	25.65	100	Nuelin-SR
Oral liq 80 mg per 15 ml	18.49	500 ml	Nuelin

Mucolytics and Expectorants

DORNASE ALFA – **Restricted** see terms below

⬇ Nebuliser soln 2.5 mg per 2.5 ml ampoule	250.00	6	Pulmozyme

➥ **Restricted** (RS1787)

Initiation – cystic fibrosis

Respiratory physician or paediatrician

Re-assessment required after 12 months

All of the following:

1 Patient has a confirmed diagnosis of cystic fibrosis; and
2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
3 Any of the following:
  3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in in the previous 12 month period; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or

3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Continuation – cystic fibrosis

Respiratory physician or paediatrician

The treatment remains appropriate and the patient continues to benefit from treatment.

Initiation – significant mucus production

Limited to 4 weeks treatment

Both:

1 Patient is an in-patient; and

2 The mucus production cannot be cleared by first line chest techniques.

Initiation – pleural emphyema

Limited to 3 days treatment

Both:

1 Patient is an in-patient; and

2 Patient diagnoses with pleural emphyema.

ELEXACAFTOR WITH TEZACAFTOR, IVACAFTOR AND IVACAFTOR – **Restricted** see terms below

⬇	Tab elexacaftor 50 mg with tezacaftor 25 mg, ivacaftor 37.5 mg (56) and ivacaftor 75 mg (28)	27,647.39	84	Trikafta
⬇	Tab elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg (56) and ivacaftor 150 mg (28)	27,647.39	84	Trikafta
⬇	Oral granules elexacaftor 80 mg with tezacaftor 40 mg, ivacaftor 60 mg (28) and ivacaftor 59.5mg (28), sachets	27,647.39	56	Trikafta
⬇	Oral granules elexacaftor 100 mg with tezacaftor 50 mg, ivacaftor 75 mg (28) and ivacaftor 75 mg (28), sachets	27,647.39	56	Trikafta

➥ **Restricted** (RS2186)

Initiation

All of the following:

1 Patient has been diagnosed with cystic fibrosis; and

2 Either:

2.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or

2.2 Patient has a sweat chloride value of at least 60 mmol/L; and

3 Either:

3.1 Patient has a heterozygous or homozygous F508del mutation; or

3.2 Patient has a mutation responsive to elexacaftor/tezacaftor/ivacaftor (see note); and

4 The treatment must be the sole funded CFTR modulator therapy for this condition; and

5 Treatment with elexacaftor/tezacaftor/ivacaftor must be given concomitantly with standard therapy for this condition.

Note: Eligible mutations are listed in the Food and Drug Administration (FDA) Trikafta prescribing information https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/212273s015lbl.pdf

IVACAFTOR – **Restricted** see terms on the next page

⬇	Tab 150 mg	29,386.00	56	Kalydeco
⬇	Oral granules 13.4 mg, sachet	29,386.00	56	Kalydeco
⬇	Oral granules 25 mg, sachet	29,386.00	56	Kalydeco
⬇	Oral granules 50 mg, sachet	29,386.00	56	Kalydeco
⬇	Oral granules 75 mg, sachet	29,386.00	56	Kalydeco

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS2187)
Initiation
All of the following:
1 Patient has been diagnosed with cystic fibrosis; and
2 Either:
2.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
2.2 Patients must have a sweat chloride value of at least 60 mmol/L; and
3 Patient must have at least one mutation on the list of CFTR mutations that produce CFTR protein and are known to be responsive to ivacaftor**; and
4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
5 The dose of ivacaftor will not exceed one tablet or one sachet twice daily.
Note: ** Mutations listed in Table 3 of the Food and Drug Administration (FDA) Ivacaftor prescribing information https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/203188s038lbl.pdf

SODIUM CHLORIDE
Nebuliser soln 7%, 90 ml bottle — 25.73 — 90 ml — Biomed

VANZACAFTOR WITH TEZACAFTOR AND DEUTIVACAFTOR – **Restricted** see terms below
⬇ Tab vanzacaftor 4 mg with tezacaftor 20 mg and deutivacaftor 50 mg — 29,029.76 — 84 — Alyftrek
⬇ Tab vanzacaftor 10 mg with tezacaftor 50 mg and deutivacaftor 125 mg — 29,029.76 — 56 — Alyftrek
➥ **Restricted** (RS2188)
Initiation
All of the following:
1 Patient has been diagnosed with cystic fibrosis; and
2 Either:
2.1 Patient has two cystic fibrosis-causing mutations in the cystic fibrosis transmembrane regulator (CFTR) gene (one from each parental allele); or
2.2 Patient has a sweat chloride value of at least 60 mmol/L; and
3 Either:
3.1 Patient has a heterozygous or homozygous F508del mutation; or
3.2 Patient has a mutation responsive to vanzacaftor/tezacaftor/deutivacaftor (see note); and
4 The treatment must be the sole funded CFTR modulator therapy for this condition; and
5 Treatment with vanzacaftor/tezacaftor/deutivacaftor must be given concomitantly with standard therapy for this condition.
Note: Eligible mutations are listed in the in the Food and Drug Administration (FDA) Alyftrek prescribing information https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/218730s002lbl.pdf

Pulmonary Surfactants

BERACTANT
Soln 200 mg per 8 ml vial

PORACTANT ALFA
Soln 120 mg per 1.5 ml vial — 425.00 — 1 — Curosurf
Soln 240 mg per 3 ml vial — 695.00 — 1 — Curosurf

Respiratory Stimulants

DOXAPRAM
Inj 20 mg per ml, 5 ml vial

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Sclerosing Agents

TALC
- Powder
- Soln (slurry) 100 mg per ml, 50 ml

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Anti-Infective Preparations			
Antibacterials			
CHLORAMPHENICOL			
Eye oint 1% – **5% DV Feb-26 to 2028**	1.55	5 g	**Devatis**
Ear drops 0.5%			
Eye drops 0.5% – **5% DV Mar-26 to 2028**	1.84	10 ml	**Chlorafast**
Eye drops 0.5%, single dose			
CIPROFLOXACIN			
Eye drops 0.3% – **5% DV Mar-25 to 2027**	10.85	5 ml	**Ciprofloxacin Teva**
FRAMYCETIN SULPHATE			
Ear/eye drops 0.5%			
GENTAMICIN SULPHATE			
Eye drops 0.3%			
PROPAMIDINE ISETHIONATE			
Eye drops 0.1%			
SODIUM FUSIDATE [FUSIDIC ACID]			
Eye drops 1%	5.29	5 g	Fucithalmic
SULPHACETAMIDE SODIUM			
Eye drops 10%			
TOBRAMYCIN			
Eye oint 0.3%	10.45	3.5 g	Tobrex
Eye drops 0.3%	11.48	5 ml	Tobrex
Antifungals			
NATAMYCIN			
Eye drops 5%			
Antivirals			
ACICLOVIR			
Eye oint 3% – **5% DV Feb-25 to 2027**	15.89	4.5 g	**ViruPOS**
Combination Preparations			
CIPROFLOXACIN WITH HYDROCORTISONE			
Ear drops ciprofloxacin 0.2% with 1% hydrocortisone	16.30	10 ml	Ciproxin HC Otic
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml			
DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE			
Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	5.39	3.5 g	Maxitrol
Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	4.50	5 ml	Maxitrol
DEXAMETHASONE WITH TOBRAMYCIN			
Eye drops 0.1% with tobramycin 0.3%	12.64	5 ml	Tobradex

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
FLUMETASONE PIVALATE WITH CLIOQUINOL			
Ear drops 0.02% with clioquinol 1%			
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN			
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml	Kenacomb

Anti-Inflammatory Preparations

Corticosteroids

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
DEXAMETHASONE			
Eye oint 0.1%	5.86	3.5 g	Maxidex
Eye drops 0.1%	4.50	5 ml	Maxidex
⬇ Ocular implant 700 mcg	1,444.50	1	Ozurdex

➥ **Restricted** (RS1606)

Initiation – Diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

All of the following:

1 Patients have diabetic macular oedema with pseudophakic lens; and
2 Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision; and
3 Either:
  3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Continuation – Diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

Both:

1 Patient's vision is stable or has improved (prescriber determined); and
2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initiation – Women of child bearing age with diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

All of the following:

1 Patients have diabetic macular oedema; and
2 Patient has reduced visual acuity of between 6/9 – 6/48 with functional awareness of reduction in vision; and
3 Patient is of child bearing potential and has not yet completed a family; and
4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Continuation – Women of child bearing age with diabetic macular oedema

Ophthalmologist

Re-assessment required after 12 months

All of the following:

1 Patient's vision is stable or has improved (prescriber determined); and
2 Patient is of child bearing potential and has not yet completed a family; and
3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
FLUOROMETHOLONE			
Eye drops 0.1%	3.09	5 ml	FML
PREDNISOLONE ACETATE			
Eye drops 0.12%			
Eye drops 1%	7.00	5 ml	Pred Forte
	6.92	10 ml	Prednisolone- AFT
PREDNISOLONE SODIUM PHOSPHATE			
Eye drops 0.5%, single dose (preservative free)	43.26	20 dose	Minims Prednisolone

Non-Steroidal Anti-Inflammatory Drugs

DICLOFENAC SODIUM			
Eye drops 0.1%			
Eye drops 0.1%, single dose – **5% DV Jul-25 to 2027**	1.85	10 dose	**Diclofenac Devatis**
	5.54	30 dose	**Diclofenac Devatis**
KETOROLAC TROMETAMOL			
Eye drops 0.5%			

Decongestants and Antiallergics

Antiallergic Preparations

LEVOCABASTINE			
Eye drops 0.05%			
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml	Lomide
OLOPATADINE			
Eye drops 0.1% – **5% DV Mar-26 to 2028**	3.39	5 ml	**Olopatadine Teva**
SODIUM CROMOGLICATE			
Eye drops 2% – **5% DV Mar-26 to 2028**	2.91	10 ml	**Allerfix**

Decongestants

NAPHAZOLINE HYDROCHLORIDE			
Eye drops 0.1% – **5% DV Jan-25 to 2027**	5.65	15 ml	**Albalon**

Diagnostic and Surgical Preparations

Diagnostic Dyes

FLUORESCEIN SODIUM			
Eye drops 2%, single dose			
Inj 10%, 5 ml vial	125.00	12	Fluorescite
Ophthalmic strips 1 mg			
FLUORESCEIN SODIUM WITH LIGNOCAINE HYDROCHLORIDE			
Eye drops 0.25% with lignocaine hydrochloride 4%, single dose			
LISSAMINE GREEN			
Ophthalmic strips 1.5 mg			
ROSE BENGAL SODIUM			
Ophthalmic strips 1%			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Irrigation Solutions

MIXED SALT SOLUTION FOR EYE IRRIGATION

Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 15 ml dropper bottle	5.00	15 ml	Balanced Salt Solution
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 250 ml			*e.g. Balanced Salt Solution*
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml bag			*e.g. Balanced Salt Solution*
Eye irrigation solution calcium chloride 0.048% with magnesium chloride 0.03%, potassium chloride 0.075%, sodium acetate 0.39%, sodium chloride 0.64% and sodium citrate 0.17%, 500 ml bottle	10.50	500 ml	Balanced Salt Solution

Ocular Anaesthetics

OXYBUPROCAINE HYDROCHLORIDE
- Eye drops 0.4%, single dose

PROXYMETACAINE HYDROCHLORIDE
- Eye drops 0.5%

TETRACAINE [AMETHOCAINE] HYDROCHLORIDE
- Eye drops 0.5%, single dose
- Eye drops 1%, single dose

Viscoelastic Substances

HYPROMELLOSE
- Inj 2%, 1 ml syringe
- Inj 2%, 2 ml syringe

SODIUM HYALURONATE [HYALURONIC ACID]

Inj 18 mg per ml, 0.85 ml syringe – **5% DV Mar-26 to 2028**	50.00	1	**Healon GV Pro**
Inj 23 mg per ml, 0.6 ml syringe – **5% DV Mar-26 to 2028**	60.00	1	**Healon 5**
Inj 10 mg per ml, 0.85 ml syringe – **5% DV Mar-26 to 2028**	28.50	1	**Healon**

SODIUM HYALURONATE [HYALURONIC ACID] WITH CHONDROITIN SULPHATE

Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.35 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.4 ml syringe	64.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.5 ml syringe and inj 10 mg sodium hyaluronate [hyaluronic acid] per ml, 0.55 ml syringe	74.00	1	Duovisc
Inj 30 mg per ml with chondroitin sulphate 40 mg per ml, 0.75 ml syringe	67.00	1	Viscoat

Other

DISODIUM EDETATE
- Inj 150 mg per ml, 20 ml ampoule
- Inj 150 mg per ml, 20 ml vial
- Inj 150 mg per ml, 100 ml vial

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
RIBOFLAVIN 5-PHOSPHATE			
Soln trans epithelial riboflavin			
Inj 0.1%			
Inj 0.1% plus 20% dextran T500			
Inj 0.1% plus hydroxypropyl methylcellulose			

Glaucoma Preparations

Beta Blockers

BETAXOLOL			
Eye drops 0.25%			
Eye drops 0.5%			
TIMOLOL			
Eye drops 0.25% – **5% DV Mar-24 to 2026**	2.42	5 ml	**Arrow-Timolol**
Eye drops 0.5% – **5% DV Mar-24 to 2026**	2.50	5 ml	**Arrow-Timolol**
➥ Eye drops 0.5%, gel forming – **Restricted:** For continuation only			

Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE			
Tab 250 mg – **5% DV Sep-25 to 2027**	13.96	100	**Medsurge**
Inj 500 mg			
BRINZOLAMIDE			
Eye drops 1% – **5% DV Dec-24 to 2027**	5.11	5 ml	**Azopt**
DORZOLAMIDE – **Restricted:** For continuation only			
➥ Eye drops 2%			
DORZOLAMIDE WITH TIMOLOL			
Eye drops 2% with timolol 0.5% – **5% DV Feb-25 to 2027**	3.58	5 ml	**Dortimopt**

Miotics

ACETYLCHOLINE CHLORIDE			
Inj 20 mg vial with diluent			
CARBACHOL			
Inj 150 mcg vial			
PILOCARPINE HYDROCHLORIDE			
Eye drops 1%	4.26	15 ml	Isopto Carpine
Eye drops 2%	5.35	15 ml	Isopto Carpine
Eye drops 4%	7.99	15 ml	Isopto Carpine
PILOCARPINE NITRATE			
Eye drops 2%, single dose			

Prostaglandin Analogues

BIMATOPROST			
Eye drops 0.03% – **5% DV Jan-25 to 2027**	5.15	3 ml	**Lumigan**
LATANOPROST			
Eye drops 0.005% – **5% DV Mar-25 to 2027**	2.08	2.5 ml	**Teva**
LATANOPROST WITH TIMOLOL			
Eye drops 0.005% with timolol 0.5% – **5% DV Mar-24 to 2026**	4.95	2.5 ml	**Arrow - Lattim**

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
TRAVOPROST			
Eye drops 0.004% – **5% DV Dec-24 to 2027**	6.80	2.5 ml	**Travatan**

Sympathomimetics

APRACLONIDINE			
Eye drops 0.5%	19.77	5 ml	Iopidine
BRIMONIDINE TARTRATE			
Eye drops 0.2% – **5% DV Mar-25 to 2027**	5.16	5 ml	**Arrow-Brimonidine**
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE			
Eye drops 0.2% with timolol 0.5% – **5% DV Dec-24 to 2027**	7.13	5 ml	**Combigan**

Mydriatics and Cycloplegics

Anticholinergic Agents

ATROPINE SULPHATE			
Eye drops 0.5%			
Eye drops 1%, single dose			
Eye drops 1% – **5% DV Feb-24 to 2026**	18.27	15 ml	**Atropt**
CYCLOPENTOLATE HYDROCHLORIDE			
Eye drops 0.5%, single dose			
Eye drops 1%	25.16	15 ml	Cyclogyl
Eye drops 1%, single dose			
TROPICAMIDE			
Eye drops 0.5%	20.52	15 ml	Mydriacyl
Eye drops 0.5%, single dose			
Eye drops 1%	24.82	15 ml	Mydriacyl
Eye drops 1%, single dose			

Sympathomimetics

PHENYLEPHRINE HYDROCHLORIDE			
Eye drops 2.5%, single dose			
Eye drops 10%, single dose			

Ocular Lubricants

CARBOMER			
Ophthalmic gel 0.2%			
CARMELLOSE SODIUM			
Eye drops 0.5%			
Eye drops 0.5%, single dose			
Eye drops 1%			
Eye drops 1%, single dose			
HYPROMELLOSE			
Eye drops 0.5%	19.50	15 ml	Methopt
Ophthalmic gel 0.3%			
HYPROMELLOSE WITH DEXTRAN			
Eye drops 0.3% with dextran 0.1%	2.30	15 ml	Poly-Tears
Eye drops 0.3% with dextran 0.1%, single dose			

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
PARAFFIN LIQUID WITH SOFT WHITE PARAFFIN			
Eye oint 42.5% with soft white paraffin 57.3%			
PARAFFIN LIQUID WITH WOOL FAT			
Eye oint 3% with wool fat 3%	3.63	3.5 g	Poly-Visc
POLYETHYLENE GLYCOL 400 AND PROPYLENE GLYCOL			
Eye drops 0.4% with propylene glycol 0.3%, 10 ml bottle			
Note: Only for use in compounding an eye drop formulation			
Eye drops 0.4% with propylene glycol 0.3% preservative free, single dose	10.06	28	Systane Unit Dose
POLYVINYL ALCOHOL WITH POVIDONE			
Eye drops 1.4% with povidone 0.6%, single dose			
RETINOL PALMITATE			
Oint 138 mcg per g	3.80	5 g	VitA-POS
SODIUM HYALURONATE [HYALURONIC ACID]			
Eye drops 1 mg per ml – **5% DV Dec-24 to 2027**	13.58	10 ml	**Hylo-Fresh**

Other Otological Preparations

ACETIC ACID WITH PROPYLENE GLYCOL
Ear drops 2.3% with propylene glycol 2.8%

DOCUSATE SODIUM
Ear drops 0.5%

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE
Tab eff 200 mg
Inj 200 mg per ml, 10 ml vial ..... 42.99 10 Hikma Acetylcysteine
Inj 200 mg per ml, 10 ml ampoule – **5% DV Apr-25 to 2027** ..... 42.99 10 **DBL Acetylcysteine**

AMYL NITRITE
Liq 98% in 3 ml capsule

DIGOXIN IMMUNE FAB
Inj 38 mg vial
Inj 40 mg vial

ETHANOL
Liq 96%

ETHANOL WITH GLUCOSE
Inj 10% with glucose 5%, 500 ml bottle

ETHANOL, DEHYDRATED
Inj 100%, 5 ml ampoule
Inj 96%

FLUMAZENIL
Inj 0.1 mg per ml, 5 ml ampoule – **5% DV Dec-24 to 2027** ..... 44.00 5 **Flumazenil-Baxter**

HYDROXOCOBALAMIN
Inj 5 g vial
Inj 2.5 g vial

NALOXONE HYDROCHLORIDE
Inj 400 mcg per ml, 1 ml ampoule – **5% DV Apr-25 to 2027** ..... 13.29 5 **DBL Naloxone Hydrochloride**

PRALIDOXIME CHLORIDE
Inj 1 g vial

PRALIDOXIME IODIDE
Inj 25 mg per ml, 20 ml ampoule

SODIUM NITRITE
Inj 30 mg per ml, 10 ml ampoule

SODIUM THIOSULFATE
Inj 250 mg per ml, 100 ml vial
Inj 250 mg per ml, 10 ml vial
Inj 250 mg per ml. 50 ml vial
Inj 500 mg per ml, 10 ml vial
Inj 500 mg per ml, 20 ml ampoule

SOYA OIL
Inj 20%, 500 ml bag
Inj 20%, 500 ml bottle

Antitoxins

BOTULISM ANTITOXIN
Inj 250 ml vial

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
DIPHTHERIA ANTITOXIN			
Inj 10,000 iu vial			

Antivenoms

RED BACK SPIDER ANTIVENOM
Inj 500 u vial

SNAKE ANTIVENOM
Inj 50 ml vial

Removal and Elimination

CHARCOAL
Oral liq 200 mg per ml ..... 59.85 — 250 ml — Carbasorb-X

DEFERASIROX – **Restricted** see terms below

⬇	Tab 125 mg dispersible	276.00	28	Exjade
⬇	Tab 250 mg dispersible	552.00	28	Exjade
⬇	Tab 500 mg dispersible	1,105.00	28	Exjade

➥ **Restricted** (RS1444)

Initiation
Haematologist
Re-assessment required after 2 years
All of the following:

1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
3 Any of the following:
  3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
  3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
  3.3 Treatment with deferiprone has resulted in arthritis; or
  3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per µL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per µL).

Continuation
Haematologist
Re-assessment required after 2 years
Either:

1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels. .

DEFERIPRONE – **Restricted** see terms below

⬇	Tab 500 mg	533.17	100	Ferriprox
⬇	Oral liq 100 mg per ml	266.59	250 ml	Ferriprox

➥ **Restricted** (RS1445)

Initiation
Patient has been diagnosed with chronic iron overload due to congenital inherited anaemia or acquired red cell aplasia.

DESFERRIOXAMINE MESILATE
Inj 500 mg vial ..... 332.88 — 10 — DBL Desferrioxamine Mesylate for Inj BP

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
DICOBALT EDETATE			
Inj 15 mg per ml, 20 ml ampoule			
DIMERCAPROL			
Inj 50 mg per ml, 2 ml ampoule			
DIMERCAPTOSUCCINIC ACID			
Cap 100 mg			*e.g. PCNZ, Optimus Healthcare, Chemet*
Cap 200 mg			*e.g. PCNZ, Optimus Healthcare, Chemet*
SODIUM CALCIUM EDETATE			
Inj 50 mg per ml, 10 ml ampoule			
Inj 200 mg per ml, 2.5 ml ampoule			
Inj 200 mg per ml, 5 ml ampoule			

Antiseptics and Disinfectants

CHLORHEXIDINE			
Soln 0.1%			
Soln 4%			
Soln 5%	15.50	500 ml	healthE
CHLORHEXIDINE WITH CETRIMIDE			
Crm 0.1% with cetrimide 0.5%			
Foaming soln 0.5% with cetrimide 0.5%			
CHLORHEXIDINE WITH ETHANOL			
Soln 0.5% with ethanol 70%			
Soln 2% with ethanol 70%			
Soln 0.5% with ethanol 70%, non-staining (pink) 25 ml	1.55	1	healthE
IODINE WITH ETHANOL			
Soln 1% with ethanol 70%			
ISOPROPYL ALCOHOL			
Soln 70%, 500 ml	5.65	1	healthE
POVIDONE-IODINE			
⬇ Vaginal tab 200 mg			
➥ **Restricted** (RS1354)			
Initiation			
Rectal administration pre-prostate biopsy.			
Oint 10%	7.40	65 g	Betadine
Soln 10%	4.99	100 ml	Riodine
Soln 5%			
Soln 7.5%			
Soln 10%,	3.83	15 ml	Riodine
	6.99	500 ml	Riodine
Pad 10%			
Swab set 10%			
POVIDONE-IODINE WITH ETHANOL			
Soln 10% with ethanol 30%			
Soln 10% with ethanol 70%			

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
SODIUM HYPOCHLORITE			
Soln			

Contrast Media

Iodinated X-ray Contrast Media

DIATRIZOATE MEGLUMINE WITH SODIUM AMIDOTRIZOATE			
Oral liq 660 mg per ml with sodium amidotrizoate 100 mg per ml, 100 ml bottle	30.00	100 ml	Gastrografin
Inj 260 mg with sodium amidotrizoate 40 mg per ml, 250 ml bottle	120.00	1	Urografin
DIATRIZOATE SODIUM			
Oral liq 370 mg per ml, 10 ml sachet	156.12	50	Ioscan
IODISED OIL			
Inj 38% w/w (480 mg per ml), 10 ml ampoule	410.00	1	Lipiodol Ultra Fluid
IODIXANOL			
Inj 270 mg per ml (iodine equivalent), 50 ml bottle	275.00	10	Visipaque
Inj 270 mg per ml (iodine equivalent), 100 ml bottle	505.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 50 ml bottle	280.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 100 ml bottle	510.00	10	Visipaque
Inj 320 mg per ml (iodine equivalent), 200 ml bottle	1,020.00	10	Visipaque
IOHEXOL			
Inj 240 mg per ml (iodine equivalent), 50 ml bottle	117.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 10 ml bottle	91.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 20 ml bottle	110.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 50 ml bottle	121.00	10	Omnipaque
Inj 300 mg per ml (iodine equivalent), 100 ml bottle	200.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 50 ml bottle	125.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 100 ml bottle	210.00	10	Omnipaque
Inj 350 mg per ml (iodine equivalent), 200 ml bottle	420.00	10	Omnipaque
Inj 350 mg per ml, 500 ml bottle	655.00	6	Omnipaque

(Omnipaque Inj 300 mg per ml (iodine equivalent), 20 ml bottle to be delisted 1 October 2026)

Non-iodinated X-ray Contrast Media

BARIUM SULPHATE			
Oral liq 400 mg per ml (40% w/v, 30% w/w), bottle	17.39	148 g	Varibar - Thin Liquid
Oral liq 400 mg per ml (40% w/v), bottle	189.15	250 ml	Varibar - Honey
	38.40	240 ml	Varibar - Nectar
	159.05	230 ml	Varibar - Pudding
Grans for oral liq 960 mg per g (96% w/w), 176 g bottle	530.00	24	Vanilla SilQ MD
Grans for oral liq 980 mg per g (98% w/w), 310 g bottle	490.00	24	Vanilla SilQ HD
Oral liq 20.9 mg per ml (2.1% w/v, 2% w/w), 450 ml bottle	97.50	12	Readi-CAT 2
Oral liq 1 mg per ml (0.1% w/v, 0.1% w/w), 450 ml bottle	15.95	1	Neulumex
	191.40	12	Neulumex
Oral liq 400 mg per ml (40% w/v, 30% w/w), 20 ml bottle	52.35	3	Tagitol V
CITRIC ACID WITH SODIUM BICARBONATE			
Powder 382.2 mg per g with sodium bicarbonate 551.3 mg per g, 4 g sachet	90.25	50 g	E-Z-Gas II

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Paramagnetic Contrast Media			
GADOBUTROL			
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 5 ml prefilled syringe	126.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 7.5 ml prefilled syringe	189.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 10 ml prefilled syringe	245.00	5	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 15 ml vial	735.00	10	Gadovist 1.0
Inj 604.72 mg per ml (equivalent to 1 mmol per ml), 65 ml bottle	3,120.00	10	Gadovist 1.0
GADOTERIC ACID			
Inj 279.30 mg per ml, 10 ml prefilled syringe			*e.g. Clariscan*
Inj 279.30 mg per ml, 10 ml vial			*e.g. Clariscan*
Inj 279.30 mg per ml, 15 ml prefilled syringe			*e.g. Clariscan*
Inj 279.30 mg per ml, 20 ml vial			*e.g. Clariscan*
Inj 279.30 mg per ml, 5 ml vial			*e.g. Clariscan*
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml prefilled syringe	172.00	10	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml bottle	25.35	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 15 ml prefilled syringe	258.00	10	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml prefilled syringe	344.00	10	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 10 ml bottle	14.30	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 20 ml bottle	28.90	1	Dotarem
Inj 279.32 mg per ml (0.5 mmol per ml), 5 ml bottle	9.10	1	Dotarem
GADOXETATE DISODIUM			
Inj 181.43 mg per ml (equivalent to 0.25 mmol per ml), 10 ml prefilled syringe	300.00	1	Primovist
MEGLUMINE GADOPENTETATE			
Inj 469 mg per ml, 10 ml prefilled syringe	95.00	5	Magnevist
Inj 469 mg per ml, 10 ml vial	185.00	10	Magnevist
MEGLUMINE IOTROXATE			
Inj 105 mg per ml, 100 ml bottle	169.15	100 ml	Biliscopin
Ultrasound Contrast Media			
PERFLUTREN			
Inj 1.1 mg per ml, 1.5 ml vial	180.00	1	Definity
	720.00	4	Definity
Diagnostic Agents			
ARGININE			
Inj 50 mg per ml, 500 ml bottle			
Inj 100 mg per ml, 300 ml bottle			
HISTAMINE ACID PHOSPHATE			
Nebuliser soln 0.6%, 10 ml vial			
Nebuliser soln 2.5%, 10 ml vial			
Nebuliser soln 5%, 10 ml vial			
MANNITOL			
Powder for inhalation			*e.g. Aridol*

⬆ Item restricted (see ➡ above); ⬇ Item restricted (see ➡ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
METHACHOLINE CHLORIDE			
Powder 100 mg			
SECRETIN PENTAHYDROCHLORIDE			
Inj 100 u vial			
Inj 80 u vial			
Inj 100 u ampoule			
SINCALIDE			
Inj 5 mcg per vial			

Diagnostic Dyes

BONNEY'S BLUE DYE			
Soln			
INDIGO CARMINE			
Inj 4 mg per ml, 5 ml ampoule			
Inj 8 mg per ml, 5 ml ampoule			
INDOCYANINE GREEN			
Inj 25 mg vial			
METHYLTHIONINIUM CHLORIDE [METHYLENE BLUE]			
Inj 5 mg per ml, 10 ml ampoule	259.57	5	Proveblue
PATENT BLUE V			
Inj 2.5%, 2 ml ampoule	440.00	5	Obex Medical
Inj 2.5%, 5 ml prefilled syringe	435.00	5	InterPharma

Irrigation Solutions

CHLORHEXIDINE WITH CETRIMIDE

⬇ Irrigation soln 0.015% with cetrimide 0.15%, 500 ml bottle

➥ **Restricted** (RS1683)

Initiation

Re-assessment required after 3 months

All of the following:

1 Patient has burns that are greater than 30% of total body surface area (BSA); and
2 For use in the perioperative preparation and cleansing of large burn areas requiring debridement/skin grafting; and
3 The use of 30 ml ampoules is impractical due to the size of the area to be covered.

Continuation

Re-assessment required after 3 months

The treatment remains appropriate for the patient and the patient is benefiting from the treatment.

Irrigation soln 0.015% with cetrimide 0.15%, 100 ml bottle			
Irrigation soln 0.015% with cetrimide 0.15%, 30 ml ampoule – **5% DV**			
Sep-25 to 2028	29.70	30	**LumaCina**
GLYCINE			
Irrigation soln 1.5%, 3,000 ml bag	96.28	4	B Braun
SODIUM CHLORIDE			
Irrigation soln 0.9%, 3,000 ml bag	80.00	4	B Braun
Irrigation soln 0.9%, 30 ml ampoule	13.25	20	InterPharma
Irrigation soln 0.9%, 1,000 ml bottle	19.50	10	Baxter Sodium Chloride 0.9%
Irrigation soln 0.9%, 250 ml bottle	24.60	12	Fresenius Kabi

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
WATER			
Irrigation soln, 3,000 ml bag	84.52	4	B Braun
Irrigation soln, 1,000 ml bottle	19.50	10	Baxter Water for Irrigation
Irrigation soln, 250 ml bottle	26.40	12	Fresenius Kabi

Surgical Preparations

BISMUTH SUBNITRATE AND IODOFORM PARAFFIN
- Paste

DIMETHYL SULFOXIDE
- Soln 50%
- Soln 99%

PHENOL
- Inj 6%, 10 ml ampoule

PHENOL WITH IOXAGLIC ACID
- Inj 12%, 10 ml ampoule

SODIUM HYDROXIDE
- Soln 10%

TROMETAMOL
- Inj 36 mg per ml, 500 ml bottle

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Cardioplegia Solutions

ELECTROLYTES

Inj 15 mmol/l sodium chloride, 9 mmol/l potassium chloride, 1 mmol/l potassium hydrogen 2-ketoglutarate, 4 mmol/l magnesium chloride, 18 mmol/l histidine hydrochloride, 180 mmol/l histidine, 2 mmol/l tryptophan, 30 mmol/l mannitol, 0.015 mmol/l calcium chloride, 1,000 ml bag — *e.g. Custodiol-HTK*

Inj aspartic acid 10.43 mg per ml, citric acid 0.22476 mg per ml, glutamic acid 11.53 mg per ml, sodium phosphate 0.1725 mg per ml, potassium chloride 2.15211 mg per ml, sodium citrate 1.80768 mg per ml, sodium hydroxide 6.31 mg per ml and trometamol 11.2369 mg per ml, 364 ml bag — *e.g. Cardioplegia Enriched Paed. Soln.*

Inj aspartic acid 8.481 mg per ml, citric acid 0.8188 mg per ml, glutamic acid 9.375 mg per ml, sodium phosphate 0.6285 mg per ml, potassium chloride 2.5 mg per ml, sodium citrate 6.585 mg per ml, sodium hydroxide 5.133 mg per ml and trometamol 9.097 mg per ml, 527 ml bag — *e.g. Cardioplegia Enriched Solution*

Inj citric acid 0.07973 mg per ml, sodium phosphate 0.06119 mg per ml, potassium chloride 2.181 mg per ml, sodium chloride 1.788 mg ml, sodium citrate 0.6412 mg per ml and trometamol 5.9 mg per ml, 523 ml bag — *e.g. Cardioplegia Base Solution*

Inj 110 mmol/l sodium, 16 mmol/l potassium, 1.2 mmol/l calcium, 16 mmol/l magnesium and 160 mmol/l chloride, 1,000 ml bag — *e.g. Cardioplegia Solution AHB7832*

Inj 143 mmol/l sodium, 16 mmol/l potassium, 16 mmol/l magnesium and 1.2 mmol/l calcium, 1,000 ml bag — *e.g. Cardioplegia Electrolyte Solution*

MONOSODIUM GLUTAMATE WITH SODIUM ASPARTATE

Inj 42.68 mg with sodium aspartate 39.48 mg per ml, 250 ml bottle

MONOSODIUM L-ASPARTATE

Inj 14 mmol per 10 ml, 10 ml

Cold Storage Solutions

SODIUM WITH POTASSIUM

Inj 29 mmol/l with potassium 125 mmol/l, 1,000 ml bag

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Extemporaneously Compounded Preparations

ACETIC ACID
Liq

ALUM
Powder BP

ARACHIS OIL [PEANUT OIL]
Liq

ASCORBIC ACID
Powder

BENZOIN
Tincture compound BP

BISMUTH SUBGALLATE
Powder

BORIC ACID
Powder

CARBOXYMETHYLCELLULOSE
Soln 1.5%

CETRIMIDE
Soln 40%

CHLORHEXIDINE GLUCONATE
Soln 20 %

CHLOROFORM
Liq BP

CITRIC ACID
Powder BP

CLOVE OIL
Liq

COAL TAR
Soln BP ..... 46.00 — 200 ml — Midwest

CODEINE PHOSPHATE
Powder

COLLODION FLEXIBLE
Liq

COMPOUND HYDROXYBENZOATE
Soln ..... 36.00 — 100 ml — Midwest

CYSTEAMINE HYDROCHLORIDE
Powder

DISODIUM HYDROGEN PHOSPHATE WITH SODIUM DIHYDROGEN PHOSPHATE
Inj 37.46 mg with sodium dihydrogen phosphate 47.7 mg in 1.5 ml ampoule

DITHRANOL
Powder

GLUCOSE [DEXTROSE]
Powder

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
GLYCERIN WITH SODIUM SACCHARIN			
Suspension	38.00	473 ml	Ora-Sweet SF
GLYCERIN WITH SUCROSE			
Suspension	38.00	473 ml	Ora-Sweet
GLYCEROL			
Liq	3.23	500 ml	healthE Glycerol BP Liquid
HYDROCORTISONE			
Powder	49.95	25 g	ABM
LACTOSE			
Powder			
MAGNESIUM HYDROXIDE			
Paste			
MENTHOL			
Crystals			
METHADONE HYDROCHLORIDE			
Powder			
METHYL HYDROXYBENZOATE			
Powder	11.00	25 g	Midwest
METHYLCELLULOSE			
Powder	44.00	100 g	Midwest
Suspension	38.00	473 ml	Ora-Plus
(Midwest Powder to be delisted 1 February 2028)			
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN			
Suspension	38.00	473 ml	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE			
Suspension	38.00	473 ml	Ora-Blend
OLIVE OIL			
Liq			
PARAFFIN			
Liq			
PHENOBARBITONE SODIUM			
Powder			
PHENOL			
Liq			
PILOCARPINE NITRATE			
Powder			
POLYHEXAMETHYLENE BIGUANIDE			
Liq			
POVIDONE K30			
Powder			
SALICYLIC ACID			
Powder			
SILVER NITRATE			
Crystals			
SODIUM BICARBONATE			
Powder BP	13.50	500 g	Midwest

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
SODIUM CITRATE			
Powder			
SODIUM METABISULFITE			
Powder			
STARCH			
Powder			
SULPHUR			
Precipitated			
Sublimed			
SYRUP			
Liq (pharmaceutical grade)	25.00	500 ml	Midwest
THEOBROMA OIL			
Oint			
TRI-SODIUM CITRATE			
Crystals			
TRICHLORACETIC ACID			
Grans			
UREA			
Powder BP			
WOOL FAT			
Oint, anhydrous			
XANTHAN			
Gum 1%			
ZINC OXIDE			
Powder			

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Food Modules

Carbohydrate

➦ **Restricted** (RS1467)
Initiation – Use as an additive
Any of the following:

1 Cystic fibrosis; or
2 Chronic kidney disease; or
3 Cancer in children; or
4 Cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
5 Faltering growth in an infant/child; or
6 Bronchopulmonary dysplasia; or
7 Premature and post premature infant; or
8 Inborn errors of metabolism.

Initiation – Use as a module
For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.
Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

CARBOHYDRATE SUPPLEMENT – **Restricted** see terms above

⬆	Powder 96 g carbohydrate per 100 g, can	6.72	400 g	Polycal

Fat

➦ **Restricted** (RS1468)
Initiation – Use as an additive
Any of the following:

1 Patient has inborn errors of metabolism; or
2 Faltering growth in an infant/child; or
3 Bronchopulmonary dysplasia; or
4 Fat malabsorption; or
5 Lymphangiectasia; or
6 Short bowel syndrome; or
7 Infants with necrotising enterocolitis; or
8 Biliary atresia; or
9 For use in a ketogenic diet; or
10 Chyle leak; or
11 Ascites; or
12 Patient has increased energy requirements, and for whom dietary measures have not been successful.

Initiation – Use as a module
For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk. .
Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

LONG-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms above

⬆	Liquid 50 g fat per 100 ml, bottle	15.38	200 ml	Calogen (neutral) Calogen (strawberry)

MEDIUM-CHAIN TRIGLYCERIDE SUPPLEMENT – **Restricted** see terms above

⬆	Liquid 95 g fat per 100 ml, bottle	37.50	500 ml	MCT Oil
⬆	Liquid 50 g fat per 100 ml, 250 ml bottle	143.65	4	Liquigen

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
WALNUT OIL – **Restricted** see terms on the previous page			
⬆ Liq			

Protein

➥ **Restricted** (RS1469)

Initiation – Use as an additive

Either:

1 Protein losing enteropathy; or
2 High protein needs.

Initiation – Use as a module

For use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk. .

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

PROTEIN SUPPLEMENT – **Restricted** see terms above			
⬆ Powder 5 g protein, 0.67 g carbohydrate and 0.6 g fat per 6.6 g, 275 g can			
⬆ Powder 6 g protein per 7 g, can	8.95	227 g	Resource Beneprotein
⬆ Powder 89 g protein, less than 1.5 g carbohydrate and 2 g fat per 100 g, can	13.82	225 g	Protifar

Other Supplements

CARBOHYDRATE AND FAT SUPPLEMENT – **Restricted** see terms below			
⬇ Powder 72.7 g carbohydrate and 22.3 g fat per 100 g, can	71.77	400 g	Duocal Super Soluble Powder

➥ **Restricted** (RS1212)

Initiation

Both:

1 Infant or child aged four years or under; and
2 Any of the following:
  2.1 Cystic fibrosis; or
  2.2 Cancer in children; or
  2.3 Faltering growth; or
  2.4 Bronchopulmonary dysplasia; or
  2.5 Premature and post premature infants.

HUMAN MILK FORTIFIER			
Powder 0.325 g protein, 0.37 g carbohydrate and 0.175 g fat per 1 g sachet	33.48	50	Human Milk Fortifier *e.g. FM 85*
Powder 0.2 g protein, 0.7 g carbohydrate and 0.02 g fat per 1 g sachet			

Food/Fluid Thickeners

NOTE:

While pre-thickened drinks and supplements have not been included in Section H, Health NZ Hospitals may continue to use such products for patients with dysphagia, provided that:

- use was established prior to 1 July 2013; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

- the product has not been specifically considered and excluded by Pharmac; and
- use of the product conforms to any applicable indication restrictions for similar products that are listed in Section H (for example, use of thickened high protein products should be in line with the restriction for high protein oral feed in Section H).

Pharmac intends to make a further decision in relation to pre-thickened drinks and supplements in the future, and will notify of any change to this situation.

CAROB BEAN GUM WITH MAIZE STARCH AND MALTODEXTRIN			
Powder	24.00	380 g	Aptamil Feed Thickener
GUAR GUM			
Powder			*e.g. Guarcol*
MAIZE STARCH			
Powder	8.29	300 g	Nutilis
MALTODEXTRIN WITH XANTHAN GUM			
Powder			*e.g. Instant Thick*
MALTODEXTRIN WITH XANTHAN GUM AND ASCORBIC ACID			
Powder			*e.g. Easy Thick*

Metabolic Products

➥ **Restricted** (RS2047)
Initiation
Either:

1 For the dietary management of inherited metabolic disease; or
2 Patient has adrenoleukodystrophy.

Supplements for Glutaric Aciduria Type 1

AMINO ACID FORMULA (WITHOUT LYSINE AND LOW TRYPTOPHAN) – **Restricted** see terms above

⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can			*e.g. GA1 Anamix Infant*
⬆ Powder 25 g protein and 51 g carbohydrate per 100 g, 500 g can			*e.g. XLYS Low TRY Maxamaid*

AMINO ACID FORMULA (WITHOUT LYSINE) – **Restricted** see terms above

⬆ Powder (neutral) 5 g protein, 5.4 g carbohydrate, 2.3 g fat and 2 g fibre per 18 g sachet	750.30	30	GA1 Anamix Junior
⬆ Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet	349.65	30	GA Explore 5
⬆ Powder, 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 3.7 g fibre per 100 g, 400 g can	260.00	400 g	GA1 Anamix Infant

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Supplements for Homocystinuria			
AMINO ACID FORMULA (WITHOUT METHIONINE) – **Restricted** see terms on the previous page			
⬆ Powder (neutral), 10 g protein, 11.5 g carbohydrate and 4.5 g fat per 36 g sachet	750.30	30	HCU Anamix Junior
⬆ Powder, 15 g protein, 3.5 g carbohydrate, 0.55 g fat per 25 g sachet	1,048.95	30	HCU Express 15
⬆ Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet	349.65	30	HCU Explore 5
⬆ Powder (neutral) 39 g protein and 34 g carbohydrate per 100 g, 500 g can	480.42	500 g	XMET Maxamum
⬆ Powder (unflavoured) 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	260.00	400 g	HCU Anamix Infant
⬆ Liquid (juicy berries), 20 g protein, 12.63 g carbohydrate and 0.46 g fat per 125 ml bottle	1,684.80	30	HCU Lophlex LQ
⬆ Liquid (orange), 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle	941.40	36	HCU Anamix Junior LQ
Supplements for MSUD and Short chain enoyl coA hydratase deficiency			
AMINO ACID FORMULA (WITHOUT ISOLEUCINE, LEUCINE AND VALINE) – **Restricted** see terms on the previous page			
⬆ Powder (neutral) 10 g protein, 11.5 g carbohydrate and 4.5 g fat per 36 g sachet	750.00	30	MSUD Anamix Junior
⬆ Powder, 15 g protein, 3.5 g carbohydrate, 0.6 g fat per 25 g sachet	1,048.95	30	MSUD Express 15
⬆ Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet	349.65	30	MSUD Explore 5
⬆ Powder (orange) 39 g protein and 34 g carbohydrate per 100 g, 500 g can	454.71	500 g	MSUD Maxamum
⬆ Powder (unflavoured) 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	260.00	400 g	MSUD Anamix Infant
⬆ Powder (unflavoured) 39 g protein and 34 g carbohydrate per 100 g, 500 g can	454.71	500 g	MSUD Maxamum
⬆ Liquid (juicy berries), 20 g protein, 12.63 g carbohydrate and 0.46 g fat per 125 ml pouch	1,684.80	30	MSUD Lophlex LQ 20
⬆ Liquid (orange) 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle	941.40	36	MSUD Anamix Junior LQ

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

Supplements for Phenylketonuria

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
AMINO ACID FORMULA (WITHOUT PHENYLALANINE) – **Restricted** see terms on page 295			
⬆ Tab 8.33 mg	99.00	75	Phlexy 10
⬆ Powder (Berry), 5.0 g protein, 14 g carbohydrate, 0 g fat per 20 g sachet	449.28	60	PKU Restore Powder
⬆ Powder (Lemon), 20 g protein, 3.9 g carbohydrate, 0.8 g fat per 34 g sachet	883.50	30	PKU Express 20
⬆ Powder (Neutral), 20 g protein, 4.8 g carbohydrate, 0.8 g fat per 34 g sachet	883.50	30	PKU Express 20
⬆ Powder (Neutral), 5.0 g protein, 5.2 g carbohydrate, 0.2 g fat per 12.5 g sachet	220.88	30	PKU Explore 5
⬆ Powder (Orange), 10 g protein, 9.8 g carbohydrate, 0.4 g fat per 25 g sachet	441.75	30	PKU Explore 10
⬆ Powder (Orange), 20 g protein, 3.9 g carbohydrate, 0.8 g fat per 34 g sachet	883.50	30	PKU Express 20
⬆ Powder (Orange), 5.0 g protein, 14 g carbohydrate, 0 g fat per 20 g sachet	449.28	60	PKU Restore Powder
⬆ Powder (Raspberry), 10 g protein, 9.8 g carbohydrate, 0.4 g fat per 25 g sachet	441.75	30	PKU Explore 10
⬆ Powder (Tropical), 20 g protein, 3.9 g carbohydrate, 0.8 g fat per 34 g sachet	883.50	30	PKU Express 20
⬆ Powder (berry) 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet	936.00	30	PKU Lophlex Powder
⬆ Powder (chocolate) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet	393.00	30	PKU Anamix Junior
⬆ Powder (neutral) 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet	936.00	30	PKU Lophlex Powder
⬆ Powder (neutral) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet	393.00	30	PKU Anamix Junior
⬆ Powder (orange) 20 g protein, 3.8 g carbohydrate and 0.23 g fibre per 28 g sachet	936.00	30	PKU Lophlex Powder
⬆ Powder (orange) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet	393.00	30	PKU Anamix Junior
⬆ Powder (unflavoured), 5 g protein, 4.8 g carbohydrate per 12.5 g sachets	234.00	30	PKU First Spoon
⬆ Powder (vanilla) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet	393.00	30	PKU Anamix Junior
⬆ Powder (Neutral), 14.3 g protein, 25 g fat per 100 g, 4 × 400 g can	715.16	1,600 g	PKU Start
⬆ Powder (orange) 39 g protein and 34 g carbohydrate per 100 g, 500 g can	320.00	500 g	XP Maxamum
⬆ Powder (unflavoured) 39 g protein and 34 g carbohydrate per 100 g, 500 g can	320.00	500 g	XP Maxamum
⬆ Powder 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	174.72	400 g	PKU Anamix Infant
⬆ Liquid 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle	13.10	1	PKU Anamix Junior LQ (Berry) PKU Anamix Junior LQ (Orange)
⬆ Liquid (juicy berries) 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 62.5 ml bottle	939.00	60	PKU Lophlex LQ 10
⬆ Liquid (juicy berries) 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle	936.00	30	PKU Lophlex LQ 20

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
⬆ Liquid (juicy orange) 20 g protein, 8.8 g carbohydrate and 0.34 g fibre per 100 ml, 125 ml bottle	936.00	30	PKU Lophlex LQ 20
⬆ Liquid (juicy tropical) 16 g protein, 7 g carbohydrate and 0.4 g fibre per 100 ml, 125 ml bottle	936.00	30	PKU Lophlex LQ 20
⬆ Liquid 6.7 g protein, 5.1 g carbohydrate and 2 g fat per 100 ml, 250 ml carton	540.00	18	Easiphen Liquid
⬆ Semi-solid 18.3 g protein, 18.5 g carbohydrate and 0.92 g fibre per 100 g, 109 g pot	1,123.20	36	PKU Lophlex Sensations 20 (berries)
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME PHENYLALANINE – **Restricted** see terms on page 295			
⬆ Powder (Neutral), 10 g protein, 0.5 g carbohydrate, 0.6 g fat per 15 g sachet	449.28	30	PKU Build 10
⬆ Powder (neutral), 15 g protein, 15 g carbohydrate, 4.5 g fat per 40 g sachet	673.92	30	Glytactin Bettermilk
⬆ Powder (unflavoured) 10 g protein, 4 g carbohydrate per 12.5 g sachet	468.00	30	PKU GMPro Mix-In
⬆ Powder 20 g protein, 1.7 g carbohydrate per 31 g sachet	898.56	30	PKU Build 20 Raspberry Lemonade PKU Build 20 Smooth
⬆ Powder 20 g protein, 1.7 g carbohydrate per 32 g sachet	898.56	30	PKU Build 20 Chocolate
⬆ Powder 20 g protein, 1.7 g carbohydrate per 33 g sachet	898.56	30	PKU Build 20 Vanilla
⬆ Powder 20 g protein, 4.9 g carbohydrate per 33.4 g sachet	936.00	30	PKU GMPro Ultra Lemonade PKU GMPro Ultra Vanilla
⬆ Powder 20 g protein, 6.0 g carbohydrate per 35 g sachet	930.00	30	PKU sphere20 Lemon
⬆ Powder 20 g protein, 6.3 g carbohydrate per 35 g sachet	930.00	30	PKU sphere20 Chocolate PKU sphere20 Red Berry PKU sphere20 Vanilla
⬆ Powder 20 g protein, 6.7 g carbohydrate per 35 g sachet	930.00	30	PKU sphere20 Banana
⬆ Liquid (Coffee Mocha), 15 g protein, 3.1 g carbohydrate, 4.6 g fat 250 ml, carton	684.45	30	PKU Glytactin RTD 15 Lite
⬆ Liquid (chocolate), 15 g protein, 22 g carbohydrate, 5.3 g fat per 250 ml, carton	684.45	30	PKU Glytactin RTD 15
⬆ Liquid (neutral),10 g protein, 8.5 g carbohydrate per 250 ml carton	280.80	18	PKU GMPro LQ
⬆ Liquid (original), 15 g protein, 22 g carbohydrate, 5.3 g fat per 250 ml, carton	684.45	30	PKU Glytactin RTD 15
⬆ Liquid (vanilla), 15 g protein, 3.3 g carbohydrate, 4.6 g fat per 250 ml, carton	684.45	30	PKU Glytactin RTD 15 Lite

Protein Free Supplements

PROTEIN FREE SUPPLEMENT CONTAINING CARBOHYDRATE, FAT WITH ADDED VITAMINS AND MINERALS – **Restricted** see terms on page 295			
⬆ Powder (neutral) nil added protein and 67 g carbohydrate per 100 g, 400 g can	49.29	400 g	Energivit

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Supplements for Tyrosinaemia			
AMINO ACID FORMULA (WITHOUT PHENYLALANINE AND TYROSINE) – **Restricted** see terms on page 295			
⬆ Powder (neutral) 36 g protein, 32 g carbohydrate and 12.5 g fat per 100 g, 36 g sachet	471.00	30	TYR Anamix Junior
⬆ Powder (neutral), 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet	349.65	30	TYR Explore 5
⬆ Powder 13.1 g protein, 49.5 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	260.00	400 g	TYR Anamix Infant
⬆ Liquid (orange) 8 g protein, 7 g carbohydrate, 3.8 g fat and 0.25 g fibre per 100 ml, 125 ml bottle	941.40	36	TYR Anamix Junior LQ
⬆ Liquid (juicy berries), 20 g protein, 12.75 g carbohydrate and 0.46 g fat and 0 g fibre per 125 ml pouch	1,684.80	30	TYR Lophlex LQ 20
GLYCOMACROPEPTIDE AND AMINO ACID CONTAINS SOME TYROSINE AND PHENYLALANINE – **Restricted** see terms on page 295			
⬆ Powder (Red Berry), 20 g protein, 6.3 carbohydrate, 1.6 g fat per 35 g sachet	1,398.60	30	TYR Sphere 20
⬆ Powder (Vanilla), 20 g protein, 6.0 g carbohydrate, 1.6 g fat per 35 g sachet	1,398.60	30	TYR Sphere 20
X-Linked Adrenoleukodystrophy Products			
GLYCEROL TRIERUCATE – **Restricted** see terms on page 295			
⬆ Liquid, 1,000 ml bottle			
GLYCEROL TRIOLEATE – **Restricted** see terms on page 295			
⬆ Liquid, bottle	131.80	500 ml	GTO Oil
Supplements for Glycogen Storage Disease			
HIGH AMYLOPECTIN CORN-STARCH – **Restricted** see terms on page 295			
⬆ Powder 0 g protein, 53 g carbohydrate, 0 g fat per 60 g sachet	241.62	30	Glycosade
Supplements for Organic Acidaemias			
AMINO ACID FORMULA (WITHOUT ISOLEUCINE, METHIONINE, THREONINE AND VALINE) – **Restricted** see terms on page 295			
⬆ Powder 13.1 g protein, 50.1 g carbohydrate, 23 g fat and 5.3 g fibre per 100 g, 400 g can	260.00	400 g	MMA/PA Anamix Infant
AMINO ACID FORMULA (WITHOUT METHIONINE, THREONINE AND VALINE) – **Restricted** see terms on page 295			
⬆ Powder (neutral), 5 g protein, 5.4 g carbohydrate, 2.3 g fat and 2.0 g fibre per 18 g sachet	750.30	30	MMA/PA Anamix Junior
⬆ Powder, 15 g protein, 3.4 g carbohydrate, 0.05 g fat per 25 g sachet	1,048.95	30	MMA/PA Express 15
⬆ Powder, 5 g protein, 5.3 g carbohydrate, 0.2 g fat per 12.5 g sachet	349.65	30	MMA/PA Explore 5
Single Dose Amino Acids			
ARGININE – **Restricted** see terms on page 295			
⬆ Powder 1.7 g protein, 1.9 g carbohydrate per 4 g sachet	211.45	30	Arginine2000
CITRULLINE – **Restricted** see terms on page 295			
⬆ Powder 0.8 g protein, 2.9 g carbohydrate per 4 g sachet	211.45	30	Citrulline1000
ISOLEUCINE – **Restricted** see terms on page 295			
⬆ Powder 0.04 g protein, 3.8 g carbohydrate per 4 g sachet	141.05	30	Isoleucine50

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
LEUCINE – **Restricted** see terms on page 295			
⬆ Powder 0.08 g protein, 3.7 g carbohydrate per 4 g sachet	141.05	30	Leucine100
PHENYLALANINE – **Restricted** see terms on page 295			
⬆ Powder 0.04 g protein, 3.8 g carbohydrate per 4 g sachet	141.05	30	Phenylalanine50
TYROSINE – **Restricted** see terms on page 295			
⬆ Powder 0.8 g protein, 2.9 g carbohydrate per 4 g sachet	211.45	30	Tyrosine1000
VALINE – **Restricted** see terms on page 295			
⬆ Powder 0.04 g protein, 3.8 g carbohydrate per 4 g sachet	141.05	30	Valine50

Other Fat Modified Products

ELEMENTAL FEED WITH HIGH MEDIUM CHAIN TRIGLYCERIDES – **Restricted** see terms on page 295			
⬆ Powder (neutral), 12.5 g protein, 60 g carbohydrate and 16.4 g fat per 100 g sachet	47.01	10	Emsogen

Essential Amino Acids

ESSENTIAL AMINO ACID FORMULA – **Restricted** see terms on page 295			
⬆ Powder (neutral) 79 g protein per 100 g, 200 g can	313.73	200 g	Essential Amino Acid Mix

Specialised Formulas

Diabetic Products

➥ **Restricted** (RS1215)

Initiation

Any of the following:

1 For patients with type I or type II diabetes suffering weight loss and malnutrition that requires nutritional support; or
2 For patients with pancreatic insufficiency; or
3 For patients who have, or are expected to, eat little or nothing for 5 days; or
4 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or
5 For use pre- and post-surgery; or
6 For patients being tube-fed; or
7 For tube-feeding as a transition from intravenous nutrition.

DIABETIC ORAL FEED 1 KCAL/ML – **Restricted** see terms above			
⬆ Liquid 4.9 g protein, 11.7 g carbohydrate, 3.8 g fat and 2 g fibre per 100 ml, 200 ml bottle	2.25	1	Diasip (strawberry) Diasip (vanilla)
LOW-GI ENTERAL FEED 1 KCAL/ML – **Restricted** see terms above			
⬆ Liquid 5 g protein, 9.6 g carbohydrate and 5.4 g fat per 100 ml, 500 ml bottle	4.65	1	Glucerna Select
⬆ Liquid 4.3 g protein, 11.3 g carbohydrate and 4.2 g fat per 100 ml, 1,000 ml bottle			*e.g. Nutrison Advanced Diason*
LOW-GI ORAL FEED 1 KCAL/ML – **Restricted** see terms above			
⬆ Liquid 7 g protein, 10.9 g carbohydrate, 2.7 g fat and 2 g fibre per 100 ml, 200 ml bottle	8.40	4	Nutren Diabetes (vanilla)

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
Elemental and Semi-Elemental Products			
➥ **Restricted** (RS1216)			
Initiation			
Any of the following:			
1 Malabsorption; or			
2 Short bowel syndrome; or			
3 Enterocutaneous fistulas; or			
4 Eosinophilic enteritis (including oesophagitis); or			
5 Inflammatory bowel disease; or			
6 Acute pancreatitis where standard feeds are not tolerated; or			
7 Patients with multiple food allergies requiring enteral feeding.			
AMINO ACID ORAL FEED – **Restricted** see terms above			
⬆ Powder 11 g protein, 62 g carbohydrate and 1 g fat per sachet, 80 g sachet	45.00	10	Vivonex TEN
AMINO ACID ORAL FEED 0.8 KCAL/ML – **Restricted** see terms above			
⬆ Liquid 2.5 g protein, 11 g carbohydrate and 3.5 g fat per 100 ml, 250 ml carton	179.46	18	Elemental 028 Extra (grapefruit) Elemental 028 Extra (pineapple & orange) Elemental 028 Extra (summer fruits)
PEPTIDE-BASED ENTERAL FEED 1 KCAL/ML – **Restricted** see terms above			
⬆ Liquid 4 g protein, 17.7 g carbohydrate and 1.7 g fat per 100 ml, 500 ml bottle	7.47	1	Nutrison Advanced Peptisorb
PEPTIDE-BASED ENTERAL FEED 1.5 KCAL/ML – **Restricted** see terms above			
⬆ Liquid 6.75 g protein, 18.4 g carbohydrate and 5.5 g fat per 100 ml, 1,000 ml bottle	22.39	1	Vital
PEPTIDE-BASED ORAL FEED – **Restricted** see terms above			
⬆ Powder 13.7 g protein, 62.9 g carbohydrate and 17.5 g fat per 100 g, 400 g can			*e.g. Peptamen Junior*
⬆ Powder 13.8 g protein, 59 g carbohydrate and 18 g fat per 100 g, 400 g can			*e.g. MCT Pepdite; MCT Pepdite 1+*
PEPTIDE-BASED ORAL FEED 1 KCAL/ML – **Restricted** see terms above			
⬆ Liquid 5 g protein, 16 g carbohydrate and 1.69 g fat per 100 ml, carton	4.95	237 ml	Peptamen OS 1.0 (Vanilla)
Fat Modified Products			
FAT-MODIFIED FEED – **Restricted** see terms below			
⬇ Powder 12.8 g protein, 68.6 g carbohydrate and 12.9 g fat per 100 g, can	62.90	400 g	Monogen
➥ **Restricted** (RS1470)			
Initiation			
Any of the following:			

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Patient has metabolic disorders of fat metabolism; or
2 Patient has a chyle leak; or
3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

Hepatic Products

➥ **Restricted** (RS1217)
Initiation
For children (up to 18 years) who require a liver transplant.

HEPATIC ORAL FEED – **Restricted** see terms above

⬆	Powder 12 g protein, 56 g carbohydrate and 22 g fat per 100 g, can	93.97	400 g	Heparon Junior

High Calorie Products

➥ **Restricted** (RS1317)
Initiation
Any of the following:
1 Patient is fluid volume or rate restricted; or
2 Patient requires low electrolyte; or
3 Both:
3.1 Any of the following:
3.1.1 Cystic fibrosis; or
3.1.2 Any condition causing malabsorption; or
3.1.3 Faltering growth in an infant/child; or
3.1.4 Increased nutritional requirements; and
3.2 Patient has substantially increased metabolic requirements.

ENTERAL FEED 2 KCAL/ML – **Restricted** see terms above

⬆	Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 500 ml bottle	6.82	1	Nutrison Concentrated
⬆	Liquid 8.4 g protein, 21.9 g carbohydrate, 9.1 g fat and 0.5 g fibre per 100 ml, 1,000 ml bottle	13.64	1	Ensure Two Cal HN RTH

ORAL FEED 2 KCAL/ML – **Restricted** see terms above

⬆	Liquid 8.4 g protein, 22.4 g carbohydrate, 8.9 g fat and 0.8 g fibre per 100 ml, 200 ml bottle	2.34	1	Two Cal HN

High Protein Products

HIGH PROTEIN ENTERAL FEED 1.25 KCAL/ML – **Restricted** see terms below

⬇	Liquid 6.3 g protein, 14.2 g carbohydrate and 4.9 g fat per 100 ml, bottle	12.00	1,000 ml	Nutrison Protein Plus

➥ **Restricted** (RS1327)
Initiation
Both:

1 The patient has a high protein requirement; and
2 Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.1 Patient has liver disease; or
2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
2.3 Patient is fluid restricted; or
2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.26 KCAL/ML – **Restricted** see terms below

⬇ Liquid 10 g protein, 10.4 g carbohydrate and 4.9 g fat per 100 ml, bottle ..... 8.67 — 500 ml — Nutrison Protein Intense

➥ **Restricted** (RS1327)

Initiation

Both:

1 The patient has a high protein requirement; and
2 Any of the following:
2.1 Patient has liver disease; or
2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
2.3 Patient is fluid restricted; or
2.4 Patient's needs cannot be more appropriately met using high calorie product.

HIGH PROTEIN ENTERAL FEED 1.28 KCAL/ML – **Restricted** see terms below

⬇ Liquid 6.3 g protein, 14.1 g carbohydrate, 4.9 g fat and 1.5 g fibre per 100 ml, bottle ..... 12.54 — 1,000 ml — Nutrison Protein Plus Multi Fibre

➥ **Restricted** (RS1327)

Initiation

Both:

1 The patient has a high protein requirement; and
2 Any of the following:
2.1 Patient has liver disease; or
2.2 Patient is obese (BMI > 30) and is undergoing surgery; or
2.3 Patient is fluid restricted; or
2.4 Patient's needs cannot be more appropriately met using high calorie product.

Infant Formulas

AMINO ACID FORMULA – **Restricted** see terms below

⬇ Powder 1.95 g protein, 8.1 g carbohydrate and 3.5 g fat per 100 ml, 400 g can			*e.g. Neocate*
⬇ Powder 13 g protein, 49 g carbohydrate and 23 g fat per 100 g, can	55.61	400 g	Neocate SYNEO
⬇ Powder 13.3 g protein, 56 g carbohydrate and 22 g fat per 100 g, can	55.61	400 g	Neocate Junior Unflavoured
⬇ Powder 13.3 g protein, 57 g carbohydrate and 24.6 g fat per 100 g, can	43.60	400 g	Alfamino
⬇ Powder 13.5 g protein, 52 g carbohydrate and 24.5 g fat per 100 g, can	55.61	400 g	Neocate Gold (Unflavoured)
⬇ Powder 14.8 g protein, 51.4 g carbohydrate and 23 g fat per 100 g, can	55.61	400 g	Neocate Junior Vanilla
⬇ Powder 15 g protein, 56 g carbohydrate and 20 g fat per 100 g, can	43.60	400 g	Alfamino Junior
⬇ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can.....	65.72	400 g	Elecare LCP (Unflavoured)
⬇ Powder 2.2 g protein, 7.8 g carbohydrate and 3.4 g fat per 100 ml, can.....	65.72	400 g	Elecare (Unflavoured) Elecare (Vanilla)

➥ **Restricted** (RS1867)

Initiation

Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; or
2 History of anaphylaxis to cows' milk protein formula or dairy products; or
3 Eosinophilic oesophagitis; or
4 Ultra-short gut; or
5 Severe Immune deficiency.

Continuation

All of the following:

1 An assessment as to whether the infant can be transitioned to a cows' milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
3 Amino acid formula is required for a nutritional deficit.

Initiation – patients who are currently funded under RS1502 or SA1557

Limited to 3 months treatment

All of the following:

1 Patient has a valid initiation or renewal approval for extensively hydrolysed formula (RS1502); and
2 Patient is unable to source funded Aptamil powder at this time; and
3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Hospital Restriction RS1502. There is no continuation criteria under this criterion.

ENTERAL LIQUID PEPTIDE FORMULA – **Restricted** see terms below

⬇ Liquid 4.2 g protein, 18.6 g carbohydrate and 6.58 g fat per 100 ml, 500 ml bottle	18.66	1	Nutrini Peptisorb Energy

➥ **Restricted** (RS1775)

Initiation

All of the following:

1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
2 Any of the following:
  2.1 Severe malabsorption; or
  2.2 Short bowel syndrome; or
  2.3 Intractable diarrhoea; or
  2.4 Biliary atresia; or
  2.5 Cholestatic liver diseases causing malabsorption; or
  2.6 Cystic fibrosis; or
  2.7 Proven fat malabsorption; or
  2.8 Severe intestinal motility disorders causing significant malabsorption; or
  2.9 Intestinal failure; or
  2.10 Both:
    2.10.1 The patient is currently receiving funded amino acid formula; and
    2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
3 Either:
  3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
  3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Continuation

Both:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula.

EXTENSIVELY HYDROLYSED FORMULA – **Restricted** see terms below

⬇ Powder 1.6 g protein, 7.5 g carbohydrate and 3.1 g fat per 100 ml, 900 g can	36.20	900 g	Allerpro Syneo 1
⬇ Powder 1.6 g protein, 7.8 g carbohydrate and 3.2 g fat per 100 ml, 900 g can	36.20	900 g	Allerpro Syneo 2
⬇ Powder 14 g protein, 53.4 g carbohydrate and 27.3 g fat per 100 g, can	18.10	450 g	Pepti-Junior

➥ **Restricted** (RS1502)

Initiation

Any of the following:

1 Both:
  1.1 Cows' milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  1.2 Either:
    1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
2 Severe malabsorption; or
3 Short bowel syndrome; or
4 Intractable diarrhoea; or
5 Biliary atresia; or
6 Cholestatic liver diseases causing malsorption; or
7 Cystic fibrosis; or
8 Proven fat malabsorption; or
9 Severe intestinal motility disorders causing significant malabsorption; or
10 Intestinal failure; or
11 For step down from Amino Acid Formula.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Continuation

Both:

1 An assessment as to whether the infant can be transitioned to a cows' milk protein or soy infant formula has been undertaken; and
2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula.

FRUCTOSE-BASED FORMULA

Powder 14.6 g protein, 49.7 g carbohydrate and 30.8 g fat per 100 g, 400 g can			*e.g. Galactomin 19*

LACTOSE-FREE FORMULA

Powder 1.3 g protein, 7.3 g carbohydrate and 3.5 g fat per 100 ml, 900 g can			*e.g. Karicare Aptamil Gold De-Lact*
Powder 1.5 g protein, 7.2 g carbohydrate and 3.6 g fat per 100 ml, 900 g can			*e.g. S26 Lactose Free*

LOW-CALCIUM FORMULA

Powder 14.8 g protein, 53.7 g carbohydrate and 26.7 g fat per 100 g and tuna fish oil (DHA), can	46.18	400 g	Locasol

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML – **Restricted** see terms on the next page

⬇ Liquid 2.6 g protein, 10.3 g carbohydrate, 5.4 g fat and 0.6 g fibre per 100 ml, 125 ml bottle	2.80	1	Infatrini

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

➥ **Restricted** (RS1614)

Initiation – Fluid restricted or volume intolerance with faltering growth

Both:

1 Either:

1.1 The patient is fluid restricted or volume intolerant; or

1.2 The patient has increased nutritional requirements due to faltering growth; and

2 Patient is under 18 months old and weighs less than 8kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

PRETERM FORMULA – **Restricted** see terms below

⬇	Liquid 2.2 g protein, 8.4 g carbohydrate and 4.4 g fat per 100 ml, bottle	0.75	100 ml	S26 LBW Gold RTF
⬇	Liquid 2.3 g protein, 8.6 g carbohydrate and 4.2 g fat per 100 ml, 90 ml bottle			*e.g. Pre Nan Gold RTF*
⬇	Liquid 2.6 g protein, 8.4 g carbohydrate and 3.9 g fat per 100 ml, 70 ml bottle			*e.g. Karicare Aptamil Gold+Preterm*

➥ **Restricted** (RS1224)

Initiation

For infants born before 33 weeks' gestation or weighing less than 1.5 kg at birth.

THICKENED FORMULA

	Powder 1.8 g protein, 8.1 g carbohydrate and 3.3 g fat per 100 ml, 900 g can			*e.g. Karicare Aptamil Thickened AR*

Ketogenic Diet Products

HIGH FAT FORMULA – **Restricted** see terms below

⬇	Powder 14.3 g protein, 2.8 g carbohydrate and 69.2 g fat per 100 g, can	36.92	300 g	Ketocal 4:1 (Unflavoured) Ketocal 4:1 (Vanilla)
⬇	Powder 15.4 g protein, 7.2 g carbohydrate and 68.6 g fat per 100 g, can	36.92	300 g	Ketocal 3:1 (Unflavoured)

➥ **Restricted** (RS1225)

Initiation

For patients with intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Paediatric Products

➦ **Restricted** (RS1473)

Initiation

Both:

1 Child is aged one to ten years; and

2 Any of the following:

2.1 The child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or

2.2 Any condition causing malabsorption; or

2.3 Faltering growth in an infant/child; or

2.4 Increased nutritional requirements; or

2.5 The child is being transitioned from TPN or tube feeding to oral feeding; or

2.6 The child has eaten, or is expected to eat, little or nothing for 3 days.

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)

e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
PAEDIATRIC ENTERAL FEED 0.76 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 2.5 g protein, 12.5 g carbohydrate, 3.3 g fat and 0.7 g fibre per 100 ml, 500 ml bottle	6.27	1	Nutrini Low Energy Multi Fibre RTH
PAEDIATRIC ENTERAL FEED 1 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, 500 ml bottle	3.32	1	Pediasure RTH
⬆ Liquid 2.7 g protein, 12.3 g carbohydrate and 4.4 g fat per 100 ml, 500 ml bottle	4.69	1	Nutrini RTH
PAEDIATRIC ENTERAL FEED 1.5 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 4.1 g protein, 18.5 g carbohydrate and 6.7 g fat per 100 ml, 500 ml bottle	7.46	1	Nutrini Energy RTH
⬆ Liquid 4.1 g protein, 18.5 g carbohydrate, 6.7 g fat and 0.8 g fibre per 100 ml, 500 ml bottle	7.14	1	Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 2.8 g protein, 11.2 g carbohydrate and 5 g fat per 100 ml, 200 ml bottle	1.33	1	Pediasure (chocolate) Pediasure (strawberry) Pediasure (vanilla)
PAEDIATRIC ORAL FEED 1.5 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 3.4 g protein, 18.8 g carbohydrate and 6.8 g fat per 100 ml, 200 ml bottle	1.90	1	Fortini (Strawberry) Fortini (Vanilla)
⬆ Liquid 4.0 g protein, 18.8 g carbohydrate, 6.8 g fat and 1.5 g fibre per 100 ml, 200 ml bottle	1.90	1	Fortini Multi Fibre (chocolate) Fortini Multi Fibre (strawberry) Fortini Multi Fibre (unflavoured) Fortini Multi Fibre (vanilla)
⬆ Liquid 4.2 g protein, 16.7 g carbohydrate and 7.5 g fat per 100 ml, 500 ml bottle	8.67	1	Pediasure Plus

Renal Products

LOW ELECTROLYTE ORAL FEED – **Restricted** see terms below			
⬇ Powder 7.5 g protein, 57.6 g carbohydrate and 25.9 g fat per 100 g, can	64.26	400 g	Kindergen

➥ **Restricted** (RS1227)

Initiation

For children (up to 18 years) with acute or chronic kidney disease.

LOW ELECTROLYTE ORAL FEED 1.8 KCAL/ML			
Liquid 8 g protein, 14.74 g carbohydrate, 9.77 g fat and 1.26 g fibre per 100 ml, 220 ml bottle	3.31	1	Nepro HP (strawberry) Nepro HP (vanilla)

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
LOW ELECTROLYTE ORAL FEED 2 KCAL/ML – **Restricted** see terms below			
⬇ Liquid 3 g protein, 25.5 g carbohydrate and 9.6 g fat per 100 ml, 237 ml bottle			
⬇ Liquid 7.5 g protein, 20 g carbohydrate and 10 g fat per 100 ml, 125 ml carton	13.72	4	Renilon 7.5 (apricot) Renilon 7.5 (caramel)
⬇ Liquid 9.1 g protein, 19 g carbohydrate and 10 g fat per 100 ml, 200 ml bottle	13.24	4	Novasource Renal (Vanilla)

➥ **Restricted** (RS1228)
Initiation
For patients with acute or chronic kidney disease.

Surgical Products

HIGH ARGININE ORAL FEED 1.4 KCAL/ML – **Restricted** see terms below			
⬇ Liquid 10.4 g protein, 8 g carbohydrate, 4.4 g fat and 0 g fibre per 100 ml, 250 ml carton	56.00	10	Impact Advanced Recovery

➥ **Restricted** (RS1231)
Initiation
Three packs per day for 5 to 7 days prior to major gastrointestinal, head or neck surgery.

PREOPERATIVE CARBOHYDRATE FEED 0.5 KCAL/ML – **Restricted** see terms below			
⬇ Oral liq 0 g protein, 12.6 g carbohydrate and 0 g fat per 100 ml, 200 ml bottle	8.64	4	preOp

➥ **Restricted** (RS1415)
Initiation
Maximum of 400 ml as part of an Enhanced Recovery After Surgery (ERAS) protocol 2 to 3 hours before major abdominal surgery.

Standard Feeds

➥ **Restricted** (RS1214)
Initiation
Any of the following:

For patients with malnutrition, defined as any of the following:

1 Any of the following:
  1.1 BMI < 18.5; or
  1.2 Greater than 10% weight loss in the last 3-6 months; or
  1.3 BMI < 20 with greater than 5% weight loss in the last 3-6 months; or
2 For patients who have, or are expected to, eat little or nothing for 5 days; or
3 For patients who have a poor absorptive capacity and/or high nutrient losses and/or increased nutritional needs from causes such as catabolism; or
4 For use pre- and post-surgery; or
5 For patients being tube-fed; or
6 For tube-feeding as a transition from intravenous nutrition; or
7 For any other condition that meets the community Special Authority criteria.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
ENTERAL FEED 1.5 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 1.5 g fibre per 100 ml, 1,000 ml bottle	8.68	1	Nutrison Energy Multi Fibre
⬆ Liquid 6.27 g protein, 20.4 g carbohydrate and 4.9 g fat per 100 ml, 1,000 ml bag	8.68	1	Ensure Plus HN RTH
⬆ Liquid 6.38 g protein, 21.1 g carbohydrate, 4.9 g fat and 1.2 g fibre per 100 ml, 1,000 ml bottle	8.68	1	Jevity HiCal RTH
⬆ Liquid 6 g protein, 18.5 g carbohydrate and 5.8 g fat per 100 ml, 1,000 ml bottle	9.00	1	Nutrison Energy
ENTERAL FEED 1 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 4 g protein, 12.3 g carbohydrate, 3.9 g fat and 1.5 g fibre per 100 ml, 1,000 ml bottle	7.21	1	Nutrison Multi Fibre
⬆ Liquid 4 g protein, 13.6 g carbohydrate and 3.4 g fat per 100 ml, 1,000 ml bottle	6.56	1	Osmolite RTH
⬆ Liquid 4 g protein, 14.1 g carbohydrate, 3.47 g fat and 1.76 g fibre per 100 ml, 1,000 ml bottle	6.56	1	Jevity RTH
⬆ Liquid 4 g protein, 12.4 g carbohydrate and 3.9 g fat per 100 ml, 1,000 ml bottle	6.90	1	Nutrison RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 5.5 g protein, 8.8 g carbohydrate, 2.5 g fat and 1.5 g fibre per 100 ml, 1,000 ml bottle	9.05	1	Nutrison 800 Complete Multi Fibre
HIGH PROTEIN ORAL FEED 2.4 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 14.6 g protein, 25.3 g carbohydrate and 9.6 g fat per 100 ml, 125 ml bottle			*e.g. Fortisip Compact Protein*
ORAL FEED – **Restricted** see terms on the previous page			
⬆ Powder 15.9 g protein, 57.4 g carbohydrate and 14 g fat per 100 g, can	40.00	800 g	Ensure (Chocolate)
		850 g	Ensure (Chocolate)
		800 g	Ensure (Vanilla)
		850 g	Ensure (Vanilla)
⬆ Powder 23 g protein, 65 g carbohydrate and 2.5 g fat per 100 g, can	15.90	840 g	Sustagen Hospital Formula (Chocolate)
			Sustagen Hospital Formula (Vanilla)
ORAL FEED 1 KCAL/ML – **Restricted** see terms on the previous page			
⬆ Liquid 3.8 g protein, 23 g carbohydrate and 12.7 g fibre per 100 ml, 237 ml carton			*e.g. Resource Fruit Beverage*

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
ORAL FEED 1.5 KCAL/ML – **Restricted** see terms on page 308			
⬆ Liquid 4 g protein and 33.5 g carbohydrate per 100 ml, 200 ml bottle	3.30	200 ml	Fortijuice (Apple) Fortijuice (Orange) Fortijuice (Strawberry)
⬆ Liquid 6 g protein, 18.4 g carbohydrate and 5.8 g fat per 100 ml, 200 ml bottle	1.76	1	Fortisip (Banana) Fortisip (Chocolate) Fortisip (Strawberry) Fortisip (Vanilla)
⬆ Liquid 6.25 g protein, 20.2 g carbohydrate and 4.92 g fat per 100 ml, 200 ml bottle	1.56	1	Ensure Plus (Banana) Ensure Plus (Chocolate) Ensure Plus (Fruit of the forest) Ensure Plus (Vanilla)
⬆ Liquid 5.5 g protein, 21.1 g carbohydrate and 4.81 g fat per 100 ml, 237 ml can	1.65	1	Ensure Plus (Vanilla)
(Ensure Plus (Vanilla) Liquid 5.5 g protein, 21.1 g carbohydrate and 4.81 g fat per 100 ml, 237 ml can to be delisted 1 July 2026)			
ORAL FEED WITH FIBRE 1.5 KCAL/ML – **Restricted** see terms on page 308			
⬆ Liquid 6 g protein, 18.4 g carbohydrate, 5.8 g fat and 2.3 g fibre per 100 ml, 200 ml bottle	1.76	1	Fortisip Multi Fibre (chocolate) Fortisip Multi Fibre (strawberry) Fortisip Multi Fibre (vanilla)

⬆ Item restricted (see ➦ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

Bacterial and Viral Vaccines

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – **Restricted** see terms below

⬇ Inj 30 IU diphtheria toxoid with 30IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units poliomyelitis virus in 0.5 ml syringe – **5% DV Dec-24 to 2027** ..... 0.00 — 10 — **Infanrix IPV**

➥ **Restricted** (RS1387)

Initiation

Any of the following:

1 A single dose for children up to the age of 7 who have completed primary immunisation; or
2 A course of up to four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
3 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
4 Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE – **Restricted** see terms below

⬇ Inj 30IU diphtheria with 40IU tetanus and 25mcg pertussis toxoids, 25mcg pertussis filamentous haemagglutinin, 8mcg pertactin, 80D-AgU polio virus, 10mcg hepatitis B antigen 10mcg H. influenzae type b with tetanus toxoid 20-40mcg in 0.5ml syringe – **5% DV Dec-24 to 2027** ..... 0.00 — 10 — **Infanrix-hexa**

➥ **Restricted** (RS2051)

Initiation

Any of the following:

1 Up to four doses for children under the age of 10 years for primary immunisation; or
2 An additional four doses (as appropriate) for (re-)immunisation of children under the age of 18 years post haematopoietic stem cell transplantation; or
3 An additional four doses (as appropriate) for (re-)immunisation of children under the age of 10 years who are post chemotherapy; pre or post splenectomy; undergoing renal dialysis and other severely immunosuppressive regimens; or
4 Up to five doses for children under the age of 10 years receiving solid organ transplantation.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Bacterial Vaccines

BACILLUS CALMETTE-GUERIN VACCINE – **Restricted** see terms below

⬇ Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial with diluent – **5% DV Dec-24 to 2027** ..... 0.00 — 10 — **BCG Vaccine AJV**

➥ **Restricted** (RS1233)

Initiation

All of the following:

For infants at increased risk of tuberculosis defined as:

1 Living in a house or family with a person with current or past history of TB; and
2 Having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

equal to 40 per 100,000 for 6 months or longer; and

3 During their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000.

Note: A list of countries with high rates of TB are available at http://www.health.govt.nz/tuberculosis (Search for Downloads) or www.bcgatlas.org/index.php

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – **Restricted** see terms below

⬇ Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous haemagglutinin and 2.5 mcg pertactin in 0.5 ml prefilled syringe – **5% DV Dec-24 to 2027** ..... 0.00 — 10 — **Boostrix**

➥ **Restricted** (RS1790)

Initiation

Any of the following:

1 A single dose for pregnant women in the second or third trimester of each pregnancy; or; or
2 A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or; or
3 A course of up to four doses is funded for children from age 7 up the age of 18 years inclusive to complete full primary immunisation; or
4 An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
5 A single dose for vaccination of patients aged from 65 years old; or
6 A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
7 For vaccination of previously unimmunised or partially immunised patients; or
8 For revaccination following immunosuppression; or
9 For boosting of patients with tetanus-prone wounds.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

HAEMOPHILUS INFLUENZAE TYPE B VACCINE – **Restricted** see terms below

⬇ Inj 10 mcg vial with diluent syringe – **5% DV Dec-24 to 2027** ..... 0.00 — 1 — **Act-HIB**

➥ **Restricted** (RS1520)

Initiation

Therapy limited to 1 dose

Any of the following:

1 For primary vaccination in children; or
2 An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre- or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or
3 For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

MENINGOCOCCAL (A, C, Y AND W-135) CONJUGATE VACCINE

⬇ Inj 10 mcg of each meningococcal polysaccharide conjugated to a total of approximately 55 mcg of tetanus toxoid carrier per 0.5 ml vial – **5% DV Dec-24 to 2027** ..... 0.00 — 1 — **MenQuadfi**

➥ **Restricted** (RS2019)

Initiation

Either:

1 Any of the following:

1.1 Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with HIV,

continued…

⬆ Item restricted (see ➥ above); ⬇ Item restricted (see ➥ below)
e.g. Brand indicates brand example only. It is not a contracted product.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

complement deficiency (acquired or inherited), functional or anatomic asplenia or pre or post solid organ transplant; or

1.2 One dose for close contacts of meningococcal cases of any group; or

1.3 One dose for person who has previously had meningococcal disease of any group; or

1.4 A maximum of two doses for bone marrow transplant patients; or

1.5 A maximum of two doses for person pre and post-immunosuppression*; or

2 Both:

2.1 Person is aged between 13 and 25 years, inclusive; and

2.2 Either:

2.2.1 One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or

2.2.2 One dose for individuals who turn 13 years of age while living in boarding school hostels.

Notes: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

⬇ Inj 5 mcg of each meningococcal polysaccharide conjugated to a total of approximately 44 mcg of tetanus toxoid carrier in 0.5 ml vial	0.00	1	Nimenrix

➥ **Restricted** (RS2037)

Initiation – Children under 12 months of age

Any of the following:

1 A maximum of three doses (dependant on age at first dose) for patients pre- and post- splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post- solid organ transplant; or

2 A maximum of three doses (dependant on age at first dose) for close contacts of meningococcal cases of any group; or

3 A maximum of three doses (dependant on age at first dose) for child who has previously had meningococcal disease of any group; or

4 A maximum of three doses (dependant on age at first dose) for bone marrow transplant patients; or

5 A maximum of three doses (dependant on age at first dose) for child pre- and post-immunosuppression*.

Notes: infants from 6 weeks to less than 6 months of age require a 2+1 schedule, infants from 6 months to less than 12 months of age require a 1+1 schedule. Refer to the Immunisation Handbook for recommended booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

MENINGOCOCCAL B MULTICOMPONENT VACCINE – **Restricted** see terms below

⬇ Inj 175 mcg per 0.5 ml prefilled syringe	0.00	1	Bexsero
		10	Bexsero

➥ **Restricted** (RS2141)

Initiation – Primary immunisation for children up to 59 months of age inclusive

Therapy limited to 3 doses

A primary course of up to three doses (dependent on age at first dose) for previously unvaccinated children up to the age of 59 months inclusive.

Initiation – High-risk individuals 5 years of age or over

Both:

1 Person is aged at least 5 years; and

2 Any of the following:

2.1 Up to two doses and a booster every five years for patients pre- and post-splenectomy; or

2.2 Up to two doses and a booster every five years for patients with functional or anatomic asplenia, HIV, complement

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

deficiency (acquired or inherited); or

2.3 Up to two doses and a booster every five years pre- or post-solid organ transplant; or

2.4 Up to two doses for close contacts of meningococcal cases of any group; or

2.5 Up to two doses for person who has previously had meningococcal disease of any group; or

2.6 Up to two doses for bone marrow transplant patients; or

2.7 Up to two doses for person pre- and post-immunosuppression* .

Initiation – Person is aged between 13 and 25 years (inclusive)

Therapy limited to 2 doses

Both:

1 Person is aged between 13 and 25 years (inclusive); and

2 Either:

2.1 Two doses for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, Youth Justice residences, or prisons; or

2.2 Two doses for individuals who turn 13 years of age while living in boarding school hostels.

Note: *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE – **Restricted** see terms below

⬇ Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4, 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5 ml syringe – **5% DV**

Dec-24 to 2027	0.00	1	**Prevenar 13**
		10	**Prevenar 13**

➥ **Restricted** (RS1936)

Initiation – Primary course for previously unvaccinated children aged under 5 years

Therapy limited to 3 doses

A primary course of three doses for previously unvaccinated children up to the age of 59 months inclusive.

Initiation – High risk individuals who have received PCV10

Therapy limited to 2 doses

Two doses are funded for high risk individuals (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10.

Initiation – High risk children aged under 5 years

Therapy limited to 4 doses

Both:

1 Up to an additional four doses (as appropriate) are funded for the (re)immunisation of high-risk children aged under 5 years; and

2 Any of the following:

2.1 on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or

2.2 primary immune deficiencies; or

2.3 HIV infection; or

2.4 renal failure, or nephrotic syndrome; or

2.5 are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or

2.6 cochlear implants or intracranial shunts; or

2.7 cerebrospinal fluid leaks; or

2.8 receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or

2.9 chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2.10 pre term infants, born before 28 weeks gestation; or
2.11 cardiac disease, with cyanosis or failure; or
2.12 diabetes; or
2.13 Down syndrome; or
2.14 who are pre-or post-splenectomy, or with functional asplenia.

Initiation – High risk individuals 5 years and over

Therapy limited to 4 doses

Up to an additional four doses (as appropriate) are funded for the (re-)immunisation of individuals 5 years and over with HIV, pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency.

Initiation – Testing for primary immunodeficiency diseases

For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – **Restricted** see terms below

⬇	Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype) – **5% DV Dec-24 to 2027**	0.00	1	**Pneumovax 23**

➥ **Restricted** (RS1587)

Initiation – High risk patients

Therapy limited to 3 doses

For patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy; or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency.

Initiation – High risk children

Therapy limited to 2 doses

Both:

1 Patient is a child under 18 years for (re-)immunisation; and
2 Any of the following:
2.1 On immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
2.2 With primary immune deficiencies; or
2.3 With HIV infection; or
2.4 With renal failure, or nephrotic syndrome; or
2.5 Who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
2.6 With cochlear implants or intracranial shunts; or
2.7 With cerebrospinal fluid leaks; or
2.8 Receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
2.9 With chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
2.10 Pre term infants, born before 28 weeks gestation; or
2.11 With cardiac disease, with cyanosis or failure; or
2.12 With diabetes; or
2.13 With Down syndrome; or
2.14 Who are pre-or post-splenectomy, or with functional asplenia.

Initiation – Testing for primary immunodeficiency diseases

For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

SALMONELLA TYPHI VACCINE – **Restricted** see terms below

⬇ Inj 25 mcg in 0.5 ml syringe

➥ **Restricted** (RS1243)

Initiation

For use during typhoid fever outbreaks.

Viral Vaccines

COVID-19 VACCINE

⬇ Inj 3 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3 ml, 0.48 ml multi-dose vial; infant vaccine, yellow cap – **5% DV Feb-26 to 30 Sep 2027** ..... 0.00 | 10 | **Comirnaty (LP.8.1)**

➥ **Restricted** (RS2171)

Initiation – initial dose

Up to three doses for previously unvaccinated children aged 6 months – 4 years at high risk of severe illness or highly immunocompromised.

Continuation – additional dose

Either:

1 One additional dose with the most current variant-matched vaccine every 6 months for highly immunocompromised children aged 6 months to 4 years; or

2 One additional dose with the most current variant-matched vaccine every 12 months for children aged 6 months to 4 years old at high risk of severe illness.

⬇ Inj 3 mcg bretovameran per 0.3 ml, 0.48 ml vial; infant vaccine, yellow cap ..... 0.00 | 10 | Comirnaty Omicron (JN.1)

➥ **Restricted** (RS2171)

Initiation – initial dose

Up to three doses for previously unvaccinated children aged 6 months – 4 years at high risk of severe illness or highly immunocompromised.

Continuation – additional dose

Either:

1 One additional dose with the most current variant-matched vaccine every 6 months for highly immunocompromised children aged 6 months to 4 years; or

2 One additional dose with the most current variant-matched vaccine every 12 months for children aged 6 months to 4 years old at high risk of severe illness.

⬇ Inj 10 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3 ml, 0.48 ml single-dose vial; paediatric vaccine, light blue cap – **5% DV Feb-26 to 30 Sep 2027** ..... 0.00 | 10 | **Comirnaty (LP.8.1)**

➥ **Restricted** (RS2172)

Initiation – initial dose

Either:

1 One dose for previously unvaccinated children aged 5-11 years old; or

2 Up to three doses for immunocompromised children aged 5-11 years old.

Continuation – additional dose

Either:

1 One additional dose with the most current variant-matched vaccine every 6 months for highly immunocompromised children aged 5 to 11 years old; or

2 One additional dose with the most current variant-matched vaccine up to every 12 months for children aged 5 to 11 years old at high-risk of severe illness.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
⬇ Inj 10 mcg bretovameran per 0.3 ml, 0.48 ml vial; paediatric vaccine, light blue cap	0.00	10	Comirnaty Omicron (JN.1)

➥ **Restricted** (RS2172)
Initiation – initial dose
Either:
1 One dose for previously unvaccinated children aged 5-11 years old; or
2 Up to three doses for immunocompromised children aged 5-11 years old.

Continuation – additional dose
Either:
1 One additional dose with the most current variant-matched vaccine every 6 months for highly immunocompromised children aged 5 to 11 years old; or
2 One additional dose with the most current variant-matched vaccine up to every 12 months for children aged 5 to 11 years old at high-risk of severe illness.

⬇ Inj 30 mcg SARS-CoV-2 spike protein (mRNA) LP.8.1 per 0.3 ml, pre-filled syringe; adult dose – **5% DV Feb-26 to 30 Sep 2027**	0.00	10	**Comirnaty (LP.8.1)**

➥ **Restricted** (RS2173)
Initiation – initial dose
Any of the following:
1 One dose for previously unvaccinated people aged 12-15 years and over 30 years old; or
2 Two doses for previously unvaccinated people aged 16-29 years old; or
3 Up to three doses for previously unvaccinated immunocompromised people from 12 years old; or
4 Up to four doses for people at risk of severe illness aged from 12-29 years.

Continuation – additional dose
Both:
1 One additional dose with the most current variant-matched vaccine every 6 months, additional dose to be given at least 6 months after last dose; and
2 Any of the following:
  2.1 Previously vaccinated people aged 30 years and over; or
  2.2 Previously vaccinated immunocompromised people from 12 years; or
  2.3 Previously vaccinated people at high-risk of severe illness from 12 years.

⬇ Inj 30 mcg bretovameran per 0.3 ml, 0.48 ml vial; adult vaccine, light grey cap	0.00	10	Comirnaty Omicron (JN.1)

➥ **Restricted** (RS2173)
Initiation – initial dose
Any of the following:
1 One dose for previously unvaccinated people aged 12-15 years and over 30 years old; or
2 Two doses for previously unvaccinated people aged 16-29 years old; or
3 Up to three doses for previously unvaccinated immunocompromised people from 12 years old; or
4 Up to four doses for people at risk of severe illness aged from 12-29 years.

Continuation – additional dose
Both:
1 One additional dose with the most current variant-matched vaccine every 6 months, additional dose to be given at least 6 months after last dose; and

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

2 Any of the following:
- 2.1 Previously vaccinated people aged 30 years and over; or
- 2.2 Previously vaccinated immunocompromised people from 12 years; or
- 2.3 Previously vaccinated people at high-risk of severe illness from 12 years.

(Comirnaty Omicron (JN.1) Inj 3 mcg bretovameran per 0.3 ml, 0.48 ml vial; infant vaccine, yellow cap to be delisted 1 June 2026)
(Comirnaty Omicron (JN.1) Inj 10 mcg bretovameran per 0.3 ml, 0.48 ml vial; paediatric vaccine, light blue cap to be delisted 1 June 2026)
(Comirnaty Omicron (JN.1) Inj 30 mcg bretovameran per 0.3 ml, 0.48 ml vial; adult vaccine, light grey cap to be delisted 1 June 2026)

HEPATITIS A VACCINE – **Restricted** see terms below

⬇	Inj 720 ELISA units in 0.5 ml syringe – **5% DV Dec-24 to 2027** 0.00	1	**Havrix Junior**
⬇	Inj 1440 ELISA units in 1 ml syringe – **5% DV Dec-24 to 2027** 0.00	1	**Havrix 1440**

➥ **Restricted** (RS1638)

Initiation

Any of the following:
1 Two vaccinations for use in transplant patients; or
2 Two vaccinations for use in children with chronic liver disease; or
3 One dose of vaccine for close contacts of known hepatitis A cases.

HEPATITIS B RECOMBINANT VACCINE

⬇	Inj 10 mcg per 0.5 ml prefilled syringe – **5% DV Dec-24 to 2027** 0.00	1	**Engerix-B**

➥ **Restricted** (RS2049)

Initiation

Any of the following:
1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
4 For HIV positive patients; or
5 For hepatitis C positive patients; or
6 For patients following non-consensual sexual intercourse; or
7 For patients prior to planned immunosuppression for greater than 28 days; or
8 For patients following immunosuppression; or
9 For solid organ transplant patients; or
10 For post-haematopoietic stem cell transplant (HSCT) patients; or
11 Following needle stick injury.

⬇	Inj 20 mcg per 1 ml prefilled syringe – **5% DV Dec-24 to 2027** 0.00	1	**Engerix-B**

➥ **Restricted** (RS2174)

Initiation

Any of the following:
1 For household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or
2 For children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or
3 For children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or
4 For HIV positive patients; or
5 For hepatitis C positive patients; or
6 For patients following non-consensual sexual intercourse; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

7 For patients prior to planned immunosuppression for greater than 28 days; or
8 For patients following immunosuppression; or
9 For solid organ transplant patients; or
10 For post-haematopoietic stem cell transplant (HSCT) patients; or
11 Following needle stick injury; or
12 For chronic kidney disease (CKD) stage 4 or 5 patients; or
13 For liver or kidney transplant patients.

HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] – **Restricted** see terms below

⬇ Inj 270 mcg in 0.5 ml syringe – **5% DV Dec-24 to 2027** ..... 0.00 | 10 | **Gardasil 9**

➥ **Restricted** (RS2038)

Initiation – Children aged 14 years and under

Therapy limited to 2 doses

Children aged 14 years and under.

Initiation – other conditions

Either:

1 Up to 3 doses for people aged 15 to 26 years inclusive; or
2 Both:
  2.1 People aged 9 to 26 years inclusive; and
  2.2 Any of the following:
    2.2.1 Up to 3 doses for confirmed HIV infection; or
    2.2.2 Up to 3 doses people with a transplant (including stem cell); or
    2.2.3 Up to 4 doses for Post chemotherapy.

Initiation – Recurrent Respiratory Papillomatosis

All of the following:

1 Either:
  1.1 Maximum of two doses for children aged 14 years and under; or
  1.2 Maximum of three doses for people aged 15 years and over; and
2 The person has recurrent respiratory papillomatosis; and
3 The person has not previously had an HPV vaccine.

INFLUENZA VACCINE

⬇ Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)..... 120.00 | 10 | Influvac Tetra (2026 formulation)

➥ **Restricted** (RS2013)

Initiation – People over 65

The patient is 65 years of age or over.

Initiation – cardiovascular disease

Any of the following:

1 Ischaemic heart disease; or
2 Congestive heart failure; or
3 Rheumatic heart disease; or
4 Congenital heart disease; or
5 Cerebro-vascular disease.

Note: hypertension and/or dyslipidaemia without evidence of end-organ disease is excluded from funding.

Initiation – chronic respiratory disease

Either:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 Asthma, if on a regular preventative therapy; or
2 Other chronic respiratory disease with impaired lung function.

Note: asthma not requiring regular preventative therapy is excluded from funding.

Initiation – Other conditions

Either:

1 Any of the following:
- 1.1 Diabetes; or
- 1.2 chronic renal disease; or
- 1.3 Any cancer, excluding basal and squamous skin cancers if not invasive; or
- 1.4 Autoimmune disease; or
- 1.5 Immune suppression or immune deficiency; or
- 1.6 HIV; or
- 1.7 Transplant recipient; or
- 1.8 Neuromuscular and CNS diseases/ disorders; or
- 1.9 Haemoglobinopathies; or
- 1.10 Is a child on long term aspirin; or
- 1.11 Has a cochlear implant; or
- 1.12 Errors of metabolism at risk of major metabolic decompensation; or
- 1.13 Pre and post splenectomy; or
- 1.14 Down syndrome; or
- 1.15 Is pregnant; or
- 1.16 Is a child 4 years of age or under (inclusive) who has been hospitalised for respiratory illness or has a history of significant respiratory illness; or

2 Patients in a long-stay inpatient mental health care unit or who are compulsorily detained long-term in a forensic unit within a Public Hospital.

Initiation – Serious mental health conditions or addiction

Any of the following:

1 schizophrenia; or
2 major depressive disorder; or
3 bipolar disorder; or
4 schizoaffective disorder; or
5 person is currently accessing secondary or tertiary mental health and addiction services.

MEASLES, MUMPS AND RUBELLA VACCINE – **Restricted** see terms below

⬇ Injection, measles virus 1,000 CCID50, mumps virus 5,012 CCID50, Rubella virus 1,000 CCID50; prefilled syringe/ampoule of diluent 0.5 ml – **5% DV Dec-24 to 2027** ..... 0.00 10 **Priorix**

➥ **Restricted** (RS1487)

Initiation – first dose prior to 12 months

Therapy limited to 3 doses

Any of the following:

1 For primary vaccination in children; or
2 For revaccination following immunosuppression; or
3 For any individual susceptible to measles, mumps or rubella.

Initiation – first dose after 12 months

Therapy limited to 2 doses

Any of the following:

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1 For primary vaccination in children; or
2 For revaccination following immunosuppression; or
3 For any individual susceptible to measles, mumps or rubella.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

POLIOMYELITIS VACCINE – **Restricted** see terms below

⬇ Inj 80 D-antigen units in 0.5 ml syringe – **5% DV Dec-24 to 2027** ..... 0.00 1 **IPOL**

➥ **Restricted** (RS1398)

Initiation

Therapy limited to 3 doses

Either:

1 For partially vaccinated or previously unvaccinated individuals; or
2 For revaccination following immunosuppression.

Note: Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

RABIES VACCINE

Inj 2.5 IU vial with diluent

ROTAVIRUS ORAL VACCINE – **Restricted** see terms below

⬇ Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator – **5% DV Dec-24 to 2027** ..... 0.00 10 **Rotarix**

⬇ Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube ..... 0.00 10 Rotarix

⬇ Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, squeezable tube (PVC free) ..... 0.00 10 Rotarix

➥ **Restricted** (RS1590)

Initiation

Therapy limited to 2 doses

Both:

1 First dose to be administered in infants aged under 14 weeks of age; and
2 No vaccination being administered to children aged 24 weeks or over.

VARICELLA VACCINE [CHICKENPOX VACCINE]

⬇ Inj 2000 PFU prefilled syringe plus vial – **5% DV Dec-24 to 2027** ..... 0.00 10 **Varilrix**

➥ **Restricted** (RS1591)

Initiation – primary vaccinations

Therapy limited to 1 dose

Either:

1 Any infant born on or after 1 April 2016; or
2 For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox).

Initiation – other conditions

Therapy limited to 2 doses

Any of the following:

1 Any of the following:
for non-immune patients:
1.1 With chronic liver disease who may in future be candidates for transplantation; or
1.2 With deteriorating renal function before transplantation; or
1.3 Prior to solid organ transplant; or
1.4 Prior to any elective immunosuppression*; or

continued…

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer

continued…

1.5 For post exposure prophylaxis who are immune competent inpatients; or

2 For patients at least 2 years after bone marrow transplantation, on advice of their specialist; or

3 For patients at least 6 months after completion of chemotherapy, on advice of their specialist; or

4 For HIV positive patients non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist; or

5 For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella; or

6 For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella; or

7 For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

Note: * immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] – **Restricted** see terms below

⬇ Inj 50 mcg per 0.5 ml vial plus vial	0.00	1	Shingrix
		10	Shingrix

➥ **Restricted** (RS2039)

Initiation – people aged 18 years and over (Shingrix)

Therapy limited to 2 doses

Any of the following:

1 Pre- and post-haematopoietic stem cell transplant or cellular therapy; or

2 Pre- or post-solid organ transplant; or

3 Haematological malignancies; or

4 People living with poorly controlled HIV infection; or

5 Planned or receiving disease modifying anti-rheumatic drugs (DMARDs – targeted synthetic, biologic, or conventional synthetic) for polymyalgia rheumatica, systemic lupus erythematosus or rheumatoid arthritis; or

6 End stage kidney disease (CKD 4 or 5);; or

7 Primary immunodeficiency.

Diagnostic Agents

TUBERCULIN PPD [MANTOUX] TEST

Inj 5 TU per 0.1 ml, 1 ml vial – **5% DV Dec-24 to 2027**	0.00	1	**Tubersol**

Optional Pharmaceuticals

NOTE:
In addition to the products expressly listed here in Part III: Optional Pharmaceuticals, a range of hospital medical devices are listed in an addendum to Part III which is available at schedule.pharmac.govt.nz. The Optional Pharmaceuticals listed in the addendum are deemed to be listed in Part III, and the Rules of the Pharmaceutical Schedule applying to products listed in Part III apply to them.

	Price (ex man. excl. GST) $	Per	Brand or Generic Manufacturer
BETA-HCG LOW SENSITIVITY URINE TEST KIT			
Note: For use in abortion services only.			
Midstream	16.28	1 test	CheckTop
BLOOD GLUCOSE DIAGNOSTIC TEST METER			
1 meter with 50 lancets, a lancing device, and 10 diagnostic test strips	20.00 10.00	1	CareSens N Premier Caresens N Caresens N POP
BLOOD GLUCOSE DIAGNOSTIC TEST STRIP			
Blood glucose test strips	10.56	50 test	CareSens N
Test strips	10.56	50 test	CareSens PRO
BLOOD KETONE DIAGNOSTIC TEST STRIP			
Test strips	15.50	10 strip	KetoSens
DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER			
Meter with 50 lancets, a lancing device, and 10 blood glucose diagnostic test strips	20.00	1	CareSens Dual
MASK FOR SPACER DEVICE			
Small	2.70	1	e-chamber Mask
PEAK FLOW METER			
Low Range	9.54	1	Mini-Wright AFS Low Range
Normal Range	9.54	1	Mini-Wright Standard
PREGNANCY TEST - HCG URINE			
Cassette – **5% DV Mar-25 to 2027**	16.00	40 test	**David One Step Cassette Pregnancy Test**
SODIUM NITROPRUSSIDE			
Test strip	22.00	50 strip	Ketostix
SPACER DEVICE			
220 ml (single patient)	3.65	1	e-chamber Turbo
510 ml (single patient)	5.95	1	e-chamber La Grande
800 ml	6.50	1	Volumatic

- Symbols -
Xaluprine ... 154
8-methoxypsoralen ... 73
- A -
A-Scabies ... 69
Abacavir sulphate ... 106
Abacavir sulphate with lamivudine ... 106
Abacavir/lamivudine Viatris ... 106
Abciximab ... 184
Abilify Maintena ... 138
Abiraterone acetate ... 175
Acarbose ... 9
Accarb ... 9
Acetazolamide ... 279
Acetec ... 45
Acetic acid
Extemporaneously Compounded Preparations ... 290
Genito-Urinary ... 75
Acetic acid with hydroxyquinoline, glycerol and ricinoleic acid ... 75
Acetic acid with propylene glycol ... 281
Acetylcholine chloride ... 279
Acetylcysteine ... 282
Aciclovir
Infections ... 109
Sensory ... 275
Aciclovir-Baxter ... 109
Acid Citrate Dextrose A ... 36
Acidex ... 5
Acipimox ... 54
Acitretin ... 72
Act-HIB ... 312
Actemra ... 235
Actinomycin D ... 152
Adalimumab (Amgevita) ... 184
Adalimumab (Humira - alternative brand) ... 194
Adapalene ... 69
Adcetris ... 204
Adenosine ... 47
Adenosine Baxter ... 47
Adrenaline
Cardiovascular ... 55
Respiratory ... 264
Adsine ... 47
Advantan ... 72
Advate ... 35
Adynovate ... 35
Aerrane ... 123
Afinitor ... 258
Aflibercept ... 201
Agents Affecting the Renin-Angiotensin System ... 45
Agents for Parkinsonism and Related Disorders ... 122
Agents Used in the Treatment of Poisonings ... 282
Ajmaline ... 47
Albalon ... 277
Albendazole ... 103
Alchemy Caspofungin ... 101
Alchemy Oxaliplatin ... 162
Alchemy Oxybutynin ... 78
Aldara ... 74
Aldurazyme ... 19
Alecensa ... 162
Alectinib ... 162
Alendronate sodium ... 116
Alendronate sodium with colecalciferol ... 116
Alfacalcidol ... 28
Alfamino ... 303
Alfamino Junior ... 303
Alfentanil ... 127
Alglucosidase alfa ... 17
Allegron ... 130
Allerfix ... 277
Allerpro Syneo 1 ... 305
Allerpro Syneo 2 ... 305
Allersoothe ... 265
Allmercap ... 154
Allopurinol ... 118
Almarytm ... 48
Alpha tocopheryl ... 28
Alpha tocopheryl acetate ... 29
Alpha-Adrenoceptor Blockers ... 47
Alphamox 125 ... 94
Alphamox 250 ... 94
Alprolix ... 34
Alprostadil ... 56
Alprostadil hydrochloride ... 56
Alteplase ... 39
Alum ... 290
Aluminium chloride ... 32
Aluminium hydroxide ... 5
Aluminium hydroxide with magnesium hydroxide and simeticone ... 5
Alyftrek ... 273
Amantadine hydrochloride ... 122
Ambrisentan ... 57
Ambrisentan Viatris ... 57
Amethocaine
Nervous ... 126
Sensory ... 278
Amgevita ... 184
Amikacin ... 90
Amiloride hydrochloride ... 51
Amiloride hydrochloride with furosemide ... 51
Amiloride hydrochloride with hydrochlorothiazide ... 51
Aminolevulinic acid hydrochloride ... 178
Aminophylline ... 271
Amiodarone hydrochloride ... 47
Amisulpride ... 136
Amitriptyline ... 129
Amlodipine ... 49
Amorolfine ... 68
Amoxicillin ... 94
Amoxicillin with clavulanic acid ... 94
Amoxiclav Devatis Forte ... 94
Amoxycillin Sandoz ... 94
Amphotericin B
Alimentary ... 26
Infections ... 98
Amphotericin Liposomal SUN ... 98
Amsacrine ... 154
Amyl nitrite ... 282
Anabolic Agents ... 79
Anaesthetics ... 123
Anagrelide hydrochloride ... 154
Analgesics ... 126
Anastrozole ... 177
Anatrole ... 177
Androgen Agonists and Antagonists ... 79
Anoro Ellipta ... 266
Antabuse ... 149
Antacids and Antiflatulents ... 5
Anti-Infective Agents ... 75
Anti-Infective Preparations
Dermatological ... 68
Sensory ... 275
Anti-Inflammatory Preparations ... 276
Antiacne Preparations ... 69
Antiallergy Preparations ... 264
Antianaemics ... 30
Antiarrhythmics ... 47
Antibacterials ... 90
Anticholinergic Agents ... 266
Anticholinesterases ... 116
Antidepressants ... 129
Antidiarrhoeals and Intestinal Anti-Inflammatory Agents ... 5
Antiepilepsy Drugs ... 131
Antifibrinolytics, Haemostatics and Local Sclerosants ... 32

Antifibrotics ... 268
Antifungals ... 98
Antihypotensives ... 48
Antimigraine Preparations ... 134
Antimycobacterials ... 102
Antinausea and Vertigo Agents ... 134
Antiparasitics ... 103
Antipruritic Preparations ... 69
Antipsychotic Agents ... 136
Antiretrovirals ... 105
Antirheumatoid Agents ... 116
Antiseptics and Disinfectants ... 284
Antispasmodics and Other Agents Altering Gut Motility ... 7
Antithrombotics ... 36
Antithymocyte globulin (equine) ... 257
Antithymocyte globulin (rabbit) ... 257
Antiulcerants ... 7
Antivirals ... 108
Anxiolytics ... 140
Anzatax ... 174
Apidra ... 10
Apidra Solostar ... 10
APO Clomipramine ... 130
APO-Atomoxetine ... 145
APO-Candesartan HCTZ 16/12.5 ... 46
APO-Candesartan HCTZ 32/12.5 ... 46
APO-Olanzapine ... 137
Apomorphine hydrochloride ... 122
Apraclonidine ... 280
Aprepitant ... 134
Apresoline ... 56
Aprotinin ... 32
Aptamil Feed Thickener ... 295
Aqueous cream ... 70
Arachis oil [Peanut oil] ... 290
Aratac ... 47
Arava ... 116
Arginine
Alimentary ... 18
Various ... 286
Arginine2000 ... 299
Argipressin [Vasopressin] ... 89
Aripiprazole ... 136, 138
Aripiprazole Sandoz ... 136
Aristocort ... 72
Arrotex-Prazosin S29 ... 47
Arrow - Clopid ... 37
Arrow - Lattim ... 279
Arrow-Amitriptyline ... 129
Arrow-Bendrofluazide ... 52
Arrow-Brimonidine ... 280
Arrow-Diazepam ... 140
Arrow-Fluoxetine ... 131
Arrow-Losartan & Hydrochlorothiazide ... 46
Arrow-Norfloxacin ... 96
Arrow-Ornidazole ... 104
Arrow-Quinapril 10 ... 45
Arrow-Quinapril 20 ... 45
Arrow-Quinapril 5 ... 45
Arrow-Roxithromycin ... 94
Arrow-Timolol ... 279
Arrow-Topiramate ... 133
Arrow-Tramadol ... 129
Arsenic trioxide ... 154
Artemether with lumefantrine ... 103
Artesunate ... 103
Articaine hydrochloride ... 124
Articaine hydrochloride with adrenaline ... 124
Asacol ... 6
Ascorbic acid
Alimentary ... 28
Extemporaneously Compounded Preparations ... 290
Aspen Adrenaline ... 55
Aspirin
Blood ... 37
Nervous ... 126
Asthalin ... 269
Atazanavir sulphate ... 107
Atazanavir Viatris ... 107
Atenolol ... 48
Atenolol Viatris ... 48
Atenolol-AFT ... 48
Atezolizumab ... 245
ATGAM ... 257
Ativan ... 140
Atomoxetine ... 145
Atorvastatin ... 53
Atovaquone with proguanil hydrochloride ... 104
Atracurium besylate ... 119
Atropine sulphate
Cardiovascular ... 47
Sensory ... 280
Atropt ... 280
Augmentin ... 94
Aurorix ... 130
Avastin ... 204
Avelox ... 95
Avonex ... 142
Axitinib ... 162
Azacitidine ... 153
Azacitidine Dr Reddy's ... 153
Azactam ... 97
Azamun ... 258
Azathioprine ... 258
Azilect ... 123
Azithromycin ... 92
Azopt ... 279
Aztreonam ... 97

- B -

Bacillus calmette-guerin (BCG) ... 258
Bacillus calmette-guerin vaccine ... 311
Baclofen ... 119
Bacterial and Viral Vaccines ... 311
Bacterial Vaccines ... 311
Balanced Salt Solution ... 278
Barium sulphate ... 285
Barrier Creams and Emollients ... 69
Basiliximab ... 202
BCG Vaccine AJV ... 311
BD PosiFlush ... 42
Beclazone 100 ... 270
Beclazone 250 ... 270
Beclazone 50 ... 270
Beclomethasone dipropionate ... 270
Bedaquiline ... 102
Bee venom ... 264
Bendamustine hydrochloride ... 151
Bendamustine Sandoz ... 151
Bendrofluazide ... 52
Bendroflumethiazide [Bendrofluazide] ... 52
Benralizumab ... 202
Benzathine benzylpenicillin ... 94
Benzatropine mesylate ... 122
Benzbromaron AL 100 ... 118
Benzbromarone ... 118
Benzetacil ... 94
Benzocaine ... 124
Benzocaine with tetracaine hydrochloride ... 124
Benzoin ... 290
Benzoyl peroxide ... 69
Benztrop ... 122
Benzydamine hydrochloride ... 25
Benzydamine hydrochloride with cetylpyridinium chloride ... 26
Benzylpenicillin sodium [Penicillin G] ... 94
Beractant ... 273
Besponsa ... 214
Beta Cream ... 71
Beta Ointment ... 71
Beta Scalp ... 73
Beta-Adrenoceptor Agonists ... 269
Beta-Adrenoceptor Blockers ... 48
Beta-hCG low sensitivity urine test kit ... 323
Betadine ... 284
Betahistine dihydrochloride ... 135

Betaine 18
Betamethasone 80
Betamethasone dipropionate 71
Betamethasone dipropionate with calcipotriol 72
Betamethasone sodium phosphate with betamethasone acetate 80
Betamethasone valerate 71, 73
Betamethasone valerate with clioquinol 72
Betamethasone valerate with sodium fusidate [Fusidic acid] 72
Betaxolol 279
Betnovate 71
Bevacizumab 203
Bevacizumab (Ocular) 204
Bexsero 313
Bezafibrate 53
Bezalip 53
Bezalip Retard 53
Bicalutamide 176
Bicillin LA 94
BiCNU 152
BiCNU S29 152
Bile and Liver Therapy 9
Biliscopin 286
Bimatoprost 279
Binarex 176
Binocrit 30
Biocon 49
Biodone 128
Biodone Extra Forte 128
Biodone Forte 128
Biotin 18
Bisacodyl 16
Bisacodyl Viatris 16
Bisacodyl-AFT 16
Bismuth subgallate 290
Bismuth subnitrate and iodoform paraffin 288
Bisoprolol fumarate 48
Bivalirudin 36
Bleomycin sulphate 152
Blood glucose diagnostic test meter 323
Blood glucose diagnostic test strip 323
Blood ketone diagnostic test strip 323
Bonney's blue dye 287
Boostrix 312
Boric acid 290
Bortezomib 155
Bortezomib Eugia 155
Bosentan 59
Bosentan Dr Reddy's 59
Botox 119
Botulism antitoxin 282
Bplex 28
Brentuximab vedotin 204
Breo Ellipta 271
Brevinor 1/28 75
Breztri Aerosphere 267
Bricanyl Turbuhaler 269
Brimonidine tartrate 280
Brimonidine tartrate with timolol maleate 280
Brinzolamide 279
Bromocriptine 122
Budesonide
Alimentary 5
Respiratory 265, 270
Budesonide Te Arai 5
Budesonide with eformoterol 271
Budesonide with glycopyrronium and eformoterol 267
Bumetanide 51
Bupafen 125
Bupivacaine hydrochloride 124
Bupivacaine hydrochloride with adrenaline 124
Bupivacaine hydrochloride with fentanyl 125
Bupivacaine hydrochloride with glucose 125
Buprenorphine Naloxone BNM 148
Buprenorphine with naloxone 148
Bupropion hydrochloride 149
Burinex 51
Buserelin 83
Buspirone hydrochloride 140
Buspirone Viatris 140
Busulfan 152

- C -

Cabergoline 82
Caffeine 145
Caffeine citrate 271
Calamine 69
Calci-Tab 500 23
Calcipotriol 72
Calcitonin 79
Calcitriol 28
Calcitriol XL 28
Calcitriol-AFT 28
Calcium carbonate 5, 23
Calcium carbonate PAI 5
Calcium Channel Blockers 49
Calcium chloride 40
Calcium folinate 174
Calcium gluconate
Blood 40
Dermatological 74
Calcium Homeostasis 79
Calcium polystyrene sulphonate 43
Calcium Resonium 43
Calogen (neutral) 293
Calogen (strawberry) 293
Candesartan cilexetil 46
Candesartan cilexetil with hydrochlorothiazide 46
Candesartan Viatris 46
Candesartan Viatris Sante 46
Candestar 46
Capecitabine 153
Capecitabine Viatris 153
Capsaicin
Musculoskeletal 121
Nervous 126
Captopril 45
Carbachol 279
Carbamazepine 131
Carbasorb-X 283
Carbimazole 88
Carbomer 280
Carboplatin 162
Carboplatin Accord 162
Carboprost trometamol 76
Carboxymethylcellulose
Alimentary 26
Extemporaneously Compounded Preparations 290
Cardinol LA 49
Cardizem CD 50
CareSens Dual 323
Caresens N 323
Caresens N POP 323
CareSens N Premier 323
CareSens PRO 323
Carglumic acid 18
Carmellose sodium 280
Carmellose sodium with pectin and gelatine 26
Carmustine 152
Carvedilol 49
Carvedilol Sandoz 49
Caspofungin 101
Catapres 51
Cefaclor 91
Cefalexin 91
Cefalexin Lupin 91
Cefalexin Sandoz 91
Cefazolin 91
Cefazolin-AFT 91
Cefepime 92
Cefepime-AFT 92
Cefotaxime 91
Cefotaxime Sandoz 91
Cefoxitin 91

Ceftaroline fosamil....92
Ceftazidime....91
Ceftazidime Kabi....91
Ceftazidime with avibactam....91
Ceftriaxone....92
Ceftriaxone-AFT....92
Cefuroxime....91
Cefuroxime Devatis....91
Celapram....130
Celebrex....120
Celecoxib....120
Celecoxib Pfizer....120
Celiprolol....49
CellCept....258
Celostea....120
Centrally-Acting Agents....50
Cephalexin ABM....91
Cerazette....76
Cerobact....94
Cerubidine....153
Cetirizine hydrochloride....265
Cetomacrogol....70
Cetomacrogol Cream AFT....70
Cetomacrogol with glycerol....70
Cetomacrogol-AFT....70
Cetrimide....290
Cetuximab....205
Champix....150
Charcoal....283
CheckTop....323
Chemotherapeutic Agents....151
Chickenpox vaccine....321
Chlorafast....275
Chloral hydrate....143
Chlorambucil....152
Chloramphenicol
  Infections....97
  Sensory....275
Chlorhexidine....284
Chlorhexidine gluconate
  Alimentary....26
  Extemporaneously Compounded Preparations....290
  Genito-Urinary....75
Chlorhexidine with cetrimide....284, 287
Chlorhexidine with ethanol....284
Chloroform....290
Chloroquine phosphate....104
Chlorothiazide....52
Chlorpheniramine maleate....265
Chlorpromazine hydrochloride....136
Chlortalidone [Chlorthalidone]....52
Chlorthalidone....52
Choice 380 7med Nsha Silver/copper Short....76
Cholestyramine....54
Choriogonadotropin alfa....84
Ciclopirox olamine....68
Ciclosporin....179
Cidofovir....109
Cilicaine VK....95
Cimetidine....8
Cinacalcet....79
Cinacalet Devatis....79
Cinchocaine hydrochloride with hydrocortisone....7
Cipla....266
Ciprofloxacin
  Infections....95
  Sensory....275
Ciprofloxacin Kabi....95
Ciprofloxacin Teva....275
Ciprofloxacin with hydrocortisone....275
Ciproxin HC Otic....275
Cisplatin....162
Cisplatin Accord....162
Citalopram hydrobromide....130
Citanest....126
Citrate sodium....36
Citric acid....290
Citric acid with magnesium carbonate hydrate and sodium picosulfate....15
Citric acid with sodium bicarbonate....285
Citrulline1000....299
Cladribine....153
Clarithromycin....93
Clexane....36
Clexane Forte....36
Clindamycin....97
Clinicians....28
Clinicians Multivit & Mineral Boost....26
Clinicians Renal Vit....27
Clobazam....131
Clobetasol propionate....71, 73
Clobetasone butyrate....71
Clofazimine....102
Clomazol
  Dermatological....68
  Genito-Urinary....75
Clomifene citrate....83
Clomipramine hydrochloride....130
Clonazepam....131, 140
Clonidine....50
Clonidine hydrochloride....51
Clonidine Teva....51
Clopidogrel....37
Clopine....136
Clopixol....137, 140
Clostridium botulinum type A toxin....119
Clotrimazole
  Dermatological....68
  Genito-Urinary....75
Clove oil....290
Clozapine....136
Clozaril....136
Clustran....134
Co-trimoxazole....98
Coal tar....290
Coal tar with salicylic acid and sulphur....72
Cocaine hydrochloride....125
Cocaine hydrochloride with adrenaline....125
Codeine phosphate
  Extemporaneously Compounded Preparations....290
  Nervous....127
Coenzyme Q10....18
Colchicine....118
Colecalciferol....28
Colestimethate....97
Colestipol hydrochloride....54
Colestyramine....54
Colestyramine - Mylan....54
Colgout....118
Colifoam....6
Colistin sulphomethate [Colestimethate]....97
Collodion flexible....290
Colloidal bismuth subcitrate....9
Colofac....7
Colomycin....97
Colony-Stimulating Factors....39
Coloxyl....15
Combigan....280
Combiprasal....266
Comirnaty (LP.8.1)....316–317
Comirnaty Omicron (JN.1)....316–317
Compound electrolytes....41, 43
Compound electrolytes with glucose [Dextrose]....41, 43
Compound hydroxybenzoate....290
Compound sodium lactate [Hartmann's solution]....41
Concerta....147
Condyline....73
Contraceptives....75
Contrast Media....285
Copaxone....142
Copper....23
Copper chloride....23

Corticorelin (ovine) 83
Corticosteroids
Dermatological 71
Hormone Preparations 80
Cosentyx 232
Cosmegen 152
Coversyl 45
COVID-19 vaccine 316
Creon 10000 14
Creon 25000 14
Creon Micro 14
Crizotinib 162
Crotamiton 69
Crystaderm 68
Cu 375 Standard 76
Curam Duo 500/125 94
Curosurf 273
Cvite 28
Cyclizine hydrochloride 135
Cyclizine lactate 135
Cyclogyl 280
Cyclonex 152
Cyclopentolate hydrochloride 280
Cyclophosphamide 152
Cycloserine 102
Cymevene 109
Cyproheptadine hydrochloride 265
Cyproterone acetate 79
Cyproterone acetate with ethinyloestradiol 75
Cystadane 18
Cysteamine hydrochloride 290
Cytarabine 153
Cytotec 7

- D -

D-Penamine 116
Dabigatran 36
Dabrafenib 163
Dacarbazine 155
Dactinomycin [Actinomycin D] 152
Daivobet 72
Daivonex 72
Dalacin C 97
Danaparoid 36
Dantrium 119
Dantrium IV 119
Dantrolene 119
Daonil 11
Dapa-Tabs 52
Dapsone 102
Daptomycin 97
Daptomycin Dr Reddy's 97
Daptomycin-AFT 97
Darunavir 107
Darunavir Viatris 107
Dasatinib 164
Dasatinib-Teva 164
Daunorubicin 153
David One Step Cassette Pregnancy Test 323
DBL Acetylcysteine 282
DBL Adrenaline 55
DBL Amikacin 90
DBL Aminophylline 271
DBL Bleomycin Sulfate 152
DBL Bortezomib 155
DBL Cefotaxime 91
DBL Dacarbazine 155
DBL Desferrioxamine Mesylate for Inj BP 283
DBL Docetaxel 174
DBL Ergometrine 76
DBL Gemcitabine 154
DBL Gentamicin 90
DBL Leucovorin Calcium 174
DBL Methotrexate Onco-Vial 154
DBL Naloxone Hydrochloride 282
DBL Pethidine Hydrochloride 129
DBL Vincristine Sulfate 175
Decongestants 269
Decongestants and Antiallergics 277
Decozol 26
Deferasirox 283
Deferiprone 283
Defibrotide 36
Definity 286
Demeclocycline hydrochloride 96
Denosumab 117
Deolate 102
Deoxycoformycin 158
Depo-Medrol 81
Depo-Provera 76
Depo-Testosterone 79
Deprim 98
Dermol 71, 73
Desferrioxamine mesilate 283
Desflurane 123
Desmopressin 89
Desmopressin acetate 89
Desmopressin-PH&T 89
Desogestrel 76
Dexamethasone
Hormone Preparations 81
Sensory 276
Dexamethasone phosphate 81
Dexamethasone with framycetin and gramicidin 275
Dexamethasone with neomycin sulphate and polymyxin B sulphate 275
Dexamethasone with tobramycin 275
Dexamfetamine sulfate 146
Dexmedetomidine 123
Dexmedetomidine Viatris 123
Dexmethsone 81
Dexrazoxane 174
Dextrose
Alimentary 9
Blood 41, 43
Extemporaneously Compounded Preparations 290
Dextrose with sodium citrate and citric acid [Acid Citrate Dextrose A] 36
DHC Continus 127
Diabetes 9
Diacomit 133
Diagnostic Agents
Vaccines 322
Various 286
Diagnostic and Surgical Preparations 277
Diamide Relief 5
Diasip (strawberry) 300
Diasip (vanilla) 300
Diatrizoate meglumine with sodium amidotrizoate 285
Diatrizoate sodium 285
Diazepam 131, 140
Diazoxide
Alimentary 9
Cardiovascular 56
Dichlorobenzyl alcohol with amylmetacresol 26
Diclofenac Devatis 277
Diclofenac Sandoz 120
Diclofenac sodium
Musculoskeletal 120
Sensory 277
Dicobalt edetate 284
Diflucan 99
Diflucortolone valerate 71
Digestives Including Enzymes 14
Digoxin 48
Digoxin immune Fab 282
Dihydrocodeine tartrate 127
Dihydroergotamine mesylate 134
Diltiazem CD Clinect 50
Diltiazem hydrochloride 50
Dimercaprol 284
Dimercaptosuccinic acid 284
Dimethicone 68–69
Dimethyl fumarate 141
Dimethyl sulfoxide 288
Dinoprostone 76
Dipentum 7

Diphemanil metilsulfate 73
Diphenoxylate hydrochloride with atropine sulphate 5
Diphtheria antitoxin 283
Diphtheria, tetanus and pertussis vaccine 312
Diphtheria, tetanus, pertussis and polio vaccine 311
Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine 311
Diprosone 71
Dipyridamole 37
Dipyridamole - Strides 37
Disodium edetate 278
Disodium hydrogen phosphate with sodium dihydrogen phosphate 290
Disopyramide phosphate 48
Disulfiram 149
Dithranol 290
Diuretics 51
Dobutamine 55
Dobutamine-hameln 55
Docetaxel 174
Docusate sodium
Alimentary 15
Sensory 281
Docusate sodium with sennosides 15
Dolutegravir 108
Dolutegravir with lamivudine 108
Domperidone 135
Domperidone Viatris 135
Donepezil hydrochloride 148
Dopamine Basi 55
Dopamine hydrochloride 55
Dornase alfa 271
Dortimopt 279
Dorzolamide 279
Dorzolamide with timolol 279
Dostinex 82
Dosulepin [Dothiepin] hydrochloride 130
Dosulepin Viatris 130
Dotarem 286
Dothiepin 130
Dovato 108
Doxapram 273
Doxazosin 47
Doxazosin Clinect 47
Doxepin hydrochloride 130
Doxine 96
Doxorubicin Ebewe 153
Doxorubicin hydrochloride 153
Doxycycline 96
DP Lotn HC 71
DP-Captopril 45
Dr Reddy's Omeprazole 8
Drofate 49
Droperidol 135
Drugs Affecting Bone Metabolism 116
Dual blood glucose and blood ketone diagnostic test meter 323
Dulaglutide 11
Dulcolax SP Drop 17
Duocal Super Soluble Powder 294
Duolin 266
DuoResp Spiromax 271
Duovisc 278
Duride 55
Durvalumab 246
Dynastat 121
Dysport 119

- E -

e-chamber La Grande 323
e-chamber Mask 323
e-chamber Turbo 323
E-Mycin 94
E-Z-Gas II 285
Easiphen Liquid 298
Econazole nitrate 68
Edrophonium chloride 116
Efavirenz 105
Efavirenz Milpharm 105
Efavirenz with emtricitabine and tenofovir disoproxil 106
Eformoterol fumarate 270
Eformoterol fumarate dihydrate 270
Eftrenonacog alfa [Recombinant factor IX] 34
Efudix 74
Elaprase 19
Elecare (Unflavoured) 303
Elecare (Vanilla) 303
Elecare LCP (Unflavoured) 303
Electral 43
Electrolytes 289
Elelyso 22
Elemental 028 Extra (grapefruit) 301
Elemental 028 Extra (pineapple & orange) 301
Elemental 028 Extra (summer fruits) 301
Elexacaftor with tezacaftor, ivacaftor and ivacaftor 272
Elidel 73
Elocon 72
Elocon Alcohol Free 72
Eltrombopag 32
Emend Tri-Pack 134
Emicizumab 33
EMLA 126
Empagliflozin 13
Empagliflozin with metformin hydrochloride 13
Emsogen 300
Emtricitabine 106
Emtricitabine with tenofovir disoproxil 111
Emtriva 106
Emulsifying ointment 70
Enalapril maleate 45
Enbrel 179
Endocrine Therapy 175
Endoxan 152
Energivit 298
Engerix-B 318
Enhertu 240
Enlafax XR 130
Enoxaparin sodium 36
Enstilar 72
Ensure (Chocolate) 309
Ensure (Vanilla) 309
Ensure Plus (Banana) 310
Ensure Plus (Chocolate) 310
Ensure Plus (Fruit of the forest) 310
Ensure Plus (Vanilla) 310
Ensure Plus HN RTH 309
Ensure Two Cal HN RTH 302
Entacapone 122
Entacapone Viatris 122
Entecavir 108
Entecavir (Rex) 108
Entrectinib 165
Entresto 24/26 46
Entresto 49/51 46
Entresto 97/103 46
Entyvio 243
Enzymes 118
Ephedrine 55
Ephedrine Aguettant 55
Ephedrine Juno 55
Epilim IV 133
Epipen 264
Epipen Jr 264
Epirubicin Ebewe 153
Epirubicin hydrochloride 153
Eplerenone 51
Epoetin alfa 30
Epoetin beta 31
Epoprostenol 63
Eptacog alfa [Recombinant factor VIIa] 34
Eptifibatide 37

Eptifibatide Viatris ... 37
Erbitux ... 205
Ergometrine maleate ... 76
Erlotinib ... 165
Ertapenem ... 90
Erythrocin IV ... 94
Erythromycin (as ethylsuccinate) ... 94
Erythromycin (as lactobionate) ... 94
Erythromycin (as stearate) ... 94
Esbriet ... 268
Escitalopram ... 131
Esmolol hydrochloride ... 49
Essential Amino Acid Mix ... 300
Estradiol TDP Mylan ... 82
Estradot ... 82
Estrogel ... 82
Etanercept ... 179
Ethambutol hydrochloride ... 102
Ethanol ... 282
Ethanol with glucose ... 282
Ethanol, dehydrated ... 282
Ethics Aspirin ... 126
Ethics Aspirin EC ... 37
Ethinyloestradiol with desogestrel ... 75
Ethinyloestradiol with levonorgestrel ... 75
Ethinyloestradiol with norethisterone ... 75
Ethosuximide ... 131
Ethyl chloride ... 125
Etomidate ... 123
Etopophos ... 155
Etoposide ... 155
Etoposide (as phosphate) ... 155
Etoricoxib ... 120
Etravirine ... 105
Eugia ... 56
Eurofolic ... 174
Evara ... 70
Evara Emulsifying Ointment ... 70
EVARA White Soft Paraffin ... 71
Everet ... 132
Everolimus ... 258
Evrysdi ... 145
Exemestane ... 177
Exjade ... 283
Extemporaneously Compounded Preparations ... 290
Eylea ... 201
Ezetimibe ... 54
Ezetimibe Sandoz ... 54
Ezetimibe with simvastatin ... 54

- F -

Factor eight inhibitor bypassing fraction ... 34
Famotidine ... 8
Famotidine Hovid MY ... 8
Faricimab ... 206
Fasenra ... 202
Fatty Emulsion Cream (Evara) ... 70
Febuxostat ... 118
Febuxostat (Teva) ... 118
FEIBA NF ... 34
Felo 10 ER ... 49
Felo 5 ER ... 49
Felodipine ... 49
Fentanyl ... 128
Fentanyl Sandoz ... 128
Ferinject ... 24
Ferric derisomaltose ... 24
Ferric subsulfate ... 33
Ferriprox ... 283
Ferro-F-Tabs ... 24
Ferro-Liquid ... 24
Ferro-tab ... 24
Ferrosig ... 24
Ferrous fumarate ... 24
Ferrous fumarate with folic acid ... 24
Ferrous gluconate with ascorbic acid ... 24
Ferrous sulfate ... 24
Ferrous sulfate with ascorbic acid ... 24
Fexaclear ... 265
Fexofenadine hydrochloride ... 265
Filgrastim ... 40
Finasteride ... 77
Fingolimod ... 142
Firazyr ... 264
Flagyl ... 104
Flagyl-S ... 104
Flamazine ... 68
Flecainide acetate ... 48
Flecainide BNM ... 48
Flecainide Controlled Release Teva ... 48
Fleet Phosphate Enema ... 16
Flixonase Hayfever & Allergy ... 265
Flixotide ... 270
Flixotide Accuhaler ... 270
Florinef ... 81
Fluanxol ... 138
Flucazole ... 99
Flucil ... 95
Flucloxacillin ... 95
Flucloxin ... 95
Fluconazole ... 99
Fluconazole-Baxter ... 99
Flucytosine ... 101
Fludara Oral ... 153
Fludarabine Ebewe ... 153
Fludarabine phosphate ... 153
Fludrocortisone acetate ... 81
Fluids and Electrolytes ... 40
Flumazenil ... 282
Flumazenil-Baxter ... 282
Flumetasone pivalate with clioquinol ... 276
Fluocortolone caproate with fluocortolone pivalate and cinchocaine ... 7
Fluorescein sodium ... 277
Fluorescein sodium with lignocaine hydrochloride ... 277
Fluorescite ... 277
Fluorometholone ... 277
Fluorouracil ... 154
Fluorouracil Accord ... 154
Fluorouracil sodium ... 74
Fluox ... 131
Fluoxetine hydrochloride ... 131
Flupenthixol decanoate ... 138
Flutamide ... 176
Flutamin ... 176
Fluticasone ... 270
Fluticasone furoate with umeclidinium and vilanterol ... 267
Fluticasone furoate with vilanterol ... 271
Fluticasone propionate ... 265
Fluticasone with salmeterol ... 271
Flynn ... 91
FML ... 277
Foban ... 68
Folic acid ... 31
Folic Acid multichem ... 31
Folic Acid Viatris ... 31
Fondaparinux sodium ... 36
Food Modules ... 293
Food/Fluid Thickeners ... 294
Forteo ... 117
Fortijuice (Apple) ... 310
Fortijuice (Orange) ... 310
Fortijuice (Strawberry) ... 310
Fortini (Strawberry) ... 307
Fortini (Vanilla) ... 307
Fortini Multi Fibre (chocolate) ... 307
Fortini Multi Fibre (strawberry) ... 307
Fortini Multi Fibre (unflavoured) ... 307
Fortini Multi Fibre (vanilla) ... 307
Fortisip (Banana) ... 310
Fortisip (Chocolate) ... 310
Fortisip (Strawberry) ... 310
Fortisip (Vanilla) ... 310
Fortisip Multi Fibre (chocolate) ... 310

Fortisip Multi Fibre (strawberry) ... 310
Fortisip Multi Fibre (vanilla) ... 310
Fosamax ... 116
Fosamax Plus ... 116
Foscarnet sodium ... 109
Fosfomycin ... 97
Framycetin sulphate ... 275
Fresofol 1% MCT/LCT ... 124
Frusemide ... 51
Fucidin ... 98
Fucithalmic ... 275
Fulvestrant ... 176
Fulvestrant EVER Pharma ... 176
Fungilin ... 26
Furosemide [Frusemide] ... 51
Furosemide-AFT ... 51
Furosemide-Baxter ... 51
Fusidic acid
  Dermatological ... 68, 72
  Infections ... 98
  Sensory ... 275

- G -

GA Explore 5 ... 295
GA1 Anamix Infant ... 295
GA1 Anamix Junior ... 295
Gabapentin ... 132
Gacet ... 127
Gadobutrol ... 286
Gadoteric acid ... 286
Gadovist 1.0 ... 286
Gadoxetate disodium ... 286
Galsulfase ... 19
Galvumet ... 11
Galvus ... 11
Ganciclovir ... 109
Gardasil 9 ... 319
Gastrodenol ... 9
Gastrografin ... 285
Gazyva ... 216
Gefitinib ... 166
Gelatine, succinylated ... 44
Gelofusine ... 44
Gemcitabine Hydrochloride ... 154
Gemtuzumab ozogamicin ... 206
Gentamicin sulphate
  Infections ... 90
  Sensory ... 275
Gestrinone ... 83
Gilenya ... 142
Ginet ... 75
Glatiramer acetate ... 142
Glaucoma Preparations ... 279
Glecaprevir with pibrentasvir ... 108
Glibenclamide ... 11
Gliclazide ... 11
Gliolan ... 178
Glipizide ... 11
Glizide ... 11
Glucagen Hypokit ... 9
Glucagon hydrochloride ... 9
Glucerna Select ... 300
Glucose [Dextrose]
  Alimentary ... 9
  Blood ... 41
  Extemporaneously Compounded Preparations ... 290
Glucose with potassium chloride ... 41
Glucose with potassium chloride and sodium chloride ... 41
Glucose with sodium chloride ... 41
Glucose with sucrose and fructose ... 9
Glycerin with sodium saccharin ... 291
Glycerin with sucrose ... 291
Glycerol
  Alimentary ... 16
  Extemporaneously Compounded Preparations ... 291
Glycerol with paraffin ... 70
Glyceryl trinitrate
  Alimentary ... 7
  Cardiovascular ... 55
Glycine ... 287
Glycoprep Orange ... 15
Glycopyrronium ... 266
Glycopyrronium bromide ... 7
Glycopyrronium with indacaterol ... 266
Glycopyrronium-AFT ... 7
Glycosade ... 299
Glytactin Bettermilk ... 298
Gonadorelin ... 83
Goserelin ... 83
Granisetron ... 135
GTO Oil ... 299

- H -

Habitrol ... 149
Habitrol (Fruit) ... 149
Habitrol (Mint) ... 149
Haem arginate ... 19
Haemophilus influenzae type B vaccine ... 312
Haldol ... 138
Haldol Concentrate ... 138
Haloperidol ... 136
Haloperidol decanoate ... 138
Hartmann's solution ... 41
Harvoni ... 108
Havrix 1440 ... 318
Havrix Junior ... 318
Haylor Syrup ... 265
HCU Anamix Infant ... 296
HCU Anamix Junior ... 296
HCU Anamix Junior LQ ... 296
HCU Explore 5 ... 296
HCU Express 15 ... 296
HCU Lophlex LQ ... 296
Healon ... 278
Healon 5 ... 278
Healon GV Pro ... 278
healthE Calamine Aqueous ... 69
healthE Dimethicone 10% ... 69
healthE Dimethicone 4% Lotion ... 68
healthE Dimethicone 5% ... 69
healthE Glycerol BP Liquid ... 291
healthE Urea Cream ... 71
Hemlibra ... 33
Heparin sodium ... 37
Heparinised saline ... 37
Heparon Junior ... 302
Hepatitis A vaccine ... 318
Hepatitis B recombinant vaccine ... 318
Herzuma ... 239
Hikma ... 47
Hikma Acetylcysteine ... 282
Hiprex ... 97
Histaclear ... 265
Histamine acid phosphate ... 286
Holoxan ... 152
Hormone Replacement Therapy ... 82
HPV ... 319
Humalog Mix 25 ... 10
Humalog Mix 50 ... 10
Human Milk Fortifier ... 294
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV] ... 319
Humatin ... 90
Humira ... 194
HumiraPen ... 194
Hyaluronic acid
  Alimentary ... 26
  Sensory ... 278, 281
Hyaluronic acid with lidocaine [lignocaine] ... 26
Hyaluronidase ... 118
Hydralazine hydrochloride ... 56
Hydralock ... 69
Hydralyte - Lemonade ... 43
Hydrapres ... 56
Hydrocortisone
  Dermatological ... 71
  Extemporaneously Compounded Preparations ... 291
  Hormone Preparations ... 81
Hydrocortisone acetate ... 6

Hydrocortisone acetate with pramoxine hydrochloride 6
Hydrocortisone and paraffin liquid and lanolin 71
Hydrocortisone butyrate 72–73
Hydrocortisone with miconazole 72
Hydrocortisone with natamycin and neomycin 72
Hydrogen peroxide 68
Hydroxocobalamin
Alimentary 28
Various 282
Hydroxocobalamin Panpharma 28
hydroxycarbamide 155
Hydroxychloroquine sulphate 116
Hydroxyurea [hydroxycarbamide] 155
Hygroton 52
Hylo-Fresh 281
Hyoscine butylbromide 7
Hyoscine Butylbromide (Adiramedica) 7
Hyoscine hydrobromide 135
Hyperuricaemia and Antigout 118
HypoPak Glucose 9
Hypromellose 278, 280
Hypromellose with dextran 280

- I -

Ibiamox 94
Ibrutinib 155
Ibuprofen 120
Ibuprofen SR BNM 120
Icatibant 264
Icatibant Lupin 264
Idarubicin hydrochloride 153
Idarucizumab 34
Idursulfase 19
Ifosfamide 152
Ilomedin 65
Iloprost 65
Imaging Agents 178
Imatinib mesilate 166
Imatinib-Rex 166
Imbruvica 155
Imfinzi 246
Imipenem with cilastatin 90
Imipenem+Cilastatin RBX 90
Imipramine Crescent 130
Imipramine hydrochloride 130
Imiquimod 74
Immune Modulators 112
Immunosuppressants 179
Impact Advanced Recovery 308
Incruse Ellipta 266
Indacaterol 270
Indapamide 52
Indigo carmine 287
Indinavir 107
Indocyanine green 287
Indometacin [Indomethacin] 120
Indomethacin 120
Infanrix IPV 311
Infanrix-hexa 311
Infatrini 305
Infliximab 207
Influenza vaccine 319
Influvac Tetra (2026 formulation) 319
Inhaled Corticosteroids 270
Inlyta 162
Inotuzumab ozogamicin 214
Inresa 25
Inspra 51
Instillagel Lido 125
Insulin aspart 10
Insulin aspart with insulin aspart protamine 10
Insulin degludec with insulin aspart 10
Insulin glargine 10
Insulin glulisine 10
Insulin isophane 10
Insulin lispro 10
Insulin lispro with insulin lispro protamine 10
Insulin neutral 10
Insulin neutral with insulin isophane 10
Intelence 105
Interferon alfa-2b 112
Interferon beta-1-alpha 142
Interferon beta-1-beta 142
Interferon gamma 112
Intra-uterine device 76
Invanz 90
Invega Sustenna 139
Invega Trinza 139
Inza 121
Iodine 88
Iodine with ethanol 284
Iodised oil 285
Iodixanol 285
Iohexol 285
Iopidine 280
Ioscan 285
Ipca-Allopurinol 118
Ipca-Bisoprolol 48
Ipca-Ciprofloxacin 95
Ipca-Donepezil 148
Ipca-Enalapril 45
Ipca-Escitalopram 131
IPCA-Frusemide 51
Ipca-Hydroxychloroquine 116
IPCA-Metoprolol 49
IPCA-Propranolol 49
Ipilimumab 247
IPOL 321
Ipratropium bromide 265–266
Ipratropium Viatris 266
Iressa 166
Irinotecan hydrochloride 156
Iron (as ferric carboxymaltose) 24
Iron (as sucrose) 24
Iron polymaltose 24
Irrigation Solutions 287
Isentress 108
Isentress HD 108
Ismo 20 55
Ismo 40 Retard 55
Isoflurane 123
Isoleucine50 299
Isoniazid 102
Isoniazid Teva 102
Isoniazid with rifampicin 102
Isoprenaline [Isoproterenol] 55
Isopropyl alcohol 284
Isoproterenol 55
Isoptin 50
Isoptin SR 50
Isopto Carpine 279
Isosorbide mononitrate 55
Isotretinoin 69
Ispaghula (psyllium) husk 15
Isradipine 50
Itch-Soothe 69
Itraconazole 99
Itraconazole Cresent 99
Itrazole 99
Ivabradine 48
Ivacaftor 272
Ivermectin 103

- J -

Jadelle 76
Jakavi 171
Jardiamet 13
Jardiance 13
Jaydess 76
Jevity HiCal RTH 309
Jevity RTH 309
Jinarc 52

- K -

Kadcyla 241
Kalydeco 272
Kenacomb 276
Kenacort-A 10 81
Kenacort-A 40 81
Kenalog in Orabase 26
Ketalar 124

Ketamine 124
Ketocal 3:1 (Unflavoured) 306
Ketocal 4:1 (Unflavoured) 306
Ketocal 4:1 (Vanilla) 306
Ketoconazole
Dermatological 68
Infections 98
Ketoprofen 120
Ketorolac trometamol 277
KetoSens 323
Ketostix 323
Keytruda 250
Kindergen 307
Kisqali 170
Klacid 93
Klacid IV 93
Klaricid 93
Kogenate FS 35
Konakion MM 35
Konakion MM Paediatric 35
Konsyl-D 15
Kuvan 21

- L -

L-ornithine L-aspartate 9
Labetalol 49
Lacosamide 132
Lactose 291
Lactulose 16
Laevolac 16
Lamictal 132
Lamivudine 106, 108
Lamivudine Viatris 106
Lamivudine/Zidovudine Viatris 106
Lamotrigine 132
Lanoxin 48
Lanoxin PG 48
Lanreotide 178
Lansoprazole 8
Lantus 10
Lantus SoloStar 10
Lanzol Relief 8
Lapatinib 166
Largactil 136
Laronidase 19
Lasix 51
Latanoprost 279
Latanoprost with timolol 279
Lax-Suppositories 16
Lax-suppositories Glycerol 16
Laxatives 15
Ledipasvir with sofosbuvir 108
Leflunomide 116
Lenalidomide (Viatris) 156
Lenalidomide Viatris 156
Lenvatinib 166
Lenvima 166
Letermovir 109
Letrole 177
Letrozole 177
Leucine100 300
Leukotriene Receptor
Antagonists 270
Leuprorelin acetate 84
Leustatin 153
Levetiracetam 132
Levetiracetam-AFT 132
Levocabastine 277
Levocarnitine 20
Levodopa with benserazide 123
Levodopa with carbidopa 123
Levodopa with carbidopa and
entacapone 123
Levomepromazine 136
Levomepromazine
hydrochloride 136
Levonorgestrel 76
Levonorgestrel -1 (Lupin) 76
Levonorgestrel BNM 76
Levosimendan 55
Levothyroxine 88
Lidocaine [Lignocaine] 125
Lidocaine [Lignocaine]
hydrochloride 125
Lidocaine [Lignocaine] hydrochloride
with adrenaline 125
Lidocaine [Lignocaine] hydrochloride
with adrenaline and tetracaine
hydrochloride 125
Lidocaine [Lignocaine] hydrochloride
with phenylephrine
hydrochloride 126
Lidocaine [Lignocaine] with
prilocaine 126
Lidocaine-Baxter 125
lignocaine
Alimentary 26
Nervous 125–126
Lincomycin 97
Linezolid 97
Linezolid Kabi 97
Lioresal Intrathecal 119
Liothyronine sodium 88
Lipid-Modifying Agents 53
Lipiodol Ultra Fluid 285
Liquigen 293
Liraglutide 12
Lisdexamfetamine dimesilate 146
Lisinopril 45
Lissamine green 277
Lithium carbonate 137
LMX4 125
Lo-Oralcon 20 ED 75
Local Preparations for Anal and
Rectal Disorders 7
Locasol 305
Locoid 72–73
Locoid Crelo 72
Locoid Lipocream 72
Lodoxamide 277
Logem 132
Lomide 277
Lomustine 152
Long-Acting Beta-Adrenoceptor
Agonists 270
Loniten 56
Loperamide hydrochloride 5
Lopinavir with ritonavir 107
Lopinavir/Ritonavir Mylan 107
Lorafix 265
Loratadine 265
Lorazepam 131, 140
Lormetazepam 143
Lorstat 53
Losartan & Hydrochlorothiazide
(Ipca) 46
Losartan Actavis 46
Losartan potassium 46
Losartan potassium with
hydrochlorothiazide 46
Lovir 109
Loxamine 131
Lucrin Depot 1-month 84
Lucrin Depot 3-month 84
LumaCina 287
Lumigan 279
Lynparza 157
Lyrica 133
Lysine acetylsalicylate [Lysine
aspirin] 38
Lysine aspirin 38

- M -

m-Eslon 128
Mabthera 222
Macrobid 97
Macrogol 3350 with ascorbic acid,
potassium chloride, sodium
chloride and citric acid with
magnesium carbonate hydrate
and sodium picosulfate 15
Macrogol 3350 with potassium
chloride and sodium chloride 15
Macrogol 3350 with potassium
chloride and sodium chloride with/
without sodium sulfate, sodium
ascorbate, ascorbic acid 15
Macrogol 3350 with potassium
chloride, sodium bicarbonate and
sodium chloride 16

Madopar 125 ... 123
Madopar 250 ... 123
Madopar 62.5 ... 123
Madopar HBS ... 123
Madopar Rapid ... 123
Mafenide acetate ... 68
Magnesium amino acid chelate ... 24
Magnesium chloride ... 24
Magnesium hydroxide
  Alimentary ... 24
  Extemporaneously Compounded Preparations ... 291
Magnesium oxide ... 25
Magnesium oxide with magnesium aspartate, magnesium amino acid chelate and magnesium citrate ... 25
Magnesium sulphate ... 25
Magnevist ... 286
Malarone ... 104
Malarone Junior ... 104
Malathion [Maldison] ... 69
Maldison ... 69
Mannitol
  Cardiovascular ... 51
  Various ... 286
Mantoux ... 322
Maprotiline hydrochloride ... 130
Marcain ... 124
Marcain Heavy ... 125
Marcain Isobaric ... 124
Marcain with Adrenaline ... 124
Marevan ... 37
Marine Blue Lotion SPF 50+ ... 74
Mask for spacer device ... 323
Maviret ... 108
Maxidex ... 276
Maxitrol ... 275
MCT Oil ... 293
Measles, mumps and rubella vaccine ... 320
Mebendazole ... 103
Mebeverine hydrochloride ... 7
Medac ... 152
Medrol ... 81
Medroxyprogesterone ... 83
Medroxyprogesterone acetate
  Genito-Urinary ... 76
  Hormone Preparations ... 82
Mefenamic acid ... 121
Mefloquine ... 104
Meglumine gadopentetate ... 286
Meglumine iotroxate ... 286
Mekinist ... 173
Melatonin ... 143
Melpha ... 152
Melphalan ... 152
Meningococcal (A, C, Y and W-135) conjugate vaccine ... 312
Meningococcal B multicomponent vaccine ... 313
MenQuadfi ... 312
Menthol ... 291
Mepivacaine hydrochloride ... 126
Mepivacaine hydrochloride with adrenaline ... 126
Mepolizumab ... 215
Mercaptopurine ... 154
Meropenem ... 91
Meropenem-AFT ... 91
Mesalazine ... 6
Mesna ... 175
Mestinon ... 116
Metabolic Disorder Agents ... 17
Metabolic Products ... 295
Metaraminol ... 56
Metformin hydrochloride ... 11
Metformin Viatris ... 11
Methacholine chloride ... 287
Methadone BNM ... 128
Methadone hydrochloride
  Extemporaneously Compounded Preparations ... 291
  Nervous ... 128
Methenamine (Hexamine) hippurate ... 97
Methohexital sodium ... 124
Methopt ... 280
Methotrexate ... 154
Methotrexate DBL Onco-Vial ... 154
Methotrexate Ebewe ... 154
Methotrexate Sandoz ... 154
Methoxsalen [8-methoxypsoralen] ... 73
Methoxyflurane ... 126
Methyl aminolevulinate hydrochloride ... 74
Methyl hydroxybenzoate ... 291
Methylcellulose ... 291
Methylcellulose with glycerin and sodium saccharin ... 291
Methylcellulose with glycerin and sucrose ... 291
Methyldopa ... 51
Methyldopa Viatris ... 51
Methylene blue ... 287
Methylnaltrexone bromide ... 16
Methylphenidate ER - Teva ... 147
Methylphenidate hydrochloride ... 147
Methylphenidate Sandoz XR ... 147
Methylprednisolone (as sodium succinate) ... 81
Methylprednisolone aceponate ... 72
Methylprednisolone acetate ... 81
Methylthioninium chloride [Methylene blue] ... 287
Methylxanthines ... 271
Metoclopramide Actavis 10 ... 135
Metoclopramide hydrochloride ... 135
Metoclopramide hydrochloride with paracetamol ... 134
Metolazone ... 52
Metoprolol IV Mylan ... 49
Metoprolol IV Viatris ... 49
Metoprolol succinate ... 49
Metoprolol tartrate ... 49
Metronidamed ... 104
Metronidazole
  Dermatological ... 68
  Infections ... 104
Metyrapone ... 83
Mexiletine hydrochloride ... 48
Miacalcic ... 79
Mianserin hydrochloride ... 130
Micolette ... 16
Miconazole ... 26
Miconazole nitrate
  Dermatological ... 68
  Genito-Urinary ... 75
Micreme ... 75
Micreme H ... 72
Microlut ... 76
Midazolam ... 144
Midazolam-Baxter ... 144
Midazolam-Pfizer ... 144
Midodrine ... 48
Midostaurin ... 168
Mifepristone ... 76
Milrinone ... 56
Milrinone-Baxter ... 56
Minerals ... 23
Mini-Wright AFS Low Range ... 323
Mini-Wright Standard ... 323
Minidiab ... 11
Minims Prednisolone ... 277
Minirin ... 89
Minirin Melt ... 89
Minocycline ... 96
Minoxidil ... 56
Mirena ... 76
Miro-Amoxicillin ... 94
Mirtazapine ... 130
Misoprostol ... 7
Mitomycin C ... 153
Mitozantrone ... 153
Mitozantrone Ebewe ... 153
Mivacurium chloride ... 119
Mixed salt solution for eye

irrigation ... 278
MMA/PA Anamix Infant ... 299
MMA/PA Anamix Junior ... 299
MMA/PA Explore 5 ... 299
MMA/PA Express 15 ... 299
Moclobemide ... 130
Modafinil ... 148
Modafinil Max Health ... 148
Molaxole ... 16
Mometasone furoate ... 72
Monofer ... 24
Monogen ... 301
Monosodium glutamate with sodium aspartate ... 289
Monosodium l-aspartate ... 289
Montelukast ... 270
Montelukast Viatris ... 270
Moroctocog alfa [Recombinant factor VIII] ... 35
Morphine hydrochloride ... 128
Morphine sulphate ... 128
Morphine tartrate ... 128
Motetis ... 122
Mouth and Throat ... 25
Movapo ... 122
Movicol ... 16
Moxifloxacin ... 95
Moxifloxacin Kabi ... 95
Mozobil ... 39
MSUD Anamix Infant ... 296
MSUD Anamix Junior ... 296
MSUD Anamix Junior LQ ... 296
MSUD Explore 5 ... 296
MSUD Express 15 ... 296
MSUD Lophlex LQ 20 ... 296
MSUD Maxamum ... 296
Mucolytics and Expectorants ... 271
Multiple Sclerosis Treatments ... 141
Multivitamin and mineral supplement ... 26
Multivitamin renal ... 27
Multivitamins ... 27
Mupirocin ... 68
Muscle Relaxants and Related Agents ... 119
Mvite ... 27
Myambutol ... 102
Mycobutin ... 103
MycoNail ... 68
Mycophenolate mofetil ... 258
Mydriacyl ... 280
Mydriatics and Cycloplegics ... 280
Mylan (24 hr release) ... 50
Mylan Clomiphen ... 83
Mylan Italy (24 hr release) ... 50
Myleran ... 152
Myloc CR ... 49
Mylotarg ... 206
Myozyme ... 17
Mytolac ... 178

- N -

Nadolol ... 49
Nadolol BNM ... 49
Naglazyme ... 19
Naloxone hydrochloride ... 282
Naltraccord ... 149
Naltrexone hydrochloride ... 149
Naphazoline hydrochloride ... 277
Naprosyn SR 1000 ... 121
Naprosyn SR 750 ... 121
Naproxen ... 121
Natalizumab ... 142
Natamycin ... 275
Natulan ... 159
Nausafix ... 135
Nausicalm ... 135
Nefopam hydrochloride ... 127
Neo-Mercazole ... 88
Neocate Gold (Unflavoured) ... 303
Neocate Junior Unflavoured ... 303
Neocate Junior Vanilla ... 303
Neocate SYNEO ... 303
Neoral ... 179
Neostigmine metilsulfate ... 116
Neostigmine metilsulfate with glycopyrronium bromide ... 116
Neosynephrine HCL ... 56
Nepro HP (strawberry) ... 307
Nepro HP (vanilla) ... 307
Neulumex ... 285
Neupogen ... 40
NeuroTabs ... 24
Nevirapine ... 105
Nevirapine Viatris ... 105
Nicardipine hydrochloride ... 50
Nicorandil ... 57
Nicotine ... 149
Nifedipine ... 50
Nifuran ... 97
Nilotinib ... 168
Nilstat
Alimentary ... 26
Genito-Urinary ... 75
Infections ... 99
Nimenrix ... 313
Nimodipine ... 50
Nimotop ... 50
Nintedanib ... 268
Niraparib ... 156
Nirmatrelvir with ritonavir ... 112
Nitazoxanide ... 104
Nitisinone ... 20
Nitisinone LogixX Pharma ... 20
Nitrates ... 55
Nitroderm TTS 10 ... 55
Nitroderm TTS 5 ... 55
Nitrofurantoin ... 97
Nitrofurantoin Clinect ... 97
Nitrolingual Pump Spray ... 55
Nivestim ... 40
Nivolumab ... 247
Nodia ... 5
Noflam 250 ... 121
Noflam 500 ... 121
Non-Steroidal Anti-Inflammatory Drugs ... 120
Nonacog gamma, [Recombinant factor IX] ... 35
Noradrenaline ... 56
Norethisterone
Genito-Urinary ... 76
Hormone Preparations ... 83
Norethisterone with mestranol ... 75
Norflex ... 119
Norfloxacin ... 96
Noriday ... 76
Noriday 28 ... 76
Normison ... 144
Nortriptyline hydrochloride ... 130
Norvir ... 107
Noumed Dexamfetamine ... 146
Noumed Isoniazid ... 102
Noumed Paracetamol ... 127
Noumed Pethidine ... 129
Noumed Phenobarbitone ... 132
Novadoz ... 152
Novasource Renal (Vanilla) ... 308
Novatretin ... 72
NovoMix 30 FlexPen ... 10
NovoRapid FlexPen ... 10
NovoSeven RT ... 34
Nozinan ... 136
Nucala ... 215
Nuelin ... 271
Nuelin-SR ... 271
Nupentin ... 132
Nusinersen ... 144
Nutilis ... 295
Nutren Diabetes (vanilla) ... 300
Nutrini Energy Multi Fibre ... 307
Nutrini Energy RTH ... 307
Nutrini Low Energy Multi Fibre RTH ... 307
Nutrini Peptisorb Energy ... 304
Nutrini RTH ... 307
Nutrison 800 Complete Multi Fibre ... 309
Nutrison Advanced Peptisorb ... 301

Nutrison Concentrated....302
Nutrison Energy....309
Nutrison Energy Multi Fibre....309
Nutrison Multi Fibre....309
Nutrison Protein Intense....303
Nutrison Protein Plus....302
Nutrison Protein Plus Multi Fibre....303
Nutrison RTH....309
Nyefax Retard....50
Nystatin
Alimentary....26
Dermatological....68
Genito-Urinary....75
Infections....99

- O -

Obinutuzumab....216
Obstetric Preparations....76
Ocrelizumab....142
Ocrevus....142
Ocrevus SC....142
Octasa....6
Octocog alfa [Recombinant factor VIII] (Advate)....35
Octocog alfa [Recombinant factor VIII] (Kogenate FS)....35
Octreotide....177
Octreotide Long-Acting....178
Ocular Lubricants....280
Oestradiol....82–83
Oestradiol valerate....82
Oestradiol with norethisterone acetate....82
Oestriol
Genito-Urinary....77
Hormone Preparations....83
Oestrogens....77
Oestrogens (conjugated equine)....82
Oestrogens with medroxyprogesterone acetate....82
Ofev....268
Oil in water emulsion....70
Oily phenol [Phenol oily]....7
Olanzapina Mylan....137
Olanzapina Mylan Pharma....137
Olanzapine....137–138
Olanzapine ODT Viatris....137
Olaparib....157
Olive oil....291
Olopatadine....277
Olopatadine Teva....277
Olsalazine....7
Omalizumab....217
Omeprazole....8
Omeprazole actavis 10....8
Omeprazole actavis 20....8
Omeprazole actavis 40....8
Omeprazole Teva....8
Omezol IV....8
Omnipaque....285
Omnitrope....84
Omnitrope AU....84
Onbrez Breezhaler....270
Oncaspar LYO....158
OncoTICE....258
Ondansetron....135
Ondansetron-AFT....135
One-Alpha....28
Opdivo....247
Optional Pharmaceuticals....323
Ora-Blend....291
Ora-Blend SF....291
Ora-Plus....291
Ora-Sweet....291
Ora-Sweet SF....291
Oralcon 30 ED....75
Oramorph....128
Oramorph CDC S29....128
Oratane....69
Ornidazole....104
Orphenadrine citrate....119
Oruvail SR....120
Oseltamivir....112
Osimertinib....168
Osmolite RTH....309
Other Cardiac Agents....55
Other Endocrine Agents....82
Other Oestrogen Preparations....83
Other Otological Preparations....281
Other Progestogen Preparations....83
Other Skin Preparations....73
Ovestin
Genito-Urinary....77
Hormone Preparations....83
Oxaliplatin....162
Oxandrolone....79
Oxazepam....140
Oxpentifylline....57
Oxybuprocaine hydrochloride....278
Oxybutynin....78
Oxycodone Amneal....129
Oxycodone hydrochloride....129
Oxycodone Lucis S29....129
Oxycodone Sandoz....129
Oxymetazoline hydrochloride....269
Oxytocin....76
Oxytocin BNM....76
Oxytocin with ergometrine maleate....77
Ozurdex....276

- P -

Pacifen....119
Pacimol....127
Paclitaxel....174
Padagis....74
Paediatric Seravit....27
Palbociclib....169
Palbociclib Pfizer....169
Paliperidone....139
Paliperidone palmitate....139
Palivizumab....218
Pamidronate disodium....116
Pamisol....116
Pamol....127
Pancreatic enzyme....14
Pancuronium bromide....119
Pantoprazole....8
Panzop Relief....8
Papaverine hydrochloride....57
Paper wasp venom....264
Para-aminosalicylic Acid....102
Paracetamol....127
Paracetamol (Ethics)....127
Paracetamol Kabi....127
Paracetamol with codeine....129
Paraffin
Alimentary....15
Dermatological....71
Extemporaneously Compounded Preparations....291
Paraffin liquid with soft white paraffin....281
Paraffin liquid with wool fat....281
Paraffin with wool fat....71
Paraldehyde....131
Parecoxib....121
Parecoxib Juno....121
Paromomycin....90
Paroxetine....131
Paser....102
Patent blue V....287
Paxam....140
Paxlovid....112
Pazopanib....170
Pazopanib ADVZ....170
Pazopanib Teva....170
Peak flow meter....323
Peanut oil....290
Pedialyte....43
Pediasure (chocolate)....307
Pediasure (strawberry)....307
Pediasure (vanilla)....307
Pediasure Plus....307
Pediasure RTH....307
Pegaspargase....158
Pegasys....113

Pegasys (S29) ... 113
Pegfilgrastim ... 40
Pegylated interferon alfa-2a ... 113
Pembrolizumab ... 250
Pemetrexed ... 154
Pemetrexed-AFT ... 154
Penicillamine ... 116
Penicillin G ... 94
Penicillin V ... 95
Pentacarinat ... 104
Pentagastrin ... 83
Pentamidine isethionate ... 104
Pentasa ... 6
Penthrox ... 126
Pentostatin [Deoxycoformycin] ... 158
Pentoxifylline [Oxpentifylline] ... 57
Peptamen OS 1.0 (Vanilla) ... 301
Pepti-Junior ... 305
Perflutren ... 286
Perhexiline maleate ... 50
Pericyazine ... 137
Perindopril ... 45
Periset ... 135
Periset ODT ... 135
Perjeta ... 220
Permethrin ... 69
Perrigo ... 74
Pertuzumab ... 220
Pertuzumab with trastuzumab ... 220
Peteha ... 103
Pethidine hydrochloride ... 129
Pevaryl ... 68
Pexsig ... 50
Pfizer Exemestane ... 177
Pharmacor Varenicline ... 150
Pheburane ... 21
Phenasen ... 154
Phenelzine sulphate ... 130
Phenindione ... 37
Phenobarbitone ... 132, 144
Phenobarbitone sodium ... 291
Phenol
  Extemporaneously Compounded Preparations ... 291
  Various ... 288
Phenol oily ... 7
Phenol with ioxaglic acid ... 288
Phenothrin ... 69
Phenoxybenzamine hydrochloride ... 47
Phenoxymethylpenicillin [Penicillin V] ... 95
Phentolamine mesylate ... 47
Phenylalanine50 ... 300
Phenylephrine hydrochloride
  Cardiovascular ... 56
  Sensory ... 280
Phenytoin ... 132
Phenytoin sodium ... 131–132
Phesgo ... 220
Phlexy 10 ... 297
Phosphorus ... 43
Phytomenadione ... 35
Picibanil ... 259
Pilocarpine hydrochloride ... 279
Pilocarpine nitrate
  Extemporaneously Compounded Preparations ... 291
  Sensory ... 279
Pimafucort ... 72
Pimecrolimus ... 73
Pine tar with trolamine laurilsulfate and fluorescein ... 73
Pinetarsol ... 73
Pioglitazone ... 11
Piperacillin with tazobactam ... 95
Pipothiazine palmitate ... 140
PipTaz-AFT ... 95
Pirfenidone ... 268
Pirfenidone Sandoz ... 268
Pituitary and Hypothalamic Hormones and Analogues ... 83
Pivmecillinam ... 98
Pizotifen ... 134
PKU Anamix Infant ... 297
PKU Anamix Junior ... 297
PKU Anamix Junior LQ (Berry) ... 297
PKU Anamix Junior LQ (Orange) ... 297
PKU Build 10 ... 298
PKU Build 20 Chocolate ... 298
PKU Build 20 Raspberry Lemonade ... 298
PKU Build 20 Smooth ... 298
PKU Build 20 Vanilla ... 298
PKU Explore 10 ... 297
PKU Explore 5 ... 297
PKU Express 20 ... 297
PKU First Spoon ... 297
PKU Glytactin RTD 15 ... 298
PKU Glytactin RTD 15 Lite ... 298
PKU GMPro LQ ... 298
PKU GMPro Mix-In ... 298
PKU GMPro Ultra Lemonade ... 298
PKU GMPro Ultra Vanilla ... 298
PKU Lophlex LQ 10 ... 297
PKU Lophlex LQ 20 ... 297–298
PKU Lophlex Powder ... 297
PKU Lophlex Sensations 20 (berries) ... 298
PKU Restore Powder ... 297
PKU sphere20 Banana ... 298
PKU sphere20 Chocolate ... 298
PKU sphere20 Lemon ... 298
PKU sphere20 Red Berry ... 298
PKU sphere20 Vanilla ... 298
PKU Start ... 297
Plasma-Lyte 148 ... 41
Plasma-Lyte 148 & 5% Glucose ... 41
Plendil ER ... 49
Plenvu ... 15
Plerixafor ... 39
Plerixafor-AFT ... 39
Pneumococcal (PCV13) conjugate vaccine ... 314
Pneumococcal (PPV23) polysaccharide vaccine ... 315
Pneumovax 23 ... 315
Podophyllotoxin ... 73
Polidocanol ... 33
Poliomyelitis vaccine ... 321
Poloxamer ... 15
Poly-Tears ... 280
Poly-Visc ... 281
Polycal ... 293
Polyethylene glycol 400 and propylene glycol ... 281
Polyflush ... 42
Polyhexamethylene biguanide ... 291
Polyvinyl alcohol with povidone ... 281
Pomalidomide ... 158
Pomolide ... 158
Poractant alfa ... 273
Posaconazole ... 99
Posaconazole Juno ... 99
Potassium chloride ... 41, 43
Potassium chloride with sodium chloride ... 42
Potassium citrate ... 77
Potassium dihydrogen phosphate ... 42
Potassium iodate
  Alimentary ... 24
  Hormone Preparations ... 88
Potassium iodate with iodine ... 24
Potassium perchlorate ... 88
Potassium permanganate ... 73
Povidone K30 ... 291
Povidone-iodine ... 284
Povidone-iodine with ethanol ... 284
Pradaxa ... 36
Pralidoxime chloride ... 282
Pralidoxime iodide ... 282
Pramipexole hydrochloride ... 123
Pravastatin ... 53
Praxbind ... 34
Praziquantel ... 103
Prazosin ... 47

Prazosin Mylan 47
Pred Forte 277
Prednisolone 81
Prednisolone acetate 277
Prednisolone hexanoate with cinchocaine hydrochoride 7
Prednisolone sodium phosphate 277
Prednisolone- AFT 277
Prednisone 81
Prednisone Clinect 81
Pregabalin 133
Pregabalin Pfizer 133
Pregnancy test - hCG urine 323
preOp 308
Presolol 49
Prevenar 13 314
Prevymis 109
Priadel 137
Prilocaine hydrochloride 126
Prilocaine hydrochloride with felypressin 126
Primaquine 104
Primidone 133
Primolut N 83
Primovist 286
Priorix 320
Probenecid 118
Procaine penicillin 95
Procarbazine hydrochloride 159
Prochlorperazine 135
Proctosedyl 7
Procyclidine hydrochloride 122
Progesterone 77
Proglicem 9
Proglycem 9
Progynova 82
Prolia 117
Promethazine hydrochloride 265
Propafenone hydrochloride 48
Propamidine isethionate 275
Propofol 124
Propranolol 49
Propylthiouracil 88
Prostin E2 76
Prostin VR 56
Protamine sulphate 37
Protifar 294
Protionamide 103
Protirelin 89
Proveblue 287
Provera 82
Provera HD 83
Proxymetacaine hydrochloride 278
Pseudoephedrine hydrochloride 269
Psoriasis and Eczema Preparations 72
PTU 88
Pulmonary Surfactants 273
Pulmozyme 271
Puri-nethol 154
Pyrazinamide 103
Pyridostigmine bromide 116
Pyridoxal-5-phosphate 20
Pyridoxine hydrochloride 28
Pyridoxine multichem 28
Pyrimethamine 104

- Q -

Quetapel 137
Quetiapine 137
Quetiapine Viatris 137
Quinapril 45
Quinine dihydrochloride 104
Qvar 270

- R -

RA-Morph 128
Rabies vaccine 321
Raltegravir potassium 108
Ramipex 123
Ramipril 45
Ramipril Mylan Generics 45
Ramipril Viatris 45
Ranbaxy-Cefaclor 91
Ranibizumab 221
Ranitidine 8
Rapamune 259
Rasagiline 123
Rasburicase 119
Readi-CAT 2 285
Reandron 1000 79
Recombinant factor IX 34–35
Recombinant factor VIIa 34
Recombinant factor VIII 35
Rectogesic 7
Red back spider antivenom 283
Redipred 81
Relenza Rotadisk 112
Relistor 16
Relonchem 270
Remdesivir 112
Remicade 207
Remifentanil 129
Remifentanil-AFT 129
Renilon 7.5 (apricot) 308
Renilon 7.5 (caramel) 308
Resonium A 43
Resource Beneprotein 294
Respiratory Stimulants 273
Retinol 27
Retinol Palmitate 281
ReTrieve 69
Retrovir 106
Retrovir IV 106
Revolade 32
Ribociclib 170
Riboflavin 20
Riboflavin 5-phosphate 279
Ricit 77
Ricovir 108
Rifabutin 103
Rifadin 103
Rifamazid 102
Rifampicin 103
Rifaximin 9
Rifinah 102
Rilutek 122
Riluzole 122
Ringer's solution 42
Rinvoq 260
Riodine 284
Risdiplam 145
Risedronate Sandoz 116
Risedronate sodium 116
Risperdal 137
Risperdal Consta 140
Risperidone 137, 140
Risperidone (Teva) 137
Risperon 137
Ritalin 147
Ritalin LA 147
Ritonavir 107
Rituximab (mabthera) 222
Rituximab (riximyo) 223
Rivaroxaban 37
Rivastigmine 148
Rivastigmine Patch BNM 10 148
Rivastigmine Patch BNM 5 148
Riximyo 223
RIXUBIS 35
Rizamelt 134
Rizatriptan 134
Robinul 7
Rocuronium bromide 119
Roma 43
Ropin 123
Ropinirole hydrochloride 123
Ropivacaine hydrochloride 126
Ropivacaine Kabi 126
Rose bengal sodium 277
Rosemont 129
Rosuvastatin 53
Rosuvastatin - Sandoz 53
Rosuvastatin Viatris 53
Rotarix 321
Rotavirus oral vaccine 321
Roxithromycin 94
Rozlytrek 165

Rubifen 147
Rubifen SR 147
Rugby 25
Rurioctocog alfa pegol [Recombinant factor VIII] 35
Ruxolitinib 171
Rydapt 168
Ryzodeg 70/30 Penfill 10

- S -

S26 LBW Gold RTF 306
Sacubitril with valsartan 46
SalAir 269
Salazopyrin 7
Salazopyrin EN 7
Salbutamol 269
Salbutamol with ipratropium bromide 266
Salicylic acid 291
Salmeterol 270
Salmonella typhi vaccine 316
Sandimmun 179
Sandomigran 134
Sandostatin LAR 178
Sapropterin Dihydrochloride 21
Scalp Preparations 73
Scandonest 3% 126
Scheriproct 7
Sclerosing Agents 274
Scopolamine Transdermal System Viatris 135
Sebizole 68
Secretin pentahydrochloride 287
Secukinumab 232
Sedatives and Hypnotics 143
Seebri Breezhaler 266
Selegiline hydrochloride 123
Selenium 25
Sennosides 16
Serc 135
Serenace 136
Seretide 271
Seretide Accuhaler 271
Serevent 270
Serevent Accuhaler 270
Sertraline 131
Setrona 131
Sevoflurane 124
Sevredol 128
Shingles vaccine 322
Shingrix 322
SII-Onco-BCG 258
Sildenafil 62
Siltuximab 234
Silver diamine fluoride 26
Silver nitrate
Dermatological 73
Extemporaneously Compounded Preparations 291
Simdax 55
Simeticone 5
Simulect 202
Simvastatin 54
Simvastatin Mylan 54
Simvastatin Viatris 54
Sincalide 287
Sinemet 123
Sinemet CR 123
Sintetica Baclofen Intrathecal 119
Sirolimus 259
Sirturo 102
Siterone 79
Slow-Lopresor 49
Snake antivenom 283
Sodibic 43
Sodium acetate 42
Sodium acid phosphate 43
Sodium alginate with magnesium alginate 5
Sodium alginate with sodium bicarbonate and calcium carbonate 5
Sodium aurothiomalate 116
Sodium benzoate 21
Sodium bicarbonate
Blood 42–43
Extemporaneously Compounded Preparations 291
Sodium calcium edetate 284
Sodium chloride
Blood 42–43
Respiratory 269, 273
Various 287
Sodium chloride with sodium bicarbonate 269
Sodium citrate
Alimentary 5
Extemporaneously Compounded Preparations 292
Sodium citrate with sodium chloride and potassium chloride 37
Sodium citrate with sodium lauryl sulphoacetate 16
Sodium citro-tartrate 78
Sodium cromoglicate
Alimentary 7
Respiratory 265
Sensory 277
Sodium dihydrogen phosphate [Sodium acid phosphate] 43
Sodium fluoride 24
Sodium fusidate [Fusidic acid]
Dermatological 68
Infections 98
Sensory 275
Sodium hyaluronate [Hyaluronic acid]
Alimentary 26
Sensory 278, 281
Sodium hyaluronate [Hyaluronic acid] with chondroitin sulphate 278
Sodium hydroxide 288
Sodium hypochlorite 285
Sodium metabisulfite 292
Sodium nitrite 282
Sodium nitroprusside
Cardiovascular 57
Optional Pharmaceuticals 323
Sodium phenylbutyrate 21
Sodium phosphate with phosphoric acid 16
Sodium picosulfate 17
Sodium polystyrene sulphonate 43
Sodium stibogluconate 104
Sodium tetradecyl sulphate 34
Sodium thiosulfate 282
Sodium valproate 133
Sodium with potassium 289
Solax 15
Solifenacin succinate 78
Solifenacin succinate Max Health 78
Solu-Cortef 81
Solu-Medrol 81
Solu-Medrol Act-O-Vial 81
Somatropin 84
Somatuline Autogel 178
Sotalol 49
Soya oil 282
Spacer device 323
Span-K 43
Spazmol 7
Specialised Formulas 300
Spinal Muscular Atrophy 144
Spinraza 144
Spiolto Respimat 266
Spiractin 52
Spiramycin 105
Spiriva 266
Spiriva Respimat 266
Spironolactone 52
Stalevo 123
Standard Feeds 308
Staphlex 95
Starch 292
Stavudine 106
Stelara 242
Sterculia with frangula 15
SteroClear 265
Stesolid 131

Stimulants / ADHD Treatments 145
Stiripentol 133
Streptomycin sulphate 90
Stromectol 103
Sucralfate 9
Sucrose 127
Sugammadex 119
Sugammadex BNM 119
Sulfadiazine silver 68
Sulfadiazine sodium 98
Sulfasalazine 7
Sulindac 121
Sulphacetamide sodium 275
Sulphur 292
Sulprix 136
Sumagran 134
Sumatriptan 134
Sunitinib 172
Sunitinib Rex 172
Sunscreen, proprietary 74
Suprane 123
Surgical Preparations 288
Sustagen Hospital Formula (Chocolate) 309
Sustagen Hospital Formula (Vanilla) 309
Suxamethonium chloride 119
Sylvant 234
Symbicort Turbuhaler 271
Symmetrel 122
Sympathomimetics 55
Synacthen 83
Synacthen Depot 83
Synagis 218
Synermox 94
Syntometrine 77
Syrup 292
Systane Unit Dose 281

- T -

Tacrolimus
Dermatological 73
Oncology 179
Tacrolimus Sandoz 179
Tafinlar 163
Tagitol V 285
Tagrisso 168
Talc 274
Taliglucerase alfa 22
Tambocor 48
Tambocor German 48
Tamoxifen citrate 177
Tamoxifen Sandoz 177
Tamsulosin hydrochloride 77
Tamsulosin-Rex 77
Tasigna 168
Tasmar 123
Taurine 23
TCu 380 Plus Normal 76
Tecentriq 245
Tecfidera 141
TEEVIR 106
Tegretol 131
Tegretol CR 131
Teicoplanin 98
Temaccord 159
Temazepam 144
Temozolomide 159
Temozolomide Taro 159
Tenecteplase 39
Tenofovir disoproxil 108
Tenofovir Disoproxil Emtricitabine Mylan 111
Tenofovir Disoproxil Emtricitabine Viatr 111
Tenofovir Disoproxil Viatris 108
Tenoxicam 121
Tensipine MR10 50
Tepadina 152
Terazosin 47
Terbinafine 102
Terbutaline 77
Terbutaline sulphate 269
Teriflunomide 142
Teriflunomide Sandoz 142
Teriparatide 117
Teriparatide - Teva 117
Terlipressin 89
Terlipressin Ever Pharma 89
Testogel 79
Testosterone 79
Testosterone cipionate 79
Testosterone esters 79
Testosterone undecanoate 79
Tetrabenazine 122
Tetracaine [Amethocaine] hydrochloride
Nervous 126
Sensory 278
Tetracosactide [Tetracosactrin] 83
Tetracosactrin 83
Tetracycline 96
Teva Lisinopril 45
Thalidomide 160
Thalomid 160
Theobroma oil 292
Theophylline 271
Thiamine hydrochloride 28
Thiamine multichem 28
Thiamine Sterop 28
Thioguanine 154
Thiopental [Thiopentone] sodium 124
Thiopentone 124
Thiotepa 152
Thiotepa-Reach 152
Thrombin 34
Thyroid and Antithyroid Preparations 88
Thyrotropin alfa 83
Ticagrelor 38
Ticagrelor Mylan 38
Ticagrelor Sandoz 38
Ticagrelor Sandoz S29 38
Ticarcillin with clavulanic acid 95
Ticlopidine 39
Tigecycline 96
Tilcotil 121
Tillomed 104
Timolol 279
Tiotropium bromide 266
Tiotropium bromide with olodaterol 266
Tivicay 108
TMP 98
Tobradex 275
Tobramycin
Infections 90
Sensory 275
Tobramycin (Viatris) 90
Tobramycin BNM 90
Tobrex 275
Tocilizumab 235
Tofranil 130
Tolcapone 123
Tolvaptan 52
Topamax 133
Topamine 26
Topicaine 125
Topical Products for Joint and Muscular Pain 121
Topiramate 133
Topiramate Actavis 133
Torbay 56
Tramadol hydrochloride 129
Tramal 100 129
Tramal 50 129
Tramal SR 100 129
Tramal SR 150 129
Tramal SR 200 129
Trametinib 173
Trandate 49
Tranexamic acid 34
Tranexamic-AFT 34
Tranylcypromine sulphate 130
Trastuzumab (Herzuma) 239
Trastuzumab deruxtecan 240
Trastuzumab emtansine 241
Travatan 280
Travoprost 280

Treatments for Dementia ... 148
Treatments for Substance Dependence ... 148
Trelegy Ellipta ... 267
Tretinoin
Dermatological ... 69
Oncology ... 160
Trexate ... 154
Tri-sodium citrate ... 292
Triamcinolone acetonide
Alimentary ... 26
Dermatological ... 72
Hormone Preparations ... 81
Triamcinolone acetonide with gramicidin, neomycin and nystatin ... 276
Triamcinolone acetonide with neomycin sulphate, gramicidin and nystatin ... 72
Triamcinolone hexacetonide ... 81
Triazolam ... 144
Trichloracetic acid ... 292
Trientine ... 23
Trientine Waymade ... 23
Trikafta ... 272
Trimethoprim ... 98
Trimethoprim with sulphamethoxazole [Co-trimoxazole] ... 98
Trisul ... 98
Trometamol ... 288
Tropicamide ... 280
Tropisetron ... 136
Trulicity ... 11
Tryzan ... 45
Tuberculin PPD [Mantoux] test ... 322
Tubersol ... 322
Two Cal HN ... 302
TYR Anamix Infant ... 299
TYR Anamix Junior ... 299
TYR Anamix Junior LQ ... 299
TYR Explore 5 ... 299
TYR Lophlex LQ 20 ... 299
TYR Sphere 20 ... 299
Tyrosine1000 ... 300
Tysabri ... 142

- U -

UK Cipla ... 269
UK Synacthen ... 83
Ultibro Breezhaler ... 266
Ultraproct ... 7
Umeclidinium ... 266
Umeclidinium with vilanterol ... 266
Univent ... 265–266
Upadacitinib ... 260
Ural ... 78
Urea
Dermatological ... 71
Extemporaneously Compounded Preparations ... 292
Urex Forte ... 51
UroFos ... 97
Urografin ... 285
Urokinase ... 39
Urologicals ... 77
Uromitexan ... 175
Ursodeoxycholic acid ... 14
Ursosan ... 14
Ustekinumab ... 242
Utrogestan ... 77

- V -

Vabysmo ... 206
Vaclovir ... 110
Valaciclovir ... 110
Valaciclovir Mylan ... 110
Valaciclovir Viatris ... 110
Valganciclovir ... 110
Valganciclovir Viatris ... 110
Valine50 ... 300
Valni Retard ... 50
Vancomycin ... 98
Vancomycin Viatris ... 98
Vanilla SilQ HD ... 285
Vanilla SilQ MD ... 285
Vanzacaftor with tezacaftor and deutivacaftor ... 273
Varenicline ... 150
Varibar - Honey ... 285
Varibar - Nectar ... 285
Varibar - Pudding ... 285
Varibar - Thin Liquid ... 285
Varicella vaccine [Chickenpox vaccine] ... 321
Varicella zoster vaccine [Shingles vaccine] ... 322
Varilrix ... 321
Vasodilators ... 56
Vasopressin ... 89
Vasopressin Agents ... 89
Vasorex ... 49
Vebulis ... 65
Vecure ... 119
Vecuronium bromide ... 119
Vedafil ... 62
Vedolizumab ... 243
Vegzelma ... 203
Veklury ... 112
Veletri ... 63
Venclexta ... 160
Venetoclax ... 160
Venlafaxine ... 130
Venofer ... 24
VENOX ... 264
Ventolin ... 269
Vepesid ... 155
Verapamil hydrochloride ... 50
Vermox ... 103
Versacloz ... 136
Vesanoid ... 160
Vexazone ... 11
Vfend ... 100
Victoza ... 12
Vigabatrin ... 133
Vigisom ... 143
Vildagliptin ... 11
Vildagliptin with metformin hydrochloride ... 11
Vimpat ... 132
Vinblastine sulphate ... 175
Vincristine sulphate ... 175
Vinorelbine ... 175
Vinorelbine Te Arai ... 175
Viral Vaccines ... 316
Viramune Suspension ... 105
ViruPOS ... 275
Viscoat ... 278
Visipaque ... 285
Vit.D3 ... 28
VitA-POS ... 281
Vital ... 301
Vitamin B complex ... 28
Vitamin B6 25 ... 28
Vitamins ... 26
Vivonex TEN ... 301
Voltaren ... 120
Voltaren D ... 120
Voltaren SR ... 120
Volumatic ... 323
Voriconazole ... 100
Vttack ... 100
Vytorin ... 54
Vyvanse ... 146

- W -

Warfarin sodium ... 37
Wart Preparations ... 73
Water
Blood ... 43
Various ... 288
White Soft Liquid Paraffin AFT ... 71
Wool fat
Dermatological ... 71
Extemporaneously Compounded Preparations ... 292

- X -

Xalkori ... 162
Xanthan ... 292
Xarelto ... 37
Xgeva ... 117

Xifaxan....9
XMET Maxamum....296
Xolair....217
XP Maxamum....297
Xylocaine....125
Xylocaine Viscous....125
Xylometazoline hydrochloride....269
Xyntha....35

- Y -

Yellow jacket wasp venom....265
Yervoy....247

- Z -

Zanamivir....112
Zarontin....131
Zavedos....153
Zavicefta....91
Zeffix....108
Zejula....156
Zematop....73
Zetlam....108
Ziagen....106
Zidovudine....106
Zidovudine with lamivudine....106
Ziextenzo....40
Zimybe....54
Zinc
  Alimentary....25
  Dermatological....69
Zinc and castor oil....70
Zinc chloride....25
Zinc oxide....292
Zinc sulphate....25
Zinc with wool fat....70
Zincaps....25
Zinforo....92
Ziprasidone....137
Zista....265
Zithromax....92
Zo-Rub HP....126
Zo-Rub Osteo....121
Zoladex....83
Zoledronic acid
  Hormone Preparations....80
  Musculoskeletal....116
Zoledronic Acid Injection Mylan....80
Zoledronic acid Viatris
  Hormone Preparations....80
  Musculoskeletal....116
Zopiclone....144
Zopiclone Actavis....144
Zostrix....121
Zostrix HP....126
Zuclopenthixol acetate....137
Zuclopenthixol decanoate....140
Zuclopenthixol hydrochloride....137
Zusdone....137
Zyban....149
Zypine....137
Zypine ODT....137
Zyprexa Relprevv....138
Zytiga....175
Zyvox....97